

# Growth of Solutions of Linear Difference Equations with Meromorphic Coefficients in Terms of Iterated $p - \phi$ Order

Anirban Bandyopadhyay<sup>1</sup>, Chinmay Ghosh<sup>2</sup>, Sanjib Kumar Datta<sup>3</sup>

<sup>1</sup>Gopalpur Primary School

Murshidabad, 742304

West Bengal, India

anirbanbanerjee010@gmail.com

<sup>2</sup>Department of Mathematics

Kazi Nazrul University

Nazrul Road, P.O.- Kalla C.H.

Asansol-713340, West Bengal, India

chinmayarp@gmail.com

<sup>2</sup>Department of Mathematics

University of Kalyani

P.O.-Kalyani, Dist-Nadia, PIN-741235,

West Bengal, India

sanjibdatta05@gmail.com

## Abstract

In this article we have studied complex linear homogeneous difference equations where the coefficients are meromorphic functions, having finite iterated  $p - \phi$  order. We have made some estimations on the growth of its nontrivial solutions. Also we have extended some of the previous results of Zhou-Zheng (2017), Belaïdi-Benkarouba (2019) and Belaïdi-Bellaama (2021).

**AMS Subject Classification** (2010) : 30D35, 39A06, 39B32.

**Keywords and phrases:** linear difference equation, meromorphic solution, iterated  $p - \phi$  order, iterated  $p - \phi$  type, iterated  $p - \phi$  exponent of convergence.

# 1 Introduction

Consider the complex difference equations

$$A_k(z)f(z + c_k) + A_{k-1}(z)f(z + c_{k-1}) + \cdots + A_1(z)f(z + c_1) + A_0(z)f(z) = 0 \quad (1)$$

and

$$A_k(z)f(z + c_k) + A_{k-1}(z)f(z + c_{k-1}) + \cdots + A_1(z)f(z + c_1) + A_0(z)f(z) = A(z) \quad (2)$$

where the coefficients  $A_0, A_1, \dots, A_k$  ( $A_k \neq 0$ ) and  $A$  ( $A \neq 0$ ) are meromorphic and  $c_k, c_{k-1}, \dots, c_1 \in \mathbb{C}$ , the set of finite complex numbers, are all distinct. Recently, study on (1) and (2) have become a topic of great interests from the perspective of Nevanlinna's theory [8, 6]. In 2006 Halburd-Korhonen [7] and in 2008 Chiang-Feng [4] independently established some results which are difference analogues of the classical logarithmic derivative lemma. After that, using those results, many authors ([11], [15], [13], [12] etc.) studied the growth properties of the solutions of linear difference equations. In 2017 and in 2019 respectively Zhou, Zheng [16] and Belaïdi, Benkarouba [1] investigated higher order linear difference equations and proved some results estimating the growth of the solutions of those equations using iterated order [10] and iterated type [3]. Again in 2021, Belaïdi and Bellaama [2] used iterated lower order [9] and iterated lower type [9] to improve some of the previous theorems.

In 2009, Chyzhykov et. al. [5] introduced  $\phi$ -order for a meromorphic function in the unit disc. Also in 2014, Shen et. al. [14] developed the concept of  $\phi$ -order of a meromorphic function in whole complex plane and applied it to the second order linear differential equation. The  $\phi$ -order is based on the function  $\phi$  which is defined as:  $\phi : [0, \infty) \rightarrow (0, \infty)$  is a nondecreasing, unbounded function which satisfies following two conditions:

- i) There exists  $k_\phi \in \mathbb{N}$ , such that  $\lim_{r \rightarrow \infty} \frac{\log_{p+1} r}{\log \phi(r)} = 0$  for all  $p \geq k_\phi$ ;
- ii)  $\lim_{r \rightarrow \infty} \frac{\log \phi(\alpha r)}{\log \phi(r)} = 1$ , for  $\alpha > 1$ .

In this paper, we first introduce the iterated  $p - \phi$  order, the iterated  $p - \phi$  type and the iterated  $p - \phi$  exponent of convergence of a meromorphic function. Since the concept of iterated  $p - \phi$  order is more generalisation than that of iterated order, we are motivated to improvise some of the results of Zhou-Zheng [16], Belaïdi-Benkarouba [1] and Belaïdi-Bellaama [2]. Here we obtain our main results for (1) or (2), when there may or may not exist exactly one coefficient having maximal iterated  $p - \phi$  order (or iterated lower  $p - \phi$  order).

Let us define for all  $r > 0$ ,

$$\log^+ r = \max(0, \log r).$$

We denote for all  $r \in \mathbb{R}$ ,

$$\exp_1 r = e^r, \quad \exp_{n+1} r = \exp(\exp_n r), \quad n \in \mathbb{N}.$$

Also we denote for all sufficiently large values of  $r$ ,

$$\log_1 r = \log^+ r, \quad \log_{n+1} r = \log^+(\log_n r), \quad n \in \mathbb{N}.$$

Further we denote

$$\exp_0 r = \log_0 r = r, \quad \exp_{-1} r = \log^+ r, \quad \log_{-1} r = e^r.$$

Moreover, in this paper, the linear measure of a set  $E \subset (0, +\infty)$  is defined by

$$m(E) = \int_0^{+\infty} \chi_E dt$$

and the logarithmic measure of a set  $F \subset (1, +\infty)$  is defined by

$$lm(F) = \int_1^{+\infty} \frac{\chi_F}{t} dt,$$

where  $\chi_G(t)$  is the characteristics function on the set  $G$ .

## 2 Basic definitions and lemmas

First we give the following definitions to indicate the rate of growth of meromorphic functions.

**Definition 1 (Iterated  $p - \phi$  order)** Let  $p \in \mathbb{N}$ . Then the iterated  $p - \phi$  order,  $\rho_p(f, \phi)$  of a meromorphic function  $f(z)$  is defined by

$$\rho_p(f, \phi) = \limsup_{r \rightarrow \infty} \frac{\log_p T(r, f)}{\log \phi(r)},$$

and when  $f$  is an entire function, we define

$$\rho_p(f, \phi) = \limsup_{r \rightarrow \infty} \frac{\log_{p+1} M_f(r)}{\log \phi(r)}.$$

**Definition 2 (Finiteness degree of the  $\phi$  order)** The finiteness degree of the  $\phi$  order of a meromorphic function  $f(z)$  is defined by

$$i_\rho(f, \phi) = \begin{cases} 0, & \text{if } \rho_1(f, \phi) = \infty, \\ \min\{j \in \mathbb{N} : \rho_j(f, \phi) < \infty\}, & \text{if there exist some } j \in \mathbb{N} \text{ with} \\ & \rho_j(f, \phi) < \infty \text{ exists,} \\ \infty, & \text{if } \rho_j(f, \phi) = \infty \text{ for all } j \in \mathbb{N}. \end{cases}$$

**Definition 3 (Iterated lower  $p - \phi$  order)** Let  $p \in \mathbb{N}$ . Then the iterated lower  $p - \phi$  order,  $\mu_p(f, \phi)$  of a meromorphic function  $f(z)$  is defined by

$$\mu_p(f, \phi) = \liminf_{r \rightarrow \infty} \frac{\log_p T(r, f)}{\log \phi(r)},$$

and when  $f$  is an entire function, we define

$$\mu_p(f, \phi) = \liminf_{r \rightarrow \infty} \frac{\log_{p+1} M_f(r)}{\log \phi(r)}.$$

**Remark 4** Let  $f$  be a meromorphic function with  $i_p(f, \phi) = p$  and the iterated  $p - \phi$  order of  $f$  be  $\rho_p(f, \phi)$ . Then it follows from the definition that, for any  $\epsilon > 0$ ,

$$T(r, f) = O\left(\exp_{p-1}\{\phi(r)\}^{\rho_p(f, \phi) + \epsilon}\right).$$

**Remark 5** Let  $f$  be a meromorphic function such that,  $T(r, f) \leq \exp_p\{\phi(r)\}^\alpha$  for some  $\alpha > 0$  and  $p \in \mathbb{N}$ , then we can say

$$i_p(f, \phi) \leq p + 1 \text{ and } \rho_{p+1}(f, \phi) \leq \alpha.$$

**Definition 6 (Iterated  $p - \phi$  type)** Let  $p \in \mathbb{N}$ . Then the iterated  $p - \phi$  type,  $\tau_p(f, \phi)$  of a meromorphic function  $f$  with nonzero finite iterated  $p - \phi$  order  $\rho_p(f, \phi)$  is defined by

$$\tau_p(f, \phi) = \limsup_{r \rightarrow \infty} \frac{\log_{p-1} T(r, f)}{\{\phi(r)\}^{\rho_p(f, \phi)}},$$

and when  $f$  is an entire function, then for  $p \geq 2$  we define

$$\tau_p(f, \phi) = \limsup_{r \rightarrow \infty} \frac{\log_p M_f(r)}{\{\phi(r)\}^{\rho_p(f, \phi)}}.$$

**Definition 7 (Iterated lower  $p - \phi$  type)** Let  $p \in \mathbb{N}$ . Then the iterated lower  $p - \phi$  type,  $t_p(f, \phi)$  of a meromorphic function  $f$  having non-zero finite iterated lower  $p - \phi$  order  $\mu_p(f, \phi)$  is defined by

$$t_p(f, \phi) = \liminf_{r \rightarrow \infty} \frac{\log_{p-1} T(r, f)}{\{\phi(r)\}^{\mu_p(f, \phi)}},$$

and when  $f$  is an entire function, then for  $p \geq 2$  we define

$$t_p(f, \phi) = \liminf_{r \rightarrow \infty} \frac{\log_p M_f(r)}{\{\phi(r)\}^{\mu_p(f, \phi)}}.$$

**Remark 8** Let  $f$  be a meromorphic function with  $i_p(f, \phi) = p$ . Also let the iterated  $p - \phi$  order and the iterated  $p - \phi$  type of  $f$  be  $\rho_p(f, \phi)$  and  $\tau_p(f, \phi)$  respectively. Then it follows from the definition that, for any  $\epsilon > 0$ ,

$$T(r, f) = O\left(\exp_{p-1}\left[\{\tau_p(f, \phi) + \epsilon\} \{\phi(r)\}^{\rho_p(f, \phi)}\right]\right).$$

**Definition 9 (Iterated  $p - \phi$  exponent of convergence)** Let  $p \in \mathbb{N}$ . Then the iterated  $p - \phi$  exponent of convergence of the sequence of poles of a meromorphic function  $f$  is defined by

$$\lambda_p\left(\frac{1}{f}, \phi\right) = \limsup_{r \rightarrow \infty} \frac{\log_p N(r, f)}{\log_q \{\phi(r)\}}.$$

Now we give some lemmas which will use to prove our main results.

**Lemma 10** [7] Suppose  $z_1 \in \mathbb{C}$ ,  $\delta < 1$  and  $\varepsilon > 0$ . Then for any nonconstant meromorphic function  $f$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) < \infty$  such that

$$m\left(r, \frac{f(z+z_1)}{f(z)}\right) = o\left(\frac{\{T(r+|z_1|, f)\}^{1+\varepsilon}}{r^\delta}\right),$$

for all  $r \notin F$ .

**Lemma 11** [6] Let  $z_1, z_2$  be nonzero complex constants. Then for any nonconstant meromorphic function  $f$ , we have

$$(1+o(1))T(r-|z_1|, f) \leq T(r, f(z+z_1)) \leq (1+o(1))T(r+|z_1|, f),$$

for sufficiently large values of  $r$ .

Consequently we have,  $\rho_p(f(z+z_1), \phi) = \rho_p(f, \phi)$  and  $\mu_p(f(z+z_1), \phi) = \mu_p(f, \phi)$ .

**Lemma 12** [7] Suppose  $z_1, z_2 \in \mathbb{C}$  ( $z_1 \neq z_2$ ),  $\delta < 1$ ,  $\varepsilon > 0$ . Then, for any nonconstant meromorphic function  $f$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) < \infty$  such that

$$m\left(r, \frac{f(z+z_1)}{f(z+z_2)}\right) = o\left(\frac{\{T(r+|z_1-z_2|+|z_2|, f)\}^{1+\varepsilon}}{r^\delta}\right),$$

for all  $r \notin F$ .

**Lemma 13** Let  $f$  be a nonconstant meromorphic function with nonzero finite iterated  $p - \phi$  order,  $\rho_p(f, \phi)$ . Then for any given  $\varepsilon > 0$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that

$$\log_{p-1} T(r, f) > \{\phi(r)\}^{\rho_p(f, \phi) - \varepsilon},$$

for  $r \in F$ .

**Proof.** By the definition of iterated  $p - \phi$  order, there exists an increasing sequence  $\{r_n\}$  ( $r_n \rightarrow \infty$ ) satisfying  $(1 + \frac{1}{n})r_n < r_{n+1}$  and

$$\lim_{r_n \rightarrow \infty} \frac{\log_p T(r_n, f)}{\log \phi(r_n)} = \rho_p(f, \phi).$$

Then for given  $\varepsilon > 0$  there exists an integer  $N_0$  such that for all  $n > N_0$ ,

$$\log_{p-1} T(r_n, f) > \{\phi(r_n)\}^{\rho_p(f, \phi) - \frac{\varepsilon}{2}}$$

Then for  $r \in [r_n, (1 + \frac{1}{n})r_n]$  and for  $n > N_0$

$$\begin{aligned} \log_{p-1} T(r, f) &> \log_{p-1} T(r_n, f) > \{\phi(r_n)\}^{\rho_p(f, \phi) - \frac{\varepsilon}{2}} \\ &> \left\{ \phi\left(\frac{n}{n+1}r\right) \right\}^{\rho_p(f, \phi) - \frac{\varepsilon}{2}} > \{\phi(r)\}^{\rho_p(f, \phi) - \varepsilon}. \end{aligned}$$

Set  $F = \cup_{n=N_0}^{\infty} [r_n, (1 + \frac{1}{n})r_n]$ , then

$$\begin{aligned} lm(F) &= \sum_{n=N_0}^{\infty} lm \left( \left[ r_n, (1 + \frac{1}{n})r_n \right] \right) = \sum_{n=N_0}^{\infty} \int_1^{+\infty} \frac{\chi_{[r_n, (1 + \frac{1}{n})r_n]}}{t} dt \\ &= \sum_{n=N_0}^{\infty} \int_{r_n}^{(1 + \frac{1}{n})r_n} \frac{1}{t} dt = \sum_{n=N_0}^{\infty} \log(1 + \frac{1}{n}) = +\infty. \end{aligned}$$

Thus Lemma 13 is proved. ■

**Lemma 14** *Let  $f$  be a nonconstant meromorphic function with nonzero finite iterated  $p - \phi$  order,  $\rho_p(f, \phi)$  and nonzero finite iterated  $p - \phi$  type,  $\tau_p(f, \phi)$ . Then for any given  $\varepsilon > 0$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that*

$$\log_{p-1} T(r, f) > (\tau_p(f, \phi) - \varepsilon) \{\phi(r)\}^{\rho_p(f, \phi)}.$$

for  $r \in F$ .

**Proof.** By the definition of iterated  $p - \phi$  order and iterated  $p - \phi$  type, there exists an increasing sequence  $\{r_n\}$  ( $r_n \rightarrow \infty$ ) satisfying  $(1 + \frac{1}{n})r_n < r_{n+1}$  and

$$\limsup_{r \rightarrow \infty} \frac{\log_{p-1} T(r_n, f)}{\{\phi(r_n)\}^{\rho_p(f, \phi)}} = \tau_p(f, \phi).$$

Then for given  $\varepsilon > 0$  there exists an integer  $N_0$  such that for all  $n > N_0$ ,

$$\log_{p-1} T(r_n, f) > \left( \tau_p(f, \phi) - \frac{\varepsilon}{2} \right) \{\phi(r_n)\}^{\rho_p(f, \phi)}$$

Then for  $r \in [r_n, (1 + \frac{1}{n})r_n]$  and for  $n > N_0$

$$\begin{aligned} \log_{p-1} T(r, f) &> \log_{p-1} T(r_n, f) > \left( \tau_p(f, \phi) - \frac{\varepsilon}{2} \right) \{\phi(r_n)\}^{\rho_p(f, \phi)} \\ &> \left( \tau_p(f, \phi) - \frac{\varepsilon}{2} \right) \left\{ \phi\left(\frac{n}{n+1}r\right) \right\}^{\rho_p(f, \phi)} > (\tau_p(f, \phi) - \varepsilon) \{\phi(r)\}^{\rho_p(f, \phi)}. \end{aligned}$$

Set  $F = \cup_{n=N_0}^{\infty} [r_n, (1 + \frac{1}{n})r_n]$ , then

$$\begin{aligned} lm(F) &= \sum_{n=N_0}^{\infty} lm \left( \left[ r_n, (1 + \frac{1}{n})r_n \right] \right) = \sum_{n=N_0}^{\infty} \int_1^{+\infty} \frac{\chi_{[r_n, (1 + \frac{1}{n})r_n]}}{t} dt \\ &= \sum_{n=N_0}^{\infty} \int_{r_n}^{(1 + \frac{1}{n})r_n} \frac{1}{t} dt = \sum_{n=N_0}^{\infty} \log(1 + \frac{1}{n}) = +\infty. \end{aligned}$$

Thus Lemma 14 is proved. ■

**Lemma 15** *Let  $f$  be a nonconstant meromorphic function with nonzero finite iterated lower  $p - \phi$  order,  $\mu_p(f, \phi)$ . Then for any given  $\varepsilon > 0$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that*

$$\log_{p-1} T(r, f) < \{\phi(r)\}^{\mu_p(f, \phi) + \varepsilon},$$

for  $r \in F$ .

**Proof.** Similar as Lemma 13. ■

**Lemma 16** *Let  $f$  be a nonconstant meromorphic function with nonzero finite iterated lower  $p - \phi$  order,  $\mu_p(f, \phi)$  and nonzero finite iterated lower  $p - \phi$  type,  $t_p(f, \phi)$ . Then for any given  $\varepsilon > 0$ , there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that*

$$\log_{p-1} T(r, f) < (t_p(f, \phi) + \varepsilon) \{\phi(r)\}^{\mu_p(f, \phi)}.$$

for  $r \in F$ .

**Proof.** Similar as Lemma 14. ■

**Lemma 17** *Let  $f$  be a nonconstant meromorphic function with  $i_p(f, \phi) = p$  and  $\rho_p(f, \phi) = \rho$ . Also let  $z_1, z_2 \in \mathbb{C}$  ( $z_1 \neq z_2$ ). Then for any  $\varepsilon > 0$  sufficiently small, we have*

$$m\left(r, \frac{f(z + z_1)}{f(z + z_2)}\right) = O\left(\exp_{p-1}\{\phi(r)\}^{\rho + \varepsilon}\right).$$

**Proof.** By using similar arguments as in the proof of ([1], Lemma 2.9) we get for some  $M > 0$ ,

$$m\left(r, \frac{f(z + z_1)}{f(z + z_2)}\right) \leq M.T(3r, f),$$

Since  $f$  has finite iterated  $p - \phi$  order,  $\rho_p(f, \phi) = \rho$ , we have for any  $\varepsilon$

$$T(r, f) \leq \exp_{p-1}\{\phi(r)\}^{\rho + \frac{\varepsilon}{2}}$$

Combining above two results we get,

$$\begin{aligned} m\left(r, \frac{f(z + z_1)}{f(z + z_2)}\right) &\leq M.T(3r, f) \leq M.\exp_{p-1}\{\phi(3r)\}^{\rho + \frac{\varepsilon}{2}} \\ &\leq M.\exp_{p-1}\{\phi(r)\}^{\rho + \varepsilon}. \end{aligned}$$

This completes the proof. ■

**Lemma 18** *Let  $f$  be a nonconstant meromorphic function with  $i_p(f, \phi) = p$  and  $\mu_p(f, \phi) = \mu$ . Also let  $z_1, z_2 \in \mathbb{C}$  ( $z_1 \neq z_2$ ). Then for any  $\varepsilon > 0$  sufficiently small, there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that*

$$m\left(r, \frac{f(z + z_1)}{f(z + z_2)}\right) = O\left(\exp_{p-1}\{\phi(r)\}^{\mu + \varepsilon}\right),$$

for  $r \in F$ .

**Proof.** As in the previous lemma for some  $M > 0$ , we get

$$m\left(r, \frac{f(z+z_1)}{f(z+z_2)}\right) \leq M.T(3r, f),$$

Since  $f$  has finite iterated lower  $p - \phi$  order,  $\mu_p(f, \phi) = \mu$ , we have from Lemma 15, for any  $\varepsilon > 0$  there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that

$$T(r, f) \leq \exp_{p-1}\{\phi(r)\}^{\mu+\frac{\varepsilon}{2}},$$

for  $r \in F$ .

Combining above two results we get,

$$\begin{aligned} m\left(r, \frac{f(z+z_1)}{f(z+z_2)}\right) &\leq M.T(3r, f) \leq M.\exp_{p-1}\{\phi(3r)\}^{\mu+\frac{\varepsilon}{2}} \\ &\leq M.\exp_{p-1}\{\phi(r)\}^{\mu+\varepsilon}, \end{aligned}$$

for  $r \in F$ .

This completes the proof. ■

**Lemma 19** *Let  $f$  be a nonconstant meromorphic function with  $i_p(f, \phi) = p$  and  $\rho_p(f, \phi) = \rho$ . Suppose  $z_1 \in \mathbb{C}$  ( $z_1 \neq 0$ ) and  $\varepsilon > 0$  be any given constant sufficiently small. Then there exists a set  $F \subset (1, +\infty)$  with  $lm(F) < \infty$  such that*

$$\left|\frac{f(z+z_1)}{f(z)}\right| \leq \exp_p\{\phi(r)\}^{\rho+\varepsilon},$$

for  $r \notin F \cup [0, 1]$ .

**Proof.** By using similar arguments as in the proof of ([1], Lemma 2.3), there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that for some  $B > 0$ ,  $\lambda > 1$ , we have

$$\left|\log\left|\frac{f(z+z_1)}{f(z)}\right|\right| \leq B\left\{T(\lambda r, f)\frac{\log^\lambda r}{r}\log T(\lambda r, f)\right\},$$

for  $r \notin F \cup [0, 1]$ . Again since,  $f$  has finite iterated  $p - \phi$  order,  $\rho_p(f, \phi) = \rho$ , we have for any given  $\varepsilon > 0$ ,

$$T(r, f) \leq \exp_{p-1}\{\phi(r)\}^{\rho+\frac{\varepsilon}{2}}$$

Combining above two results we get

$$\begin{aligned} \left|\log\left|\frac{f(z+z_1)}{f(z)}\right|\right| &\leq B\exp_{p-1}\{\phi(\lambda r)\}^{\rho+\frac{\varepsilon}{2}}\frac{\log^\lambda r}{r}\exp_{p-2}\{\phi(\lambda r)\}^{\rho+\frac{\varepsilon}{2}} \\ &\leq \exp_{p-1}\{\phi(r)\}^{\rho+\varepsilon}, \text{ for } r \notin F \cup [0, 1]. \end{aligned}$$

$$\text{Therefore, } \left|\frac{f(z+z_1)}{f(z)}\right| \leq \exp_p\{\phi(r)\}^{\rho+\varepsilon}, \text{ for } r \notin F \cup [0, 1].$$

Thus Lemma 19 is proved. ■



**Lemma 20** *Let  $f$  be a nonconstant meromorphic function with  $i_\rho(f, \phi) = p$  and  $\rho_p(f, \phi) = \rho$ . Suppose  $z_1, z_2 \in \mathbb{C}$  ( $z_1 \neq z_2$ ) and  $\varepsilon > 0$  be any given constant sufficiently small. Then there exists a set  $F \subset (1, +\infty)$  with  $lm(F) < \infty$  such that*

$$\left| \frac{f(z + z_1)}{f(z + z_2)} \right| \leq \exp_p \{ \phi(r) \}^{\rho + \varepsilon},$$

for  $r \notin F \cup [0, 1]$ .

**Proof.** We can write

$$\left| \frac{f(z + z_1)}{f(z + z_2)} \right| = \left| \frac{f(z + z_2 + z_1 - z_2)}{f(z + z_2)} \right|.$$

By Lemma 11, we have  $\rho_p(f(z + z_2), \phi) = \rho_p(f, \phi) = \rho$ . Again since  $z_1 - z_2 \neq 0$ , by Lemma 19, there exists a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that for given any  $\varepsilon > 0$ , we get

$$\left| \frac{f(z + z_2 + z_1 - z_2)}{f(z + z_2)} \right| \leq \exp_p \{ \phi(r) \}^{\rho + \varepsilon}, \text{ for } r \notin F \cup [0, 1].$$

Thus Lemma 20 is proved. ■

### 3 Main Results

We now state and prove our main theorems. In Theorem 21 and Theorem 22 we consider homogeneous linear difference equation (1) with exactly one coefficient having maximal iterated  $p - \phi$  order and iterated lower  $p - \phi$  order respectively.

**Theorem 21** *Let  $A_j(z)$  ( $j = 0, 1, \dots, k$ ) be meromorphic functions and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n - 1\}$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) such*

that

$$\lambda_p \left( \frac{1}{A_m}, \phi \right) < \rho_p(A_m, \phi) < +\infty,$$

and  $\max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} < \rho_p(A_m, \phi)$

then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A_m, \phi)$ .

**Proof.** Let  $f$  be a meromorphic solution of (1) and put  $c_0 = 0$ . We divide (1) by  $f(z + c_m)$  and we get

$$-A_m(z) = \sum_{j=0, j \neq m}^k A_j(z) \frac{f(z + c_j)}{f(z + c_m)}. \quad (3)$$

Now from Lemma 12 for any  $\varepsilon > 0$  we get,

$$m\left(r, \frac{f(z+c_j)}{f(z+c_m)}\right) = o\left(\frac{\{T(r+3C, f)\}^{1+\varepsilon}}{r^\delta}\right), j = 0, 1, \dots, k, j \neq m,$$

for  $r \notin F_1$ , where  $lm(F_1) < +\infty$  and  $C = \max\{|c_j| : j = 0, 1, \dots, k\}$ .

Using the above result, from (3) we get,

$$\begin{aligned} & T(r, A_m) \\ &= m(r, A_m) + N(r, A_m) \\ &\leq \sum_{j=0, j \neq m}^k m(r, A_j) + \sum_{j=0, j \neq m}^k m\left(r, \frac{f(z+c_j)}{f(z+c_m)}\right) + N(r, A_m) + O(1) \quad (4) \\ &\leq \sum_{j=0, j \neq m}^k T(r, A_j) + o\left(\frac{\{T(r+3C, f)\}^{1+\varepsilon}}{r^\delta}\right) + N(r, A_m) + O(1) \\ &\leq \sum_{j=0, j \neq m}^k T(r, A_j) + \{T(2r, f)\}^2 + N(r, A_m) + O(1) \quad (5) \end{aligned}$$

for  $r \notin F_1$ .

We denote,  $\rho = \rho_p(A_m, \phi)$ ,  $\rho_1 = \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k; j \neq m\}$  and  $\lambda = \lambda_p\left(\frac{1}{A_m}, \phi\right)$ .

By Lemma 13, for that  $\varepsilon$  there exists a set  $F_2 \subset (1, +\infty)$  with  $lm(F_2) = +\infty$  such that

$$T(r, A_m) > \exp_{p-1}\{\phi(r)\}^{\rho-\varepsilon} \quad (6)$$

for  $r \in F_2$ . Also for  $j \neq m$ , for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1}\{\phi(r)\}^{\rho_1+\varepsilon} \quad (7)$$

for sufficiently large  $r$ .

Again by the definition of  $\lambda_p\left(\frac{1}{A_m}, \phi\right)$ , we have for that  $\varepsilon$  and for sufficiently large  $r$

$$N(r, A_m) \leq \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon}. \quad (8)$$

Now, using the relations (6)-(8) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min\{\rho - \rho_1, \rho - \lambda\}$ , we have from (5)

$$\begin{aligned} \exp_{p-1}\{\phi(r)\}^{\rho-\varepsilon} &\leq k \cdot \exp_{p-1}\{\phi(r)\}^{\rho_1+\varepsilon} + \{T(2r, f)\}^2 \\ &\quad + \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon} + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 \geq O\left(\exp_{p-1}\{\phi(r)\}^{\rho-\varepsilon}\right) \end{aligned}$$

for sufficiently large  $r$  and  $r \in F_2 \setminus F_1$ .

Which implies,  $i_p(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$ . ■

**Theorem 22** Let  $A_j(z)$  ( $j = 0, 1, \dots, k$ ) be meromorphic functions and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\}$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) such

that

$$\lambda_p\left(\frac{1}{A_m}, \phi\right) < \mu_p(A_m, \phi) < +\infty,$$

$$\text{and } \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} < \mu_p(A_m, \phi)$$

then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu_p(A_m, \phi)$ .

**Proof.** Let  $\varepsilon > 0$  be any arbitrary constant. As in the previous theorem, from (5), we get,

$$T(r, A_m) \leq \sum_{j=0, j \neq m}^k T(r, A_j) + \{T(2r, f)\}^2 + N(r, A_m) + O(1) \quad (9)$$

for  $r \notin F_1$ , where  $lm(F_1) < +\infty$ .

We denote,  $\mu = \mu_p(A_m, \phi)$ ,  $\rho_1 = \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k; j \neq m\}$  and  $\lambda = \lambda_p\left(\frac{1}{A_m}, \phi\right)$ .

Then for that  $\varepsilon$  and, we have

$$T(r, A_m) > \exp_{p-1}\{\phi(r)\}^{\mu-\varepsilon} \quad (10)$$

for sufficiently large  $r$ . Also for  $j \neq m$ , for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1}\{\phi(r)\}^{\rho_1+\varepsilon} \quad (11)$$

for sufficiently large  $r$ .

Again by the definition of  $\lambda_p\left(\frac{1}{A_m}, \phi\right)$ , we have for that  $\varepsilon$  and for sufficiently large  $r$

$$N(r, A_m) \leq \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon}. \quad (12)$$

Now, using the relations (10)-(12) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min\{\mu - \rho_1, \mu - \lambda\}$ , we have from (9)

$$\begin{aligned} \exp_{p-1}\{\phi(r)\}^{\mu-\varepsilon} &\leq k \cdot \exp_{p-1}\{\phi(r)\}^{\rho_1+\varepsilon} + \{T(2r, f)\}^2 \\ &\quad + \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon} + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 \geq O\left(\exp_{p-1}\{\phi(r)\}^{\mu-\varepsilon}\right) \end{aligned}$$

for  $r \notin F_1$ .

Which implies,  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu = \mu_p(A_m, \phi)$ . ■

In Theorem 23 and Theorem 24 we consider nonhomogeneous linear difference equation (2) which may have more than one coefficient with the maximal iterated  $p - \phi$  order and iterated lower  $p - \phi$  order respectively. For these we need to consider the iterated  $p - \phi$  type or iterated lower  $p - \phi$  type among the coefficients having maximal iterated  $p - \phi$  order.

**Theorem 23** Let  $A_j(z) (j = 0, 1, \dots, k)$  and  $A(z)$  be meromorphic functions and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\}$ . If there exists an  $A_m(z) (0 \leq m \leq k)$  such that

$$\lambda_p\left(\frac{1}{A_m}, \phi\right) < \rho_p(A_m, \phi) < \infty,$$

$$\max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} \leq \rho_p(A_m, \phi)$$

$$\text{and } \max\{\tau_p(A_j, \phi) : \rho_p(A_j, \phi) = \rho_p(A_m, \phi), j = 0, 1, \dots, k, j \neq m\} < \tau_p(A_m, \phi) < \infty,$$

then the following cases can be arise.

i) If  $\rho_p(A, \phi) < \rho_p(A_m, \phi)$ , or  $\rho_p(A, \phi) = \rho_p(A_m, \phi)$  and  $\tau_p(A, \phi) \neq \tau_p(A_m, \phi)$ , then every solution  $f(z)$  ( $f \neq 0$ ) of (2) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A_m, \phi)$ .

ii) If  $\rho_p(A, \phi) > \rho_p(A_m, \phi)$  then every solution  $f(z)$  ( $f \neq 0$ ) of (2) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A, \phi)$ .

**Proof.** Let  $f$  be a meromorphic solution of (2) and put  $c_0 = 0$ . We divide (2) by  $f(z + c_m)$  and we get

$$-A_m(z) = \sum_{j=0, j \neq m}^k A_j(z) \frac{f(z + c_j)}{f(z + c_m)} - \frac{A(z)}{f(z + c_m)}. \quad (13)$$

Now from Lemma 12 and Lemma 11, for any  $\varepsilon > 0$  we get,

$$m\left(r, \frac{f(z + c_j)}{f(z + c_m)}\right) \leq o\left(\frac{\{T(r + 3C, f)\}^{1+\varepsilon}}{r^\delta}\right), j = 0, 1, \dots, k, j \neq m,$$

$$\begin{aligned} \text{and } m\left(r, \frac{1}{f(z + c_m)}\right) &\leq T\left(r, \frac{1}{f(z + c_m)}\right) \\ &= T(r, f(z + c_m)) + O(1) \\ &\leq (1 + O(1))T(r + C, f), \end{aligned}$$

for  $r \notin F_1$ , where  $lm(F_1) < +\infty$  and  $C = \max\{|c_j| : j = 0, 1, \dots, k\}$ .

Using above results in (13) we get,

$$\begin{aligned}
& T(r, A_m) \\
&= m(r, A_m) + N(r, A_m) \\
&\leq \sum_{j=0, j \neq m}^k m(r, A_j) + \sum_{j=0, j \neq m}^k m\left(r, \frac{f(z+c_j)}{f(z+c_m)}\right) + m(r, A) \\
&\quad + m\left(r, \frac{1}{f(z+c_m)}\right) + N(r, A_m) + O(1) \\
&\leq \sum_{j=0, j \neq m}^k T(r, A_j) + o\left(\frac{\{T(r+3C, f)\}^{1+\varepsilon}}{r^\delta}\right) + T(r, A) \\
&\quad + (1+O(1))T(r+C, f) + N(r, A_m) + O(1) \\
&\leq \sum_{j=0, j \neq m}^k T(r, A_j) + 3\{T(2r, f)\}^2 + T(r, A) + N(r, A_m) + O(1) \quad (14)
\end{aligned}$$

for sufficiently large  $r$  and  $r \notin F_1$ .

Now denote,  $\rho = \rho_p(A_m, \phi)$ ,  $\rho_1 = \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, \rho_p(A_j, \phi) < \rho\}$ ,  $\tau = \tau_p(A_m, \phi)$ ,  $\tau_1 = \max\{\tau_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m, \rho_p(A_j, \phi) = \rho\}$  and  $\lambda = \lambda_p\left(\frac{1}{A_m}, \phi\right)$ .

Now from lemma 14 for that  $\varepsilon$  there exists a set  $F_2 \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that

$$T(r, A_m) > \exp_{p-1}[(\tau - \varepsilon)\{\phi(r)\}^\rho] \quad (15)$$

for sufficiently large  $r$  and  $r \in F_2$ .

Again if  $\rho_p(A_j, \phi) < \rho$ , then for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1}\left[\{\phi(r)\}^{\rho_1+\varepsilon}\right] \quad (16)$$

for sufficiently large  $r$ .

If  $\rho_p(A_j, \phi) = \rho$ ,  $j \neq m$ , then for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1}[(\tau_1 + \varepsilon)\{\phi(r)\}^\rho] \quad (17)$$

for sufficiently large  $r$ .

Also by the definition of  $\lambda_p\left(\frac{1}{A_m}, \phi\right)$ , we have for the above  $\varepsilon$  and for sufficiently large  $r$

$$N(r, A_m) \leq \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon}. \quad (18)$$

(i) Case 1: For the first part of the proof, we take  $\rho_p(A, \phi) < \rho$ , then for that  $\varepsilon$  we have

$$T(r, A) \leq \exp_{p-1}\left[\{\phi(r)\}^{\rho_p(A, \phi)+\varepsilon}\right] \quad (19)$$

for sufficiently large  $r$ .

Now, using the relations (15)-(19) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\rho - \rho_1, \tau - \tau_1, \rho - \lambda, \rho - \rho_p(A, \phi)\}$  we have from (14)

$$\begin{aligned} \exp_{p-1} [(\tau - \varepsilon) \{\phi(r)\}^\rho] &\leq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho_1 + \varepsilon} \right] \right) + O \left( \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\rho] \right) \\ &+ 3 \{T(2r, f)\}^2 + \exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) + \varepsilon} \right] + \exp_{p-1} \left[ \{\phi(r)\}^{\lambda + \varepsilon} \right] + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 \geq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho + \varepsilon} \right] \right), \end{aligned}$$

for sufficiently large  $r$  and  $r \in F_2 \setminus F_1$ .

Which implies,  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$ .

Case 2: Next we suppose that  $\rho_p(A, \phi) = \rho$  and  $\tau_p(A, \phi) < \tau$ , then for the above  $\varepsilon$  we have

$$T(r, A) \leq \exp_{p-1} [\{\tau_p(A, \phi) + \varepsilon\} \{\phi(r)\}^\rho] \quad (20)$$

for sufficiently large  $r$ .

Now, using relations (15)-(18) and (20) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\rho - \rho_1, \tau - \tau_1, \rho - \lambda, \tau - \tau_p(A, \phi)\}$ , we have from (14)

$$\begin{aligned} \exp_{p-1} [(\tau - \varepsilon) \{\phi(r)\}^\rho] &\leq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho_1 + \varepsilon} \right] \right) + O \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\rho] \\ &+ 3 \{T(2r, f)\}^2 + \exp_{p-1} [\{\tau_p(A, \phi) + \varepsilon\} \{\phi(r)\}^\rho] + \exp_{p-1} \left[ \{\phi(r)\}^{\lambda + \varepsilon} \right] + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 > O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho + \varepsilon} \right] \right) \end{aligned}$$

for sufficiently large  $r$  and  $r \in F_2 \setminus F_1$ .

Which implies,  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$ .

Case 3: Next we take,  $\rho_p(A, \phi) = \rho$  and  $\tau_p(A, \phi) > \tau$ , then by Lemma 14, and for the above  $\varepsilon$  we have

$$T(r, A) > \exp_{p-1} [\{\tau_p(A, \phi) - \varepsilon\} \{\phi(r)\}^\rho] \quad (21)$$

for  $r \in F_3$ , where  $lm(F_3) = +\infty$ .

Again by the definition of  $\tau_p(A_m, \phi)$ , we have for the above  $\varepsilon$  and for sufficiently large  $r$

$$T(r, A_m) \leq \exp_{p-1} [(\tau + \varepsilon) \{\phi(r)\}^\rho]. \quad (22)$$

Now from (2) and using Lemma 11 we get

$$\begin{aligned}
A(z) &= \sum_{j=0}^k A_j(z)f(z + c_j) \\
\Rightarrow T(r, A) &\leq \sum_{j=0, j \neq m}^k T(r, A_j) + T(r, A_m) + \sum_{j=0}^k T(r, f(z + c_j)) + O(1) \\
\Rightarrow T(r, A) &\leq \sum_{j=0, j \neq m}^k T(r, A_j) + T(r, A_m) + (k+1)(1+o(1))T(r+C, f) + O(1) \\
\Rightarrow T(r, A) &\leq \sum_{j=0, j \neq m}^k T(r, A_j) + T(r, A_m) + 2(k+1)T(2r, f) + O(1) \quad (23)
\end{aligned}$$

for sufficiently large  $r$

Now, using relations (16), (17), (21), (22) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\rho - \rho_1, \tau - \tau_1, \tau_p(A, \phi) - \tau\}$ , we have from (23)

$$\begin{aligned}
&\exp_{p-1} [(\tau_p(A, \phi) - \varepsilon)(\phi(r))^\rho] \leq O \left[ \exp_{p-1} \left[ \{\phi(r)\}^{\rho_1 + \varepsilon} \right] \right] \\
+ O(\exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\rho]) + \exp_{p-1} [(\tau + \varepsilon) \{\phi(r)\}^\rho] + 2(k+1)T(2r, f) + O(1) \\
\Rightarrow T(2r, f) &\geq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho + \varepsilon} \right] \right)
\end{aligned}$$

for sufficiently large  $r$  and  $r \in F_3$ .

It follows that,  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$ .

(ii) For the second part of the theorem, we take  $\rho_p(A, \phi) > \rho$ . Then from lemma 13 for the above  $\varepsilon$  we have

$$T(r, A) \geq \exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) - \varepsilon} \right] \quad (24)$$

for sufficiently large  $r$  with  $r \in F_4$ , where  $lm(F_4) = +\infty$ .

Again for the above  $\varepsilon$  we have

$$T(r, A_m) \leq \exp_{p-1} \left[ \{\phi(r)\}^{\rho + \varepsilon} \right], \quad (25)$$

for sufficiently large  $r$ .

Now, using relations (16), (17), (24), (25) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\rho - \rho_1, \tau - \tau_1, \rho_p(A, \phi) - \rho\}$ , we have from (23)

$$\begin{aligned}
&\exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) - \varepsilon} \right] \leq O \left[ \exp_{p-1} \left[ \{\phi(r)\}^{\rho_1 + \varepsilon} \right] \right] + O(\exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\rho]) \\
&+ \exp_{p-1} \left[ \{\phi(r)\}^{\rho + \varepsilon} \right] + 2(k+1)T(2r, f) + O(1) \\
\Rightarrow T(2r, f) &\geq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) - \varepsilon} \right] \right)
\end{aligned}$$

for sufficiently large  $r$  and  $r \in F_4$ .

Hence we have  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A, \phi)$ . ■

**Theorem 24** Let  $A_j(z) (j = 0, 1, \dots, k)$  and  $A(z)$  be meromorphic functions and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\}$ . If there exists an  $A_m(z) (0 \leq m \leq k)$  such that

$$\begin{aligned} \lambda_p \left( \frac{1}{A_m}, \phi \right) &< \mu_p(A_m, \phi) < \infty, \\ \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} &\leq \mu_p(A_m, \phi) \\ \text{and } \max\{\tau_p(A_j, \phi) : \rho_p(A_j, \phi) = \mu_p(A_m, \phi), j = 0, 1, \dots, k, j \neq m\} &< t_p(A_m, \phi) < \infty, \end{aligned}$$

then the following cases can be arise.

i) If  $\rho_p(A, \phi) < \mu_p(A_m, \phi)$ , or  $\rho_p(A, \phi) = \mu_p(A_m, \phi)$  and  $\tau_p(A, \phi) < t_p(A_m, \phi)$ , or  $\mu_p(A, \phi) = \mu_p(A_m, \phi)$  and  $t_p(A, \phi) > t_p(A_m, \phi)$ , then every solution  $f(z) (f \neq 0)$  of (2) satisfies  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu_p(A_m, \phi)$ .

ii) If  $\mu_p(A, \phi) > \mu_p(A_m, \phi)$  then every solution  $f(z) (f \neq 0)$  of (2) satisfies  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu_p(A, \phi)$ .

**Proof.** Let  $\varepsilon > 0$  be any arbitrary constant. As in the previous theorem, from (14), we get,

$$T(r, A_m) \leq \sum_{j=0, j \neq m}^k T(r, A_j) + 3 \{T(2r, f)\}^2 + T(r, A) + N(r, A_m) + O(1) \quad (26)$$

for sufficiently large  $r$  and  $r \notin F_1$ .

Now denote,  $\mu = \mu_p(A_m, \phi)$ ,  $\rho_1 = \max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, \rho_p(A_j, \phi) < \mu\}$ ,  $t = t_p(A_m, \phi)$ ,  $\tau_1 = \max\{\tau_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m, \rho_p(A_j, \phi) = \mu\}$  and  $\lambda = \lambda_p\left(\frac{1}{A_m}, \phi\right)$ .

Then by the definition of  $t_p(A_m, \phi)$ , for that  $\varepsilon > 0$  we have

$$T(r, A_m) \geq \exp_{p-1} [(t - \varepsilon) \{\phi(r)\}^\mu] \quad (27)$$

for sufficiently large  $r$ .

Again if  $\rho_p(A_j, \phi) < \mu$ , then for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1} [\{\phi(r)\}^{\mu-\varepsilon}] \quad (28)$$

for sufficiently large  $r$ .

And if  $\rho_p(A_j, \phi) = \rho$ ,  $j \neq m$ , then for that  $\varepsilon$  we have

$$T(r, A_j) \leq \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\mu] \quad (29)$$

for sufficiently large  $r$ .

Also by the definition of  $\lambda_p\left(\frac{1}{A_m}, \phi\right)$ , we have for the above  $\varepsilon$  and for sufficiently large  $r$

$$N(r, A_m) \leq \exp_{p-1} \{\phi(r)\}^{\lambda+\varepsilon}. \quad (30)$$



(i) Case 1: First we take,  $\rho_p(A, \phi) < \mu$ , then for that  $\varepsilon$  we have

$$T(r, A) \leq \exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) + \varepsilon} \right] \quad (31)$$

for sufficiently large  $r$ .

Now, using the relations (27)-(31) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\mu - \rho_1, t - \tau_1, \mu - \lambda, \mu - \rho_p(A, \phi)\}$  we have from (26)

$$\begin{aligned} \exp_{p-1} [(\tau - \varepsilon) \{\phi(r)\}^\mu] &\leq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\mu - \varepsilon} \right] \right) + O \left( \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\mu] \right) \\ &\quad + 3 \{T(2r, f)\}^2 + \exp_{p-1} \left[ \{\phi(r)\}^{\rho_p(A, \phi) + \varepsilon} \right] + \exp_{p-1} \left[ \{\phi(r)\}^{\lambda + \varepsilon} \right] + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 \geq O \left( \exp_{p-1} [\{\phi(r)\}^\mu] \right) \end{aligned}$$

for sufficiently large  $r$  and  $r \notin F_1$ .

Which implies,  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu = \mu_p(A_m, \phi)$ .

Case 2: Next we suppose that  $\rho_p(A, \phi) = \mu$  and  $\tau_p(A, \phi) < t$ , then for the above  $\varepsilon$  we have

$$T(r, A) \leq \exp_{p-1} [\{\tau_p(A, \phi) + \varepsilon\} \{\phi(r)\}^\mu] \quad (32)$$

for sufficiently large  $r$ .

Now, using relations (27)-(30) and (32) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\mu - \rho_1, t - \tau_1, \mu - \lambda, t - \tau_p(A, \phi)\}$ , we have from (26)

$$\begin{aligned} \exp_{p-1} [(\tau - \varepsilon) \{\phi(r)\}^\mu] &\leq O \left( \exp_{p-1} \left[ \{\phi(r)\}^{\mu - \varepsilon} \right] \right) + O \left( \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\mu] \right) \\ &\quad + 3 \{T(2r, f)\}^2 + \exp_{p-1} [\{\tau_p(A, \phi) + \varepsilon\} \{\phi(r)\}^\mu] + \exp_{p-1} \left[ \{\phi(r)\}^{\lambda + \varepsilon} \right] + O(1) \\ &\Rightarrow \{T(2r, f)\}^2 \geq O \left( \exp_{p-1} [(\tau - \varepsilon) \{\phi(r)\}^\mu] \right) \end{aligned}$$

for sufficiently large  $r$  and  $r \notin F_1$ .

Which implies,  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu = \mu_p(A_m, \phi)$ .

Case 3: Next we take,  $\mu_p(A, \phi) = \mu$  and  $t_p(A, \phi) > t$ , then for the above  $\varepsilon$  we have

$$T(r, A) > \exp_{p-1} [\{t_p(A, \phi) - \varepsilon\} \{\phi(r)\}^\mu] \quad (33)$$

for sufficiently large  $r$ .

Again by the definition of  $t_p(A_m, \phi)$ , and using Lemma 16, for the above  $\varepsilon$  we have

$$T(r, A_m) \leq \exp_{p-1} [(t + \varepsilon) \{\phi(r)\}^\mu], \quad (34)$$

for sufficiently large  $r$  with  $r \in F_2$ , where  $lm(F_2) = +\infty$ .

Again as in previous Theorem, from (23) we get

$$\Rightarrow T(r, A) \leq \sum_{j=0, j \neq m}^k T(r, A_j) + T(r, A_m) + 2(k+1)T(2r, f) + O(1) \quad (35)$$

for sufficiently large  $r$

Now, using relations (28), (29), (33), (34) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\mu - \rho_1, t - \tau_1, t_p(A, \phi) - t\}$ , we have from (35)

$$\begin{aligned} & \exp_{p-1} [\{t_p(A, \phi) - \varepsilon\} \{\phi(r)\}^\mu] \leq O \left( \exp_{p-1} [\{\phi(r)\}^{\mu-\varepsilon}] \right) \\ & + O \left( \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\mu] \right) + O \left( \exp_{p-1} [(t + \varepsilon) \{\phi(r)\}^\mu] \right) + (2k + 2)T(2r, f) + O(1) \\ & \Rightarrow T(2r, f) \geq O \left( \exp_{p-1} [\{t_p(A, \phi) - \varepsilon\} \{\phi(r)\}^\mu] \right) \end{aligned}$$

for sufficiently large  $r$  and  $r \in F_2$ .

It follows that,  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu = \mu_p(A_m, \phi)$ .

(ii) For the second part of the theorem, we take  $\mu_p(A, \phi) > \mu$ . Then for the above  $\varepsilon$  we have

$$T(r, A) \geq \exp_{p-1} [\{\phi(r)\}^{\mu_p(A, \phi) - \varepsilon}] \quad (36)$$

Again by using Lemma 15, for the above  $\varepsilon$  we have

$$T(r, A_m) \leq \exp_{p-1} [\{\phi(r)\}^{\mu + \varepsilon}], \quad (37)$$

for sufficiently large  $r$  with  $r \in F_3$ , where  $lm(F_3) = +\infty$ .

Now, using relations (28), (29), (36) and (37) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\mu - \rho_1, t - \tau_1, \mu_p(A, \phi) - \mu\}$ , we have from (35)

$$\begin{aligned} & \exp_{p-1} [\{\phi(r)\}^{\mu_p(A, \phi) - \varepsilon}] \leq O \left( \exp_{p-1} [\{\phi(r)\}^{\mu - \varepsilon}] \right) + O \left( \exp_{p-1} [(\tau_1 + \varepsilon) \{\phi(r)\}^\mu] \right) \\ & + \exp_{p-1} [\{\phi(r)\}^{\mu + \varepsilon}] + (2k + 2)T(2r, f) + O(1) \\ & \Rightarrow T(2r, f) \geq O \left( \exp_{p-1} [\{\phi(r)\}^{\mu_p(A, \phi) - \varepsilon}] \right) \end{aligned}$$

for sufficiently large  $r$  and  $r \in F_3$ .

It follows that,  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu_p(A, \phi)$ .

This completes the proof. ■

Next we state two corollaries considering the homogeneous linear difference equation (1) which directly follows from Theorem 23 and Theorem 24.

**Corollary 25** *Let  $A_j(z)$  ( $j = 0, 1, \dots, k$ ) be meromorphic coefficients of (1) and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\}$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) such that*

$$\lambda_p \left( \frac{1}{A_m}, \phi \right) < \rho_p(A_m, \phi) < \infty,$$

$$\max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} \leq \rho_p(A_m, \phi)$$

$$\text{and } \max\{\tau_p(A_j, \phi) : \rho_p(A_j, \phi) = \rho_p(A_m, \phi), j = 0, 1, \dots, k, j \neq m\} < \tau_p(A_m, \phi) < \infty,$$

*then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A_m, \phi)$ .*

**Proof.** Putting  $A = 0$  in the Theorem 23 we get the required result. ■

**Corollary 26** *Let  $A_j(z)$  ( $j = 0, 1, \dots, k$ ) and  $A(z)$  be meromorphic functions and  $p = \max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\}$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) such that*

$$\lambda_p \left( \frac{1}{A_m}, \phi \right) < \mu_p(A_m, \phi) < \infty,$$

$$\max\{\rho_p(A_j, \phi) : j = 0, 1, \dots, k, j \neq m\} \leq \mu_p(A_m, \phi)$$

$$\text{and } \max\{\tau_p(A_j, \phi) : \rho_p(A_j, \phi) = \mu_p(A_m, \phi), j = 0, 1, \dots, k, j \neq m\} < t_p(A_m, \phi) < \infty,$$

*then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\mu_p(f, \phi) \geq \mu_p(A_m, \phi)$ .*

**Proof.** Putting  $A = 0$  in the Theorem 24 we get the required result. ■

Our Theorem 27 and Theorem 28 are improvements of the theorems due to Belaïdi-Benkarouba ([1], Theorem 1.1 and Theorem 1.2). Here we modify the hypothesis by giving some conditions on the Characteristic functions of the coefficients of (1) rather than taking coefficient having maximal iterated  $p - \phi$  order.

**Theorem 27** *Let  $A_0(z), A_1(z), \dots, A_k(z)$  be meromorphic coefficients of (1) satisfying  $\max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\} = p$  and  $\max\{\rho_p(A_j, \phi) : 0 \leq j \leq k\} \leq \rho$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) with  $\lambda_p \left( \frac{1}{A_m}, \phi \right) < \rho_p(A_m, \phi)$  and a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$  such that for some constants  $a, b$  ( $0 \leq b < a$ ) and  $\delta$  ( $0 < \delta < \rho$ ) sufficiently small with*

$$T(r, A_m) \geq \exp_{p-1} \left[ a \{\phi(r)\}^{\rho-\delta} \right], \quad (38)$$

$$\text{and } T(r, A_j) \leq \exp_{p-1} \left[ b \{\phi(r)\}^{\rho-\delta} \right], \quad j = 0, 1, \dots, k; j \neq m, \quad (39)$$

*as  $r \rightarrow \infty$  for  $r \in F$ , then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A_m, \phi)$ .*

**Proof.** First we shall show that  $\rho_p(A_m, \phi) = \rho$ . If not let us assume that  $\rho > \rho_p(A_m, \phi) = \sigma$  (say). Then for given any  $\varepsilon > 0$  and for sufficiently large  $r$ , we have

$$T(r, A_m) \leq \exp_{p-1} \left[ \{\phi(r)\}^{\sigma+\varepsilon} \right]$$

Again by (38), there exists  $\delta$  ( $0 < \delta < \rho$ ), such that

$$T(r, A_m) \geq \exp_{p-1} \left[ a \{\phi(r)\}^{\rho-\delta} \right]$$

as  $r \rightarrow \infty$  for  $r \in F$ .

Now choosing  $\varepsilon < \rho - \sigma - 2\delta$  and combining the above two inequalities, we obtain for  $r \in F$

$$\exp_{p-1} \left[ a \{\phi(r)\}^{\rho-\delta} \right] \leq T(r, A_m) \leq \exp_{p-1} \left[ \{\phi(r)\}^{\sigma+\varepsilon} \right],$$

which is a contradiction as  $r \rightarrow \infty$ .

Hence  $\rho_p(A_m, \phi) = \rho$ .

Now let  $f$  be a nonzero meromorphic solution of (1) and put  $c_0 = 0$ . If possible let us assume that  $\rho > \rho_p(f, \phi) = \kappa$  (say). Then by Lemma 17, for given any  $\varepsilon > 0$  and for sufficiently large  $r$ , we have

$$m\left(r, \frac{f(z+c_j)}{f(z+c_m)}\right) = O\left(\exp_{p-1}\left[\{\phi(r)\}^{\kappa+\varepsilon}\right]\right), \text{ for } j \neq m. \quad (40)$$

Also by the definition of  $\lambda_p\left(\frac{1}{A_m}, \phi\right) = \lambda$ , (say) we have for the above  $\varepsilon$  and for sufficiently large  $r$

$$N(r, A_m) \leq \exp_{p-1}\{\phi(r)\}^{\lambda+\varepsilon}. \quad (41)$$

By using similar arguments as in Theorem 1, from (4) we get,

$$\begin{aligned} T(r, A_m) &= m(r, A_m) + N(r, A_m) \\ &\leq \sum_{j=0, j \neq m}^k m(r, A_j) + \sum_{j=0, j \neq m}^k m\left(r, \frac{f(z+c_j)}{f(z+c_m)}\right) + N(r, A_m) + O(1) \end{aligned} \quad (42)$$

for  $r \notin F_1$ , where  $lm(F_1) < +\infty$

Now, using the relations (38)-(41) and choosing  $\varepsilon$  such that  $0 < \varepsilon < \frac{1}{2} \min\{\rho - \kappa - 2\delta, \rho - \lambda - 2\delta\}$ , we have from (42)

$$\begin{aligned} \exp_{p-1}\left[\{a\phi(r)\}^{\rho-\delta}\right] &\leq k \exp_{p-1}\left[b\{\phi(r)\}^{\rho-\delta}\right] \\ &\quad + \exp_{p-1}\left[\{\phi(r)\}^{\kappa+\varepsilon}\right] + \exp_{p-1}\left[\{\phi(r)\}^{\lambda+\varepsilon}\right] + O(1). \end{aligned}$$

for  $r \in F \setminus F_1$ .

Which implies

$$(a-b)\{\phi(r)\}^{\rho-\delta} \leq \{\phi(r)\}^{\kappa+\varepsilon} + \{\phi(r)\}^{\lambda+\varepsilon} + O(1), \text{ for } r \in F \setminus F_1.$$

Since  $(a-b) > 0$ , the above expression implies

$$1 \leq \frac{\{\phi(r)\}^{\kappa+\varepsilon-\rho+\delta} + \{\phi(r)\}^{\lambda+\varepsilon-\rho+\delta}}{(a-b)} + \frac{O(1)}{(a-b)\{\phi(r)\}^{\rho-\delta}} \rightarrow 0 \text{ as } r \rightarrow +\infty,$$

which is a contradiction.

Thus we have,  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$  as well as  $i_\rho(f, \phi) \geq p$ . ■

Our next theorem i.e. Theorem 28 is the entire version of the previous theorem.

**Theorem 28** *Let  $A_0(z), A_1(z), \dots, A_k(z)$  be entire coefficients of (1) satisfying  $\max\{i_\rho(A_j, \phi) : j = 0, 1, \dots, n-1\} = p$  and  $\max\{\rho_p(A_j, \phi) : 0 \leq j \leq k\} \leq \rho$ . If there exists an  $A_m(z)$  ( $0 \leq m \leq k$ ) and a set  $F \subset (1, +\infty)$  with  $lm(F) = +\infty$*

such that for some constants  $a, b$  ( $0 \leq b < a$ ) and  $\delta$  ( $0 < \delta < \rho$ ) sufficiently small with

$$|A_m(z)| \geq \exp_p \left[ a \{\phi(r)\}^{\rho-\delta} \right] \quad (43)$$

$$\text{and } |A_j(z)| \leq \exp_p \left[ b \{\phi(r)\}^{\rho-\delta} \right], \quad j = 0, 1, \dots, k; \quad j \neq l, \quad (44)$$

as  $r \rightarrow \infty$  for  $r \in F$ , then every solution  $f(z)$  ( $f \neq 0$ ) of (1) satisfies  $i_\rho(f, \phi) \geq p$  and  $\rho_p(f, \phi) \geq \rho_p(A_m, \phi)$ .

**Proof.** First we shall show that  $\rho_p(A_m, \phi) = \rho$ . If not let us assume that  $\rho > \rho_p(A_m, \phi) = \sigma$  (say). Then for given any  $\varepsilon > 0$  and for sufficiently large  $r$ , we have

$$|A_m(z)| \leq \exp_p \left[ \{\phi(r)\}^{\sigma+\varepsilon} \right]$$

Again by (43), there exists  $\delta$  ( $0 < \delta < \rho$ ), such that

$$|A_m(z)| \geq \exp_p \left[ a \{\phi(r)\}^{\rho-\delta} \right]$$

as  $r \rightarrow \infty$  for  $r \in F$ .

Now choosing  $\varepsilon < \rho - \sigma - 2\delta$  and combining the above two inequalities, we obtain for  $r \in F$

$$\exp_p \left[ a \{\phi(r)\}^{\rho-\delta} \right] \leq |A_m(z)| \leq \exp_p \left[ \{\phi(r)\}^{\sigma+\varepsilon} \right],$$

which is a contradiction as  $r \rightarrow \infty$ .

Hence  $\rho_p(A_m, \phi) = \rho$ .

Now let  $f$  be a nonzero meromorphic solution of (1) and put  $c_0 = 0$ . If possible let us assume that  $\rho > \rho_p(f, \phi) = \kappa$  (say).

Now divide (1) by  $f(z + c_m)$  we get

$$\begin{aligned} -A_m(z) &= \sum_{j=0, j \neq m}^k A_j(z) \frac{f(z + c_j)}{f(z + c_m)}. \\ \Rightarrow -1 &= \sum_{j=0, j \neq l}^k \frac{A_j(z) f(z + c_j)}{A_m(z) f(z + c_m)} \\ \Rightarrow 1 &\leq \sum_{j=0, j \neq l}^k \left| \frac{A_j(z) f(z + c_j)}{A_m(z) f(z + c_m)} \right| \end{aligned} \quad (45)$$

Choose  $\varepsilon > 0$  such that  $0 < \varepsilon < \frac{1}{2} \min \{\rho - \kappa - 2\delta, \rho - \lambda - 2\delta\}$ . Then by Lemma 20, for that  $\varepsilon$  there exists a subset  $F_1 \subset (1, +\infty)$  with  $lm(F_1) < +\infty$  such that for all  $r \notin F_1 \cup [0, 1]$ , we have

$$\left| \frac{f(z + c_j)}{f(z + c_m)} \right| \leq \exp_p \left[ \{\phi(r)\}^{\kappa+\varepsilon} \right] < \exp_p \left[ \{\phi(r)\}^{\rho-2\delta} \right], \quad \text{for } j \neq m. \quad (46)$$

Thus for  $r \in F \setminus (F_1 \cup [0, 1])$ , substituting (43), (44) and (46) in (45), we obtain

$$1 \leq k \frac{\exp_p \left[ b \{\phi(r)\}^{\rho-\delta} \right]}{\exp_p \left[ a \{\phi(r)\}^{\rho-\delta} \right]} \exp_p \left[ \{\phi(r)\}^{\rho-2\delta} \right] \rightarrow 0 \text{ as } r \rightarrow \infty.$$

Which is a contradiction.

Hence we get  $\rho_p(f, \phi) \geq \rho = \rho_p(A_m, \phi)$  as well as  $i_\rho(f, \phi) \geq p$ . ■

## References

- [1] B. Belaïdi and Y. Benkarouba, Some properties of meromorphic solutions of higher order linear difference equations, Ser. A Appl. Math. Inform. Mech. 11(2), (2019) 75–95.
- [2] B. Belaïdi and R. Bellaama, Study of the growth properties of meromorphic solutions of higher-order linear difference equations, Arab. J. Math. (2021). <https://doi.org/10.1007/s40065-021-00324-2>.
- [3] T. B. Cao, J. F. Xu and Z. X. Chen, On the meromorphic solutions of linear differential equations on the complex plane. J. Math. Anal. Appl. 364(1), (2010) 130–142.
- [4] Y. M. Chiang and S. J. Feng, On the Nevanlinna characteristic of  $f(z + \eta)$  and difference equations in the complex plane, Ramanujan J. 16(1), (2008) 105–129.
- [5] I. Chyzhykov, J. Heittokangas and J. Rättyä: Finiteness of  $\phi$ -order of solutions of linear differential equations in the unit disc. J. Anal. Math., Vol. 109, (2009) 163–198.
- [6] A. Goldberg and I. Ostrovskii, Value Distribution of Meromorphic functions, Transl. Math. Monogr., vol. 236, Amer. Math. Soc., Providence RI, 2008.
- [7] R. G. Halburd and R. J. Korhonen, Difference analogue of the lemma on the logarithmic derivative with applications to difference equations, J. Math. Anal. Appl. 314(2), (2006) 477–487.
- [8] W. K. Hayman, Meromorphic Functions, Oxford Mathematical Monographs. Clarendon Press, Oxford, 1964.
- [9] H. Hu and X. M. Zheng, Growth of solutions to linear differential equations with entire coefficients. Electron. J. Differ. Equ. 2012 (226), (2012) 1–15.
- [10] L. Kinnunen, Linear differential equations with solutions of finite iterated order. Southeast Asian Bull. Math. 22 (4) (1998) 385–405.

- [11] I. Laine and C. C. Yang, Clunie theorems for difference and  $q$ -difference polynomials. *J. Lond. Math. Soc.* (2) 76(3), (2007) 556–566.
- [12] Z. Latreuch and B. Belaïdi, Growth and oscillation of meromorphic solutions of linear difference equations. *Mat. Vesnik* 66(2), (2014) 213–222.
- [13] H. Liu and Z. Mao, On the meromorphic solutions of some linear difference equations. *Adv. Differ. Equ.* 2013(133), (2013) 1–12.
- [14] X. Shen, J. Tu and H. Y. Xu: Shen et al.: Complex oscillation of a second-order linear differential equation with entire coefficients of  $[p, q] - \phi$  order. *Advances in Difference Equations* 2014, 2014:200.
- [15] X. M. Zheng and J. Tu, Growth of meromorphic solutions of linear difference equations, *J. Math. Anal. Appl.* 384 (2), (2011) 349–356.
- [16] Y. P. Zhou and X. M. Zheng, Growth of meromorphic solutions to homogeneous and non-homogeneous linear (differential-)difference equations with meromorphic coefficients. *Electron. J. Differ. Equ.* 2017(34), (2017) 1-15.