

Graphs with Sudoku number $n - 1$

Alexey Pokrovskiy *

June 20, 2022

Abstract

Recently Lau-Jeyaseeli-Shiu-Arumugam introduced the concept of the “Sudoku colourings” of graphs — partial $\chi(G)$ -colourings of G that have a unique extension to a proper $\chi(G)$ -colouring of all the vertices. They introduced the Sudoku number of a graph as the minimal number of coloured vertices in a Sudoku colouring. They conjectured that a connected graph has Sudoku number $n - 1$ if, and only if, it is complete. In this note we prove that this is true.

1 Introduction

All colourings in this note are vertex-colourings. A vertex colouring of a graph is *proper* if adjacent vertices receive different colours. The chromatic number, $\chi(G)$ is the minimal number of colours in a proper colouring of G . A *partial* proper colouring of a graph is a colouring of some subset of $V(G)$ which doesn’t give adjacent vertices the same colour. We say that a colouring ϕ extends a partial colouring ψ if all vertices coloured in ψ receive the same colour in ϕ . A variety of tasks can be encoded as taking a partial proper colouring in a graph and then extending it to a full proper colouring of all the vertices.

For example the well known “Sudoku puzzle” can be encoded in this form. Consider a graph G_{Sudoku} on 27 vertices that are identified with the cells in a 9×9 grid. Add an edge between any two vertices in the same column, between any two vertices in the same row, and between any two vertices in the same 3×3 box. It is easy to see that a proper 9-colouring of G_{Sudoku} exactly corresponds to filling in 9×9 array according to the rules of Sudoku puzzles. Thus a Sudoku puzzle can be summarized as “you are given a partial colouring of G_{Sudoku} and need to complete it to a proper colouring of all the vertices of G_{Sudoku} ”.

One of the conventions for designing Sudoku puzzles is that there should always be precisely one way of filling in the 9×9 array i.e. there should always exist one solution, and there shouldn’t exist multiple solutions. This motivates the definition of a Sudoku colouring of a graph. Lau-Jeyaseeli-Shiu-Arumugam defined it as follows in [1].

Definition 1. *A Sudoku colouring of a graph G is a partial proper $\chi(G)$ -colouring of G which has precisely one extension to a $\chi(G)$ -colouring of G .*

Sudoku colourings of G_{Sudoku} are thus in one-to-one correspondence with uncompleted Sudoku puzzles (that have unique completion). But we can also investigate Sudoku colourings of general graphs.

Lau-Jeyaseeli-Shiu-Arumugam [1] defined the Sudoku number of G , $sn(G)$ as the smallest number of coloured vertices in a Sudoku colouring of G . The motivation for this is that $sn(G_{\text{Sudoku}})$ now asks for the minimum number of clues (i.e. non-blank entries) in a Sudoku puzzle with unique solution. This number has been determined as $sn(G_{\text{Sudoku}}) = 17$ by McGuire, Tugemann, and Civario using a computer-assisted proof [2].

*Email: dr.alexey.pokrovskiy@gmail.com. Address: Department of Mathematics, University College London, UK.

For general graphs, Lau-Jeyaseeli-Shiu-Arumugam [1] determined $sn(G)$ for various classes of graphs G and obtained bounds for other classes. For example, they showed that $sn(G) = 1$ if, and only if, G is connected and bipartite. On the other extreme, they showed that $sn(G) \leq |G| - 1$ for all graphs and conjectured that for connected graphs, equality holds if, and only if, G is complete. Here we show that this is the case.

Theorem 2. *A connected graph has $sn(G) = |G| - 1$ if, and only if, G is complete.*

The backwards direction already appears in [1] (see Corollary 3.4), so we focus on proving the statement “let G be a connected graph with $sn(G) = |G| - 1$. Then G is complete”. This amounts to showing that non-complete graphs have partial proper $\chi(G)$ -colourings with ≥ 2 uncoloured vertices which have a unique extension to a full proper $\chi(G)$ -colouring of G .

2 Proofs

In a k -coloured graph our set of colours will always be $[k] = \{1, \dots, k\}$. For a colouring ϕ and vertices u_1, \dots, u_k , we use $\phi - u_1 - \dots - u_k$ to mean the partial colouring formed by uncolouring the vertices u_1, \dots, u_k . For a partial colouring ϕ and a set of vertices S , we define $\phi(S) := \{\phi(s) : s \in S\}$ to mean the set of colours appearing in S . We use $N(v)$ to denote the set of vertices connected to v by an edge (our graphs are simple, so this never includes v itself).

The following is the tool we use for constructing Sudoku colourings in this note. It gives two kinds of Sudoku colourings with two uncoloured vertices.

Lemma 3. *Let ψ be a partial proper $\chi(G)$ -colouring with exactly two uncoloured vertices u, v . Suppose that either of the following holds:*

- (i) *uv is a nonedge and $|\phi(N(u))| = |\phi(N(v))| = \chi(G) - 1$.*
- (ii) *uv is an edge, $|\phi(N(u))| = \chi(G) - 1$, $|\phi(N(v))| = \chi(G) - 2$, and $\phi(N(v)) \subset \phi(N(u)$.*

Then ψ is a Sudoku colouring.

Proof. In case (i), there is precisely one colour missing from $N(u)$ and precisely one colour missing from $N(v)$. In a proper $\chi(G)$ -colouring, u and v must receive exactly these colours, and so the extension is unique.

In case (ii), there are two colours c, d missing from $N(v)$ and one of these colours (say c), is missing from $N(u)$. To complete the colouring u must receive colour c , and v must receive colour d , so the extension is unique. \square

The following definition is crucial for us.

Definition 4. *Let ϕ be a proper $\chi(G)$ -colouring of G . A vertex is **full** if it is adjacent to vertices of all colours (aside from its own) i.e. if $\phi(N(v)) = [\chi(G)] \setminus \phi(v)$ (or equivalently if $|\phi(N(v))| = \chi(G) - 1$).*

In graphs with Sudoku number $|G| - 1$, it turns out that the full vertices form a complete subgraph.

Lemma 5. *Let $sn(G) = |G| - 1$ and let ϕ be a proper $\chi(G)$ -colouring of G . Then any two full vertices are connected by an edge.*

Proof. Let u, v be full and suppose for contradiction that uv is not an edge in G . Consider the partial colouring $\psi := \phi - u - v$. Since u, v are full, we have $|\phi(N(u))| = |\phi(N(v))| = \chi(G) - 1$. Since uv is a non-edge, the neighbours of u, v all remain coloured in ψ and so $|\psi(N(u))| = |\psi(N(v))| = \chi(G) - 1$. Thus, by Lemma 3 (i), ψ is a Sudoku colouring. It has two uncoloured vertices, contradicting $sn(G) = |G| - 1$. \square

We say that a proper k -colouring of G is c -minimal if it has as few colour c vertices as possible (for a k -colouring of G). The following lemma shows that there are a lot of full vertices around all c -coloured vertices in a c -minimal colouring.

Lemma 6. *Let ϕ be a 1-minimal proper $\chi(G)$ -colouring of G . Let v be a vertex with $\phi(v) = 1$. Then for every colour $c = 2, \dots, \chi(G)$, the vertex v has at least one colour c neighbour u with u full. In particular, v is full.*

Proof. First notice that it is impossible that v has no colour c neighbours — indeed otherwise, we could recolour v with colour c to get a proper colouring with one fewer colour 1 vertex (contradicting 1-minimality). This proves the “in particular v is full” part — since we’ve shown that every colour other than $\phi(v) = 1$ appears on $N(v)$.

Now let the set of colour c neighbours of v be $\{u_1, \dots, u_k\}$. Suppose for contradiction that none of these are full — equivalently there are colours $c_1, \dots, c_k \in [\chi(G)] \setminus c$ with c_i missing from $N(u_i)$. Note that $c_i \neq 1$ for all i , since $v \in N(u_i)$ and $\phi(v) = 1$. Note that $\{u_1, \dots, u_k\}$ is an independent set since all these vertices have colour c and the colouring is proper.

Now recolour u_i by c_i for each i and recolour v by c . Notice that this colouring is proper. To show this, we need to check that the recoloured vertices v, u_1, \dots, u_k have different colours to all their neighbours (everywhere else the colouring remains proper just because ϕ was proper). Indeed v has no colour c neighbours since $\{u_1, \dots, u_k\}$ was the set of all colour c neighbours of v and these have all been recoloured by colours $c_i \neq c$. Vertex u_i has no colour c_i neighbour since it initially had no colour c_i neighbours and the only neighbour of u_i that was recoloured was v (which received colour $c \neq c_i$).

But the new colouring we have has one fewer colour 1 vertex, contradicting 1-minimality. \square

Applying the above lemma to a graph with Sudoku number $|G| - 1$ gives even more structure in a minimal colouring.

Lemma 7. *Let $sn(G) = |G| - 1$ and let ϕ be a 1-minimal proper $\chi(G)$ -colouring of G . Then there is precisely one colour 1 vertex. Additionally, this vertex v is full and has $|N(v)| = \chi(G) - 1$.*

Proof. Let v be a colour 1 vertex. By Lemma 6 v is full. There can’t be another colour 1 vertex z — because otherwise z would also be full, which would give two disconnected full vertices (contradicting Lemma 5).

Suppose that $|N(v)| > \chi(G) - 1$. Then, by the pigeonhole principle there must be some colour c which occurs more than once on $N(v)$. By Lemma 6 v has some full colour c neighbour u . Let w be some other colour c neighbour of v . Now let $\psi := \phi - u - v$. Note that $\psi(N(v)) = \phi(N(v)) = [\chi(G)] \setminus 1$ (the first equality holds because precisely one neighbour u of v was uncoloured, but that neighbour u had colour c which is still present at w . The second equality holds because v is full and has colour 1). Also $\psi(N(u)) = \phi(N(u)) \setminus 1 = [\chi(G)] \setminus \{1, c\}$ (the first equality holds because precisely one neighbour v or u was uncoloured, and that neighbour had colour 1 which isn’t present anywhere else in the graph. The second equality holds because $\phi(N(u)) = [\chi(G)] \setminus c$ since u is full and has colour c). Thus by Lemma 3 (ii), ψ is a Sudoku colouring with two uncoloured vertices, contradicting $sn(G) = |G| - 1$. \square

We are ready to prove our main theorem.

Proof of Theorem 2. The backwards direction already appears in [1] (see Corollary 3.4), so it remains to prove that every connected G with $sn(G) = |G| - 1$ is complete. To that end, let G be connected with $sn(G) = |G| - 1$.

Consider a 1-minimal proper $\chi(G)$ -colouring of G . By Lemma 7, there is precisely one colour 1 vertex, call it v . Also by Lemma 7, v is full and $|N(v)| = \chi(G) - 1$ — which, using the definition of “full”, implies that all neighbours of v have different colours. By Lemma 6, we have that all neighbours of v are full. Now we have that all vertices in $v \cup N(v)$ are full, and so Lemma 5 tells us that $v \cup N(v)$ is complete.

Unless G is complete, then, by connectedness, there is some vertex w outside $v \cup N(v)$ with a neighbour u in $N(v)$. Now construct a colouring ψ by recolouring w by colour 1 and uncolouring u, v . First notice that this is a proper (partial) colouring — indeed w has no colour 1 neighbours since v is the unique colour 1 vertex and $w \notin N(v)$. We have that $\psi(N(v)) = \phi(N(v)) \setminus \phi(u) = [\chi(G)] \setminus \{1, \phi(u)\}$ (the first equality holds because the neighbour u of v was uncoloured and $\phi(u)$ doesn't appear anywhere else on $N(v)$ since the neighbours of v have different colours. The second equality holds because $\phi(N(v)) = [\chi(G)] \setminus 1$ since v is full and has colour 1). Also $\psi(N(u)) = [\chi(G)] \setminus \{\phi(u)\}$ (colour 1 is present on $N(u)$ in ψ because $\psi(w) = 1$ and $w \in N(u)$). All colours in $[\chi(G)] \setminus \{1, \phi(u)\}$ are present on $N(u)$ in ψ because the vertices of $N(v) \setminus u$ have exactly these colours in ϕ , u is connected to all of them, and their colours don't change going from ϕ to ψ . Thus by Lemma 3 (ii), ψ is a Sudoku colouring with two uncoloured vertices, contradicting $sn(G) = |G| - 1$. \square

References

- [1] Gee-Choon Lau, J. Maria Jeyaseeli, Wai-Chee Shiu, and S. Arumugam. Sudoku number of graphs. *arXiv:2206.08106*, 2022.
- [2] Gary McGuire, Bastian Tugemann, and Gilles Civario. There is no 16-clue sudoku: Solving the sudoku minimum number of clues problem via hitting set enumeration. *Experimental Mathematics*, 23(2):190–217, 2014.