

THE POSET OF SPECHT IDEALS FOR HYPEROCTAHEDRAL GROUPS

SEBASTIAN DEBUS, PHILIPPE MOUSTROU, CORDIAN RIENER, AND HUGUES VERDURE

ABSTRACT. Specht polynomials classically realize the irreducible representations of the symmetric group. The ideals defined by these polynomials provide a strong connection with the combinatorics of Young tableaux and have been intensively studied by several authors. We initiate similar investigations for the ideals defined by the Specht polynomials associated to the hyperoctahedral group B_n . We introduce a bidominance order on bipartitions which describes the poset of inclusions of these ideals and study algebraic consequences on general B_n -invariant ideals and varieties, which can lead to computational simplifications.

CONTENTS

1. Introduction	1
2. \mathcal{S}_n -Specht ideals	3
2.1. Definitions	3
2.2. Inclusions and applications	4
3. Definition and first properties of B_n -Specht ideals	4
3.1. Definitions	4
3.2. Relations between B_n and \mathcal{S}_n Specht polynomials	5
3.3. Existing orders on bipartitions	6
4. The poset of bipartitions	7
5. The posets of Specht ideals and varieties	11
6. Orbit types	15
7. Applications to B_n -invariant ideals	18
7.1. Specht ideals in B_n -invariant ideals	18
7.2. Connections with Representation Theory	20
8. Conclusion and open questions	22
References	22

1. INTRODUCTION

Symmetries provide beautiful connections between algebra, geometry and efficient computations: on the one hand, the symmetries of geometrical objects can be described with the algebraic language of group theory, while on the other hand algebraic problems affording additional structure can be solved more efficiently once symmetry is appropriately taken into consideration. A particular incarnation of these phenomena occurs when studying algebraic systems of polynomial equations whose solution set is invariant under a group action. In this set-up, when looking at the corresponding polynomial ideal, the machinery of invariant and representation theory can be employed to gain information about the solutions of the initial system, and to simplify its resolution.

Key words and phrases. bipartitions, Specht polynomials, hyperoctahedral group, invariant ideals.

This work has been supported by European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 813211 (POEMA) and the Tromsø Research foundation grant agreement 17matteCR.

This kind of questions have been extensively studied in the literature for the *symmetric group* \mathcal{S}_n , acting on the polynomial ring $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ over a field \mathbb{K} by permuting variables. In particular, it has been observed in different computational tasks that the understanding of this action can lead to substantial algorithmic improvements (see for example [Fau+20; Rie+13; RS18; HS21; BR21b; Mou+22; BR21a; KLM20]). These improvements mostly build on the fact that in this situation, both representation and invariant theory are classically understood, and are closely related to the combinatorics of partitions and Young Tableaux. More precisely, the irreducible representations of \mathcal{S}_n are in bijection with the partitions of n , through a construction due to Specht: for every partition, one can define a polynomial whose \mathcal{S}_n -orbit spans an irreducible \mathcal{S}_n -module, called *Specht module* [Spe37b]. This motivates the study of *Specht ideals*, the ideals generated by such modules, since they can be seen as building blocks of the action of the symmetric group on a polynomial ring. The study of these objects has shown to be fruitful from various aspects, and the connection between these ideals and the combinatorics of partitions turns out to be deeper: not only there is a bijection between \mathcal{S}_n -Specht ideals and partitions of n , but this correspondence respects the poset structures. First results were proven by Haiman and Woo (see Woo’s doctoral thesis [Woo05]), and then independently revisited and extended for algorithmic purposes in [MRV21]. In turn, this combinatorial understanding also provides information on these ideals from the point of view of commutative algebra: for instance they all are radical [MOY21], and the partitions for which they are Cohen-Macaulay are understood [Yan21]. The study of these ideals has also paved algorithmic ways to simplify calculations for \mathcal{S}_n -closed ideals and their corresponding varieties. They allow to understand the symmetry of the coordinates of the points in the variety, which in turn gives information on their dimension. This information can then be used to design more efficient algorithms by reducing the number of variables.

In this article, we initiate a similar study for the action of the *hyperoctahedral group* B_n on a polynomial ring $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$. The field \mathbb{K} is assumed to be infinite, although many results remain valid for finite fields. In this representation, this group can be seen as the group generated by permutations of variables and sign switches of variables, namely maps sending \mathbf{x}_i to $-\mathbf{x}_i$. The group is isomorphic to the Weyl group of type B and appears in several different areas, as hyperplane arrangements ([AM17, Section 6.7], [Abe+20]), representation theory [CS93; Che93; Mus93], and has applications in the study of non negative even symmetric polynomials [CLR87; Har99] and optimization [Dos+17]. Similarly to the case of permutations, this situation is profoundly connected to combinatorics. In this case, instead of partitions, the irreducible representations of B_n are in bijection with *bipartitions* of n . Furthermore, polynomial generators of the irreducible B_n -modules can be constructed in a similar way [Spe37a]. We aim at a first investigation of the corresponding ideals with the goal to extend the connections between algebra and combinatorics as far as possible. In contrast to the \mathcal{S}_n -case where there is a natural order on partitions, several orders are possible on bipartitions [AMP81; DJM95]. However, while in the \mathcal{S}_n case the poset of the standard order on partitions reflects the corresponding poset of ideal inclusions, none of the previously studied orders of bipartitions satisfies this property. Therefore, we define another order on bipartitions. After studying the basic properties of this order, we are able to show that it indeed translates well to the ideal inclusion. Similarly to the case of \mathcal{S}_n , this combinatorial connection finds consequences for the corresponding varieties. In addition to the inclusion of varieties we are able to give a complete characterization in terms of orbit types of the points in these varieties. Further, this gives information on the possible orbit types of points in general B_n -invariant varieties, allowing for complexity reduction in the resolution of B_n -closed polynomial systems.

The paper is structured as follows: Section 2 overviews the situation of \mathcal{S}_n -Specht ideals. We initiate the study of B_n -Specht ideals in Section 3 with definitions and natural connections to the \mathcal{S}_n case. In Section 4 we define our order for bipartitions and study its combinatorial properties. Following this, we show equivalence between our poset of bipartitions and the posets of Specht ideals and varieties in Section 5. In Section 6, study possible decompositions of Specht

varieties in terms of orbit types. Finally, we extend our study to general B_n -invariant ideals in Section 7, before concluding the paper with closing remarks and open questions in Section 8.

2. \mathcal{S}_n -SPECHT IDEALS

2.1. Definitions. A *partition* $\lambda = (\lambda_1, \dots, \lambda_l)$ of n is a sequence of non-increasing non negative integers such that $\sum_{i \geq 1} \lambda_i = n$. We write $\lambda \vdash n$ when λ is a partition of n and say that \emptyset is the unique partition of 0. The *size* of a partition λ is $|\lambda| = \sum_{i \geq 1} \lambda_i$. The *length* of a non-empty partition $\lambda = (\lambda_1, \lambda_2, \dots) \vdash n$ is the maximal $l \in \mathbb{N}_0$ with $\lambda_l > 0$, while the length of \emptyset is 0. We denote the length by $\text{len}(\lambda)$. For any partition $\lambda = (\lambda_1, \dots, \lambda_l) \vdash n$ we use the convention that $\lambda_s = 0$ for every $s \geq l + 1$.

Let $\lambda, \mu \vdash n$ be partitions of the same size. Then λ *dominates* μ if and only if $\sum_{j=1}^k \lambda_j \geq \sum_{j=1}^k \mu_j$ for any k . We denote domination by $\mu \trianglelefteq \lambda$. A partition λ can be represented via its (*Young*) *diagram*, i.e., the ordered sequence of boxes from the left to the right and the top to the bottom, where the i -th line contains λ_i many boxes. We say that the associated diagram has *shape* λ . A *tableau* of shape λ is a filling of a diagram of shape λ with all the numbers $[n] = \{1, \dots, n\}$.

Then, we write $\text{sh}(T) = \lambda$ if T is a tableau of shape λ . For instance, $S = \begin{array}{|c|c|c|c|} \hline 9 & 3 & 6 & 4 \\ \hline 2 & 1 & 8 & \\ \hline 5 & 7 & & \\ \hline \end{array}$ is a tableau of shape $(4, 3, 2)$. A *generalized* tableau is a filling of a diagram with elements in \mathbb{K} . The *conjugate partition* λ^\vee of a partition λ is the partition whose diagram is the one obtained from the diagram of λ by interchanging the rows and columns.

For a sequence (i_1, \dots, i_m) of natural numbers, we define the associated *Vandermonde polynomial* in the variables $\mathbf{x}_{i_1}, \dots, \mathbf{x}_{i_m}$ as

$$\Delta_{(i_1, \dots, i_m)}(\mathbf{x}) = \prod_{j < k \in [m]} (\mathbf{x}_{i_j} - \mathbf{x}_{i_k}),$$

while $\Delta_{(i)} = \prod_{\emptyset} (\mathbf{x}_{i_j} - \mathbf{x}_{i_k}) = 1$.

Definition 2.1. Let T be a tableau of shape $\lambda \vdash n$ with m columns and let T_i be the sequence of natural numbers containing the entries of the i -th column of T from above to below. Then, the associated \mathcal{S}_n Specht polynomial $\text{sp}_T(\mathbf{x})$ is the product of all the column Vandermonde polynomials of the columns, i.e.,

$$\text{sp}_T(\mathbf{x}) = \prod_{j=1}^m \Delta_{T_j}.$$

For the tableau S of shape $(4, 3, 2)$ above, we have

$$\begin{aligned} \text{sp}_S(\mathbf{x}) &= \Delta_{(9,2,5)}(\mathbf{x}) \Delta_{(3,1,7)}(\mathbf{x}) \Delta_{(6,8)}(\mathbf{x}) \Delta_{(4)}(\mathbf{x}) \\ &= (\mathbf{x}_9 - \mathbf{x}_2)(\mathbf{x}_9 - \mathbf{x}_5)(\mathbf{x}_2 - \mathbf{x}_5)(\mathbf{x}_3 - \mathbf{x}_1)(\mathbf{x}_3 - \mathbf{x}_7)(\mathbf{x}_1 - \mathbf{x}_7)(\mathbf{x}_6 - \mathbf{x}_8). \end{aligned}$$

Definition 2.2. Let λ be a partition of n . We define the \mathcal{S}_n -Specht ideal

$$I_\lambda = \langle \text{sp}_T(\mathbf{x}) : T \text{ is a tableau of shape } \lambda \rangle \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$$

and the \mathcal{S}_n -Specht variety

$$V_\lambda = \{a \in \mathbb{K}^n : f(a) = 0 \text{ for all } f \in I_\lambda\} \subset \mathbb{K}^n$$

associated to λ .

The group \mathcal{S}_n acts transitively on the set of tableaux of shape λ , where an element $\sigma \in \mathcal{S}_n$ acts on a tableau T by replacing every entry i in a box by $\sigma(i)$. Thus, the \mathcal{S}_n -Specht ideal I_λ is the ideal generated by the \mathcal{S}_n -orbit of a Specht polynomial of a tableau of shape λ .

Definition 2.3. For a partition $\lambda \vdash n$ we write $\mathcal{S}_\lambda = \mathcal{S}_{\lambda_1} \times \mathcal{S}_{\lambda_2} \times \dots \subset \mathcal{S}_n$ and define the \mathcal{S}_n -orbit set $H_\lambda = \{z \in \mathbb{K}^n : \text{Stab}_{\mathcal{S}_n}(z) \simeq \mathcal{S}_\lambda\}$. If $z \in H_\lambda$ we call λ the \mathcal{S}_n -orbit type of z .

The orbit set of any partition is non-empty and the H_λ 's define a set partition of \mathbb{K}^n . For instance, $H_{(3,2,2,1)}$ is the \mathcal{S}_n orbit of the set


$$\{(a_1, a_1, a_1, a_2, a_2, a_3, a_3, a_4) \in \mathbb{K}^n : a_i \neq a_j, \forall i \neq j\}.$$

2.2. Inclusions and applications. The dominance order for integer partitions is well studied and understood. We recall that if (P, \leq) is a poset and $p, q \in P$ then p covers q if and only if $p \neq q$, $q \leq p$, and for any $r \in P$, $q \leq r \leq p$ implies $r \in \{p, q\}$. Brylawski studied the lattice of integer partitions of n with respect to the dominance order and classified the covering relations ([Bry73, Proposition 2.3]). Let $\lambda, \mu \vdash n$ be partitions. Then, $\mu \leq \lambda$ is a covering if and only if λ is of the form

$$\lambda = (\mu_1, \dots, \mu_{i-1}, \mu_i + 1, \mu_{i+1}, \dots, \mu_{j-1}, \mu_j - 1, \mu_{j+1}, \dots, \mu_l),$$

and either $j = i + 1$ or $\mu_i = \mu_{j-1}$ (and $\mu_{i-1} > \mu_i$ and $\mu_j > \mu_{j+1}$ to ensure that μ is a partition). In particular, the diagram of shape μ can be obtained from the diagram of shape λ via moving one box from the end of row i to row j .

Example 2.4. The following are two coverings of partitions displayed by their diagrams.



The following theorem shows the equivalences of the posets of partitions with respect to dominance order, and the posets of \mathcal{S}_n -Specht ideals and varieties with respect to inclusion.

Theorem 2.5 ([MRV21], Theorem 1). *Let λ and μ be partitions of n . Let I_λ, I_μ denote their associated \mathcal{S}_n -Specht ideals and V_λ, V_μ their associated \mathcal{S}_n -Specht varieties. Then, the following assertions are equivalent:*

- (1) *The partition λ dominates μ , i.e. $\lambda \geq \mu$;*
- (2) *The \mathcal{S}_n -Specht ideal I_λ contains the \mathcal{S}_n -Specht ideal I_μ , i.e. $I_\lambda \supset I_\mu$;*
- (3) *The \mathcal{S}_n -Specht variety V_λ is contained in the \mathcal{S}_n -Specht variety V_μ , i.e. $V_\lambda \subset V_\mu$.*

The \mathcal{S}_n -Specht varieties can be decomposed using \mathcal{S}_n orbit sets.

Theorem 2.6 ([MRV21], Corollary 1). *Let $\mu \vdash n$ be a partition. Then, the associated \mathcal{S}_n -Specht variety is*

$$V_\mu = \left(\bigcup_{\lambda \leq \mu} H_\lambda \right)^c = \bigcup_{\lambda \not\leq \mu} H_\lambda.$$

This characterization already shows that in general $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]/I_\lambda$ is not Cohen-Macaulay for a \mathcal{S}_n -Specht ideal I_λ , since the varieties are not equidimensional. Yanagawa classified the few cases when a Specht ideal is Cohen-Macaulay.

Theorem 2.7 ([Yan21], Corollary 4.4). *The ring $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]/I_\lambda$ is Cohen-Macaulay if and only if λ is one of the following form*

- (1) $\lambda = (n - d, 1, \dots, 1)$;
- (2) $\lambda = (n - d, d)$;
- (3) $\lambda = (a, a, 1)$.

The authors in [Woo05; MOY21] prove that a \mathcal{S}_n -Specht ideal is radical. Their proof uses Theorem 2.6, i.e., that \mathcal{S}_n -Specht varieties can be written as disjoint unions of \mathcal{S}_n orbit sets, and the non-emptiness of any orbit set H_λ .

3. DEFINITION AND FIRST PROPERTIES OF B_n -SPECHT IDEALS

3.1. Definitions. A *bipartition* of n is a pair (λ, μ) , where $\lambda \vdash n_1$, $\mu \vdash n_2$ are partitions and $n_1 + n_2 = n$. We denote the set of all bipartitions of n by BP_n . A (*Young*) *bidiagram* of a bipartition (λ, μ) is the pair of diagrams of shape λ and μ . A *bitableau* is a filling of a bidiagram with all the numbers in $[n]$. We write $\text{sh}(T, S) = (\lambda, \mu)$ if (T, S) is a bitableau of shape (λ, μ) .

For example, $(T', S') = \left(\begin{array}{|c|c|} \hline 4 & 3 \\ \hline 2 & \\ \hline 5 & \\ \hline \end{array}, \begin{array}{|c|} \hline 6 \\ \hline 1 \\ \hline \end{array} \right)$ is a bitableau of shape $((2, 1, 1), (1, 1))$. When we consider representatives of B_n -orbits of points, we do not need to distinguish between the

signs of coordinates. Thus, we write $\mathbf{x}^2 = (\mathbf{x}_1^2, \dots, \mathbf{x}_n^2)$ and analogously $z^2 = (z_1^2, \dots, z_n^2)$ for points $z \in \mathbb{K}^n$. A *generalized* bitableau is a filling of a bidiagram with elements in \mathbb{K} .

Definition 3.1. Let (T, S) be a bitableau and let T_i, S_i be the sequences of natural numbers containing the entries of the i -th column of T and S from above to below. Then, the associated B_n Specht polynomial is

$$\text{sp}_{(T,S)}(\mathbf{x}) = \text{sp}_T(\mathbf{x}^2) \text{sp}_S(\mathbf{x}^2) \prod_{k \in S} \mathbf{x}_k = \prod_{i \geq 1} \Delta_{T_i}(\mathbf{x}^2) \prod_{j \geq 1} \Delta_{S_j}(\mathbf{x}^2) \prod_{k \in S} \mathbf{x}_k$$

where the notation sp_T is naturally adapted in this context to a Tableau T which is not necessarily filled with the integers $1, \dots, k$. For the bitableau (T', S') of shape $((2, 1, 1), (1, 1))$ above, we have

$$\text{sp}_{(T,S)}(\mathbf{x}) = \Delta_{(4,2,5)}(\mathbf{x}^2) \cdot \Delta_{(3)}(\mathbf{x}^2) \cdot \Delta_{(6,1)}(\mathbf{x}^2) \cdot \mathbf{x}_6 \mathbf{x}_1 = (\mathbf{x}_4^2 - \mathbf{x}_2^2)(\mathbf{x}_4^2 - \mathbf{x}_5^2)(\mathbf{x}_2^2 - \mathbf{x}_5^2)(\mathbf{x}_6^2 - \mathbf{x}_1^2) \mathbf{x}_1 \mathbf{x}_6.$$

From now on, if not specified, Specht polynomials will stand for B_n -Specht polynomials.

Definition 3.2. Let (λ, μ) be a bipartition of n . We define the B_n -Specht ideal

$$I_{(\lambda, \mu)} = \langle \text{sp}_{(T,S)}(\mathbf{x}) : (T, S) \text{ is a bitableau of shape } (\lambda, \mu) \rangle \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$$

and the B_n -Specht variety

$$V_{(\lambda, \mu)} = \{z \in \mathbb{K}^n : f(z) = 0 \text{ for all } f \in I_{(\lambda, \mu)}\} \subset \mathbb{K}^n$$

associated to (λ, μ) .


Again, the B_n -Specht ideal $I_{(\lambda, \mu)}$ is the ideal generated by the \mathcal{S}_n orbit of a Specht polynomial of a bitableau of shape (λ, μ) . We observe that switching of signs of variables in a Specht polynomial $\text{sp}_{(T,S)}$ returns $\pm \text{sp}_{(T,S)}$. Thus, $I_{(\lambda, \mu)}$ also equals the B_n orbit of $\text{sp}_{(T,S)}$.

3.2. Relations between B_n and \mathcal{S}_n Specht polynomials.


Definition 3.3. Let $\lambda \vdash n_1, \mu \vdash n_2$ be partitions. Then, the glueing of λ and μ is the partition $\lambda \uplus \mu = (\lambda_1 + \mu_1, \lambda_2 + \mu_2, \dots) \vdash n_1 + n_2$. The concatenation $\lambda \vee \mu \vdash n_1 + n_2$ is the partition obtained by rearranging $(\lambda_1, \dots, \lambda_s, \mu_1, \dots, \mu_t)$ in decreasing order.

The glueing $\lambda \uplus \mu$ defines indeed again a partition. Since $\lambda_i \geq \lambda_{i+1}$ and $\mu_i \geq \mu_{i+1}$ we have $\lambda_i + \mu_i \geq \lambda_{i+1} + \mu_{i+1}$ for any i .

Example 3.4. The glueing of the partitions $(3, 2, 2), (4, 1)$ with diagrams



is the partition $(7, 3, 2)$ with diagram



Lemma 3.5. Let λ, μ be two partitions. Then, we have

$$(\lambda \uplus \mu)^\perp = \lambda^\perp \vee \mu^\perp.$$

As a consequence, the columns of the diagram of shape $\lambda \uplus \mu$ are in bijection with the columns in the bidiagram of shape (λ, μ) .

Proof. Let $\rho = (\lambda^\perp \vee \mu^\perp)^\perp$. Then

$$\begin{aligned} \rho_k &= |\{j, (\lambda^\perp \vee \mu^\perp)_j \geq k\}| &= |\{j, (\lambda^\perp)_j \geq k\} + \#\{j, (\mu^\perp)_j \geq k\}| \\ &= ((\lambda^\perp)^\perp)_k + ((\mu^\perp)^\perp)_k \\ &= \lambda_k + \mu_k \\ &= (\lambda \uplus \mu)_k. \end{aligned}$$

The last claim of the statement follows, since the columns of $\lambda \uplus \mu$ are the rows of its conjugation. \square

The previous Lemma provides a natural connection between bitableaux of shape (λ, μ) and Tableaux of shape $\lambda \uplus \mu$. Concretely, let (T, S) be a bitableau of shape (λ, μ) . Then, we can consider the tableau $T \uplus S$ of shape $\lambda \uplus \mu$, where the columns of $T \uplus S$ are filled like the columns of T and S . When two columns in $\lambda \uplus \mu$ have the same length, they are ordered by their occurrence in the bitableau (T, S) from the left to the right. For instance, for $(T^*, S^*) = \left(\begin{array}{|c|c|c|c|} \hline 1 & 2 & 10 & 9 \\ \hline 4 & 8 & 7 & \\ \hline 6 & & & \\ \hline \end{array}, \begin{array}{|c|} \hline 3 \\ \hline 5 \\ \hline \end{array} \right)$

we have $T^* \uplus S^* = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 10 & 3 & 9 \\ \hline 4 & 8 & 7 & 5 & \\ \hline 6 & & & & \\ \hline \end{array}$. Since this map is invertible we get:

Proposition 3.6. *The tableaux of shape $\lambda \uplus \mu$ are in 1:1 correspondence with the bitableaux of shape (λ, μ) . A bijection is given by $(T, S) \mapsto T \uplus S$.*

The lemma below describes the connection between \mathcal{S}_n and B_n -Specht polynomials. In particular, it motivates the definition of the same operations on the bidiagram of shape (λ, μ) and on the diagram of its glueing $\lambda \uplus \mu$ via moving some of the boxes in a diagram in section 4.

Lemma 3.7. *Let $(\lambda, \mu) \in \text{BP}_n$ be a bipartition and let (T, S) be a bitableau of shape (λ, μ) . Then*

$$\text{sp}_{(T,S)}(\mathbf{x}_1, \dots, \mathbf{x}_n) = \text{sp}_{T \uplus S}(\mathbf{x}_1^2, \dots, \mathbf{x}_n^2) \prod_{j \in S} \mathbf{x}_j.$$

Proof. It is an immediate consequence of Proposition 3.6, since the Specht polynomials are defined as product of Vandermonde polynomials on the columns of the glued partition. \square

3.3. Existing orders on bipartitions. As mentioned in the introduction, partial orders on the set of bipartitions of n have been studied by several authors. Let $(\lambda, \mu), (\lambda', \mu') \in \text{BP}_n$ be bipartitions. For instance the following statements define partial orders:

$$(3.1) \quad \begin{cases} \sum_{j \leq k} \lambda'_j \leq \sum_{j \leq k} \lambda_j, & \text{for all } k, \text{ and} \\ |\lambda'| + \sum_{j \leq k} \mu'_j \leq |\lambda| + \sum_{j \leq k} \mu_j, & \text{for all } k. \end{cases}$$

was introduced in [DJM95] to study Hecke algebras of type B_n , and was recently proven to occur naturally in the field of spin group theory [Xia18]. Ariki generalized their order to multipartitions to study Hecke algebras of type $G(m, 1, n)$ [Ari01]. The partial order

$$(3.2) \quad \begin{cases} |\lambda'| < |\lambda|, & \text{or} \\ |\lambda'| = |\lambda|, & \text{and } \lambda' \trianglelefteq \lambda, \mu' \trianglelefteq \mu. \end{cases}$$

was formalized in [AMP81] to construct B_n -irreducible representations based on a more general procedure valid for finite groups. The partial orders (3.1) and (3.2) are not equivalent. For instance, in (3.1) the bipartitions $((2), (1))$ and $((1, 1, 1), \emptyset)$ are not comparable, while the latter is larger than the first in (3.2).

Moreover, these orders do not capture inclusions of ideals and varieties. Namely, for both orders, we have the following ordering of bipartitions of $n = 2$:

$$((2), \emptyset) > ((1, 1), \emptyset) > ((1), (1)),$$

while the corresponding ideals are

$$I_{((1,1),\emptyset)} = \langle \mathbf{x}_1^2 - \mathbf{x}_2^2 \rangle \supsetneq I_{((1),(1))} = \langle \mathbf{x}_1, \mathbf{x}_2 \rangle \supsetneq I_{(2),\emptyset} = \langle 1 \rangle.$$

In the next section, we introduce a new order on bipartitions that will capture inclusion of Specht ideals.

4. THE POSET OF BIPARTITIONS

In this section we introduce our new order for bipartitions:

Definition 4.1. Let $(\lambda, \mu), (\lambda', \mu') \in \text{BP}_n$ be bipartitions of n . We say that (λ, μ) bidominates (λ', μ') if and only if

$$\sum_{j=1}^k (\lambda'_j + \mu'_j) \leq \sum_{j=1}^k (\lambda_j + \mu_j), \quad \text{and}$$

$$\sum_{j=1}^{k-1} (\lambda'_j + \mu'_j) + \lambda'_k \leq \sum_{j=1}^{k-1} (\lambda_j + \mu_j) + \lambda_k$$

for all positive integer k . If (λ, μ) bidominates (λ', μ') we write $(\lambda', \mu') \trianglelefteq (\lambda, \mu)$. We call \trianglelefteq the bidominance order.

We point out that the first condition is just a condition on the glueing of the bipartitions, i.e.,

$$\lambda' \uplus \mu' \trianglelefteq \lambda \uplus \mu.$$

Example 4.2. The following bipartitions of 8 are comparable: $((2, 1, 1), (3, 1)) \trianglelefteq ((3, 2), (2, 1))$, since

$$2 \leq 3, \quad 5 \leq 5, \quad 6 \leq 7, \quad 7 \leq 8, \quad 8 \leq 8.$$

However, the bipartitions $((2), (1, 1))$ and $(\emptyset, (4))$ are not comparable, since $2 > 0$ but $3 < 4$.

Although we use the same symbol for dominance and bidominance, this should not create any confusion, as they are defined on sets with empty intersection. We identify bipartitions with their associated bidiagrams and speak about *boxes* in a bipartition.

It follows from the definition that our bidominance order is a partial order on BP_n , and the previous example shows that it is not a total order.

Before proving our main theorem in the next section, we need a better understanding of our poset of bipartitions.

The smallest element in $(\text{BP}_n, \trianglelefteq)$ is $(\emptyset, (1, \dots, 1))$, while the largest element is $((n), \emptyset)$. The following theorem characterizes the covering relations in the poset $(\text{BP}_n, \trianglelefteq)$. It turns out that there are four different cases, that are illustrated in Example 4.4.

Theorem 4.3. Let $(\lambda, \mu), (\lambda', \mu') \in \text{BP}_n$ be bipartitions and let $i = \min\{j \in [n] : (\lambda_j, \mu_j) \neq (\lambda'_j, \mu'_j)\}$. Then, (λ, μ) covers (λ', μ') if and only if one of the following statements is true:

- (1) $\mu = \mu'$, λ covers λ' with respect to the dominance order on partitions with $\lambda'_i = \lambda_i - 1$, and for k such that $\lambda'_k = \lambda_k + 1$, we have $\mu_{i-1} = \mu_i = \dots = \mu_k$;
- (2) $\lambda = \lambda'$, μ covers μ' with respect to the dominance order on partitions, with $\mu'_i = \mu_i - 1$, and for k such that $\mu'_k = \mu_k + 1$, we have $\lambda_i = \lambda_{i+1} = \dots = \lambda_{k+1}$;
- (3) $\lambda \neq \lambda'$, $\mu \neq \mu'$ and $\lambda_i > \lambda'_i$. If k is maximal with $\lambda_i = \lambda_k$, then $\mu_i = \mu_k$, $(\lambda'_j, \mu'_j) = (\lambda_j - 1, \mu_j + 1)$ for any integer $i \leq j \leq k$, and $(\lambda'_j, \mu'_j) = (\lambda_j, \mu_j)$ otherwise;
- (4) $\lambda \neq \lambda'$, $\mu \neq \mu'$, $\lambda_i = \lambda'_i$ (and therefore $\mu_i > \mu'_i$). If k is maximal with $\mu_i = \mu_k$, then $\lambda_{i+1} = \lambda_{k+1}$, $(\mu'_j, \lambda'_{j+1}) = (\mu_j - 1, \lambda_{j+1} + 1)$ for any integer $i \leq j \leq k$ and there is equality otherwise.

The example below shows instances for all the covering cases of bipartitions. The boxes that are moved are colored in red.

Example 4.4. • An example of a covering of type (1), where $i = 2$ and $k = 4$:

$$\left(\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \color{red}{\square} & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right), \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) \triangleright \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right), \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right).$$

- An example of a covering of type (2), where $i = 1$ and $k = 4$:

$$\left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right) \triangleright \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \square \\ \hline \end{array} \right).$$

- An example of a covering of type (3), where $i = 2$:

$$\left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right) \triangleright \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \color{red}{\square} \\ \hline \end{array} \right).$$

- An example of a covering of type (4), where $i = 1$:

$$\left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \color{red}{\square} & \color{red}{\square} \\ \hline \square & \color{red}{\square} & \color{red}{\square} \\ \hline \square & \color{red}{\square} & \color{red}{\square} \\ \hline \square & \color{red}{\square} & \color{red}{\square} \\ \hline \square & \color{red}{\square} & \color{red}{\square} \\ \hline \end{array} \right) \triangleright \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right).$$

One can think of the cases (3) and (4) as moving a *partial column* from the left diagram in the bidiagram (λ, μ) to the right side and staying in the same row, or from moving a partial column from the right diagram to the left and going down one row.

Now we present a proof of the theorem.

Proof of Theorem 4.3. We start the proof by showing that operations (1) – (4) define covering relations. Suppose that (λ', μ') is obtained from (λ, μ) by one of these operations. We need to show that (λ, μ) covers (λ', μ') , that is, if (λ^*, μ^*) is such that $(\lambda, \mu) \triangleright (\lambda^*, \mu^*) \triangleright (\lambda', \mu')$, then either $(\lambda^*, \mu^*) = (\lambda, \mu)$ or $(\lambda^*, \mu^*) = (\lambda', \mu')$. Since the proofs for operations (2) and (4) are respectively similar to (1) and (3), we will focus on these two operations.

- (A) Suppose that (λ', μ') is obtained from (λ, μ) by operation (1). In particular, $\mu = \mu'$, and there exists $1 < i < k$ such that $\lambda'_i = \lambda_i - 1$, $\lambda'_k = \lambda_k + 1$, $\lambda'_j = \lambda_j$ for $j \neq i, k$ and $\mu_{i-1} = \dots = \mu_k$. It is not difficult to show, by taking the difference between two consecutive partial sums, that

$$\forall j \in \{1, \dots, i-1, k+1, \dots\}, \lambda_j = \lambda_j^* = \lambda'_j \text{ and } \mu_j = \mu_j^* = \mu'_j,$$

as well as $\mu_k^* = \mu_k = \mu'_k$. Since $\mu_{i-1}^* = \mu_{i-1} = \mu_k = \mu_k^*$, this implies that $\mu_{i-1}^* = \dots = \mu_k^*$ as well. In turn, this means that

$$\bar{\lambda} = (\lambda_i, \dots, \lambda_k) \triangleright \bar{\lambda}^* = (\lambda_i^*, \dots, \lambda_k^*) \triangleright (\lambda'_i, \dots, \lambda'_k) = \bar{\lambda}'.$$

By hypothesis, $\bar{\lambda}$ covers $\bar{\lambda}'$, so that either $\bar{\lambda}^* = \bar{\lambda}$ or $\bar{\lambda}^* = \bar{\lambda}'$. In turn, this shows that (λ, μ) covers (λ', μ') .

- (B) Now, we assume that (λ', μ') is obtained from (λ, μ) by operation (3). This means, that there exists $i \leq k$ such that $\lambda_i = \lambda_k > \lambda_{k+1}$, $\mu_{i-1} > \mu_i = \mu_k$ and $\lambda'_j = \lambda_j - 1$, $\mu'_j = \mu_j + 1$, for $i \leq j \leq k$ and otherwise $\lambda'_j = \lambda_j$ and $\mu'_j = \mu_j$.

As above we observe that

$$\forall j < i \text{ or } j > k, \lambda_j = \lambda_j^* = \lambda'_j \text{ and } \mu_j = \mu_j^* = \mu'_j.$$

In the same way, it is easy to show that

$$\forall i \leq j \leq k, \lambda_j + \mu_j \geq \lambda_j^* + \mu_j^* \geq \lambda'_j + \mu'_j = \lambda_j + \mu_j,$$

and

$$\forall i \leq j \leq k, \lambda_j \geq \lambda_j^* \geq \lambda'_j = \lambda_j - 1.$$

Together, this implies that

$$\forall i \leq j \leq k, \lambda_j^* \in \{\lambda_j, \lambda_j - 1\} \text{ and } \mu_j^* \in \{\mu_j, \mu_j + 1\}.$$

- (a) Assume first that $\lambda_i^* = \lambda_i - 1$. Then for $i \leq j \leq k$,

$$\lambda_j - 1 \leq \lambda_j^* \leq \lambda_i^* = \lambda_i - 1 = \lambda_j - 1$$

which implies $\lambda_j^* = \lambda_j - 1$ and $\mu_j^* = \mu_j + 1$, that is, $(\lambda^*, \mu^*) = (\lambda', \mu')$.

(b) On the other hand, if $\lambda_i^* = \lambda_i$ or equivalently $\mu_i^* = \mu_i$, then for $i \leq j \leq k$, we have

$$\mu_j \leq \mu_j^* \leq \mu_i^* = \mu_i = \mu_j$$

which implies that $\mu_j^* = \mu_j$ and $\lambda_j^* = \lambda_j$, that is $(\lambda^*, \mu^*) = (\lambda, \mu)$.

Thus, they describe a covering relation in the poset $(\text{BP}_n, \trianglelefteq)$.

Now, we prove the converse. Let (λ, μ) and (λ', μ') be two different bipartitions of n and assume that $(\lambda', \mu') \trianglelefteq (\lambda, \mu)$. We show that there exists a bipartition (λ^*, μ^*) of n that can be obtained from (λ, μ) through one of the cases (1)-(4), and $(\lambda', \mu') \trianglelefteq (\lambda^*, \mu^*) \trianglelefteq (\lambda, \mu)$. Let $i \in \mathbb{N}$ be minimal with $(\lambda_i, \mu_i) \neq (\lambda'_i, \mu'_i)$.

(A) We consider first the case where $\lambda_i > \lambda'_i$ and show that we can obtain (λ^*, μ^*) using one of the operations (1),(3), or (4). Let $k \in \mathbb{N}$ be maximal with $\lambda_i = \lambda_k$.

We begin our analysis with distinguishing between the following two cases. Either there exists a $p \in \mathbb{N}$ such that $i \leq p \leq k$ and $\mu_p < \mu_{p-1}$ or not, with the convention that $\mu_1 < \mu_0$.

(a) First, we assume that there exists such a p and we fix the minimal p with this property. Then μ_p is the first place after μ_{i-1} where we can put a box to still obtain a partition. Let $q \in \mathbb{N}$ be minimal such that $p \leq q \leq k$ and $\mu_q = \dots = \mu_k$. We define (λ^*, μ^*) as the bipartition of n with $(\lambda_j^*, \mu_j^*) = (\lambda_j - 1, \mu_j + 1)$ for every $q \leq j \leq k$ and otherwise $(\lambda_j^*, \mu_j^*) = (\lambda_j, \mu_j)$. We observe easily that $(\lambda^*, \mu^*) \trianglelefteq (\lambda, \mu)$ and $\lambda^* \uplus \mu^* = \lambda \uplus \mu \triangleright \lambda' \uplus \mu'$, and we are just left with verifying

$$\lambda_t^* + \sum_{j=1}^{t-1} (\lambda_j^* + \mu_j^*) \geq \lambda'_t + \sum_{j=1}^{t-1} (\lambda'_j + \mu'_j),$$

for any $t \in \mathbb{N}$. However, this is clear for any $t < q$ and $t > k$. If $q \leq t \leq k$, we have

$$\lambda'_t \leq \lambda'_i \leq \lambda_i - 1 = \lambda_t - 1 = \lambda_t^*$$

and

$$\sum_{j=1}^{t-1} (\lambda_j^* + \mu_j^*) = \sum_{j=1}^{t-1} (\lambda_j + \mu_j) \geq \sum_{j=1}^{t-1} (\lambda'_j + \mu'_j)$$

so that we also have

$$\lambda_t^* + \sum_{j=1}^{t-1} (\lambda_j^* + \mu_j^*) \geq \lambda'_t + \sum_{j=1}^{t-1} (\lambda'_j + \mu'_j).$$

This is operation (3).

(b) Next, we assume that no such p exists. In particular, $i > 1$ and $\mu_{i-1} = \dots = \mu_k$. We consider the closest possible free place in the bidiagram (λ, μ) , namely we take $r > k$ to be the minimal integer with $\lambda_r < \lambda_k - 1$ or $(\lambda_r = \lambda_k - 1$ and $\mu_r < \mu_k)$. Such a r always exists, since we allow ourselves to extend the partitions with empty rows. If it did not exist, that would mean that $\lambda_k = 1$ and $\mu_k = 0$. By definition of k and i , this would mean that $\lambda'_j = \lambda_j$ for $1 \leq j < i$, $\mu'_j = \mu_j$ for $1 \leq j < i$, that $1 = \lambda_k = \dots = \lambda_i > \lambda'_i = 0$. Also, we have $0 = \mu_k = \dots = \mu_{i-1} = \mu'_{i-1}$. That means that the sizes of the bipartitions (λ, μ) and (λ', μ') are different, which is absurd. We proceed again with a case distinction.

(i) Let us start with assuming that $\lambda_r = \lambda_k - 1$ and $\mu_r < \mu_k$. We define $\mu_j^* = \mu_j - 1$ and $\lambda_{j+1}^* = \lambda_{j+1} + 1$ for all $k \leq j \leq r - 1$, while $\mu_j^* = \mu_j$ and $\lambda_{j+1}^* = \lambda_{j+1}$ for any other $j \in \mathbb{N}_0$. Since $\lambda_{k+1} < \lambda_k$ and $\mu_{r-1} = \mu_k > \mu_r$, (λ^*, μ^*) is a bipartition. Clearly $(\lambda^*, \mu^*) \trianglelefteq (\lambda, \mu)$. By construction we have

$$\lambda_t^* + \sum_{j=1}^{t-1} (\lambda_j^* + \mu_j^*) = \lambda_t + \sum_{j=1}^{t-1} (\lambda_j + \mu_j) \geq \lambda'_t + \sum_{j=1}^{t-1} (\lambda'_j + \mu'_j)$$

for any $t \in \mathbb{N}$. Also, for $t < k$ or $t \geq r$, we have

$$\sum_{j=1}^t (\lambda_j + \mu_j) = \sum_{j=1}^t (\lambda_j^* + \mu_j^*)$$

Thus, it remains to show this inequality holds for $k \leq t < r$. The following inequalities follow from the definitions of k , i and r :

$$\begin{aligned} \lambda_k^* &= \lambda_k = \lambda_i \geq \lambda'_i + 1 \geq \lambda'_k + 1, \\ \lambda_t^* &= \lambda_t + 1 = \lambda_k - 1 + 1 \geq \lambda'_k + 1 \geq \lambda'_t + 1 \quad \text{for } k < t \leq r \\ \mu_t^* + 1 &= \mu_t = \mu_k = \mu_{i-1} = \mu'_{i-1} \geq \mu'_t \quad \text{for } k \leq t < r. \end{aligned}$$

This shows that $\lambda^* \uplus \mu^* \supseteq \lambda' \uplus \mu'$, and we obtain (λ^*, μ^*) from (λ, μ) by operation (4).

(ii) Finally, we assume that $\lambda_r < \lambda_k - 1$. In particular, $\mu_{i-1} = \dots = \mu_{r-1}$ and $\lambda_{r-1} = \dots = \lambda_{k+1} = \lambda_k - 1 = \dots = \lambda_i - 1$. We distinguish between two cases.

- First, assume that $\mu_{r-1} > \mu_r$. We define $\lambda_r^* = \lambda_r + 1 \leq \lambda_{r-1}$ and $\mu_{r-1}^* = \mu_{r-1} - 1 \geq \mu_r$, while $\mu_j^* = \mu_j$ and $\lambda_j^* = \lambda_j$ otherwise. Then (λ^*, μ^*) is a well-defined bipartition, and as usual, $(\lambda^*, \mu^*) \trianglelefteq (\lambda, \mu)$. Also, proving that $(\lambda', \mu') \trianglelefteq (\lambda^*, \mu^*)$ is straightforward, except maybe proving that

$$\sum_{j=1}^{r-1} (\lambda_j^* + \mu_j^*) \geq \sum_{j=1}^{r-1} (\lambda'_j + \mu'_j).$$

But as previously, we have:

$$\begin{aligned} \lambda_j^* &= \lambda_j = \lambda'_j && \text{for } 1 \leq j < i \\ \mu_j^* &= \mu_j = \mu'_j && \text{for } 1 \leq j < i \\ \lambda_j^* &= \lambda_j = \lambda_i \geq \lambda'_i + 1 \geq \lambda'_j + 1 && \text{for } i \leq j \leq k \\ \lambda_j^* &= \lambda_j = \lambda_k - 1 = \lambda_i - 1 \geq \lambda'_i \geq \lambda'_j && \text{for } k < j \leq r - 1 \\ \mu_j^* &= \mu_j = \mu_{i-1} = \mu'_{i-1} \geq \mu'_j && \text{for } i \leq j \leq r - 2 \\ \mu_{r-1}^* &= \mu_{i-1} - 1 = \mu'_{i-1} - 1 \geq \mu'_{r-1} - 1 \end{aligned}$$

All together, this gives

$$\sum_{j=1}^{r-1} (\lambda_j^* + \mu_j^*) \geq \sum_{j=1}^{r-1} (\lambda'_j + \mu'_j) + (k - i) \geq \sum_{j=1}^{r-1} (\lambda'_j + \mu'_j)$$

as wanted. This also means that we obtain (λ^*, μ^*) from (λ, μ) by operation (4).

- We are left with the case $\mu_r = \mu_{r-1}$. By a previous remark, this means that $\mu_{i-1} = \dots = \mu_r$. We set $\mu^* = \mu$ and $\lambda_k^* = \lambda_k - 1$, $\lambda_r^* = \lambda_r + 1$, and otherwise $\lambda_j^* = \lambda_j$. Then, by assumption (λ^*, μ^*) is a bipartition and $(\lambda^*, \mu^*) \trianglelefteq (\lambda, \mu)$. It is also straightforward to show that $(\lambda^*, \mu^*) \supseteq (\lambda', \mu')$ except maybe the partial sums inequalities in rows k to r . Since $\lambda_k^* + \lambda_r^* = \lambda_k + \lambda_r$, we only need to look at rows k to $r - 1$. To this purpose, we remark that:

$$\begin{aligned} \lambda_j^* &= \lambda_j = \lambda'_j && \text{for } 1 \leq j < i \\ \mu_j^* &= \mu_j = \mu'_j && \text{for every } j \\ \lambda_j^* &= \lambda_j = \lambda_i \geq \lambda'_i + 1 \geq \lambda'_j + 1 && \text{for } i \leq j < k \\ \lambda_k^* &= \lambda_k - 1 = \lambda_i - 1 \geq \lambda'_i \geq \lambda'_k && \\ \lambda_j^* &= \lambda_j = \lambda_k - 1 = \lambda_i - 1 \geq \lambda'_i \geq \lambda'_j && \text{for } k < j \leq r - 1 \end{aligned}$$

Then for any $k \leq j < r$, we have

$$\sum_{t=1}^j (\lambda_t^* + \mu_t^*) \geq \sum_{t=1}^j (\lambda'_t + \mu'_t) + (k - i) \geq \sum_{t=1}^j (\lambda'_t + \mu'_t)$$

and in the same way

$$\lambda_j^* + \sum_{t=1}^{j-1} (\lambda_t^* + \mu_t^*) \geq \lambda_j' + \sum_{t=1}^{j-1} (\lambda_t' + \mu_t').$$

Thus we obtain (λ^*, μ^*) from (λ, μ) by operation (1).

- (B) It remains to deal with the case $\lambda_i = \lambda_i'$, and $\mu_i > \mu_i'$. It can easily be deduced from the previous case, by noticing the following: let $\rho = (\mu_1, \mu_1, \mu_2, \dots)$ and $\rho' = (\mu_1, \mu_1', \mu_2', \dots)$, then (ρ, λ) and (ρ', λ') are bipartitions of $n + \mu_1$ such that $(\rho, \lambda) \succeq (\rho', \lambda')$. If j is minimal such that $(\rho_j, \lambda_j) \neq (\rho_j', \lambda_j')$, then $j = i + 1$ and $\rho_j = \mu_i > \mu_i' = \rho_j'$. From what we have just seen, there exists (ρ^*, λ^*) , obtained from (ρ, λ) by operations (1), (3) and (4), such that $(\rho, \lambda) \succeq (\rho^*, \lambda^*) \succeq (\rho', \lambda')$. It is then clear that $\rho_1^* = \rho_1 = \rho_1' = \mu_1$. Let $\mu^* = (\rho_2^*, \rho_3^*, \dots)$. It is clear that (λ^*, ρ^*) is a bipartition and obviously $(\lambda, \mu) \succeq (\lambda^*, \mu^*) \succeq (\lambda', \mu')$. Moreover if we obtained (μ^*, ρ^*) from (μ, ρ) by operations (1), (3) or (4) respectively, we obtain (λ^*, μ^*) from (λ, μ) by operations (2), (4) and (3) respectively.

□

It is in general not true that if λ is a partition covering λ' , then (λ, μ) covers (λ', μ) , as the following example shows:

Example 4.5. Consider $(\lambda, \mu) = ((3, 3, 2, 1), (2, 2, 2, 1))$ and $(\lambda', \mu') = ((3, 2, 2, 2), (2, 2, 2, 1))$. Indeed, it is $\mu_1 = \mu_2 = \mu_3 > \mu_4$. Thus, there exist bipartitions which lie in between.

Since the poset of partitions for the standard order is a lattice, it is natural to ask whether this holds for our order on bipartitions. However, this is not the case already for $n = 4$. Consider $a = ((2), (1, 1))$ and $b = ((2, 2), \emptyset)$. Now take $c = ((2, 1, 1), \emptyset)$. It is covered by both a and b , so if a and b have a meet, that is a greatest lower bound, it has to be c . However, for $d = (\emptyset, (2, 2))$, $d < a$ and $d < b$, but c and d are not comparable. Similarly, one can ask if the poset (BP_n, \preceq) is graded, i.e., any maximal chain has equal length. However, already for $n = 3$ the poset is non-graded since there exist maximal chains of length 6 and 7.

5. THE POSETS OF SPECHT IDEALS AND VARIETIES

In this section, we state and prove our main theorem:

Theorem 5.1. Let (λ, μ) and (θ, ω) be bipartitions of n . Let $I_{(\lambda, \mu)}, I_{(\theta, \omega)}$ denote their associated Specht ideals and $V_{(\lambda, \mu)}, V_{(\theta, \omega)}$ their associated Specht varieties. Then, the following assertions are equivalent:

- (1) The bipartition (λ, μ) bidominates (θ, ω) , i.e. $(\lambda, \mu) \succeq (\theta, \omega)$;
- (2) The B_n -Specht ideal $I_{(\lambda, \mu)}$ contains the B_n -Specht ideal $I_{(\theta, \omega)}$, i.e. $I_{(\lambda, \mu)} \supset I_{(\theta, \omega)}$;
- (3) The B_n -Specht variety $V_{(\lambda, \mu)}$ is contained in the B_n -Specht variety $V_{(\theta, \omega)}$, i.e. $V_{(\lambda, \mu)} \subset V_{(\theta, \omega)}$.

We start with the first implication, namely that a dominance of bipartitions implies the containment of the corresponding B_n -Specht ideals.

Proposition 5.2. Let $(\lambda, \mu), (\lambda', \mu') \in \text{BP}_n$ be bipartitions of n and let $(\lambda', \mu') \preceq (\lambda, \mu)$. Then, $I_{(\lambda', \mu')} \subset I_{(\lambda, \mu)}$.

Proof. It is sufficient to prove the theorem in the four covering cases in 4.3.

In cases (1) and (2), we have in particular $(\lambda' \preceq \lambda$ and $\mu' = \mu)$ or $(\mu' \preceq \mu$ and $\lambda' = \lambda)$, and the result follows from the proof of ([MRV21, Theorem 1]), combined with the definition of B_n -Specht polynomials.

Now, we consider case (3). In this case, to go from (λ', μ') to (λ, μ) , we remove a number a of boxes from a column U_1 in μ' , that will be added to a column U_2 in λ' . We can restrict

our attention to these two columns. Let $b = |U_2|$, we then have $|U_1| = a + |U_2| = a + b$. Let $A = \{1, \dots, a\}$, $B_1 = \{a + 1, \dots, a + b\}$ and $B_2 = \{a + b + 1, \dots, a + 2b\}$. Up to permutation, it suffices to show that the polynomial

$$P(\mathbf{x}) = \Delta_{B_2}(\mathbf{x}^2) \Delta_{A \cup B_1}(\mathbf{x}^2) \prod_{i \in A \cup B_1} \mathbf{x}_i$$

is in the ideal generated by polynomials of the form $\Delta_S(\mathbf{x}^2) \Delta_{\bar{S}}(\mathbf{x}^2) \prod_{i \in \bar{S}} \mathbf{x}_i$, where $\{1, \dots, a + 2b\}$ is the disjoint union of S and \bar{S} , and $|S| = a + b$. Let us consider

$$Q(\mathbf{x}) = \Delta_{A \cup B_2}(\mathbf{x}^2) \Delta_{B_1}(\mathbf{x}^2) \prod_{i \in B_1} \mathbf{x}_i,$$

which is a polynomial of the expected form, and

$$\begin{aligned} \tilde{Q}(\mathbf{x}) &= Q(\mathbf{x}) \prod_{i \in A} \mathbf{x}_i \\ &= \Delta_{A \cup B_2}(\mathbf{x}^2) \Delta_{B_1}(\mathbf{x}^2) \prod_{i \in A \cup B_1} \mathbf{x}_i, \end{aligned}$$

Note that we have:

$$\deg(P) = b(b - 1) + (a + b)(a + b - 1) + a + b = 2b^2 + a^2 + b(2a - 1)$$

and

$$\deg(Q) = (a + b)(a + b - 1) + b(b - 1) + b = 2b^2 + a^2 + b(2a - 1) - a$$

so that P and \tilde{Q} have the same degree. We are going to show that P is a combination of $\epsilon(\sigma)\sigma \cdot \tilde{Q}$ for σ 's in $G = \mathcal{S}_{A \cup B_1}$. Note that we can rewrite

$$\Delta_{A \cup B_2}(\mathbf{x}^2) = \Delta_A(\mathbf{x}^2) \Delta_{B_2}(\mathbf{x}^2) \prod_{i \in A} R(\mathbf{x}_i^2)$$

where

$$R(y) = \prod_{j \in B_2} (y - \mathbf{x}_j^2).$$

Then, since $\prod_{i \in A \cup B_1} \mathbf{x}_i$ and $\Delta_{B_2}(\mathbf{x}^2)$ are invariant by G , we can factor them out and focus on the remaining terms, then looking at

$$P^*(\mathbf{x}) = \Delta_{A \cup B_1}(\mathbf{x}^2)$$

and

$$Q^*(\mathbf{x}) = \Delta_A(\mathbf{x}^2) \Delta_{B_1}(\mathbf{x}^2) \prod_{i \in A} R(\mathbf{x}_i^2).$$

Also consider the subgroup $H = \mathcal{S}_A \times \mathcal{S}_{B_1}$ of G . Then, for $\tau_1 \in \mathcal{S}_A$, $\tau_2 \in \mathcal{S}_{B_1}$, we have $\tau_1 \tau_2 (\Delta_A(\mathbf{x}^2)) = \epsilon(\tau_1) (\Delta_A(\mathbf{x}^2))$ and $\tau_1 \tau_2 (\Delta_{B_1}(\mathbf{x}^2)) = \epsilon(\tau_2) (\Delta_{B_1}(\mathbf{x}^2))$, and because $\prod_{i \in A} R(\mathbf{x}_i^2)$ is H -invariant, we get

$$\epsilon(\tau_1 \tau_2) \tau_1 \tau_2 Q^* = Q^*,$$

allowing us to consider the sum

$$\bar{Q} = \sum_{\sigma \in G/H} \epsilon(\sigma) \sigma Q^*,$$

and we claim that $P^* = \bar{Q}$.

First, we show that P^* divides \bar{Q} , namely that for every $i \neq j \in A \cup B_1$, $\mathbf{x}_i^2 - \mathbf{x}_j^2$ divides \bar{Q} . Since for every $\sigma \in G$, $\sigma \bar{Q} = \pm \bar{Q}$, and G acts transitively on pairs (i, j) , it is enough to check that $\mathbf{x}_1^2 - \mathbf{x}_2^2$ divides \bar{Q} . We hence have to show that \bar{Q} vanishes when imposing $\mathbf{x}_1^2 = \mathbf{x}_2^2$. To see this, first observe that the terms in the sum are in correspondence with set partitions $K \cup \bar{K}$ of $A \cup B_1$, where $|K| = a$. Indeed, up to permutation by elements of H , we only need to choose where to send the subset $A = \{1, \dots, a\}$. Now, if σ sends 1 and 2 in the same subset, the corresponding Vandermonde determinant in $\epsilon(\sigma) \sigma Q^*$ vanishes whenever $\mathbf{x}_1^2 = \mathbf{x}_2^2$. We then only need to focus on partitions where 1 and 2 are not in the same subset. There are two kinds of such partitions: those with $1 \in K$ and $2 \in \bar{K}$, and those with $2 \in K$ and $1 \in \bar{K}$. The transposition (12)

naturally induces a bijection between these sets of partitions. If $\sigma \in G$ is a representative for a partition of the first kind, then $(12)\sigma$ is a representative for the corresponding partition of the second kind. When $\mathbf{x}_1^2 = \mathbf{x}_2^2$, we have $(12)\sigma Q^*(x) = \sigma Q^*(x)$, and because $\epsilon((12)\sigma) = -\epsilon(\sigma)$, the two corresponding terms cancel out.

Then, we need to check that P^* and \bar{Q} have the same leading term with respect to the lexicographical ordering. The leading term of P^* is $\mathbf{x}_1^{2(a+b-1)}\mathbf{x}_2^{2(a+b-2)}\cdots\mathbf{x}_{a+b-1}^2$. Then, for $\sigma \in G$ sending $\{1, \dots, k\}$ onto K , the partial degree of σQ^* in \mathbf{x}_1 is

$$\begin{cases} 2(a-1) + 2b & \text{if } 1 \in K \\ 2(b-1) & \text{if } 1 \notin K \end{cases}$$

and therefore σQ^* can give a contribution to the leading term of \bar{Q} only if $1 \in K$. By the same argument, 2 has to be in K , and in the end, $K = \{1, \dots, a\}$: Indeed, assume there is a minimal $i \leq a$ with $i \notin K$. Then, the leading term of σQ^* is of the form $\mathbf{x}_1^{2(a+b-1)}\mathbf{x}_2^{2(a+b-2)}\cdots\mathbf{x}_{j-1}^{2(a+b-j+1)}\mathbf{x}_j^{2(b-1)}m$ where m is a monomial in the variables $\mathbf{x}_{j+1}, \dots, \mathbf{x}_{a+b}$. Since, $j \leq a$, then $2(b-1) < 2(a+b-j)$, and therefore the leading monomial of σQ^* is strictly lower than that of Q^* . Thus, the leading term of \bar{Q} is exactly the leading term of Q^* , which is $\mathbf{x}_1^{2(a+b-1)}\mathbf{x}_2^{2(a+b-2)}\cdots\mathbf{x}_{a+b-1}^2$, as expected. This concludes the covering case (3).

The proof for the covering case (4) is very similar. In this situation, to go from (λ', μ') to (λ, μ) , we remove a boxes from a column U_1 in λ' , before adding them to a column U_2 in μ' , with $|U_2| = b$ and $|U_1| = a + b + 1$. We can apply the previous argument, where this time $A = \{1, \dots, a\}$, $B_1 = \{a + 1, \dots, a + b + 1\}$, $B_2 = \{a + b + 2, \dots, a + 2b + 1\}$,

$$P = \Delta_{A \cup B_1}(\mathbf{x}^2)\Delta_{B_2}(\mathbf{x}^2) \prod_{i \in B_2} \mathbf{x}_i$$

$$Q = \Delta_{B_1}(\mathbf{x}^2)\Delta_{A \cup B_2}(\mathbf{x}^2) \prod_{i \in A \cup B_2} \mathbf{x}_i$$

and

$$\begin{aligned} \tilde{Q} &= \left(\prod_{j \in A} \mathbf{x}_j \right) Q \\ &= \Delta_{B_1}(\mathbf{x}^2)\Delta_{A \cup B_2}(\mathbf{x}^2) \prod_{i \in A} \mathbf{x}_i^2 \prod_{i \in B_2} \mathbf{x}_i. \end{aligned}$$

□

The second implication of 5.1 is clear, it remains to prove that (3) implies (1):

Proposition 5.3. *Let $(\lambda, \mu), (\theta, \omega)$ be bipartitions and $V_{(\lambda, \mu)} \subset V_{(\theta, \omega)}$. Then, $(\lambda, \mu) \succeq (\theta, \omega)$.*

To prove this implication, we will consider two types of points in \mathbb{K}^n :

Lemma 5.4. *Let $(\theta, \omega) \in \text{BP}_n$ and $\Lambda = \theta \uplus \omega$ be a partition of n . Consider the point*

$$z = \left(\underbrace{a_1, \dots, a_1}_{\Lambda_1}, \underbrace{a_2, \dots, a_2}_{\Lambda_2}, \dots, \underbrace{a_m, \dots, a_m}_{\Lambda_m} \right)$$

with $a_i^2 \neq a_j^2$ if $i \neq j$ and $a_i \neq 0$ if $i \leq \text{len}(\omega)$.

- i) $z \notin V_{(\theta, \omega)}$.
- ii) If $(\lambda, \mu) \in \text{BP}_n$ is a bipartition such that $z \in V_{(\lambda, \mu)}$, then $\lambda \uplus \mu \succeq \Lambda$.

Proof. i) Let (T, S) be the generalized bitableau of shape (θ, ω) which has the filling

$$\left(\begin{array}{cccc} \boxed{a_1} & \boxed{a_1} & \cdots & \boxed{a_1} \\ \boxed{a_2} & \boxed{a_2} & \cdots & \\ \vdots & & & \\ \vdots & & & \\ \boxed{a_h} & \cdots & & \end{array} , \begin{array}{cccc} \boxed{a_1} & \boxed{a_1} & \cdots & \cdots & \boxed{a_1} \\ \boxed{a_2} & \boxed{a_2} & \cdots & \cdots & \\ \vdots & & & & \\ \boxed{a_l} & \cdots & & & \end{array} \right).$$

i.e., the i -th row of both T and S contains only a_i 's. The assumption $a_i \neq 0$ for $i \leq \text{len}(\omega)$ ensures that S contains no 0 entry, and by construction the squares of column entries are pairwise different. Thus, $z \in V_{(\theta, \omega)}^c$.

- ii) By assumption, there is a bitableau (T, S) of shape (λ, μ) such that $\text{sp}_{(T, S)}(z) \neq 0$. Then, according to Lemma 3.7, there is a tableau $U = T \uplus S$ of shape $\lambda \uplus \mu$ such that $\text{sp}_U(z^2) \neq 0$. Therefore z^2 does not belong to the \mathcal{S}_n -Specht variety $V_{\lambda \uplus \mu}$, and ([MRV21, Prop 1.ii]) gives

$$\Lambda = \Lambda(z) = \Lambda(z^2) \leq \lambda \uplus \mu,$$

which proves the Lemma. □

Lemma 5.5. *Let $(\theta, \omega) \in \text{BP}_n$ be a bipartition of n . Consider the point*

$$z = \left(\underbrace{0, \dots, 0}_{\theta_1}, \underbrace{a_1, \dots, a_1}_{\omega_1 + \theta_2}, \underbrace{a_2, \dots, a_2}_{\omega_2 + \theta_3}, \dots, \underbrace{a_m, \dots, a_m}_{\omega_m + \theta_{m+1}} \right)$$

with $a_i^2 \neq a_j^2$ if $i \neq j$ and $a_i \neq 0$. Then:

- i) $z \notin V_{(\theta, \omega)}$.
ii) If $(\lambda, \mu) \in \text{BP}_n$ is a bipartition such that $z \notin V_{(\lambda, \mu)}$, then

$$\sum_{j=1}^{k-1} (\lambda_j + \mu_j) + \lambda_k \geq \sum_{j=1}^{k-1} (\theta_j + \omega_j) + \theta_k$$

for any integer $k \geq 1$.

Proof. i) Let (T, S) be the generalized bitableau of shape (θ, ω) which has the filling

$$\left(\begin{array}{cccc} \boxed{0} & \boxed{0} & \cdots & \boxed{0} \\ \boxed{a_1} & \boxed{a_1} & \cdots & \\ \vdots & & & \\ \boxed{a_h} & \cdots & & \end{array} , \begin{array}{cccc} \boxed{a_1} & \boxed{a_1} & \cdots & \boxed{a_1} \\ \boxed{a_2} & \boxed{a_2} & \cdots & \\ \vdots & & & \\ \vdots & & & \\ \boxed{a_l} & \cdots & & \end{array} \right)$$

i.e., θ_i contains only a_{i-1} 's and ω_i contains only a_i 's, where $a_0 = 0$. We observe that no entry in S equals 0 and the squares of column entries are pairwise different. Thus, we have $z \in V_{(\theta, \omega)}^c$.

- ii) By assumption, there exists a bitableau (T, S) of shape (λ, μ) such that

$$0 \neq \text{sp}_{(T, S)}(z) = \text{sp}_T(z^2) \text{sp}_S(z^2) \cdot \prod_{j \in S} z_j.$$

Let (T^*, S^*) be the generalized bitableau obtained from (T, S) by replacing i with a_i in any box. This means that the zeros of z are written in T^* and no column in T^* or S^* contains entries with equal squares. Since permutation of the column entries can only change the sign of $\text{sp}_{(T, S)}(z)$, we can assume that the entries in every column in (T^*, S^*) are sorted increasingly by the indices of the a_i 's from above to below, and with

$a_0 = 0$.

We obtain that all the 0's must be written in the first row of T^* which implies $\lambda_1 \geq \theta_1$. Now, for an integer $k \geq 1$ the a_k 's in (T^*, S^*) must be written in different columns in the generalized bitableau (T^*, S^*) . Since the entries in (T^*, S^*) are written with increasing indices in each column from the top to the bottom, we know that the a_j 's with $0 \leq j \leq k$ must be written within the first k rows of S and the first $k+1$ -rows in T . Thus, by the pigeon hole principle, we have

$$\sum_{j=1}^k (\lambda_j + \mu_j) + \lambda_{k+1} \geq \sum_{j=1}^k (\theta_j + \omega_j) + \theta_{k+1}.$$

□

Now, we can prove Proposition 5.3:

Proof of Proposition 5.3. The assumption is equivalent to $V_{(\theta, \omega)}^c \subset V_{(\lambda, \mu)}^c$. We have to prove that $\lambda \uplus \mu \supseteq \theta \uplus \omega$ and that $\sum_{j=1}^{k-1} (\lambda_j + \mu_j) + \lambda_k \geq \sum_{j=1}^{k-1} (\theta_j + \omega_j) + \theta_k$ for every integer $k \geq 1$.

For the first claim, consider the point

$$z = \underbrace{(a_1, \dots, a_1)}_{\theta_1 + \omega_1}, \underbrace{(a_2, \dots, a_2)}_{\theta_2 + \omega_2}, \dots, \underbrace{(a_m, \dots, a_m)}_{\theta_m + \omega_m}$$

with $a_i^2 \neq a_j^2$ for $i \neq j$, and $a_i \neq 0$ if $i \leq \text{len}(\omega)$. According to i) in Lemma 5.4, $z \in V_{(\theta, \omega)}^c$. By assumption, we then have $z \in V_{(\lambda, \mu)}^c$, and ii) in Lemma 5.4 gives

$$\lambda \uplus \mu \supseteq \Lambda(z) = \theta \uplus \omega.$$

For the second claim, consider the point

$$z = \underbrace{(0, \dots, 0)}_{\theta_1}, \underbrace{(a_1, \dots, a_1)}_{\omega_1 + \theta_2}, \underbrace{(a_2, \dots, a_2)}_{\omega_2 + \theta_3}, \dots, \underbrace{(a_m, \dots, a_m)}_{\omega_m + \theta_{m+1}}$$

with $a_i^2 \neq a_j^2$ for $i \neq j$, and $a_i \neq 0$. According to i) in Lemma 5.5, $z \in V_{(\theta, \omega)}^c$. By assumption, we then have $z \in V_{(\lambda, \mu)}^c$, and ii) in Lemma 5.5 gives

$$\sum_{j=1}^{k-1} (\lambda_j + \mu_j) + \lambda_k \geq \sum_{j=1}^{k-1} (\theta_j + \omega_j) + \theta_k$$

for any integer $k \geq 1$.

□

6. ORBIT TYPES

In this section we define orbit types of elements in \mathbb{K}^n with respect to the action of the hyperoctahedral group. Compared with the \mathcal{S}_n -orbit types, they allow a finer set decomposition of \mathbb{K}^n since one distinguishes whether coordinates are 0 or not. This leads to a set partition of the B_n -Specht varieties based on the combinatorics of the poset $(\text{BP}_n, \trianglelefteq)$.

Recall that if $z = (a_1, \dots, a_n) \in \mathbb{K}^n$, then the \mathcal{S}_n -orbit type of z is the unique partition $\Lambda(z) = (\Lambda_1, \dots, \Lambda_l) \vdash n$ such that $\text{Stab}_{\mathcal{S}_n}(z) \simeq \mathbb{Z}/\Lambda_1\mathbb{Z} \times \dots \times \mathbb{Z}/\Lambda_l\mathbb{Z}$, or equivalently, there exists b_1, \dots, b_l pairwise distinct such that $z \in \mathcal{S}_n \cdot \underbrace{(b_1, \dots, b_1)}_{\Lambda_1}, \dots, \underbrace{(b_l, \dots, b_l)}_{\Lambda_l}$.

Definition 6.1. Let $\lambda = (\lambda_1, \dots, \lambda_m) \vdash n$ be a partition and $t \in \mathbb{N}_0$. Let $j = \min\{i : \lambda_i < t\}$ with the convention that $j = m+1$ if $t = 0$. Then the t -cut of λ is the bipartition (ρ, σ) defined as $\rho = (t, \dots, t, \lambda_j, \dots, \lambda_m)$ and $\sigma = (\lambda_1 - t, \dots, \lambda_{j-1} - t)$. We denote it by $\text{cut}(\lambda, t)$.

We have $\text{cut}(\lambda, 0) = (\emptyset, \lambda)$ while $\text{cut}(\lambda, t) = (\lambda, \emptyset)$ for any $t \geq \lambda_1$. We observe that if $\text{cut}(\lambda, t) = (\sigma, \rho)$, then $\sigma \uplus \rho = \lambda$.

We are now ready to define the B_n -orbit type of a point in \mathbb{K}^n and the notion of B_n -orbit set:

Definition 6.2. Let $z \in \mathbb{K}^n$. The B_n -orbit type of z is

$$\Omega(z) = \text{cut}(\Lambda(z^2), t_z)$$

where t_z is the number of 0 coordinates of z .

For $(\lambda, \mu) \in \text{BP}_n$, the B_n -orbit set associated to (λ, μ) is then

$$H_{(\lambda, \mu)} = \{z \in \mathbb{K}^n : \Omega(z) = (\lambda, \mu)\}.$$

We observe that B_n -orbits sets might be empty. The non-empty B_n -orbits sets correspond to bipartitions (λ, μ) such that $\lambda_1 = \dots = \lambda_{\text{len}(\mu)+1}$. Moreover, if (λ, μ) is such that $H_{(\lambda, \mu)} \neq \emptyset$, then any point $z \in H_{(\lambda, \mu)}$ is of the following form: let $m = \text{len}(\mu)$. Then there exists non-zero elements $a_1, \dots, a_l \in \mathbb{K}$, with distinct squares such that

$$z = \sigma \cdot \underbrace{(a_1, \dots, a_1)}_{\lambda_1 + \mu_1}, \dots, \underbrace{(a_m, \dots, a_m)}_{\lambda_1 + \mu_m}, \underbrace{(0, \dots, 0)}_{\lambda_1}, \underbrace{(a_{m+1}, \dots, a_{m+1})}_{\lambda_{m+2}}, \dots, \underbrace{(a_l, \dots, a_l)}_{\lambda_{l+1}}$$

for some $\sigma \in B_n$. It is straightforward that $l = m$ if $\text{len}(\lambda) \leq m$ (which implies $\lambda = \emptyset$) and $l = \text{len}(\lambda) - 1$ if $\text{len}(\lambda) > m$.

Example 6.3. We present the orbit types of \mathbb{K}^3 . Let $a, b, c \in \mathbb{K}^*$ be such that they have distinct squares.

z	$\Omega(z)$
$(0, 0, 0)$	$((3), \emptyset)$
$(\pm a, 0, 0)$	$((2, 1), \emptyset)$
$(\pm a, \pm b, 0)$	$((1, 1, 1), \emptyset)$
$(\pm a, \pm a, 0)$	$((1, 1), (1))$
$(\pm a, \pm a, \pm a)$	$(\emptyset, (3))$
$(\pm a, \pm a, \pm b)$	$(\emptyset, (2, 1))$
$(\pm a, \pm b, \pm c)$	$(\emptyset, (1, 1, 1))$

The remaining bipartitions $((2), (1)), ((1), (2)), ((1), (1, 1))$ of 3 have an empty orbit set.

The following proposition follows from the previous definitions and comments:

Proposition 6.4. Let $z \in \mathbb{K}^n$ and $(\lambda, \mu) = \Omega(z)$. Then $\Lambda(z^2) = \lambda \uplus \mu$. Moreover, the B_n -orbit sets define a set partition of \mathbb{K}^n , namely $\mathbb{K}^n = \uplus_{(\lambda, \mu) \in \text{BP}_n} H_{(\lambda, \mu)}$.

Proposition 6.5. Let $z \in \mathbb{K}^n$, and $(\lambda, \mu) \in \text{BP}_n$. Then:

- (1) $z \notin V_{\Omega(z)}$,
- (2) $z \notin V_{(\lambda, \mu)} \Rightarrow (\lambda, \mu) \supseteq \Omega(z)$.

Proof. Let $(\theta, \omega) = \Omega(z)$ and $m = \text{len}(\omega)$. Then there exists $\sigma \in B_n$ and $a_1, \dots, a_l \in \mathbb{K}^*$ with distinct squares such that $z = \sigma z'$ with

$$z' = \underbrace{(a_1, \dots, a_1)}_{\theta_1 + \omega_1}, \dots, \underbrace{(a_m, \dots, a_m)}_{\theta_1 + \omega_m}, \underbrace{(0, \dots, 0)}_{\lambda_1}, \underbrace{(a_{m+1}, \dots, a_{m+1})}_{\theta_{m+2}}, \dots, \underbrace{(a_l, \dots, a_l)}_{\theta_{l+1}}.$$

Since the Specht varieties are invariant under the action of B_n , we can assume that $z = z'$. With this shape, we can apply Lemma 5.4 to z which gives immediately (1), and partly (2): if $z \notin V(\lambda, \mu)$, then $\lambda \uplus \mu \supseteq \theta \uplus \omega$. It remains to prove that if $z \notin V(\lambda, \mu)$, then $\sum_{j=1}^{k-1} (\lambda_j + \mu_j) + \lambda_k \geq \sum_{j=1}^{k-1} (\theta_j + \omega_j) + \theta_k$ for any integer $k \geq 1$. To do so, it is enough to observe that since $\theta_1 = \dots = \theta_{m+1}$, the point

$$z'' = \underbrace{(0, \dots, 0)}_{\theta_1}, \underbrace{(a_1, \dots, a_1)}_{\omega_1 + \theta_2}, \underbrace{(a_2, \dots, a_2)}_{\omega_2 + \theta_3}, \dots, \underbrace{(a_m, \dots, a_m)}_{\omega_m + \theta_{m+1}}$$

is in the same orbit, and we can apply Lemma 5.5 to conclude the proof. \square

As a consequence of our previous results, we get a decomposition of B_n -Specht varieties in terms of orbit sets:

Theorem 6.6.

$$V_{(\lambda, \mu)} = \left(\bigcup_{(\theta, \omega) \in \text{BP}_n, (\theta, \omega) \trianglelefteq (\lambda, \mu)} H_{(\theta, \omega)} \right)^c = \bigcup_{(\theta, \omega) \in \text{BP}_n, (\theta, \omega) \not\trianglelefteq (\lambda, \mu)} H_{(\theta, \omega)}.$$

Proof. The collection $\{H_{(\theta, \omega)} : (\theta, \omega) \in \text{BP}_n\}$ defines a set partition of \mathbb{K}^n by definition, which explains the second equality.

In order to prove the first one, we first assume that $z \notin V_{(\lambda, \mu)}$. By part (2) in Proposition 6.5 we obtain that $(\lambda, \mu) \trianglerighteq \Omega(z)$. Thus $z \in \bigcup_{(\theta, \omega) \in \text{BP}_n, (\theta, \omega) \trianglelefteq (\lambda, \mu)} H_{(\theta, \omega)}$. Conversely, let $z \in \bigcup_{(\theta, \omega) \trianglelefteq (\lambda, \mu)} H_{(\theta, \omega)}$. In other words, $\Omega(z) \trianglelefteq (\lambda, \mu)$. Then, part (1) in Proposition 6.5 implies $z \notin V_{\Omega(z)}$. On the other hand, by Theorem 5.1, $V_{(\lambda, \mu)} \subset V_{\Omega(z)}$. Therefore $z \notin V_{(\lambda, \mu)}$. \square

Example 6.7. We calculate the Specht variety corresponding with the bipartition $((1, 1), (2))$ using Theorem 6.6. The bipartitions (θ, ω) encoding non-empty orbit sets such that $(\theta, \omega) \not\trianglelefteq ((1, 1), (2))$ are the bipartitions in

$$\Omega = \{((2, 2), \emptyset), ((2, 1, 1), \emptyset), (\emptyset, (4)), ((3, 1), \emptyset), ((4), \emptyset)\} \subset \text{BP}_4.$$

Then,

$$\begin{aligned} V_{((1, 1), (2))} &= \bigcup_{(\lambda, \mu) \in \Omega} H_{(\lambda, \mu)} \\ &= H_{((2, 2), \emptyset)} \cup H_{((2, 1, 1), \emptyset)} \cup H_{(\emptyset, (4))} \cup H_{((3, 1), \emptyset)} \cup H_{((4), \emptyset)}, \end{aligned}$$

which means

$$V_{((1, 1), (2))} = B_4 \cdot \{(0, 0, a, a), (0, 0, a, b), (a, a, a, a), (0, 0, 0, a), (0, 0, 0, 0) : a, b \in \mathbb{K}_{>0}\}.$$

One might look for a more natural orbit-type, only involving the number of zeroes of a point, and the \mathcal{S}_n -orbit-type of the remaining non-zero squared coordinates. Indeed, our previous decomposition can be reformulated in such a way, and it can be obtained either using S_n -invariance and results of [MRV21], or as a consequence of our previous results on bipartitions. We just briefly describe here the latter approach because such a point of view, even if it gives a natural decomposition, does not give information on inclusions of B_n -Specht varieties, which will be needed for our applications in the next section.

If $z = (a_1, \dots, a_n) \in \mathbb{K}^n$ is a point, then

$$\text{Stab}_{\mathcal{S}_n}(a_1^2, \dots, a_n^2) \simeq \mathcal{S}_{\Lambda_1} \times \dots \times \mathcal{S}_{\Lambda_l} \times \mathcal{S}_t$$

where $\Lambda_1 \geq \dots \geq \Lambda_l$ and t is the number of zero coordinates of z . We could have defined the orbit type of z as

$$\Lambda(z) = (t, (\Lambda_1, \dots, \Lambda_l)).$$

Then, there is a bijection φ between the set of pairs (t, Λ) where $n \geq t \geq 0$ and $\Lambda \vdash n - t$ and the set of bipartitions (λ, μ) such that $H_{(\lambda, \mu)} \neq \emptyset$ given by

$$(t, \Lambda) \mapsto \overline{\text{cut}((\Lambda_1, \Lambda_2, \dots, \Lambda_l, t), t)}$$

where $\overline{(\Lambda_1, \Lambda_2, \dots, \Lambda_l, t)}$ denotes the partition obtained by rearranging $(\Lambda_1, \Lambda_2, \dots, \Lambda_l, t)$ in non increasing order.

Now, for $t \leq 0$ and $\Lambda \vdash n - t$, if we denote by $G_{t, \Lambda} = \{z \in \mathbb{K}^n; \Lambda(z) = (t, \Lambda)\}$, we have by construction

$$G_{t, \Lambda} = H_{\varphi(t, \Lambda)}.$$

Moreover, φ preserves the orders in the following sense: for t fixed, $\varphi(t, \Lambda) \trianglelefteq \varphi(t, \Lambda')$ in our poset of bipartitions if and only if $\Lambda \trianglelefteq \Lambda'$ in the poset of partitions. Also, it is obvious that if $t > \lambda_1$, then $z \in V_{(\lambda, \mu)}$.

As a consequence, our decomposition in Theorem 6.6 becomes in this context

$$(V_{(\lambda, \mu)})^c = \bigcup_{t=1}^{\lambda_1} \bigcup_{\substack{\Lambda, \\ \phi(t, \Lambda) \trianglelefteq (\lambda, \mu)}} G_{t, \Lambda}.$$

Actually, if one fixes $0 \leq t \leq \lambda_1$, one can prove that

$$\phi(t, \Lambda) \trianglelefteq (\lambda, \mu) \Leftrightarrow \Lambda \trianglelefteq \lambda^{(t)} \uplus \mu,$$

where $\lambda^{(t)}$ is defined as follows: if $s = \max\{i; \lambda_i \geq k\}$, then

$$\lambda^{(t)} = (\lambda_1, \dots, \lambda_{s-1}, \lambda_s + \lambda_{s+1} - k, \lambda_{s+2}, \dots, \lambda_\ell).$$

Finally, we can reformulate the decomposition as:

$$(V_{(\lambda, \mu)})^c = \bigcup_{t=1}^{\lambda_1} \bigcup_{\Lambda \trianglelefteq \lambda^{(t)} \uplus \mu} G_{(t, \Lambda)}.$$

7. APPLICATIONS TO B_n -INVARIANT IDEALS

7.1. Specht ideals in B_n -invariant ideals. The main result in the article [MRV21] studies the \mathcal{S}_n -Specht ideals contained in an ideal I which is globally invariant under the action of \mathcal{S}_n . We give here the main ideas of this statement, before extending it to the case of B_n .

For P a polynomial in an \mathcal{S}_n -invariant ideal $I \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ of degree d , we denote by $\text{wt}(P_d)$ the number of variables appearing in the highest component P_d of P . Moreover, for a monomial m of degree d in P , the partial degrees of m induce a partition (k_1, \dots, k_ℓ) , and under the assumption $\text{wt}(P_d) + d \leq n$, we have in particular $\text{wt}(P_d) + \ell \leq n$, and therefore we can define the partition

$$\mu(m) = (k_1 + 1, k_2 + 1, \dots, k_\ell + 1, \underbrace{1, \dots, 1}_{n-d-\ell})$$

of n . It is then proved ([MRV21, Theorem 4]) that for every monomial $m \in \text{Mon}(P_d)$, the ideal I contains every sp_T for which $\text{sh}(T) \trianglelefteq \mu(m)^\perp$.

The proof works as follows: up to permutation of the variables, we may assume that

$$m = \mathbf{x}_1^{k_1} \mathbf{x}_2^{k_2} \dots \mathbf{x}_\ell^{k_\ell}$$

and since $\text{wt}(P_d) + d \leq n$, there exists $d = k_1 + \dots + k_\ell$ many variables in $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ that do not appear in P_d . More precisely, we can take I_1, \dots, I_ℓ , disjoint subsets of $\{1, \dots, n\}$ such that for any $1 \leq i \leq \ell$, there are k_i elements in I_i , and none of them appears in P_d . Then, if for $1 \leq i \leq \ell$, $J_i = \{i\} \cup I_i$, we can prove

$$(7.1) \quad \Delta_{J_1} \dots \Delta_{J_\ell} = \frac{k}{k_1! \dots k_\ell!} \sum_{\sigma \in S_{J_1} \times \dots \times S_{J_\ell}} \epsilon(\sigma) \sigma(\Delta_{I_1} \dots \Delta_{I_\ell} P)$$

where Δ_I is the Vandermonde polynomial of the ordered set I and $k \neq 0$.

Now, we want to generalize this result to B_n -invariant ideals. First, we need to associate a bipartition to a given monomial:

Definition 7.1. *Let m be a monomial in $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$. There exist unique sets I_1 and I_2 such that we can write m as*

$$m = \prod_{i \in I_1} \mathbf{x}_i^{2k_i} \prod_{i \in I_2} \mathbf{x}_i^{2r_i} \prod_{i \in I_2} \mathbf{x}_i$$

and $k_i \neq 0$. Denote $\ell = |I_1|$, $d_1 = \sum_{i \in I_1} k_i$, $s = |I_2|$, and $d_2 = \sum_{i \in I_2} r_i$. The sets $\{k_i, i \in I_1\}$ and $\{r_i, i \in I_2\}$ respectively induce partitions $(\lambda_1, \dots, \lambda_\ell)$ of d_1 and (μ_1, \dots, μ_s) of d_2 . If moreover we assume that $\ell + s + d_1 + d_2 \leq n$, we can define a bipartition $\Gamma(m)$ of n by

$$\Gamma(m) = (\tilde{\lambda}, \tilde{\mu}) = ((\lambda_1 + 1, \lambda_2 + 1, \dots, \lambda_\ell + 1, \underbrace{1, \dots, 1}_{n-(\ell+s+d_1+d_2)}), (\mu_1 + 1, \mu_2 + 1, \dots, \mu_s + 1)).$$

Finally, we define

$$\Gamma^*(m) = (\tilde{\lambda}^\perp, \tilde{\mu}^\perp) = (n - (s + d_1 + d_2), \ell, \tilde{\lambda}_3^\perp, \dots), (s, \tilde{\mu}_2^\perp, \dots),$$

which is a bipartition of n as well.

With this notion, we get, for B_n -invariant ideals:

Theorem 7.2. *Let $I \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ be a B_n -invariant ideal, and let $P \in I$. Assume that m is a monomial in the homogeneous component of highest degree of P . Using the notation of Definition 7.1, assume that $\text{wt}(P_d) + d_1 + d_2 \leq n$. Then, we have the ideal inclusion*

$$I_{\Gamma^*(m)} \subset I.$$

Proof. Up to permutation, we may assume that

$$m = \prod_{i=1}^{\ell} \mathbf{x}_i^{2k_i} \prod_{i=\ell+1}^{\ell+s} \mathbf{x}_i^{2r_i} \prod_{i=\ell+1}^{\ell+s} \mathbf{x}_i$$

and that the coefficient of m in P is 1. Let $\epsilon_i \in B_n$ the map changing \mathbf{x}_i in $-\mathbf{x}_i$. Then, the polynomial

$$\frac{P - \epsilon_i P}{2}$$

is a polynomial in I whose terms are exactly the terms of P having an odd degree in \mathbf{x}_i , and therefore divisible by \mathbf{x}_i . After applying this transformation for every $i \in \{\ell + 1, \dots, \ell + s\}$, we may substitute P with a polynomial of the form

$$\tilde{P}(\mathbf{x}) \prod_{i=\ell+1}^{\ell+s} \mathbf{x}_i,$$

containing m in its leading term, and where every term in \tilde{P} has even degree in \mathbf{x}_i , for every $i \in \{\ell + 1, \dots, \ell + s\}$. Further, for $i \notin \{\ell + 1, \dots, \ell + s\}$, we can apply the transformation

$$\frac{\tilde{P}(\mathbf{x}) + \epsilon_i \tilde{P}(\mathbf{x})}{2}$$

to get a polynomial which is still in I , but its terms are exactly the terms of P having an even degree in \mathbf{x}_i . In the end, we may assume that P is of the form:

$$P(\mathbf{x}) = Q(\mathbf{x}^2) \prod_{i=\ell+1}^{\ell+s} \mathbf{x}_i,$$

where m is still a monomial of the leading term.

Now we can apply a strategy similar to the one described for \mathcal{S}_n -invariant ideals. Since $\ell + s + d_1 + d_2 \leq \text{wt}(P_d) + d_1 + d_2 \leq n$, there exists $d_1 + d_2 = k_1 + \dots + k_\ell + r_1 + \dots + r_s$ many variables in $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ that do not appear in P_d , and we can take $I_1, \dots, I_\ell, I_{\ell+1}, \dots, I_{\ell+s}$, disjoint subsets of $\{1, \dots, n\}$ such that for any $1 \leq i \leq \ell$, there are k_i elements in I_i , for any $\ell + 1 \leq i \leq \ell + s$, there are r_i elements in I_i , and none of them appears in P_d . Then, for $1 \leq i \leq \ell + s$, denote

$$J_i = \{i\} \cup I_i,$$

and $\tilde{\Delta}_J(\mathbf{x}) = \Delta_J(\mathbf{x}^2)$ the Vandermonde polynomial associated with the variables \mathbf{x}_i^2 for $i \in J$. Consider

$$R(\mathbf{x}) = P(\mathbf{x}) \tilde{\Delta}_{I_1} \cdots \tilde{\Delta}_{I_{\ell+s}} \prod_{i \in I_{\ell+1} \times \cdots \times I_{\ell+s}} \mathbf{x}_i$$

We then have

$$\begin{aligned}
\sum_{\sigma \in S_{J_1} \times \cdots \times S_{J_{\ell+s}}} \epsilon(\sigma) \sigma(R(\mathbf{x})) &= \sum_{\sigma \in S_{J_1} \times \cdots \times S_{J_{\ell+s}}} \epsilon(\sigma) \sigma(P(\mathbf{x}) \tilde{\Delta}_{I_1} \cdots \tilde{\Delta}_{I_{\ell+s}} \prod_{i \in I_{\ell+1} \times \cdots \times I_{\ell+s}} \mathbf{x}_i) \\
&= \sum_{\sigma \in S_{J_1} \times \cdots \times S_{J_{\ell+s}}} \epsilon(\sigma) \sigma \left(Q(\mathbf{x}^2) \tilde{\Delta}_{I_1} \cdots \tilde{\Delta}_{I_{\ell+s}} \prod_{i \in J_{\ell+1} \times \cdots \times J_{\ell+s}} \mathbf{x}_i \right) \\
&= \prod_{i \in J_{\ell+1} \times \cdots \times J_{\ell+s}} \mathbf{x}_i \sum_{\sigma \in S_{J_1} \times \cdots \times S_{J_{\ell}}} \epsilon(\sigma) \sigma(Q(\mathbf{x}^2) \tilde{\Delta}_{I_1} \cdots \tilde{\Delta}_{I_{\ell+s}}) \\
&= k_1! \cdots k_{\ell}! \cdot r_{\ell+1}! \cdots r_{\ell+s}! \tilde{\Delta}_{J_1} \cdots \tilde{\Delta}_{J_{\ell}} \tilde{\Delta}_{J_{\ell+1}} \cdots \tilde{\Delta}_{J_{\ell+s}} \prod_{i \in J_{\ell+1} \times \cdots \times J_{\ell+s}} \mathbf{x}_i
\end{aligned}$$

where the last equality follows from (7.1). Therefore, I contains the polynomial

$$\tilde{\Delta}_{J_1} \cdots \tilde{\Delta}_{J_{\ell}} \tilde{\Delta}_{J_{\ell+1}} \cdots \tilde{\Delta}_{J_{\ell+s}} \prod_{i \in J_{\ell+1} \times \cdots \times J_{\ell+s}} \mathbf{x}_i$$

which is one of the generators of $I_{\Gamma^*(m)}$. By symmetry, we get the theorem. \square

Corollary 7.3. *Let $I \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ be a B_n -invariant ideal, and let $P \in I$. Assume that m is a monomial in the leading term of P . Using the notation of Definition 7.1, assume that $\text{wt}(P_d) + d_1 + d_2 \leq n$. Then, we have the inclusion of varieties*

$$V(I) \subset V(I_{\Gamma^*(m)})$$

and $V(I) \cap H_{(\theta, \omega)} = \emptyset$ for any bipartition $(\theta, \omega) \in \text{BP}_n$ bidominated by $\Gamma^*(m)$.

Proof. The inclusion of varieties follows immediately from the set inclusion in Theorem 7.2, while the second claim follows from the set partition of Specht varieties in Theorem 6.6. \square

We illustrate how this can be applied:

Example 7.4. *Let $P = \mathbf{x}_2 \mathbf{x}_3 (\mathbf{x}_1^2 - 1) \in \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_4]$. The polynomial P contains a unique monomial $m = \mathbf{x}_1^2 \mathbf{x}_2 \mathbf{x}_3$ of highest degree. Using the notation of Definition 7.1 we obtain $\text{wt}(P_4) = 3, l = 1, s = 2, d_1 = 1, d_2 = 0, \Gamma(m) = ((2), (1, 1))$, and $\Gamma^*(m) = ((1, 1), (2))$. Let I denote the ideal that is generated by the B_n orbit of P . Then, by Corollary 7.3 it must be $V(I) \subset V_{((1, 1), (2))}$. Thus, we have*

$$V(I) \subset B_4 \cdot \{(0, 0, a, a), (0, 0, a, b), (a, a, a, a), (0, 0, 0, a), (0, 0, 0, 0) : a, b \in \mathbb{K}_{>0}\}.$$

We observe that $V(I) = \{(1, 1, 1, 1), (0, 0, 0, a), (0, 0, 0, 0) : a \in \mathbb{K}\}$. Thus $V(I)$ contains already points of three of the five possible orbit types.

7.2. Connections with Representation Theory. We assume that $\mathbb{K} = \mathbb{C}$, or $\mathbb{K} = \mathbb{R}$ if G is a real reflection group. Let G be a finite group acting linearly on the polynomial ring $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$. The polynomials fixed by this action form a finitely generated subalgebra $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$. Moreover, each finite group admits - up to isomorphism - a finite number of irreducible $\mathbb{K}[G]$ -modules and the action on $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ can be decomposed into isotypic components, i.e., we have a decomposition of the form

$$\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n] = \bigoplus_{\chi} \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{\chi},$$

where χ runs over the pairwise non-isomorphic representations and each isotypic component $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{\chi}$ contains only pairwise isomorphic $\mathbb{K}[G]$ -submodules. Notice, that with this notion the invariant polynomials $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$ correspond to the trivial representation. Clearly, $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ has the structure of a $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$ -module. Finally, let $J_+ \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ be the ideal generated by invariant polynomials of positive degree. Then the algebra $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_G := \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]/J_+$ is called the *coinvariant algebra*. Whereas these algebras can be defined and studied for all finite groups, it was shown by Chevalley ([Che55, Theorem (B)]) and Sephard-Todd ([ST54]) that for finite reflection groups, these algebras are very regular and moreover that they characterize finite reflection groups by what is now known as Chevalley–Shephard–Todd theorem:

Theorem 7.5 ([Che55; ST54]). *Let G be a finite group. Then, the following are equivalent*

- (1) G is a group generated by reflections.
- (2) The algebra of polynomial invariants $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$ is a (free) polynomial algebra.
- (3) The algebra $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ is a free module over $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$
- (4) $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_G$ affords the regular representation of G , i.e.,

$$\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_G \simeq \bigoplus_{\chi} \dim(\chi) \chi,$$

where χ runs over the pairwise non-isomorphic representations of G .

Let $(\lambda, \mu) \in \text{BP}_n$ be a bipartition, denote by $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)}$ the isotypic component corresponding to (λ, μ) , and by $I_{(\lambda, \mu)}$ the associated Specht ideal. Notice that by Theorem 7.5, $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)}$ is also a finitely generated $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^G$ module, generated by $s_{\lambda, \mu}$ many elements, where $s_{\lambda, \mu}$ denotes the dimension of the corresponding irreducible representation. It follows from ([MY98, Theorem 1 (2)]) that $s_{\lambda, \mu}$ is in fact equal to the number of standard bitableaux of shape λ, μ . We note the following Proposition:

Proposition 7.6. *Let d be minimal with*

$$\mathcal{V} := \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)} \cap \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{\leq d} \neq \emptyset.$$

Then, the multiplicity of an irreducible representation of type (λ, μ) in \mathcal{V} is 1. This unique irreducible representation is given by the B_n -Specht polynomials of shape (λ, μ) . Moreover, $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)}$ is contained in the ideal generated by this unique irreducible representation, i.e.,

$$\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)} \subset I_{(\lambda, \mu)}.$$

Proof. Since $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]_{(\lambda, \mu)}$ is a direct sum of irreducible B_n -modules isomorphic to the standard B_n -Specht module, $W_{(\lambda, \mu)}$, it is enough to show that for every Specht polynomial $Q = \text{sp}_{(T, S)}$ with $\text{sh}(T, S) = (\lambda, \mu)$, and every B_n -isomorphism ϕ , the polynomial

$$P = \phi(Q)$$

is divisible by Q . First, for every $i \in \{1, \dots, n\}$, if ϵ_i is the map replacing \mathbf{x}_i with $-\mathbf{x}_i$, since ϕ respects the action of B_n , we must have

$$\epsilon_i P = \begin{cases} -P & \text{if } i \in S \\ P & \text{if } i \notin S \end{cases},$$

which implies that P is of the form

$$P(\mathbf{x}) = \tilde{P}(\mathbf{x}^2) \prod_{i \in S} \mathbf{x}_i.$$

Then, for every τ switching two elements in a same column of T or S , we must have $\tau \tilde{P} = -\tilde{P}$, so that P is divisible by Q . \square

Remark 7.7. *We remark that the statement about multiplicity 1 of an irreducible representation in Proposition 7.6 does not apply in general. Consider the real reflection group $D_n \subset B_n$ which is generated by all permutations and those maps that switch an even number of signs. Then the B_n -irreducible representation of type (λ, μ) and (μ, λ) remain D_n -irreducible if $\lambda \neq \mu$, but are D_n -isomorphic. By Theorem 5.1 we have*

$$I_{((2), (1, 1))} \not\subset I_{((1, 1), (2))} \quad \text{and} \quad I_{((1, 1), (2))} \not\subset I_{((2), (1, 1))}.$$

Thus, no polynomial in the B_n -orbit of $(\mathbf{x}_3^2 - \mathbf{x}_4^2)\mathbf{x}_3\mathbf{x}_4$ divides the polynomial $(\mathbf{x}_1^2 - \mathbf{x}_2^2)\mathbf{x}_3\mathbf{x}_4$ although

$$\langle \text{sp}_{(T, S)} : \text{sh}(T, S) = ((2), (1, 1)) \rangle_{\mathbb{K}} \simeq_{D_n} \langle \text{sp}_{(T, S)} : \text{sh}(T, S) = ((1, 1), (2)) \rangle_{\mathbb{K}}.$$

The D_n -irreducible representation $((2), (1, 1))$ occurs for the first time in $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_4]_{\leq 4}$ but with multiplicity 2.

Combining Corollary 7.3 with Proposition 7.6 we obtain the following.

Theorem 7.8. *Let $I \subset \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]$ a B_n -invariant ideal satisfying the conditions of 7.3. Consider the the associated coordinate ring $R_I = \mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]/I$. Then, viewed as a $\mathbb{K}[B_n]$ -module, R_I does not contain any irreducible $\mathbb{K}[B_n]$ -submodule which is isomorphic to $W_{\lambda, \mu}$. Moreover, R_I is a finite $\mathbb{K}[\mathbf{x}_1, \dots, \mathbf{x}_n]^{B_n}$ module of rank bounded by $\sum_{(\theta, \omega) \triangleleft (\lambda, \mu)} s_{(\theta, \omega)}^2$.*

8. CONCLUSION AND OPEN QUESTIONS

We initiated in this article the investigation of a class of polynomial ideals which are naturally linked to the action of a group on a polynomial ring. Our results provide an analogue of the relation of the combinatorics of integer partitions and \mathcal{S}_n -Specht ideals to bipartitions and B_n -Specht ideals. The present work shows that it is indeed possible to derive an analogous connection between combinatorics and algebra for the case of the hyperoctahedral group as was observed in the case of the symmetric group. Both groups are finite reflection groups, and they thus share important similarities from a view point of invariant theory and representation theory. Our results here lead to the natural question, if similar relations between integer (bi)-partitions and ideals can be derived for other (pseudo)-reflection groups. Indeed, in [MY98] a similar basis of the coinvariant algebra is provided for complex reflection groups of type $G(r, p, n)$, where $r, p, n \in \mathbb{Z}_{\geq 1}$ and $p \mid n$. We recover $G(1, 1, n) \simeq \mathcal{S}_n$, $G(2, 1, n) \simeq B_n$, and $G(2, 2, n) \simeq D_n$. It seems plausible to envision similar results to the ones presented here in these cases as well. More precisely, that there is a partial order on r -multipartitions, which are linked to the irreducible representations of the complex reflection group $G(r, 1, n)$, which transfers to the inclusion of the $G(r, 1, n)$ -Specht ideals and their corresponding varieties. Furthermore, it remains to investigate if the B_n -Specht ideals also have similarly nice algebraic properties as their \mathcal{S}_n counter parts. Indeed, it is known that the \mathcal{S}_n -Specht ideals are radical (see [MOY21, Theorem 1.1] and [Woo05, Proposition 4]). Both proofs rely on the understanding of the \mathcal{S}_n -Specht varieties in terms of orbit sets, i.e., Theorem 6.6, and crucially depend on the property that any \mathcal{S}_n -orbit set is non-empty which is not true for B_n -Specht varieties. Nevertheless, computational evidence for small number of variables motivates the conjecture, similar to the \mathcal{S}_n situation ([Lie21; Woo05; MOY21; Rie21]).

Conjecture 8.1. *The B_n -Specht ideals are radical. Moreover, for a bipartition $(\lambda, \mu) \in \text{BP}_n$ the B_n -Specht polynomials $\{\text{sp}_{(T, S)} : (T, S) \text{ is a bitableau of shape } (\theta, \omega) \triangleleft (\lambda, \mu)\}$ form a universal Gröbner basis of $I_{(\lambda, \mu)}$.*

Finally, Yanagawa [Yan21] classified the partitions for which the associated \mathcal{S}_n -Specht ideals are Cohen-Macaulay and it would be interesting to derive a similar characterization of bipartitions (λ, μ) for which the corresponding B_n -Specht ideals have this property.

REFERENCES

- [Abe+20] T. Abe, T. Horiguchi, M. Masuda, S. Murai, and T. Sato. “Hessenberg varieties and hyperplane arrangements”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 2020.764 (2020), pp. 241–286.
- [AM17] M. Aguiar and S. Mahajan. *Topics in hyperplane arrangements*. Vol. 226. American Mathematical Soc., 2017.
- [AMP81] E. Al-Aamily, A. Morris, and M. Peel. “The representations of the Weyl groups of type B_n ”. In: *Journal of Algebra* 68.2 (1981), pp. 298–305.
- [Ari01] S. Ariki. “On the classification of simple modules for cyclotomic Hecke algebras of type $G(m, 1, n)$ and Kleshchev multipartitions”. In: *Osaka Journal of Mathematics* 38.4 (2001), pp. 827–837.
- [BR21a] S. Basu and C. Riener. “Vandermonde varieties, mirrored spaces, and the cohomology of symmetric semi-algebraic sets”. In: *Foundations of Computational Mathematics* (2021), pp. 1–68.

- [BR21b] G. Blekherman and C. Riener. “Symmetric non-negative forms and sums of squares”. In: *Discrete & Computational Geometry* 65.3 (2021), pp. 764–799.
- [Bry73] T. Brylawski. “The lattice of integer partitions”. In: *Discrete mathematics* 6.3 (1973), pp. 201–219.
- [Che55] C. Chevalley. “Invariants of finite groups generated by reflections”. In: *American Journal of Mathematics* 77.4 (1955), pp. 778–782.
- [Che93] W. Y. Chen. “Induced cycle structures of the hyperoctahedral group”. In: *SIAM Journal on Discrete Mathematics* 6.3 (1993), pp. 353–362.
- [CLR87] M.-D. Choi, T.-Y. Lam, and B. Reznick. “Even symmetric sextics”. In: *Mathematische Zeitschrift* 195.4 (1987), pp. 559–580.
- [CS93] W. Y. Chen and R. P. Stanley. “Derangements on the n-cube”. In: *Discrete Mathematics* 115.1-3 (1993), pp. 65–75.
- [DJM95] R. Dipper, G. James, and E. Murphy. “Hecke algebras of type B_n at roots of unity”. In: *Proceedings of the London Mathematical Society* 3.3 (1995), pp. 505–528.
- [Dos+17] M. Dostert, C. Guzmán, F. M. de Oliveira Filho, and F. Vallentin. “New upper bounds for the density of translative packings of three-dimensional convex bodies with tetrahedral symmetry”. In: *Discrete Comput. Geom.* 58.2 (2017), pp. 449–481. ISSN: 0179-5376.
- [Fau+20] J.-C. Faugère, G. Labahn, M. S. E. Din, É. Schost, and T. X. Vu. “Computing critical points for invariant algebraic systems”. In: *arXiv preprint arXiv:2009.00847* (2020).
- [Har99] W. R. Harris. “Real even symmetric ternary forms”. In: *Journal of Algebra* 222.1 (1999), pp. 204–245.
- [HS21] A. Heaton and I. Shankar. “An SOS counterexample to an inequality of symmetric functions”. In: *Journal of Pure and Applied Algebra* 225.8 (2021), p. 106656.
- [KLM20] A. Kurpisz, S. Leppänen, and M. Mastrolilli. “Sum-of-squares hierarchy lower bounds for symmetric formulations”. In: *Mathematical Programming* 182.1 (2020), pp. 369–397.
- [Lie21] A. Lien. “Symmetric ideals”. MA thesis. UiT Norges arktiske universitet, 2021.
- [Mou+22] P. Moustrou, H. Naumann, C. Riener, T. Theobald, and H. Verdure. “Symmetry reduction in AM/GM-based optimization”. In: *SIAM Journal on Optimization* 32.2 (2022), pp. 765–785.
- [MOY21] S. Murai, H. Ohsugi, and K. Yanagawa. “A note on the reducedness and Gröbner bases of Specht ideals”. In: *arXiv preprint arXiv:2111.05525* (2021).
- [MRV21] P. Moustrou, C. Riener, and H. Verdure. “Symmetric ideals, Specht polynomials and solutions to symmetric systems of equations”. In: *Journal of Symbolic Computation* 107 (2021), pp. 106–121.
- [Mus93] C. Musili. “Representations of the Hyperoctahedral Group B_n ”. In: *Representations of Finite Groups*. Springer, 1993, pp. 197–220.
- [MY98] H. Morita and H.-F. Yamada. “Higher Specht polynomials for the complex reflection group $G(r, p, n)$ ”. In: *Hokkaido mathematical journal* 27.3 (1998), pp. 505–515.
- [Rie+13] C. Riener, T. Theobald, L. J. Andrén, and J. B. Lasserre. “Exploiting symmetries in SDP-relaxations for polynomial optimization”. In: *Mathematics of Operations Research* 38.1 (2013), pp. 122–141.
- [Rie21] C. Riener. “Specht Ideals, their varieties and connections to symmetric ideals”. In: *MFO-RIMS Tandem Workshop: Symmetries on Polynomial Ideals and Varieties*. Vol. Workshops 2021. Oberwolfach Reports, 2021.
- [RS18] C. Riener and M. Safey el Din. “Real root finding for equivariant semi-algebraic systems”. In: *Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*. 2018, pp. 335–342.
- [Spe37a] W. Specht. “Darstellungstheorie der Hyperoktaedergruppe”. In: *Mathematische Zeitschrift* 42.1 (1937), pp. 629–640.

- [Spe37b] W. Specht. “Zur Darstellungstheorie der symmetrischen Gruppe”. In: *Mathematische Zeitschrift* 42.1 (1937), pp. 774–779.
- [ST54] G. C. Shephard and J. A. Todd. “Finite Unitary Reflection Groups”. In: *Canadian Journal of Mathematics* 6 (1954), pp. 274–304.
- [Woo05] A. Woo. *Ideals of the polynomial ring generated by irreducible symmetric group representations and Ellingsrud-Strømme cells on the Hilbert scheme*. University of California, Berkeley, 2005.
- [Xia18] J. Xia. “A partial order on bipartitions from the generalized Springer correspondence”. In: *Journal of Combinatorial Algebra* 2.3 (2018), pp. 301–309.
- [Yan21] K. Yanagawa. “When is a Specht ideal Cohen–Macaulay?” In: *Journal of Commutative Algebra* 13.4 (2021), pp. 589–608.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UiT - THE ARCTIC UNIVERSITY OF NORWAY, 9037 TROMSØ, NORWAY

Email address: `sebastian.debus@uit.no`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR 5219, UT2J, 31058 TOULOUSE, FRANCE

Email address: `philippe.moustrou@math.univ-toulouse.fr`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UiT - THE ARCTIC UNIVERSITY OF NORWAY, 9037 TROMSØ, NORWAY

Email address: `cordian.riener@uit.no`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UiT - THE ARCTIC UNIVERSITY OF NORWAY, 9037 TROMSØ, NORWAY

Email address: `hugues.verdure@uit.no`