

Topological twists of massive SQCD, Part I

Johannes Aspmann, Elias Furrer, Jan Manschot

School of Mathematics, Trinity College, Dublin 2, Ireland

Hamilton Mathematical Institute, Trinity College, Dublin 2

E-mail: aspmannj@maths.tcd.ie, furrere@maths.tcd.ie,
manschot@maths.tcd.ie

ABSTRACT: We consider topological twists of four-dimensional $\mathcal{N} = 2$ supersymmetric QCD with gauge group $SU(2)$ and $N_f \leq 3$ fundamental hypermultiplets. The twists are labelled by a choice of background fluxes for the flavour group, which provides an infinite family of topological partition functions. In this Part I, we demonstrate that in the presence of such fluxes the theories can be formulated for arbitrary gauge bundles on a compact four-manifold. Moreover, we consider arbitrary masses for the hypermultiplets, which introduce new intricacies for the evaluation of the low-energy path integral on the Coulomb branch. We develop techniques for the evaluation of these path integrals. In the forthcoming Part II, we will deal with the explicit evaluation.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Special geometry and SW theories | 4 |
| 2.1 | Field content | 4 |
| 2.2 | Seiberg-Witten geometry | 5 |
| 2.3 | Characteristic functions on the Coulomb branch | 8 |
| 2.4 | Fundamental domains | 11 |
| 2.5 | Monodromies | 18 |
| 3 | The UV theory on a four-manifold | 20 |
| 3.1 | Aspects of four-manifolds | 20 |
| 3.2 | Topological twisting with background fluxes | 21 |
| 3.3 | Correlation functions and moduli spaces | 25 |
| 4 | The effective theory on a four-manifold | 27 |
| 4.1 | Hypermultiplets and background fields | 27 |
| 4.2 | Sum over fluxes | 28 |
| 4.3 | Observables and contact terms | 30 |
| 5 | The u-plane integral | 31 |
| 5.1 | Definition of the integrand | 31 |
| 5.2 | Monodromy transformations of the integrand | 33 |
| 6 | Integration over fundamental domains | 37 |
| A | Modular forms | 41 |
| A.1 | Theta functions and Eisenstein series | 42 |
| A.2 | Siegel-Narain theta function | 43 |
| B | Class \mathcal{S} representation | 44 |
| C | Winding numbers | 45 |

1 Introduction

Correlations functions of topologically twisted quantum field theories provide many insights into non-perturbative aspects of quantum field theory as well as the geometry of four-manifolds [1–3]. We consider topologically twisted $\mathcal{N} = 2$ supersymmetric Yang-Mills theories with additional matter multiplets on a compact four-manifold, which were introduced in [4–9]. After the work by Seiberg and Witten on the full non-perturbative solution [3, 10, 11], these theories have received much attention in physics [12–26] and mathematics [27–38].¹

We consider in this article topological twists of $\mathcal{N} = 2$ QCD with gauge group $SU(2)$ and matter multiplets in the fundamental representation of the gauge group. By including background fluxes for the flavour group, we obtain an infinite family of topological theories [16]. The choice of a background flux makes it possible to formulate topologically twists for $\mathcal{N} = 2$ SQCD for arbitrary 't Hooft fluxes, or first Chern classes of the gauge bundle. This is similar to the topological twist of $\mathcal{N} = 2^*$ $SU(2)$ gauge theory, which requires a non-vanishing background flux on a non-spin four-manifold [26]. We moreover develop techniques to determine correlation functions for arbitrary values of the masses of the hypermultiplets.

The starting point of our approach is the low-energy effective field theory on the Coulomb branch. This phase of the theory contributes for a compact four-manifold X with the topological condition that $b_2^+(X) = 1$ [12]. In this way, the classical Donaldson invariants can be derived starting from the Seiberg-Witten (SW) solution to $\mathcal{N} = 2$ supersymmetric Yang-Mills theory with gauge group $SU(2)$. The Coulomb branch integral (or u -plane integral) reduces to an integral over zero modes [12], and reads schematically

$$\Phi = \int_{\mathcal{B}} da \wedge d\bar{a} \rho(a) \Psi(a, \bar{a}), \quad (1.1)$$

where \mathcal{B} is the Coulomb branch with local coordinates a and \bar{a} , $\rho(a)$ contains the couplings to the background and $\Psi(a, \bar{a})$ is a sum over fluxes of the unbroken $U(1)$ gauge group. For simplicity, we have suppressed the dependence on the metric and not included observables here. For the pure $SU(2)$ theory, the Coulomb branch integral can be formulated and evaluated for arbitrary four-manifolds, without a requirement for Kähler or toric properties.

Recently, progress has been made on evaluating these u -plane integrals using a change of variables from a to the running coupling τ . As a result, the integration domain becomes a fundamental domain $\mathcal{F} \subseteq \mathbb{H}$ in the upper half-plane \mathbb{H} for the running coupling [24, 26, 43–51]. The integral then takes the form

$$\Phi = \int_{\mathcal{F}} d\tau \wedge d\bar{\tau} \nu(\tau) \Psi(\tau, \bar{\tau}), \quad (1.2)$$

¹For reviews, see for example [39–42].

where the measure factor $\nu(\tau)$ further contains the Jacobian for the change of variables from a to τ . The domain \mathcal{F} is a modular fundamental domain in previous analyses, corresponding to the duality group $\Gamma^0(4)$ for the pure $SU(2)$ theory [12, 47, 49], $\Gamma(2)$ for the $\mathcal{N} = 2^*$ theory [26] and similarly $\Gamma(2)$ and $\Gamma_0(4)$ for the theories with two and three massless flavours [43].

As mentioned above, we aim to apply this approach to $\mathcal{N} = 2$ supersymmetric $SU(2)$ theories with $N_f \leq 3$ hypermultiplets in the fundamental representation. Topological correlators of these asymptotically free theories have been considered in various papers before, in particular the formulation of the low energy path integral in [12, 13], SW contributions for four-manifolds with $b_2^+ > 1$ [12, 20, 21, 25], the u -plane integral for \mathbb{P}^2 [43], and the calculation of the partition function of the AD theory within the $N_f = 1$ theory [24]. Since no background fluxes are included in these works, the 't Hooft flux necessarily matches the second Stiefel-Whitney class of the four-manifold, $w_2(E) = w_2(X)$, since the twisted hypermultiplets are not well-defined otherwise.

Extending to generic 't Hooft fluxes, and application of the above approach (1.2) to fundamental hypermultiplets with generic masses, gives rise to several new aspects. In particular:

1. The fundamental domain of the effective coupling constant becomes more intricate for massive theories, and does for generic masses not correspond to a modular fundamental domain for a subgroup of $PSL(2, \mathbb{Z})$. The domain contains generically a set of branch points, and branch cuts starting from these points [52–54]. These aspects have to be dealt with appropriately.
2. We couple the hypermultiplets to background fluxes \mathbf{k}_j for the flavour group to formulate the theories for arbitrary 't Hooft fluxes. This gives rise to additional couplings in (1.1) and (1.2),

$$\prod_{j,k=1}^{N_f} \exp\left(-2\pi i \frac{\partial^2 F}{\partial m_j \partial m_k} B(\mathbf{k}_j, \mathbf{k}_k)\right), \quad (1.3)$$

where F is the prepotential of the massive theory, and $B(\cdot, \cdot)$ is the quadratic form associated to the intersection form on the middle homology $H_2(X, \mathbb{Z})$ of X . Such couplings were suggested by Shapere and Tachikawa [55], and are also essential for the formulation of the $\mathcal{N} = 2^*$ Yang-Mills theory on a non-spin four-manifold [26]. Similarly to [26], we also deduce a non-holomorphic coupling to \mathbf{k}_j . Moreover, for arriving at a single-valued integrand, we fix an ambiguity in the quadratic terms of the prepotential. These terms have appeared earlier in the literature in the context of singularities of the SW differential and winding numbers [11, 56].

3. Special points on the Coulomb branch give rise to superconformal theories, such as the Argyres-Douglas (AD) theories [57, 58] and the massless $N_f = 4$ theory [11]. Their topological partition functions and correlators can be found by considering them in certain mass deformations. The case of $N_f = 1$ is analysed in [42].

The paper is organised as follows. In Section 2, we present the Seiberg-Witten solution of $SU(2)$ $\mathcal{N} = 2$ SQCD in flat space, focusing on the fundamental domains for the effective coupling, which we illustrate in several interesting examples. In Section 3, we formulate the topological twist by coupling the hypermultiplets to external fluxes, such that the topological field theory is well-defined for arbitrary 't Hooft flux and non-spin manifolds. The topological low-energy effective theory coupled to N_f background fluxes is then modelled in Section 4 as a $SU(2) \times U(1)^{N_f}$ theory, with the matter fields corresponding to frozen $U(1)$ factors. This allows to compute the path integral explicitly as an integral over the u -plane. In Section 5, we formulate the u -plane integral as an integral over the fundamental domains. We prove that the single-valuedness under monodromies holds for a specific choice of magnetic winding numbers. Finally, in Section 6 we demonstrate that such integrals may be evaluated using mock modular forms, and we show that they localise at the cusps, elliptic points and interior singularities of the fundamental domains.

A following paper, Part II, will be dedicated to explicit analyses and computations for specific examples. There, we will give a detailed discussion on the contribution from the AD points for each type, as briefly mentioned in point 3 above, as well as calculating the u -plane integrals for a number of examples.

2 Special geometry and SW theories

In this Section, we review aspects of the non-perturbative solution for the low energy effective theory of $\mathcal{N} = 2$ SQCD with gauge group $SU(2)$ and $0 \leq N_f \leq 3$ fundamental hypermultiplets [10, 11]. See [59] for a review. Throughout, we let Λ_{N_f} denote the scale of the theory with N_f hypermultiplets having masses m_j , $j = 1, \dots, N_f$, and a the mass of the W-boson on the Coulomb branch.

2.1 Field content

The $\mathcal{N} = 2$ theories we consider contain a vector multiplet and $N_f \leq 3$ hypermultiplets. The fields in these multiplets form representations of $\text{Spin}(4) = SU(2)_+ \times SU(2)_-$ and $SU(2)_R$, which we denote by $(\mathbf{k}, \mathbf{l}, \mathbf{m})$, with \mathbf{k}, \mathbf{l} and \mathbf{m} dimensions of the representations.

The vector multiplet consists of a gauge field A_μ , complex scalar field ϕ , and a pair of Weyl fermions $\Psi_\alpha^I, \bar{\Psi}_{\dot{\alpha}}^I$. This multiplet transforms under the adjoint representation of the gauge group G . The representation of $SU(2)_+ \times SU(2)_- \times SU(2)_R$

formed by the bosonic fields is,

$$(\mathbf{2}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{1}), \quad (2.1)$$

while the representation for the fermions is

$$(\mathbf{1}, \mathbf{2}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{1}, \mathbf{2}). \quad (2.2)$$

The hypermultiplet consist of a pair of complex scalar fields, q and \tilde{q} , and Weyl fermions, λ_α , $\bar{\lambda}_{\dot{\alpha}}$, χ_α and $\bar{\chi}_{\dot{\alpha}}$. We fix the gauge group $G = \text{SU}(2)$, and let the hypermultiplets transform under the fundamental representation of this group. With the same notation as above, the bosonic fields of this multiplet form the representation,

$$(\mathbf{1}, \mathbf{1}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{2}), \quad (2.3)$$

while the fermions form the representation

$$(\mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1}). \quad (2.4)$$

2.2 Seiberg-Witten geometry

The Seiberg-Witten geometry underlies the Coulomb branch of $\mathcal{N} = 2$ gauge theory. The Coulomb branch is the phase of the theory where $\text{SU}(2)$ is broken to $\text{U}(1)$ by a vacuum expectation value (vev) of the vector multiplet scalar ϕ . The vev is semi-classically parametrised by a complex parameter a ,

$$\phi = \begin{pmatrix} a & 0 \\ 0 & -a \end{pmatrix}, \quad (2.5)$$

up to gauge transformations. In particular, $a \rightarrow -a$ is a gauge transformation. The gauge invariant order parameter is the Coulomb branch expectation value of the theory in \mathbb{R}^4 ,

$$u = \frac{1}{16\pi} \langle \text{Tr}(\phi^2) \rangle_{\mathbb{R}^4}. \quad (2.6)$$

The non-perturbative effective action of $\mathcal{N} = 2$ SQCD is characterised by the prepotential $F(a, \mathbf{m})$, with \mathbf{m} the mass vector $\mathbf{m} = (m_1, \dots, m_{N_f})$. The semi-classical part of F reads [56, 60–62]

$$\begin{aligned} F(a, \mathbf{m}) &= \frac{2i}{\pi} a^2 \log(a/\Lambda_{N_f}) - \frac{1}{2} \sum_{j=1}^{N_f} \left(n_j \frac{m_j}{\sqrt{2}} a + \frac{3}{8} \frac{i}{\pi} m_j^2 \right) \\ &\quad - \frac{i}{4\pi} \sum_{j=1}^{N_f} \left(a + \frac{m_j}{\sqrt{2}} \right)^2 \log\left(\left(a + \frac{m_j}{\sqrt{2}} \right) / \Lambda_{N_f} \right) + \left(a - \frac{m_j}{\sqrt{2}} \right)^2 \log\left(\left(a - \frac{m_j}{\sqrt{2}} \right) / \Lambda_{N_f} \right) \\ &\quad + \dots, \end{aligned} \quad (2.7)$$

where the ... indicate further non-perturbative corrections.

The $n_j \in \mathbb{Z}$ in (2.7) are the *magnetic winding numbers* of the periods $a_D := \frac{\partial F}{\partial a}$ dual to a [56, 60, 63]. These numbers seem to be only rarely discussed in the literature beyond these references.² Generally, the theory allows for N_f electric winding numbers for a and N_f magnetic winding numbers for a_D . These appear in the massive $N_f > 0$ theories since the Seiberg-Witten differentials now have poles with nonzero residues [60]. See also Appendix C. The choice (2.7) of the prepotential corresponds to fixing the electric winding numbers to be zero, or equivalently fixing the monodromy at infinity to map $a \rightarrow e^{\pi i} a$. Compare for example with [60, Eq. (2.17)]. In Section 5, we will discuss that the single-valuedness of the u -plane integral requires $n_j \equiv -1 \pmod{4}$.

We introduce the period a_D dual to a , and the parameters $m_{D,j}$ dual to m_j by

$$a_D = \frac{\partial F}{\partial a}, \quad m_{D,j} = \sqrt{2} \frac{\partial F}{\partial m_j}. \quad (2.8)$$

These parameters are further combined into the $(2 + 2N_f)$ -dimensional vector Π ,

$$\Pi = \begin{pmatrix} a_D \\ a \\ m_{D,1} \\ \frac{m_1}{\sqrt{2}} \\ \vdots \\ m_{D,N_f} \\ \frac{m_{N_f}}{\sqrt{2}} \end{pmatrix}. \quad (2.9)$$

This vector forms a local system over the u -plane. The elements of the vector form the symplectic form,

$$\omega_{N_f} = da_D \wedge da + \frac{1}{\sqrt{2}} \sum_{j=1}^{N_f} dm_{D,j} \wedge dm_j. \quad (2.10)$$

The effective gauge coupling is related to the prepotential through

$$\tau = \frac{\partial^2 F}{\partial a^2}. \quad (2.11)$$

We also introduce the couplings v_j and w_{jk} with $j, k \in 1, \dots, N_f$,³

$$v_j = \sqrt{2} \frac{\partial^2 F}{\partial a \partial m_j}, \quad w_{jk} = 2 \frac{\partial^2 F}{\partial m_j \partial m_k}. \quad (2.12)$$

²Nekrasov's partition function gives a specific choice upon expanding the function $\gamma_{\hbar}(x; \Lambda)$ in the perturbative part [62, 64].

³If we consider F as a function of $(a, \frac{1}{\sqrt{2}} m_1, \dots, \frac{1}{\sqrt{2}} m_{N_f})$, then the dual parameters are encoded in the Jacobian $\mathbf{J}_F = (a_D, \mathbf{m}_D)$, while the couplings are the elements of the Hessian $\mathbf{H}_F = \begin{pmatrix} \tau & \mathbf{v}^T \\ \mathbf{v} & \mathbf{w} \end{pmatrix}$.

The derivative of the prepotential with respect to the scale Λ_{N_f} provides the order parameter u (2.6) on the Coulomb branch,

$$u = -\frac{4\pi i}{4 - N_f} \Lambda_{N_f} \frac{\partial F}{\partial \Lambda_{N_f}}. \quad (2.13)$$

Using the different relations introduced above together with the fact that the prepotential satisfies the homogeneity equation [65–67]

$$2F = \Lambda_{N_f} \frac{\partial F}{\partial \Lambda_{N_f}} + \sum_{j=1}^{N_f} m_j \frac{\partial F}{\partial m_j} + a \frac{\partial F}{\partial a}, \quad (2.14)$$

we can find some non-trivial relations between the respective quantities. For instance, from the perturbative prepotential (2.7), we deduce that the leading terms of u are

$$u = -2a^2 + \frac{2}{4 - N_f} \sum_{j=1}^{N_f} m_j^2 + \dots \quad (2.15)$$

The weak-coupling limit then becomes in our conventions⁴

$$\begin{cases} \tau \rightarrow i\infty, \\ a \rightarrow \infty, \\ u \rightarrow -\infty. \end{cases} \quad (2.16)$$

The Seiberg-Witten (SW) solution provides a family of elliptic curves parametrised by the order parameter u and the masses m_i , whose complex structure corresponds to the running coupling $\tau = \frac{\theta}{\pi} + \frac{8\pi i}{g^2}$. For the theories of interest in this paper, the curves are given by [11]

$$\begin{aligned} N_f = 0: \quad & y^2 = x^3 - ux^2 + \frac{1}{4}\Lambda_0^4 x, \\ N_f = 1: \quad & y^2 = x^2(x - u) + \frac{1}{4}m\Lambda_1^3 x - \frac{1}{64}\Lambda_1^6, \\ N_f = 2: \quad & y^2 = (x^2 - \frac{1}{64}\Lambda_2^4)(x - u) + \frac{1}{4}m_1 m_2 \Lambda_2^2 x - \frac{1}{64}(m_1^2 + m_2^2)\Lambda_2^4, \\ N_f = 3: \quad & y^2 = x^2(x - u) - \frac{1}{64}\Lambda_3^2(x - u)^2 - \frac{1}{64}(m_1^2 + m_2^2 + m_3^2)\Lambda_3^2(x - u) \\ & + \frac{1}{4}m_1 m_2 m_3 \Lambda_3 x - \frac{1}{64}(m_1^2 m_2^2 + m_2^2 m_3^2 + m_1^2 m_3^2)\Lambda_3^2. \end{aligned} \quad (2.17)$$

The family of SW curves (2.17) are Jacobian rational elliptic surfaces with singular fibres [68–70]. These surfaces are well studied in the mathematical literature [71–73]. Rational in this context means that g_2 and g_3 are polynomials in u of degree at most

⁴Note that this differs slightly from some of the previous literature. However, it is shown in [52] to be the unique limit consistent with the RG flow.

4 and 6, respectively [74]. In Appendix B, we summarise the class \mathcal{S} representation of the SW curves [75].

The curve for N_f hypermultiplets reduces to the curve for $N_f - 1$ hypermultiplets upon decoupling a hypermultiplet in the double scaling limit [76]

$$m_{N_f} \rightarrow \infty, \quad \Lambda_{N_f} \rightarrow 0, \quad m_{N_f} \Lambda_{N_f}^{4-N_f} = \Lambda_{N_f-1}^{4-(N_f-1)}. \quad (2.18)$$

The curves (2.17) provide the exact results for the vevs of the scalar a and its dual a_D as period integrals. We have explicitly,

$$a = \int_A \lambda, \quad a_D = \int_B \lambda, \quad (2.19)$$

with λ the SW differential, for which we refer to [11, 56, 60].

2.3 Characteristic functions on the Coulomb branch

In this subsection, we discuss various characteristic functions on the Coulomb branch, which are instrumental for determining the u -plane integral. They are:

- the physical discriminant Δ_{N_f} ,
- the order parameter u ,
- the period da/du , and
- the derivatives $du/d\tau$ and $da/d\tau$.

The physical discriminant Δ_{N_f}

We define the physical discriminant Δ_{N_f} as the monic polynomial

$$\Delta_{N_f} = \prod_{j=1}^{N_f+2} (u - u_j), \quad (2.20)$$

where u_j for $j = 1, \dots, N_f + 2$ are the singular points of the effective theory. We let $j = 1, \dots, N_f$ label the singular points where one of the matter hypermultiplets becomes massless; and $j = N_f + 1, N_f + 2$ denote the strong coupling singularities where a monopole and a dyon, respectively, becomes massless.

The discriminant is a polynomial of degree $2 + N_f$ in u . It can also be determined directly from the SW curve [52],

$$\Delta_{N_f} = (-1)^{N_f} \Lambda_{N_f}^{2N_f-8} (g_2^3 - 27g_3^2), \quad (2.21)$$

where g_2 and g_3 are the Weierstraß invariants of the SW curve. Up to the sign and scale, the physical discriminant Δ_{N_f} equals the mathematical discriminant $g_2^3 - 27g_3^2$ for these theories.

As the masses are tuned, some of the singularities on the Coulomb branch can collide. If we consider Δ_{N_f} as a polynomial in u , its discriminant $D(\Delta_{N_f})$ vanishes if and only if two roots coincide. It is straightforward to show that for $N_f \leq 3$,

$$D(\Delta_{N_f}) = \left(D_{N_f}^{\text{AD}}\right)^3 \prod_{i < j} (m_i - m_j)^2 (m_i + m_j)^2. \quad (2.22)$$

This factorises the locus in mass space where singularities collide into two orthogonal components: The first component is the Argyres-Douglas (AD) locus given by the polynomial equation $D_{N_f}^{\text{AD}} = 0$, where mutually *non-local* singularities collide [52, 58, 77]. The other component is characterised by the equations $m_i = \pm m_j$, and one can check that this gives rise to mutually *local* singularities colliding. Here, the flavour symmetry gets enhanced and a Higgs branch opens up [11]. This extends the analysis of [52].

Given a mass configuration $\mathbf{m} = (m_1, \dots, m_{N_f})$, we can denote by k_l the weight (or multiplicity) of the l -th singularity, and by $\mathbf{k}(\mathbf{m}) = (k_1, k_2, \dots)$ the vector of those weights. Since the Coulomb branch $\mathcal{B}_{N_f}(\mathbf{m})$ contains $2 + N_f$ singularities aside from weak coupling $u = \infty$, it is clear that $\mathbf{k}(\mathbf{m})$ provides a partition of $2 + N_f$. This in turn partitions the mass space $\mathbb{C}^{N_f} \ni \mathbf{m}$ into finitely many regions where $\mathbf{k}(\mathbf{m})$ is locally constant. As an example, in Fig. 1 we plot the contours of (2.22) for $N_f = 2$ in the real $\mathbf{m} = (m_1, m_2)$ plane.

The possible singularity structures of the rational elliptic curves (2.17) are classified in Persson's list of allowed configurations of singular fibres [78, 79]. From Kodaira's classification, it follows that any solution to (2.22) gives rise to a singularity on the Coulomb branch of Kodaira type I_k , II , III , or IV . As described in [52, 70], the solutions to $0 = D_{N_f}^{\text{AD}}$ give rise to AD points of Kodaira type II , III and IV . The second component $0 = \prod_{i < j} (m_i - m_j)(m_i + m_j)$ can be studied in more detail. These are $2(N_f - 1)$ independent equations. Whenever one of the factors vanishes, the SW surface contains an I_k singularity with $k \geq 2$. For $N_f = 2$, the only possibility is I_2 , while for $N_f = 3$ singularities of type I_2 , I_3 and I_4 are possible. The point in the Coulomb branch \mathcal{B}_{N_f} corresponding to an I_k singularity with $k \geq 2$ intersects with a Higgs branch of quaternionic dimension $k - 1 \geq 1$ [11]. Further merging these I_k singularities with a mutually non-local singularity does not affect the Higgs branch, such that the points with a III or IV singularity also intersect with a Higgs branch of quaternionic dimension one or two, respectively, while the points with AD theories of type II do not intersect with any Higgs branch.

Order parameter u

The order parameter u , introduced in (2.6), is invariant under monodromies on the Coulomb branch. By bringing the SW curve to Weierstraß form, u can be expressed in terms of the running coupling $\tau \in \mathbb{H}$ (2.11) [52, 54, 80–82]. As in [52], we restrict the domain for $\tau \in \mathbb{H}$ to $\mathcal{F}(\mathbf{m}) \subseteq \mathbb{H}$, such that u is 1-to-1. In specific cases, u can




Figure 1: Partitioning of the real $\mathbf{m} = (m_1, m_2)$ plane in $N_f = 2$, in units of $\Lambda_2 = 1$. On the AD component the Coulomb branch $\mathcal{B}_2(\mathbf{m})$ contains an AD point of Kodaira type II (blue) or III (green). On the other component (orange), mutually local singularities collide. If \mathbf{m} is varied along a continuous path that does not cross the partitioning $0 = D(\Delta_{N_f})$, the weight vector $\mathbf{k}(\mathbf{m})$ is constant.

be expressed in terms of modular forms. We follow the conventions of [52, Section 2.2] in picking the solutions. We discuss the order parameters in more detail in the following Subsection 2.4.

The period da/du

The period $\frac{da}{du}$ can be related to the complex structure τ of the curve through [81]

$$\frac{da}{du} = \frac{1}{6} \sqrt{\frac{g_2 E_6}{g_3 E_4}}, \quad (2.23)$$

where the Weierstraß invariants g_2, g_3 are polynomials in u , and $E_4 = E_4(\tau)$, $E_6 = E_6(\tau)$ are the Eisenstein series of modular weights four and six respectively, defined in Eq. (A.7). The square root in (2.23) leads to an ambiguity in the sign of da/du . This can be resolved by taking a branch of the square root, for example at weak coupling. On the other hand, da/du is not a single-valued function on the u -plane. It changes by a sign under monodromies at weak coupling, and transforms with a modular weight under monodromies around strong coupling singularities.

The derivatives $du/d\tau$ and $da/d\tau$

The derivative $du/d\tau$ is related to the period $\frac{da}{du}$ through [52, 83]

$$\frac{du}{d\tau} = -\frac{16\pi i}{4 - N_f} \frac{\Delta_{N_f}}{P_{N_f}^M} \left(\frac{da}{du} \right)^2, \quad (2.24)$$

where $P_{N_f}^M$ is a polynomial in u . For $N_f \leq 3$, it reads

$$P_{N_f}^M = \frac{6}{4 - N_f} (-1)^{N_f} \Lambda_{N_f}^{2N_f - 8} (2g_2 g_3' - 3g_2' g_3), \quad (2.25)$$

where the prime denotes differentiation with respect to u .

When u is known in terms of modular forms, it is also straightforward to determine $\frac{du}{d\tau}$ by differentiating modular forms [52]. For generic masses \mathbf{m} , we can compute the q -series of u and thus $\frac{du}{d\tau}$ to any order. We will not explicitly need $du/d\tau$ for evaluating the u -plane integral, but through

$$\frac{da}{d\tau} = \frac{da}{du} \frac{du}{d\tau} \quad (2.26)$$

we can use it to evaluate $da/d\tau$. This derivative appears as the Jacobian for the change of integration variables from the periods a to the couplings τ .

2.4 Fundamental domains

The recent progress in computing u -plane integrals has been enabled by mapping the u -plane to a modular fundamental domain, on which the u -plane integrand can be related to mock modular forms and thus be efficiently evaluated [24, 26, 43–51]. It has been known since the 1990s that the u -planes for $\mathcal{N} = 2$ SQCD with $N_f = 0, 2, 3$ massless hypermultiplets are modular and correspond to fundamental domains for congruence subgroups of $\text{PSL}(2, \mathbb{Z})$ [80]. On the other hand, the generic mass case including the peculiar role of massless $N_f = 1$ have remained elusive.

The order parameter u for a given mass $\mathbf{m} = (m_1, \dots, m_{N_f})$ can be considered as a function

$$u : \mathbb{H} \rightarrow \mathcal{B}_{N_f}(\mathbf{m}), \quad (2.27)$$

where $\mathcal{B}_{N_f}(\mathbf{m})$ is the Coulomb branch of the theory with N_f hypermultiplets of mass \mathbf{m} . In [52], it was found that for $N_f \geq 1$ and generic masses the duality group does not act on τ by fractional linear transformations. This prevents the preimage $u^{-1}(\mathcal{B}_{N_f}(\mathbf{m}))$ from being a modular fundamental domain for a subgroup of $\text{PSL}(2, \mathbb{Z})$. Instead, we can define a fundamental domain $\mathcal{F}_{N_f}(\mathbf{m})$ through the equivalence relation $\tau \sim \tau' \Leftrightarrow u(\tau) = u(\tau')$, such that $u : \mathcal{F}_{N_f}(\mathbf{m}) \rightarrow \mathcal{B}_{N_f}(\mathbf{m})$ is bijective. In [52, 54], such fundamental domains for the effective gauge coupling have been found explicitly for $\mathcal{N} = 2$ SQCD with $N_f \leq 4$ generic masses. They decompose as

$$\mathcal{F}_{N_f}(\mathbf{m}) = \bigcup_{j=1}^n \alpha_j \mathcal{F}, \quad (2.28)$$

where $\mathcal{F} = \text{PSL}(2, \mathbb{Z}) \backslash \mathbb{H}$ is the key-hole fundamental domain of $\text{PSL}(2, \mathbb{Z})$, and $\alpha_j \in \text{PSL}(2, \mathbb{Z})$ are $n \leq 6$ maps that are locally constant as functions of the masses \mathbf{m} . More precisely, $\mathcal{F}_{N_f}(\mathbf{m})$ is constant on any of the finitely many connected components of $\{\mathbf{m} \in \mathbb{C}^{N_f} \mid D(\Delta_{N_f}) = 0\}$, where $D(\Delta_{N_f})$ is given in (2.22). We also call n the index of the domain $\mathcal{F}_{N_f}(\mathbf{m})$.

The geometry of $\mathcal{F}_{N_f}(\mathbf{m})$ agrees with that of the Coulomb branch $\mathcal{B}_{N_f}(\mathbf{m})$. Namely, $\mathcal{B}_{N_f}(\mathbf{m})$ is the complex plane with $2 + N_f$ singular points removed. In (2.28), those singularities are reflected in the cusps $\alpha_j(\infty)$, where the number of α_j with the same $\alpha_j(\infty)$ (giving the *width* of the cusp) agrees with the multiplicity of the singularity $u(\alpha_j(\infty))$.

The domains $\mathcal{F}(\mathbf{m}) := \mathcal{F}_{N_f}(\mathbf{m})$ are endowed with further data. For generic masses \mathbf{m} , there are N_f pairs of branch points on $\mathcal{F}(\mathbf{m})$ that are connected by branch cuts. The branch points are determined by the non-removable singularities of the rational function $\Delta_{N_f}(u)/P_{N_f}^M(u)$, as it appears in Matone's relation (2.24). They correspond to the nontrivial zeros of the discriminant of the sextic equation associated with the SW curve [52]. As the masses are varied, the branch points move on continuous paths in the fundamental domain.

When two branch points collide, there appears an elliptic point, which is a further dedicated point on the domain $\mathcal{F}(\mathbf{m})$. These elliptic points correspond to the superconformal Argyres-Douglas (AD) points [57, 58]. By Kodaira's classification of singular fibres, they are always in the $\text{PSL}(2, \mathbb{Z})$ orbit of i or $e^{\pi i/3}$. Their presence is also responsible for the index n of the domain to be reduced. The possible values are $n = 2, 3, 4$ if $\mathcal{F}(\mathbf{m})$ contains such an elliptic point, and $n = 6$ otherwise.

The $N_f = 4$ theory furthermore contains a UV coupling τ_{UV} . Since $u(\tau) \rightarrow \infty$ for $\tau \rightarrow \tau_{\text{UV}}$, it can be viewed as a puncture of the fundamental domain, and has to be excluded from the integration domain.

Let us review the explicit construction of $\mathcal{F}(\mathbf{m})$ in a few important examples. Many more examples can be found in [52, 54]. Here, we are mainly considering the masses to be small compared to the period a , such that the mass singularities discussed in previous sections can be considered as strong coupling singularities, or in other words as cusps on the real line of the fundamental domains. When the masses are increased it is more natural to think of the mass singularities as weakly coupled.

The pure SU(2) $N_f = 0$ theory

It is well-known that the pure SU(2) Coulomb branch corresponds to the fundamental domain for the congruence subgroup $\Gamma^0(4)$ of $\text{PSL}(2, \mathbb{Z})$. The order parameter can be explicitly determined in terms of a modular function for the duality group [84],

$$\frac{u}{\Lambda_0^2} = -\frac{1}{2} \frac{\vartheta_2^4 + \vartheta_3^4}{\vartheta_2^2 \vartheta_3^2}, \quad (2.29)$$

where the Jacobi theta functions ϑ_i are defined in Appendix A.1. A fundamental domain for the pure SU(2) theory is given in Fig. 2. Since the domain is modular,

the branch points are absent.




Figure 2: Fundamental domain of $\Gamma^0(4)$. This is the duality group of the pure $SU(2)$ theory. The two cusps on the real line correspond to the strong coupling singularities of the gauge theory, while the cusp at $\tau = i\infty$ corresponds to weak coupling. Boundaries of the same colour are identified.

The massless $N_f = 1$ theory

From the massless $N_f = 1$ SW curve, one directly finds [80]

$$\frac{u}{\Lambda_1^2} = -\frac{3}{2^{\frac{7}{3}}} \frac{E_4^{\frac{1}{2}}}{(E_4^{\frac{3}{2}} - E_6)^{\frac{1}{3}}}, \quad (2.30)$$

where E_4 and E_6 are the Eisenstein series defined in (A.7). This function also appears as an order parameter in pure $SU(3)$ SW theory [53] as well as in the description of certain elliptically fibred Calabi-Yau spaces [85]. Due to the presence of the square root, this function is not modular. The massless $N_f = 1$ curve has three distinct strongly coupled singularities, which become the three cusps of the fundamental domain aside from the weakly coupled region $u \rightarrow \infty$. The fundamental domain is plotted in Fig. 3. Note that, this is not the fundamental domain of any subgroup of $PSL(2, \mathbb{Z})$.

When a positive mass m is turned on, the branch points move in the fundamental domain emerging from those of the massless case. They move as a function of the mass as shown in [52], and depicted in Fig. 4. This further incorporates the hypermultiplet decoupling on the level of the fundamental domain.

The equal mass $N_f = 2$ theory

For $N_f = 2$ with $m_1 = m_2 = m$, the order parameter can be determined explicitly [52],

$$\frac{u}{\Lambda_2^2} = -\frac{\vartheta_4^8 + \vartheta_2^4 \vartheta_3^4 + (\vartheta_2^4 + \vartheta_3^4) \sqrt{16 \frac{m^2}{\Lambda_2^2} \vartheta_2^4 \vartheta_3^4 + \vartheta_4^8}}{8 \vartheta_2^4 \vartheta_3^4}. \quad (2.31)$$




Figure 3: Fundamental domain for massless $N_f = 1$.

The curve has a double singularity at $u_* = m^2 + \frac{\Lambda_2^2}{8}$ and two simple singularities at $u_{\pm} = -\frac{\Lambda_2^2}{8} \pm m\Lambda_2$. When m is varied, it can be determined how the branch point τ_{bp} moves in the fundamental domain. When m is not zero and not equal to $m_{\text{AD}} = \frac{\Lambda_2}{2}$, the singularities u_* , u_+ and u_- are distinct and the fundamental domain $\mathcal{F}(m, m)$ is given in Fig. 5. For any such mass, there is a pair of branch points in $\mathcal{F}(m, m)$, which are connected by a branch cut. Near a branch point τ_{bp} , u reads

$$u(\tau) = u_{\text{bp}} + c(\tau_{\text{bp}}) (\tau - \tau_{\text{bp}})^{1/2}, \quad (2.32)$$

for some constant $c(\tau_{\text{bp}})$.

The massless limit in $N_f = 2$

In the limit where $m_1 = m_2 = m \rightarrow 0$, one finds from (2.31) that

$$\frac{u}{\Lambda_2^2} = -\frac{1}{8} \frac{\vartheta_3^4 + \vartheta_4^4}{\vartheta_2^4} \quad (2.33)$$

This expression is a modular function for the principal congruence subgroup $\Gamma(2)$ of $\text{PSL}(2, \mathbb{Z})$. The holomorphy of (2.33) is explained by the fact that the pair of branch points for $m \neq 0$ collides in the limit $m \rightarrow 0$, which annihilates the branch cut. This is because the mutually local singularities $u_{\pm} \rightarrow -\frac{\Lambda_2^2}{8}$ collide, and thus the singularities corresponding to $\tau \rightarrow 0$ and $\tau \rightarrow 2$ in Fig. 5 are identified. The fundamental domain of the massless theory is thus the modular fundamental domain for the duality group $\Gamma(2)$, as depicted in Fig. 6.

The AD limit in $N_f = 2$

When $m \rightarrow m_{\text{AD}} = \frac{\Lambda_2}{2}$, the singularities u_* and u_+ collide and form the superconformal AD point (A_1, A_3) of Kodaira type III. As studied in detail in [52, 70, 86], this AD point forms an elliptic point of the fundamental domain. Since three mutually non-local singularities collide for this mass, the index of the domain $\mathcal{F}(m_{\text{AD}}, m_{\text{AD}})$




Figure 4: Choice of branch cuts (zigzag lines) for varying (real) mass in $N_f = 1$. Lines with the same color are identified. Starting with a small mass in Figure (a), we cut along the arcs of radius 1 around $\tau = 1$ and $\tau = 2$. At the AD mass we can use the identifications of the different boundaries to reorganise the domain in Figure (b) to the one of Figure (c). When we increase the mass further the cuts of Figure (c) move upwards as in Figure (d) eventually reaching infinity and disappearing, leaving us with the domain in Fig. 2 of the pure theory.

is reduced by 3, giving a domain of index $6 - 3 = 3$. By taking the limit from (2.31), one finds that the square root resolves by virtue of the Jacobi identity (A.4). This restores holomorphy and even modularity. One finds that the order parameter

$$\frac{u}{\Lambda_2^2} = -\frac{1}{64} \left(\frac{16\vartheta_4^8}{\vartheta_2^4\vartheta_3^4} + 40 \right) \quad (2.34)$$

is a modular function for the congruence subgroup $\Gamma_0(2)$ of $\text{PSL}(2, \mathbb{Z})$ [52]. Its elliptic fixed point $\tau_{\text{AD}} = 1 + i$ is precisely the AD point. We depict the corresponding fundamental domain in Fig. 7.

The $N_f = 4$ theory




Figure 5: Fundamental domain $\mathcal{F}_2(m, m)$ of the massive $\mathbf{m} = (m, m)$ $N_f = 2$ theory. The dashed lines correspond to the paths of the branch points from zero to infinite mass. For given positive mass m , the two branch points are identified under TST^{-1} , such that there is only one branch point $u_{\text{bp}} = 2m^2 - \frac{\Lambda^2}{8}$ on the u -plane.




Figure 6: Fundamental domain for $\Gamma(2)$, the duality group of the massless $N_f = 2$ theory.

In $N_f = 4$, the fundamental domain for the effective coupling τ can be found analogously to the asymptotically free cases. One notable new feature is the additional singularity of the theory when τ approaches the UV coupling τ_{UV} . For example, with the mass configuration $\mathbf{m} = (m, m, 0, 0)$ one finds [54]

$$u(\tau, \tau_{\text{UV}}) = -\frac{m^2}{3} \vartheta_3(\tau_{\text{UV}})^4 \frac{\lambda(\tau_0)^2 + 2(\lambda(\tau) - 1)\lambda(\tau_0) - \lambda(\tau)}{\lambda(\tau_0) - \lambda(\tau)}, \quad (2.35)$$

where $\lambda = \frac{\vartheta_2^4}{\vartheta_3^4}$. This function is a meromorphic $\Gamma(2)$ modular form of weight 2 in




Figure 7: Fundamental domain of the $N_f = 2$ theory with equal AD mass $m_1 = m_2 = m_{\text{AD}} = \frac{\Lambda_2}{2}$. It is the modular fundamental domain for the congruence subgroup $\Gamma_0(2)$ of $\text{PSL}(2, \mathbb{Z})$. The AD point $\tau_{\text{AD}} = 1 + i$ is the elliptic fixed point of $\Gamma_0(2)$. The three singularities collided in u_{AD} results in a reduction of the index from 6 to 3.

τ_{UV} for fixed τ , and of weight 0 in τ for fixed τ_{UV} . Furthermore, it has mixed weight $(0, 2)$ under $\text{PSL}(2, \mathbb{Z})$ acting on τ and τ_{UV} simultaneously. As such, it is an example of a *bimodular* form for the triple $(\Gamma(2), \Gamma(2); \text{PSL}(2, \mathbb{Z}))$ [26, 54]. Since $u(\tau, \tau_{\text{UV}})$ has a pole as $\tau \rightarrow \tau_{\text{UV}}$, the fundamental domain $\mathcal{F}_4(\mathbf{m})$ has a puncture at τ_{UV} . This is depicted in Fig. 8.




Figure 8: Fundamental domain of the $N_f = 4$ theory with mass $\mathbf{m} = (m, m, 0, 0)$. The six singularities on the Coulomb branch \mathcal{B}_4 are described by the three cusps, each of width 2. The I_0^* singularity corresponding to $u = \infty$ sits at $\tau = \tau_{\text{UV}}$.

2.5 Monodromies

This section determines the monodromies around the $N_f + 2$ monodromies. We leave the winding numbers $n_j, j = 1, \dots, N_f$, for a_D generic. Starting with the monodromy around infinity, $a \rightarrow e^{\pi i} a$, we deduce from the (2.7) that the vector Π transforms as $\Pi \rightarrow \mathbf{M}_\infty \Pi$, with \mathbf{M}_∞ given by

$$\mathbf{M}_\infty = \begin{pmatrix} -1 & 4 - N_f & 0 & -n_1 & \cdots & 0 & -n_{N_f} \\ 0 & -1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & n_1 & 1 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & & & \ddots & & \\ 0 & n_{N_f} & 0 & 0 & \cdots & 1 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}. \quad (2.36)$$

The monodromy matrix \mathbf{M}_∞ is in $\text{SL}(2 + 2N_f, \mathbb{Z})$, while it acts on the couplings by a symplectic transformation, i.e. it preserves the symplectic form (2.10). This can be checked by requiring that any monodromy \mathbf{M}_∞ satisfies $\mathbf{M}^T \mathbf{J} \mathbf{M} = \mathbf{J}$, with

$$\mathbf{J} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{\oplus N_f + 1}. \quad (2.37)$$

The action on the couplings τ (2.11), v_j and w_{jk} (2.12) is thus

$$\mathbf{M}_\infty : \begin{cases} \tau \rightarrow \tau + N_f - 4, \\ v_j \rightarrow -v_j - n_j, \\ w_{jk} \rightarrow w_{jk} + \delta_{jk}, \end{cases} \quad (2.38)$$

with δ_{jk} the Kronecker delta.

If we assume that the mass m_j is large, we can also deduce the monodromies around $a = \frac{m_j}{\sqrt{2}}, j = 1, \dots, N_f$ from the perturbative prepotential (2.7). For a encircling $\frac{m_1}{\sqrt{2}}$ counterclockwise, $\Pi \rightarrow \mathbf{M}_1 \Pi$, we find for the monodromy matrix \mathbf{M}_1 ,

$$\mathbf{M}_1 = \begin{pmatrix} 1 & 1 & 0 & -1 & \cdots & 0 & 0 \\ 0 & 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & -1 & 1 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & & & \ddots & & \\ 0 & 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}, \quad (2.39)$$

while the \mathbf{M}_j for other values of j are given by permutations. Its action on the couplings is

$$\mathbf{M}_j : \begin{cases} \tau \rightarrow \tau + 1, \\ v_k \rightarrow v_k - \delta_{jk}, \\ w_{kl} \rightarrow w_{kl} + \delta_{kl}\delta_{jl}. \end{cases} \quad (2.40)$$

To visualise the monodromies with large $\text{Im}(\tau)$ for $N_f = 1$, we refer to Fig. 4(d). In this regime of the masses, there is one monodromy with periodicity 3 and one with periodicity 1.

Besides the monodromies \mathbf{M}_∞ and \mathbf{M}_j , there are monodromies \mathbf{M}_m and \mathbf{M}_d around the points where a monopole and a dyon becomes massless, respectively. By requiring that the electro-magnetic charges of the massless particles are $(n_m, n_e) = (1, 0)$ and $(1, -2)$, respectively, we can fix the upper left blocks of the monodromies. We fix the remaining entries by assuming that the masses remains invariant, $m_j \rightarrow m_j$, and that the other periods only change by a multiple of the vanishing cycle at the corresponding cusp, together with the requirement that

$$\mathbf{M}_\infty = \mathbf{M}_m \mathbf{M}_d \prod_{j=1}^{N_f} \mathbf{M}_j. \quad (2.41)$$

For $N_f = 1$ and $n_1 = n$, this gives for \mathbf{M}_m ,

$$\mathbf{M}_m = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & -(n+1)/2 \\ (n+1)/2 & 0 & 1 & (n+1)^2/4 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.42)$$

This acts on the couplings as

$$\mathbf{M}_m : \begin{cases} \tau \rightarrow \frac{\tau}{-\tau+1}, \\ v \rightarrow \frac{v+(n+1)\tau/2}{-\tau+1}, \\ w \rightarrow w + \frac{(v+(n+1)/2)^2}{-\tau+1}. \end{cases} \quad (2.43)$$

The monodromy \mathbf{M}_d around the dyon singularity for $N_f = 1$ is

$$\mathbf{M}_d = \begin{pmatrix} -1 & 4 & 0 & -n-1 \\ -1 & 3 & 0 & -(n+1)/2 \\ (n+1)/2 & -n-1 & 1 & (n+1)^2/4 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (2.44)$$

This acts on the couplings as

$$\mathbf{M}_d : \begin{cases} \tau \mapsto \frac{-\tau+4}{-\tau+3}, \\ v \mapsto \frac{v+(n+1)\tau-n-1}{-\tau+3}, \\ w \mapsto w + \frac{(v+(n+1)/2)^2}{-\tau+3}. \end{cases} \quad (2.45)$$

We can note that all the above monodromy matrices leave the symplectic form (2.10) invariant and are independent of the masses.

For a small mass m , the fourth “hypermultiplet” cusp of the fundamental domain for $N_f = 1$ lies naturally near the real axis, $\tau \rightarrow 1$. See for example Fig. 4(a). Having determined \mathbf{M}_∞ , \mathbf{M}_m and \mathbf{M}_d , we can easily determine the monodromy $\tilde{\mathbf{M}}_1$ in this regime as

$$\tilde{\mathbf{M}}_1 = \mathbf{M}_m^{-1} \mathbf{M}_\infty \mathbf{M}_d^{-1} = \begin{pmatrix} 0 & 1 & 0 & (1-n)/2 \\ -1 & 2 & 0 & (1-n)/2 \\ (n-1)/2 & (1-n)/2 & 1 & (n-1)^2/4 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (2.46)$$

Thus for $n = -1$, the massless particle has charge $\pm(-1, 1, 0, 1)$.

We get similar monodromies for $N_f = 2, 3$. The action on the running couplings τ are the same for all N_f , by construction. The transformations of v_j and w_{jk} also take the same form for all N_f and can be summarised as

$$\begin{aligned} \mathbf{M}_m : & \begin{cases} v_j & \rightarrow \frac{v_j + (n_j + 1)\tau/2}{-\tau + 1}, \\ w_{jk} & \rightarrow w_{jk} + \frac{(v_j + (n_j + 1)/2)(v_k + (n_k + 1)/2)}{-\tau + 1}, \end{cases} \\ \mathbf{M}_d : & \begin{cases} v_j & \rightarrow \frac{v_j + (n_j + 1)\tau - n_j - 1}{-\tau + 3}, \\ w_{jk} & \rightarrow w_{jk} + \frac{(v_j + (n_j + 1)/2)(v_k + (n_k + 1)/2)}{-\tau + 3}. \end{cases} \end{aligned} \quad (2.47)$$

3 The UV theory on a four-manifold

We review various aspects of the formulation of the UV theory on a compact smooth four-manifold.

3.1 Aspects of four-manifolds

We let X be a smooth, compact, oriented Riemannian four-manifold, with Euler number $\chi = \chi(X)$ and signature $\sigma = \sigma(X) = b_2^+ - b_2^-$. The u -plane integral is non-vanishing only for four-manifolds X with $b_2^+ \leq 1$. In this article, we consider manifolds with $b_2^+ = 1$. Such four-manifolds admit a linear complex structure \mathcal{J} on the tangent space TX_p at each point p of X . The complex structure varies smoothly on X , such that TX is a complex bundle. We introduce furthermore the canonical class $K_X = -c_1(TX)$ of X , with $c_1(TX)$ the first Chern class of TX . For a manifold X with $(b_1, b_2^+) = (0, 1)$, we have that

$$K_X^2 = 8 + \sigma(X). \quad (3.1)$$

The middle cohomology $H^2(X, \mathbb{Z})$ of X gives rise to the uni-modular lattice L . More precisely, we identify L with the natural embedding of $H^2(X, \mathbb{Z})$ in $H^2(X, \mathbb{Z}) \otimes$

\mathbb{R} , which mods out the torsion of $H^2(X, \mathbb{Z})$. A characteristic element $K \in L$ is an element which satisfies $\mathbf{l}^2 + B(K, \mathbf{l}) \in 2\mathbb{Z}$ for all $\mathbf{l} \in L$. The Riemann-Roch theorem demonstrates that the canonical class K_X of X is a characteristic element of L . The Wu formula furthermore shows that any characteristic vector K of L is a lift of $w_2(X)$.

The quadratic form Q of the lattice L for a 4-manifold with $(b_1, b_2^+) = (0, 1)$ can be brought to a simple standard form depending on whether Q is even or odd [87]. This divides such manifolds into two classes, for which the evaluation of their u -plane integrals needs to be done separately [49, 88]. The period point $J \in H^2(X, \mathbb{R})$ is defined as the unique class in the forward light cone of $H^2(X, \mathbb{R})$ that satisfies $J = *J$ and $J^2 = 1$.

All four-manifolds without torsion and even intersection form admit a Spin structure. More generally, for any oriented four-manifold one can define a Spin^c-structure. The group Spin^c(4) can be defined as pairs of unitary 2×2 matrices with coinciding determinant,

$$\text{Spin}^c(4) = \{(u_1, u_2) \in U(2) \times U(2) \mid \det u_1 = \det u_2\}. \quad (3.2)$$

There exists a short exact sequence

$$1 \longrightarrow U(1) \longrightarrow \text{Spin}^c(4) \longrightarrow SO(4) \longrightarrow 1. \quad (3.3)$$

A Spin^c-structure \mathfrak{s} on a four-manifold X is then a reduction of the structure group of the tangent bundle on X , i.e. $SO(4)$, to the group $\text{Spin}^c(4)$. The different Spin^c-structures correspond to the inequivalent ways of choosing transition functions of the tangent bundle such that the cocycle condition is satisfied. The Spin^c-structure defines two rank two hermitian vector bundles W^\pm . We let $c(\mathfrak{s})$ be the first Chern class of the determinant bundles, $c(\mathfrak{s}) := c_1(\det W^\pm) \in H^2(X, \mathbb{Z})$.

If \mathfrak{s} is the canonical Spin^c structure associated to an almost complex structure on X , then $c(\mathfrak{s})^2 = 2\chi + 3\sigma$. More generally,

$$c_1(\mathfrak{s})^2 \equiv \sigma \pmod{8}. \quad (3.4)$$

3.2 Topological twisting with background fluxes

We discuss in this section topological twisting of theories with fundamental hypermultiplets including background fluxes. The discussion is parallel to the case of $\mathcal{N} = 2^*$ [26], where the hypermultiplet is in the adjoint representation of the gauge group.

We let $(E \rightarrow X, \nabla)$ be a principal $SU(2)/\mathbb{Z}_2 \cong SO(3)$ -bundle with connection ∇ . The second Stiefel-Whitney class $w_2(E) \in H^2(X, \mathbb{Z}_2)$ measures the obstruction to lift E to an $SU(2)$ bundle, which will exist locally but not globally if $w_2(E) \neq 0$. We denote a lift of $w_2(E)$ to the middle cohomology lattice L by $\bar{w}_2(E) \in L$, and

define the 't Hooft flux $\boldsymbol{\mu} = \bar{w}_2(E)/2 \in L/2$. The instanton number of the principal bundle is defined as $k = -\frac{1}{4} \int_X p_1(E)$ and satisfies $k \in -\boldsymbol{\mu}^2 + \mathbb{Z}$, where p_1 is the first Pontryagin class.

To formulate the theories with N_f fundamental hypermultiplets on a compact four-manifold, we perform a topological twist. Coupling the four-dimensional $\mathcal{N} = 2$ SU(2) theory to background fields means choosing two sets of data:

- A principal SU(2)_R R-symmetry bundle, with connection ∇_R ,
- and a principal bundle \mathcal{L} with connection for global symmetries (the flavour symmetries) [26].

The relevant twist for the $\mathcal{N} = 2$ supersymmetry algebra in four dimensions is the Donaldson-Witten twist. This twist is the local identification of the SU(2)₊ with the diagonal subgroup of the SU(2)₊ × SU(2)_R factor of the spin lift of the local spin group Spin(4) \cong SU(2)₊ × SU(2)₋ [1]. Alternatively, one can view the fields as sections of a non-trivial R-symmetry bundle, isomorphic to the spin bundle S^+ . Application of this to the representations of the vector multiplet (2.1) and (2.4) gives:

$$\begin{aligned} \text{bosons:} & \quad (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}), \\ \text{fermions:} & \quad (\mathbf{2}, \mathbf{2}) \oplus (\mathbf{3}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}). \end{aligned} \tag{3.5}$$

Thus the bosons remain unchanged, a vector and a complex scalar, while the fermions reorganise to a vector, self-dual two-form and real scalar, which we denote as ψ , χ and η , respectively. We note that none of these fields are spinors, and can thus be considered on a non-spin four-manifold. The original supersymmetry generators also transform in the representations for the fermions above. Thus the theory contains a scalar fermionic supercharge $\mathcal{Q} = \epsilon^{\dot{A}\dot{B}} \bar{\mathcal{Q}}_{\dot{A}\dot{B}}$, whose cohomology provides the operators in the topological theory [1].

For the fields of a hypermultiplet, (2.3) and (2.4), one finds

$$\begin{aligned} \text{bosons:} & \quad (\mathbf{1}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{2}), \\ \text{fermions:} & \quad (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}). \end{aligned} \tag{3.6}$$

Thus hypermultiplet bosons become spinors, i.e. sections of the spin bundle S^+ , while the fermions are sections of S^+ and S^- . Thus the twisted hypermultiplets can in this case only be formulated on four-manifolds which are spin, i.e. $w_2(X) = 0$ [12, 16].

However, if the hypermultiplets are charged under a gauge field or flux, the product of these bundles with S^\pm may be a Spin^c bundle, W^+ or W^- [16, 26, 39]. The latter are defined for arbitrary four-manifolds. For example, an almost complex structure on X determines two canonical Spin^c bundles $W^\pm \simeq S^\pm \otimes K_X^{-1/2}$ with K_X the canonical class determined by the almost complex structure. Since the

hypermultiplets are in the fundamental, two-dimensional representation of $SU(2)$, the topologically twisted hypermultiplets are well-defined on a non-spin four-manifold if $\boldsymbol{\mu} = -K_X/2$ [12].

Let us state this also in terms of the gauge bundle E . To this end, we label the two components of the fundamental, two-dimensional representation of $SU(2)$ by \pm . The two components are sections of a line bundle $\mathcal{L}_E^{\pm 1/2}$ with $c_1(\mathcal{L}_E) = \bar{w}_2(E)$. Of course, the square root $\mathcal{L}_E^{1/2}$ only exists if $w_2(E) \in 2L$. On the other hand, the physical requirement is that $S^+ \otimes \mathcal{L}_E^{1/2}$ is well defined, or $\bar{w}_2(X) + \bar{w}_2(E) \in 2L$. Therefore, the obstructions can cancel each other for a suitable choice of $w_2(E)$. Thus the topological twisted theory is not well-defined for an arbitrary choice of 't Hooft flux $\boldsymbol{\mu} := \frac{1}{2}\bar{w}_2(E)$; but rather $\boldsymbol{\mu}$ has to satisfy $\boldsymbol{\mu} = \frac{1}{2}\bar{w}_2(X) \pmod{L}$ [12], or

$$\bar{w}_2(X) = \bar{w}_2(E) \pmod{2L}. \quad (3.7)$$

To consider more general 't Hooft fluxes $\boldsymbol{\mu}$ or equivalently $w_2(E)$, we can couple the j 'th hypermultiplet to a background flux or line bundle \mathcal{L}_j , with \mathcal{L}_j possibly different for each j . We let $\mathcal{E}_j = \mathcal{L}_E \otimes \mathcal{L}_j$. Then the requirement that $S^\pm \otimes \mathcal{E}_j^{\pm 1/2}$ is globally well-defined is that

$$c_1(\mathcal{E}_j) \in \bar{w}_2(X) + 2L, \quad (3.8)$$

which can be satisfied for any $\bar{w}_2(E)$ for a suitable choice of \mathcal{L}_j . Thus we can formulate the u -plane integral for arbitrary $\bar{w}_2(E)$, if we require that the background fluxes satisfy

$$c_1(\mathcal{L}_j) \in \bar{w}_2(X) + \bar{w}_2(E) + 2L, \quad (3.9)$$

for each j . This is consistent with (3.7) for $c_1(\mathcal{L}_j) = 0$.

The Chern classes $c_1(\mathcal{L}_j)$ can also be seen as the splitting classes of the $\text{Spin}(2N_f)$ principal bundle \mathcal{L} . The Chern class of \mathcal{L} reads

$$c(\mathcal{L}) = \sum_{l=0}^2 c_l(\mathcal{L}) = \prod_{j=1}^{N_f} (1 + c_1(\mathcal{L}_j)). \quad (3.10)$$

The scalar generators of the equivariant cohomology of $\text{Spin}(2N_f)$ are the masses m_j , which generate the N_f -dimensional Cartan subalgebra of $\text{Spin}(2N_f)$. The gauge bundle E_k is also $\text{Spin}(2N_f)$ equivariant. For generic masses, the flavour group is $U(1)^{N_f}$, and is enhanced for special loci of the masses, for example to $U(N_f)$ for equal masses [11].

The \mathcal{Q} -fixed equations are the non-Abelian monopole equations with N_f matter fields in the fundamental representation. For generic gauge group G and with

representation R , these equations read [15]

$$\begin{aligned} \left(F_{\dot{\alpha}\dot{\beta}}^a\right)^+ + \frac{i}{2} \sum_{j=1}^{N_f} \bar{M}_{(\dot{\alpha}}^j T^a M_{\dot{\beta})}^j &= 0, \\ \not{D}M^j = \sum_{\mu} \sigma^{\mu} D_{\mu}M^j &= 0, \end{aligned} \tag{3.11}$$

where T^a is a generator of the Lie algebra in the representation R . Including the sum over matrix elements, we have

$$M_{(\alpha}^j T^a M_{\beta)}^j = \sum_{k,l} (M^j)_{(\alpha}^k (T^a)^{kl} (M^j)_{\beta)}^l. \tag{3.12}$$

We denote the moduli space of solutions to (3.11) by $\mathcal{M}_{k,\mathcal{L}_j}^{Q,N_f}$, and leave the dependence on the 't Hooft flux $\boldsymbol{\mu}$ and the metric J implicit. For $N_f = 4$ on $X = \mathbb{CP}^2$, such moduli spaces are studied in [32].

The moduli spaces $\mathcal{M}_{k,\mathcal{L}_j}^{Q,N_f}$ is non-compact for vanishing masses [25, 36, 89]. This is improved upon turning on masses and localizing with respect to the $U(1)^{N_f}$ flavour symmetry, $M_{\alpha}^j \rightarrow e^{i\varphi_j} M_{\alpha}^j$, which leave invariant the \mathcal{Q} -fixed equations (3.11). There are two components:

- the instanton component, with $F^+ = 0$ and $M^j = 0$, $j = 1, \dots, N_f$. The moduli space for this component is denoted \mathcal{M}_k^i . Since the hypermultiplet fields vanish, this component is associated to the Coulomb branch.
- the abelian or monopole component, for which a $U(1)$ subgroup of the flavour group acts as pure gauge. Here the connection is reducible, and a $U(1)$ subgroup of the $SU(2)$ gauge group is preserved. For generic masses, there are N_f such components, where M^{ℓ} is upper or lower triangular for some ℓ , and $M^j = 0$ for all $j \neq \ell$. The moduli space of this component is denoted $\mathcal{M}_k^{a,j}$, $j = 1, \dots, N_f$. Since some of the hypermultiplet fields are non-vanishing, this component is associated to the Higgs branch [22, 90].

The instanton component \mathcal{M}_k^i is non-compact due to point-like instantons. This can be cured using the Uhlenbeck compactification or algebraic-geometric compactifications. We assume that the physical path integral chooses a specific compactification, whose details are however not manifest at the level of the low energy effective field theory other than that the compactification must be in agreement with the correlation functions.

The topological twist for $\mathcal{N} = 2$ supersymmetric QCD can be further made dependent on a cocycle $\zeta_{\alpha\beta\gamma}^{\text{gauge}}$ representing the 't Hooft flux, and $\zeta_{\alpha\beta\gamma}^s$ a cocycle representing $w_2(X)$ (the cocycles are the $U(1)$ valued functions measuring the obstruction of the cocycle condition for transition functions) [26]. Without additional

line bundles, $\bar{w}_2(X) = \bar{w}_2(E)$ is equivalent to the cocycle $\zeta_{\alpha\beta\gamma}^{\text{gauge}} \zeta_{\alpha\beta\gamma}^s$ being trivialisable. We leave it for future work to explore whether the invariants depend on the choice of trivialisation.

3.3 Correlation functions and moduli spaces

The \mathcal{Q} -fixed equations (3.11) include a Dirac equation for each hypermultiplet $j = 1, \dots, N_f$ in the fundamental representation. The corresponding index bundle W_k^j defines an element of the K -group of \mathcal{M}_k^i . Its virtual rank $\text{rk}(W_k^j)$ is the formal difference of two infinite dimensions. It is given by an index theorem and reads

$$\text{rk}(W_k^j) = -k + \frac{1}{4}(c_1(\mathcal{L}_j)^2 - \sigma) \in \mathbb{Z}, \quad (3.13)$$

where $c_1(\mathcal{L}_j)$ is the first Chern class of the bundle \mathcal{L}_j . Note that the rhs is *not* an integer for an arbitrary $c_1(\mathcal{L}_j) \in H^2(X, \mathbb{Z})$. To verify that the rhs is integral for the $c_1(\mathcal{L}_j)$'s satisfying (3.9), we rewrite $\text{rk}(W_k^j)$ as

$$\text{rk}(W_k^j) = -(k + \boldsymbol{\mu}^2) - c_1(\mathcal{L}_j) \cdot \boldsymbol{\mu} + \frac{1}{4}((c_1(\mathcal{L}_j) + 2\boldsymbol{\mu})^2 - \sigma). \quad (3.14)$$

Then the first term on the rhs is an integer since $k \in -\frac{1}{4}w_2(E)^2 + \mathbb{Z}$ for an $\text{SO}(3)$ bundle. The second term is an integer because $c_1(\mathcal{L}_j) \cdot \boldsymbol{\mu} = (\bar{w}_2(X) - 2\boldsymbol{\mu}) \cdot \boldsymbol{\mu} \pmod{\mathbb{Z}} \in \mathbb{Z}$, and the third term is an integer using (3.4) and the fact that $c_1(\mathcal{L}_j) + 2\boldsymbol{\mu}$ equals the characteristic class of a Spin^c -structure \mathfrak{s}_j by (3.9),

$$c_1(\mathcal{L}_j) + 2\boldsymbol{\mu} = c(\mathfrak{s}_j), \quad (3.15)$$

for each j .

The mass m_j is the equivariant parameter of the $U(1)$ flavour symmetry associated to the j 'th hypermultiplet. The equivariant Chern class of W_k^j reads in terms of the splitting class x_l ,

$$c(W_k^j) = \prod_{l=0}^{-\text{rk}(W_k^j)} (x_l + m_j) = m_j^{-\text{rk}(W_k^j)} \sum_l \frac{c_l(W_k^j)}{m_j^l}. \quad (3.16)$$

We abbreviate $c_l(W_k^j)$ to $c_{l,j}$, and let $c(W_k) = \prod_{j=1}^{N_f} c(W_k^j)$.

The moduli space $\mathcal{M}_{k,\boldsymbol{\mu},\mathcal{L}_j}^{\mathcal{Q}}$ for N_f hypermultiplets corresponds to the vanishing locus of the obstructions for the existence of N_f zero modes of the Dirac operator. As a result, the virtual complex dimension of the moduli space $\mathcal{M}_{k,\boldsymbol{\mu},\mathcal{L}_j}^{\mathcal{Q}}$ is that of the instanton moduli space plus the sum of (typically negative) ranks of the index bundles W_k^j , $\text{vdim}(\mathcal{M}_{k,\boldsymbol{\mu},\mathcal{L}_j}^{\mathcal{Q},N_f}) = \text{vdim}(\mathcal{M}_k^{\mathcal{Q}})_{N_f=0} + \sum_{j=1}^{N_f} \text{rk}(W_k^j)$ [17, 26, 31, 36]. This gives

$$\text{vdim}(\mathcal{M}_{k,\boldsymbol{\mu},\mathcal{L}_j}^{\mathcal{Q},N_f}) = (4 - N_f)k + \frac{1}{4} \left(-3\chi - (3 + N_f)\sigma + \sum_{j=1}^{N_f} c_1(\mathcal{L}_j)^2 \right). \quad (3.17)$$

It is argued in [13] that the inclusion of massive matter amounts to inserting an integral of the equivariant Euler class of the Dirac index bundle over the moduli space. Therefore, the correlation functions are the generating functions for the intersection numbers of the standard Donaldson observables and the Poincaré duals to the Chern classes of the various vector bundles.

The correlation functions on X in the theory with N_f massive fundamental hypermultiplets are conjectured to be

$$\langle \mathcal{O}_1 \dots \mathcal{O}_p \rangle = \sum_k \Lambda_{N_f}^{\text{vdim}(\mathcal{M}_{k, \mathcal{L}_j}^{Q, N_f})} \int_{\mathcal{M}_{k, \mathcal{L}_j}^{Q, N_f}} c(W_k) \omega_1 \wedge \dots \wedge \omega_p, \quad (3.18)$$

where $\omega_i = \mu(\mathcal{O}_i)$ are the Donaldson classes associated to the physical observable \mathcal{O}_i , and $c(M)$ is the Euler class of the matter bundle [91–94]. Localising to the fixed point locus in $\mathcal{M}_{k, \mathcal{L}_j}^{Q, N_f}$ with respect to $U(1)^{N_f}$ gives

$$\langle \mathcal{O}_1 \dots \mathcal{O}_p \rangle = \sum_k \Lambda_{N_f}^{\text{vdim}(\mathcal{M}_{k, \mathcal{L}_j}^{Q, N_f})} \int_{\mathcal{M}_k^i \cup \mathcal{M}_k^a} \left(\prod_{j=1}^{N_f} m_j^{-\text{rk}(W_k^j)} \sum_l \frac{\sigma_{j,l}}{m_j^l} \right) \omega_1 \wedge \dots \wedge \omega_p. \quad (3.19)$$

where the integral is over the union $\mathcal{M}_k^i \cup \mathcal{M}_k^a$ of the instanton component \mathcal{M}_k^i [13, Eq. (5.13)] and the monopole component \mathcal{M}_k^a [26, 95]. The equation together with the dimension of the moduli spaces (3.17) demonstrate a selection rule for observables together with powers of Λ_{N_f} and m_j .

In the decoupling limit $m_{N_f} \rightarrow \infty$, $\Lambda_{N_f} \rightarrow 0$ (2.18), the only contribution for $j = N_f$ is from $l = 0$, $c_{0, N_f} = 1$. The powers of m_j and Λ_{N_f} work out such that the correlation functions reduce to those of the theory with $N_f - 1$ hypermultiplets [21]

$$\langle \mathcal{O}_1 \dots \mathcal{O}_p \rangle_{N_f} \rightarrow \left(\frac{\Lambda_{N_f}}{\Lambda_{N_f-1}} \right)^\alpha \langle \mathcal{O}_1 \dots \mathcal{O}_p \rangle_{N_f-1}, \quad (3.20)$$

We deduce from (3.19) that

$$\begin{aligned} \alpha &= \text{vdim}(\mathcal{M}_k^{Q, N_f}) + (4 - N_f) \text{rk}(W_k^{N_f}) \\ &= \frac{1}{4} \left(-3\chi - 7\sigma + (5 - N_f) c_1(\mathcal{L}_{N_f})^2 + \sum_{j=1}^{N_f-1} c_1(\mathcal{L}_j)^2 \right). \end{aligned} \quad (3.21)$$

The overall factor can be accounted for by an overall renormalization in the decoupling limit.

The correlation function has a smooth massless limit $m_j \rightarrow 0$, for which only terms with the top Chern classes contribute. These are given by

$$\langle \mathcal{O}_1 \dots \mathcal{O}_p \rangle = \sum_k \prod_{j=1}^{N_f} \Lambda_{N_f}^{\text{vdim}(\mathcal{M}_{k, \mathcal{L}_j}^{Q, N_f})} \int_{\mathcal{M}_k^i \cup \mathcal{M}_k^a} c_{l_j, j} \omega_1 \wedge \dots \wedge \omega_p \quad (3.22)$$

with $l_j = -\text{rk}(W_k^j)$ for each $c_{i,j}$. For a non-vanishing result, the degree of $\omega_1 \dots \omega_p$ must equal $\text{vdim}(\mathcal{M}_{k,\mathcal{L}_j}^{Q,N_f})$.

4 The effective theory on a four-manifold

We consider in this Section the low energy effective field theory on a four-manifold. We derive the semi-classical action of the theory coupled to background U(1) fields. As in previous cases [12, 26, 55, 96], the final expression takes the form of a Siegel-Narain theta series multiplied by a measure factor.

4.1 Hypermultiplets and background fields

The effective theory coupled to N_f background fluxes can be modelled as that of a theory with gauge group $\text{SU}(2) \times \text{U}(1)^{N_f}$, where the fields of the U(1) factors have been frozen in a special way [26, 97]. To derive the precise form, we recall the low-energy effective Lagrangian for the r multiplets $(\phi^J, \eta^J, \chi^J, \psi^J, F^J)$ of the topologically twisted $\text{U}(1)^r$ SYM theory [98]. Since the u -plane integral reduces to an integral over zero-modes [12], it suffices to only include the zero-modes in the Lagrangian. For simply connected four-manifolds, there is no contribution from the one-form fields ψ^J . The Lagrangian is then given in terms of the prepotential $F(\{a^J\})$ and its derivatives to the vevs $\langle \phi^J \rangle = a^J$, as

$$\begin{aligned} \mathcal{L} = & \frac{i}{16\pi} (\bar{\tau}_{JK} F_+^J \wedge F_+^K + \tau_{JK} F_-^J \wedge F_-^K) - \frac{1}{8\pi} y_{JK} D^J \wedge D^K \\ & + \frac{i\sqrt{2}}{16\pi} \bar{\mathcal{F}}_{JKL} \eta^J \chi^K \wedge (D + F_+)^L, \end{aligned} \quad (4.1)$$

with $y_{JK} = \text{Im}(\tau_{JK})$, $\tau_{JK} = \partial_J \partial_K F(\{a^J\})$ and $\mathcal{F}_{JKL} = \partial_J \partial_K \partial_L F(\{a^J\})$. It is left invariant by the BRST operator Q , which acts on the zero modes as

$$\begin{aligned} [Q, A^J] &= \psi^J = 0, & [Q, \psi^J] &= 4\sqrt{2} da^J, \\ [Q, a^J] &= 0, & [Q, \bar{a}^J] &= \sqrt{2} i \eta^J, \\ [Q, \eta^J] &= 0, & [Q, \chi^J] &= i(F_+ - D_+)^J, \\ [Q, D^J] &= (d\psi^J)_+ = 0. \end{aligned} \quad (4.2)$$

Using this operator, we can write \mathcal{L} as the sum of a topological, holomorphic term and a Q -exact term,

$$\mathcal{L} = \frac{i}{16\pi} \tau_{JK} F^J \wedge F^K + \{Q, W\}, \quad (4.3)$$

with

$$W = -\frac{i}{8\pi} y_{JK} \chi^J (F_+ + D)^K. \quad (4.4)$$

The low-energy theory of $\text{SU}(2)$ gauge theory with N_f hypermultiplets coupled to N_f background fluxes can then be modelled by the above rank r description with

$r = N_f + 1$. We identify $F(\{a^J\})$ with $F(a, \mathbf{m})$. We let the indices J, K run from 0 to N_f and identify the index 0 with the unbroken $U(1)$ of the $SU(2)$ gauge group and the indices $j, k, l = 1, \dots, N_f$ with that of the frozen $U(1)^{N_f}$ factors. We further set $\phi^0 := \phi$ for any field ϕ . We will proceed by using lower indices for j, k, l , except where the summation convention is explicitly used, to avoid confusion with powers of the fields.

The masses of the hypermultiplets are the vevs of the frozen scalar fields of the corresponding vector multiplets, $\frac{m_j}{\sqrt{2}} = \langle \phi_j \rangle = a_j$ [97]. We set $[F_j] = 4\pi \mathbf{k}_j$ with

$$\mathbf{k}_j = c_1(\mathcal{L}_j)/2 \in L/2. \quad (4.5)$$

To make the BRST variations of the fields from the frozen $U(1)$ factors vanish, we set $\eta_j = \chi_j = 0$, as well as $D^j = F_+^j$. With these identifications, the Lagrangian becomes

$$\begin{aligned} \mathcal{L} = & \frac{i}{16\pi} \tau_{JK} F^J \wedge F^K + \frac{1}{8\pi} y_{00} F_+ \wedge F_+ - \frac{1}{8\pi} y_{00} D \wedge D \\ & + \frac{i\sqrt{2}}{16\pi} \bar{\mathcal{F}}_{000} \eta \chi \wedge (D + F_+) + \frac{i\sqrt{2}}{8\pi} \bar{\mathcal{F}}_{00j} \eta \chi \wedge F_+^j \\ & + \frac{1}{4\pi} y_{0j} (F_+ - D) \wedge F_+^j. \end{aligned} \quad (4.6)$$

Integrating over D, η and χ in the standard way [12, 26, 98], we end up with

$$\begin{aligned} & \int dD d\eta d\chi e^{-\int_X \mathcal{L}} \\ & = \frac{\partial}{\partial \bar{a}} \left(i\sqrt{y_{00}} B \left(F + \frac{y_{0j}}{y_{00}} F^j, J \right) \right) e^{-\int_X \mathcal{L}_0}, \end{aligned} \quad (4.7)$$

where

$$\begin{aligned} \mathcal{L}_0 = & \frac{i}{16\pi} \tau_{JK} F^J \wedge F^K + \frac{1}{8\pi} y_{00} F_+ \wedge F_+ + \frac{y_{0j}}{4\pi} F_+ \wedge F_+^j + \frac{1}{8\pi} \frac{y_{0j} y_{0k}}{y_{00}} F_+^j \wedge F_+^k \\ = & \frac{i}{16\pi} (\bar{\tau} F_+ \wedge F_+ + \tau F_- \wedge F_-) + \frac{i}{8\pi} (v_j F_- \wedge F_-^j + \bar{v}_j F_+ \wedge F_+^j) \\ & + \frac{i}{16\pi} w_{jk} F^j \wedge F^k + \frac{y}{8\pi} \text{Im}(v_j) \text{Im}(v_k) F_+^j \wedge F_+^k, \end{aligned} \quad (4.8)$$

and we identified $\tau := \tau_{00}$, $y = \text{Im}(\tau) = y_{00}$, $v_j := \tau_{0j}$ and $w_{jk} := \tau_{jk}$. Thus the coupling w_{jk} is holomorphic, but the coupling v_j is *non-holomorphic*. This is similar to the couplings for $\mathcal{N} = 2^*$ [26].

4.2 Sum over fluxes

The path integral includes a sum over fluxes $\mathbf{k} = [F]/4\pi \in L/2$. After summing the exponentiated action (4.7) over the fluxes \mathbf{k} and multiplying by $\frac{d\bar{a}}{d\bar{\tau}}$, we find that this takes the form

$$\sum_{\mathbf{k} \in L + \boldsymbol{\mu}} \int dD d\eta d\chi e^{-\int_X \mathcal{L}} = \left(\prod_{j,k=1}^{N_f} C_{jk}^{B(\mathbf{k}_j, \mathbf{k}_k)} \right) \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}). \quad (4.9)$$

The couplings C_{jk} are given in terms of w_{jk} (2.12) by

$$C_{jk} = e^{-\pi i w_{jk}}, \quad (4.10)$$

for $j, k = 1, \dots, N_f$. Such couplings were first put forward in [55], and were also crucial in [26].

The term Ψ_{μ}^J is an example of a Siegel-Narain theta function. It reads explicitly

$$\begin{aligned} \Psi_{\mu}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}) &= e^{-2\pi y \mathbf{b}_+^2} \sum_{\mathbf{k} \in L + \mu} \partial_{\bar{\tau}} (4\pi i \sqrt{y} B(\mathbf{k} + \mathbf{b}, J)) \\ &\times (-1)^{B(\mathbf{k}, K)} q^{-\mathbf{k}_-^2/2} \bar{q}^{\mathbf{k}_+^2/2} e^{-2\pi i B(\mathbf{z}, \mathbf{k}_-) - 2\pi i B(\bar{\mathbf{z}}, \mathbf{k}_+)}, \end{aligned} \quad (4.11)$$

and discussed in more detail in Appendix A.2. The elliptic variable reads in terms of v_j and \mathbf{k}_j ,

$$\mathbf{z} = \sum_{j=1}^{N_f} v_j \mathbf{k}_j, \quad \text{and} \quad \mathbf{b} = \frac{\text{Im}(\mathbf{z})}{y}, \quad (4.12)$$

thus inducing a non-holomorphic dependence on v_j . Furthermore, K appearing in the fourth root of unity $(-1)^{B(\mathbf{k}, K)}$ is a characteristic vector of L . Note that Ψ_{μ}^J changes by the sign $(-1)^{B(\mu, K - K')}$ upon replacing K by a different characteristic vector K' [12, 26, 99].

For $N_f = 0$, this phase can be understood as arising from integrating out massive fermionic modes [96]. It also appears naturally in decoupling the adjoint hypermultiplet in the analogous function for $\mathcal{N} = 2^*$ [26]. For $N_f > 0$, the constant part of the couplings v_j (2.12) effectively contribute to the phase, such that the total phase reads

$$e^{\pi i B(\mathbf{k}, K)} \prod_{j=1}^{N_f} e^{\pi i n_j B(\mathbf{k}_j, \mathbf{k})}, \quad (4.13)$$

with n_j the magnetic winding numbers. For arbitrary $n_j \in \mathbb{Z}$, the phase is an eighth root of unity. It would be interesting to understand this phase from integrating out massive modes.

We deduce from (4.13) that the summand of Ψ_{μ}^J changes by a phase

$$e^{\pi i (n'_j - n_j) B(\mathbf{k}_j, \mathbf{k})} \quad (4.14)$$

if the winding numbers n_j are replaced by n'_j . Since $\mathbf{k}_j \in K/2 - \mu \pmod{L}$ (see (3.9)) and $\mathbf{k} \in L + \mu$, this phase is 1 if $n'_j - n_j = 0 \pmod{4}$. We can therefore restrict to $n_j \in \mathbb{Z}_4$. For specific choices of μ and \mathbf{k}_j , the n_j can lie in a subgroup of \mathbb{Z}_4 .

The modular transformations of Ψ_{μ}^J are discussed in Appendix A.2, which are crucial input for single-valuedness of the u -plane integrand. We will demonstrate in Section 5.2 that the u -plane integrand is single-valued if we impose further constraints on the winding numbers n_j .

Finally, if the theory is considered on a curved background, topological couplings arise in the effective field theory [96]. These terms couple to the Euler characteristic and the signature of the four-manifold X , respectively denoted A and B . These take the form [12, 96],

$$A = \alpha \left(\frac{du}{da} \right)^{1/2}, \quad B = \beta \Delta_{N_f}^{1/8}. \quad (4.15)$$

Here, Δ_{N_f} is the physical discriminant incorporating the singularities of the effective theory, while $\frac{du}{da}$ is the (reciprocal of) the periods of the SW curves as introduced in Section 2. Both can be determined directly from the SW curve, as described in Section 2.3. The prefactors α and β are independent of u , but can be functions of other moduli such as the masses \mathbf{m} , the dynamical scale Λ_{N_f} or the UV coupling τ_{UV} . However, it turns out that for the theories with fundamental matter they are independent of the masses and only depend on the scale [21, 100]. They satisfy several constraints from holomorphy, RG flow, homogeneity and dimensional analysis, and can in principle be fixed for any Lagrangian theory from a computation in the Ω -background [21, 24, 70, 100].

4.3 Observables and contact terms

The observables in the topologically twisted theories are the point observable or 0-observable u , as well as d -observables supported on a d -dimensional submanifold of X . The d -observables are only non-vanishing if the submanifold corresponds to a non-trivial homology class. For $b_1 = 0$, the d -observables with d odd therefore do not contribute.

To introduce the surface observable, let $\mathbf{x} \in H_2(X, \mathbb{Q})$. Then the surface observable reads in terms of the UV fields,

$$I(\mathbf{x}) = \frac{1}{4\pi^2} \int_{\mathbf{x}} \text{Tr} \left[\psi \wedge \psi - \frac{1}{\sqrt{2}} \phi F \right]. \quad (4.16)$$

In the effective infrared theory, this operator becomes,

$$\tilde{I}(\mathbf{x}) = \frac{i}{\sqrt{2}\pi} \int_{\mathbf{x}} \frac{1}{32} \frac{d^2 u}{da^2} \psi \wedge \psi - \frac{\sqrt{2}}{4} \frac{du}{da} (F_- + D). \quad (4.17)$$

Generating functions of correlation functions are obtained by inserting

$$e^{p u / \Lambda_{N_f}^2 + \tilde{I}(\mathbf{x}) / \Lambda_{N_f}} \quad (4.18)$$

in the path integral. The surface observable leads to a change in the argument of the sum over fluxes (4.11),

$$\mathbf{z} \rightarrow \mathbf{z} + \frac{\mathbf{x}}{2\pi \Lambda_{N_f}} \frac{du}{da}, \quad \bar{\mathbf{z}} \rightarrow \bar{\mathbf{z}}. \quad (4.19)$$

and to analytically continue \mathbf{b} (4.12) to the complex number by setting $\mathbf{b} = (\mathbf{z} - \bar{\mathbf{z}})/(2iy)$.

The inclusion of the surface observable also gives rise to a contact term [2, 12, 93], which in particular ensures that the u -plane integrand is single-valued. For $0 \leq N_f \leq 3$, the contact term is $\exp(\mathbf{x}^2 G_{N_f})$ with [13, 23, 101]

$$G_{N_f} = -\frac{1}{24 \Lambda_{N_f}^2} E_2 \left(\frac{du}{da} \right)^2 + \frac{1}{3 \Lambda_{N_f}^2} \left(u + \frac{\Lambda_3^2}{64} \delta_{N_f,3} \right), \quad (4.20)$$

while for $N_f = 4$ it is given by [23, 102]

$$G_{N_f=4} = -\frac{1}{24 \Lambda_4^2} E_2 \left(\frac{du}{da} \right)^2 + \frac{u}{3 \Lambda_4^2} E_2(\tau_{UV}) + \frac{1}{18 \Lambda_4^2} \llbracket m_1^2 \rrbracket E_4(\tau_{UV}). \quad (4.21)$$

This expression (4.20) is valid for the theories with N_f arbitrary hypermultiplet masses. The reason for it is the following [23, 98, 103]: G is guaranteed to be $\bar{\mathcal{Q}}$ -closed and hence locally holomorphic. First, notice that $\frac{\partial F}{\partial \tau_0} = \frac{u}{4}$, where $\Lambda^{4-N_f} =: e^{\pi i \tau_0}$ for the asymptotically free theories ($N_f \leq 3$) and $\tau_0 = \tau_{UV}$ for $N_f = 4$. The real part of the exponential prefactor of Ψ_{μ}^J can be added to G to give a monodromy-invariant contribution \hat{G} which multiplies the intersection \mathbf{x}^2 . From the action of a duality transformation on \hat{G} it can be inferred that

$$G_{N_f} = -\frac{4i}{\pi \Lambda_{N_f}^2} \frac{\partial^2 F}{\partial \tau_0^2}. \quad (4.22)$$

The expressions (4.20) follow by direct computation. A more general scheme to fix the contact terms is proposed in [13]. Contact terms can also be derived from the corresponding Whitham hierarchies [101, 104]. In the presence of surface observables, there are additional mixed contact terms $\frac{\partial^2 \mathcal{F}}{\partial \tau_0 \partial m}$ for the external fluxes $\{\mathbf{k}_j\}$ as encountered in [26] for the $\mathcal{N} = 2^*$ theory.

5 The u -plane integral

In this section, we set up the u -plane integral schematically given in (1.2), and demonstrate that it is well-defined on the integration domain for any μ with appropriate background fluxes. The case $\mu = \bar{w}_2(X)/2$ and $\mathbf{k}_j = 0$ was analyzed in [12].

5.1 Definition of the integrand

As discussed in the previous sections, the u -plane integral on a closed four-manifold X with $(b_1, b_2^+) = (0, 1)$ depends on many parameters. We summarise:

- The scale Λ_{N_f} and masses $\mathbf{m} = (m_1, \dots, m_{N_f})$ of the theory. See Section 2.
- The magnetic winding numbers n_j , $j = 1, \dots, N_f$. See Section 2.2.

- The four-manifold X , in particular its signature $\sigma = \sigma(X)$, Euler characteristic $\chi = \chi(X)$, period point J and intersection form Q . See Section 3.1.
- The 't Hooft flux $\boldsymbol{\mu}$, and the external fluxes $\{\mathbf{k}_j\} = (\mathbf{k}_1, \dots, \mathbf{k}_{N_f})$. See Section 3.3.
- The fugacities for the point and surface observables p and \mathbf{x} . See Section 4.3.

In terms of these parameters, the u -plane path integral reduces to the following finite dimensional integral over $\mathcal{F}_{N_f}(\mathbf{m})$,

$$\boxed{\Phi_{\boldsymbol{\mu}, \{\mathbf{k}_j\}}^J(p, \mathbf{x}, \mathbf{m}, \Lambda_{N_f}) = \mathcal{K}_{N_f} \int_{\mathcal{F}_{N_f}(\mathbf{m})} d\tau \wedge d\bar{\tau} \nu(\tau; \{\mathbf{k}_j\}) \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}) e^{2pu + \mathbf{x}^2 G_{N_f}}.} \quad (5.1)$$

We summarise the different elements on the rhs:

- \mathcal{K}_{N_f} is an overall normalisation factor. For $N_f = 0$, it is fixed by matching to known Donaldson invariants. Due to $\chi + \sigma = 4$, there is an ambiguity [24]

$$(\mathcal{K}_{N_f}, \alpha, \beta) \sim (\zeta^{-4} \mathcal{K}_{N_f}, \zeta \alpha, \zeta \beta), \quad (5.2)$$

with α and β the u -independent prefactors in (4.15).

- The integration domain $\mathcal{F}_{N_f}(\mathbf{m})$ in (5.1) is crucially the fundamental domain of the effective gauge coupling. As discussed in Section 2.4, this domain requires new aspects compared to integration domains for earlier discussions of u -plane integrals. The evaluation of integrals over $\mathcal{F}_{N_f}(\mathbf{m})$ will be discussed in more detail in Section 6.
- ν is the “measure factor” [12, 13, 21, 26, 96]

$$\nu(\tau; \{\mathbf{k}_j\}) = \frac{da}{d\tau} A^\chi B^\sigma \prod_{i,j=1}^{N_f} C_{ij}^{B(\mathbf{k}_i, \mathbf{k}_j)}. \quad (5.3)$$

It combines the topological couplings (4.15) and the couplings to the background fluxes (4.10) with the Jacobian $\frac{da}{d\tau}$ of the change of variables from a to τ .

- The function $\Psi_{\boldsymbol{\mu}}^J$ arises from the sum over $U(1)$ fluxes. It is a Siegel-Narain theta function (4.11) and discussed in detail in Section 4.2. The elliptic parameter \mathbf{z} of the Siegel-Narain theta function reads

$$\begin{aligned} \mathbf{z} &= \frac{\mathbf{x}}{2\pi \Lambda_{N_f}} \frac{du}{da} + \sum_{j=1}^{N_f} v_j \mathbf{k}_j, \\ \bar{\mathbf{z}} &= \sum_{j=1}^{N_f} \bar{v}_j \mathbf{k}_j. \end{aligned} \quad (5.4)$$

- Finally, G_{N_f} is the contact term, discussed in more detail in Section 4.3.

While the path integral set up in Section 4 integrates the exponentiated action over the local coordinates a and \bar{a} , in (5.1) we have changed variables to τ and $\bar{\tau}$. This change of variables $(a, \bar{a}) \rightarrow (\tau, \bar{\tau})$ is valid as long as the Jacobian is nonsingular in the integration region. Since the coordinates a and \bar{a} are holomorphic and anti-holomorphic respectively, the Jacobian is diagonal and the functional determinant accordingly reads $\frac{da}{d\tau} \frac{d\bar{a}}{d\bar{\tau}}$. We thus need to show that $\frac{da}{d\tau}$ is not singular away from isolated points in $\mathcal{F}_{N_f}(\mathbf{m})$, which in (5.1) we remove implicitly from the integration domain.

Using $\frac{da}{du} = \frac{da}{du} \frac{du}{d\tau}$, we can study the singular points in detail. First, it is shown in [52] that the singularities of $\frac{da}{d\tau}$ are in one-to-one correspondence with the branch points. In fact, both $\frac{du}{d\tau} = 0$ and $\frac{du}{d\tau} = \infty$ are realised as branch points in $\mathcal{N} = 2$ SQCD. In the following Section 6, we remove a small circle in $\mathcal{F}_{N_f}(\mathbf{m})$ around the branch points, and show that they do not give an extra contribution. Furthermore, the solutions to $\frac{du}{da} = 0$ are shown to be the Argyres-Douglas (AD) points. We exclude them from the integration region, and study their contribution also in Section 6. Finally, we know that $\eta^{24} \propto \left(\frac{da}{du}\right)^{12} \Delta_{N_f}$ [52], with η the Dedekind eta-function as defined in (A.5). Since $\eta \neq 0$ and Δ_{N_f} does not have poles, we find that $\frac{da}{du}$ never vanishes. This agrees with the fact that $\frac{da}{du}$ is the period of a holomorphic differential and therefore is never zero.

We conclude that the functional determinant is singular in \mathbb{H} precisely at the branch points and AD points, however with the proper exclusion of those as done in the following Section, it is non-singular and the change of variables is well-defined. This furthermore conveniently solves the problem that there is no natural integration region in (a, \bar{a}) space [12].

5.2 Monodromy transformations of the integrand

We continue by explicitly verifying that the u -plane integral is single-valued around the singular points of the moduli space. We find that this puts a constraint on the magnetic winding numbers n_j , in addition to the constraints on the background fluxes \mathbf{k}_j discussed in Section 3.3.

Monodromy around infinity

Let us determine how the u -plane integrand transforms under the monodromy around infinity. As a function of the effective coupling τ , the measure factor (5.3) is proportional to $\frac{da}{d\tau} \left(\frac{du}{da}\right)^{\frac{\chi}{2}} \Delta^{\frac{\sigma}{8}}$ times the product over the couplings C_{ij} . We take the monodromy at infinity to be oriented as $u \rightarrow e^{2\pi i} u$ and $a \rightarrow e^{\pi i} a$, as in Section 2.5. Then this path also encircles all singularities u_j , which are the roots of the physical discriminant, $\Delta = \prod_{j=1}^{N_f+2} (u - u_j)$. We thus have that $\Delta \rightarrow e^{2\pi i(N_f+2)} \Delta$, and hence

$$\Delta^{\frac{\sigma}{8}} \rightarrow e^{\pi i(N_f+2)\sigma/4} \Delta^{\frac{\sigma}{8}} \quad (5.5)$$

Next, since $u \rightarrow e^{2\pi i}$ and $a \rightarrow e^{\pi i} a$ we find $\frac{du}{da} \rightarrow e^{\pi i} \frac{du}{da}$, and therefore

$$\left(\frac{du}{da}\right)^{\frac{\chi}{2}} \rightarrow e^{\pi i \chi/2} \left(\frac{du}{da}\right)^{\frac{\chi}{2}}. \quad (5.6)$$

For $\frac{da}{d\tau}$ we have that $a \rightarrow e^{\pi i} a$, while $d\tau \rightarrow d\tau$, and thus

$$\frac{da}{d\tau} \rightarrow -\frac{da}{d\tau}. \quad (5.7)$$

From (2.38) we recall that $w_{ij} \rightarrow w_{ij} + \delta_{ij}$, such that with the definition (4.10) we find $C_{ij} \rightarrow e^{-\pi i \delta_{ij}} C_{ij}$. The couplings C_{ij} transform in the measure factor as

$$\prod_{i,j=1}^{N_f} C_{ij}^{B(\mathbf{k}_i, \mathbf{k}_j)} \rightarrow e^{-\pi i \sum_j \mathbf{k}_j^2} \prod_{i,j=1}^{N_f} C_{ij}^{B(\mathbf{k}_i, \mathbf{k}_j)}. \quad (5.8)$$

Combining (5.5), (5.6), (5.7), (5.8), and using $\chi = 4 - \sigma$, we obtain

$$\nu \rightarrow -e^{\pi i N_f \sigma/4} e^{-\pi i \sum_j \mathbf{k}_j^2} \nu. \quad (5.9)$$

This phase for $\mathbf{k}_j = 0$ can be checked directly by taking q -expansions from the SW curves, for generic masses.

From (2.38) we recall that under the monodromy around infinity $v_j \rightarrow -v_j - n_j$, and thus

$$\mathbf{z} \rightarrow -\mathbf{z} - \sum_{j=1}^{N_f} n_j \mathbf{k}_j. \quad (5.10)$$

For the sum over fluxes we can now deduce using (A.12) that

$$\begin{aligned} & \Psi_{\boldsymbol{\mu}}^J \left(\tau + N_f - 4, -\mathbf{z} - \sum_{j=1}^{N_f} n_j \mathbf{k}_j \right) \\ &= e^{\pi i (N_f - 4)(\boldsymbol{\mu}^2 - \boldsymbol{\mu} \cdot \mathbf{K})} \Psi_{\boldsymbol{\mu}}^J \left(\tau, -\mathbf{z} - \sum_{j=1}^{N_f} n_j \mathbf{k}_j + (N_f - 4) \left(\boldsymbol{\mu} - \frac{\mathbf{K}}{2} \right) \right), \end{aligned} \quad (5.11)$$

where we suppressed the dependence on the anti-holomorphic parts. Recall from (3.9) that

$$c_1(\mathcal{L}_j) \equiv K - 2\boldsymbol{\mu} \pmod{2L}. \quad (5.12)$$

and as such we can express $\mathbf{k}_j = c_1(\mathcal{L}_j)/2$ as

$$\mathbf{k}_j = \frac{K}{2} - \boldsymbol{\mu} + \boldsymbol{\ell}_j, \quad \boldsymbol{\ell}_j \in L. \quad (5.13)$$

We have then

$$\mathbf{k}_j^2 = \frac{\sigma}{4} - K \cdot \boldsymbol{\mu} + \boldsymbol{\mu}^2 - 2\boldsymbol{\mu} \cdot \boldsymbol{\ell}_j \pmod{2\mathbb{Z}}, \quad (5.14)$$

where we used that K is a characteristic vector of L , and $K^2 = \sigma \pmod{8}$. Using (A.12) and substitution of (5.13) in this expression, (5.11) equals

$$e^{\pi i(N_f-4)(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K)} \Psi_{\boldsymbol{\mu}}^J\left(\tau, -\mathbf{z} - \sum_{j=1}^{N_f} n_j \boldsymbol{\ell}_j + (N_f - 4 + \sum_j n_j)(\boldsymbol{\mu} - \frac{K}{2})\right). \quad (5.15)$$

Our aim is to write this as a phase times $\Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z})$. The constraints on the winding numbers should be independent of $\boldsymbol{\mu}$ and \mathbf{k}_j , since the prepotential is. From (A.15), we therefore get the first constraint

$$\sum_j n_j = N_f \pmod{2}. \quad (5.16)$$

Using identity (A.15), $2\boldsymbol{\mu}^2 - K \cdot \boldsymbol{\mu} \in \mathbb{Z}$ and $4(\boldsymbol{\mu} - \frac{K}{2}) \in 2L$, this simplifies to

$$\begin{aligned} & e^{\pi i(N_f-4)(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K)+2\pi i\boldsymbol{\mu}\cdot\sum_j n_j \boldsymbol{\ell}_j-2\pi i(N_f+\sum_j n_j)(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K/2)} \Psi_{\boldsymbol{\mu}}^J(\tau, -\mathbf{z}) \\ &= -e^{\pi i N_f(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K)+2\pi i\boldsymbol{\mu}\cdot\sum_j n_j \boldsymbol{\ell}_j-2\pi i(N_f+\sum_j n_j)(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K/2)} \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}) \\ &= -e^{-\pi i N_f \boldsymbol{\mu}^2+2\pi i\boldsymbol{\mu}\cdot\sum_j n_j \boldsymbol{\ell}_j-2\pi i\sum_j n_j(\boldsymbol{\mu}^2-\boldsymbol{\mu}\cdot K/2)} \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}). \end{aligned} \quad (5.17)$$

Finally using (5.14), we can express the phase in terms of \mathbf{k}_j ,

$$\mathbf{M}_{\infty} : \quad \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}) \rightarrow -e^{-\pi i N_f \boldsymbol{\mu}^2 - \pi i \sum_j n_j (\mathbf{k}_j^2 + \boldsymbol{\mu}^2 - \sigma/4)} \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}). \quad (5.18)$$

By multiplying (5.9) with (5.18), we find

$$\nu(\tau; \{\mathbf{k}_j\}) \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}) \rightarrow e^{-\pi i \sum_j (n_j+1) \mathbf{k}_j^2 + \frac{\pi i}{4} \sum_j (\sigma - 4\boldsymbol{\mu}^2)(n_j+1)} \nu(\tau; \{\mathbf{k}_j\}) \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}). \quad (5.19)$$

Combining (3.15) with (3.4), we have that $4(\mathbf{k}_j + \boldsymbol{\mu})^2 \equiv \sigma \pmod{8}$ for every $j = 1, \dots, N_f$. We insert this into the second exponential of (5.19), such that

$$\mathbf{M}_{\infty} : \quad \nu(\tau; \{\mathbf{k}_j\}) \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}) \rightarrow e^{2\pi i \boldsymbol{\mu} \cdot \sum_j (n_j+1) \mathbf{k}_j} \nu(\tau; \{\mathbf{k}_j\}) \Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z}), \quad (5.20)$$

and the u -plane integrand is invariant under T^{N_f-4} if and only if $\boldsymbol{\mu} \sum_j (n_j+1) \mathbf{k}_j \in \mathbb{Z}$. Using (5.13) and the fact that K is a characteristic vector of L , we find

$$n_j = 1 \pmod{2} \quad (5.21)$$

for all $j = 1, \dots, N_f$, which implies the above constraint (5.16).

Monodromy M_j

Let us determine how the integrand transforms under the monodromy M_j around the mass singularity $m_j/\sqrt{2}$. Since the mass singularity corresponds to a singularity u_j on the u -plane, we have that $(u - u_j) \rightarrow e^{2\pi i}(u - u_j)$. This implies that $\Delta = (u - u_j) \prod_{i \neq j}^{2+N_f} (u - u_i) \rightarrow e^{2\pi i} \Delta$, such that $\Delta^{\frac{\sigma}{8}} \rightarrow e^{\pi i \sigma/4} \Delta^{\frac{\sigma}{8}}$. The transformation

of $\frac{da}{du}$ can be determined from (2.23): While $u \rightarrow u_j$, both g_2 and g_3 remain finite and nonzero (otherwise u_j would be an AD point). This implies that $\frac{g_3}{g_2}$ contains no factors of $(u - u_j)$, and thus $\frac{du}{da} \rightarrow \frac{du_j}{da}$. Similarly, we have that $\frac{da}{d\tau} \rightarrow \frac{da}{d\tau}$. From (2.40) we finally have that $w_{ik} \rightarrow w_{ik} + \delta_{ij}\delta_{ik}$. We combine

$$\mathbf{M}_j : \quad \nu \rightarrow e^{\pi i \sigma/4} e^{-\pi i \mathbf{k}_j^2} \nu. \quad (5.22)$$

For the monodromy around the mass singularity $m_j/\sqrt{2}$, we find for Ψ_μ^J with (A.12)

$$\begin{aligned} \Psi_\mu^J(\tau + 1, \mathbf{z} - \mathbf{k}_j) &= e^{\pi i(\mu^2 - \mu \cdot K)} \Psi_\mu^J(\tau, \mathbf{z} - \boldsymbol{\ell}_j) \\ &= e^{\pi i(\mu^2 - \mu \cdot K) + 2\pi i B(\mu, \boldsymbol{\ell}_j)} \Psi_\mu^J(\tau, \mathbf{z}) \\ &= e^{-\pi i \sigma/4 + \pi i \mathbf{k}_j^2} \Psi_\mu^J(\tau, \mathbf{z}). \end{aligned} \quad (5.23)$$

The phases thus cancel precisely,

$$\mathbf{M}_j : \quad \nu(\tau + 1) \Psi_\mu^J(\tau + 1, \mathbf{z} - \mathbf{k}_j) = \nu(\tau) \Psi_\mu^J(\tau, \mathbf{z}), \quad (5.24)$$

without any constraints.

Monodromy \mathbf{M}_m

For the monopole singularity in $N_f = 1$ we find that

$$\begin{aligned} \Psi_\mu^J\left(\frac{\tau}{-\tau+1}, \frac{v\mathbf{k}_1 + (n+1)/2\tau\mathbf{k}_1}{-\tau+1}\right) &= e^{\pi i(n+1)\frac{v}{-\tau+1}\mathbf{k}_1^2 + \pi i\frac{(n+1)^2}{4}\frac{\tau}{-\tau+1}\mathbf{k}_1^2} (-1)^{(n+1)\mathbf{k}_1 \cdot K/2} \\ &\times \Psi_{\mu+(n+1)\mathbf{k}_1/2}^J\left(\frac{\tau}{-\tau+1}, \frac{v\mathbf{k}_1}{-\tau+1}\right), \end{aligned} \quad (5.25)$$

where we have used (A.16). Then using (A.13), we arrive at

$$\begin{aligned} \Psi_\mu^J\left(\frac{\tau}{-\tau+1}, \frac{v\mathbf{k}_1 + (n+1)/2\tau\mathbf{k}_1}{-\tau+1}\right) &= (-\tau + 1)^{b_2/2} (-\bar{\tau} + 1)^2 e^{\pi i(n+1)\mathbf{k}_1 \cdot K/2 - (n+1)^2\mathbf{k}_1^2/4} e^{-\pi i \sigma/4} \\ &\times \exp\left[\pi i \frac{(v + (n+1)/2)^2}{-\tau + 1} \mathbf{k}_1^2\right] \Psi_{\mu+(n+1)\mathbf{k}_1/2}^J(\tau, \mathbf{z}). \end{aligned} \quad (5.26)$$

Since Ψ_μ^J is required to transform to itself up to an overall factor, we must demand that $(n+1)\mathbf{k}_1/2 \in L$. Therefore for $\mathbf{k}_1 \in L/2$, we find the requirement that $n = -1 \in \mathbb{Z}_4$. This simplifies the transformations considerably, and we find

$$\Psi_\mu^J(\tau, \mathbf{z}) \rightarrow (-\tau + 1)^{b_2/2} (-\bar{\tau} + 1)^2 e^{-\pi i \sigma/4} e^{\pi i \mathbf{k}_1^2 \frac{v^2}{-\tau+1}} \Psi_\mu^J(\tau, \mathbf{z}). \quad (5.27)$$

The \mathbf{k}_j -independent part of the measure factor transforms precisely as under \mathbf{M}_j (see (5.22)), as the same argument holds. However, due to the transformation $\tau \rightarrow \frac{\tau}{-\tau+1}$, the measure also picks up its modular weight $\frac{\sigma}{2} + 1$. From (2.43) we furthermore find the transformation of C_{11} , such that

$$\nu(\tau, \mathbf{k}_1) \rightarrow e^{\pi i \sigma/4} e^{-\pi i \mathbf{k}_1^2 \frac{v^2}{-\tau+1}} (-\tau + 1)^{\frac{\sigma}{2} + 1} \nu(\tau, \mathbf{k}_1), \quad (5.28)$$

where we have already used $n = -1 \in \mathbb{Z}_4$. If we multiply (5.27) and (5.28) with $d\tau \wedge d\bar{\tau}$ (which has modular weight $(-2, -2)$), then

$$\mathbf{M}_m : \quad d\tau \wedge d\bar{\tau} \nu(\tau, \mathbf{k}_1) \Psi_\mu^J(\tau, \mathbf{z}) \rightarrow d\tau \wedge d\bar{\tau} \nu(\tau, \mathbf{k}_1) \Psi_\mu^J(\tau, \mathbf{z}), \quad (5.29)$$

where we have used $\sigma + b_2 = 2$. Thus, the u -plane integrand is also invariant under \mathbf{M}_m .

For $N_f > 1$ we find the same condition, namely that $n_j = -1 \pmod 4$ for all j .

Monodromy \mathbf{M}_d

Given the relation (2.41), it is not necessary to explicitly check single-valuedness of the integrand under this monodromy, as it is a product of the above monodromies.

To conclude this section, let us stress the constraints for the winding number n_j , such that the u -plane integral is invariant under all monodromies in $N_f \leq 3$. To this end, we need to satisfy the constraints $n_j = 1 \pmod 2$ (5.21) from \mathbf{M}_∞ , and $n_j = -1 \pmod 4$ for \mathbf{M}_m . Since the latter is the stronger constraint, we require

$$n_j = -1 \pmod 4, \quad (5.30)$$

for all $j = 1, \dots, N_f$.

6 Integration over fundamental domains

As discussed in Sections 2 and 5, u -plane integrals for massive $\mathcal{N} = 2$ theories with fundamental hypermultiplets include new aspects. This section discusses how to evaluate such integrals (5.1). More concretely, we aim to define and evaluate integrals of the form

$$\mathcal{I}_f = \int_{\mathcal{F}(\mathbf{m})} d\tau \wedge d\bar{\tau} y^{-s} f(\tau, \bar{\tau}), \quad (6.1)$$

with $s \leq 1$. The domain $\mathcal{F}(\mathbf{m})$ is the fundamental domain for the effective coupling constant as discussed in Section 2.4, and f a non-holomorphic function of weight $(2-s, 2-s)$ arising from the topologically twisted Yang-Mills theory. For $\mathcal{F}(\mathbf{m})$ a fundamental domain of a congruence subgroup, such integrals (6.1) have been studied in the context of theta lifts of weakly holomorphic modular forms and harmonic Maass forms [105–107] as well as one-loop amplitudes in string theory [108–110].

We assume that the integrand $y^{-s} f(\tau, \bar{\tau})$ can be expressed as

$$\partial_{\bar{\tau}} \widehat{h}(\tau, \bar{\tau}) = y^{-s} f(\tau, \bar{\tau}), \quad (6.2)$$

for a suitable function $\widehat{h}(\tau, \bar{\tau})$ using mock modular forms. This was indeed the case in [26, 47, 48], and will be demonstrated for massive $\mathcal{N} = 2$ theories with fundamental hypermultiplets. The integral \mathcal{I}_f then reads

$$\mathcal{I}_f = - \int_{\partial\mathcal{F}(\mathbf{m})} d\tau \widehat{h}(\tau, \bar{\tau}), \quad (6.3)$$

with $\partial\mathcal{F}(\mathbf{m})$ the boundary of $\mathcal{F}(\mathbf{m})$. We will carry this out evaluation in Part II [88].

There are a number of aspects to be addressed in order to evaluate integrals over $\mathcal{F}(\mathbf{m})$:

1. Identifications of boundary components of $\mathcal{F}(\mathbf{m})$ due to monodromies on the u -plane.
2. Contributions from the cusps, that is $\tau \rightarrow i\infty$ or $\tau \rightarrow \gamma(i\infty) \in \mathbb{Q}$ for an element $\gamma \in \text{PSL}(2, \mathbb{Z})$.
3. Contributions from a singular point in the interior of $\mathcal{F}(\mathbf{m})$.
4. Contributions from an elliptic point $p \in \mathbb{H}$ of $\text{PSL}(2, \mathbb{Z})$.
5. Branch points and branch cuts.

We will discuss these aspects 1.–5. in the following.

1. *Identifications*

The modular transformation induced by monodromies identify components of the boundary of the fundamental domain $\partial\mathcal{F}(\mathbf{m})$ pairwise. Their contributions to the integral (6.3) vanish, which is, for example, familiar from deriving valence formulas for modular forms [111, Fig. 2]. See Fig. 4 for an example.

2. *Cusps*

Contributions near the cusps require a regularisation [12, 48]. Such regularisations have been developed in the context of string amplitudes [108–110] and analytic number theory [111–113].

Let us first consider the cusp $\tau \rightarrow i\infty$. To regularise the divergence, one introduces a cut-off $\text{Im } \tau = Y \gg 1$, and takes the limit $Y \rightarrow \infty$ after evaluation. We require that f near $i\infty$ has a Fourier expansion of the form⁵

$$f(\tau, \bar{\tau}) = \sum_{m \gg -\infty, n \geq 0} c(m, n) q^m \bar{q}^n. \quad (6.4)$$

Then the function \hat{h} has the form,

$$\hat{h}(\tau, \bar{\tau}) = h(\tau) + 2^s \int_{-\bar{\tau}}^{i\infty} \frac{f(\tau, -v)}{(-i(v + \tau))^s} dv, \quad (6.5)$$

where $h(\tau)$ is a weakly holomorphic q -series, with expansion

$$h(\tau) = \sum_{m \gg -\infty} d(m) q^m. \quad (6.6)$$

⁵Also if f does not satisfy this requirement, the integral can be regularised as explained in [48, 113]. We do not need this regularisation for the correlators in this paper.

The cusp $\tau \rightarrow i\infty$ then contributes

$$[\mathcal{I}_f]_\infty = w_\infty d(0), \quad (6.7)$$

with $d(0)$ the constant term of $h(\tau)$ (6.6), and w_∞ the width of the cusp $\mathcal{F}(\mathbf{m})$ at $i\infty$. For $N_f \leq 3$, w_∞ is $4 - N_f$ [52].

The other cusps can be treated in a similar fashion using modular transformations. We label the n_c cusps in $\mathcal{F}(\mathbf{m})$ by $j = 1, \dots, n_c$. If the cusp is on the horizontal axis at $-\frac{d_j}{c_j} \in \mathbb{Q}$ with relative prime $(c_j, d_j) \in \mathbb{Z}^2$, we can map the cusp to $i\infty$ by a modular transformation

$$\gamma_j = \begin{pmatrix} a_j & b_j \\ c_j & d_j \end{pmatrix}. \quad (6.8)$$

We let $\tau_j = \gamma_j \tau$. Then the holomorphic part $h_j(\tau_j)$ of $(c_j \tau + d_j)^{-2} \widehat{h}(\gamma \tau_j, \gamma \bar{\tau}_j)$ can be expanded for τ near $-\frac{d_j}{c_j}$ as

$$h_j(\tau_j) = \sum d_j(n) q_j^n, \quad q_j = e^{2\pi i \tau_j}. \quad (6.9)$$

As a result, the cusp j contributes

$$[\mathcal{I}_f]_j = w_j d_j(0). \quad (6.10)$$

3. Singular points in the interior of $\mathcal{F}(\mathbf{m})$

The integrand can be singular at a point τ_s in the interior of $\mathcal{F}(\mathbf{m})$. Such singularities appear typically for deformations of superconformal theories, such as the $\mathcal{N} = 2^*$ theory and the $N_f = 4$ theory, where the UV coupling τ_{UV} gives rise to such a singularity [26, 54]. See Fig. 8 for an example. We require that the expansion of f near such a singularity reads,

$$f(\tau, \bar{\tau}) = \sum_{m \gg -\infty, n \geq 0} c_s(m, n) (\tau - \tau_s)^m (\bar{\tau} - \bar{\tau}_s)^n. \quad (6.11)$$

Then, the anti-derivative $\widehat{h}(\tau, \bar{\tau})$ has similar expansion,

$$\widehat{h}(\tau, \bar{\tau}) = \sum_{m \gg -\infty, n \geq 0} d_s(m, n) (\tau - \tau_s)^m (\bar{\tau} - \bar{\tau}_s)^n. \quad (6.12)$$

The contour integral for a small contour around τ_s ,

$$C_\varepsilon(\tau_s) = \{ \tau = \tau_s + \varepsilon e^{i\varphi}, \varphi \in [0, 2\pi) \}, \quad (6.13)$$

is bounded for such a function. Moreover, in the limit $\varepsilon \rightarrow 0$, the contour integral is finite. We define the “residue” of a non-holomorphic function $g(\tau, \bar{\tau})$

$$\text{nRes}_{\tau=\tau_s} [g(\tau, \bar{\tau})] = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0} \oint_{C_\varepsilon(\tau_s)} g(\tau, \bar{\tau}) d\tau. \quad (6.14)$$

For the expansion (6.11) this evaluates to

$$[\mathcal{I}_f]_s = 2\pi i \operatorname{nRes}_{\tau=\tau_s} \left[\widehat{h}(\tau, \bar{\tau}) \right] = d_s(-1, 0), \quad (6.15)$$

with $d_s(-1, 0)$ the coefficient in the expansion (6.12).

4. Elliptic points

For $\mathcal{N} = 2$ QCD, AD points are the elliptic points of the duality group, and lie on the boundary of $\mathcal{F}(\mathbf{m})$. See Fig. 7 for an example. The elliptic points are $\alpha = e^{\pi i/3}$ and i , and their images under $\operatorname{PSL}(2, \mathbb{Z})$. Contour integrals around such points can be regularised using a cut-off ε . We assume that the anti-derivative \widehat{h} has the following expansion near an elliptic point τ_e ,

$$\widehat{h}(\tau, \bar{\tau}) = \sum_{m \gg -\infty, n \geq 0} d_e(m, n) (\tau - \tau_e)^m (\bar{\tau} - \bar{\tau}_e)^n. \quad (6.16)$$

As a result, the boundary arc around τ_{AD} in \mathbb{H} is a fraction of 2π , which needs to be properly accounted for. These neighbourhoods have an angle $\frac{2\pi}{k_e}$, with $k_e = 2$ for $\tau_e = i$, and $k_e = 6$ for $\tau_e = \alpha$ [114]. Furthermore, it is important how many images of \mathcal{F} in $\mathcal{F}(\mathbf{m})$ coincide at the elliptic point. We denote this number by n_e . For $\mathcal{N} = 2$ SQCD, we found examples with $n_e = 2$ and 4 for $\tau_e \sim \alpha$, while for $\tau_e \sim i$, $n_e = 1$ [52]. The contribution from an elliptic point is then,

$$[\mathcal{I}_f]_e = 2\pi i \frac{n_e}{k_e} \operatorname{nRes}_{\tau=\tau_e} \left[\widehat{h}(\tau, \bar{\tau}) \right] = \frac{n_e}{k_e} d_e(-1, 0), \quad (6.17)$$

5. Branch points and cuts

Branch points and cuts are a new aspect compared to previous analyses (see for instance Fig. 4 and 5). We will demonstrate that their contribution vanishes for the integrands of interest.

We assume that the integrand f satisfies

$$\widehat{h}(\tau, \bar{\tau}) = (\tau - \tau_{\text{bp}})^n g(\tau, \bar{\tau}), \quad (6.18)$$

with $n \in \mathbb{Z}/2$ and $n \geq -1/2$, $g(\tau, \bar{\tau})$ being a real analytic function near τ_{bp} . This assumption is satisfied for the twisted Yang-Mills theories [88]. To treat this type of singularity, we remove a δ neighbourhood and analyse the $\delta \rightarrow 0$ limit. Let C_δ be the contour

$$C_\delta = \{\tau_{\text{bp}} + \delta e^{i\theta} \mid \theta \in (0, 2\pi)\} \quad (6.19)$$

around τ_{bp} with radius $\delta > 0$. Therefore, on the contour $|y^{-s} f|$ is bounded by

$$|\widehat{h}| \leq \delta^n K \quad (6.20)$$

for some $K > 0$. The integral around the branch point therefore vanishes in the limit,

$$\begin{aligned} \mathcal{I}_f^{\text{bp}} &= \lim_{\delta \rightarrow 0} \int_{C_\delta} \widehat{h} |d\tau| \leq \lim_{\delta \rightarrow 0} \int_0^{2\pi} \delta^n K \delta d\theta \\ &= \lim_{\delta \rightarrow 0} 2\pi K \delta^{n+1} = 0. \end{aligned} \tag{6.21}$$

The branch points necessarily give rise to branch cuts. For the purpose of integration, we remove a neighbourhood with distance r from the cut, and take the limit $r \rightarrow 0$ after determining the integral. Since the value of the integrand is finite near the branch cut, the contribution to the integral vanishes.

Summary

Combining all the contributions discussed above, we find

$$\mathcal{I}_f = \sum_{j=1}^n w_j d_j(0) + \sum_s d_s(-1, 0) + \sum_e \frac{n_e}{k_e} d_e(-1, 0). \tag{6.22}$$

This formula generalises [12] for the pure $N_f = 0$ theory on a smooth four-manifold X that admits a metric of positive scalar curvature, [49, Equation (5.10)] for the pure theory on generic X , [26, Equation (4.88)] for the $\mathcal{N} = 2^*$ theory on X , and [43] for the massless $N_f = 2$ and $N_f = 3$ theories on $X = \mathbb{C}\mathbb{P}^2$.

Acknowledgments

We are happy to thank Cyril Closset, Horia Magureanu, Greg Moore and Samson Shatashvili for correspondence and discussions. JM thanks Greg Moore for collaboration on related projects. JA is supported by the Government of Ireland Postgraduate Scholarship Programme GOIPG/2020/910 of the Irish Research Council. EF is supported by the TCD Provost’s PhD Project Award. JM is supported by the Laureate Award 15175 “Modularity in Quantum Field Theory and Gravity” of the Irish Research Council.

A Modular forms

In this Appendix, we collect some properties of modular forms for subgroups of $\text{PSL}(2, \mathbb{Z})$. For further reading, see [111, 114–118].

A.1 Theta functions and Eisenstein series

We make use of modular forms for the congruence subgroups $\Gamma_0(n)$ and $\Gamma^0(n)$ of $\mathrm{PSL}(2, \mathbb{Z})$. These subgroups are defined as

$$\begin{aligned}\Gamma_0(n) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}) \mid c \equiv 0 \pmod{n} \right\}, \\ \Gamma^0(n) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}) \mid b \equiv 0 \pmod{n} \right\},\end{aligned}\tag{A.1}$$

and are related by conjugation with the matrix $\mathrm{diag}(n, 1)$. We furthermore define the *principal congruence subgroup* $\Gamma(n)$ as the subgroup of $\mathrm{SL}(2, \mathbb{Z}) \ni A$ with $A \equiv \mathbb{1} \pmod{n}$. A subgroup Γ of $\mathrm{SL}(2, \mathbb{Z})$ is called a congruence subgroup if it contains $\Gamma(n)$ for some $n \in \mathbb{N}$.

The above introduced congruence subgroups host a number of interesting modular forms. The Jacobi theta functions $\vartheta_j : \mathbb{H} \rightarrow \mathbb{C}$, $j = 2, 3, 4$, are defined as

$$\vartheta_2(\tau) = \sum_{r \in \mathbb{Z} + \frac{1}{2}} q^{r^2/2}, \quad \vartheta_3(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2/2}, \quad \vartheta_4(\tau) = \sum_{n \in \mathbb{Z}} (-1)^n q^{n^2/2},\tag{A.2}$$

with $q = e^{2\pi i \tau}$. These functions transform under $T, S \in \mathrm{SL}(2, \mathbb{Z})$ as

$$\begin{aligned}S : \quad & \vartheta_2(-1/\tau) = \sqrt{-i\tau} \vartheta_4(\tau), \\ & \vartheta_3(-1/\tau) = \sqrt{-i\tau} \vartheta_3(\tau), \\ & \vartheta_4(-1/\tau) = \sqrt{-i\tau} \vartheta_2(\tau), \\ T : \quad & \vartheta_2(\tau + 1) = e^{\frac{\pi i}{4}} \vartheta_2(\tau), \\ & \vartheta_3(\tau + 1) = \vartheta_4(\tau), \\ & \vartheta_4(\tau + 1) = \vartheta_3(\tau).\end{aligned}\tag{A.3}$$

They furthermore satisfy the Jacobi abstruse identity

$$\vartheta_2^4 + \vartheta_4^4 = \vartheta_3^4.\tag{A.4}$$

The Dedekind eta function $\eta : \mathbb{H} \rightarrow \mathbb{C}$ is defined as the infinite product

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n), \quad q = e^{2\pi i \tau}.\tag{A.5}$$

It transforms under the generators of $\mathrm{SL}(2, \mathbb{Z})$ as

$$\begin{aligned}S : \quad & \eta(-1/\tau) = \sqrt{-i\tau} \eta(\tau), \\ T : \quad & \eta(\tau + 1) = e^{\frac{\pi i}{12}} \eta(\tau),\end{aligned}\tag{A.6}$$

and relates to the Jacobi theta series as $\eta^3 = \frac{1}{2} \vartheta_2 \vartheta_3 \vartheta_4$.

Eisenstein series

We let $\tau \in \mathbb{H}$ and define $q = e^{2\pi i\tau}$. Then the Eisenstein series $E_k : \mathbb{H} \rightarrow \mathbb{C}$ for even $k \geq 2$ are defined as the q -series

$$E_k(\tau) = 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n, \quad (\text{A.7})$$

where B_k are the Bernoulli numbers and $\sigma_k(n) = \sum_{d|n} d^k$ is the divisor sum. For $k \geq 4$ even, E_k is a modular form of weight k for $\text{SL}(2, \mathbb{Z})$. Any modular form for $\text{SL}(2, \mathbb{Z})$ can be related to the Jacobi theta functions (A.2) by

$$E_4 = \frac{1}{2}(\vartheta_2^8 + \vartheta_3^8 + \vartheta_4^8), \quad E_6 = \frac{1}{2}(\vartheta_2^4 + \vartheta_3^4)(\vartheta_3^4 + \vartheta_4^4)(\vartheta_4^4 - \vartheta_2^4). \quad (\text{A.8})$$

With our normalisation (A.7), the j -invariant can be written as

$$j = 1728 \frac{E_4^3}{E_4^3 - E_6^2} = 256 \frac{(\vartheta_3^8 - \vartheta_3^4 \vartheta_4^4 + \vartheta_4^8)^3}{\vartheta_2^8 \vartheta_3^8 \vartheta_4^8}. \quad (\text{A.9})$$

A.2 Siegel-Narain theta function

Let L be an n -dimensional uni-modular lattice with signature $(1, n-1)$. For the application to the u -plane integral, $n = b_2(X)$. Let K be a characteristic vector of L . Its defining property is $\mathbf{l}^2 = \mathbf{l} \cdot K \pmod{2}$ for every $\mathbf{l} \in L$. Furthermore, we have that $\boldsymbol{\mu} \in L/2$.

We consider the Siegel-Narain theta function $\Psi_{\boldsymbol{\mu}}^J : \mathbb{H} \times \mathbb{C} \rightarrow \mathbb{C}$ defined in the main text in (4.11). We repeat it here for convenience,

$$\begin{aligned} \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}) &= e^{-2\pi y \mathbf{b}_+^2} \sum_{\mathbf{k} \in L + \boldsymbol{\mu}} \partial_{\bar{\tau}} (4\pi i \sqrt{y} B(\mathbf{k} + \mathbf{b}, J)) \\ &\times (-1)^{B(\mathbf{k}, K)} q^{-\mathbf{k}_-^2/2} \bar{q}^{\mathbf{k}_+^2/2} e^{-2\pi i B(\mathbf{z}, \mathbf{k}_-)} e^{-2\pi i B(\bar{\mathbf{z}}, \mathbf{k}_+)}, \end{aligned} \quad (\text{A.10})$$

where J is a normalized positive vector in $L \otimes \mathbb{R}$, $\mathbf{k}_+ = B(\mathbf{k}, J) J$, $\mathbf{k}_- = \mathbf{k} - \mathbf{k}_+$ and $\mathbf{b} = \text{Im}(\mathbf{z})/y$. The transformations under the generators S and T of $\text{PSL}(2, \mathbb{Z})$ are most easily determined if we shift $\boldsymbol{\mu} \rightarrow \boldsymbol{\mu} + K/2$. One finds [47, 49]

$$\begin{aligned} S : \quad & \Psi_{\boldsymbol{\mu}+K/2}^J(-1/\tau, -1/\bar{\tau}, \mathbf{z}/\tau, \bar{\mathbf{z}}/\bar{\tau}) = -i(-i\tau)^{n/2} (i\bar{\tau})^2 \\ & \times e^{-\pi i \mathbf{z}^2/\tau + \pi i K^2/2} (-1)^{B(\boldsymbol{\mu}, K)} \Psi_{K/2}^J(\tau, \bar{\tau}, \mathbf{z} - \boldsymbol{\mu}, \bar{\mathbf{z}} - \boldsymbol{\mu}), \\ T : \quad & \Psi_{\boldsymbol{\mu}+K/2}^J(\tau + 1, \bar{\tau} + 1, \mathbf{z}, \bar{\mathbf{z}}) = \\ & e^{\pi i(\boldsymbol{\mu}^2 - K^2/4)} \Psi_{\boldsymbol{\mu}+K/2}^J(\tau, \bar{\tau}, \mathbf{z} + \boldsymbol{\mu}, \bar{\mathbf{z}} + \boldsymbol{\mu}). \end{aligned} \quad (\text{A.11})$$

Using these transformations, one finds for the periodicity in τ ,

$$\Psi_{\boldsymbol{\mu}}^J(\tau + 1, \bar{\tau} + 1, \mathbf{z}, \bar{\mathbf{z}}) = e^{\pi i(\boldsymbol{\mu}^2 - B(\boldsymbol{\mu}, K))} \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z} + \boldsymbol{\mu} - K/2, \bar{\mathbf{z}} + \boldsymbol{\mu} - K/2) \quad (\text{A.12})$$

and for $S^{-1}T^{-k}S = \begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix}$,

$$\Psi_{\boldsymbol{\mu}}^J\left(\frac{\tau}{k\tau+1}, \frac{\bar{\tau}}{k\bar{\tau}+1}, \frac{\mathbf{z}}{k\tau+1}, \frac{\bar{\mathbf{z}}}{k\bar{\tau}+1}\right) = (k\tau+1)^{\frac{n}{2}}(k\bar{\tau}+1)^2 e^{-\frac{\pi i k \mathbf{z}^2}{k\tau+1}} e^{\frac{\pi i}{4} k K^2} \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}). \quad (\text{A.13})$$

We furthermore list the following transformations for \mathbf{z} :

- For the reflection $\mathbf{z} \rightarrow -\mathbf{z}$,

$$\Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, -\mathbf{z}, -\bar{\mathbf{z}}) = -e^{2\pi i B(\boldsymbol{\mu}, K)} \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}). \quad (\text{A.14})$$

- For shifting $\mathbf{z} \rightarrow \mathbf{z} + \boldsymbol{\nu}$ with $\boldsymbol{\nu} \in L$,

$$\Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z} + \boldsymbol{\nu}, \bar{\mathbf{z}} + \boldsymbol{\nu}) = e^{-2\pi i B(\boldsymbol{\nu}, \boldsymbol{\mu})} \Psi_{\boldsymbol{\mu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}). \quad (\text{A.15})$$

- For shifting $\mathbf{z} \rightarrow \mathbf{z} + \boldsymbol{\nu}\tau$ with $\boldsymbol{\nu} \in L \otimes \mathbb{R}$,

$$\Psi_{\boldsymbol{\mu}}^J(\tau, \mathbf{z} + \boldsymbol{\nu}\tau) = e^{2\pi i B(\mathbf{z}, \boldsymbol{\nu})} q^{\boldsymbol{\nu}^2/2} (-1)^{-B(\boldsymbol{\nu}, K)} \Psi_{\boldsymbol{\mu}+\boldsymbol{\nu}}^J(\tau, \bar{\tau}, \mathbf{z}, \bar{\mathbf{z}}). \quad (\text{A.16})$$

We can restrict to $\boldsymbol{\nu} \in L/2$, if the characteristic $\boldsymbol{\mu} + \boldsymbol{\nu}$ is required to be in $L/2$.

B Class \mathcal{S} representation

A different parametrisation of the SW curves (2.17) is the class \mathcal{S} representation. This representation gives the SW differential in a canonical form

$$\lambda^2 = p_{N_f}(z, u, \Lambda_{N_f}, \mathbf{m}) dz^2, \quad (\text{B.1})$$

where the Laurent polynomials p_{N_f} read [75, Section 10]

$$\begin{aligned} p_0 &= \frac{\Lambda_0^2}{z^3} + \frac{2u}{z^2} + \frac{\Lambda_0^2}{z}, \\ p_1 &= \frac{\Lambda_1^2}{z^3} + \frac{3u}{z^2} + \frac{2\Lambda_1 m}{z} + \Lambda_1^2, \\ p_2 &= \frac{\Lambda_2^2}{z^4} + \frac{2\Lambda_2 m_1}{z^3} + \frac{4u}{z^2} + \frac{2\Lambda_2 m_2}{z} + \Lambda_2^2. \end{aligned} \quad (\text{B.2})$$

The corresponding elliptic curves can be found as $x^2 = z^4 p_{N_f}$, which is quartic for $N_f = 0, 1$ and cubic for $N_f = 2$. By comparing invariants of the SW curves (2.17) with those of the class \mathcal{S} curve (B.1), one finds the dictionary

$$\begin{aligned} N_f = 0 : \quad & u_{\mathcal{S}} = u_{\text{SW}}, \\ N_f = 1 : \quad & u_{\mathcal{S}} = \frac{4}{3} u_{\text{SW}}, \quad m_{\mathcal{S}} = 2m_{\text{SW}}, \\ N_f = 2 : \quad & u_{\mathcal{S}} = 4u_{\text{SW}}, \quad m_{\mathcal{S},i} = 4m_{\text{SW},i}. \end{aligned} \quad (\text{B.3})$$

These relations merely amount to a rescaling of the parametrisation of the Coulomb branch, and in particular leave its geometry invariant. For this reason, we proceed above with using the SW curves (2.17).

C Winding numbers

In this Appendix, we discuss the winding numbers appearing in (2.7) in some more detail. As mentioned in Section 2.2, generally the theory admits N_f electric winding numbers and N_f magnetic winding numbers. While in the main text we set the electric winding numbers to zero, in this Appendix we keep the possibility open for them to be nonzero.

The $\mathcal{N} = 2$ supersymmetry algebra requires that the central charge Z is a linear combination of conserved charges. The $U(1)$ conserved charges S_i of the hypermultiplets must appear in Z as follows [11],

$$Z = n_m a_D + n_e a + \sum_{j=1}^{N_f} S_j \frac{m_j}{\sqrt{2}}. \quad (\text{C.1})$$

We note that Z is an inner product of periods $(a_D, a, \frac{1}{\sqrt{2}}\mathbf{m})$ and conserved charges (n_m, n_e, \mathbf{S}) . The periods are given by contour integrals

$$a = \int_{\gamma_1} \lambda, \quad a_D = \int_{\gamma_2} \lambda, \quad (\text{C.2})$$

where γ_1, γ_2 generate $H_1(E, \mathbb{Z})$, and λ is a meromorphic 1-form, the SW differential. In massive SQCD, λ has poles with nonzero residues. If the contours γ_i are deformed across poles of λ , then a and a_D pick up contributions from the residues. By invariance of (C.1), these jumps are linear combinations of the masses,

$$\sum_{j=1}^{N_f} \Delta S_j \frac{m_j}{\sqrt{2}}, \quad (\text{C.3})$$

i.e. the global charges of the hypermultiplets are shifted as $S_j \rightarrow S_j + \Delta S_j$. From the residue theorem, the shift in the periods gives

$$2\pi i \sum_{i=1}^{N_f} n_i \text{Res}(\lambda, x_i), \quad (\text{C.4})$$

where x_i are the N_f poles of λ , and $n_i \in \mathbb{Z}$ are the winding numbers of the contour deformation. Since both a_D and a are given by contour integrals of the same differential, a deformation of either γ_1 or γ_2 or both results in such a shift.

Example. In order to see how n_i and ΔS_j are related, let us study $N_f = 2$ as an example. In $N_f = 2$, the S_j are integers for fundamental particles and half-integers for monopoles [11]. The two poles of λ are $x^\pm = \pm \frac{1}{8}\Lambda_2^2$, with residue [11, (15.4)]

$$2\pi i \text{Res}(\lambda, x^\pm) = \pm \frac{m_1 + m_2}{2\sqrt{2}}. \quad (\text{C.5})$$

Then,

$$\sum_{j=1}^2 \Delta S_j \frac{m_j}{\sqrt{2}} = 2\pi i \sum_{i=1}^2 n_i \text{Res}(\lambda, x_i) = \sum_{j=1}^2 \frac{n_1 - n_2}{2} \frac{m_j}{\sqrt{2}}, \quad (\text{C.6})$$

such that $\Delta S_j = \frac{n_1 - n_2}{2} \in \mathbb{Z}/2$ for both $j = 1, 2$. Constraints on the n_i entail constraints on the change ΔS_j . For instance, if both n_1 and n_2 are odd, or if both are even, then ΔS_1 and ΔS_2 are integers. In the main text of this paper, we find such a constraint.

For generic $N_f \leq 3$, by construction of the SW curves the residues of λ are linear combinations of the masses

$$2\pi i \text{Res}(\lambda, x_i) = \sum_{j=1}^{N_f} l_{ij} \frac{m_j}{\sqrt{2}}, \quad (\text{C.7})$$

with $i = 1, \dots, N_f$, and $l_{ij} \in \mathbb{Z}/2$ [11, (17.1)]. Then we can compute

$$\sum_{j=1}^{N_f} \Delta S_j \frac{m_j}{\sqrt{2}} = 2\pi i \sum_{i=1}^{N_f} n_i \text{Res}(\lambda, x_i) = \sum_{i=1}^{N_f} \sum_{j=1}^{N_f} n_i l_{ij} \frac{m_j}{\sqrt{2}} = \sum_{j=1}^{N_f} \frac{m_j}{\sqrt{2}} \sum_{i=1}^{N_f} l_{ij} n_i. \quad (\text{C.8})$$

We find that the change in the abelian global charges ΔS_j is given by a half-integral linear combination of winding numbers,

$$\Delta S_j = \sum_{i=1}^{N_f} l_{ij} n_i. \quad (\text{C.9})$$

In order to relate the winding numbers n_i to those for the periods a_i and $a_{D,i}$ as found in [56], we can expand

$$\mathbf{M}_\infty^{\text{Ohta}} : \quad a_D \rightarrow -a_D + (4 - N_f)a - \frac{1}{\sqrt{2}} \left(\frac{n_{N_f}^{a_D}}{n_{N_f}^a} - \frac{4 - N_f}{2} \right) \sum_{j=1}^{N_f} n_j^a m_j, \quad (\text{C.10})$$

with n_j^a being the coefficients of $-\frac{1}{2\sqrt{2}}m_j$ in the constant term of a , and similarly $n_j^{a_D}$ those for a_D [56, (5.2)]⁶. We are aiming to compare this to (2.36),

$$\mathbf{M}_\infty : \quad a_D \rightarrow -a_D + (4 - N_f)a - \frac{1}{\sqrt{2}} \sum_{j=1}^{N_f} n_j m_j. \quad (\text{C.11})$$

Since in (C.10) we divide and multiply by n_j^a , we cannot directly relate the two expressions by setting $n_j^a = 0$. However, if we relate first

$$n_j^a = \frac{n_{N_f}^a}{n_{N_f}^{a_D}} n_j^{a_D}, \quad j = 1, \dots, N_f - 1, \quad (\text{C.12})$$

⁶In [56], the n_j^a are called n_j and $n_j^{a_D}$ are called n'_j .

then we find

$$\mathbf{M}_\infty^{\text{Ohta}} : \quad a_D \rightarrow -a_D + (4 - N_f)a - \frac{1}{\sqrt{2}} \sum_{j=1}^{N_f} \left(n_j^{a_D} - \frac{4 - N_f}{2} n_j^a \right) m_j. \quad (\text{C.13})$$

In this case, it is well-defined to set $n_j^a = 0$, such that the action of $\mathbf{M}_\infty^{\text{Ohta}}$ and \mathbf{M}_∞ coincide for $n_j = n_j^{a_D}$, as anticipated.

The $N_f - 1$ conditions (C.12) reproduce the constraints on the winding numbers in [56, (5.3)]: For $N_f = 1$, the condition is empty. For $N_f = 2$, it agrees precisely with Ohta. For $N_f = 3$ finally, the $N_f - 1 = 2$ equations (C.12) are equivalent to those found in [56]. It seems however that our result $n_j^a = 0, n_j^{a_D} = -1$ (5.30) is merely one particular of the infinitely many solutions to the geometric constraint (C.12).⁷ Thus it appears that the formulation of the theory on a compact four-manifold introduces further constraints. It would be interesting to understand whether it is possible to introduce non-vanishing electric winding numbers in the u -plane integral, and whether this leads to different correlation functions.

References

- [1] E. Witten, *Topological Quantum Field Theory*, *Commun. Math. Phys.* **117** (1988) 353.
- [2] E. Witten, *Supersymmetric Yang-Mills theory on a four manifold*, *J. Math. Phys.* **35** (1994) 5101–5135, [[hep-th/9403195](#)].
- [3] E. Witten, *Monopoles and four manifolds*, *Math. Res. Lett.* **1** (1994) 769–796, [[hep-th/9411102](#)].
- [4] J. P. Yamron, *Topological actions from twisted supersymmetric theories*, *Physics Letters B* **213** (1988) 325 – 330.
- [5] D. Anselmi and P. Fré, *Topological twist in four dimensions, r-duality and hyperinstantons*, *Nuclear Physics B* **404** (08, 1993) 288–320.
- [6] D. Anselmi and P. Fré, *Topological sigma-models in four dimensions and triholomorphic maps*, *Nuclear Physics B* **416** (03, 1994) 255–300.
- [7] D. Anselmi and P. Fré, *Gauged hyperinstantons and monopole equations*, *Physics Letters B* **347** (1995) 247 – 254.
- [8] M. Alvarez and J. Labastida, *Breaking of topological symmetry*, *Physics Letters B* **315** (1993) 251 – 257.
- [9] M. Alvarez and J. Labastida, *Topological matter in four-dimensions*, *Nucl. Phys. B* **437** (1995) 356–390, [[hep-th/9404115](#)].

⁷One other simple solution is $n_j^a = n$ and $n_j^{a_D} = \tilde{n}$, with arbitrary $n, \tilde{n} \in \mathbb{Z}$.

- [10] N. Seiberg and E. Witten, *Electric - magnetic duality, monopole condensation, and confinement in $N=2$ supersymmetric Yang-Mills theory*, *Nucl. Phys.* **B426** (1994) 19–52, [[hep-th/9407087](#)].
- [11] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in $N=2$ supersymmetric QCD*, *Nucl. Phys.* **B431** (1994) 484–550, [[hep-th/9408099](#)].
- [12] G. W. Moore and E. Witten, *Integration over the u plane in Donaldson theory*, *Adv. Theor. Math. Phys.* **1** (1997) 298–387, [[hep-th/9709193](#)].
- [13] A. Losev, N. Nekrasov and S. L. Shatashvili, *Issues in topological gauge theory*, *Nucl. Phys.* **B534** (1998) 549–611, [[hep-th/9711108](#)].
- [14] J. Labastida and M. Marino, *A topological lagrangian for monopoles on four-manifolds*, *Physics Letters B* **351** (03, 1995) 146–152.
- [15] J. M. F. Labastida and M. Marino, *NonAbelian monopoles on four manifolds*, *Nucl. Phys. B* **448** (1995) 373–398, [[hep-th/9504010](#)].
- [16] S. Hyun, J. Park and J.-S. Park, *Spin- c Topological QCD*, *Nucl. Phys.* **B453** (1995) 199–224, [[hep-th/9503201](#)].
- [17] S. Hyun, J. Park and J.-S. Park, *$N=2$ supersymmetric QCD and four manifolds: 1. The Donaldson and Seiberg-Witten invariants*, [hep-th/9508162](#).
- [18] J. Labastida and C. Lozano, *Mass perturbations in twisted $n = 4$ supersymmetric gauge theories*, *Nuclear Physics B* **518** (05, 1998) 37–58.
- [19] R. Dijkgraaf, J.-S. Park and B. J. Schroers, *$N=4$ supersymmetric Yang-Mills theory on a Kahler surface*, [hep-th/9801066](#).
- [20] H. Kanno and S.-K. Yang, *Donaldson-Witten functions of massless $N=2$ supersymmetric QCD*, *Nucl. Phys. B* **535** (1998) 512–530, [[hep-th/9806015](#)].
- [21] M. Mariño, G. Moore and G. Peradze, *Superconformal invariance and the geography of four-manifolds*, *Communications in Mathematical Physics* **205** (1999) 691–735.
- [22] M. Marino, G. W. Moore and G. Peradze, *Four manifold geography and superconformal symmetry*, [math/9812042](#).
- [23] M. Marino and G. Moore, *Integrating over the coulomb branch in $n = 2$ gauge theory*, *Nuclear Physics B - Proceedings Supplements* **68** (1998) 336 – 347.
- [24] G. W. Moore and I. Nidaiev, *The Partition Function Of Argyres-Douglas Theory On A Four-Manifold*, [1711.09257](#).
- [25] M. Dedushenko, S. Gukov and P. Putrov, *Vertex algebras and 4-manifold invariants*, in *Nigel Hitchin’s 70th Birthday Conference*, vol. 1, pp. 249–318, 5, 2017. [1705.01645](#). DOI.
- [26] J. Manschot and G. W. Moore, *Topological correlators of $SU(2)$, $\mathcal{N} = 2^*$ SYM on four-manifolds*, [2104.06492](#).
- [27] L. Göttsche, H. Nakajima and K. Yoshioka, *Donaldson = seiberg-witten from mochizuki’s formula and instanton counting*, [1001.5024](#).

- [28] P. M. N. Feehan and T. G. Leness, *PU(2) monopoles. I. Regularity, Uhlenbeck compactness, and transversality*, *J. Diff. Geom.* **49** (1998) 265–410, [[dg-ga/9710032](#)].
- [29] P. M. N. Feehan and T. G. Leness, *PU(2) monopoles and relations between four manifold invariants*, *Topology and its Applications* **88** (1998) 111–145, [[dg-ga/9709022](#)].
- [30] P. M. N. Feehan and T. G. Leness, *PU(2) monopoles. 2. Top level Seiberg-Witten moduli spaces and Witten’s conjecture in low degrees*, [dg-ga/9712005](#).
- [31] P. M. N. Feehan and T. G. Leness, *SO(3) monopoles, level one Seiberg-Witten moduli spaces, and Witten’s conjecture in low degrees*, *Topology Appl.* **124** (2002) 221–326, [[math/0106238](#)].
- [32] A. L. Gorodentsev and M. I. Leenson, *How to calculate the correlation function in twisted SYM $N=2$, $N_f = 4$ QFT on projective plane*, [alg-geom/9604011](#).
- [33] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. II. K-theoretic partition function*, [math/0505553](#).
- [34] H. Nakajima and K. Yoshioka, *Perverse coherent sheaves on blowup, iii: Blow-up formula from wall-crossing*, *Kyoto Journal of Mathematics* **51** (2011) 263–335.
- [35] M. Marcolli, *Seiberg-Witten gauge theory*. Hindustan Book Agency, 2011.
- [36] J. Bryan and R. Wentworth, *The multi-monopole equations for Kähler surfaces*, *Turkish J. Math.* **20** (1996) 119–128.
- [37] V. Pidstrigach and A. Tyurin, *Localisation of the donaldson’s invariants along seiberg-witten classes*, *arXiv preprint dg-ga/9507004* (1995) .
- [38] C. Okonek and A. Teleman, *Recent developments in seiberg-witten theory and complex geometry*, *arXiv preprint alg-geom/9612015* (1996) .
- [39] J. Labastida and M. Marino, *Topological quantum field theory and four manifolds*. Springer Netherlands, 2005.
- [40] M. Marino, *Topological quantum field theory and four manifolds*, in *3rd European Congress of Mathematics: Shaping the 21st Century*, 8, 2000. [hep-th/0008100](#).
- [41] H. Nakajima and K. Yoshioka, *Lectures on instanton counting*, in *CRM Workshop on Algebraic Structures and Moduli Spaces Montreal, Canada, July 14-20, 2003*, 2003. [math/0311058](#).
- [42] G. Moore, “Lectures on the physical approach to donaldson and seiberg-witten invariants of four-manifolds.” URL: <https://www.physics.rutgers.edu/~gmoore/SCGP-FourManifoldsNotes-2017.pdf>, 2017.
- [43] A. Malmendier and K. Ono, *SO(3)-Donaldson invariants of \mathbb{P}^2 and Mock Theta Functions*, *Geom. Topol.* **16** (2012) 1767–1833, [[0808.1442](#)].

- [44] M. Griffin, A. Malmendier and K. Ono, *SU(2)-Donaldson invariants of the complex projective plane*, *Forum Math.* **27** (2015) 2003–2023, [[1209.2743](#)].
- [45] A. Malmendier and K. Ono, *Moonshine and Donaldson invariants of \mathbb{P}^2* , [1207.5139](#).
- [46] A. Malmendier, *Donaldson invariants of $\mathbb{P}^1 \times \mathbb{P}^1$ and Mock Theta Functions*, *Commun. Num. Theor. Phys.* **5** (2011) 203–229, [[1008.0175](#)].
- [47] G. Korpas and J. Manschot, *Donaldson-Witten theory and indefinite theta functions*, *JHEP* **11** (2017) 083, [[1707.06235](#)].
- [48] G. Korpas, J. Manschot, G. Moore and I. Nidaiev, *Renormalization and BRST Symmetry in Donaldson–Witten Theory*, *Annales Henri Poincare* **20** (2019) 3229–3264, [[1901.03540](#)].
- [49] G. Korpas, J. Manschot, G. W. Moore and I. Nidaiev, *Mocking the u-plane integral*, *Research in the Mathematical Sciences* **8** (2021) 43.
- [50] J. Aspmann, E. Furrer, G. Korpas, Z.-C. Ong and M.-C. Tan, *The u-plane integral, mock modularity and enumerative geometry*, *Lett. Math. Phys.* **112** (2022) 30, [[2109.04302](#)].
- [51] G. Korpas, *Mock modularity and surface defects in topological $\mathcal{N} = 2$ super Yang-Mills theory*, *Phys. Rev. D* **105** (2022) 026025.
- [52] J. Aspmann, E. Furrer and J. Manschot, *Cutting and gluing with running couplings in $\mathcal{N} = 2$ QCD*, *Phys. Rev. D* **105** (2022) 025021, [[2107.04600](#)].
- [53] J. Aspmann, E. Furrer and J. Manschot, *Elliptic Loci of SU(3) Vacua*, *Annales Henri Poincare* **22** (2021) 2775–2830, [[2010.06598](#)].
- [54] J. Aspmann, E. Furrer and J. Manschot, *Four flavours, triality, and bimodular forms*, *Phys. Rev. D* **105** (2022) 025017, [[2110.11969](#)].
- [55] A. D. Shapere and Y. Tachikawa, *Central charges of $N=2$ superconformal field theories in four dimensions*, *JHEP* **09** (2008) 109, [[0804.1957](#)].
- [56] Y. Ohta, *Prepotentials of $n=2$ su(2) yang–mills theories coupled with massive matter multiplets*, *Journal of Mathematical Physics* **38** (Feb, 1997) 682–696.
- [57] P. C. Argyres and M. R. Douglas, *New phenomena in SU(3) supersymmetric gauge theory*, *Nucl. Phys.* **B448** (1995) 93–126, [[hep-th/9505062](#)].
- [58] P. C. Argyres, M. R. Plesser, N. Seiberg and E. Witten, *New $N=2$ superconformal field theories in four-dimensions*, *Nucl. Phys.* **B461** (1996) 71–84, [[hep-th/9511154](#)].
- [59] A. Klemm, *On the geometry behind $N=2$ supersymmetric effective actions in four-dimensions*, in *33rd Karpacz Winter School of Theoretical Physics: Duality - Strings and Fields*, 5, 1997. [hep-th/9705131](#).
- [60] Y. Ohta, *Prepotential of $N=2$ SU(2) Yang-Mills gauge theory coupled with a massive matter multiplet*, *J. Math. Phys.* **37** (1996) 6074–6085, [[hep-th/9604051](#)].

- [61] E. D'Hoker, I. Krichever and D. Phong, *The effective prepotential of $n = 2$ supersymmetric $su(nc)$ gauge theories*, *Nuclear Physics B* **489** (Mar, 1997) 179–210.
- [62] N. A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831–864, [[hep-th/0206161](#)].
- [63] Y. Ohta, *Non-perturbative solutions to $n=2$ supersymmetric yang-mills theories -progress and perspective-*, 1999.
- [64] N. Nekrasov and A. Okounkov, *Seiberg-Witten theory and random partitions*, *Prog. Math.* **244** (2006) 525–596, [[hep-th/0306238](#)].
- [65] J. Sonnenschein, S. Theisen and S. Yankielowicz, *On the relation between the holomorphic prepotential and the quantum moduli in susy gauge theories*, *Physics Letters B* **367** (Jan, 1996) 145–150.
- [66] T. Eguchi and S.-K. Yang, *Prepotentials of $N=2$ supersymmetric gauge theories and soliton equations*, *Mod. Phys. Lett.* **A11** (1996) 131–138, [[hep-th/9510183](#)].
- [67] E. D'Hoker, I. Krichever and D. Phong, *The Renormalization group equation in $N=2$ supersymmetric gauge theories*, *Nucl. Phys. B* **494** (1997) 89–104, [[hep-th/9610156](#)].
- [68] A. Malmendier, *The Signature of the Seiberg-Witten surface*, *Surveys Diff. Geom.* **15** (2010) 255–278, [[0802.1363](#)].
- [69] M. Caorsi and S. Cecotti, *Special Arithmetic of Flavor*, *JHEP* **08** (2018) 057, [[1803.00531](#)].
- [70] C. Closset and H. Magureanu, *The U -plane of rank-one $4d \mathcal{N} = 2$ KK theories*, [2107.03509](#).
- [71] T. Shioda, *On elliptic modular surfaces*, *J. Math. Soc. Japan* **24** (01, 1972) 20–59.
- [72] M. Schuett and T. Shioda, *Elliptic surfaces*, 2009.
- [73] R. S. Maier, *On Rationally Parametrized Modular Equations*, *arXiv Mathematics e-prints* (Nov., 2006) math/0611041, [[math/0611041](#)].
- [74] R. Miranda, *An overview of algebraic surfaces*, *Lecture Notes in Pure and Appl. Math.* (1997) 157–217.
- [75] D. Gaiotto, G. W. Moore and A. Neitzke, *Wall-crossing, Hitchin Systems, and the WKB Approximation*, [0907.3987](#).
- [76] T. Eguchi, *Seiberg-Witten Theory and S-Duality*, pp. 103–120. Springer Netherlands, Dordrecht, 1999.
- [77] T. Eguchi, K. Hori, K. Ito and S.-K. Yang, *Study of $n = 2$ superconformal field theories in 4 dimensions*, *Nuclear Physics B* **471** (1996) 430 – 442.
- [78] U. Persson, *Configurations of kodaira fibers on rational elliptic surfaces*, *Mathematische Zeitschrift* **205** (1990) 1–47.

- [79] R. Miranda, *Persson's list of singular fibers for a rational elliptic surface*, *Mathematische Zeitschrift* **205** (1990) 191–211.
- [80] W. Nahm, *On the Seiberg-Witten approach to electric - magnetic duality*, [hep-th/9608121](#).
- [81] A. Brandhuber and S. Stieberger, *Periods, coupling constants and modular functions in $N=2$ $SU(2)$ SYM with massive matter*, *Int. J. Mod. Phys. A* **13** (1998) 1329–1344, [[hep-th/9609130](#)].
- [82] M.-x. Huang and A. Klemm, *Holomorphicity and Modularity in Seiberg-Witten Theories with Matter*, *JHEP* **07** (2010) 083, [[0902.1325](#)].
- [83] M. Matone, *Instantons and recursion relations in $N=2$ SUSY gauge theory*, *Phys. Lett.* **B357** (1995) 342–348, [[hep-th/9506102](#)].
- [84] M. Matone, *Koebe $1/4$ theorem and inequalities in $n=2$ supersymmetric qcd*, *Phys. Rev. D* **53** (Jun, 1996) 7354–7358.
- [85] A. Klemm, J. Manschot and T. Wotschke, *Quantum geometry of elliptic Calabi-Yau manifolds*, [1205.1795](#).
- [86] H. Magureanu, *Seiberg-Witten geometry, modular rational elliptic surfaces and BPS quivers*, [2203.03755](#).
- [87] S. K. Donaldson and P. B. Kronheimer, *The geometry of four-manifolds / S.K. Donaldson and P.B. Kronheimer*. Clarendon Press ; Oxford University Press Oxford : New York, 1990.
- [88] J. Aspman, E. Furrer and J. Manschot, “Topological twists of massive sqcd, part ii.” to appear, 2022.
- [89] G. W. Moore, N. Nekrasov and S. Shatashvili, *Integrating over Higgs branches*, *Commun. Math. Phys.* **209** (2000) 97–121, [[hep-th/9712241](#)].
- [90] G. Moore, N. Nekrasov and S. Shatashvili, *Integrating over higgs branches*, *Communications in Mathematical Physics* **209** (2000) 97–121.
- [91] V. Mathai and D. G. Quillen, *Superconnections, Thom classes and equivariant differential forms*, *Topology* **25** (1986) 85–110.
- [92] M. Atiyah and L. Jeffrey, *Topological Lagrangians and cohomology*, *J. Geom. Phys.* **7** (1990) 119–136.
- [93] A. Losev, N. Nekrasov and S. Shatashvili, *Testing seiberg-witten solution*, 1998.
- [94] S. Cordes, G. W. Moore and S. Ramgoolam, *Lectures on 2-d Yang-Mills theory, equivariant cohomology and topological field theories*, *Nucl. Phys. B Proc. Suppl.* **41** (1995) 184–244, [[hep-th/9411210](#)].
- [95] C. Vafa and E. Witten, *A Strong coupling test of S duality*, *Nucl. Phys.* **B431** (1994) 3–77, [[hep-th/9408074](#)].
- [96] E. Witten, *On S duality in Abelian gauge theory*, *Selecta Math.* **1** (1995) 383, [[hep-th/9505186](#)].

- [97] A. E. Nelson and N. Seiberg, *R symmetry breaking versus supersymmetry breaking*, *Nucl. Phys. B* **416** (1994) 46–62, [[hep-ph/9309299](#)].
- [98] M. Marino and G. W. Moore, *The Donaldson-Witten function for gauge groups of rank larger than one*, *Commun. Math. Phys.* **199** (1998) 25–69, [[hep-th/9802185](#)].
- [99] C. Córdova and T. T. Dumitrescu, *Candidate Phases for $SU(2)$ Adjoint QCD_4 with Two Flavors from $\mathcal{N} = 2$ Supersymmetric Yang-Mills Theory*, [1806.09592](#).
- [100] J. Manschot, G. W. Moore and X. Zhang, *Effective gravitational couplings of four-dimensional $\mathcal{N} = 2$ supersymmetric gauge theories*, [1912.04091](#).
- [101] M. Mariño, *The uses of whitham hierarchies*, *Progress of Theoretical Physics Supplement* **135** (1999) 29–52.
- [102] J. M. F. Labastida and C. Lozano, *Duality in twisted $N=4$ supersymmetric gauge theories in four-dimensions*, *Nucl. Phys.* **B537** (1999) 203–242, [[hep-th/9806032](#)].
- [103] M. Marino and F. Zamora, *Duality symmetry in softly broken $N=2$ gauge theories*, *Nucl. Phys. B* **533** (1998) 373–405, [[hep-th/9804038](#)].
- [104] K. Takasaki, *Integrable hierarchies and contact terms in u -plane integrals of topologically twisted supersymmetric gauge theories*, *International Journal of Modern Physics A* **14** (Mar, 1999) 1001–1013.
- [105] R. E. Borcherds, *Automorphic forms with singularities on Grassmannians*, *Invent. Math.* **132** (1998) 491 (1998) , [[alg-geom/9609022](#)].
- [106] K. Bringmann, N. Diamantis and S. Ehlen, *Regularized inner products and errors of modularity*, *International Mathematics Research Notices* **2017** (2017) 7420–7458.
- [107] J. H. Bruinier and J. Funke, *On two geometric theta lifts*, *Duke Mathematical Journal* **125** (2004) 45–90.
- [108] W. Lerche, A. N. Schellekens and N. P. Warner, *Lattices and Strings*, *Phys. Rept.* **177** (1989) 1.
- [109] L. J. Dixon, V. Kaplunovsky and J. Louis, *Moduli dependence of string loop corrections to gauge coupling constants*, *Nucl. Phys.* **B355** (1991) 649–688.
- [110] J. A. Harvey and G. W. Moore, *Algebras, BPS states, and strings*, *Nucl. Phys.* **B463** (1996) 315–368, [[hep-th/9510182](#)].
- [111] G. H. J.H. Bruinier, G. van der Geer and D. Zagier, *The 1-2-3 of Modular Forms*. Springer-Verlag Berlin Heidelberg, 2008, [10.1007/978-3-540-74119-0](#).
- [112] H. Petersson, *Konstruktion der Modulformen und der zu gewissen Grenzkreisgruppen gehörigen automorphen Formen von positiver reeller Dimension und die vollständige Bestimmung ihrer Fourierkoeffizienten*, *S.-B. Heidelberger Akad. Wiss. Math.-Nat. Kl.* (1950) 417–494.
- [113] K. Bringmann, N. Diamantis and S. Ehlen, *Regularized inner products and errors of modularity*, *International Mathematics Research Notices* **2017** (2017) 7420–7458.

- [114] F. Diamond and J. Shurman, *A First Course in Modular Forms*, vol. 228 of *Graduate Texts in Mathematics*. Springer-Verlag New York, 1 ed., 2005.
- [115] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q -series*, vol. 102. American Mathematical Society, cbms regional conference series in mathematics ed., 2004.
- [116] D. Zagier, *Introduction to modular forms; From Number Theory to Physics*. Springer, Berlin (1992), pp. 238-291, 1992.
- [117] D. Schultz, “Notes on modular forms.” URL:
<https://faculty.math.illinois.edu/~schult25/ModFormNotes.pdf>.
- [118] N. Koblitz, *Introduction to Elliptic Curves and Modular Forms*. Graduate Texts in Mathematics. Springer New York, 1993.