

HODGE FILTRATIONS ON TEMPERED HODGE MODULES

DOUGAL DAVIS AND KARI VILONEN

ABSTRACT. In this companion paper to [DV], we show that the Hodge filtration of a tempered Hodge module is generated by the lowest piece of its Hodge filtration. As a consequence, we prove the main conjecture of [SV] in the special case of tempered representations of real reductive Lie groups.

1. INTRODUCTION

In [SV], Schmid and the second author initiated a program and formulated a series of conjectures postulating the existence of a (mixed) Hodge structure on irreducible and standard Harish-Chandra modules. The conjectured Hodge structures arise by taking global sections of appropriate mixed Hodge modules on a flag variety. These conjectures are expected to help in understanding the unitary dual of reductive Lie groups. In [DV], we confirmed that results about unitarity can indeed be obtained from the mixed Hodge modules by deriving a key theorem in [ALTV] as a consequence of a stronger result.

In this paper, we verify this conjecture in the case of tempered Harish-Chandra modules by computing the Hodge filtration explicitly. To be slightly more precise, recall that by Beilinson-Bernstein localization we can associate to Harish-Chandra modules their corresponding Harish-Chandra sheaves. Thus we can talk about tempered sheaves etc. As is explained in [DV], for example, when the infinitesimal character is real the irreducible modules carry a canonical Hodge module structure. In general this Hodge structure can be complicated. However, in this paper we prove the following result, which is Theorem 3.1 in the main body of the paper:

Theorem. *The Hodge filtration of a tempered Hodge module is generated by the lowest piece of its Hodge filtration as a filtered \mathcal{D} -module.*

In other words, the Hodge filtration on a tempered Hodge module is, in some sense, as simple as possible.

As is explained in section 2, tempered Hodge modules are mixtures of tempered Hodge modules for spherical principal series for split groups and Hodge modules associated to closed orbits. In the latter case, the proof follows easily from the definition of push-forward for filtered \mathcal{D} -modules. In the former case a crucial ingredient is the fact that the minimal K -types lie in the lowest piece of the Hodge filtration, one of the main results in [DV].

As a consequence of the theorem above we obtain

Date: September 20, 2022.

2020 Mathematics Subject Classification. 14F10; 22E46; 32S35.

DD was supported by the EPSRC programme grant EP/R034826/1 and the ARC grant FL200100141.

KV was supported in part by the ARC grants DP180101445, FL200100141 and the Academy of Finland.

Theorem. *The main conjecture in [SV] holds in the tempered case.*

This result is stated as Theorem 3.2 and we refer the reader there for a more precise statement.

The authors thank Wilfried Schmid for his contributions to this paper.

2. TEMPERED HARISH-CHANDRA SHEAVES

It is convenient for us to follow the basic set up of [DV], which we recall here briefly. We will work in the context of Harish-Chandra modules. Let us fix a complex reductive group G and an involution θ of G . We write $K = G^\theta$ for the fixed point group. We will always use lower case Gothic letters to denote the corresponding Lie algebras. On the level of Lie algebras we have the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ into eigenspaces of θ . If B is a Borel and N is its unipotent radical then $H = B/N$, the universal Cartan, is independent of the choice of B and comes equipped with a canonical root system. In what follows we will always consider the roots in B to be negative.

We write $\mathrm{HC}(\mathfrak{g}, K)$ for the category of Harish-Chandra modules of the pair (\mathfrak{g}, K) , and $\mathrm{HC}(\mathfrak{g}, K)_\lambda$ for the full subcategory of Harish-Chandra modules with infinitesimal character χ_λ associated with $\lambda \in \mathfrak{h}^*$ under the Harish-Chandra homomorphism. For convenience, we deviate from the notation in [DV] and use Harish-Chandra's convention for the Harish-Chandra homomorphism. In particular, $\mathrm{HC}(\mathfrak{g}, K)_\rho$ contains the trivial representation, and $\lambda = 0$ corresponds to the most singular infinitesimal character and is the center of the action of the Weyl group W .

Let us write $\mathcal{B} = G/B$ for the flag manifold of G . Associated to λ we have the sheaf of twisted differential operators \mathcal{D}_λ on \mathcal{B} . We write $\mathrm{HC}(\mathcal{D}_\lambda, K)$ for the category of Harish-Chandra sheaves, i.e., for the category of K -equivariant \mathcal{D}_λ -modules on \mathcal{B} . If the parameter λ is dominant then, according to Beilinson-Bernstein, each irreducible Harish-Chandra module M is obtained as global sections of a unique irreducible Harish-Chandra sheaf \mathcal{M} . We call an irreducible Harish-Chandra sheaf *tempered* if the associated representation is. The tempered Harish-Chandra modules were first classified in [KZ]. A geometric classification, which we recall below, is given in [HMSW].

An irreducible Harish-Chandra sheaf \mathcal{M} is an intermediate extension of a rank one equivariant λ -twisted local system γ on a K -orbit Q , i.e., $\mathcal{M} = j_{!*}\gamma$ for $j : Q \rightarrow \mathcal{B}$ the inclusion. We call $\mathcal{M} = j_{!*}\gamma$ *clean* if it coincides with $j_!\gamma$ and hence with $j_*\gamma$. Associated to the orbit Q , we may choose, uniquely up to K -conjugacy, a θ -stable Cartan T with a fixed point in Q . We decompose \mathfrak{t} under θ into its eigenspaces as $\mathfrak{t} = \mathfrak{a} \oplus \mathfrak{c}$ with \mathfrak{a} being the (-1) -eigenspace and \mathfrak{c} the $(+1)$ -eigenspace. Identifying T with H via the chosen fixed point, γ is specified by a one dimensional Harish-Chandra module for the pair (\mathfrak{h}, T^θ) , i.e., by a pair $\lambda \in \mathfrak{h}^*$, $\Lambda : T^\theta \rightarrow \mathbb{C}^*$ such that $d\Lambda + \rho = \lambda|_{\mathfrak{c}}$.

We now impose the condition that the infinitesimal character $\lambda \in \mathfrak{h}_{\mathbb{R}}^*$ is real; here $\mathfrak{h}_{\mathbb{R}}^* = \mathbb{R} \otimes_{\mathbb{Z}} \mathbb{X}^*(H)$ for $\mathbb{X}^*(H) = \mathrm{Hom}(H, \mathbb{C}^*)$ the character lattice. All other tempered representations are obtained by moving the parameter $\nu = \lambda|_{\mathfrak{a}^*}$ in the imaginary direction. Under this condition, the classification states that \mathcal{M} is tempered if and only if

$$\lambda|_{\mathfrak{a}} = 0 \quad \text{and} \quad \mathcal{M} = j_{!*}\gamma \text{ is clean.}$$

Thus, in particular, $d\Lambda = \lambda - \rho$ is integral, and hence cleanness of \mathcal{M} imposes a further condition on the orbit Q : for any complex positive root α , the root $\theta\alpha$ must also be positive.

Let us conclude by remarking that orbits Q with the above property have the following simple geometric structure. Let $Z_{\mathfrak{g}}(\mathfrak{c}) = \mathfrak{l} \oplus \mathfrak{c}$ and let us write

$$(2.1) \quad \mathfrak{v} = \bigoplus_{\substack{\alpha \in \Phi^-(\mathfrak{g}, \mathfrak{t}) \\ \alpha|_{\mathfrak{c}} \neq 0}} \mathfrak{g}_{\alpha}.$$

This gives us a parabolic

$$\mathfrak{q}_L = \mathfrak{l} \oplus \mathfrak{c} \oplus \mathfrak{v}.$$

Let us first consider the projection

$$p : \mathcal{B} \rightarrow \mathcal{B}_{Q_L},$$

where \mathcal{B}_{Q_L} denotes the generalized flag manifold of parabolics of type \mathfrak{q}_L . Then \mathfrak{q}_L is θ -stable and hence the image of Q in \mathcal{B}_{Q_L} is closed. The fiber of p is the flag manifold \mathcal{B}_L of L . Furthermore, \mathfrak{a} is the Lie algebra of a maximal torus in L and hence (L, θ) is split and the orbit $Q \cap \mathcal{B}_L$ is the open orbit in \mathcal{B}_L .

3. TEMPERED HODGE MODULES

As in [DV, SV], we work in this section in the context of twisted mixed Hodge modules. In the general set-up in [DV] we work in the context of complex Hodge theory as in [SaSch], but in the tempered case we treat here working with the mixed Hodge modules of Saito [S1, S2] would also suffice.

The category $\mathrm{HC}(\mathcal{D}_{\lambda}, K)$ has a mixed Hodge module version $\mathrm{MHM}(\mathcal{D}_{\lambda}, K)$ (denoted by $\mathrm{MHM}_{\lambda-\rho}(K \setminus \mathcal{B})$ in [DV]) if the parameter $\lambda \in \mathfrak{h}_{\mathbb{R}}^*$, which we assume from now on.

An irreducible Harish-Chandra sheaf $\mathcal{M} = j_{!*}\gamma$ supported on an orbit Q has a unique lift to $\mathrm{MHM}(\mathcal{D}_{\lambda}, K)$ of weight $\dim Q$ with Hodge filtration given by

$$F_p \gamma = \begin{cases} 0 & p < 0, \\ \gamma & p \geq 0. \end{cases}$$

Assume that the infinitesimal character λ is dominant and real. We can now state our main result:

Theorem 3.1. *Let us write $c = \mathrm{codim} Q$. Then the Hodge filtration of an irreducible tempered Harish-Chandra sheaf $j_{!*}\gamma$ supported on Q is generated by $F_c j_{!*}\gamma$.*

We remark that Theorem 3.1 is a local statement about the generation of the Hodge filtration of a tempered Harish-Chandra sheaf as a \mathcal{D}_{λ} -module. In the case of spherical principal series for split groups only, we also prove the global statement that the Hodge filtration of the corresponding tempered Harish-Chandra module is generated by its lowest piece (Theorem 4.1), but we do not know this in general.

We also prove the following result, which is a special case of a conjecture of Schmid and the second author [SV].

Theorem 3.2. *In the context of Theorem 3.1, let S be the polarization on $j_{!*}\gamma$ and $\Gamma(S)$ the induced Hermitian form on the irreducible tempered (\mathfrak{g}, K) -module $V := \Gamma(j_{!*}\gamma)$ (see e.g., [DV, §4.3]). Then the form*

$$\Gamma(S)|_{F_p V \cap (F_{p-1} V)^{\perp}}$$

is $(-1)^{p-c}$ -definite for all p .

4. TEMPERED HODGE MODULES FOR SPLIT GROUPS

In this section we consider the special case of a Hodge module \mathcal{M} associated with a tempered spherical principal series representation of a split group. Such an \mathcal{M} is an intermediate extension of a local system on the open orbit Q . By the considerations of section 2, we know that the infinitesimal character is zero (i.e., \mathcal{M} is a \mathcal{D}_0 -module) and that the extension is clean. Thus, \mathcal{M} is self dual as a Hodge module and hence as a filtered \mathcal{D}_0 -module. We also recall that $\mathcal{D}_0^{op} = \mathcal{D}_0$, so the filtered dual is given simply by the filtered derived hom

$$\mathbb{D}(\mathcal{M}, F) = \mathrm{RHom}_{(\mathcal{D}_0, F)}((\mathcal{M}, F), (\mathcal{D}_0, F))[\dim \mathcal{B}]$$

without the need for a twist by a line bundle.

The corresponding spherical Harish-Chandra module $M = \Gamma(\mathcal{B}, \mathcal{M})$ is isomorphic, as a (\mathfrak{g}, K) -module, to $U(\mathfrak{g})_0 \otimes_{U(\mathfrak{k})} \mathbb{C}_0$, where $U(\mathfrak{g})_0$ is the quotient of $U(\mathfrak{g})$ through which the center acts by infinitesimal character χ_0 and \mathbb{C}_0 is the trivial representation of K . We have a corresponding description of \mathcal{M} as

$$\mathcal{M} = \mathcal{D}_0 \otimes_{U(\mathfrak{k})} \mathbb{C}_0.$$

Using this description, we may view \mathcal{M} as a filtered module with the filtration induced from \mathcal{D}_0 . We denote this filtration by $F'_\bullet \mathcal{M}$. This filtration is generated by the rank one \mathcal{O}_X -module $F'_0 \mathcal{M}$. Let us identify \mathfrak{g} with \mathfrak{g}^* , write \mathcal{N} for the nilpotent cone in \mathfrak{g} and write $\mu : T^*\mathcal{B} \rightarrow \mathcal{N}$ for the moment map. We then have

Theorem 4.1. *The filtration F' on \mathcal{M} coincides with the Hodge filtration F and $\mathrm{gr}_\bullet^F \mathcal{M} = \mu^*(\mathcal{O}_{\mathfrak{p}})$. Furthermore, $\mathrm{H}^k(\mathcal{B}, \mathrm{gr}_\bullet^F \mathcal{M}) = 0$ for $k > 0$ and $\Gamma(\mathcal{B}, \mathrm{gr}_\bullet^F \mathcal{M}) = \mathcal{O}_{\mathcal{N} \cap \mathfrak{p}}$.*

The rest of this section is devoted to the proof of this theorem. We begin with

Lemma 4.2. *The filtered module (\mathcal{M}, F') is filtered self dual and $\mathrm{gr}_\bullet^{F'} \mathcal{M} = \mu^*(\mathcal{O}_{\mathfrak{p}})$ is Cohen-Macaulay.*

It is perhaps helpful to recall that the associated graded of the Hodge filtration of any Hodge module is Cohen-Macaulay. The Cohen-Macaulay condition is necessary to have a good notion of filtered dual.

Proof. We begin with some preparatory statements. Observe that we have $\mu^{-1}(\mathfrak{p}) = T_K^* \mathcal{B}$, the union of conormal bundles of K -orbits. Thus, because the group is split, $\mu^{-1}(\mathfrak{p})$ is of pure codimension

$$\mathrm{codim}_{T^*\mathcal{B}} \mu^{-1}(\mathfrak{p}) = \dim K.$$

We conclude that the scheme $\mu^{-1}(\mathfrak{p})$, being the preimage of a codimension $\dim K$ complete intersection under a map of smooth varieties, is itself a complete intersection and its structure sheaf $\mu^*(\mathcal{O}_{\mathfrak{p}})$ has the Koszul resolution

$$(4.1) \quad \cdots \rightarrow \mathcal{O}_{T^*\mathcal{B}}\{i\} \otimes \wedge^i \mathfrak{k} \rightarrow \mathcal{O}_{T^*\mathcal{B}}\{i-1\} \otimes \wedge^{i-1} \mathfrak{k} \rightarrow \cdots \rightarrow \mathcal{O}_{T^*\mathcal{B}}\{1\} \otimes \mathfrak{k} \rightarrow \mathcal{O}_{T^*\mathcal{B}},$$

where $\{\cdot\}$ denotes a grading shift.

Let us now consider the filtered module (\mathcal{M}, F') . It has the following filtered resolution

$$(4.2) \quad \cdots \rightarrow \mathcal{D}_0\{i\} \otimes \wedge^i \mathfrak{k} \rightarrow \mathcal{D}_0\{i-1\} \otimes \wedge^{i-1} \mathfrak{k} \rightarrow \cdots \rightarrow \mathcal{D}_0\{1\} \otimes \mathfrak{k} \rightarrow \mathcal{D}_0.$$

Here $\mathcal{D}_0\{i\}$ denotes the filtered module \mathcal{D}_0 with filtration shifted so that it begins in degree i . To verify that the complex above is a resolution we pass to the associated graded complex. The graded resolution coincides with (4.1). Thus we conclude that (4.2) is a filtered resolution of (\mathcal{M}, F') and, furthermore that $\mathrm{gr}_{\bullet}^{F'} = \mu^*(\mathcal{O}_{\mathfrak{p}})$

We now form the filtered dual $\mathbb{D}(\mathcal{M}, F')$ by making use of the resolution above. We obtain, writing $n = \dim(X) = \dim \mathfrak{k}$:

$$\cdots \rightarrow \mathcal{D}_0\{-i\} \otimes \wedge^i \mathfrak{k}^* \rightarrow \cdots \rightarrow \mathcal{D}_0\{-n+1\} \otimes \wedge^{n-1} \mathfrak{k}^* \rightarrow \mathcal{D}_0\{-n\} \otimes \wedge^n \mathfrak{k}^*.$$

Thus, we conclude that $\mathbb{D}(\mathcal{M}, F') = (\mathcal{M}, F'\{-n\})$, i.e., that (\mathcal{M}, F') is filtered self dual. \square

Lemma 4.3. *If $F'_0\mathcal{M} \subset F_0\mathcal{M}$ then $F'_i\mathcal{M} = F_i\mathcal{M}$ for all i .*

Proof. Our hypothesis implies that the identity map is a morphism of filtered modules $(\mathcal{M}, F') \rightarrow (\mathcal{M}, F)$, i.e., that $F'_i\mathcal{M} \subset F_i\mathcal{M}$ for all i . As both (\mathcal{M}, F') and (\mathcal{M}, F) are self dual filtered modules we obtain a morphism of filtered modules $(\mathcal{M}, F) \rightarrow (\mathcal{M}, F')$, i.e., that $F_i\mathcal{M} \subset F'_i\mathcal{M}$ for all i . Thus $F'_i\mathcal{M} = F_i\mathcal{M}$ for all i . \square

Proof of Theorem 4.1. By the above lemmas, it remains to show that $F'_0\mathcal{M} \subset F_0\mathcal{M}$. The module $F'_0\mathcal{M}$ is generated by the minimal K -type \mathbb{C}_0 . But by [DV, Theorem 4.5] the minimal K -type lies in $F_0\mathcal{M}$, so this concludes the proof of the first part of the theorem.

We will now give a proof of the second part. We will show that the global sections of the Koszul resolution (4.1) give a resolution for $\mathcal{O}_{\mathcal{N} \cap \mathfrak{p}}$. Since $H^k(\mathcal{B}, \mathcal{O}_{T^*\mathcal{B}}) = 0$ for $k > 0$, this implies the statement.

As the group is split the Cartan subspace \mathfrak{a} of \mathfrak{p} is also a Cartan for \mathfrak{g} and the little Weyl group of \mathfrak{a} is the full Weyl group of \mathfrak{g} . Thus the restriction map

$$(4.3) \quad \mathbb{C}[\mathfrak{g}]^G \xrightarrow{\sim} \mathbb{C}[\mathfrak{p}]^K$$

is an isomorphism. Consider the diagram

$$\begin{array}{ccc} \mathfrak{p} & \longrightarrow & \mathfrak{g} \\ \downarrow & & \downarrow \\ \mathrm{Spec} \mathbb{C}[\mathfrak{p}]^K & \xlongequal{\quad} & \mathrm{Spec} \mathbb{C}[\mathfrak{g}]^G. \end{array}$$

By Kostant [K, Theorems 0.1 and 0.2] and Kostant-Rallis [KR, Theorems 14 and 15], the vertical maps are flat and their scheme-theoretic fibers \mathcal{N} and $\mathcal{N} \cap \mathfrak{p}$ are reduced complete intersections. We see therefore that

$$\mathcal{O}_{\mathcal{N} \cap \mathfrak{p}} = \mathcal{O}_{\mathcal{N}} \otimes_{\mathcal{O}_{\mathfrak{g}}}^{\mathbb{L}} \mathcal{O}_{\mathfrak{p}}$$

has the graded Koszul resolution

$$(4.4) \quad \cdots \rightarrow \mathcal{O}_{\mathcal{N}}\{i\} \otimes \wedge^i \mathfrak{k} \rightarrow \mathcal{O}_{\mathcal{N}}\{i-1\} \otimes \wedge^{i-1} \mathfrak{k} \rightarrow \cdots \rightarrow \mathcal{O}_{\mathcal{N}}\{1\} \otimes \mathfrak{k} \rightarrow \mathcal{O}_{\mathcal{N}}.$$

But (4.4) is given by global sections of (4.1) so we are done. \square

5. PROOF OF THEOREM 3.1

Let us now consider a general irreducible tempered Hodge module \mathcal{M} . By the considerations in section 2, we have $\mathcal{M} = j_{!*}\gamma$ where γ is a clean local system on an orbit Q with the special properties specified in that section. Using the notation in section 2 we write $S = p(Q)$ for the closed K -orbit in the partial flag variety. Then

$\bar{Q} = p^{-1}(S)$ and we have a K -equivariant smooth fibration $\bar{p} : \bar{Q} \rightarrow S$. We further write $i : \bar{Q} \rightarrow \mathcal{B}$ for the closed embedding and $\tilde{j} : Q \rightarrow \bar{Q}$ for the open embedding.

Let us start by considering $\mathcal{N} = \tilde{j}_*(\gamma)$. Let us consider the fiber \mathcal{B}_L of \bar{p} and the restrictions $\mathcal{N}|_{\mathcal{B}_L}$ and $\gamma_L = \gamma|_{Q \cap \mathcal{B}_L}$. As the restriction is non-characteristic, we have $\mathcal{N}|_{\mathcal{B}_L} = \tilde{j}_{!*}\gamma_L$. So, after we adjust the cohomological shift and the weights, the Hodge module $\mathcal{N}|_{\mathcal{B}_L}$ is the tempered spherical principal series sheaf considered in section 4 for (\mathfrak{l}, K_L) , and so we know that its Hodge filtration is generated by $F_0\mathcal{N}|_{\mathcal{B}_L}$. Thus the same is true for \mathcal{N} and we conclude that its Hodge filtration is generated by $F_0\mathcal{N}$.

Let us write \mathcal{J} for the ideal sheaf of \bar{Q} in \mathcal{B} . As $\mathcal{M} = i_*\mathcal{N}$ and i is an inclusion of a closed smooth subvariety

$$\mathcal{M} = i_*\mathcal{N} = (\mathcal{D}_{\mathcal{B},\lambda}/\mathcal{D}_{\mathcal{B},\lambda}\mathcal{J}) \otimes_{\mathcal{O}_{\bar{Q}}} \mathcal{N} \otimes_{\mathcal{O}_{\bar{Q}}} \omega_{\bar{Q}/\mathcal{B}}.$$

Let us write $c = \text{codim } Q$. Then, by the definition of filtered proper push-forwards, we have

$$F_p\mathcal{M} = F_p i_*\mathcal{N} = \sum_{r+k \leq p-c} (F_k \mathcal{D}_{\mathcal{B},\lambda} / F_k \mathcal{D}_{\mathcal{B},\lambda} \mathcal{J}) \otimes_{\mathcal{O}_{\bar{Q}}} F_r \mathcal{N} \otimes_{\mathcal{O}_{\bar{Q}}} \omega_{\bar{Q}/\mathcal{B}}.$$

From this formula and the fact that the Hodge filtration of \mathcal{N} is generated by $F_0\mathcal{N}$ we conclude that the Hodge filtration of \mathcal{M} is generated by $F_c\mathcal{M}$.

6. PROOF OF THEOREM 3.2

We will prove Theorem 3.2 by appealing to the known unitarity of tempered Harish-Chandra modules. We first recall the relationship between compact and non-compact real forms for G and the corresponding invariant Hermitian forms, cf., [ALTV, §12].

Recall that we have fixed a complex reductive group G with an involution θ . We choose also a compact real form $U_{\mathbb{R}} \subset G$ invariant under θ . Denoting by $\sigma_c : G \rightarrow G$ the complex conjugation with respect to $U_{\mathbb{R}}$, the involution θ determines another real form $G_{\mathbb{R}} = G^{\theta\sigma_c}$. The corresponding real Lie algebras are $\mathfrak{u}_{\mathbb{R}} = \mathfrak{g}^{\sigma_c}$ and $\mathfrak{g}_{\mathbb{R}} = \mathfrak{g}^{\theta\sigma_c}$. A key property of tempered Harish-Chandra modules is that they always admit positive definite $\mathfrak{g}_{\mathbb{R}}$ -invariant Hermitian forms.

The link between the $\mathfrak{g}_{\mathbb{R}}$ -invariant forms and the polarizations is as follows. First, since the flag variety \mathcal{B} and the universal Cartan H are canonically associated with G , the involution $\theta : G \rightarrow G$ induces compatible involutions on both. We will write $\delta : H \rightarrow H$ for the induced involution on H . We remark that δ preserves the positive roots in $\mathbb{X}^*(H)$, and if $Q \subset \mathcal{B}$ is a closed K -orbit and T is a θ -stable Cartan with a fixed point in Q , then δ agrees with $\theta|_T$ under the corresponding identification $T \cong H$. The involution $\theta : \mathcal{B} \rightarrow \mathcal{B}$ also lifts non-canonically to the H -torsor $\tilde{\mathcal{B}}$, intertwining the action of δ on H ; we fix such a lift in what follows.

For any $(\mathcal{D}_{\lambda}, K)$ -module \mathcal{M} , the pullback $\theta^*\mathcal{M}$ is a $(\mathcal{D}_{\delta\lambda}, K)$ -module, and we have an isomorphism

$$\Gamma(\mathcal{M}) \xrightarrow{\sim} \Gamma(\theta^*\mathcal{M})$$

intertwining $\theta : U(\mathfrak{g}) \rightarrow U(\mathfrak{g})$. If $\mathcal{M} = j_{!*}\gamma$ is an irreducible tempered module and $\lambda \in \mathfrak{h}_{\mathbb{R}}^*$ is real, then we have $\delta\lambda = \lambda$ and $\theta^*j_{!*}\gamma \cong j_{!*}\gamma$. We will fix such an isomorphism so that the induced map

$$\theta : \Gamma(j_{!*}\gamma) \rightarrow \Gamma(j_{!*}\gamma)$$

is equal to the identity on the (unique) minimal K -type. In particular, the above map is an involution.

Suppose now that $\lambda \in \mathfrak{h}_{\mathbb{R}}^*$, and $j_{!*}\gamma$ is an irreducible tempered $(\mathcal{D}_{\lambda}, K)$ -module, equipped with its polarization S as a pure Hodge module. We therefore have the $\mathfrak{u}_{\mathbb{R}}$ -invariant form $\Gamma(S)$ on $\Gamma(j_{!*}\gamma)$; it follows immediately that the form

$$\langle u, v \rangle := \Gamma(S)(u, \theta v)$$

is $\mathfrak{g}_{\mathbb{R}}$ -invariant. Since the tempered representation $\Gamma(j_{!*}\gamma)$ is unitary as a $\mathfrak{g}_{\mathbb{R}}$ -module and $\Gamma(S)$ is positive definite on the minimal K -type by [DV, Theorem 4.3 and Proposition 4.7], we deduce the following.

Proposition 6.1. *For $\epsilon = \pm 1$, the polarization $\Gamma(S)$ is ϵ -definite on the ϵ -eigenspace $\Gamma(j_{!*}\gamma)^{\epsilon\theta}$.*

We now claim the following.

Proposition 6.2. *The associated graded of the Hodge filtration on $\Gamma(j_{!*}\gamma)^{\theta}$ (resp., $\Gamma(j_{!*}\gamma)^{-\theta}$) is concentrated in degrees $2k + c$ (resp., $2k + 1 + c$) for $k \in \mathbb{Z}_{\geq 0}$.*

Proof of Theorem 3.2. Follows immediately from Propositions 6.1 and 6.2. \square

Proof of Proposition 6.2. Consider first the case of a tempered spherical principal series for a split group as in section 4. By Theorem 4.1, we have

$$\mathrm{gr}_{\bullet}^F \Gamma(j_{!*}\gamma) = \mathcal{O}_{\mathcal{N} \cap \mathfrak{p}}$$

is naturally a graded quotient of $\mathrm{Sym}(\mathfrak{p})$. Since θ acts on \mathfrak{p} with eigenvalue -1 by definition the result in this case follows.

For the general tempered case, consider as in section 5 the smooth fibration

$$\bar{p}: \bar{Q} \rightarrow S$$

and the Hodge module $\tilde{j}_{!*}\gamma$ on \bar{Q} . We have

$$(6.1) \quad \Gamma(\mathcal{B}, \mathrm{gr}_{\bullet}^F \tilde{j}_{!*}\gamma) = \Gamma(S, \mathcal{F}\{c\}),$$

where

$$\mathcal{F} = \mathrm{Sym}(\mathcal{N}_{S/\mathcal{B}_{Q_L}}) \otimes \omega_{S/\mathcal{B}_{Q_L}} \otimes \bar{p}_* \mathrm{gr}_{\bullet}^F \tilde{j}_{!*}\gamma,$$

$\{c\}$ denotes a grading shift, and \bar{p}_* the sheaf-theoretic pushforward. Since S is closed, it is fixed by θ pointwise, so θ acts on the sheaf \mathcal{F} . Since $\bar{p}_* \tilde{j}_{!*}\gamma$ is fiberwise a tempered spherical principal series for L , the $+1$ (resp., -1) eigenspace of $\mathrm{gr}_{\bullet}^F \tilde{j}_{!*}\gamma$ is concentrated in only even (resp., odd) degrees as shown above. Moreover, θ acts on the normal bundle $\mathcal{N}_{S/\mathcal{B}_{Q_L}}$ with eigenvalue -1 , so the above is also true for \mathcal{F} . The proposition now follows by (6.1). \square

REFERENCES

- [ALTV] Adams, Jeff; van Leeuwen, Marc; Trapa, Peter; Vogan, David, *Unitary representations of real reductive groups*, Astérisque 417 (2020).
- [DV] Davis, Dougal; Vilonen, Kari, *Mixed Hodge modules and real groups*, arXiv:2202.08797.
- [HMSW] Hecht Henryk; Miličić, Dragan; Schmid, Wilfried; Wolf, Joseph, *Localization and standard modules for real semisimple Lie groups II: irreducibility, vanishing theorems, and classification*, manuscript.
- [K] Kostant, Bertram, *Lie group representations on polynomial rings*, Amer. J. Math. 86 (1963), 327–402.
- [KR] Kostant, Bertram; Rallis, Steven, *Orbits and representations associated with symmetric spaces*, Amer. J. Math. 93 (1971), 753–809.

- [KZ] Knapp, Anthony W.; Zuckerman, Gregg J., *Classification of irreducible tempered representations of semisimple groups I, II*, Annals of Math., 11 (1982) 389–501.
- [SaSch] Sabbah, Claude; Schnell, Christian, *Mixed Hodge Module Project*, <http://www.cmls.polytechnique.fr/perso/sabbah/MHMPProject/mhm.html>.
- [S1] Saito, Morihiko, *Modules de Hodge polarisables*, Publ. Res. Inst. Math. Sci. 24 (1988), no. 6, 849–995.
- [S2] Saito, Morihiko, *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. 26 (1990), no. 2, 221–333.
- [SV] Schmid, Wilfried; Vilonen, Kari, *Hodge theory and unitary representations of reductive Lie groups*. In: *Frontiers of Mathematical Sciences*, International Press, Somerville MA (2011), 397–420.

(DD) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA

Email address: dougal.davis1@unimelb.edu.au

(KV) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA, ALSO DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI, HELSINKI, FINLAND

Email address: kari.vilonen@unimelb.edu.au, kari.vilonen@helsinki.fi