

Existence and Minimax Theorems for Adversarial Surrogate Risks in Binary Classification

Natalie S. Frank

NF1066@NYU.EDU

Courant Institute

New York University

New York, NY 10012, USA

Editor: Kevin Murphy and Bernhard Schölkopf

Abstract

Adversarial training is one of the most popular methods for training methods robust to adversarial attacks, however, it is not well-understood from a theoretical perspective. We prove existence, regularity, and minimax theorems for adversarial surrogate risks. Our results explain some empirical observations on adversarial robustness from prior work and suggest new directions in algorithm development. Furthermore, our results extend previously known existence and minimax theorems for the adversarial classification risk to surrogate risks.

Keywords: Adversarial Learning, Direct Method of the Calculus of Variations, Minimax Theorems, Wasserstein metric, Adversarial Bayes risk

1. Introduction

Deep neural networks are state-of-the-art methods for a variety of machine learning tasks including image classification and speech recognition. However, a concerning problem with these models is their susceptibility to *adversarial attacks*: small perturbations introduced at test-time can cause incorrect classification by the network (Szegedy et al., 2013). This issue has security implications; for instance, (Gu et al., 2017) show that a yellow sticker can cause a neural net to misclassify a stop sign. Self-driving cars often utilize such neural nets. Furthermore, one can find adversarial examples that generalize to other neural nets; these sort of attacks are called *transfer attacks*. In other words, an adversarial example generated for one neural net will also be an adversarial example for a different neural net trained for the same classification problem (Tramèr et al., 2017; Demontis et al., 2018; Kurakin et al., 2017; Rozsa et al., 2016; Papernot et al., 2016). This phenomenon shows that access to a specific neural net is not necessary for generating adversarial examples. One method for defending against such adversarial perturbations is *adversarial training*, in which a neural net is trained on such adversarially perturbed data points (Kurakin et al., 2017; Madry et al., 2019; Wang et al., 2021). However, adversarial training is not well understood from a theoretical perspective.

Several prior works study properties of the *adversarial classification risk*, which is the expected number of classification errors under adversarial perturbations. (Awasthi et al., 2021; Bungert et al., 2021; Pydi and Jog, 2021) all prove existence of the adversarial Bayes classifier. However, because minimizing the empirical adversarial classification risk is com-

putationally intractable, the adversarial training procedure minimizes adversarial perturbations of a surrogate risks. In order to gain further insight on adversarial training procedure, this paper extends some of these results in prior work to these surrogate risks. We first prove existence theorems for supremum-based adversarial surrogate risks. One of these theorems implies that for perturbations in the 2-norm, there always exists a minimizer f for which the principal curvatures of the level sets is bounded by the inverse of the perturbation radius. In fact, one of our intermediate results states that reducing the curvature of the level sets of a function will always reduce the adversarial risk. This observation explains the experiments of (Moosavi-Dezfooli et al., 2018) who show that cross-sections of robust decision surfaces have lower curvature. Moreover, this observation inspired research on regularizers that induce adversarial robustness. However, the regularity which we prove for robust functions is somewhat different than the Jacobian regularization previously studied in the context of adversarial learning (Ross and Doshi-Velez, 2017; Moosavi-Dezfooli et al., 2018; Yu et al., 2018; Jakubovitz and Giryes, 2018). In Section 4 we suggest a new regularizer that might induce this regularity. Finding alternative regularizers for adversarial robustness remains an open question.

Next, we use one of these existence theorems to prove a minimax relation characterizing the adversarial learning problem. This idea was first pioneered by Pydi and Jog (2021) who rephrased the adversarial learning problem as a game. In short, such minimax theorems recast the adversarial learning problem as a game between two players: a *learner*, who aims to find a function with minimal adversarial risk, and an *adversary*, who tries to pick the maximal adversarial perturbation. Our minimax theorem states that the player to choose first does not have an advantage. These results suggest an explanation for transfer attacks on adversarially trained networks, which we discuss in Section 5. (Demontis et al., 2018) discussed an explanation of transfer attacks on neural nets trained using standard methods, but did not extend their analysis to the adversarial training setting.

Both our minimax and existence theorems implications for understating statistical properties of surrogate losses. Recall that one minimizes a surrogate risk because minimizing the original risk is computationally intractable. If a sequence of functions which minimizes the surrogate risk also minimizes the classification risk, then that surrogate is referred to as a *consistent risk*. Similarly, if a sequence of functions which minimizes the adversarial surrogate risk also minimizes the adversarial classification risk, then that surrogate is referred to as an *adversarially consistent risk*. Much prior work studies the consistency of surrogate risks (Bartlett et al., 2006; Lin, 2004; Steinwart, 2007; Philip M. Long, 2013; Mingyuan Zhang, 2020; Zhang, 2004). Follow-up work used the minimax and existence theorems to characterize adversarially consistent supremum-based risks for binary classification with distributions absolutely continuous with respect to Lebesgue measure (Frank and Niles-Weed, 2022). Specifically, Frank and Niles-Weed (2022) uses our theorems to reduce adversarial consistency for supremum-based risks to (non-adversarial) consistency. These results show that our theorems about the structure of surrogate risks can improve our understanding of adversarial learning algorithms.

We will now discuss the outline of the paper. The problem setup and notation are presented in Section 3. We state our existence results at the beginning of Section 4.4 and provide the proofs in the following subsections. Next, we prove our minimax result

in Section 5. The proofs of some technical lemmas are delayed to the Appendix. The arguments in Appendices B through D all depend on the results in B.2.

2. Related Works

This paper extends prior work on the adversarial Bayes classifier. We start by highlighting three papers which inspired our direction and proof techniques. Awasthi et al. (2021); Pydi and Jog (2021); Bungert et al. (2021) all proved the existence of the adversarial Bayes classifier using different methods. Notably, (Pydi and Jog, 2021) first proved a minimax theorem for the adversarial risk, (Bungert et al., 2021) studied minimal and maximal adversarial Bayes classifiers, and (Awasthi et al., 2021) introduced the concept of pseudo-certifiably robustness. Our analysis of the existence of minimizers to surrogate risks combines their various approaches. (Pydi and Jog, 2021; Bungert et al., 2021) also introduced promising mathematical techniques not discussed in this paper—most notably (Pydi and Jog, 2021) use Choquet capacities and Strassen’s theorem from optimal transport to study the adversarial risk, (Bungert et al., 2021) used the nonlocal perimeter from PDEs to prove the existence of the adversarial Bayes classifier. All three of these papers study properties of the $\epsilon, -\epsilon$ operations applied to sets; we study the operations S_ϵ, I_ϵ which we show are generalizations of $\epsilon, -\epsilon$ to functions.

Next we discuss some papers which prove further results on the adversarial Bayes classifier. Yang et al. (2020) studied the adversarial Bayes classifier in the context of non-parametric methods. The papers (Pydi and Jog, 2019; Bhagoji et al., 2019) further introduced methods from optimal transport to study adversarial learning. The methods of (Bungert et al., 2021) are inspired by the use of the non-local perimeter in PDEs. Papers such as (Cesaroni and Matteo, 2017; Cesaroni et al., 2018; Chambolle et al., 2012, 2015; Bellettini, 2004) use the non-local perimeter and other methods to study variational problems. Lastly, (Frank, 2022; Trillos and Murray, 2020) give necessary and sufficient conditions describing the boundary of the adversarial Bayes classifier.

Many papers study implementing and improving the adversarial training algorithm. (Wang et al., 2021; Kurakin et al., 2017; Madry et al., 2019) study the adversarial training algorithm and improving the runtime. Two particularly popular attacks used in adversarial training are the FGSM attack (Goodfellow et al., 2014) and the PGD attack (Madry et al., 2019). More recent popular variants of this algorithm include (Shafahi et al., 2019; Xie et al., 2018; Kannan et al., 2018). This list of references is far from complete, there are many other promising attack and defense algorithms.

3. Background and Notation

3.1 Adversarial Classification

This paper studies binary classification on \mathbb{R}^d . The two classes are -1 and $+1$. Data is distributed according to a distribution \mathcal{D} on $\mathbb{R}^d \times \{-1, +1\}$. We denote the marginal distributions according to the class label as $\mathbb{P}_0(S) = \mathcal{D}(S \times \{-1\})$ and $\mathbb{P}_1(S) = \mathcal{D}(S \times \{+1\})$.

To classify points in \mathbb{R}^d , algorithms typically learn a real-valued function f and then classify points \mathbf{x} according to the sign of f , where we define sign as

$$\text{sign } \alpha = \begin{cases} +1 & \text{if } \alpha > 0 \\ -1 & \text{if } \alpha \leq 0 \end{cases}.$$

The sign of 0 was chosen arbitrarily. The *classification error* of a function f is then

$$R(f) = \int \mathbf{1}_{f(\mathbf{x}) \leq 0} d\mathbb{P}_1 + \int \mathbf{1}_{f(\mathbf{x}) > 0} d\mathbb{P}_0. \quad (1)$$

Notice that finding minimizers to R is straightforward: define the measure $\mathbb{P} = \mathbb{P}_0 + \mathbb{P}_1$ and let $\eta = d\mathbb{P}_1/d\mathbb{P}$. Then the risk R can be re-written as

$$R(f) = \int \eta(\mathbf{x}) \mathbf{1}_{f(\mathbf{x}) \leq 0} + (1 - \eta(\mathbf{x})) \mathbf{1}_{f(\mathbf{x}) > 0} d\mathbb{P}.$$

Hence a minimizer of R must minimize the function $\alpha \mapsto \eta(\mathbf{x}) \mathbf{1}_{\alpha \leq 0} + (1 - \eta(\mathbf{x})) \mathbf{1}_{\alpha > 0}$ point-wise.

This paper analyzes the *evasion attack*, in which an adversary perturbs each input before it is evaluated by the function f . In this attack model, the adversary knows both the function f and the true label of the data point. To constrain the adversary, we assume that perturbations are bounded by ϵ in a norm $\|\cdot\|$. Thus a point \mathbf{x} with label +1 is misclassified if there is a perturbation \mathbf{h} with $\|\mathbf{h}\| \leq \epsilon$ for which $f(\mathbf{x} + \mathbf{h}) \leq 0$ and a point \mathbf{x} with label -1 is misclassified if there is a perturbation \mathbf{h} with $\|\mathbf{h}\| \leq \epsilon$ for which $f(\mathbf{x} + \mathbf{h}) > 0$. Therefore, the *adversarial classification risk* is

$$R^\epsilon(f) = \int \sup_{\|\mathbf{h}\| \leq \epsilon} \mathbf{1}_{f(\mathbf{x} + \mathbf{h}) \leq 0} d\mathbb{P}_1 + \int \sup_{\|\mathbf{h}\| \leq \epsilon} \mathbf{1}_{f(\mathbf{x} + \mathbf{h}) > 0} d\mathbb{P}_0 \quad (2)$$

which is the expected proportion of errors under subject to the adversarial evasion attack.

Previous works study minimizers of R^ϵ and the nature of this minimization problem. Notice that establishing the existence of minimizers to R^ϵ is non-trivial— one can re-write R^ϵ in terms of \mathbb{P}_0 and \mathbb{P}_1 but the resulting integrand cannot be minimized in a pointwise fashion. (Awasthi et al., 2021; Bungert et al., 2021; Pydi and Jog, 2021) all establish the existence of the adversarial Bayes classifier. Awasthi et al. (2021) further show that level sets of f satisfy the *pseudo-certifiably robustness* property:

Definition 1 Fix a perturbation radius $\epsilon > 0$. We say that a set A is pseudo-certifiably robust at a point \mathbf{x} with radius ϵ if either $\mathbf{x} \in A$ and there exists a ball $B_\epsilon(\mathbf{y})$ with $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ and $B_\epsilon(\mathbf{y}) \subset A$ or $\mathbf{x} \in A^C$ and there exists a ball $B_\epsilon(\mathbf{y})$ with $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ and $B_\epsilon(\mathbf{y}) \subset A^C$. We say a set A is pseudo-certifiably robust if it is pseudo-certifiably robust with radius ϵ at every point.

In other words, a set is pseudo-certifiably robust if for every point $\mathbf{x} \in \mathbb{R}^d$, there is a ball $\overline{B_\epsilon(\mathbf{y})}$ for which $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ and $B_\epsilon(\mathbf{y})$ is classified the same as \mathbf{x} . A pseudo-certifiably set is fairly regular, in fact, Theorem 1 of Lewicka and Peres (2020) implies that the boundary of a set which is pseudo-certifiably robust with respect to perturbations in the 2-norm is a $C^{1,1}$ manifold. Awasthi et al. (2021) show that there always exists a minimizer to R^ϵ with this type of regularity:

Theorem 2 *There always exists a minimizer to R^ϵ for which the sets $\{\mathbf{x}: f(\mathbf{x}) > 0\}$ and $\{\mathbf{x}: f(\mathbf{x}) \leq 0\}$ are pseudo-certifiably robust.*

Next, we will discuss Theorem 7.1 of (Pydi and Jog, 2021) which suggests a new interpretation of adversarial examples. To start, we introduce the ∞ -Wasserstein metric on measures. Let \mathbb{P}, \mathbb{Q} be two Borel measures with $\mathbb{P}(\mathbb{R}^d) = \mathbb{Q}(\mathbb{R}^d)$. A *coupling* is a measure on the product space $\mathbb{R}^d \times \mathbb{R}^d$ with marginals \mathbb{P}, \mathbb{Q} . We denote the set of all couplings with marginals \mathbb{P}, \mathbb{Q} by $\Pi(\mathbb{P}, \mathbb{Q})$. The ∞ -Wasserstein distance with respect to a norm $\|\cdot\|$ is defined as

$$W_\infty(\mathbb{P}, \mathbb{Q}) = \inf_{\pi \in \Pi(\mathbb{P}, \mathbb{Q})} \text{ess sup}_{(\mathbf{x}, \mathbf{x}') \sim \pi} \|\mathbf{x} - \mathbf{x}'\|$$

Pydi and Jog (2021) prove the following minimax result for the classification risk:

Theorem 3 *Assume that $\mathbb{P}_0, \mathbb{P}_1$ are absolutely continuous with respect to Lebesgue measure. Let*

$$\hat{R}(f, \mathbb{P}_1, \mathbb{P}_0) = \int \mathbf{1}_{f(\mathbf{x}) \leq 0} d\mathbb{P}_1 + \int \mathbf{1}_{f(\mathbf{x}) > 0} d\mathbb{P}_0$$

Then

$$R_*^\epsilon = \inf_f \sup_{\substack{d(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ d(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \hat{R}(f, \mathbb{P}'_1, \mathbb{P}'_0) = \sup_{\substack{d(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ d(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \inf_f \hat{R}(f, \mathbb{P}'_1, \mathbb{P}'_0) \quad (3)$$

and furthermore equality is attained for some Borel measurable \hat{f} and $\hat{\mathbb{P}}_1, \hat{\mathbb{P}}_0$ with $W_\infty(\hat{\mathbb{P}}_0, \mathbb{P}_0) \leq \epsilon$ and $W_\infty(\hat{\mathbb{P}}_1, \mathbb{P}_1) \leq \epsilon$.

This theorem states that the adversarial learning problem can be thought of as a game between the learner who selects a function f and the adversary who chooses perturbations. Furthermore, the player to pick first does not have an advantage. Next, the fact that equality holds in (3) implies that the optimal function \hat{f} actually minimizes $\hat{R}(\cdot, \hat{\mathbb{P}}_1, \hat{\mathbb{P}}_0)$. Comparing with (1), this statement implies that \hat{f} must minimize the *standard* classification risk with respect to the adversarially perturbed distribution $\hat{\mathbb{P}}_0, \hat{\mathbb{P}}_1$. The goal of this paper is to generalize Theorems 2 and 3 to surrogate risks. Due to the mathematical considerations present in Theorem 3, Lebesgue measure will play a special role in this work. We will denote Lebesgue measure by μ throughout this paper.

3.2 Surrogate Risks

We will now introduce surrogate risks. Typically, minimizing the empirical version of the risk (1) is computationally intractable; in fact, Ben-David et al. (2003) show its NP-hard in general. Algorithms in practice minimize the surrogate

$$R_\phi(f) = \int \phi(f) d\mathbb{P}_1 + \int \phi(-f) d\mathbb{P}_0 \quad (4)$$

as a proxy for R . To define a classifier, one then thresholds f at zero. Notice that because $\text{sign } 0 = -1$, R_ϕ is not a generalization of R ; specifically, $R \neq R_{\ell_{0/1}}$ for $\ell_{0/1} = \mathbf{1}_{f(\mathbf{x}) \leq 0}$. There are many reasonable choices for ϕ —one typically chooses an upper bound on the zero-one loss which is easy to optimize. We make the following assumption on ϕ :

Assumption 1 ϕ is decreasing, non-negative, and lower lower continuous.

In order to find minimizers of R_ϕ , we rewrite the risk as

$$R_\phi(f) = \int \eta(\mathbf{x})\phi(f(\mathbf{x})) + (1 - \eta(\mathbf{x}))\phi(-f(\mathbf{x}))d\mathbb{P}$$

Thus the minimizer of R_ϕ must minimize $C_\phi(\eta, \cdot)$ pointwise, where

$$C_\phi(\eta, \alpha) = \eta\phi(\alpha) + (1 - \eta)\phi(-\alpha) \quad (5)$$

However, a minimizer of $C_\phi(\eta, \cdot)$ may not always exist for some choices of η and ϕ — consider for instance the exponential loss $\phi(\alpha) = e^{-\alpha}$ at $\eta = 1$: Then $C_\phi(1, \alpha) = e^{-\alpha}$ which decreases to 0 as $\alpha \rightarrow \infty$. To avoid this technicality, we assume that all our functions assume values in the extended real numbers $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$.

Similarly, because minimizing the adversarial classification risk is computationally intractable, a common approach is to instead minimize the surrogate risk

$$R_\phi^\epsilon(f) = \int \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(f(\mathbf{x} + \mathbf{h}))d\mathbb{P}_1 + \int \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(-f(\mathbf{x} + \mathbf{h}))d\mathbb{P}_0. \quad (6)$$

We will now discuss some notation particularly useful in the analysis of this risk. The operations of finding the supremums and infimums of functions over ϵ -balls are especially important. For a function f , we define

$$S_\epsilon(f)(\mathbf{x}) = \sup_{\|\mathbf{h}\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}) \quad (7) \quad I_\epsilon(f)(\mathbf{x}) = \inf_{\|\mathbf{h}\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}) \quad (8)$$

Using this notation, one can re-write the risk R_ϕ^ϵ as

$$R_\phi^\epsilon(f) = \int S_\epsilon(\phi \circ f)d\mathbb{P}_1 + \int S_\epsilon(\phi \circ -f)d\mathbb{P}_0 = \int \phi(I_\epsilon(f))d\mathbb{P}_1 + \int \phi \circ (-S_\epsilon(f))d\mathbb{P}_0$$

3.3 Measurability

In order to define the integral in (6), the functions $S_\epsilon(\phi \circ f)$ and $S_\epsilon(\phi \circ -f)$ must be measurable. To illustrate this concern, Pydi and Jog (2021) show that for every ϵ , there is a Borel set C for which the function $S_\epsilon(\mathbf{1}_C)(\mathbf{x})$ is not Borel measurable. In order to define the adversarial surrogate risk in (6), we assume that f is measurable with respect to a σ -algebra called the *universal σ -algebra* $\mathcal{U}(\mathbb{R}^d)$. We use the treatment of (Nishiura, 2010) for our definitions. Let $\mathcal{B}(\mathbb{R}^d)$ be the Borel σ algebra, and denote the completion of $\mathcal{B}(\mathbb{R}^d)$ with respect to a Borel measure ν by $\mathcal{L}_\nu(\mathcal{B}(\mathbb{R}^d))$. Let $\mathcal{M}(\mathbb{R}^d)$ be the set of all Borel measures on \mathbb{R}^d . Then the universal σ -algebra $\mathcal{U}(\mathbb{R}^d)$ is

$$\mathcal{U}(\mathbb{R}^d) = \bigcap_{\nu \in \mathcal{M}(\mathbb{R}^d)} \mathcal{L}_\nu(\mathbb{R}^d).$$

In other words, the universal σ -algebra is the sigma-algebra of sets which are measurable with respect to the completion of every Borel measure. We prove the following theorem in Appendix A.

Theorem 4 *Let f be a universally measurable function. Then the function $S_\epsilon(f)(\mathbf{x})$ defined by*

$$S_\epsilon(f)(\mathbf{x}) = \sup_{\|\mathbf{h}\| \leq \epsilon} f(\mathbf{x} + \mathbf{h})$$

is universally measurable as well.

In the remainder of the paper, unless otherwise noted, we assume that all functions f are universally measurable and that the sigma algebra for measure spaces is $\mathcal{U}(\mathbb{R}^d)$.

4. Existence Results

4.1 Overview of existence results

In this section, we describe our existence results. In short, the surrogate adversarial risk R_ϕ^ϵ always admits a minimizing function with nice properties. To start, we discuss the challenges in proving this theorem. Establishing the existence of minimizers in the non-adversarial scenario is straightforward because minimizers to the standard risks R and R_ϕ are found in a pointwise manner. For instance, for the standard risk R_ϕ , one can construct a minimizer in a pointwise manner by setting $f(\mathbf{x})$ equal to the minimizer of $C_\phi(\eta(\mathbf{x}), \alpha)$ defined in (5). However, this approach fails for the risks (2) and (6) because the loss at each point depends on many nearby points. We will prove a theorem which states that if ϕ is decreasing, then there always exists a minimizer to (6) and furthermore, there is always a minimizer with nice regularity properties. We call this notion of regularity *pseudo-certifiably robustness for functions*.

Definition 5 *We say that a function f is pseudo-certifiably robust if the level sets of f ($\{\mathbf{x}: f(\mathbf{x}) \leq a\}$, $\{\mathbf{x}: f(\mathbf{x}) < a\}$, $\{\mathbf{x}: f(\mathbf{x}) \geq a\}$, $\{\mathbf{x}: f(\mathbf{x}) > a\}$) are all pseudo-certifiably robust for every $a \in \mathbb{R}$.*

In the appendix, we actually prove that to show a function is pseudo-certifiably robust, it suffices to show that just one of the level sets $\{\mathbf{x}: f(\mathbf{x}) \leq a\}$, $\{\mathbf{x}: f(\mathbf{x}) < a\}$, $\{\mathbf{x}: f(\mathbf{x}) \geq a\}$, $\{\mathbf{x}: f(\mathbf{x}) > a\}$ is pseudo-certifiably robust (Lemma 40 of Appendix C).

We now present our first existence theorem.

Theorem 6 *Let ϵ be the perturbation radius, and let f_n be a minimizing sequence of R_ϕ^ϵ .*

- 1) *Then there exists a subsequence f_{n_j} which converges pointwise to a minimizer f of R_ϕ^ϵ*
- 2) *Let f be any minimizer. Then there is a pseudo-certifiably robust minimizer g of R_ϕ^ϵ for which $S_\epsilon(\phi \circ g) = S_\epsilon(\phi \circ f)$ \mathbb{P}_1 -a.e. and $S_\epsilon(\phi \circ -g) = S_\epsilon(\phi \circ -f)$ \mathbb{P}_0 -a.e.*
- 3) *Assume that $\mathbb{P}_1, \mathbb{P}_0$ are absolutely continuous with respect to Lebesgue measure and let g a pseudo-certifiably robust minimizer of R_ϕ^ϵ . Then there exists a Borel-measurable minimizer h to R_ϕ^ϵ which assumes its supremum and infimum over closed ϵ -balls. Furthermore, this minimizer is pseudo-certifiably robust and satisfies $S_\epsilon(\phi \circ h) = S_\epsilon(\phi \circ g)$ \mathbb{P}_1 -a.e., $S_\epsilon(\phi \circ -h) = S_\epsilon(\phi \circ -g)$ \mathbb{P}_0 -a.e.*

This theorem is proved in the next three subsections. Now we discuss the significance of this theorem. First, item 1) states that every minimizing sequence of R_ϕ^ϵ has a convergent subsequence, or in other words, a minimizing sequence of R_ϕ^ϵ is sequentially compact in the topology of pointwise convergence on $\overline{\mathbb{R}}$ -valued functions. Intuitively, this result implies that a sequence of minimizers does not ‘escape to infinity’.

Next, item 2) implies that there always exists a minimizer g with well-behaved level sets. Note specifically that if a is a local minimum or maximum of g , and furthermore there is a point \mathbf{x} with $g(\mathbf{x}) = a$, then there is a ball $B_\epsilon(\mathbf{y})$ that contains \mathbf{x} for which $B_\epsilon(\mathbf{y}) \subset g^{-1}(a)$. Notice that such a regularity result does not hold for minimizers of the standard surrogate risk (4).

In fact, Lewicka and Peres (2020) show that the pseudo-certifiable robustness criterion with respect to the 2-norm is equivalent to a bound on curvature. The following statement is a consequence of their Theorem 1, and a formal derivation is presented in (Frank, 2022).

Theorem 7 *Let A be a set that is pseudo-certifiably robust with respect to the 2-norm with balls of radius ϵ . Then the principal curvatures of the surface ∂A exist μ -a.e. and are bounded by $1/\epsilon$.*

Therefore, 2) of Theorem 6 implies that there always exist minimizers to R_ϕ^ϵ for which the principal curvatures of level sets are bounded by ϵ . Furthermore, intermediate results (Lemmas 12 and 13 in Section 4.3) imply that if g is any function, there is always another pseudo-certifiably robust function h with smaller adversarial ϕ -risk. The function h is g with level sets “smoothed out”.

This statement explains the experimental results of (Moosavi-Dezfooli et al., 2018), who demonstrate that cross-sections of empirical minimizers of the adversarial risk tend to have lower curvature than minimizers of the standard risk.

Consider a twice differentiable function f . Notice that at each point \mathbf{x} , the pseudo-certifiable robustness criterion implies bounds on the eigenvalues of the Hessian $\nabla^2 f$ in the directions *perpendicular* to ∇f , but not in the direction of ∇f . Previous results from image analysis prove that regularization by $\|\nabla f(\mathbf{x})\|_2^2$ tends to reduce the eigenvalues of $\nabla^2 f(\mathbf{x})$. Prior work applied this principle to adversarial robustness: (Ross and Doshi-Velez, 2017; Moosavi-Dezfooli et al., 2018; Yu et al., 2018; Jakubovitz and Giryes, 2018) use the Jacobian regularization $\|\nabla f(\mathbf{x})\|_2^2$ to improve the robustness of neural nets. The observation that pseudo-certifiable robustness does not imply smoothing in the direction of the gradient suggests alternative regularizers. In the field of calculus of variations, it is known that regularization by $\|\nabla f(\mathbf{x})\|_2$ induces bounds on the curvatures in every direction except the direction of the gradient. One application for this regularizer is denoising images (Sapiro, 2001). Perhaps this regularization term could improve adversarial robustness as well.

Even though the above discusses focused on the 2-norm, the results of (Frank, 2022) hint that one may be able to generalize this analysis to other situations. Specifically, Frank (2022) explain that in certain situations, one can define principle curvatures in terms of general strictly convex norms.

Next we discuss the implications of 3). Notice that since we minimize R_ϕ^ϵ over *universally measurable* functions, minimizing over Borel functions could result in a different value of the risk. In fact, the proof of item 1) does not rely on the fact that we are working

in the universal sigma algebra, so minimizers still even when working over just the Borel sigma algebra. In contrast, the proof of item 2) does rely on the fact that minimization is with respect to universally measurable functions. Item 3) implies that when $\mathbb{P}_0, \mathbb{P}_1$ are absolutely continuous with respect to Lebesgue measure, minimizing over universally measurable functions is equivalent to minimizing over Borel measurable functions and additionally the conclusions of Item 2) still hold. Combining Items 2) and 3) results in the following statement: for any minimizer f of R_ϕ^ϵ , there is another minimizer h which is pseudo-certifiably robust, assumes its suprema and infima over closed ϵ -balls, and satisfies $S_\epsilon(\phi \circ f) = S_\epsilon(\phi \circ h)$ \mathbb{P}_1 -a.e., $S_\epsilon(\phi \circ -f) = S_\epsilon(\phi \circ -h)$ \mathbb{P}_0 -a.e. This last criterion suggests that the functions f and h are not very different. The fact that $\phi \circ h, \phi \circ -h$ assume their optima on closed ϵ -balls is central in the proof the minimax theorem (Theorem 26) in the next section.

We next describe the concept of minimal and maximal minimizers inspired by (Bungert et al., 2021). They define minimal and maximal minimizers for the risk R^ϵ in terms of sets— for this risk, a minimizer f is *minimal* if for every other minimizer g with $\mathbf{1}_{\{x: g(x) > 0\}} \leq \mathbf{1}_{\{x: f(x) > 0\}}$ \mathbb{P} -a.e., then in fact $\mathbf{1}_{\{x: g(x) > 0\}} = \mathbf{1}_{\{x: f(x) > 0\}}$. Similarly, a risk is *maximal* if for every other minimizer g with $\mathbf{1}_{\{x: g(x) > 0\}} \geq \mathbf{1}_{\{x: f(x) > 0\}}$ \mathbb{P} -a.e., then in fact $\mathbf{1}_{\{x: g(x) > 0\}} = \mathbf{1}_{\{x: f(x) > 0\}}$ \mathbb{P} -a.e. Proposition 3.23 of (Bungert et al., 2021) shows that there always exist minimal and maximal minimizers to R^ϵ .

Definition 8 *The function f is a minimal minimizer to R_ϕ^ϵ if every other minimizer g with $g \leq f$ \mathbb{P} -a.e. satisfies $f = g$ \mathbb{P} -a.e. Similarly, f is a maximal minimizer if every other minimizer g of R_ϕ^ϵ with $g \geq f$ \mathbb{P} -a.e. satisfies $f = g$ \mathbb{P} -a.e.*

We prove that there always exist maximal and minimal minimizers of R_ϕ^ϵ .

Theorem 9 *There exist minimal and maximal minimizers of R_ϕ^ϵ .*

Again, the proof of this theorem does not rely on the use of the universal sigma-algebra.

We now compare our new theorems to previously published existence theorems for the risk R^ϵ . (Awasthi et al., 2021) proved that there is always a pseudo-certifiably robust minimizer to the adversarial classification risk (2). The authors Pydi and Jog (2021) also proved existence of a minimizer using Choquet capacities. They also realized that a condition similar to Item 2) implies the minimax theorem for classification (Theorem 3). Lastly, Bungert et al. (2021) prove existence of a minimizer to a version of R^ϵ defined with perturbations in open balls by studying the nonlocal perimeter functional. They were also first to realize the existence of minimal and maximal solutions and we adopt their approach in this matter.

4.2 Proof of Item 1) of Theorem 6

We now prove 1) of Theorem 6. To start, we prove a useful lemma about convergent subsequences of functions (Lemma 11). The proof utilizes the following result from (Rockafellar and Wets, 1998) about sequences of sets:

Lemma 10 (Rockafellar and Wets (1998)) *Let A_n be a sequence of sets for which $\limsup A_n \neq \emptyset$. Then there is a subsequence A_{n_k} of A_n for which $\liminf A_{n_k} = \limsup A_{n_k}$ and this set is non-empty.*

Notice that if $\limsup A_n = \liminf A_n = \emptyset$, then for every subsequence A_{n_k} , $\liminf A_{n_k} = \limsup A_{n_k} = \emptyset$. Thus one can always choose a subsequence A_{n_k} of A_n for which the \liminf and the \limsup match. If the \liminf and the \limsup of a sequence of sets A_n match, then we say that A_n is *convergent*. Notably, this result about convergent subsequences of sets implies a similar result for sequences of functions.

Lemma 11 *Let f_n be a sequence of $\overline{\mathbb{R}}$ -valued functions. Then there exists subsequence of f_n which converges pointwise.*

This lemma is likely well-known in the analysis literature, the authors would appreciate a pointer to a reference from the reviewers if they are aware of one.

Proof Let $\{q_j\}_{j=1}^\infty$ be an enumeration of the set of rational numbers. We will construct a sequence of functions g_k for which the set $\{\mathbf{x}: g_k(\mathbf{x}) \geq q_j\}$ is convergent. We proceed via a diagonalization argument. For convenience, denote $V_j(f) = \{\mathbf{x}: f(\mathbf{x}) \geq q_j\}$. Notice that if $q_j < q_k$, then $V_j(f) \supset V_k(f)$.

To start, sequences of functions $\{f_n^j\}$ are constructed inductively. Using Lemma 10, pick a subsequence n_k for which $\liminf_k V_1(f_{n_k}) = \limsup_k V_1(f_{n_k})$ and set $f_k^1 = f_{n_k}$. Next, given the sequence f_n^j , we define f_k^{j+1} inductively: pick a subsequence n_k for which $\liminf_k V_{j+1}(f_{n_k}^j) = \limsup_k V_{j+1}(f_{n_k}^j)$ and set $f_k^{j+1} = f_{n_k}^j$.

Now let $g_n = f_n^n$. Notice that for $n \geq j$, g_n is a subsequence of f_n^j , and thus $\liminf_n V_j(g_n) = \limsup_n V_j(g_n)$. Define U_j as $U_j := \liminf_n V_j(g_n) = \limsup_n V_j(g_n)$. Now define the function f as

$$f(\mathbf{x}) = \sup\{q_j: \mathbf{x} \in U_j\}$$

Then we will show that $\lim_{n \rightarrow \infty} g_n(\mathbf{x}) = f(\mathbf{x})$.

We will show that for $q_j < f(\mathbf{x})$, there is a sufficiently large n for which $g_n(\mathbf{x}) \geq q_j$, and similarly, for any $q_j > f(\mathbf{x})$, there is a sufficiently large n for which $g_n(\mathbf{x}) \leq q_j$.

First consider $q_j < f(\mathbf{x})$. Then the fact that $\mathbf{x} \in U_j = \liminf_n V_j(g_n)$ implies that for sufficiently large n , $\mathbf{x} \in \{\mathbf{z}: g_n(\mathbf{z}) \geq q_j\}$, and thus for all sufficiently large n , $g_n(\mathbf{x}) \geq q_j$.

Now pick $q_j > f(\mathbf{x})$. Then the fact that $\mathbf{x} \notin U_j = \limsup_n V_j(g_n)$ implies that for sufficiently large n , $\mathbf{x} \notin \{\mathbf{z}: g_n(\mathbf{z}) \geq q_j\}$, and thus for all sufficiently large n , $g_n(\mathbf{x}) < q_j$.

Therefore, $g_n(\mathbf{x}) \rightarrow f(\mathbf{x})$ pointwise. ■

We now prove item 1) of Theorem 6.

Proof [Proof of 1) of Theorem 6] Let f_n be a sequence minimizing R_ϕ^ϵ . Now using Lemma 11, pick the subsequence f_{n_k} of f_n so that f_{n_k} that converges pointwise. By passing to another subsequence $f_{n_{k_j}}$, we can assume that $\int S_\epsilon(\phi \circ f_{n_{k_j}}) d\mathbb{P}_1$, $\int S_\epsilon(\phi \circ -f_{n_{k_j}}) d\mathbb{P}_0$ both converge. For convenience, we drop the triple subscript and denote $g_j := f_{n_{k_j}}$. Let f be the pointwise limit of the g_j .

Next, let h_n be any sequence of functions. We observe that $\liminf_{n \rightarrow \infty} S_\epsilon(h_n) \geq S_\epsilon(\liminf_{n \rightarrow \infty} h_n)$:

$$\liminf_{n \rightarrow \infty} \sup_{\|\mathbf{h}\| \leq \epsilon} h_n(\mathbf{x} + \mathbf{h}) = \sup_N \inf_{n \geq N} \sup_{\|\mathbf{h}\| \leq \epsilon} h_n(\mathbf{x} + \mathbf{h}) \geq \sup_{\|\mathbf{h}\| \leq \epsilon} \sup_N \inf_{n \geq N} h_n(\mathbf{x} + \mathbf{h}) = \sup_{\|\mathbf{h}\| \leq \epsilon} \liminf_{n \rightarrow \infty} h_n(\mathbf{x} + \mathbf{h}) \quad (9)$$

Then Fatou's lemma and Equation 9 with h_k as $\phi \circ g_j$ and $\phi \circ -g_j$ imply that

$$\begin{aligned}
 \lim_{j \rightarrow \infty} R_\phi^\epsilon(g_j) &\geq \int \liminf_{j \rightarrow \infty} S_\epsilon(\phi \circ g_j) d\mathbb{P}_1 + \int \liminf_{j \rightarrow \infty} S_\epsilon(\phi \circ -g_j) d\mathbb{P}_0 && \text{(Fatou's Lemma)} \\
 &\geq \int S_\epsilon(\liminf_{j \rightarrow \infty} \phi \circ g_j) d\mathbb{P}_1 + \int S_\epsilon(\liminf_{j \rightarrow \infty} \phi(-g_j)) d\mathbb{P}_0 && \text{(Equation 9)} \\
 &\geq \int S_\epsilon(\phi(f)) d\mathbb{P}_1 + \int S_\epsilon(\phi(-f)) d\mathbb{P}_0 = R_\phi^\epsilon(f) && \text{(lower semi-continuity of } \phi)
 \end{aligned}$$

Therefore, f must be a minimizer. ■

4.3 Proof of Item 2) of Theorem 6

We will now discuss the proof of Item 2) of Theorem 6. This result relies on the fact that applying the S_ϵ, I_ϵ operation in succession decreases the risk R_ϕ^ϵ .

Lemma 12 *Let f be a measurable function and define*

$$L_\epsilon(f)(\mathbf{x}) = S_\epsilon(I_\epsilon(f))$$

$$U_\epsilon(f)(\mathbf{x}) = I_\epsilon(S_\epsilon(f))$$

Then the following hold:

- 1) $L_\epsilon(f), U_\epsilon(f)$ are universally measurable
- 2) $L_\epsilon(f) \leq f$ and $U_\epsilon(f) \geq f$
- 3) $S_\epsilon(L_\epsilon(f))(\mathbf{x}) \leq S_\epsilon(f)(\mathbf{x})$ and $I_\epsilon(L_\epsilon(f)) = I_\epsilon(f)$
- 4) $S_\epsilon(U_\epsilon(f))(\mathbf{x}) = S_\epsilon(f)(\mathbf{x})$ and $I_\epsilon(U_\epsilon(f))(\mathbf{x}) \geq I_\epsilon(f)(\mathbf{x})$
- 5) $R_\phi^\epsilon(L_\epsilon(f)) \leq R_\phi^\epsilon(f)$ and $R_\phi^\epsilon(U_\epsilon(f)) \leq R_\phi^\epsilon(f)$

The operation L_ϵ, U_ϵ are denoted with the letters L, U because of item 2): $L_\epsilon(f)$ is a lower bound on f and $U_\epsilon(f)$ is an upper bound on f .

Notably, this lemma implies that if f is a minimizer of R_ϕ^ϵ , then $U_\epsilon(L_\epsilon(f))$ and $L_\epsilon(U_\epsilon(f))$ are minimizers of R_ϕ^ϵ as well. It turns out that $U_\epsilon(L_\epsilon(f))$ and $L_\epsilon(U_\epsilon(f))$ have pseudo-certifiably robust level sets:

Lemma 13 *Let f be any function. Then $L_\epsilon(U_\epsilon(f))$ and $U_\epsilon(L_\epsilon(f))$ are pseudo-certifiably robust.*

Lemma 13 and 5) of Lemma 12 show that if f is any function, then $R_\phi^\epsilon(L_\epsilon(U_\epsilon(f))) \leq R_\phi^\epsilon(f)$, and furthermore these operations make the level sets of f pseudo-certifiably robust.

These two observations suffice to prove 2) of Theorem 6. We prove Lemma 12 below, and delay the proof of Lemma 13 to Appendix C as the argument is rather technical. Afterwards we will prove 2) of Theorem 6.

Proof [Proof of Lemma 12] To start, that $L_\epsilon(f), U_\epsilon(f)$ are universally measurable follows from applying Theorem 4 twice.

We now show 2). To start, note that we can write the operation L_ϵ as

$$L_\epsilon(f)(\mathbf{x}) = \sup_{\|\mathbf{h}_1\| \leq \epsilon} \inf_{\|\mathbf{h}_2\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2) \quad (10)$$

Thus

$$L_\epsilon(f)(\mathbf{x}) = \sup_{\|\mathbf{h}_2\| \leq \epsilon} \inf_{\|\mathbf{h}_1\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2) \leq \sup_{\|\mathbf{h}_2\| \leq \epsilon} \inf_{\mathbf{h}_1 = -\mathbf{h}_2} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2) = f(\mathbf{x})$$

One can show the result for U_ϵ by writing $U_\epsilon(f) = -L_\epsilon(-f)$.

Next we discuss the proofs of Items 3) and 4). Notice that if $h \leq g$, then $S_\epsilon(h) \leq I_\epsilon(g)$ and $I_\epsilon(h) \leq I_\epsilon(g)$. Thus 2) immediately implies that $S_\epsilon(L_\epsilon(f)) \leq S_\epsilon(f)$ and $I_\epsilon(U_\epsilon(f)) \geq I_\epsilon(f)$.

Next, analogous to (10), the U_ϵ can be written as

$$U_\epsilon(f)(\mathbf{x}) = \inf_{\|\mathbf{h}_2\| \leq \epsilon} \sup_{\|\mathbf{h}_1\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2)$$

Thus

$$\begin{aligned} \inf_{\|\mathbf{h}\| \leq \epsilon} L_\epsilon(f)(\mathbf{x} + \mathbf{h}) &= \inf_{\|\mathbf{h}_3\| \leq \epsilon} \sup_{\|\mathbf{h}_1\| \leq \epsilon} \inf_{\|\mathbf{h}_2\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2 + \mathbf{h}_3) \\ &\geq \inf_{\|\mathbf{h}_3\| \leq \epsilon} \sup_{\mathbf{h}_1 = -\mathbf{h}_3} \inf_{\|\mathbf{h}_2\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2 + \mathbf{h}_3) = \inf_{\|\mathbf{h}\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}) \end{aligned}$$

Similarly,

$$\begin{aligned} \inf_{\|\mathbf{h}\| \leq \epsilon} L_\epsilon(f)(\mathbf{x} + \mathbf{h}) &= \inf_{\|\mathbf{h}_3\| \leq \epsilon} \sup_{\|\mathbf{h}_1\| \leq \epsilon} \inf_{\|\mathbf{h}_2\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2 + \mathbf{h}_3) \\ &\leq \inf_{\|\mathbf{h}_3\| \leq \epsilon} \sup_{\|\mathbf{h}_1\| \leq \epsilon} \inf_{\mathbf{h}_2 = -\mathbf{h}_1} f(\mathbf{x} + \mathbf{h}_1 + \mathbf{h}_2 + \mathbf{h}_3) = \inf_{\|\mathbf{h}_3\| \leq \epsilon} f(\mathbf{x} + \mathbf{h}_3) \end{aligned}$$

We can show the corresponding statement for $U_\epsilon(f)$ in item 4) by writing $U_\epsilon(f) = -L_\epsilon(-f)$.

We will now show 5). Note that because ϕ is decreasing, we can re-write the risk in (6) as

$$R_\phi^\epsilon(f) = \int \phi(I_\epsilon(f)) d\mathbb{P}_0 + \int \phi(-S_\epsilon(f)) d\mathbb{P}_1$$

Since ϕ is decreasing, it follows that increasing $I_\epsilon(f)$ and decreasing $S_\epsilon(f)$ decreases the risk. Thus 3) and 4) imply 5). ■

Now using Lemmas 12 and 13, we prove 2) of Theorem 6.

Proof [Proof of 2) of Theorem 6] Let f be any minimizer. Let $h = L_\epsilon(f)$ and $g = U_\epsilon(h) = U_\epsilon(L_\epsilon(f))$. Then by Lemma 12, $\int S_\epsilon(\phi(h)) d\mathbb{P}_1 = \int S_\epsilon(\phi(f)) d\mathbb{P}_1$ and $\int S_\epsilon(\phi(-h)) d\mathbb{P}_0 \leq \int S_\epsilon(\phi(-f)) d\mathbb{P}_0$. Therefore, $R_\phi^\epsilon(h) \leq R_\phi^\epsilon(f)$. However, because f is a minimizer of R_ϕ^ϵ , the risks of h and f must be equal: $R_\phi^\epsilon(h) = R_\phi^\epsilon(f)$. The relation $\int S_\epsilon(\phi(h)) d\mathbb{P}_1 = \int S_\epsilon(\phi(f)) d\mathbb{P}_1$ implies that $\int S_\epsilon(\phi(-h)) d\mathbb{P}_0 = \int S_\epsilon(\phi(-f)) d\mathbb{P}_0$. However, Lemma 12 implies that $S_\epsilon(\phi(-h(\mathbf{x}))) \leq S_\epsilon(\phi(-f(\mathbf{x})))$, and therefore $S_\epsilon(\phi(-h(\mathbf{x}))) = S_\epsilon(\phi(-f(\mathbf{x})))$ \mathbb{P}_0 -a.e.

The same argument applied h and g demonstrates that g is also a minimizer of R_ϕ^ϵ and furthermore, $S_\epsilon(\phi(-g(\mathbf{x}))) = S_\epsilon(\phi(-h(\mathbf{x})))$ everywhere and $S_\epsilon(\phi(g(\mathbf{x}))) = S_\epsilon(\phi(h(\mathbf{x})))$ \mathbb{P}_1 -a.e.

Because $S_\epsilon(\phi(-h(\mathbf{x}))) = S_\epsilon(\phi(-f(\mathbf{x})))$ \mathbb{P}_0 -a.e. and $S_\epsilon(\phi(h(\mathbf{x}))) = S_\epsilon(\phi(f(\mathbf{x})))$ everywhere, the equalities $S_\epsilon(\phi(-g(\mathbf{x}))) = S_\epsilon(\phi(-f(\mathbf{x})))$ and $S_\epsilon(\phi(g(\mathbf{x}))) = S_\epsilon(\phi(f(\mathbf{x})))$ hold \mathbb{P}_0 -a.e. and \mathbb{P}_1 a.e. respectively. ■

4.4 Proof of Items 3) of Theorem 6

We first provide a brief overview of the proof. To prove Item 3), we need to argue that certain minimizers assume their suprema and infima over closed ϵ -balls. The L_ϵ, U_ϵ produce such functions:

Lemma 14 *For any function g , $U_\epsilon(L_\epsilon(g))$ assumes its supremum on every closed ball radius ϵ and $L_\epsilon(U_\epsilon(g))$ assumes its infimum on every closed ball radius ϵ .*

The proof of Item 3) hinges on this observation. We prove this lemma in Appendix B. However, this lemma does not produce a minimizer that assumes its supremum and infimum over ϵ -balls *simultaneously*. On the other hand, the L_ϵ, U_ϵ operations preserve this property for functions:

Lemma 15 *Let f be a lower semi-continuous function. Then $I_\epsilon(f)$ and $S_\epsilon(f)$ are both lower semicontinuous.*

Therefore, if f is lower semi-continuous, $U_\epsilon(L_\epsilon(f))$ assumes both its infimum and supremum over closed ϵ -balls. Hence it remains to find a lower semi-continuous minimizer of R_ϕ^ϵ . We use properties of porous sets to show the following result:

Lemma 16 *Let g be a pseudo-certifiably robust minimizer, and define $h = \liminf_{\mathbf{y} \rightarrow \mathbf{x}} g(\mathbf{y})$. Then if $\mathbb{P}_0, \mathbb{P}_1$ are absolutely continuous with respect to Lebesgue measure, then h is a minimizer as well.*

Lastly, we argue that if f is a pseudo-certifiably robust function, then $\liminf_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ is as well.

Lemma 17 *Let f be a pseudo-certifiably robust function. Then $\liminf_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ is also pseudo-certifiably robust.*

This result is proved in Appendix D.

Now using these lemmas, we prove Item 3) of Theorem 6. Lemmas 15 and 16 are proved later in this section.

Proof [Proof of 3) of Theorem 6] Define $h(\mathbf{x}) = \liminf_{\mathbf{y} \rightarrow \mathbf{x}} g(\mathbf{y})$. Then by Lemma 16, h is also a minimizer of R_ϕ^ϵ . Next, Lemma 12 implies that $U_\epsilon(L_\epsilon(h))$ is also a minimizer of R_ϕ^ϵ and Lemma 17 implies that $U_\epsilon(L_\epsilon(h))$ is pseudo-certifiably robust. Next, Lemma 14 implies that $U_\epsilon(L_\epsilon(h))$ assumes its suprema over closed ϵ -balls and Lemma 15 implies that $U_\epsilon(L_\epsilon(h))$ is lower semi-continuous, so it assumes its infima over closed ϵ -balls. The function




Figure 1: The sets A , A^ϵ and $A^{-\epsilon}$ for perturbations in the 2-norm. Figure credit: Awasthi et al. (2021)

$U_\epsilon(L_\epsilon(h))$ is Borel measurable because it is lower semi-continuous. We will now discuss the proof of the fact that $S_\epsilon(\phi \circ h) = S_\epsilon(\phi \circ g)$ \mathbb{P}_1 -a.e., $S_\epsilon(\phi \circ -h) = S_\epsilon(\phi \circ -g)$ \mathbb{P}_0 -a.e. Notice that $h \leq g$, so that $S_\epsilon(\phi \circ -h) \leq S_\epsilon(\phi \circ -g)$. The proof now follows by the argument in Item 2) that $S_\epsilon(\phi \circ f) = S_\epsilon(\phi \circ g)$ \mathbb{P}_1 -a.e., $S_\epsilon(\phi \circ -f) = S_\epsilon(\phi \circ -g)$ \mathbb{P}_0 -a.e. ■

4.4.1 THE $^\epsilon$ AND $^{-\epsilon}$ OPERATIONS

Before proving Lemmas 15 and 16, we introduce some notation that elucidates the relationship between $S_\epsilon(f)$, $L_\epsilon(f)$, and level sets of f .

For a set A , we define A^ϵ as the sets of points that can fall in the set A after an additive perturbation of size at most ϵ . Notably, the set A^ϵ can be expressed in terms of a Minkowski sum \oplus with the closed ϵ -ball $\overline{B_\epsilon(\mathbf{0})}$: $A^\epsilon = \{\mathbf{x} : \exists \mathbf{h} \text{ with } \|\mathbf{h}\| \leq \epsilon \text{ and } \mathbf{x} + \mathbf{h} \in A\} = A \oplus \overline{B_\epsilon(\mathbf{0})}$. As illustrated in Figure 1, the operation $^\epsilon$ thickens the boundary of a set by ϵ . We will now introduce an operation for “removing” ϵ from the boundary of A which denote by $A^{-\epsilon}$. Formally, we define this operation as $A^{-\epsilon} = ((A^C)^\epsilon)^C$. $A^{-\epsilon}$ is the set of points which cannot be perturbed outside of A by perturbations in $\overline{B_\epsilon(\mathbf{0})}$. The operations $^\epsilon, ^{-\epsilon}$ can be thought of as the “set versions” of the S_ϵ and I_ϵ operations—in fact $S_\epsilon(\mathbf{1}_A) = \mathbf{1}_{A^\epsilon}$ and $I_\epsilon(\mathbf{1}_A) = \mathbf{1}_{A^{-\epsilon}}$.

Awasthi et al. (2021) provide an alternative interpretation of the $^\epsilon$ and $^{-\epsilon}$ operations:

Lemma 18 *Alternative characterizations of the $^\epsilon, ^{-\epsilon}$ operations are*

$$A^\epsilon = \{\mathbf{x} \in \mathbb{R}^d : \overline{B_\epsilon(\mathbf{x})} \cap A \neq \emptyset\}$$

$$A^{-\epsilon} = \{\mathbf{x} : \overline{B_\epsilon(\mathbf{x})} \subset A\}$$

Due to this lemma, level sets of $I_\epsilon(f)$ and $S_\epsilon(f)$ can be expressed in terms of the level sets of f and the $^\epsilon, ^{-\epsilon}$ operations. Notice that $S_\epsilon(f) > a$ iff there is a $\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}$ for which $f(\mathbf{y}) > a$. In other words, $S_\epsilon(f)(\mathbf{x}) > a$ iff $\overline{B_\epsilon(\mathbf{x})}$ intersects $f^{-1}((a, \infty])$, which is equivalent to $\mathbf{x} \in f^{-1}((a, \infty])^\epsilon$ by Lemma 18. Therefore,

$$(S_\epsilon(f))^{-1}((a, \infty]) = f^{-1}((a, \infty])^\epsilon. \quad (11)$$

Similarly, $S_\epsilon(f)(\mathbf{x}) \leq a$ iff for every $\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}$, $f(\mathbf{y}) \leq a$. In other words, $S_\epsilon(f)(\mathbf{x}) \leq a$ iff $B_\epsilon(\mathbf{x}) \subset f^{-1}([-\infty, a])$, which is equivalent to $\mathbf{x} \in f^{-1}([-\infty, a])^{-\epsilon}$. Therefore,

$$S_\epsilon(f)^{-1}([-\infty, a]) = f^{-1}([-\infty, a])^{-\epsilon} \quad (12)$$

Awasthi et al. (2021) studied topological properties of the sets $A^\epsilon, A^{-\epsilon}$ in terms of the topological properties of A . We use these properties together with (12) to understand the semi-continuity of $S_\epsilon(f)$ and $I_\epsilon(f)$. Specifically they show the following two lemmas:

Lemma 19 *Let A be any set. Then $\overline{A^\epsilon} = \overline{A}^\epsilon$.*

Lemma 20 *For any set A , ∂A^ϵ has Lebesgue measure zero.*

Lemma 19 actually implies that the $^\epsilon, ^{-\epsilon}$ operations preserve closed sets:

Lemma 21 *If A is closed, then A^ϵ and $A^{-\epsilon}$ are both closed.*

Proof Let A be a closed set. Lemma 19 implies that $\overline{A^\epsilon} = \overline{A}^\epsilon = A^\epsilon$, and therefore the set A^ϵ must be closed.

We will now show that $A^{-\epsilon}$ is also closed. First, because A is closed, A^C is open. Now $(A^C)^\epsilon$ can be written as a union of open sets:

$$(A^C)^\epsilon = \bigcup_{\mathbf{x} \in \overline{B_\epsilon(\mathbf{x})}} \{\mathbf{x}\} \oplus A^C.$$

Therefore, $(A^C)^\epsilon$ is open and $A^{-\epsilon} = ((A^C)^\epsilon)^C$ is closed. ■

4.4.2 PROOF OF LEMMAS 15 AND 16

We start with proving Lemma 15

Proof [Proof of Lemma 15] Let f be lower-semicontinuous. Then the sublevel sets $f^{-1}([-\infty, a])$ are closed. Then by (12) and Lemma 21, the sublevel set $S_\epsilon(f)^{-1}([-\infty, a])$ is closed. Therefore $S_\epsilon(f)$ is lower semi-continuous.

Now note that because f is lower-semicontinuous, f assumes its infimum over every closed ϵ ball. Therefore, $I_\epsilon(f)(\mathbf{x}) \leq a$ iff there is a $\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}$ with $f(\mathbf{y}) = a$, or in other words, $\overline{B_\epsilon(\mathbf{x})}$ intersects $f([-\infty, a])$. Hence $\mathbf{x} \in (I_\epsilon(f))^{-1}([-\infty, a])$ iff $\mathbf{x} \in f([-\infty, a])^\epsilon$ and therefore $(I_\epsilon(f))^{-1}([-\infty, a]) = f([-\infty, a])^\epsilon$. This set is then closed by Lemma 21, and therefore $I_\epsilon(f)$ is lower semi-continuous. ■

Note that the relation $(I_\epsilon(f))^{-1}([-\infty, a]) = f([-\infty, a])^\epsilon$ is *false* in general.

Before proving Lemma 16, characterize the level sets of the lower envelope $\limsup_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ in terms of the level sets of f .

Lemma 22 *Let F be a function whose level sets are pseudo-certifiably robust and let $G(\mathbf{x}) = \limsup_{\mathbf{y} \rightarrow \mathbf{x}} F(\mathbf{y})$. Then*

$$G^{-1}([a, \infty]) = \overline{F^{-1}([a, \infty])} \quad (13)$$

This lemma is proved in Appendix D. In the proof of Lemma 16, we apply this lemma with F as $\phi \circ f$ and $\phi \circ -f$. In order to apply Lemma 22, we will need to show that $\phi \circ f$ and $\phi \circ -f$ are pseudo-certifiably robust as well.

Lemma 23 *Let $\phi: \bar{\mathbb{R}} \rightarrow \bar{\mathbb{R}}$ be a decreasing function and let $f: \mathbb{R}^d \rightarrow \bar{\mathbb{R}}$ be a pseudo-certifiably robust function. Then $\phi \circ f$ is pseudo-certifiably robust as well.*

We prove this lemma in Appendix C.

Now using these two results, we prove Lemma 16:

Proof [Proof of Lemma 16] First note that because $h \leq g$,

$$\int S_\epsilon(\phi \circ -g) d\mathbb{P}_0 \geq \int S_\epsilon(\phi \circ -h) d\mathbb{P}_0$$

We will now show that $\int S_\epsilon(\phi \circ g) d\mathbb{P}_1 = \int S_\epsilon(\phi \circ h) d\mathbb{P}_1$ and this relation will complete the proof. We use a well-known formula from probability that expresses the expectation of a non-negative integrable function in terms of its level sets:

$$\int f d\mathbb{Q} = \int \mathbb{Q}(f(\mathbf{x}) > a) da \tag{14}$$

The integral in terms of a above is with respect to Lebesgue measure.

One can write

$$\begin{aligned} & \int S_\epsilon(\phi(g)) d\mathbb{P}_1 \\ &= \int \mathbb{P}_1(S_\epsilon(\phi(g)) > a) da && \text{(Equation 14)} \\ &= \int \mathbb{P}_1((\phi(g))^{-1}((a, \infty])^\epsilon) da && \text{(Equation 11)} \\ &= \int \mathbb{P}_1(\overline{(\phi(g))^{-1}((a, \infty])^\epsilon}) da && \text{(Lemma 20)} \\ &= \int \mathbb{P}_1(\overline{(\phi(g))^{-1}((a, \infty])^\epsilon}) da && \text{(Lemma 19)} \\ &= \int \mathbb{P}_1((\limsup \phi(g))^{-1}((a, \infty])^\epsilon) da && \text{(Lemma 22)} \\ &= \int \mathbb{P}_1(S_\epsilon(\limsup \phi(g))^{-1}((a, \infty])) da && \text{(Equation 11)} \\ &= \int \mathbb{P}_1(S_\epsilon(\phi(\liminf g))^{-1}((a, \infty])) da && (\phi \text{ decreasing}) \\ &= \int S_\epsilon(\phi(\liminf g)) d\mathbb{P}_1 \\ &= \int S_\epsilon(\phi(h)) d\mathbb{P}_1 \end{aligned}$$

■

4.5 Proof of Theorem 9

To start, we outline the proof of Theorem 9. Bungert et al. (2021) show that there exist minimal and maximal minimizers to the adversarial classification risk in Proposition 3.23. Our theorem simply generalizes this result to $\overline{\mathbb{R}}$ -valued functions, and we apply the same proof strategy as (Bungert et al., 2021). We first show that the set of minimizers of R_ϕ^ϵ is closed under monotonic limits. Next, let ψ be the sigmoid function and define a risk

$$\overline{R}(f) = \int \psi(f(\mathbf{x})) d\mathbb{P}.$$

Thus, for any sequence of minimizers f_n which maximize \overline{R} , the limit $\sup_n f_n$ must a minimizer as well. Lastly, we argue that $\sup f_n$ should be the largest minimizer \mathbb{P} -a.e.

Theorem 9 follows from the fact that if f, g are two minimizers, so are $\max(f, g)$ and $\min(f, g)$, which is analogous to Proposition 3.21 of (Bungert et al., 2021). To show this result, Bungert et al. (2021) prove that $R^\epsilon(\max(f, g)) + R^\epsilon(\min(f, g)) \leq R^\epsilon(f) + R^\epsilon(g)$. We prove an analogous statement for the risk R_ϕ^ϵ .

Lemma 24 *Let f_1 and f_2 be two universally measurable functions. Then*

$$R_\phi^\epsilon(\max(f_1, f_2)) + R_\phi^\epsilon(\min(f_1, f_2)) \leq R_\phi^\epsilon(f_1) + R_\phi^\epsilon(f_2) \quad (15)$$

Proof Notice that

$$\begin{aligned} & \sup_{\|\mathbf{h}\| \leq \epsilon} \phi \left(\max_{i \in \{1,2\}} f_i(\mathbf{x} + \mathbf{h}) \right) + \sup_{\|\mathbf{h}\| \leq \epsilon} \phi \left(\min_{i \in \{1,2\}} f_i(\mathbf{x} + \mathbf{h}) \right) \\ &= \sup_{\|\mathbf{h}\| \leq \epsilon} \min_{i \in \{1,2\}} \phi(f_i(\mathbf{x} + \mathbf{h})) + \sup_{\|\mathbf{h}\| \leq \epsilon} \max_{i \in \{1,2\}} \phi(f_i(\mathbf{x} + \mathbf{h})) \quad (\phi \text{ decreasing}) \\ &\leq \min_{i \in \{1,2\}} \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(f_i(\mathbf{x} + \mathbf{h})) + \max_{i \in \{1,2\}} \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(f_i(\mathbf{x} + \mathbf{h})) \quad (\text{swapping min and sup}) \\ &= \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(f_1(\mathbf{x} + \mathbf{h})) + \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(f_2(\mathbf{x} + \mathbf{h})) \end{aligned} \quad (16)$$

Similar reasoning implies

$$\sup_{\|\mathbf{h}\| \leq \epsilon} \phi \left(\max_{i \in \{1,2\}} -f_i(\mathbf{x} + \mathbf{h}) \right) + \sup_{\|\mathbf{h}\| \leq \epsilon} \phi \left(\min_{i \in \{1,2\}} -f_i(\mathbf{x} + \mathbf{h}) \right) \leq \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(-f_1(\mathbf{x} + \mathbf{h})) + \sup_{\|\mathbf{h}\| \leq \epsilon} \phi(-f_2(\mathbf{x} + \mathbf{h})) \quad (17)$$

Integrating (16) with respect to \mathbb{P}_1 and (17) with respect to \mathbb{P}_0 and subsequently summing the results produces (15). ■

This lemma implies that if f, g are two minimizers, so are $\max(f, g)$ and $\min(f, g)$:

Lemma 25 *If f, g are two minimizers of R_ϕ^ϵ , then so are $\min(f, g)$ and $\max(f, g)$.*

Proof If $R_{\phi,*}^\epsilon = R_\phi^\epsilon(f) = R_\phi^\epsilon(g)$, then Lemma 24 implies

$$R_\phi^\epsilon(\max(f, g)) + R_\phi^\epsilon(\min(f, g)) \leq 2R_{\phi,*}^\epsilon \quad (18)$$

Thus we must have that either $R_\phi^\epsilon(\max(f, g)) \leq R_{\phi,*}^\epsilon$ or $R_\phi^\epsilon(\min(f, g)) \leq R_{\phi,*}^\epsilon$, which implies that either $R_\phi^\epsilon(\max(f, g)) = R_{\phi,*}^\epsilon$ or $R_\phi^\epsilon(\min(f, g)) = R_{\phi,*}^\epsilon$. WLOG assume that $R_\phi^\epsilon(\max(f, g)) = R_{\phi,*}^\epsilon$. Then (18) is equivalent to

$$R_{\phi,*}^\epsilon + R_\phi^\epsilon(\min(f, g)) \leq 2R_{\phi,*}^\epsilon$$

and therefore $R_\phi^\epsilon(\min(f, g))$ also equals $R_{\phi,*}^\epsilon$. \blacksquare

Lemma 25 now proves Theorem 9.

Proof [Proof of Theorem 9] Let S be the set of all minimizers of R_ϕ^ϵ . We will start by showing that S is closed under increasing limits. Let f_n be an increasing sequence in S , and define

$$f(\mathbf{x}) = \lim_{n \rightarrow \infty} f_n(\mathbf{x})$$

Then (9) implies that $R_{\phi,*}^\epsilon = R_\phi^\epsilon(f)$:

$$\begin{aligned} R_{\phi,*}^\epsilon &= \lim_{n \rightarrow \infty} R_\phi^\epsilon(f_n) = \liminf_{n \rightarrow \infty} R_\phi^\epsilon(f_n) \\ &\geq \int \liminf_{n \rightarrow \infty} S_\epsilon(\phi(f_n)) d\mathbb{P}_1 + \int \liminf_{n \rightarrow \infty} S_\epsilon(\phi(-f_n)) d\mathbb{P}_0 && \text{(Fatou's Lemma)} \\ &\geq \int S_\epsilon(\liminf_{n \rightarrow \infty} \phi(f_n)) d\mathbb{P}_1 + \int S_\epsilon(\liminf_{n \rightarrow \infty} \phi(-f_n)) d\mathbb{P}_0 && \text{(Equation 9)} \\ &= R_\phi^\epsilon(f) \end{aligned}$$

Therefore, f is also a minimizer of R_ϕ^ϵ .

Now let ψ be the sigmoid function $\psi(\alpha) = 1/(1 + e^{-\alpha})$ and define a risk

$$\bar{R}(f) = \int \psi(f(\mathbf{x})) d\mathbb{P}.$$

Notice that for any function taking values in \mathbb{R}^* , $\bar{R}(f)$ is always finite. Furthermore, since ψ is strictly increasing, if $f(\mathbf{x}) > g(\mathbf{x})$, then $\psi(f(\mathbf{x})) > \psi(g(\mathbf{x}))$. Let f_n be a sequence in S that maximizes \bar{R} . Then Lemma 25 implies that if we define $g_n = \max_{1 \leq k \leq n} f_k$ then each g_n is also a minimizer of R_ϕ^ϵ , and the fact that S is closed under monotone limits implies that $f^M := \limsup_{n \rightarrow \infty} g_n$ is also a minimizer of R_ϕ^ϵ . Furthermore, because $g_n \geq f_n$, the sequence g_n also maximizes R_ϕ^ϵ . Therefore,

$$\sup_{f \in S} \bar{R}(f) = \lim_{n \rightarrow \infty} \bar{R}(g_n) = \bar{R}(\limsup g_n) = \bar{R}(f^M) \quad (19)$$

The second-to-last equality follows from the monotone convergence theorem and the fact that the sigmoid function ψ is increasing.

We will now show that f^M is in fact a maximal solution \mathbb{P} -a.e. For contradiction, assume that there is a function $g \in S$ for which $g > f^M$ on a set of positive measure. Then Lemma 25 implies that $\max(g, f^M)$ is also a minimizer of R_ϕ^ϵ ; however, $\bar{R}(\max(g, f^M)) > \bar{R}(f^M)$ which contradicts (19).

One can prove the existence of minimal solutions by an analogous argument. \blacksquare

5. A Minimax Theorem

In this section, we introduce a minimax theorem for surrogate risks which suggests a new interpretation of the adversarial training procedure. This result extends Theorem 3 to surrogate risks.

Theorem 26 *Let \mathcal{F} be the set of all universally measurable functions on \mathbb{R}^d and assume that $\mathbb{P}_0, \mathbb{P}_1$ are absolutely continuous with respect to Lebesgue measure. Let*

$$\tilde{R}_\phi(f, \mathbb{P}_1, \mathbb{P}_0) = \int \phi(f(\mathbf{x}))d\mathbb{P}_1 + \int \phi(-f(\mathbf{x}))d\mathbb{P}_0$$

Then

$$R_\phi^{\epsilon,*} = \inf_{f \in \mathcal{F}} \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0) = \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \inf_{f \in \mathcal{F}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0). \quad (20)$$

Let f^* be a minimizer of R_ϕ^ϵ satisfying (3) of Theorem 6. Then equality is attained for this f^* and $\mathbb{P}_1^*, \mathbb{P}_0^*$ with $W_\infty(\mathbb{P}_i^*, \mathbb{P}_i) \leq \epsilon$. Furthermore, the measures \mathbb{P}_i^* are given by $\mathbb{P}_i^* = \mathbb{P}_i \circ \varphi_i^{-1}$, where φ_i is a universally measurable function with $\|\varphi_i(\mathbf{x}) - \mathbf{x}\| \leq \epsilon$ and $\inf_{\|\mathbf{y} - \mathbf{x}\| \leq \epsilon} f(\mathbf{y}) = f(\varphi_1(\mathbf{x}))$ and $\sup_{\|\mathbf{y} - \mathbf{x}\| \leq \epsilon} f(\mathbf{y}) = f(\varphi_0(\mathbf{x}))$.

As highlighted by Pydi and Jog (2021), such a minimax result states that minimizing R_ϕ^ϵ can be interpreted as a game between two players: a learner who chooses a function f , and an adversary which chooses perturbations. Theorem 26 states that picking first does not give either player an advantage.

The fact that the optimum is assumed at $\mathbb{P}_1^*, \mathbb{P}_0^*$ with $\mathbb{P}_i^* = \mathbb{P}_i \circ \varphi_i^{-1}$ implies that just like there is a optimal function f^* that minimizes the risk R_ϕ^ϵ , there are *optimal adversarial perturbations*. The measures $\mathbb{P}_0^*, \mathbb{P}_1^*$ give the distribution with respect to these optimal adversarial perturbations and the minimizer f^* of R_ϕ^ϵ must in fact minimize $\tilde{R}_\phi(\cdot, \mathbb{P}_1^*, \mathbb{P}_0^*)$. Notice that $\tilde{R}_\phi(\cdot, \mathbb{P}_1^*, \mathbb{P}_0^*)$ is simply the standard surrogate risk R_ϕ of (4) with respect to the measures $\mathbb{P}_0^*, \mathbb{P}_1^*$. Therefore, f^* minimizes the *standard* surrogate risk with respect to the distribution of optimal adversarial perturbations.

This interpretation of our minimax theorem sheds light on the phenomenon of transfer attacks on adversarially trained neural nets. This theorem suggests that adversarial examples are a property of the data distribution rather than a specific model, which could explain why attacks transfer to other models. Because neural networks are highly expressive model classes, one would hope that these classes can achieve adversarial error close to $R_{\phi,*}^\epsilon$. If g is a neural net with $R_\phi^\epsilon(g) \approx R_\phi^\epsilon(f^*) = R_{\phi,*}^\epsilon$, one would expect that adversarial examples for f^* would be adversarial examples for g as well.

5.1 Proof of Theorem 26

Item 3) of Theorem 6 is essential for the proof of Theorem 26. With 3) of Theorem 6 in hand, the proof of Theorem 26 reduces to the argument of Pydi and Jog (2021) for Theorem 3.

To start, we introduce a theorem for measurable selections which will be central to the proofs in this Section. This theorem is a consequence of Proposition 7.50 of (Bertsekas and Shreve, 1996).

Theorem 27 *Let X, Y be Borel sets and assume that $D \subset X \times Y$ is also Borel. Let D_x denote*

$$D_x = \{y: (x, y) \in D\}$$

and

$$\text{Proj}_X(D) := \{x: (x, y) \in D\}$$

Let f be a function mapping D to $\overline{\mathbb{R}}$, $f: D \rightarrow \overline{\mathbb{R}}$ and define

$$f^*(x) = \inf_{y \in D_x} f(x, y)$$

Let

$$I = \{x \in \text{Proj}_X(D) \mid \text{for some } y_x \in D_x, f(x, y_x) = f^*(x)\}$$

Then for any $\delta > 0$, there is a universally measurable $\varphi: \text{Proj}_X(D) \rightarrow Y$ for which $f(x, \varphi(x)) = f^*(x)$ for all $x \in I$ and

$$f(x, \varphi(x)) \leq \begin{cases} f(x) + \delta & \text{if } x \notin I, f^*(x) > -\infty \\ -1/\delta & \text{if } x \notin I, f^*(x) = -\infty \end{cases}$$

Note that by applying this theorem to $-f$, we can draw the same conclusions for supremums. The following lemma relates the integral of a supremum to the supremum over measures of an integral. This result generalizes Lemma 5.1 of (Pydi and Jog, 2021).

Lemma 28 *Let \mathbb{Q} be an arbitrary Borel measure and let F be a non-negative Borel measurable function which assumes its supremum on every closed ϵ -ball. Then*

$$\int \sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \epsilon} F(\mathbf{x}') d\mathbb{Q} = \sup_{W_\infty(\mathbb{Q}', \mathbb{Q}) \leq \epsilon} \int F f(\mathbf{x}) d\mathbb{Q}' \quad (21)$$

Furthermore, the supremum on the right hand side is assumed by a measure given by $\mathbb{Q} \circ \varphi^{-1}$ for a universally measurable map φ . This map φ satisfies

$$\sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \epsilon} F(\mathbf{x}') = F(\varphi(\mathbf{x}))$$

Proof Pydi and Jog (2021) proved that for any measure \mathbb{Q}' with $W_\infty(\mathbb{Q}, \mathbb{Q}') \leq \epsilon$,

$$\mathbb{Q}'(A) \leq \mathbb{Q}(A^\epsilon) \quad (22)$$

We reproduce their proof here for completeness. If $W_\infty(\mathbb{Q}, \mathbb{Q}') \leq \epsilon$, then there is a coupling $\lambda \in \Pi(\mathbb{Q}', \mathbb{Q})$ for which if (\mathbf{x}, \mathbf{y}) is sampled according to λ , then $\|\mathbf{x} - \mathbf{y}\| \leq \epsilon$ a.e. Thus for any set A , $\lambda(A^C \times A^\epsilon) = 0$. Therefore,

$$\mathbb{Q}'(A) = \lambda(A \times \mathbb{R}^d) = \lambda(A \times A^\epsilon) \leq \lambda(\mathbb{R}^d \times A^\epsilon) = \mathbb{Q}(A^\epsilon)$$

Next, by (22) and Equation 11,

$$\int F d\mathbb{Q}' = \int \mathbb{Q}'(\{\mathbf{x}: F(\mathbf{x}) > a\}) da \leq \int \mathbb{Q}(\{\mathbf{x}: F(\mathbf{x}) > a\}^\epsilon) da = \int \sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \epsilon} F(\mathbf{x}') d\mathbb{Q}$$

and hence

$$\sup_{W_\infty(\mathbb{Q}', \mathbb{Q}) \leq \epsilon} \int F d\mathbb{Q}' \leq \int \sup_{\|\mathbf{x} - \mathbf{x}'\| \leq \epsilon} F(\mathbf{x}') d\mathbb{Q} \quad (23)$$

We will now show the opposite inequality. Theorem 27 implies that there is a universally measurable $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ for which $\|\mathbf{x} - \varphi(\mathbf{x})\| \leq \epsilon$ and $\sup_{\|\mathbf{x} - \mathbf{x}'\| \leq \epsilon} F(\mathbf{x}') = F(\varphi(\mathbf{x}))$. Thus, because $W_\infty(\mathbb{Q} \circ \varphi^{-1}, \mathbb{Q}) \leq \epsilon$,

$$\int \sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \epsilon} F(\mathbf{x}') d\mathbb{Q} = \int F(\varphi(\mathbf{x})) d\mathbb{Q} = \int F d\mathbb{Q} \circ \varphi^{-1} \leq \sup_{W_\infty(\mathbb{Q}', \mathbb{Q}) \leq \epsilon} \int \phi(f(\mathbf{x})) d\mathbb{Q}'$$

Therefore, we have shown that (23) is actually an equality and furthermore equality holds at $\mathbb{Q}' = \mathbb{Q} \circ \varphi^{-1}$. ■

Finally, using the measurable selection theorem and Lemma 28, we prove Theorem 26.

Proof [Proof of Theorem 26]

To start, Theorem 6 proves that there exists a Borel measurable minimizer f to R_ϕ^ϵ which achieves its infimum and supremum over closed ϵ -balls. Therefore, we can assume that $\mathbb{P}_0, \mathbb{P}_1$ are Borel measures and we can restrict our search for the optimal f to the Borel σ -algebra. With this fact in mind, the first equality of (20) is just Lemma 28. The inequality

$$\inf_{f \in \mathcal{F}} \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0) \geq \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \inf_{f \in \mathcal{F}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0) \quad (24)$$

follows from switching the infimum and supremum. Next, fix \mathbb{P}'_0 and \mathbb{P}'_1 with $W_\infty(\mathbb{P}'_i, \mathbb{P}_i) \leq \epsilon$. Theorem 6 proves that there exists a minimizer to R_ϕ^ϵ with distribution given by \mathbb{P}'_0 and \mathbb{P}'_1 , so in fact there is an $f_{\mathbb{P}'_0, \mathbb{P}'_1}^*$ for which

$$\tilde{R}_\phi(f_{\mathbb{P}'_1, \mathbb{P}'_0}^*, \mathbb{P}'_0, \mathbb{P}'_1) \leq \inf_{f \in \mathcal{F}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0)$$

Thus if we take supremums over $\mathbb{P}'_0, \mathbb{P}'_1$ on both sides of this equation,

$$\sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \tilde{R}_\phi(f_{\mathbb{P}'_0, \mathbb{P}'_1}^*, \mathbb{P}'_1, \mathbb{P}'_0) \leq \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \inf_{f \in \mathcal{F}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0)$$

now we can replace the $f_{\mathbb{P}'_0, \mathbb{P}'_1}^*$ on the left with an infimum over functions:

$$R_\phi^{\epsilon, *} = \inf_{f \in \mathcal{F}} \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0) \leq \sup_{\substack{W_\infty(\mathbb{P}'_0, \mathbb{P}_0) \leq \epsilon \\ W_\infty(\mathbb{P}'_1, \mathbb{P}_1) \leq \epsilon}} \inf_{f \in \mathcal{F}} \tilde{R}_\phi(f, \mathbb{P}'_1, \mathbb{P}'_0)$$

This inequality combined with (24) produces (20).

Lastly, by Lemma 28 and item 3) of Theorem 6, the infimum and supremums are assumed. Furthermore, Lemma 28 implies that the supremum is assumed by \mathbb{P}_i^* with $\mathbb{P}_i^* = \mathbb{P}_i \circ \varphi_i^{-1}$ for which φ_i are universally measurable with $\|\varphi_i(\mathbf{x}) - \mathbf{x}\| \leq \epsilon$ and $\inf_{\|\mathbf{y} - \mathbf{x}\| \leq \epsilon} f(\mathbf{y}) = f(\varphi_1(\mathbf{x}))$ and $\sup_{\|\mathbf{y} - \mathbf{x}\| \leq \epsilon} f(\mathbf{y}) = f(\varphi_0(\mathbf{x}))$. ■

6. Conclusion

We initiated the study of minimizers and minimax relations for adversarial surrogate risks. Specifically, we proved that there always exists a minimizer to the adversarial surrogate risk when perturbing in a closed ϵ -ball, and furthermore, there always exists minimizers which are well-behaved. Our minimax theorem provides an interpretation of the adversarial learning problem as a game between two players. Our theory helps explain the phenomenon of transfer attacks and the reduced curvature of robust classifiers, as seen in experiments. We hope the insights gained from this research will assist in the development of algorithms for training classifiers robust to adversarial perturbations.

References

- P. Awasthi, N. S. Frank, and M. Mohri. On the existence of the adversarial bayes classifier (extended version). *arxiv*, 2021.
- P. L. Bartlett, M. I. Jordan, and J. D. McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473), 2006.
- G. Bellettini. Anisotropic and crystalline mean curvature flow. *MSRI publications*, 2004.
- S. Ben-David, N. Eiron, and P. M. Long. On the difficulty of approximately maximizing agreements. *Journal of Computer System Sciences*, 2003.
- D. P. Bertsekas and S. E. Shreve. *Stochastic Optimal Control: The Discrete-Time Case*. Athena Scientific, 1996.
- A. N. Bhagoji, D. Cullina, and P. Mittal. Lower bounds on adversarial robustness from optimal transport. In *Advances in Neural Information Processing Systems*, pages 7498–7510, 2019.
- L. Bungert, N. G. Trillos, and R. Murray. The geometry of adversarial training in binary classification. *arxiv*, 2021.
- A. Cesaroni and N. Matteo. Isoperimetric problems for a nonlocal perimeter of minkowski type. *arXiv preprint arXiv:1709.05284*, 2017.
- A. Cesaroni, D. Serena, and N. Matteo. Minimizers for nonlocal perimeters of minkowski type. *SpringerLink*, 2018.
- A. Chambolle, M. Morini, and M. Ponsiglione. A non-local mean curvature flow and its semi-implicit time-discrete approximation. *SIAM Journal on Mathematical Analysis*, 2012.
- A. Chambolle, M. Morini, and M. Ponsiglione. Nonlocal curvature flows. *Archive for Rational Mechanics and Analysis*, 218(3):1263–1329, 2015.
- A. Demontis, M. Melis, M. Pintor, M. Jagielski, B. Biggio, A. Oprea, C. Nita-Rotaru, and F. Roli. Why do adversarial attacks transfer? explaining transferability of evasion and poisoning attacks, 2018.

- N. Frank and J. Niles-Weed. The consistency of adversarial training for binary classification. *arxiv*, 2022.
- N. S. Frank. The structure of the adversarial bayes classifier in one dimension. *arxiv*, 2022.
- I. J. Goodfellow, J. Shlens, and C. Szegedy. Explaining and harnessing adversarial examples, 2014.
- T. Gu, B. Dolan-Gavitt, and S. Garg. Badnets: Identifying vulnerabilities in the machine learning model supply chain. *CoRR*, 2017.
- D. Jakubovitz and R. Giryes. Improving DNN robustness to adversarial attacks using jacobian regularization. *ECCV*, 2018.
- H. Kannan, A. Kurakin, and I. J. Goodfellow. Adversarial logit pairing. *CoRR*, 2018.
- A. Kurakin, I. Goodfellow, and S. Bengio. Adversarial machine learning at scale, 2017.
- M. Lewicka and Y. Peres. Which domains have two-sided supporting unit spheres at every boundary point? *Expositiones Mathematicae*, 38(4):548–558, 2020.
- Y. Lin. A note on margin-based loss functions in classification. *Statistics & Probability Letters*, 68(1):73–82, 2004.
- A. Madry, A. Makelov, L. Schmidt, D. Tsipras, and A. Vladu. Towards deep learning models resistant to adversarial attacks, 2019.
- S. A. Mingyuan Zhang. Consistency vs. h-consistency: The interplay between surrogate loss functions and the scoring function class. *NeurIps*, 2020.
- S. Moosavi-Dezfooli, A. Fawzi, J. Uesato, and P. Frossard. Robustness via curvature regularization, and vice versa. 2018.
- T. Nishiura. *Absolute Measurable Spaces*. Cambridge University Press, 2010.
- N. Papernot, P. D. McDaniel, I. J. Goodfellow, S. Jha, Z. B. Celik, and A. Swami. Practical black-box attacks against deep learning systems using adversarial examples. *CoRR*, abs/1602.02697, 2016. URL <http://arxiv.org/abs/1602.02697>.
- R. A. S. Philip M. Long. Consistency versus realizable h-consistency for multiclass classification. *ICML*, 2013.
- M. S. Pydi and V. Jog. Adversarial risk via optimal transport and optimal couplings. *arXiv preprint arXiv:1912.02794*, 2019.
- M. S. Pydi and V. Jog. The many faces of adversarial risk. *Neural Information Processing Systems*, 2021.
- R. T. Rockafellar and R. J. Wets. *Variational Analysis*. Springer, 1998.

- A. S. Ross and F. Doshi-Velez. Improving the adversarial robustness and interpretability of deep neural networks by regularizing their input gradients. *Proceedings of the AAAI Conference on Artificial Intelligence*, 2017.
- A. Rozsa, M. Günther, and T. E. Boult. Are accuracy and robustness correlated? *CoRR*, 2016.
- G. Sapiro. *Geometric Partial Differential Equations and Image Analysis*. Cambridge University Press, 2001.
- A. Shafahi, M. Najibi, A. Ghiasi, Z. Xu, J. P. Dickerson, C. Studer, L. S. Davis, G. Taylor, and T. Goldstein. Adversarial training for free! *CoRR*, 2019.
- I. Steinwart. How to compare different loss functions and their risks. *Constructive Approximation*, 2007.
- C. Szegedy, W. Zaremba, I. Sutskever, J. Bruna, D. Erhan, I. Goodfellow, and R. Fergus. Intriguing properties of neural networks. *arXiv preprint arXiv:1312.6199*, 2013.
- F. Tramèr, N. Papernot, I. Goodfellow, D. Boneh, and P. McDaniel. The space of transferable adversarial examples, 2017.
- N. G. Trillos and R. Murray. Adversarial classification: Necessary conditions and geometric flows, 2020.
- Y. Wang, X. Ma, J. Bailey, J. Yi, B. Zhou, and Q. Gu. On the convergence and robustness of adversarial training, 2021.
- C. Xie, Y. Wu, L. van der Maaten, A. L. Yuille, and K. He. Feature denoising for improving adversarial robustness. *CoRR*, 2018.
- Y.-Y. Yang, C. Rashtchian, Y. Wang, and K. Chaudhuri. Robustness for non-parametric classification: A generic attack and defense. *Proceedings of Machine Learning Research*, 2020.
- F. Yu, C. Liu, Y. Wang, L. Zhao, and X. Chen. Interpreting adversarial robustness: A view from decision surface in input space. *ICLR*, 2018.
- T. Zhang. Statistical behavior and consistency of classification methods based on convex risk minimization. *The Annals of Statistics*, 2004.

A. Measurability–Proof of Theorem 4

We will now we prove Theorem 4 using (11). This result is a consequence of a theorem of Awasthi et al. (2021):

Theorem 29 (Awasthi et al., 2021) *If $A \in \mathcal{U}(\mathbb{R}^d)$, then $A^\epsilon \in \mathcal{U}(\mathbb{R}^d)$ as well.*

Theorem 4 follows from Equation 11 and Theorem 29:

Theorem 4 *Let f be a universally measurable function. Then the function $S_\epsilon(f)(\mathbf{x})$ defined by*

$$S_\epsilon(f)(\mathbf{x}) = \sup_{\|\mathbf{h}\| \leq \epsilon} f(\mathbf{x} + \mathbf{h})$$

is universally measurable as well.

Proof If f is universally measurable, then the set $f^{-1}((a, \infty])$ is universally measurable as well. Equation 11 states that $(S_\epsilon(f))^{-1}((a, \infty]) = f^{-1}((a, \infty])^\epsilon$ and Theorem 29 then implies that this set is measurable. ■

Note that because we can write $I_\epsilon(f) = -S_\epsilon(-f)$, it immediately follows that $I_\epsilon(f)$ is universally measurable for any universally measurable f .

B. Assuming Local Extrema— Proof of Lemma 14

We start with a brief overview of the proof of Lemma 14. The criterion that h assumes its local supremum can be expressed as a set containment involving level sets of h . Because functions assume values in $\overline{\mathbb{R}}$ throughout this paper, infinite intervals will include the endpoints $\pm\infty$. Let g be any function and let $h = U_\epsilon(L_\epsilon(g))$. Set $a = \sup_{\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}} h(\mathbf{y})$. The function h assumes its supremum on $\overline{B_\epsilon(\mathbf{x})}$ iff there is a point in $h^{-1}([a, \infty])$ that is also in $\overline{B_\epsilon(\mathbf{x})}$, or in other words, $h^{-1}([a, \infty])$ intersects $\overline{B_\epsilon(\mathbf{x})}$. By Lemma 18, this condition is equivalent to $\mathbf{x} \in h^{-1}([a, \infty])^\epsilon$.

However, for any $c < a$, $\overline{B_\epsilon(\mathbf{x})}$ must intersect $h^{-1}([c, \infty])$, so $\mathbf{x} \in h^{-1}([c, \infty])^\epsilon$. Therefore, $\mathbf{x} \in \bigcap_{c < a} h^{-1}([c, \infty])^\epsilon$.

The proof of Lemma 14 proceeds by arguing that $\bigcap_{c < a} h^{-1}([c, \infty])^\epsilon = h^{-1}([a, \infty])^\epsilon$. Properties of the ${}^\epsilon, {}^{-\epsilon}$ operations are central to this result, as are properties of the inverse images of $S_\epsilon(g), I_\epsilon(g)$. We discuss the ${}^\epsilon, {}^{-\epsilon}$ operations in B.1 and inverse images of $S_\epsilon(g), I_\epsilon(g)$ in B.2. Finally, we prove Lemma 14 in B.3.

B.1 Fundamental Properties of the ${}^{-\epsilon}, {}^\epsilon$ Operations

We begin with three fundamental lemmas describing the ${}^\epsilon, {}^{-\epsilon}$ operations. The first characterizes how these operations interacts with unions and intersections. Awasthi et al. (2021) proved this lemma for countable unions and intersections. The same proof applies for uncountable unions and intersections as well, which we sketch below for completeness.

Lemma 30 (Awasthi et al., 2021) *Let \mathcal{I} be an indexing set for a collection of sets $\{A_i\}_{i \in \mathcal{I}}$. Then the following set relations hold:*

$$\bigcup_{i \in \mathcal{I}} A_i^\epsilon = \left[\bigcup_{i \in \mathcal{I}} A_i \right]^\epsilon \quad (25) \qquad \bigcap_{i \in \mathcal{I}} A_i^{-\epsilon} = \left[\bigcap_{i \in \mathcal{I}} A_i \right]^{-\epsilon} \quad (26)$$

$$\bigcap_{i \in \mathcal{I}} A_i^\epsilon \supset \left[\bigcap_{i \in \mathcal{I}} A_i \right]^\epsilon \quad (27) \qquad \bigcup_{i \in \mathcal{I}} A_i^{-\epsilon} \subset \left[\bigcup_{i \in \mathcal{I}} A_i \right]^{-\epsilon} \quad (28)$$

Proof We start by proving (25). Recall that for a set A , $A^\epsilon = A \oplus \overline{B_\epsilon(\mathbf{0})} = \bigcup_{\mathbf{a} \in A} \overline{B_\epsilon(\mathbf{a})}$. Therefore

$$\bigcup_{i \in \mathcal{I}} A_i^\epsilon = \bigcup_{i \in \mathcal{I}} \bigcup_{\mathbf{a} \in A_i} \overline{B_\epsilon(\mathbf{a})} = \bigcup_{\mathbf{a} \in \bigcup_{i \in \mathcal{I}} A_i} \overline{B_\epsilon(\mathbf{a})} = \left[\bigcup_{i \in \mathcal{I}} A_i \right]^\epsilon.$$

Relation (26) follows from by applying (25) to the collection of sets $\{A_i^C\}_{i \in \mathcal{I}}$ and then taking complements. We will now show (27). Let \mathbf{x} be a point in $(\bigcap_{i \in \mathcal{I}} A_i)^\epsilon$. Then $\mathbf{x} = \mathbf{a} + \mathbf{h}$ for some $\mathbf{a} \in \bigcap_{i \in \mathcal{I}} A_i$ and $\mathbf{h} \in \overline{B_\epsilon(\mathbf{0})}$. Because \mathbf{a} is in each A_i , $\mathbf{x} = \mathbf{a} + \mathbf{h} \in A_i^\epsilon$. Therefore $\mathbf{x} \in \bigcap_{i \in \mathcal{I}} A_i^\epsilon$.

Relation (28) follows from by applying (27) to the collection of sets $\{A_i^C\}_{i \in \mathcal{I}}$ and then taking complements. ■

The containments in (27) and (28) are in general strict. However, Awasthi et al. (2021) found a situation for which the containment in (27) is actually an equality:

Lemma 31 (Awasthi et al., 2021) *Let B_n be a decreasing sequence of sets ($B_{n+1} \subset B_n$).*

Then if we define $C_n = ((B_n^{-\epsilon})^{2\epsilon})^{-\epsilon}$, then $\bigcap_{n=1} C_n$ is pseudo-certifiably robust at every point and the C_n satisfy

$$\bigcap_{n=1} C_n^\epsilon = \left(\bigcap_{n=1} C_n \right)^\epsilon, \quad \bigcap_{n=1} C_n^{-\epsilon} = \left(\bigcap_{n=1} C_n \right)^{-\epsilon}.$$

Now we discuss how the $^\epsilon$ and $^{-\epsilon}$ operations interact when applied in succession.

Lemma 32 (Awasthi et al., 2021) *Let A be any set. Then $((A^\epsilon)^{-\epsilon})^\epsilon = A^\epsilon$ and $((A^{-\epsilon})^\epsilon)^{-\epsilon} = A^{-\epsilon}$.*

This Lemma is in fact just Items 3) and 4) of Lemma 12 applied to the indicator function $\mathbf{1}_A$.

B.2 Inverse Images of Level Sets of $S_\epsilon(f)$ and $I_\epsilon(f)$

Next we present two simple arguments that assist in finding inverse images of level sets of $I_\epsilon(f)$ and $S_\epsilon(f)$.

Lemma 33 *Let F be any function. Then the following hold:*

$$F^{-1}([a, \infty]) = \bigcap_{c < a} F^{-1}((c, \infty]) \quad (29) \quad F^{-1}((a, \infty]) = \bigcup_{c > a} F^{-1}([c, \infty]) \quad (30)$$

Proof Notice that

$$[a, \infty] = \bigcap_{c < a} (c, \infty] \quad \text{and} \quad (a, \infty] = \bigcup_{c < a} [c, \infty].$$

Therefore (29) and (30) hold. ■

We will start with a short argument that describes how the $^\epsilon$ and $^{-\epsilon}$ operations interact with intersections and inverse images of level sets. We will use this lemma and analogous arguments several times throughout this appendix.

Lemma 34 *Let f be a function. then the following hold:*

$$\bigcap_{c < a} f^{-1}((c, \infty])^\epsilon = \bigcap_{c < a} f^{-1}([c, \infty])^\epsilon \quad (31)$$

$$\bigcup_{c < a} f^{-1}((c, \infty])^\epsilon = \bigcup_{c < a} f^{-1}([c, \infty])^\epsilon \quad (32)$$

Proof Note that $[c, \infty] \supset (c, \infty]$ implies that $f^{-1}([c, \infty])^\epsilon \supset f^{-1}((c, \infty])^\epsilon$. Similarly, because (31) involves an intersection over all $c < a$, there always exists a c' with $c < c' < a$ and thus there is always a c' for which $f^{-1}((c', \infty])^\epsilon \supset f^{-1}([c, \infty])^\epsilon \supset f^{-1}((c, \infty])^\epsilon$. Intersecting over all $c < c' < a$ and $c < a$ results in

$$\bigcap_{c' < a} f^{-1}((c', \infty])^\epsilon \supset \bigcap_{c < a} f^{-1}([c, \infty])^\epsilon \supset \bigcap_{c < a} f^{-1}((c, \infty])^\epsilon$$

and therefore

$$\bigcap_{c < a} f^{-1}((c, \infty])^\epsilon = \bigcap_{c < a} f^{-1}([c, \infty])^\epsilon.$$

Similarly, taking unions of $f^{-1}((c', \infty])^\epsilon \supset f^{-1}([c, \infty])^\epsilon \supset f^{-1}((c, \infty])^\epsilon$ over all $c < c' < a$ and $c < a$ results in

$$\bigcup_{c' < a} f^{-1}((c', \infty])^\epsilon \supset \bigcup_{c < a} f^{-1}([c, \infty])^\epsilon \supset \bigcup_{c < a} f^{-1}((c, \infty])^\epsilon$$

which implies (32). ■

Lemma 35 *Define $I_\epsilon(f), S_\epsilon(f)$ as in Equations 7 and 8. Then*

- 1) $(S_\epsilon(f))^{-1}((a, \infty]) = f^{-1}((a, \infty])^\epsilon$
- 2) $(S_\epsilon(f))^{-1}([-\infty, a]) = f^{-1}([-\infty, a])^{-\epsilon}$
- 3) $(I_\epsilon(f))^{-1}([a, \infty]) = f^{-1}([a, \infty])^{-\epsilon}$
- 4) $(I_\epsilon(f))^{-1}([-\infty, a]) = f^{-1}([-\infty, a])^\epsilon$
- 5) $(S_\epsilon(f))^{-1}([a, \infty]) = \bigcap_{c < a} f^{-1}((c, \infty])^\epsilon = \bigcap_{c < a} f^{-1}([c, \infty])^\epsilon$
- 6) $(I_\epsilon(f))^{-1}((a, \infty]) = \bigcup_{c > a} f^{-1}([c, \infty])^{-\epsilon} = \bigcup_{c > a} f^{-1}((c, \infty])^{-\epsilon}$

Proof Item 1) is simply Equation 11 and 2) is simply Equation 12.

Now 4) follows from applying 1) to $-f$:

$$\begin{aligned} I_\epsilon(f)(\mathbf{x}) &< a \\ \Leftrightarrow -S_\epsilon(-f) &< a \\ \Leftrightarrow S_\epsilon(-f) &> -a \\ \Leftrightarrow \mathbf{x} &\in ((-f)^{-1}([-\infty, -a]))^\epsilon && \text{(by item 1)} \\ \Leftrightarrow \mathbf{x} &\in f^{-1}([-\infty, a])^\epsilon \end{aligned}$$

Similarly, 3) follows from 2).

We now show the first equality of Item 5) and Item 6). Lemma 33 and Item 1) imply that

$$(S_\epsilon(f))^{-1}([a, \infty]) = \bigcap_{c < a} S_\epsilon(f)^{-1}([c, \infty]) = \bigcap_{c < a} S_\epsilon(f)^{-1}((c, \infty]) = \bigcap_{c < a} f^{-1}((c, \infty])^\epsilon$$

Similarly, 6) follows from (30) and 3)

The second equality of 5) and 6) follow from Lemma 34. ■

B.3 Proof of Lemma 14

Next we use Lemma 35 to find functions which achieve their supremums/infimums on balls.

Lemma 14 *For any function g , $U_\epsilon(L_\epsilon(g))$ assumes its supremum on every closed ball radius ϵ and $L_\epsilon(U_\epsilon(g))$ assumes its infimum on every closed ball radius ϵ .*

Proof We will prove that for every function g , if we set $h = U_\epsilon(L_\epsilon(g))$, then h assumes its supremum on every ball radius ϵ . The for $L_\epsilon(U_\epsilon(g))$ follows from writing

$$\inf_{\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}} L_\epsilon(U_\epsilon(g))(\mathbf{y}) = - \sup_{\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}} U_\epsilon(L_\epsilon(-g))(\mathbf{y}).$$

Let $a = \sup_{\mathbf{y} \in \overline{B_\epsilon(\mathbf{x})}} h(\mathbf{y})$. The function h assumes its supremum on $\overline{B_\epsilon(\mathbf{x})}$ iff $h^{-1}([a, \infty])$ intersects $\overline{B_\epsilon(\mathbf{x})}$, or in other words, $\mathbf{x} \in h^{-1}([a, \infty])^\epsilon$. However, we know that for any $c < a$, $\overline{B_\epsilon(\mathbf{x})}$ intersects $h^{-1}([c, \infty])$ and thus $\mathbf{x} \in h^{-1}([c, \infty])^\epsilon$. Therefore $\mathbf{x} \in \bigcap_{c < a} h^{-1}([c, \infty])^\epsilon$. The result then follows from a chain of set equalities. We present the argument below and justify three of the equalities ((33),(34), (35)) at the end of the proof.

$$\begin{aligned} \mathbf{x} &\in \bigcap_{c < a} h^{-1}([c, \infty])^\epsilon \\ &= \bigcap_{c < a} \left((I_\epsilon(S_{2\epsilon}(I_\epsilon(g))))^{-1}([c, \infty]) \right)^\epsilon \\ &= \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right)^\epsilon && \text{(3) of Lemma 35} \\ &= \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right)^\epsilon && \text{(33)} \\ &= \bigcap_{c < a} \left(\left((I_\epsilon(g)^{-1}((c, \infty]))^{2\epsilon} \right)^{-\epsilon} \right)^\epsilon && \text{(1) of Lemma 35} \\ &= \bigcap_{c < a} \left(\left((I_\epsilon(g)^{-1}([c, \infty]))^{2\epsilon} \right)^{-\epsilon} \right)^\epsilon && \text{(34)} \\ &= \bigcap_{c < a} \left(\left((g^{-1}([c, \infty]))^{-\epsilon} \right)^{2\epsilon} \right)^{-\epsilon} && \text{(3) of Lemma 35} \\ &= \left(\bigcap_{c < a} \left((g^{-1}([c, \infty]))^{-\epsilon} \right)^{2\epsilon} \right)^{-\epsilon} && \text{(Lemma 31)} \\ &= h^{-1}([a, \infty])^\epsilon && \text{(35)} \end{aligned}$$

The equalities (33) and (34) follow from reasoning similar to the proof of Lemma 34.

Let F be any function. For any $c < a$, there is a c' in strictly less than c . Then $(c, \infty] \subset [c, \infty] \subset (c', \infty]$ and hence

$$F^{-1}((c, \infty]) \subset F^{-1}([c, \infty]) \subset F^{-1}((c', \infty]). \quad (36)$$

We now pick $F = S_{2\epsilon}(I_\epsilon(g))$ and apply the $^{-\epsilon}$ and $^\epsilon$ operations to conclude that

$$\left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right)^\epsilon \subset \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right)^\epsilon \subset \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c', \infty]))^{-\epsilon} \right)^\epsilon.$$

Because this relation holds for every $c' < c$,

$$\left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right)^\epsilon \subset \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right)^\epsilon \subset \bigcap_{c' < c} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c', \infty]))^{-\epsilon} \right)^\epsilon$$

and by intersecting over $c < a$, we can conclude that

$$\begin{aligned} \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right)^\epsilon &\subset \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right)^\epsilon \subset \bigcap_{c < a} \bigcap_{c' < c} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c', \infty]))^{-\epsilon} \right)^\epsilon \\ &= \bigcap_{c' < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c', \infty]))^{-\epsilon} \right)^\epsilon \end{aligned}$$

and therefore

$$\bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right)^\epsilon = \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right)^\epsilon. \quad (37)$$

This equality implies (33).

Similar reasoning implies (34). Lemma 34 implies that

$$\bigcap_{c < a} ((I_\epsilon(g))^{-1}((c, \infty]))^{2\epsilon} = \bigcap_{c < a} ((I_\epsilon(g))^{-1}([c, \infty]))^{2\epsilon}. \quad (38)$$

Lemma 32 then implies

$$\bigcap_{c < a} \left(\left(((I_\epsilon(g))^{-1}((c, \infty]))^{2\epsilon} \right)^{-\epsilon} \right)^\epsilon = \bigcap_{c < a} \left(\left(((I_\epsilon(g))^{-1}([c, \infty]))^{2\epsilon} \right)^{-\epsilon} \right)^\epsilon. \quad (39)$$

Equation 39 implies (34).

The equality (35) follows by arguing that

$$\bigcap_{c < a} \left(\left((g^{-1}([c, \infty]))^{-\epsilon} \right)^{2\epsilon} \right)^{-\epsilon} = h^{-1}([a, \infty])$$

by using a similar chain of reasoning in reverse:

$$\begin{aligned}
 & \bigcap_{c < a} \left(\left((g^{-1}([c, \infty]))^{-\epsilon} \right)^{2\epsilon} \right)^{-\epsilon} \\
 &= \bigcap_{c < a} \left((I_\epsilon(g)^{-1}([c, \infty]))^{2\epsilon} \right)^{-\epsilon} && \text{(3) of Lemma 35} \\
 &= \left(\bigcap_{c < a} (I_\epsilon(g)^{-1}([c, \infty]))^{2\epsilon} \right)^{-\epsilon} && \text{(Equation 26)} \\
 &= \left(\bigcap_{c < a} (I_\epsilon(g)^{-1}((c, \infty]))^{2\epsilon} \right)^{-\epsilon} && \text{(Equation 38)} \\
 &= \bigcap_{c < a} \left((I_\epsilon(g)^{-1}((c, \infty]))^{2\epsilon} \right)^{-\epsilon} && \text{(Equation 26)} \\
 &= \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}((c, \infty]))^{-\epsilon} \right) && \text{(1) of Lemma 35} \\
 &= \bigcap_{c < a} \left((S_{2\epsilon}(I_\epsilon(g))^{-1}([c, \infty]))^{-\epsilon} \right) && \text{(Lemma 34)} \\
 &= \bigcap_{c < a} (I_\epsilon(S_{2\epsilon}(I_\epsilon(g))))^{-1}([c, \infty]) && \text{(3) of Lemma 35} \\
 &= (I_\epsilon(S_{2\epsilon}(I_\epsilon(g))))^{-1} \left(\bigcap_{c < a} [c, \infty] \right) = (I_\epsilon(S_{2\epsilon}(I_\epsilon(g))))^{-1}([a, \infty]) \\
 &= h^{-1}([a, \infty])
 \end{aligned}$$

■

C. Constructing Pseudo-Certifiably Robust Functions— Proof of Lemmas 13 and 23

We begin by outlining the proof of Lemmas 13 and then we discuss the proof of Lemma 23. Awasthi et al. (2021) prove the following result:

Lemma 36 *Let A be any set. Then $(A^{2\epsilon})^{-\epsilon}$ and $(A^{-2\epsilon})^\epsilon$ are both pseudo-certifiably robust.*

This lemma states that the $^\epsilon, ^{-\epsilon}$ operations have a regularizing effect on sets.

Next, we state an analogous result for functions. Specifically, we show that $S_\epsilon(I_{2\epsilon}(f))$, $I_\epsilon(S_{2\epsilon}(f))$ are pseudo-certifiably robust. Applying this result to the functions $I_\epsilon(f)$, $S_\epsilon(f)$ then proves Lemma 13.

Lemma 37 *Let f be any function. Then the level sets of $S_\epsilon(I_{2\epsilon}(f))$, $I_\epsilon(S_{2\epsilon}(f))$ are pseudo-certifiably robust.*

Just like Lemma 36, Lemma 37 states that the operations S_ϵ, I_ϵ have a regularizing effect on functions when applied in succession. Notice that choosing $f = \mathbf{1}_A$ reduces Lemma 37 to Lemma 36. Thus Lemma 37 is a generalization of Lemma 36. Notably, one of our intermediate results states that monotone intersections and unions of pseudo-certifiably robust sets are also pseudo-certifiably robust. This statement suffices to show that if ϕ is decreasing, and f is pseudo-certifiably robust, then $\phi \circ f$ is pseudo-certifiably robust as well.

We use several results about pseudo-certifiably robust sets to prove Lemma 37 which we state below. Note that the proof of this statement also utilizes Lemma 35 proved in Appendix B.2.

To start, we present an argument from (Awasthi et al., 2021). This result is useful when reasoning about pseudo-certifiably robust sets, so we reproduce the proof here for clarity.

Lemma 38 (Awasthi et al., 2021) *Let \mathbf{y}_n be a sequence of points which approaches \mathbf{y} . Let $\mathbf{z} \in B_\epsilon(\mathbf{y})$. Then for all sufficiently large n , $\mathbf{z} \in B_\epsilon(\mathbf{y}_n)$.*

Proof Let $\delta = \|\mathbf{z} - \mathbf{y}\|$. Notice that this quantity is less than ϵ . Now pick N sufficiently large so that $\|\mathbf{y} - \mathbf{y}_n\| < \epsilon - \delta$. Then if $n \geq N$

$$\|\mathbf{z} - \mathbf{y}_n\| \leq \|\mathbf{z} - \mathbf{y}\| + \|\mathbf{y} - \mathbf{y}_n\| < \delta + \epsilon - \delta = \epsilon$$

■

Next, we demonstrate that monotone unions and intersections preserve the pseudo-certifiably robustness property.

Lemma 39 *If A_n is a decreasing sequence of pseudo-certifiably robust sets, then $\bigcap_{n \geq 1} A_n$ is pseudo-certifiably robust as well.*

Similarly, if B_n is an increasing sequence of pseudo-certifiably robust sets, then $\bigcup_{n \geq 1} B_n$ is pseudo-certifiably robust as well.

Proof Note that it suffices to show the statement for the decreasing A_n — if $\bigcap_{n \geq 1} A_n$ is pseudo-certifiably robust, so is its complement and therefore $\bigcup_{n \geq 1} A_n^C$ is pseudo-certifiably robust as well. Because A_n is decreasing, A_n^C is increasing.

We now prove the statement for the decreasing A_n . First consider $\mathbf{x} \in \left(\bigcap_{n \geq 1} A_n\right)^C$. Then the point \mathbf{x} must be in A_m^C for some m and therefore there exists a \mathbf{y} for which $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ and $B_\epsilon(\mathbf{y}) \subset A_m^C$. This observation implies that

$$B_\epsilon(\mathbf{y}) \subset A_m^C \subset \bigcup_{n \geq 1} A_n^C = \left(\bigcap_{n \geq 1} A_n\right)^C$$

Now pick \mathbf{x} in $\bigcap_{n \geq 1} A_n$. Then for each n , there exists a \mathbf{y}_n for which $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y}_n)}$ and $B_\epsilon(\mathbf{y}_n) \subset A_n$. Now since $\{\mathbf{y}_n\} \subset \overline{B_\epsilon(\mathbf{x})}$, \mathbf{y}_n must have a convergent subsequence $\{\mathbf{y}_{n_k}\}$ in $\overline{B_\epsilon(\mathbf{x})}$. For convenience, we will drop the double subscript and denote $\{\mathbf{y}_{n_k}\}$ as \mathbf{y}_n . Let $\mathbf{y} = \lim_{n \rightarrow \infty} \mathbf{y}_n$. Because $\|\mathbf{x} - \mathbf{y}\| \leq \epsilon$, $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$. We will now show that $B_\epsilon(\mathbf{y}) \subset A_n$ for all n , and then this containment will imply that $B_\epsilon(\mathbf{y}) \subset \bigcap_{n \geq 1} A_n$. Lemma 38 implies that for any $\mathbf{z} \in B_\epsilon(\mathbf{y})$, there is an N for which $n \geq N$ implies that $\mathbf{z} \in B_\epsilon(\mathbf{y}_n)$. Because the ball $B_\epsilon(\mathbf{y}_n)$ is contained in A_n , $\mathbf{z} \in A_n$ for all $n \geq N$. However, since A_n is decreasing, we must have that $\mathbf{z} \in A_n$ for $n < N$ as well. \blacksquare

A consequence of this lemma is that in order to verify that a function f is pseudo-certifiably robust, it suffices to show that only one of the level sets $f^{-1}([a, \infty])$, $f^{-1}((a, \infty))$, $f^{-1}([-\infty, a])$, $f^{-1}((-\infty, a))$ is pseudo-certifiably robust.

Lemma 40 *The following are equivalent:*

- a) $f^{-1}([a, \infty])$ is pseudo-certifiably robust
- b) $f^{-1}((a, \infty))$ is pseudo-certifiably robust
- c) $f^{-1}([-\infty, a])$ is pseudo-certifiably robust
- d) $f^{-1}((-\infty, a))$ is pseudo-certifiably robust

Proof Note that a set is pseudo-certifiably robust iff its complement is pseudo-certifiably robust. This observation implies that a) and d) are equivalent and b) and c) are equivalent. We will now show that b) implies a). By Lemma 33, $f^{-1}([a, \infty]) = \bigcap_{c < a} f^{-1}((c, \infty))$. Because the rationals are dense in the set $\{c : c < a\}$ the intersection above can actually be expressed as a countable intersection:

$$f^{-1}([a, \infty]) = \bigcap_{\substack{c < a \\ c \in \mathbb{Q}}} f^{-1}((c, \infty))$$

Therefore, b) and Lemma 39 imply that $f^{-1}([a, \infty])$ is pseudo-certifiably robust.

We now show that a) implies b). Again, Lemma 33 and the density of the rationals imply that

$$f^{-1}((a, \infty)) = \bigcup_{\substack{c > a \\ c \in \mathbb{Q}}} f^{-1}([c, \infty])$$

Therefore, a) and Lemma 39 imply that $f^{-1}((a, \infty])$ is pseudo-certifiably robust. ■

Proof [Proof of Lemma 37]

We first argue that it suffices to show that the level sets of $S_\epsilon(I_{2\epsilon}(f))$ are pseudo-certifiably robust for any f . Since $S_\epsilon(I_{2\epsilon}(f)) = -I_\epsilon(S_{2\epsilon}(-f))$, if the level sets of $S_\epsilon(I_{2\epsilon}(-f))$ are pseudo-certifiably robust then so are the level sets of $I_\epsilon(S_{2\epsilon}(f))$.

We now show that $S_\epsilon(I_{2\epsilon}(f))^{-1}((a, \infty])$ is pseudo-certifiably robust. Lemma 40 will then imply that all the other level sets of $S_\epsilon(I_{2\epsilon}(f))$ are pseudo-certifiably robust.

$$\begin{aligned} S_\epsilon(I_{2\epsilon}(f))^{-1}((a, \infty]) &= (I_{2\epsilon}(f)^{-1}((a, \infty]))^\epsilon && \text{(1) of Lemma 35} \\ &= \left(\bigcup_{c>a} f^{-1}((a, \infty])^{-2\epsilon} \right)^\epsilon && \text{(6) of Lemma 35} \\ &= \bigcup_{c>a} (f^{-1}((a, \infty])^{-2\epsilon})^\epsilon && \text{((25) of Lemma 30)} \end{aligned}$$

Note that the union over $c > a$ can actually be written as a countable union. Thus Lemmas 36 and 39 imply that this set is pseudo-certifiably robust. ■

Finally, we prove Lemma 13.

Lemma 13 *Let f be any function. Then $L_\epsilon(U_\epsilon(f))$ and $U_\epsilon(L_\epsilon(f))$ are pseudo-certifiably robust.*

Proof Apply Lemma 37 to $S_\epsilon(f), I_\epsilon(f)$. ■

Lastly, we prove Lemma 23:

Lemma 23 *Let $\phi: \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ be a decreasing function and let $f: \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$ be a pseudo-certifiably robust function. Then $\phi \circ f$ is pseudo-certifiably robust as well.*

Proof To start, we show that $(\phi \circ f)^{-1}([a, \infty])$ is a pseudo-certifiably robust set.

Notice that because ϕ is decreasing, if $f(\mathbf{x}) \leq c$, then $(\phi \circ f)(\mathbf{x}) \geq a$ iff $\phi(c) \geq a$. Therefore,

$$(\phi \circ f)^{-1}([a, \infty]) = \bigcup_{c: \phi(c) \geq a} f^{-1}([-\infty, c]) \quad (40)$$

Now because ϕ is decreasing, the set $\{c: \phi(c) \geq a\} = \phi^{-1}([\infty, a])$ must be an interval of the form $[-\infty, r]$ or $[\infty, r)$ for some $r \in \overline{\mathbb{R}}$. Because the rationals are dense in $[-\infty, r]$ and $[-\infty, r)$, the union in (40) can actually be expressed as a countable union of increasing sets. Therefore, Lemma 39 implies that $(\phi \circ f)^{-1}([a, \infty])$ is pseudo-certifiably robust. Lemma 40 then implies that all the other level sets of $\phi \circ f$ are pseudo-certifiably robust. ■

D. Upper Envelopes and Pseudo-certifiably Robust Functions— Proof of Lemmas 17 and 22

We briefly outline of the proofs of Lemmas 17 and 22. The key observation in Lemma 22 is that if A_n are decreasing pseudo-certifiably robust sets, then $\overline{\bigcap_{n \geq 1} A_n} = \bigcap_{n \geq 1} \overline{A_n}$. This result allows us to commute closures with intersections when computing the inverse image of $\limsup_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$. This fact together with Lemma 35 of Appendix B.2 then implies the result. We also use Lemmas 38 and 40 of Appendix C in the proofs of this section.

Lemma 17 follows from Lemma 22 and the observation that if A is pseudo-certifiably robust, then so is \overline{A} .

D.1 Proof of Lemma 22

Lemma 41 *We have the following expression for \overline{A} :*

$$\overline{A} = \bigcap_{\delta > 0} A^\delta \tag{41}$$

Proof

We will first show $\overline{A} \subset \bigcap_{\delta > 0} A^\delta$. Consider any point \mathbf{a} in \overline{A} . Then for any $\delta > 0$, $\overline{B_\delta(\mathbf{a})}$ intersects A . Lemma 18 then implies that \mathbf{a} is in A^δ . Therefore $\mathbf{a} \in \bigcap_{\delta > 0} A^\delta$.

It remains to show the other inclusion. Specifically, we will show $\overline{A^C} \subset (\bigcap_{\delta > 0} A^\delta)^C = \bigcup_{\delta > 0} (A^\delta)^C$. Pick $\mathbf{x} \in \overline{A^C}$. Then since $\overline{A^C}$ is open, there is a $\delta > 0$ for which $\overline{B_\delta(\mathbf{x})}$ is disjoint from A . By Lemma 18, $\overline{B_\delta(\mathbf{x})}$ is disjoint from A is equivalent to $\mathbf{x} \notin A^\delta$. Thus $\mathbf{x} \in (A^\delta)^C$. ■

In the next lemma, we only require half of the pseudo-certifiably robust assumption, namely we require that for every $\mathbf{x} \in A_n$, there is a ball $\overline{B_\epsilon(\mathbf{y})}$ with $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ contained in A_n but we don't impose this requirement for A_n^C . This lemma implies that pseudo-certifiably robust sets interact nicely with topological closures.

Lemma 42 *Let A_n be a decreasing sequence of pseudo-certifiably robust sets. Then*

$$\overline{\bigcap_{n \geq 1} A_n} = \bigcap_{n \geq 1} \overline{A_n}$$

In fact, in the proof of this result, we only use half of the pseudo-certifiably robustness property: we use that for every $\mathbf{x} \in A_n$, there is a ball $\overline{B_\epsilon(\mathbf{y})}$ with $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$ and $B_\epsilon(\mathbf{y}) \subset A_n$. [[todo: make sure we can replace the hypothesis with pcr](#)]

Proof To start, $\bigcap_{n \geq 1} \overline{A_n}$ is a closed set containing $\bigcap_{n \geq 1} A_n$ so $\overline{\bigcap_{n \geq 1} A_n}$ is contained in $\bigcap_{n \geq 1} \overline{A_n}$.

We will now show the other inclusion. Let $\mathbf{x} \in \bigcap_{n \geq 1} \overline{A_n}$. Then for each n , there is a sequence $\mathbf{x}_{n,k}$ for which $\mathbf{x}_{n,k} \in A_n$ and $\mathbf{x}_{n,k}$ approaches \mathbf{x} . Because each A_n is pseudo-certifiably robust, there is a $\mathbf{y}_{n,k}$ for which $\mathbf{x}_{n,k} \in \overline{B_\epsilon(\mathbf{y}_{n,k})}$ and $B_\epsilon(\mathbf{y}_{n,k}) \subset A_n$. Let k_n be the index for which $\|\mathbf{x}_{n,k_n} - \mathbf{x}\| \leq 2^{-n}$. Now consider the diagonal sequence \mathbf{y}_{n,k_n} . By the definition of k_n ,

$$\|\mathbf{x} - \mathbf{y}_{n,k_n}\| \leq \|\mathbf{x} - \mathbf{x}_{n,k_n}\| + \|\mathbf{x}_{n,k_n} - \mathbf{y}_{n,k_n}\| \leq 2^{-n} + \epsilon$$

and therefore $\limsup_{n \rightarrow \infty} \|\mathbf{y}_{n,k_n} - \mathbf{x}\| \leq \epsilon$. Therefore, the sequence $\{\mathbf{y}_{n,k_n}\}$ has a limit point \mathbf{y} in $\overline{B_\epsilon(\mathbf{x})}$. Now Lemma 38 implies that for any $\mathbf{z} \in B_\epsilon(\mathbf{y})$, there is an N for which $n \geq N$ implies that $\mathbf{z} \in B_\epsilon(\mathbf{y}_{n,k_n})$. Because the ball $B_\epsilon(\mathbf{y}_{n,k_n})$ is contained in A_n , $\mathbf{z} \in A_n$ for all $n \geq N$. However, since A_n is decreasing, we must have that $\mathbf{z} \in A_n$ for $n < N$ as well. Therefore $B_\epsilon(\mathbf{y})$ is contained in $\bigcap_{n \geq 1} A_n$. Since $\mathbf{x} \in \overline{B_\epsilon(\mathbf{y})}$, $\mathbf{x} \in \overline{\bigcap_{n \geq 1} A_n}$. ■

Lemma 22 *Let F be a function whose level sets are pseudo-certifiably robust and let $G(\mathbf{x}) = \limsup_{\mathbf{y} \rightarrow \mathbf{x}} F(\mathbf{y})$. Then*

$$G^{-1}([a, \infty]) = \overline{F^{-1}([a, \infty])} \quad (13)$$

Proof Note that we can write $G(\mathbf{x}) = \inf_{\delta} \sup_{\overline{B_\delta(\mathbf{x})}} F(\mathbf{y})$. Therefore

$$\begin{aligned} \{\mathbf{x}: G(\mathbf{x}) \geq a\} &= \bigcap_{\delta > 0} \{\mathbf{x}: \sup_{\mathbf{y} \in \overline{B_\delta(\mathbf{x})}} F(\mathbf{y}) \geq a\} \\ &= \bigcap_{\delta > 0} S_\delta(F)^{-1}([a, \infty]) \\ &= \bigcap_{\delta > 0} \bigcap_{c > a} F^{-1}([c, \infty])^\delta && \text{(Lemma 35)} \\ &= \bigcap_{c > a} \overline{F^{-1}([c, \infty])} && \text{(Lemma 41)} \\ &= \bigcap_{c > a} F^{-1}([c, \infty]) && \text{(Lemma 42)} \\ &= \overline{F^{-1}([a, \infty])} \end{aligned}$$

■

D.2 Proof of Lemma 17

We now show that the boundary of a set does not affect whether its pseudo-certifiably robust.

Lemma 43 *Let A be a pseudo-certifiably robust set and let $C \subset \partial A$. Then $A - C$ is also pseudo-certifiably robust.*

Proof To start, if $B_\epsilon(\mathbf{y}) \subset A$, then $B_\epsilon(\mathbf{y}) \subset A - C$. Therefore, if $\mathbf{a} \in A - C$ and $\overline{B_\epsilon(\mathbf{y})}$ is a ball for which $\mathbf{a} \in \overline{B_\epsilon(\mathbf{y})}$ with $B_\epsilon(\mathbf{y}) \subset A$, then $B_\epsilon(\mathbf{y}) \subset A$ as well. Next, if $B_\epsilon(\mathbf{y}) \subset A^C$, then $B_\epsilon(\mathbf{y}) \subset (A - C)^C$. Therefore, if $\mathbf{a} \in A^C$, and $\overline{B_\epsilon(\mathbf{y})}$ is a ball for which $\mathbf{a} \in \overline{B_\epsilon(\mathbf{y})}$ with $B_\epsilon(\mathbf{y}) \subset A^C$, then $B_\epsilon(\mathbf{y}) \subset A^C$ as well. Thus it remains to show that if $\mathbf{c} \in C$, then there is a ball $B_\epsilon(\mathbf{y})$ for which $B_\epsilon(\mathbf{y}) \subset (A - C)^C$ and $\mathbf{c} \in \overline{B_\epsilon(\mathbf{y})}$. Consider a sequence of points \mathbf{a}_i in A^C which converges to \mathbf{c} . By the pseudo-certifiably robustness of A , one can find a sequence of points \mathbf{y}_i for which $\mathbf{a}_i \in \overline{B_\epsilon(\mathbf{y}_i)}$ and $B_\epsilon(\mathbf{y}_i) \subset A^C$. Because the \mathbf{y}_i s are in a bounded set for sufficiently large i , there is a subsequence \mathbf{y}_{i_k} which converges to a point \mathbf{y} . Lemma 38 then implies that $B_\epsilon(\mathbf{y}) \subset A^C$. Next we will show that \mathbf{y} is distance

at most ϵ from \mathbf{c} so $\mathbf{c} \in \overline{B_\epsilon(\mathbf{y})}$. Pick $\delta > 0$ and k large enough so that $\|\mathbf{a}_{i_k} - \mathbf{a}\| \leq \delta$ and $\|\mathbf{y}_{i_k} - \mathbf{y}\| \leq \delta$. Then

$$\|\mathbf{y} - \mathbf{c}\| \leq \|\mathbf{y} - \mathbf{y}_{i_k}\| + \|\mathbf{y}_{i_k} - \mathbf{a}_{i_k}\| + \|\mathbf{a}_{i_k} - \mathbf{c}\| \leq \delta + \epsilon + \delta = \epsilon + 2\delta$$

Because δ was arbitrary, $\|\mathbf{y} - \mathbf{c}\| \leq \epsilon$. ■

Notice that because a set A is pseudo-certifiably robust iff A^C is pseudo-certifiably robust, this lemma also implies that if C is an arbitrary subset of the boundary $A \cup C$ is also pseudo-certifiably robust.

Lemma 44 *If A is a pseudo-certifiably robust set, then so are \overline{A} and $\text{int } A$.*

Proof This statement is a consequence of Lemma 43. ■

Next, we prove that the \liminf operation preserves pseudo-certifiable robustness.

Lemma 17 *Let f be a pseudo-certifiably robust function. Then $\liminf_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ is also pseudo-certifiably robust.*

Proof Note that a function h pseudo-certifiably robust iff $-h$ is pseudo-certifiably robust as well. We show that $g(\mathbf{x}) := \limsup_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ is pseudo-certifiably robust by studying its level sets. One can then conclude that $\liminf_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y})$ is pseudo-certifiably robust by writing $\liminf_{\mathbf{y} \rightarrow \mathbf{x}} f(\mathbf{y}) = -\limsup_{\mathbf{y} \rightarrow \mathbf{x}} -f(\mathbf{y})$. Lemma 22 implies that

$$g^{-1}([a, \infty)) = \overline{f^{-1}([a, \infty])}$$

This set is pseudo-certifiably robust by Lemma 44. Lemma 40 then implies that f is pseudo-certifiably robust. ■