

Reconfiguration of Vertex Colouring and Forbidden Induced Subgraphs

Manoj Belavadi ^{*} Kathie Cameron [†] Owen Merkel [‡]

March 7, 2023

Abstract

The reconfiguration graph of the k -colourings, denoted $\mathcal{R}_k(G)$, is the graph whose vertices are the k -colourings of G and two colourings are adjacent in $\mathcal{R}_k(G)$ if they differ in colour on exactly one vertex. In this paper, we investigate the connectivity and diameter of $\mathcal{R}_{k+1}(G)$ for a k -colourable graph G restricted by forbidden induced subgraphs. We show that $\mathcal{R}_{k+1}(G)$ is connected for every k -colourable H -free graph G if and only if H is an induced subgraph of P_4 or $P_3 + P_1$. We also start an investigation into this problem for classes of graphs defined by two forbidden induced subgraphs. We show that if G is a k -colourable $(2K_2, C_4)$ -free graph, then $\mathcal{R}_{k+1}(G)$ is connected with diameter at most $4n$. Furthermore, we show that $\mathcal{R}_{k+1}(G)$ is connected for every k -colourable (P_5, C_4) -free graph G .

Keywords: reconfiguration graph, forbidden induced subgraph, k -colouring, k -mixing, frozen colouring.

1 Introduction

Let G be a finite simple graph with vertex-set $V(G)$ and edge-set $E(G)$. We use $n = |V(G)|$ to denote the number of vertices of G when the context is clear. For a positive integer k , a k -colouring of G is a mapping $\alpha: V(G) \rightarrow \{1, 2, \dots, k\}$ such that $\alpha(u) \neq \alpha(v)$ whenever $uv \in E(G)$. We say that G is k -colourable if it admits a k -colouring and the *chromatic number* of G , denoted $\chi(G)$, is the smallest integer k such that G is k -colourable. The set of vertices that are assigned the same colour is called a *colour class*. We say that the colour classes of a colouring α of G matches the colour classes of a colouring β of G if both colourings induce the same colour classes.

The *reconfiguration graph of the k -colourings*, denoted $\mathcal{R}_k(G)$, is the graph whose vertices are the k -colourings of G and two colourings are joined by an edge if they differ in colour on exactly one vertex. We say that G is k -mixing if $\mathcal{R}_k(G)$ is connected and the *k -recolouring diameter* of G is the diameter of $\mathcal{R}_k(G)$. Given two k -colourings α and β of G , deciding whether there exists a path between the two colourings in $\mathcal{R}_k(G)$ was proved to be PSPACE-complete for all $k > 3$ [5]. The problem remains PSPACE-complete for graphs with bounded bandwidth and hence bounded treewidth [13].

^{*}Department of Mathematics, Wilfrid Laurier University, Waterloo, ON, Canada, N2L 3C5. Email: mbelavadi@wlu.ca. ORCID: 0000-0002-3153-2339. Research supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2016-06517.

[†]Department of Mathematics, Wilfrid Laurier University, Waterloo, ON, Canada, N2L 3C5. Email: kcameron@wlu.ca. ORCID: 0000-0002-0112-2494. Research supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2016-06517.

[‡]Department of Mathematics, Wilfrid Laurier University, Waterloo, ON, Canada, N2L 3C5. Email: owenmerkel@gmail.com. ORCID: 0000-0002-8839-6243. Research supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2016-06517.

A k -colouring of a graph G is called *frozen* if it is an isolated vertex in $\mathcal{R}_k(G)$. In other words, for every vertex $v \in V(G)$, each of the k colours appears in the closed neighbourhood of v . One way to show that a graph G is not k -mixing is to exhibit a frozen k -colouring of G . Since every k -colouring of K_k is frozen, it is common to study $\mathcal{R}_{k+1}(G)$ for a k -colourable graph G . For the rest of this paper, we assume that $\ell \geq \chi(G) + 1$.

A graph G is H -free if no induced subgraph of G is isomorphic to H . For a collection of graphs \mathcal{H} , G is \mathcal{H} -free if G is H -free for every $H \in \mathcal{H}$. Let P_n , C_n , and K_n denote the path, cycle, and complete graph on n vertices, respectively. For two vertex-disjoint graphs G and H , the *disjoint union* of G and H , denoted by $G+H$, is the graph with vertex-set $V(G) \cup V(H)$ and edge-set $E(G) \cup E(H)$. For a positive integer r , we use rG to denote the graph obtained from the disjoint union of r copies of G . The goal of this paper is to answer the following question.

Question 1. *For which H is every H -free graph ℓ -mixing?*

Question 1 has been answered completely when H is a 3-vertex graph, namely $3K_1$, $P_2 + P_1$, P_3 , and K_3 . The last author [11] showed that every $3K_1$ -free graph is ℓ -mixing and the ℓ -recolouring diameter is at most $4n$. The class of $(P_2 + P_1)$ -free graphs is a subclass of P_4 -free graphs, and so we refer to the results on P_4 -free graphs in this case (see Theorem 1). A graph is P_3 -free if it is the disjoint union of cliques and it is well known that every P_3 -free graph is ℓ -mixing (see e.g. [4]). Bonamy and Bousquet [3] showed that the ℓ -recolouring diameter of a P_3 -free graph is at most $2n$. Cereceda, van den Heuvel, and Johnson [6] showed that for all $\ell \geq 3$, there is a bipartite graph that is not ℓ -mixing (see Figure 3). Since bipartite graphs are a subclass of K_3 -free graphs, it follows that not every K_3 -free graph is ℓ -mixing. See Table 1 for a summary of these results.

H	Always ℓ -mixing	Upper bound on the ℓ -recolouring diameter
$3K_1$	YES [11]	$4n$ [11]
$P_2 + P_1$	YES [3]	$4n$ [1] (Theorem 1)
P_3	YES [4]	$2n$ [3]
K_3	NO [6]	∞ [6] (see Figure 3)

Table 1: Summary of recolouring an H -free graph for a 3-vertex graph H .

Next we survey results on Question 1 when H is a 4-vertex graph. There are 11 graphs on 4 vertices (see Figure 1). Bonamy and Bousquet [3] showed that every P_4 -free graph G is ℓ -mixing and the ℓ -recolouring diameter is at most $2 \cdot \chi(G) \cdot n$. Biedl, Lubiw, and Merkel [1] investigated a class of graphs that generalizes P_4 -free graphs. The following theorem was not directly stated in [1] but follows from the proof of Theorem 1 in [1] and improves the bound on the ℓ -recolouring diameter of a P_4 -free graph.

Theorem 1 ([1]). *The ℓ -recolouring diameter of a P_4 -free graph is at most $4n$.*

Feghali and Merkel [8] showed that for all $p \geq 1$, there exists a k -colourable $2K_2$ -free graph with a frozen $(k+p)$ -colouring. This result together with the result of Bonamy and Bousquet [3] answers Question 1 when H is a path.

Theorem 2 ([8]). *Every P_t -free graph is ℓ -mixing if and only if $t \leq 4$.*

The connectivity and diameter of $\mathcal{R}_\ell(G)$ has also been investigated for classes of graphs defined by more than one forbidden induced subgraph. Bonamy, Johnson, Lignos, Patel, and Paulusma [4] showed that every chordal graph and every chordal bipartite graph is ℓ -mixing and the ℓ -recolouring diameter is at most $2n^2$. Feghali and Fiala [7] showed that every co-chordal graph is ℓ -mixing and the ℓ -recolouring diameter is at most $2n^2$. They also showed that every




Figure 1: The 11 graphs on 4 vertices.

3-colourable (P_5 , $\text{co-}P_5$, C_5)-free graph is ℓ -mixing and the ℓ -recolouring diameter is at most $2n^2$ [7]. Biedl, Lubiw, and Merkel [1] showed that every P_4 -sparse graph is ℓ -mixing and the ℓ -recolouring diameter is at most $4n^2$. The last author [11] showed that for all $p \geq 1$, there exists a k -colourable weakly chordal graph that is not $(k + p)$ -mixing.

Our contributions

We completely answer Question 1 with the following theorem.

Theorem 3. *Every H -free graph is ℓ -mixing if and only if H is an induced subgraph of P_4 or $P_3 + P_1$.*

We use the following theorems in the proof of Theorem 3.

Theorem 4. *For all $p \geq 1$, there exists a k -colourable $(4K_1, C_4, \text{claw})$ -free graph that is not $(k + p)$ -mixing.*

Theorem 5. *For all $p \geq 1$, there exists a k -colourable $(K_4, \text{diamond}, \text{paw}, \text{co-claw}, \text{co-diamond})$ -free graph that is not $(k + p)$ -mixing.*

Theorem 6. *Every $(P_3 + P_1)$ -free graph is ℓ -mixing and the ℓ -recolouring diameter is at most $6n$.*

The proof of Theorem 6 leads to a polynomial-time algorithm to find a path of length at most $6n$ between any two ℓ -colourings of G . See Table 2 for a summary of these results.

H	Always ℓ -mixing	Upper bound on the ℓ -recolouring diameter
$4K_1$	NO	∞ (Theorem 4)
co-diamond	NO	∞ (Theorem 5)
$2K_2$	NO [8]	∞ [8]
$P_3 + P_1$	YES	$6n$ (Theorem 6)
claw	NO	∞ (Theorem 4)
P_4	YES [3]	$4n$ [1] (Theorem 1)
co-claw	NO	∞ (Theorem 5)
paw	NO	∞ (Theorem 5)
C_4	NO	∞ (Theorem 4)
diamond	NO	∞ (Theorem 5)
K_4	NO	∞ (Theorem 5)

Table 2: Summary of recolouring an H -free graph for a 4-vertex graph H .

We also start an investigation into classes of graphs defined by two forbidden induced subgraphs.

Theorem 7. *Every $(2K_2, C_4)$ -free graph is ℓ -mixing and the ℓ -recolouring diameter is at most $4n$.*

Note that the class of split graphs, equivalently the class of chordal and co-chordal graphs, is a subclass of $(2K_2, C_4)$ -free graphs. Theorem 7 improves the upper bound on the ℓ -recolouring diameter for a split graph from $2n^2$ to $4n$. We also investigate the superclass of (P_5, C_4) -free graphs and prove the following.

Theorem 8. *Every (P_5, C_4) -free graph is ℓ -mixing.*

We note that there are P_5 -free graphs and C_4 -free graphs that are not ℓ -mixing.

The rest of the paper is organized as follows. In Section 2 we give definitions and terminology used throughout the paper. We prove Theorems 4 and 5 in Section 3 and we prove Theorem 6 in Section 4. We prove Theorem 3 in Section 5 and we prove Theorems 7 and 8 in Section 6. We end with some open problems in Section 7.

2 Preliminaries

For a graph G , the *complement* of G is the graph with vertex-set $V(G)$ such that the edges of the complement are exactly the non-edges of G . A *component* of G is a maximal connected subgraph and an *anticomponent* of G is a component of the complement of G . For a vertex $v \in V(G)$, the *open neighbourhood* of v is the set of vertices adjacent to v in G . The *closed neighbourhood* of v is the set of vertices adjacent to v in G together with v . For $X, Y \subseteq V(G)$, we say that X is *complete* to Y if every vertex in X is adjacent to every vertex in Y . If no vertex of X is adjacent to a vertex of Y , we say that X is *anticomplete* to Y . Let G and H be vertex-disjoint graphs and let $v \in V(G)$. By *substituting* H for the vertex v of G , we mean taking the graph $G - v$ and adding an edge between every vertex of H and every vertex of $G - v$ that is adjacent to v in G .

For a colouring α of G and $X \subseteq V(G)$, we say that the colour c *appears* in X if $\alpha(x) = c$ for some $x \in X$, and we use $\alpha(X)$ for the set of colours appearing in X and $|\alpha(X)|$ for the number of colours appearing on X . A k -colouring of a graph G is called *frozen* if it is an isolated vertex in the recolouring graph $\mathcal{R}_k(G)$. In other words, for every vertex $v \in V(G)$, each of the k colours appears in the closed neighbourhood of v .

3 Frozen colourings

We use frozen colourings to prove Theorems 4 and 5, answering Question 1 in the negative for 8 out of the 11 cases when H is a 4-vertex graph. To prove Theorem 4, we construct a family of graphs $\{G_p \mid p \geq 1\}$ such that G_p is a $2p$ -colourable $(4K_1, C_4, \text{claw})$ -free graph that has a frozen $3p$ -colouring.

Lemma 1. *For $p \geq 1$, let G_p be the graph obtained from C_6 by substituting the complete graph K_p into each vertex. Then G_p is $(4K_1, C_4, \text{claw})$ -free, is $2p$ -colourable, and has a frozen $3p$ -colouring.*

Proof. See Figure 2 for a $2p$ -colouring and frozen $3p$ -colouring of G_p . Notice that the vertices of G_p can be partitioned into 3 cliques, and so G_p must be $4K_1$ -free. Also notice that every vertex of G_p is bisimplicial. That is, the closed neighbourhood of every vertex can be partitioned into




Figure 2: A $2p$ -colouring of G_p and a frozen $3p$ -colouring.




Figure 3: A frozen p -colouring of the graph B_p . [6]

two cliques, and so G_p must be claw-free. Finally, suppose G_p contains an induced C_4 . Then G_p contains an induced P_3 , call it P , with vertices w, x, y in order. Notice that each vertex of P must belong to a distinct copy of K_p . But then any other vertex of G_p that is adjacent to both w and x must also be adjacent to y , a contradiction. \square

To prove Theorem 5, we use the family of bipartite graphs $\{B_p \mid p \geq 3\}$ introduced by Cereceda, van den Heuvel, and Johnson [6]. For $p \geq 3$, the graph B_p is 2-colourable, and has a frozen p -colouring.

Lemma 2. *For $p \geq 3$, let B_p be the graph obtained from the complete bipartite graph $K_{p,p}$ by deleting the edges of a perfect matching. Then B_p is $(K_4, \text{diamond}, \text{paw}, \text{co-claw}, \text{co-diamond})$ -free, 2-colourable, and has a frozen p -colouring.*

Proof. See Figure 3 for a frozen p -colouring of B_p . Since B_p is bipartite, it is also K_3 -free and so it must be $(K_4, \text{diamond}, \text{paw}, \text{co-claw})$ -free. To show that B_p is co-diamond-free, we show that the complement of B_p is diamond-free. The complement of B_p consists of two disjoint cliques with p vertices, call them Q_1 and Q_2 , and the edges between Q_1 and Q_2 are a perfect matching. Suppose by contradiction that the complement of B_p contains an induced diamond D . Let w, x be the adjacent vertices of degree 3 in D and let y, z be the non-adjacent vertices of degree 2 in D . Since each vertex in Q_i , for $i = 1, 2$, can be adjacent to at most one vertex in the other clique, either $x, w, y \in Q_1$ or $x, w, z \in Q_1$. Without loss of generality, let $x, w, y \in Q_1$. Since z is not adjacent to y , z must be in Q_2 . But then $z \in Q_2$ must be adjacent to two vertices in Q_1 , a contradiction. \square

4 Recolouring $(P_3 + P_1)$ -free graphs

In this section, we investigate the connectivity and diameter of $\mathcal{R}_\ell(G)$ for a $(P_3 + P_1)$ -free graph G . Our strategy takes advantage of the anticomponents of G . Since there are all possible edges between the anticomponents of a graph, in any colouring of the graph, the colours that appear on each anticomponent must be pairwise distinct. Thus if we can recolour a vertex in some anticomponent of G , that recolouring step would extend to a recolouring step for all of G .

Given two arbitrary ℓ -colourings α and β of G , we show how to recolour α into β using the following strategy. First fix an arbitrary $\chi(G)$ -colouring γ of G by finding an optimal colouring of each anticomponent of G . Next recolour α into a $\chi(G)$ -colouring α' of G by recolouring each anticomponent A of G so that the colour classes of α'_A match the colour classes of γ_A . Similarly recolour β into a $\chi(G)$ -colouring β' of G by recolouring each anticomponent A of G so that the colour classes of β'_A match the colour classes of γ_A . Once we have the colourings α' and β' , we use the following Renaming Lemma.

Lemma 3 (Renaming Lemma [3]). *Let α' and β' be two k -colourings of G that induce the same partition of vertices into colour classes and let $\ell \geq k + 1$. Then α' can be recoloured into β' in $\mathcal{R}_\ell(G)$ by recolouring each vertex at most 2 times.*

Olariu [12] proved the following useful result characterizing the anticomponents of a $(P_3 + P_1)$ -free graph.

Theorem 9 ([12]). *Each connected component of a paw-free graph is either K_3 -free or $(P_2 + P_1)$ -free.*

Therefore each anticomponent of a $(P_3 + P_1)$ -free graph is either $3K_1$ -free or P_3 -free. The last author proved the following theorem on recolouring $3K_1$ -free graphs.

Theorem 10 ([11]). *Every $3K_1$ -free graph is ℓ -mixing and the ℓ -recolouring diameter is at most $4n$.*

The proof of Theorem 10 uses the following lemma which we will use in the proof of Theorem 6.

Lemma 4 ([11]). *Let γ be a $\chi(G)$ -colouring of a $3K_1$ -free graph G . Any ℓ -colouring of G can be recoloured into a $\chi(G)$ -colouring γ' such that the colour classes of γ' match the colour classes of γ by recolouring each vertex at most once.*

We are now ready to prove Theorem 6.

Proof of Theorem 6. Let G be a k -colourable $(P_3 + P_1)$ -free graph and let α and β be two ℓ -colourings of G with $\ell \geq k + 1$. Fix a $\chi(G)$ -colouring γ of G . We note that this colouring can be found in polynomial time by optimally colouring each anticomponent of G [10].

Let A_1, A_2, \dots, A_p be the anticomponents of G . Since the anticomponents of any graph are pairwise complete, the colours used on each anticomponent are pairwise distinct, and thus $\chi(G) = \sum_{1 \leq i \leq p} \chi(A_i)$. We proceed by recolouring each anticomponent A_i of G so that the colouring of A_i matches the colour classes of A_i in γ . The anticomponent that is selected to be recoloured is determined based on the following claim.

Claim 1. *In any ℓ -colouring α of G , either some colour does not appear in α or there is some anticomponent A of G such that α uses at least $\chi(A) + 1$ colours on A .*

Proof of claim: Suppose α colours each anticomponent A_i of G with $\chi(A_i)$ colours. Since $\chi(G) = \sum_{1 \leq i \leq p} \chi(A_i)$, only $\chi(G)$ colours appear in α and since $\ell \geq \chi(G) + 1$, some colour does not appear in α . This completes the proof of the Claim 1.

If some colour c is not being used, we take some anticomponent A_i of G that has not been recoloured. Recolour A_i as described below depending on whether A_i is P_3 -free or $3K_1$ -free using c together with the $\chi(A_i)$ colours appearing on A_i .

Now suppose that there is some anticomponent A_i where the current colouring uses at least $\chi(A_i) + 1$ colours on A_i . If A_i is P_3 -free then since each component of A_i is a clique, we use the Renaming Lemma to recolour each component of A_i to match the colour classes of γ by recolouring each vertex at most twice. If A_i is $3K_1$ -free, then we use Lemma 4 to recolour the current colouring of A_i to match the colour classes of γ .

We have recoloured α into a $\chi(G)$ -colouring α' whose colour classes correspond to the colour classes of γ by recolouring each vertex at most twice. Similarly, we recolour β into a $\chi(G)$ -colouring β' whose colour classes correspond to those of γ by recolouring each vertex at most twice. By the Renaming Lemma, α' can be recoloured into β' by recolouring each vertex at most twice. Thus, we have found a path from α to β in $\mathcal{R}_\ell(G)$ by recolouring each vertex at most 6 times. \square

5 Proof of Theorem 3

Theorem 3 follows from the following lemma.

Lemma 5. *There is no graph H on five vertices such that every H -free graph G is $(\chi(G)+1)$ -mixing.*

Proof. There are 34 graphs on 5 vertices. We partition them into four groups: Group 1 consists of the 21 graphs which are not bipartite. Group 2 consists of the 7 graphs which are bipartite and contain the co-diamond. Group 3 consists of P_5 . Group 4 consists of the remaining 5 graphs, which are $5K_1$, $K_{1,4}$, claw+ K_1 , $K_{2,3}$ and the banner, where the banner consists of the cycle C_4 on four vertices together with a vertex joined to exactly one vertex of the C_4 .

The graph B_p is bipartite and, as mentioned in Lemma 2, is co-diamond-free, and thus is H -free for any non-bipartite graph H and for any graph H containing the co-diamond. By Lemma 2, B_p has a frozen p -colouring, for every $p \geq 3$.

Theorem 2 [8] says that not every P_5 -free graph is $(\chi+1)$ -mixing.

Lemma 1 says that there are frozen colourings of the graph G_p which is $(4K_1, C_4, \text{claw})$ -free, and thus also $(5K_1, K_{1,4}, \text{claw}+K_1, K_{2,3}, \text{banner})$ -free. Thus, for every graph H in Group 4, there is an H -free graph which is not $(\chi+1)$ -mixing. \square

6 Recolouring (P_5, C_4) -free graphs

A *blow-up* of a graph G with vertices v_1, v_2, \dots, v_n is any graph H that can be obtained by substituting a clique A_i for v_i in G , one at a time for all $i \in [n]$. A *clique cutset* Q of a graph G is a clique in G such that $G-Q$ has more components than G .

A *split graph* is one whose vertex-set can be partitioned into a clique and an independent set. Equivalently, split graphs are the class of $(2K_2, C_4, C_5)$ -free graphs or the class of graphs which are both chordal and cochordal. Blázsik et al [2] gave the following characterization of $(2K_2, C_4)$ -free graphs.

Theorem 11 ([2]). *A graph G is $(2K_2, C_4)$ -free if and only if its vertex-set can be partitioned into three possibly empty sets, Q, C and I such that Q is a complete graph, I is an independent set and C (if nonempty) induces a C_5 , and C is complete to Q and anticomplete to I .*

First we consider the subclass $(2K_2, C_4)$ -free graphs and prove Theorem 7.

Proof of Theorem 7. Let γ_1 and γ_2 be any two colourings of a $(2K_2, C_4)$ -free graph G . We prove the theorem by providing a path between the two colourings of length at most $4n$. First we obtain two colourings α and β from γ_1 and γ_2 , respectively, such that α and β induce the same partition of the vertices into colour classes. We consider two cases.

Case 1: When G has an induced C_5 . By Theorem 11, $V(G)$ can be partitioned into three subsets C , Q , and I , where C induces a C_5 , Q is a clique and I is an independent set. Since C is complete to Q and anticomplete to I , we have $\chi(G) = 3 + |Q|$ and $\gamma_i(C) \cap \gamma_i(Q) = \emptyset$, for $i = 1, 2$. Let $C = \{v_1, v_2, \dots, v_5\}$ such that $v_j v_{j+1} \in E(G)$ for all $j \pmod{5}$. Since $\ell \geq \chi(G) + 1 = 4 + |Q|$, if γ_i uses three colours on C then there is a colour that does not appear on $C \cup Q$ and we recolour a vertex of C with this colour to use at least four colours on C . There exist three vertices, say v_1, v_2, v_3 , such that $\gamma_i(v_1) \neq \gamma_i(v_2) \neq \gamma_i(v_3)$ and $\gamma_i(v_1) \neq \gamma_i(v_3)$, for $i = 1, 2$. We define colourings α and β , from γ_1 and γ_2 , respectively, as follows;

$$\alpha(v) = \begin{cases} \gamma_1(v_1) & \text{if } v = v_4 \\ \gamma_1(v_2) & \text{if } v = v_5 \\ \gamma_1(v_1) & \text{if } v \in I \\ \gamma_1(v) & \text{if } v \in \{v_1, v_2, v_3\} \cup Q; \end{cases} \quad \beta(v) = \begin{cases} \gamma_2(v_1) & \text{if } v = v_4 \\ \gamma_2(v_2) & \text{if } v = v_5 \\ \gamma_2(v_1) & \text{if } v \in I \\ \gamma_2(v) & \text{if } v \in \{v_1, v_2, v_3\} \cup Q; \end{cases}$$

Case 2: When G is C_5 -free. Since G is $(2K_2, C_4, C_5)$ -free, G is a split graph. Hence, $V(G)$ can be partitioned into subsets Q and I , where Q is a maximum clique and I is an independent set. Let $Q = \{u_1, u_2, \dots, u_p\}$. We partition I into subsets I_1, I_2, \dots, I_p , where $I_j = \{v \in I \mid v \text{ is non-adjacent to } u_j \text{ and adjacent to } u_1, \dots, u_{j-1}\}$, for all $j \in [p]$. We define colourings α and β , from γ_1 and γ_2 , respectively, as follows:

$$\alpha(v) = \begin{cases} \gamma_1(v_j) & \text{if } v \in I_j \\ \gamma_1(v) & \text{if } v \in Q; \end{cases} \quad \beta(v) = \begin{cases} \gamma_2(v_j) & \text{if } v \in I_j \\ \gamma_2(v) & \text{if } v \in Q; \end{cases}$$

Therefore, in both cases, we obtain α and β from γ_1 and γ_2 , respectively, by recolouring a vertex at each step in at most n steps. Since α and β induce the same partition of the vertices into colour classes and use $\chi(G)$ colours, by the Renaming Lemma there exists a path between the colourings of length at most $2n$ in $R_\ell(G)$. Hence there exists a path between γ_1 and γ_2 in $R_\ell(G)$ of length at most $4n$. \square

A clique cutset Q in a graph G is called a *tight clique cutset* if there exists a component H of $G-Q$ which is complete to Q , and then H is called a *tight component*.

Lemma 6. *Suppose G is a graph with a tight clique cutset Q and let H be a tight component of $G-Q$. Let $V_2 = V(G) \setminus V(H)$. Let α and β be two ℓ -colourings of G such that $\alpha_{V_2} = \beta_{V_2}$. Suppose H is ℓ_1 -mixing for all $\ell_1 \geq \chi(H) + 1$. Then there exists a path between α and β in $R_\ell(G)$.*

Proof. Let $V_1 = V(H)$ and $V_2 = V(G) \setminus V(H)$. Since $\ell > \chi(G) \geq \chi(H) + |Q|$ and $\alpha(Q) = \beta(Q)$, there exist at least ℓ_1 colours that do not appear in $\alpha(Q)$. Let A be the set of all colours that does not appear in $\alpha(Q)$.

Since H is ℓ_1 -mixing, there exists a path between α_1 , the restriction of α on V_1 , and β_1 , the restriction of β on V_1 , in $R_{\ell_1}(H)$ such that every colouring in the path uses only colours in A . We know that none of the colours in A appear in $\alpha(Q)$, so we can extend every colouring in the path by colouring every vertex $v \in V_2$ with the colour $\alpha(v) = \beta(v)$. \square

Lemma 7. *Suppose G has a tight clique cutset Q . If every induced subgraph F of G is $\chi(F) + c$ -mixing, for all $c \geq 1$, then G is ℓ -mixing.*

Proof. Suppose G has a tight clique cutset Q and H is a tight component of $G-Q$. Let $V_1 = V(H)$, $V_2 = V(G) \setminus V_1$, $G_2 = G[V_2]$, and $\ell_2 \geq \chi(G_2) + 1$.

Let α be an ℓ -colouring of G and let γ be a χ -colouring of G . We shall prove that there exists a path between α and γ in $R_\ell(G)$. Since G_2 is ℓ_2 -mixing, there exists a path, say P , between α_2 , the restriction of α on V_2 , and γ_2 , the restriction of γ on V_2 , in $R_{\ell_2}(G_2)$.

Let ϵ_2 and ψ_2 be any two colourings adjacent in the path P . Note that any ℓ -colouring of G_2 can be extended to an ℓ -colouring of G .

Claim 2. *There exists a path between any two ℓ -colourings ϵ and ψ of G , whose restrictions on V_2 are ϵ_2 and ψ_2 , respectively.*

By Lemma 6, it is sufficient to prove that there exists a path between ϵ and some colouring ψ whose restriction on V_2 is ψ_2 . We may also assume that ϵ_1 , the restriction of ϵ on V_1 , uses $\chi(H)$ colours.

Since ϵ_2 and ψ_2 are adjacent in the path P , there is a unique vertex $v \in V_2$ such that $\epsilon_2(v) \neq \psi_2(v)$. If $v \notin Q$ or $\psi(v) \notin \epsilon(V_1)$, we can recolour v in ϵ with the colour $\psi(v)$ to obtain ψ .

Let $v \in Q$ and $\psi(v) \in \epsilon(V_1)$. Since $\ell > \chi(H) + |Q|$, there exists a colour, say r , that does not appear in $\epsilon(V_1 \cup Q)$. Starting with ϵ recolour every vertex in V_1 coloured $\psi(v)$ with the colour r , then recolour v with the colour $\psi(v)$ to obtain the colouring ψ . \square

Proof of Theorem 8. Let G be a connected (P_5, C_4) -free graph. The proof is by induction on the number of vertices. If there is no induced C_5 in G , then G is a chordal graph and hence $R_\ell(G)$ is connected [4]. If there is an induced C_5 in G , then by the structure of (P_5, C_4) -free graphs given in [9] there exists a blowup of C_5 , say C , and a clique Q such that C is complete to Q and C is a component of $G-Q$. Hence the proof follows from Lemma 7. \square

Note that there are other classes of graphs that admit a tight clique cutset, for example, chordal graphs.

7 Conclusion

We have proved a dichotomy theorem: The reconfiguration graph of the $(k+1)$ -colourings of a graph G , $R_{k+1}(G)$, is connected for every k -colourable H -free graph G if and only if H is an induced subgraph of P_4 or $P_3 + P_1$. It is interesting to note that these are exactly the family of H -free graphs for which the colouring problem is polynomial-time solvable [10]. Next it would be natural to do try to do the same for (H_1, H_2) -free graphs.

One might first consider graphs on four vertices. There are 11 such graphs, however, by the dichotomy result, we do not need to consider P_4 or $P_3 + P_1$. Of the remaining 36 pairs of the other 9 graphs on four vertices, the recolouring diameter is infinite for 3 pairs by Theorem 4 and for 10 other pairs by Theorem 5, and finite for $(2K_2, C_4)$ -free graphs. Further, it is easy to see that C_6 has a frozen 3-colouring. The only graphs on four vertices which are induced subgraphs of C_6 are P_4 , $P_3 + P_1$ and $2K_2$. Thus it is of particular interest to study the connectivity of the reconfiguration graph of the k -colourings of $(2K_2, H)$ -free graphs, where H is a graph with at least four vertices and is not P_4 , $P_3 + P_1$ or C_4 .

Theorem 8 states that the class of (P_5, C_4) -free graphs is ℓ -mixing, however the ℓ -recolouring diameter is still unknown. It is interesting to find other classes of (H_1, H_2) -free graphs which are ℓ -mixing and the ℓ -recolouring diameter is linear or polynomial. In [8], Feghali and Merkel asked if there is a k -colourable graph G such that $\mathcal{R}_{k+1}(G)$ is disconnected but every component of $\mathcal{R}_{k+1}(G)$ has at least two vertices. When $k = 2$, L. Cereceda *et.al* [6] proved that there exist bipartite graphs that are not 3-mixing and do not admit a frozen 3-colouring, for example C_8 . Furthermore, let H be any graph with $V(H) = \{v_1, v_2, \dots, v_n\}$ which admits a frozen

ℓ -colouring, $\ell \geq 3$. We can construct a connected graph G from H by adding vertices u_1, \dots, u_i , $1 < i \leq n$, and edges $v_j u_j$ for all $j \in [i]$. Note that $\chi(G) = \chi(H)$ and G contains H as an induced subgraph, but G is not ℓ -mixing and does not admit a frozen ℓ -colouring. We ask the following:

Question 2. *Is there a graph G which is not ℓ -mixing, $\ell \geq 4$, and does not contain an induced subgraph which admits a frozen ℓ -colouring?*

References

- [1] T. Biedl, A. Lubiw, O. Merkel. Building a larger class of graphs for efficient reconfiguration of vertex colouring. In J. Nešetřil, G. Perarnau, J. Rué, and O. Serra, editors, *Extended Abstracts EuroComb 2021*, pages 286–292, Cham, 2021. Springer International Publishing.
- [2] Z. Blázsik, M. Hujter, A. Pluhár, and Z. Tuza. Graphs with no induced C_4 and $2K_2$. *Discrete Mathematics*, 115:51–55, 1993.
- [3] M. Bonamy, N. Bousquet. Recoloring graphs via tree decompositions. *European Journal of Combinatorics*, 69:200–213, 2018.
- [4] M. Bonamy, M. Johnson, I. Lignos, V. Patel, D. Paulusma. Reconfiguration graphs for vertex colourings of chordal and chordal bipartite graphs. *Journal of Combinatorial Optimization*, 27:132–143, 2014.
- [5] P. Bonsma, L. Cereceda. Finding Paths between graph colourings: PSPACE-completeness and superpolynomial distances. *Theoretical Computer Science*, 410:5215–5226, 2009.
- [6] L. Cereceda, J. van den Heuvel, M. Johnson. Connectedness of the graph of vertex-colourings. *Discrete Mathematics*, 308:913–919, 2008.
- [7] C. Feghali, J. Fiala. Reconfiguration graph for vertex colourings of weakly chordal graphs. *Discrete Mathematics*, 343:111733, 2020.
- [8] C. Feghali, O. Merkel. Mixing colourings in $2K_2$ -free graphs. [arXiv:2108.00001](https://arxiv.org/abs/2108.00001) [math.CO], 2021.
- [9] J. L. Fouquet, V. Giakoumakis, F. Marie, and H. Thuillier. On graphs without P_5 and \bar{P}_5 . *Discrete Mathematics*, 146:33–44, 1995.
J. L. Fouquet, V. Giakoumakis, F. Marie, and H. Thuillier. On graphs without P_5 and \bar{P}_5 . *Discrete Mathematics*, 146:33–44, 1995.
- [10] D. Král’, J. Kratochvíl, Z. Tuza, G. J. Woeginger. Complexity of coloring graphs without forbidden induced subgraphs. *Lecture Notes in Computer Science*, vol. 2204, 254–262, 2001.
- [11] O. Merkel. Recolouring weakly chordal graphs and the complement of triangle-free graphs. *Discrete Mathematics*, 345:112708, 2022.
- [12] S. Olariu. Paw-free graphs. *Information Processing Letters*, 28:53–54, 1988.
- [13] M. Wrochna. Reconfiguration in bounded bandwidth and tree-depth. *Journal of Computer and System Sciences*, 93:1–10, 2018.