

(UN)KNOTTED NULLHOMOTOPIC SPHERES AND DISKS IN
4-MANIFOLDS PRODUCED VIA KNOT SURGERY.

YOUNES BENYAHIA, OLIVIERO MALECH AND RAFAEL TORRES

Abstract: For each knot $K \subset S^3$ and a 4-manifold X that satisfies certain conditions, we use Fintushel-Stern's knot surgery to construct a nullhomotopic 2-sphere $S_K \subset X \# S^2 \times S^2$ such that S_K bounds a topologically embedded handlebody, i.e., S_K is topologically isotopic to an unknotted 2-sphere contained in a 4-ball. If there is a diffeomorphism of pairs $(X \# S^2 \times S^2, S_{K_1}) \rightarrow (X \# S^2 \times S^2, S_{K_2})$, then the knots K_1 and K_2 have the same Alexander polynomial. In particular, the construction in this note yields an infinite set of topologically unknotted nullhomotopic 2-spheres in $X \# S^2 \times S^2$ and an infinite set of topologically equivalent 2-disks in $X \# S^2 \times S^2 \setminus D^4$ that are smoothly inequivalent, and for which the diffeomorphism types of the embeddings are detected by the Alexander polynomial. Examples of knotted behaviour of 2-spheres in 4-manifolds that is detected by the Reidemeister/Turaev torsion of lens spaces are also provided.

1. INTRODUCTION

Fintushel-Stern's knot surgery [15] put on display the existence of as many inequivalent smooth structures on certain 4-manifolds as there are knots in the 3-sphere with different Alexander polynomial. In [14], they showed that the latter knot invariant also detects changes on the diffeomorphism type of embeddings of homologically essential surfaces of positive genus in 4-manifolds through Seiberg-Witten theory. The purpose of this note is to point out that knot surgery also establishes an injective map between knots in S^3 with different Alexander polynomial and nullhomotopic smoothly inequivalent 2-spheres that bound a topologically embedded handlebody. The interest of our examples lies within the difficulty of finding nullhomotopic smoothly knotted and topologically unknotted 2-spheres inside closed 4-manifolds. We say that two surfaces S and S' smoothly embedded in a 4-manifold X are smoothly inequivalent if there is no diffeomorphism $f : X \rightarrow X$ such that $f(S) = S'$, and that they are topologically equivalent if there is such a homeomorphism. The fundamental group of the complement of $\nu(S)$ in X is called the 2-knot group and the signature of X is denoted by $\sigma(X)$ throughout the paper.

Theorem A. *Let X be a closed connected 4-manifold with non-zero Seiberg-Witten invariant $\text{SW}_X \neq 0$ and $b_2(X) \geq |\sigma(X)| + 6$. Suppose there is a pair of disjoint smoothly embedded 2-tori $\{T_i \hookrightarrow X : i = 1, 2\}$ of self-intersection $[T_i]^2 = 0$ for which the inclusion induced group homomorphism $\pi_1(X \setminus \nu(T_1) \sqcup \nu(T_2)) \rightarrow \pi_1(X) = \{1\}$ is an isomorphism.*

There is an infinite set

$$(1.1) \quad \{S_K : K \subset S^3\}$$

2020 *Mathematics Subject Classification.* Primary 57K45, 57R55, 57K14; Secondary 57R40, 57R52.

of 2-spheres smoothly embedded in $X\#S^2 \times S^2$ with the following properties.

- (1) For every knot $K \subset S^3$, the corresponding element in (1.1) is nullhomotopic with infinite cyclic 2-knot group $\pi_1(X\#S^2 \times S^2 \setminus \nu(S_K)) = \mathbb{Z}$, and it bounds a topologically embedded solid handlebody in $X\#S^2 \times S^2$, i.e., S_K is topologically isotopic to an unknotted 2-sphere contained in a 4-ball.
- (2) If there is a diffeomorphism of pairs

$$(1.2) \quad (X\#S^2 \times S^2, S_{K_i}) \rightarrow (X\#S^2 \times S^2, S_{K_j}),$$

then the symmetric Alexander polynomials of the knots K_i and K_j coincide, i.e., $\Delta_{K_i}(t) = \Delta_{K_j}(t)$.

- (3) There is an infinite subset of (1.1) made of pairwise smoothly inequivalent nullhomotopic 2-spheres in $X\#S^2 \times S^2$ that are topologically isotopic to the unknotted 2-sphere.
- (4) Surgery to $X\#S^2 \times S^2$ on each nullhomotopic 2-sphere in the set of Item (4) produces an infinite set $\{X_K^{\mathbb{Z}} : K \subset S^3\}$ that of smooth 4-manifolds that are all homeomorphic to $X\#S^2 \times S^2 \# S^1 \times S^3$, and whose Seiberg-Witten invariant viewed as a Laurent polynomial is $\mathcal{SW}_{X_K^{\mathbb{Z}}} = \mathcal{SW}_X \cdot \Delta_K(t)$ for $t = \exp(2[T_1])$. These inequivalent smooth structures are obtained by performing knot surgery.
- (5) There is an infinite set

$$(1.3) \quad \{D_K : K \subset S^3\}$$

made of smoothly embedded 2-disks in $X\#S^2 \times S^2 \setminus D^4$ that are pairwise topologically equivalent as well as pairwise smoothly inequivalent.

Kim [28], Kim-Ruberman [29, 30] and Hoffman-Sunukjian [27] had previously used knot surgery to study knotted surfaces in closed 4-manifolds besides Fintushel-Stern [12, Section 5]. None of their examples are nullhomotopic 2-spheres like those of Theorem A. For smooth knottings of surfaces in the 4-ball the reader is directed towards Hayden [24], Hayden-S. Kim-Miller-Park-Sundberg [25] Hayden-Kjuchukova-Krishna-Miller-Powell-Sunukjian [26], Juhász-Miller-Zemke [23] and Oba [33]. Conway-Piccirillo-Powell [11] showed that every 2-handlebody with a 3-sphere boundary contains a pair of smoothly knotted disks [11].

There is no shortage in the literature of 4-manifolds that satisfy the hypothesis of Theorem A including elliptic surfaces; see Remark 1. A minimal symplectic 4-manifold with small topology constructed by Baldridge-Kirk [7, Theorem 18] yields the following example.

Example B. *There is an infinite set of smoothly embedded nullhomotopic 2-spheres in a 4-manifold homeomorphic to $4\mathbb{C}\mathbb{P}^2 \# \overline{6\mathbb{C}\mathbb{P}^2}$ that are pairwise smoothly inequivalent and each element is topologically isotopic to the unknotted 2-sphere inside a 4-ball. There is an injective map between elements in this set and knots in the 3-sphere with different Alexander polynomials. Surgery on each of these 2-spheres yields infinitely many inequivalent smooth structures on*

$$(1.4) \quad 4\mathbb{C}\mathbb{P}^2 \# \overline{6\mathbb{C}\mathbb{P}^2} \# S^1 \times S^3,$$

that are obtained by performing Fintushel-Stern's knot surgery.

There is an infinite set of smoothly embedded 2-disks in a 4-manifold homeomorphic to $4\mathbb{C}\mathbb{P}^2 \# \overline{6\mathbb{C}\mathbb{P}^2} \setminus D^4$ that are pairwise topologically equivalent and pairwise smoothly inequivalent.

Fintushel-Stern's knot surgery and stabilization results of Akbulut [1] and Baykur [8] along with results of Sunukjian [38] on TOP isotopy classes of embeddings of surfaces immediately provide examples of homologically primitive 2-spheres embedded in $X \# S^2 \times S^2$ that are smoothly knotted and topologically unknotted as the next proposition summarizes; cf. Auckly-Kim-Melvin-Ruberman [4], Akbulut [3], [5].

Proposition C. *Let X be a closed connected 4-manifold with non-zero Seiberg-Witten invariant $SW_X \neq 0$ and that contains a smoothly embedded 2-torus $T \hookrightarrow X$ of self-intersection $[T]^2 = 0$. There is an infinite set*

$$(1.5) \quad \{S_K : K \subset S^3\}$$

made of homologically primitive and topologically isotopic 2-spheres smoothly embedded in $X \# S^2 \times S^2$ with trivial 2-knot group and for which performing surgery along each element produces an infinite set $\{X_K^{\{1\}} : K \subset S^3\}$ of pairwise non-diffeomorphic smooth structures on X that are constructed using Fintushel-Stern's knot surgery. In particular, the set (1.5) is made of pairwise smoothly inequivalent 2-spheres.

By removing a small 4-ball from $X \# S^2 \times S^2$ that intersects each element in (1.5), one obtains an infinite set of 2-disks smoothly embedded 2-disks in a punctured $X \# S^2 \times S^2$ that are pairwise topologically equivalent and pairwise smoothly inequivalent.

While it is still unknown if the 4-sphere contains topologically unknotted 2-spheres that are smoothly inequivalent, certain 4-manifolds with small topology do present this behavior as we now exemplify. This example follows from work of Fintushel-Stern [16], Baykur-Sunukjian [9], and Sunukjian [38].

Example D. *There is an infinite set of smoothly embedded primitive 2-spheres in $2\mathbb{C}\mathbb{P}^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ that are pairwise smoothly inequivalent and pairwise topologically isotopic, and whose 2-knot group is the trivial group. The corresponding statement for 2-disks smoothly embedded in a punctured $2\mathbb{C}\mathbb{P}^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ holds.*

The following result points out other knotting phenomena of 2-spheres in 4-manifolds that is detected using three-dimensional invariants, in this case either the Reidemeister or Turaev torsion of lens spaces.

Theorem E. *To each pair of lens spaces $L(p, q)$ and $L(p, q')$ there is an associated spun 4-manifold Σ_p that is a \mathbb{Q} -homology 4-sphere with finite cyclic fundamental group $\pi_1 \Sigma_p = \mathbb{Z}/p$ that contains smoothly embedded nullhomologous 2-spheres $S_{p,q}$ and $S_{p,q'}$ whose complements*

$$\Sigma_p \setminus \nu(S_{p,q}) \text{ and } \Sigma_p \setminus \nu(S_{p,q'})$$

are homotopy equivalent if and only if $q = \pm l^2 q' \pmod{p}$. Their complements are homeomorphic and there is a homeomorphism of pairs

$$(1.6) \quad (\Sigma_p, S_{p,q}) \rightarrow (\Sigma_p, S_{p,q'})$$

if and only if $q = \pm q'$ or $qq' = \pm 1 \pmod{p}$.

In particular, the nullhomologous 2-spheres are topologically equivalent if and only if the Reidemeister torsion/Turaev torsion of the associated lens spaces agree.

Work of Pao [34], Plotnick [36], Turaev [44] and the classification of lens spaces due to Reidemeister [37], Franz [17] and Whitehead [45] ([32, §2]) are the key ingredients in the proof of Theorem E.

The paper has been organized in the following way. We first manufacture infinitely many inequivalent smooth structures on $X \# S^2 \times S^2 \# S^1 \times S^3$ in Section 2.1 using Fintushel-Stern's knot surgery [15], a generalized fiber sum [20, Definition 7.1.11] with the Kodaira-Thurston manifold [43] and gluing results of their Seiberg-Witten invariants of Taubes [40, 41]. Their homeomorphism class is determined with Hambleton-Teichner's classification of closed orientable 4-manifolds with infinite cyclic fundamental group [22]. Doing surgery along the loops whose homotopy class generates the fundamental group of these smooth 4-manifolds provides us with an infinite set of pairwise inequivalent nullhomotopic 2-spheres, which is parametrized by knots in S^3 with different Alexander polynomial. A stabilization result of Akbulut [1] and Baykur [8] regarding knot surgery allows us to conclude that these 2-knots are all contained in $X \# S^2 \times S^2$. These tasks are taken care of in Section 2.2, Section 2.4 and Section 2.5. A summary is found in Section 2.6. The proof of Proposition C and Example D are given in Section 2.7, and Section 2.8, respectively. The proof of Theorem E is given in Section 2.9.

2. PROOFS.

Unless stated otherwise, we assume in the sequel that X is a closed smooth connected oriented 4-manifold such that

- $b_2(X) \geq |\sigma(X)| + 6$,
- its Seiberg-Witten invariant is non-zero $\text{SW}_X \neq 0$, and
- it contains a pair of smoothly embedded 2-tori $\{T_i \hookrightarrow X : i = 1, 2\}$ of self-intersection $[T_i]^2 = 0$ for which the inclusion induced homomorphism

$$(2.1) \quad \pi_1(X \setminus \nu(T_1) \sqcup \nu(T_2)) \rightarrow \pi_1(X) = \{1\}$$

is a group isomorphism.

Remark 1. *Symplectic 4-manifolds for which T_1 and T_2 are symplectic submanifolds are prototypes of 4-manifolds that satisfy these properties [19, 7].*

2.1. Infinitely many inequivalent smooth structures. We construct infinitely many inequivalent smooth structures on a 4-manifold with infinite cyclic fundamental group, which shares with X the same Euler characteristic, signature and second Stiefel-Whitney class.

Proposition 1. *There is an infinite set*

$$(2.2) \quad \{X_K^{\mathbb{Z}} : K \subset S^3\}$$

of pairwise non-diffeomorphic 4-manifolds such that every element is homeomorphic to the connected sum

$$(2.3) \quad X \# S^2 \times S^2 \# S^1 \times S^3$$

and the value of its Seiberg-Witten invariant seen as a Laurent polynomial is

$$(2.4) \quad \text{SW}_{X_K^{\mathbb{Z}}} = \text{SW}_X \cdot \Delta_K(t),$$

where $t = \exp(2[T_1])$.

Regarding the value of the Seiberg-Witten invariant (2.4), we are denoting by $[T_1]$ the class in $H_2(X_K^{\mathbb{Z}}; \mathbb{Z})$ that is induced by the torus $T_1 \subset X$ [15, p. 367].

Proof. Denote the Kodaira-Thurston 4-manifold by N [43], which is a T^2 -bundle over T^2 and let $T \subset N$ be a section. Every member of the set (2.2) is constructed from X by applying Fintushel-Stern's knot surgery on the 2-torus T_1 and a generalized fiber sum [20, Definition 7.1.11] with N along T_2 and T , i.e.,

$$(2.5) \quad X_K^{\mathbb{Z}} := (X \setminus (\nu(T_1) \sqcup \nu(T_2))) \cup ((S^1 \times E(K)) \sqcup (N \setminus \nu(T))),$$

where $E(K) = S^3 \setminus \nu(K)$ is the complement of a knot $K \subset S^3$. The two gluing maps in (2.5) are described as follows. Let $\{x_i, y_i\}$ be loops whose homotopy classes generate $\pi_1 T_i$, and let μ_i be a curve isotopic to $\{pt\} \times \partial D^2 \subset T_i \times D^2 = \nu(T_i)$ for $i = 1, 2$. Let $\{m_i, l_i, \mu_i\}$ be a basis for $H_1(\partial\nu(T_i))$, where m_i and l_i are the pushoffs of x_i and y_i into $\partial\nu(T_i)$, respectively. Let μ_K and λ_K be the meridian and the longitude of the knot $K \subset S^3$, and let the triple $\{[S^1], \mu_K, \lambda_K\}$ be a basis for $H_1(S^1 \times \partial E(K))$. The knot surgery part of (2.5) is a generalized fiber sum with gluing diffeomorphism

$$(2.6) \quad \varphi_K : \partial\nu(T_1) \rightarrow S^1 \times \partial E(K)$$

determined by

$$(2.7) \quad (\varphi_K)_*(m_1) = [S^1], (\varphi_K)_*(l_1) = [\mu_K], (\varphi_K)_*(\mu_1) = \lambda_K.$$

The gluing map for the generalized fiber sum in (2.5) is a diffeomorphism

$$(2.8) \quad \varphi : \partial\nu(T_2) \rightarrow \partial\nu(T)$$

that is determined by

$$(2.9) \quad (\varphi)_*(m_2) = m_T, (\varphi)_*(l_2) = l_T, (\varphi)_*(\mu_2) = \mu_T.$$

The basis $\{m_T, l_T, \mu_T\}$ for $H_1(\partial\nu(T))$ used in (2.9) arises from a canonical framing of the section T of the Kodaira-Thurston manifold N . In particular, if there is a symplectic form on X for which the 2-tori $\{T_1, T_2\}$ are symplectic submanifolds as in Remark 1, and K is a fibered knot, the maps (2.6) and (2.8) are such that there will be symplectic structure on $X_K^{\mathbb{Z}}$ [19], [20, Remark 10.3.5].

The Seifert-van Kampen theorem tells us that the fundamental group of (2.5) is $\pi_1(X_K^{\mathbb{Z}}) = \mathbb{Z}$ for every knot $K \subset S^3$ using the hypothesis (2.1); cf. [7, Section 2.2]. The Euler characteristic is $\chi(X_K^{\mathbb{Z}}) = \chi(X)$, the signature is $\sigma(X_K^{\mathbb{Z}}) = \sigma(X)$, and the second Stiefel-Whitney class $w_2(X_K^{\mathbb{Z}}) = w_2(X)$. Since we assumed that $b_2(X) \geq |\sigma(X)| + 6$, a result of Hambleton-Teichner [22, Corollary 3] (cf. [21]) allows us to conclude that $X_K^{\mathbb{Z}}$ is homeomorphic to $X \# S^2 \times S^2 \# S^1 \times S^3$ for every knot K in the 3-sphere.

Even if the diffeomorphism type of (2.5) were not determined by the description above, the 4-manifolds that are obtained in this way will have the same Seiberg-Witten invariant. The arguments used to conclude on the value (2.4) are well documented in the bibliography and are as follows; see [15, Section 2] and [12, Section 1]. We think of the construction (2.5) in two steps: first the generalized fiber sum of X and N along T_2 and T , and then an application of knot surgery using the 2-torus T_1 . The Seiberg-Witten invariant of the Kodaira-Thurston manifold is $SW_N = 1$ by a result of Taubes [40, Main Theorem] since it admits a symplectic structure. Another result of Taubes [41] and Park [35] assures that the Seiberg-Witten invariant of the generalized fiber sum $X \setminus \nu(T_2) \cup N \setminus \nu(T)$ does

not vanish. The 4-manifold (2.5) is obtained from this generalized fiber sum by applying Fintushel-Stern's knot surgery along the 2-torus $T_1 \subset X \setminus \nu(T_2) \cup N \setminus \nu(T)$. The claim regarding (2.4) follows from [15, 39]; cf. [12, Section 2 and Section 3]. \square

2.2. Framing and nullhomotopic 2-spheres in $X \# S^2 \times S^2$. Take any element $X_K^{\mathbb{Z}}$ of the infinite set (2.2), and let $\alpha \subset X_K^{\mathbb{Z}}$ be a loop whose homotopy class generates the group $\pi_1(X_K^{\mathbb{Z}}) = \mathbb{Z}$. Its tubular neighborhood is $\nu(\alpha) = S^1 \times D^3$, and we specify the choice of framing as follows. The loop α is contained in a 2-torus with trivial tubular neighborhood that is smoothly embedded in the complement of T inside N , i.e.,

$$(2.10) \quad \alpha \subset T^2 \times D^2 \subset N \setminus \nu(T) \subset X_K^{\mathbb{Z}},$$

and the framing of α is the one that is induced by the product structure of the $T^2 \times D^2$ in (2.10); see Section 2.3. Use this framing and construct the simply connected 4-manifold (cf. [13])

$$(2.11) \quad X_K := (X_K^{\mathbb{Z}} \setminus \nu(\alpha)) \cup (D^2 \times S^2),$$

and consider the belt 2-sphere

$$(2.12) \quad S_K := \{0\} \times S^2 \subset D^2 \times S^2 \subset X_K.$$

If the second Stiefel-Whitney class satisfies $w_2(X_K^{\mathbb{Z}}) \neq 0$, then the diffeomorphism type of (2.11) does not depend on the choice of framing [20, Proposition 5.2.4]. If $w_2(X_K^{\mathbb{Z}}) = 0$ on the other hand, there is a Spin-structure on $X_K^{\mathbb{Z}}$ that induces a Spin-structure on the submanifolds $X_K^{\mathbb{Z}} \setminus \nu(\alpha)$ and $\partial(X_K^{\mathbb{Z}} \setminus \nu(\alpha))$. The Spin-structure on the latter restricts to the Spin-structure on $S^1 \times \{pt\} \subset S^1 \times S^2 = \partial(X_K^{\mathbb{Z}} \setminus \nu(\alpha))$ that Spin-bounds. The gluing diffeomorphism that is used to identify the pieces in (2.11) is the identity map, and the Spin-structure on $X_K^{\mathbb{Z}} \setminus \nu(\alpha)$ matches the unique Spin-structure on $D^2 \times S^2$ and yields a Spin-structure on (2.11).

Lemma 1. *The 4-manifold X_K defined in (2.11) is homeomorphic to $X \# S^2 \times S^2$ and the 2-sphere (2.12) is nullhomotopic with infinite cyclic 2-knot group for every choice of knot $K \subset S^3$. The 4-manifold $X_K^{\mathbb{Z}}$ is recovered by applying surgery to X_K along S_K , i.e.,*

$$(2.13) \quad X_K^{\mathbb{Z}} = X_K \setminus \nu(S_K) \cup S^1 \times D^3.$$

We shall see that X_K is diffeomorphic to $X \# S^2 \times S^2$ for every knot $K \subset S^3$.

2.3. Result of surgery to the Kodaira-Thurston manifold along a homotopically essential loop. We now identify the loop involved in the surgery (2.11) within the Kodaira-Thurston manifold N [43], [6, Section 2.4], [7, Section 2]. As a building block in the construction of the infinite set $\{X_K^{\mathbb{Z}} : K \subset S^3\}$ of Proposition 1, N is the total space of a T^2 -bundle over T^2 . It can be seen as a product $N = S^1 \times Y_{D_a}$, where Y_{D_a} is the mapping torus of a Dehn twist $D_a : T^2 \rightarrow T^2$ around an embedded loop $\alpha_a \subset T^2$ for $\{\alpha_a, \alpha_b\}$ smoothly embedded loops whose homotopy classes compose a basis for $\pi_1 T^2 = \mathbb{Z}a \oplus \mathbb{Z}b$. We write $T^2 = \alpha_a \times \alpha_b$. The 3-manifold Y_{D_a} is the total space of an $\alpha_a \times \alpha_b$ -bundle over a circle α_y with $H_1(Y_{D_a}) = \mathbb{Z}b \oplus \mathbb{Z}y$. Notice that the loop α_a is homologically trivial in Y_{D_a} . Write the T^2 -bundle over T^2 structure of $N = S^1 \times Y_{D_a} = \alpha_x \times Y_{D_a}$ as

$$(2.14) \quad \alpha_a \times \alpha_b \hookrightarrow N \rightarrow \alpha_x \times \alpha_y$$

where our notation emphasizes the loops contained in the 2-torus fiber and the 2-torus section and that generate the first homology group $H_1(N) = \mathbb{Z}x \oplus \mathbb{Z}y \oplus \mathbb{Z}b$. The fiber 2-torus $\alpha_a \times \alpha_b$ is geometrically dual to the 2-torus section $T := \alpha_x \times \alpha_y$, which is the submanifold used in the construction (2.8) of inequivalent smooth structures in Proposition 1. There is another pair of geometrically dual 2-tori inside N that we denote by $\alpha_x \times \alpha_b$ and $\alpha_y \times \alpha_a$ that are disjoint from both the fiber and the section; see [6, Section 2] for a detailed description of these submanifolds.

We equip N with a symplectic structure for which the 2-torus $\alpha_x \times \alpha_b \subset N$ is a Lagrangian submanifold, and use the unique Lagrangian framing when we perform surgery along this submanifold; this is explained in detail in [6, Section 2.1]. Let m and l be the Lagrangian pushoffs of α_x and α_b , respectively, and let μ be the meridian of $\alpha_x \times \alpha_b$, i.e., a curve isotopic to $\{pt\} \times \partial D^2 \subset \partial\nu(\alpha_x \times \alpha_b)$. The triple $\{m, l, \mu\}$ is a basis for the group $H_1(\partial\nu(\alpha_x \times \alpha_b))$; see [6, Section 2.1] for further details. Define a 4-manifold

$$(2.15) \quad \hat{N} = N \setminus \nu(\alpha_x \times \alpha_b) \cup_{\varphi_0} T^2 \times D^2,$$

where the diffeomorphism

$$(2.16) \quad \varphi_0 : \partial\nu(\alpha_x \times \alpha_b) \rightarrow T^2 \times \partial D^2$$

that is used to identify the boundaries in the construction (2.15) sends the loop $\{pt\} \times \alpha_b \times \{pt\} \subset \partial\nu(\alpha_x \times \alpha_b)$ to $\{pt\} \times \{pt\} \times \partial D \subset S^1 \times S^1 \times D^2$; this is also known as a multiplicity zero log transform [20, p. 83]. That is, (2.16) is determined by $\varphi_0(l)_* = [\{pt\} \times \partial D^2]$

We identify the diffeomorphism type of a pair of 4-manifolds that are constructed from the Kodaira-Thurston manifold in the following lemma.

Lemma 2. *Let \hat{N} be the 4-manifold in (2.15) that is obtained from the Kodaira-Thurston manifold $N = S^1 \times Y_{D_a}$ by performing a torus surgery on $\alpha_x \times \alpha_b$ along the loop α_b using the gluing map (2.16). There is a diffeomorphism*

$$(2.17) \quad \hat{N} \approx T^2 \times S^2.$$

Let N^ be the 4-manifold that is obtained from the Kodaira-Thurston manifold by performing surgery along the loop α_b with respect to the framing induced by the $T^2 \times D^2$ product structure on the tubular neighborhood of the 2-torus $\alpha_x \times \alpha_b \subset N$. There is a diffeomorphism*

$$(2.18) \quad N^* \approx T^2 \times S^2 \# S^2 \times S^2.$$

Proof. The existence of the diffeomorphism (2.17) follows from a standard argument; see Baldridge-Kirk in [7, Proof Lemma 2]. The 4-manifold \hat{N} is diffeomorphic to the product $S^1 \times M^3$ of the circle with a 3-manifold M^3 that fibers over the circle with monodromy given by the composition $D_a \circ D_b^0$ of Dehn twists D_a^1 and D_b^0 along loops α_a and α_b in T^2 , respectively. The notation on the Dehn twists is explained by seeing them as a (1, 1)-Dehn surgery along α_a and a 0-surgery along α_b to the 3-torus $\alpha_y \times \alpha_a \times \alpha_b$. The resulting 3-manifold M^3 has infinite cyclic fundamental group; see [6, Section 2.2]. It is immediate to see that M^3 is diffeomorphic to $S^1 \times S^2$ and we conclude that $\hat{N} = S^1 \times (S^1 \times S^2) = T^2 \times S^2$.

We now prove the existence of the diffeomorphism (2.18). Notice first that the 4-manifold N^* splits off a copy of $S^2 \times S^2$, i.e., there is a diffeomorphism

$$(2.19) \quad N^* := N \setminus \nu(\alpha_b) \cup D^2 \times S^2 \approx X \# S^2 \times S^2,$$

where X is a closed 4-manifold with fundamental group $\pi_1 = \mathbb{Z} \oplus \mathbb{Z}$ generated by the loops α_x and α_y . The latter claim follows from the Seifert-van Kampen theorem. To see that the decomposition (2.19) holds, a well-known argument is employed and which already appears in the work of Gompf and Fintushel-Stern [13, §5]. There is a pair of geometrically dual 2-tori $\alpha_x \times \alpha_b$ and $\alpha_y \times \alpha_a$ of self-intersection zero contained in N . Surgery along α_b with the framing induced by the product structure $\alpha_b \subset \alpha_x \times \alpha_b = T^2 \times D^2$, turns the pair of 2-tori into a pair of geometrically dual 2-spheres S_1 and S_2 . These two 2-spheres form the $S^2 \times S^2$ piece in the connected sum in (2.19).

The job now is to see that X is diffeomorphic to $\hat{N} = T^2 \times S^2$ in order to conclude the existence of the diffeomorphism (2.18). The 4-manifold X is obtained by performing surgery along either of the 2-spheres S_1 or S_2 in (2.19). That is, one carves a $D^2 \times S^2$ out of N^* and caps off the boundary with a copy of $S^1 \times D^3$ in order to get rid of the $S^2 \times S^2$ piece and obtain X . We use an argument due to Moishezon to identify the diffeomorphism type of X . As a corollary of Moishezon's more general argument [31, Lemma 13], [18, proof of Lemma 3] we know that we can also build X from N by carving out the tubular neighborhood $\nu(\alpha_x \times \alpha_b) = S^1 \times S^1 \times D^2$ and gluing this 2-torus back in $N \setminus \nu(\alpha_x \times \alpha_b)$ using a diffeomorphism that sends the loop $\{pt\} \times \alpha_b \times \{pt\}$ to $\{pt\} \times \{pt\} \times \partial D \subset S^1 \times S^1 \times D^2$; a handlebody argument to show this is given by Gompf in [18, Lemma 3]. This is exactly the gluing diffeomorphism (2.16) used in the construction of \hat{N} in (2.15). We conclude that $X = \hat{N}$ and the diffeomorphism (2.17) implies that X is diffeomorphic to $\hat{N} \approx T^2 \times S^2$. We conclude that $N^* \approx T^2 \times S^2 \# S^2 \times S^2$. \square

2.4. Diffeomorphism type of the ambient 4-manifold. Results of Akbulut [1] and Baykur [8] allow us to upgrade the conclusion of Lemma 1 to cover the diffeomorphism type of X_K .

Proposition 2. *The 4-manifold X_K defined in (2.11) is diffeomorphic to the connected sum $X \# S^2 \times S^2$ for every knot $K \subset S^3$.*

Notice that the connected sum of X and $S^2 \times S^2$ can be written as

$$(2.20) \quad \begin{aligned} X \# S^2 \times S^2 &\approx (X \setminus \nu(T_2)) \cup (T^2 \times D^2 \# S^2 \times S^2) \approx \\ &\approx (X \# S^2 \times S^2 \setminus \nu(T_2)) \cup (T^2 \times D^2), \end{aligned}$$

where the gluing map

$$(2.21) \quad \varphi : \partial\nu(T_2) \rightarrow \partial(T^2 \times D^2)$$

is given by

$$(2.22) \quad (\varphi)_*(m_2) = [S^1 \times \{\theta_2\} \times \{\theta_3\}], \quad (\varphi)_*(l_2) = [\{\theta_1\} \times S^1 \times \{\theta_3\}], \quad (\varphi)_*(\mu_2) = [\{(\theta_1, \theta_2)\} \times \partial D^2]$$

for $(\theta_1, \theta_2, \theta_3) \in S^1 \times S^1 \times \partial D^2 = T^2 \times D^2$.

Proof. A deeper inspection into the constructions (2.5) and (2.11) reveals the diffeomorphisms

$$(2.23) \quad \begin{aligned} X_K &\approx (X \setminus \nu(T_1) \sqcup \nu(T_2)) \cup (S^1 \times E(K)) \sqcup (T^2 \times D^2 \# S^2 \times S^2) \approx \\ &\approx ((X \setminus \nu(T_1)) \cup (S^1 \times E(K))) \# S^2 \times S^2 \approx X_K^{\{1\}} \# S^2 \times S^2 \end{aligned}$$

where

$$(2.24) \quad X_K^{\{1\}} := X \setminus \nu(T_1) \cup S^1 \times E(K)$$

is the simply connected 4-manifold obtained from X by applying knot surgery along the 2-torus $T_1 \subset X$ with gluing diffeomorphism (2.6). The diffeomorphism (2.23) exists by Lemma 2 and (2.20). A result of Akbulut [1, Theorem 2.2] and Baykur [8, Theorem 1] says

$$(2.25) \quad X_K^{\{1\}} \# S^2 \times S^2 \approx X \# S^2 \times S^2$$

for every knot $K \subset S^3$. This concludes the proof of the proposition. \square

2.5. Topologically isotopic and unknotted 2-knots. Let us discuss the knotted behavior of the 2-spheres that were constructed in the previous sections.

Proposition 3. *Let \mathcal{K} be a collection of knots $\{K \subset S^3\}$ with different Alexander polynomial. There is an infinite set of nullhomotopic 2-spheres*

$$(2.26) \quad \{S_K : K \in \mathcal{K}\}$$

smoothly embedded in $X \# S^2 \times S^2$ that are topologically isotopic and topologically bound a solid handlebody, and for which performing surgery along each element yields the infinite set (2.2) of inequivalent smooth structures on the connected sum $X \# S^2 \times S^2 \# S^1 \times S^3$. In particular, the infinite set (2.26) consists of pairwise inequivalent nullhomotopic 2-spheres.

Every element in the set (2.26) bounds a topologically embedded solid handlebody in $X \# S^2 \times S^2$, and it is topologically unknotted.

Proof. As Lemma 1 indicates, carving out a tubular neighborhood $\nu(S_{K_i})$ from $X \# S^2 \times S^2$ and gluing back an $S^1 \times D^3$ yields a 4-manifold $X_{K_i}^{\mathbb{Z}}$ of the infinite set (2.2). Since the infinite set (2.2) is made of pairwise non-diffeomorphic 4-manifolds, this implies that the infinite set (2.26) is made of pairwise inequivalent 2-spheres. Indeed, suppose there were a diffeomorphism of $X \# S^2 \times S^2$ that takes two different elements S_{K_i} and S_{K_j} of (2.26) to each other. Applying surgery along each of these 2-spheres in a copy of $X \# S^2 \times S^2$ would then yield a pair of diffeomorphic 4-manifolds $X_{K_i}^{\mathbb{Z}}$ and $X_{K_j}^{\mathbb{Z}}$, which contradicts Proposition 1. A result of Sunukjian [38, Theorem 7.2] states that every element of (2.26) is topologically unknotted. \square

2.6. Proof of Theorem A. The infinite set of nullhomotopic 2-spheres (1.1) inside $X \# S^2 \times S^2$ is produced in Proposition 1, Section 2.2 and Proposition 2. The claim of Item (3) is proven in Section 2.5. The claim of Item (4) follows from Lemma 1, Proposition 1 and Proposition 2. We now prove the claim of Item (5) regarding the existence of the infinite set (1.3) inside a punctured $X \# S^2 \times S^2$. Remove a small 4-ball from $X \# S^2 \times S^2$ in a way that it punctures the nullhomotopic 2-sphere S_K . This way we get a 2-disk $D_K \subset X \# S^2 \times S^2 \setminus D^4$ for each element in (1.1). This produces the infinite set (1.3) of smoothly inequivalent 2-disks since the existence of a diffeomorphism of pairs

$$(X \# S^2 \times S^2 \setminus D^4, D_{K_i}) \rightarrow (X \# S^2 \times S^2 \setminus D^4, D_{K_j})$$

is equivalent to the existence of a diffeomorphism of pairs

$$(X \# S^2 \times S^2, S_{K_i}) \rightarrow (X \# S^2 \times S^2, S_{K_j})$$

for any pair of 2-spheres S_{K_i} and S_{K_j} in the infinite set of Item (4). The existence of the homeomorphism of pairs follows from a result of Conway-Powell [10, Theorem 1.3]. This concludes the proof of the theorem. \square

2.7. Proof of Proposition C: homologically essential examples. Fintushel-Stern's knot surgery [15], [12, Theorem 4] produces an infinite set of pairwise inequivalent smooth structures

$$(2.27) \quad \{X_K^{\{1\}} : K \in \mathcal{K}\}$$

on the 4-manifold X , where \mathcal{K} is a set of knots in the 3-sphere with different Alexander polynomials. There is a diffeomorphism $X_K^{\{1\}} \# S^2 \times S^2 \approx X \# S^2 \times S^2$ for any knot $K \subset S^3$ by results of Akbulut [1] and Baykur [8]. Take the factor 2-sphere $S := \{pt\} \times S^2 \subset S^2 \times S^2$ and denote by S_K the image under its canonical embedding in $X_K^{\{1\}} \# S^2 \times S^2$ for a fixed knot $K \subset S^3$. It is a homologically essential smoothly embedded 2-sphere $S_K \subset X_K^{\{1\}} \# S^2 \times S^2$ whose 2-knot group is $\pi_1(X_K^{\{1\}} \# S^2 \times S^2 \setminus \nu(S_K)) = \pi_1(X_K^{\{1\}} \# S^2 \times S^2) = \{1\}$. Surgery on S_K produces the 4-manifold $X_K^{\{1\}}$ in (2.27). In particular, the existence of a diffeomorphism of pairs $(X \# S^2 \times S^2, S_{K_i}) \rightarrow (X \# S^2 \times S^2, S_{K_j})$, would imply that the Alexander polynomial of K_i and K_j coincide. A result of Sunukjian [38, Theorem 7.1] says that the 2-spheres in (1.5) are topologically isotopic and topologically unknotted. \square

2.8. About Example D. Fintushel-Stern [16, Theorem 1.1] showed that by applying surgery to a nullhomologous 2-torus $T \subset \mathbb{C}\mathbb{P}^2 \# 3\overline{\mathbb{C}\mathbb{P}^2}$, one constructs an infinite set $\{Y_n : n \in \mathbb{Z}\}$ of pairwise non-diffeomorphic irreducible 4-manifolds, which are all homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 3\overline{\mathbb{C}\mathbb{P}^2}$. Results of Baykur-Sunukjian say that $Y_n \# S^2 \times S^2$ is diffeomorphic to $2\mathbb{C}\mathbb{P}^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ for every $n \in \mathbb{Z}$ [9, Corollary 16]. Take the factor 2-sphere $S := \{pt\} \times S^2 \subset S^2 \times S^2$ and denote by S_n the image under its canonical embedding in $Y_n \# S^2 \times S^2 = 2\mathbb{C}\mathbb{P}^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$. The remaining part of the argument proceeds verbatim to the proof given in Section 2.7. \square

2.9. Proof of Theorem E. Results of Turaev [44, Theorems 1.4, 1.5] indicate that the lens spaces $L(p, q)$ and $L(p, q')$ are homeomorphic if and only if their product with the circle $S^1 \times L(p, q)$ and $S^1 \times L(p, q')$ are homeomorphic. Consider the spun 4-manifold

$$(2.28) \quad \Sigma_p := S^1 \times (L(p, q) \setminus D^3) \cup (D^2 \times S^2),$$

which is a \mathbb{Q} -homology 4-sphere with fundamental group $\pi_1 \Sigma_p = \mathbb{Z}/p$. Results of Pao [34] and Plotnik [36] imply that there is a unique diffeomorphism type regardless of the choice of variable q in the lens space $L(p, q)$ and the framing used in (2.28). The belt 2-sphere $S_{p,q} := \{0\} \times S^2 \subset D^2 \times S^2$ in (2.28) is a nullhomologous 2-sphere with 2-knot group $\pi_1(\Sigma_p \setminus \nu(S_{p,q})) = \mathbb{Z} \times \mathbb{Z}/p$. Notice that there is a homeomorphism of pairs $(\Sigma_p, S_{p,q}) \rightarrow (\Sigma_p, S_{p,q'})$ if and only if the lens spaces $L(p, q)$ and $L(p, q')$ in the construction (2.28) are homeomorphic. The homeomorphism classes of lens spaces are detected by their Reidemeister torsion [32, §2.7] or Turaev torsion [44, Lemma 1.2]. \square

REFERENCES

- [1] S. Akbulut, *Variations on Fintushel-Stern knot surgery on 4-manifolds*, Turk. J. Math 26 (2002), 81 - 92.
- [2] S. Akbulut, *Nash homotopy spheres are standard*, in Proceedings of the 17th. Gökova Geometry-Topology Conference 2010, 135 - 144, GGT, Gökova, 2010.
- [3] S. Akbulut, *Isotoping 2-spheres in 4-manifolds*, in Proceedings of the 21st. Gökova Geometry-Topology Conference 2014, 264 - 266, GGT, Gökova, 2014.
- [4] D. Auckly, H. J. Kim, P. Melvin and D. Ruberman, *Stable isotopy in four dimensions*, Jour. London Math. Soc. 91 (2015), 439 - 463.
- [5] D. Auckly, H. J. Kim, P. Melvin, D. Ruberman and H. Schwartz, *Isotopy of surfaces in 4-manifolds after a single stabilization* Adv. in Math. 341 (2018), 609 - 615.
- [6] S. Baldrige and P. Kirk, *Symplectic 4-manifolds with arbitrary fundamental group near the Bogomolov-Miyaoka-Yau line*, J. Symplectic Geom. 4 (2006), 63 - 70.
- [7] S. Baldrige and P. Kirk, *Constructions of small symplectic 4-manifolds using Luttinger surgery*, J. Diff. Geom. 82 (2008), 919 - 940.
- [8] R. I. Baykur, *Dissolving knot surgered 4-manifolds by classical cobordism arguments*, J. Knot Theory and its Ramifications, Vol. 27, No. 05, 1871001 (2018).
- [9] R. I. Baykur and N. Sunukjian, *Round handles, logarithmic transforms and smooth 4-manifolds*, J. Topol. 6 (2013), 49 - 63.
- [10] A. Conway and M. Powell, *Embedded surfaces with infinite cyclic knot group*, to appear Geom. Topol. arXiv:2009.13461v3, 2020.
- [11] A. Conway, L. Piccirillo and M. Powell, *4-manifolds with boundary and fundamental group \mathbb{Z}* arXiv:2205.12774, 2022.
- [12] R. A. Fintushel, *Knot surgery revisited*, (English summary) Floer homology, gauge theory, and low-dimensional topology, 195 - 224, Clay Math. Proc., 5, Amer. Math. Soc., Providence, RI, 2006.
- [13] R. A. Fintushel and R. J. Stern, *A fake 4-manifold with $\pi_1 = \mathbb{Z}$ and $b^+ = 4$* , Turk J. Math 18 (1994), 1 - 6.
- [14] R. A. Fintushel and R. J. Stern, *Surfaces in 4-manifolds*, Math. Res. Lett. 4 (1997), 907 - 914.
- [15] R. A. Fintushel and R. J. Stern, *Knots, links, and 4-manifolds*, Invent. Math. 134 (1998), 363 - 400.
- [16] R. A. Fintushel and R. J. Stern, *Surgery on nullhomologous tori*, Geom. Topol. Monographs 18 (2012), 61 - 81.
- [17] W. Franz, *Abbildungsklassend und Fixpunktklassen dreidimensionaler Linseräume*, J. Reine Angew. Math. 185 (1943), 65 - 77.
- [18] R. E. Gompf, *Sums of elliptic surfaces*, J. Differential Geom. 34 (1991), 93 - 114.
- [19] R. E. Gompf, *A new construction of symplectic manifolds*, Ann. of Math. 142 (1995), 527 - 595.
- [20] R. E. Gompf and A. I. Stipsicz, *4-Manifolds and Kirby Calculus*, Graduate Studies in Mathematics, 20. Amer. Math. Soc., Providence, RI, 1999. xv + 557 pp.
- [21] I. Hambleton, M. Kreck and P. Teichner, *Topological 4-manifolds with geometrically 2-dimensional fundamental groups*, J. Topol. Anal. 1 (2009), 123 - 151.
- [22] I. Hambleton and P. Teichner, *A non-extendable hermitian form over $\mathbb{Z}[\mathbb{Z}]$* , Manuscripta Math. 93 (1997), 435 - 442.
- [23] A. Juhász, M. Miller and I. Zemke, *Transverse invariants and exotic surfaces in the 4-ball*, Geom. Topol. 25 (2021), 2963 - 3012.
- [24] K. Hayden, *Exotically knotted disks and complex curves*, arXiv: 2003.13681v2, 2020.
- [25] K. Hayden, S. Kim, M. Miller, J. Park, I. Sundberg, *Seifert surfaces in the 4-ball*, arXiv2205.15283, 2022.
- [26] K. Hayden, A. Kjukhukova, S. Krishna, M. Miller, M. Powell, N. Sunukjian, *Brunnian exotic surface links in the 4-ball*, arXiv:2106.13776, 2021.
- [27] N. R. Hoffman and N. S. Sunukjian, *Null-homologous exotic tori*, Alg. Geom. Topol. 20 (2020), 2677 - 2685.
- [28] H. J. Kim, *Knotted surfaces in 4-manifolds by knot surgery and stabilization*, Trans. Amer. Math. Soc. 371 (2019), 8405 - 8427.

- [29] H. J. Kim and D. Ruberman, *Topological triviality of smoothly knotted surfaces in 4-manifolds*, Trans. Amer. Math. Soc. 360 (2008), 5869 - 5881.
- [30] H. J. Kim and D. Ruberman, *Smooth surfaces with non-simply-connected complements*, Alg. & Geom. Topol. 8 (2008), 2263 - 2287.
- [31] B. Moishezon, *Complex surfaces and connected sums of complex projective planes*, Lecture Notes in Math. 603, Springer-Berlin, 1977.
- [32] L. Nicolaescu, *Notes on the Reidemeister torsion*, de Gruyter Studies in Mathematics 30, 2003.
- [33] T. Oba, *Surfaces in the 4-disk with the same boundary and fundamental group*, Math. Res. Lett. 27 (2020), 265 - 279.
- [34] P. S. Pao, *The topological structure of 4-manifolds with effective torus actions. I*, Trans. Amer. Math. Soc 18 (1943), 12 - 20.
- [35] B. D. Park, *A gluing formula for the Seiberg-Witten invariant along T^3* , Michigan Math. J. 50 (2002), 593 - 611.
- [36] S. P. Plotnik, *Equivariant intersection forms, knots in S^4 , and rotations in 2-spheres*, Trans. Amer. Math. Soc. 296 (1986), 543 - 575.
- [37] K. Reidemeisterin, *Homotopieringe und Linseräume*, Hamburger Abhandlungen 11 (1935).
- [38] N. S. Sunukjian, *Surfaces in 4-manifolds: concordance, isotopy, and surgery*, Int. Math. Res. Not. 17 (2015), 7950 - 7978.
- [39] N. S. Sunukjian, *A note on knot surgery*, J. Knot Theory Ramifications 24 (2015), no. 9, 1520003, 5 pp.
- [40] C. H. Taubes, *The Seiberg-Witten invariants and symplectic forms*, Math. Res. Lett. 1 (1994), 809 - 822.
- [41] C. H. Taubes, *The Seiberg-Witten invariants and 4-manifolds with essential tori*, Geom. Topol. 5 (2001), 441- 519.
- [42] C. H. Taubes, *The geometry of the Seiberg-Witten invariants*, in Surveys in differential geometry. Vol. III, Cambridge, MA, 1996, 299 - 339, Int. Press, Boston, MA.
- [43] W. P. Thurston, *Some simple examples of symplectic manifolds*, Proc. Amer. Math. Soc. 55 (1976), 467 - 468.
- [44] V. G. Turaev, *Towards the topological classification of geometric 3-manifolds*. Topology and geometry, Rohlin Seminar, 291 - 323, Lecture Notes in Math., 1346, Springer, Berlin, 1988.
- [45] G. H. Whitehead, *On incidence matrices, nuclei and homotopy types*, Ann. of Math. 42 (1941), 1197 - 1237.

SCUOLA INTERNAZIONALE SUPERIORI DI STUDI AVANZATI (SISSA), VIA BONOMEA 265, 34136, TRIESTE, ITALY

Email address: {ybenyahi, omalech, rtorres}@sisssa.it