

## SIGNED CLASP NUMBERS AND FOUR-GENUS BOUNDS

CHARLES LIVINGSTON

ABSTRACT. There exist knots having positive and negative four-dimensional clasp numbers zero but having four-genus, and hence clasp number, arbitrarily large. Such examples were first constructed by Allison Miller, answering a question of Juhász-Zemke. Further examples are constructed here, complementing those of Miller in that they include examples of infinite order in concordance in addition to elements of order two.

For a knot  $K \subset S^3$ , the four-dimensional clasp numbers  $c(K)$ ,  $c^+(K)$ , and  $c^-(K)$ , are the minimum number of double points, positive double points, and negative double points, respectively, among all generically immersed disks in  $B^4$  bounded by  $K$ . A simple geometric construction shows that the four-genus satisfies  $g_4(K) \leq c(K)$ .

It is relatively easy to see that  $c(K) \geq c^+(K) + c^-(K)$ ; for the figure eight knot,  $c(K) = 1$  and  $c^\pm(K) = 0$ . Juhász-Zemke [9] asked whether the difference  $c(K) - (c^+(K) + c^-(K))$  can be arbitrarily large. An affirmative answer was given by Allison Miller in [10]. For every integer  $g$ , Miller constructed strongly negative amphicheiral knots that satisfy the conditions that  $c^+(K) = c^-(K) = 0$  and  $g_4(K) \geq g$ . Here we use two-bridge knots to provide further examples. We also show that the two-bridge knot  $B(13, 2)$  has  $c^\pm(B(13, 2)) = 0$  and that  $g_4(nB(13, 2))$  goes to infinity as  $n$  increases.

## 1. FOUR-GENUS BOUNDS FROM CASSON-GORDON INVARIANTS

For a knot  $K$ , let  $M(K)$  denote its 2-fold branched cover and let  $\chi: H_1(M(K)) \rightarrow \mathbb{Z}_p$  be a homomorphism, where  $p$  is a prime number. In [3], Casson and Gordon defined two rational-valued invariants,  $\sigma(K, \chi)$  and  $\sigma_1\tau(K, \chi)$ , and proved that  $\sigma_1\tau(K, \chi)$  provides an obstruction to a knot being slice. The invariant  $\sigma(K, \chi)$  has the advantage of being more computable in some situations, making the following result useful.

**Theorem 1.** *If  $M(K)$  is a lens space, then for all  $\chi$ ,  $|\sigma(K, \chi) - \sigma_1\tau(K, \chi)| \leq 1$ .*

A theorem of Gilmer [6, Theorem 1] relates the values of  $\sigma_1\tau(K, \chi)$  to the four-genus. In general, the linking form on  $M(K)$  provides an identification of  $H_1(M(K))$  with  $\text{hom}(H_1(M(K)), \mathbb{Q}/\mathbb{Z})$ . In the case that  $H_1(M(K)) \cong (\mathbb{Z}_p)^n \oplus T$ , where the order of  $T$  is prime to  $p$ , this gives an identification of  $\text{hom}(H_1(M(K)), \mathbb{Z}_p)$  with  $(\mathbb{Z}_p)^n \subset H_1(M(K))$ . That identification is used in the following statement.

**Theorem 2.** *Let  $K$  be knot for which  $H_1(M(K)) \cong (\mathbb{Z}_p)^n \oplus T$ , where the order of  $T$  is prime to  $p$ . If  $g_4(K) \leq n/2$  and the classical signature of  $K$  satisfies  $\sigma(K) = 0$ , then there is a subspace  $\mathcal{M} \subset (\mathbb{Z}_p)^n \subset H_1(M(K))$  of dimension at least  $(n - 2g_4(K))/2$  such that for all  $\chi \in \mathcal{M}$ ,*

$$|\sigma_1\tau(K, \chi)| \leq 4g_4(K).$$

## 2. EXAMPLE 1

We consider the two-bridge knots  $B_m = B(2m, 2m)$ . The paper [3] provides the necessary background about such knots and their lens space covers, as do standard knot theory references [2, 11]. For  $m = 1$ , this is the figure eight knot. In this section we prove the following theorem. In general,  $M(B_m) = L(a, b)$ , where  $\frac{a}{b} = 2m + \frac{1}{2m}$ .

**Theorem 3.** *For all  $m$ ,  $B_m$  is negative amphicheiral and has  $c^+(B_m) = c^-(B_m) = 0$ . For every  $g$ , there is a finite set of integers  $m_i$  such that  $g_4(\#_i B_{m_i}) \geq g$ .*

The knot  $B_m$  bounds a Seifert surface built from  $B^2$  by adding two bands, one with  $m$  positive full twists and one with  $m$  negative full twists. It can be unknotted with either  $m$  negative crossing changes or  $m$

---

This work was supported by a grant from the National Science Foundation, NSF-DMS-1505586.

positive crossing changes. Thus,  $c^+(B_m)$  and  $c^-(B_m)$  are both 0. A diagram makes it evident that it is negative amphicheiral. Thus, our focus is on the four-genus.

The two-fold branched cover  $M(B_m)$  is the lens space  $L(4m^2 + 1, 2m)$ . It follows from [1] that  $M(B_m)$  is given by surgery on the Hopf link with surgery coefficients  $2m$  and  $-2m$  and surgery coefficient matrix

$$B_m = \begin{pmatrix} 2m & 1 \\ 1 & -2m \end{pmatrix}.$$

We will restrict to the case that  $m = 5k + 1$  so that  $M(B_m) = L(5(20k^2 + 8k + 1), 10k + 2)$ . We then further restrict to the case of  $k \equiv 0 \pmod{5}$  to ensure that  $H_1(M(B_m)) = \mathbb{Z}_5 \oplus \mathbb{Z}_n$  where  $\gcd(5, n) = 1$ . This ensures that every  $\chi: H_1(M(B_m)) \rightarrow \mathbb{Z}_5$  is of the form  $r\chi_1$  for some fixed  $\chi$ . We write  $r\chi_1 = \chi_r$ .

We wish to bound the values of  $\sigma_1\tau(B_m, \chi_r)$ . Since  $\sigma_1\tau(K, \chi) = \sigma_1\tau(K, -\chi)$  for all  $\chi$  and  $\sigma_1\tau(K, \chi_0) = 0$ , we can restrict our attention to  $\chi_1$  and  $\chi_2$ . Since  $B_m \# B_m = B_m \# -B_m$  is slice, the additivity of  $\sigma_1\tau$ , proved in [7], along with the fact it obstructs sliceness implies that  $\sigma_1\tau(B_m, \chi_1) = -\sigma_1\tau(B_m, \chi_2)$ . By changing indices if needed, we can assume that  $\sigma_1\tau(B_m, \chi_1) \geq 0$ .

In [3] the value of  $\sigma(K, \chi)$  for two-bridge knots was given in terms of trigonometric sums or in terms of a count of lattice points in a triangle. We use a simpler formula, first presented in [5, Remark (b), page 70] (see also [4, 8]), which implies that for the map  $\chi: H_1(M(B_m)) \rightarrow \mathbb{Z}_5$  taking value 2 on the first meridian and 1 on the second, we have

$$\sigma(B_m, \chi_1) = \frac{2}{25} (2 \ 1) \begin{pmatrix} 2m & 1 \\ 1 & -2m \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} - 1 = \frac{8k+1}{5}.$$

(We leave it to the reader to check that this  $\chi$  is well-defined, using the fact that  $m$  is of the form  $5k + 1$ .) It follows from Theorem 1 that the values of  $\sigma_1\tau(B_m, \chi_1)$  goes to infinity as  $m$  becomes large. Thus we can select a subset of  $\{B_m\}$ , which we denote  $\{C_k\}$ , with  $C_1 = B_1$  and such that for  $k > 1$ :

$$(1) \quad \sigma_1\tau(C_k, \chi_1) > \sum_{i < k} \sigma_1\tau(C_i, \chi_1) + 4k.$$

Since  $\sigma_1\tau(C_i, \chi_r)$  is maximized with  $r = 1$ , this gives the following inequality, in which the  $r_i$  are arbitrary.

$$(2) \quad \sigma_1\tau(C_k, \chi_1) > \sum_{i < k} \sigma_1\tau(C_i, \chi_{r_i}) + 4k.$$

For a given  $g$ , consider the connected sum  $C = \#_{k=1}^{4g} C_k$ . The 5-torsion in  $H_1(M(C))$  is  $(\mathbb{Z}_5)^{4g}$ . If  $g_4(C) \leq g$ , then by Theorem 2 there is a subgroup  $\mathcal{M} \subset (\mathbb{Z}_5)^{4g}$  of rank at least  $g$  such that for all  $\chi \in \mathcal{M}$ ,  $|\sigma_1\tau(C, \chi)| \leq 4g$ . Given the ranks, we can assume that some such  $\chi$  is nonvanishing on the summand corresponding to some  $C_i$ , with  $i \geq g$ . We let  $a$  be the maximal such  $i$ . By taking a multiple if need be, we can assume that the restriction of  $\chi$  to  $C_a$  is  $\chi_1$ . Then by Equation 2,  $\sigma_1\tau(C, \chi) \geq 4a > 4g$ , giving the desired contradiction.

### 3. EXAMPLE 2

Our second set of examples are not amphicheiral, but have the property that  $g_4(nK)$  goes to infinity as  $n$  goes to infinity. We let  $K_1 = B(13, 2)$ ; this is the three-twisted double of the unknot,  $D_+(U, 3)$ . Its Seifert form and a presentation of the homology of  $M(K_1)$  are given by

$$V = \begin{pmatrix} -1 & 0 \\ 1 & 3 \end{pmatrix}, \quad V + V^T = \begin{pmatrix} -2 & 1 \\ 1 & 6 \end{pmatrix}.$$

**Theorem 4.** *Let  $K_1 = B(13, 2) = D_+(U, 3)$ . Let  $K_n = \#_n K_1$  be the  $n$ -fold connected sum. Then  $c^+(K_n) = 0 = c^-(K_n)$  and for  $n$  even,  $g_4(K_n) \geq n/82$ .*

The rest of this section presents the proof of Theorem 4.

The clasp in the doubled knot can be removed by changing a positive crossing to negative, yielding the unknot. The three-twisted band in the standard genus one Seifert surface can be converted to a two-twisted band by changing a negative crossing to a positive crossing, yielding the Stevedore's knot,  $6_1 = D_+(U, 2)$ , which is a ribbon knot. This implies that  $c^+(K_1) = 0 = c^-(K_1)$ . The signed clasp numbers are subadditive, so  $c^+(K_n) = 0 = c^-(K_n)$  for all  $n$ . All that remains is to provide the lower bound on  $g_4(K_n)$ . Henceforth, we assume  $n$  is even.

Using either the trigonometric sum from [3] or the matrix formula from [5] used above, one computes that the set of values of  $\sigma(K_1, \chi_r)$  that arise for the two-bridge knot  $B(13, 2)$  are, for  $1 \leq r \leq 6$ , as listed below; the values for  $7 \leq r \leq 12$  repeat these.

$$\left\{ -\frac{9}{13}, -\frac{23}{13}, -\frac{29}{13}, -\frac{27}{13}, -\frac{17}{13}, \frac{1}{13} \right\}.$$

Using Theorem 1 this yields two estimates, which we state in a lemma.

**Lemma 5.** *There exists a value of  $r$ , which we denote  $r_m$ , such that  $\sigma_1\tau(K_1, \chi_{r_m}) \leq -\frac{16}{13}$ . For all  $r$  we have  $\sigma_1\tau(K_1, \chi_r) \leq \frac{14}{13}$ .*

The proof of Theorem 4 is completed with an application of Theorem 2. We assume that  $g_4(K_n) < n/82$ . In particular,  $g_4(K_n) < n/2$ , so Theorem 2 can be applied.

**Lemma 6.** *There exists an element vector  $\vec{c} \in \mathcal{M} \subset (\mathbb{Z}_{13})^n$  for which  $(n - 2g_4(K_n))/2$  of its coordinates equal  $r_m$ .*

*Proof.* The subspace  $\mathcal{M}$  contains at least  $k = (n - 2g_4(K_n))/n$  independent vectors. Use these to form a  $k \times n$  of rank  $k$ . Some  $k \times k$  submatrix is nonsingular. That submatrix can be reduced to the identity using row operations. Summing the rows then yields a vector in  $\mathcal{M}$  with  $k$  of its coordinates all equal to 1. This vector can then be multiplied by  $r_m$  to form  $\vec{c}$ .  $\square$

Applying the additivity of  $\tau$  to the bound given in Theorem 2 along with the bounds of Lemma 5 yields

$$\frac{16(n - 2g_4(K_n))}{26} - \frac{14}{13} \left( n - \binom{n - 2g_4(K_n)}{2} \right) \leq 4g_4(K_n).$$

An algebraic simplification yields  $n \leq 82g_4(K_n)$ , contradicting our assumption that  $g_4(K_n) < n/82$ .

**Stronger examples.** The knot  $K_1$  used above is the first in the family of doubles of torus knots:  $J_n = D_+(T(n, n+1), n(n+1)+1)$ . We leave it to the reader to check that  $D_+(T(n, n+1), n(n+1))$  is slice. Using these examples one obtains much better bounds than  $n/82$ ; it appears that for large  $n$  the bound approaches  $n/2$ .

*Acknowledgements.* I appreciate the valuable comments provided by Pat Gilmer.

#### REFERENCES

- [1] Selman Akbulut and Robion Kirby, *Branched covers of surfaces in 4-manifolds*, Math. Ann. **252** (1979/80), no. 2, 111–131.
- [2] Gerhard Burde, Heiner Zieschang, and Michael Heusener, *Knots*, extended, De Gruyter Studies in Mathematics, vol. 5, De Gruyter, Berlin, 2014.
- [3] A. J. Casson and C. McA. Gordon, *Cobordism of classical knots*, à la recherche de la topologie perdue, 1986, pp. 181–199. With an appendix by P. M. Gilmer.
- [4] David Cimasoni and Vincent Florens, *Generalized Seifert surfaces and signatures of colored links*, Trans. Amer. Math. Soc. **360** (2008), no. 3, 1223–1264.
- [5] Patrick M. Gilmer, *Configurations of surfaces in 4-manifolds*, Trans. Amer. Math. Soc. **264** (1981), no. 2, 353–380.
- [6] ———, *On the slice genus of knots*, Invent. Math. **66** (1982), no. 2, 191–197.
- [7] ———, *Slice knots in  $S^3$* , Quart. J. Math. Oxford Ser. (2) **34** (1983), no. 135, 305–322.
- [8] Patrick M. Gilmer and Stepan Yu. Orevkov, *Signatures of real algebraic curves via plumbing diagrams*, J. Knot Theory Ramifications **27** (2018), no. 3, 1840003, 33.
- [9] Andras Juhasz and Ian Zemke, *New heegaard floer slice genus and clasp number bounds*, arXiv:2007.07106 (2020).
- [10] A. N. Miller, *Amphichiral knots with large 4-genus*, Bull. of the London Math. Soc. **54** (2022), no. 2, 624–634.
- [11] Dale Rolfsen, *Knots and links*, Publish or Perish, Inc., Berkeley, Calif., 1976. Mathematics Lecture Series, No. 7.

CHARLES LIVINGSTON: DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405  
*Email address:* livingst@indiana.edu