

# On-Site Potential Creates Complexity in Systems with Disordered Coupling

I. Gershenzon,<sup>1</sup> B. Lacroix-A-Chez-Toine,<sup>2,1</sup> O. Raz,<sup>1</sup> E. Subag,<sup>3</sup> and O. Zeitouni<sup>3</sup>

<sup>1</sup>*Department of Physics of Complex Systems, Weizmann Institute of Science, Rehovot 7610001, Israel*

<sup>2</sup>*Department of Mathematics, Kings College London, London WC2R 2LS, United Kingdom*

<sup>3</sup>*Department of Mathematics, Weizmann Institute of Science, Rehovot 76100, Israel*

We calculate the average number of critical points  $\bar{\mathcal{N}}$  of the energy landscape in a spin system with disordered two-body interactions and an on-site potential. Without the on-site potential,  $\bar{\mathcal{N}}$  is at most linear in the system size. Nonetheless, we find that introducing a weak on-site potential can increase  $\bar{\mathcal{N}}$  to exponential in system size. Our results begin to explain the non-linearity driven transition from trivial to complex energy landscapes and are of relevance to glassy systems, nonlinear oscillator networks and many-body interacting systems.

*Introduction* The interplay between coupling, disorder and non-linearity is of interest in diverse areas of science. In physics, notable examples occur in spin and structural glasses [1–6], many-body localisation [7, 8], nonlinear wave propagation in disordered media [9–14], “dirty” superconductors [15, 16], coupled oscillator networks [17–20], atomic spin gases [21, 22] and in other systems [23–28]. Non-linearity, coupling and disorder are also commonly found in sociological models [29–31], epidemiology [32–36], ecological systems [37–39], computer-science [40–43] and many others [44].

The energy of disordered, non-linear complex systems commonly exhibits multiple critical point. The number of critical points, their classification, and their distribution in space can profoundly influence both static and dynamic properties of the system [2, 25, 38, 45–53]. Of particular interest is the scaling of the disorder averaged number of critical points with the system size [2]. This is characterized by the (annealed) complexity,

$$\Sigma = \lim_{N \rightarrow \infty} \frac{1}{N} \log \bar{\mathcal{N}}, \quad (1)$$

where  $\bar{\mathcal{N}}$  is the disorder averaged number of fixed points. Early works used replica methods to derive approximate expressions for  $\Sigma_{\text{tot}}$  in spin-glass models [1, 2, 52, 54–57] and for disordered nonlinear optical systems [9, 11]. Recently, rigorous random matrix methods were applied to count critical points in many models, including spin glasses [46, 58–67], ecological systems [38], neural networks [42, 68] and others [47, 69].

Several works studied the complexity in confined disordered models, where a global confining potential term is added to a disordered energy landscape [58, 59, 65, 70–72]. Others studied models with mean-field coupling between nonlinear (soft) spins [55]. However, in many physical problems one is interested in coupled many body systems with an on-site (nonlinear) potential.

In this work, we study the annealed complexity of a disordered bilinear (“two body interaction”) potential summed with weak single-particle nonlinear potentials (“on-site potential”) given in Eq. 2. This study is motivated by the following: The bilinear form potential has at most  $N$  critical points, thus zero complexity. On the other hand, on-site potentials can lead to high complexity

landscapes, e.g., for a double-well potential. It is therefore natural to ask: at which ratio between the two terms does the energy landscape become complex? To answer this, we derive a perturbative expression of the annealed complexity in orders of the relative strength of the on-site potential.

We find that in our model even a weak on-site term can lead to non-zero complexity, namely an exponential number of fixed points. In contrast, when the potential is close to being linear (i.e. an external magnetic field) the total complexity is zero to leading order. As the potential is gradually changed from non-linear to linear, a sharp transition occurs between the *trivial phase* with zero complexity and the *complex phase* with positive complexity. Technically, the calculations are facilitated by deriving a result on the disorder average of the modulus of a matrix determinant, where the matrix is the sum of a random matrix and a small deterministic diagonal matrix [65].

The model analysed in this work is inspired by the form of the energy function of the discrete nonlinear Schrodinger equation [73] as well as by “soft” spin models [55]:

$$H(\sigma) = \sum_{i,j=1}^N J_{ij} \sigma_i \sigma_j + \kappa \sum_{i=1}^N u(\sigma_i) \equiv \phi(\sigma) + \kappa U(\sigma) \quad (2)$$

where  $\sigma = (\sigma_1, \dots, \sigma_N)$  is the vector whose components  $\sigma_i$  are continuous real valued “spin” DOFs constrained to the  $N$ -sphere i.e.  $\sum_i \sigma_i^2 = N$ ,  $J_{ij}$  is a random coupling matrix modelling disordered coupling,  $u(\sigma_i)$  is a deterministic “on-site” nonlinear potential (with bounded derivatives), and  $\kappa$  is the deterministic potential strength. The stochastic part of  $H$  is denoted by  $\phi(\sigma)$  and the deterministic one by  $\kappa U(\sigma)$ . We choose the coupling matrix  $J$  to be distributed according to the Gaussian orthogonal ensemble [74] - a disordered mean-field coupling. This choice along with the spherical constraints allows for the identification of the coupling term in  $H$  with the 2-spin spherical model [60].

To evaluate the average total number of fixed points of the model we use the Kac-Rice formula on the sphere

(as given in [60]):

$$\overline{\mathcal{N}}_{\text{tot}} = \int_{\sqrt{N}\mathbb{S}^{N-1}} d\sigma_{N-1} \overline{|\det(\nabla^2 H)| \delta(\nabla H)} \quad (3)$$

where  $\nabla H$  and  $\nabla^2 H$  are the covariant derivative and Hessian matrix of  $H$  on the sphere, respectively. In what follows we evaluate the annealed total complexity  $\Sigma_{\text{tot}}$  defined in (1) in the perturbative regime of weak on site potential, i.e.  $\kappa \ll 1$ .

*Main Results* We calculate the total annealed complexity  $\Sigma_{\text{tot}}$ , and the annealed complexity of fixed extensive index  $k = Na$  (number of negative eigenvalues of the Hessian)  $\Sigma_a$  as a functional of the potential  $u(x)$ . The complexity  $\Sigma_{\text{tot}}$  is given by:

$$\Sigma_{\text{tot}}[u] = \kappa^2 \max(0, \Theta[u]) + O(\kappa^3), \quad (4)$$

with

$$\Theta[u] = \frac{1}{4} \int dx \frac{e^{-x^2/2}}{\sqrt{2\pi}} (u''(x)^2 - u'(x)^2). \quad (5)$$

For simplicity, we denote by  $x$  the argument of the on-site potential  $u$ . The complexity is second order in  $\kappa$  and has two distinct solution branches. To leading order, the added complexity is zero in one of these branches, and depends on the potential  $u(x)$  in the other branch. The dominant branch is the one with the largest added complexity. The choice of  $u(x)$  dictates the sign of  $\Theta[u]$  and consequently which branch dominates.

The complexity of fixed extensive index  $k = aN$  (with  $a \in (0, 1)$ ) is given by:

$$\Sigma_a[u] = \kappa^2 \left(1 - \frac{\eta^2(a)}{4}\right) \Theta[u] + O(\kappa^3) \quad (6)$$

where  $\eta(a)$  is the solution of:

$$\frac{1}{2\pi} \int_{-2}^{\eta(a)} dx \sqrt{4 - x^2} = a. \quad (7)$$

The dependence of  $\Sigma_a[u]$ , to leading order in  $\kappa$ , on the index  $a$  is plotted in Fig.(1) for the two solution branches discussed above. In the positive complexity branch most critical points are saddle points of index  $N/2$ , i.e. with  $N/2$  positive directions. This complexity decreases smoothly until reaching zero for critical points with  $k = o(N)$  or  $k = N - o(N)$  where the little  $o$  notation is used. In the zero total complexity branch, only saddle points of index  $k = o(N)$  or  $k = N - o(N)$  typically exist.

Although our main motivation is to analyze perturbations of the 2-spin spherical model, the same analysis can be readily applied for a general  $p$ -spin spherical model. The average total complexity  $\Sigma_{\text{tot}}^{(p>2)}$  reads in that case

$$\Sigma_{\text{tot}}^{(p>2)}[u] = \frac{1}{2} \log(p-1) + \kappa^2 \Theta^{(p)}[u] + O(\kappa^3) \quad (8)$$




FIG. 1. The complexity of critical points of a given index  $\Sigma_a$  as a function of the index scaled by  $N$ . The complexity is normalized by its maximal value in absolute value. In blue,  $\Sigma_a$  is the plotted vs.  $a$  in the case where  $\Theta[u]$  is positive. In red, we depict the case where  $\Theta[u]$  is negative

where  $\Theta^{(p)}[u]$  appears in Eq. (9) of the SI.

Note that the unperturbed complexity is consistent with [60]. As in the 2-spin spherical model, the correction to the total complexity is  $O(\kappa^2)$ . However, there is only a single solution branch and the total complexity is always positive (in the regime of weak  $\kappa$ ). Moreover, the on-site potential can either increase or decrease the total complexity depending on its form, in contrast to the 2-spin spherical model, where it can only be increased. Specifically, we also recover the results of [72] in the regime of small external magnetic field. In the SI Eq. (9) we also provide an expression for the complexity  $\Sigma_a^{(p>2)}[u]$  of the  $p$ -spin model with extensive fixed index  $k = Na$  for any integer  $p$ .

Returning to our main model (i.e.  $p = 2$ ), let us consider the physical interpretation of the system's state in each of the branches. We start by identifying the families of potentials which give rise to each of the branches. We consider a general case of a confining potential given by a power series, convergent for all  $x \in \mathbb{R}$ :

$$u(x) = \sum_{n=1}^{\infty} a_n x^n \quad (9)$$

$\Theta[u]$  for this power series is given by:

$$\Theta \left[ \sum_{n=1}^{\infty} a_n x^n \right] = -\frac{a_1^2}{4} - a_1 \sum_{n \geq 3} A_n a_n + \sum_{n \geq 3} B_n a_n^2 + \sum_{m \geq 3} \sum_{n \geq 3} D_{n,m} a_m a_n \quad (10)$$

where  $A_n, B_n, D_{m,n} > 0$ , and their expressions can be found in the SI (appendix D). Note that, as expected,




FIG. 2. The complexity  $\Sigma_{\text{tot}}$  second order correction as a function of the magnetic field strength, scaled by the critical strength, for the case of an on-site potential with  $\overline{u'_0} = 0$

the term  $a_2$  does not affect the complexity since it corresponds to a constant function on the sphere. We draw the following conclusions from this expression: (i) if  $a_1 = 0$ , then  $\Theta[u]$  is positive (see proof in SI) and thus the total complexity is positive as well (see 4). (ii) If the potential is linear (only  $a_1$  is nonzero) then  $\Theta[u]$  is negative and the total complexity is zero to leading order.

Let us consider the case where the external field component  $a_1$  is strong enough s.t the complexity is zero to leading order. In this regime, our results are similar to those in [70, 72] where “trivial topology” of the landscape was found under a sufficiently strong external magnetic field. In the trivial topology phase, the energy landscape has only two critical points (a minimum and a maximum) [72]. Our results for the complexity of a given index  $\Sigma_a$ , in this case, are consistent with this picture: the complexity is non-negative only for indices  $k = o(N)$  (or  $k = N - o(N)$ ), which represent critical points that are a minimum (maximum) in most directions. From the perspective of the spin value distribution, we show (see eq. (59) in SI) that the distribution is skewed in the direction of the applied magnetic field s.t. for  $u(x) = a_1 x$  this leads to an  $O(\kappa)$  shift in the disorder averaged magnetization  $\bar{\sigma} \sim \kappa a_1$ . This shift directly shows the alignment of the spins along the external field at the maxima or minima of the model.

Let us now focus on the case motivating this work, the addition of an on-site potential with  $a_1 = 0$ , namely no magnetic field component. As stated above, in this case the addition to the complexity is always positive. The added complexity grows like  $\kappa^2$ . Somewhat surprisingly, this means that the energy landscape has an exponential number of critical points even for small on-site potential, and most of them – irrespective of the form of  $u(x)$  – are saddles with  $a = \frac{1}{2}$ .

Having explored both system states, we now turn to discuss the transition between them. As shown in (10), driving the system between the states can be achieved by changing the magnetic field strength  $a_1$ . Assume that for a specific potential  $u_0(x)$  the added complexity  $\Theta[u_0]$  is

positive, e.g. any potential with  $u'_0(0) = 0$ . Next, we add an external field of strength  $h$  s.t.  $u(x) = u_0(x) + hx$ . A simple calculation shows:

$$\Theta[u] = \Theta[u_0] - \frac{h^2}{4} - \frac{h}{2} \int \frac{dx}{\sqrt{2\pi}} e^{-x^2/2} u'_0(x). \quad (11)$$

It is seen that for large enough field strength, i.e.  $h > h_{c+}$  or  $h < h_{c-}$ ,  $\Theta[u]$  drops below zero. At  $h_c$  there is a transition to the zero complexity branch.  $h_c$  is found by equating the above expression to zero:

$$h_{c\pm} = \pm \sqrt{\overline{u'_0} + 4\Theta[u_0]} - \overline{u'_0} \quad (12)$$

where the overbar denotes averaging w.r.t the standard Gaussian distribution. It is also seen that the form of this transition is independent of the form of  $u_0(x)$ , and that the critical exponent of the complexity is unity.

*Derivation Outline* The full derivation of our results is quite lengthy and technical. We therefore provide only the outline of the calculation of the complexity defined in (1) for the  $p = 2$  case. The full derivation for this and other cases is given in the SI. We start the calculation of (3) by evaluating the disorder average  $|\overline{\det(\nabla^2 H)}| \overline{\delta(\nabla H)}$  at a fixed position  $\sigma$  on the sphere. The average is taken over the joint distribution of the covariant Hessian and gradient of  $\phi(\sigma)$ . First, we use the independence of  $\nabla\phi$  and  $\nabla\phi^2$  to factor out the expectation as follows

$$|\overline{\det(\nabla^2 H)}| \overline{\delta(\nabla H)} = \overline{\delta(\nabla H)} |\overline{\det(\nabla^2 H)}|. \quad (13)$$

To evaluate the expectation values in Eq. (13) we start by writing the distribution of the random matrix  $\nabla^2\phi$  and  $\nabla\phi$  (see SI appendix A). Equipped with this pdf, the integral  $\overline{\delta(\nabla H)}$  is easily carried out to give:

$$\overline{\delta(\nabla H(\sigma))} = \left(\frac{1}{2\pi p}\right)^{N/2} \exp\left(-\frac{\kappa^2}{2p} \|\nabla U(\sigma)\|^2\right). \quad (14)$$

The evaluation of  $|\overline{\det \nabla^2 H(\sigma)}|$  is more involved and constitutes the main technical contribution of this work. The challenge is due to  $\nabla^2 H$  being a sum of non-commuting matrices: the random matrix  $\nabla^2\phi$  and the diagonal deterministic matrix  $\kappa\nabla^2 U$  (see sup. mat. for explicit expressions). We address this difficulty by resorting to a perturbative evaluation of the eigenvalue distribution of  $\nabla^2 H$  in the small  $\kappa$  regime. This distribution is used in turn to calculate the determinant modulus as described below.

To evaluate  $|\overline{\det \nabla^2 H(\sigma)}|$  we use recent results (Proposition 5.3 in [65], Theorem 4.1 in [75]), to formally express the expectation value in terms of the eigenvalue distributions of  $\nabla^2\phi$  and  $\kappa\nabla^2 U$ ,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\overline{\det \nabla^2 H(\sigma)}| = \sup_s \int d\lambda \nu_{\nabla^2 H(\sigma)}(\lambda) \log |\lambda - s| - \frac{s^2}{2p^2}, \quad (15)$$

where  $\nu_{\nabla^2 H(\sigma)}$  is the eigenvalue distribution of  $\nabla^2 H(\sigma)$ . This distribution, in the limit of large  $N$ , converges to the free convolution, denoted  $\boxplus$ , of the eigenvalue distributions of  $\nabla^2 \phi$  and  $\kappa \nabla^2 U$  [65, 74],

$$\nu_{\nabla^2 H}(\lambda) = (\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U})(\lambda), \quad (16)$$

where  $\nu_{\kappa \nabla^2 U}(\lambda)$  is the eigenvalue distribution of  $\kappa \nabla^2 U$  and  $\rho_{\text{sc}}(\lambda)$  is the Wigner semicircle distribution with variance  $V_p \equiv p(p-1)$ :

$$\rho_{\text{sc}}(\lambda) = \frac{1}{2\pi V_p} \sqrt{4V_p - \lambda^2}. \quad (17)$$

Intuitively, the free convolution is the random matrix analogue to the convolution operation used to calculate the probability density of a sum of independent random variables. It is non-linear and generally the result cannot be expressed in a closed form. However, in the regime of small  $\kappa$  we are able to find a perturbative expression for  $\nu_{\nabla^2 H}(\lambda)$  by solving perturbatively the Pastur integral equation [76]. This results, up to  $O(\kappa^3)$ , in a shifted and widened semi-circular distribution given by:

$$\nu_{\nabla^2 H}(\lambda) = \frac{1}{\sqrt{1 + \kappa^2 V_{\nabla^2 U}}} \rho_{\text{sc}} \left( \frac{\lambda - \kappa \sqrt{V_p} m_{\nabla^2 U}}{\sqrt{1 + \kappa^2 V_{\nabla^2 U}}} \right) + O(\kappa^3), \quad (18)$$

with  $m_{\nabla^2 U}$  and  $V_{\nabla^2 U}$  the mean and variance of the eigenvalue distribution of  $\frac{1}{\sqrt{V_p}} \nabla^2 U$ , respectively. Now we can proceed to derive a closed form expression for  $|\overline{\det(\nabla^2 H)}|$  by substitution of the above expression for  $\nu_{\nabla^2 H}(\lambda)$  in Eq. 15 and optimization w.r.t  $s$ :

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\overline{\det \nabla^2 H(\sigma)}| = \frac{1}{2} (-1 + \log 2) - \frac{1}{4} \kappa^2 m_{\nabla^2 U}^2 + \kappa^2 V_{\nabla^2 U} \begin{cases} |\eta_2| - \frac{1}{2} & |\eta_2(\sigma)| \geq 2 \\ \frac{1}{2} + \frac{1}{4} \eta_2^2 & |\eta_2(\sigma)| < 2 \end{cases}, \quad (19)$$

where we ignored terms which are  $O(\kappa)^3$  and  $\eta_2 = \frac{m_{\nabla^2 U}/\kappa}{V_{\nabla^2 U}}$ . Note that the result contains two branches. The form of the supremum equation and its solution branches is similar to the one found in [72] as was discussed in detail following the results statement in the previous section.

Now we turn to the evaluation of the integration over the  $N$ -sphere in (3):

$$\overline{\mathcal{N}_{\text{tot}}} = \left( \frac{1}{2\pi p} \right)^{\frac{N}{2}} \int_{\mathbb{S}^{N-1}} d\sigma \exp \left( -\frac{\kappa^2}{2p} \|\nabla U\|^2 \right) |\overline{\det \nabla^2 H}|. \quad (20)$$

The approach we adopt for the evaluation of this integral is to pass from spatial integration over the surface of the sphere to functional integration over the empirical distribution of  $\sigma_i$  which we denote by  $\mu$ , i.e.

$$\mu(x) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \delta(x - \sigma_i) \quad (21)$$

Utilizing the Coulomb gas technique [77] we arrive at the following functional integral:

$$\overline{\mathcal{N}_{\text{tot}}} = \frac{C_N}{\mathcal{Z}} \int D[\mu] \delta \left( \int x^2 \mu(x) dx - 1 \right) e^{NS[\mu]} \quad (22)$$

where  $C_N = \left( \frac{1}{2\pi p} \right)^{N/2} S_{N-1}$ , with  $S_{N-1}$  the surface area of the  $N$ -sphere of radius  $\sqrt{N}$ , and where

$$\begin{aligned} S[\mu] &= \int dx \mu(x) \frac{1}{N} \log |\overline{\det \nabla^2 H}[\mu]| \\ &\quad + \int dx \mu(x) \left( -\log \mu - \frac{\kappa^2}{2p} \|\nabla U\|^2[\mu] \right) \\ \mathcal{Z} &= \int D[\mu] \delta \left( \int x^2 \mu(x) dx - 1 \right) e^{-N \int dx \mu \log \mu}. \end{aligned}$$

$|\overline{\det \nabla^2 H}[\mu]|$  and  $\|\nabla U\|^2[\mu]$  are obtained from the corresponding  $\sigma$  dependent expressions and the usual replacement rules of functional integration (see eqs. (44)-(46) in SI). Note that the delta function constraining the second moment of  $\mu$  is a consequence of the spherical integration. In the limit of large  $N$  we solve the functional integral by the saddle point method w.r.t to  $\mu$  under the spherical and normalization constraints:

$$\begin{aligned} \frac{\delta S}{\delta \mu} + \alpha + \beta x^2 &= 0 \\ \int \mu(x) dx &= 1, \quad \int x^2 \mu(x) dx = 1 \end{aligned} \quad (23)$$

where  $\alpha$  and  $\beta$  are Lagrange multipliers enforcing the above mentioned constraints. The expression for  $\log |\overline{\det \nabla^2 H}|$  is a perturbative one, and we solve the variation problem separating orders of  $\kappa$ . Since the saddle point solution for  $\log |\overline{\det \nabla^2 H}[\mu]|$  has two distinct branches, we obtain two branches of solutions for each order of  $\mu$ ,  $\alpha$  and  $\beta$ . Closed form expressions for both branches are given in the SI. Substitution the optimal  $\mu$  in the expression for  $S[\mu]$  gives the complexity for each branch, since its non-trivial part is given by  $\max_{\mu} S[\mu]$ . Consequently, the value of the critical point complexity is given by  $\Sigma = \max(\Sigma_1, \Sigma_2)$ . The results for the complexity are given in the main results section above.

*Discussion* We found that introducing an ‘‘on-site’’ non-linearity to a disordered mean-field coupled system can significantly increase the complexity of the energy landscape of the system. These results are immediately applicable to physical systems with ‘‘on-site’’ weak non-linearity and disordered long range coupling such as ‘‘soft-spin’’ glass models [1, 2, 52, 54], coupled oscillator networks [17–19] and atomic spin gasses [21].

In many physics problems one considers only short range (local) deterministic coupling, e.g. in the many body localisation setup. In such problems, disorder is commonly introduced ‘‘on-site’’ as opposed to the coupling disorder present in this work. A possible direction to address these problem is by considering random matrix

ensembles modelling short-range coupling such as banded matrix models [65, 78].

Additional questions for future research include extending the results of this work beyond the perturbative regime to explore the next transition in the complexity conjectured to take place for bi-stable potentials. In addition, it is of interest to study the quenched complexity, possibly by carrying out a second moment analysis of the number of critical points in order to determine the extent of concentration of the number of critical points [62].

*Acknowledgments* O. R. is the incumbent of the Shlomo and Michla Tomarin career development chair, and is supported by the Abramson Family Center for

Young Scientists, the Israel Science Foundation Grant No. 950/19 and by the Minerva foundation. IG is grateful to Amy Entin for her help with writing this manuscript. B. LACT is supported by the EPSRC Grant EP/V002473/1 Random Hessians and Jacobians: theory and applications. O. Z. was partially supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 692452) and by the Israel Science Foundation grant 421/20. E. S. is the incumbent of the Skirball Chair in New Scientists, and is supported by the Israel Science Foundation grant 2055/21.

- 
- [1] T. Castellani and A. Cavagna, Spin-glass theory for pedestrians, *Journal of Statistical Mechanics: Theory and Experiment*, 215 (2005).
- [2] M. Mézard, G. Parisi, and M. Virasoro, *Spin glass theory and beyond: An Introduction to the Replica Method and Its Applications*, Vol. 9 (World Scientific Publishing Company, 1987).
- [3] H. T. Diep, *Frustrated Spin Systems, 2nd Edition*, 2nd ed. (World Scientific Publishing Co., 2013) pp. 1–617.
- [4] L. Gartner and E. Lerner, Nonlinear modes disentangle glassy and Goldstone modes in structural glasses, *SciPost Physics* **1**, 016 (2016).
- [5] E. A. Goremychkin, R. Osborn, B. D. Rainford, R. T. MacAluso, D. T. Adroja, and M. Koza, Spin-glass order induced by dynamic frustration, *Nature Physics* **4**, 10.1038/nphys1028 (2008).
- [6] K. Kim, M. S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G. D. Lin, L. M. Duan, and C. Monroe, Quantum simulation of frustrated Ising spins with trapped ions, *Nature* **465**, 590 (2010).
- [7] K. S. Tikhonov and A. D. Mirlin, From Anderson localization on random regular graphs to many-body localization, *Annals of Physics* **435**, 168525 (2021).
- [8] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Reviews of Modern Physics* **91**, 10.1103/RevModPhys.91.021001 (2019).
- [9] F. Antenucci, A. Crisanti, and L. Leuzzi, Complex spherical 2+4 spin glass: A model for nonlinear optics in random media, *Physical Review A - Atomic, Molecular, and Optical Physics* **91**, 053816 (2015).
- [10] A. Zyuzin and B. Spivak, Propagation of nonlinear waves in disordered media, *Journal of the Optical Society of America B* **21**, 177 (2004).
- [11] C. Conti and L. Leuzzi, Complexity of waves in nonlinear disordered media, *Physical Review B - Condensed Matter and Materials Physics* **83**, 10.1103/PhysRevB.83.134204 (2011).
- [12] N. Ghofraniha, I. Viola, F. Di Maria, G. Barbarella, G. Gigli, L. Leuzzi, and C. Conti, Experimental evidence of replica symmetry breaking in random lasers, *Nature Communications* **6**, 10.1038/ncomms7058 (2015).
- [13] S. W. Shaw and C. Pierre, Normal Modes for Non-Linear Vibratory Systems, *Journal of Sound and Vibration* **164**, 85 (1993).
- [14] S. Fishman, Y. Krivolapov, and A. Soffer, The nonlinear Schrödinger equation with a random potential: results and puzzles, *Nonlinearity* **25**, R53 (2012).
- [15] G. Tarnopolsky, C. Li, D. G. Joshi, and S. Sachdev, Metal-insulator transition in a random Hubbard model, *Physical Review B* **101**, 205106 (2020).
- [16] T. R. Kirkpatrick and D. Belitz, Suppression of superconductivity by disorder, *Physical Review Letters* **68**, 3232 (1992).
- [17] T. Inagaki, Y. Haribara, K. Igarashi, T. Sonobe, S. Tamate, T. Honjo, A. Marandi, P. L. McMahon, T. Umeki, K. Enbutsu, O. Tadanaga, H. Takenouchi, K. Aihara, K. I. Kawarabayashi, K. Inoue, S. Utsunomiya, and H. Takesue, A coherent Ising machine for 2000-node optimization problems, *Science* **354**, 603 (2016).
- [18] Z. Wang, A. Marandi, K. Wen, R. L. Byer, and Y. Yamamoto, Coherent Ising machine based on degenerate optical parametric oscillators, *Physical Review A - Atomic, Molecular, and Optical Physics* **88**, 10.1103/PhysRevA.88.063853 (2013).
- [19] M. Nixon, E. Ronen, A. A. Friesem, and N. Davidson, Observing geometric frustration with thousands of coupled lasers, *Physical Review Letters* **110**, 10.1103/PhysRevLett.110.184102 (2013).
- [20] S. Gopalakrishnan, B. L. Lev, and P. M. Goldbart, Frustration and glassiness in spin models with cavity-mediated interactions, *Physical Review Letters* **107**, 277201 (2011).
- [21] Y. Horowicz, O. Katz, O. Raz, and O. Firstenberg, Critical dynamics and phase transition of a strongly interacting warm spin gas, *Proceedings of the National Academy of Sciences of the United States of America* **118**, 10.1073/PNAS.2106400118 (2021).
- [22] C. Lippe, T. Klas, J. Bender, P. Mischke, T. Niederprüm, and H. Ott, Experimental realization of a 3D random hopping model, *Nature Communications* **12**, 10.1038/s41467-021-27243-2 (2021).
- [23] C. Fort, L. Fallani, V. Guarrera, J. E. Lye, M. Modugno, D. S. Wiersma, and M. Inguscio, Effect of optical disorder and single defects on the expansion of a Bose-Einstein condensate in a one-dimensional waveguide, *Physical Review Letters* **95**, 10.1103/PhysRevLett.95.170410 (2005).
- [24] P. Lugan, D. Clément, P. Bouyer, A. Aspect, M. Lewenstein, and L. Sanchez-Palencia, Ultracold Bose gases in 1D disorder: From Lifshits glass to Bose-Einstein

- condensate, *Physical Review Letters* **98**, 10.1103/PhysRevLett.98.170403 (2007).
- [25] D. C. Tsui, Nobel lecture: Interplay of disorder and interaction in two-dimensional electron gas in intense magnetic fields, *Reviews of Modern Physics* **71**, 891 (1999).
- [26] D. G. Joshi, C. Li, G. Tarnopolsky, A. Georges, and S. Sachdev, Deconfined Critical Point in a Doped Random Quantum Heisenberg Magnet, *Physical Review X* **10**, 021033 (2020).
- [27] J. C. Eilbeck, P. S. Lomdahl, and A. C. Scott, The discrete self-trapping equation, *Physica D: Nonlinear Phenomena* **16**, 318 (1985).
- [28] M. Goldstein, Viscous liquids and the glass transition: A potential energy barrier picture, *The Journal of Chemical Physics* **51**, 10.1063/1.1672587 (1969).
- [29] A. Namatame and S.-H. Chen, *Agent-Based Modeling and Network Dynamics* (Oxford University Press, 2016).
- [30] G. Palla, A. L. Barabási, and T. Vicsek, Community dynamics in social networks, *Fluctuation and Noise Letters* **7**, 0 (2007).
- [31] H. Ebel, J. Davidsen, and S. Bornholdt, Dynamics of social networks, *Complexity* **8**, 24 (2002).
- [32] L. Sattenspiel and K. Dietz, A structured epidemic model incorporating geographic mobility among regions, *Mathematical Biosciences* **128**, 71 (1995).
- [33] R. Pastor-Satorras, C. Castellano, P. Van Mieghem, and A. Vespignani, Epidemic processes in complex networks, *Reviews of Modern Physics* **87**, 10.1103/RevModPhys.87.925 (2015).
- [34] F. Fu, D. I. Rosenbloom, L. Wang, and M. A. Nowak, Imitation dynamics of vaccination behaviour on social networks, *Proceedings of the Royal Society B: Biological Sciences* **278**, 42 (2011).
- [35] A. W. Roddam, *Mathematical Epidemiology of Infectious Diseases: Model Building, Analysis and Interpretation*, *International Journal of Epidemiology* **30**, 10.1093/ije/30.1.186 (2001).
- [36] H. W. Hethcote, *Mathematics of infectious diseases*, *SIAM Review* **42**, 10.1137/S0036144500371907 (2000).
- [37] R. M. May, *Stability and Complexity in Model Ecosystems* (Princeton Landmarks In Biology), Princeton University Press **6**, 1 (2001).
- [38] Y. V. Fyodorov and B. A. Khoruzhenko, Nonlinear analogue of the May-Wigner instability transition, *Proceedings of the National Academy of Sciences of the United States of America* **113**, 10.1073/pnas.1601136113 (2016).
- [39] S. Allesina and S. Tang, The stability-complexity relationship at age 40: a random matrix perspective, *Population Ecology* **57**, 63 (2015).
- [40] Y. Bahri, J. Kadmon, J. Pennington, S. S. Schoenholz, J. Sohl-Dickstein, and S. Ganguli, *Statistical Mechanics of Deep Learning* (2020).
- [41] M. Baity-Jesi, L. Sagun, M. Geiger, S. Spigler, G. Ben Arous, C. Cammarota, Y. Lecun, M. Wyart, and G. Biroli, Comparing dynamics: deep neural networks versus glassy systems, *Journal of Statistical Mechanics: Theory and Experiment* **2019**, 10.1088/1742-5468/ab3281 (2019).
- [42] S. Becker, Y. Zhang, and A. A. Lee, Geometry of Energy Landscapes and the Optimizability of Deep Neural Networks, *Physical Review Letters* **124**, 108301 (2020).
- [43] M. Mézard and A. Montanari, *Information, Physics, and Computation*, *Information, Physics, and Computation* **9780198570837**, 1 (2009).
- [44] M. E. Newman, A. L. Barabási, and D. J. Watts, *The Structure and Dynamics of Networks*, Vol. 9781400841356 (Princeton University Press, 2011).
- [45] G. Ben Arous, Y. V. Fyodorov, and B. A. Khoruzhenko, Counting equilibria of large complex systems by instability index, *Proceedings of the National Academy of Sciences of the United States of America* **118**, 10.1073/pnas.2023719118 (2021).
- [46] V. Ros, G. Biroli, and C. Cammarota, Complexity of energy barriers in mean-field glassy systems, *EPL* **126**, 10.1209/0295-5075/126/20003 (2019).
- [47] B. Lacroix-A-Chez-Toine and Y. V. Fyodorov, Counting equilibria in a random non-gradient dynamics with heterogeneous relaxation rates, *Journal of Physics A: Mathematical and Theoretical* **55**, 10.1088/1751-8121/ac564a (2022).
- [48] U. Buchenau, Energy landscape-A key concept in the dynamics of liquids and glasses, *Journal of Physics Condensed Matter* **15**, 10.1088/0953-8984/15/11/319 (2003).
- [49] G. B. Arous, E. Subag, and O. Zeitouni, Geometry and Temperature Chaos in Mixed Spherical Spin Glasses at Low Temperature: The Perturbative Regime, *Communications on Pure and Applied Mathematics* **73**, 10.1002/cpa.21875 (2020).
- [50] K. Dahmen and J. P. Sethna, Hysteresis, avalanches, and disorder-induced critical scaling: A renormalization-group approach, *Physical Review B - Condensed Matter and Materials Physics* **53**, 10.1103/PhysRevB.53.14872 (1996).
- [51] Y. N. Dauphin, R. Pascanu, C. Gulcehre, K. Cho, S. Ganguli, and Y. Bengio, Identifying and attacking the saddle point problem in high-dimensional non-convex optimization, in *Advances in Neural Information Processing Systems*, Vol. 4 (2014).
- [52] A. Cavagna, I. Giardina, and G. Parisi, Role of saddles in mean-field dynamics above the glass transition, *Journal of Physics A: Mathematical and General* **34**, 10.1088/0305-4470/34/26/302 (2001).
- [53] M. Goldstein, Viscous liquids and the glass transition. V. Sources of the excess specific heat of the liquid, *The Journal of Chemical Physics* **64**, 10.1063/1.432063 (1976).
- [54] A. Cavagna, I. Giardina, and G. Parisi, Stationary points of the Thouless-Anderson-Palmer free energy, *Physical Review B - Condensed Matter and Materials Physics* **57**, 11251 (1998).
- [55] M. L. Rosinberg and T. Munakata, Hysteresis and complexity in the mean-field random-field Ising model: The soft-spin version, *Physical Review B - Condensed Matter and Materials Physics* **79**, 174207 (2009).
- [56] J. Kurchan, Replica trick to calculate means of absolute values: Applications to stochastic equations, *Journal of Physics A: General Physics* **24**, 10.1088/0305-4470/24/21/011 (1991).
- [57] R. Monasson, Structural glass transition and the entropy of the metastable states, *Physical Review Letters* **75**, 10.1103/PhysRevLett.75.2847 (1995).
- [58] Y. V. Fyodorov, Complexity of random energy landscapes, glass transition, and absolute value of the spectral determinant of random matrices, *Physical Review Letters* **92**, 10.1103/PhysRevLett.92.240601 (2004).
- [59] Y. V. Fyodorov and I. Williams, Replica symmetry breaking condition exposed by random matrix calculation of landscape complexity, *Journal of Statistical Physics* **129**, 1081 (2007).

- [60] A. Auffinger, G. Ben Arous, and J. Černý, Random matrices and complexity of spin glasses, *Communications on Pure and Applied Mathematics* **66**, 165 (2013).
- [61] A. Auffinger and G. Ben Arous, Complexity of random smooth functions on the high-dimensional sphere, *Annals of Probability* **41**, 10.1214/13-AOP862 (2013).
- [62] E. Subag, The complexity of spherical p-spin models—a second moment approach, *Annals of Probability* **45**, 10.1214/16-AOP1139 (2017).
- [63] B. McKenna, *Non-invariant Random Matrices and Landscape Complexity*, Ph.D. thesis, New York University, New York (2021).
- [64] B. McKenna, Complexity of bipartite spherical spin glasses, arXiv:2105.05043 10.48550/arxiv.2105.05043 (2021).
- [65] G. B. Arous, P. Bourgade, and B. McKenna, Landscape complexity beyond invariance and the elastic manifold, arXiv:2105.05051 10.48550/arxiv.2105.05051 (2021).
- [66] E. Subag, The free energy of spherical pure p-spin models – computation from the TAP approach, arXiv:2101.04352 (2021).
- [67] E. Subag, Free energy landscapes in spherical spin glasses, arXiv:1804.10576 (2018).
- [68] A. Choromanska, M. Henaff, M. Mathieu, G. Ben Arous, and Y. LeCun, The loss surfaces of multilayer networks, in *Journal of Machine Learning Research*, Vol. 38 (2015).
- [69] Y. V. Fyodorov and P. Le Doussal, Manifolds in a high-dimensional random landscape: Complexity of stationary points and depinning, *Physical Review E* **101**, 10.1103/PHYSREVE.101.020101 (2020).
- [70] Y. V. Fyodorov and P. Le Doussal, Topology Trivialization and Large Deviations for the Minimum in the Simplest Random Optimization, *Journal of Statistical Physics* **154**, 466 (2014).
- [71] A. Dembo and O. Zeitouni, Matrix optimization under random external fields, *J. Stat. Phys.* **159**, 1306 (2015).
- [72] D. Belius, J. Černý, S. Nakajima, and M. A. Schmidt, Triviality of the Geometry of Mixed p-Spin Spherical Hamiltonians with External Field, *Journal of Statistical Physics* **186**, 10.1007/S10955-021-02855-6 (2022).
- [73] P. G. Kevrekidis, K. Rasmussen, and A. R. Bishop, The discrete nonlinear schrödinger equation: A survey of recent results, *International Journal of Modern Physics B* **15**, 2833 (2001).
- [74] G. W. Anderson, A. Guionnet, and O. Zeitouni, *An Introduction to Random Matrices* (Cambridge University Press, 2009).
- [75] G. Ben Arous, P. Bourgade, and B. McKenna, Exponential growth of random determinants beyond invariance, arXiv:2105.05000 (2021).
- [76] P. Biane, On the free convolution with a semi-circular distribution, *Indiana University Mathematics Journal* **46**, 10.1512/iumj.1997.46.1467 (1997).
- [77] F. J. Dyson, Statistical theory of the energy levels of complex systems. I, *Journal of Mathematical Physics* **3**, 10.1063/1.1703773 (1962).
- [78] P. Bourgade, L. Erdős, H. T. Yau, and J. Yin, Universality for a class of random band matrices, *Advances in Theoretical and Mathematical Physics* **21**, 739 (2017).
- [79] A. J. Bray and D. S. Dean, Statistics of Critical Points of Gaussian Fields on Large-Dimensional Spaces, *Physical Review Letters* **98**, 150201 (2007).
-

# Supplemental Information: On-Site Potential Creates Complexity in Systems with Disordered Coupling

## Part I

### Setup

Define an “energy landscape” on the  $N$ -sphere composed of a “pure”  $p$ -spin Gaussian field  $\phi(\sigma)$  and an “on-site” deterministic potential  $U(\sigma)$ :

$$H(\sigma) = \phi(\sigma) + \kappa U(\sigma) = \phi(\sigma) + \kappa \sum_{i=1}^N u(\sigma_i) \quad (\text{S1})$$

where  $\sigma \in \sqrt{N}\mathbb{S}^{N-1}$ ,  $\kappa$  is the “on-site” potential strength parameter, and  $u(x)$  is the potential function, assumed throughout this work to be smooth and of bounded derivatives. The Gaussian field on the sphere is defined by the following correlation function corresponding to the “pure”  $p$ -spin spherical model:

$$\overline{\phi(\sigma)\phi(\sigma')} = N \left( \frac{\sigma \cdot \sigma'}{N} \right)^p. \quad (\text{S2})$$

To evaluate the average total number of critical points on the sphere, we use the spherical analogue of the Kac-Rice formula (as given in [60])

$$\overline{\mathcal{N}_{\text{tot}}} = \int_{\sqrt{N}\mathbb{S}^{N-1}} d\sigma_{N-1} \overline{|\det(\nabla^2 H)| \delta(\nabla H)} \quad (\text{S3})$$

where  $\nabla H$  and  $\nabla^2 H$  are respectively the covariant derivative and Hessian matrix of  $H$  on the sphere, and  $\overline{X}$  denotes averaging a quantity  $X$  over the random field distribution. Defining the index as the number of negative eigenvalues of the Hessian, the average number of critical points of index  $k$  is similarly given by

$$\overline{\mathcal{N}_k} = \int_{\sqrt{N}\mathbb{S}^{N-1}} d\sigma_{N-1} \overline{|\det(\nabla^2 H)| \delta(\nabla H) \delta(I(\nabla^2 H) - k)}, \quad (\text{S4})$$

where  $I(\cdot)$  is the index function; see [79]. In this work, we are mainly interested in the annealed (that is, averaged over the Gaussian field  $\phi$ ) complexity of the number of critical points. The total complexity and the complexity of critical points of extensive index  $k = [aN]$  (with  $a \in (0, 1)$ ) are defined respectively as:

$$\Sigma_{\text{tot}} = \Sigma_{\text{tot}}[u(x)] = \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{\mathcal{N}_{\text{tot}}} \quad (\text{S5})$$

$$\Sigma_a = \Sigma_a[u(x)] = \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{\mathcal{N}_{aN}} \quad (\text{S6})$$

where the limit in the second equation is taken such that  $a$  is fixed.

## Part II

### Main Results

Our main results are given by the statements in Sections I and II. The proofs occupy the rest of this report.

#### I. COMPLEXITY OF THE MEAN TOTAL NUMBER OF CRITICAL POINTS

The total annealed complexity of a  $p$ -spin model perturbed by a weak “on-site” potential,  $\Sigma_{\text{tot}}^{(p)}$ , as a functional of the potential  $u(\cdot)$ , is given by:

$$\Sigma_{\text{tot}}^{(p=2)}[u] = \kappa^2 \max\left(0, \Theta^{(2)}[u]\right) + O(\kappa^3) \quad (\text{S7})$$

$$\Sigma_{\text{tot}}^{(p>2)}[u] = \frac{1}{2} \log(p-1) + \kappa^2 \Theta^{(p)}[u] + O(\kappa^3) \quad (\text{S8})$$



where

$$\begin{aligned} \Theta^{(p)}[u] &= \frac{1}{2p} \int dx \frac{e^{-x^2/2}}{\sqrt{2\pi}} \left( \frac{1}{p-1} u''(x)^2 - u'(x)^2 \right) \\ &\quad + \frac{p-2}{2p(p-1)} \left( \int \frac{e^{-x^2/2}}{\sqrt{2\pi}} u''(x) dx \right)^2. \end{aligned} \quad (\text{S9})$$

Note that for  $p = 2$ , Eq. S9 specializes to

$$\Theta^{(p=2)}[u] = \frac{1}{4} \int dx \frac{e^{-x^2/2}}{\sqrt{2\pi}} (u''(x)^2 - u'(x)^2). \quad (\text{S10})$$

## II. COMPLEXITY OF THE MEAN NUMBER OF CRITICAL POINTS OF A GIVEN INDEX

The annealed complexity of critical points of (extensive) index  $k = \lfloor aN \rfloor$  of a  $p$ -spin model perturbed by a weak “on-site” potential,  $\Sigma_a^{(p)}$ , as a functional of the potential  $u$ , is given by

$$\Sigma_a^{(p)}[u] = \frac{1}{2} \log(p-1) - \frac{p-2}{4p} \eta^2(a) + \kappa^2 \left( 1 - \eta^2(a) \frac{p-1}{p^2} \right) \Theta^{(p)}[u] + O(\kappa^3) \quad (\text{S11})$$

where  $\eta(a) \in (-2, 2)$  is defined by

$$\frac{1}{2\pi} \int_{-2}^{\eta(a)} \sqrt{4-x^2} dx = a. \quad (\text{S12})$$

## Part III

# Results Derivation

## III. EVALUATION OF THE MEAN TOTAL NUMBER OF CRITICAL POINTS

### A. Preliminaries

We start the calculation of (S3) by evaluating the disorder average  $\overline{|\det(\nabla^2 H(\sigma))| \delta(\nabla H(\sigma))}$  at a fixed position  $\sigma$  on the sphere. The average is taken over the joint distribution of the covariant Hessian and gradient of the field  $\phi$ . See Appendix A for detailed description of the joint distribution of the fields  $\phi$ ,  $\nabla\phi$ ,  $\nabla^2\phi$  adopted from [60]. We note here, for the sake of clarity, the main properties of this distribution which are instrumental for our calculation:

1. At each point  $\sigma$ , the field  $\phi$  and its derivatives are mean zero jointly normally distributed random variables.
2. At any point  $\sigma$ ,  $\phi(\sigma)$  and  $\nabla^2\phi(\sigma)$  are independent of  $\nabla\phi(\sigma)$ .
3. The components of  $\nabla\phi(\sigma)$  are independent. The variance of each component is  $\text{Var}((\nabla\phi)_m) = p$ .
4. The random matrix  $\nabla^2\phi$  is distributed like  $\sqrt{p(p-1)}M + \xi p \mathbf{I}_{N-1}$  where  $M \sim \sqrt{\frac{N-1}{N}} \text{GOE}_{N-1}$ ,  $\xi \sim \mathcal{N}(0, \frac{1}{N})$ ,  $\mathbf{I}_{N-1}$  is the identity matrix of size  $(N-1) \times (N-1)$ . We denote by  $\text{GOE}_n$  the Gaussian orthogonal ensemble of dimension  $n$  [74] with an off-diagonal variance of  $\frac{1}{n}$  i.e. if  $A \sim \text{GOE}_n$  then  $\mathbb{E}[A_{ij}^2] = \frac{1+\delta_{ij}}{n}$ .

First, we use the independence of  $\nabla\phi$  and  $\nabla\phi^2$  to factorize the expectation in (S3) as follows:

$$\overline{|\det(\nabla^2 H)| \delta(\nabla H)} = \overline{\delta(\nabla H)} \times \overline{|\det(\nabla^2 H)|}. \quad (\text{S13})$$

The average  $\overline{\delta(\nabla H(\sigma))}$  is easily carried out and gives:

$$\begin{aligned} \overline{\delta(\nabla H(\sigma))} &= \prod_{m=1}^{N-1} \overline{\delta((\nabla\phi(\sigma))_m + \kappa(\nabla U(\sigma))_m)} = \prod_{m=1}^{N-1} \sqrt{\frac{1}{2\pi p}} \exp\left(-\frac{\kappa^2}{2p} (\nabla U(\sigma))_m^2\right) \\ &= \left(\frac{1}{2\pi p}\right)^{(N-1)/2} \exp\left(-\frac{\kappa^2}{2p} \|\nabla U(\sigma)\|^2\right), \end{aligned} \quad (\text{S14})$$

where in the first equality we used the gradient component independence and in the second the normal distribution of each component.

We evaluate  $|\det(\nabla^2 H)|$  by using the distribution in 4 as follows [60]:

$$\begin{aligned} \overline{|\det(\nabla^2 H)|} &= \overline{|\det(\nabla^2 \phi + \kappa \nabla^2 U)|} \\ &= \sqrt{\frac{N}{2\pi p^2}} \int_{\mathbb{R}} \exp\left(-N \frac{s^2}{2p^2}\right) \overline{|\det \sqrt{p(p-1)M + \kappa \nabla^2 U + sI_{N-1}}|} ds, \end{aligned} \quad (\text{S15})$$

where  $M \sim \sqrt{\frac{N-1}{N}} \text{GOE}_{N-1}$  and it is understood that the averaging is carried out over the matrix  $M$ . In the second equality we have used the distribution of  $\nabla^2 \phi$  defined in item 4 above. Next, we employ the same method as in Proposition 5.3 of [65] and use Theorem 4.1 in [75] to get

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \left( \overline{|\det \nabla^2 H(\sigma)|} \right) = \sup_s \left( \int d\lambda (\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U})(\lambda) \log |\lambda - s| - \frac{s^2}{2p^2} \right), \quad (\text{S16})$$

where  $\boxplus$  is the free-convolution [74],  $\rho_{\text{sc}}$  is the Wigner semicircle of variance  $p(p-1)$  and  $\nu_{\kappa \nabla^2 U}$  is the large  $N$  limit of the empirical spectral distribution of  $\kappa \nabla^2 U$ , i.e.

$$\rho_{\text{sc}}(\lambda) = \frac{1}{2\pi p(p-1)} \sqrt{4p(p-1) - \lambda^2} \mathbb{1}_{|\lambda| < 2\sqrt{p(p-1)}} \quad (\text{S17})$$

$$\nu_{\kappa \nabla^2 U}(\lambda) = \lim_{N \rightarrow \infty} \frac{1}{N-1} \sum_{m=1}^{N-1} \delta(\lambda - \kappa \lambda_m(\nabla^2 U)) \quad (\text{S18})$$

where  $\lambda_m(\nabla^2 U)$  with  $m = 1, \dots, N-1$  are the eigenvalues of the matrix  $\nabla^2 U$ .

## B. Free convolution evaluation

The free convolution operation is non-linear and generally the result cannot be expressed in a closed form. However, in the regime of small  $\kappa$ , we are able to solve the Pastur integral equation [76] and find a perturbative expression for  $\nu_{\nabla^2 H}(\lambda)$ . Let  $G(z)$  be the Stieltjes transform [74] of  $\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U}$ . Since one of the distributions in the convolution is the semicircle, one obtains a relatively simple implicit equation for  $G$  (the Pastur equation) [76]:

$$G(z) = \int \frac{\nu_{\kappa \nabla^2 U}(\lambda) d\lambda}{z - p(p-1)G(z) - \lambda}, \quad z \in \mathbf{C}_+. \quad (\text{S19})$$

In Appendix B, the perturbative solution of this equation is detailed up to order  $O(\kappa^2)$ . Next, the inverse transform is utilized to calculate  $\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U}$  from  $G$ . Up to  $O(\kappa^3)$ , this results in a shifted and widened semi-circular distribution given by

$$\begin{aligned} (\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U})(\lambda) &= O(\kappa^3) + \frac{1}{2\pi p(p-1)(1 + \kappa^2 V_{\nabla^2 U})} \times \\ &\sqrt{-\left(\lambda - \kappa \sqrt{p(p-1)} m_{\nabla^2 U}\right)^2 + 4p(p-1)(1 + \kappa^2 V_{\nabla^2 U})}, \end{aligned} \quad (\text{S20})$$

where we have denoted the expectation and variance of the spectrum of  $\frac{1}{\sqrt{p(p-1)}} \nabla^2 U(\sigma)$  by  $m_{\nabla^2 U}$  and  $V_{\nabla^2 U}$ , respectively, i.e.,

$$\begin{aligned} m_{\nabla^2 U}(\sigma) &\equiv \frac{1}{\sqrt{p(p-1)}} \int \lambda \nu_{\nabla^2 U(\sigma)}(\lambda) d\lambda \\ V_{\nabla^2 U}(\sigma) &\equiv \frac{1}{p(p-1)} \int (\lambda - m_{\nabla^2 U(\sigma)})^2 \nu_{\nabla^2 U(\sigma)}(\lambda) d\lambda. \end{aligned} \quad (\text{S21})$$

### C. Evaluation of the supremum

Equipped with an explicit, simple expression for the spectral distribution of  $\nabla^2 H(\sigma)$ , we can evaluate the integral in (S16) explicitly:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)| &= \sup_s \left[ \Phi \left( \frac{\kappa \sqrt{p(p-1)} m_{\nabla^2 U} + s}{\sqrt{p(p-1)}(1 + \kappa^2 V_{\nabla^2 U})} \right) + \frac{(\kappa \sqrt{p(p-1)} m_{\nabla^2 U} + s)^2}{4p(p-1)(1 + \kappa^2 V_{\nabla^2 U})} \right. \\ &\quad \left. + \frac{1}{2} (-1 + \log(p(p-1)(1 + \kappa^2 V_{\nabla^2 U}))) - \frac{s^2}{2p^2} \right] + O(\kappa^3), \end{aligned} \quad (\text{S22})$$

where

$$\begin{aligned} \Phi(\eta) &= \int_{-2}^2 \frac{d\lambda}{2\pi} \sqrt{4 - \lambda^2} \log |\lambda - \eta| - \frac{\eta^2}{4} + \frac{1}{2} \\ &= -\Theta_H(|\eta| - 2) \left( \frac{|\eta| \sqrt{\eta^2 - 4}}{4} + \log \frac{|\eta| - \sqrt{\eta^2 - 4}}{2} \right) \end{aligned} \quad (\text{S23})$$

and  $\Theta_H$  denotes the Heaviside step function.

The change of variables

$$\eta = \frac{\kappa \sqrt{p(p-1)} m_{\nabla^2 U} + s}{\sqrt{p(p-1)}(1 + \kappa^2 V_{\nabla^2 U})} \quad (\text{S24})$$

and some algebra give:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)| &= -\frac{1}{2} + \frac{1}{2} \log(p(p-1)) \\ &+ \sup_{\eta} \left[ \Phi(\eta) - \frac{p-2}{4p} \eta^2 + \kappa \frac{p-1}{p} m_{\nabla^2 U} \eta + \frac{\kappa^2}{2p} ((p - (p-1)\eta^2) V_{\nabla^2 U} - (p-1) m_{\nabla^2 U}^2) \right] + O(\kappa^3). \end{aligned} \quad (\text{S25})$$

Optimization with respect to  $\eta$  has to be carried out separately for the case  $p = 2$  and  $p > 2$  due to the vanishing of the  $O(1)$  and  $\eta^2$  term at  $p = 2$ .

#### 1. Optimization for the case of $p > 2$

For this case, the optimal  $\eta$  is given by:

$$\eta_{(p>2)}^*(\sigma) = \kappa m_{\nabla^2 U} \frac{2(p-1)}{p-2} + O(\kappa^2). \quad (\text{S26})$$

Substitution of  $\eta_{(p>2)}^*$  into (S25) gives the following:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)| = \frac{1}{2} (-1 + \log p(p-1)) + \kappa^2 \left( \frac{1}{2} V_{\nabla^2 U} + m_{\nabla^2 U}^2 \frac{(p-1)}{2(p-2)} \right) + O(\kappa^3). \quad (\text{S27})$$

#### 2. Optimization for $p = 2$

For the case of  $p = 2$ , the  $-\eta^2$  term is absent, leading to a different solution form. Define:

$$\eta_1(\sigma) = \text{sign}(m_{\nabla^2 U}) \left( 2 + \frac{1}{4} \kappa^2 m_{\nabla^2 U}^2 \right) + O(\kappa^3) \quad (\text{S28})$$

$$\eta_2(\sigma) = \frac{m_{\nabla^2 U} / \kappa}{V_{\nabla^2 U}} + O(\kappa) \quad (\text{S29})$$

As (S25) is defined in a piece-wise manner (see definition of  $\Phi(\eta)$  above) we take the derivative of (S25) w.r.t  $\eta$  and look for extrema in each domain separately. In the domain  $|\eta| < 2$  we have:

$$\frac{d}{d\eta} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)| = \begin{cases} \frac{\kappa}{2} m_{\nabla^2 U} - \eta \frac{\kappa^2}{2} V_{\nabla^2 U} + O(\kappa^3) & |\eta| < 2 \\ -\frac{1}{2} \sqrt{\eta^2 - 4} + \frac{\kappa}{2} m_{\nabla^2 U} - \eta \frac{\kappa^2}{2} V_{\nabla^2 U} + O(\kappa^3) & |\eta| > 2 \end{cases} \quad (\text{S30})$$

We see from the above that a maximum can be found in both domains. Thus, depending on the value of  $\eta_2$ , we have either one or two maxima values s.t. the set of maxima values  $\eta_{\max}$  is given by:

$$\eta_{\max} = \begin{cases} \{\eta_1, \eta_2\} & |\eta_2| < 2 \\ \eta_1 & |\eta_2| \geq 2 \end{cases} \quad (\text{S31})$$

where  $\eta_1$  is in the domain  $|\eta| > 2$  and  $\eta_2$ , if exists as an extremum, is found in  $|\eta| < 2$ . The values of the function at each of the maxima are given by:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)|_{\eta=\eta_1} = \frac{1}{2} (-1 + \log 2) - \frac{1}{4} \kappa^2 m_{\nabla^2 U}^2 + \kappa^2 V_{\nabla^2 U} \left( |\eta_2| - \frac{1}{2} \right) \quad (\text{S32})$$

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)|_{\eta=\eta_2} = \frac{1}{2} (-1 + \log 2) - \frac{1}{4} \kappa^2 m_{\nabla^2 U}^2 + \kappa^2 V_{\nabla^2 U} \left( \frac{1}{2} + \frac{1}{4} \eta_2^2 \right) \quad (\text{S33})$$

Observing the above expressions and noting that  $|\eta_2| \leq 2$  (since  $\eta_2$  is a solution in the domain  $|\eta| < 2$ ) we see that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)|_{\eta=\eta_1} \leq \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)|_{\eta=\eta_2}. \quad (\text{S34})$$

Combining all of the above we find that the optimal  $\eta^*$  and the supremum of the log-determinant are found to be:

$$\eta_{(p=2)}^*(\sigma) = \begin{cases} \eta_1(\sigma) & , \quad |\eta_2| \geq 2 \\ \eta_2(\sigma) & , \quad |\eta_2| < 2 \end{cases}, \quad (\text{S35})$$

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H(\sigma)|_{\eta=\eta^*} = \\ \frac{1}{2} (-1 + \log 2) - \frac{1}{4} \kappa^2 m_{\nabla^2 U}^2 + \kappa^2 V_{\nabla^2 U} \begin{cases} |\eta_2| - \frac{1}{2} & , \quad |\eta_2(\sigma)| \geq 2 \\ \frac{1}{2} + \frac{1}{4} \eta_2^2 & , \quad |\eta_2(\sigma)| < 2 \end{cases}. \end{aligned} \quad (\text{S36})$$

#### D. Integration over the $N$ -sphere

Having completed the evaluation of the disorder average  $\overline{|\det(\nabla^2 H(\sigma))| \delta(\nabla H(\sigma))}$ , we proceed to evaluate the integration over the surface of the sphere in (S3), i.e.,

$$\overline{\mathcal{N}_{\text{tot}}} = \left( \frac{1}{2\pi p} \right)^{N/2} \int_{\sqrt{N}\mathbb{S}^{N-1}} d\sigma_{N-1} \exp \left( -\frac{\kappa^2}{2p} \|\nabla U(\sigma)\|^2 + \log |\det \nabla^2 H(\sigma)| \right). \quad (\text{S37})$$

The approach we adopt for the evaluation of this integral is to pass from spatial integration over the surface of the sphere to functional integration over the empirical distribution of the coordinates  $\sigma_i$ , which we denote by  $\mu$ , that is  $\mu(x) = N^{-1} \sum_{i=1}^N \delta(x - \sigma_i)$ . First, the integration over the sphere is converted as follows to an integration over  $\mathbb{R}^N$ , such that for any smooth function on the sphere,

$$\int_{\sqrt{N}\mathbb{S}^{N-1}} f(\sigma) d\sigma_{N-1} = \int_{\mathbb{R}^N} f(\sigma) \delta \left( \sum_{i=1}^N \sigma_i^2 - N \right) d^N \sigma. \quad (\text{S38})$$

This allows us to pass from integration over  $\mathbb{R}^N$  to functional integration over  $\mu$ . Noting that for any function  $f$  on the sphere, invariant under permutations of the coordinates, we have  $f(\sigma) = f[\mu]$  and therefore applying the Coulomb

gas technique [77] leads to the following rule for passing between integration on the sphere to integration over the space of probability measures on the real line:

$$\int_{\sqrt{N}\mathbb{S}^{N-1}} d\sigma_{N-1} f(\sigma) \sim \frac{S_{N-1}(\sqrt{N})}{\mathcal{Z}} \int D[\mu] \exp\left(-N \int dx \mu \log \mu\right) \delta\left(N \int \mu(x) x^2 dx - N\right) f[\mu] \quad (\text{S39})$$

with  $S_{N-1}(\sqrt{N}) = \frac{2}{\Gamma(N/2)}(\pi N)^{N/2}$  being the surface area of the  $N$ -sphere of radius  $\sqrt{N}$  and

$$\mathcal{Z} = \int D[\mu] \exp\left(-N \int dx \mu \log \mu\right) \delta\left(N \int \mu(x) x^2 dx - N\right). \quad (\text{S40})$$

Applying this functional integration procedure to the expression for  $\overline{\mathcal{N}_{\text{tot}}}$  results in the following:

$$\begin{aligned} \overline{\mathcal{N}_{\text{tot}}} &= \frac{C_N}{\mathcal{Z}} \int D[\mu] \delta\left(\int dx \mu(x) x^2 - 1\right) \times \\ &\exp N \left( \frac{1}{N} \log |\overline{\nabla^2 H}[\mu]| - \frac{\kappa^2}{2pN} \|\nabla U[\mu]\|^2 - \int dx \mu \log \mu \right) \end{aligned} \quad (\text{S41})$$

where  $C_N = \left(\frac{1}{2\pi p}\right)^{N/2} S_{N-1}(\sqrt{N})$ . To derive explicit expressions for  $\log |\overline{\nabla^2 H}[\mu]|$  and  $\|\nabla U[\mu]\|^2$ , we first calculate the covariant derivatives  $\nabla U$  and  $\nabla^2 U$  by exploiting the ‘‘single spin’’ structure of  $U = \sum u(\sigma_m)$ : the covariant derivative and Hessian on the sphere of  $U$  are given by:

$$(\nabla U)_m = u'(\sigma_m) - \frac{\sigma_m}{N} \sum_l \sigma_l u'(\sigma_l) \quad (\text{S42})$$

and

$$(\nabla^2 U)_{mn} \approx \delta_{mn} \left( u''(\sigma_m) - \frac{1}{N} \sum_l \sigma_l u'(\sigma_l) \right) \quad (\text{S43})$$

where the approximation in the last equality constitutes neglecting terms of rank  $O(1)$  as described in Appendix B. Using these expressions, we get:

$$\|\nabla U[\mu]\|^2 = \int \mu(x) u'^2(x) dx - \left( \int x u'(x) \mu(x) dx \right)^2 \quad (\text{S44})$$

$$m_{\nabla^2 U}[\mu] = \frac{1}{\sqrt{p(p-1)}} \left( \int u''(x) \mu(x) dx - \int x u'(x) \mu(x) dx \right) \quad (\text{S45})$$

$$V_{\nabla^2 U}[\mu] = \frac{1}{p(p-1)} \left( \int u''^2(x) \mu(x) dx - \left( \int u''(x) \mu(x) dx \right)^2 \right) \quad (\text{S46})$$

where the explicit form of  $\log |\overline{\nabla^2 H}[\mu]|$  is found by replacing  $m_{\nabla^2 U}(\sigma) \rightarrow m_{\nabla^2 U}[\mu]$  and  $V_{\nabla^2 U}(\sigma) \rightarrow V_{\nabla^2 U}[\mu]$ . Since the integral in the numerator of  $\overline{\mathcal{N}_{\text{tot}}}$  is of the form  $\int D[\mu] \exp NS[\mu]$  it is evaluated by the saddle point technique in the large  $N$  limit. We start by writing down the functional derivative of the exponent w.r.t  $\mu$ :

$$\begin{aligned} \frac{\delta S}{\delta \mu} &= \alpha + \beta x^2 - \log \mu - 1 + \lim_{N \rightarrow \infty} \frac{1}{N} \frac{\delta \log |\det \nabla^2 H[\mu]|}{\delta \mu(x)} \\ &- \kappa^2 \frac{1}{2p} \left( u'^2(x) - 2 \left( \int y u'(y) \mu(y) dy \right) x u'(x) \right) = 0 \end{aligned} \quad (\text{S47})$$

where  $\alpha$  and  $\beta$  are Lagrange multipliers associated with the constraints on the normalization and second moment of  $\mu$ . We solve the above equation for  $\mu$  and the Lagrange multipliers  $\alpha$ ,  $\beta$  by expanding each in a perturbative expansion as follows:

$$\begin{aligned}\mu(x) &= \mu_0(x) + \kappa\mu_1(x) + \kappa^2\mu_2(x) + O(\kappa^3) \\ \beta &= \beta_0 + \kappa\beta_1 + \kappa^2\beta_2 + O(\kappa^3) \\ \alpha &= \alpha_0 + \kappa\alpha_1 + \kappa^2\alpha_2 + O(\kappa^3).\end{aligned}\tag{S48}$$

As the expression for  $\log |\overline{\det \nabla^2 H}[\mu]|$  is different for the cases  $p > 2$  and  $p = 2$  (which includes two cases of its own) we solve the saddle point equations for these cases separately.

### 1. The case of $p = 2$

Since for the case of  $p = 2$  the log-determinant is defined in a piecewise form, we evaluate the integral over the measure  $\mu$  separately for each case as well as for the boundary between the cases i.e.

$$\overline{\mathcal{N}}_{\text{tot}} = \int_{\mathcal{M}} D[\mu] \exp NS[\mu] = \int_{\mathcal{M}_<} D[\mu] \exp NS_{<}[\mu] + \int_{\mathcal{M}_{\geq}} D[\mu] \exp NS_{\geq}[\mu]\tag{S49}$$

where  $\mathcal{M}$  is the space of measures with a unit second moment,  $\mathcal{M}_<$ ,  $\mathcal{M}_{\geq}$  denote the sets  $\{\mu \in \mathcal{M} : \eta_2[\mu] < 2\}$ ,  $\{\mu \in \mathcal{M} : \eta_2[\mu] \geq 2\}$ , respectively, and  $S_{<}[\mu]$ ,  $S_{\geq}[\mu]$  denote the exponent in the cases  $\eta_2 < 2$  and  $\eta_2 \geq 2$ , respectively. Next, we evaluate each of the integrals by the saddle point method. Thus,  $\Sigma_{\text{tot}}$  is given by:

$$\Sigma_{\text{tot}} = \max \left\{ \sup_{\mu \in \mathcal{M}_<} S_{<}[\mu] \quad , \quad \sup_{\mu \in \mathcal{M}_{\geq}} S_{\geq}[\mu] \right\}\tag{S50}$$

In addition, we evaluate the integral on the boundary between the two branches to make sure that the global maximum is indeed found.

*a. The case of  $|\eta_2| \geq 2$*  For this case, the absolute value of the determinant of the Hessian and its functional derivative are given by:

$$\begin{aligned}\lim_{N \rightarrow \infty} \frac{1}{N} \log |\overline{\det \nabla^2 H}[\mu]| \\ = \frac{1}{2} (-1 + \log 2) + \kappa m_{\nabla^2 U}[\mu] - \frac{\kappa^2}{4} (m_{\nabla^2 U}[\mu] + 2V_{\nabla^2 U}[\mu]) + O(\kappa^3)\end{aligned}\tag{S51}$$

$$\begin{aligned}\lim_{N \rightarrow \infty} \frac{1}{N} \frac{\delta \log |\overline{\det \nabla^2 H}[\mu]|}{\delta \mu(x)} \\ = \frac{\kappa}{\sqrt{2}} (u''(x) - xu'(x)) - \frac{\kappa^2}{4} \left( u''(x)^2 + 2u''(x) \int \mu_0(y)u''(y) dy \right) + O(\kappa^3).\end{aligned}\tag{S52}$$

Substitution of this expression, along with the perturbative expansions given above in the saddle point condition in eq. (S47) gives the following equations:

$$O(\kappa^0) : \alpha_0 + \beta_0 x^2 - \log(\mu_0(x)) = 0\tag{S53}$$

$$O(\kappa^1) : \alpha_1 + \beta_1 x^2 - \frac{\mu_1(x)}{\mu_0(x)} + \frac{u''(x) - xu'(x)}{\sqrt{2}} = 0\tag{S54}$$

$$\begin{aligned}O(\kappa^2) : \alpha_2 + \beta_2 x^2 - \frac{\mu_2(x)}{\mu_0(x)} + \frac{\mu_1(x)^2}{2\mu_0(x)^2} - \frac{u''(x)^2 + u'(x)^2}{4} \\ - \frac{u''(x) - xu'(x)}{2} \int \mu_0(y)u''(y) dy = 0.\end{aligned}\tag{S55}$$

Note that we have used the identity for the law  $\mu_0(x) = \exp(-x^2/2)/\sqrt{2\pi}$ :

$$\int \mu_0(y)u''(y) dy = \int \mu_0(y)yu'(y) dy,\tag{S56}$$

and that the Lagrange multiplier expansion terms  $\alpha_l, \beta_l$  are determined by the constraints:

$$\begin{aligned} \int \mu_l(x) dx &= \delta_{l0} \\ \int x^2 \mu_l(x) dx &= \delta_{l0}. \end{aligned} \tag{S57}$$

Solving these equations gives the following expressions for  $\mu$ :

$$\mu_0(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \tag{S58}$$

$$\mu_1(x) = \frac{\mu_0(x)}{\sqrt{2}} \left( \frac{1-x^2}{2} \int \mu_0(y) y^2 (u''(y) - y u'(y)) dy + u''(x) - x u'(x) \right) \tag{S59}$$

$$\begin{aligned} \mu_2(x) = & \mu_0(x) \left( \alpha_2 x^2 + \beta_2 - \frac{u'(x)^2 + u''(x)^2}{4} - \frac{u''(x) - x u'(x)}{2} \left( \int \mu_0(y) u''(y) dy \right) \right) \\ & - \frac{\mu_1(x)^2}{2\mu_0(x)}, \end{aligned} \tag{S60}$$

where  $\alpha_2$  and  $\beta_2$  have a cumbersome expression, obtained self-consistently by imposing that (S57) hold. Using this expression, we can evaluate the average log-determinant:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H[\mu]| &= \frac{1}{2} (-1 + \log 2) + \frac{\kappa^2}{\sqrt{2}} \int dx \mu_1(x) (u''(x) - x u'(x)) \\ & - \frac{\kappa^2}{4} \left[ \int dx \mu_0(x) u''(x)^2 - \left( \int dx \mu_0(x) u''(x) \right)^2 \right] + O(\kappa^3) \end{aligned}$$

Next, we can substitute the expressions we found for the log-determinant and  $\mu$  in the expression for the annealed complexity to get

$$\begin{aligned} \overline{\Sigma}_{\eta \geq 2}^{(p=2)} &= \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{\mathcal{N}}_{\text{tot}} \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \log C_N - \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z} \\ &+ \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H[\mu]| - \int dx \mu \log \mu - \frac{\kappa^2}{4} \|\nabla U[\mu]\|^2. \end{aligned}$$

Let us first use that in the large  $N$  limit

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log C_N = \lim_{N \rightarrow \infty} \frac{1}{N} \log S_{N-1} - \frac{1}{2} \log(2\pi p) = \frac{1}{2} (1 - \log(p)). \tag{S61}$$

We may then evaluate the quantity involving  $\mathcal{Z}$  by the Laplace method, as

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z} \\ &= \max_{\mu, \alpha, \beta} \left[ - \int dx \mu(x) \log \mu(x) + \alpha \left( \int dx \mu(x) - 1 \right) + \beta \left( \int dx x^2 \mu(x) - 1 \right) \right]. \end{aligned} \tag{S62}$$

One can easily check that the maximum is reached for  $\mu(x) = \mu_0(x)$  and thus

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z} = - \int dx \mu_0(x) \log \mu_0(x). \tag{S63}$$

The entropic term can simply be developed up to second order in  $\kappa$ , yielding

$$\begin{aligned}
-\int dx \mu(x) \ln \mu(x) &= -\int dx \mu_0(x) \ln \mu_0(x) - \kappa \int dx \mu_1(x) \left(1 - \frac{1}{2} \ln(2\pi) - \frac{x^2}{2}\right) \\
&\quad - \kappa^2 \int dx \left[ \mu_2(x) \left(1 - \frac{1}{2} \ln(2\pi) - \frac{x^2}{2}\right) + \frac{\mu_1(x)^2}{2\mu_0(x)} \right] + O(\kappa^3) \\
&= -\int dx \mu_0(x) \ln \mu_0(x) - \frac{\kappa^2}{2} \int dx \frac{\mu_1(x)^2}{\mu_0(x)} + O(\kappa^3) \\
&= -\int dx \mu_0(x) \ln \mu_0(x) - \frac{\kappa^2}{2\sqrt{2}} \int dx \mu_1(x) (u''(x) - xu'(x)) + O(\kappa^3),
\end{aligned}$$

where we have used the explicit expression of  $\mu_0(x)$  as well as (S57). Finally, as the term  $-\kappa^2 \|\nabla U[\mu]\|^2/4$  is already of order  $\kappa^2$ , it can directly be evaluated for  $\mu_0$  for the purpose of our perturbative computation and reads

$$-\frac{\kappa^2}{4} \|\nabla U[\mu]\|^2 = -\frac{\kappa^2}{4} \left[ \int dx \mu_0(x) u'(x)^2 - \left( \int dx \mu_0(x) x u'(x) \right)^2 \right] + O(\kappa^3) \quad (\text{S64})$$

Gathering all terms and some straightforward algebra lead to the following complexity:

$$\begin{aligned}
\Sigma_{\eta \geq 2}^{(p=2)} &= \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{\mathcal{N}_{\text{tot}}} = \\
&\quad \frac{\kappa^2}{2} \left( -\frac{1}{4} \left( \int x^2 \mu_0(x) (u''(x) - xu'(x)) dx \right)^2 + \left( \int dx \mu_0(x) u''(x) \right)^2 \right. \\
&\quad \left. + \frac{1}{2} \int dx (x^2 - 1) \mu_0(x) u'(x)^2 - \int dx x \mu_0(x) u'(x) u''(x) \right) + O(\kappa^3).
\end{aligned} \quad (\text{S65})$$

By using the identity  $\overline{f'(x)}_{\mu_0} = \overline{xf(x)}_{\mu_0}$  (as  $\mu_0$  is the Gaussian) we get:

$$-\frac{1}{4} \left( \int x^2 \mu_0(x) (u''(x) - xu'(x)) dx \right)^2 + \left( \int dx \mu_0(x) u''(x) \right)^2 = 0 \quad (\text{S66})$$

$$\frac{1}{2} \int dx (x^2 - 1) \mu_0(x) u'(x)^2 - \int dx x \mu_0(x) u'(x) u''(x) = 0 \quad (\text{S67})$$

thus we are left with

$$\Sigma_{\eta \geq 2}^{(p=2)} = 0 + O(\kappa^3). \quad (\text{S68})$$

Next, to check the nature of the extremum, we calculate the second functional derivative of  $S[\mu]$  evaluated at  $\mu^*$ :

$$\left. \frac{\delta^2 S[\mu]}{\delta \mu(x) \delta \mu(y)} \right|_{\mu^*} = -\delta(x-y) \left( \frac{1}{\mu_0(x)} + O(\kappa) \right) + \frac{\kappa^2}{2} (u''(x)u''(y) + xu'(x)yu'(y)) + O(\kappa^3) \quad (\text{S69})$$

We consider the RHS as an operator in  $L^2(\mu_0)$ . By assumption, both  $u''$  and  $xu'$  are bounded in  $L^2(\mu_0)$ , and using that  $\kappa^2$  is small, we deduce that this operator is negative definite. In particular, the extremum we found is a (local) maximum.

Finally, we derive conditions for self-consistency of this solution by requiring that  $|\eta_2[\mu^*]| > 2$ . Substitution of  $\mu^*$  in  $\eta_2[\mu^*]$  gives:

$$\begin{aligned}
\eta_2[\mu^*] &= \frac{m_{\nabla^2 U}[\mu^*]}{\kappa V_{\nabla^2 U}[\mu^*]} = \frac{m_{\nabla^2 U}[\mu_1]}{V_{\nabla^2 U}[\mu_0]} + O(\kappa) \\
&= \frac{\frac{1}{2} \left( \int x^2 \mu_0(x) (u''(x) - xu'(x)) dx \right)^2 + \int \mu_0(x) (u''(x) - xu'(x))^2 dx}{\int \mu_0(x) u''(x)^2 dx - \left( \int \mu_0(x) u''(x) dx \right)^2}
\end{aligned} \quad (\text{S70})$$

The requirement  $\eta_2[\mu^*] > 2$  gives after some algebra and use of (S66) (we select  $m_{\nabla^2 U} > 0$  without loss of generality):

$$\eta_2[\mu^*] > 2 \iff \int \mu_0(x) (u''(x)^2 - u'(x)^2) dx < 0 \quad (\text{S71})$$

Thus this solution is self-consistent when the complexity of the other ( $\eta_2 < 2$ ) branch has negative complexity i.e.  $\Theta^{(2)}[u] < 0$ , see (S9). This is consistent with the picture of the  $\eta_2 > 2$  branch corresponding to zero leading order added complexity.



b. *The case of  $|\eta_2| < 2$*  For this case, the absolute value of the determinant of the Hessian and its functional derivative are given by:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H [\mu]| &= \frac{1}{2} (-1 + \log 2) \\ &+ \frac{1}{4} \kappa^2 \left( 2V_{\nabla^2 U} [\mu] + \frac{m_{\nabla^2 U} [\mu]^2}{\kappa^2 V_{\nabla^2 U} [\mu]} + m_{\nabla^2 U} [\mu]^2 \right) + O(\kappa^3). \end{aligned} \quad (\text{S72})$$

Then,

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \frac{\delta \log |\det \nabla^2 H [\mu]|}{\delta \mu(x)} &= \kappa \frac{m_{\nabla^2 U} [\mu_1]}{2V_{\nabla^2 U} [\mu_0]} (u''(x) - xu'(x)) + \\ &\frac{\kappa^2}{2} \left( \frac{m_{\nabla^2 U} [\mu_2]}{V_{\nabla^2 U} [\mu_0]} - \frac{m_{\nabla^2 U} [\mu_1] V_{\nabla^2 U} [\mu_1]}{(V_{\nabla^2 U} [\mu_0])^2} \right) (u''(x) - xu'(x)) \\ &- \kappa^2 \frac{m_{\nabla^2 U} [\mu_1]^2 (u''(x)^2 - 2u''(x) \int \mu_0(y) u''(y) dy)}{4(V_{\nabla^2 U} [\mu_0])^2} \\ &+ \frac{\kappa^2}{2} \left( u''(x)^2 - 2u''(x) \left( \int \mu_0(y) u''(y) dy \right) \right) + O(\kappa^3) + o_N(1). \end{aligned} \quad (\text{S73})$$

Substitution of this expression, along with the perturbative expansions for  $\mu$  in (S47) gives the following equations:

$$\begin{aligned} O(\kappa^0) : \alpha_0 + \beta_0 x^2 - \log(\mu_0(x)) &= 0 \\ O(\kappa^1) : \alpha_1 + \beta_1 x^2 - \frac{\mu_1(x)}{\mu_0(x)} + \frac{m_{\nabla^2 U} [\mu_1]}{2V_{\nabla^2 U} [\mu_0]} (u''(x) - xu'(x)) &= 0 \\ O(\kappa^2) : \alpha_2 + \beta_2 x^2 - \frac{\mu_2(x)}{\mu_0(x)} + \left( \frac{m_{\nabla^2 U} [\mu_2]}{2V_{\nabla^2 U} [\mu_0]} - \int \mu_0(y) u''(y) dy \right) (u''(x) - xu'(x)) \\ &+ \frac{1}{2} (u''(x)^2 - u'(x)^2) = 0 \end{aligned} \quad (\text{S74})$$

where we have used that all first order correction terms vanish in the second order equation, as can easily be deduced from the first order equation. The Lagrange multiplier expansion terms  $\alpha_l, \beta_l$  are determined by (S57) as in the calculation of the previous case. Solving these equations gives the following:

$$\begin{aligned} \mu_0(x) &= \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \\ \mu_1(x) &= 0, \\ \frac{\mu_2}{\mu_0}(x) \\ &= \alpha_2 x^2 + \beta_2 + \frac{1}{2} (u''(x)^2 - u'(x)^2) + (u''(x) - xu'(x)) \left( \frac{m_{\nabla^2 U} [\mu_2]}{2V_{\nabla^2 U} [\mu_0]} - \int \mu_0(y) u''(y) dy \right). \end{aligned}$$

Next, we can substitute the expression for  $\mu$  in the expression for the number of critical points, use that  $\mu_1 = 0$  and use the expansions derived above to find

$$\begin{aligned} \overline{\mathcal{N}_{\text{tot}}} &= C_N \exp N \left( \frac{1}{N} \log |\det \nabla^2 H [\mu]| - \frac{\kappa^2}{4} \|\nabla U [\mu_0]\|^2 + O(\kappa^3) + o_N(1) \right) \\ &= C_N \exp N \left( \frac{1}{2} (-1 + \log 2) + \kappa^2 \frac{1}{2} V_{\nabla^2 U} [\mu_0] - \frac{\kappa^2}{4} \|\nabla U [\mu_0]\|^2 + O(\kappa^3) + o_N(1) \right). \end{aligned} \quad (\text{S75})$$

Gathering all terms and some straightforward algebra lead to the following complexity:

$$\begin{aligned} \Sigma_{\eta_2 < 2}^{(p=2)} &= \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{\mathcal{N}_{\text{tot}}} = \kappa^2 \frac{1}{4} \int dx \mu_0(x) (u''(x)^2 - u'(x)^2) + O(\kappa^3) \\ &= \kappa^2 \Theta^{(2)} [u(x)] + O(\kappa^3) + o_N(1) \end{aligned} \quad (\text{S76})$$

To verify the type of extremum we found in this case, we proceed to evaluate the contribution of the integral on the boundary  $\eta_2 = 2$  in what follows.

c. *The case of  $|\eta_2| = 2$*  Since  $\log \det |\nabla^2 H|$  and  $\partial_{\eta_2} \log \det |\nabla^2 H|$  are continuous w.r.t to  $\eta_2$  at  $\eta_2 = 2$  (see (S36)), we can pick either one of the branches for the following calculation. We pick the expressions for the case  $\eta_2 > 2$ :

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H [\mu]| &= \frac{1}{2} (-1 + \log 2) + \kappa m_{\nabla^2 U} [\mu] + \frac{1}{4} \kappa^2 (m_{\nabla^2 U} [\mu] - 2V_{\nabla^2 U} [\mu]) \\ &+ \gamma (m_{\nabla^2 U} - \kappa 2V_{\nabla^2 U}) + O(\kappa^3 + o_N(1)). \end{aligned} \quad (\text{S77})$$

where we have introduced an additional Lagrange multiplier  $\gamma$  to enforce the constraint  $\eta_2[\mu] = 2$ . Taking the functional derivative of  $S[\mu]$  under this constraint and consecutive substitution of perturbative expansions give the following equations for the zero and first order in  $\kappa$ :

$$\begin{aligned} O(\kappa^0) : \alpha_0 + \beta_0 x^2 + \gamma_0 \frac{1}{\sqrt{2}} (u''(x) - xu'(x)) - \log(\mu_0(x)) &= 0 \\ O(\kappa^1) : \alpha_1 + \beta_1 x^2 - \frac{\mu_1(x)}{\mu_0(x)} + \left( \frac{1}{\sqrt{2}} + \gamma_1 \right) (u''(x) - xu'(x)) &+ \gamma_0 \sqrt{2} (-u''(x)^2 + u''(x) \overline{u''}_{\mu_0}) = 0. \end{aligned} \quad (\text{S78})$$

Solving these equations, under the three constraints on  $\mu$  (total mass equal to one, unit second moment and  $\eta_2[\mu] = 2$ ) gives the following solutions:

$$\begin{aligned} \mu_0(x) &= \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \\ \frac{\mu_1}{\mu_0}(x) &= \frac{1}{\sqrt{2}} \left( 1 + \frac{\Theta^{(2)}[u(x)]}{m_{\nabla^2 U}[\mu_{1, \eta_2 > 2}]} \right) \left( \frac{1}{2} (x^2 - 1) \int \mu_0(y) y^2 (u''(y) - yu'(y)) dy + u''(x) - xu'(x) \right). \end{aligned} \quad (\text{S79})$$

where we found that  $\gamma = \kappa \frac{\Theta^{(2)}[u(x)]}{\sqrt{2} m_{\nabla^2 U}[\mu_{1, \eta_2 > 2}]} + O(\kappa^2)$ . and denoted by  $\mu_{1, \eta_2 > 2}$  the first order correction to  $\mu^*$  in the case  $\eta_2 > 2$ . Substitution of these solutions in  $S[\mu]$  results in the following complexity:

$$\Sigma_{\eta_2=2} = \kappa^2 \left( \Theta^{(2)}[u] \left( 1 - \frac{c}{\sqrt{2}} \right) - \frac{c}{4} \left( \Theta^{(2)}[u] \right)^2 \right) + O(\kappa^3) \quad (\text{S80})$$

where we have denoted by  $c > 0$  the following expression:

$$c = 1 + \frac{\left( \overline{y^2 (u''(y) - yu'(y))}_{\mu_0} \right)^2}{\frac{1}{2} \left( \overline{y^2 (u''(y) - yu'(y))}_{\mu_0} \right)^2 + \left( \overline{u''(y) - yu'(y)}_{\mu_0} \right)^2}. \quad (\text{S81})$$

Now we are in a position to compare the complexity resulting from the boundary  $\eta_2 = 2$  with the complexity of the extrema found for the case  $\eta_2 < 2$ . As  $c > 0$  we can see that in the case where  $\Theta^{(2)} > 0$  we have that  $\Sigma_{\eta_2 < 2} > \Sigma_{\eta_2=2}$ . Thus, we conclude that the extremum found for the case  $\eta_2 < 2$  is a maximum.

## 2. The case of $p > 2$

For this case, the log determinant is given by:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log |\det \nabla^2 H [\mu]| &= \\ \frac{1}{2} \log p(p-1) + \kappa^2 \left( \frac{1}{2} V_{\nabla^2 U} [\mu] + m_{\nabla^2 U} [\mu]^2 \frac{(p-1)^2}{4p(p-2)} \right) &+ O(\kappa^3). \end{aligned} \quad (\text{S82})$$

It is clear from the previous cases that  $m_{\nabla^2 U} = O(\kappa)$  and thus, to leading order in  $\kappa$ , the complexity is given by:

$$\begin{aligned} \Sigma^{(p>2)} &= \frac{1}{2} \log(p-1) + \kappa^2 \left( \frac{1}{2} V_{\nabla^2 U} [\mu_0] - \frac{1}{2p} \|\nabla U [\mu_0]\|^2 \right) + O(\kappa^3) \\ &= \frac{1}{2} \log(p-1) + \kappa^2 \Theta^{(p)} [u(x)] + O(\kappa^3). \end{aligned} \quad (\text{S83})$$

#### IV. EVALUATION OF THE MEAN NUMBER OF CRITICAL POINTS OF A GIVEN INDEX

Here we aim to evaluate the number of critical points of a given index  $k$  (number of negative eigenvalues of the Hessian). We recall (S4). Taking identical steps as for the case of  $\overline{N}_{\text{tot}}$ , we arrive at the average value  $\overline{|\det(\nabla^2 H)| \delta(I(\nabla^2 H) - k)}$ . This is evaluated as follows:

$$\begin{aligned} \overline{|\det(\nabla^2 H)| \delta(I(\nabla^2 H) - k)} &= \overline{|\det(\nabla^2 \phi + \kappa \nabla^2 U)| \delta(I(\nabla^2 \phi + \kappa \nabla^2 U) - k)} \\ &= \int_{\mathbb{R}} \frac{\exp\left(-\frac{Ns^2}{2p^2}\right)}{\sqrt{2\pi p^2/N}} \overline{|\det \tilde{M} + \kappa \nabla^2 U + sI_{N-1}| \delta\left(I\left(\tilde{M} + \kappa \nabla^2 U + sI_{N-1}\right) - k\right)} \end{aligned} \quad (\text{S84})$$

with  $\tilde{M} \sim \sqrt{p(p-1)\frac{N-1}{N}} \text{GOE}_{N-1}$ . We first evaluate the GOE averaging as in [65] and follow by the integration over  $\mathbb{R}$  to get:

$$\begin{aligned} \overline{|\det \nabla^2 H(\sigma)| \delta(I(\nabla^2 H) - k)} \\ = \exp N \left( \int dx (\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U})(x) \log |x - s^*(a)| - \frac{s^*(a)^2}{2p^2} + o_N(1) \right), \end{aligned}$$

where we have explicitly assumed that  $k$  is extensive i.e. the ratio  $a = k/N$  is fixed when the large  $N$  limit is taken. Also,  $s^*(a)$  is determined by the index constraint:

$$\int_{-\infty}^0 (\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U})(x + s^*(a)) dx = a. \quad (\text{S85})$$

Note that the above expression is very similar in form to the one considered in the case of the total complexity with the difference that here  $s$  is set by the index whereas in the previous case the complexity is the supremum over  $s$ . Thus, we can utilize the same expressions derived in the previous section to give:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log \overline{|\det \nabla^2 H(\sigma)| \delta(I(\nabla^2 H) - k)} &= \Phi(\eta) + \frac{1}{2} \log p(p-1) \\ &- \frac{p-2}{4p} \eta^2 + \kappa \frac{p-1}{p} m_{\nabla^2 U} \eta + \kappa^2 \left( \frac{V_{\nabla^2 U}}{2} \left(1 - \frac{(p-1)\eta^2}{p}\right) - m_{\nabla^2 U}^2 \frac{p-1}{2p} \right) + O(\kappa^3) \end{aligned} \quad (\text{S86})$$

where  $\eta$  is defined as above. Thus, the choice of index  $k = aN$  fixes the parameter  $\eta$ .

As in the case of the total complexity, we proceed to carry out the integration over the  $N$ -sphere by passing to functional integration over the measure  $\mu$ . The functional derivative of  $\frac{1}{N} \log |\det \nabla^2 H[\mu]|$  is given by:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \frac{\delta \log |\det \nabla^2 H[\mu]|}{\delta \mu(x)} &= \kappa \sqrt{\frac{p-1}{p^3}} \eta (u''(x) - xu'(x)) \\ &+ \frac{\kappa^2}{p(p-1)} \left( -\frac{1}{2} \left( \frac{p-1}{p} \eta^2 - 1 \right) u''(x)^2 + u''(x) \left( \frac{p-1}{p} \eta^2 - 1 \right) \int \mu_0(y) u''(y) dy \right) + O(\kappa^3). \end{aligned} \quad (\text{S87})$$

Substitution of this expression, along with the perturbative expansions given above in eq. (S47) gives the following equations:

$$\begin{aligned} O(\kappa^0) : \alpha_0 + \beta_0 x^2 - \log(\mu_0(x)) &= 0 \\ O(\kappa^1) : \alpha_1 + \beta_1 x^2 - \frac{\mu_1(x)}{\mu_0(x)} + \sqrt{\frac{p-1}{p^3}} \eta (u''(x) - xu'(x)) &= 0. \end{aligned} \quad (\text{S88})$$

Solving these equations, along with the constraints as above, gives the following solutions:

$$\begin{aligned} \mu_0(x) &= \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \\ \mu_1(x) &= \mu_0(x) \sqrt{\frac{p-1}{p^3}} \eta \left( \frac{1}{2} (x^2 - 1) \int \mu_0(y) y^2 dy (u''(y) - yu'(y)) + u''(x) - xu'(x) \right). \end{aligned} \quad (\text{S89})$$

Here we focus only on the solutions up to  $O(\kappa)$  since  $\mu_2$  does not affect the complexity to leading order as found for the total complexity. Substitution of these results in the expression for the log-determinant leads to the following expression for the number of critical points:

$$\begin{aligned}\overline{\mathcal{N}}_a &= C_N \exp N \left( \frac{1}{N} \log |\det \nabla^2 H [\mu]| - \kappa^2 \frac{1}{2} \int dx \frac{\mu_1^2}{\mu_0} - \frac{\kappa^2}{2p} \|\nabla U [\mu_0]\|^2 + O(\kappa^3) \right) \\ &= C_N \exp N \left( \Phi(\eta) + \frac{1}{2} \log p(p-1) - \frac{p-2}{4p} \eta^2 \right. \\ &\quad \left. + \kappa^2 \left( \frac{p-1}{p} \eta m_{\nabla^2 U} [\mu_1] + \frac{1}{2} \left( 1 - \frac{\eta^2(p-1)}{p} \right) V_{\nabla^2 U} [\mu_0] - \frac{1}{2} \int dx \frac{\mu_1^2}{\mu_0} - \frac{1}{2p} \|\nabla U [\mu_0]\|^2 \right) \right. \\ &\quad \left. + O(\kappa^3), \right.\end{aligned}\tag{S90}$$

where as before  $C_N = (\frac{1}{2\pi p})^{N/2} S_{N-1}(\sqrt{N})$ . Next, we observe that  $\mu_{1,k} = \sqrt{\frac{p-1}{p^3}} \eta \mu_{1,\eta>2}$  (where  $\mu_{1,\eta>2}$  denotes the first order term  $\mu_1$  for the case  $\eta_2 > 2$  and  $p = 2$ ) and make this substitution to get:

$$\begin{aligned}\Sigma_a^{(p)}[u] &= \lim_{N \rightarrow \infty} \frac{1}{N} \ln \overline{\mathcal{N}}_a = \Phi(\eta) + \frac{1}{2} \log p(p-1) - \frac{p-2}{4p} \eta^2 + \\ &\quad \kappa^2 \left( \frac{2(p-1)}{p^3} \eta^2 m_{\nabla^2 U} [\mu_{1,\eta>2}^{(2)}] + \frac{1}{p(p-1)} \left( 1 - \frac{\eta^2(p-1)}{p} \right) V_{\nabla^2 U}^{(2)} [\mu_0] \right. \\ &\quad \left. - \frac{p-1}{p^3} \eta^2 \int dx \frac{(\mu_{1,\eta>2}^{(2)})^2}{\mu_0} - \frac{1}{2p} \|\nabla U [\mu_0]\|^2 \right) + O(\kappa^3).\end{aligned}\tag{S91}$$

Simple algebraic manipulations give the following expression:

$$\begin{aligned}\Sigma_a^{(p)}[u] &= \lim_{N \rightarrow \infty} \frac{1}{N} \ln \overline{\mathcal{N}}_a \\ &= \frac{1}{2} \log(p-1) - \frac{p-2}{4p} \eta^2 + \kappa^2 \left( 1 - \eta^2 \frac{p-1}{p^2} \right) \Theta^{(p)}[u] + O(\kappa^3)\end{aligned}\tag{S92}$$

where we have used:

$$m_{\nabla^2 U}^{(2)} [\mu_{1,\eta>2}] - \frac{1}{2} \int dx \frac{(\mu_{1,\eta>2}^{(2)})^2}{\mu_0} - \frac{1}{2} V_{\nabla^2 U}^{(2)} [\mu_0] - \frac{1}{4} \|\nabla U [\mu_0]\|^2 = 0\tag{S93}$$

and

$$\Theta^{(p)}[u] = \frac{1}{2} V_{\nabla^2 U}^{(p)} [\mu_0] - \frac{1}{2p} \|\nabla U [\mu_0]\|^2.\tag{S94}$$

where  $m_{\nabla^2 U}^{(p)}$  and  $V_{\nabla^2 U}^{(p)}$  denote the normalized mean and variance of  $\nu_{\nabla^2 U}$ , respectively, for the case of a  $p$ -spin model (see (S46))

## Appendix A: Distribution of Random Fields

In [60] Lemma 1, the covariances of the field  $\phi$ , and its covariant derivatives  $\nabla\phi$ , and  $\nabla^2\phi$  at a given point were derived for the of  $p$ -spin spherical models and are given by:

$$\begin{aligned}\mathbb{E}[\phi^2] &= N \\ \mathbb{E}[(\nabla\phi)_m \phi] &= \mathbb{E}[(\nabla\phi)_m (\nabla^2\phi)_{np}] = 0 \\ \mathbb{E}[(\nabla\phi)_m (\nabla\phi)_n] &= p\delta_{mn} \\ \mathbb{E}[\phi (\nabla^2\phi)_{mn}] &= -p\delta_{mn} \\ \mathbb{E}[(\nabla^2\phi)_{mn} (\nabla^2\phi)_{lp}] &= \frac{p(p-1)}{N} (\delta_{ml}\delta_{np} + \delta_{mp}\delta_{nl}) + \frac{p^2}{N} \delta_{mn}\delta_{lp}\end{aligned}\tag{A1}$$

Note that here  $m = 1, \dots, N-1$  and the coordinates of the gradient and the Hessian are with respect to an orthonormal system in the tangent space to the sphere at  $p$ .

### Appendix B: $N$ -Sphere covariant Hessian for Single Particle Potential

The covariant gradient on the sphere can be written concisely as a Euclidean gradient followed by a projection to the sphere tangent surface:

$$\nabla_{\mathbb{S}^N} = (I_N - \hat{\sigma} \otimes \hat{\sigma}) \cdot \nabla_{\mathbb{R}^N} \quad (\text{B1})$$

where we have denoted for clarity  $\nabla_{\mathbb{S}^N}$  as the covariant gradient on the sphere and  $\nabla_{\mathbb{R}^N}$  the Euclidean gradient in  $\mathbb{R}^N$ . Also,  $I_N$  denotes the unit operator in  $\mathbb{R}^N$  and  $\hat{\sigma}$  is the radial unit vector. The covariant Hessian is a result of two repeated applications of the covariant Hessian:

$$\nabla_{\mathbb{S}^N}^2 U = (I_N - \hat{\sigma} \otimes \hat{\sigma}) \cdot \nabla_{\mathbb{R}^N} ((I_N - \hat{\sigma} \otimes \hat{\sigma}) \cdot \nabla_{\mathbb{R}^N} U) \quad (\text{B2})$$

expanding this expression we get

$$\nabla_{\mathbb{S}^N}^2 U = (I_N - \hat{\sigma} \otimes \hat{\sigma}) \cdot (\nabla_{\mathbb{R}^N}^2 U - \nabla_{\mathbb{R}^N} (\hat{\sigma} \otimes \hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U)). \quad (\text{B3})$$

We now turn to evaluate the last term on the right hand side in (B3) for the case of single particle  $U$ :

$$\begin{aligned} (\nabla_{\mathbb{R}^N} (\hat{\sigma} \otimes \hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U))_{ij} &= \frac{\partial}{\partial x_i} \sum_m \frac{x_j x_m}{\sum_n x_n^2} \frac{\partial U}{\partial x_m} = \\ &= \sum_m \left( \frac{\delta_{ij} x_m + x_j \delta_{im}}{\sum_n x_n^2} - 2 \frac{x_i x_j x_m}{(\sum_n x_n^2)^2} \right) \frac{\partial U}{\partial x_m} + \sum_m \frac{x_j x_m}{\sum_n x_n^2} \frac{\partial^2 U}{\partial x_m \partial x_i} = \\ &= \frac{1}{\sqrt{\sum_n x_n^2}} \left( \delta_{ij} \sum_m \hat{\sigma}_m \frac{\partial U}{\partial x_m} + \hat{\sigma}_j \frac{\partial U}{\partial x_i} - 2 (\hat{\sigma} \otimes \hat{\sigma})_{ij} \sum_m \hat{\sigma}_m \frac{\partial U}{\partial x_m} \right) + \sum_m (\hat{\sigma} \otimes \hat{\sigma})_{jm} \frac{\partial^2 U}{\partial x_m \partial x_i} \end{aligned} \quad (\text{B4})$$

where we have used the following identity:

$$\frac{\partial}{\partial x_i} \left( \frac{x_j x_m}{\sum_n x_n^2} \right) = \frac{\delta_{ij} x_m + x_j \delta_{im}}{\sum_n x_n^2} - 2 \frac{x_i x_j x_m}{(\sum_n x_n^2)^2}. \quad (\text{B5})$$

Now, we can write this expression in vector notation as follows:

$$\nabla_{\mathbb{R}^N} (\hat{\sigma} \otimes \hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U) = \frac{1}{\sqrt{N}} ((\hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U) I_N + \hat{\sigma} \otimes \nabla_{\mathbb{R}^N} U - 2 (\hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U) \hat{\sigma} \otimes \hat{\sigma}) + (\hat{\sigma} \otimes \hat{\sigma}) \cdot \nabla_{\mathbb{R}^N}^2 U \quad (\text{B6})$$

where we have used the fact that the derivative is calculated on  $\sqrt{N}\mathbb{S}^{N-1}$  i.e  $\sum_n x_n^2 = N$ . For the purpose of the asymptotic calculation carried out in this work all terms of rank  $O(1)$  can be neglected. Observation of the above expressions yields that:

$$\nabla_{\mathbb{S}^N}^2 U = \nabla_{\mathbb{R}^N}^2 U - \frac{1}{\sqrt{N}} (\hat{\sigma} \cdot \nabla_{\mathbb{R}^N} U) I_N + M_1 \quad (\text{B7})$$

where  $M_1$  denotes a matrix of rank  $O(1)$ .

### Appendix C: Evaluating The Density of the Empirical Spectral Distribution of $\nabla^2 H = \nabla^2 \phi + \kappa \nabla^2 U$

Denoting by  $G(z)$  the Stieltjes transform of  $\rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U}$ ,  $G$  is known to obey the following implicit formula [76] Proposition 2:

$$G(z) = \int \frac{\nu_{\kappa \nabla^2 U}(x) dx}{z - t^2 G(z) - x} = \int \frac{\frac{1}{\kappa} \nu_{\nabla^2 U}(\frac{x}{\kappa}) dx}{z - t^2 G(z) - x}, \quad z \in \mathbf{C}_+ \quad (\text{C1})$$

where  $t^2 = p(p-1)$  and we have used the identity  $\nu_{\kappa \nabla^2 U} = \frac{1}{\kappa} \nu_{\nabla^2 U}(\frac{x}{\kappa})$ . To retrieve  $\nu_{\nabla^2 H} = \rho_{\text{sc}} \boxplus \nu_{\kappa \nabla^2 U}$  one employs the inverse transform:

$$\psi(x) = \frac{1}{\pi} \lim_{\epsilon \rightarrow 0^+} \text{Im} G(x + i\epsilon). \quad (\text{C2})$$

a. *Evaluating the Free Convolution with a Small Perturbation*

Here, we employ perturbation theory in  $\kappa$  to solve the implicit equation (C1). First we make the change of variables  $s = x/\kappa$  followed by the change of variables  $s = m_{\nabla^2 U} + s'$  with  $m_{\nabla^2 U}$  denoting the expectation value of  $\nu_{\nabla^2 U}$ ,

$$\int \frac{\frac{1}{\kappa} \nu_{\nabla^2 U} \left(\frac{x}{\kappa}\right) dx}{z - t^2 G(z) - x} = \int \frac{\nu_{\nabla^2 U}(s) ds}{z - t^2 G(z) - \kappa s} = \int \frac{\nu_{\nabla^2 U}(m_{\nabla^2 U} + s') ds'}{z - t^2 G(z) - \kappa(m_{\nabla^2 U} + s')}. \quad (\text{C3})$$

Now we can expand the integral in a power series in  $\kappa$  as follows:

$$\begin{aligned} \int \frac{\nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds'}{z - t^2 G(z) - \kappa(m_{\nabla^2 U} + s')} &= \frac{1}{z - t^2 G(z) - \kappa m_{\nabla^2 U}} \int \frac{\nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds'}{1 - \kappa \frac{s'}{z - t^2 G(z) - \kappa m_{\nabla^2 U}}} \\ &= \frac{1}{z - t^2 G(z) - \kappa m_{\nabla^2 U}} + \frac{\kappa^2}{(z - t^2 G(z) - \kappa m_{\nabla^2 U})^3} \int s'^2 \nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds' + O(\kappa^3) \end{aligned} \quad (\text{C4})$$

where we have used the simple fact that  $\int s' \nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds' = 0$ . Reordering (C1) results in:

$$(z - \kappa m_{\nabla^2 U}) G - t^2 G^2 - 1 = \frac{\kappa^2}{(z - \kappa m_{\nabla^2 U} - t^2 G)^2} \int s'^2 \nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds' + O(\kappa^3). \quad (\text{C5})$$

We solve this perturbative equation, up to order  $\kappa^2$ , by introducing the following ansatz:

$$G(z) = \frac{z - \kappa m_{\nabla^2 U} + \sqrt{(z - \kappa m_{\nabla^2 U})^2 - 4T^2}}{2T^2} \quad (\text{C6})$$

where we expand the support  $T$  in a power series:

$$T^2 = t^2 + \kappa T_1 + \kappa^2 T_2 + O(\kappa^3). \quad (\text{C7})$$

Substitution in the equation above results in:

$$\begin{aligned} T_1 &= 0 \\ T_2 &= \int s'^2 \nu_{\nabla^2 U}(s' + m_{\nabla^2 U}) ds' \equiv V_{\nabla^2 U}. \end{aligned} \quad (\text{C8})$$

It is seen that up to second order in  $\kappa$  the free convolution is a slightly wider semicircle distribution with a width of  $t^2 + \kappa^2 V_{\nabla^2 U}$ :

$$\psi(x) = 1_{x \in [-2T + \kappa m_{\nabla^2 U}, 2T + \kappa m_{\nabla^2 U}]} \frac{1}{2\pi(t^2 + \kappa^2 V_{\nabla^2 U})} \sqrt{-(x - \kappa m_{\nabla^2 U})^2 + 4(t^2 + \kappa^2 V_{\nabla^2 U})}. \quad (\text{C9})$$

#### Appendix D: Positivity of $\Theta^{(2)}[f(x)]$ for potentials with $f'(0) = 0$

Here we show that  $\Theta^{(2)}[u(x)]$  is positive when  $u'(0) = 0$  or equivalently  $a_1 = 0$  in the Taylor expansion of  $u(x)$ :

$$u(x) = \sum_{n=1}^{\infty} a_n x^n \quad (\text{D1})$$

Explicit integration gives

$$\Theta \left[ \sum_{n=1}^{\infty} a_n x^n \right] = -\frac{a_1^2}{4} - a_1 \sum_{n \geq 3} A_n a_n + \sum_{n \geq 3} B_n a_n^2 + \sum_{m \geq 3} \sum_{n \geq 3} D_{n,m} a_m a_n \quad (\text{D2})$$

where we have denoted:

$$A_n = \frac{1}{4\sqrt{2\pi}} 2^{n/2} c_{n+1} n \Gamma\left(\frac{n}{2}\right) \quad (\text{D3})$$

$$B_n = \frac{1}{4\sqrt{2\pi}} 2^{n-\frac{3}{2}} (n-2)^2 n^2 \Gamma\left(n - \frac{3}{2}\right) \quad (\text{D4})$$

$$D_{n,m} = \frac{1}{4\sqrt{2\pi}} (m-2)m(n-2)n 2^{\frac{1}{2}(m+n-3)} c_{n+m} \Gamma\left(\frac{1}{2}(m+n-3)\right) \quad (\text{D5})$$

with  $c_n = ((-1)^n + 1)$ .

Next, we show that  $\Theta[u] > 0$  for  $a_1 = 0$ . As the term  $\sum_{n \geq 3} B_n a_n^2$  is obviously positive it is left to show that  $\sum_{m \geq 3} \sum_{n \geq 3} D_{n,m} a_m a_n > 0$ . Using the inequality  $\Gamma\left(\frac{1}{2}(m+n-3)\right) > \Gamma\left(\frac{1}{2}(m-3/2)\right) \Gamma\left(\frac{1}{2}(n-3/2)\right)$  we can write:

$$\begin{aligned} \sum_{m \geq 3} \sum_{n \geq 3} D_{n,m} a_m a_n &= \sum_{m \geq 3} \sum_{n \geq 3} m(m-2)n(n-2)2^{\frac{1}{2}(m+n-3)} c_{n+m} \Gamma\left(\frac{1}{2}(m+n-3)\right) a_m a_n > \\ &\left( \sum_{n \geq 3} a_n n(n-2)2^{\frac{1}{2}(n-3/2)} \Gamma\left(\frac{1}{2}(n-3/2)\right) \right)^2 \\ &+ \left( \sum_{n \geq 3} a_n n(n-2)(-1)^n 2^{\frac{1}{2}(n-3/2)} \Gamma\left(\frac{1}{2}(n-3/2)\right) \right)^2 \end{aligned} \quad (\text{D6})$$

and we have the desired result.