

# LAGRANGIAN TRACES FOR THE JOHNSON FILTRATION OF THE HANDLEBODY GROUP

QUENTIN FAES

ABSTRACT. In this paper, we define trace-like operators on a subspace of the space of derivations of the free Lie algebra generated by the first homology group  $H$  of a surface  $\Sigma$ . This definition depends on the choice of a Lagrangian of  $H$ , and we call these operators the *Lagrangian traces*. We suppose that  $\Sigma$  is the boundary of a handlebody with first homology group  $H'$ , and we show that the Lagrangian traces corresponding to the Lagrangian  $\text{Ker}(H \rightarrow H')$  vanish on the image by the Johnson homomorphisms of the elements of the Johnson filtration that extend to the handlebody.

## CONTENTS

1.	The Lagrangian traces	3
2.	Magnus representations for the handlebody group	4
2.1.	Morita's trace map and the Magnus representation	4
2.2.	The handlebody case	4
	References	6

Let  $\Sigma := \Sigma_{g,1}$  be a compact connected oriented surface of genus  $g$  with one boundary component, and denote by  $\mathcal{M} := \mathcal{M}(\Sigma)$  the mapping class group of  $\Sigma$ . Let  $V_g$  be a handlebody whose boundary minus a disk is identified with  $\Sigma$ . The handlebody group  $\mathcal{A}$  is the subgroup of mapping classes of  $\mathcal{M}$  that admit an extension to the handlebody. It is a non-normal subgroup of the mapping class group with infinite index, hence its study as a subgroup of  $\mathcal{M}$  is difficult.

It is known, by the Dehn-Nielsen theorem, that  $\mathcal{M}$  can be studied via its action on the fundamental group  $\pi := \pi_1(\Sigma_{g,1})$  of the surface, which is a free group on  $2g$  generators. Let us denote  $H := H_1(\Sigma_{g,1})$ . Similarly, let  $\pi' := \pi_1(V_g)$  and  $H' := H_1(V_g)$  denote respectively the fundamental group and the first homology group of the handlebody. The abelian groups  $H$  and  $H'$  are the abelianization of  $\pi$  and  $\pi'$ , respectively. We also denote  $\mathbb{A}$  the kernel of the projection  $\pi \rightarrow \pi'$  and  $A$  the kernel of the projection  $H \rightarrow H'$ . We have the following commutative exact diagram, induced by the inclusion  $\Sigma_{g,1} \subset V_g$ :

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \mathbb{A} & \longrightarrow & \pi & \longrightarrow & \pi' & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & A & \longrightarrow & H & \longrightarrow & H' & \longrightarrow & 0.
\end{array}$$

The group  $\mathcal{A}$  actually coincides with the subgroup of  $\mathcal{M}$  that preserves  $\mathbb{A}$  [3].

Recall that the intersection form of  $\Sigma$  induces a symplectic form  $\omega : H \otimes H \rightarrow \mathbb{Z}$ , hence  $H$  is naturally isomorphic to  $H^*$  via the map  $x \mapsto \omega(x, -)$ . The action of  $\mathcal{M}$  on the homology of the surface preserves the symplectic form  $\omega$ , inducing a representation  $\rho_0 : \mathcal{M} \rightarrow \text{Sp}(H)$  whose kernel is, by definition, the Torelli group  $\mathcal{I}$ . The duality  $H \simeq H^*$  actually restricts to an isomorphism between the lagrangian  $A$  and  $H'^*$  (see [1, Section 4.2]). We fix a basis  $(\alpha_1, \alpha_2 \dots \alpha_g, \beta_1, \beta_2 \dots \beta_g)$  of  $\pi$ , such that:

- it induces a symplectic basis of  $H$ .
- $\mathbb{A}$  is normally generated by the  $(\alpha_i)_{1 \leq i \leq g}$
- the image  $(\beta'_i)_{1 \leq i \leq g}$  of the family  $(\beta_i)_{1 \leq i \leq g}$  in  $\pi'$  form a basis of  $\pi'$ .

Such a choice of basis for  $\pi$  is induced by the curves drawn in Figure 1. Notice that the curves  $\alpha_i$  bound disks in the handlebody.




FIGURE 1. A basis of  $\pi$  adapted to the inclusion  $\Sigma \subset V_g$ .

The  $k$ -th term  $J_k$  of the Johnson filtration is defined as the subgroup of  $\mathcal{M}$  of elements acting trivially on  $\pi$  modulo the commutators of length  $k+1$ . In particular the first term of this filtration is  $\mathcal{I}$ . The Johnson filtration  $(J_k)_{k \geq 1}$  of the mapping class group is a separating filtration of  $\mathcal{M}$  and its associated graded space  $\text{Gr}^J := \bigoplus_{k \geq 1} \frac{J_k}{J_{k+1}}$  has been extensively studied. The latter is in particular a graded Lie ring. The so-called Johnson homomorphisms  $(\tau_k)_{k \geq 1}$  induce an embedding  $\tau$  of the graded space associated to the Johnson filtration to the graded Lie ring  $D(H) := \bigoplus_{k \geq 1} D_k(H)$  of symplectic derivations of  $\mathcal{L}(H)$ , the free graded Lie algebra generated by  $H$ . Recall that a derivation of the free Lie algebra can be identified via the restriction to  $H$ , with an element of  $H^* \otimes \mathcal{L}(H) \simeq H \otimes \mathcal{L}(H)$ . The space  $D_k(H)$  then corresponds to the kernel of the bracket from  $H \otimes \mathcal{L}_{k+1}(H)$  to  $\mathcal{L}_{k+2}(H)$ . The determination of the cokernel of  $\tau$  is a question of major importance. In [9], Morita defined a family of trace-like operators that detect a part of this cokernel (but not all of it).

In this paper, we are interested in the intersection of the Johnson filtration with  $\mathcal{A}$ . We set  $\mathcal{A}_k := \mathcal{A} \cap J_k$ . The  $\mathcal{A}J$  filtration  $(\mathcal{A}_k)_{k \geq 1}$  is also a separating filtration of  $\mathcal{A}$ . Hence we would like to understand its associated graded  $\text{Gr}^{\mathcal{A}J}$ , or, equivalently, the subspace  $\tau_k(\mathcal{A}_k) \subset D_k(H)$ . Note that the study of the  $\mathcal{A}J$  filtration is important for the study of Heegaard splittings and finite-type invariants of integral homology 3-spheres [11, 12, 2]. In [10], Omori gave a generating set of the first term of the  $\mathcal{A}J$  filtration. In [6], Levine noticed that the image of the restriction of  $\tau_k$  to  $\mathcal{A}_k$  is included in the kernel  $\mathcal{G}_k$  of the natural projection  $D_k(H) \rightarrow D_k(H')$ , where  $D_k(H') := \text{Ker}(H' \otimes \mathcal{L}_{k+1}(H') \rightarrow \mathcal{L}_{k+2}(H'))$ . Hence the Johnson homomorphisms induce a well-defined embedding  $\text{Gr}^{\mathcal{A}J} \rightarrow \bigoplus_{k \geq 1} \mathcal{G}_k$ . A computation of Morita in [8] shows that  $\mathcal{G}_1 = \tau_1(\mathcal{A}_1)$ . We will show that this equality does not hold in general. In [1], the author defined a family of trace-like operators  $(\text{Tr}_k^A)_{k \geq 1}$ , defined on  $\mathcal{G}_k$ , and valued in  $S^k(H')$ . We shall refer to these maps as the *Lagrangian traces*. In the same paper, using a variation formula for the Casson invariant from [8], the author shows that  $\tau_2(\mathcal{A}_2) = \text{Ker}(\text{Tr}_2^A) \cap \text{Im}(\tau_2)$ . Here, adapting the strategy of Morita in [9] to the handlebody case, we prove the following theorem:

**Theorem A.** *Let  $g \geq 2$ , and  $k \geq 2$ , then the map  $\text{Tr}_k^A$  vanishes on  $\tau_k(\mathcal{A}_k) \subset \mathcal{G}_k$ .*

The proof of Theorem A uses the Magnus representation of the mapping class group, and also its reduction to the handlebody case. We will recall only what is needed here, and refer to [13] for a survey on the subject. We define in Section 2 two Magnus “representations”  $r^A : \mathcal{A} \rightarrow \text{GL}(g, \mathbb{Z}[\pi'])$  and its abelian version  $r^{A, \text{a}} : \mathcal{A} \rightarrow \text{GL}(g, \mathbb{Z}[H'])$ . These are representations only in a twisted sense. For any group  $G$ , we shall denote  $I_G$  the augmentation ideal of  $\mathbb{Z}[G]$ . For short, we set  $I := I_\pi$ .

**Theorem B.** *Let  $g \geq 2$ , and  $\varphi \in \mathcal{A}_1$ , then  $\det(r^{A, \text{a}}(\varphi)) \in H' \subset \mathbb{Z}[H']$  and*

$$\det(r^{A, \text{a}}(\varphi)) = \text{Tr}_1^A(\tau_1(\varphi)).$$

---

This research has been supported by the project “AlMaRe” (ANR-19-CE40-0001-01) of the ANR and the project “ITIQ-3D” of the Région Bourgogne Franche-Comté. The IMB receives support from the EIPHI Graduate School (ANR-17-EURE-0002)

Consequently, the determinant of the Magnus representation is trivial on  $\mathcal{A}_k$ , for  $k \geq 2$ .

In the first section, we review the definition of the trace-like operators  $(\text{Tr}_k^A)_{k \geq 1}$ , and study their properties. In the second section, we will prove Theorem B and Theorem A by considering the magnus representation of the handlebody group, and relating it to the A-traces. Throughout the paper, we shall detail the connections with Morita's work in [9].

**Acknowledgment.** The author would like to thank Gwénaél Massuyeau for his corrections, as well as Renaud Detcherry and Jules Martel for discussions about the Magnus representation.

## 1. THE LAGRANGIAN TRACES

Before recalling the definition of the Lagrangian trace  $\text{Tr}_k^A$ , let us recall a few facts about the Johnson homomorphisms  $(\tau_k)_{k \geq 1}$  and the spaces  $D_k(H)$ . The  $k$ -th Johnson homomorphism of  $\varphi \in J_k$  associates to the class  $\{x\} \in H$  of an element  $x$  in  $\pi$ , the class of  $\phi(x)x^{-1}$  in  $\Gamma_{k+1}\pi/\Gamma_{k+2}\pi \simeq \mathcal{L}_{k+1}(H)$ . This defines a map  $\tau_k : J_k \rightarrow \text{Hom}(H, \mathcal{L}_{k+1}(H))$ . The target space can be identified with the space of derivations of degree  $k$  of the free Lie algebra. Furthermore, one can prove that such a derivation sends  $\omega$ , the bivector dual to the symplectic form, to 0. Such an element will be called a *symplectic* derivation. These form a graded subalgebra  $D(H)$ . Any element in  $D_k(H)$  has a *Morita trace*, which coincides with the evaluation:

$$D_k(H) \subset H^* \otimes \mathcal{L}_{k+1}(H) \longrightarrow H^* \otimes T_{k+1}(H) \longrightarrow H^* \otimes H \otimes T_k(H) \longrightarrow T_k(H) \longrightarrow S^k(H)$$

where  $\mathcal{L}(H)$  embeds in the associative tensor algebra  $T(H)$  via the rule  $[a, b] = a \otimes b - b \otimes a$ , and  $H^* \otimes H \rightarrow \mathbb{Z}$  is the evaluation. Morita's trace map actually vanishes on the image of the Johnson homomorphism [9]. Let us recall the definition of the trace-like operators defined in [1]. As we recalled, there is a canonical isomorphism between  $A$  and  $(H')^*$  given by the intersection pairing

$$\omega' : A \otimes H' \longrightarrow \mathbb{Z}$$

which is induced by the symplectic form  $\omega$ . We denote  $(\omega')^{1,2}$  the contraction of the first two tensors in  $A \otimes T(H')$  by  $\omega'$ . An element of  $\mathcal{G}_k$  naturally induces an element of  $A \otimes \mathcal{L}_{k+1}(H')$ . Indeed  $A \otimes \mathcal{L}_{k+1}(H')$  is the kernel of  $H \otimes \mathcal{L}_{k+1}(H') \rightarrow H' \otimes \mathcal{L}_{k+1}(H')$ . Then the map  $\text{Tr}_k^A$  is defined by

$$\text{Tr}_k^A : \mathcal{G}_k \longrightarrow A \otimes \mathcal{L}_{k+1}(H') \xrightarrow{i} A \otimes T_{k+1}(H') \xrightarrow{(\omega')^{1,2}} T_k(H') \longrightarrow S^k(H').$$

**Remark 1.1.** We actually have  $A \simeq H'^*$ , hence an element of  $\mathcal{G}_k$  naturally induces an element of  $H'^* \otimes \mathcal{L}_{k+1}(H')$ , i.e. a derivation of  $\mathcal{L}(H')$ . The Lagrangian traces that we define here are hence a handlebody version of the traces defined by Morita.

Note that  $\mathcal{G}$  is a graded Lie subalgebra of  $D(H)$ . This subalgebra is not stable by the action of  $\text{Sp}(H)$ , but it is stable by  $\rho_0(\mathcal{A})$ , the image of  $\mathcal{A}$  in  $\text{Sp}(H)$ . This is precisely the subgroup of  $\text{Sp}(H)$  that preserves  $A \subset H$ , hence it acts on  $H'$ , and on  $S^k(H')$ .

**Lemma 1.2.** *The linear maps  $(\text{Tr}_k^A)_{k \geq 1}$  are  $\rho_0(\mathcal{A})$ -equivariant.*

*Proof.* It follows from the fact that an element of  $\rho_0(\mathcal{A})$  preserves the intersection form  $\omega'$ .  $\square$

Note also that the graded space associated to the  $\mathcal{A}J$ -filtration is stable by conjugation by elements of  $\mathcal{A}$ , and the imbedding  $\text{Gr}^{\mathcal{A}J} \rightarrow \bigoplus_{k \geq 1} \mathcal{G}_k$  induced by  $\tau$  is  $\mathcal{A}$ -equivariant with respect to the actions we described.

For the sake of completeness, we describe how the Lagrangian traces behave with respect to the brackets in  $\mathcal{G}$ . As a consequence of [7, Lemma 3.2], we have the following result:

**Proposition 1.3.** *The map  $\text{Tr}^A$  vanishes on brackets of derivations. For any  $\delta, \eta \in \mathcal{G}$ , we have*

$$\text{Tr}^A([\delta, \eta]) = 0.$$

We shall not use this proposition in the sequel.

**Remark 1.4.** Unlike the trace-like operators defined by Morita, the Lagrangian traces do not necessarily vanish in even degree.

## 2. MAGNUS REPRESENTATIONS FOR THE HANDLEBODY GROUP

**2.1. Morita's trace map and the Magnus representation.** We begin by reviewing the case of the mapping class group, following [9]. In this section we work with use the notation  $\gamma_i = \alpha_i$  and  $\gamma_{g+i} = \beta_i$  for  $1 \leq i \leq g$ . Recall that the projections  $(\beta'_i)$  of the  $\beta_i$ s in  $\pi'$  induce a basis of  $\pi'$  and that the elements  $\alpha_i$  normally generate the kernel of the projection  $\pi \rightarrow \pi'$ . We consider Fox derivatives  $(\frac{\partial}{\partial \gamma_i} : \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\pi])_{1 \leq i \leq 2g}$  with respect with the elements  $\gamma_i$  of the basis. By definition, these are  $\mathbb{Z}$ -linear maps which verify  $\frac{\partial \gamma_i}{\partial \gamma_j} = \delta_{i,j}$  for  $1 \leq i, j \leq 2g$  and  $\frac{\partial uv}{\partial \gamma_i} = \frac{\partial u}{\partial \gamma_i} + u \frac{\partial v}{\partial \gamma_i}$  for any  $u, v \in \pi$ . We also consider Fox derivatives for the basis chosen for  $\pi'$ .

To an element  $\varphi$  in  $\mathcal{M}$ , we associate the *Fox matrix*  $r(\varphi)$ :

$$\begin{aligned} r : \mathcal{M} &\longrightarrow \mathrm{GL}(2g, \mathbb{Z}[\pi]) \\ \varphi &\longmapsto \left( \frac{\partial \varphi(\gamma_j)}{\partial \gamma_i} \right)_{1 \leq i, j \leq 2g} \end{aligned}$$

where for any  $x \in \mathbb{Z}[\pi]$ ,  $\bar{x}$  is the element obtained by applying the anti-automorphism induced by the inversion in  $\pi$ . The abelianization map from  $\mathbb{Z}[\pi]$  to  $\mathbb{Z}[H]$  then gives a map

$$r^a : \mathcal{M} \longrightarrow \mathrm{GL}(2g, \mathbb{Z}[H])$$

usually called the *Magnus representation* of the mapping class group. Nevertheless, this is *not* a representation in the usual sense because it is a crossed homomorphism. When the map  $r^a$  is restricted to the Torelli group  $\mathcal{I}$ , it becomes a representation. Recall that  $D_1(H) \simeq \Lambda^3(H)$  and define  $C$  to be the contraction map sending  $a \wedge b \wedge c$  to  $\omega(a, b)c + \omega(b, c)a + \omega(c, a)b$ . Morita [9, Prop. 6.15] proved the following proposition:

**Proposition 2.1.** *Let  $g \geq 2$ , and  $\varphi \in \mathcal{I}$ , then  $\det(r^a(\varphi)) \in H \subset \mathbb{Z}[H]$  and*

$$\det(r^a(\varphi)) = 2C(\tau_1(\varphi)).$$

Consequently, the determinant of the Magnus representation is trivial on  $J_k$ , for  $k \geq 2$ .

Furthermore, Morita proved that the reduction  $r_k(\varphi)$  of the Fox matrix of  $\varphi$  in  $\mathrm{GL}(2g, \mathbb{Z}[\pi]/I^{k+1})$  can be recovered from the  $k$ -th Johnson homomorphism. Any element in  $D_k(H)$  sends an element of  $H$  to an element in  $\Gamma_{k+1}\pi/\Gamma_{k+2}\pi$ . The Fox derivatives of such an element relatively to the  $\gamma'_i$ s yield a well-defined vector in  $(I^k/I^{k+1})^{2g}$ . Indeed a Fox derivation sends an element of  $\Gamma_k\pi$  to an element of  $I^{k-1}$ . For  $d \in D_k(H)$ , we set

$$\|d\| := \left( \frac{\partial d(\{\gamma_j\})}{\partial \gamma_i} \right) \in \mathrm{GL}(2g, I^k/I^{k+1})$$

where  $\{\gamma_j\}$  is the class of  $\gamma_j$  in  $H$ . The following formula was proven by Morita:

$$(2.1) \quad \forall \varphi \in J_k, r_k(\varphi) = \mathrm{Id} + \overline{\|\tau_k(\varphi)\|}$$

For an element  $d \in D_k(H)$ , Morita's trace map was originally defined as the trace of  $\|d\|$  after reduction in  $I_H^k/I_H^{k+1}$  (the projection from  $\pi$  to  $H$  sends  $I$  to  $I_H$ ). From formula (2.1) it is clear that

$$\det(r^a(\varphi)) \equiv 1 + \mathrm{Tr}(\overline{\|\tau_k(\varphi)\|}) \pmod{I_H^{k+1}}.$$

Then it follows from Proposition 2.1 that Morita's trace map vanishes on the image of the Johnson homomorphisms. Note here that  $\mathrm{Tr}$  is valued in  $I_H^k/I_H^{k+1}$  which can be identified with  $S^k(H)$ .

**2.2. The handlebody case.** We will now adapt Morita's strategy of proof to the case of the handlebody group. We first give an equivalent definition for the A-trace  $\mathrm{Tr}_k^A$ . Let  $d$  be an element of  $\mathcal{G}_k := \mathrm{Ker}(D_k(H) \rightarrow D_k(H'))$ . As explained in Section 1, the class of  $d$  in  $H^* \otimes \mathcal{L}_{k+1}(H')$  sends  $A$  to 0. Equivalently, an element of  $\mathcal{G}_k$  induces an element of  $H'^* \otimes \mathcal{L}_{k+1}(H')$ . Similarly to the definition of  $\|d\|$ , we define

$$\|d\|^A := \left( \frac{\partial d(b'_j)}{\partial \beta'_i} \right) \in \mathrm{GL}(g, I_{\pi'}^k/I_{\pi'}^{k+1}).$$

It is clear that this is actually the reduction in  $\pi'$  of the  $g \times g$  right-down submatrix of  $\|d\|$ . We denote  $\|d\|^{A,a} \in \mathrm{GL}(g, I_{H'}^k/I_{H'}^{k+1})$ . Once again, note that  $I_{H'}^k/I_{H'}^{k+1}$  can be identified with  $S^k(H')$ .

As a consequence [9, Prop. 6.3], we have:

**Lemma 2.2.** *For any  $k \geq 1$ , for any  $d \in \mathcal{G}_k$ ,  $\mathrm{Tr}_k^A(d) = \mathrm{Tr}(\|d\|^{A,\mathfrak{a}}) \in S^k(H')$ .*

We now define the Magnus representation in the handlebody case. We are in fact interested in the right-down  $g \times g$  part of the usual Fox matrix, after reduction in  $\mathbb{Z}[\pi']$ . Set

$$\begin{aligned} r^A : \mathcal{A} &\longrightarrow \mathrm{GL}(g, \mathbb{Z}[\pi']) \\ \varphi &\longmapsto \left( \frac{\partial \varphi(b_j)}{\partial b_i'} \right)_{1 \leq i, j \leq 2g} \end{aligned}$$

and  $r^{A,\mathfrak{a}} : \mathcal{A} \rightarrow \mathrm{GL}(g, \mathbb{Z}[H'])$  its abelian version. It is actually the composition of  $\mathcal{A} \rightarrow \mathrm{Aut}(\pi')$  with the usual Magnus representation of the group  $\mathrm{Aut}(\pi')$ . The map  $r^A$  is a crossed homomorphism in the following sense:

$$(2.2) \quad \forall \varphi, \psi \in \mathcal{A}, r^A(\varphi\psi) = r^A(\varphi)(\varphi \cdot r^A(\psi))$$

where  $\varphi \cdot r^A(\psi)$  is the matrix obtained from  $r^A(\psi)$  after applying  $\varphi$  to all the coefficients.

**Remark 2.3.** The mapping  $\mathrm{GL}(2g, \mathbb{Z}[\pi]) \rightarrow \mathrm{GL}(g, \mathbb{Z}[\pi'])$  (resp. its abelian version), which consist in projecting in  $\pi'$  (resp.  $H'$ ) the coefficients of the right-down matrix is of course not a homomorphism. This explains why the Lagrangian traces are not reductions of the Morita trace maps (see [1, Remark 4.4]).

We are now ready to prove Theorem B.

*Proof of Theorem B.* First, the map  $\det \circ r^{A,\mathfrak{a}}$  is  $\mathcal{A}$ -equivariant. Indeed, if  $\psi$  is an element of  $\mathcal{A}$ , and  $\varphi$  is at least in  $\mathcal{A}_1$ , we deduce from (2.2) that

$$r^{A,\mathfrak{a}}(\psi\varphi\psi^{-1}) = r^{A,\mathfrak{a}}(\psi)(\psi \cdot r^{A,\mathfrak{a}}(\varphi))(\psi \cdot r^{A,\mathfrak{a}}(\psi^{-1})),$$

because  $\varphi$  acts trivially on  $H'$ . We conclude by observing that

$$\mathrm{Id} = r^{A,\mathfrak{a}}(\psi\psi^{-1}) = r^{A,\mathfrak{a}}(\psi)(\psi \cdot r^{A,\mathfrak{a}}(\psi^{-1})),$$

hence  $r^{A,\mathfrak{a}}(\psi\varphi\psi^{-1})$  is conjugated to  $\psi \cdot r^{A,\mathfrak{a}}(\varphi)$ . The map  $\mathrm{Tr}_1^A \circ \tau_1$  is also  $\mathcal{A}$ -equivariant, by Lemma 1.2. According to [10], the subgroup  $\mathcal{A}_1$  is normally generated by the map  $\varphi := T_c \circ T_d^{-1}$  described in Figure 2. Note that  $\varphi$  is an *annulus twist* in the sense of [3] and a BP map in the sense of [5]. Indeed  $c$  and  $d$  cobound an annulus in the handlebody, and a surface in  $\Sigma$ , which ensures that  $\varphi$  is both an element of  $\mathcal{A}$  and  $\mathcal{I}$ . We set  $\tilde{c} := \beta_2[\alpha_1, \beta_1]$  a lift of the curve  $c$  in  $\pi$ . The action of  $\varphi$  on  $\pi$  is

$$\begin{aligned} \alpha_1 &\mapsto \tilde{c}\alpha_1\tilde{c}^{-1} \\ \alpha_2 &\mapsto \alpha_2\beta_2\tilde{c}^{-1} \\ \beta_1 &\mapsto \tilde{c}\beta_1\tilde{c}^{-1} \\ \beta_2 &\mapsto \tilde{c}\beta_2\tilde{c}^{-1}, \end{aligned}$$

hence we compute directly that

$$r^{A,\mathfrak{a}}(\varphi) = \begin{pmatrix} b_2'^{-1} & 0 \\ 1 - b_1'^{-1} & 1 \end{pmatrix}.$$

We deduce that  $\det(r^{A,\mathfrak{a}}(\varphi)) = b_2'^{-1} \in \mathbb{Z}[H']$ , and, switching to the additive notation in  $H'$ , we obtain an element  $-b_2' \in H'$ . By equivariance, as the map  $\det \circ r^{A,\mathfrak{a}}$  is a multiplicative map, we deduce that it sends the subgroup  $\mathcal{A}_1$  in  $H'$ . Furthermore, by [4], we know that  $\tau_1(\varphi) = a_1 \wedge b_1 \wedge b_2$ . Hence,  $\mathrm{Tr}_1^A(\tau_1(\varphi)) = -b_2'$ , and we deduce that, on  $\mathcal{A}_1$ ,

$$\det \circ r^{A,\mathfrak{a}} = \mathrm{Tr}_1^A \circ \tau_1.$$

As the first Johnson homomorphism vanishes on  $\mathcal{A}_k$ , for  $k \geq 2$ , we deduce that the map  $\det \circ r^{A,\mathfrak{a}}$  is trivial on these subgroups.  $\square$

We can now prove Theorem A.

*Proof of Theorem A.* The argument is similar to [9]. Indeed, we set  $r_k^A$  the reduction of  $r^A$  modulo  $I_{\pi'}^{k+1}$ . Clearly, as in [9][Prop. 6.13], we have the following equality:

$$(2.3) \quad \forall \varphi \in \mathcal{A}_k, r_k^A(\varphi) = \mathrm{Id} + \overline{\|\tau_k(\varphi)\|^A}.$$




FIGURE 2. Two curves  $c$  and  $d$  such that  $T_c \circ T_d^{-1}$  normally generates  $\mathcal{A}_1$ .

Recall that  $\varphi \in \mathcal{A}_k$  implies that  $\tau_k(\varphi) \in \mathcal{G}_k$ , hence the right-hand side of the equation makes sense. We can deduce equation (2.3) from equation (2.1), by reducing the coefficients of the right-down  $g \times g$  matrix of  $r_k(\varphi)$ , or directly using the very definition of the Johnson homomorphisms. After projection of the coefficients in  $\mathbb{Z}[H']$ , we obtain

$$\det(r^{\mathfrak{a}, \mathcal{A}}(\varphi)) \equiv 1 + \text{Tr}(\overline{\|\tau_k(\varphi)\|^{\mathcal{A}}}) \text{ mod } I_{H'}^{k+1}.$$

Using Theorem B and Lemma 2.2, we infer that, for any  $\varphi \in \mathcal{A}_k$ ,  $\text{Tr}_k^{\mathcal{A}}(\tau_k(\varphi)) \equiv 0 \text{ mod } I_{H'}^{k+1}$ .  $\square$

#### REFERENCES

- [1] Quentin Faes. The handlebody group and the images of the second Johnson homomorphism. To appear in *Algebraic & Geometric Topology*, arXiv:2001.09825, preprint 2021.
- [2] Quentin Faes. Triviality of the  $J_4$ -equivalence among homology 3-spheres. To appear in *Trans. Amer. Math. Soc.*, arXiv:2105.14253, preprint 2021.
- [3] Sebastian Hensel. A primer on handlebody groups. In *Handbook of group actions. V*, volume 48 of *Adv. Lect. Math. (ALM)*, pages 143–177. Int. Press, Somerville, MA, 2020.
- [4] Dennis Johnson. An abelian quotient of the mapping class group I. *Mathematische annalen*, 249(3):225–242, 1980.
- [5] Dennis L. Johnson. Homeomorphisms of a surface which act trivially on homology. *Proc. Amer. Math. Soc.*, 75(1):119–125, 1979.
- [6] Jerome Levine. The Lagrangian filtration of the mapping class group and finite-type invariants of homology spheres. *Mathematical Proceedings of the Cambridge Philosophical Society*, 141(2):303–315, 2006.
- [7] Gwénaél Massuyeau and Takuya Sakasai. Morita’s trace maps on the group of homology cobordisms. *J. Topol. Anal.*, 12(3):775–818, 2020.
- [8] Shigeyuki Morita. Casson’s invariant for homology 3-spheres and characteristic classes of surface bundles I. *Topology*, 28(3):305–323, 1989.
- [9] Shigeyuki Morita. Abelian quotients of subgroups of the mapping class group of surfaces. *Duke Mathematical Journal*, 70(3):699–726, 1993.
- [10] Genki Omori. A small normal generating set for the handlebody subgroup of the Torelli group. *Geometriae Dedicata*, 201(1):353–367, 2019.
- [11] Wolfgang Pitsch. Integral homology 3-spheres and the Johnson filtration. *Transactions of the American Mathematical Society*, 360(6):2825–2847, 2008.
- [12] Ricard Riba. The Rohlin invariant and  $\mathbb{Z}/2$ -valued invariants of homology spheres, arxiv:2010.11766, preprint 2020.
- [13] Takuya Sakasai. A survey of Magnus representations for mapping class groups and homology cobordisms of surfaces. In *Handbook of Teichmüller theory. Volume III*, volume 17 of *IRMA Lect. Math. Theor. Phys.*, pages 531–594. Eur. Math. Soc., Zürich, 2012.