

# MINIMAL GROUPS OF GIVEN REPRESENTATION DIMENSION

JONATHAN COHEN

ABSTRACT. For a finite group  $G$ , let  $\text{rdim}(G)$  denote the smallest dimension of a faithful, complex linear representation of  $G$ . It is clear that  $\text{rdim}(H) \leq \text{rdim}(G)$  for any subgroup  $H$  of  $G$ . We consider  $G$  with the property that  $\text{rdim}(H) < \text{rdim}(G)$  whenever  $H$  is a proper subgroup of  $G$ , in particular proving a classification of such groups when  $G$  is abelian or  $\text{rdim}(G) \leq 3$ .

## 1. INTRODUCTION

We say a finite group  $G$  has representation dimension  $n$ , written  $\text{rdim}(G)$ , if there is a faithful representation  $\rho : G \rightarrow \text{GL}_n(\mathbb{C})$  and no faithful representation of dimension  $n - 1$ . Clearly  $\text{rdim}(H) \leq \text{rdim}(G)$  for every subgroup  $H$  of  $G$ . In this article we consider the class of groups  $G$  such that  $\text{rdim}(H) < \text{rdim}(G)$  for all proper subgroups  $H$  of  $G$ ; we call these groups *minimally faithful of degree*  $\text{rdim}(G)$ .

If  $G$  is abelian or  $\text{rdim}(G) = 2$ , such groups are straightforward to classify; see Lemma 2.2 and Proposition 3.1. The main result of this paper is a classification of minimally faithful groups of degree 3. We summarize this result here; let  $C_n$  be a cyclic group of order  $n$  and  $Q_8$  the quaternion group of order 8.

**Theorem 1.1.** *Let  $G$  be a minimally faithful group of degree 3. Then  $G$  is one of the following:*

- a)  $C_p \times C_p \times C_p$  for a prime  $p$ ,
- b)  $(C_p \times C_p) \rtimes C_{3^k}$  for a prime  $p \equiv -1 \pmod{3}$ ,  $k \geq 1$ , with center  $C_{3^{k-1}}$ ,
- c)  $C_p \rtimes C_{3^k}$  for a prime  $p \equiv 1 \pmod{3}$ ,  $k \geq 1$ , with center  $C_{3^{k-1}}$ ,
- d)  $C_{3^k} \rtimes C_3$  for  $k \geq 2$ , with center  $C_{3^{k-1}}$ ,
- e) The Heisenberg group of order 27,
- f)  $Q_8 \times C_p \times C_p$  for an odd prime  $p$ ,
- g)  $(C_{2^k} \rtimes C_2) \times C_p \times C_p$  where  $k \geq 2$ ,  $p$  is an odd prime, and  $C_2$  acts on  $C_{2^k}$  by  $g \mapsto g^{1+2^{k-1}}$ ,

h)  $(C_q \rtimes C_{2^m}) \times C_p \times C_p$  where  $m \geq 1$ ,  $q, p$  are distinct odd primes, and a generator of  $C_{2^m}$  acts by inversion on  $C_q$ ,

i)  $(C_p \rtimes C_{2^m}) \times C_2$  where  $m > 1$ ,  $p$  is an odd prime, and a generator of  $C_{2^m}$  acts by inversion on  $C_p$ ,

j)  $Q_8 \times C_2$ ,

k) A nonabelian 2-group whose index-2 subgroups are all abelian with two invariant factors.

## 2. PRELIMINARIES

We write  $D_{2n}$  for the dihedral group of order  $2n$ ,  $A_n$  for the alternating group of degree  $n$ , and  $S_n$  for symmetric group of degree  $n$ . Let  $G^1 = [G, G]$  be the derived group of  $G$ ,  $G^2 = [G^1, G^1]$  for the derived group of  $G^1$ , and so on. The group  $\text{GL}_n(\mathbb{C})$  is the group of  $n \times n$  invertible matrices over the complex numbers, and its derived group  $\text{SL}_n(\mathbb{C})$  is the subgroup of matrices with determinant 1. The group  $\text{PGL}_n(\mathbb{C})$  is the quotient of  $\text{GL}_n(\mathbb{C})$  by its center. For a subset  $S$  of a group  $G$  we write  $C_G(S)$  for the centralizer of  $S$  in  $G$  and  $N_G(S)$  for its normalizer. We write  $\text{Aut}(G)$  for the automorphism group of a group  $G$  and  $\text{Inn}(G) \cong G/Z(G)$  for the group of inner automorphisms.

**Definition 2.1.** The group  $G$  is *minimally faithful of degree  $n$*  if  $\text{rdim}(G) = n$  and  $\text{rdim}(H) < n$  for all proper subgroups  $H$  of  $G$ .

Our main results classify minimally faithful groups of degree 2 and 3. First we show that the case of abelian minimally faithful groups is straightforward.

**Lemma 2.2.** *An abelian group  $G$  is minimally faithful of degree  $n$  if and only if it is an elementary abelian  $p$ -group of rank  $n$ , for some prime  $p$ .*

*Proof.* If  $G = C_{n_1} \times \cdots \times C_{n_k}$  then  $\text{rdim}(G) \leq k$ , with equality if and only if we have written  $G$  with a minimal number of factors. Let the  $p$ -syllow subgroup of  $G$  be written as  $\prod_{i=1}^{k_p} C_{p^{a_i}}$ . The smallest number of factors in an expression for  $G$  is  $\max_p(k_p)$ . If  $p$  is such that  $k_p = n$ , then since  $C_p^n \subset G$  and  $\text{rdim}(C_p^n) = n = \text{rdim}(G)$ , the conclusion follows from minimal faithfulness. The converse is clear.  $\square$

We mention an easy and useful fact: if  $G$  is a finite group and  $n$  is coprime to  $|Z(G)|$  then  $\text{rdim}(G \times C_n) = \text{rdim}(G)$ . This follows from the fact that any representation of  $G$  can be extended to  $G \times C_n$  by having  $C_n$  act by scalars.

Next, we recall a particular component of the classification of finite nonabelian groups containing only abelian proper subgroups.

**Proposition 2.3** ([MM03]). If  $G$  is a finite nonabelian group with only abelian proper subgroups then either  $G$  is a  $p$ -group or else  $G = Q \rtimes C_{p^a}$  where  $Q$  is an elementary abelian  $q$ -group for a prime  $q \neq p$ , in which case we have  $|Q| \equiv 1 \pmod{p}$ , and  $Z(G) = C_{p^{a-1}}$ .

### 3. MINIMALLY FAITHFUL OF DEGREE 2

**Proposition 3.1.** Suppose  $G$  is a nonabelian group and is minimally faithful of degree 2. Then either  $G = Q_8$  or  $G = \langle a, b \mid a^p = b^{2^m} = bab^{-1}a = 1 \rangle = C_p \rtimes C_{2^m}$  for some odd prime  $p$  and some  $m \geq 1$ .

*Proof.* A faithful representation  $G \rightarrow \text{GL}_2(\mathbb{C})$  must be irreducible, since otherwise it is a sum of two 1-dimensional representations, which would imply that  $G$  is abelian. Thus  $G$  has even order. Let  $P$  be a 2-Sylow subgroup of  $G$ .

Suppose  $G$  is not a 2-group. Then  $\text{rdim}(P) = 1$  so  $P$  is cyclic. By Cayley's normal 2-complement theorem,  $G = K \rtimes P$  for some subgroup  $K$  of odd order. Since  $\text{rdim}(K) = 1$ ,  $K$  is cyclic. Let  $x \in P$  be a generator. If  $x$  centralized  $K$  then  $G$  would be abelian, and if  $x^2$  did not centralize  $K$  then the proper subgroup generated by  $K$  and  $x^2$  would not be cyclic. Therefore  $x$  acts on  $K$  by an automorphism of order 2. We may decompose  $K = K^+ \times K^-$  where  $xkx^{-1} = k^{\pm 1}$  for all  $k \in K^{\pm}$ . Since  $G$  is nonabelian,  $K^-$  is nontrivial. If  $h \in K^-$  has prime order  $p$  then  $x$  and  $h$  generate a nonabelian group, which must equal  $G$ . Therefore  $G = C_p \rtimes C_{2^m}$  as claimed. These groups can be realized inside  $\text{GL}_2(\mathbb{C})$ : let  $H$  be the cyclic group of order  $2^{m-1}p$  generated by  $x^2$  and  $h$ , take any faithful character  $\chi : H \rightarrow \mathbb{C}^\times$ , and then  $\text{Ind}_H^G(\chi)$  is a faithful 2-dimensional representation.

Finally, suppose  $G$  is a 2-group. Then  $G$  has a cyclic maximal (index-2) subgroup. The  $p$ -groups with cyclic maximal subgroups have been classified; if  $G \neq Q_8$  then one readily sees that  $G$  contains  $C_2 \times C_2$ , and if  $G = Q_8$  then it is easy to check that  $G$  is minimally faithful of degree 2.  $\square$

#### 4. DEGREE 3

We now begin the classification of minimally faithful groups of degree 3, first proving some easy elementary results.

**Lemma 4.1.** *Suppose  $G$  is a nonabelian group.*

- a) *If  $G$  has odd order or  $Z(G)$  is not cyclic then  $\text{rdim}(G) \geq 3$ .*
- b) *If  $\text{rdim}(G) = 3$  and  $G$  has odd order then 3 divides  $|G|$  and  $Z(G)$  is cyclic.*

*Proof.* a) Since  $G$  is nonabelian,  $\text{rdim}(G) \geq 2$ . If  $G$  has odd order or  $Z(G)$  is not cyclic then a faithful representation  $\rho : G \rightarrow \text{GL}_2(\mathbb{C})$  would be reducible (the latter case because of Schur's lemma), hence would factor as a direct sum of two 1-dimensional representations, so  $1 \neq [G, G] \subset \ker(\rho)$ , a contradiction.

b) If  $\text{rdim}(G) = 3$  then a faithful representation  $G \rightarrow \text{GL}_3(\mathbb{C})$  must be irreducible: it cannot have a 2-dimensional irreducible summand since  $|G|$  is odd and cannot be a sum of 1-dimensional representations since  $G$  is nonabelian. Thus 3 divides  $|G|$ , and by Schur's lemma  $Z(G)$  is cyclic. □

**Corollary 4.2.** *Suppose  $G$  is a nonabelian group and is minimally faithful of degree 3. Let  $H$  be a proper subgroup of  $G$ .*

- a) *If  $H$  is nonabelian then  $Z(H)$  is cyclic.*
- b) *If  $H$  has odd order then  $H$  is abelian. In particular, if  $G$  has odd order then  $H$  is abelian.*
- c) *If  $H$  is abelian then  $H$  has at most 2 invariant factors.*
- d) *The order of  $G$  is divisible by 2 or 3 (or both).*

*Proof.* The first three parts are immediate since  $\text{rdim}(H) \leq 2$ . For the last, observe that if  $G$  has order coprime to 6 then any representation  $G \rightarrow \text{GL}_3(\mathbb{C})$  must be a direct sum of three 1-dimensional representations. □

**Lemma 4.3.** *Suppose that  $G$  is minimally faithful of degree 3. If  $Z(G) \cap [G, G]$  has even order then any faithful representation  $\rho : G \rightarrow \text{GL}_3(\mathbb{C})$  is reducible and  $[G, G]$  is isomorphic to a subgroup of  $\text{SL}_2(\mathbb{C})$ .*

*Proof.* Suppose  $z \in Z(G) \cap [G, G]$  has order 2. If  $\rho$  is irreducible then  $\rho(z) = -I_3$  by Schur's lemma. On the other hand,  $\rho([G, G]) \subset \text{SL}_3(\mathbb{C})$  and  $-I_3 \notin \text{SL}_3(\mathbb{C})$ . So  $\rho$  is

reducible,  $G \cong \rho(G) \subset \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$ , and  $\rho([G, G])$  is contained in the derived group of  $\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$ , which is  $\mathrm{SL}_2(\mathbb{C}) \times \{1\}$ .  $\square$

**Theorem 4.4.** *Let  $G$  be minimally faithful of degree 3. Then  $[G, G]$  is abelian.*

*Proof.* We may assume  $G$  is nonabelian. Let  $G^1 := [G, G]$ .

**STEP 1:**  $G \neq G^1$ .

Suppose  $G = G^1$ . Then any faithful representation  $\rho : G \rightarrow \mathrm{GL}_3(\mathbb{C})$  has image in  $\mathrm{SL}_3(\mathbb{C})$ . We refer to the classification of finite subgroups of  $\mathrm{SL}_3(\mathbb{C})$ , up to  $\mathrm{GL}_3(\mathbb{C})$ -conjugacy, found in [Serr14]. If  $\rho$  is reducible then (up to conjugacy)  $\rho(G)$  is contained in the image of  $g \mapsto (g, \det(g)^{-1}) : \mathrm{GL}_2(\mathbb{C}) \rightarrow (\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})) \cap \mathrm{SL}_3(\mathbb{C})$ , which would imply  $\mathrm{rdim}(G) = 2$ . Therefore  $\rho$  is irreducible. Among the subgroups of  $\mathrm{SL}_3(\mathbb{C})$  that act irreducibly and primitively on  $\mathbb{C}^3$ , the groups  $A_5$ ,  $PSL_2(7)$ , and  $3A_6$  (the nontrivial triple cover of  $A_6$ ) all contain  $A_4$ , while the others all have a nonabelian 3-Sylow subgroup (and are not themselves 3-groups). Since  $\mathrm{rdim}(A_4) = 3$ , none of these groups are minimally faithful of degree 3. The same issue occurs if  $G = H \cdot Z(\mathrm{SL}_3(\mathbb{C}))$  where  $H = A_5$  or  $PSL_2(7)$ . Therefore  $G$  acts irreducibly and imprimitively on  $\mathbb{C}^3$ .

Thus  $\rho(G)$  is generated by a finite subgroup  $D$  of the diagonal matrices and a cyclic permutation  $p = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$ , or else is generated by such a group together with a matrix  $Q = \begin{bmatrix} 0 & a & 0 \\ b & 0 & 0 \\ 0 & 0 & c \end{bmatrix}$  with  $abc = -1$ . In the both cases  $D$  is a normal subgroup of  $\rho(G)$ . In the first case  $\rho(G)/D = C_3$ , so  $G \neq G^1$ . In the second case, since  $Q^2 \in D$  and  $QpQ^{-1} \in Dp^{-1}$ , we have  $\rho(G)/D = S_3$ , which admits a nontrivial character, so  $G \neq G^1$ .

**STEP 2:**  $G$  is solvable.

If  $G$  is not solvable then  $G^1$  is not solvable. Since  $G^1 \neq G$  we can regard  $G^1$  as a subgroup of  $\mathrm{GL}_2(\mathbb{C})$ . Then  $G^2 := [G^1, G^1] \subset \mathrm{SL}_2(\mathbb{C})$  is not solvable. The unique nonsolvable subgroup of  $\mathrm{SL}_2(\mathbb{C})$  is the binary icosahedral group, which we will denote by  $BI_{120}$ . Thus  $Z(G^2) \cong Z(BI_{120})$  has order 2 which implies  $Z(G^2) \subset Z(G) \cap [G, G]$ . By lemma 4.3,  $\rho(G)$  may be regarded as a subgroup of  $\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$  and  $\rho(G^1) \subset \mathrm{SL}_2(\mathbb{C}) \times \{1\}$  so  $G^1 \cong BI_{120}$ . If  $(M, t) \in \rho(G \setminus G^1)$  where  $M \in \mathrm{GL}_2(\mathbb{C})$ ,  $t \in \mathbb{C}^\times$ , and  $(g, 1) \in \rho(G^1)$ , then  $(MgM^{-1}, 1) \in \rho(G^1)$ . so  $(M, 1)$  and  $\rho(G^1)$  generate a finite

subgroup of  $\mathrm{GL}_2(\mathbb{C}) \times \{1\}$  which contains a copy of  $BI_{120}$ . The only such groups are generated by  $BI_{120}$  and a subgroup of scalar matrices, by [Ngu08].

Therefore, since  $(M, t) \notin \rho(G^1)$ ,  $(M, t)$  commutes with  $\rho(G^1)$ , which implies  $M$  is a scalar matrix by Schur's lemma. So  $G = Z(G)G^1$  with  $Z(G) \cap G^1$  of order 2. If  $Z(G)$  is cyclic then  $\mathrm{rdim}(G) = 2$  since we can let  $Z(G)$  act by scalars on  $\mathbb{C}^2$  in a realization of  $BI_{120} \subset \mathrm{SL}_2(\mathbb{C})$ . Suppose  $Z(G)$  is not cyclic. Since  $Q_8$  is the 2-Sylow subgroup of  $BI_{120}$ , we have the nonabelian subgroup  $Q_8Z(G)$  with noncyclic center. Thus  $\mathrm{rdim}(Q_8Z(G)) = 3$ . But  $Q_8Z(G) \neq G$  since  $Q_8$  is not centralized by  $BI_{120}$ , contradicting minimality of  $G$ .

**STEP 3:** If  $G^1$  is not abelian then  $G^1 = Q_8$  or  $G^1 = Q_8 \rtimes C_3$ .

Suppose  $G^1$  is nonabelian. Note  $G^2 = [G^1, G^1] \subset \mathrm{SL}_2(\mathbb{C})$ . The finite subgroups of  $\mathrm{SL}_2(\mathbb{C})$  are  $C_n$ ,  $BD_{4n} := \langle A, B : A^n = B^2, B^4 = 1, BAB^{-1} = A^{-1} \rangle$  (dicyclic or binary dihedral groups of order  $4n$ ,  $n \geq 2$ ),  $Q_8 \rtimes C_3$  (the binary tetrahedral group),  $BO_{48}$  (the binary octahedral group), and  $BI_{120}$  (the binary icosahedral group). Their derived groups are, respectively,  $\{1\}$ ,  $C_{2n}$  (respectively,  $C_n$ ) if  $n$  is odd (respectively, even),  $Q_8$ ,  $Q_8 \rtimes C_3$ , and  $BI_{120}$ . The finite subgroups of  $PGL_2(\mathbb{C})$  are  $C_n$ ,  $D_{2n}$ ,  $A_4$ ,  $S_4$ , and  $A_5$ . In particular,  $G^1/Z(G^1)$  and  $G^2/(Z(G^2) \cap \{\pm I_2\}) = [G^1/Z(G^1), G^1/Z(G^1)]$  are such subgroups of  $PGL_2(\mathbb{C})$ . If  $Z(G^2)$  has a unique element  $z$  of order 2 then  $z \in Z(G) \cap [G, G]$  and lemma 4.3 applies. This holds except when  $G^2$  is cyclic of odd order.

Case 1: Since  $G$  is solvable,  $G^2 \neq BI_{120}$ .

Case 2: If  $G^2 = BO_{48}$  then  $G^2/Z(G^2) \cong S_4$ , so  $G^1/Z(G^1) \subset PGL_2(\mathbb{C})$  must contain  $S_4$ , hence equal  $S_4$ . Thus  $G^2/Z(G^2) \subset [S_4, S_4] = A_4$ , a contradiction.

Case 3: If  $G^2 = Q_8 \rtimes C_3$  then  $G^1 \subset \mathrm{SL}_2(\mathbb{C})$  equals  $BO_{48}$ , the only subgroup of  $\mathrm{SL}_2(\mathbb{C})$  with the correct commutator subgroup. We claim that  $BO_{48}$  cannot be realized as a commutator subgroup of any group, hence cannot equal  $G^1$ . We have  $\mathrm{Inn}(BO_{48}) = BO_{48}/(\pm I_2) = S_4$  and  $\mathrm{Aut}(BO_{48}) = S_4 \times C_2$ . But then we cannot have  $\mathrm{Inn}(BO_{48}) \subset \mathrm{Aut}(BO_{48})^1 = A_4$ , which needs to occur since the image of the  $G^1$  under the natural map  $G \rightarrow \mathrm{Aut}(G^1)$  equals  $\mathrm{Inn}(G^1)$  and lies in  $\mathrm{Aut}(G^1)^1$ .

Case 4: If  $G^2 = BD_{4n}$  with  $n > 2$  then  $G^1 \subset \mathrm{SL}_2(\mathbb{C})$  cannot exist because  $BD_{4n}$  is not the commutator subgroup of any finite subgroup of  $\mathrm{SL}_2(\mathbb{C})$ . If  $n = 2$  then  $G^2 = Q_8$  and  $G^1 = Q_8 \rtimes C_3$ , since the binary tetrahedral group is the only subgroup of  $\mathrm{SL}_2(\mathbb{C})$  with the correct commutator subgroup.

Case 5i: If  $G^2 = C_n$  is cyclic of even order  $n > 1$  then  $G^1 \subset \mathrm{SL}_2(\mathbb{C})$  and has derived group  $C_n$ . Therefore  $G^1 = BD_{4n}$ . Realizing  $G$  as a subgroup of  $\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$  (using lemma 4.3), let  $H$  be the subgroup of  $\mathrm{GL}_2(\mathbb{C})$  generated by the first components of elements of  $G$ . Clearly  $H^1 = G^1$ . If  $n \geq 4$  then  $H/Z(H) \subset \mathrm{PGL}_2(\mathbb{C})$  cannot have the dihedral group  $D_{2n}$  as its derived group. Since  $G^1/Z(G^1) = D_{2n}$  this is a contradiction. If  $n = 2$ , then  $G^1 = Q_8$  since this is the unique subgroup of  $\mathrm{SL}_2(\mathbb{C})$  with commutator subgroup of order 2.

Case 5ii: Now assume  $G^2 = C_n$  is cyclic of odd order  $n > 1$ . Since  $G^1$  is a nonabelian subgroup of  $G$ ,  $Z(G^1)$  is cyclic. Suppose  $Z(G^1)$  has even order. Then its unique element of order 2 will lie in  $Z(G)$ , so lemma 4.3 applies and  $G^1 \subset \mathrm{SL}_2(\mathbb{C})$ . The only subgroups of  $\mathrm{SL}_2(\mathbb{C})$  with cyclic derived group are  $BD_{4k}$ , but  $BD_{4k}^1 = C_{2k}$  if  $k$  is odd and  $BD_{4k} = C_k$  if  $k$  is even, so the odd-order cyclic group  $G^2$  is not a derived group of  $G^1 \subset \mathrm{SL}_2(\mathbb{C})$ , which is absurd. Thus  $Z(G^1)$  has odd order. Now  $G^1 \subset \mathrm{GL}_2(\mathbb{C})$  is nonabelian so  $G^1/Z(G^1) \subset \mathrm{PGL}_2(\mathbb{C})$ , and  $(G^1/Z(G^1))^1 = G^2$  so  $G^1/Z(G^1) = D_{2n}$ , the dihedral group of order  $2n$ . From the classification of finite subgroups of  $\mathrm{GL}_2(\mathbb{C})$  in [Ngu08],  $G^1 = C_r \times D_{2n} = C_r \times (G^2 \rtimes C_2)$  where  $r$  is odd (all the other subgroups contain  $-I_2$ ). But then  $G^1/G^2 = C_{2r}$  and  $G^2 = C_n$  are both cyclic, which implies that  $G^1$  is cyclic and  $G^2$  is trivial [proof:  $G/C_G(G^2)$  is a subgroup of  $\mathrm{Aut}(G^2)$ , which is abelian, hence  $G^1 \subset C_G(G^2)$ , so  $G^2 \subset Z(G^1)$ . But then  $G^1/G^2$  being cyclic implies  $G^1/Z(G^1)$  is cyclic, so  $G^1$  is cyclic].

**STEP 4:**  $G^1 \neq Q_8$  and  $G^1 \neq Q_8 \times C_3$ .

Suppose  $G^1$  is either of these two groups. We have  $\mathrm{Aut}(Q_8) \cong S_4 \cong \mathrm{Aut}(Q_8 \times C_3)$ . Therefore if  $P$  is a  $p$ -Sylow subgroup of  $G$  with  $p \geq 5$  then  $P \subset C_G(G^1)$ . Since  $G^1$  is nonabelian, and  $P \subset Z(PG^1)$ , ( $P$  is abelian since it is a proper subgroup of odd order)  $P$  is cyclic. Since  $\rho(G^1) \subset \mathrm{GL}_2(\mathbb{C}) \times \{1\}$  is nonabelian and  $\rho(G) \subset \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$ , Schur's lemma implies that  $\rho(C_G(G^1))$  is a group of matrices of the form  $\mathrm{diag}(x, x, y)$ , so  $P \subset Z(G)$ . Because  $P \cap G^1 = 1$ , we have  $G/G^1 \cong F \times P$  for an abelian group  $F$ . If  $N$  is the preimage of  $F$  in  $G$ , then  $G = N \times P$ . Clearly  $|P|$  and  $|Z(N)|$  are coprime, so  $\mathrm{rdim}(N \times P) = \mathrm{rdim}(P)$ . Therefore  $P$  is trivial and the only primes that may divide  $|G|$  are 2 and 3.

Since  $G^1$  contains a unique element  $z$  of order 2, we have  $z \in Z(G)$  and Schur's lemma forces  $\rho(z) = \mathrm{diag}(-1, -1, 1)$ . Let  $K = \ker(\rho(G) \rightarrow \mathrm{GL}_2(\mathbb{C}))$ , where the map is the projection onto the first factor. The group  $K$  is nontrivial since  $\mathrm{rdim}(G) > 2$  and  $K \subset \rho(Z(G))$  since  $K$  consists of elements of the form  $\mathrm{diag}(1, 1, y)$ . If  $k \in K$  had order 2 then  $k$  and  $\rho(z)$  would generate a noncyclic subgroup of  $\rho(Z(G))$ . Thus

$G^1Z(G)$  would be nonabelian with noncyclic center, hence would equal  $G$ . But then  $G^1 = (G^1Z(G))^1 \subset G^2Z(G)$ , which is false for both  $G^1 = Q_8$  and  $G^1 = Q_8 \rtimes C_3$ . Therefore  $K$  has odd order, hence is a 3-group, and if  $k \in K$  has order 3 then  $k = \text{diag}(1, 1, \omega)$  for a primitive 3rd root of unity  $\omega$ .

Suppose there exists a scalar matrix  $xI_3 \in \rho(Z(G))$  for some  $1 \neq x \in \mathbb{C}^\times$ . Replacing  $x$  by a power if necessary, we can assume that  $x = -1$  or  $x = \omega$ . In any case,  $xI_3$ ,  $\rho(z)$ , and  $\text{diag}(1, 1, \omega)$  will generate a noncyclic subgroup of  $\rho(Z(G))$ , yielding the same contradiction as above. Therefore  $\rho(G)$  contains no nontrivial scalar matrices. If we write  $\rho = \sigma \oplus \chi : G \rightarrow \text{GL}_2(\mathbb{C}) \times \text{GL}_1(\mathbb{C})$ , then the representation  $\sigma \otimes \chi^{-1} : G \rightarrow \text{GL}_2(\mathbb{C})$  has trivial kernel, since  $\sigma(g)\chi^{-1}(g) = I_2$  if and only if  $\sigma(g) = \chi(g)I_2$  if and only if  $\rho(g) = \chi(g)I_3$  is a scalar matrix. Thus  $\text{rdim}(G) < 3$ , a contradiction.  $\square$

Now we will break into cases depending on the rank of the 2-Sylow subgroup of the abelian group  $G^1 = [G, G]$ . First we prove two auxiliary results.

**Lemma 4.5.** *Suppose  $G$  is minimally faithful of degree 3, with even order and cyclic 2-Sylow subgroups. Then  $G = (C_q \rtimes C_{2^m}) \times C_p \times C_p$  where  $q, p$  are distinct odd primes,  $m \geq 1$ , and a generator of  $C_{2^m}$  acts by inversion on  $C_q$ .*

*Proof.* Let  $P$  be a 2-Sylow subgroup of  $G$  and assume  $P$  is cyclic. By Cayley's normal 2-complement theorem,  $G \cong K \rtimes P$ , where  $K$  is a normal 2-complement. Since  $K \neq G$  and has odd order,  $K$  is abelian. Let  $\phi : P \rightarrow \text{Aut}(K)$  be the homomorphism defining  $G$ , let  $x$  be a generator of  $P$  and  $n$  be the order of  $\phi(x)$ . Finally, let  $H$  be the subgroup generated by  $x^2$  and  $K$ , which is the unique subgroup of index 2 in  $G$ .

If  $n = 2$  then we claim  $\text{rdim}(G) \neq 3$  or  $Z(G)$  is not cyclic. In this case  $H$  is abelian. If  $K$  is cyclic then so is  $H$ , and there is a faithful character  $\chi : H \rightarrow \mathbb{C}^\times$ . But then  $\text{Ind}_H^G(\chi)$  is a faithful 2-dimensional representation of  $G$ . Thus  $K$  is not cyclic. Since  $K$  has odd order and  $x$  acts on  $K$  by an automorphism of order 2, we can decompose  $K = K^+ \times K^-$  where  $K^\pm := \{k \in K : xkx^{-1} = k^{\pm 1}\}$ . Since  $K^+$  is centralized by  $H$  and  $x$ , it lies in  $Z(G)$ . Clearly  $K^-$  is normalized by  $P$ , so

$$G \cong K^+ \times (K^- \rtimes P).$$

Suppose  $K^-$  is not cyclic, so it has two invariant factors. For independent generators  $a$  and  $b$  of  $K^-$  we have  $xax^{-1} = a^{-1}$  and  $xbx^{-1} = b^{-1}$ . Suppose  $\rho : G \rightarrow \text{GL}_3(\mathbb{C})$  was faithful. Since  $a$  and  $a^{-1}$  are conjugate in  $G$ , the set of eigenvalues of  $\rho(a)$  is equal to the set of its reciprocals. Since  $a$  has odd order, a nontrivial eigenvalue of  $\rho(a)$  cannot



be equal to its inverse. Thus the eigenvalues of  $\rho(a)$  are  $\zeta_a^{\pm 1}, 1$ , for some root of unity  $\zeta_a$  of order equal to the order of  $a$  in  $G$ . Similar considerations hold for  $\rho(b)$ . Since  $K^-$  is not cyclic there is a common prime dividing the orders of  $a$  and of  $b$ . This, the commutativity of  $K$ , and the assumption that  $\rho$  is faithful means we can assume without loss of generality that  $\rho(a) = \text{diag}(\zeta_a, \zeta_a^{-1}, 1)$  and  $\rho(b) = \text{diag}(\zeta_b, 1, \zeta_b^{-1})$  (we cannot put the 1 in the same entry for both since then  $\rho$  would not be faithful). But there is no matrix in  $\text{GL}_3(\mathbb{C})$  that simultaneously conjugates both of the matrices  $\rho(a)$  and  $\rho(b)$  to their inverses: the centralizers of either matrix consist solely of diagonal matrices, so any matrix that conjugates  $\rho(b)$  to  $\rho(b)^{-1}$  is a diagonal matrix

times  $\begin{bmatrix} & & 1 \\ & 1 & \\ 1 & & \end{bmatrix}$ , but no such matrix can conjugate  $\rho(a)$  to its inverse. Therefore

$\rho(x)$  cannot be defined, so either  $K^-$  is not cyclic or  $\text{rdim}(G) > 3$ . So  $K^-$  is cyclic. Observe that  $\text{rdim}(K^- \rtimes P) = 2$ : let  $N$  be the cyclic group  $\langle K^-, x^2 \rangle$  and  $\chi : N \rightarrow \mathbb{C}^\times$  a faithful character. Then  $\text{Ind}_N^{K^- \rtimes P}(\chi)$  is a faithful 2-dimensional representation. Therefore  $K^+$  is nontrivial. If  $K^+$  is cyclic then  $\text{rdim}(G) = \text{rdim}(K^- \rtimes P) = 2$ . Thus  $K^+$ , and hence  $Z(G)$ , is not cyclic. There exists an odd prime  $p$  such that  $C_p \times C_p \subset K^+$ ; the group  $(C_p \times C_p) \times (K^- \rtimes P)$  is nonabelian with noncyclic center, so has representation dimension at least 3, so equals  $G$  by minimality. Noting that every subgroup of  $K^-$  is normalized by  $P$  by definition, if  $q$  is an odd prime such that  $C_q \subset K^-$ , then  $(C_p \times C_p) \times (C_q \rtimes P)$  is nonabelian with noncyclic center, hence equals  $G$ .

If  $n \geq 4$  then we claim  $\text{rdim}(G) > 3$ . First assume  $n = 4$ . Suppose there is a cyclic subgroup  $L \subset K$  which is normalized by  $x$  on which conjugation by  $x$  is an automorphism of order 4. We claim that  $\text{rdim}(PL) > 3$  (in fact it equals 4). If  $y$  generates the cyclic group  $L$  then  $xyx^{-1} = y^t$  where  $t^2 \equiv -1 \pmod{p}$  for some prime  $p$  dividing  $|L|$ . Thus  $y$  is conjugate its inverse by  $x^2$ . If  $\rho : PL \rightarrow \text{GL}_3(\mathbb{C})$  is a faithful representation then, as above, the eigenvalues of  $\rho(y)$  are  $\zeta^{\pm 1}, 1$  for some root of unity  $\zeta$  of order equal to  $|L|$ . But since  $y$  is conjugate to  $y^t$ ,  $\rho(y^t)$  has the same eigenvalues as  $\rho(y)$ , which forces  $\zeta^t = \zeta^{-1}$  or  $\zeta^t = \zeta$  or  $\zeta^t = 1$ . These are all absurd, so  $L$  does not exist. Since  $x^2$  acts on  $K$  by an automorphism of order 2, we decompose  $K = K^+ \times K^-$  where  $K^\pm := \{k \in K : x^2 k x^{-2} = k^{\pm 1}\}$ . If  $K^-$  is not cyclic then the proof of the  $n = 2$  case shows that the group generated by  $x^2$  and  $K^-$  has representation dimension greater than 3. If  $K^-$  is trivial then  $x$  acts on  $K$  by an element of order 2, contradicting the assumption that  $n = 4$ . So  $K^-$  is cyclic and nontrivial. Note that  $K^-$  is normalized by  $x$  since  $x^2(x k x^{-1})x^{-2} = x k^{-1} x^{-1} = (x k x^{-1})^{-1}$  for  $k \in K^-$ . But now  $K^-$  satisfies the conditions of the group  $L$  above, so cannot exist. Finally,

assume  $n \geq 8$ . Then  $x^2$  acts on  $K$  by an automorphism of order 4, so the previous case shows that  $\text{rdim}(H) > 3$ , hence  $\text{rdim}(G) > 3$  (note that the proof of the  $n = 4$  case made no use of the minimality of  $G$  to derive  $\text{rdim}(G) > 3$ ).  $\square$

**Lemma 4.6.** *Suppose  $G$  is a nonabelian 2-group containing only abelian proper subgroups and  $\text{rdim}(G) = 2$ . Then either  $G = Q_8$  or  $G = C_{2^k} \rtimes C_2$  where  $k \geq 2$  and  $C_2$  acts on  $C_{2^k}$  by  $g \mapsto g^{1+2^{k-1}}$ .*

*Proof.* If  $G$  only has cyclic proper subgroups then  $G$  is minimally faithful of degree 2, so by our earlier classification  $G = Q_8$ . If  $G \supset V = C_2 \times C_2$  then we can choose an index-2 subgroup  $H \subset G$  with  $H \supset V$ . The group  $H$  is abelian by assumption, and  $V$  is the characteristic subgroup of  $H$  generated by elements of order 2, so  $N_G(V) = G$ . The map  $G \rightarrow \text{Aut}(V) \cong S_3$  factors through  $G/H$  and is nontrivial because  $Z(G)$  is cyclic. Thus there exists a unique  $1 \neq z \in V \cap Z(G)$ . Let  $v$  and  $vz$  be the other nontrivial elements of  $V$ , and let  $g \in G \setminus H$ . We have  $vgvg^{-1} = z$ . If  $g^2 = 1$  then  $g$  and  $V$  generate the nonabelian group  $C_4 \rtimes C_2$ , which must therefore equal  $G$ . If  $g^2 \neq 1$ , then some power of  $g^{2^{k-1}}$  is nontrivial and lies in  $V$ , hence is equal to  $z$  since  $g$  does not commute with  $v$  or  $vz$ . This yields the groups  $\langle g, v : g^{2^k} = v^2 = 1, vgv = g^{1+2^{k-1}} \rangle$ , where  $k \geq 2$ . Notice these groups have representation dimension 2 since we can induce a faithful character of the index-2 cyclic subgroup generated by  $g$  up to  $G$ .  $\square$

**Theorem 4.7.** *Let  $G$  be nonabelian and minimally faithful of degree 3, and let  $0 \leq r \leq 2$  be the rank of the 2-Sylow subgroup of  $G^1$ .*

- A) *If  $r = 2$  then  $G = (C_2 \times C_2) \rtimes C_{3^k}$  for some  $k \geq 1$ .*
- B) *If  $r = 1$  then either*
  - i)  *$G$  is a 2-group,*
  - ii)  *$G = Q_8 \times C_p \times C_p$  for an odd prime  $p$ , or*
  - iii)  *$G = (C_{2^k} \rtimes C_2) \times C_p \times C_p$  for an odd prime  $p$ , where  $k \geq 2$  and  $C_2$  acts on  $C_{2^k}$  by  $g \mapsto g^{1+2^{k-1}}$ .*
- C) *If  $r = 0$  then either*
  - i)  *$G$  has odd order,*
  - ii)  *$G$  is the group appearing in lemma 4.5, or*
  - iii)  *$G = (C_p \rtimes C_{2^m}) \times C_2$  where  $m > 1$ ,  $p$  is an odd prime, and a generator of  $C_{2^m}$  acts by inversion on  $C_p$ .*

*Proof.* Let  $\rho : G \rightarrow \mathrm{GL}_3(\mathbb{C})$  be a faithful representation of  $G$ .

A) The elements of order exactly 2 inside  $G^1$  generate a subgroup  $V \cong C_2 \times C_2$ ; it is characteristic in  $G^1$  so  $N_G(V) = G$ . Since  $\mathrm{Aut}(V) = S_3$ , the index  $[G : C_G(V)]$  divides 6. The centralizer  $C_G(V)$  is abelian since  $V \subset Z(C_G(V))$  and  $V$  is not cyclic. If  $\rho$  was reducible then  $G^1$  would be an abelian subgroup of  $\mathrm{SL}_2(\mathbb{C})$ , hence cyclic. But  $G^1 \supset V$ , so  $\rho$  is irreducible. Thus 3 divides  $|G|$  and  $G$  contains a nontrivial 3-Sylow subgroup  $P$ . If  $P \subset C_G(V)$  then  $[G : C_G(V)] = 2$ , since the image of the map  $G \rightarrow \mathrm{Aut}(V)$  could contain no element of order 3. But if  $G$  has an index-2 abelian subgroup then every irreducible representation of  $G$  has degree dividing 2, contradicting the irreducibility of  $\rho$ . Thus  $P$  does not centralize  $V$ . Let  $x \in P \setminus C_G(V)$ . Then  $x$  and  $V$  generate a group isomorphic to  $R := (C_2 \times C_2) \rtimes C_{3^k}$ . Every proper subgroup of  $R$  is abelian of rank at most 2, so it suffices to show that  $\mathrm{rdim}(R) = 3$ .

The group  $R$  has a faithful 3-dimensional representation: let  $\chi$  be a character of the index-3 abelian subgroup  $H := C_2 \times C_2 \times C_{3^{k-1}}$  with  $\ker(\chi)$  of order 2, then  $\rho = \mathrm{Ind}_H^R(\chi)$  is readily checked to be faithful. We claim that the group  $R$  has no faithful 2-dimensional representation. The subgroup  $C_2 \times C_2$  is abelian so we can assume it acts diagonally on  $\mathbb{C}^2$ , but the only diagonal  $2 \times 2$  matrices of order 2 are  $\mathrm{diag}(1, -1)$ ,  $\mathrm{diag}(-1, 1)$ ,  $\mathrm{diag}(-1, -1)$ . The third matrix must lie in the center but no order-2 element of  $R$  is central. So  $\mathrm{rdim}(R) = 3$  and  $R = G$ .

B) We will first show that either  $G$  is a 2-group or  $G = P \times C_p \times C_p$  for a 2-Sylow group  $P$  and odd prime  $p$ . Since  $r = 1$ ,  $G^1$  has a nontrivial cyclic 2-Sylow subgroup. Hence  $G^1$  contains a unique element  $z$  of order 2, lemma 4.3 applies:  $\rho$  is reducible and  $\rho(G^1) \subset \mathrm{SL}_2(\mathbb{C}) \times \{1\}$ , so  $G^1$  is cyclic. Therefore  $\rho(z) = \mathrm{diag}(-1, -1, 1)$ , where without loss of generality we have assumed  $\rho(G) \subset \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$ .

Suppose that  $|G^1| = 2$ . Then  $G^1 \subset Z(G)$ , and  $G = P \times K$  where  $K$  is abelian of odd order [proof: let  $a \in G$  and let  $b \in G$  of odd order. If  $aba^{-1} = bz$ , then the left side has odd order and the right side even order. Thus  $aba^{-1}b^{-1} = 1$ ]. If  $K \supset C_p \times C_p$  for an odd prime  $p$  then  $P \times C_p \times C_p$  is nonabelian with noncyclic center, so has representation dimension 3 and must equal  $G$ . Otherwise  $K$  is cyclic, which implies  $\mathrm{rdim}(P \times K) = \mathrm{rdim}(P)$ , so by minimality  $G = P$ .

Now suppose  $g$  generates the cyclic group  $G^1$  and has (even) order greater than two. Since  $G^1$  is abelian we can assume  $\rho(G^1)$  consists of diagonal matrices. Thus  $\rho(g) = \mathrm{diag}(x, x^{-1}, 1)$  for some  $x \neq \pm 1$  and  $\rho(C_G(G^1))$  is contained in the abelian group of diagonal matrices. If  $n \in G \setminus C_G(G^1)$ , then  $\rho(ngn^{-1}) = \mathrm{diag}(y, y^{-1}, 1)$ , since it must lie in  $\rho(G^1)$ . The eigenvalues of  $\rho(g)$  and  $\rho(ngn^{-1})$  are the same but

$g \neq ngn^{-1}$  so  $y = x^{-1}$ . Thus  $n$  acts on  $G^1$  by inversion. Since  $n$  was an arbitrary element of  $G \setminus C_G(G^1)$ , we have  $[G : C_G(G^1)] = 2$ .

Write  $C_G(G^1) = Q \times K = Q \times K^+ \times K^-$  where  $Q$  is the 2-sylow subgroup of  $C_G(G^1)$ ,  $K$  is its subgroup of odd-order elements, and  $K^\pm := \{k \in K : nkn^{-1} = k^{\pm 1}\}$ , noting that the action of  $n$  on  $K$  is by an element of order 2 since  $[G : C_G(G^1)] = 2$ . Both  $Q$  and  $K^\pm$  are normal in  $G$ , with  $Q$  being characteristic in the abelian group  $C_G(G^1)$  and  $K^\pm$  being normalized by definition. Replacing  $n$  by an odd power if necessary, we can assume that the order of  $n$  is a power of 2. Let  $P = \langle Q, n \rangle$ , so  $P$  is a 2-Sylow subgroup of  $G$  and

$$G = K^+ \times (K^- \rtimes P).$$

If  $k \in K^-$  then  $nkn^{-1}k^{-1} = k^{-2} \in G^1$ , but  $k$  has odd order so  $K^- \subset G^1$ . Every nontrivial commutator has the form  $ana^{-1}n^{-1}$  for some  $a \in Q \times K^-$ . If  $Q \subset Z(P)$  then the nontrivial commutators would all have odd order, contradicting our initial assumption. Thus  $P$  is nonabelian.

If  $K^+$  is not cyclic then it contains  $C_p \times C_p$  for an odd prime  $p$ , so  $C_p \times C_p \times P$  is a nonabelian subgroup of  $G$  with noncyclic center, so has representation dimension 3 and must equal  $G$ . If  $K^+$  is cyclic and nontrivial then  $\text{rdim}(G) = \text{rdim}(K^- \rtimes P)$  since  $K^- \cap Z(G) = \{1\}$ , but  $\text{rdim}(K^- \rtimes P) < \text{rdim}(G)$  by minimality so this is absurd.

Now assume that  $K^+$  is trivial. We will show that  $Z(P)$  is not cyclic, which forces  $G = P$ . Let  $L$  be the kernel of the projection  $\rho(G) \rightarrow \text{GL}_2(\mathbb{C})$ . So  $L \subset \rho(Z(G))$  is those matrices of the form  $\text{diag}(1, 1, y)$ . If  $L$  was trivial then  $\text{rdim}(G) < 3$ . Since  $\rho(K^-) \subset \rho(G^1) \subset \text{SL}_2(\mathbb{C}) \times \{1\}$ , we have  $\rho(K^-) \cap \rho(L) = \{1\}$ . Hence  $|L|$  is a nontrivial power of 2, so  $L \ni \text{diag}(1, 1, -1)$ , which together with  $\rho(z) = \text{diag}(-1, -1, 1)$  creates a noncyclic subgroup of  $\rho(Z(G) \cap P) \subset \rho(Z(P))$ .

We have shown that if  $G$  is not a 2-group then  $G = Q \times C_p \times C_p$  for a 2-group  $Q$  and odd prime  $p$ . Suppose  $G$  is not a 2-group. Clearly the 2-Sylow subgroup  $Q$  of  $G$  must be nonabelian. If  $Q$  contained a nonabelian proper subgroup  $Q_1$ , then  $Q_1 \times C_p \times C_p$  would be nonabelian with noncyclic center, hence would have representation dimension 3, contradicting minimality of  $G$ . Thus every proper subgroup of  $Q$  is abelian. Since  $\text{rdim}(Q) = 2$ , lemma 4.6 gives the two possible structures of  $Q$ , hence of  $G$ . That these groups both have a faithful representation of dimension 3 is simple: let  $\rho = \sigma \oplus \chi : G \rightarrow \text{GL}_3(\mathbb{C})$  where  $\sigma : Q \times C_p \rightarrow \text{GL}_2(\mathbb{C})$  and  $\chi : C_p \rightarrow \text{GL}_1(\mathbb{C})$  are faithful.

C) Let  $P$  be a 2-Sylow subgroup of  $G$ , so  $P$  is abelian since  $G^1$  has odd order. If  $P$  is trivial then  $|G|$  is odd. If  $P$  is cyclic then lemma 4.5 applies. So we may assume  $P$  has two invariant factors. If  $g \in G$  has odd order then  $g$  and  $G^1$  generate an odd-order subgroup of  $G$ . Since  $P$  is nontrivial, any odd-order subgroup of  $G$  is proper, hence abelian. Thus  $g \in C_G(G^1)$  and  $C_G(G^1)$  contains all elements of  $G$  of odd order. Therefore  $G = PC_G(G^1)$ . We have  $P \supset V \cong C_2 \times C_2$  since  $P$  is not cyclic.

Suppose that there exists  $v \in V \setminus Z(G)$ . From this we will derive a contradiction. Since  $G^1$  is abelian of odd order,  $G^1 = K^+ \times K^-$  where  $K^\pm := \{k \in G^1 : vkv = k^{\pm 1}\}$ . Note  $K^-$  is nontrivial since  $v \notin Z(G)$ . If  $1 \neq k \in K^-$  then  $\rho(k) \in \rho(G^1) \subset \mathrm{SL}_3(\mathbb{C})$  is conjugate to its inverse, so its eigenvalues are  $\zeta^{\pm 1}, 1$  for some odd root of unity  $\zeta \neq 1$ . Choosing a basis so that  $\rho(k) = \mathrm{diag}(\zeta, \zeta^{-1}, 1)$ , we see  $\rho(C_G(k))$  is contained in the abelian subgroup of diagonal matrices. Thus  $C_G(G^1) = C_G(k)$  (both consist under  $\rho$  of the diagonal matrices in  $\rho(G)$ ) is an abelian normal subgroup of  $G$ . A conjugate of  $\rho(k)$  under  $\rho(G)$  lies in  $\rho(G^1)$  so it is diagonal with the same eigenvalues. The permutations of the diagonal entries of  $\rho(k)$  induces a homomorphism  $G \rightarrow S_3$  with kernel  $C_G(G^1)$ . So  $[G : C_G(G^1)]$  divides 6 and is divisible by 2. Since  $[G : C_G(G^1)] = [P : P \cap C_G(G^1)]$  is a power of 2, we have  $[G : C_G(G^1)] = 2$ . Now  $v \notin C_G(G^1)$  and  $v$  has order 2 so  $G = C_G(G^1) \rtimes \langle v \rangle$ . Let  $x \in C_G(G^1) \cap P$  be the generator of the nontrivial and cyclic 2-Sylow subgroup of  $C_G(G^1)$ . If  $T$  is the subgroup of  $C_G(G^1)$  consisting of odd-order elements then  $G = \langle x \rangle \times (T \rtimes \langle v \rangle)$ . Since  $Z(T \rtimes \langle v \rangle)$  has odd order,  $\mathrm{rdim}(G) = \mathrm{rdim}(T \rtimes \langle v \rangle)$ . But  $T \rtimes \langle v \rangle$  is a proper subgroup of  $G$ , contradicting minimality of  $G$ .

Thus  $V \subset Z(G)$ . In particular,  $Z(G)$  is not cyclic so by Schur's lemma  $\rho$  is reducible. We can assume without loss of generality that  $\rho(G) \subset \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$ , hence  $\rho(G^1) \subset \mathrm{SL}_2(\mathbb{C}) \times \{1\}$ . In particular,  $G^1$  is cyclic. Let  $g \in G^1$  be nontrivial, so that we may assume  $\rho(g) = \mathrm{diag}(\zeta, \zeta^{-1}, 1)$  for some odd root of unity  $\zeta \neq 1$ . It follows that  $\rho(C_G(G^1))$  consists of the diagonal matrices in  $\rho(G)$ , hence  $C_G(G^1)$  is abelian. Since  $G = PC_G(G^1)$  is nonabelian and  $P$  is abelian, there exists  $y \in P \setminus C_G(G^1)$ . Since  $\rho(ygy^{-1})$  has the same eigenvalues as  $\rho(g)$ , is not equal to  $\rho(g)$ , and lies in  $\rho(G^1)$ , we must have  $ygy^{-1} = g^{-1}$ . The subgroup of  $G$  generated by  $y, g$ , and  $V$  is nonabelian with noncyclic center, so must be equal to  $G$ . The same is true if we replace  $g$  by  $g^i \neq 1$ , so by minimality  $g$  has prime order  $p$ . The group  $H$  generated by  $y$  and  $g$  is isomorphic to  $C_p \rtimes C_{2^m}$ . Since some power of  $y$  lies in  $V$ , the group  $G$ , which is generated by  $H$  and  $V$ , must be isomorphic to  $(C_p \rtimes C_{2^m}) \times C_2$ . If  $m = 1$  this group has representation dimension 2, so  $m > 1$ . A faithful 3-dimensional representation can be constructed similarly to the previous case.  $\square$

Remark. The proof of part b) broke into the cases  $|G^1| = 2$  and  $|G^1| \geq 4$ , and showed that  $|G^1|$  is always a power of 2. The final classification reveals that in fact we must have  $|G^1| = 2$  in the case  $r = 1$ .

Next we shall show that if  $G$  is a minimally faithful 2-group of degree 3 then  $Z(G)$  is not cyclic. This will be useful in the subsequent result that classifies the minimally faithful 2-groups of degree 3.

**Lemma 4.8.** *Suppose  $G$  is a nonabelian 2-group and minimally faithful of degree 3. Then  $Z(G)$  is not cyclic.*

*Proof.* Suppose  $Z(G)$  is cyclic and  $\pi : G \rightarrow \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C})$  is a faithful 3-dimensional representation, noting that no 3-dimensional representation of  $G$  is irreducible since  $G$  is a 2-group. Define the two subgroups

$$N = \ker(G \xrightarrow{\pi} \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C}) \rightarrow \mathrm{GL}_1(\mathbb{C}))$$

$$K = \ker(G \xrightarrow{\pi} \mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C}) \rightarrow \mathrm{GL}_2(\mathbb{C})).$$

Both  $K$  and  $N$  are nontrivial since  $\mathrm{rdim}(G) = 3$ , while  $N \cap K = 1$ . Clearly  $K \subset Z(G)$  since  $\pi(K)$  consists of matrices of the form  $\mathrm{diag}(1, 1, c) \in Z(\mathrm{GL}_2(\mathbb{C}) \times \mathrm{GL}_1(\mathbb{C}))$ . Let  $z \in Z(G)$  be the unique element of order 2. For any subgroup  $M \subset G$  we have  $z \in M$  iff  $M \cap Z(G) \neq 1$ . Since  $K \subset Z(G)$ ,  $z \in K$ , so  $z \notin N$  and  $N \cap Z(G) = 1$ . Since  $N$  is a 2-group it has nontrivial center. The subgroup  $Z(G)N \cong Z(G) \times N$  therefore has noncyclic center, and must therefore be abelian (it cannot equal  $G$  since  $Z(G)$  is cyclic). So  $N$  is abelian, and in fact  $N$  must be cyclic: otherwise  $N \supset C_2^2$  and  $Z(G)N \supset C_2^3$ . Let  $n \in N$  be the unique element in  $N$  of order 2. Since  $N$  is a normal subgroup of  $G$ ,  $n \in Z(G)$ ,  $n \neq z$ , a contradiction.  $\square$

**Proposition 4.9.** Let  $G$  be a nonabelian 2-group.

a) If  $G$  is minimally faithful of degree 3 and contains a nonabelian proper subgroup then  $G = Q_8 \times C_2$ .

b) Suppose  $G$  contains only abelian proper subgroups. Then  $G$  is minimally faithful of degree 3 if and only if every index-2 subgroup of  $G$  has two invariant factors.

*Proof.* a) Let  $N$  be a proper nonabelian subgroup of  $G$ , and  $Z = Z(G)$ , which we have shown is not cyclic. Since  $NZ$  is a nonabelian group with noncyclic center,  $\mathrm{rdim}(NZ) \geq 3$  and  $G = NZ$ . This remains true if we replace  $Z$  by any noncyclic subgroup of  $Z$ . So there exists a subgroup  $V \cong C_2 \times C_2 \subset Z$  with  $G = NV$ . If

$N \cap V$  was trivial then any element in  $N$  of order 2 would, with  $V$ , generate  $C_2^3$ . Thus  $N \cap V$  has order 2, since  $V \not\subset N$ . This implies  $G \cong N \times C_2$ . If  $N$  contained a nonabelian proper subgroup  $N_1$ , then  $N_1 \times C_2$  would be nonabelian with noncyclic center, hence would have representation dimension 3 and contradicting minimality of  $G$ . Since  $\text{rdim}(N) = 2$  and  $N \not\cong C_2 \times C_2$ , lemma 4.6 forces  $N = Q_8$ .

b) Let  $H \subset G$  be a subgroup of index 2. If  $H$  is cyclic then there is a faithful character  $\chi : H \rightarrow \mathbb{C}^\times$ , and then  $\text{Ind}_H^G(\chi)$  is a faithful 2-dimensional representation of  $G$ , which is impossible. The subgroup  $H$  cannot have more than 2 invariant factors since  $G \not\cong C_2^3$ . The first direction is proved.

Now assume that  $G$  is a nonabelian 2-group whose index-2 subgroups are all abelian with two invariant factors. Let  $H \subset G$  of index 2, so  $H = C_{2^r} \times C_{2^s} = \langle x, y \rangle$  where  $r \geq s > 0$ . Let  $a \in G \setminus H$ . Since proper subgroups of  $G$  are abelian,  $a$  must centralize every proper subgroup of  $H$  that it normalizes. So every characteristic subgroup of  $H$  is contained in  $Z(G)$ , and thus  $C_{2^{r-1}} \times C_{2^s}$  is central if  $r > s$  and  $C_{2^{s-1}} \times C_{2^{s-1}}$  is central if  $r = s$ . So  $x^2, y^2 \in Z(G)$  if  $r = s$  and  $x^2, y \in Z(G)$  if  $r > s$ .

If  $a^2 = 1$  then  $\langle a, x^{2^{r-1}}, y^{2^{s-1}} \rangle = C_2^3$ . This is a proper subgroup of the nonabelian group  $G$ , contradicting our assumption that all proper subgroups have at most 2 invariant factors. So  $a^2 \neq 1$ . Now suppose  $r = s = 1$ . Since  $1 \neq a^2 \in H = C_2 \times C_2$ , we have  $a^4 = 1$  and  $|G| = [G : H]|H| = 8$ . The subgroup generated by  $a$  has index 2 in  $G$ , contradicting the assumption that all index-2 subgroups have two invariant factors. Thus either  $r \geq s > 1$  or  $r > s = 1$ . In either case  $Z(G) \cap H$  contains all elements of  $H$  order 2, hence  $Z(G)$  is not cyclic and  $\text{rdim}(G) \geq 3$ . All that remains is to construct a faithful 3-dimensional representation of  $G$ , since our assumption implies  $\text{rdim}(H) = 2$  for all maximal subgroups  $H$  of  $G$ .

We may assume without loss of generality that  $x \notin Z(G)$ ; this is automatic if  $r > s$ . We have  $axa^{-1} = x^i y^j$  for some  $i, j \geq 0$ . Since  $x^2 = ax^2 a^{-1} = (x^i y^j)^2 = x^{2i} y^{2j}$ , we have  $j \in \{0, 2^{s-1}\}$  and  $i \in \{1, 1 + 2^{r-1}\}$ . In any case,  $z := axa^{-1}x^{-1}$  is of order 2, so  $z \in Z(G)$ . Since  $x$  and  $a$  do not commute, they generate  $G$ . Therefore  $G/\langle z \rangle$  is abelian, and  $z$  generates  $[G, G]$ .

Let  $\chi : H \rightarrow \mathbb{C}^\times$  with  $\chi(z) = -1$ . Then  $\rho := \text{Ind}_H^G(\chi)$  is a 2-dimensional representation of  $G$ . It is clear that  $\ker(\rho) \cap aH$  is empty, so  $\ker(\rho) \subset H$ . If  $\ker(\rho)$  was not cyclic then it would contain the three elements in  $H$  of order 2, but  $z \notin \ker(\rho)$ . We have  $\ker(\rho) \neq 1$  since  $\text{rdim}(G) \geq 3$ . Let  $k$  be the unique element of order 2 in  $\ker(\rho)$  and  $\nu : G \rightarrow \mathbb{C}^\times$  with  $\nu(k) = -1$ ; such a  $\nu$  exists since the image of  $k$  in  $G/[G, G] = G/\langle z \rangle$  is nontrivial. The representation  $\rho \oplus \nu$  is faithful.  $\square$

The 2-groups of the form described here have been classified - see [MM03] and [M01]. For example, the only group of order at most 16 on this list is the nontrivial semidirect product  $C_4 \rtimes C_4$ . Since the actual construction of these groups would involve a large detour, we content ourselves to provide the relevant reference.

All that remains is to classify the odd-order groups that are minimally faithful of degree 3. This naturally breaks into the cases of 3-groups and not 3-groups.

**Proposition 4.10.** Suppose  $G$  is a nonabelian 3-group and minimally faithful of degree 3. Then either

- a)  $G$  is the Heisenberg group of order 27, or
- b)  $G = \langle a, b \mid a^{3^k} = b^3 = 1, b^{-1}ab = a^{1+3^{k-1}} \rangle = C_{3^k} \rtimes C_3$  for some  $k \geq 2$ .

*Proof.* Let  $H$  be an index-3 subgroup, so  $H$  is abelian and normal. If every index-3 subgroup of  $G$  is cyclic then every proper subgroup is cyclic, which would force  $G$  to be cyclic since it has odd order. So we may assume  $H$  is not cyclic. Let  $B \subset H$  be the subgroup generated by elements of order 3, so  $B \cong C_3 \times C_3$ . Since  $Z(G)$  is cyclic,  $B \not\subset Z(G)$ . Let  $a \in G \setminus C_G(B)$ . Since  $a$  and  $B$  generate a nonabelian group, they generate  $G$ . Since  $B$  is characteristic in  $H$ ,  $a$  normalizes  $B$ . We can choose generators  $z$  and  $b$  of  $B$  so that  $az = za$  and  $aba^{-1} = bz$ . In particular,  $G$  is generated by  $a$  and  $b$ . Suppose  $a$  has order  $3^k$ . Since  $a^3, z \in Z(G)$  and  $Z(G)$  is cyclic, either  $k = 1$  or else  $k \geq 2$  and  $z = a^{3^{k-1}}$ . The former case yields the Heisenberg group, which is easily shown to be minimally faithful of degree 3. The latter case yields the groups  $C_{3^k} \rtimes C_3$  with the claimed presentations. Every subgroup is clearly abelian of rank at most 2. Let  $\chi : \langle a \rangle \rightarrow \mathbb{C}^\times$  be a faithful character. Then  $\text{Ind}_{\langle a \rangle}^G(\chi)$  is a faithful 3-dimensional representation of  $G$ .  $\square$

**Proposition 4.11.** Suppose  $G$  is a nonabelian group of odd order, is not a 3-group, and is minimally faithful of degree 3. Then there exists a prime  $p \geq 5$  such that either

- a)  $G = C_p \rtimes C_{3^n}$  if  $p \equiv 1 \pmod{3}$ , where  $C_{3^{n-1}}$  is the center, or
- b)  $G = (C_p \times C_p) \rtimes C_{3^n}$  if  $p \equiv -1 \pmod{3}$ , where  $C_{3^{n-1}}$  is the center.

*Proof.* We know that 3 divides  $|G|$  and every proper subgroup of  $G$  is abelian. From proposition 2.3, we have  $|G| = 3^a p^b$  for some prime  $p > 3$ . Let  $P$  be a  $p$ -Sylow subgroup and  $Q$  a 3-Sylow subgroup. We know one of them is cyclic and the other normal and elementary abelian (of rank at most 2). Since  $|\text{Aut}(C_3^2)| = 2^4 \cdot 3$  and  $|\text{Aut}(C_3)| = 2$ , the subgroup  $Q$  cannot be normalized by  $P$  without being centralized



by  $P$ . Thus  $Q$  is cyclic and  $P$  is elementary abelian. A generator  $x$  of  $Q$  must act on  $P$  by an automorphism of order 3, in order that  $G$  contains no proper nonabelian subgroups.

Suppose  $P = C_p$ . Then 3 divides  $|\text{Aut}(P)| = p - 1$  and  $G = C_p \rtimes C_{3^n}$ . One obtains  $\text{rdim}(G) = 3$  by inducing a faithful character of the index-3 cyclic subgroup  $C_p \cdot Z(G) \cong C_{3^{n-1}p}$ .

Suppose  $P = C_p \times C_p$ . Then 3 divides  $|\text{Aut}(P)| = (p + 1)p(p - 1)^2$  and  $G = (C_p \times C_p) \rtimes C_{3^n}$ . If  $p \equiv 1 \pmod{3}$ , let  $\alpha, \alpha^2 \in \mathbb{Z}/p\mathbb{Z}$  be the two roots of  $t^2 + t + 1 = 0 \pmod{p}$ . Then  $P = P_0 \times P_1 \times P_2$  where  $P_i := \{y \in P : xyx^{-1} = y^{\alpha^i}\}$ . If there exists  $1 \neq y \in P_1$  or  $1 \neq y \in P_2$  then the subgroup generated by  $x$  and  $y$  will necessarily be a nonabelian proper subgroup of  $G$ . But if  $P = P_0$  then  $G$  is abelian. Thus  $p \equiv -1 \pmod{3}$ . We need only show that  $G$  has a faithful 3-dimensional representation. Let  $H = C_p \times C_p \times C_{3^{n-1}}$  and  $\chi : H \rightarrow \mathbb{C}^\times$  have kernel of order  $p$ . Let  $\rho := \text{Ind}_H^G(\chi)$ . Then  $\dim(\rho) = [G : H] = 3$ . It is easy to see that  $\ker(\rho) \subset H$ , and from there that  $\ker(\rho) \subset \ker(\chi)$ . If  $1 \neq y \in \ker(\rho)$  then  $xyx^{-1} \notin \ker(\chi)$  since  $xyx^{-1}$  does not lie in the subgroup generated by  $y$ . So  $xyx^{-1} \notin \ker(\rho)$ , which contradicts the fact that  $\ker(\rho)$  is a normal subgroup of  $G$ . Thus  $\rho$  is faithful even though  $\chi$  is not.  $\square$

## 5. FURTHER WORK

There are several natural questions that one can pose about the class of minimally faithful finite groups  $G$  of degree  $n$ , for  $n > 3$ . Are they always solvable? Which  $p$ -groups are minimally faithful? Which  $G$  can be realized inside  $\text{SL}_n(\mathbb{C})$ ? It is perhaps computationally intractable to obtain a full classification for substantially larger  $n$ .

In another direction, the notion of minimally faithful groups of degree  $n$  can be readily defined over an arbitrary field  $F$  instead of  $\mathbb{C}$ . There is still much that can be said for minimally faithful groups of degree 2; we state the results here without proof. Any finite subgroup of  $\text{GL}_1(F)$  is cyclic. It follows from this and simple calculations in  $\text{GL}_2(\overline{F})$  that the only candidates for nonabelian minimally faithful groups of degree 2 are a subset of those for  $\mathbb{C}$ , namely  $Q_8$  and  $C_p \rtimes C_{2^n}$ ; these work if and only if  $F$  contain a 4th root of unity and a  $2^n p$ th root of unity, respectively. In particular, if  $F$  has characteristic 2 then all minimally faithful groups of degree 2 over  $F$  are abelian. An abelian minimally faithful group over  $F$  need not be elementary abelian. For example, one can have  $G = C_{p^a}$  with  $F$  of characteristic  $p$  and containing a  $p^{a-1}$ -root of unity but not a  $p^a$ th root of unity, and such that the polynomial  $(t^{p^a} - 1)/(t^{p^{a-1}} - 1)$  has an irreducible quadratic factor. A root  $r$  of such a quadratic factor generates a quadratic extension  $F(r)/F$ , and multiplication by  $r$ , as an element of  $\text{Aut}_F(F(r)) \cong \text{GL}_2(F)$ ,

generates a faithful representation of  $C_{p^a}$ . This arises, for example, if  $F = \mathbb{Q}$  and  $p^a = 4$ . Similarly, if  $F$  has characteristic  $p$  then  $C_p$  is not a subgroup of  $\mathrm{GL}_1(F)$  but is a subgroup of  $\mathrm{GL}_2(F)$  via the matrices  $\begin{bmatrix} 1 & i \\ 0 & 1 \end{bmatrix}$ ,  $0 \leq i \leq p-1$ . Finally, one has the groups  $C_p \times C_p$ , which will be minimally faithful of degree 2 if and only if  $F$  contains a  $p$ th root of unity.

One could also broaden the class of groups and representations considered beyond the finite class. If  $G$  is an arbitrary group, one can define its representation dimension to be the minimum cardinality  $\kappa$  such that  $G$  admits a faithful representation on a vector space of dimension  $\kappa$ . Then one could define  $G$  to be minimally faithful of degree  $\kappa$  if, for all proper subgroups  $H$  of  $G$ , the representation dimension of  $H$  is a strictly smaller cardinal than  $\kappa$ . For example, if  $F$  is a finite field and  $G = \mathbb{Z}_{p^\infty}$  is the Prüfer  $p$ -group (the group of  $p$ -power torsion in  $\mathbb{C}^\times$ ) then  $G$  has infinite representation dimension over  $F$ , while every proper subgroup  $H$  of  $G$  is finite (and cyclic), so  $H$  has finite representation dimension over  $F$ . Thus  $G$  is minimally faithful over any finite field.

#### REFERENCES

- [M01] Miller, G. A. "Determination of all the groups of order  $p^m$  which contain the abelian group of type  $(m-2, 1)$ ,  $p$  being any prime." Transactions of the American Mathematical Society 2.3 (1901): 259-272.
- [MM03] Miller, George A., and Halcott C. Moreno. "Non-abelian groups in which every subgroup is abelian." Transactions of the American Mathematical Society 4.4 (1903): 398-404.
- [Ngu08] Nguyen, K. A., M. Van Der Put, and Jakob Top. "Algebraic subgroups of  $\mathrm{GL}_2(\mathbb{C})$ ." Indagationes Mathematicae 19.2 (2008): 287-297.
- [Serr14] Serrano, J. "Finite subgroups of  $SL(2, \mathbb{C})$  and  $SL(3, \mathbb{C})$ ." Notes, Warwick (2014).