

BLOW-UP OF THE CRITICAL NORM FOR A SUPERCRITICAL SEMILINEAR HEAT EQUATION

HIDEYUKI MIURA AND JIN TAKAHASHI

ABSTRACT. We consider the scaling critical Lebesgue norm of blow-up solutions to the semilinear heat equation $u_t = \Delta u + |u|^{p-1}u$ in an arbitrary smooth domain of \mathbf{R}^n . In the range $p > p_S := (n+2)/(n-2)$, we show that the critical norm must be unbounded near the blow-up time, where the type I blow-up condition is not imposed. The range $p > p_S$ is optimal in view of the existence of type II blow-up solutions with bounded critical norm for $p = p_S$.

CONTENTS

1. Introduction	2
1.1. Background	2
1.2. Main theorem	3
1.3. Organization of this paper	6
1.4. Notation	6
2. Gradient estimate	6
3. Localized weighted energy	10
3.1. Definition and change of variables	10
3.2. Quasi-monotonicity	13
4. ε -regularity	22
5. Proof of main theorem	33
5.1. Existence of blow-up limit	33
5.2. Blow-up analysis	36
5.3. Proof of localized statement	44
5.4. Completion of proof	46
Appendix A. Regularity estimates	46
Appendix B. Compactness results	53
Acknowledgments	53
References	54

1. INTRODUCTION

1.1. Background. We study blow-up solutions of the following semilinear heat equation:

$$(1.1) \quad \begin{cases} u_t = \Delta u + |u|^{p-1}u, & x \in \Omega, t > 0, \\ u(x, t) = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), & x \in \Omega. \end{cases}$$

Here $p > 1$, Ω is a smooth domain in \mathbf{R}^n with $n \geq 1$ and $u_0 \in L^q(\Omega)$ with $q \geq 1$. The boundary condition is not present if $\Omega = \mathbf{R}^n$.

The equation in (1.1) has attracted much attention as one of the simplest model for scaling invariant nonlinear parabolic equations. For each solution u , the rescaled function $u_\lambda(x, t) := \lambda^{2/(p-1)}u(\lambda x, \lambda^2 t)$ ($\lambda > 0$) also satisfies the equation. This implies that $L^{q_c}(\Omega)$ with

$$q_c := \frac{n(p-1)}{2}$$

is the scaling critical Lebesgue space for (1.1). The critical space plays a crucial role in well-posedness. If $u_0 \in L^{q_c}(\Omega)$ and $q_c > 1$, it is well-known [10, 93, 94] that there exists a unique classical L^{q_c} -solution u of (1.1) with the maximal existence time $T \in (0, \infty]$. For the definition of classical L^q -solution, see Remark 1.2. The solution u is smooth for $t \in (0, T)$, belongs to $C([0, T]; L^{q_c}(\Omega)) \cap C((0, T); L^\infty(\Omega))$ and admits the following blow-up criterion: If $T < \infty$, then $\lim_{t \rightarrow T} \|u(\cdot, t)\|_{L^\infty(\Omega)} = \infty$. This leads to the critical norm blow-up problem.

Problem. If $T < \infty$, does the following property hold?

$$(CNB) \quad \lim_{t \rightarrow T} \|u(\cdot, t)\|_{L^{q_c}(\Omega)} = \infty.$$

The problem is stated in Brezis and Cazenave [10, Open problem 7] and Quittner and Souplet [79, OP 2.1, Section 55], and many sufficient conditions of (CNB) are known, see [9, 32, 37, 55, 56, 58, 67, 96] and [79, Section 16]. Among others, Mizoguchi and Souplet [67] recently proved that (CNB) holds whenever the blow-up is type I. Here the blow-up of u is called type I if the blow-up rate is bounded by a spatially homogeneous solution up to the coefficient, that is, $\limsup_{t \rightarrow T} (T - t)^{1/(p-1)} \|u(\cdot, t)\|_{L^\infty(\Omega)} < \infty$. This rate is natural in view of the scaling. Note that the blow-up is called type II if it is not of type I. Up to now, it has been proved that the blow-up is always type I under the conditions that either $p < p_S$ and Ω is convex [38, 39], or $p < p_S$ and u is nonnegative [78]. Here p_S is the Sobolev critical exponent given by

$$p_S := \begin{cases} \frac{n+2}{n-2} & \text{for } n \geq 3, \\ \infty & \text{for } n = 1, 2. \end{cases}$$

Therefore, under such conditions, (CNB) holds for any blow-up solutions in the subcritical range $p < p_S$. For related results concerning sufficient conditions for type I blow-up, we refer [15, 16, 18, 33, 37, 38, 39, 45, 55, 57, 60, 66, 74, 75, 77, 78, 95] and [79, Section 23].

The situation is different in the case $p \geq p_S$. Type II blow-up solutions were constructed in a number of papers, see Remark 1.7 for a brief review. In particular,

the recent development [21, 23, 24, 40, 41, 52, 84] provides type II blow-up solutions satisfying

$$\sup_{0 < t < T} \|u(\cdot, t)\|_{L^{q_c}(\Omega)} < \infty$$

for $p = p_S$ and $3 \leq n \leq 6$, see [67, Section 4] for computations of the L^{q_c} norm of the solutions constructed in [21, 84]. This demonstrates that type I assumption in [67] is indeed necessary if $p = p_S$ and $3 \leq n \leq 6$. For $p = p_S$ and $n \geq 7$, no counter-examples are known. In contrast with the case $p = p_S$, all the known type II blow-up solutions do satisfy (CNB) in the range $p > p_S$. Taking the above results into account, we expect that (CNB) holds whenever $p > p_S$.

1.2. Main theorem. Our main result shows that, in the optimal range $p > p_S$, the critical norm of all finite time blow-up solutions must be infinite without assuming nonnegativity, monotonicity, symmetry, convexity or the type of blow-up.

Theorem 1.1. *Let $n \geq 3$, $p > p_S$, Ω be any smooth domain in \mathbf{R}^n and u be a classical L^{q_c} -solution of (1.1) with $u_0 \in L^{q_c}(\Omega)$. If the maximal existence time $T > 0$ is finite, then*

$$\limsup_{t \rightarrow T} \|u(\cdot, t)\|_{L^{q_c}(\Omega)} = \infty.$$

We give comments on the proof, and then we list remarks concerning the statement of this theorem and related results including other scaling invariant nonlinear evolution equations.

As in the earlier work [67] and also [31, 90] for related equations, the proof of Theorem 1.1 consists of two parts: (i) the blow-up (rescaling and compactness) procedure and (ii) the analysis of the blow-up limit. However, some new ingredients are necessary for the proof. In (i), a concentration theorem of the L^{q_c} norm near a blow-up point plays a crucial role for the nondegeneracy of the blow-up limit in [67]. Unfortunately, due to the absence of the type I assumption, we could not rely on the concentration theorem. In order to circumvent this difficulty, we will prove an ε -regularity theorem (Theorem 4.1), which guarantees the energy concentration near a blow-up point. Theorem 4.1 is motivated by similar ε -regularity theorems in [13]. In comparison, we do not need to assume that the solution is globally defined in time and has a certain lower bound of the energy. In (ii), in contrast to [67], the smoothness of our blow-up limit is unclear even before the final time. To overcome this issue, we invoke the monotonicity estimate similar to [37], which plays a key role to identify the blow-up limit. We note that the uses of the ε -regularity theorem and the monotonicity estimate are partially inspired by the related works [85, 90] for the harmonic map heat flow, but several modifications are needed for adapting to our problem, owing to the lack of the coercivity of the energy and the difference of the scaling critical spaces.

Remark 1.2 (Classical L^q -solution). For $1 \leq q \leq \infty$, the definition of classical L^q -solution is as follows (see [79, Definition 15.1]). Let $u_0 \in L^q(\Omega)$ and $T \in (0, \infty]$. We say that $u \in C([0, T]; L^q(\Omega))$ is a classical L^q -solution of (1.1) if $u \in C^{2,1}(\Omega \times (0, T)) \cap C(\bar{\Omega} \times (0, T))$, $u(\cdot, 0) = u_0$ and u is a classical solution of (1.1) for $t \in (0, T)$. If Ω is unbounded, we also impose $u \in L_{\text{loc}}^\infty((0, T); L^\infty(\Omega))$. If $q = \infty$, then $u \in C([0, T]; L^q(\Omega))$ is replaced with $u \in C((0, T); L^\infty(\Omega))$ and $\lim_{t \rightarrow 0} \|u(\cdot, t) - e^{t\Delta} u_0\|_{L^\infty(\Omega)} = 0$, where $e^{t\Delta}$ is the Dirichlet heat semigroup in Ω . We note that the maximal existence time $T_{\max} = T_{\max}(u_0) \in (0, \infty]$ can be defined for classical L^q -solutions, see [79, Proposition 16.1 (i), (ii)] for the definition.

Remark 1.3 (Uniqueness). By [93, 94] and [10, Theorem 1], a unique classical $L^{q_c}(\Omega)$ -solution of (1.1) exists for each $u_0 \in L^{q_c}(\Omega)$ with $q_c > 1$. Moreover, if we further assume $q_c > p$, the unconditional uniqueness [10, Theorem 4] holds for mild solutions, that is, solutions of the corresponding integral equation

$$u(\cdot, t) = e^{t\Delta}u_0 + \int_0^t e^{(t-s)\Delta}|u(\cdot, s)|^{p-1}u(\cdot, s)ds \quad \text{for } t \in (0, T)$$

in $C([0, T]; L^{q_c}(\Omega))$. Note that $q_c > p$ is equivalent to $p > n/(n-2)$ and is satisfied for $p > p_S$. Hence, under the assumption of Theorem 1.1, the condition $u_0 \in L^{q_c}(\Omega)$ implies the existence of a unique mild solution in $C([0, T]; L^{q_c}(\Omega))$. This solution is also a classical L^{q_c} -solution, since $q_c > 1$.

Remark 1.4 (Other Lebesgue spaces). We recall the case $u_0 \in L^q(\Omega)$ with $q \neq q_c$. For $1 \leq q < q_c$, there are results of the nonexistence and nonuniqueness of solutions, see [5, 11, 42, 94] and [74, Section 6]. By [33, Theorem 2.4], there exist solutions satisfying $T_{\max} < \infty$ and

$$\sup_{0 < t < T_{\max}} \|u(\cdot, t)\|_{L^q(\Omega)} < \infty.$$

For $q > q_c$, it is known [10, 93, 94] that (1.1) has a unique classical L^q -solution u . By [10, Corollary 13], it is also known that if $T_{\max} < \infty$,

$$\lim_{t \rightarrow T_{\max}} \|u(\cdot, t)\|_{L^q(\Omega)} = \infty.$$

Remark 1.5 (Application to backward self-similar solutions). Let $\Omega = \mathbf{R}^n$. We call the solution u of (1.1) is backward self-similar if it is of the form $u(x, t) = (T-t)^{-1/(p-1)}U(x/\sqrt{T-t})$ for some $T > 0$ and some profile function $U \in C^2(\mathbf{R}^n)$. Theorem 1.1 immediately shows the following corollary on the Liouville type theorem for backward self-similar solutions u of (1.1) with $p > p_S$. It slightly refines the result [67, Corollary 2] for $p > 1$ in the case $p > p_S$ by removing the assumption that U is bounded.

Corollary 1.6. *Let $n \geq 3$, $p > p_S$ and u be a backward self-similar solution of (1.1). If the profile function U of u belongs to $L^{q_c}(\mathbf{R}^n)$, then $u \equiv 0$.*

Remark 1.7 (Brief review of type II blow-up). Type II blow-up solutions were first found by Herrero and Velázquez [43, 44] in the Joseph–Lundgren supercritical range $p > p_{JL} := (n - 2\sqrt{n-1})/(n - 4 - 2\sqrt{n-1}) (> p_S)$ with $n \geq 11$. For a refined construction, see [63]. See also [57, 64, 65] for the Lepin supercritical range $p > p_L := (n - 4)/(n - 10) (> p_{JL})$. The critical cases $p = p_{JL}$ and $p = p_L$ were handled in [80, 81], see also [68]. In [69], the case where $p = p_S$, $n = 3$ and a suitably shrinking $\Omega = \Omega(t)$ was handled. Remark that the above type II blow-up solutions are radially symmetric. In the range $p_S < p < p_{JL}$, it was proved [55] that all radially symmetric blow-up solutions are of type I if either $\Omega = \mathbf{R}^n$ or Ω is a ball. For the existence of non-radial type II blow-up solutions, see [14, 17] for $p > p_{JL}$, [22] for $p = p_* := (n+1)/(n-3)$ and $n \geq 7$, and [19] for $p = 3$ and $5 \leq n \leq 7$. We note that p_* is the so-called second critical exponent (after [20]) and satisfies $p_S < p_* < p_{JL}$ for $n \geq 11$.

In the Sobolev critical case $p = p_S$, type II blow-up solutions were formally found by Filippas, Herrero and Velázquez [32] for $3 \leq n \leq 6$ and were rigorously constructed in [84] for $n = 4$. The recent development of the inner-outer gluing method refined the construction, see [23] for $n = 3$, [24, 52] for $n = 4$, [21, 40] for

$n = 5$ and [41] for $n = 6$. On the other hand, it was proved [15] for $n \geq 7$ that there is no type II blow-up solution if u_0 is close to the Aubin-Talenti function. Recently, it was also shown [92] that all interior blow-up solutions are of type I provided that $n \geq 7$, $u \geq 0$ and either $\Omega = \mathbf{R}^n$ or Ω is a convex bounded domain.

Remark 1.8 (Critical norm blow-up for the Navier-Stokes equations). Theorem 1.1 corresponds to the pioneering work of Escauriaza, Seregin and Sverak [31] for the three-dimensional Navier-Stokes equations. They showed the blow-up of the critical norm in the sense that

$$(1.2) \quad \text{if } T_{\max} < \infty, \text{ then } \limsup_{t \rightarrow T_{\max}} \|u(\cdot, t)\|_{L^3} = \infty.$$

The limit superior condition was later improved to the limit condition in [83]. In the case of the domains with boundary, the condition (1.2) was verified for the flat boundary [82] and for general cases [62]. These results were also refined to the limit condition for the flat case [7, 54] and for general cases [2]. On the other hand, the L^3 norm in (1.2) was further refined to the Lorentz norm [73] and the Besov norm [1, 35]. Actually, our norm in Theorem 1.1 can be replaced by the Lorentz norm $L^{q_c, r}$ with $r < \infty$, but we do not pursue this issue here. We also refer [25, 26] for the critical norm blow-up in higher dimensions $n \geq 4$.

Recently, Tao [86] proved that

$$\text{if } T_{\max} < \infty, \text{ then } \limsup_{t \rightarrow T_{\max}} \frac{\|u(\cdot, t)\|_{L^3}}{(\log \log \log(1/(T_{\max} - t)))^c} = \infty$$

for some constant $c > 0$. See [6, 72] for further developments in this subject. By analogy, it is expected that there is a general quantitative blow-up criterion for the semilinear heat equation (1.1) with $p > p_S$. This direction seems interesting and also challenging.

Remark 1.9 (Critical norm blow-up for other equations). Wang [90] studied the critical norm of the harmonic map heat flow between compact Riemannian manifolds without boundaries in the energy supercritical dimension $n \geq 3$. It was shown that if $T_{\max} < \infty$, then

$$\limsup_{t \rightarrow T_{\max}} \|\nabla u(\cdot, t)\|_{L^n} = \infty.$$

One of the key ideas in the proof is the monotonicity formula of Struwe [85].

For nonlinear dispersive equations with power nonlinearities, there are also many works on the blow-up of the critical norm of the form

$$\limsup_{t \rightarrow T_{\max}} \|u(\cdot, t)\|_{\dot{H}^{s_c}} = \infty.$$

Here s_c is the scaling critical exponent for each of the equations. Kenig and Merle [46] showed that blow-up solutions of the cubic defocusing Schrödinger equation in \mathbf{R}^3 must satisfy the above condition with $s_c = 1/2$. Their method is based on the concentration compactness procedure and the rigidity theorem. A similar method is applicable to the defocusing supercritical nonlinearity, see [48]. In the radial case, Merle and Raphaël [59] gave an explicit lower bound of the critical norm in some energy subcritical range. Similar results were also obtained for the nonlinear wave equation starting from [47]. For the focusing case, see [29, 30].

Remark 1.10 (Related results for supercritical elliptic equations). In the proof of Theorem 1.1, the obtained blow-up limit \bar{u} is a weak solution of the semilinear

heat equation and satisfies a monotonicity estimate. In addition, the singular set of $\bar{u}(\cdot, t)$ consists of finitely many points for each t . A similar situation can be found for the so-called stationary solutions of the semilinear elliptic equation $-\Delta u = |u|^{p-1}u$ for $p > p_S$, see [27, 70, 71, 91]. In this context, a weak solution u of the elliptic equation is called a stationary solution if u is a critical point of the corresponding energy functional with respect to domain variations.

1.3. Organization of this paper. This paper is organized as follows. In Section 2, we derive a key gradient estimate. In Section 3, we define a localized weighted energy and prove its quasi-monotonicity. In Section 4, we prove an ε -regularity theorem by analyzing the energy. In Section 5, we construct and examine a blow-up limit with the aid of the ε -regularity, and then we prove Theorem 1.1. In Appendix A, we give regularity estimates used in Section 5. In Appendix B, we recall an Aubin-Lions type compactness result also used in Section 5.

1.4. Notation. For $x \in \mathbf{R}^n$, we often write $x = (x', x_n)$ with $x' \in \mathbf{R}^{n-1}$ and $x_n \in \mathbf{R}$. Set $\mathbf{R}_+^n := \{x \in \mathbf{R}^n; x' \in \mathbf{R}^{n-1}, x_n > 0\}$. We denote by χ_A and $|A|$ the characteristic function and the Lebesgue measure of a measurable set A , respectively. For $r > 0$ and $(x, t) \in \mathbf{R}^{n+1}$, we write

$$\begin{aligned} B_r(x) &:= \{y \in \mathbf{R}^n; |x - y| < r\}, & B_r &:= B_r(0), \\ \Omega_r(x) &:= \Omega \cap B_r(x), & \Omega_r &:= \Omega_r(0), \\ P_r(x, t) &:= B_r(x) \times (t - r^2, t), & P_r &:= P_r(0, 0), \\ Q_r(x, t) &:= \Omega_r(x) \times (t - r^2, t), & Q_r &:= Q_r(0, 0). \end{aligned}$$

For $\rho > 0$ and $(x, t) \in \mathbf{R}^{n+1}$, we write

$$\begin{aligned} B_\rho^+(x) &:= \mathbf{R}_+^n \cap B_\rho(x), & B_\rho^+ &:= \mathbf{R}_+^n \cap B_\rho(0), \\ Q_\rho^+(x, t) &:= B_\rho^+(x) \times (t - \rho^2, t), & Q_\rho^+ &:= Q_\rho^+(0, 0). \end{aligned}$$

We denote by $G_\Omega = G_\Omega(x, y, t)$ the Dirichlet heat kernel in Ω . Set

$$K_{(\tilde{x}, \tilde{t})}(x, t) := (\tilde{t} - t)^{-\frac{n}{2}} e^{-\frac{|x - \tilde{x}|^2}{4(\tilde{t} - t)}}$$

for $x, \tilde{x} \in \mathbf{R}^n$ and $t < \tilde{t}$. The critical exponents are defined by

$$p_S := \frac{n+2}{n-2}, \quad q_c := \frac{n(p-1)}{2}, \quad q_* := \frac{n(p-1)}{p+1}.$$

Note that each conditions $q_c > p+1$ and $q_* > 2$ is equivalent to $p > p_S$. In what follows, we always assume $n \geq 3$ and $p > p_S$.

2. GRADIENT ESTIMATE

Let $R > 0$ and Ω be any smooth domain in \mathbf{R}^n with $0 \in \bar{\Omega}$. As will be seen in Section 5, the proof of Theorem 1.1 is based on the study of the localized problem

$$(2.1) \quad \begin{cases} u_t = \Delta u + |u|^{p-1}u & \text{in } \Omega_R \times (-1, 0), \\ u = 0 & \text{on } (\partial\Omega \cap B_R) \times (-1, 0), \\ u \text{ is } C^{2,1} & \text{on } (\bar{\Omega}_R \cap B_R) \times (-1, 0), \end{cases}$$

under the assumption that there exists $M > 0$ satisfying

$$(2.2) \quad \sup_{-1 < t < 0} \|u(\cdot, t)\|_{L^{q_c}(\Omega_R)} \leq M.$$

Here the boundary condition in (2.1) is ignored if $\partial\Omega \cap B_R = \emptyset$.

In this section, we show a key gradient estimate which is a basic tool to control an energy defined in Section 3. The method to estimate a term from $|u|^{p-1}u$ is based on Meyer [61, Theorem 18.1] (see also [87, Proposition 1.5]).

Proposition 2.1. *If u satisfies (2.1) and (2.2), then there exists a constant $C > 0$ depending on R such that*

$$\sup_{-3/4 < t < 0} \|\nabla u(\cdot, t)\|_{L^{q^*, \infty}(\Omega_{3R/4})} \leq C(M + M^p).$$

Remark 2.2. It seems to the authors that $L^{q^*, \infty}$ cannot be replaced with L^{q^*} in the left-hand side. On the other hand, we may improve L^{q_c} to $L^{q_c, \infty}$ in the right-hand side, but the above inequality is sufficient for later use.

Proof of Proposition 2.1. In the same spirit of [34, Proposition A.1], we derive a localized integral equation, and then we estimate each of the terms. Let $\phi \in C_0^\infty(\mathbf{R}^n)$ satisfy $0 \leq \phi \leq 1$ in \mathbf{R}^n , $\phi = 0$ in $\mathbf{R}^n \setminus B_{15R/16}$ and $\phi = 1$ in $B_{7R/8}$. Set $v(x, t) := u(x, t)\phi(x)$. Then v belongs to $C^{2,1}(\bar{\Omega} \times (-1, 0))$ and satisfies

$$\begin{cases} v_t - \Delta v = \phi|u|^{p-1}u - 2\nabla\phi \cdot \nabla u - u\Delta\phi & \text{in } \Omega \times (-1, 0), \\ v = 0 & \text{on } \partial\Omega \times (-1, 0). \end{cases}$$

Thus,

$$(2.3) \quad \begin{aligned} u(x, t) &= \int_{\Omega} G_{\Omega}(x, y, t + 7/8)\phi(y)u(y, -7/8)dy \\ &+ \int_{-7/8}^t \int_{\Omega} G_{\Omega}(x, y, t - s)\phi(y)|u(y, s)|^{p-1}u(y, s)dyds \\ &- \int_{-7/8}^t \int_{\Omega} G_{\Omega}(x, y, t - s)(2\nabla\phi \cdot \nabla u + u\Delta\phi)dyds \end{aligned}$$

for $x \in \Omega_{3R/4}$ and $-7/8 < t < 0$, where $G_{\Omega} = G_{\Omega}(x, y, t)$ is the Dirichlet heat kernel in Ω . Since $G_{\Omega}(x, y, t) = 0$ for $y \in \partial\Omega$ and $u\nabla\phi = 0$ on $\partial\Omega$, integrating by parts in the third term in the right-hand side yields

$$(2.4) \quad \begin{aligned} u(x, t) &= \int_{\Omega} G_{\Omega}(x, y, t + 7/8)\phi(y)u(y, -7/8)dy \\ &+ \int_{-7/8}^t \int_{\Omega} G_{\Omega}(x, y, t - s)(\phi|u|^{p-1}u + u\Delta\phi)dyds \\ &+ 2 \int_{-7/8}^t \int_{\Omega} \nabla_y G_{\Omega}(x, y, t - s) \cdot \nabla\phi(y)u(y, s)dyds \end{aligned}$$

for $x \in \Omega_{3R/4}$ and $-7/8 < t < 0$.

By the smoothness of Ω , the following estimate holds (see [50, Theorem IV.16.3] for instance): There exists a constant $C > 0$ such that

$$(2.5) \quad |\nabla_x^j G_{\Omega}(x, y, t)| \leq CK_j(x - y, t) \quad (j = 0, 1, 2)$$

for $x, y \in \Omega$ and $0 < t < 1$, where

$$(2.6) \quad K_j(x, t) := t^{-\frac{n}{2} - \frac{j}{2}} e^{-\frac{|x|^2}{4t}} \quad (j = 0, 1, 2).$$

Remark that the constant in (2.5) and (2.6) depends only on n , Ω and the length of the time interval $(0, 1)$. We prepare an estimate of $\partial_{x_i}\partial_{y_j}G_\Omega(x, y, t)$. By the semigroup property and (2.5), we have

$$\begin{aligned} |\partial_{x_i}\partial_{y_j}G_\Omega(x, y, t)| &= \left| \int_\Omega \partial_{x_i}G_\Omega(x, z, t/2)\partial_{y_j}G_\Omega(z, y, t/2)dz \right| \\ &\leq C \int_\Omega K_1(x-z, t/2)K_1(z-y, t/2)dz \\ &\leq Ct^{-1} \int_{\mathbf{R}^n} G(x-z, Ct/8)G(z-y, Ct/8)dz \\ &= Ct^{-1}G(x-y, Ct/4) \leq CK_2(x-y, t), \end{aligned}$$

where $G(x, t) := (4\pi t)^{-n/2}e^{-|x|^2/(4t)}$ and we changed the constant C in (2.6). Then by differentiating the integral equation (2.4) and using $K_1(x, t) \leq K_2(x, t)$ for $x \in \Omega$ and $0 < t < 1$, we see that

$$\begin{aligned} (2.7) \quad |\nabla u(x, t)| &\leq C \int_{\mathbf{R}^n} K_1(x-y, t+7/8)|u(y, -7/8)|\chi_{\Omega_R}(y)dy \\ &\quad + C \int_{-7/8}^t \int_{\Omega_R} K_1(x-y, t-s)|u(y, s)|^p dy ds \\ &\quad + C \int_{-7/8}^t \int_{\mathbf{R}^n} K_2(x-y, t-s)|u|\chi_{\overline{\Omega_{15R/16}} \setminus \Omega_{7R/8}} dy ds \\ &=: CU_1(x, t) + CU_2(x, t) + CU_3(x, t) \end{aligned}$$

for $x \in \Omega_{3R/4}$ and $-7/8 < t < 0$. Remark that each U_i is defined for all $x \in \mathbf{R}^n$ and $-7/8 < t < 0$.

For U_1 , from $q_* < q_c$ and the same argument to prove the L^{q_c} - L^{q_c} estimate for the heat semigroup (see [36, Section 1.1.3] for instance), it follows that

$$\begin{aligned} \|U_1(\cdot, t)\|_{L^{q_*\infty}(\Omega_{3R/4})} &\leq C\|U_1(\cdot, t)\|_{L^{q_c}(\mathbf{R}^n)} \\ &\leq C(t+7/8)^{-1/2}\|u(\cdot, -7/8)\chi_{\Omega_R}\|_{L^{q_c}(\mathbf{R}^n)} \\ &\leq C\|u(\cdot, -7/8)\|_{L^{q_c}(\Omega_R)} \leq CM \end{aligned}$$

for $-3/4 < t < 0$. Then,

$$(2.8) \quad \sup_{-3/4 < t < 0} \|U_1(\cdot, t)\|_{L^{q_*\infty}(\Omega_{3R/4})} \leq CM.$$

For U_3 , since $K_2(x-y, t-s)\chi_{\overline{\Omega_{15R/16}} \setminus \Omega_{7R/8}}(y) \leq C\chi_{\Omega_R}(y)$ for $x \in \Omega_{3R/4}$, $y \in \mathbf{R}^n$ and $-7/8 < s < t < 0$, we have

$$U_3(x, t) \leq C \int_{-7/8}^t \int_{\Omega_R} |u| dy ds \leq CM$$

for $x \in \Omega_{3R/4}$ and $-7/8 < t < 0$. Thus,

$$(2.9) \quad \sup_{-3/4 < t < 0} \|U_3(\cdot, t)\|_{L^{q_*\infty}(\Omega_{3R/4})} \leq CM.$$

We estimate U_2 . This part is a modification of [61, Theorem 18.1]. Let $-7/8 < t < 0$. By the change of variables, we have

$$U_2(x, t) \leq \tilde{U}(x; t), \quad \tilde{U} = \tilde{U}(x; t) := \int_0^\infty S(x, s; t) ds,$$

$$S = S(x, s; t) := \chi_{(0, t+7/8)}(s) \int_{\Omega_R} K_1(x-y, s) |u(y, t-s)|^p dy.$$

For $\lambda > 0$ and $\tau > 0$, define $E := \{x \in \Omega_R; \tilde{U}(x) > \lambda\}$ and

$$\tilde{U}(x) = \left(\int_0^\tau + \int_\tau^\infty \right) S(x, s) ds =: \tilde{U}_1(x) + \tilde{U}_2(x).$$

We estimate the Lebesgue measure $|E|$ of E . By the same argument to prove the $L^{q_c/p}$ - L^∞ estimate for the heat semigroup, we have

$$\begin{aligned} \|S(\cdot, s)\|_{L^\infty(\Omega_R)} &\leq C s^{-\frac{np}{2q_c} - \frac{1}{2}} \chi_{(0, t+7/8)}(s) \| |u(\cdot, t-s)|^p \|_{L^{q_c/p}(\Omega_R)} \\ &\leq C M^p s^{-\frac{np}{2q_c} - \frac{1}{2}} \end{aligned}$$

for any $s > 0$, where $C > 0$ is independent of t . Then,

$$\tilde{U}_2(x) \leq \int_\tau^\infty S(x, s) ds \leq C M^p \int_\tau^\infty s^{-\frac{np}{2q_c} - \frac{1}{2}} ds = C' M^p \tau^{-\frac{p+1}{2(p-1)}},$$

where $C' > 0$ is independent of t . For $\lambda > 0$, we choose τ so large that

$$(2.10) \quad C' M^p \tau^{-\frac{p+1}{2(p-1)}} = \frac{\lambda}{2}.$$

Then $\tilde{U}_2 \leq \lambda/2$. By setting $E_1 := \{x \in \Omega_R; \tilde{U}_1(x) > \lambda/2\}$ and using $\tilde{U}_2 \leq \lambda/2$ and $\tilde{U} = \tilde{U}_1 + \tilde{U}_2$, we see that $E \subset E_1$.

From the same argument to prove the $L^{q_c/p}$ - $L^{q_c/p}$ estimate, it follows that

$$\begin{aligned} \|S(\cdot, s)\|_{L^{\frac{q_c}{p}, \infty}(\Omega_R)} &\leq \|S(\cdot, s)\|_{L^{\frac{q_c}{p}}(\Omega_R)} \\ &\leq C s^{-\frac{1}{2}} \chi_{(0, t+7/8)}(s) \| |u(\cdot, t-s)|^p \|_{L^{\frac{q_c}{p}}(\Omega_R)} \leq C M^p s^{-\frac{1}{2}} \end{aligned}$$

for any $s > 0$, where $C > 0$ is independent of t . Thus,

$$\|\tilde{U}_1\|_{L^{\frac{q_c}{p}, \infty}(\Omega_R)} \leq \int_0^\tau \|S(\cdot, s)\|_{L^{\frac{q_c}{p}, \infty}(\Omega_R)} ds \leq C M^p \tau^{\frac{1}{2}}.$$

This together with the definition of $L^{q_c/p, \infty}$ shows that

$$\int_{E_1} \tilde{U}_1(x) dx \leq |E_1|^{1-\frac{p}{q_c}} \|\tilde{U}_1\|_{L^{\frac{q_c}{p}, \infty}(\Omega_R)} \leq C M^p |E_1|^{1-\frac{p}{q_c}} \tau^{\frac{1}{2}}.$$

On the other hand, the definition of E_1 gives $\int_{E_1} \tilde{U}_1(x) dx \geq (\lambda/2) |E_1|$. By $E \subset E_1$ and (2.10), we obtain

$$|E| \leq |E_1| \leq C \lambda^{-\frac{q_c}{p}} M^{q_c} \tau^{\frac{q_c}{2p}} = C M^{p q^*} \lambda^{-q^*},$$

and so $\lambda |\{x \in \Omega_R; \tilde{U}(x) > \lambda\}|^{1/q^*} \leq C M^p$ for any $\lambda > 0$. This implies that $\|\tilde{U}(\cdot; t)\|_{L^{q^*, \infty}(\Omega_R)} \leq C M^p$ for $-7/8 < t < 0$, where $C > 0$ is independent of t . Hence by the definition of \tilde{U} and M^p , we see that

$$\sup_{-3/4 < t < 0} \|U_2(\cdot, t)\|_{L^{q^*, \infty}(\Omega_{3R/4})} \leq \sup_{-3/4 < t < 0} \|\tilde{U}(\cdot; t)\|_{L^{q^*, \infty}(\Omega_R)} \leq C M^p.$$

Combining this inequality, (2.7), (2.8) and (2.9), we obtain the desired inequality. \square

3. LOCALIZED WEIGHTED ENERGY

Let u satisfy (2.1) and (2.2). In this section, we define a localized weighted energy of u analogous to Giga, Matsui and Sasayama [38, 39] and prove its quasi-monotonicity without assuming the convexity of Ω . Our computations to prove quasi-monotonicity is in the same spirit of Chou and Du [12], but the details are different. Note that, among the results in this section, we will refer only Lemma 3.1 and Proposition 3.2 in the subsequent sections.

Let $n \geq 3$, $p > p_S$ and Ω be any smooth domain in \mathbf{R}^n with $0 \in \overline{\Omega}$. We fix $R > 0$ as one of the following:

- (R1) In the case $0 \in \Omega$, we fix $0 < R < 1/2$ such that $\overline{B_R} \subset \Omega$.
- (R2) $\left\{ \begin{array}{l} \text{In the case } \partial\Omega \neq \emptyset \text{ and } 0 \in \partial\Omega, \text{ we fix } 0 < R < 1/2 \text{ such that} \\ \text{there exists } f \in C_0^\infty(\mathbf{R}^{n-1}) \text{ satisfying } f(0) = 0, \nabla' f(0) = 0, \\ \|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})} \leq 1/2 \text{ and } \Omega_R = \{x \in B_R; x_n > f(x')\} \\ \text{by relabeling and reorienting the coordinates axes if necessary.} \end{array} \right.$

Here $\nabla' f$ is the gradient on \mathbf{R}^{n-1} . Remark that the existence of f in (R2) is guaranteed by the smoothness of Ω .

3.1. Definition and change of variables. We define a localized weighted energy E and show its boundedness by using Proposition 2.1. To obtain quasi-monotonicity, we locally straighten the boundary. After that, we introduce backward similarity variables and derive the corresponding representation of E .

Let $\varphi \in C^\infty([0, \infty))$ satisfy $\varphi(z) = 1$ for $0 \leq z \leq 1/2$, $0 \leq \varphi(z) \leq 1$ for $1/2 < z < 1$, $\varphi(z) = 0$ for $z \geq 1$ and $|\varphi'(z)| \leq 4$ for $z \geq 0$. For $x, \tilde{x} \in \mathbf{R}^n$, $\tilde{t} > t$ and $r > 0$, we set $\phi_r = \phi_{\tilde{x}, r}(x) := \varphi(|x - \tilde{x}|/r)$ and

$$K = K_{(\tilde{x}, \tilde{t})}(x, t) := (\tilde{t} - t)^{-\frac{n}{2}} e^{-\frac{|x - \tilde{x}|^2}{4(\tilde{t} - t)}}.$$

For $\tilde{x} \in \overline{\Omega_{R/4}}$, $-1 < t < 0$ and $\tilde{t} > t$, define a localized weighted energy by

$$\begin{aligned} E(t) &= E_{(\tilde{x}, \tilde{t})}(t; \phi_{\tilde{x}, R/4}) \\ &:= (\tilde{t} - t)^{\frac{p+1}{p-1}} \int_{\Omega_R} \left(\frac{|\nabla u(x, t)|^2}{2} - \frac{|u(x, t)|^{p+1}}{p+1} + \frac{|u(x, t)|^2}{2(p-1)(\tilde{t} - t)} \right) \\ &\quad \times K_{(\tilde{x}, \tilde{t})}(x, t) \phi_{\tilde{x}, R/4}^2(x) dx. \end{aligned}$$

Remark that u is defined on $(\overline{\Omega_R} \cap B_R) \times (-1, 0)$, but we allow the choice of the reference time \tilde{t} as $\tilde{t} > 0$, that is, we can choose \tilde{t} after the existence time of u . Note that

$$(3.1) \quad \text{supp } \phi_{\tilde{x}, R/4} \subset B_{R/2} \quad \text{for } \tilde{x} \in \overline{\Omega_{R/4}}.$$

We check the boundedness of E .

Lemma 3.1. *There exists $C > 0$ such that*

$$|E_{(\tilde{x}, \tilde{t})}(t; \phi_{\tilde{x}, R/4})| \leq C(M + M^p)^2$$

for any $\tilde{x} \in \overline{\Omega_{R/4}}$, $-1 < t < 0$ and $\tilde{t} > t$.

Proof. From the Hölder inequality for the Lorentz spaces (see [49, Proposition 2.1]), (3.1), Proposition 2.1 and a direct computation, it follows that

$$(3.2) \quad \begin{aligned} & \int_{\Omega_R} |\nabla u|^2 K_{(\tilde{x}, \tilde{t})} \phi_{\tilde{x}, R/4}^2 dx \\ & \leq (\tilde{t} - t)^{-\frac{n}{2}} \|\nabla u(\cdot, t)\|_{L^{q_*}, \infty(\Omega_{R/2})}^2 \left\| e^{-\frac{|\cdot - \tilde{x}|^2}{8(\tilde{t} - t)}} \right\|_{L^{\frac{2q_*}{q_* - 2}, 2}(\mathbf{R}^n)}^2 \\ & \leq C(M + M^p)^2 (\tilde{t} - t)^{-\frac{p+1}{p-1}} \end{aligned}$$

for $-1 < t < 0$ and $\tilde{t} > t$. The Hölder inequality and (2.2) show that

$$\begin{aligned} \int_{\Omega_R} |u|^{p+1} K_{(\tilde{x}, \tilde{t})} \phi_{\tilde{x}, R/4}^2 dx & \leq (\tilde{t} - t)^{-\frac{n}{2}} \|u\|_{L^{q_c}(\Omega_{R/2})}^{p+1} \left(\int_{\mathbf{R}^n} e^{-\frac{|x - \tilde{x}|^2}{C(\tilde{t} - t)}} dx \right)^{1 - \frac{p+1}{q_c}} \\ & \leq CM^{p+1} (\tilde{t} - t)^{-\frac{p+1}{p-1}}, \\ \int_{\Omega_R} |u|^2 K_{(\tilde{x}, \tilde{t})} \phi_{\tilde{x}, R/4}^2 dx & \leq CM^2 (\tilde{t} - t)^{-\frac{2}{p-1}}, \end{aligned}$$

for $-1 < t < 0$ and $\tilde{t} > t$. The lemma follows from the above estimates. \square

We locally straighten the boundary. In the case (R2), we define smooth maps $\Phi = (\Phi_1, \dots, \Phi_n)$ and $\Psi = (\Psi_1, \dots, \Psi_n)$ by

$$\begin{cases} \xi_i = x_i =: \Phi_i(x), & x_i = \xi_i =: \Psi_i(\xi), \\ \xi_n = x_n - f(x') =: \Phi_n(x), & x_n = \xi_n + f(\xi') =: \Psi_n(\xi), \end{cases}$$

for $i = 1, \dots, n-1$. To handle the case (R1) in a unified way, we also set

$$\begin{cases} \xi_i = x_i =: \Phi_i(x), & x_i = \xi_i =: \Psi_i(\xi), \\ \xi_n = x_n =: \Phi_n(x), & x_n = \xi_n =: \Psi_n(\xi), \end{cases}$$

for (R1). We note that the maps Φ and Ψ for (R1) are identity maps and they are obtained by setting $f \equiv 0$ in the definitions of Φ and Ψ for (R2).

We write $\Phi(x) = \xi$ and $\Psi(\xi) = x$. Set

$$(3.3) \quad \hat{u}(\xi, t) := u(\Psi(\xi), t).$$

Then, direct computations show that

$$(3.4) \quad \begin{aligned} \nabla_x u(x, t) & = (\partial_{\xi_1} \hat{u} - (\partial_{\xi_n} \hat{u}) \partial_{\xi_1} f, \dots, \partial_{\xi_{n-1}} \hat{u} - (\partial_{\xi_n} \hat{u}) \partial_{\xi_{n-1}} f, \partial_{\xi_n} \hat{u}) \\ & = (\nabla' \hat{u} - (\partial_{\xi_n} \hat{u}) \nabla' f, \partial_{\xi_n} \hat{u}) =: \widehat{\nabla} \hat{u}(\xi, t) \end{aligned}$$

and that

$$\begin{aligned} \Delta_x u(x, t) & = \Delta_\xi \hat{u} - 2 \sum_{i=1}^{n-1} (\partial_{\xi_i} \partial_{\xi_n} \hat{u}) \partial_{\xi_i} f + \partial_{\xi_n}^2 \hat{u} \sum_{i=1}^{n-1} (\partial_{\xi_i} f)^2 - \partial_{\xi_n} \hat{u} \sum_{i=1}^{n-1} \partial_{\xi_i}^2 f \\ & = \Delta_\xi \hat{u} - 2 \nabla'_\xi (\partial_{\xi_n} \hat{u}) \cdot \nabla'_\xi f + (\partial_{\xi_n}^2 \hat{u}) |\nabla'_\xi f|^2 - (\partial_{\xi_n} \hat{u}) \Delta'_\xi f, \end{aligned}$$

where ∇' and Δ' are the gradient and Laplacian on \mathbf{R}^{n-1} with respect to the first $(n-1)$ components, respectively. Remark that the terms in the right-hand sides are evaluated at $(\xi, t) = (\Phi(x), t)$. Since u satisfies (2.1), we see that \hat{u} satisfies

$$(3.5) \quad \begin{cases} \hat{u}_t - \hat{A} \hat{u} = |\hat{u}|^{p-1} \hat{u} & \text{in } \Phi(\Omega_R) \times (-1, 0), \\ \hat{u} = 0 & \text{on } \Phi(\partial\Omega \cap B_R) \times (-1, 0). \end{cases}$$

Here, by abbreviations of the subscripts ξ and ξ_n , we set

$$\hat{A}\hat{u} := \Delta\hat{u} - 2\nabla'(\partial_n\hat{u}) \cdot \nabla'f + (\partial_n^2\hat{u})|\nabla'f|^2 - (\partial_n\hat{u})\Delta'f.$$

For $\tilde{x} \in \overline{\Omega_{R/4}}$, we write $\tilde{\xi} := \Phi(\tilde{x})$. Note that $\tilde{x} = \Psi(\tilde{\xi})$. By (3.4), the change of variables $x = \Psi(\xi)$ and the definition of K , we have

$$(3.6) \quad E(t) = (\tilde{t} - t)^{\frac{p+1}{p-1} - \frac{n}{2}} \int_{\Phi(\Omega_R)} \left(\frac{|\widehat{\nabla}\hat{u}|^2}{2} - \frac{|\hat{u}|^{p+1}}{p+1} + \frac{\hat{u}^2}{2(p-1)(\tilde{t} - t)} \right) \\ \times e^{-\frac{|\Psi(\xi) - \Psi(\tilde{\xi})|^2}{4(\tilde{t} - t)}} \phi_{\Psi(\tilde{\xi}), R/4}^2(\Psi(\xi)) d\xi.$$

We introduce the backward similarity variables

$$\eta := \frac{\xi - \tilde{\xi}}{(\tilde{t} - t)^{1/2}}, \quad \tau := -\log(\tilde{t} - t).$$

Then the rescaled functions are given by

$$(3.7) \quad w(\eta, \tau) := e^{-\frac{1}{p-1}\tau} \hat{u}(\tilde{\xi} + e^{-\frac{1}{2}\tau}\eta, \tilde{t} - e^{-\tau}),$$

$$(3.8) \quad g(\eta', \tau) := e^{\frac{1}{2}\tau} f(\tilde{\xi}' + e^{-\frac{1}{2}\tau}\eta').$$

Note that

$$\xi = \tilde{\xi} + e^{-\frac{1}{2}\tau}\eta, \quad \tilde{t} - t = e^{-\tau}, \\ \hat{u}(\xi, t) = (\tilde{t} - t)^{-\frac{1}{p-1}} w((\tilde{t} - t)^{-\frac{1}{2}}(\xi - \tilde{\xi}), -\log(\tilde{t} - t)).$$

Since \hat{u} satisfies (3.5), we see that w solves

$$(3.9) \quad \begin{cases} w_\tau + \frac{1}{2}\eta \cdot \nabla w + \frac{1}{p-1}w - Aw - |w|^{p-1}w = 0, \\ \eta \in \Omega(\tau), \tau \in (-\log(\tilde{t} + 1/2), -\log\tilde{t}), \\ w = 0, \quad \eta \in e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi}), \end{cases}$$

where the time interval $(-\log(\tilde{t} + 1), -\log\tilde{t})$ has been shortened to $(-\log(\tilde{t} + 1/2), -\log\tilde{t})$ and

$$Aw := \Delta w - 2\nabla'(\partial_n w) \cdot \nabla'g + (\partial_n^2 w)|\nabla'g|^2 - (\partial_n w)\Delta'g$$

by abbreviations of the subscripts η and η_n . In addition,

$$\Omega(\tau) := \{\eta \in \mathbf{R}^n; \tilde{\xi} + e^{-\tau/2}\eta \in \Phi(\Omega_R)\} = e^{\tau/2}(\Phi(\Omega_R) - \tilde{\xi}).$$

By using the backward similarity variables, (3.6) can be written as

$$E(t) = \int_{\Omega(\tau)} \left(\frac{|\widehat{\nabla}w|^2}{2} - \frac{|w|^{p+1}}{p+1} + \frac{w^2}{2(p-1)} \right) \\ \times \exp\left(-\frac{1}{4}e^\tau |\Psi(\tilde{\xi} + e^{-\tau/2}\eta) - \Psi(\tilde{\xi})|^2\right) \\ \times \varphi^2\left(\frac{4}{R} |\Psi(\tilde{\xi} + e^{-\tau/2}\eta) - \Psi(\tilde{\xi})|\right) d\eta,$$

where $\widehat{\nabla}w := (\nabla'w - (\partial_n w)\nabla'g, \partial_n w)$. We observe that

$$(3.10) \quad \begin{aligned} & |\Psi(\tilde{\xi} + e^{-\tau/2}\eta) - \Psi(\tilde{\xi})|^2 \\ &= e^{-\tau} |(\eta', \eta_n + g(\eta', \tau) - g(0, \tau))|^2 \\ &= e^{-\tau} (|\eta|^2 + 2(g(\eta', \tau) - g(0, \tau))\eta_n + (g(\eta', \tau) - g(0, \tau))^2). \end{aligned}$$

Then by setting

$$(3.11) \quad \mathcal{E}(\tau) := \int_{\Omega(\tau)} \left(\frac{|\widehat{\nabla}w|^2}{2} - \frac{|w|^{p+1}}{p+1} + \frac{w^2}{2(p-1)} \right) \rho \psi^2 d\eta,$$

$$(3.12) \quad \begin{aligned} \rho = \rho(\eta, \tau) := & \exp \left(-\frac{1}{4} (|\eta|^2 + 2(g(\eta', \tau) - g(0, \tau))\eta_n \right. \\ & \left. + (g(\eta', \tau) - g(0, \tau))^2) \right), \end{aligned}$$

$$(3.13) \quad \psi = \psi(\eta, \tau) := \varphi \left(\frac{4|\Psi(\tilde{\xi} + e^{-\tau/2}\eta) - \Psi(\tilde{\xi})|}{R} \right),$$

we see that $E(t) = \mathcal{E}(\tau)$ with $\tau = -\log(\tilde{t} - t)$. By direct computations, we note that

$$(3.14) \quad \left\{ \begin{aligned} \partial_i \rho &= -\frac{1}{2} (\eta_i + (\eta_n + g(\eta', \tau) - g(0, \tau))\partial_i g) \rho \quad (i = 1, \dots, n-1), \\ \partial_i^2 \rho &= -\frac{1}{2} (1 + (\partial_i g)^2 + (\eta_n + g(\eta', \tau) - g(0, \tau))\partial_i^2 g) \rho \\ &\quad + \frac{1}{4} (\eta_i^2 + 2(\eta_n + g(\eta', \tau) - g(0, \tau))\eta_i \partial_i g) \rho \\ &\quad + \frac{1}{4} (\eta_n + g(\eta', \tau) - g(0, \tau))^2 (\partial_i g)^2 \rho \quad (i = 1, \dots, n-1), \\ \partial_n \rho &= -\frac{1}{2} (\eta_n + g(\eta', \tau) - g(0, \tau)) \rho, \\ \partial_n^2 \rho &= -\frac{1}{2} \rho + \frac{1}{4} (\eta_n + g(\eta', \tau) - g(0, \tau))^2 \rho \end{aligned} \right.$$

and that

$$(3.15) \quad \rho_\tau = -\frac{1}{2} \partial_\tau (g(\eta', \tau) - g(0, \tau)) (\eta_n + g(\eta', \tau) - g(0, \tau)) \rho.$$

3.2. Quasi-monotonicity. We prove the quasi-monotonicity of E . This property plays a crucial role in the proof of ε -regularity and blow-up analysis.

Proposition 3.2. *Fix $R > 0$ such that either (R1) or (R2) holds. Let u be a solution of (2.1) satisfying (2.2). Then there exists $C' > 0$ depending only on n, p, Ω and R such that*

$$(3.16) \quad \begin{aligned} & E_{(\tilde{x}, \tilde{t})}(t; \phi_{\tilde{x}, R/4}) + \frac{1}{2} \int_{t'}^t (\tilde{t} - s)^{\frac{2}{p-1} - \frac{n}{2} - 1} \\ & \quad \times \int_{\Phi(\Omega_R)} \left| \frac{\hat{u}}{p-1} + \frac{(\xi - \Phi(\tilde{x})) \cdot \nabla \hat{u}}{2} - (\tilde{t} - s) \hat{u}_s \right|^2 \\ & \quad \times e^{-\frac{|\Psi(\xi) - \tilde{x}|^2}{4(\tilde{t} - s)}} \phi_{\tilde{x}, R/4}^2(\Psi(\xi)) d\xi ds \\ & \leq E_{(\tilde{x}, \tilde{t})}(t'; \phi_{\tilde{x}, R/4}) + C'(M + M^p)^2 (\tilde{t} - t') \end{aligned}$$

for any $\tilde{x} \in \overline{\Omega_{R/4}}$, $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$.

To prove this, we compute the derivative of \mathcal{E} .

Lemma 3.3. *The derivative of $\mathcal{E}(\tau)$ satisfies*

$$\frac{d}{d\tau}\mathcal{E}(\tau) = - \int w_\tau^2 \rho \psi^2 - \frac{1}{4}\mathcal{B}(\tau) + \mathcal{R}(\tau) + \mathcal{S}(\tau).$$

Here

$$\begin{aligned} \mathcal{B}(\tau) := & \int_{e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})} (\nabla w \cdot \nu)^2 \rho \psi^2 \\ & \times (\nu \cdot \eta)(1 - 2\nu_n \nabla' g \cdot \nu' + |\nabla' g|^2 \nu_n^2) dS(\eta), \end{aligned}$$

$\nu = (\nu', \nu_n)$ is the outward unit normal, dS is the surface area element,

$$\begin{aligned} \mathcal{R}(\tau) := & \int \frac{w^2}{p-1} \rho \psi \psi_\tau - \int \frac{2|w|^{p+1}}{p+1} \rho \psi \psi_\tau - \int w \rho \nabla w \cdot (\psi \nabla \psi)_\tau \\ & - \int w \psi \psi_\tau \nabla \cdot (\rho \nabla w) - 2 \int w_\tau \rho \psi \nabla w \cdot \nabla \psi \\ & + 2 \int w_\tau (\partial_n w) \rho \psi \nabla' \psi \cdot \nabla' g + 2 \int w (\partial_n w) \rho (\psi \nabla' \psi)_\tau \cdot \nabla' g \\ & + 2 \int w \psi \psi_\tau \nabla' (\rho \partial_n w) \cdot \nabla' g + 2 \int w (\partial_n w) \rho \psi \psi_\tau \Delta' g \\ & + 2 \int w_\tau \rho \psi (\partial_n \psi) \nabla' w \cdot \nabla' g - \int w (\partial_n w) \rho (\psi (\partial_n \psi))_\tau |\nabla' g|^2 \\ & - 2 \int w_\tau (\partial_n w) \rho \psi (\partial_n \psi) |\nabla' g|^2 - \int w \partial_n (\rho \partial_n w) \psi \psi_\tau |\nabla' g|^2 \\ & + \frac{1}{2} \int w_\tau (\partial_n w) \rho \psi^2 (g(\eta', \tau) - g(0, \tau) - \eta' \cdot \nabla' g) \end{aligned}$$

and

$$\mathcal{S}(\tau) := \int_{\Omega(\tau)} \left(\frac{|\widehat{\nabla} w|^2}{2} - \frac{|w|^{p+1}}{p+1} + \frac{w^2}{2(p-1)} \right) \rho_\tau \psi^2 d\eta$$

with the abbreviation $\int(\cdots) = \int_{\Omega(\tau)}(\cdots)d\eta$.

Proof. In what follows, we perform integration by parts several times, and so we observe the boundary value of $w\psi$. We claim that

$$(3.17) \quad w\psi = 0 \quad \text{on } \partial\Omega(\tau).$$

To prove this, we note that

$$\partial\Omega(\tau) = e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi}) \cup e^{\tau/2}(\Phi(\Omega \cap \partial B_R) - \tilde{\xi}).$$

For $\eta \in e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})$, by the boundary condition in (3.9), we obtain $w(\eta, \tau) = 0$. On the other hand, for $\eta \in e^{\tau/2}(\Phi(\Omega \cap \partial B_R) - \tilde{\xi})$, we have $\Phi(x) = \xi = e^{-\tau/2}\eta + \tilde{\xi} \in \Phi(\Omega \cap \partial B_R)$. Thus $x \in \Omega \cap \partial B_R$ and $|x| = R$. This together with $\tilde{x} \in \overline{\Omega_{R/4}}$ gives $|x - \tilde{x}| \geq 3R/4$. Therefore $\psi(\eta, \tau) = \varphi(4|x - \tilde{x}|/R) = 0$ for $\eta \in e^{\tau/2}(\Phi(\Omega \cap \partial B_R) - \tilde{\xi})$. Hence (3.17) holds.

For simplicity, we write $\int(\cdots) = \int_{\Omega(\tau)}(\cdots)d\eta$ when no confusion can arise. By (3.17), we see that

$$(3.18) \quad \begin{aligned} & \frac{d}{d\tau} \int \left(-\frac{|w|^{p+1}}{p+1} + \frac{w^2}{2(p-1)} \right) \rho\psi^2 \\ &= \int \left(-\frac{|w|^{p+1}}{p+1} + \frac{w^2}{2(p-1)} \right) \rho_\tau\psi^2 \\ & \quad + \int \frac{ww_\tau}{p-1} \rho\psi^2 - \int ww_\tau|w|^{p-1} \rho\psi^2 + \mathcal{R}_0, \end{aligned}$$

where

$$\mathcal{R}_0 := \int \frac{w^2}{p-1} \rho\psi\psi_\tau - \int \frac{2|w|^{p+1}}{p+1} \rho\psi\psi_\tau.$$

On the other hand, we set

$$(3.19) \quad \begin{aligned} \frac{d}{d\tau} \int \frac{|\widehat{\nabla}w|^2}{2} \rho\psi^2 &= \frac{1}{2} \frac{d}{d\tau} \int |\nabla w|^2 \rho\psi^2 - \frac{d}{d\tau} \int (\partial_n w) \rho\psi^2 \nabla' w \cdot \nabla' g \\ & \quad + \frac{1}{2} \frac{d}{d\tau} \int (\partial_n w)^2 \rho\psi^2 |\nabla' g|^2 + \int \frac{|\widehat{\nabla}w|^2}{2} \rho_\tau\psi^2 \\ &=: \frac{1}{2} \frac{dI_1}{d\tau} - \frac{dI_2}{d\tau} + \frac{1}{2} \frac{dI_3}{d\tau} + \int \frac{|\widehat{\nabla}w|^2}{2} \rho_\tau\psi^2. \end{aligned}$$

We estimate the derivatives of I_1 , I_2 and I_3 . Integration by parts and (3.17) show that

$$\begin{aligned} \frac{1}{2} \frac{dI_1}{d\tau} &= -\frac{d}{d\tau} \left(\int (w\psi\nabla\psi \cdot \nabla w) \rho + \frac{1}{2} \int w\psi^2 \nabla \cdot (\rho\nabla w) \right) \\ &= -\int (w\psi\nabla\psi \cdot \nabla w)_\tau \rho - \frac{1}{2} \int w_\tau \psi^2 \nabla \cdot (\rho\nabla w) \\ & \quad - \int w\psi\psi_\tau \nabla \cdot (\rho\nabla w) - \frac{1}{2} \int w\psi^2 \nabla \cdot (\rho\nabla w_\tau). \end{aligned}$$

Integrating by parts twice and (3.17) yield

$$\begin{aligned} -\frac{1}{2} \int w\psi^2 \nabla \cdot (\rho\nabla w_\tau) &= -\frac{1}{2} \int w_\tau \psi^2 \nabla \cdot (\rho\nabla w) - \int w_\tau \rho\psi \nabla w \cdot \nabla \psi \\ & \quad + \int w\rho\psi \nabla w_\tau \cdot \nabla \psi + \frac{1}{2} \int_{\partial\Omega(\tau)} w_\tau \rho\psi^2 \nabla w \cdot \nu dS. \end{aligned}$$

This implies that

$$(3.20) \quad \frac{1}{2} \frac{dI_1}{d\tau} = -\int w_\tau \psi^2 \nabla \cdot (\rho\nabla w) + \frac{1}{2} \int_{\partial\Omega(\tau)} w_\tau \rho\psi^2 \nabla w \cdot \nu dS + \mathcal{R}_1,$$

where

$$\begin{aligned} \mathcal{R}_1 &:= -\int (w\psi\nabla\psi \cdot \nabla w)_\tau \rho - \int w\psi\psi_\tau \nabla \cdot (\rho\nabla w) \\ & \quad - \int w_\tau \rho\psi \nabla w \cdot \nabla \psi + \int w\rho\psi \nabla w_\tau \cdot \nabla \psi. \end{aligned}$$

By the same argument of [37, Proposition 2.1] with $\Phi(\partial\Omega \cap B_R) \subset \mathbf{R}_+^n$, we can see that

$$(3.21) \quad \begin{aligned} & \frac{1}{2} \int_{\partial\Omega(\tau)} w_\tau \rho \psi^2 \nabla w \cdot \nu dS \\ &= -\frac{1}{4} \int_{e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})} (\nabla w \cdot \nu)^2 \rho \psi^2 (\nu \cdot \eta) dS. \end{aligned}$$

Indeed, from the boundary conditions in (3.5) and (3.9), it follows that

$$0 = \hat{u}_t = (\tilde{t} - t)^{-\frac{1}{p-1}-1} \left(\nabla w \cdot \frac{y}{2} + w_\tau \right),$$

and so $w_\tau = -\nabla w \cdot (\eta/2)$ on $e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})$. By the boundary condition in (3.9), we also have $\nabla w \cdot \nu^\perp = 0$ on the boundary, where ν^\perp is the unit tangent vector corresponding to ν . This implies $\nabla w = (\nabla w \cdot \nu)\nu$. Thus

$$(3.22) \quad w_\tau = -\frac{1}{2}(\nabla w \cdot \nu)(\nu \cdot \eta), \quad \eta \in e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi}).$$

Recall that $\psi(\eta, \tau) = 0$ for $\eta \in e^{\tau/2}(\Phi(\Omega \cap \partial B_R) - \tilde{\xi})$ by the proof of (3.17). Then (3.21) follows. For later use, since $\nabla w = (\nabla w \cdot \nu)\nu$, we note that

$$(3.23) \quad \nabla' w = (\nabla w \cdot \nu)\nu', \quad \partial_n w = (\nabla w \cdot \nu)\nu_n.$$

We consider I_2 . Integration by parts gives

$$\begin{aligned} -\frac{dI_2}{d\tau} &= \int w_\tau \psi^2 \nabla'(\rho \partial_n w) \cdot \nabla' g + \int w \psi^2 \nabla'(\rho \partial_n w_\tau) \cdot \nabla' g \\ &\quad + \int w_\tau (\partial_n w) \rho \psi^2 \Delta' g + \int w (\partial_n w_\tau) \rho \psi^2 \Delta' g + 2 \int w (\partial_n w) \rho \psi \psi_\tau \Delta' g \\ &\quad + 2 \int \rho (w (\partial_n w) \psi \nabla' \psi)_\tau \cdot \nabla' g + 2 \int w \psi \psi_\tau \nabla'(\rho \partial_n w) \cdot \nabla' g. \end{aligned}$$

Again by integrating by parts twice, we can see that

$$(3.24) \quad \begin{aligned} -\frac{dI_2}{d\tau} &= \int w_\tau \psi^2 \nabla'(\rho \partial_n w) \cdot \nabla' g + \int w_\tau \psi^2 \partial_n(\rho \nabla' w) \cdot \nabla' g \\ &\quad + \int w_\tau (\partial_n w) \rho \psi^2 \Delta' g - \int_{\partial\Omega(\tau)} w_\tau \rho \psi^2 \nu_n \nabla' w \cdot \nabla' g dS + \mathcal{R}_2, \end{aligned}$$

where

$$\begin{aligned} \mathcal{R}_2 &:= 2 \int \rho (w (\partial_n w) \psi \nabla' \psi)_\tau \cdot \nabla' g + 2 \int w \psi \psi_\tau \nabla'(\rho \partial_n w) \cdot \nabla' g \\ &\quad + 2 \int w (\partial_n w) \rho \psi \psi_\tau \Delta' g - 2 \int w (\partial_n w_\tau) \rho \psi \nabla' \psi \cdot \nabla' g \\ &\quad + 2 \int w_\tau \rho \psi (\partial_n \psi) \nabla' w \cdot \nabla' g. \end{aligned}$$

From the same computations as in the proof of (3.21), it follows that

$$(3.25) \quad \begin{aligned} & - \int_{\partial\Omega(\tau)} w_\tau \rho \psi^2 \nu_n \nabla' w \cdot \nabla' g dS \\ &= \frac{1}{2} \int_{e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})} (\nabla w \cdot \nu)^2 \rho \psi^2 (\nu \cdot \eta) \nu_n \nabla' g \cdot \nu' dS. \end{aligned}$$

Indeed, by (3.23), we have $\nabla' w \cdot \nabla' g = (\nabla w \cdot \nu)(\nabla' g \cdot \nu')$ on $e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})$. This together with (3.22) shows the above relation.

We examine I_3 . Again by the integration by parts, we have

$$\begin{aligned} \frac{1}{2} \frac{dI_3}{d\tau} &= -\frac{1}{2} \int w_\tau \psi^2 \partial_n(\rho \partial_n w) |\nabla' g|^2 - \frac{1}{2} \int w \psi^2 \partial_n(\rho \partial_n w_\tau) |\nabla' g|^2 \\ &\quad - \int (\psi(\partial_n \psi) w \partial_n w)_\tau \rho |\nabla' g|^2 - \int w \psi \psi_\tau \partial_n(\rho \partial_n w) |\nabla' g|^2. \end{aligned}$$

Again by integrating by parts twice, we also have

$$(3.26) \quad \begin{aligned} \frac{1}{2} \frac{dI_3}{d\tau} &= - \int w_\tau \psi^2 \partial_n(\rho \partial_n w) |\nabla' g|^2 \\ &\quad + \frac{1}{2} \int_{\partial\Omega(\tau)} w_\tau (\partial_n w) \rho \psi^2 |\nabla' g|^2 \nu_n dS + \mathcal{R}_3, \end{aligned}$$

where

$$\begin{aligned} \mathcal{R}_3 &:= - \int (w \partial_n w \psi (\partial_n \psi))_\tau \rho |\nabla' g|^2 - \int w \partial_n(\rho \partial_n w) \psi \psi_\tau |\nabla' g|^2 \\ &\quad + \int w (\partial_n w_\tau) \rho \psi (\partial_n \psi) |\nabla' g|^2 - \int w_\tau (\partial_n w) \rho \psi (\partial_n \psi) |\nabla' g|^2. \end{aligned}$$

By (3.22) and (3.23), we see that

$$(3.27) \quad \begin{aligned} &\frac{1}{2} \int_{\partial\Omega(\tau)} w_\tau (\partial_n w) \rho \psi^2 |\nabla' g|^2 \nu_n dS \\ &= -\frac{1}{4} \int_{e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})} (\nabla w \cdot \nu)^2 \rho \psi^2 (\nu \cdot \eta) \nu_n^2 |\nabla' g|^2 dS. \end{aligned}$$

Combining (3.18), (3.19), (3.20), (3.21), (3.24), (3.25), (3.26) and (3.27), we obtain

$$\frac{d}{d\tau} \mathcal{E}(\tau) = \mathcal{J} - \frac{1}{4} \mathcal{B} + \sum_{i=0}^3 \mathcal{R}_i + \mathcal{S},$$

where \mathcal{B} and \mathcal{S} are given in the statement of this lemma and

$$\begin{aligned} \mathcal{J} &:= \int w_\tau \psi^2 \left(\frac{w\rho}{p-1} - w|w|^{p-1}\rho - \nabla \cdot (\rho \nabla w) + \nabla'(\rho \partial_n w) \cdot \nabla' g \right. \\ &\quad \left. + \partial_n(\rho \nabla' w) \cdot \nabla' g + \rho \partial_n w \Delta' g - \partial_n(\rho \partial_n w) |\nabla' g|^2 \right). \end{aligned}$$

From (3.9) and $\nabla'(\partial_n w) = \partial_n \nabla' w$, it follows that

$$\begin{aligned} \mathcal{J} &= \int w_\tau \psi^2 \left(-w_\tau \rho - \frac{1}{2} \rho \eta \cdot \nabla w - \nabla \rho \cdot \nabla w + \partial_n \rho \nabla' w \cdot \nabla' g \right. \\ &\quad \left. + \partial_n w \nabla' \rho \cdot \nabla' g - \partial_n w (\partial_n \rho) |\nabla' g|^2 \right). \end{aligned}$$

By using (3.14), we have

$$\begin{aligned} &-\frac{1}{2} \rho \eta \cdot \nabla w - \nabla \rho \cdot \nabla w + \partial_n \rho \nabla' w \cdot \nabla' g = \frac{1}{2} (\partial_n w) \rho (g(\eta', \tau) - g(0, \tau)), \\ &\partial_n w \nabla' \rho \cdot \nabla' g - \partial_n w (\partial_n \rho) |\nabla' g|^2 = -\frac{1}{2} (\partial_n w) \rho \eta' \cdot \nabla' g. \end{aligned}$$

Thus,

$$\mathcal{J} = - \int w_\tau^2 \rho \psi^2 + \frac{1}{2} \int w_\tau (\partial_n w) \rho \psi^2 (g(\eta', \tau) - g(0, \tau) - \eta' \cdot \nabla' g).$$

Then by direct computations of $\sum \mathcal{R}_i$, we see that

$$\mathcal{J} + \sum_{i=0}^3 \mathcal{R}_i = - \int w_\tau^2 \rho \psi^2 + \mathcal{R}.$$

The proof is complete. \square

We observe \mathcal{B} .

Lemma 3.4. $\mathcal{B}(\tau) \geq 0$.

Proof. In (R1), we have $\mathcal{B} = 0$, since the domain of integration is far from the boundary. Thus, we consider the case (R2). Since $\Phi(\partial\Omega \cap B_R) \subset \mathbf{R}_+^n$ and $\tilde{\xi}_n = \Phi_n(\tilde{x}) = \tilde{x}_n - f(\tilde{x}') \geq 0$ for $\tilde{x} \in \overline{\Omega_{R/4}}$ by the choice of f , we see that $\nu \cdot \eta \geq 0$ for $\eta \in e^{\tau/2}(\Phi(\partial\Omega \cap B_R) - \tilde{\xi})$. In addition, since $|\nu'| \leq 1$, we have

$$\begin{aligned} & 1 - 2\nu_n \nabla' g \cdot \nu' + |\nabla' g|^2 \nu_n^2 \\ &= (1 - (\nabla' g \cdot \nu') \nu_n)^2 + (|\nabla' g|^2 - (\nabla' g \cdot \nu')^2) \nu_n^2 \geq 0. \end{aligned}$$

Then the lemma follows. \square

We estimate \mathcal{R} .

Lemma 3.5. \mathcal{R} satisfies

$$\mathcal{R}(\tau) \leq \frac{1}{2} \int w_\tau^2 \psi^2 \rho + C\tilde{\mathcal{R}}_1(\tau) + C\tilde{\mathcal{R}}_2(\tau)$$

for $\tau = -\log(\tilde{t} - t)$ with $-1/2 < t < 0$ and $1 > \tilde{t} > t$, where

$$\begin{aligned} (3.28) \quad \tilde{\mathcal{R}}_1(\tau) &:= \int (w^2 + |\nabla w|^2 + |w|^{p+1})(|\nabla \psi|^2 + |\psi_\tau| + \psi_\tau^2 + |\nabla \psi_\tau|) \rho, \\ \tilde{\mathcal{R}}_2(\tau) &:= \int |\nabla w|^2 \psi^2 e^{-\frac{|\eta|^2}{16}} e^{-\tau}. \end{aligned}$$

Proof. We only consider the case (R2), since the case (R1) is simpler. Indeed, in (R1), we can show that $\mathcal{R}(\tau) \leq (1/2) \int w_\tau^2 \psi^2 \rho + C\tilde{\mathcal{R}}_1(\tau)$.

By (3.8), the choice of f and $\tau > -\log(\tilde{t} + 1/2) > -\log(3/2)$, we have

$$(3.29) \quad \|\nabla' g\|_{L^\infty(\mathbf{R}^{n-1})} \leq \frac{1}{2}, \quad \|(\nabla')^2 g\|_{L^\infty(\mathbf{R}^{n-1})} \leq C e^{-\frac{1}{2}\tau} \leq C.$$

By (3.12) and (3.29), we also have

$$(3.30) \quad \rho(\eta, \tau) \leq \exp\left(-\frac{1}{4} \left(|\eta'|^2 + \frac{1}{2}\eta_n^2 - (g(\eta') - g(0))^2\right)\right) \leq e^{-\frac{|\eta|^2}{8}},$$

where $g(\eta') := g(\eta', \tau)$ and $g(0) := g(0, \tau)$. These inequalities together with integration by parts, Cauchy's inequality, $0 \leq \psi \leq 1$ and (3.29) show that

$$\begin{aligned} \mathcal{R}(\tau) &\leq \frac{1}{4} \int w_\tau^2 \rho \psi^2 + C \tilde{\mathcal{R}}_1(\tau) - \int w \psi \psi_\tau \nabla \cdot (\rho \nabla w) \\ &\quad + 2 \int w \psi \psi_\tau \nabla'(\rho \partial_n w) \cdot \nabla' g - \int w \psi \psi_\tau \partial_n(\rho \partial_n w) |\nabla' g|^2 \\ &\quad + \frac{1}{2} \int w_\tau (\partial_n w) \rho \psi^2 (g(\eta') - g(0) - \eta' \cdot \nabla' g) \\ &\leq \frac{1}{2} \int w_\tau^2 \rho \psi^2 + C \tilde{\mathcal{R}}_1(\tau) + C \int (\partial_n w)^2 \rho \psi^2 |\eta'|^4 e^{-\tau}. \end{aligned}$$

Then the lemma follows from (3.30) and $|\eta'|^4 e^{-|\eta|^2/8} \leq C e^{-|\eta|^2/16}$. \square

By Lemmas 3.3, 3.4 and 3.5, we see that

$$(3.31) \quad \mathcal{E}(\tau) + \frac{1}{2} \int_{\tau'}^\tau \int_{\Omega(\sigma)} w_\tau^2 \rho \psi^2 d\eta d\sigma \leq \mathcal{E}(\tau') + C \int_{\tau'}^\tau (\tilde{\mathcal{R}}_1 + \tilde{\mathcal{R}}_2 + |\mathcal{S}|)$$

for $\tau' = -\log(\tilde{t} - t')$ and $\tau = -\log(\tilde{t} - t)$ with $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$, where $\int_{\tau'}^\tau \tilde{\mathcal{R}}_i := \int_{\tau'}^\tau \tilde{\mathcal{R}}_i(\sigma) d\sigma$ for $i = 1, 2$. Note that this inequality holds for both (R1) and (R2). Let us estimate $\int_{\tau'}^\tau (\tilde{\mathcal{R}}_1 + \tilde{\mathcal{R}}_2)$.

Lemma 3.6. $\int_{\tau'}^\tau \tilde{\mathcal{R}}_1$ and $\int_{\tau'}^\tau \tilde{\mathcal{R}}_2$ satisfy

$$\int_{\tau'}^\tau \tilde{\mathcal{R}}_1(\sigma) d\sigma + \int_{\tau'}^\tau \tilde{\mathcal{R}}_2(\sigma) d\sigma \leq C(M + M^p)^2 (\tilde{t} - t')$$

for $\tau' = -\log(\tilde{t} - t')$ and $\tau = -\log(\tilde{t} - t)$ with $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$.

Proof. Let us first consider $\int_{\tau'}^\tau \tilde{\mathcal{R}}_1(\sigma) d\sigma$. We give precise proof only for

$$\int_{\tau'}^\tau \tilde{\mathcal{R}}_1^0(\sigma) d\sigma := \int_{\tau'}^\tau \int_{\Omega(\sigma)} |\nabla w|^2 |\nabla \psi|^2 \rho d\eta d\sigma,$$

since the other terms in the integration of (3.28) can be handled in the same way. Direct computations and $\|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})} \leq 1/2$ yield

$$\begin{aligned} |\nabla w(\eta, \tau)|^2 &= e^{-(\frac{2}{p-1}+1)\tau} |\nabla \hat{u}(\tilde{\xi} + e^{-\frac{1}{2}\tau} \eta, \tilde{t} - e^{-\tau})|^2 \\ &\leq C e^{-(\frac{2}{p-1}+1)\tau} |\nabla u(\tilde{\xi} + e^{-\frac{1}{2}\tau} \eta, \tilde{t} - e^{-\tau})|^2 \end{aligned}$$

and

$$\begin{aligned} |\nabla \psi(\eta, \tau)|^2 &= \frac{16e^{-\tau}}{R^2} \left| \varphi' \left(\frac{4|\Psi(\tilde{\xi} + e^{-\tau/2} \eta) - \Psi(\tilde{\xi})|}{R} \right) \right|^2 \\ &\quad \times \frac{|(\eta' + (\eta_n + g(\eta') - g(0)) \nabla' g, \eta_n + g(\eta') - g(0))|^2}{|(\eta', \eta_n + g(\eta') - g(0))|^2} \\ &\leq C e^{-\tau} \chi_{(\frac{1}{2}, 1)} \left(\frac{4|\Psi(\tilde{\xi} + e^{-\tau/2} \eta) - \Psi(\tilde{\xi})|}{R} \right), \end{aligned}$$

where $g(\eta') = g(\eta', \tau)$ and $g(0) = g(0, \tau)$. By (3.10) and (3.29), we can observe that

$$\begin{aligned} & \left\{ \eta; \frac{4|\Psi(\tilde{\xi} + e^{-\sigma/2}\eta) - \Psi(\tilde{\xi})|}{R} \geq \frac{1}{2} \right\} \\ & \subset \left\{ \eta; \frac{4}{R}(|\eta|^2 + \eta_n^2 + 2(g(\eta') - g(0))^2)e^{-\frac{\sigma}{2}} \geq \frac{1}{2} \right\} \\ & \subset \left\{ \eta; \frac{8}{R}e^{-\frac{\sigma}{2}}|\eta| \geq \frac{1}{2} \right\} \subset \left\{ \eta; |\eta| \geq \frac{R}{16}e^{\frac{\sigma}{2}} \right\} \end{aligned}$$

for $\tau' < \sigma < \tau$. Thus, on the set in the left-hand side, we have

$$\exp\left(-\frac{|\eta|^2}{8}\right) \leq \exp\left(-\frac{R^2 e^\sigma}{2048}\right).$$

From the above inequalities, $|\varphi'| \leq 4$ and (3.30), it follows that

$$\begin{aligned} \int_{\tau'}^{\tau} \tilde{\mathcal{R}}_1^0(\sigma) d\sigma & \leq C \int_{\tau'}^{\tau} \int_{\Omega(\sigma)} e^{-(\frac{2}{p-1}+2)\sigma} |\nabla u(\tilde{\xi} + e^{-\frac{\sigma}{2}}\eta, \tilde{t} - e^{-\sigma})|^2 \\ & \quad \times \chi_{(\frac{1}{2}, 1)} \left(\frac{4|\Psi(\tilde{\xi} + e^{-\sigma/2}\eta) - \Psi(\tilde{\xi})|}{R} \right) e^{-\frac{R^2}{2048}e^\sigma} d\eta d\sigma \\ & = C \int_{t'}^t \int_{\Omega_R} |\nabla u|^2 \chi_{(\frac{1}{2}, 1)} \left(\frac{4|x - \tilde{x}|}{R} \right) e^{(\frac{n}{2} - \frac{2}{p-1} + 1)\sigma - \frac{R^2}{2048}e^\sigma} dx ds. \end{aligned}$$

The conditions $\tilde{t} < 1$ and $t' > -1/2$ imply $\tau' = -\log(\tilde{t} - t') \geq -\log(3/2)$. Then, with the aid of a double exponential in the integrand, we observe that

$$e^{(\frac{n}{2} - \frac{2}{p-1} + 1)\sigma - \frac{R^2}{2048}e^\sigma} \leq C \quad \text{for } -\log(3/2) \leq \sigma < \infty.$$

Then by (3.1), we have

$$\int_{\tau'}^{\tau} \tilde{\mathcal{R}}_1^0(\sigma) d\sigma \leq C \int_{t'}^t \int_{\Omega_{R/2}} |\nabla u|^2 dx ds \leq C \int_{t'}^t \|\nabla u\|_{L^{2,2}(\Omega_{R/2})}^2 ds.$$

From the Hölder inequality for the Lorentz spaces and (4.1), it follows that

$$\begin{aligned} \int_{\tau'}^{\tau} \int_{\Omega(\sigma)} |\nabla w|^2 |\nabla \psi|^2 \rho dy d\sigma & \leq C \int_{t'}^t \|\nabla u\|_{L^{q^*, \infty}(\Omega_{R/2})}^2 ds \\ & \leq C(M + M^p)^2 (t - t') \leq C(M + M^p)^2 (\tilde{t} - t'). \end{aligned}$$

Computations similar to the above show that

$$\int_{\tau'}^{\tau} \tilde{\mathcal{R}}_1(\sigma) d\sigma \leq C(M + M^p)^2 (\tilde{t} - t').$$

We next estimate $\int_{\tau'}^{\tau} \tilde{\mathcal{R}}_2(\sigma) d\sigma$. By the change of variables and the same computations as in (3.2), we have

$$\begin{aligned} \int_{\tau'}^{\tau} \tilde{\mathcal{R}}_2(\sigma) d\sigma &\leq C \int_{\tau'}^{\tau} \int_{\Omega(\tau)} e^{-(\frac{2}{p-1}+2)\sigma} |\nabla u(\tilde{\xi} + e^{-\frac{\sigma}{2}}\eta, \tilde{t} - e^{-\sigma})|^2 \\ &\quad \times \varphi^2 \left(\frac{4|\Psi(\tilde{\xi} + e^{-\sigma/2}\eta) - \Psi(\tilde{\xi})|}{R} \right) e^{-\frac{|\eta|^2}{16}} d\eta d\sigma \\ &= C \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{\frac{2}{p-1}+1} \int_{\Omega_R} |\nabla u|^2 (\tilde{t} - s)^{-\frac{p}{2}} e^{-\frac{|x-\tilde{x}|^2}{16(\tilde{t}-s)}} \phi_{\tilde{x}, R/4}^2 dx ds \\ &\leq C(M + M^p)^2 \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{\frac{2}{p-1}+1 - \frac{p+1}{p-1}} ds \leq C(M + M^p)^2 (\tilde{t} - t'). \end{aligned}$$

Hence the lemma follows. \square

We estimate $\int_{\tau'}^{\tau} |\mathcal{S}|$.

Lemma 3.7. $\int_{\tau'}^{\tau} |\mathcal{S}(\sigma)| d\sigma$ satisfies

$$\int_{\tau'}^{\tau} |\mathcal{S}(\sigma)| d\sigma \leq C(M + M^p)^2 (\tilde{t} - t')$$

for $\tau' = -\log(\tilde{t} - t')$ and $\tau = -\log(\tilde{t} - t)$ with $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$.

Proof. From (3.15), (3.29), (3.30) and $\|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})} \leq 1/2$, it follows that

$$\begin{aligned} |\rho_\tau| &= \frac{1}{4} |e^{\frac{\tau}{2}} (f(\tilde{\xi}' + e^{-\frac{\tau}{2}}\eta') - f(\tilde{\xi}')) - \nabla' f(\tilde{\xi}' + e^{-\frac{\tau}{2}}\eta') \cdot \eta'| \\ &\quad \times |\eta_n + g(\eta', \tau) - g(0, \tau)| \rho \\ &\leq C|\eta'| (|\eta_n| + |\eta'|) e^{-\frac{|\eta|^2}{8}} \leq C e^{-\frac{|\eta|^2}{16}}. \end{aligned}$$

Then by the change of variables and the same computations as in Lemma 3.1, we obtain

$$\begin{aligned} \int_{\tau'}^{\tau} |\mathcal{S}(\sigma)| d\sigma &\leq C \int_{\tau'}^{\tau} \int_{\Omega(\tau)} (|\widehat{\nabla} w|^2 + |w|^{p+1} + w^2) e^{-\frac{|\eta|^2}{16}} \psi^2 d\eta d\sigma \\ &\leq C(M + M^p)^2 \int_{t'}^{\tilde{t}} ds \leq C(M + M^p)^2 (\tilde{t} - t'). \end{aligned}$$

Then the lemma follows. \square

We are now in a position to prove Proposition 3.2.

Proof of Proposition 3.2. By (3.31) and Lemmas 3.6 and 3.7, we have

$$(3.32) \quad \mathcal{E}(\tau) + \frac{1}{2} \int_{\tau'}^{\tau} \int_{\Omega(\sigma)} w_\tau^2 \rho \psi^2 d\eta d\sigma \leq \mathcal{E}(\tau') + C(M + M^p)^2 (\tilde{t} - t')$$

for $\tau' = -\log(\tilde{t} - t')$ and $\tau = -\log(\tilde{t} - t)$ with $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$. Observe that $\mathcal{E}(\tau) = E(t)$ and

$$w_\tau = (\tilde{t} - t)^{\frac{1}{p-1}} \left(-\frac{\hat{u}}{p-1} - \frac{(\xi - \tilde{\xi}) \cdot \nabla \hat{u}}{2} + (\tilde{t} - t) \hat{u}_t \right).$$

Then the desired inequality follows, and the proof is complete. \square

4. ε -REGULARITY

In this section, we prove an ε -regularity theorem. Note that, throughout the contents of this section, we will cite Theorem 4.1, Remark 4.2 and Lemma 4.6 to prove our main result.

Let u satisfy (2.1) and (2.2). Note that Proposition 2.1 gives

$$(4.1) \quad \sup_{-3/4 < t < 0} (\|u(\cdot, t)\|_{L^{q_c}(\Omega_{3R/4})} + \|\nabla u(\cdot, t)\|_{L^{q^*, \infty}(\Omega_{3R/4})}) \leq C(M + M^p).$$

Chou, Du and Zheng [13, Theorems 2 and 2'] proved ε -regularity theorems for global-in-time solutions (or borderline solutions) of (2.1) in convex domains. In [12, Theorem 2], the assumption on convexity was removed, see also [28, Proposition 4.2]. By the nontrivial modifications of [12, 13] with the aid of Blatt and Struwe [8], we show the following theorem for local-in-time solutions:

Theorem 4.1. *Let $n \geq 3$, $p > p_S$ and Ω be any smooth domain in \mathbf{R}^n with $0 \in \overline{\Omega}$. Fix $R > 0$ such that either (R1) or (R2) holds. Let u be a solution of (2.1) satisfying (2.2). Then there exist mutually independent constants ε_0 , δ_0 and θ with $0 < \varepsilon_0, \theta < 1$ and $0 < \delta_0 < R$ depending only on n, p, M, Ω and R such that the following holds: If there exists $0 < \delta \leq \delta_0$ such that*

$$(4.2) \quad \delta^{\frac{4}{p-1}-n} \iint_{Q_\delta} (|\nabla u|^2 + |u|^{p+1}) dxdt \leq \varepsilon_0,$$

then

$$(4.3) \quad \|u\|_{L^\infty(Q_{\theta\delta})} \leq C(\theta\delta)^{-\frac{2}{p-1}}.$$

Here C is a positive constant depending only on n, p, M, Ω and R and independent of $\varepsilon_0, \delta, \delta_0$ and θ .

Remark 4.2. Theorem 4.1 remains to hold for weak solutions satisfying an estimate of the form (4.9) in Lemma 4.6. In particular, we may apply it to the blow-up limit of certain rescaled solutions in Section 5.

In the rest of this section, C denotes a constant depending only on n, p, M, Ω and R . Each C may have different values also within the same line. Unless otherwise stated, we always assume either (R1) or (R2) and handle each of the cases in a unified way. In addition, we always assume that u is a solution of (2.1) satisfying (2.2).

We show an analog of the preliminary ε -regularity result [13, Lemma 3] by applying [8, Proposition 4.1].

Lemma 4.3. *Let $0 < R_1 < R$. Then there exist $\varepsilon_1 > 0$ and $C > 0$ depending only on n, p and Ω such that the following holds for each $0 < R_1 < R$: If*

$$(4.4) \quad (r/2)^{\frac{4}{p-1}-n} \iint_{Q_{r/2}(x_1, t_1)} |u(x, t)|^{p+1} dxdt \leq \varepsilon_1$$

for any $Q_r(x_1, t_1)$ with $Q_r(x_1, t_1) \subset Q_{2R_1}$, then $\|u\|_{L^\infty(Q_{R_1/4})} \leq CR_1^{-2/(p-1)}$.

Proof. Let $0 < R_1 < R$. We first prove the following statement: There exist $\varepsilon'_1 > 0$ and $C > 0$ depending only on n, p and Ω such that if

$$r^{\frac{4}{p-1}-n} \iint_{P_r(x'_1, t'_1) \cap Q_{R_1}} |u(x, t)|^{p+1} dxdt \leq \varepsilon'_1$$

for any $(x'_1, t'_1) \in Q_{R_1}$ and $r > 0$, then $\|u\|_{L^\infty(Q_{R_1/4})} \leq CR_1^{-2/(p-1)}$. To prove this, we set $\tilde{u}(y, s) := R_1^{2/(p-1)}u(R_1y, R_1^2s)$. Remark that \tilde{u} satisfies $\tilde{u}_s = \Delta_y \tilde{u} + |\tilde{u}|^{p-1}\tilde{u}$ in Q_1 , since $R_1 < R < 1$. Then by [8, Proposition 4.1], there exist $\varepsilon'_1 > 0$ and $C > 0$ depending only on n, p and Ω such that if \tilde{u} satisfies

$$\sup_{(\tilde{x}'_1, \tilde{t}'_1) \in Q_1, \tilde{r} > 0} \tilde{r}^{\frac{4}{p-1}-n} \iint_{P_{\tilde{r}}(\tilde{x}'_1, \tilde{t}'_1) \cap Q_1} |\tilde{u}(y, s)|^{p+1} dy ds \leq \varepsilon'_1,$$

then $\|\tilde{u}\|_{L^\infty(Q_{1/4})} \leq C$. By setting $\tilde{r} := r/R_1$, $\tilde{x}'_1 := x'_1/R_1$ and $\tilde{t}'_1 := t'_1/R_1^2$ and by the change of variables $x = R_1y$ and $t = R_1^2s$, we see that the inequality

$$\sup_{(x'_1, t'_1) \in Q_{R_1}, r > 0} r^{\frac{4}{p-1}-n} \iint_{P_r(x'_1, t'_1) \cap Q_{R_1}} |u(x, t)|^{p+1} dx dt \leq \varepsilon'_1$$

gives $\|u\|_{L^\infty(Q_{R_1/4})} \leq CR_1^{-2/(p-1)}$. Then the statement follows.

We next show the lemma. Set $\varepsilon_1 := 2^{4/(p-1)-n}\varepsilon'_1$. Note that $\varepsilon_1 < \varepsilon'_1$, since $4/(p-1) - n < 0$. Let $(x'_1, t'_1) \in Q_{R_1}$. In the case $r \leq R_1/2$, by $Q_{2r}(x'_1, t'_1) \subset Q_{2R_1}$ and (4.4) with r replaced by $2r$, we have

$$\iint_{P_r(x'_1, t'_1) \cap Q_{R_1}} |u|^{p+1} dx dt \leq \iint_{Q_r(x'_1, t'_1)} |u|^{p+1} dx dt \leq \varepsilon_1 r^{n-\frac{4}{p-1}} \leq \varepsilon'_1 r^{n-\frac{4}{p-1}}.$$

In the case $r > R_1/2$, by (4.4) again, we also have

$$\iint_{P_r(x'_1, t'_1) \cap Q_{R_1}} |u|^{p+1} dx dt \leq \iint_{Q_{R_1}} |u|^{p+1} dx dt \leq \varepsilon_1 R_1^{n-\frac{4}{p-1}} \leq \varepsilon'_1 r^{n-\frac{4}{p-1}}.$$

Hence the statement in the first part shows $\|u\|_{L^\infty(Q_{R_1/4})} \leq CR_1^{-2/(p-1)}$. \square

In the rest of this section, we prove Theorem 4.1 by using Lemma 4.3 and estimates of E . First of all, we replace δ with $A\delta$ in the assumption (4.2), where $A > 1$ is a large constant and $0 < \delta < 1$ is a small constant depending on A . Namely, we assume

$$(4.5) \quad \delta^{\frac{4}{p-1}-n} \iint_{Q_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) dx dt \leq A^{n-\frac{4}{p-1}} \varepsilon_0.$$

Here we take the constants A, ε_0, δ and θ as

$$(4.6) \quad A > 3, \quad 0 < \varepsilon_0 < 1, \quad 0 < \delta < \frac{R}{16A} < \frac{1}{16}, \quad 0 < \theta < \frac{1}{32A},$$

which will be specified later, see (4.17). For $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$, set

$$I_r = I_r(x_1, t_1) := (r/2)^{\frac{4}{p-1}-n} \iint_{Q_{r/2}(x_1, t_1)} |u|^{p+1} dx dt,$$

To apply Lemma 4.3, we examine the validity of the following statement:

$$(4.7) \quad I_r(x_1, t_1) \leq \varepsilon_1 \quad \text{for any } Q_r(x_1, t_1) \subset Q_{8A\theta\delta},$$

where ε_1 is given in Lemma 4.3. Note that

$$0 < r < 8A\theta\delta < \frac{1}{4}\delta < \delta < \frac{R}{16} < \frac{1}{16}.$$

Once we prove (4.7), Lemma 4.3 guarantees $\|u\|_{L^\infty(Q_{A\theta\delta})} \leq C(A\theta\delta)^{-2/(p-1)}$. Therefore our temporary task is to estimate I_r .

Proposition 4.4. *There exists a constant $\tilde{C} > 0$ depending only on n, p, M, Ω and R such that*

$$I_r(x_1, t_1) \leq \tilde{C}(h \circ \tilde{h}) \left(e^{-A^2/\tilde{C}} + A^{n-\frac{4}{p-1}} + \delta^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8s}} + \delta^2 \right)$$

for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$, where $h(s) := s + s^{1/2}$ and $\tilde{h}(s) := s + s^{2/(p+1)}$ for $s \geq 0$.

We prove this proposition by means of several lemmas.

Lemma 4.5. *There exists $C > 0$ such that*

$$(4.8) \quad I_r \leq C \int_{t_1 - \frac{r^2}{4}}^{t_1} (t_1 + r^2 - s)^{\frac{2}{p-1}} \int_{\Omega_R} |u|^{p+1} K_{(x_1, t_1+r^2)}(x, s) \phi_{x_1, R/8}^2 dx ds$$

for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$.

Proof. Recall that

$$I_r = (r/2)^{\frac{4}{p-1}-n} \int_{t_1 - \frac{r^2}{4}}^{t_1} \int_{\Omega_{r/2}(x_1)} |u|^{p+1} dx ds.$$

For $(x, s) \in Q_{r/2}(x_1, t_1)$, we have $|x - x_1| \leq r/2 < R/32$ and $t_1 - (r^2/4) < s < t_1$. Thus,

$$\begin{aligned} K_{(x_1, t_1+r^2)}(x, s) \phi_{x_1, R/8}^2(x) &= (t_1 + r^2 - s)^{-\frac{n}{2}} e^{-\frac{|x-x_1|^2}{4(t_1+r^2-s)}} \varphi^2 \left(\frac{8|x-x_1|}{R} \right) \\ &\geq (5/4)^{-\frac{n}{2}} r^{-n} e^{-\frac{1}{16}} \varphi^2(1/4) \geq Cr^{-n}. \end{aligned}$$

By $r^{4/(p-1)} \leq (t_1 + r^2 - s)^{2/(p-1)}$ for $t_1 - r^2/4 < s < t_1$, the lemma follows. \square

To estimate the right-hand side of (4.8), we prepare the following lemma by using Proposition 3.2.

Lemma 4.6. *There exists $C > 0$ such that*

$$(4.9) \quad \begin{aligned} &\int_{t'}^{\tilde{t}} (\tilde{t} - s)^{\frac{2}{p-1}} \int_{\Omega_R} |u(x, s)|^{p+1} K_{(\tilde{x}, \tilde{t})}(x, s) \phi_{\tilde{x}, R/4}^2(x) dx ds \\ &\leq C \left(\log \frac{\tilde{t} - t'}{\tilde{t} - t} \right)^{\frac{1}{2}} \left(E_{(\tilde{x}, \tilde{t})}(t'; \phi_{\tilde{x}, R/4}) - E_{(\tilde{x}, \tilde{t})}(t; \phi_{\tilde{x}, R/4}) + C(\tilde{t} - t') \right)^{\frac{1}{2}} \\ &\quad + C_p \left(E_{(\tilde{x}, \tilde{t})}(t'; \phi_{\tilde{x}, R/4}) + C(\tilde{t} - t') \right) \log \frac{\tilde{t} - t'}{\tilde{t} - t} + Ch(\tilde{t} - t') \end{aligned}$$

for any $\tilde{x} \in \overline{\Omega_{R/4}}$, $-1/2 < t' < t < 0$ and $1 > \tilde{t} > t$, where $C_p := 2(p+1)/(p-1)$.

Remark 4.7. We do not know the sign of $E_{(\tilde{x}, \tilde{t})}$. If we choose the reference time \tilde{t} as $\tilde{t} = 0$ (the existence time of u), then we can prove $E_{(\tilde{x}, 0)}(t) \geq -C(\tilde{t} - t)$ as in [13, Lemma 2]. However, we allow $\tilde{t} > 0$, and so it seems to the authors that this kind of lower estimate for $E_{(\tilde{x}, \tilde{t})}$ cannot be expected. We will give a lower estimate of E by using $p > p_S$, see Lemma 4.9.

Proof of Lemma 4.6. We handle (R1) and (R2) in a unified way. Define w, \mathcal{E}, ρ and ψ by (3.7), (3.11), (3.12) and (3.13), respectively. Then the left-hand side of the desired inequality equals

$$\int_{\tau'}^{\tau} \int_{\Omega(\sigma)} |w|^{p+1} \rho \psi^2 d\eta d\sigma.$$

We write $f(\cdots) = \int_{\Omega(\tau)}(\cdots)d\eta$ and $\iint(\cdots) = \int_{\tau'}^{\tau} \int_{\Omega(\sigma)}(\cdots)d\eta d\sigma$ unless otherwise stated. From (3.9) and (3.11), it follows that

$$\begin{aligned}
\frac{1}{2} \frac{d}{d\tau} \int w^2 \rho \psi^2 &= \int w w_\tau \rho \psi^2 + \int w^2 \psi \psi_\tau - 2\mathcal{E}(\tau) + 2\mathcal{E}(\tau) \\
&= -2\mathcal{E}(\tau) + \frac{p-1}{p+1} \int |w|^{p+1} \rho \psi^2 - \frac{1}{2} \int w \rho \psi^2 \nabla w \cdot \eta + \int w \rho \psi^2 \Delta w \\
&\quad - 2 \int w \rho \psi^2 \nabla'(\partial_n w) \cdot \nabla' g + \int w(\partial_n^2 w) \rho \psi^2 |\nabla' g|^2 \\
&\quad - \int w(\partial_n w) \rho \psi^2 \Delta' g + \int |\nabla w|^2 \rho \psi^2 - 2 \int (\partial_n w) \rho \psi^2 \nabla' w \cdot \nabla' g \\
&\quad + \int (\partial_n w)^2 \rho \psi^2 |\nabla' g|^2 + \int w^2 \rho \psi \psi_\tau.
\end{aligned}$$

Integrating by parts gives

$$\frac{1}{2} \frac{d}{d\tau} \int w^2 \rho \psi^2 = -2\mathcal{E}(\tau) + \frac{p-1}{p+1} \int |w|^{p+1} \rho \psi^2 + \tilde{\mathcal{J}} + \tilde{\mathcal{R}}_0,$$

where $\tilde{\mathcal{J}} = \tilde{\mathcal{J}}(\tau)$ and $\tilde{\mathcal{R}}_0 = \tilde{\mathcal{R}}_0(\tau)$ are given by

$$\begin{aligned}
\tilde{\mathcal{J}} &:= -\frac{1}{2} \int w \rho \psi^2 \nabla w \cdot \eta - \int w \psi^2 \nabla w \cdot \nabla \rho + 2 \int w(\partial_n w) \psi^2 \nabla' \rho \cdot \nabla' g \\
&\quad + \int w(\partial_n w) \rho \psi^2 \Delta' g - \int w(\partial_n w)(\partial_n \rho) \psi^2 |\nabla' g|^2, \\
\tilde{\mathcal{R}}_0 &:= -2 \int w \rho \psi \nabla w \cdot \nabla \psi + 4 \int w(\partial_n w) \rho \psi \nabla' \psi \cdot \nabla' g \\
&\quad - 2 \int w(\partial_n w) \rho \psi (\partial_n \psi) |\nabla' g|^2 + \int w^2 \rho \psi \psi_\tau.
\end{aligned}$$

Since $w \nabla w = \nabla(w^2)/2$ and $w \partial_n w = \partial_n(w^2)/2$, integrating by parts again shows that

$$\begin{aligned}
\tilde{\mathcal{J}} &= \int w^2 \psi^2 \left(\frac{\eta}{4} \cdot \nabla \rho + \frac{n}{4} \rho + \frac{1}{2} \Delta \rho - \partial_n(\nabla' \rho) \cdot \nabla' g \right. \\
&\quad \left. - \frac{1}{2} (\partial_n \rho) \Delta' g + \frac{1}{2} (\partial_n^2 \rho) |\nabla' g|^2 \right) \\
&\quad + \frac{1}{2} \int w^2 \rho \psi \nabla \psi \cdot \eta + \int w^2 \psi \nabla \psi \cdot \nabla \rho - 2 \int w^2 \psi (\partial_n \psi) \nabla' \rho \cdot \nabla' g \\
&\quad - \int w^2 \rho \psi (\partial_n \psi) \Delta' g + \int w^2 (\partial_n \rho) \psi (\partial_n \psi) |\nabla' g|^2.
\end{aligned}$$

From (3.14), (3.29), (3.30) and direct computations, it follows that

$$\begin{aligned}
& \frac{\eta}{4} \cdot \nabla \rho + \frac{n}{4} \rho + \frac{1}{2} \Delta \rho - \partial_n(\nabla' \rho) \cdot \nabla' g - \frac{1}{2}(\partial_n \rho) \Delta' g + \frac{1}{2}(\partial_n^2 \rho) |\nabla' g|^2 \\
&= \frac{1}{8}(\eta_n + g(\eta') - g(0))(g(\eta') - g(0) - \eta' \cdot \nabla' g) \rho \\
&\quad + \frac{1}{8}(\eta_n + g(\eta') - g(0))^2 (1 - 2|\nabla' g|^2) \rho \\
&\geq -\frac{1}{8} |\eta_n + g(\eta') - g(0)| |g(\eta') - g(0) - \eta' \cdot \nabla' g| \rho \\
&\geq -C |\eta|^3 e^{-\frac{\tau}{2}} e^{-\frac{|\eta|^2}{8}} \geq -C e^{-\frac{\tau}{2}} e^{-\frac{|\eta|^2}{16}}.
\end{aligned}$$

Thus,

$$\tilde{\mathcal{J}} \geq -C \int w^2 \psi^2 e^{-\frac{\tau}{2}} e^{-\frac{|\eta|^2}{16}} - C \int w^2 |\nabla \psi| e^{-\frac{|\eta|^2}{16}}.$$

On the other hand, we see that

$$\tilde{\mathcal{R}}_0 \geq -C \int (w^2 + |\nabla w|^2) (|\nabla \psi| + |\psi_\tau|) e^{-\frac{|\eta|^2}{16}}.$$

Hence by $\mathcal{E}(\tau) = E(t)$ and Proposition 3.2, we obtain

$$\begin{aligned}
& \frac{1}{2} \frac{d}{d\tau} \int w^2 \rho \psi^2 \\
&\geq -2\mathcal{E}(\tau) + \frac{p-1}{p+1} \int |w|^{p+1} \rho \psi^2 - C \bar{\mathcal{R}}_1 - C \bar{\mathcal{R}}_2 \\
&\geq -2(E(t') + C''(\tilde{t} - t')) + \frac{p-1}{p+1} \int |w|^{p+1} \rho \psi^2 - C \bar{\mathcal{R}}_1 - C \bar{\mathcal{R}}_2,
\end{aligned}$$

where $C'' := C'(M + M^p)^2$, C' is given in Proposition 3.2 and

$$\begin{aligned}
\bar{\mathcal{R}}_1 &:= \int (w^2 + |\nabla w|^2) (|\nabla \psi| + |\psi_\tau|) e^{-\frac{1}{16}|\eta|^2}, \\
\bar{\mathcal{R}}_2 &:= \int w^2 \psi^2 e^{-\frac{\tau}{2}} e^{-\frac{|\eta|^2}{16}}.
\end{aligned}$$

Set $C_p = 2(p+1)/(p-1)$. Then the above estimates imply that

$$\int |w|^{p+1} \rho \psi^2 \leq \frac{C_p}{4} \frac{d}{d\tau} \int w^2 \rho \psi^2 + C_p (E(t') + C''(\tilde{t} - t')) + C \bar{\mathcal{R}}_1 + C \bar{\mathcal{R}}_2.$$

We integrate this inequality over $\sigma \in (\tau', \tau)$ with $\tau' = -\log(\tilde{t} - t')$ and $\tau = -\log(\tilde{t} - t)$. Then,

$$\begin{aligned}
\iint |w|^{p+1} \rho \psi^2 &\leq \frac{C_p}{4} (\mathcal{K}(\tau) - \mathcal{K}(\tau')) + C_p (E(t') + C''(\tilde{t} - t')) \log \frac{\tilde{t} - t'}{\tilde{t} - t} \\
&\quad + C \int_{\tau'}^{\tau} \bar{\mathcal{R}}_1(\sigma) d\sigma + C \int_{\tau'}^{\tau} \bar{\mathcal{R}}_2(\sigma) d\sigma,
\end{aligned}$$

where

$$\mathcal{K}(\tau) := \int w^2(\eta, \tau) \rho(\eta) \psi^2(\eta, \tau) d\eta.$$

By computations similar to the proof of Lemma 3.6, we have

$$\begin{aligned} \int_{\tau'}^{\tau} \overline{\mathcal{R}}_1(\sigma) d\sigma &\leq C(\tilde{t} - t'), \\ \int_{\tau'}^{\tau} \overline{\mathcal{R}}_2(\sigma) d\sigma &= C \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{\frac{2}{p-1} - \frac{1}{2}} \int_{\Omega_R} |u|^2 (\tilde{t} - s)^{-\frac{n}{2}} e^{-\frac{|x-\tilde{x}|^2}{16(\tilde{t}-s)}} \phi_{\tilde{x}, R/4}^2 dx ds \\ &\leq C \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{-\frac{1}{2}} ds \leq C(\tilde{t} - t')^{\frac{1}{2}}. \end{aligned}$$

We estimate $|\mathcal{K}(\tau) - \mathcal{K}(\tau')|$. By computations similar to Lemma 3.6 again, we see that

$$\begin{aligned} |\mathcal{K}(\tau) - \mathcal{K}(\tau')| &= \left| \int_{\tau'}^{\tau} \mathcal{K}'(\sigma) d\sigma \right| = \left| \iint (2ww_{\tau} \rho \psi^2 + 2w^2 \rho \psi \psi_{\tau}) \right| \\ &\leq 2 \iint |w| |w_{\tau}| \rho \psi^2 + C(\tilde{t} - t'). \end{aligned}$$

The Hölder inequality and (3.32) with $\mathcal{E}(\tau) = E(t)$ yield

$$\begin{aligned} \iint |w| |w_{\tau}| \rho \psi^2 &\leq \left(\iint w^2 \rho \psi^2 \right)^{\frac{1}{2}} \left(\iint w_{\tau}^2 \rho \psi^2 \right)^{\frac{1}{2}} \\ &\leq \sqrt{2} \left(\iint w^2 \rho \psi^2 \right)^{\frac{1}{2}} (E(t') - E(t) + C''(\tilde{t} - t'))^{\frac{1}{2}}. \end{aligned}$$

Recall (3.30). Then computations similar to the estimate of $\overline{\mathcal{R}}_2$ give

$$\begin{aligned} \iint w^2 \rho \psi^2 &\leq \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{\frac{2}{p-1} - 1} \int_{\Omega} |u|^2 (\tilde{t} - s)^{-\frac{n}{2}} e^{-\frac{|x-\tilde{x}|^2}{8(\tilde{t}-s)}} \phi_{\tilde{x}, R/4}^2 dx ds \\ &\leq C \int_{t'}^{\tilde{t}} (\tilde{t} - s)^{-1} ds \leq C \log \frac{\tilde{t} - t'}{\tilde{t} - t}. \end{aligned}$$

Thus,

$$|\mathcal{K}(\tau) - \mathcal{K}(\tau')| \leq C \left(\log \frac{\tilde{t} - t'}{\tilde{t} - t} \right)^{\frac{1}{2}} (E(t') - E(t) + C''(\tilde{t} - t'))^{\frac{1}{2}} + C(\tilde{t} - t').$$

The above estimates show that $\int_{\tau'}^{\tau} \int_{\Omega(\sigma)} |w|^{p+1} \rho \psi^2 d\eta d\sigma$ is bounded by the right-hand side of the desired inequality. Then the lemma follows. \square

We estimate the right-hand side of (4.8). Since $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$, there is a possibility that $t_1 = 0$. Hence we first assume $t_1 < 0$, and then we consider the case $t_1 = 0$.

Assume $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$. Then by $x_1 \in \overline{\Omega_{R/4}}$, $-1/2 < t_1 - (r^2/4) < t_1 < 0$ and $1 > t_1 + r^2 > t_1$, we see that Lemma 4.6 (with $\phi_{x_1, R/4}$ replaced by $\phi_{x_1, R/8}$) gives

$$\begin{aligned} (4.10) \quad I_r &\leq C \left(E_{(x_1, t_1+r^2)} \left(t_1 - \frac{r^2}{4} \right) - E_{(x_1, t_1+r^2)}(t_1) + Cr^2 \right)^{\frac{1}{2}} \\ &\quad + C_p \left(E_{(x_1, t_1+r^2)} \left(t_1 - \frac{r^2}{4}; \phi_{x_1, R/8} \right) + Cr^2 \right) + Ch(r^2) \end{aligned}$$

for $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$. Then, we give estimates of $E_{(x_1, t_1+r^2)}(t_1 - r^2/4)$ and $E_{(x_1, t_1+r^2)}(t_1)$, respectively.

Lemma 4.8. *There exists $C > 0$ independent of t_1 such that*

$$E_{(x_1, t_1+r^2)} \left(t_1 - \frac{r^2}{4}; \phi_{x_1, R/8} \right) \leq E_{(x_1, t_1+r^2)} (-4\delta^2; \phi_{x_1, R/8}) + C\delta^2$$

for $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$.

Proof. By $x_1 \in \overline{\Omega_{R/4}}$, $-1/2 < -4\delta^2 < t_1 - r^2/4 < 0$, $1 > t_1 + r^2 > t_1 - r^2/4$ and Proposition 3.2 (with $\phi_{x_1, R/4}$ replaced by $\phi_{x_1, R/8}$), we see that

$$E_{(x_1, t_1+r^2)} \left(t_1 - \frac{r^2}{4}; \phi_{x_1, R/8} \right) \leq E_{(x_1, t_1+r^2)}(-4\delta^2; \phi_{x_1, R/8}) + C(t_1 + r^2 + 4\delta^2).$$

This implies the desired inequality. \square

Lemma 4.9. *There exists $C > 0$ independent of t_1 such that*

$$E_{(x_1, t_1+r^2)}(t_1; \phi_{x_1, R/8}) \geq -C(\delta^{\frac{n}{2} - \frac{p+1}{p-1}} + e^{-\frac{1}{8\delta}})$$

for $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$.

Remark 4.10. Since $p > p_S$, we have $(n/2) - (p+1)/(p-1) > 0$.

Proof of Lemma 4.9. By the definition of E and $\text{supp } \phi_{x_1, R/8} \subset B_{R/2}$, we have

$$\begin{aligned} & -E_{(x_1, t_1+r^2)}(t_1; \phi_{x_1, R/8}) \\ & \leq r^{\frac{2(p+1)}{p-1}} \int_{\Omega_{R/2}} \frac{|u(x, t_1)|^{p+1}}{p+1} K_{(x_1, t_1+r^2)}(x, t_1) dx \\ & = \frac{1}{p+1} r^{\frac{2(p+1)}{p-1} - n} \left(\int_{\Omega_{R/2} \cap B_{\sqrt{\delta}}(x_1)} + \int_{\Omega_{R/2} \setminus B_{\sqrt{\delta}}(x_1)} \right) |u|^{p+1} e^{-\frac{|x-x_1|^2}{4r^2}} dx \\ & =: \frac{1}{p+1} r^{\frac{2(p+1)}{p-1} - n} (I_1 + I_2). \end{aligned}$$

The Hölder inequality and (4.1) imply that

$$\begin{aligned} I_1 & \leq C \left(\int_{B_{\sqrt{\delta}}(x_1)} e^{-\frac{|x-x_1|^2}{Cr^2}} dx \right)^{1 - \frac{2(p+1)}{n(p-1)}} \\ & = C \left(r^n \int_0^{\sqrt{\delta}} \eta^{n-1} e^{-\frac{\eta^2}{C}} d\eta \right)^{1 - \frac{2(p+1)}{n(p-1)}} \leq Cr^{n - \frac{2(p+1)}{p-1}} \delta^{\frac{n}{2} - \frac{p+1}{p-1}}. \end{aligned}$$

On the other hand, by (2.2), we have

$$\begin{aligned} I_2 & = \int_{\Omega_{R/2} \setminus B_{\sqrt{\delta}}(x_1)} |u(x, t_1)|^{p+1} e^{-\frac{|x-x_1|^2}{8r^2}} e^{-\frac{|x-x_1|^2}{8r^2}} dx \\ & \leq e^{-\frac{\delta}{8r^2}} \int_{\Omega_{R/2} \setminus B_{\sqrt{\delta}}(x_1)} |u(x, t_1)|^{p+1} e^{-\frac{|x-x_1|^2}{8r^2}} dx \\ & \leq Ce^{-\frac{\delta}{8r^2}} \left(\int_{\mathbf{R}^n} e^{-\frac{|x-x_1|^2}{Cr^2}} dx \right)^{1 - \frac{2(p+1)}{n(p-1)}} \leq Cr^{n - \frac{2(p+1)}{p-1}} e^{-\frac{\delta}{8r^2}}. \end{aligned}$$

Since $r < \delta$, we have $e^{-\delta/(8r^2)} \leq e^{-1/(8\delta)}$. Then the lemma follows. \square

By using the above lemmas, we can estimate I_r by using $E_{(x_1, t_1+r^2)}(-4\delta^2)$. Note that we also consider the case $t_1 = 0$ in the following lemma:

Lemma 4.11. *There exists $C > 0$ such that*

$$I_r \leq Ch \left(E_{(x_1, t_1+r^2)}(-4\delta^2; \phi_{x_1, R/8}) + C(\delta^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8\delta}} + \delta^2) \right) + Ch(\delta^2)$$

for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$.

Proof. By combining (4.10) and Lemmas 4.8 and 4.9, we see that

$$\begin{aligned} I_r &\leq C \left(E_{(x_1, t_1+r^2)}(-4\delta^2) + C\delta^2 + C(\delta^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8\delta}}) + Cr^2 \right)^{\frac{1}{2}} \\ &\quad + C \left(E_{(x_1, t_1+r^2)}(-4\delta^2) + C\delta^2 + Cr^2 \right) + Ch(r^2) \\ &\leq Ch \left(E_{(x_1, t_1+r^2)}(-4\delta^2) + C\delta^2 + C(\delta^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8\delta}}) + Cr^2 \right) + Ch(r^2) \end{aligned}$$

for $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$. By using $r < \delta$, we obtain the desired inequality for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 < 0$. Since the constants in the right-hand side are independent of t_1 , by letting $t_1 \rightarrow 0$, we see that the desired inequality also holds for $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$ with $t_1 = 0$. \square

We estimated the right-hand side of (4.8), and then obtained Lemma 4.11. To prove Proposition 4.4, we estimate $E_{(x_1, t_1+r^2)}(-4\delta^2)$ by using some spatial integrals of $|\nabla u|^2$ and $|u|^{p+1}$, where $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$.

Lemma 4.12. *There exists $C > 0$ such that*

$$E_{(x_1, t_1+r^2)}(-4\delta^2; \phi_{x_1, R/8}) \leq C\tilde{h}(\delta^{\frac{4}{p-1}+2} J_r(t)) + C\delta^2$$

for any $-9\delta^2 \leq t \leq -4\delta^2$ and $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$, where h and \tilde{h} are given in Proposition 4.4 and

$$\begin{aligned} J_r(t) &= J_r(t; x_1, t_1, R) \\ &:= \int_{\Omega_R} (|\nabla u(x, t)|^2 + |u(x, t)|^{p+1}) K_{(x_1, t_1+r^2)}(x, t) \phi_{x_1, R/8}^2(x) dx. \end{aligned}$$

Proof. We claim that

$$E_{(x_1, t_1+r^2)}(-4\delta^2) \leq E_{(x_1, t_1+r^2)}(t) + C\delta^2$$

for $-9\delta^2 \leq t \leq -4\delta^2$. If $t = -4\delta^2$, this inequality is clear. Then it suffices to consider the case $-9\delta^2 \leq t < -4\delta^2$. Since $x_1 \in \overline{\Omega} \cap B_{R/4}$, $-1/2 < t < -4\delta^2 < 0$ and $1 > t_1 + r^2 > -4\delta^2$, Proposition 3.2 shows the claim. Indeed,

$$E_{(x_1, t_1+r^2)}(-4\delta^2) \leq E_{(x_1, t_1+r^2)}(t) + C(t_1 + r^2 - t) \leq E_{(x_1, t_1+r^2)}(t) + C\delta^2.$$

From the definition of E , it follows that

$$\begin{aligned} E_{(x_1, t_1+r^2)}(t) &\leq \frac{1}{2}(t_1 + r^2 - t)^{\frac{p+1}{p-1}} \int_{\Omega_R} |\nabla u|^2 K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx \\ &\quad + \frac{1}{2}(t_1 + r^2 - t)^{\frac{2}{p-1}} \int_{\Omega_R} |u|^2 K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx \\ &\leq C\delta^{\frac{2(p+1)}{p-1}} \int_{\Omega_R} |\nabla u|^2 K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx \\ &\quad + C\delta^{\frac{4}{p-1}} \int_{\Omega_R} |u|^2 K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx. \end{aligned}$$

By the Hölder inequality, $\phi_{x_1, R/8} \leq 1$ and $\int_{\mathbf{R}^n} K dx = (4\pi)^{n/2}$, we have

$$\begin{aligned} E_{(x_1, t_1+r^2)}(t) &\leq C\delta^{\frac{4}{p-1}+2} \int_{\Omega_R} |\nabla u|^2 K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx \\ &\quad + C \left(\delta^{\frac{4}{p-1}+2} \int_{\Omega_R} |u|^{p+1} K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^{p+1} dx \right)^{\frac{2}{p+1}} \\ &\leq C\tilde{h} \left(\delta^{\frac{4}{p-1}+2} \int_{\Omega_R} (|\nabla u|^2 + |u|^{p+1}) K_{(x_1, t_1+r^2)} \phi_{x_1, R/8}^2 dx \right). \end{aligned}$$

Hence the desired inequality follows. \square

We estimate $J_r(t; x_1, t_1, R)$ uniformly for x_1, t_1 and r .

Lemma 4.13. *There exists $C > 0$ such that*

$$J_r(t; x_1, t_1, R) \leq C J_0(t; 0, 20\delta^2, 2R)$$

for any $-9\delta^2 \leq t \leq -4\delta^2$ and $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$.

Proof. By the definition of J_r , it suffices to show that

$$(4.11) \quad K_{(x_1, t_1+r^2)}(x, t) \leq C K_{(0, 20\delta^2)}(x, t),$$

$$(4.12) \quad \phi_{x_1, R/8}^2(x) \leq C \phi_{0, R/4}^2(x),$$

for $x \in \Omega_R$ and $-9\delta^2 \leq t \leq -4\delta^2$. Since $(x_1, t_1) \in Q_{4A\theta\delta}$ and $A\theta < 1/4$, we have

$$\begin{aligned} \frac{K_{(x_1, t_1+r^2)}(x, t)}{K_{(0, 20\delta^2)}(x, t)} &= \left(\frac{20\delta^2 - t}{t_1 + r^2 - t} \right)^{\frac{n}{2}} \exp \left(-\frac{|x - x_1|^2}{4(t_1 + r^2 - t)} + \frac{|x|^2}{4(20\delta^2 - t)} \right) \\ &\leq \left(\frac{20\delta^2 - t}{-(4A\theta\delta)^2 - t} \right)^{\frac{n}{2}} \exp \left(-\frac{|x - x_1|^2}{4(\delta^2 - t)} + \frac{|x|^2}{4(20\delta^2 - t)} \right) \\ &\leq C \exp \left(-\frac{|x - x_1|^2}{40\delta^2} + \frac{|x|^2}{96\delta^2} \right). \end{aligned}$$

If $|x| \geq 10\delta$, then

$$|x - x_1| \geq |x| - |x_1| \geq 10\delta - 8A\theta\delta \geq 10\delta - \delta = 9\delta,$$

and so

$$\frac{|x|}{|x - x_1|} \leq \frac{|x - x_1| + |x_1|}{|x - x_1|} \leq 1 + \frac{8A\theta\delta}{9\delta} \leq \frac{10}{9}.$$

Hence, if $|x| \geq 10\delta$, then

$$\frac{K_{(x_1, t_1+r^2)}(x, s)}{K_{(0, 20\delta^2)}(x, t)} \leq C \exp \left(-\frac{|x - x_1|^2}{40\delta^2} + \frac{25|x - x_1|^2}{24 \cdot 81\delta^2} \right) \leq C.$$

On the other hand, if $|x| < 10\delta$, then

$$\frac{K_{(x_1, t_1+r^2)}(x, t)}{K_{(0, 20\delta^2)}(x, t)} \leq C \exp \left(-\frac{|x - x_1|^2}{40\delta^2} + \frac{25}{24} \right) \leq C e^{\frac{25}{24}}.$$

Therefore, we obtain (4.11).

To check (4.12), we prove $\varphi(8|x - x_1|/R) \leq C\varphi(4|x|/R)$. In the case $|x - x_1| \leq R/8$, we have

$$|x| \leq |x - x_1| + |x_1| \leq \frac{R}{8} + 8A\theta\delta \leq \frac{3}{16}R,$$

and so $\varphi(4|x|/R) \geq \varphi(3/4) > 0$ for $|x - x_1| \leq R/8$. Thus,

$$\varphi(8|x - x_1|/R) \leq 1 \leq \frac{\varphi(4|x|/R)}{\varphi(3/4)}.$$

In the case $|x - x_1| \geq R/8$, we see that $\varphi(8|x - x_1|/R) = 0 \leq \varphi(4|x|/R)$. Hence (4.12) holds. \square

By Lemmas 4.12 and 4.13, to obtain a bound of $E_{(x_1, t_1+r^2)}(-4\delta^2)$, it suffices to estimate $J_0(t; 0, 20\delta^2, 2R)$ at some $t = \hat{t}$.

Lemma 4.14. *There exists $-9\delta^2 \leq \hat{t} \leq -4\delta^2$ such that*

$$\delta^{\frac{4}{p-1}+2} J_0(\hat{t}; 0, 20\delta^2, 2R) \leq CA^{n-\frac{4}{p-1}} \varepsilon_0 + Ce^{-A^2/C}.$$

Proof. By the mean value theorem, the lemma will be proved once we show

$$\delta^{\frac{4}{p-1}} \int_{-9\delta^2}^{-4\delta^2} J_0(s; 0, 20\delta^2, 2R) ds \leq CA^{n-\frac{4}{p-1}} \varepsilon_0 + Ce^{-A^2/C}.$$

To prove this, we show that

$$(4.13) \quad \begin{aligned} J_1 &:= \delta^{\frac{4}{p-1}} \int_{-9\delta^2}^{-4\delta^2} \int_{\Omega_R \cap B_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) K_{(0, 20\delta^2)} \phi_{0, R/4}^2 dx dt \\ &\leq CA^{n-\frac{4}{p-1}} \varepsilon_0, \end{aligned}$$

$$(4.14) \quad \begin{aligned} J_2 &:= \delta^{\frac{4}{p-1}} \int_{-9\delta^2}^{-4\delta^2} \int_{\Omega_R \setminus B_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) K_{(0, 20\delta^2)} \phi_{0, R/4}^2 dx dt \\ &\leq Ce^{-A^2/C}. \end{aligned}$$

For (4.13), we recall our assumption (4.5), that is,

$$\delta^{\frac{4}{p-1}-n} \int_{-(A\delta)^2}^0 \int_{\Omega_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) dx ds \leq A^{n-\frac{4}{p-1}} \varepsilon_0.$$

Since $A > 3$, we have

$$\delta^{\frac{4}{p-1}-n} \int_{-9\delta^2}^{-4\delta^2} \int_{\Omega_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) dx ds \leq A^{n-\frac{4}{p-1}} \varepsilon_0.$$

This gives (4.13). Indeed,

$$\begin{aligned} J_1 &\leq \delta^{\frac{4}{p-1}} \int_{-9\delta^2}^{-4\delta^2} \int_{\Omega_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) (20\delta^2 - s)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4(20\delta^2 - s)}} dx ds \\ &\leq C \delta^{\frac{4}{p-1}-n} \int_{-9\delta^2}^{-4\delta^2} \int_{\Omega_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) dx ds \leq CA^{n-\frac{4}{p-1}} \varepsilon_0. \end{aligned}$$

We show (4.14). For $x \in \Omega_R \setminus B_{A\delta}$ and $-9\delta^2 \leq s \leq -4\delta^2$, we see that

$$\begin{aligned} \frac{K_{(0, 20\delta^2)}(x, s)}{K_{(0, 21\delta^2)}(x, s)} &= \left(\frac{21\delta^2 - s}{20\delta^2 - s} \right)^{\frac{n}{2}} \exp \left(-\frac{|x|^2}{4} \frac{\delta^2}{(20\delta^2 - s)(21\delta^2 - s)} \right) \\ &\leq Ce^{-A^2/C}. \end{aligned}$$

Thus,

$$\begin{aligned}
(4.15) \quad J_2 &\leq C e^{-A^2/C} \int_{-9\delta^2}^{-4\delta^2} \delta^{\frac{4}{p-1}} \int_{\Omega_R \setminus B_{A\delta}} (|\nabla u|^2 + |u|^{p+1}) \\
&\quad \times K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx ds \\
&\leq C e^{-A^2/C} \int_{-9\delta^2}^{-4\delta^2} (21\delta^2 - s)^{\frac{2}{p-1}} \int_{\Omega_R} (|\nabla u|^2 + |u|^{p+1}) \\
&\quad \times K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx ds.
\end{aligned}$$

We estimate the term including $|u|^{p+1}$. Since $0 \in \overline{\Omega_{R/4}}$, $-1/2 < -9\delta^2 < -4\delta^2 < 0$ and $1 > 21\delta^2 > -4\delta^2$, Lemmas 4.6 and 3.1 give

$$\begin{aligned}
(4.16) \quad &\int_{-9\delta^2}^{-4\delta^2} (21\delta^2 - s)^{\frac{2}{p-1}} \int_{\Omega_R} |u|^{p+1} K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx ds \\
&\leq C \left(\log \frac{30}{25} \right)^{\frac{1}{2}} (E_{(0,21\delta^2)}(-9\delta^2) - E_{(0,21\delta^2)}(-4\delta^2) + C(30\delta^2))^{\frac{1}{2}} \\
&\quad + C_p (E_{(0,21\delta^2)}(-9\delta^2) + C(30\delta^2)) \log \frac{30}{25} + Ch(30\delta^2) \leq C.
\end{aligned}$$

For the term including $|\nabla u|^2$, the definition of E and Lemma 3.1 imply that

$$\begin{aligned}
&\int_{\Omega_R} |\nabla u|^2 K_{(0,21\delta^2)} \phi_{0,R/4}^4 dx \\
&\leq 2(21\delta^2 - s)^{-\frac{p+1}{p-1}} E_{(0,21\delta^2)}(s) + \frac{2}{p+1} \int_{\Omega_R} |u|^{p+1} K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx \\
&\leq C \delta^{-\frac{2(p+1)}{p-1}} + C \int_{\Omega_R} |u|^{p+1} K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx
\end{aligned}$$

for $-9\delta^2 < s < -4\delta^2$. By multiplying $(21\delta^2 - s)^{2/(p-1)}$, integrating over $s \in (-9\delta^2, -4\delta^2)$ and using (4.16), we see that

$$\begin{aligned}
&\int_{-9\delta^2}^{-4\delta^2} (21\delta^2 - s)^{\frac{2}{p-1}} \int_{\Omega_R} |\nabla u|^2 K_{(0,21\delta^2)} \phi_{0,R/4}^2 dx ds \\
&\leq C \delta^{-\frac{2(p+1)}{p-1} + \frac{4}{p-1} + 2} + C \leq C.
\end{aligned}$$

This together with (4.15) and (4.16) implies (4.14). As explained in the first part of this proof, the lemma follows. \square

We are now in a position to prove Proposition 4.4.

Proof of Proposition 4.4. By combining Lemmas 4.12, 4.13 and 4.14, we obtain

$$E_{(x_1, t_1+r_2)}(-4\delta^2; \phi_{x_1, R/8}) \leq C \tilde{h}(A^{n-\frac{4}{p-1}} \varepsilon_0 + e^{-A^2/C}) + C\delta^2.$$

for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$. This together with Lemma 4.11 There exists $C > 0$ such that

$$I_r \leq Ch \left(C \tilde{h}(A^{n-\frac{4}{p-1}} \varepsilon_0 + e^{-A^2/C}) + C(\delta^{\frac{n}{2} - \frac{p+1}{p-1}} + e^{-\frac{1}{8\delta}} + \delta^2) \right) + Ch(\delta^2)$$

for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$. Then the desired inequality follows. \square

Finally, we complete the proof of Theorem 4.1.

Proof of Theorem 4.1. As explained in the first part of this subsection, it sufficed to prove that the statement (4.7) holds under the assumption (4.5). Let A , ε_0 , δ_0 and θ satisfy (4.6) with δ replaced by δ_0 . Let $Q_r(x_1, t_1) \subset Q_{8A\theta\delta_0}$. Then by Proposition 4.4, we have

$$\begin{aligned} I_r(x_1, t_1) &\leq \tilde{C}'(h \circ \tilde{h}) \left(e^{-A^2/\tilde{C}} \right) + \tilde{C}'(h \circ \tilde{h}) \left(A^{n-\frac{4}{p-1}} \varepsilon_0 \right) \\ &\quad + \tilde{C}'(h \circ \tilde{h}) \left(\delta_0^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8\delta_0}} + \delta_0^2 \right), \end{aligned}$$

where $\tilde{C}' > 0$ is a constant depending only on n , p , M , Ω and R . By taking (4.6) into account, we choose constants A , ε_0 , δ_0 and θ satisfying the following conditions:

$$(4.17) \quad \begin{cases} A > 3 & \text{with } \tilde{C}'(h \circ \tilde{h}) \left(e^{-A^2/\tilde{C}} \right) < \frac{\varepsilon_1}{3}, \\ 0 < \varepsilon_0 < 1 & \text{with } \tilde{C}'(h \circ \tilde{h}) \left(A^{n-\frac{4}{p-1}} \varepsilon_0 \right) < \frac{\varepsilon_1}{3}, \\ 0 < \delta_0 < \frac{R}{16A} & \text{with } \tilde{C}'(h \circ \tilde{h}) \left(\delta_0^{\frac{n}{2}-\frac{p+1}{p-1}} + e^{-\frac{1}{8\delta_0}} + \delta_0^2 \right) < \frac{\varepsilon_1}{3}, \\ 0 < \theta < \frac{1}{32A}. \end{cases}$$

Here ε_1 is given in Lemma 4.3. Remark that ε_0 , δ_0 and θ can be chosen independently.

Finally, we assume that there exists $0 < \delta \leq \delta_0$ such that (4.5) holds. Then we obtain $I_r(x_1, t_1) \leq \varepsilon_1$ for any $Q_r(x_1, t_1) \subset Q_{8A\theta\delta}$. Hence the statement (4.7) holds under the assumption (4.5), and so Lemma 4.3 shows that $\|u\|_{L^\infty(Q_{A\theta\delta})} \leq C(A\theta\delta)^{-2/(p-1)}$. Recall that we rewrote the assumptions in the first part of this subsection. Then we can conclude that the original assumption (4.2) implies (4.3). The proof of Theorem 4.1 is complete. \square

5. PROOF OF MAIN THEOREM

We first prove a localized statement of Theorem 1.1. At the level of strategy, the proof is based on [31] and [89, Proof of Theorem 9.8] for the Navier-Stokes equations, and [53, Chapter 8] and [85, 90] for the harmonic map heat flow. Indeed, we use compactness, backward uniqueness and unique continuation. However, the analysis in each of the steps seems to be more involved.

Theorem 5.1 (Localized statement). *Let $n \geq 3$, $p > ps$, Ω be any smooth domain in \mathbf{R}^n and u be a classical L^{q_c} -solution of (1.1) with $u_0 \in L^{q_c}(\Omega)$. If the maximal existence time $T > 0$ is finite and u has a blow-up point a on $\overline{\Omega}$, then for any $r > 0$,*

$$\limsup_{t \rightarrow T} \|u(\cdot, t)\|_{L^{q_c}(\Omega_r(a))} = \infty.$$

Note that if Ω is bounded, then Theorem 1.1 immediately follows from this theorem. The unbounded case will be considered in Subsection 5.4.

5.1. Existence of blow-up limit. To prove Theorem 5.1, we set up notation, and then we define rescaled solutions and take the limit. Let u be a classical L^{q_c} -solution of (1.1). Assume that $a \in \overline{\Omega}$ is a blow-up point of u . By translation and

scaling, we may assume that $a = 0$ and u satisfies

$$\begin{cases} u_t = \Delta u + |u|^{p-1}u & \text{in } \Omega_{R_1} \times (-1, 0), \\ u = 0 & \text{on } (\partial\Omega \cap B_{R_1}) \times (-1, 0), \end{cases}$$

for some $0 < R_1 < 1$. Here we assume that $t = 0$ is the blow-up time. Suppose, contrary to Theorem 5.1, that

$$(5.1) \quad \sup_{-1 < t < 0} \|u(\cdot, t)\|_{L^{q_c}(\Omega_{R_2})} \leq M$$

for some $M > 0$ and $0 < R_2 < 1$. Fix $0 < R < \min\{R_1, R_2\}$ so small that (R1) and (R2) hold. Note that u satisfies (2.1) and (2.2).

From the parabolic regularity estimates in Lemma A.2, it follows that

$$\|u\|_{W^{1,l}(-1/4, 0; L^r(\Omega_{R/2}))} \leq C(M + M^p)$$

for $1 \leq l < \infty$ and $1 \leq r \leq q_c/p$. Hence after a redefinition of a zero set in the time interval, we may assume $u \in C([-1/4, 0]; L^r(\Omega_{R/2}))$. This together with the uniform bound (2.2) gives

$$(5.2) \quad u \in C_{\text{weak}}([-1/4, 0]; L^{q_c}(\Omega_{R/2})).$$

By Theorem 4.1, there exist $\varepsilon_0 > 0$ and $0 < \delta_k < 1/2$ with $\delta_k \rightarrow 0$ as $k \rightarrow \infty$ such that

$$(5.3) \quad \delta_k^{\frac{4}{p-1}-n} \iint_{Q_{\delta_k}} (|\nabla u|^2 + |u|^{p+1}) dx dt > \varepsilon_0.$$

To take a blow-up limit of u , we define rescaled solutions and derive the corresponding equations. In the case (R1), we define

$$u_k(x, t) := \delta_k^{\frac{2}{p-1}} u(\delta_k x, \delta_k^2 t).$$

Then,

$$\begin{cases} (u_k)_t - \Delta u_k = |u_k|^{p-1}u_k & \text{in } \delta_k^{-1}\Omega_R \times (-\delta_k^{-2}, 0), \\ u_k = 0 & \text{on } \delta_k^{-1}(\partial\Omega \cap B_R) \times (-\delta_k^{-2}, 0). \end{cases}$$

In the case (R2), we define

$$(5.4) \quad u_k(x, t) := \delta_k^{\frac{2}{p-1}} \hat{u}(\delta_k x, \delta_k^2 t), \quad f_k(x') := \delta_k^{-1} f(\delta_k x'),$$

where \hat{u} is defined by (3.3). Then u_k satisfies

$$(5.5) \quad \begin{cases} (u_k)_t - A_k u_k = |u_k|^{p-1}u_k & \text{in } \delta_k^{-1}\Phi(\Omega_R) \times (-\delta_k^{-2}, 0), \\ u_k = 0 & \text{on } \delta_k^{-1}\Phi(\partial\Omega \cap B_R) \times (-\delta_k^{-2}, 0), \end{cases}$$

where

$$A_k u_k(x, t) := \Delta u_k - 2\nabla'(\partial_{x_n} u_k) \cdot \nabla' f_k + (\partial_{x_n}^2 u_k) |\nabla' f_k|^2 - (\partial_{x_n} u_k) \Delta' f_k.$$

In what follows, we take a limit of u_k and study its properties. Since the case (R1) is easier to handle than the case (R2), we focus on (R2). For $\rho > 0$, we set $B_\rho^+ := \mathbf{R}_+^n \cap B_\rho$. To take a limit of u_k , we give estimates of rescaled solutions uniformly for $k \geq k_\rho$. Here $k_\rho \geq 1$ is taken so that $0 < \delta_k < 1/2$, $\Psi(B_{3\rho\delta_k}^+) \subset \Omega_{R/2}$,

$\Psi(B_{3\rho\delta_k}) \subset B_{R/2}$ for all $k \geq k_\rho$. Since $\nabla' f(0) = 0$, $f \in C_0^\infty(\mathbf{R}^{n-1})$ and $\delta_k \rightarrow 0$ as $k \rightarrow \infty$, we have

$$(5.6) \quad |\nabla' f_k(x')| = |\nabla' f(\delta_k x')| \rightarrow 0,$$

$$(5.7) \quad \|\Delta' f_k\|_{L^\infty(\mathbf{R}^{n-1})} \leq \delta_k \|\Delta' f\|_{L^\infty(\mathbf{R}^{n-1})} \rightarrow 0.$$

We first give uniform estimates of u_k and ∇u_k .

Lemma 5.2. *Assume (R2). Let $\rho > 0$. Then there exists a constant $C > 0$ such that the rescaled functions u_k satisfy*

$$\begin{aligned} \sup_{-1 < t < 0} \|u_k(\cdot, t)\|_{L^{q_c}(B_\rho^+)} &\leq M, \\ \sup_{-1 < t < 0} \|\nabla u_k(\cdot, t)\|_{L^{q^*, \infty}(B_\rho^+)} &\leq C(M + M^p), \end{aligned}$$

for all $k \geq k_\rho$, where C depends on R and is independent of ρ and k .

Proof. By the change of variables and the choice of k_ρ , we can easily see the inequality for u_k . From the following computation

$$\begin{aligned} |\nabla u_k(x, t)| &= \delta_k^{\frac{2}{p-1}+1} |\nabla u(\Psi(\delta_k x), \delta_k^2 t) + \partial_{x_n} u(\Psi(\delta_k x), \delta_k^2 t) \nabla' f(\delta_k x')| \\ &\leq \delta_k^{\frac{2}{p-1}+1} (1 + \|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})}) |\nabla u(\Psi(\delta_k x), \delta_k^2 t)| \end{aligned}$$

and Proposition 2.1, it follows that

$$\begin{aligned} \|\nabla u_k(\cdot, t)\|_{L^{q^*, \infty}(B_\rho^+)} &\leq (1 + \|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})}) \|\nabla u(\cdot, \delta_k^2 t)\|_{L^{q^*, \infty}(\Omega_{R/2})} \\ &\leq C(M + M^p) \end{aligned}$$

for $-1 < t < 0$. Then the lemma follows. \square

Remark 5.3. For $\rho > 0$, by setting $u_k \equiv 0$ for $k < k_\rho$, we can consider that the obtained inequalities in Lemma 5.2 hold for all $k \geq 1$.

We next give a uniform parabolic regularity estimate.

Lemma 5.4. *Assume (R2). Let $1 \leq l < \infty$, $1 \leq r \leq q_c/p$ and $\rho > 0$. Then there exists $C > 0$ such that the rescaled functions u_k satisfy*

$$\|(u_k)_t\|_{L^l(-1, 0; L^r(B_\rho^+))} + \|\nabla^2 u_k\|_{L^l(-1, 0; L^r(B_\rho^+))} \leq C(M + M^p)$$

for all $k \geq k_\rho$, where C depends on ρ and R and is independent of k .

Proof. By direct computations, we have

$$\begin{aligned} &\|(u_k)_t\|_{L^l(-1, 0; L^r(B_\rho^+))} + \|\nabla^2 u_k\|_{L^l(-1, 0; L^r(B_\rho^+))} \\ &= \delta_k^{\frac{2}{p-1}+2-\frac{n}{r}-\frac{2}{l}} (\|\hat{u}_t\|_{L^l(-\delta_k^2, 0; L^r(B_{\rho\delta_k}^+))} + \|\nabla^2 \hat{u}\|_{L^l(-\delta_k^2, 0; L^r(B_{\rho\delta_k}^+))}). \end{aligned}$$

This together with Lemma A.1 shows the lemma. \square

Finally in this subsection, we show the existence of a blow-up limit. Let $\rho > 0$. To avoid technicalities due to the corner of ∂B_ρ^+ , we prepare a smooth domain \mathcal{B}_ρ satisfying $B_{\rho/2}^+ \subset \mathcal{B}_\rho \subset B_\rho^+$.

Lemma 5.5. *Assume (R2). There exist a subsequence still denoted by u_k and a function \bar{u} defined on $\mathbf{R}_+^n \times [-1, 0]$ satisfying the following statements:*

- (i) $u_k \rightarrow \bar{u}$ strongly in $L^\infty(-1, 0; W^{1,2}(\mathcal{B}_\rho))$ for each $\rho > 0$ as $k \rightarrow \infty$.
- (ii) $u_k \rightarrow \bar{u}$ strongly in $L^\infty(-1, 0; L^{p+1}(\mathcal{B}_\rho))$ for each $\rho > 0$ as $k \rightarrow \infty$.

- (iii) $\|\bar{u}\|_{L^\infty(-1,0;L^{q_c}(\mathbf{R}_+^n))} \leq M$.
- (iv) $\|\nabla\bar{u}\|_{L^\infty(-1,0;L^{q^*,\infty}(\mathbf{R}_+^n))} \leq C(M + M^p)$ with a constant $C > 0$ depending on R .

Here M is the constant in (5.1).

Proof. Let $1 < r < \min\{2, q_c/p\}$. By a consequence of an Aubin-Lions type compactness result (see Lemma B.2), we have

$$W^{1,5}(-1,0;L^r(\mathcal{B}_\rho)) \cap L^5(-1,0;W^{2,r}(\mathcal{B}_\rho)) \hookrightarrow C([-1,0];W^{1,r}(\mathcal{B}_\rho))$$

and this embedding is compact. Therefore, the uniform bounds in Lemma 5.4 show that there exists a subsequence still denoted by $\{u_k\}$ satisfying $u_k \rightarrow \bar{u}$ in $C([-1,0];W^{1,r}(\mathcal{B}_\rho))$ as $k \rightarrow \infty$. By Lemma 5.2, the rescaled functions u_k also satisfy the following uniform bounds:

$$(5.8) \quad \begin{aligned} & \sup_{-1 < t < 0} \|u_k(\cdot, t)\|_{L^{q_c}(\mathcal{B}_\rho)} \leq M, \\ & \sup_{-1 < t < 0} \|\nabla u_k(\cdot, t)\|_{L^{q^*,\infty}(\mathcal{B}_\rho)} \leq C(M + M^p), \end{aligned}$$

for all $k \geq k_\rho$, where C depends on R and is independent of ρ and k . Hence the standard interpolations give

$$\begin{aligned} \|u_k - \bar{u}\|_{L^\infty(-1,0;L_x^{p+1})} &\leq \|u_k - \bar{u}\|_{L^\infty(-1,0;L_x^r)}^{\theta_1} \|u_k - \bar{u}\|_{L^\infty(-1,0;L_x^{q_c})}^{1-\theta_1} \\ &\leq CM \|u_k - \bar{u}\|_{L^\infty(-1,0;L_x^r)}^{\theta_1}, \\ \|\nabla(u_k - \bar{u})\|_{L^\infty(-1,0;L_x^2)} &\leq \|\nabla(u_k - \bar{u})\|_{L^\infty(-1,0;L_x^r)}^{\theta_2} \|\nabla(u_k - \bar{u})\|_{L^\infty(-1,0;L_x^{q^*,\infty})}^{1-\theta_2} \\ &\leq C(M + M^p) \|\nabla(u_k - \bar{u})\|_{L^\infty(-1,0;L_x^r)}^{\theta_2}, \end{aligned}$$

where $L_x^r := L^r(\mathcal{B}_\rho)$ and $L_x^{q^*,\infty} := L^{q^*,\infty}(\mathcal{B}_\rho)$. These imply (i) and (ii). By the lower semicontinuity of the weak limit, we see that

$$\|\bar{u}(\cdot, t)\|_{L^{q_c}(\mathcal{B}_\rho)} \leq \liminf_{k \rightarrow \infty} \|u_k(\cdot, t)\|_{L^{q_c}(\mathcal{B}_\rho)} \leq M$$

for $-1 < t < 0$. Thus, letting $\rho \rightarrow \infty$ gives (iii). Since the constant C in (5.8) is independent of ρ , we also obtain (iv). The proof is complete. \square

Remark 5.6. By Lemma 5.4, up to subsequence still denoted by u_k , we can check that $(u_k)_t \rightarrow \bar{u}_t$ weakly in $L^r(\mathcal{B}_\rho \times (-1,0))$ for each $\rho > 0$ as $k \rightarrow \infty$.

5.2. Blow-up analysis. We continue to study the case (R2). For (R1), see Remark 5.13. By using Lemma 5.5, we show several properties of the blow-up limit \bar{u} . For $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t < 0$ and $\tilde{t} > t$, define a global weighted energy by

$$\overline{E}_{(\tilde{x}, \tilde{t})}(t) := (\tilde{t} - t)^{\frac{p+1}{p-1}} \int_{\mathbf{R}_+^n} \left(\frac{|\nabla \bar{u}|^2}{2} - \frac{|\bar{u}|^{p+1}}{p+1} + \frac{\bar{u}^2}{2(p-1)(\tilde{t} - t)} \right) K_{(\tilde{x}, \tilde{t})}(x, t) dx.$$

Note that the same computations as in Lemma 3.1 together with Lemma 5.5 (iii) and (iv) yield

$$(5.9) \quad |\overline{E}_{(\tilde{x}, \tilde{t})}(t)| \leq C(M + M^p)^2$$

for any $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t < 0$ and $\tilde{t} > t$.

We first show that \overline{E} is a scaling limit of E .

Lemma 5.7. *Assume (R2). Then*

$$E_{(\delta_k \tilde{x}, \delta_k^2 \tilde{t})}(\delta_k^2 t; \phi_{\delta_k \tilde{x}, R/4}) \rightarrow \overline{E}_{(\tilde{x}, \tilde{t})}(t) \quad \text{as } k \rightarrow \infty$$

for $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t < 0$ and $\tilde{t} > t$.

Proof. Fix $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t < 0$ and $\tilde{t} > t$. We take k so large that $\tilde{x} \in \delta_k^{-1} \overline{\Omega_{R/4}}$, $-(1/2)\delta_k^{-2} < t < 0$ and $\delta_k^{-2} > \tilde{t} > t$. By the definition of E and the change of variables, we have

$$\begin{aligned} & E_{(\delta_k \tilde{x}, \delta_k^2 \tilde{t})}(\delta_k^2 t; \phi_{\delta_k \tilde{x}, R/4}) \\ &= (\tilde{t} - t)^{\frac{p+1}{p-1} - \frac{n}{2}} \int_{\delta_k^{-1} \Phi(\Omega_R)} \left(\frac{|\widehat{\nabla} u_k|^2}{2} - \frac{|u_k|^{p+1}}{p+1} + \frac{|u_k|^2}{2(p-1)(\tilde{t}-t)} \right) \\ & \quad \times \exp\left(-\frac{|\delta_k^{-1} \Psi(\delta_k x) - \tilde{x}|^2}{4(\tilde{t}-t)}\right) \varphi^2\left(\frac{4}{R} |\Psi(\delta_k x) - \delta_k \tilde{x}|\right) dx, \end{aligned}$$

where $\widehat{\nabla} u_k := (\nabla' u_k - (\partial_n u_k) \nabla' f_k, \partial_n u_k)$. For simplicity of the presentation, we assume that $\Psi(x) = x$, that is, $\partial\Omega \cap B_R$ is flat. In addition, we focus on the convergence of the first term, since the other terms can be estimated in the same manner. Namely, we prove that

$$\begin{aligned} & \int_{\delta_k^{-1} \Omega_R} |\nabla u_k|^2 e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} \varphi^2\left(\frac{4\delta_k|x-\tilde{x}|}{R}\right) dx - \int_{\mathbf{R}_+^n} |\nabla \bar{u}|^2 e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} dx \\ &= \int_{\delta_k^{-1} \Omega_R} (|\nabla u_k|^2 - |\nabla \bar{u}|^2) e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} \varphi^2 dx \\ & \quad + \int_{\delta_k^{-1} \Omega_R} |\nabla \bar{u}|^2 e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} (\varphi^2 - 1) dx \\ & \quad - \int_{\mathbf{R}_+^n \setminus \delta_k^{-1} \Omega_R} |\nabla \bar{u}|^2 e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} dx =: J_1^k + J_2^k - J_3^k \rightarrow 0 \end{aligned} \tag{5.10}$$

as $k \rightarrow \infty$. By the Hölder inequality for the Lorentz spaces (see [49, Proposition 2.1]) and Lemma 5.5 (iv), we can see that $J_2^k \rightarrow 0$ and $J_3^k \rightarrow 0$.

Let us estimate J_1^k . For $R' > 0$, by dividing the domain of integration, we set

$$\begin{aligned} |J_1^k| &\leq \left(\int_{B_{R'}(\tilde{x})} + \int_{\mathbf{R}^n \setminus B_{R'}(\tilde{x})} \right) \left| |\nabla u_k|^2 - |\nabla \bar{u}|^2 \right| e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} \chi_{\delta_k^{-1} \Omega_R} dx \\ &=: J_{\text{in}}^k + J_{\text{out}}^k. \end{aligned}$$

We consider J_{out}^k . By computations similar to (3.2) together with Lemma 5.2, Remark 5.3 and Lemma 5.5 (iv), we see that

$$\begin{aligned}
J_{\text{out}}^k &\leq \sum_{m=1}^{\infty} \int_{(\mathbf{R}^n \setminus B_{R'}(\tilde{x})) \cap (B_m(\tilde{x}) \setminus \overline{B_{m-1}(\tilde{x})})} \left| |\nabla u_k|^2 - |\nabla \bar{u}|^2 \right| e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} \chi_{\delta_k^{-1}\Omega_R} dx \\
&\leq \sum_{m=1}^{\infty} e^{-\frac{(R')^2+(m-1)^2}{12(\tilde{t}-t)}} \int_{B_m(\tilde{x})} (|\nabla u_k| \chi_{\delta_k^{-1}\Omega_R} + |\nabla \bar{u}|)^2 e^{-\frac{|x-\tilde{x}|^2}{12(\tilde{t}-t)}} dx \\
&\leq C \sum_{m=1}^{\infty} (\tilde{t}-t)^{-\frac{p+1}{p-1}} e^{-\frac{(R')^2+(m-1)^2}{12(\tilde{t}-t)}} (\|\nabla u_k| \chi_{\delta_k^{-1}\Omega_R} + |\nabla \bar{u}|\|_{L^{q^*,\infty}(B_m(\tilde{x}))}^2) \\
&\leq C(M + M^p)^2 (\tilde{t}-t)^{-\frac{p+1}{p-1}} e^{-\frac{(R')^2}{12(\tilde{t}-t)}} \sum_{m=1}^{\infty} e^{-\frac{(m-1)^2}{4(\tilde{t}-t)}} \rightarrow 0
\end{aligned}$$

as $R' \rightarrow \infty$, where $\overline{B_0(\tilde{x})} := \emptyset$. Remark that the above convergence is uniformly for k . Therefore, for any $\varepsilon > 0$, there exists $R' > 0$ independent of k such that

$$|J_1^k| \leq J_{\text{in}}^k + \varepsilon \leq \int_{B_{R'}} \left| |\nabla u_k|^2 - |\nabla \bar{u}|^2 \right| \chi_{\delta_k^{-1}\Omega_R} dx + \varepsilon.$$

Then by Lemma 5.5 (i) and (iv), we obtain $|J_1^k| \rightarrow 0$, and so (5.10) also holds. The other terms can be handled in the same way. \square

We next estimate an integral concerning $|\bar{u}|^{p+1}$ by using $\overline{E}_{(\tilde{x}, \tilde{t})}$. More precisely, we prove the analog of Lemma 4.6 for the blow-up limit \bar{u} .

Lemma 5.8. *Assume (R2). Then there exists a constant $C > 0$ such that*

$$\begin{aligned}
&\int_{t'}^t (\tilde{t}-s)^{\frac{p+1}{p-1}} \int_{\mathbf{R}_+^n} |\bar{u}|^{p+1} K_{(\tilde{x}, \tilde{t})}(x, s) dx ds \\
&\leq C \left(\log \frac{\tilde{t}-t'}{\tilde{t}-t} \right)^{\frac{1}{2}} (\overline{E}_{(\tilde{x}, \tilde{t})}(t') - \overline{E}_{(\tilde{x}, \tilde{t})}(t) + C(\tilde{t}-t'))^{\frac{1}{2}} \\
&\quad + C_p (\overline{E}_{(\tilde{x}, \tilde{t})}(t') + C(\tilde{t}-t')) \log \frac{\tilde{t}-t'}{\tilde{t}-t} + Ch(\tilde{t}-t)
\end{aligned}$$

for $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t' < t < 0$ and $1 > \tilde{t} > t$. Here $C_p := 2(p+1)/(p-1)$.

Proof. Define $\tilde{\varphi} \in C^\infty([0, \infty))$ so that $\tilde{\varphi}' \leq 0$, $0 \leq \tilde{\varphi} \leq 1$, $\tilde{\varphi}(z) = 0$ for $z > 2$ and $\tilde{\varphi}(z) = 1$ for $0 \leq z < 1$. For $\tilde{R} > 1$, set $\tilde{\varphi}_{\tilde{R}}(z) := \tilde{\varphi}(z/\tilde{R})$. For $\tilde{x} \in \overline{\mathbf{R}_+^n}$, $-1 < t < 0$ and $1 > \tilde{t} > t$, we define

$$\begin{aligned}
E_k(t) &:= (\tilde{t}-t)^{\frac{p+1}{p-1} - \frac{n}{2}} \int_{\delta_k^{-1}\Phi(\Omega_R)} \left(\frac{|\widehat{\nabla} u_k(\xi, t)|^2}{2} - \frac{|u_k|^{p+1}}{p+1} + \frac{(\tilde{t}-t)^{-1}|u_k|^2}{2(p-1)} \right) \\
&\quad \times e^{-\frac{|\Psi_k(\xi) - \Psi_k(\tilde{\xi}_k)|^2}{4(\tilde{t}-t)}} \varphi^2 \left(\frac{4\delta_k}{\tilde{R}} |\Psi_k(\xi) - \Psi_k(\tilde{\xi}_k)| \right) \tilde{\varphi}_{\tilde{R}}^2(|\xi|) d\xi,
\end{aligned}$$

where $\Psi_k(\xi) := \delta_k^{-1}\Psi(\delta_k\xi)$, $\Phi_k(x) := \delta_k^{-1}\Phi(\delta_kx)$ and $\tilde{\xi}_k := \Phi_k(\tilde{x})$. We note that $\Psi_k(\tilde{\xi}_k) = \tilde{x}$ and that the right-hand side of this definition coincides with $E_{(\delta_k\tilde{x}, \delta_k^2\tilde{t})}(\delta_k^2t; \phi_{\delta_k\tilde{x}, R/4})$ if we replace $\tilde{\varphi}_{\tilde{R}}$ with 1.

By using the backward similarity variables

$$\eta := \frac{\xi - \tilde{\xi}_k}{(\tilde{t} - t)^{1/2}}, \quad \tau := -\log(\tilde{t} - t),$$

we define

$$\begin{aligned} w_k(\eta, \tau) &:= e^{-\frac{1}{p-1}\tau} u_k(\tilde{\xi}_k + e^{-\frac{1}{2}\tau}\eta, \tilde{t} - e^{-\tau}), \\ g_k(\eta', \tau) &:= e^{\frac{1}{2}\tau} f_k(\tilde{\xi}'_k + e^{-\frac{1}{2}\tau}\eta'). \end{aligned}$$

Then w_k satisfies

$$\begin{cases} (w_k)_\tau + \frac{1}{2}\eta \cdot \nabla w_k + \frac{1}{p-1}w_k - A_k w_k - |w_k|^{p-1}w_k = 0, \\ \eta \in \Omega_k(\tau) := e^{\tau/2}(\Phi_k(\delta_k^{-1}\Omega_R) - \tilde{x}), \quad \tau \in (-\log(\tilde{t} + 1/2), -\log\tilde{t}), \\ w_k = 0, \quad \eta \in e^{\tau/2}(\Phi_k(\delta_k^{-1}(\partial\Omega \cap B_R)) - \tilde{\xi}_k), \end{cases}$$

where

$$A_k w_k := \Delta w_k - 2\nabla'(\partial_n w_k) \cdot \nabla' g_k + (\partial_n^2 w_k)|\nabla' g_k|^2 - (\partial_n w_k)\Delta' g_k.$$

Setting $\widehat{\nabla} w_k := (\nabla' w_k - (\partial_n w_k)\nabla' g_k, \partial_n w_k)$ and

$$\begin{aligned} \mathcal{E}_k(\tau) &:= \int_{\Omega_k(\tau)} \left(\frac{|\widehat{\nabla} w_k|^2}{2} - \frac{|w_k|^{p+1}}{p+1} + \frac{w_k^2}{2(p-1)} \right) \rho_k \psi_k^2 d\eta, \\ \rho_k = \rho_k(\eta, \tau) &:= \exp \left(-\frac{1}{4}(|\eta|^2 + 2(g_k(\eta', \tau) - g_k(0, \tau))\eta_n \right. \\ &\quad \left. + (g_k(\eta', \tau) - g_k(0, \tau))^2) \right), \\ \psi_k = \psi_k(\eta, \tau) &:= \varphi \left(\frac{4|\Psi_k(\tilde{\xi}_k + e^{-\tau/2}\eta) - \Psi_k(\tilde{\xi}_k)|}{R} \right) \tilde{\varphi}_{\tilde{R}}^2(|\tilde{\xi}_k + e^{-\frac{\tau}{2}}\eta|), \end{aligned}$$

we can check that $E_k(t) = \mathcal{E}_k(\tau)$ with $\tau = -\log(\tilde{t} - t)$. In addition, we can observe that the above situation is almost the same as in Section 3, except for the appearance of $\varphi_{\tilde{R}}$.

By the same computations as in Section 3 and by $|\tilde{\varphi}'_{\tilde{R}}| + |\tilde{\varphi}''_{\tilde{R}}| \leq C(\tilde{R}^{-1} + \tilde{R}^{-2}) \leq C$ for $\tilde{R} > 1$, we obtain

$$\mathcal{E}_k(\tau) + \frac{1}{2} \int_{\tau'}^{\tau} \int_{\Omega_k(\tau)} (w_k)_\tau \rho_k \psi_k^2 d\eta d\sigma \leq \mathcal{E}_k(\tau') + C(\tilde{t} - t'),$$

where $C > 0$ is independent of k and \tilde{R} . Then the same computations as in Lemma 4.6 yield

$$\begin{aligned} &\int_{t'}^t (\tilde{t} - s)^{\frac{p+1}{p-1} - \frac{n}{2}} \int_{\delta_k^{-1}\Phi(\Omega_R)} |u_k|^{p+1} \exp \left(-\frac{|\delta_k^{-1}\Psi(\delta_k x) - \tilde{x}|^2}{4(\tilde{t} - s)} \right) \\ &\quad \times \varphi^2 \left(\frac{4}{R} |\Psi(\delta_k x) - \delta_k \tilde{x}| \right) \tilde{\varphi}_{\tilde{R}}^2(|x|) dx ds \\ &\leq C \left(\log \frac{\tilde{t} - t'}{\tilde{t} - t} \right)^{\frac{1}{2}} (E_k(t') - E_k(t) + C(\tilde{t} - t'))^{\frac{1}{2}} \\ &\quad + C_p (E_k(t') + C(\tilde{t} - t')) \log \frac{\tilde{t} - t'}{\tilde{t} - t} + Ch(\tilde{t} - t'), \end{aligned}$$

where $C_p := 2(p+1)/(p-1)$ and $C > 0$ is independent of k and \tilde{R} . By Lemma 5.5 (ii), the same argument as in Lemma 5.7 and letting $k \rightarrow \infty$, we have

$$\begin{aligned} & \int_{t'}^t (\tilde{t} - s)^{\frac{p+1}{p-1}} \int_{\mathbf{R}_+^n} |\bar{u}|^{p+1} K_{(\tilde{x}, \tilde{t})}(x, s) \tilde{\varphi}_{\tilde{R}}^2(|x|) dx ds \\ & \leq C \left(\log \frac{\tilde{t} - t'}{\tilde{t} - t} \right)^{\frac{1}{2}} (\bar{E}_{\tilde{\varphi}_{\tilde{R}}}(t') - \bar{E}_{\tilde{\varphi}_{\tilde{R}}}(t) + C(\tilde{t} - t'))^{\frac{1}{2}} \\ & \quad + C_p (\bar{E}_{\tilde{\varphi}_{\tilde{R}}}(t') + C(\tilde{t} - t')) \log \frac{\tilde{t} - t'}{\tilde{t} - t} + Ch(\tilde{t} - t'), \end{aligned}$$

where

$$\bar{E}_{\tilde{\varphi}_{\tilde{R}}}(t) := (\tilde{t} - t)^{\frac{p+1}{p-1}} \int_{\mathbf{R}_+^n} \left(\frac{|\nabla \bar{u}|^2}{2} - \frac{|\bar{u}|^{p+1}}{p+1} + \frac{\bar{u}^2}{2(p-1)(\tilde{t} - t)} \right) K_{(\tilde{x}, \tilde{t})} \tilde{\varphi}_{\tilde{R}}^2 dx.$$

Letting $\tilde{R} \rightarrow \infty$ with the aid of the monotone convergence theorem to the left-hand side and the Lebesgue dominated convergence theorem to the right-hand side, we obtain the desired inequality. \square

Remark 5.9. As stated in Remark 4.2, the ε -regularity (Theorem 4.1) is also valid for \bar{u} , since \bar{u} satisfies the analog of Lemma 4.6.

We next show a monotonicity estimate of $\bar{E}_{(0,0)}$.

Lemma 5.10. *Assume (R2). Then there exists $C > 0$ such that*

$$\begin{aligned} \bar{E}_{(0,0)}(t') - \bar{E}_{(0,0)}(t) & \geq C^{-1} \frac{(-t)^{4/(p-1)}}{(-t')^{2p/(p-1)}} \\ & \quad \times \left(\int_{t'}^t \int_{\mathbf{R}_+^n} \left(\frac{\bar{u}}{p-1} + \frac{x \cdot \nabla \bar{u}}{2} + s \bar{u}_s \right) K_{(0,0)}(x, s) \zeta(x, s) dx ds \right)^2 \end{aligned}$$

for any $-1 < t' < t < 0$ and $\zeta \in C_0^\infty(\mathbf{R}_+^n \times (-1, 0))$ with $|\zeta| \leq 1$.

Proof. Let $-1 < t' < t < 0$ and let $\zeta \in C_0^\infty(\mathbf{R}_+^n \times (-1, 0))$ satisfy $|\zeta| \leq 1$. We take k large as in the proof of Lemma 5.7. From (3.16) and the change of variables, it follows that

$$\begin{aligned} (5.11) \quad & E_{(0,0)}(\delta_k^2 t; \phi_{0, R/4}) + \frac{1}{2} \int_{t'}^t (-s)^{\frac{2}{p-1} - \frac{n}{2} - 1} \\ & \quad \times \int_{\delta_k^{-1} \Phi(\Omega_R)} \left| \frac{u_k}{p-1} + \frac{x \cdot \nabla u_k}{2} + s(u_k)_s \right|^2 \\ & \quad \times \exp \left(-\frac{|\delta_k^{-1} \Psi(\delta_k x)|^2}{4(-s)} \right) \varphi^2 \left(\frac{4}{R} |\Psi(\delta_k x)| \right) dx ds \\ & \leq E_{(0,0)}(\delta_k^2 t'; \phi_{0, R/4}) + C' \delta_k^2 (-t') \end{aligned}$$

for $-(1/2)\delta_k^{-2} < t' < t < 0$. For simplicity of the presentation, we assume $\Psi(x) = x$. Then the second term in the left-hand side of (5.11) can be estimated as

$$\begin{aligned} \mathcal{I} &:= \int_{t'}^t (-s)^{\frac{2}{p-1}-\frac{n}{2}-1} \int_{\delta_k^{-1}\Omega_R} \left| \frac{u_k}{p-1} + \frac{x \cdot \nabla u_k}{2} + s(u_k)_s \right|^2 \\ &\quad \times \exp\left(-\frac{|x|^2}{4(-s)}\right) \varphi^2\left(\frac{4\delta_k}{R}|x|\right) dx ds \\ &\geq \iint_{\delta_k^{-1}\Omega_R \times (t', t)} |F(u_k)|^2 \zeta^2(x, s) d\mu_k(x, s). \end{aligned}$$

Here $F(u_k)$ and μ_k are defined by

$$\begin{aligned} F(u_k) &:= \frac{u_k}{p-1} + \frac{x \cdot \nabla u_k}{2} + s(u_k)_s, \\ d\mu_k(x, s) &:= (-s)^{\frac{2}{p-1}-1} K_{(0,0)}(x, s) \varphi^2\left(\frac{4\delta_k}{R}|x|\right) dx ds. \end{aligned}$$

We compute that

$$\mu_k(\delta_k^{-1}\Omega_R \times (t', t)) \leq \int_{t'}^t (-s)^{\frac{2}{p-1}-1} \int_{\mathbf{R}^n} K_{(0,0)}(x, s) dx ds \leq C(-t')^{\frac{2}{p-1}},$$

where C depends only on n and p and is independent of k . Then by Jensen's inequality, we see that

$$\begin{aligned} \mathcal{I} &\geq \frac{1}{\mu_k(\delta_k^{-1}\Omega_R \times (t', t))} \left(\iint_{\delta_k^{-1}\Omega_R \times (t', t)} F(u_k) \zeta d\mu_k(x, s) \right)^2 \\ &\geq C^{-1}(-t')^{-\frac{2}{p-1}} \left(\iint_{\delta_k^{-1}\Omega_R \times (t', t)} F(u_k) \zeta d\mu_k(x, s) \right)^2. \end{aligned}$$

From this and (5.11), it follows that

$$\begin{aligned} E_{(0,0)}(\delta_k^2 t; \phi_{0,R/4}) + C^{-1}(-t')^{-\frac{2}{p-1}} \left(\iint_{\delta_k^{-1}\Omega_R \times (t', t)} F(u_k) \zeta d\mu_k \right)^2 \\ \leq E_{(0,0)}(\delta_k^2 t'; \phi_{0,R/4}) + C' \delta_k^2 (-t'). \end{aligned}$$

From Remark 5.6, Lemma 5.5 (i) and (ii) and computations similar to the derivation of (5.10), it follows that

$$\iint_{\delta_k^{-1}\Omega_R \times (t', t)} F(u_k) \zeta d\mu_k(x, s) \rightarrow \int_{t'}^t (-s)^{\frac{2}{p-1}-1} \int_{\mathbf{R}_+^n} F(\bar{u}) K_{(0,0)} \zeta dx ds$$

as $k \rightarrow \infty$. This together with Lemma 5.7 and letting $k \rightarrow \infty$ implies that

$$\begin{aligned} \overline{E}_{(0,0)}(t') - \overline{E}_{(0,0)}(t) \\ \geq C^{-1}(-t')^{-\frac{2}{p-1}} \left(\int_{t'}^t (-s)^{\frac{2}{p-1}-1} \int_{\mathbf{R}_+^n} F(\bar{u}) K_{(0,0)} \zeta dx ds \right)^2 \\ \geq C^{-1}(-t')^{-\frac{2p}{p-1}} (-t)^{\frac{4}{p-1}} \left(\int_{t'}^t \int_{\mathbf{R}_+^n} F(\bar{u}) K_{(0,0)} \zeta dx ds \right)^2. \end{aligned}$$

Then the desired inequality follows. \square

The monotonicity estimate gives the following equality based on the argument of [85, Theorem 8.1].

Lemma 5.11. *Assume (R2). Then,*

$$\frac{\bar{u}(x, t)}{p-1} + \frac{x \cdot \nabla \bar{u}}{2} + t\bar{u}_t = 0 \quad \text{a.e. in } \mathbf{R}_+^n \times (-1, 0).$$

Proof. For $0 < r < 1/2$, we set

$$\Phi(r) := \int_{-4r^2}^{-r^2} \bar{E}_{(0,0)}(t) \frac{dt}{-t}.$$

We note that the well-definedness of Φ is guaranteed by (5.9). By Lemmas 5.7 and 3.1, the Lebesgue dominated convergence theorem and the change of variables, we have

$$\Phi(r) = \lim_{k \rightarrow \infty} \int_{-4r^2}^{-r^2} E_{(0,0)}(\delta_k^2 t) \frac{dt}{-t} = \lim_{k \rightarrow \infty} \int_{-4\delta_k^2 r^2}^{-\delta_k^2 r^2} E_{(0,0)}(s) \frac{ds}{-s}.$$

We claim that $\Phi(r)$ is independent of r . To show this, we set

$$H(\tilde{r}) := \int_{-4\tilde{r}^2}^{-\tilde{r}^2} E_{(0,0)}(s) \frac{ds}{-s} = \int_{-4}^{-1} E_{(0,0)}(\tilde{r}^2 \theta) \frac{d\theta}{-\theta}$$

for $0 < \tilde{r} < 1/2$ and check that $\lim_{\tilde{r} \rightarrow 0} H(\tilde{r})$ exists. Let $\varepsilon > 0$. Since Lemma 3.1 yields the boundedness of H , we see that $H_\varepsilon := \inf_{0 < \tilde{r} < \varepsilon} H(\tilde{r})$ is finite. Then there exists $0 < \tilde{r}_\varepsilon < \varepsilon$ such that $H(\tilde{r}_\varepsilon) \leq H_\varepsilon + \varepsilon$. From Proposition 3.2 and the negativity of θ , it follows that

$$\begin{aligned} \limsup_{\tilde{r} \rightarrow 0} H(\tilde{r}) &\leq \limsup_{\tilde{r} \rightarrow 0} \int_{-4}^{-1} (E_{(0,0)}(\tilde{r}_\varepsilon^2 \theta) + C'(M + M^p)^2 (-\tilde{r}_\varepsilon^2 \theta)) \frac{d\theta}{-\theta} \\ &= H(\tilde{r}_\varepsilon) + 3C'(M + M^p)^2 \tilde{r}_\varepsilon^2 \leq H_\varepsilon + \varepsilon + C(M + M^p)^2 \varepsilon^2. \end{aligned}$$

Thus, letting $\varepsilon \rightarrow 0$ gives $\limsup_{\tilde{r} \rightarrow 0} H(\tilde{r}) \leq \liminf_{\tilde{r} \rightarrow 0} H(\tilde{r})$. Hence the limit of $H(\tilde{r})$ exists, and so $\Phi(r) = \lim_{\tilde{r} \rightarrow 0} H(\tilde{r})$. Then the claim follows.

From the claim and Lemma 5.10, it follows that

$$\begin{aligned} 0 &= \int_{r_2}^{r_1} \Phi'(s) ds = \int_{r_2}^{r_1} \frac{2}{s} (\bar{E}_{(0,0)}(-4s^2) - \bar{E}_{(0,0)}(-s^2)) ds \\ &\geq C^{-1} \int_{r_2}^{r_1} \frac{9-5p}{s^{\frac{9-5p}{p-1}}} \left(\int_{-4s^2}^{-s^2} \int_{\mathbf{R}_+^n} \left(\frac{\bar{u}}{p-1} + \frac{x \cdot \nabla \bar{u}}{2} + t\bar{u}_t \right) K_{(0,0)} \zeta dx dt \right)^2 ds \end{aligned}$$

for any $0 < r_1 < r_2 < 1/2$ and $\zeta \in C_0^\infty(\mathbf{R}_+^n \times (-1, 0))$ with $|\zeta| \leq 1$. Thus,

$$\int_{-4s^2}^{-s^2} \int_{\mathbf{R}_+^n} \left(\frac{\bar{u}}{p-1} + \frac{x \cdot \nabla \bar{u}}{2} + t\bar{u}_t \right) K_{(0,0)}(x, t) \zeta(x, t) dx dt = 0$$

for a.e. $s \in (0, 1/2)$. Hence by the fundamental lemma of the calculus of variations, the lemma follows. \square

Finally in this subsection, we prove a partial regularity result for \bar{u} by using the ε -regularity. For $t \in (-1, 0)$, define the singular set of $\bar{u}(\cdot, t)$ by

$$\Sigma(t) := \{x \in \mathbf{R}_+^n; \bar{u}(\cdot, t) \notin L^\infty(B_\rho^+(x)) \text{ for all } \rho > 0\}.$$

By modifying the argument of Wang [90, Lemma 3.3], we show that $\Sigma(t)$ consists of finitely many points.

Lemma 5.12. *Assume (R2). Then the singular set $\Sigma(t)$ consists of finitely many points for each $t \in (-1/4, 0)$. More precisely, there exists a constant $C > 0$ depending on R such that the cardinality $\text{card}(\Sigma(t))$ of $\Sigma(t)$ satisfies*

$$\text{card}(\Sigma(t)) \leq C(M + M^p)^{q_c} \varepsilon_0^{-\frac{q_c}{2}}$$

for each $t \in (-1/4, 0)$. Here ε_0 is given in Theorem 4.1 (see also Remark 5.9) and C is independent of t and M .

Proof. Let $t_0 \in (0, -1/4)$. Since $\text{card}(\Sigma(t_0)) = 0$ if $\Sigma(t_0) = \emptyset$, it suffices to consider the case $\Sigma(t_0) \neq \emptyset$. Let $(x_0, t_0) \in \Sigma(t_0)$. We examine the L^{q_c} -norm of \bar{u} near (x_0, t_0) . The contraposition of the ε -regularity (see Theorem 4.1 and Remark 5.9), it follows that

$$\varepsilon_0 < (\rho/2)^{\frac{4}{p-1}-n} \iint_{Q_{\rho/2}^+(x_0, t_0)} (|\nabla \bar{u}|^2 + |\bar{u}|^{p+1}) dx dt \quad \text{for } 0 < \rho \leq 2\delta_0.$$

Lemma A.4 with translation gives

$$\iint_{Q_{\rho/2}^+(x_0, t_0)} |\nabla \bar{u}|^2 dx dt \leq C \rho^{n-\frac{4}{p-1}-\frac{4}{q_c}} (1 + M^{p-1})^2 \|\bar{u}\|_{L^{q_c}(Q_{\rho/2}^+(x_0, t_0))}^2,$$

where $C > 0$ is independent of ρ . The Hölder inequality also gives

$$\begin{aligned} & \iint_{Q_{\rho/2}^+(x_0, t_0)} |\bar{u}|^{p+1} dx dt \\ & \leq C \rho^{(n+2)(1-\frac{p+1}{q_c})} \left(\iint_{Q_{\rho/2}^+(x_0, t_0)} |\bar{u}|^{q_c} dx dt \right)^{\frac{p-1}{q_c}} \|\bar{u}\|_{L^{q_c}(Q_{\rho/2}^+(x_0, t_0))}^2 \\ & \leq C \rho^{(n+2)(1-\frac{p+1}{q_c})+\frac{2(p-1)}{q_c}} M^{p-1} \|\bar{u}\|_{L^{q_c}(Q_{\rho/2}^+(x_0, t_0))}^2. \end{aligned}$$

These estimates imply that

$$\varepsilon_0 \leq C(1 + M^{p-1})^2 \rho^{-\frac{4}{q_c}} \|\bar{u}\|_{L^{q_c}(Q_{\rho/2}^+(x_0, t_0))}^2.$$

Hence there exists a constant $C > 0$ depending on R and independent of ρ , t_0 and M such that

$$\varepsilon_0^{\frac{q_c}{2}} \leq C(1 + M^{p-1})^{q_c} \rho^{-2} \int_{t_0-\rho^2}^{t_0} \int_{B_{\rho/2}^+(x_0)} |\bar{u}|^{q_c} dx dt \quad \text{for } 0 < \rho \leq 2\delta_0.$$

Let $S \subset \Sigma(t_0)$ be any finite set. We write $S = \{x_1, \dots, x_N\}$ and choose $0 < \rho_0 \leq \delta_0$ ($< 1/2$) such that $\{B_{\rho_0}^+(x_i)\}_{1 \leq i \leq N}$ is pairwise disjoint. Then we have

$$\varepsilon_0^{\frac{q_c}{2}} \leq C(1 + M^{p-1})^{q_c} \rho_0^{-2} \int_{t_0-\rho_0^2}^{t_0} \int_{B_{\rho_0}^+(x_i)} |\bar{u}|^{q_c} dx dt$$

for each $1 \leq i \leq N$, and so we also have

$$\begin{aligned} N \varepsilon_0^{\frac{q_c}{2}} & \leq C(1 + M^{p-1})^{q_c} \rho_0^{-2} \int_{t_0-\rho_0^2}^{t_0} \left(\sum_{i=1}^N \int_{B_{\rho_0}^+(x_i)} |\bar{u}|^{q_c} dx \right) dt \\ & \leq C(1 + M^{p-1})^{q_c} \rho_0^{-2} \int_{t_0-\rho_0^2}^{t_0} \int_{\mathbf{R}_+^n} |\bar{u}|^{q_c} dx dt \leq C(M + M^p)^{q_c}, \end{aligned}$$

where $C > 0$ is independent of ρ_0 , t_0 , M and N . Hence $\text{card}(S) \leq C(M + M^p)^{q_c} \varepsilon_0^{-q_c/2}$. Remark that the right-hand side of this inequality does not depend on

the choice of S . Therefore we can conclude that $\text{card}(\Sigma(t_0)) \leq C(M + MP)^{q_c} \varepsilon_0^{-q_c/2}$. The proof is complete. \square

Remark 5.13. In the case (R1), Lemma 5.5 also holds with \mathcal{B}_ρ and \mathbf{R}_+^n replaced by B_ρ and \mathbf{R}^n , respectively. In particular, the blow-up limit \bar{u} exists. The analogs to the ε -regularity (Theorem 4.1), the equality in Lemma 5.11 and the partial regularity (Lemma 5.12) also hold for \bar{u} in the case (R1).

5.3. Proof of localized statement. We are now in a position to prove Theorem 5.1.

Proof of Theorem 5.1. We show this theorem by contradiction under the condition supposed in Subsection 5.1. We focus on the case (R2), since (R1) is easier. From the lower bound (5.3) together with (3.3), (3.4) and (5.4), it follows that

$$\begin{aligned} \varepsilon_0 &< \delta_k^{\frac{4}{p-1}-n} \int_{-\delta_k^2}^0 \int_{\Phi(\Omega_{\delta_k})} (|\widehat{\nabla} \hat{u}|^2 + |\hat{u}|^{p+1}) d\xi dt \\ &= \int_{-1}^0 \int_{\delta_k^{-1} \Phi(\Omega_{\delta_k})} (|\widehat{\nabla} u_k|^2 + |u_k|^{p+1}) dx dt. \end{aligned}$$

Using $\|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})} \leq 1/2$ gives $\delta_k^{-1} \Phi(\Omega_{\delta_k}) \subset B_{\sqrt{2}}^+$. Hence by (5.6) and Lemma 5.5, we see that

$$(5.12) \quad \int_{-1}^0 \int_{B_{\sqrt{2}}^+} (|\nabla \bar{u}|^2 + |\bar{u}|^{p+1}) dx dt > \varepsilon_0.$$

We will show that $\bar{u} \equiv 0$ a.e. in $\mathbf{R}_+^n \times (-1, 0)$, contrary to (5.12).

By the argument of [53, Lemma 8.5.3] (see also [85, 90]), we see that Lemma 5.11 implies the following two scenarios:

- (i) \bar{u} is time-independent, and thus homogeneous.
- (ii) \bar{u} is time-dependent, and thus self-similar.

We examine these scenarios and deduce a contradiction to (5.12).

- (i) In this case, \bar{u} can be written as

$$\bar{u}(x, t) = U \left(\frac{x}{|x|} \right) |x|^{-\frac{2}{p-1}}$$

for some function U defined on the hemisphere \mathbf{S}_+^{n-1} . By Lemma 5.5 (iii), we have

$$\int_{\mathbf{R}_+^n} |\bar{u}|^{q_c} dx = \int_0^\infty r^{-1} dr \int_{\mathbf{S}_+^n} |U|^{q_c} dS \leq M^{q_c} < \infty.$$

Therefore U must vanish a.e. in \mathbf{S}_+^n . This contradicts (5.12).

- (ii) In this case, there exists a profile function $\bar{U} \in L^{q_c}(\mathbf{R}_+^n)$ such that

$$u(x, t) = (-t)^{-\frac{1}{p-1}} \bar{U}(z), \quad z := \frac{x}{\sqrt{-t}},$$

where \bar{U} satisfies the equation

$$(5.13) \quad \Delta \bar{U} - \frac{1}{2} z \cdot \nabla \bar{U} - \frac{1}{p-1} \bar{U} + |\bar{U}|^{p-1} \bar{U} = 0, \quad z \in \mathbf{R}_+^n.$$

We claim that there exist constants $\tilde{R}, C > 1$ satisfying

$$(5.14) \quad \begin{cases} \bar{u}(\cdot, 0) = 0 & \text{a.e. in } \mathbf{R}_+^n, \\ |\bar{u}| \leq C & \text{a.e. in } (\mathbf{R}_+^n \setminus B_{\tilde{R}}) \times [-1/9, 0]. \end{cases}$$

Recall from the proof of Lemma 5.5 that $u_k \rightarrow \bar{u}$ in $C([-1, 0]; W^{1,r}(\mathcal{B}_\rho))$ for any $\rho > 0$ with some $1 < r < \min\{2, q_c/p\}$ as $k \rightarrow \infty$. Then, for $x_0 \in \mathbf{R}^n$ and $\delta > 0$, we have

$$\int_{B_1^+(x_0)} |\bar{u}(x, 0)|^r dx \leq 2^{r-1} \delta + 2^{r-1} \int_{B_1^+(x_0)} |u_k(x, 0)|^r dx$$

for sufficiently large k . By returning the original variables, we also have

$$\begin{aligned} \int_{B_1^+(x_0)} |u_k(x, 0)|^r dx &= \delta_k^{\frac{2r}{p-1}-n} \int_{\Psi(B_{\delta_k}^+(\delta_k x_0))} |u(x, 0)|^r dx \\ &\leq \delta_k^{\frac{2r}{p-1}-n} \int_{\Psi(B_{(1+|x_0|)\delta_k}^+(0))} |u(x, 0)|^r dx \\ &\leq C(1 + |x_0|)^{n - \frac{2r}{p-1}} \|u(\cdot, 0)\|_{L^{q_c}(\Psi(B_{(1+|x_0|)\delta_k}^+(0)))}^r \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$, where u belongs to $C_{\text{weak}}([-1/4, 0]; L^{q_c}(\Omega_{R/2}))$ in our situation, see (5.2). Hence we obtain the former equality in (5.14).

We prove the latter inequality in (5.14). Let $\tilde{\varepsilon} > 0$. We claim that there exists a constant $\tilde{R} > 1$ depending on $\tilde{\varepsilon}$ and M satisfying

$$(5.15) \quad \iint_{Q_{1/2}^+(x_1, 0)} (|\nabla \bar{u}|^2 + |\bar{u}|^{p+1}) dx dt \leq \tilde{\varepsilon}$$

for any $x_1 \in \mathbf{R}_+^n \setminus B_{\tilde{R}}^+(0)$, where $Q_{1/2}^+(x_1, 0) = B_{1/2}^+(x_1) \times (-1/4, 0)$. Indeed, for $\tilde{\varepsilon}' > 0$, we can choose $\tilde{R}' > 1$ by Lemma 5.5 (iii) so large that

$$\int_{-1}^0 \int_{\mathbf{R}_+^n \setminus B_{\tilde{R}'}(0)} |\bar{u}|^{q_c} dx dt \leq \tilde{\varepsilon}'.$$

From the Hölder inequality, it follows that

$$\iint_{Q_1^+(x_1, 0)} |\bar{u}|^{p+1} dx dt \leq C \iint_{Q_1^+(x_1, 0)} |\bar{u}|^{q_c} dx dt \leq C \tilde{\varepsilon}'$$

for any $x_1 \in \mathbf{R}_+^n \setminus B_{\tilde{R}'}(0)$, and so Lemma A.4 with $\rho = 1$ yields

$$\begin{aligned} \iint_{Q_{1/2}^+(x_1, 0)} |\nabla \bar{u}|^2 dx dt &\leq C(1 + M^{p-1})^2 \left(\iint_{Q_1^+(x_1, 0)} |\bar{u}|^{q_c} dx dt \right)^{\frac{2}{q_c}} \\ &\leq C(1 + M^{p-1})^2 \tilde{\varepsilon}^{\frac{2}{q_c}}. \end{aligned}$$

Thus choosing $\tilde{\varepsilon}'$ small gives (5.15), and so the claim follows. Hence by translation and Theorem 4.1 (see also Remark 5.9), we see that $|\bar{u}| \leq C\delta_0^{-2/(p-1)}$ in $B_{\delta_0}^+(x_1) \times [-1/9, 0]$ for any $x_1 \in \mathbf{R}_+^n \setminus B_{\tilde{R}}^+(0)$, where C and δ_0 are constants independent of x_1 . This proves the latter inequality in (5.14).

From (5.14) and the backward uniqueness theorem [31, Theorem 5.1], it follows that

$$\bar{u} \equiv 0 \quad \text{on } (\mathbf{R}_+^n \setminus B_{\tilde{R}}) \times [-1/9, 0].$$

In particular, $\bar{u}(x, -1/9) = 9^{-1/(p-1)}\bar{U}(3x) = 0$ for $x \in \mathbf{R}_+^n \setminus B_R^+$. Since \bar{U} satisfies (5.13) and is smooth except for a finite set by Lemma 5.12, the unique continuation theorem for elliptic equations in a connected set (see [4] for instance) concludes that $\bar{U} \equiv 0$ in \mathbf{R}_+^n . This contradicts (5.12), and hence the proof of the localized statement is complete. \square

5.4. Completion of proof. Finally, we prove Theorem 1.1.

Proof of Theorem 1.1. If Ω is bounded, then Theorem 1.1 immediately follows from Theorem 5.1. In what follows, we consider the case where Ω is unbounded. To obtain a contradiction, suppose that

$$\sup_{0 < t < T} \|u(\cdot, t)\|_{L^{q_c}(\Omega)} \leq M$$

for some $M > 0$. Let $\varepsilon > 0$ and $a \in \Omega$. Then by the same argument as in the derivation of (5.15), there exists a constant $\tilde{R} > 0$ depending on ε , a and M satisfying

$$\int_{T/2}^T \int_{\Omega_{1/2}(\tilde{x})} (|u|^{p+1} + |\nabla u|^2) dx dt \leq \varepsilon$$

for any $\tilde{x} \in \Omega \setminus B_{\tilde{R}}(a)$. Therefore, in the same way as in the proof of (5.14), we see that u is bounded on $(\Omega \setminus B_{\tilde{R}'}(a)) \times (T/3, T)$ for some $\tilde{R}' > 0$. This implies that there exists at least one blow-up point $a' \in \overline{\Omega_{\tilde{R}'}}(a)$. Hence Theorem 5.1 shows that

$$\limsup_{t \rightarrow T} \|u(\cdot, t)\|_{L^{q_c}(\Omega_r(a'))} = \infty$$

for any $r > 0$, a contradiction. The proof of Theorem 1.1 is complete. \square

APPENDIX A. REGULARITY ESTIMATES

We give parabolic regularity estimates for solutions of (2.1) and a gradient estimate for the blow-up limit obtained in Lemma 5.5. Let $n \geq 3$, $p > p_S$ and Ω be any smooth domain in \mathbf{R}^n with $0 \in \overline{\Omega}$. Fix $R > 0$ such that either (R1) or (R2) holds. Let u satisfy (2.1) and (2.2). Define \hat{u} by (3.3). For $\rho > 0$, we take $0 < \delta' < 1/2$ such that $\Psi(B_{3\rho\delta'}^+) \subset \Omega_{R/2}$ and $\Psi(B_{3\rho\delta'}) \subset B_{R/2}$. We first give parabolic regularity estimates. Remark that we mainly focus on the case (R2), since (R1) is easier.

Lemma A.1. *Assume (R2). Let $1 \leq l < \infty$ and $1 \leq r \leq q_c/p$. Then there exists $C > 0$ depending on R and ρ such that*

$$\|\hat{u}_t\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} + \|\nabla^2 \hat{u}\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \leq C \delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{p}{q_c})} (M + M^p)$$

for any $0 < \delta < \delta'$, where C is independent of δ .

Proof. By (3.3), we have

$$|\nabla^2 \hat{u}(x, t)| \leq C(|(\nabla u)(\Psi(x), t)| + |(\nabla^2 u)(\Psi(x), t)|),$$

where $C > 0$ depends on $\|\nabla f\|_{L^\infty(\mathbf{R}^{n-1})}$ and $\|\nabla^2 f\|_{L^\infty(\mathbf{R}^{n-1})}$. Remark that the choice of δ_R guarantees $\Psi(B_{\rho\delta}^+) \subset \Omega_{3R/4}$, and so Proposition 2.1 is applicable in

$\Psi(B_{\delta R}^+)$. Then by $r \leq q_c/p < q_*$, the Hölder inequality in the Lorentz spaces (see for instance [49, Proposition 2.1]) and Proposition 2.1, we see that

$$\begin{aligned}
& \|\nabla u(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \\
\text{(A.1)} \quad & \leq C\delta^{n(\frac{1}{r} - \frac{1}{q_*})} \|\nabla u(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^{q_*, \infty}(B_{\rho\delta}^+))} \\
& \leq C\delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{1}{q_*})} (M + M^p) \leq C\delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{p}{q_c})} (M + M^p).
\end{aligned}$$

We estimate $\|\nabla^2 u_1(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))}$. Let $\phi \in C_0^\infty(\mathbf{R}^n)$ satisfy $0 \leq \phi \leq 1$ in \mathbf{R}^n , $\phi = 0$ in $\mathbf{R}^n \setminus B_{3\rho}$ and $\phi = 1$ in $B_{2\rho}$. Set $\tilde{\phi}(x) := \phi(\delta^{-1}\Phi(x))$ and $v(x, t) := u(x, t)\tilde{\phi}(x)$. Then by the same computation as in the derivation of (2.3), we have

$$\begin{aligned}
u(x, t) &= \int_{\Omega} G_{\Omega}(x, y, t + 2\delta^2) \tilde{\phi}(y) u(y, -2\delta^2) dy \\
&+ \int_{-2\delta^2}^t \int_{\Omega} G_{\Omega}(x, y, t - s) \tilde{\phi} |u|^{p-1} u dy ds \\
&- \int_{-2\delta^2}^t \int_{\Omega} G_{\Omega}(x, y, t - s) (2\nabla \tilde{\phi} \cdot \nabla u + u \Delta \tilde{\phi}) dy ds
\end{aligned}$$

for $x \in \Psi(B_{\rho\delta}^+)$ and $-2\delta^2 < t < 0$, and so

$$\begin{aligned}
|\nabla^2 u| &\leq C \int_{\mathbf{R}^n} K_2(x - y, t + 2\delta^2) |u(y, -2\delta^2)| \chi_{\Omega_R \cap \Psi(B_{3\rho\delta})}(y) dy \\
&+ C \int_{-2\delta^2}^t \int_{\mathbf{R}^n} K_2(x - y, t - s) |u(y, s)|^p \chi_{\Omega_R \cap \Psi(B_{3\rho\delta})} dy ds \\
&+ C \int_{-2\delta^2}^t \int_{\mathbf{R}^n} K_2\left(\frac{|u|}{\delta^2} + \frac{|\nabla u|}{\delta}\right) \chi_{\Omega_R \cap \Psi(\overline{B_{3\rho\delta}} \setminus B_{2\rho\delta})} dy ds \\
&=: CV_1(x, t) + CV_2(x, t) + CV_3(x, t)
\end{aligned}$$

for $x \in \Psi(B_{\rho\delta}^+)$ and $-2\delta^2 < t < 0$, where K_2 is defined by (2.6).

First, we estimate V_1 . By $\|\nabla' f\|_{L^\infty(\mathbf{R}^{n-1})} \leq 1/2$ in (R2), we have

$$|\Psi(x) - \Psi(y)|^2 \geq |x' - y'|^2 + \frac{1}{2}(x_n - y_n)^2 - (f(x') - f(y'))^2 \geq \frac{1}{4}|x - y|^2.$$

This together with the change of variables shows that

$$\begin{aligned}
& V_1(\Psi(x), t) \\
\text{(A.2)} \quad &= \int_{\mathbf{R}^n} K_2(\Psi(x) - \Psi(y), t + 2\delta^2) |u(\Psi(y), -2\delta^2)| \chi_{\Phi(\Omega_R) \cap B_{3\rho\delta}} dy \\
&\leq \int_{\mathbf{R}^n} K_2((x - y)/2, t + 2\delta^2) |u(\Psi(y), -2\delta^2)| \chi_{\Phi(\Omega_R) \cap B_{3\rho\delta}} dy.
\end{aligned}$$

Then Young's inequality gives

$$\begin{aligned}
\|V_1(\Psi(\cdot), t)\|_{L^{q_c}(B_{\rho\delta}^+)} &\leq C(t + 2\delta^2)^{-1} \|u(\Psi(\cdot), -2\delta^2)\|_{L^{q_c}(\Phi(\Omega_R) \cap \Psi(B_{3\rho\delta}))} \\
&\leq C\delta^{-2} M
\end{aligned}$$

for $-\delta^2 < t < 0$. Therefore, the Hölder inequality shows that

$$\begin{aligned}
\|V_1(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \\
\text{(A.3)} \quad &\leq C\delta^{n(\frac{1}{r} - \frac{1}{q_c})} \|V_1(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^{q_c}(B_{\rho\delta}^+))} \leq C\delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{p}{q_c})} M.
\end{aligned}$$

Concerning the term V_2 , the Hölder inequality gives

$$\|V_2(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \leq C\delta^{n(\frac{1}{r} - \frac{p}{4c})} \|V_2(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^{\frac{4c}{p}}(B_{\rho\delta}^+))}.$$

In the same way as in (A.2), we have

$$V_2(\Psi(x), t) \leq \int_{-2\delta^2}^t \int_{\mathbf{R}^n} K_2((x-y)/2, t-s) |u(\Psi(y), s)|^p \chi_{\Phi(\Omega_R) \cap B_{3\rho\delta}} dy ds.$$

By the same argument to prove the maximal regularity for the heat kernel (see [51, Theorem 7.3]), we see that

$$\begin{aligned} & \|V_2(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^{\frac{4c}{p}}(B_{\rho\delta}^+))} \\ & \leq C \| |u(\Psi(\cdot), \cdot)|^p \chi_{\Phi(\Omega_R) \cap B_{3\rho\delta}} \|_{L^l(-\delta^2, 0; L^{\frac{4c}{p}}(\mathbf{R}^n))} \\ & = C \|u(\Psi(\cdot), \cdot)\|_{L^{pl}(-\delta^2, 0; L^{4c}(\Phi(\Omega_R) \cap B_{3\rho\delta}))}^p \leq \delta^{\frac{2}{l}} M^p, \end{aligned}$$

so that

$$(A.4) \quad \|V_2(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \leq C\delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{p}{4c})} M^p.$$

Finally, we consider V_3 . Again in the same way as in (A.2), we have

$$\begin{aligned} V_3(\Psi(x), t) & \leq \int_{-2\delta^2}^t \int_{\mathbf{R}^n} K_2((x-y)/2, t-s) \\ & \quad \times (\delta^{-2}|u| + \delta^{-1}|\nabla u|)(\Psi(y), s) \chi_{\Phi(\Omega_R) \cap (\overline{B_{3\rho\delta}} \setminus B_{2\rho\delta})} dy ds. \end{aligned}$$

for $x \in B_{\rho\delta}^+$ and $-2\delta^2 < t < 0$. We observe that there exists $C > 0$ satisfying

$$K_2((x-y)/2, t-s) \leq C\delta^{-n-2}$$

for $x \in B_{\rho\delta}^+$, $y \in \overline{B_{3\rho\delta}} \setminus B_{2\rho\delta}$ and $-2\delta^2 < s < t < 0$. Then by the change of variables, the Hölder inequality, $\Omega_R \cap \Psi(B_{3\rho\delta}) \subset \Omega_{3R/4}$ and Proposition 2.1, we see that

$$\begin{aligned} & V_3(\Psi(x), t) \\ & \leq C\delta^{-n-2} \int_{-2\delta^2}^t \int_{\Phi(\Omega_R) \cap B_{3\rho\delta}} (\delta^{-2}|u(\Psi(y), s)| + \delta^{-1}|\nabla u(\Psi(y), s)|) dy ds \\ & \leq C\delta^{-n-2} (M\delta^{n(1-\frac{1}{4c})} + (M+M^p)\delta^{n(1-\frac{1}{4^*})+1}) \leq C\delta^{-\frac{2p}{4c}} (M+M^p) \end{aligned}$$

for $x \in B_{\rho\delta}^+$ and $-2\delta^2 < t < 0$. Hence we obtain

$$\|V_3(\Psi(\cdot), \cdot)\|_{L^l(-\delta^2, 0; L^r(B_{\rho\delta}^+))} \leq C\delta^{\frac{2}{l} + n(\frac{1}{r} - \frac{p}{4c})} (M+M^p).$$

By combining this inequality, (A.1), (A.3) and (A.4), we obtain the desired estimate for \hat{u}_t . Then the desired estimate for \hat{u}_t can be obtained by using the equation in (3.5). \square

Lemma A.2. *Assume either (R1) or (R2). Let $1 \leq l < \infty$ and $1 \leq r \leq qc/p$. Then there exists $C > 0$ depending on R such that*

$$\|u_t\|_{L^l(-1/4, 0; L^r(\Omega_{R/2}))} + \|\nabla^2 u\|_{L^l(-1/4, 0; L^r(\Omega_{R/2}))} \leq C(M+M^p).$$

Proof. By easy modifications of Lemma A.1, we can see that

$$\|\nabla^2 u\|_{L^l(-1/4, 0; L^r(\Omega_{R/2}))} \leq C(M+M^p).$$

Then by the equation in (2.1), we obtain the desired inequality. \square

Let us next show a gradient estimate for the blow-up limit \bar{u} obtained in Lemma 5.5. To estimate $\nabla \bar{u}$, we derive a localized integral equation for \bar{u} .

Lemma A.3. *Assume (R2). Let $\phi \in C_0^\infty(\mathbf{R}^n)$ satisfy $0 \leq \phi \leq 1$ in \mathbf{R}^n , $\phi = 0$ in $\mathbf{R}^n \setminus B_{3/5}$ and $\phi = 1$ in $B_{4/5}$. Set $\phi_\rho(x) := \phi(x/\rho)$ for $0 < \rho < 1$. Then \bar{u} satisfies*

$$\begin{aligned} \bar{u}(x, t) &= \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, \rho^2/4) \phi_\rho(y) \bar{u}(y, t - \rho^2/4) dy \\ &\quad + \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, t-s) (\phi_\rho |\bar{u}|^{p-1} \bar{u} + \bar{u} \Delta \phi_\rho) dy ds \\ &\quad + 2 \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} \nabla_y G_{\mathbf{R}_+^n}(x, y, t-s) \cdot \nabla \phi_\rho(y) \bar{u}(y, s) dy ds \end{aligned}$$

for a.e. $x \in B_{\rho/2}^+$ and $-\rho^2/4 < t < 0$.

Proof. Let us convert our problem to the one in \mathbf{R}_+^n . Let $\psi \in C_0^\infty(\mathbf{R}^n)$ satisfy $0 \leq \psi \leq 1$ in \mathbf{R}^n , $\psi = 0$ in $\mathbf{R}^n \setminus B_{4R/5}$ and $\psi = 1$ in $B_{3R/5}$. Set $\psi_k(x) := \psi(\Psi(\delta_k x))$ and $v_k := \phi_\rho \psi_k u_k$. Note that $\psi_k = 0$ in $\mathbf{R}^n \setminus \delta_k^{-1} \Phi(B_{4R/5})$ and $\psi_k = 1$ in $\delta_k^{-1} \Phi(B_{3R/5})$. Then by (5.5), we see that

$$\begin{cases} (v_k)_t - \Delta v_k = \phi_\rho \psi_k |u_k|^{p-1} u_k - \psi_k u_k \Delta \phi_\rho - 2\psi_k \nabla \phi_\rho \cdot \nabla u_k + \mathcal{R}_k \\ \text{in } \mathbf{R}_+^n \times (-\delta_k^{-2}, 0), \\ v_k = 0 \quad \text{on } \partial \mathbf{R}_+^n \times (-\delta_k^{-2}, 0), \end{cases}$$

where

$$\begin{aligned} \mathcal{R}_k &:= \phi_\rho \psi_k (-2\nabla'(\partial_{x_n} u_k) \cdot \nabla' f_k + (\partial_{x_n}^2 u_k) |\nabla' f_k|^2 - (\partial_{x_n} u_k) \Delta' f_k) \\ &\quad - (2u_k \nabla \psi_k \cdot \nabla \phi_\rho + \phi_\rho u_k \Delta \psi_k + 2\phi_\rho \nabla \psi_k \cdot \nabla u_k). \end{aligned}$$

Thus,

$$\begin{aligned} u_k(x, t) &= \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, \rho^2/4) \phi_\rho(y) \psi_k(y) u_k(y, t - \rho^2/4) dy \\ &\quad + \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, t-s) \phi_\rho \psi_k |u_k|^{p-1} u_k dy ds \\ &\quad - \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(\psi_k u_k \Delta \phi_\rho + 2\psi_k \nabla \phi_\rho \cdot \nabla u_k + \mathcal{R}_k) dy ds \\ &=: V_1^k(x, t) + V_2^k(x, t) + V_3^k(x, t) \end{aligned}$$

for $x \in B_{\rho/2}^+$, $-\rho^2/4 < t < 0$ and $k \geq k_\rho$, where k_ρ is given by the first part of Subsection 5.1. Lemma 5.5 (ii) shows that $u_k(\cdot, t)$ converges to $\bar{u}(\cdot, t)$ in $L^1(B_{\rho/2}^+)$ for each $-\rho^2/4 < t < 0$ as $k \rightarrow \infty$.

We show that the right-hand side of the integral equation for a subsequence of u_k still denoted by u_k converges to the one in the following integral equation:

$$\begin{aligned}
\bar{u}(x, t) &= \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, \rho^2/4) \phi_\rho(y) \bar{u}(y, t - \rho^2/4) dy \\
&\quad + \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, t-s) \phi_\rho |\bar{u}|^{p-1} \bar{u} dy ds \\
&\quad - \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(x, y, t-s) (\bar{u} \Delta \phi_\rho + 2 \nabla \phi_\rho \cdot \nabla \bar{u}) dy ds \\
&=: \bar{V}_1(x, t) + \bar{V}_2(x, t) + \bar{V}_3(x, t)
\end{aligned} \tag{A.5}$$

for a.e. $x \in B_{\rho/2}^+$ and $-\rho^2/4 < t < 0$. For V_1^k , from (2.5), $0 \leq \psi_k \leq 1$, the Hölder inequality and Lemma 5.5 (ii) and (iii), it follows that

$$\begin{aligned}
\|V_1^k(\cdot, t) - \bar{V}_1(\cdot, t)\|_{L^1(B_{\rho/2}^+)} &\leq C \rho^{-n} \int_{B_{\rho/2}^+} \int_{\mathbf{R}_+^n} \phi_\rho |\psi_k u_k - \bar{u}| dy dx \\
&\leq C \|u_k(\cdot, t - \rho^2/4) - \bar{u}(\cdot, t - \rho^2/4)\|_{L^1(B_\rho^+)} \\
&\quad + C \|(\psi_k - 1) \bar{u}(\cdot, t - \rho^2/4)\|_{L^1(B_\rho^+)} \rightarrow 0
\end{aligned}$$

as $k \rightarrow \infty$. For V_2^k , by computations similar to that of V_1^k with Young's inequality, we see that, as $k \rightarrow \infty$,

$$\begin{aligned}
\|V_2^k(\cdot, t) - \bar{V}_2(\cdot, t)\|_{L^1(B_{\rho/2}^+)} &\leq C \int_{t-\rho^2/4}^t \| |\psi_k| |u_k|^{p-1} u_k - |\bar{u}|^{p-1} \bar{u} \|_{L^1(B_\rho^+)} ds \\
&\leq C \int_{t-\rho^2/4}^t \| (|u_k|^{p-1} + |\bar{u}|^{p-1}) |u_k - \bar{u}| \|_{L^1(B_\rho^+)} ds \\
&\quad + C \int_{t-\rho^2/4}^t \| (\psi_k - 1) |\bar{u}|^p \|_{L^1(B_\rho^+)} ds \rightarrow 0.
\end{aligned}$$

For V_3^k , we focus on the most subtle term $\phi_\rho \psi_k \nabla'(\partial_{x_n} u_k) \cdot \nabla' f_k$ in \mathcal{R}_k , that is, we prove that

$$\tilde{V}_3^k(\cdot, t) := \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} G_{\mathbf{R}_+^n}(\cdot, y, t-s) \phi_\rho \psi_k \nabla'(\partial_{x_n} u_k) \cdot \nabla' f_k dy ds \rightarrow 0$$

in $L^1(B_{\rho/2}^+)$ for each $-\rho^2/4 < t < 0$ as $k \rightarrow \infty$. From integration by parts and (2.5), it follows that

$$\begin{aligned}
|\tilde{V}_3^k(x, t)| &\leq C \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} K_1(x-y, t-s) \phi_\rho \psi_k |\partial_{x_n} u_k| |\nabla' f_k| dy ds \\
&\quad + C \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} K_0 |\nabla'(\phi_\rho \psi_k)| |\partial_{x_n} u_k| |\nabla' f_k| dy ds \\
&\quad + C \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} K_0 \phi_\rho \psi_k |\partial_{x_n} u_k| |\Delta' f_k| dy ds.
\end{aligned}$$

By Young's inequality, $|\nabla\phi_\rho| \leq C$, $|\nabla\psi_k| \leq C\delta_k \leq C$, Hölder inequality and $\int_{t-\rho^2/4}^t (t-s)^{-(1/2)\cdot(4/3)} ds \leq C$, we have

$$\begin{aligned} \|\tilde{V}_3^k(\cdot, t)\|_{L^1(B_{\rho/2}^+)} &\leq C \left(\int_{t-\rho^2/4}^t \|\nabla u_k(\cdot, s)\| \|\nabla' f_k\|_{L^1(B_\rho^+)}^4 ds \right)^{1/4} \\ &\quad + C \int_{t-\rho^2/4}^t \|\nabla u_k(\cdot, s)\| (|\nabla' f_k| + |\Delta' f_k|)_{L^1(B_\rho^+)} ds. \end{aligned}$$

Hence from the Hölder inequality for the Lorentz spaces (see [49, Proposition 2.1]), (5.8), (5.6), (5.7) and the Lebesgue dominated convergence theorem, it follows that

$$\|\tilde{V}_3^k(\cdot, t)\|_{L^1(B_{\rho/2}^+)} \leq C(M + M^p) \|\nabla' f_k + |\Delta' f_k|\|_{L^{\frac{q_*}{q_*-1}}(B_\rho^+)} \rightarrow 0$$

for each $-\rho^2/4 < t < 0$ as $k \rightarrow \infty$. The other terms in V_3^k can be handled more easily. Hence we obtain (A.5). This implies the desired equality. \square

We give a gradient estimate for \bar{u} .

Lemma A.4. *Assume (R2). Then there exists a constant $C > 0$ independent of ρ such that*

$$\|\nabla\bar{u}\|_{L^2(Q_{\rho/2}^+)} \leq C\rho^{\frac{n}{2} - \frac{2}{p-1} - \frac{2}{q_c}} (1 + M^{p-1}) \|\bar{u}\|_{L^{q_c}(Q_\rho^+)}$$

for any $0 < \rho \leq 1$.

Proof. By Lemma A.3, $\nabla\phi_\rho(x) = \rho^{-1}\nabla\phi(x/\rho)$, $\Delta\phi_\rho(x) = \rho^{-2}\Delta\phi(x/\rho)$ and similar computations to (2.7), there exists $C > 0$ independent of ρ such that

$$\begin{aligned} |\nabla\bar{u}| &\leq C \int_{B_\rho^+} K_1(x-y, \rho^2/4) |\bar{u}(y, t-\rho^2/4)| dy \\ &\quad + C \int_{t-\rho^2/4}^t \int_{B_\rho^+} K_1(x-y, t-s) |\bar{u}(y, s)|^p dy ds \\ &\quad + C \int_{t-\rho^2/4}^t \int_{\mathbf{R}_+^n} \left(\frac{K_1}{\rho^2} + \frac{K_2}{\rho} \right) |\bar{u}| \chi_{\overline{B_{4\rho/5}} \setminus B_{3\rho/5}} dy ds \\ &=: CW_1 + CW_2 + C\rho^{-2}W_3 + C\rho^{-1}W_4 \end{aligned}$$

for a.e. $x \in B_{\rho/2}^+$ and $-\rho^2/4 < t < 0$, where K_1 and K_2 are given by (2.6).

By the Hölder inequality, we have

$$\begin{aligned} W_1(x, t) &\leq \|\bar{u}(\cdot, t-\rho^2/4)\|_{L^{q_c}(B_\rho^+)} \left(\int_{B_\rho^+} K_1(x-y, \rho^2/4)^{\frac{q_c}{q_c-1}} dy \right)^{1-\frac{1}{q_c}} \\ &\leq C\rho^{-1-\frac{n}{q_c}} \|\bar{u}(\cdot, t-\rho^2/4)\|_{L^{q_c}(B_\rho^+)}, \\ \|W_1\|_{L^2(Q_{\rho/2}^+)} &\leq C\rho^{-\frac{n}{q_c} + \frac{n}{2} - \frac{2}{q_c}} \left(\int_{-\rho^2/4}^0 \int_{L^{q_c}(B_\rho^+)} |\bar{u}(x, t-\rho^2/4)|^{q_c} dx dt \right)^{\frac{1}{q_c}} \\ &\leq C\rho^{\frac{n}{2} - \frac{2}{p-1} - \frac{2}{q_c}} \|\bar{u}\|_{L^{q_c}(Q_\rho^+)}. \end{aligned}$$

We estimate W_2 . We consider the cases $q_c/p \geq 2$ and $q_c/p < 2$, respectively. If $q_c/p \geq 2$, then the Hölder inequality gives

$$\|W_2\|_{L^2(Q_{\rho/2}^+)} \leq C\rho^{\frac{n+2}{2}(1-\frac{2}{q_c})} \|W_2\|_{L^{\frac{q_c}{p}}(Q_{\rho/2}^+)}.$$

From the same argument to prove the $L^{q_c/p}$ - $L^{q_c/p}$ estimate and the change of variables, it follows that

$$\begin{aligned} \|W_2(\cdot, t)\|_{L^{\frac{q_c}{p}}(B_{\rho/2}^+)} &\leq C \int_{t-\rho^2/4}^t (t-s)^{-\frac{1}{2}} \|\bar{u}(\cdot, s)\|_{L^{\frac{q_c}{p}}(\mathbf{R}^n)}^p \chi_{B_\rho^+} ds \\ &= C \int_0^{\rho^2/4} \tau^{-\frac{1}{2}} \|\bar{u}(\cdot, t-\tau)\|_{L^{q_c}(B_\rho^+)}^p d\tau. \end{aligned}$$

Thus, by the change of variables again and by $(-\tau - \rho^2/4, -\tau) \subset (-\rho^2, 0)$ for $0 < \tau < \rho^2/4$, we have

$$\begin{aligned} \|W_2\|_{L^{\frac{q_c}{p}}(Q_{\rho/2}^+)} &\leq C \int_0^{\rho^2/4} \tau^{-\frac{1}{2}} \left(\int_{-\rho^2/4}^0 \int_{B_\rho^+} |\bar{u}(\cdot, t-\tau)|^{q_c} dx dt \right)^{\frac{p}{q_c}} d\tau \\ &\leq C \int_0^{\rho^2/4} \tau^{-\frac{1}{2}} \left(\int_{-\rho^2}^0 \int_{B_\rho^+} |\bar{u}(\cdot, s)|^{q_c} dx ds \right)^{\frac{p}{q_c}} d\tau \\ &\leq C \rho^{1+\frac{2(p-1)}{q_c}} M^{p-1} \|\bar{u}\|_{L^{q_c}(Q_\rho^+)}, \end{aligned}$$

and so

$$\|W_2\|_{L^2(Q_{\rho/2}^+)} \leq C \rho^{\frac{n}{2} - \frac{2}{p-1} - \frac{2}{q_c}} M^{p-1} \|\bar{u}\|_{L^{q_c}(Q_\rho^+)}.$$

If $q_c/p < 2$, then the same argument to prove the $L^{q_c/p}$ - L^2 estimate yields

$$\begin{aligned} \|W_2(\cdot, t)\|_{L^2(B_\rho^+)} &\leq C \int_{t-\rho^2/4}^t (t-s)^{-\frac{1}{2} - \frac{n}{2}(\frac{p}{q_c} - \frac{1}{2})} \|\bar{u}(\cdot, s)\|_{L^{\frac{q_c}{p}}(\mathbf{R}^n)}^p \chi_{B_\rho^+} ds \\ &\leq C \int_{\mathbf{R}} |t-s|^{-1+\alpha} \|\bar{u}(\cdot, s)\|_{L^{q_c}(B_\rho^+)}^p \chi_{(-\rho^2, 0)}(s) ds \\ &\leq C M^{p-1} \int_{\mathbf{R}} |t-s|^{-1+\alpha} \|\bar{u}(\cdot, s)\|_{L^{q_c}(B_\rho^+)} \chi_{(-\rho^2, 0)} ds, \end{aligned}$$

where

$$\alpha := \frac{1}{2} - \frac{n}{2} \left(\frac{p}{q_c} - \frac{1}{2} \right).$$

Note that $0 < \alpha < 1/2$ by $p > p_S$ and $q_c/p < 2$. From the Hardy-Littlewood-Sobolev inequality and the Hölder inequality with $2/(2\alpha + 1) < q_c$, it follows that

$$\begin{aligned} \|W_2\|_{L^2(Q_{\rho/2}^+)} &= \left\| \|W_2(\cdot, t)\|_{L^2(B_\rho^+)} \chi_{(-\rho^2/4, 0)} \right\|_{L^2(\mathbf{R})} \\ &\leq C M^{p-1} \left\| \|\bar{u}(\cdot, t)\|_{L^{q_c}(B_\rho^+)} \chi_{(-\rho^2, 0)} \right\|_{L^{\frac{2}{2\alpha+1}}(\mathbf{R})} \\ &\leq C \rho^{\frac{n}{2} - \frac{2}{p-1} - \frac{2}{q_c}} M^{p-1} \|\bar{u}\|_{L^{q_c}(Q_\rho^+)}. \end{aligned}$$

We consider W_3 and W_4 . By the definitions of K_1 and K_2 in (2.6), there exists a constant $C > 0$ independent of ρ such that

$$\rho^{-2} K_1(x-y, t) + \rho^{-1} K_2(x-y, t) \leq C \rho^{-n-3}$$

for $x \in B_{\rho/2}$, $y \in \overline{B_{4\rho/5}} \setminus B_{3\rho/5}$ and $t > 0$. Hence by the Hölder inequality, we have

$$\begin{aligned} \rho^{-2}|W_3(x,t)| + \rho^{-1}|W_4(x,t)| &\leq C\rho^{-n-3+(n+2)(1-\frac{1}{q_c})}\|\bar{u}\|_{L^{q_c}(Q_\rho^+)}, \\ \rho^{-2}\|W_3\|_{L^2(Q_{\rho/2}^+)} + \rho^{-1}\|W_4\|_{L^2(Q_{\rho/2}^+)} &\leq C\rho^{\frac{n}{2}-\frac{2}{p-1}-\frac{2}{q_c}}\|\bar{u}\|_{L^{q_c}(Q_\rho^+)}. \end{aligned}$$

The above estimates show the desired inequality. \square

APPENDIX B. COMPACTNESS RESULTS

We recall an Aubin-Lions type compactness result from [79, Proposition 51.3] and [76, Proposition 2.1]. See also [3] for more general statement. Note that a pair of Banach spaces (E_0, E_1) is called an interpolation couple if there exists a locally convex space E such that $E_0, E_1 \hookrightarrow E$.

Proposition B.1. *Let (E_0, E_1) be an interpolation couple. Assume that E_1 is compactly embedded in E_0 . Let $1 \leq p_0, p_1 < \infty$, $0 < \theta < 1$, $1/p_\theta = (1-\theta)/p_0 + \theta/p_1$ and $s < 1 - \theta$. Then,*

$$W^{1,p_0}(0, T; E_0) \cap L^{p_1}(0, T; E_1) \hookrightarrow W^{s,p_\theta}(0, T; (E_0, E_1)_{\theta,p_\theta})$$

and this embedding is compact.

We give a consequence of this result in a form which is used to prove Lemma 5.5.

Lemma B.2. *Let $1 < r < \infty$ and let \mathcal{B} be a smooth bounded domain in \mathbf{R}^n . Then,*

$$W^{1,5}(-1, 0; L^r(\mathcal{B})) \cap L^5(-1, 0; W^{2,r}(\mathcal{B})) \hookrightarrow C([-1, 0]; W^{1,r}(\mathcal{B}))$$

and this embedding is compact.

Proof. We write $L_x^r := L^r(\mathcal{B})$, $W_x^{2,r} := W^{2,r}(\mathcal{B})$ and so on. Since $W_x^{2,r} \hookrightarrow L_x^r$ is compact, Proposition B.1 with $p_0 = p_1 = 5$, $\theta = 2/3$ and $s = 1/4$ gives

$$W^{1,5}(-1, 0; L_x^r) \cap L^5(-1, 0; W_x^{2,r}) \hookrightarrow W^{1/4,5}(-1, 0; (L_x^r, W_x^{2,r})_{2/3,5}).$$

By [88, page 327], we have $(L_x^r, W_x^{2,r})_{2/3,5} = B_{r,5,x}^{4/3} \hookrightarrow W_x^{1,r}$. Then the Sobolev embedding in time yields

$$W^{1/4,5}(-1, 0; (L_x^r, W_x^{2,r})_{2/3,5}) \hookrightarrow C([-1, 0]; W_x^{1,r}),$$

and hence the lemma follows. \square

ACKNOWLEDGMENTS

The authors would like to express their gratitude to Professors Kazuhiro Ishige, Hiroaki Kikuchi and Mitsuo Higaki for their valuable comments and to Professor Shi-Zhong Du for sending [12] to the authors.

The first author was supported in part by JSPS KAKENHI Grant Numbers 16H06339, 17K05312, 21H00991 and 21H04433. The second author was supported in part by JSPS KAKENHI Grant Numbers 19K14567 and 22H01131.

REFERENCES

- [1] D. Albritton, Blow-up criteria for the Navier-Stokes equations in non-endpoint critical Besov spaces. *Anal. PDE* 11 (2018), no. 6, 1415–1456.
- [2] D. Albritton, T. Barker, Localised necessary conditions for singularity formation in the Navier-Stokes equations with curved boundary. *J. Differential Equations* 269 (2020), no. 9, 7529–7573.
- [3] H. Amann, Compact embeddings of vector-valued Sobolev and Besov spaces. *Glas. Mat. Ser. III* 35 (55) (2000), no. 1, 161–177.
- [4] N. Aronszajn, A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. *J. Math. Pures Appl.* 36 (1957), 235–249.
- [5] P. Baras, Non-unicité des solutions d’une equation d’évolution non-linéaire. *Ann. Fac. Sci. Toulouse Math.* 5 (1983), no. 3-4, 287–302.
- [6] T. Barker, C. Prange, Quantitative regularity for the Navier-Stokes equations via spatial concentration. *Comm. Math. Phys.* 385 (2021), no. 2, 717–792.
- [7] T. Barker, G. Seregin, A necessary condition of potential blowup for the Navier-Stokes system in half-space. *Math. Ann.* 369 (2017), no. 3-4, 1327–1352.
- [8] S. Blatt, M. Struwe, An analytic framework for the supercritical Lane-Emden equation and its gradient flow. *Int. Math. Res. Not. IMRN* 2015, no. 9, 2342–2385.
- [9] H. Brezis, 1979, unpublished manuscript.
- [10] H. Brezis, T. Cazenave, A nonlinear heat equation with singular initial data. *J. Anal. Math.* 68 (1996), 277–304.
- [11] T. Cazenave, F. Dickstein, I. Naumkin, F. B. Weissler, Sign-changing self-similar solutions of the nonlinear heat equation with positive initial value. *Amer. J. Math.* 142 (2020), no. 5, 1439–1495.
- [12] K.-S. Chou, S.-Z. Du, On global partial regularity for borderline solutions of semilinear parabolic problems. *Trends in partial differential equations*, 41–52, *Adv. Lect. Math. (ALM)*, 10, Int. Press, Somerville, MA, 2010.
- [13] K.-S. Chou, S.-Z. Du, G.-F. Zheng, On partial regularity of the borderline solution of semilinear parabolic problems. *Calc. Var. Partial Differential Equations* 30 (2007), no. 2, 251–275.
- [14] C. Collot, Nonradial type II blow up for the energy-supercritical semilinear heat equation. *Anal. PDE* 10 (2017), no. 1, 127–252.
- [15] C. Collot, F. Merle, P. Raphaël, Dynamics near the ground state for the energy critical nonlinear heat equation in large dimensions. *Comm. Math. Phys.* 352 (2017), no. 1, 215–285.
- [16] C. Collot, F. Merle, P. Raphaël, Stability of ODE blow-up for the energy critical semilinear heat equation. *C. R. Math. Acad. Sci. Paris* 355 (2017), no. 1, 65–79.
- [17] C. Collot, F. Merle, P. Raphaël, Strongly anisotropic type II blow up at an isolated point. *J. Amer. Math. Soc.* 33 (2020), no. 2, 527–607.
- [18] C. Collot, P. Raphaël, J. Szeftel, On the stability of type I blow up for the energy supercritical heat equation. *Mem. Amer. Math. Soc.* 260 (2019), no. 1255, v+97 pp.
- [19] M. del Pino, C.-C. Lai, M. Musso, J. Wei, Y. Zhou, New type II Finite time blow-up for the energy supercritical heat equation, preprint, arXiv:2006.00716.
- [20] M. del Pino, M. Musso, F. Pacard, Bubbling along boundary geodesics near the second critical exponent. *J. Eur. Math. Soc. (JEMS)* 12 (2010), no. 6, 1553–1605.
- [21] M. del Pino, M. Musso, J. Wei, Type II blow-up in the 5-dimensional energy critical heat equation. *Acta Math. Sin. (Engl. Ser.)* 35 (2019), no. 6, 1027–1042.
- [22] M. del Pino, M. Musso, J. Wei, Geometry driven type II higher dimensional blow-up for the critical heat equation. *J. Funct. Anal.* 280 (2021), no. 1, Paper No. 108788, 49 pp.
- [23] M. del Pino, M. Musso, J. Wei, Q. Zhang, Y. Zhang, Type II Finite time blow-up for the three dimensional energy critical heat equation, preprint, arXiv:2002.05765.
- [24] M. del Pino, M. Musso, J. Wei, Y. Zhou, Type II finite time blow-up for the energy critical heat equation in \mathbf{R}^4 . *Discrete Contin. Dyn. Syst.* 40 (2020), no. 6, 3327–3355.
- [25] H. Dong, D. Du, The Navier-Stokes equations in the critical Lebesgue space. *Comm. Math. Phys.* 292 (2009), no. 3, 811–827.
- [26] H. Dong, K. Wang, Interior and boundary regularity for the Navier-Stokes equations in the critical Lebesgue spaces. *Discrete Contin. Dyn. Syst.* 40 (2020), no. 9, 5289–5323.
- [27] S.-Z. Du, The singular set of stationary solution for semilinear elliptic equations with supercritical growth. *J. Differential Equations* 256 (2014), no. 7, 2392–2405.

- [28] S.-Z. Du, Energy non-collapsing and refined blowup for a semilinear heat equation. *J. Differential Equations* 266 (2019), no. 9, 5942–5970.
- [29] T. Duyckaerts, C. Kenig, F. Merle, Scattering for radial, bounded solutions of focusing supercritical wave equations. *Int. Math. Res. Not. IMRN* 2014, no. 1, 224–258.
- [30] T. Duyckaerts, J. Yang, Blow-up of a critical Sobolev norm for energy-subcritical and energy-supercritical wave equations. *Anal. PDE* 11 (2018), no. 4, 983–1028.
- [31] L. Escauriaza, G. A. Seregin, V. Sverak, $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Uspekhi Mat. Nauk* 58 (2003), no. 2(350), 3–44; translation in *Russian Math. Surveys* 58 (2003), no. 2, 211–250.
- [32] S. Filippas, M. A. Herrero, J. J. L. Velázquez, Fast blow-up mechanisms for sign-changing solutions of a semilinear parabolic equation with critical nonlinearity. *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* 456 (2000), no. 2004, 2957–2982.
- [33] A. Friedman, B. McLeod, Blow-up of positive solutions of semilinear heat equations. *Indiana Univ. Math. J.* 34 (1985), no. 2, 425–447.
- [34] H. Fujita, On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$. *J. Fac. Sci. Univ. Tokyo Sect. I* 13 (1966), 109–124.
- [35] I. Gallagher, G. S. Koch, F. Planchon, Blow-up of critical Besov norms at a potential Navier-Stokes singularity. *Comm. Math. Phys.* 343 (2016), no. 1, 39–82.
- [36] M.-H. Giga, Y. Giga, J. Saal, Nonlinear partial differential equations. Asymptotic behavior of solutions and self-similar solutions. *Progress in Nonlinear Differential Equations and their Applications*, 79. Birkhäuser Boston, Ltd., Boston, MA, 2010.
- [37] Y. Giga, R. V. Kohn, Characterizing blowup using similarity variables. *Indiana Univ. Math. J.* 36 (1987), no. 1, 1–40.
- [38] Y. Giga, S. Matsui, S. Sasayama, Blow up rate for semilinear heat equations with subcritical nonlinearity. *Indiana Univ. Math. J.* 53 (2004), no. 2, 483–514.
- [39] Y. Giga, S. Matsui, S. Sasayama, On blow-up rate for sign-changing solutions in a convex domain. *Math. Methods Appl. Sci.* 27 (2004), no. 15, 1771–1782.
- [40] J. Harada, A higher speed type II blowup for the five dimensional energy critical heat equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 37 (2020), no. 2, 309–341.
- [41] J. Harada, A type II blowup for the six dimensional energy critical heat equation. *Ann. PDE* 6 (2020), no. 2, Paper No. 13, 63 pp.
- [42] A. Haraux, F. B. Weissler, Nonuniqueness for a semilinear initial value problem. *Indiana Univ. Math. J.* 31 (1982), no. 2, 167–189.
- [43] M. A. Herrero, J. J. L. Velázquez, A blow up result for semilinear heat equations in the supercritical case, unpublished manuscript.
- [44] M. A. Herrero, J. J. L. Velázquez, Explosion de solutions d'équations paraboliques semilinéaires supercritiques. *C. R. Acad. Sci. Paris Sér. I Math.* 319 (1994), no. 2, 141–145.
- [45] K. Ishige, N. Mizoguchi, Blow-up behavior for semilinear heat equations with boundary conditions. *Differential Integral Equations* 16 (2003), no. 6, 663–690.
- [46] C. Kenig, F. Merle, Scattering for $\dot{H}^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions. *Trans. Amer. Math. Soc.* 362 (2010), no. 4, 1937–1962.
- [47] C. Kenig, F. Merle, Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications. *Amer. J. Math.* 133 (2011), no. 4, 1029–1065.
- [48] R. Killip, M. Visan, Energy-supercritical NLS: critical H^s -bounds imply scattering. *Comm. Partial Differential Equations* 35 (2010), no. 6, 945–987.
- [49] H. Kozono, M. Yamazaki, Uniqueness criterion of weak solutions to the stationary Navier-Stokes equations in exterior domains. *Nonlinear Anal.* 38 (1999), no. 8, 959–970.
- [50] O. A. Ladyženskaja, V. A. Solonnikov, N. N. Ural'ceva, Linear and quasilinear equations of parabolic type. *Translations of Mathematical Monographs*, Vol. 23 American Mathematical Society, Providence, R.I. 1968.
- [51] P. G. Lemarié-Rieusset, Recent developments in the Navier-Stokes problem. *Chapman & Hall/CRC Research Notes in Mathematics*, 431. Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [52] T. Li, L. Sun, S. Wang, A slow blow up solution for the four dimensional energy critical semi linear heat equation, preprint, arXiv: 2204.11201.
- [53] F. Lin, C. Wang, The analysis of harmonic maps and their heat flows. *World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ*, 2008.

- [54] Y. Maekawa, H. Miura, C. Prange, Local energy weak solutions for the Navier-Stokes equations in the half-space. *Comm. Math. Phys.* 367 (2019), no. 2, 517–580.
- [55] H. Matano, F. Merle, On nonexistence of type II blowup for a supercritical nonlinear heat equation. *Comm. Pure Appl. Math.* 57 (2004), no. 11, 1494–1541.
- [56] H. Matano, F. Merle, Classification of type I and type II behaviors for a supercritical nonlinear heat equation. *J. Funct. Anal.* 256 (2009), no. 4, 992–1064.
- [57] H. Matano, M. Merle, Threshold and generic type I behaviors for a supercritical nonlinear heat equation. *J. Funct. Anal.* 261 (2011), no. 3, 716–748.
- [58] J. Matos, Blow up of critical and subcritical norms in semilinear heat equations. *Adv. Differential Equations* 3 (1998), no. 4, 497–532.
- [59] F. Merle, P. Raphaël, Blow up of the critical norm for some radial L^2 super critical nonlinear Schrödinger equations. *Amer. J. Math.* 130 (2008), no. 4, 945–978.
- [60] F. Merle, P. Raphaël, J. Szeftel, On strongly anisotropic type I blowup. *Int. Math. Res. Not. IMRN* 2020, no. 2, 541–606.
- [61] Y. Meyer, Wavelets, paraproducts, and Navier-Stokes equations. *Current developments in mathematics, 1996* (Cambridge, MA), 105–212, Int. Press, Boston, MA, 1997.
- [62] A. S. Mikhailov, T. N. Shilkin, $L_{3,\infty}$ -solutions to the 3D-Navier-Stokes system in the domain with a curved boundary. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 336 (2006), *Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts.* 37, 133–152, 276; reprinted in *J. Math. Sci. (N.Y.)* 143 (2007), no. 2, 2924–2935.
- [63] N. Mizoguchi, Type-II blowup for a semilinear heat equation. *Adv. Differential Equations* 9 (2004), no. 11-12, 1279–1316.
- [64] N. Mizoguchi, Boundedness of global solutions for a supercritical semilinear heat equation and its application. *Indiana Univ. Math. J.* 54 (2005), no. 4, 1047–1059.
- [65] N. Mizoguchi, Rate of type II blowup for a semilinear heat equation. *Math. Ann.* 339 (2007), no. 4, 839–877.
- [66] N. Mizoguchi, Nonexistence of type II blowup solution for a semilinear heat equation. *J. Differential Equations* 250 (2011), no. 1, 26–32.
- [67] N. Mizoguchi, Ph. Souplet, Optimal condition for blow-up of the critical L^q norm for the semilinear heat equation. *Adv. Math.* 355 (2019), 106763, 24 pp.
- [68] A. Mukai, Y. Seki, Refined construction of type II blow-up solution for semilinear heat equations with Joseph-Lundgren supercritical nonlinearity. *Discrete Contin. Dyn. Syst.* 41 (2021), no. 10, 4847–4885.
- [69] Y. Naito, T. Suzuki, Existence of type II blowup solutions for a semilinear heat equation with critical nonlinearity. *J. Differential Equations* 232 (2007), no. 1, 176–211.
- [70] F. Pacard, Partial regularity for weak solutions of a nonlinear elliptic equation. *Manuscripta Math.* 79 (1993), no. 2, 161–172.
- [71] F. Pacard, Convergence and partial regularity for weak solutions of some nonlinear elliptic equation: the supercritical case. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 11 (1994), no. 5, 537–551.
- [72] S. Palasek, Improved quantitative regularity for the Navier-Stokes equations in a scale of critical spaces. *Arch. Ration. Mech. Anal.* 242 (2021), no. 3, 1479–1531.
- [73] N. C. Phuc, The Navier-Stokes equations in nonendpoint borderline Lorentz spaces. *J. Math. Fluid Mech.* 17 (2015), no. 4, 741–760.
- [74] P. Poláčik, P. Quittner, Ph. Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems. II. Parabolic equations. *Indiana Univ. Math. J.* 56 (2007), no. 2, 879–908.
- [75] C.-C. Poon, Blow-up behavior for semilinear heat equations in nonconvex domains. *Differential Integral Equations* 13 (2000), no. 7-9, 1111–1138.
- [76] P. Quittner, Continuity of the blow-up time and a priori bounds for solutions in superlinear parabolic problems. *Houston J. Math.* 29 (2003), no. 3, 757–799.
- [77] P. Quittner, Liouville theorems for scaling invariant superlinear parabolic problems with gradient structure. *Math. Ann.* 364 (2016), no. 1-2, 269–292.
- [78] P. Quittner, Optimal Liouville theorems for superlinear parabolic problems. *Duke Math. J.* 170 (2021), no. 6, 1113–1136.
- [79] P. Quittner, Ph. Souplet, Superlinear parabolic problems. Blow-up, global existence and steady states. Second edition. *Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser/Springer, Cham*, 2019.

- [80] Y. Seki, Type II blow-up mechanisms in a semilinear heat equation with critical Joseph-Lundgren exponent. *J. Funct. Anal.* 275 (2018), no. 12, 3380–3456.
- [81] Y. Seki, Type II blow-up mechanisms in a semilinear heat equation with Lepin exponent. *J. Differential Equations* 268 (2020), no. 3, 853–900.
- [82] G. Seregin, On smoothness of $L_{3,\infty}$ -solutions to the Navier-Stokes equations up to boundary. *Math. Ann.* 332 (2005), no. 1, 219–238.
- [83] G. Seregin, A certain necessary condition of potential blow up for Navier-Stokes equations. *Comm. Math. Phys.* 312 (2012), no. 3, 833–845.
- [84] R. Schweyer, Type II blow-up for the four dimensional energy critical semi linear heat equation. *J. Funct. Anal.* 263 (2012), no. 12, 3922–3983.
- [85] M. Struwe, On the evolution of harmonic maps in higher dimensions. *J. Differential Geom.* 28 (1988), no. 3, 485–502.
- [86] T. Tao, Quantitative bounds for critically bounded solutions to the Navier-Stokes equations. *Nine mathematical challenges—an elucidation*, 149–193, *Proc. Sympos. Pure Math.*, 104, Amer. Math. Soc., Providence, RI, 2021.
- [87] E. Terraneo, Non-uniqueness for a critical non-linear heat equation. *Comm. Partial Differential Equations* 27 (2002), no. 1-2, 185–218.
- [88] H. Triebel, *Interpolation theory, function spaces, differential operators*. North-Holland Mathematical Library, 18. North-Holland Publishing Co., Amsterdam-New York, 1978.
- [89] T.-P. Tsai, *Lectures on Navier-Stokes equations*. Graduate Studies in Mathematics, 192. American Mathematical Society, Providence, RI, 2018.
- [90] C. Wang, Heat flow of harmonic maps whose gradients belong to $L_x^n L_t^\infty$. *Arch. Ration. Mech. Anal.* 188 (2008), no. 2, 351–369.
- [91] K. Wang, J. Wei, Analysis of blow-up locus and existence of weak solutions for nonlinear supercritical problems. *Int. Math. Res. Not. IMRN* 2015, no. 10, 2634–2670.
- [92] K. Wang, J. Wei, Refined blowup analysis and nonexistence of Type II blowups for an energy critical nonlinear heat equation, preprint, arXiv: 2101.07186.
- [93] F. B. Weissler, Semilinear evolution equations in Banach spaces. *J. Functional Analysis* 32 (1979), no. 3, 277–296.
- [94] F. B. Weissler, Local existence and nonexistence for semilinear parabolic equations in L^p . *Indiana Univ. Math. J.* 29 (1980), no. 1, 79–102.
- [95] F. B. Weissler, An L^∞ blow-up estimate for a nonlinear heat equation. *Comm. Pure Appl. Math.* 38 (1985), no. 3, 291–295.
- [96] F. B. Weissler, L^p -energy and blow-up for a semilinear heat equation. *Nonlinear functional analysis and its applications, Part 2* (Berkeley, Calif., 1983), 545–551, *Proc. Sympos. Pure Math.*, 45, Part 2, Amer. Math. Soc., Providence, RI, 1986.

DEPARTMENT OF MATHEMATICAL AND COMPUTING SCIENCE, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1 OOKAYAMA, MEGURO-KU, TOKYO 152-8552, JAPAN

Email address: miura@is.titech.ac.jp

DEPARTMENT OF MATHEMATICAL AND COMPUTING SCIENCE, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1 OOKAYAMA, MEGURO-KU, TOKYO 152-8552, JAPAN

Email address: takahashi@c.titech.ac.jp