

ANNIHILATOR OF (\mathfrak{g}, K) -MODULES OF $O(p, q)$ ASSOCIATED TO THE FINITE-DIMENSIONAL REPRESENTATION OF \mathfrak{sl}_2

TAKASHI HASHIMOTO

ABSTRACT. Let \mathfrak{g} denote the complexified Lie algebra of $G = O(p, q)$ and K a maximal compact subgroup of G . In the previous paper, we constructed (\mathfrak{g}, K) -modules associated to the finite-dimensional representation of \mathfrak{sl}_2 of dimension $m + 1$, which we denote by $M^+(m)$ and $M^-(m)$. The aim of this paper is to show that the annihilator of $M^\pm(m)$ is the Joseph ideal if and only if $m = 0$. We shall see that an element of the symmetric of square $S^2(\mathfrak{g})$ that is given in terms of the Casimir elements of \mathfrak{g} and the complexified Lie algebra of K plays a critical role in the proof of the main result.

1. INTRODUCTION

For a complex simple Lie algebra \mathfrak{g} , Joseph constructed the so-called Joseph ideal in [8], which is a completely prime two-sided primitive ideal in the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} and has the associated variety of minimal dimension. He also proved that it is unique if \mathfrak{g} is not of type A (see also [2]). An irreducible unitary representation of a simple Lie group is called a minimal representation if its annihilator in $U(\mathfrak{g})$ is equal to the Joseph ideal.

The best known example of the minimal representation is the oscillator (or Segal-Shale-Weil) representation of the metaplectic group, the double cover of the symplectic group $\mathrm{Sp}(n, \mathbb{R})$. It was shown in [5] that the canonical quantization of the moment map on real symplectic vector spaces gives rise to the underlying (\mathfrak{g}, K) -module of the oscillator representation of real reductive Lie groups G , where \mathfrak{g} denotes the complexified Lie algebra of G and K a maximal compact subgroup of G .

In the case of $G = O(p, q)$, the indefinite orthogonal group, applying this method to the symplectic vector space $((\mathbb{C}^{p+q})_{\mathbb{R}}, \omega)$ with $\omega(z, w) = \mathrm{Im}(z^* I_{p,q} w)$, on which G acts via matrix multiplication in a Hamiltonian way with moment map (see below for details), and making use of the fact that $(O(p, q), \mathrm{SL}_2(\mathbb{R}))$ is a dual pair in $\mathrm{Sp}(p+q, \mathbb{R})$, the author constructed (\mathfrak{g}, K) -modules $M^+(m)$ and $M^-(m)$ of $O(p, q)$ associated to the finite-dimensional representation of \mathfrak{sl}_2 of dimension $m + 1$ for any non-negative integer m in [6]. Note that $M^+(0) = M^-(0)$ by definition and $M^+(m)$ is naturally isomorphic to $M^-(m)$ in fact. Moreover, under the condition

$$p \geq 2, q \geq 2, p + q \in 2\mathbb{N} \quad \text{and} \quad m + 3 \leq \frac{p + q}{2}, \quad (1.1)$$

Date: June 23, 2022.

2010 Mathematics Subject Classification. Primary: 22E46, 17B20, 17B10.

Key words and phrases. indefinite orthogonal group, minimal representation, annihilator, Joseph ideal, Casimir element.

he obtained the K -type formula of $M^\pm(m)$, thereby showed that they are irreducible and that $M^\pm(0)$ is the underlying (\mathfrak{g}, K) -module of the minimal representation of $O(p, q)$. Although it is well known that the Gelfand-Kirillov dimension of the minimal representation of $O(p, q)$ is equal to $p+q-3$ (see, e.g., [9, 12]), we remark that the K -type formula of $M^\pm(m)$ mentioned above implies that they all have the Gelfand-Kirillov dimension equal to $p+q-3$, whereas they have the Bernstein degree equal to $4(m+1)\binom{p+q-4}{p-2}$.

The original object of this project is to clarify the relation between our (\mathfrak{g}, K) -modules $M^\pm(m)$ and the minimal representation of $O(p, q)$ more directly. In fact, we shall see that the annihilator in the universal enveloping algebra $U(\mathfrak{g})$ of $M^\pm(m)$, denoted by $\mathcal{I}^\pm(m)$ in this paper, is the Joseph ideal if and only if $m=0$, which is the main result of this paper (Theorem 3.6). We use a criterion due to Garfinkle [3] (see also [2]) to determine whether $\mathcal{I}^\pm(m)$ is the Joseph ideal or not. We will see that an element of the symmetric square $S^2(\mathfrak{g})$, which we denote by Ξ below, plays a critical role in the proof of our main result. We remark that Ξ is given in terms of the Casimir elements of \mathfrak{g} and \mathfrak{k} , where \mathfrak{k} is the complexified Lie algebra of the maximal compact subgroup K .

The rest of the paper is organized as follows. In Section 2 we briefly review the construction of (\mathfrak{g}, K) -modules $M^\pm(m)$ of $O(p, q)$ mentioned above as well as their properties needed to prove our main results. In Section 3 we prove our main result using the criterion due to Garfinkle. In order to keep the present paper self-contained, we include some standard facts about irreducible decomposition of $S^2(\mathfrak{o}_n)$ in the Appendix.

Notation. Let \mathbb{N} denote the set of non-negative integers. For positive integers p and q , we set $[p] := \{1, 2, \dots, p\}$ and $p+[q] := \{p+1, p+2, \dots, p+q\}$.

2. (\mathfrak{g}, K) -MODULES OF $O(p, q)$ ASSOCIATED TO THE FINITE-DIMENSIONAL REPRESENTATION OF \mathfrak{sl}_2

Throughout this paper, we denote by G the indefinite orthogonal group $O(p, q)$ realized by

$$O(p, q) = \{g \in \mathrm{GL}_n(\mathbb{R}); {}^t g I_{p,q} g = I_{p,q}\},$$

where $n = p+q$ and $I_{p,q} = \begin{bmatrix} 1_p & \\ & -1_q \end{bmatrix}$. Let $\mathfrak{g} = \mathfrak{o}(p, q) \otimes \mathbb{C}$, the complexified Lie algebra of G . Thus,

$$\mathfrak{g} = \{X \in \mathfrak{gl}_n(\mathbb{C}); {}^t X I_{p,q} + I_{p,q} X = O\}.$$

Needless to say, our Lie algebra \mathfrak{g} is \mathbb{C} -isomorphic to the Lie algebra \mathfrak{o}_n consisting of all complex $n \times n$ alternating matrices:

$$\mathfrak{o}_n = \{X \in \mathfrak{gl}_n(\mathbb{C}); {}^t X + X = O\}.$$

Let K be a maximal compact subgroup of G given by

$$K = \left\{ \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} \in G; a \in O(p), d \in O(q) \right\} \simeq O(p) \times O(q).$$

Denote the Lie algebra of K and its complexification by \mathfrak{k}_0 and \mathfrak{k} respectively. Then $\mathfrak{k} \simeq \mathfrak{o}_p \oplus \mathfrak{o}_q$.

Let us fix a nondegenerate invariant bilinear form B defined both on \mathfrak{g} and on \mathfrak{o}_n as follows:

$$B(X, Y) = \frac{1}{2} \operatorname{tr}(XY). \quad (2.1)$$

Let $\{X_{i,j}\}_{i,j \in [n]}$ be the generators of \mathfrak{g} given by

$$X_{i,j} = \epsilon_j E_{i,j} - \epsilon_i E_{j,i} \quad (2.2)$$

for $i, j = 1, 2, \dots, n$, where $E_{i,j}$ stands for the matrix unit and

$$\epsilon_i = \begin{cases} 1 & \text{if } i \in [p], \\ -1 & \text{if } i \in p + [q]. \end{cases} \quad (2.3)$$

Then one sees $X_{j,i} = -X_{i,j}$ and

$$\mathfrak{k} = \bigoplus_{\substack{i,j \in [p] \\ i < j}} \mathbb{C}X_{i,j} \oplus \bigoplus_{\substack{i,j \in p+[q] \\ i < j}} \mathbb{C}X_{i,j}.$$

In addition, if one sets

$$\mathfrak{p} = \bigoplus_{\substack{i \in [p] \\ j \in p+[q]}} \mathbb{C}X_{i,j},$$

then $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ is the complexified Cartan decomposition. Finally, the dual generators $\{X_{i,j}^\vee\}_{i,j}$ to $\{X_{i,j}\}_{i,j}$ with respect to B , i.e., those satisfying $B(X_{i,j}, X_{k,l}^\vee) = \delta_{i,k} \delta_{j,l}$, are given by

$$X_{i,j}^\vee = \begin{cases} -X_{i,j} & \text{if } i, j \in [p], \text{ or if } i, j \in p + [q], \\ X_{i,j} & \text{otherwise.} \end{cases} \quad (2.4)$$

Let V denote the vector space \mathbb{R}^{p+q} equipped with a quadratic form η defined by

$$\eta(v) = {}^t v I_{p,q} v \quad (v \in \mathbb{R}^{p+q}).$$

Then G acts on V by matrix multiplication, leaving η invariant. It is well known that the space $\mathbb{C}[V]$ of polynomial functions on V decomposes as follows (see e.g. [4]): for $v = {}^t(x_1, \dots, x_p, y_1, \dots, y_q) \in V$, one sees that

$$\begin{aligned} \mathbb{C}[V] &\simeq \mathbb{C}[x_1, \dots, x_p] \otimes \mathbb{C}[y_1, \dots, y_q] \\ &\simeq \bigoplus_{k=0}^{\infty} \left(\mathbb{C}[r_x^2] \otimes \mathcal{H}^k(\mathbb{R}^p) \right) \otimes \bigoplus_{l=0}^{\infty} \left(\mathbb{C}[r_y^2] \otimes \mathcal{H}^l(\mathbb{R}^q) \right) \\ &\simeq \bigoplus_{k,l=0}^{\infty} \mathcal{H}^k(\mathbb{R}^p) \otimes \mathcal{H}^l(\mathbb{R}^q) \otimes \mathbb{C}[r_x^2, r_y^2], \end{aligned} \quad (2.5)$$

where $r_x^2 = x_1^2 + \dots + x_p^2$, $r_y^2 = y_1^2 + \dots + y_q^2$ and $\mathcal{H}^k(\mathbb{R}^p)$ denotes the space of homogeneous harmonic polynomials on \mathbb{R}^p of degree k and so on.

Taking (2.5) into account, let us introduce our space \mathcal{E} of functions on V by

$$\mathcal{E} := \bigoplus_{k,l=0}^{\infty} \mathcal{H}^k(\mathbb{R}^p) \otimes \mathcal{H}^l(\mathbb{R}^q) \otimes \mathbb{C}[[r_x^2, r_y^2]], \quad (2.6)$$

where $\mathbb{C}[[r_x^2, r_y^2]]$ denotes the ring of formal power series in r_x^2 and r_y^2 , and the direct sum is algebraic; we assume that the radial part should be a formal power series rather than a polynomial so that we can find a highest weight vector or a lowest weight vector with respect to the \mathfrak{sl}_2 -action (2.10) given below.

Definition 2.1. For an element $f \in \mathcal{E}$ of the form $f = h_1(x)h_2(y)\phi(r_x^2, r_y^2)$ with $h_1 \in \mathcal{H}^k(\mathbb{R}^p)$, $h_2 \in \mathcal{H}^l(\mathbb{R}^q)$ and $\phi(r_x^2, r_y^2) \in \mathbb{C}[[r_x^2, r_y^2]]$, we set $\kappa_+ = \kappa_+(f) = k + p/2$, $\kappa_- = \kappa_-(f) = l + q/2$, and

$$\kappa(f) = (\kappa_+, \kappa_-) = \left(k + \frac{p}{2}, l + \frac{q}{2}\right), \quad (2.7)$$

which we call the K -type of $f = h_1(x)h_2(y)\phi(r_x^2, r_y^2)$ in this paper by abuse.

Furthermore, for non-negative integers $k, l \geq 0$ given, we say that an element of \mathcal{E} is K -homogeneous with K -type $(\kappa_+, \kappa_-) = (k + p/2, l + q/2)$ if it is a linear combination of elements whose K -types are all equal to (κ_+, κ_-) .

Denoting the ring of polynomial coefficient differential operators on V by $\mathcal{PD}(V)$, let us define a representation $\pi : U(\mathfrak{g}) \rightarrow \mathcal{PD}(V)$ by

$$\pi(X_{i,j}) = \begin{cases} x_i \partial_{x_j} - x_j \partial_{x_i} & \text{if } i, j \in [p], \\ -y_{i'} \partial_{y_{j'}} + y_{j'} \partial_{y_{i'}} & \text{if } i, j \in p + [q], \\ -\sqrt{-1} (x_i y_{j'} + \partial_{x_i} \partial_{y_{j'}}) & \text{if } i \in [p], j \in p + [q], \\ \sqrt{-1} (x_j y_{i'} + \partial_{x_j} \partial_{y_{i'}}) & \text{if } i \in p + [q], j \in [p], \end{cases} \quad (2.8)$$

where we set $i' := i - p$, $i \in p + [q]$, for short. It is easy to verify that π indeed defines a representation of $U(\mathfrak{g})$. It is also clear that the action of \mathfrak{k}_0 lifts to the one of K , i.e., if $X \in \mathfrak{k}_0$ then

$$(\pi(X)f)(v) = \left. \frac{d}{dt} \right|_{t=0} f(\exp(-tX)v) \quad (f \in \mathcal{E}, v \in V). \quad (2.9)$$

Remark 2.2. It was shown in [6] that if one regards V as a Lagrangian subspace of the symplectic vector space $((\mathbb{C}^{p+q})_{\mathbb{R}}, \omega)$ given by

$$V = \langle e_1, e_2, \dots, e_p, \sqrt{-1} e_{p+1}, \sqrt{-1} e_{p+2}, \dots, \sqrt{-1} e_{p+q} \rangle_{\mathbb{R}},$$

where $e_i = {}^i(0, \dots, 1, \dots, 0) \in \mathbb{C}^{p+q}$ and

$$\omega(z, w) = \text{Im}(z^* I_{p,q} w) \quad (z, w \in \mathbb{C}^{p+q}),$$

then one obtains the representation π given in (2.8) via the canonical quantization of the moment map on $((\mathbb{C}^{p+q})_{\mathbb{R}}, \omega)$.

Define elements H, X^+, X^- of $\mathcal{PD}(V)$ by

$$H = -E_x - \frac{p}{2} + E_y + \frac{q}{2}, \quad X^+ = -\frac{1}{2}(\Delta_x + r_y^2), \quad X^- = \frac{1}{2}(r_x^2 + \Delta_y), \quad (2.10)$$

where

$$\begin{aligned} E_x &= \sum_{i \in [p]} x_i \partial_{x_i}, & r_x^2 &= \sum_{i \in [p]} x_i^2, & \Delta_x &= \sum_{i \in [p]} \partial_{x_i}^2, \\ E_y &= \sum_{j \in [q]} y_j \partial_{y_j}, & r_y^2 &= \sum_{j \in [q]} y_j^2, & \Delta_y &= \sum_{j \in [q]} \partial_{y_j}^2. \end{aligned} \quad (2.11)$$

Then they satisfy that

$$[H, X^+] = 2X^+, \quad [H, X^-] = -2X^-, \quad [X^+, X^-] = H.$$

Namely, $\mathfrak{g}' := \mathbb{C}\text{-span}\{H, X^+, X^-\}$ forms a Lie subalgebra of $\mathcal{PD}(V)$ isomorphic to \mathfrak{sl}_2 . Moreover, \mathfrak{g}' forms the commutant of \mathfrak{g} in $\mathcal{PD}(V)$:

$$[Y, Z] = 0 \quad \text{for all } Y \in \pi(\mathfrak{g}) \text{ and } Z \in \mathfrak{g}'.$$

In view of the fact that $(O(p, q), \text{SL}(2, \mathbb{R}))$ is a dual pair in $\text{Sp}(n, \mathbb{R})$, we introduce (\mathfrak{g}, K) -modules of $O(p, q)$ associated to the finite-dimensional representation of \mathfrak{sl}_2 as follows.

Definition 2.3 ([6]). Given $m \in \mathbb{N}$, we define (\mathfrak{g}, K) -modules $M^\pm(m)$ by

$$M^+(m) := \{f \in \mathcal{E}; Hf = mf, X^+f = 0, (X^-)^{m+1}f = 0\},$$

$$M^-(m) := \{f \in \mathcal{E}; Hf = -mf, X^-f = 0, (X^+)^{m+1}f = 0\}.$$

Note that, by definition, each $f \in M^+(m)$ (resp. $f \in M^-(m)$) is a highest weight vector (resp. lowest weight vector) with respect to the $\mathfrak{g}' = \mathfrak{sl}_2$ action. It is easy to see that $M^+(m)$ is isomorphic to $M^-(m)$ for any $m \in \mathbb{N}$, though $M^+(0) = M^-(0)$.

In order to describe elements of $M^\pm(m)$ explicitly, let us introduce a convergent power series Ψ_α on \mathbb{C} for $\alpha \in \mathbb{C} \setminus (-\mathbb{N})$ by

$$\Psi_\alpha(u) = \sum_{j=0}^{\infty} \frac{(-1)^j}{j! (\alpha)_j} u^j \quad (u \in \mathbb{C})$$

with $(\alpha)_j = \Gamma(\alpha + j)/\Gamma(\alpha) = \alpha(\alpha + 1) \cdots (\alpha + j - 1)$. Note that it is related to the Bessel function J_ν of the first kind. In fact, one has

$$\Psi_\alpha(u) = \Gamma(\alpha) u^{-(\alpha-1)/2} J_{\alpha-1}(2u^{1/2}). \quad (2.12)$$

Now we set

$$\psi_\alpha := \Psi_\alpha(r_x^2 r_y^2 / 4) \quad (2.13)$$

for brevity. We will find it convenient to write $\rho_x := r_x^2/2$, $\rho_y := r_y^2/2$. Then a typical element $f \in M^+(m)$ is given by

$$f = h_1(x) h_2(y) \rho_y^{\mu_-} \psi_{\kappa_+} \quad (2.14)$$

with $h_1 \in \mathcal{H}^k(\mathbb{R}^p)$, $h_2 \in \mathcal{H}^l(\mathbb{R}^q)$ and $\mu_- = \frac{1}{2}(m + \kappa_+ - \kappa_-) \in \mathbb{N}$, while a typical one $f \in M^-(m)$ is given by

$$f = h_1(x) h_2(y) \rho_x^{\mu_+} \psi_{\kappa_-} \quad (2.15)$$

with $h_1 \in \mathcal{H}^k(\mathbb{R}^p)$, $h_2 \in \mathcal{H}^l(\mathbb{R}^q)$ and $\mu_+ = \frac{1}{2}(m - \kappa_+ + \kappa_-) \in \mathbb{N}$ (see [6, Proposition 3.2]). A general element of $M^+(m)$ (resp. $M^-(m)$) is a linear combination of these elements given in (2.14) (resp. (2.15)).

Remark 2.4. All K -homogeneous elements of $M^+(m)$ with K -type (κ_+, κ_-) have the same radial part $\rho_y^{\mu_-} \psi_{\kappa_+} \in \mathbb{C}[[r_x^2, r_y^2]]$ with $\mu_- = \frac{1}{2}(m + \kappa_+ - \kappa_-)$. Similarly, all K -homogeneous elements of $M^-(m)$ with K -type (κ_+, κ_-) have the same radial part $\rho_x^{\mu_+} \psi_{\kappa_-}$ with $\mu_+ = \frac{1}{2}(m - \kappa_+ + \kappa_-)$.

Now we collect a few facts from [6] that we need in order to prove our main result.

First, we recall the action of \mathfrak{p} on $M^\pm(m)$ ([6, Lemma 4.1]). Let $X_{i,p+j}$, $i \in [p]$, $j \in [q]$, be a basis of \mathfrak{p} given in (2.2). Then, for $f = h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} \in M^+(m)$ with $\kappa(f) = (\kappa_+, \kappa_-)$ and $\mu_- = \frac{1}{2}(m + \kappa_+ - \kappa_-)$, one sees that

$$\begin{aligned} -\sqrt{-1} \pi(X_{i,p+j})f &= \frac{\kappa_- + \mu_- - 1}{\kappa_- - 1} (\partial_{x_i} h_1) (\partial_{y_j} h_2) \rho_y^{\mu_-} \psi_{\kappa_+ - 1} \\ &\quad + \mu_- (\partial_{x_i} h_1) (y_j h_2)^\dagger \rho_y^{\mu_- - 1} \psi_{\kappa_+ - 1} \\ &\quad + \frac{\kappa_+ - \kappa_- - \mu_-}{\kappa_+ (\kappa_- - 1)} (x_i h_1)^\dagger (\partial_{y_j} h_2) \rho_y^{\mu_- + 1} \psi_{\kappa_+ + 1} \\ &\quad + \frac{\kappa_+ - \mu_- - 1}{\kappa_+} (x_i h_1)^\dagger (y_j h_2)^\dagger \rho_y^{\mu_-} \psi_{\kappa_+ + 1}, \end{aligned} \quad (2.16a)$$

and, for $f = h_1 h_2 \rho_x^{\mu_+} \psi_{\kappa_-} \in M^-(m)$ with $\kappa(f) = (\kappa_+, \kappa_-)$ and $\mu_+ = \frac{1}{2}(m - \kappa_+ + \kappa_-)$,

$$\begin{aligned} -\sqrt{-1} \pi(X_{i,p+j})f &= \frac{\kappa_+ + \mu_+ - 1}{\kappa_+ - 1} (\partial_{x_i} h_1) (\partial_{y_j} h_2) \rho_x^{\mu_+} \psi_{\kappa_- - 1} \\ &\quad + \frac{\kappa_- - \kappa_+ - \mu_+}{\kappa_- (\kappa_+ - 1)} (\partial_{x_i} h_1) (y_j h_2)^\dagger \rho_x^{\mu_+ + 1} \psi_{\kappa_- + 1} \\ &\quad + \mu_+ (x_i h_1)^\dagger (\partial_{y_j} h_2) \rho_x^{\mu_+ - 1} \psi_{\kappa_- - 1} \\ &\quad + \frac{\kappa_- - \mu_+ - 1}{\kappa_-} (x_i h_1)^\dagger (y_j h_2)^\dagger \rho_x^{\mu_+} \psi_{\kappa_- + 1}. \end{aligned} \quad (2.16b)$$

Here, in general, for a homogeneous polynomial $P = P(x_1, \dots, x_n)$ on \mathbb{R}^n of degree d , we set

$$P^\dagger := P - \frac{\rho}{2d + n - 4} \Delta P,$$

with $\Delta = \sum_{i=1}^n \partial_{x_i}^2$ and $\rho = (1/2) \sum_{i=1}^n x_i^2$. Note that if $\Delta^2 P = 0$ then P^\dagger is harmonic and that if $h = h(x_1, \dots, x_n)$ is harmonic then $\Delta(x_i h) = 2\partial_{x_i} h$ and $\Delta^2(x_i h) = 0$.

Next, the following facts describe the K -types of $M^\pm(m)$ and when they are irreducible ([6, Theorem 4.6]).

Facts 2.5. *Assume that $p \geq 1$, $q \geq 1$ and $p + q \in 2\mathbb{N}$. Let $m \in \mathbb{N}$ be a non-negative integer satisfying $m + 3 \leq (p + q)/2$. Then one has the following.*

(1) *The K -type formula of $M^\pm(m)$ is given by*

$$M^\pm(m)|_K \simeq \bigoplus_{\substack{k, l \geq 0 \\ k + \frac{p}{2} - (l + \frac{q}{2}) \in \Lambda_m}} \mathcal{H}^k(\mathbb{R}^p) \otimes \mathcal{H}^l(\mathbb{R}^q), \quad (2.17)$$

where $\Lambda_m = \{-m, -m + 2, -m + 4, \dots, m - 2, m\}$.

(2) *Suppose further that $p, q \geq 2$. Then $M^\pm(m)$ are irreducible (\mathfrak{g}, K) -modules.*

Note in particular that if the K -type of $f \in M^\pm(m)$ is $\kappa(f) = (\kappa_+, \kappa_-)$, then it follows from (2.17) that

$$|\kappa_+ - \kappa_-| \leq m. \quad (2.18)$$

Henceforth in the rest of the paper, we assume that

$$p \geq 2, q \geq 2, p + q \in 2\mathbb{N} \quad \text{and} \quad m + 3 \leq \frac{p + q}{2} \quad (2.19)$$

so that $M^\pm(m)$ is irreducible.

3. ANNIHILATOR OF $M^\pm(m)$

In order to translate objects in terms of \mathfrak{o}_n into ones in terms of \mathfrak{g} , and vice versa, let us fix an isomorphism Φ between \mathfrak{g} and \mathfrak{o}_n given by

$$\Phi : \mathfrak{g} \xrightarrow{\sim} \mathfrak{o}_n, \quad X \mapsto I_{p,q}^{1/2} X I_{p,q}^{-1/2}, \quad (3.1)$$

where $I_{p,q}^{1/2} = \text{diag}(1, \dots, 1, \sqrt{-1}, \dots, \sqrt{-1})$ and $I_{p,q}^{-1/2}$ is its inverse.

Let $\{M_{i,j}\}$ be generators of \mathfrak{o}_n given by

$$M_{i,j} = E_{i,j} - E_{j,i} \quad (3.2)$$

for $i, j = 1, 2, \dots, n$. Then, one has

$$\Phi(X_{i,j}) = \begin{cases} M_{i,j} & \text{if } i, j \in [p], \\ -M_{i,j} & \text{if } i, j \in p + [q], \\ \sqrt{-1} M_{i,j} & \text{otherwise.} \end{cases} \quad (3.3)$$

We extend Φ to the isomorphism from $\bigoplus_{j=0}^{\infty} T^j(\mathfrak{g})$ onto $\bigoplus_{j=0}^{\infty} T^j(\mathfrak{o}_n)$, where $T^j(\mathfrak{g}) := \mathfrak{g} \otimes \dots \otimes \mathfrak{g}$ (j factors) etc..

Let $U(\mathfrak{g})$ denote the universal enveloping algebra of \mathfrak{g} , and $U_j(\mathfrak{g})$ its subspace spanned by products of at most j elements of \mathfrak{g} , with $U_{-1}(\mathfrak{g}) = 0$. It is well known that the associated graded algebra $\text{gr } U(\mathfrak{g}) = \bigoplus_{j=0}^{\infty} U_j(\mathfrak{g})/U_{j-1}(\mathfrak{g})$ is isomorphic to the symmetric algebra $S(\mathfrak{g}) = \bigoplus_{j=0}^{\infty} S^j(\mathfrak{g})$. Let $\sigma : \text{gr } U(\mathfrak{g}) \rightarrow S(\mathfrak{g})$ be the algebra isomorphism, and σ_j its j -th piece:

$$\sigma_j : U_j(\mathfrak{g})/U_{j-1}(\mathfrak{g}) \xrightarrow{\sim} S^j(\mathfrak{g}). \quad (3.4)$$

Let $\gamma : S(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ be the linear isomorphism called *symmetrization*, and γ_j its j -th piece

$$\gamma_j : S^j(\mathfrak{g}) \rightarrow U(\mathfrak{g}), \quad (3.5)$$

which is an injective linear map onto a vector-space complement to $U_{j-1}(\mathfrak{g})$ in $U_j(\mathfrak{g})$. Then the following diagram is commutative:

$$\begin{array}{ccc} & & U_j(\mathfrak{g}) \\ & \nearrow^{1:1} & \downarrow p_j \\ S^j(\mathfrak{g}) & \xleftarrow{\sim \sigma_j} & U_j(\mathfrak{g})/U_{j-1}(\mathfrak{g}), \end{array}$$

where p_j is the canonical projection. In particular, $(\sigma_j \circ p_j \circ \gamma_j)(v) = v$ for all $v \in S^j(\mathfrak{g})$.

In what follows, given a finite-dimensional vector space U over \mathbb{C} , we will identify $S^j(U)$ (resp. $\wedge^j U$) with the subspace of symmetric (resp. alternating) tensors in $U^{\otimes j}$.

For $u \in U(\mathfrak{g})$, we sometimes denote $\pi(u)$ by \hat{u} for brevity. Let $\Omega_{\mathfrak{g}}$, $\Omega_{\mathfrak{o}_p}$ and $\Omega_{\mathfrak{o}_q}$ be the Casimir elements of \mathfrak{g} , \mathfrak{o}_p and \mathfrak{o}_q respectively¹, which are given by

$$\Omega_{\mathfrak{g}} = \sum_{\substack{i,j \in [n] \\ i < j}} X_{i,j} X_{i,j}^{\vee}, \quad \Omega_{\mathfrak{o}_p} = \sum_{\substack{i,j \in [p] \\ i < j}} X_{i,j} X_{j,i}, \quad \Omega_{\mathfrak{o}_q} = \sum_{\substack{i,j \in [q] \\ i < j}} X_{i,j} X_{j,i}. \quad (3.6)$$

¹We regard \mathfrak{o}_p and \mathfrak{o}_q as subalgebras of $\mathfrak{k} \simeq \mathfrak{o}_p \oplus \mathfrak{o}_q$ canonically.

Then the corresponding operators represented in $\mathcal{PD}(V)$ are given by

$$\begin{aligned} \hat{\Omega}_{\mathfrak{g}} &= (E_x - E_y)^2 + (p - q)(E_x - E_y) - 2(E_x + E_y) \\ &\quad - \left(r_x^2 r_y^2 + r_x^2 \Delta_x + r_y^2 \Delta_y + \Delta_x \Delta_y \right) - pq, \end{aligned} \quad (3.7a)$$

$$\hat{\Omega}_{\mathfrak{o}_p} = E_x^2 + (p - 2)E_x - r_x^2 \Delta_x, \quad (3.7b)$$

$$\hat{\Omega}_{\mathfrak{o}_q} = E_y^2 + (q - 2)E_y - r_y^2 \Delta_y. \quad (3.7c)$$

It is well known that the Casimir operator $\hat{\Omega}_{\mathfrak{g}'}$ of $\mathfrak{g}' = \mathfrak{sl}_2$ given by

$$\hat{\Omega}_{\mathfrak{g}'} = H^2 + 2(X^+ X^- + X^- X^+)$$

satisfies the relation

$$\hat{\Omega}_{\mathfrak{g}} = \hat{\Omega}_{\mathfrak{g}'} - \frac{1}{4}(p + q)^2 + (p + q)$$

(see [7, 11]), which implies that

$$\hat{\Omega}_{\mathfrak{g}}|_{M^\pm(m)} = m(m + 2) - \frac{1}{4}(p + q)^2 + (p + q) \quad (3.8)$$

since $\hat{\Omega}_{\mathfrak{g}'}$ acts on $M^\pm(m)$ by the scalar $m(m + 2)$.

Let α denote the highest root of \mathfrak{g} relative to any positive root system. Then $S^2(\mathfrak{g})$ decomposes as $S^2(\mathfrak{g}) \simeq V_{2\alpha} \oplus W$, where $V_{2\alpha}$ is the irreducible \mathfrak{g} -module with highest weight 2α and W is the \mathfrak{g} -invariant complement of $V_{2\alpha}$ (see (A.8) below).

Theorem 3.1 (Garfinkle). *Let \mathcal{I} be an ideal of infinite codimension in $U(\mathfrak{g})$ and set $I = \bigoplus_{j=0}^{\infty} (\mathcal{I} \cap U_j(\mathfrak{g})) / (\mathcal{I} \cap U_{j-1}(\mathfrak{g})) \subset \text{gr } U(\mathfrak{g})$. Then \mathcal{I} is the Joseph ideal if and only if $\sigma(I) \cap S^2(\mathfrak{g}) = W$.*

In fact, it is shown that Theorem 3.1 holds true for all complex simple Lie algebras not of type A in [3, 2].

Using the notation in Appendix A below, the \mathfrak{g} -invariant complement W is given by

$$W = E_{\emptyset}^{\mathfrak{g}} \oplus E_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}^{\mathfrak{g}} \oplus E_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{\mathfrak{g}}, \quad (3.9)$$

where

$$E_{\lambda}^{\mathfrak{g}} = \Phi^{-1}(E_{\lambda}^{\mathfrak{o}_n}) \quad (\lambda \in \{\emptyset, \begin{smallmatrix} \square \\ \square \end{smallmatrix}, \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}\}). \quad (3.10)$$

The following lemma is trivial, and its proof is left to the reader.

Lemma 3.2. *Define an element $Q \in T^2(\mathfrak{g})$ by $Q = \sum_{i,j} X_{i,j} \otimes X_{i,j}^{\vee}$. Then Q is a symmetric tensor and satisfies $Q = \Phi^{-1}(Q^{\mathfrak{o}_n})$ and $\gamma_2(Q) = 2\Omega_{\mathfrak{g}}$.*

Following [1], let us introduce an element $\Xi^{\mathfrak{o}_n} \in S^2(\mathfrak{o}_n)$, or $\Xi \in S^2(\mathfrak{g})$, that plays a critical role in the proof of our main result:

$$\Xi^{\mathfrak{o}_n} := \frac{1}{2} \sum_{i,j} \eta^{ij} S_{ij}^{\mathfrak{o}_n} = \frac{1}{2} \left(\sum_{i \in [p]} S_{ii}^{\mathfrak{o}_n} - \sum_{i \in p+[q]} S_{ii}^{\mathfrak{o}_n} \right), \quad (3.11)$$

where η^{ij} denotes the (i, j) -th entry of $I_{p,q}^{-1} = I_{p,q}$.

Lemma 3.3. *Set $\Xi := \Phi^{-1}(\Xi^{0n}) \in S^2(\mathfrak{g})$. Then it satisfies that*

$$\gamma_2(\Xi) = \Omega_{\mathfrak{v}_p} - \Omega_{\mathfrak{v}_q} - \frac{p-q}{p+q}\Omega_{\mathfrak{g}}. \quad (3.12)$$

Proof. Since $S_{ii}^{0n} = \sum_k M_{i,k} \otimes M_{k,i} - \frac{1}{n}Q^{0n}$, one sees

$$\begin{aligned} \Xi^{0n} &= \frac{1}{2} \sum_k \left(\sum_{i \in [p]} - \sum_{i \in p+[q]} \right) M_{i,k} \otimes M_{k,i} - \frac{p-q}{2n} Q^{0n} \\ &= \frac{1}{2} \left(\sum_{i,k \in [p]} + \sum_{\substack{i \in [p] \\ k \in p+[q]}} \right) M_{i,k} \otimes M_{k,i} - \frac{1}{2} \left(\sum_{\substack{i \in p+[q] \\ k \in [p]}} + \sum_{i,k \in p+[q]} \right) M_{i,k} \otimes M_{k,i} - \frac{p-q}{2n} Q^{0n} \\ &= \frac{1}{2} \left(\sum_{i,k \in [p]} - \sum_{i,k \in p+[q]} \right) M_{i,k} \otimes M_{k,i} - \frac{p-q}{2n} Q^{0n}. \end{aligned}$$

Therefore, it follows from (3.3) that

$$\begin{aligned} \Xi &= \frac{1}{2} \left(\sum_{i,k \in [p]} X_{i,k} \otimes X_{k,i} - \sum_{i,k \in p+[q]} (-X_{i,k}) \otimes (-X_{k,i}) \right) - \frac{p-q}{2n} \Phi^{-1}(Q^{0n}) \\ &= \sum_{\substack{i,k \in [p] \\ i < k}} X_{i,k} \otimes X_{k,i} - \sum_{\substack{i,k \in p+[q] \\ i < k}} X_{i,k} \otimes X_{k,i} - \frac{p-q}{2n} Q. \end{aligned} \quad (3.13)$$

Now, the result immediately follows from Lemma 3.2. \square

Proposition 3.4. *For a K -homogeneous $f \in M^\pm(m)$ with $\kappa(f) = (\kappa_+, \kappa_-)$, one has*

$$\hat{\Omega}_{\mathfrak{v}_p} f = \left((\kappa_+ - 1)^2 - \frac{(p-2)^2}{4} \right) f, \quad (3.14)$$

$$\hat{\Omega}_{\mathfrak{v}_q} f = \left((\kappa_- - 1)^2 - \frac{(q-2)^2}{4} \right) f. \quad (3.15)$$

Proof. We only show the case of $M^+(m)$ here. The other case can be proved exactly in the same manner.

First, we show (3.14). It suffices to prove it for a typical element $f = h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} \in M^+(m)$ with $h_1 \in \mathcal{H}^k(\mathbb{R}^p)$, $h_2 \in \mathcal{H}^l(\mathbb{R}^q)$ and $\mu_- = (1/2)(m + \kappa_+ - \kappa_-)$, where $(\kappa_+, \kappa_-) = (k + \frac{p}{2}, l + \frac{q}{2})$. Recall that for a homogeneous polynomial $h = h(x_1, \dots, x_n)$ on \mathbb{R}^n of degree d and for a smooth function $\phi(u)$ in a single variable u , we have

$$\Delta(h\phi(\rho)) = (2d+n)h\phi'(\rho) + 2h\rho\phi''(\rho), \quad (3.16)$$

where $\Delta = \sum_{i=1}^n \partial_{x_i}^2$ and $\rho = (1/2) \sum_{i=1}^n x_i^2$ (cf. [6, Lemma 3.2]). Thus, for $f = h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+}$, we have

$$\begin{aligned} \Delta_x f &= h_2 \rho_y^{\mu_-} \Delta_x (h_1 \Psi_{\kappa_+}(\rho_x \rho_y)) \\ &= (2k+p)h_1 h_2 \rho_y^{\mu_-+1} \psi'_{\kappa_+} + 2h_1 h_2 \rho_x \rho_y^{\mu_-+2} \psi''_{\kappa_+}, \end{aligned}$$

where we set $\psi'_{\kappa_+} = \Psi'(\rho_x \rho_y)$, $\psi''_{\kappa_+} = \Psi''(\rho_x \rho_y)$. Similarly, one sees that

$$E_x f = k h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} + 2h_1 h_2 \rho_x \rho_y^{\mu_-+1} \psi'_{\kappa_+},$$

$$\begin{aligned} E_x^2 f &= k^2 h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} + 4(k+1) h_1 h_2 \rho_x \rho_y^{\mu_-+1} \psi'_{\kappa_+} \\ &\quad + 4h_1 h_2 \rho_x^2 \rho_y^{\mu_-+2} \psi''_{\kappa_+}. \end{aligned}$$

Therefore,

$$\begin{aligned} \hat{\Omega}_{0,p} f &= (E_x^2 + (p-2)E_x - 2\rho_x \Delta_x) h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} \\ &= (k^2 + (p-2)k) h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} \\ &= \left(\left(k + \frac{p-2}{2} \right)^2 - \frac{(p-2)^2}{4} \right) f. \end{aligned}$$

Now, substituting $\kappa_+ = k + p/2$, one obtains (3.14).

Next, we turn to (3.15). Similar calculations yield

$$\begin{aligned} \Delta_y f &= h_1 \Delta_x (h_2 \rho_y^{\mu_-} \Psi_{\kappa_+}(\rho_x \rho_y)) \\ &= ((2l+q)\mu_- + 2\mu_-(\mu_- - 1)) h_1 h_2 \rho_y^{\mu_- - 1} \psi_{\kappa_+} \\ &\quad + ((2l+q) + 4\mu_-) h_1 h_2 \rho_x \rho_y^{\mu_-} \psi'_{\kappa_+} + 2h_1 h_2 \rho_x^2 \rho_y^{\mu_-+1} \psi''_{\kappa_+}, \\ E_y f &= (l + 2\mu_-) h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} + 2h_1 h_2 \rho_x \rho_y^{\mu_-+1} \psi'_{\kappa_+}, \\ E_y^2 f &= (l + 2\mu_-)^2 h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+} + 4(l + 2\mu_- + 1) h_1 h_2 \rho_x \rho_y^{\mu_-+1} \psi'_{\kappa_+} \\ &\quad + 4h_1 h_2 \rho_x^2 \rho_y^{\mu_-+2} \psi''_{\kappa_+}, \end{aligned}$$

and

$$\hat{\Omega}_{0,q} f = ((l + 2\mu_-)^2 + (q-2)(l + 2\mu_-) - 2\mu_-(2l + q + 2\mu_- - 2)) f$$

for $f = h_1 h_2 \rho_y^{\mu_-} \psi_{\kappa_+}$. Substituting $\kappa_- = l + q/2$, $\mu_- = (1/2)(m + \kappa_+ - \kappa_-)$, one obtains (3.15). \square

Corollary 3.5. *Set $\hat{\Xi} := \pi(\gamma_2(\Xi)) \in \mathcal{PD}(V)$. Then, for a K -homogeneous $f \in M^\pm(m)$ with $\kappa(f) = (\kappa_+, \kappa_-)$, one has $\hat{\Xi}f = \lambda_{\kappa(f)} f$, where $\lambda_{\kappa(f)}$ is a scalar given by*

$$\lambda_{\kappa(f)} = (\kappa_+ - \kappa_-)(\kappa_+ + \kappa_- - 2) - \frac{p-q}{p+q} m(m+2). \quad (3.17)$$

In particular, $\lambda_{\kappa(f)} = 0$ if $m = 0$.

Proof. The first statement immediately follows from Lemma 3.3 and Proposition 3.4. As for the second, we note that if $m = 0$ then $\kappa_+ = \kappa_-$ by (2.18). \square

In what follows, let $\mathcal{I}^\pm(m)$ denote the annihilator in $U(\mathfrak{g})$ of our (\mathfrak{g}, K) -module $M^\pm(m)$ for short:

$$\mathcal{I}^\pm(m) := \text{Ann}_{U(\mathfrak{g})} M^\pm(m).$$

Correspondingly, set $I^\pm(m) := \bigoplus_{j=0}^{\infty} (\mathcal{I}^\pm(m) \cap U_j(\mathfrak{g})) / (\mathcal{I}^\pm(m) \cap U_{j-1}(\mathfrak{g}))$.

Theorem 3.6. *The annihilator $\mathcal{I}^\pm(m)$ is the Joseph ideal if and only if $m = 0$.*

Proof. Recall that our \mathfrak{g} -invariant complement $W \subset S^2(\mathfrak{g})$ is given by

$$W = E_{\emptyset}^{\mathfrak{g}} \oplus E_{\begin{array}{c} \square \\ \square \\ \square \end{array}}^{\mathfrak{g}} \oplus E_{\begin{array}{c} \square \\ \square \end{array}}^{\mathfrak{g}}.$$

We must show that $W = \sigma(I^\pm(m)) \cap S^2(\mathfrak{g})$ if and only if $m = 0$ by Theorem 3.1. However, the \mathfrak{g} -invariance of both sides reduces the problem to showing that $W \subset \sigma(I^\pm(m)) \cap S^2(\mathfrak{g})$ if and only if $m = 0$.

Since $\Omega_{\mathfrak{g}} \in U_2(\mathfrak{g})$ acts on $M^\pm(m)$ by the scalar given by (3.8), say λ_0 ,

$$Q = \sigma_2(p_2(2(\Omega_{\mathfrak{g}} - \lambda_0))) \in \sigma(I^\pm(m)),$$

and hence we have

$$E_{\emptyset}^{\mathfrak{g}} \subset \sigma(I^\pm(m)) \cap S^2(\mathfrak{g}). \quad (3.18)$$

Next, a simple calculation shows that $\pi(\Phi^{-1}(S_{ijkl}^{\mathfrak{g}_n})) = 0$ in $\mathcal{PD}(V)$ for all $i < j < k < l$. Namely, we have

$$E_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}^{\mathfrak{g}} \subset \sigma(I^\pm(m)) \cap S^2(\mathfrak{g}). \quad (3.19)$$

Finally, we show that

$$E_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}^{\mathfrak{g}} \subset \sigma(I^\pm(m)) \cap S^2(\mathfrak{g}) \quad \text{if and only if} \quad m = 0. \quad (3.20)$$

Since $E_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}^{\mathfrak{g}}$ is irreducible, in order that (3.20) holds, it suffices to show

$$\Xi \in \sigma(I^\pm(m)) \cap S^2(\mathfrak{g}) \quad \text{if and only if} \quad m = 0. \quad (3.21)$$

Assume that Ξ is in $\sigma(I^\pm(m)) \cap S^2(\mathfrak{g})$. Then there exists an element $u \in \mathcal{S}^\pm(m) \cap U_2(\mathfrak{g})$ satisfying $\Xi = \sigma_2(p_2(u))$ and

$$\pi(u)f = 0 \quad \text{for all } f \in M^\pm(m). \quad (3.22)$$

Since $\Xi = \sigma_2(p_2(\gamma_2(\Xi)))$, there exists an element $u_1 \in U_1(\mathfrak{g})$ such that $\gamma_2(\Xi) = u + u_1$. Moreover, u_1 can be written as $u_1 = \lambda + Y$ with $\lambda \in \mathbb{C}$ and $Y \in \mathfrak{g}$ since $U_1(\mathfrak{g}) = \mathbb{C} \oplus \mathfrak{g}$. Now, it follows from (3.22) and Corollary 3.5 that

$$\begin{aligned} \pi(Y)f &= (\lambda_{\kappa(f)} - \lambda)f \\ &= \left((\kappa_+ - \kappa_-)(\kappa_+ + \kappa_- + 2) - \frac{p-q}{p+q}m(m+2) - \lambda \right) f \end{aligned} \quad (3.23)$$

for all K -homogeneous $f \in M^\pm(m)$ with $\kappa(f) = (\kappa_+, \kappa_-)$. In view of the right-hand side of (3.23), the \mathfrak{p} -part action given in (2.16) forces this Y to be in \mathfrak{k} . Thus, letting $Y = Y_1 + Y_2$, $Y_1 \in \mathfrak{o}_p$, $Y_2 \in \mathfrak{o}_q$, be the decomposition corresponding to $\mathfrak{k} = \mathfrak{o}_p \oplus \mathfrak{o}_q$, one sees that (3.23) is equivalent to the fact that $\pi(Y_1)$ acts on $\mathcal{H}^k(\mathbb{R}^p)$ by a scalar and $\pi(Y_2)$ acts on $\mathcal{H}^l(\mathbb{R}^q)$ by a scalar, where $\kappa_+ = k + \frac{p}{2}$ and $\kappa_- = l + \frac{q}{2}$. If $m = 0$ then $\Xi \in \sigma(I^\pm(m))$; in fact, one can take $Y = 0$ and $\lambda = 0$ since $\lambda_{\kappa(f)} = 0$. On the other hand, if $m \neq 0$, this is impossible, and hence $\Xi \notin \sigma(I^\pm(m))$. Therefore, (3.21) holds. \square

APPENDIX A. IRREDUCIBLE DECOMPOSITION OF $S^2(\mathfrak{o}_n)$

Let E denote the natural representation of \mathfrak{o}_n , i.e., $E = \mathbb{C}^n$ equipped with the standard bilinear form β_0 , where $\beta_0(u, v) = {}^t uv$ ($u, v \in E$). Then,

$$\varphi : \wedge^2 E \rightarrow \mathfrak{o}_n, \quad v \wedge w \mapsto -\iota(\cdot)(v \wedge w)$$

provides an isomorphism between \mathfrak{o}_n -modules, where ι denotes the contraction with respect to β_0 . Hence, we have $S^2(\wedge^2 E) \simeq S^2(\mathfrak{o}_n)$.

It is well known that any irreducible representation of \mathfrak{o}_n is realized in the subspace of traceless tensors of $E^{\otimes k}$ for some $k \in \mathbb{N}$ and is parameterized by Young diagram. Noting that $\wedge^2 E = E_{\square}$, it follows that $\wedge^2 E \otimes \wedge^2 E$ decomposes as

$$\wedge^2 E \otimes \wedge^2 E = E_{\emptyset} \oplus E_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \oplus E_{\square\square} \oplus E_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \oplus E_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} \oplus E_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}, \quad (\text{A.1})$$

which can be verified by calculating Littlewood-Richardson coefficients in terms of universal characters (see e.g. [10]). Note also that $S^2(\wedge^2 E)$ is equal to the direct sum of the first four factors in (A.1):

$$E_{\emptyset}, \quad E_{\begin{array}{|c|} \hline \square \\ \hline \end{array}}, \quad E_{\square\square}, \quad E_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}. \quad (\text{A.2})$$

One can easily verify that $E_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}$ is the highest weight module with highest weight 2α , where α is the highest root of \mathfrak{o}_n .

Next, let us describe the first three irreducible summands in (A.2) concretely. Let $\{e_i; i = 1, \dots, n\}$ denote the standard basis of \mathbb{C}^n , i.e., $e_i = {}^t(0, \dots, \overset{i\text{-th}}{1}, \dots, 0)$. Then, one has

$$\begin{aligned} E_{\emptyset} &= \left\langle \sum_{i=1}^n e_i^2 \right\rangle_{\mathbb{C}}, \\ E_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} &= \left\langle e_i \wedge e_j \wedge e_k \wedge e_l; 1 \leq i < j < k < l \leq n \right\rangle_{\mathbb{C}}, \\ E_{\square\square} &= \left\langle e_i e_j - \frac{1}{n} \delta_{i,j} \sum_{k=1}^n e_k^2; i, j \in [n] \right\rangle_{\mathbb{C}}. \end{aligned} \quad (\text{A.3})$$

Thus, if one defines injective O_n -equivariant linear maps by

$$\begin{aligned} \varphi_{\emptyset} : E_{\emptyset} &\rightarrow S^2(\mathfrak{o}_n), & \sum_{i=1}^n e_i^2 &\mapsto Q^{\mathfrak{o}_n}, \\ \varphi_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} : E_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} &\rightarrow S^2(\mathfrak{o}_n), & e_i \wedge e_j \wedge e_k \wedge e_l &\mapsto S_{ijkl}^{\mathfrak{o}_n}, \\ \varphi_{\square\square} : E_{\square\square} &\rightarrow S^2(\mathfrak{o}_n), & e_i e_j - \frac{1}{n} \delta_{ij} \sum_{k=1}^n e_k^2 &\mapsto S_{ij}^{\mathfrak{o}_n}, \end{aligned} \quad (\text{A.4})$$

with

$$Q^{\mathfrak{o}_n} = \sum_{i,k \in [n]} M_{i,k} \otimes M_{k,i} = - \sum_{i,k \in [n]} M_{i,k} \otimes M_{i,k}, \quad (\text{A.5})$$

$$\begin{aligned} S_{ijkl}^{\mathfrak{o}_n} &= \frac{1}{2} (M_{i,j} \otimes M_{k,l} - M_{i,k} \otimes M_{j,l} + M_{i,l} \otimes M_{j,k} \\ &\quad + M_{k,l} \otimes M_{i,j} - M_{j,l} \otimes M_{i,k} + M_{j,k} \otimes M_{i,l}), \end{aligned} \quad (\text{A.6})$$

$$S_{ij}^{\mathfrak{o}_n} = \frac{1}{2} \sum_{k \in [n]} (M_{i,k} \otimes M_{k,j} + M_{k,j} \otimes M_{i,k}) - \frac{1}{n} \delta_{i,j} Q^{\mathfrak{o}_n}, \quad (\text{A.7})$$

then one sees that the irreducible decomposition of $S^2(\mathfrak{o}_n)$ is given by

$$S^2(\mathfrak{o}_n) = E_{\emptyset}^{\mathfrak{o}_n} \oplus E_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}^{\mathfrak{o}_n} \oplus E_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{\mathfrak{o}_n} \oplus E_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}^{\mathfrak{o}_n}, \quad (\text{A.8})$$

where

$$\begin{aligned} E_{\emptyset}^{\mathfrak{o}_n} &= \langle Q^{\mathfrak{o}_n} \rangle_{\mathbb{C}}, \\ E_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}^{\mathfrak{o}_n} &= \langle S_{ijkl}^{\mathfrak{o}_n}; 1 \leq i < j < k < l \leq n \rangle_{\mathbb{C}}, \\ E_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{\mathfrak{o}_n} &= \langle S_{ij}^{\mathfrak{o}_n}; i, j \in [n] \rangle_{\mathbb{C}}, \end{aligned} \quad (\text{A.9})$$

and $E_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{\mathfrak{o}_n}$ is the image of $E_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{\mathfrak{o}_n}$ under the isomorphism $S^2(\wedge^2 E) \simeq S^2(\mathfrak{o}_n)$.

REFERENCES

1. B. Binegar and R. Zierau, *Unitarization of a singular representation of $SO(p, q)$* , Comm. Math. Phys. **138** (1991), no. 2, 245–258. MR 1108044
2. W.-T. Gan and G. Savin, *Uniqueness of Joseph ideal*, Math. Res. Lett. **11** (2004), no. 5-6, 589–597. MR 2106228
3. D. Garfinkle, *A new construction of the joseph ideal*, (1982), Thesis (Ph.D.), Massachusetts Institute of Technology. MR 2941017
4. R. Goodman and N. Wallach, *Representations and invariants of the classical groups*, Encyclopedia of Mathematics and its Applications, vol. 68, Cambridge University Press, Cambridge, 1998. MR 1606831
5. T. Hashimoto, *The moment map on symplectic vector space and oscillator representation*, Kyoto J. Math. **57** (2017), no. 3, 553–583. MR 3685055
6. ———, *(\mathfrak{g}, K) -module of $O(p, q)$ associated with the finite-dimensional representation of \mathfrak{sl}_2* , Internat. J. Math. **32** (2021), no. 2, Paper No. 2150009, 23. MR 4223528
7. R. Howe, *On some results of Strichartz and of Rallis and Schiffman*, J. Funct. Anal. **32** (1979), no. 3, 297–303. MR 538856 (80f:22014)
8. A. Joseph, *The minimal orbit in a simple Lie algebra and its associated maximal ideal*, Ann. Sci. École Norm. Sup. (4) **9** (1976), no. 1, 1–29. MR 404366
9. T. Kobayashi and B. Ørsted, *Analysis on the minimal representation of $O(p, q)$. II. Branching laws*, Adv. Math. **180** (2003), no. 2, 513–550. MR 2020551 (2004k:22018b)
10. S. Okada, *Representation theory of the classical groups and combinatorics II*, Baihukan, 2006, (in japanese).
11. S. Rallis and G. Schiffmann, *Weil representation. I. Intertwining distributions and discrete spectrum*, Mem. Amer. Math. Soc. **25** (1980), no. 231, iii+203. MR 567800 (81j:22007)
12. C.-B. Zhu and J.-S. Huang, *On certain small representations of indefinite orthogonal groups*, Represent. Theory **1** (1997), 190–206 (electronic). MR 1457244

CENTER FOR DATA SCIENCE EDUCATION, TOTTORI UNIVERSITY, 4-101, KOYAMA-MINAMI, TOTTORI, 680-8550, JAPAN

Email address: thashi@tottori-u.ac.jp