

# UNIQUENESS OF QUASI-ROOTS IN RIGHT-ANGLED ARTIN GROUPS

EON-KYUNG LEE AND SANG-JIN LEE

ABSTRACT. We introduce the notion of quasi-roots and study their uniqueness in right-angled Artin groups.

## 1. INTRODUCTION

In a group, if the equation  $g^n = h$  holds, then  $g$  is called an  $n$ th root of  $h$ .

The uniqueness of roots in groups has been studied by many authors, e.g. [1]. Mal'cev proved that roots are unique (i.e.  $g_1^n = g_2^n$  implies  $g_1 = g_2$ ) in a torsion-free locally nilpotent group [17]. It is well-known that centralizers of nontrivial elements of a torsion-free hyperbolic group are necessarily infinite cyclic [3, 7], hence roots are unique. In the Artin groups of finite type  $\mathbf{A}_n$ ,  $\mathbf{B}_n = \mathbf{C}_n$  and affine type  $\tilde{\mathbf{A}}_{n-1}$ ,  $\tilde{\mathbf{C}}_{n-1}$ , roots are unique up to conjugacy [8, 13, 16]. For the uniqueness of roots in general mapping class groups, see [2]. Pure braid groups and right-angled Artin groups are biorderable, hence roots are unique [6, 9, 18].

In this article, we introduce the notion of quasi-roots and study the uniqueness of quasi-roots in right-angled Artin groups.

**1.1. Right-angled Artin groups.** Before stating our results, we recall basic notions in right-angled Artin groups. See [4, 10, 19] for further details.

Throughout this article all graphs are finite and simple. For a graph  $\Gamma$ , let  $V(\Gamma)$  and  $E(\Gamma)$  denote the vertex set and the edge set of  $\Gamma$ , respectively. For a graph  $\Gamma$ , the *right-angled Artin group*  $A(\Gamma)$  on  $\Gamma$  is defined by the presentation

$$A(\Gamma) = \langle v \in V(\Gamma) \mid [v_i, v_j] = 1 \text{ if } \{v_i, v_j\} \in E(\Gamma) \rangle.$$

In this article, we use the opposite convention

$$G(\Gamma) = \langle v \in V(\Gamma) \mid [v_i, v_j] = 1 \text{ if } \{v_i, v_j\} \notin E(\Gamma) \rangle.$$

In other words,  $G(\Gamma) = A(\Gamma^c)$ , where  $\Gamma^c$  denotes the complement graph of  $\Gamma$ .

Each element in  $V(\Gamma) \cup V(\Gamma)^{-1}$  is called a *letter*. A *word* means a finite sequence of letters. For two words  $w_1$  and  $w_2$ , the notation  $w_1 \equiv w_2$  means that  $w_1$  and  $w_2$  coincide as sequences of letters. For example, if  $w_1 \equiv v_1 v_1 v_2 v_2^{-1}$  and  $w_2 \equiv v_1 v_1$ , where  $v_1, v_2 \in V(\Gamma)$ ,

---

2010 *Mathematics Subject Classification.* 20F36, 20F65.

*Key words and phrases.* Right-angled Artin group, quasi-root, normal form.

then  $w_1 \not\equiv w_2$ . A word  $w'$  is called a *subword* of a word  $w$  if  $w \equiv w_1 w' w_2$  for possibly empty words  $w_1$  and  $w_2$ .

An element  $g$  in  $G(\Gamma)$  can be expressed as a word  $w$ . The word  $w$  is called *reduced* if  $w$  is a shortest word among all words representing  $g$ . In this case, the length of  $w$  is called the *word length* of  $g$ , denoted by  $|g|$ .

The *support* of  $g \in G(\Gamma)$ , denoted by  $\text{supp}(g)$ , is defined as the set of all generators  $v \in V(\Gamma)$  such that  $v$  or  $v^{-1}$  appears in a reduced word representing  $g$ . It is known that  $\text{supp}(g)$  is well-defined. For example, see [10, 11, 14, 15].

Let  $w$  be a (possibly non-reduced) word in  $G(\Gamma)$ . A subword  $v^{\pm 1} w_1 v^{\mp 1}$ ,  $v \in V(\Gamma)$ , of  $w$  is called a *cancellation* of  $v$  in  $w$  if each generator in  $\text{supp}(w_1)$  commutes with  $v$ . If, furthermore, no letter in  $w_1$  is equal to  $v$  or  $v^{-1}$ , it is called an *innermost cancellation* of  $v$  in  $w$ . It is known that  $w$  is reduced if and only if  $w$  has no innermost cancellation [12, 14, 15].

Abusing notation, we shall sometimes regard a word as the group element represented by that word.

**Definition 1.1** (disjointly commute). We say that  $g_1, g_2 \in G(\Gamma)$  *disjointly commute*, denoted by  $g_1 \rightleftharpoons g_2$ , if  $\text{supp}(g_1) \cap \text{supp}(g_2) = \emptyset$  and each  $v_1 \in \text{supp}(g_1)$  commutes with each  $v_2 \in \text{supp}(g_2)$  in  $G(\Gamma)$ , i.e.  $\{v_1, v_2\} \notin E(\Gamma)$ .

**Definition 1.2** (strongly non-split). An element  $g \in G(\Gamma)$  is *non-split* if  $\text{supp}(g)$  spans a connected subgraph of  $\Gamma$ . Otherwise, it is *split*. We say that  $g$  is *strongly non-split* if  $g$  is non-split and there is no generator  $v \in V(\Gamma)$  that disjointly commutes with  $g$ .

**Example 1.3.** Let  $P_5 = \overset{\bullet}{v_1} - \overset{\bullet}{v_2} - \overset{\bullet}{v_3} - \overset{\bullet}{v_4} - \overset{\bullet}{v_5}$  be the path graph on five vertices  $v_1, \dots, v_5$ . The right-angled Artin group  $G(P_5)$  has a presentation

$$G(P_5) = \langle v_1, \dots, v_5 \mid [v_i, v_j] = 1 \text{ if } |i - j| \geq 2 \rangle.$$

Let  $g_1 = v_2 v_3 v_5^{-2}$ ,  $g_2 = v_2 v_3$  and  $g_3 = v_2 v_3 v_4$ . Then  $\text{supp}(g_1) = \{v_2, v_3, v_5\}$ ,  $\text{supp}(g_2) = \{v_2, v_3\}$  and  $\text{supp}(g_3) = \{v_2, v_3, v_4\}$ . The element  $g_1$  is split because  $\text{supp}(g_1)$  spans the disconnected subgraph  $\overset{\bullet}{v_2} - \overset{\bullet}{v_3} \quad \overset{\bullet}{v_5}$  of  $\Gamma$ . The element  $g_2$  is non-split but it is not strongly non-split because  $v_5$  disjointly commutes with  $g_2$ . The element  $g_3$  is strongly non-split because  $\text{supp}(g_3) = \{v_2, v_3, v_4\}$  spans the connected subgraph  $\overset{\bullet}{v_2} - \overset{\bullet}{v_3} - \overset{\bullet}{v_4}$  of  $\Gamma$  and because no generator disjointly commutes with  $g_3$ .

**Definition 1.4** (geodesic decomposition). For elements  $g, g_1, \dots, g_k \in G(\Gamma)$ , the decomposition  $g = g_1 \cdots g_k$  is a *geodesic decomposition* of  $g$  if  $|g| = |g_1| + \cdots + |g_k|$ .

**Definition 1.5** (cyclic conjugate). For  $g \in G(\Gamma)$  with a geodesic decomposition  $g = g_1 g_2$ , the element  $g_2 g_1$  is a *cyclic conjugate* of  $g$ . For a word  $w \equiv w_1 w_2$ , the word  $w_2 w_1$  is a *cyclic conjugate* of  $w$ .

**Definition 1.6** (cyclically reduced). An element  $g \in G(\Gamma)$  is *cyclically reduced* if  $g$  has the minimum word length in its conjugacy class.

Servatius [19, Proposition on p.38] proved that for any  $g \in G(\Gamma)$  there exists a geodesic decomposition  $g = u^{-1}hu$  for  $u, h \in G(\Gamma)$  such that  $h$  is cyclically reduced.

**Definition 1.7** (primitive). A nontrivial element  $g \in G(\Gamma)$  is *primitive* if  $g$  is not a nontrivial power of another element, i.e.  $g = u^n$  never holds for any  $n \geq 2$  and  $u \in G(\Gamma)$ . Similarly, a nonempty word  $w$  is *primitive* if  $w$  is not a nontrivial power of another word.

1.2. **Quasi-roots.** We define quasi-roots as follows.

**Definition 1.8.** Let  $0 \leq \lambda < \frac{1}{2}$  and  $N \geq 2$ . For  $h \in G(\Gamma)$ , an element  $g \in G(\Gamma)$  is a  $(\lambda, N)$ -*quasi-root* of  $h$  if there exists a geodesic decomposition

$$(1) \quad h = ag^nb = a \cdot \underbrace{g \cdot g \cdots g}_n \cdot b$$

for some  $n \geq N$  and  $a, b \in G(\Gamma)$  with  $|a|, |b| \leq \lambda|h|$ . The above decomposition is called a  $(\lambda, N)$ -*quasi-root decomposition* of  $h$ .

We are interested in the uniqueness of quasi-roots: given an element  $h \in G(\Gamma)$ , we want to find conditions under which we can guarantee that a quasi-root of  $h$  is uniquely determined up to conjugacy.

Here are immediate observations concerning the quasi-root decomposition (1).

- (i) If  $|a| = |b| = 0$ , then (1) becomes  $h = g^n$ , hence  $g$  is an  $n$ th root of  $h$ .
- (ii) The condition  $N \geq 2$  is natural because we cannot expect any kind of uniqueness of quasi-roots for  $n = 1$  (and  $|a| + |b| > 0$ ). Meanwhile, since  $n \geq N \geq 2$  and  $g^n = g \cdots g$  is geodesic,  $g$  is cyclically reduced. (See Lemma 2.1.)
- (iii) The condition  $\lambda < \frac{1}{2}$  is natural. If  $|a| \geq \frac{1}{2}|h|$  or  $|b| \geq \frac{1}{2}|h|$  is allowed, then  $g$  is not uniquely determined up to conjugacy. For example, let  $h = v_1^n v_2^n$ , where  $v_1$  and  $v_2$  are different generators. Then  $h = v_1^n \cdot 1^n \cdot v_2^n = 1 \cdot v_1^n \cdot v_2^n = v_1^n \cdot v_2^n \cdot 1$ , hence  $g$  is possibly one of  $1, v_1$  and  $v_2$ . However, the elements  $1, v_1$  and  $v_2$  are not conjugate to each other.

The following is the main result of this article.

**Theorem 1.9.** Let  $\Gamma$  be a finite connected graph, and let  $0 \leq \lambda < \frac{1}{2}$  and  $N \geq \frac{2|V(\Gamma)|+1}{1-2\lambda}$ . If there are two  $(\lambda, N)$ -quasi-root decompositions

$$h = a_1 g_1^{n_1} b_1 = a_2 g_2^{n_2} b_2$$

such that  $g_1$  and  $g_2$  are strongly non-split and primitive, then the quasi-roots  $g_1$  and  $g_2$  are conjugate such that  $a_1 g_1 a_1^{-1} = a_2 g_2 a_2^{-1}$  and  $b_1^{-1} g_1 b_1 = b_2^{-1} g_2 b_2$ .

The constant  $N$  depends only on  $\lambda$  and  $|V(\Gamma)|$ . Hence we obtain the following.

**Corollary 1.10.** Let  $\Gamma$  be a finite connected graph. For any  $\lambda \in [0, \frac{1}{2})$ , there exists  $N \geq 3$  such that strongly non-split and primitive  $(\lambda, N)$ -quasi-roots are unique up to conjugacy.

Concerning the main theorem, we remark the following.

- (i) The strongly non-splitness of  $g_i$  is a necessary condition for the uniqueness of quasi-roots. For example, consider  $G(P_5)$  in Example 1.3. Let  $h = (v_2^m v_3^m v_5)^n$ . Observe that  $h = 1 \cdot (v_2^m v_3^m v_5)^n \cdot 1 = v_5^n \cdot (v_2^m v_3^m)^n \cdot 1 = 1 \cdot (v_2^m v_3^m)^n \cdot v_5^n$ . Therefore, if  $m$  is large enough so that  $(2m + 1) \geq 1/\lambda$ , then  $n \leq \lambda(2m + 1)n = \lambda|h|$ . Hence both  $v_2^m v_3^m v_5$  and  $v_2^m v_3^m$  are  $(\lambda, n)$ -quasi-roots of  $h$ . However,  $v_2^m v_3^m v_5$  and  $v_2^m v_3^m$  are not conjugate. Notice that  $v_2^m v_3^m v_5$  is split and that  $v_2^m v_3^m$  is non-split but not strongly non-split.
- (ii) The primitiveness of  $g_i$  is not a restriction. Without that condition, one may conclude that there exists an element  $u$  such that any  $(\lambda, N)$ -quasi-root is conjugate to a power of  $u$ .

Note that we cannot expect a naive generalization of the main theorem to mapping class groups. For example, consider the  $n$ -strand Artin braid group  $B_n$ . Let  $\delta = \sigma_{n-1} \cdots \sigma_1$  and  $\epsilon = \delta\sigma_1$ , where  $\sigma_1, \dots, \sigma_{n-1}$  are the standard generators of  $B_n$ . Then  $\delta^n = \epsilon^{n-1}$ . For any  $m \geq 1$ , the element  $h = \delta^{mn} = \epsilon^{m(n-1)}$  has two primitive (quasi-)roots  $\delta$  and  $\epsilon$  which are not conjugate to each other. In mapping class groups, pseudo-Anosov elements may behave like strongly non-split elements in right-angled Artin groups. Hence the following questions would be interesting.

**Question 1.11.** In mapping class groups, is it true that primitive pseudo-Anosov quasi-roots are unique up to conjugacy?

**Question 1.12.** In torsion-free hyperbolic groups, is it true that primitive quasi-roots are unique up to conjugacy?

The results in this article seem insufficient to build a finite-time algorithm for finding a quasi-root or deciding the existence of a quasi-root. Hence the following questions would be also interesting.

**Question 1.13** (Quasi-root decision problem).

Given a graph  $\Gamma$  and  $(\lambda, N, h) \in [0, 1/2) \times [2, \infty) \times G(\Gamma)$ , decide whether or not  $h$  has a  $(\lambda, N)$ -quasi-root.

**Question 1.14** (Quasi-root search problem).

Given a graph  $\Gamma$  and  $(\lambda, N, h) \in [0, 1/2) \times [2, \infty) \times G(\Gamma)$ , where  $h$  is assumed to have a  $(\lambda, N)$ -quasi-root, find a  $(\lambda, N)$ -quasi-root  $g$  of  $h$  together with a  $(\lambda, N)$ -quasi-root decomposition  $h = ag^nb$ .

## 2. SD-CONICAL ELEMENTS

Crisp, Godelle and Wiest introduced a normal form of elements and the notion of pyramidal elements in order to solve the conjugacy problem in right-angled Artin groups in linear-time [5]. We use their normal form and a variation of pyramidal elements, called

SD-conical elements. In this section we define SD-conical elements and explore their properties.

The following equivalences are used implicitly in many works. We include a proof for readers' convenience.

**Lemma 2.1.** *The following are equivalent for  $g \in G(\Gamma)$ .*

- (i)  $g$  is cyclically reduced.
- (ii) There is no geodesic decomposition as  $g = u^{-1}hu$  for  $u, h \in G(\Gamma)$  with  $u \neq 1$ .
- (iii)  $g^n = gg \cdots g$  is geodesic (i.e.  $|g^n| = n|g|$ ) for all  $n \geq 2$ .
- (iv)  $g^n = gg \cdots g$  is geodesic (i.e.  $|g^n| = n|g|$ ) for some  $n \geq 2$ .
- (v) Every cyclic conjugate of  $g$  has the same word length as  $g$ , i.e. for any geodesic decomposition  $g_1g_2$  of  $g$ , the decomposition  $g_2g_1$  is also geodesic.

*Proof.* “(i)  $\Leftrightarrow$  (ii)” is known by Servatius [19, Proposition on p.38]. The implications “(iii)  $\Rightarrow$  (iv)  $\Rightarrow$  (v)  $\Rightarrow$  (ii)” are obvious. Therefore it is enough to show “(ii)  $\Rightarrow$  (iii)”.

Assume that  $g^n = gg \cdots g$  is not geodesic for some  $n \geq 2$ . Let  $w$  be a reduced word representing  $g$ . Then  $w^n = ww \cdots w$  has an innermost cancellation. Since  $w$  is reduced, this cancellation occurs between two consecutive  $w$ 's, i.e. there are words  $w_1, \dots, w_4$  and a letter  $x \in V(\Gamma)^{\pm 1}$  such that

$$w \equiv w_1xw_2 \equiv w_3x^{-1}w_4, \quad x \equiv w_2, \quad x^{-1} \equiv w_3.$$

Since  $x \equiv w_2$ , we have  $x \notin \text{supp}(w_2)$ , hence the subword  $w_3x^{-1}$  must be a prefix of  $w_1$ . Therefore  $w_1 \equiv w_3x^{-1}w_5$  for some word  $w_5$ . Now we have

$$w \equiv w_3x^{-1}w_5xw_2, \quad x \equiv w_2, \quad x^{-1} \equiv w_3.$$

Then  $g$  has a geodesic decomposition as  $g = x^{-1}hx$ , where  $h = w_3w_5w_2$ . This contradicts (ii).  $\square$

**Definition 2.2** (set of starting generators). For  $g \in G(\Gamma)$ , the *set of starting generators*, denoted by  $S(g)$ , is the set of all generators  $v \in V(\Gamma)$  such that either  $v$  or  $v^{-1}$  is the first letter of a reduced word representing  $g$ , i.e.

$$S(g) = \{v \in V(\Gamma) \mid g = v^\epsilon h \text{ is geodesic for some } \epsilon = \pm 1 \text{ and } h \in G(\Gamma)\}.$$

**Definition 2.3** (conical element and apex). For  $g \in G(\Gamma)$ , if  $S(g)$  consists of a single generator, say  $v_0$ , then we say that  $g$  is *conical* or  $v_0$ -conical. The generator  $v_0$  is called the *apex* of  $g$ , denoted by  $\text{apex}(g)$ .

Note that a conical element is necessarily non-split.

The following proposition is a quantitative version of Proposition 2.18 in [5]. Though it can be proved in almost the same way, we include a sketchy proof for better readability.

**Proposition 2.4.** *Let  $g \in G(\Gamma)$  be non-split and cyclically reduced, and let  $v_0 \in \text{supp}(g)$ . Then  $g$  is conjugate to a  $v_0$ -conical element  $p \in G(\Gamma)$  by  $a, b \in G(\Gamma)$  as*

$$g = apa^{-1} = b^{-1}pb,$$

where  $g^k = ab$  holds and is geodesic for some  $0 \leq k \leq |V(\Gamma)| - 1$ . In particular, if  $n \geq |V(\Gamma)|$ , then  $g^n$  has a geodesic decomposition as  $g^n = ap^{n-k}b$ .

*Sketchy proof.* If  $g$  is  $v_0$ -conical, there is nothing to prove because we can take  $k = 0$  and  $a = b = 1$ . So we assume that  $g$  is not  $v_0$ -conical.

We follow the argument in [5]. For any non-split element  $h \in G(\Gamma)$  and for any  $v \in \text{supp}(h)$ , there exists a unique geodesic decomposition as  $h = tp$  such that  $p$  is  $v$ -conical and  $v \notin \text{supp}(t)$  [5, Lemma 2.16].

Starting from  $g_0 = g$ , we inductively do the following for  $i = 0, 1, 2, \dots$ : (i) determine the geodesic decomposition  $g_i = t_i \cdot p_i$ , where  $p_i$  is  $v_0$ -conical and  $v_0 \notin \text{supp}(t_i)$ ; (ii) take the conjugation  $g_{i+1} = t_i^{-1}g_it_i = p_it_i$ .

Then  $g_j$  is  $v_0$ -conical for some  $1 \leq j \leq |V(\Gamma)| - 1$ . (In fact, the proof of [5, Proposition 2.18] says that one can take  $j \leq \max\{\text{dist}_{\Gamma_1}(v_0, v) \mid v \in \text{supp}(g)\}$ , where  $\Gamma_1$  is the subgraph of  $\Gamma$  spanned by  $\text{supp}(g)$ . Notice that  $\max\{\text{dist}_{\Gamma_1}(v_0, v) \mid v \in \text{supp}(g)\} \leq |V(\Gamma)| - 1$ .) Let  $k$  be the smallest integer such that  $g_k$  is  $v_0$ -conical. Then  $1 \leq k \leq |V(\Gamma)| - 1$ .

In the decomposition  $g_k = t_k \cdot p_k$ , we have  $t_k = 1$  and  $g_k = p_k$ . Keeping track of the above procedure, we get

$$\begin{aligned} g_0 &= g = t_0 \cdot p_0, \\ g_1 &= t_0^{-1}g_0t_0 = p_0t_0 = t_1 \cdot p_1, \\ g_2 &= t_1^{-1}g_1t_1 = p_1t_1 = t_2 \cdot p_2, \\ &\dots \\ g_k &= t_{k-1}^{-1}g_{k-1}t_{k-1} = p_{k-1}t_{k-1} = t_k p_k = p_k. \end{aligned}$$

Let  $p = g_k = p_k$ ,  $a = t_0 \cdots t_{k-1}$  and  $b = p_{k-1} \cdots p_0$ . Then the following hold

$$|g| = |g_0| = \cdots = |g_{k-1}| = |g_k| = |p|,$$

$$|g_i^m| = m|g_i| \text{ for } 0 \leq i \leq k, m \geq 1$$

because  $g$  is cyclically reduced and each  $g_{i+1}$  is a cyclic conjugate of  $g_i$ , and hence also cyclically reduced for  $0 \leq i \leq k - 1$ .

Using the identities  $p_it_i = t_{i+1}p_{i+1}$  for  $0 \leq i \leq k - 1$ , we obtain

$$g^k = (t_0p_0)^k = \underbrace{(t_0p_0)(t_0p_0) \cdots (t_0p_0)}_k = (t_0t_1 \cdots t_{k-1})(p_{k-1} \cdots p_1p_0) = ab,$$

$$ga = t_0p_0t_0t_1 \cdots t_{k-1} = t_0 \cdots t_{k-1}t_k p_k = ap,$$

$$bg = p_{k-1} \cdots p_1p_0t_0p_0 = t_k p_k p_{k-1} \cdots p_1p_0 = pb,$$

namely  $g^k = ab$  and  $g = apa^{-1} = b^{-1}pb$ .

On the other hand,

$$|a| + |b| \leq |t_0| + \cdots + |t_{k-1}| + |p_0| + \cdots + |p_{k-1}| = |g_0| + \cdots + |g_{k-1}| = k|g| = |g^k|,$$

where the first equality holds because each  $g_i = t_i \cdot p_i$  is a geodesic decomposition. Since  $|g^k| \leq |a| + |b|$  is obvious, we get  $|a| + |b| = |g^k|$ , namely  $g^k = ab$  is a geodesic decomposition.

The identity  $g^n = ap^{n-k}b$  follows from  $g^k = ab$  and  $g = b^{-1}pb$ . Observe

$$|a| + |b| + |p^{n-k}| = k|g| + (n-k)|p| = n|g| = |g^n|,$$

hence  $g^n = ap^{n-k}b$  is a geodesic decomposition.  $\square$

**Definition 2.5** ( $v_0$ -conical conjugate). Given a non-split and cyclically reduced element  $g \in G(\Gamma)$  and a generator  $v_0 \in \text{supp}(g)$ , the procedure in the proof of Proposition 2.4 uniquely determines a  $v_0$ -conical element  $p$ , called the  $v_0$ -conical conjugate of  $g$ .

The conical conjugate  $p$  in the above definition is also non-split and cyclically reduced because  $p$  is obtained from  $g$  by iterated cyclic conjugations and cyclic conjugations preserve non-splitness and cyclically reducedness. In particular,  $|g| = |p|$ .

**Definition 2.6** (CGW-normal form). Suppose  $V(\Gamma)$  is endowed with a linear order.

- (i) A reduced word  $w \equiv v_{i_1}^{\epsilon_1} \cdots v_{i_k}^{\epsilon_k}$  on  $V(\Gamma)^{\pm 1}$  representing an element  $g \in G(\Gamma)$  is *initially normal* if either  $w$  is trivial or  $v_{i_1}$  is the largest element of  $S(g)$ .
- (ii) A reduced word  $w$  is *normal* if all its suffixes are initially normal.
- (iii) Any element  $g \in G(\Gamma)$  has a unique normal representative word [5, Proposition 2.6]. We call this normal word the *CGW-normal form* of  $g$ , and denote it by  $\sigma(g)$ .

Note that all the subwords of a normal word are normal words.

One can understand  $\sigma(g)$  as follows. Extend the linear order on  $V(\Gamma)$  to  $V(\Gamma)^{\pm 1}$  so that each generator  $v$  is the immediate predecessor of  $v^{-1}$ . (Namely, if the order on  $V(\Gamma)$  is  $v_1 < \cdots < v_r$ , then the order on  $V(\Gamma)^{\pm 1}$  is  $v_1 < v_1^{-1} < v_2 < v_2^{-1} < \cdots < v_r < v_r^{-1}$ .) Then  $\sigma(g)$  is the largest in the lexicographic order among all the reduced words representing  $g$ .

**Definition 2.7** (SD-conical). Suppose that  $V(\Gamma)$  is endowed with a linear order  $<$ .

- (i) A conical element  $g$  is *pyramidal* if  $\text{apex}(g)$  is the smallest element of  $\text{supp}(g)$ .
- (ii) A conical element  $g$  is *SD-conical* if  $v_0 = \text{apex}(g)$  does not commute with any generator smaller than  $v_0$ , i.e. if  $v < v_0$ , then  $\{v, v_0\} \in E(\Gamma)$ . (From a graph theoretical point of view, the vertex  $v_0$  dominates all the vertices  $v \in V(\Gamma)$  smaller than  $v_0$ . SD stands for *smaller vertex dominating*.)

**Example 2.8.** Consider  $G(P_5)$  in Example 1.3. Define a linear order on  $V(P_5)$  by  $v_1 < v_2 < v_3 < v_4 < v_5$ . Let  $g = v_2v_4^{-1}v_3^{-1}v_5$ . Then  $S(g) = \{v_2, v_4\}$ . The following are reduced words representing  $g$ .

$$w_1 \equiv v_2v_4^{-1}v_3^{-1}v_5, \quad w_2 \equiv v_4^{-1}v_2v_5v_3^{-1}, \quad w_3 \equiv v_4^{-1}v_5v_2v_3^{-1}, \dots$$

The word  $w_1$  is not initially normal because  $v_2$  is not the largest element of  $S(g)$ . The word  $w_2$  is initially normal, but it is not normal because the suffix  $v_2v_5v_3^{-1}$  is not initially normal. The word  $w_3$  is normal. Hence  $\sigma(g) \equiv w_3 \equiv v_4^{-1}v_5v_2v_3^{-1}$ .

The element  $g = v_2v_4^{-1}v_3^{-1}v_5 = v_4^{-1}v_5v_2v_3^{-1}$  is not conical because  $S(g)$  is not a singleton. The conjugate  $g_1 = (v_4^{-1}v_5)^{-1}g(v_4^{-1}v_5) = v_2v_3^{-1}v_4^{-1}v_5$  is pyramidal because  $S(g_1) = \{v_2\}$  and  $v_2 = \text{apex}(g_1)$  is the smallest element of  $\text{supp}(g_1) = \text{supp}(g) = \{v_2, v_3, v_4, v_5\}$ . Moreover,  $g_1$  is SD-conical because  $v_1$  is the only generator smaller than  $v_2 = \text{apex}(g_1)$  but it does not commute with  $v_2$ .

In the same way, one can see that  $h_1 = v_4v_5$  is pyramidal but not SD-conical, and that  $h_2 = v_2v_1v_3v_4$  is SD-conical but not pyramidal.

The notions of normal form, pyramidal and SD-conical are defined via a linear order on  $V(\Gamma)$ . Whenever such terms are used, it is assumed that  $V(\Gamma)$  is endowed with some linear order even though it is not specified.

**Lemma 2.9.** *Let  $g_1g_2$  be geodesic for  $g_1, g_2 \in G(\Gamma)$ .*

- (i) *If  $g_2$  is SD-conical, then  $\sigma(g_1g_2) \equiv \sigma(g_1)\sigma(g_2)$ .*
- (ii) *If  $g_1$  is SD-conical and strongly non-split, then  $g_1g_2$  is also SD-conical and strongly non-split.*

*Proof.* In the proof of Proposition 2.20 in [5], the following is observed.

If  $v_1^{\epsilon_1}, v_2^{\epsilon_2}$  are letters and  $w$  is a reduced word such that the words  $v_1^{-\epsilon_1}w$  and  $wv_2^{\epsilon_2}$  are reduced but the word  $v_1^{-\epsilon_1}wv_2^{\epsilon_2}$  is not, then  $v_1^{\epsilon_1} = v_2^{\epsilon_2}$  and  $v_1 \equiv w$ .

Using this, we get the following.

*Claim.* For  $v \in V(\Gamma)$ , if  $v \in S(g_1g_2)$  and  $v \notin S(g_1)$ , then  $v \in S(g_2)$  and  $v \equiv g_1$ .

*Proof of Claim.* Let  $w_1$  and  $w_2$  be reduced words representing  $g_1$  and  $g_2$ , respectively. Since  $v \in S(g_1g_2)$  and  $v \notin S(g_1)$ , we can write  $w_2 \equiv w'_2v_2^{\epsilon_2}w''_2$  such that  $v \notin S(w_1w'_2)$  and  $v \in S(w_1w'_2v_2^{\epsilon_2})$ , i.e.  $v^{-\epsilon}w_1w'_2$  is reduced but  $v^{-\epsilon}w_1w'_2v_2^{\epsilon_2}$  is not. Since  $g_1g_2$  is geodesic,  $w_1w'_2v_2^{\epsilon_2}$  is reduced. From the above observation,  $v \equiv w_1w'_2$  and  $v^\epsilon = v_2^{\epsilon_2}$ . Hence  $v \equiv g_1$  and  $v \in S(g_2)$ .  $\square$

(i) Our proof is similar to the one in [5, Proposition 2.20], where it is shown that if  $p$  is pyramidal and cyclically reduced, then  $\sigma(p)\sigma(p)$  is normal, hence  $\sigma(p^2) \equiv \sigma(p)\sigma(p)$ .

Assume  $\sigma(g_1g_2) \not\equiv \sigma(g_1)\sigma(g_2)$ , i.e.  $\sigma(g_1)\sigma(g_2)$  is not normal. Then there exists a suffix of  $\sigma(g_1)\sigma(g_2)$  that is not initially normal. Since  $\sigma(g_1)$  and  $\sigma(g_2)$  are normal, we can write

$$\sigma(g_1) \equiv w'_1v_1^{\epsilon_1}w_1, \quad \sigma(g_2) \equiv w_2v_2^{\epsilon_2}w'_2$$

such that  $v_1^{\epsilon_1}w_1w_2$  is initially normal but  $v_1^{\epsilon_1}w_1w_2v_2^{\epsilon_2}$  is not. Therefore there exists  $v_0^{\epsilon_0}$  such that

$$v_0 > v_1, \quad v_0 \notin S(v_1^{\epsilon_1}w_1w_2), \quad v_0 \in S(v_1^{\epsilon_1}w_1w_2v_2^{\epsilon_2}).$$



By the above claim,  $v_0 \equiv v_1^{\varepsilon_1} w_1 w_2$  and  $v_0 \in S(v_2^{\varepsilon_2}) = \{v_2\}$ . Therefore,

$$v_0 \equiv v_1, \quad v_0 \equiv w_1, \quad v_0 \equiv w_2, \quad v_0 = v_2.$$

Assume  $w_2$  is the empty word. Since  $g_2$  is SD-conical, we have  $v_0 = v_2 = \text{apex}(g_2)$  and  $v_0$  does not commute with any generator smaller than  $v_0$ . This is a contradiction because  $v_0 \equiv v_1$  and  $v_0 > v_1$ . Therefore  $w_2$  is not the empty word, hence  $S(w_2)$  is not empty.

Let  $v'_2 \in S(w_2)$ . Since  $v_2 = v_0 \equiv w_2$ , both  $v'_2$  and  $v_0$  belong to  $S(w_2 v_2^{\varepsilon_2})$ . Moreover,  $v'_2 \neq v_0$  because  $v'_2 \in S(w_2)$  and  $v_0 \equiv w_2$  imply  $v'_2 \in \text{supp}(w_2)$  and  $v_0 \notin \text{supp}(w_2)$ , respectively. Since  $S(w_2 v_2^{\varepsilon_2}) \subset S(g_2)$ , we have  $|S(g_2)| \geq 2$ , which is a contradiction because  $g_2$  is conical.

(ii) We first show that  $g_1 g_2$  is conical. Assume  $|S(g_1 g_2)| \geq 2$ . Let  $v_1 = \text{apex}(g_1)$ . Then  $v_1 \in S(g_1) \subset S(g_1 g_2)$ . Because  $|S(g_1 g_2)| \geq 2$ , there exists  $v_2 \in S(g_1 g_2) \setminus \{v_1\}$ . Since  $g_1$  is conical and  $v_2 \neq v_1$ , we have  $v_2 \notin S(g_1)$ . By the claim, we have  $v_2 \equiv g_1$ . This contradicts that  $g_1$  is strongly non-split. Therefore  $g_1 g_2$  is conical, and hence non-split.

Since  $S(g_1) \subset S(g_1 g_2)$  and  $|S(g_1 g_2)| = 1$ , we have  $S(g_1 g_2) = S(g_1)$ , hence  $\text{apex}(g_1 g_2) = \text{apex}(g_1)$ . Because  $g_1$  is SD-conical,  $\text{apex}(g_1 g_2) = \text{apex}(g_1)$  does not commute with any generator smaller than itself. Therefore  $g_1 g_2$  is SD-conical.

The element  $g_1 g_2$  is strongly non-split because  $\text{supp}(g_1 g_2) \supset \text{supp}(g_1)$  and  $g_1$  is strongly non-split.  $\square$

**Proposition 2.10.** *For  $n \geq 1$ , if  $ag^nb$  is geodesic and  $g$  is strongly non-split and SD-conical, then  $\sigma(ag^nb) \equiv \sigma(a)\sigma(g)^{n-1}\sigma(gb) \equiv \sigma(a)\underbrace{\sigma(g) \cdots \sigma(g)}_{n-1}\sigma(gb)$ .*

*Proof.* Because  $g$  is strongly non-split and SD-conical, by Lemma 2.9(ii),  $g^k b = g \cdot g^{k-1} b$  is strongly non-split and SD-conical for all  $1 \leq k \leq n$ . By Lemma 2.9(i),  $\sigma(ag^nb) \equiv \sigma(a)\sigma(g^nb)$  and  $\sigma(g^k b) \equiv \sigma(g \cdot g^{k-1} b) \equiv \sigma(g)\sigma(g^{k-1} b)$  for  $2 \leq k \leq n$ . By an induction on  $k$ , we can show  $\sigma(g^k b) \equiv \underbrace{\sigma(g) \cdots \sigma(g)}_{k-1}\sigma(gb)$  for all  $1 \leq k \leq n$ . Therefore

$$\sigma(ag^nb) \equiv \sigma(a)\sigma(g^nb) \equiv \sigma(a)\underbrace{\sigma(g) \cdots \sigma(g)}_{n-1}\sigma(gb). \quad \square$$

**Lemma 2.11.** *Let  $g_1, g_2 \in G(\Gamma)$  be strongly non-split. Then there exists a linear order on  $V(\Gamma)$  together with vertices  $v_1 \in \text{supp}(g_1)$  and  $v_2 \in \text{supp}(g_2)$  such that both the  $v_1$ -conical conjugate of  $g_1$  and the  $v_2$ -conical conjugate of  $g_2$  are strongly non-split and SD-conical.*

*Proof.* First, suppose  $\text{supp}(g_1) \cap \text{supp}(g_2) \neq \emptyset$ . Choose any  $v_0 \in \text{supp}(g_1) \cap \text{supp}(g_2)$ . Give a linear order on  $V(\Gamma)$  such that  $v_0$  is the smallest element of  $V(\Gamma)$ . Let  $p_1$  and  $p_2$  be the  $v_0$ -conical conjugates of  $g_1$  and  $g_2$ , respectively. Then  $\text{apex}(p_1) = \text{apex}(p_2) = v_0$  is the smallest element of  $V(\Gamma)$ , hence  $p_1$  and  $p_2$  are SD-conical.

Now suppose  $\text{supp}(g_1) \cap \text{supp}(g_2) = \emptyset$ . Since  $g_1$  is strongly non-split, there exists  $(v_1, v_2) \in \text{supp}(g_1) \times \text{supp}(g_2)$  such that  $\{v_1, v_2\} \in E(\Gamma)$ . Give a linear order on  $V(\Gamma)$

such that  $v_1 = \min V(\Gamma)$  and  $v_2 = \min (V(\Gamma) - \{v_1\})$ . For  $i = 1, 2$ , let  $p_i$  be the  $v_i$ -conjugate of  $g_i$ . Then  $p_1$  is SD-conical because  $\text{apex}(p_1) = v_1$  is the smallest element of  $V(\Gamma)$ . And  $p_2$  is SD-conical because  $\text{apex}(p_2) = v_2$  is smaller than any generator other than  $v_1$  and  $\{v_1, v_2\} \in E(\Gamma)$ .

In both cases above,  $p_1$  and  $p_2$  are strongly non-split because  $\text{supp}(p_i) = \text{supp}(g_i)$  for  $i = 1, 2$ .  $\square$

### 3. UNIQUENESS OF QUASI-ROOTS

In this section, we prove Theorem 1.9 in three steps.

**3.1. Step 1: Quasi-roots of words.** Let  $X$  be a set of symbols. In this subsection, a letter means an element of  $X^{\pm 1}$  and a word means a word on  $X^{\pm 1}$ . In Proposition 3.3, we will prove the uniqueness of primitive quasi-roots up to conjugacy for words.

**Lemma 3.1.** *Let  $\{x_n\}_{n \geq 1}$  and  $\{y_n\}_{n \geq 1}$  be sequences of letters of period  $p \geq 1$  and  $q \geq 1$ , respectively, such that the first  $p + q$  terms coincide, i.e.*

$$\begin{aligned} x_{n+p} &= x_n & \text{for } n \geq 1, \\ y_{n+q} &= y_n & \text{for } n \geq 1, \\ x_n &= y_n & \text{for } 1 \leq n \leq p + q. \end{aligned}$$

Then  $x_n = y_n$  for all  $n \geq 1$  and the sequence  $\{x_n\}_{n \geq 1}$  has period  $\gcd(p, q)$ .

*Proof.* For  $1 \leq n \leq p$ , we have

$$x_n = y_n = y_{n+q} = x_{n+q}.$$

Because  $\{x_n\}_{n \geq 1}$  has period  $p$ , the identity  $x_n = x_{n+q}$  holds for all  $n \geq 1$ . Therefore  $\{x_n\}_{n \geq 1}$  has period  $q$ , hence it has period  $\gcd(p, q)$ . By the same argument  $\{y_n\}$  has period  $\gcd(p, q)$ . Because  $\{x_n\}$  and  $\{y_n\}$  are sequences of period  $\gcd(p, q)$  whose first  $p + q$  terms are identical, we have  $x_n = y_n$  for all  $n \geq 1$ .  $\square$

**Corollary 3.2.** *Let  $w_1 \equiv x_1 \cdots x_p$  and  $w_2 \equiv y_1 \cdots y_q$  be two primitive words of lengths  $p$  and  $q$ , respectively. Suppose that there exist powers  $w_1^{m_1}$  and  $w_2^{m_2}$  such that  $w_1^2$  is a prefix of  $w_2^{m_2}$  and  $w_2^2$  is a prefix of  $w_1^{m_1}$ . Then  $p = q$  and  $w_1 \equiv w_2$ .*

*Proof.* Consider the infinite words

$$\begin{aligned} w_1^* &\equiv w_1 w_1 \cdots \equiv x_1 \cdots x_p x_1 \cdots x_p x_1 \cdots x_p \cdots, \\ w_2^* &\equiv w_2 w_2 \cdots \equiv y_1 \cdots y_q y_1 \cdots y_q y_1 \cdots y_q \cdots. \end{aligned}$$

From the hypothesis, the first  $p + q$  letters of  $w_1^*$  and  $w_2^*$  coincide.

Let  $d = \gcd(p, q)$ . By Lemma 3.1,  $w_1^*$  and  $w_2^*$  coincide and both have period  $d$ . Namely,  $u = x_1 \cdots x_d$  is a common root of  $w_1$  and  $w_2$ , i.e.  $w_1 \equiv u^a$  and  $w_2 \equiv u^b$ , where  $a = p/d$  and  $b = q/d$ . Since  $w_1$  and  $w_2$  are primitive, we have  $a = b = 1$ , hence  $p = q$  and  $w_1 \equiv w_2$ .  $\square$




FIGURE 1. The word  $w \equiv a_1 w_1^{m_1} b_1 \equiv a_2 w_2^{m_2} b_2$

**Proposition 3.3.** *Let  $w$  be a word with the following two decompositions*

$$w \equiv a_1 w_1^{m_1} b_1 \equiv a_2 w_2^{m_2} b_2,$$

where  $m_i \geq 2$  and  $w_i, a_i, b_i$  are words for  $i = 1, 2$ . Suppose that  $w_1$  and  $w_2$  are primitive and that there exist non-negative constants  $A$  and  $B$  such that

$$|a_i| \leq A, \quad |b_i| \leq B, \quad |w| - (A + B) \geq 2|w_i|$$

for  $i = 1, 2$ . Then  $w_1$  and  $w_2$  are cyclically conjugate. Furthermore,  $a_1 w_1 a_1^{-1} = a_2 w_2 a_2^{-1}$  and  $b_1^{-1} w_1 b_1 = b_2^{-1} w_2 b_2$  when the words are regarded as elements of a free group.

*Proof.* Without loss of generality, we may assume  $|a_1| \geq |a_2|$ . Then we have a decomposition  $w_2 \equiv w_2' w_2''$  such that  $a_1 \equiv a_2 w_2^k w_2'$  for some  $k \geq 0$ . (Notice that  $|a_1| + |b_2| + 2|w_2| \leq A + B + 2|w_2| \leq |w| = |a_2| + |b_2| + m_2|w_2|$  and hence  $|a_1| \leq |a_2| + (m_2 - 2)|w_2|$ .) See Figure 1. Therefore

$$w \equiv a_2 w_2^{m_2} b_2 \equiv a_2 w_2^k w_2' (w_2'' w_2')^{m_2 - k - 1} w_2'' b_2 \equiv a_1 (w_2'' w_2')^{m_2 - k - 1} w_2'' b_2.$$

Let  $z$  be the subword of  $w$  obtained by removing the first  $|a_1|$  letters and the last  $|B|$  letters from  $w$ . From the hypothesis that  $|w| - (A + B) \geq 2|w_i|$ , we have  $|z| \geq 2|w_i|$  for  $i = 1, 2$ . Notice that  $z$  is a prefix of both  $w_1^{m_1}$  and  $(w_2'' w_2')^{m_2 - k - 1} w_2''$ .

Since  $z$  is a prefix of  $w_1^{m_1}$ , it is of the form

$$z \equiv w_1 \cdots w_1 w_1' \equiv w_1^{n_1} w_1'$$

for some  $n_1 \geq 2$  and some prefix  $w_1'$  of  $w_1$ . On the other hand, since  $z$  is prefix of  $(w_2'' w_2')^{m_2 - k - 1} w_2''$ , it is of the form

$$z \equiv (w_2'' w_2')^{n_2} w_2''$$

for some  $n_2 \geq 2$  and some prefix  $w_2''$  of  $w_2'' w_2'$ .

Notice that  $w_2'' w_2'$  is primitive because it is a cyclic conjugate of the primitive word  $w_2$ . The words  $w_1$  and  $w_2'' w_2'$  now satisfy the hypothesis of Corollary 3.2. Therefore  $w_1 \equiv w_2'' w_2'$ , hence  $w_1$  and  $w_2$  are cyclically conjugate.

When the words are regarded as elements of a free group,

$$\begin{aligned} a_1 w_1 a_1^{-1} &= (a_2 w_2^k w_2')(w_2'' w_2')(a_2 w_2^k w_2')^{-1} = a_2 w_2^k w_2' w_2'' w_2^{-k} a_2^{-1} \\ &= a_2 w_2^k w_2 w_2^{-k} a_2^{-1} = a_2 w_2 a_2^{-1}. \end{aligned}$$

Using the same argument, we obtain  $b_1^{-1} w_1 b_1 = b_2^{-1} w_2 b_2$ .  $\square$

### 3.2. Step 2: SD-conical quasi-roots.

**Proposition 3.4.** *Let  $\Gamma$  be a connected graph, and let*

$$h = a_1 p_1^{n_1} b_1 = a_2 p_2^{n_2} b_2$$

*be two geodesic decompositions of  $h \in G(\Gamma)$ , where  $n_i \geq 3$  and  $a_i, b_i, p_i \in G(\Gamma)$  for  $i = 1, 2$ . Suppose that  $p_1$  and  $p_2$  are strongly non-split, SD-conical and primitive and that there exist non-negative constants  $A$  and  $B$  such that*

$$|a_i| \leq A, \quad |b_i| \leq B, \quad |h| - (A + B) \geq 3|p_i|$$

*for  $i = 1, 2$ . Then the CGW-normal forms  $\sigma(p_1)$  and  $\sigma(p_2)$  are cyclically conjugate. Furthermore,  $a_1 p_1 a_1^{-1} = a_2 p_2 a_2^{-1}$  and  $b_1^{-1} p_1 b_1 = b_2^{-1} p_2 b_2$ .*

*Proof.* Without loss of generality, we may assume  $|p_1| \geq |p_2|$ .

Since  $a_1 p_1^{n_1} b_1$  and  $a_2 p_2^{n_2} b_2$  are geodesic decompositions and since  $p_1$  and  $p_2$  are strongly non-split and SD-conical, the following holds by Proposition 2.10.

$$(2) \quad \sigma(h) \equiv \sigma(a_1) \sigma(p_1)^{n_1-1} \sigma(p_1 b_1) \equiv \sigma(a_2) \sigma(p_2)^{n_2-1} \sigma(p_2 b_2)$$

Let  $A' = A$  and  $B' = B + |p_1|$ . Then

$$|h| - (A' + B') = |h| - (A + B) - |p_1| \geq 3|p_1| - |p_1| = 2|p_1| \geq 2|p_2|.$$

Applying Proposition 3.3 to (2), we get the desired result.  $\square$

### 3.3. Step 3: Proof of Theorem 1.9.

**Proposition 3.5.** *Let  $\Gamma$  be a connected graph, and let*

$$(3) \quad h = a_1 g_1^{n_1} b_1 = a_2 g_2^{n_2} b_2$$

*be two geodesic decompositions of  $h \in G(\Gamma)$ , where  $n_i \geq 1$  and  $a_i, b_i, g_i \in G(\Gamma)$  for  $i = 1, 2$ . Suppose that  $g_1$  and  $g_2$  are strongly non-split and primitive and that there exist non-negative constants  $A$  and  $B$  such that*

$$|a_i| \leq A, \quad |b_i| \leq B, \quad |h| - (A + B) \geq (2|V(\Gamma)| + 1)|g_i|$$

*for  $i = 1, 2$ . Then  $g_1$  and  $g_2$  are conjugate such that  $a_1 g_1 a_1^{-1} = a_2 g_2 a_2^{-1}$  and  $b_1^{-1} g_1 b_1 = b_2^{-1} g_2 b_2$ .*

*Proof.* Let  $V = |V(\Gamma)|$ . Since  $n_i|g_i| = |g_i^{n_i}| = |h| - (|a_i| + |b_i|) \geq |h| - (A + B) \geq (2V + 1)|g_i|$ , we have  $n_i \geq 2V + 1 \geq 2$ . Since  $a_i g_i^{n_i} b_i$  is geodesic,  $g_i^{n_i}$  is also geodesic, hence  $g_i$  is cyclically reduced by Lemma 2.1.

By Lemma 2.11, we can choose a linear order on  $V(\Gamma)$  together with vertices  $v_1 \in \text{supp}(g_1)$  and  $v_2 \in \text{supp}(g_2)$  such that the  $v_i$ -conical conjugate  $p_i$  of  $g_i$  is strongly non-split and SD-conical for  $i = 1, 2$ . Note that  $|g_i| = |p_i|$ .

By Proposition 2.4, we have the following geodesic decompositions

$$(4) \quad g_1^{n_1} = c_1 p_1^{n_1 - k_1} d_1, \quad g_2^{n_2} = c_2 p_2^{n_2 - k_2} d_2$$

for some  $0 \leq k_1, k_2 \leq V - 1$ , where the elements  $c_i, d_i \in G(\Gamma)$  are such that  $g_i^{k_i} = c_i d_i$  is geodesic and  $g_i = c_i p_i c_i^{-1} = d_i^{-1} p_i d_i$  for  $i = 1, 2$ . Since  $g_i^{k_i} = c_i d_i$  is geodesic, we have

$$|c_i|, |d_i| \leq |g_i^{k_i}| \leq k_i |g_i| \leq (V - 1)|g_i|$$

for  $i = 1, 2$ . Combining (3) and (4), we have the following two decompositions of  $h$ :

$$(5) \quad h = a_1 c_1 p_1^{n_1 - k_1} d_1 b_1 = a_2 c_2 p_2^{n_2 - k_2} d_2 b_2.$$

They are geodesic decompositions because both  $h = a_i g_i^{n_i} b_i$  and  $g_i^{n_i} = c_i p_i^{n_i - k_i} d_i$  are geodesic.

Let  $r = \max\{|g_1|, |g_2|\}$ ,  $A' = A + (V - 1)r$  and  $B' = B + (V - 1)r$ . Then for  $i = 1, 2$

$$|a_i c_i| \leq |a_i| + |c_i| \leq A + (V - 1)|g_i| \leq A',$$

$$|d_i b_i| \leq |d_i| + |b_i| \leq B + (V - 1)|g_i| \leq B',$$

$$\begin{aligned} |h| - (A' + B') &= |h| - (A + B) - 2(V - 1)r \\ &\geq (2V + 1)r - 2(V - 1)r = 3r \\ &\geq 3|g_i| = 3|p_i|. \end{aligned}$$

Since  $p_i$  is conjugate to the primitive element  $g_i$ , it is also primitive. Since  $n_i \geq 2V + 1$  and  $k_i \leq V - 1$ , one has  $n_i - k_i \geq V + 2 \geq 3$ . Applying Proposition 3.4 to (5), we get

$$a_1 c_1 p_1 c_1^{-1} a_1^{-1} = a_2 c_2 p_2 c_2^{-1} a_2^{-1}, \quad b_1^{-1} d_1^{-1} p_1 d_1 b_1 = b_2^{-1} d_2^{-1} p_2 d_2 b_2.$$

From the identity  $g_i = c_i p_i c_i^{-1} = d_i^{-1} p_i d_i$ , we obtain

$$a_1 g_1 a_1^{-1} = a_2 g_2 a_2^{-1}, \quad b_1^{-1} g_1 b_1 = b_2^{-1} g_2 b_2.$$

□

*Proof of Theorem 1.9.* We are given  $0 \leq \lambda < \frac{1}{2}$ ,  $N \geq \frac{2|V(\Gamma)|+1}{1-2\lambda}$  and two geodesic decompositions of  $h$

$$(6) \quad h = a_1 g_1^{n_1} b_1 = a_2 g_2^{n_2} b_2,$$

where  $n_i \geq N$ ,  $a_i, b_i, g_i \in G(\Gamma)$ ,  $|a_i|, |b_i| \leq \lambda|h|$ , and  $g_i$  is strongly non-split and primitive for  $i = 1, 2$ .

Since  $|h| \geq |g_i^{n_i}| = n_i|g_i| \geq N|g_i|$ , one has  $|g_i| \leq |h|/N$  for  $i = 1, 2$ . Let  $A = B = \lambda|h|$ . Then  $|a_i| \leq A$ ,  $|b_i| \leq B$  and

$$|h| - (A + B) = (1 - 2\lambda)|h| \geq (2|V(\Gamma)| + 1)\frac{|h|}{N} \geq (2|V(\Gamma)| + 1)|g_i|$$

for  $i = 1, 2$ . Applying Proposition 3.5, we get the desired results.  $\square$

#### ACKNOWLEDGEMENTS

The first author was partially supported by NRF-2018R1D1A1B07043291. The second author was partially supported by NRF-2018R1D1A1B07043268.

#### REFERENCES

- [1] G. Baumslag, *Some aspects of groups with unique roots*, Acta Math. **104** (1960), 217–303.
- [2] C. Bonatti and L. Paris, *Roots in the mapping class groups*, Proc. Lond. Math. Soc. **98** (2009), 471–503.
- [3] M.R. Bridson and A. Häflicher, *Metric spaces of non-positive curvature*, Grundlehren Math. Wiss. **319**, Springer-Verlag Berlin Heidelberg, 1999.
- [4] R. Charney, *An introduction to right-angled Artin groups*, Geom. Dedicata **125** (2007), 141–158.
- [5] J. Crisp, E. Godelle and B. Wiest, *The conjugacy problem in subgroups of right-angled Artin groups*, J. Topology **2** (2009), 442–460.
- [6] G. Duchamp and J.-Y. Thibon, *Simple orderings for free partially commutative groups*, Internat. J. Algebra Comput. **2** (1992), 351–355.
- [7] E. Ghys and P. de la Harpe, *Sur les groupes hyperboliques d’après Mikhael Gromov*, Progress in Mathematics **83**, Birkhäuser, Boston, 1990.
- [8] J. González-Meneses, *The  $n$ th root of a braid is unique up to conjugacy*, Algebr. Geom. Topol. **3** (2003), 1103–1118.
- [9] D. Kim and D. Rolfsen, *An ordering for groups of pure braids and fibre-type hyperplane arrangements*, Canad. J. Math. **55** (2003), 822–838.
- [10] S.-h. Kim and T. Koberda, *Embedability between right-angled Artin groups*, Geom. Topol. **17** (2013), 493–530.
- [11] S.-h. Kim and T. Koberda, *The geometry of the curve complex of a right-angled Artin group*, Internat. J. Algebra Comput. **24** (2014), 121–169.
- [12] S.-h. Kim and T. Koberda, *Anti-trees and right-angled Artin subgroups of braid group*, Geom. Topol. **19** (2015), 3289–3306.
- [13] E.-K. Lee and S.-J. Lee, *Uniqueness of roots up to conjugacy for some affine and finite type Artin groups*, Math. Z. **265** (2010), 571–587.
- [14] E.-K. Lee and S.-J. Lee, *Path lifting properties and embedding between RAAGs*, J. Algebra **448** (2016), 575–594.
- [15] E.-K. Lee and S.-J. Lee, *Embeddability of right-angled Artin groups on complements of trees*, Internat. J. Algebra Comput. **28** (2018), 381–394.
- [16] G.S. Makanin, *On normalizers in the braid group*, Mat. Sb. **86** (1971), 171–179.
- [17] A.I. Mal’cev, *Nilpotent torsion-free groups*, Izv. Akad. Nauk SSSR Ser. Mat. **13** (1949), 201–212.
- [18] D. Rolfsen and J. Zhu, *Braids, orderings and zero divisors*, J. Knot Theory Ramifications **7** (1998), 837–841.
- [19] H. Servatius, *Automorphisms of graph groups*, J. Algebra **126** (1989), 34–60.

DEPARTMENT OF MATHEMATICS AND STATISTICS, SEJONG UNIVERSITY, SEOUL, KOREA  
*Email address:* `eonkyung@sejong.ac.kr`

DEPARTMENT OF MATHEMATICS, KONKUK UNIVERSITY, SEOUL, KOREA  
*Email address:* `sangjin@konkuk.ac.kr`