

# PULLBACK FORMULAS FOR ARITHMETIC CYCLES

BENJAMIN HOWARD

ABSTRACT. On an orthogonal Shimura variety, one has a collection of special cycles in the Gillet-Soulé arithmetic Chow group. We describe how these cycles behave under pullback to an embedded orthogonal Shimura variety of lower dimension. The bulk of the paper is devoted to cases in which the special cycles intersect the embedded Shimura variety improperly, for which we construct logarithmic expansions of certain Green currents on the deformation to the normal bundle of the embedding.

## 1. INTRODUCTION

On an orthogonal Shimura variety  $M$ , one has a systematic supply of special cycles coming from embeddings of smaller orthogonal Shimura varieties. These cycles are the subject of a series of conjectures of Kudla [Kud04], whose central theme is that they should be geometric analogues of the coefficients of Siegel theta functions. For example, it is known by work of Bruinier and Raum [BWR15] that these cycles are the coefficients of a Siegel modular form valued in the Chow group of  $M$ .

In order to do arithmetic intersection theory with these cycles, one must endow them with Green currents. One construction of such Green currents was done by Garcia-Sankaran [GS19], using ideas of Bismut [Bis90] and Bismut-Gillet-Soulé [BGS90], and the special cycles endowed with these currents define arithmetic special cycles in the Gillet-Soulé arithmetic Chow group of  $M$ . The goal of this paper is to show that these arithmetic special cycles behave nicely under pullbacks via embeddings  $M_0 \rightarrow M$  of smaller orthogonal Shimura varieties, in the sense that the pullback of an arithmetic special cycle on  $M$  is a prescribed linear combination of arithmetic special cycles on  $M_0$ .

When an arithmetic cycle intersects  $M_0$  properly, its pullback can be defined in a naive way, and is easy to compute directly from the definitions. Unfortunately, the intersections that arise in our setting are very rarely proper. For improper intersections, Gillet and Soulé define pullbacks using the moving lemma, which is poorly suited to any kind of explicit calculation. One doesn't have any natural choice of rationally trivial cycle with which to move the special cycle, and even if one did, replacing an arithmetic special

---

This research was supported in part by NSF grant DMS-2101636.

cycle by a rationally equivalent one would destroy all the nice properties of the special cycle and Green current that one started with.

Our approach to treating improper intersections is to use the work of Hu [Hu99], which gives an alternate definition of arithmetic pullbacks based on Fulton's deformation to the normal cone approach to intersection theory. One can always specialize a cycle on  $M$  to a cycle on the normal cone to  $M_0 \subset M$ , which, in our setting, is just (the total space of) the normal bundle  $N_{M_0/M} \rightarrow M_0$ . Hu shows that there is an analogous specialization of Green currents. The core of this paper is the calculation of the specializations of Garcia-Sankaran Green currents to  $N_{M_0/M}$ , or at least the calculation of enough of them to deduce the pullback formula.

**1.1. Statement of the main result.** Fix a quadratic space  $V$  of dimension  $n + 2 \geq 3$  over a totally real number field  $F$ . Assume that  $V$  has signature  $(n, 2)$  at one embedding  $\sigma : F \rightarrow \mathbb{R}$ , but is positive definite at every other embedding.

From  $V$  one can construct a Shimura datum  $(G, \mathcal{D})$  in which  $G$  is the restriction of scalars to  $\mathbb{Q}$  of either  $\mathrm{SO}(V)$  or  $\mathrm{GSpin}(V)$ , and  $\mathcal{D}$  is a hermitian symmetric domain of dimension  $n$ . Fixing a sufficiently small compact open subgroup  $K \subset G(\mathbb{A}_f)$ , we obtain a quasi-projective Shimura variety  $M$  over the reflex field  $\sigma(F) \subset \mathbb{C}$  with complex points

$$M(\mathbb{C}) = G(\mathbb{Q}) \backslash \mathcal{D} \times G(\mathbb{A}_f) / K.$$

For the rest of the introduction we assume that  $V$  is anisotropic, so that  $M$  is projective.

Fix a positive integer  $d \leq n+1$ . Given the data of a nonsingular symmetric matrix  $T \in \mathrm{Sym}_d(F)$  and a  $K$ -fixed Schwartz function  $\varphi \in \mathcal{S}(\widehat{V}^d)$ , one can define a special cycle  $Z(T, \varphi)$  on  $M$  of codimension  $d$ , as in [Kud04]. After fixing a positive definite  $v \in \mathrm{Sym}_d(\mathbb{R})$ , Garcia-Sankaran [GS19] construct a smooth form  $\mathfrak{g}^\circ(T, v, \varphi)$  of type  $(d-1, d-1)$  on  $M(\mathbb{C}) \setminus Z(T, \varphi)(\mathbb{C})$ . This form is locally integrable on  $M(\mathbb{C})$ , and its associated current satisfies the Green equation

$$dd^c[\mathfrak{g}^\circ(T, v, \varphi)] + \delta_{Z(T, \varphi)} = [\omega^\circ(T, v, \varphi)]$$

for a smooth form  $\omega^\circ(T, v, \varphi)$ . We obtain an arithmetic cycle class

$$(1.1.1) \quad \widehat{C}_M(T, v, \varphi) = (Z(T, \varphi), \mathfrak{g}^\circ(T, v, \varphi)) \in \widehat{\mathrm{CH}}^d(M)$$

in the Gillet-Soulé arithmetic Chow group.

Because our main focus is on the Green currents, in this paper we work exclusively with the arithmetic Chow group of the canonical model over the reflex field. No integral models will appear.

Following [GS19], in §4.2 we extend the definition of (1.1.1) to all  $T \in \mathrm{Sym}_d(F)$ . For the purposes of this introduction, we say only that this extension makes use of a special hermitian line bundle

$$\widehat{\omega}^{-1} \in \widehat{\mathrm{Pic}}(M) \cong \widehat{\mathrm{CH}}^1(M).$$

For example, in the degenerate case of the zero matrix  $0_d \in \text{Sym}_d(F)$ ,

$$\widehat{C}(0_d, v, \varphi) = \varphi(0) \cdot \underbrace{[\widehat{\omega}^{-1} \cdots \widehat{\omega}^{-1}]_d} + (0, -\log(\det(v)) \cdot \Omega^{d-1})$$

where the  $\cdots$  on the right hand side is iterated arithmetic intersection, and  $\Omega^{d-1}$  is the  $d-1$  exterior power of the Chern form of  $\widehat{\omega}^{-1}$ .

Now suppose that our quadratic space is presented as an orthogonal direct sum  $V = V_0 \oplus W$  with  $W$  totally positive definite and  $\dim(V_0) \geq 3$ . In particular,  $V_0$  has signature  $(n_0, 2)$  at the real embedding  $\sigma : F \rightarrow \mathbb{R}$ , and is positive definite at all other embeddings. As such, it has its own Shimura datum  $(G_0, \mathcal{D}_0)$ , and a choice of compact open  $K_0 \subset G_0(\mathbb{A}_f)$  determines a Shimura variety  $M_0$  over  $\sigma(F) \subset \mathbb{C}$  with its own family of arithmetic special cycles

$$\widehat{C}_{M_0}(T, v, \varphi_0) \in \widehat{\text{CH}}^d(M_0).$$

The inclusion  $V_0 \rightarrow V$  induces an embedding of Shimura data  $(G_0, \mathcal{D}_0) \rightarrow (G, \mathcal{D})$ . We choose  $K_0$  and  $K$  in such a way that  $K_0 \subset G_0(\mathbb{A}_f) \cap K$ , and so that the induced  $i_0 : M_0 \rightarrow M$  is a closed immersion. Our main result, stated in the text as Corollary 4.3.3, is the following pullback formula for arithmetic special cycles.

**Theorem A.** *The pullback  $i_0^* : \widehat{\text{CH}}^d(M) \rightarrow \widehat{\text{CH}}^d(M_0)$  satisfies*

$$i_0^* \widehat{C}_M(T, v, \varphi) = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \sum_j \psi_j(y) \cdot \widehat{C}_{M_0}(T_0, v, \varphi_{0,j})$$

for any  $T \in \text{Sym}_d(F)$ , positive definite  $v \in \text{Sym}_d(\mathbb{R})$ , and  $K$ -fixed

$$\varphi = \sum_j \varphi_{0,j} \otimes \psi_j \in S(\widehat{V}_0^d) \otimes S(\widehat{W}^d) \cong S(\widehat{V}^d).$$

If one forgets Green currents and works with the usual Chow groups of  $M$  and  $M_0$ , similar pullback formulas are proved in [Kud21].

We have not attempted to exhaust the methods, which can presumably be pushed much further. For example, one would like a similar statement for noncompact Shimura varieties and integral models, as well as a formula expressing the intersection of two arithmetic special cycles as a linear combination of other arithmetic special cycles. There should be similar results for the Shimura varieties associated to quadratic spaces with signature  $(n, 2)$  at several archimedean places. One could also try to prove similar formulas for other Green currents, such as those of Funke-Hofmann [FH21]. The author hopes to address some of these questions in future work.

**1.2. Outline of the paper.** In §2 we recall what we need from Hu's thesis [Hu99]. Suppose  $X_0 \rightarrow X$  is a closed immersion of complex manifolds. If  $G$  is a Green current for an analytic cycle  $Z$  on  $X$ , one would like to construct a

Green current  $\sigma_{X_0/X}(G)$  for the specialization  $\sigma_{X_0/X}(Z)$  of  $Z$  to the normal bundle  $N_{X_0/X}$ .

To see how this works, denote by  $\tilde{X}$  the deformation of the normal bundle  $N_{X_0/X}$ . It comes with a holomorphic function  $\tau : \tilde{X} \rightarrow \mathbb{C}$  whose fiber over  $t \in \mathbb{C}$  we denote by  $\tilde{X}_t$ . If  $t \neq 0$  there is a canonical closed immersion  $j_t : X \cong \tilde{X}_t \subset \tilde{X}$ , and we obtain a current  $j_{t*}G$  on  $\tilde{X}$ . In essence, Hu's idea is to look for a logarithmic expansion

$$j_{t*}G = \sum_{i \geq 0} G_i(t) \cdot (\log |t|)^i$$

whose coefficients  $G_i(t)$  are currents on  $\tilde{X}$ , with the property that each function  $t \mapsto G_i(t)$  extends continuously to  $t = 0$ , and define  $\sigma_{X_0/X}(G)$  in terms of the current  $G_0(0)$ . In this generality such a logarithmic expansion need not exist. If it does exist the logarithmic expansion will not be unique, but  $\sigma_{X_0/X}(G)$  is independent of the choice.

In Theorems 2.2.5 and 2.3.1 we quote two results of Hu. The first guarantees the existence of logarithmic expansions (and hence specializations to the normal bundle) for certain currents on  $X$ . The second shows that if  $X$  is a projective variety over a number field, one can use the specialization of cycles and Green currents to define a morphism from the arithmetic Chow group of  $X$  to the arithmetic Chow group of  $N_{X_0/X}$ .

Now suppose  $L$  a hermitian line bundle on  $X$ . In §3 we recall from [GS19] a construction that takes a tuple  $s = (s_1, \dots, s_d)$  of global sections of  $L^\vee$  and produces a Green current  $\mathfrak{g}^\circ(s)$  for the analytic cycle  $Z(s)$  defined by  $s_1 = \dots = s_d = 0$ . The central problem is to understand the specialization  $\sigma_{X_0/X}(\mathfrak{g}^\circ(s))$ . For our applications it is enough to assume that  $s = (p, q)$  is the concatenation of tuples  $p = (p_1, \dots, p_k)$  and  $q = (q_1, \dots, q_\ell)$ , arranged so that the cycle  $Z(p)$  meets  $X_0$  properly, while  $X_0 \subset Z(q)$ .

In essence, our strategy is to show that the star product formula

$$(1.2.1) \quad \mathfrak{g}^\circ(s) = \mathfrak{g}^\circ(p) \star \mathfrak{g}^\circ(q_1) \star \dots \star \mathfrak{g}^\circ(q_\ell)$$

of Garcia-Sankaran implies the analogous star product formula

$$(1.2.2) \quad \sigma_{X_0/X}(\mathfrak{g}^\circ(s)) = \sigma_{X_0/X}(\mathfrak{g}^\circ(p)) \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q_1)) \star \dots \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q_\ell))$$

for specializations, and then compute each specialization on the right individually. As  $Z(p)$  intersects  $X_0$  properly, the specialization  $\sigma_{X_0/X}(\mathfrak{g}^\circ(p))$  is easy to compute. To compute the specialization of  $\mathfrak{g}^\circ(q_i)$  one must do more work, but the idea is imitate the construction of the current with  $X$  replaced by  $\tilde{X}$ , and use the resulting current on the deformation to the normal bundle to find an explicit logarithmic expansion for  $\mathfrak{g}^\circ(q_i)$ .

It is not obvious to the author that Hu's specialization to the normal bundle is compatible with  $\star$  products in general; that is to say, deducing (1.2.2) from (1.2.1) seems to require using particular properties of the Green currents  $\mathfrak{g}^\circ(s)$ . After passing to arithmetic Chow groups the compatibility of specialization with star products follows from Theorem 2.3.1, but on the

level of arithmetic cycles (that is, before passing to their rational equivalence classes) things are more complicated. When we apply the calculations described above to the case of orthogonal Shimura varieties, the complex manifold  $X$  is not the Shimura variety  $M(\mathbb{C})$ , but rather the hermitian symmetric domain  $\mathcal{D}$  that uniformizes it. As  $\mathcal{D}$  does not have an arithmetic Chow group in any useful sense, our calculations must be carried out before passing to rational equivalence classes of arithmetic cycles.

To prove the compatibility of specializations and star products we need to show that the Green currents in question admit logarithmic expansions of an especially nice form; this is essentially Lemma 3.3.3, which is the core of the proof of Proposition 3.3.7. While Hu’s proof of Theorem 2.3.1 provides a general construction of logarithmic expansions, the expansions one gets from this method are quite unpleasant. For example, if one starts with a current  $G$  that is represented by a locally integrable form, the currents  $G_i(t)$  produced by Hu’s construction will typically not have this form. It is essential to our method that we find logarithmic expansions for  $\mathfrak{g}^\circ(s)$  that are better behaved than those obtained by tracing through Hu’s proof of Theorem 2.3.1.

Finally, in §4 we define the precise arithmetic cycle classes  $\widehat{C}(T, v, \varphi)$  appearing in Theorem A, and apply the general constructions of the preceding sections to the case of orthogonal Shimura varieties. The strategy for proving Theorem A is then to use the explicit calculation of specializations of cycles and Green currents to show that the arithmetic cycles appearing in the equality of that theorem become equal after pullback via the bundle map  $N_{M_0/M} \rightarrow M_0$ . By Proposition 2.3.3, they must have been equal before the pullback as well.

The technical calculations in §3 are carried out in the setting of an arbitrary complex manifold  $X$ , and don’t involve orthogonal Shimura varieties (or any Shimura varieties) at all. Because of this, there is no difficulty in extending Theorem A to include, for example, Shimura varieties of type  $U(n, 1)$ . We leave the formulation of the theorem in this setting as an exercise for the reader.

As a final remark, we note that something like specializations to the normal bundle of Green currents were computed in [AGHMP17], but for Bruinier’s [Bru02] Green functions for special divisors. When working only with arithmetic divisors, the situation is much simpler, and one doesn’t really need specialization to the normal bundle at all. The codimension one arithmetic Chow group can be identified with the group of hermitian line bundles, and pullback then agrees with the naive notion of pullback of line bundles. One can compute arithmetic pullbacks (even in cases of improper intersection) more directly using this interpretation. This is the approach taken in [BHY15], which is the unitary Shimura variety analogue of [AGHMP17]. To compute pullbacks for higher codimension arithmetic Chow groups, the author knows of no method other than the specialization to the normal bundle approach developed here.

**1.3. Acknowledgements.** The author thanks Jan Bruinier, Luis Garcia, José Burgos Gil, and Siddarth Sankaran for helpful discussions, some of which took place during the AIM workshop *Arithmetic intersection theory on Shimura varieties*, January 4-8, 2021. The author thanks AIM for facilitating these conversations.

## 2. ARITHMETIC INTERSECTION THEORY

In this section we recall some results from Hu's thesis [Hu99], and restate them in the precise form they will be needed later.

**2.1. Logarithmic differential forms.** We recall some definitions and results from [Bur94]. Let  $X$  be a complex manifold of dimension  $n = \dim(X)$ .

**Definition 2.1.1.** If  $Z \subset X$  is any analytic subset (i.e., a reduced closed analytic subspace), a *resolution of singularities*

$$(2.1.1) \quad r : (X', Z') \rightarrow (X, Z)$$

is a complex manifold  $X'$  together with a proper surjection  $r : X' \rightarrow X$ , such that  $Z' = r^{-1}(Z)$  is a divisor with normal crossings, and  $r$  restricts to an isomorphism  $X' \setminus Z' \cong X \setminus Z$ .

*Remark 2.1.2.* A resolution of singularities always exists by Theorem 3.3.5 of [Kol07], extended to analytic spaces as in §3.4.4 of *loc. cit.*. See also [Wlo09]. For quasi-projective varieties, this is Hironaka's theorem.

Denote by

$$E_X^\bullet = \bigoplus_{k \geq 0} E_X^k$$

the graded  $\mathbb{C}$ -algebra of smooth differential forms on  $X$ . For  $\phi \in E_X^\bullet$ , let

$$\phi_{[k]} \in E_X^k$$

be its component in degree  $k$ . Let  ${}_c E_X^\bullet \subset E_X^\bullet$  be the graded subspace of compactly supported forms, and denote by  $D_X^k$  be the space of currents dual to  ${}_c E_X^{2n-k}$ . There is a canonical injection  $E_X^k \rightarrow D_X^k$ , denoted  $g \mapsto [g]$ , defined by

$$(2.1.2) \quad [g](\phi) = \int_X g \wedge \phi.$$

When no confusion can arise, we sometimes omit the brackets, and write  $g$  both for the form and its associated current.

Given a divisor with normal crossings  $Z \subset X$ , let

$$E_X^\bullet(\log Z) \subset E_{X \setminus Z}^\bullet$$

be the graded subalgebra of forms with logarithmic growth along  $Z$  as in §1.2 of [Bur94]: in local coordinates on  $X$  such that  $Z$  is given by the equation

$z_1 \cdots z_m = 0$ , the forms of logarithmic growth are generated, as an algebra over the ring of smooth forms, by

$$\log |z_i|, \frac{dz_i}{z_i}, \frac{d\bar{z}_i}{\bar{z}_i} \quad \text{for } 1 \leq i \leq m.$$

Now let  $Z \subset X$  be an analytic subset of codimension  $d$ . A choice of resolution of singularities (2.1.1) determines a subspace

$$E_X^\bullet(\log Z) \subset E_{X \setminus Z}^\bullet,$$

consisting of those forms whose pullbacks to  $E_{X' \setminus Z'}$  have logarithmic growth along the normal crossing divisor  $Z'$ . Although the notation hides it, this subspace genuinely depends on the choice of resolution of singularities.

Denote by

$$E_X^\bullet(\text{null } Z) \subset E_X^\bullet$$

the graded subspace of forms whose pullback to (the smooth locus of)  $Z$  vanishes, and let  ${}_c E_X^\bullet(\text{null } Z)$  be the graded subspace of compactly supported such forms. The inclusion above induces a canonical surjection

$$D_X^\bullet \rightarrow D_{X/Z}^\bullet,$$

where  $D_{X/Z}^k$  is the space of currents dual to  ${}_c E_X^{2n-k}(\text{null } Z)$ .

**Proposition 2.1.3** (Burgos). *For any  $g \in E_X^\bullet(\log Z)$  and  $\phi \in E_X^\bullet(\text{null } Z)$ , the form  $g \wedge \phi$  is locally integrable on  $X$ . The integral (2.1.2) defines an injection*

$$E_X^\bullet(\log Z) \xrightarrow{g \mapsto [g]} D_{X/Z}^\bullet$$

satisfying  $\partial[g] = [\partial g]$ , and similarly for  $\bar{\partial}$ .

*Proof.* See Corollaries 3.7 and 3.8 in [Bur94].  $\square$

*Remark 2.1.4.* If  $k < 2d$  then any form in  $E_X^{2n-k}$  has trivial pullback to  $Z$ , and hence  $D_{X/Z}^k = D_X^k$ . In particular, we obtain an injection

$$E_X^k(\log Z) \xrightarrow{g \mapsto [g]} D_X^k.$$

For  $g \in E_X^k(\log Z)$  with  $k < 2d - 1$  we have  $\partial[g] = [\partial g]$  in  $D_X^{k+1}$ , and similarly for  $\bar{\partial}$ .

**Definition 2.1.5.** Suppose that  $U$  is a smooth quasi-projective complex variety. By a *smooth compactification* of  $U$  we mean a smooth projective variety  $U^*$ , a divisor with normal crossings  $\partial U^* \subset U^*$ , and an isomorphism  $i : U \cong U^* \setminus \partial U^*$ . The smooth compactifications of  $U$  form a cofiltered category in a natural way, allowing us to form the graded subalgebra

$$E_{\log}^\bullet(U) = \varinjlim_{(U^*, \partial U^*, i)} E_{U^*}^\bullet(\log \partial U^*) \subset E_U^\bullet$$

of forms with logarithmic singularities along  $\infty$ ; see Definition 1.2 of [Bur97] and the discussion surrounding it.

*Remark 2.1.6.* Of special interest is the case in which  $X$  is a smooth quasi-projective complex variety,  $Z \subset X$  is a closed subvariety of codimension  $d$ , and we take  $U = X \setminus Z$ . In this case, for any

$$g \in E_{\log}^k(X \setminus Z)$$

there is a resolution of singularities (2.1.1) such that  $g \in E_X^k(\log Z)$ . When  $k < 2d$ , Remark 2.1.4 therefore provides us with an injection

$$E_{\log}^k(X \setminus Z) \xrightarrow{g \rightarrow [g]} D_X^k.$$

In the usual way, the complex structure on  $X$  induces bigradings

$$E_X^k = \bigoplus_{p+q=k} E_X^{p,q} \quad \text{and} \quad D_X^k = \bigoplus_{p+q=k} D_X^{p,q},$$

and similarly for the other spaces of forms and currents appearing above.

**2.2. Specialization to the normal bundle.** For a closed immersion of schemes  $X_0 \subset X$  one has the normal cone  $C_{X_0/X} \rightarrow X_0$ . If  $X_0 \subset X$  is a regular immersion, the normal cone agrees with (the total space of) the normal bundle  $N_{X_0/X} \rightarrow X_0$ . These constructions, as well as the deformation to the normal cone, generalize in an obvious way to a closed immersion of complex analytic spaces; the necessary technical details are in [AM86].

Now suppose that  $X_0 \subset X$  is a closed immersion of complex manifolds. Denote by  $\tilde{X}$  the deformation to  $N_{X_0/X} = C_{X_0/X}$ . By construction, it comes with morphisms

$$(2.2.1) \quad X \xleftarrow{\pi} \tilde{X} \xrightarrow{\tau} \mathbb{C}$$

such that  $\pi$  identifies every fiber  $\tilde{X}_t = \tau^{-1}(t)$  with

$$\tilde{X}_t \cong \begin{cases} X & \text{if } t \neq 0 \\ N_{X_0/X} & \text{if } t = 0. \end{cases}$$

When  $t \neq 0$  we denote by

$$j_t : X \cong \tilde{X}_t \hookrightarrow \tilde{X}$$

the inclusion, and similarly for  $j_0 : N_{X_0/X} \cong \tilde{X}_0 \hookrightarrow \tilde{X}$ .

Let  $Z \subset X$  be any equidimensional analytic subset, endowed with its reduced complex analytic structure. The *strict transform*

$$(2.2.2) \quad \tilde{Z} \subset \tilde{X}$$

of  $Z$  is defined as the deformation to the normal cone  $C_{(Z \times_X X_0)/Z}$ . It is again reduced (although  $Z \times_X X_0$  and the normal cone  $C_{(Z \times_X X_0)/Z}$  need not be), and can be characterized as the union of all irreducible components of  $\pi^{-1}(Z)$  not contained in  $N_{X_0/X}$ . Equivalently, (2.2.2) is the closure of  $\pi^{-1}(Z) \setminus \tilde{X}_0$  in  $\tilde{X}$ .

By an *analytic cycle* on a complex manifold we mean a locally finite formal  $\mathbb{Z}$ -linear combination of irreducible analytic subsets, all of the same



codimension. Being reduced and equidimensional, we may view  $\tilde{Z}$  as an analytic cycle on  $\tilde{X}$  and form, for every  $t \in \mathbb{C}$ , the analytic cycle

$$(2.2.3) \quad \tilde{Z}_t = \tilde{Z} \cdot \tilde{X}_t$$

on  $\tilde{X}$  supported on the fiber  $\tilde{X}_t$ . Here the proper intersection of analytic cycles on the right is taken in the sense of Draper [Dra69]. Of special interest is the analytic cycle (2.2.3) at  $t = 0$ .

**Definition 2.2.1.** The analytic cycle

$$(2.2.4) \quad \sigma_{X_0/X}(Z) \stackrel{\text{def}}{=} \tilde{Z}_0$$

on  $N_{X_0/X} = \tilde{X}_0$  is the *specialization to the normal bundle* of  $Z$ .

Having defined  $\tilde{Z}$ ,  $\tilde{Z}_t$ , and  $\sigma_{X_0/X}(Z)$  for a reduced analytic subset  $Z \subset X$ , extend the definitions linearly to all analytic cycles  $Z$  on  $X$ .

*Remark 2.2.2.* The results of [Kin71] allow us to characterize the family of cycles (2.2.3) using continuity of fiber integration. If  $t \neq 0$  then  $\tilde{Z}_t$  is simply the pushforward of  $Z$  under  $j_t : X \cong \tilde{X}_t \hookrightarrow \tilde{X}$ . The cycle  $\tilde{Z}_0$ , which may be nonreduced, is then uniquely determined by the continuity at  $t = 0$  of

$$\delta_{\tilde{Z}_t}(\phi) = \int_{\tilde{Z}_t} \phi,$$

for every  $\phi \in {}_cE_{\tilde{X}}^{2\dim(Z)}$ .

*Remark 2.2.3.* In the case where  $X$  and  $Z$  are the complex analytic spaces associated to finite type schemes over  $\mathbb{C}$ , the analytic intersection (2.2.3) agrees with the proper intersection of cycles in the algebraic sense of [Ful84] or [Sou92], and the cycle (2.2.4) agrees with the specialization to the normal cone in the sense [Ful84].

**Definition 2.2.4.** Fix a  $G \in D_X^k$ , and note that every  $t \in \mathbb{C} \setminus \{0\}$  determines a current

$$j_{t*}G \in D_{\tilde{X}}^{k+2}.$$

We say that  $G$  admits a *logarithmic expansion along  $X_0$*  if there is a sequence of functions  $G_0, G_1, G_2, \dots : \mathbb{C} \rightarrow D_{\tilde{X}}^{k+2}$  with the following properties.

- (1) For all  $t \in \mathbb{C} \setminus \{0\}$  we have

$$j_{t*}G = \sum_{i \geq 0} G_i(t) \cdot (\log |t|)^i,$$

and the sum is locally finite: for every compact subset  $K \subset \tilde{X}$  there is an integer  $M_K$  such that  $G_i(t)(\phi) = 0$  for all  $i > M_K$ , all  $t \in \mathbb{C} \setminus \{0\}$ , and all  $\phi \in E_{\tilde{X}}^{2n-k}$  with support contained in  $K$ .

- (2) For every  $i \geq 0$  and every  $\phi \in {}_cE_{\tilde{X}}^{2n-k}$ , the function  $t \mapsto G_i(t)(\phi)$  is continuous at  $t = 0$ , and is Hölder continuous at  $t = 0$  if  $i > 0$ .
- (3) Each  $G_i(0)$  lies in the image of  $j_{0*} : D_{\tilde{X}_0}^k \rightarrow D_{\tilde{X}}^{k+2}$ .

The following result is slightly weaker than Theorem 3.2.2 of [Hu99]; see Remark 2.2.11 below. It provides a general criterion for the existence of logarithmic expansions.

**Theorem 2.2.5** (Hu). *If  $Z \subset X$  is an equidimensional analytic subset of codimension  $d$ , and  $g \in E_X^k(\log Z)$  (for some choice of resolution of singularities) is locally integrable on  $X$ , then the current  $[g] \in D_X^k$  of Remark 2.1.4 admits a logarithmic expansion along  $X_0$ . If  $X$  is compact, there exists a logarithmic expansion with  $G_i = 0$  for  $i \gg 0$ .*

*Remark 2.2.6.* Hu works on smooth quasi-projective complex varieties, but the same proof works for complex manifolds. The only difference is that in the quasi-projective case one can use the existence of smooth compactifications of  $X$  and  $\tilde{X}$  to prove the existence of finite (not just locally finite) logarithmic expansions. See Remark 2.2.11 below.

*Remark 2.2.7.* Suppose we are given functions  $f_0, \dots, f_m : \mathbb{C} \rightarrow \mathbb{C}$  with  $f_0$  continuous at 0, and  $f_1, \dots, f_m$  Hölder continuous at 0. An easy induction on  $m$ , as in Lemma 3.1.5 of [Hu99], shows that if

$$\lim_{t \rightarrow 0} \sum_{i=0}^m f_i(t) \cdot (\log |t|)^i = 0,$$

then  $f_i(0) = 0$  for all  $i$ .

The functions  $G_i$  in Definition 2.2.4, when they exist, are not uniquely determined. However, it follows from Remark 2.2.7 that their values  $G_i(0)$  at  $t = 0$  are. This allows us to make the following definition.

**Definition 2.2.8.** If  $G \in D_X^k$  admits a logarithmic expansion along  $X_0$ , its *specialization to the normal bundle* is the current

$$\sigma_{X_0/X}(G) \in D_{N_{X_0/X}}^k$$

characterized by  $j_{0*}\sigma_{X_0/X}(G) = G_0(0)$ .

As a trivial example, if  $g \in E_X^k$  then  $\pi^*g$  is a smooth form on  $\tilde{X}$ , and

$$j_{t*}[g](\phi) = \int_{\tilde{X}_t} \pi^*g \wedge \phi$$

for all  $t \neq 0$ . By Remark 2.2.2, the right hand side is a continuous function of  $t \in \mathbb{C}$ , and we obtain a logarithmic expansion of  $[g]$  by setting  $G_0(t) = \pi^*g \wedge \delta_{\tilde{X}_t}$  and  $G_i(t) = 0$  for  $i > 0$ . In particular,

$$\sigma_{X_0/X}(g) = [j_0^*\pi^*g] = [\pi_0^*i_0^*g],$$

where  $\pi_0 : N_{X_0/X} \rightarrow X_0$  is the bundle map and  $i_0 : X_0 \rightarrow X$  is the inclusion.

*Remark 2.2.9.* If  $G \in D_X^k$  admits a logarithmic expansion along  $X_0$ , then so does  $\partial G$ , and

$$\partial\sigma_{X_0/X}(G) = \sigma_{X_0/X}(\partial G).$$

The same holds with  $\partial$  replaced by  $\bar{\partial}$ . This is a formal consequence of the definitions; see Theorem 3.1.6 of [Hu99].

The following proposition connects Definitions 2.2.1 and 2.2.8. The proof is extracted from the second proof of Theorem 3.2.3 in [Hu99].

**Proposition 2.2.10.** *Suppose  $X_0 \subset X$  is a closed complex submanifold,  $Z$  is a codimension  $d$  analytic cycle on  $X$ , and  $G \in D_X^{d-1, d-1}$  satisfies the Green equation*

$$dd^c G + \delta_Z = [\omega]$$

for some  $\omega \in E_X^{d, d}$ . If  $G$  admits a logarithmic expansion along  $X_0$ , then its specialization to the normal bundle satisfies the Green equation

$$dd^c \sigma_{X_0/X}(G) + \delta_{\sigma_{X_0/X}(Z)} = [\pi_0^* i_0^* \omega].$$

Here  $i_0 : X_0 \rightarrow X$  is the inclusion and  $\pi_0 : N_{X_0/X} \rightarrow X_0$  is the bundle map.

*Proof.* When  $t \neq 0$ , we may push forward the Green equation for  $G$  via  $j_t : X \rightarrow \tilde{X}$ . This yields the equality

$$dd^c j_{t*} G + \delta_{\tilde{Z}_t} = \pi^* \omega \wedge \delta_{\tilde{X}_t}$$

of currents on  $\tilde{X}$ . Replacing  $j_{t*} G$  by a logarithmic expansion results in

$$\left( dd^c G_0(t) + \delta_{\tilde{Z}_t} - \pi^* \omega \wedge \delta_{\tilde{X}_t} \right) + \sum_{i>0} dd^c G_i(t) \cdot (\log |t|)^i = 0,$$

and it follows from Remarks 2.2.2 and 2.2.7 that

$$dd^c G_0(0) + \delta_{\tilde{Z}_0} - \pi^* \omega \wedge \delta_{\tilde{X}_0} = 0.$$

The claim now follows using  $\pi^* \omega \wedge \delta_{\tilde{X}_0} = j_{0*} [j_0^* \pi^* \omega] = j_{0*} [\pi_0^* i_0^* \omega]$ .  $\square$

*Remark 2.2.11.* In Hu's version of Theorem 2.2.5 it is assumed that  $X$  is a quasi-projective variety, and that  $g \in E_{\log}^k(X \setminus Z)$ . These extra assumptions are not used in the proof in any essential way. However, the first guarantees the existence of a smooth compactification of  $X$ . Using this, Hu proves a stronger result than what we have stated.

After choosing a smooth compactification  $X \subset X^*$ , Hu constructs a smooth compactification  $\tilde{X} \subset \tilde{X}^*$ , a diagram

$$X^* \xleftarrow{\pi} \tilde{X}^* \xrightarrow{\tau} \mathbb{C}$$

extending (2.2.1), and a finite expansion of currents

$$(2.2.5) \quad \pi^* g \wedge \delta_{\tilde{X}^*} = \sum_{i=0}^M G_i(t) \cdot (\log |t|)^i$$

in the space  $D_{\tilde{X}^*/\partial\tilde{X}^*}^{k+2}$ . The inclusion  ${}_c E_{\tilde{X}}^\bullet \rightarrow {}_c E_{\tilde{X}^*}^\bullet(\text{null } \partial\tilde{X}^*)$  induces a surjection

$$D_{\tilde{X}^*/\partial\tilde{X}^*}^\bullet \rightarrow D_{\tilde{X}}^\bullet,$$

and applying this map to both sides of (2.2.5) yields a finite logarithmic expansion of  $[g]$ . The refined logarithmic expansion (2.2.5) contains more information than a logarithmic expansion in our sense. Using it, Hu is able to construct a smooth compactification  $N_{X_0/X} \subset N_{X_0/X}^*$  and a distinguished lift of  $\sigma_{X_0/X}(g)$  under the surjection

$$D_{N_{X_0/X}^*/\partial N_{X_0/X}}^\bullet \rightarrow D_{N_{X_0/X}}^\bullet.$$

Although we will not need such a lift, the benefits of having one are explained in Remark 2.3.2.

**2.3. Arithmetic Chow groups.** We will use the arithmetic Chow groups defined in §3 of [GS90], but only in the simple case of varieties over a field  $F$  with a chosen real embedding  $\sigma : F \rightarrow \mathbb{R}$ . If we let  $c \in \text{Aut}(\mathbb{C}/\mathbb{R})$  be complex conjugation, the triple  $(F, \{\sigma\}, c)$  is an arithmetic ring, and any smooth quasi-projective variety  $X$  over  $F$  is an arithmetic variety over  $(F, \{\sigma\}, c)$  in the sense of *loc. cit.*

Let  $X_{\mathbb{R}} = X \otimes_{F,\sigma} \mathbb{R}$  be the base change of  $X$  to  $\mathbb{R}$ , and regard  $X(\mathbb{C}) = X_{\mathbb{R}}(\mathbb{C})$  as a complex manifold. Define a real vector space

$$E_X^{d,d} = \{\omega \in E_{X(\mathbb{C})}^{d,d} : \omega \text{ is real and } c^*\omega = (-1)^d\omega\},$$

where now  $c : X(\mathbb{C}) \rightarrow X(\mathbb{C})$  is complex conjugation, and similarly

$$D_X^{d,d} = \{G \in D_{X(\mathbb{C})}^{d,d} : G \text{ is real and } c^*G = (-1)^dG\}.$$

A codimension  $d$  arithmetic cycle on  $X$  is a pair  $(Z, G)$  in which  $Z$  is a codimension  $d$  cycle in the usual sense, and

$$G \in \tilde{D}_X^{d-1,d-1} \stackrel{\text{def}}{=} \frac{D_X^{d-1,d-1}}{\text{Im}(\partial) + \text{Im}(\bar{\partial})},$$

satisfies the Green equation  $dd^c G + \delta_{Z(\mathbb{C})} = [\omega]$  for some  $\omega \in E_X^{d,d}$ . Denote by  $\widehat{Z}^d(X)$  the abelian group of all such pairs. The arithmetic Chow group is the quotient

$$\widehat{\text{CH}}^d(X) = \widehat{Z}^d(X)/(\text{rational equivalence}).$$

Now assume that  $X$  is projective, and that  $X_0 \subset X$  is a smooth closed subvariety. Let  $(Z, G)$  be any codimension  $d$  arithmetic cycle, and set  $U = X \setminus Z$ . Recalling Definition 2.1.5, define a real vector space

$$E_{\log}^{d,d}(U) = \{g \in E_{\log}^{d,d}(U(\mathbb{C})) : g \text{ is real and } F_\infty^*g = (-1)^d g\}.$$

There is a canonical map

$$E_{\log}^{d-1,d-1}(U) \xrightarrow{g \mapsto [g]} D_X^{d-1,d-1},$$

and Theorem 4.4 of [Bur94] implies the existence of a unique lift of  $G$  to

$$g \in \tilde{E}_{\log}^{d-1,d-1}(U) \stackrel{\text{def}}{=} \frac{E_{\log}^{d-1,d-1}(U)}{\text{Im}(\partial) + \text{Im}(\bar{\partial})}.$$

Theorem 2.2.5 therefore implies that the current  $G = [g]$  admits a logarithmic expansion along  $X_0$ . Combining this with Remark 2.2.9 and Proposition 2.2.10, we obtain an arithmetic cycle

$$(\sigma_{X_0/X}(Z), \sigma_{X_0/X}(G)) \in \widehat{Z}^d(N_{X_0/X}).$$

This defines a homomorphism

$$(2.3.1) \quad \widehat{Z}^d(X) \rightarrow \widehat{Z}^d(N_{X_0/X}).$$

The following is slightly weaker than what is proved in §4.1 of [Hu99]; see Remark 2.3.2 below.

**Theorem 2.3.1** (Hu). *Still assuming that  $X$  is projective, the homomorphism (2.3.1) descends to*

$$\widehat{\text{CH}}^d(X) \rightarrow \widehat{\text{CH}}^d(N_{X_0/X}),$$

and this map agrees with the composition

$$\widehat{\text{CH}}^d(X) \xrightarrow{i_0^*} \widehat{\text{CH}}^d(X_0) \xrightarrow{\pi_0^*} \widehat{\text{CH}}^d(N_{X_0/X}).$$

Here  $i_0 : X_0 \rightarrow X$  is the inclusion,  $\pi_0 : N_{X_0/X} \rightarrow X_0$  is the bundle map, and  $i_0^*$  and  $\pi_0^*$  are the induced pullbacks on arithmetic Chow groups.

*Remark 2.3.2.* Assuming only that  $X$  is quasi-projective, there are canonical maps

$$\widehat{Z}^d(X, \mathcal{D}_{\log}) \rightarrow \widehat{Z}^d(X) \quad \text{and} \quad \widehat{\text{CH}}^d(X, \mathcal{D}_{\log}) \rightarrow \widehat{\text{CH}}^d(X),$$

where the domains are the  $\mathcal{D}_{\log}$  arithmetic cycles and Chow groups of [BGKK07]. These agree with those of [Bur97], and both maps are isomorphisms if  $X$  is projective. Hu proves the existence of a distinguished lift of (2.3.1) to

$$(2.3.2) \quad \widehat{Z}^d(X, \mathcal{D}_{\log}) \rightarrow \widehat{Z}^d(N_{X_0/X}, \mathcal{D}_{\log}),$$

which then descends to a map on  $\mathcal{D}_{\log}$  arithmetic groups. This descent agrees with the composition

$$\widehat{\text{CH}}^d(X, \mathcal{D}_{\log}) \xrightarrow{i_0^*} \widehat{\text{CH}}^d(X_0, \mathcal{D}_{\log}) \xrightarrow{\pi_0^*} \widehat{\text{CH}}^d(N_{X_0/X}, \mathcal{D}_{\log}).$$

Even when  $X$  is projective, this is stronger than Theorem 2.3.1 (because  $N_{X_0/X}$  is not projective). The construction of the lift (2.3.2) is subtle, but the key ingredient is the lift of  $\sigma_{X_0/X}(g)$  mentioned at the end of Remark 2.2.11.

**Proposition 2.3.3.** *The pullback  $\pi_0^*$  in Theorem 2.3.1 is injective.*

*Proof.* A similar statement is found in [Bur97], but for the  $\mathcal{D}_{\log}$  arithmetic Chow groups of Remark 2.3.2. The proof for Gillet-Soulé arithmetic Chow

groups follows the same idea. By Theorem 3.3.5 of [GS90] there is commutative diagram with exact rows

$$\begin{array}{ccccccc}
\mathrm{CH}^{d,d-1}(X_0) & \longrightarrow & \tilde{E}_{X_0}^{d-1,d-1} & \longrightarrow & \widehat{\mathrm{CH}}^d(X_0) & \longrightarrow & \mathrm{CH}^d(X_0) \\
\downarrow & & \downarrow & & \downarrow \pi_0^* & & \downarrow \\
\mathrm{CH}^{d,d-1}(N_{X_0/X}) & \longrightarrow & \tilde{E}_{N_{X_0/X}}^{d-1,d-1} & \longrightarrow & \widehat{\mathrm{CH}}^d(N_{X_0/X}) & \longrightarrow & \mathrm{CH}^d(N_{X_0/X}).
\end{array}$$

The first and last vertical arrows are isomorphisms by Theorem 8.3 of [Gil81]. The second vertical arrow is injective, and hence the third is as well.  $\square$

### 3. SPECIAL GREEN CURRENTS

Given a closed immersion of complex manifolds  $X_0 \subset X$ , the constructions of Garcia-Sankaran [GS19], Bismut [Bis90], and Bismut-Gillet-Soulé [BGS90], provide a systematic way to produce Green currents for certain cycles on  $X$ . Theorem 2.2.5 can be applied to these currents to prove the existence of logarithmic expansions, but this abstract existence theorem is not sharp enough for our purposes.

The goal of this section is to construct explicit logarithmic expansions for these currents, and so effectively compute their specializations to  $N_{X_0/X}$ .

**3.1. Construction of Green forms.** Let  $X$  be a complex manifold, and let  $L$  be a holomorphic line bundle on  $X$ . We use the same symbol for both the total space  $L \rightarrow X$ , viewed as a complex manifold fibered over  $X$ , and for its sheaf of holomorphic sections.

Let  $h(-, -)$  be a hermitian metric on  $L$ . If  $s$  is any local holomorphic section of  $L$ , abbreviate  $h(s) = h(s, s)$ . The *Chern form* of  $L$  is the  $(1, 1)$  form defined locally by

$$\mathrm{ch}(L) = \frac{1}{2\pi i} \partial \bar{\partial} \log h(s).$$

We denote again by  $h$  the induced metric on the dual bundle  $L^\vee$ .

Fix an integer  $1 \leq d \leq \dim(X)$  and a tuple  $s = (s_1, \dots, s_d)$  with  $s_i \in H^0(X, L^\vee)$ , and abbreviate

$$h(s) = h(s_1) + \dots + h(s_d).$$

Denote by  $Z(s) \subset X$  the (possibly nonreduced) analytic subspace defined by  $s_1 = \dots = s_d = 0$ .

**Definition 3.1.1.** Fix a point  $x \in Z(s)$ , trivialize  $L$  in a neighborhood of  $x$ , and use this to view  $s_{1,x}, \dots, s_{d,x} \in \mathcal{O}_{X,x}$  as germs of holomorphic functions at  $x$ . We say that  $s$  is:

- *regular at  $x$*  if  $s_{1,x}, \dots, s_{d,x} \in \mathcal{O}_{X,x}$  is a regular sequence in the sense of commutative algebra;
- *smooth at  $x$*  if  $s_{1,x}, \dots, s_{d,x}$  are linearly independent in  $\mathfrak{m}_{X,x}/\mathfrak{m}_{X,x}^2$ , where  $\mathfrak{m}_{X,x} \subset \mathcal{O}_{X,x}$  is the maximal ideal.

The tuple  $s = (s_1, \dots, s_d)$  is *regular* or *smooth* if it has this property at every point of  $Z(s)$ .

*Remark 3.1.2.* Regularity of  $s$  at  $x$  is equivalent to all irreducible components of  $Z(s)$  passing through  $x$  having codimension  $d$  in  $X$ , as both are equivalent to  $\mathcal{O}_{Z(s),x}$  being Cohen-Macaulay of dimension  $\dim(X) - d$ .

*Remark 3.1.3.* Smoothness of  $s$  at  $x$  is equivalent to  $Z(s)$  being nonsingular (that is, a complex manifold) of codimension  $d$  in some open neighborhood of  $x$ , as both are equivalent to  $\mathcal{O}_{Z(s),x}$  being regular of dimension  $\dim(X) - d$ .

*Remark 3.1.4.* If  $s = (s_1, \dots, s_d)$  is regular or smooth at a point  $x$ , the same is true of all tuples obtained by reordering the components of  $s$ , and of all tuples  $(s_1, \dots, s_r)$  with  $1 \leq r \leq d$ .

Regularity of  $s$  is equivalent<sup>1</sup> to the corresponding morphism of vector bundles  $s : L^{\oplus d} \rightarrow \mathcal{O}_X$  being regular in the sense of §2.1.1 of [GS19]. Therefore, if  $s$  is regular, the constructions (2.5) and (2.12) of *loc. cit.* define forms

$$\varphi^\circ(s) \in E_X^\bullet \quad \text{and} \quad \nu^\circ(s) \in E_X^\bullet.$$

Both have trivial components in odd degree, and their components in even degree  $2p$  have type  $(p, p)$ . Abbreviate

$$\omega^\circ(s) = (-2\pi i)^{-d} \cdot \varphi^\circ(s)_{[2d]} \in E_X^{d,d}.$$

We will not recall the detailed construction of the forms above, as we only need the degree  $2d$  component of  $\varphi^\circ(s)$  and the degree  $2d - 2$  component of  $\nu^\circ(s)$ . Explicit formulas for these can be found in [GS19] and [Gar18]. If  $d = 1$  then

$$(3.1.1) \quad \varphi^\circ(s)_{[2]} = 2\pi i e^{-2\pi h(s)} \left( \text{ch}(L) - i \frac{\partial h(s) \wedge \bar{\partial} h(s)}{h(s)} \right)$$

and

$$(3.1.2) \quad \nu^\circ(s)_{[0]} = e^{-2\pi h(s)}.$$

If  $d > 1$  then

$$(3.1.3) \quad \varphi^\circ(s_1, \dots, s_d)_{[2d]} = \varphi^\circ(s_1)_{[2]} \wedge \cdots \wedge \varphi^\circ(s_d)_{[2]}$$

and

$$(3.1.4) \quad \nu^\circ(s_1, \dots, s_d)_{[2d-2]} = \sum_{j=1}^d \nu^\circ(s_j)_{[0]} \wedge \varphi^\circ(s_1, \dots, \widehat{s}_j, \dots, s_d)_{[2d-2]}.$$

Strictly speaking, the above formulas are given in [GS19] and [Gar18] only for specific hermitian line bundles on hermitian symmetric domains associated

---

<sup>1</sup>This equivalence, and the claims of Remarks 3.1.2, 3.1.3, and 3.1.4, all follow from basic properties of regular sequences and complex analytic spaces, as found in [Mat89] and [Fis76].

to orthogonal and unitary groups, but the derivations of these formulas hold verbatim in our more general setting.

As explained in [GS19], results of Bismut [Bis90] and Bismut-Gillet-Soulé [BGS90] can be used to produce Green currents for the cycles  $Z(s) \subset X$  defined above. We need a slight strengthening of those results.

**Proposition 3.1.5.** *If  $s$  is regular, the integral*

$$(3.1.5) \quad \mathfrak{g}^\circ(s) = \left(\frac{-1}{2\pi i}\right)^{d-1} \int_1^\infty \nu^\circ(\sqrt{u} \cdot s)_{[2d-2]} \frac{du}{u}$$

defines a smooth form on  $X \setminus Z(s)$  with

$$(3.1.6) \quad \mathfrak{g}^\circ(s) = \frac{a(s)}{h(s)^{d-1}} + b(s) \cdot \log(h(s))$$

for some  $a(s), b(s) \in E_X^{d-1, d-1}$ . If  $s$  is smooth, then

$$(3.1.7) \quad \mathfrak{g}^\circ(s) \in E_X^{d-1, d-1}(\log Z(s))$$

with respect to the resolution of singularities of  $(X, Z(s))$  obtained by blowing up along  $Z(s) \subset X$ , and the associated current (Remark 2.1.4) satisfies the Green equation

$$dd^c[\mathfrak{g}^\circ(s)] + \delta_{Z(s)} = [\omega^\circ(s)].$$

*Proof.* First assume  $d = 1$ , so that  $s$  is a nonzero section of  $L^\vee$ . Plugging (3.1.2) into (3.1.5) yields

$$(3.1.8) \quad \mathfrak{g}^\circ(s) = \int_1^\infty e^{-2\pi u h(s)} \frac{du}{u} = E_1(2\pi h(s)),$$

where

$$(3.1.9) \quad E_1(x) = \int_1^\infty e^{-xu} \frac{du}{u} = -\log(x) - \gamma - \sum_{k=1}^\infty \frac{(-x)^k}{k \cdot k!}.$$

This gives a more precise version of (3.1.6), which will be essential later.

Now suppose  $d > 1$ . For each  $1 \leq j \leq d$  abbreviate

$$\eta(s_j) = -i \cdot \frac{\partial h(s_j) \wedge \bar{\partial} h(s_j)}{h(s_j)} \in E_X^{1,1},$$

so that (3.1.1) becomes

$$\varphi^\circ(s_j)_{[2]} = 2\pi i e^{-2\pi h(s_j)} (\text{ch}(L) + \eta(s_j)),$$

and (3.1.3) and (3.1.4) imply

$$\nu^\circ(s)_{[2d-2]} = (2\pi i)^{d-1} e^{-2\pi h(s)} \sum_{j=1}^d \underbrace{(\text{ch}(L) + \eta(s_1)) \wedge \cdots \wedge (\text{ch}(L) + \eta(s_d))}_{\text{omit } j^{\text{th}} \text{ factor}}.$$



Expanding out the wedge products in each term, we rewrite this as

$$(3.1.10) \quad \nu^\circ(s)_{[2d-2]} = e^{-2\pi h(s)} \sum_{k=0}^{d-1} \eta_k(s),$$

in which each  $\eta_k(s) \in E_X^{d-1, d-1}$  is (up to multiplication by a constant) the wedge product of  $\text{ch}(L)^{d-k-1}$  with a sum of  $k$ -fold wedges of  $\eta(s_1), \dots, \eta(s_d)$ . For any  $t \in \mathbb{C}$  we have  $\eta(ts_j) = |t|^2 \eta(s_j)$ , and hence

$$(3.1.11) \quad \eta_k(ts) = |t|^{2k} \eta_k(s).$$

Plugging (3.1.10) into (3.1.5) results in

$$(-2\pi i)^{d-1} \mathfrak{g}^\circ(s) = \sum_{k=0}^{d-1} \eta_k(s) \int_1^\infty u^k e^{-2\pi u h(s)} \frac{du}{u}.$$

If  $k > 0$ , a calculus exercise shows that

$$(3.1.12) \quad \int_1^\infty u^k e^{-ux} \frac{du}{u} = \frac{e^{-x} \cdot (k-1)!}{x^k} \cdot \sum_{i=0}^{k-1} \frac{x^i}{i!},$$

and so there are polynomials  $P_k(x)$  satisfying

$$(-2\pi i)^{d-1} \mathfrak{g}^\circ(s) = \eta_0(s) \cdot E_1(2\pi h(s)) + e^{-2\pi h(s)} \sum_{k=1}^{d-1} \frac{\eta_k(s)}{h(s)^k} \cdot P_k(h(s)).$$

The equality (3.1.6) follows immediately.

Assuming now that  $s$  is smooth, (3.1.7) follows from these explicit formulas by a local calculation on the blowup of  $Z(s) \subset X$ . For the Green equation, see Proposition 2.2 of [GS19].  $\square$

**3.2. The star product formula.** Suppose  $G_1$  and  $G_2$  are currents on  $X$  satisfying the Green equations

$$dd^c G_i + \delta_{Z_i} = [\omega_i]$$

for analytic cycles  $Z_1$  and  $Z_2$  of codimensions  $d_1$  and  $d_2$  intersecting properly. If  $G_2 = [g_2]$  for some locally integrable form  $g_2$  on  $X$ , we define

$$G_1 \star G_2 = \delta_{Z_1} \wedge G_2 + G_1 \wedge \omega_2 \in D_X^{d_1+d_2-1, d_1+d_2-1},$$

provided that the integral

$$(\delta_{Z_1} \wedge G_2)(\phi) = \int_{Z_1} g_2 \wedge \phi$$

converges for all  $\phi \in {}_c E_X^\bullet$  of the appropriate degree.

*Remark 3.2.1.* Note that we understand the star product to be a current on  $X$ , not an element of the space of currents modulo currents of the form  $\partial a + \bar{\partial} b$ . Because of this, the star product is neither commutative nor associative, and in fact it may be that  $G_1 \star G_2$  is defined while  $G_2 \star G_1$  is not.

*Remark 3.2.2.* Keeping the previous remark in mind, we caution the reader that we are using the convention for star products opposite to [Sou92] and [GS19]: our  $G_1 \star G_2$  is their  $G_2 \star G_1$ .

*Remark 3.2.3.* The expression  $G_1 \star (G_2 \star G_3)$  does not make sense, as  $G_2 \star G_3$  is not represented by a locally integrable form (even if  $G_2$  and  $G_3$  are). We therefore understand

$$\begin{aligned} G_1 \star G_2 \star G_3 &= (G_1 \star G_2) \star G_3 \\ G_1 \star G_2 \star G_3 \star G_4 &= ((G_1 \star G_2) \star G_3) \star G_4 \\ &\vdots \end{aligned}$$

provided that each star product on the right is defined.

Fix a smooth tuple  $s = (s_1, \dots, s_d)$  with  $s_i \in H^0(X, L^\vee)$ . If we write  $d = k + \ell$  with  $k, \ell > 0$ , and express  $s = (p, q)$  as the concatenation of the smooth tuples

$$p = (s_1, \dots, s_k) \quad \text{and} \quad q = (s_{k+1}, \dots, s_d),$$

then  $Z(s) = Z(p) \times_X Z(q)$  as analytic spaces.

**Lemma 3.2.4.** *If  $G(p) \in D_X^{k-1, k-1}$  is any Green current for  $Z(p)$ , the star product  $G(p) \star \mathfrak{g}^\circ(q)$  is defined.*

*Proof.* The pullback of  $\mathfrak{g}^\circ(q)$  to  $Z(p)$  is the form  $\mathfrak{g}^\circ(q|_{Z(p)})$  obtained by applying the construction of Proposition 3.1.5 to the smooth  $\ell$ -tuple

$$q|_{Z(p)} = (s_{k+1}|_{Z(p)}, \dots, s_d|_{Z(p)})$$

of sections of  $L^\vee|_{Z(p)}$  on the complex manifold  $Z(p)$ . In particular, this pullback is locally integrable.  $\square$

**Theorem 3.2.5** (Garcia-Sankaran). *We have the equality of currents*

$$\mathfrak{g}^\circ(s) = \mathfrak{g}^\circ(p) \star \mathfrak{g}^\circ(q) - \partial[A(p; q)] - \bar{\partial}[B(p; q)]$$

on  $X$  for locally integrable forms

$$\begin{aligned} A(p; q) &= \left( \frac{-1}{2\pi i} \right)^{d-1} \int_{1 < v < u < \infty} \bar{\partial}(\nu^\circ(\sqrt{u}p)_{[2k-2]}) \wedge \nu^\circ(\sqrt{v}q)_{[2\ell-2]} \frac{du}{u} \frac{dv}{v} \\ B(p; q) &= \left( \frac{-1}{2\pi i} \right)^{d-1} \int_{1 < v < u < \infty} \nu^\circ(\sqrt{u}p)_{[2k-2]} \wedge \partial(\nu^\circ(\sqrt{v}q)_{[2\ell-2]}) \frac{du}{u} \frac{dv}{v}. \end{aligned}$$

Moreover, for an appropriate choice of resolution of singularities, these forms lie in  $E_X^\bullet(\log Z(p) \cup Z(q))$ .

*Proof.* Except for the final claim, this is Theorem 2.16 of [GS19], modified as per Remark 3.2.2. For those authors  $X$  is a particular hermitian symmetric domain, but the same argument works in our more general context.

We sketch the proof of the final claim. Abbreviate  $\hbar = 2\pi\hbar$ . As in (3.1.10), we expand

$$\nu^\circ(s)_{[2d-2]} = e^{-\hbar(s)} \sum_{r=0}^{d-1} \eta_r(s)$$

Plugging this expansion into the definitions of  $A(p; q)$  and  $B(p; q)$ , and noting that  $\eta_0(p)$  and  $\eta_0(q)$  are closed, we find that

$$\begin{aligned} (-2\pi i)^{d-1} A(p; q) &= \sum_{\substack{0 < a < k \\ 0 \leq b < \ell}} F_{a,b}(\hbar(p), \hbar(q)) \cdot \bar{\partial} \eta_a(p) \wedge \eta_b(q) \\ (3.2.1) \quad &- \sum_{\substack{0 \leq a < k \\ 0 \leq b < \ell}} F_{a+1,b}(\hbar(p), \hbar(q)) \cdot \bar{\partial} \hbar(p) \wedge \eta_a(p) \wedge \eta_b(q) \end{aligned}$$

$$\begin{aligned} (-2\pi i)^{d-1} B(p; q) &= \sum_{\substack{0 \leq a < k \\ 0 < b < \ell}} F_{a,b}(\hbar(p), \hbar(q)) \cdot \eta_a(p) \wedge \partial \eta_b(q) \\ (3.2.2) \quad &- \sum_{\substack{0 \leq a < k \\ 0 \leq b < \ell}} F_{a,b+1}(\hbar(p), \hbar(q)) \cdot \eta_a(p) \wedge \partial \hbar(q) \wedge \eta_b(q), \end{aligned}$$

in which we have set

$$\begin{aligned} F_{a,b}(x, y) &= \int_{1 < v < u < \infty} u^a v^b e^{-ux} e^{-vy} \frac{du}{u} \frac{dv}{v} \\ &= \int_1^\infty v^{a+b} e^{-vy} \left( \int_1^\infty u^a e^{-uvx} \frac{du}{u} \right) \frac{dv}{v}. \end{aligned}$$

If  $a, b > 0$ , then (3.1.12) applies to the inner integral, leaving

$$(3.2.3) \quad F_{a,b}(x, y) = \sum_{i=0}^{a-1} \frac{(a-1)!}{x^{a-i} \cdot i!} \int_1^\infty v^{b+i} e^{-v(x+y)} \frac{dv}{v}.$$

Applying (3.1.12) once again leaves

$$F_{a,b}(x, y) = e^{-x-y} \sum_{i=0}^{a-1} \frac{\text{poly}(x, y)}{x^{a-i} \cdot (x+y)^{b+i}},$$

where in each term  $\text{poly}(x, y)$  is some polynomial (depending on  $i$ ) in  $x$  and  $y$  whose exact value is irrelevant to us. If  $a > 0$  and  $b = 0$  one argues in the same way, except that the integral appearing in the  $i = 0$  term of (3.2.3) is  $E_1(x+y)$ . Thus

$$F_{a,0}(x, y) = \frac{E_1(x+y) \cdot (a-1)!}{x^a} + e^{-x-y} \sum_{i=1}^{a-1} \frac{\text{poly}(x, y)}{x^{a-i} \cdot (x+y)^i}.$$

If  $a = 0$  and  $b > 0$  then, again using (3.1.12), rewrite  $F_{0,b}(x, y)$  as

$$\int_1^\infty \left( \int_1^\infty v^b e^{-v(y+ux)} \frac{dv}{v} \right) \frac{du}{u} = \sum_{i=0}^{b-1} \frac{(b-1)!}{i!} \int_1^\infty \frac{e^{-(y+ux)}}{(y+ux)^{b-i}} \frac{du}{u}.$$

The integral on the right can again be evaluated using elementary methods, and one finds that  $F_{0,b}(x, y)$  has the form

$$e^{-y} E_1(x) \cdot \frac{\text{poly}(y)}{y^b} + E_1(x+y) \cdot \frac{\text{poly}(y)}{y^b} + \sum_{j=1}^{b-1} \frac{e^{-(x+y)} \cdot \text{poly}(x, y)}{y^{b-j}(x+y)^j}.$$

Substituting these formulas for  $F_{a,b}(x, y)$  into (3.2.1) and (3.2.2), one finds long but explicit formulas making clear the singularities of  $A(p; q)$  and  $B(p; q)$ . From these, a term-by-term calculation shows that both are locally integrable. For the final claim of the proposition, one constructs resolutions of singularities

$$(X', D') \rightarrow (Y', E') \xrightarrow{r'} (X, Z(p))$$

and

$$(X'', D'') \rightarrow (Y'', E'') \xrightarrow{r''} (X, Z(q))$$

by first taking  $Y'$  and  $Y''$  to be the blowups of  $X$  along  $Z(p)$  and  $Z(q)$ , respectively. Then let  $X'$  and  $X''$  be the blowups of  $Y'$  and  $Y''$  along the preimages of  $Z(s) = Z(p) \cap Z(q)$  under  $r'$  and  $r''$ . Now take any resolution of singularities of the analytic subspace  $D' \times_X D'' \subset X' \times_X X''$ . The result is a resolution of singularities of  $(X, Z(p) \cup Z(q))$ , and by explicit calculations in local coordinates one can show that  $A(p; q)$  and  $B(p; q)$  lie in  $E_X^\bullet(\log Z(p) \cup Z(q))$ . The tedious details are left to the reader.  $\square$

As a special case of Theorem 3.2.5,

$$\begin{aligned} \mathfrak{g}^\circ(s_1, \dots, s_d) &= \mathfrak{g}^\circ(s_1, \dots, s_{d-1}) \star \mathfrak{g}^\circ(s_d) \\ &\quad - \partial[A(s_1, \dots, s_{d-1}; s_d)] - \bar{\partial}[B(s_1, \dots, s_{d-1}; s_d)]. \end{aligned}$$

Repeated application of this results in

$$(3.2.4) \quad \mathfrak{g}^\circ(s) = \mathfrak{g}^\circ(s_1) \star \dots \star \mathfrak{g}^\circ(s_d) - \partial[\mathfrak{a}(s)] - \bar{\partial}[\mathfrak{b}(s)]$$

for locally integrable forms

$$\begin{aligned} \mathfrak{a}(s) &= \sum_{r=2}^d A(s_1, \dots, s_{r-1}; s_r) \wedge \omega^\circ(s_{r+1}) \wedge \dots \wedge \omega^\circ(s_d) \\ \mathfrak{b}(s) &= \sum_{r=2}^d B(s_1, \dots, s_{r-1}; s_r) \wedge \omega^\circ(s_{r+1}) \wedge \dots \wedge \omega^\circ(s_d). \end{aligned}$$

**3.3. Explicit logarithmic expansions.** We now return to the setting of §2.2, so that  $X_0 \subset X$  is a closed complex submanifold, but now assume that  $X_0$  is presented to us in a particular way: there is a holomorphic vector bundle  $N \rightarrow X$  of dimension  $\dim(X) - \dim(X_0)$  and a section

$$u \in H^0(X, N)$$

such that  $X_0 \subset X$  is defined (as an analytic space) by the equation  $u = 0$ .

This presentation of  $X_0 \subset X$  identifies

$$(3.3.1) \quad N_{X_0/X} \cong N|_{X_0}.$$

Indeed, if we denote by  $\mathcal{I} \subset \mathcal{O}_X$  the ideal sheaf of holomorphic functions vanishing along  $X_0$ , then evaluation at  $u$  defines an isomorphism  $N^\vee \cong \mathcal{I}$ . Restricting this to  $X_0$  yields an isomorphism  $N|_{X_0}^\vee \cong \mathcal{I}/\mathcal{I}^2$  of vector bundles on  $X_0$ , and the normal bundle to  $X_0 \subset X$  is (by definition) the dual of the right hand side.

Viewing points of the total space  $N \rightarrow X$  as pairs  $(x, v_x)$  consisting of a point  $x \in X$  and a vector  $v_x \in N_x$  in the fiber at  $x$ , the deformation to the normal bundle  $X_0 \subset X$  can be identified with the subset

$$\tilde{X} \subset N \times \mathbb{C}$$

of triples  $(x, v_x, t)$  consisting of a point  $(x, v_x) \in N$ , and a scalar  $t \in \mathbb{C}$  satisfying  $t \cdot v_x = u_x$ . The morphisms

$$X \xleftarrow{\pi} \tilde{X} \xrightarrow{\tau} \mathbb{C}$$

of (2.2.1) are given by  $\pi(x, v_x, t) = x$  and  $\tau(x, v_x, t) = t$ . This is essentially McPherson's description of the deformation to the normal bundle, as in Remark 5.1.1 of [Ful84].

As in §3.1, fix a line bundle  $L \rightarrow X$  with a hermitian metric  $h$ . Any morphism of holomorphic vector bundles  $y : N \rightarrow L^\vee$ , determines a section

$$(3.3.2) \quad q = y(u) \in H^0(X, L^\vee)$$

vanishing along  $X_0$ . We call this the *degenerating section* determined by  $y$ . Like any vector bundle,  $\pi_0 : N_{X_0/X} \rightarrow X_0$  acquires a tautological section

$$v_0 \in H^0(N_{X_0/X}, \pi_0^* N_{X_0/X})$$

after pullback via its own bundle map. Setting  $L_0 = L|_{X_0}$ , we may restrict  $y : N \rightarrow L^\vee$  to a morphism

$$N_{X_0/X} = N|_{X_0} \rightarrow L^\vee|_{X_0} = L_0^\vee$$

and pull back by  $\pi_0$ . This results in a morphism

$$\pi_0^* y : \pi_0^* N_{X_0/X} \rightarrow \pi_0^* L_0^\vee,$$

and hence a section

$$(3.3.3) \quad \sigma_{X_0/X}(q) = (\pi^* y)(v_0) \in H^0(N_{X_0/X}, \pi_0^* L_0^\vee).$$

This is the *specialization to the normal bundle* of  $q$ . The connection between a degenerating section and its specialization is clarified by the following lemma.

**Lemma 3.3.1.** *For any  $q = y(u)$  as above, there is a unique section*

$$\tilde{q} \in H^0(\tilde{X}, \pi^* L^\vee)$$

*satisfying  $\tau \cdot \tilde{q} = \pi^* q$ . The pullback of  $\tilde{q}$  to  $N_{X_0/X} = \tilde{X}_0$  is  $\sigma_{X_0/X}(q)$ .*

*Proof.* There is a tautological section  $v \in H^0(\tilde{X}, \pi^* N)$  whose fiber at a point  $(x, v_x, t) \in \tilde{X}$  is  $v_x$ . This section satisfies  $\tau \cdot v = \pi^* u$ , and its restriction to  $N_{X_0/X}$  is  $v_0$ . Denoting by  $\pi^* y : \pi^* N \rightarrow \pi^* L^\vee$  the pullback of  $y$ , the section

$$\tilde{q} = (\pi^* y)(v) \in H^0(\tilde{X}, \pi^* L^\vee)$$

has the stated properties.  $\square$

Now fix a smooth (Definition 3.1.1) tuple  $s = (s_1, \dots, s_d)$  with  $s_i \in H^0(X, L^\vee)$ , and assume  $s = (p, q)$  is the concatenation of

$$p = (p_1, \dots, p_k) \quad \text{and} \quad q = (q_1, \dots, q_\ell),$$

satisfying the following properties:

- (1) The tuple  $p|_{X_0}$  formed from the restrictions

$$p_1|_{X_0}, \dots, p_k|_{X_0} \in H^0(X_0, L_0^\vee)$$

is again smooth; equivalently, the analytic subspace

$$Z(p|_{X_0}) = Z(p) \times_X X_0 \subset X_0$$

is smooth of codimension  $k$ .

- (2) The sections  $q_1, \dots, q_\ell \in H^0(X, L^\vee)$  are the degenerating sections determined by morphisms  $N \rightarrow L^\vee$  as above. Denote by  $Z(\tilde{q}) \subset \tilde{X}$  and

$$Z(\sigma_{X_0/X}(q)) \subset N_{X_0/X}$$

the analytic subspaces defined by the vanishing of the components of  $\tilde{q} = (\tilde{q}_1, \dots, \tilde{q}_d)$  and

$$\sigma_{X_0/X}(q) = (\sigma_{X_0/X}(q_1), \dots, \sigma_{X_0/X}(q_d)).$$

In particular,  $Z(p)$  intersects  $X_0$  transversely, while  $X_0 \subset Z(q)$ . We allow the possibility that  $s = p$  or  $s = q$ . Note that the tuples  $p$  and  $q$  are again smooth, by Remark 3.1.4. We consider the specializations of  $Z(s)$ ,  $Z(p)$ , and  $Z(q)$  to  $N_{X_0/X}$ .

**Proposition 3.3.2.** *We have the equality*

$$\sigma_{X_0/X}(Z(s)) = \sigma_{X_0/X}(Z(p)) \cdot \sigma_{X_0/X}(Z(q)),$$

*where the intersection is the proper intersection of [Dra69], and*

$$\sigma_{X_0/X}(Z(p)) = \pi_0^* Z(p|_{X_0}).$$

Moreover, the tuples  $\tilde{q}$  and  $\sigma_{X_0/X}(q)$  are smooth, satisfy

$$\sigma_{X_0/X}(Z(q)) = Z(\sigma_{X_0/X}(q)) = Z(\sigma_{X_0/X}(q_1)) \cdots Z(\sigma_{X_0/X}(q_\ell)),$$

and all three of these cycles are equal to  $Z(\tilde{q}) \cdot N_{X_0/X}$ .

*Proof.* Let  $\Delta \subset \mathbb{C}$  be the open unit disk, and abbreviate  $n = \dim(X)$  and  $m = \dim(X) - \dim(X_0)$ . The smoothness of  $s = (p, q)$  implies that we may find a coordinate neighborhood in  $X$  near a point of  $X_0$  of the form

$$U \cong \Delta^n = \{(z_1, \dots, z_n) : z_i \in \Delta\}$$

in such a way that

- the line bundle  $L$  is trivial on  $U$ ,
- $U_0 = X_0 \cap U$  is defined by the vanishing of  $z_1, \dots, z_m$ ,
- $p_1 = z_{m+1}, \dots, p_k = z_{m+k}$ ,
- $q_1 = z_1, \dots, q_\ell = z_\ell$ .

The deformation to the normal bundle of  $U_0 \subset U$  is identified with

$$\tilde{U} = \{(z_1, \dots, z_n, w_1, \dots, w_m, t) \in \Delta^n \times \mathbb{C}^m \times \mathbb{C} : z_i = tw_i, \forall 1 \leq i \leq m\},$$

and  $\tilde{q}_i = w_i$  for all  $1 \leq i \leq \ell$ . The normal bundle itself is identified with

$$N_{U_0/U} = \{(0, \dots, 0, z_{m+1}, \dots, z_n, w_1, \dots, w_m, 0) \in \Delta^n \times \mathbb{C}^m \times \mathbb{C}\},$$

and  $\sigma_{X_0/X}(q_i) = w_i$  for all  $1 \leq i \leq \ell$ . The strict transforms of  $Z(p)$  and  $Z(q)$  are defined by (respectively) the vanishing of  $z_{m+1}, \dots, z_{m+k}$  and the vanishing of  $w_1, \dots, w_\ell$ . Their specializations to  $N_{U_0/U}$  are defined by the same equations. All parts of the proposition follow immediately from computations in these local coordinates.  $\square$

Now we turn to the Green current

$$\mathfrak{g}^\circ(s) \in E_X^{d-1, d-1}(\log Z(s))$$

of Proposition 3.1.5, and the similar currents  $\mathfrak{g}^\circ(p)$  and  $\mathfrak{g}^\circ(q)$ . The following lemmas are the key to understanding their logarithmic expansions along  $X_0$ , and hence their specializations to  $N_{X_0/X}$ .

**Lemma 3.3.3.** *There are forms  $a, b, c \in E_{\tilde{X}}^{\ell-1, \ell-1}$  such that*

$$(3.3.4) \quad \mathfrak{g}^\circ(q) = j_t^* \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) + c \cdot \log |\tau| \right)$$

for all  $t \in \mathbb{C} \setminus \{0\}$ . If we define currents

$$G_0(t) = \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) \right) \wedge \delta_{\tilde{X}_t}$$

and  $G_1(t) = c \wedge \delta_{\tilde{X}_t}$  on  $\tilde{X}$ , then

$$j_{t*}[\mathfrak{g}^\circ(q)] = G_0(t) + G_1(t) \log |t|$$

is a logarithmic expansion of  $\mathfrak{g}^\circ(q)$  along  $X_0$ .

*Proof.* The smoothness of  $\tilde{q}$  allows us to apply the constructions of §3.1 to obtain Green forms  $\mathfrak{g}^\circ(\tilde{q})$  and  $\mathfrak{g}^\circ(\tau\tilde{q})$  for the cycles  $Z(\tilde{q}) \subset \tilde{X}$  and

$$Z(\tilde{q}) \setminus N_{X_0/X} \subset \tilde{X} \setminus N_{X_0/X},$$

respectively. Recalling that  $\tau\tilde{q} = \pi^*q$ , for  $t \neq 0$  these are related by

$$\mathfrak{g}^\circ(q) = j_t^* \pi^* \mathfrak{g}^\circ(q) = j_t^* \mathfrak{g}^\circ(\tau\tilde{q}).$$

As in the proof Proposition 3.1.5, we may write

$$(3.3.5) \quad (-2\pi i)^{\ell-1} \mathfrak{g}^\circ(\tilde{q}) = \eta_0(\tilde{q}) \cdot E_1(2\pi h(\tilde{q})) + e^{-2\pi h(\tilde{q})} \sum_{j=1}^{\ell-1} \frac{\eta_j(\tilde{q})}{h(\tilde{q})^j} \cdot P_j(h(\tilde{q})),$$

where  $P_j$  is a polynomial and  $\eta_j(\tilde{q})$  is a smooth form on  $\tilde{X}$  satisfying the homogeneity property (3.1.11). If we replace  $\tilde{q}$  by  $\tau\tilde{q}$  in (3.3.5), pull back by  $j_t : X \rightarrow \tilde{X}$ , and use

$$j_t^* \left( \frac{\eta_k(\tau\tilde{q})}{h(\tau\tilde{q})^j} \right) = j_t^* \left( \frac{\eta_k(\tilde{q})}{h(\tilde{q})^j} \right),$$

we find that  $\mathfrak{g}^\circ(q) = j_t^* \mathfrak{g}^\circ(\tau\tilde{q}) = (-2\pi i)^{1-\ell} j_t^* \Psi$  where

$$\Psi = \eta_0(\tilde{q}) \cdot E_1(2\pi|\tau|^2 h(\tilde{q})) + e^{-2\pi|\tau|^2 h(\tilde{q})} \sum_{j=1}^{\ell-1} \frac{\eta_j(\tilde{q})}{h(\tilde{q})^j} \cdot P_j(h(\tau\tilde{q})).$$

The equality (3.3.4) follows easily from this and (3.1.9).

Applying  $j_{t*}$  to both sides of (3.3.4) yields

$$j_{t*}[\mathfrak{g}^\circ(q)] = \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) + c \cdot \log |t| \right) \wedge \delta_{\tilde{X}_t}.$$

To show that this is a logarithmic expansion, one must verify the continuity and Hölder continuity at  $t = 0$  of  $G_0(t)(\varphi)$  and  $G_1(t)(\varphi)$ , respectively, for any smooth compactly supported form  $\varphi$  on  $\tilde{X}$ . Using a partition of unity argument, we may reduce to the case in which the support of  $\varphi$  is contained in a coordinate neighborhood

$$\begin{aligned} \tilde{U} &= \{(z_1, \dots, z_n, w_1, \dots, w_m, t) \in \Delta^n \times \mathbb{C}^m \times \mathbb{C} : z_i = tw_i, \forall 1 \leq i \leq m\} \\ &\subset \{(z_{m+1}, \dots, z_n, w_1, \dots, w_m, t) \in \mathbb{C}^{n-m} \times \mathbb{C}^m \times \mathbb{C}\} \end{aligned}$$

chosen as in the proof of Proposition 3.3.2. In particular,  $\tilde{q}_i = w_i$  for all  $1 \leq i \leq \ell$ , and the function  $h(\tilde{q})$  has the form

$$H(z, w, t) = f_1(z, w, t) \cdot |w_1|^2 + \dots + f_\ell(z, w, t) \cdot |w_\ell|^2$$

for smooth compactly supported  $f_1, \dots, f_\ell : \mathbb{C}^{n-m} \times \mathbb{C}^m \times \mathbb{C} \rightarrow \mathbb{R}^{>0}$ .

The continuity of  $G_0(t)(\varphi)$  now amounts to the continuity in  $t$  of

$$\int_{\mathbb{C}^{n-m} \times \mathbb{C}^m} \frac{g(z, w, t)}{H(z, w, t)^{\ell-1}} \cdot \mu \quad \text{and} \quad \int_{\mathbb{C}^{n-m} \times \mathbb{C}^m} g(z, w, t) \cdot \log H(z, w, t) \cdot \mu$$



for any smooth compactly supported function  $g(z, w, t)$  on  $\mathbb{C}^{n-m} \times \mathbb{C}^m \times \mathbb{C}$ , where

$$\mu = dz_{m+1} \wedge d\bar{z}_{m+1} \wedge \cdots \wedge dz_n \wedge d\bar{z}_n \wedge dw_1 \wedge d\bar{w}_1 \wedge \cdots \wedge dw_m \wedge d\bar{w}_m.$$

The smoothness (hence Hölder continuity) of  $G_1(t)(\varphi)$  amounts to the smoothness in  $t$  of

$$\int_{\mathbb{C}^{n-m} \times \mathbb{C}^m} g(z, w, t) \cdot \mu.$$

These are routine calculus exercises, left to the reader.  $\square$

For the Green currents  $\mathfrak{g}^\circ(q_i)$  associated to the individual components of  $q = (q_1, \dots, q_\ell)$ , one has a more precise version of Lemma 3.3.3.

**Lemma 3.3.4.** *For  $1 \leq i \leq \ell$  there is a smooth function  $f_i$  on  $\tilde{X}$  such that*

$$\mathfrak{g}^\circ(q_i) = j_t^* \left( -\log(2\pi e^\gamma h(\tilde{q}_i)) + |\tau|^2 f_i - 2 \log |\tau| \right)$$

for all  $t \in \mathbb{C} \setminus \{0\}$ . If we define currents

$$G_0 = \left( -\log(2\pi e^\gamma h(\tilde{q}_i)) + |\tau|^2 f_i \right) \wedge \delta_{\tilde{X}_t}$$

and  $G_1 = -2\delta_{\tilde{X}_t}$  on  $N_{X_0/X}$ , then

$$j_{t*}[\mathfrak{g}^\circ(q_i)] = G_0 + G_1 \cdot \log |t|$$

is a logarithmic expansion of  $\mathfrak{g}^\circ(q_i)$  along  $X_0$ .

*Proof.* The proof is the same as that of Lemma 3.3.3, except that one replaces (3.3.5) with the simpler equality  $\mathfrak{g}^\circ(\tilde{q}_i) = E_1(2\pi h(\tilde{q}_i))$  of (3.1.8).  $\square$

**Proposition 3.3.5.** *We have the equality of currents*

$$\sigma_{X_0/X}(\mathfrak{g}^\circ(p)) = [\pi_0^* \mathfrak{g}^\circ(p|_{X_0})] \in D_{N_{X_0/X}}^{k-1, k-1}$$

where  $\pi_0 : N_{X_0/X} \rightarrow X_0$  is the bundle map, and there are smooth forms  $a_0$  and  $b_0$  on  $N_{X_0/X}$  such that

$$\sigma_{X_0/X}(\mathfrak{g}^\circ(q)) = \frac{a_0}{h(\sigma_{X_0/X}(q))^{\ell-1}} + b_0 \log h(\sigma_{X_0/X}(q)) \in D_{N_{X_0/X}}^{\ell-1, \ell-1},$$

For the individual components of  $q = (q_1, \dots, q_\ell)$  we have the exact formula

$$\sigma_{X_0/X}(\mathfrak{g}^\circ(q_i)) = -\log h(\sigma_{X_0/X}(q_i)) - \log(2\pi e^\gamma) \in D_{N_{X_0/X}}^{0,0}.$$

*Proof.* For  $\mathfrak{g}^\circ(p)$ , note that the tuple  $p$  remains smooth (as one can check in the local coordinates of Proposition 3.3.2) after pullback via any arrow in

$$\begin{array}{ccc} N_{X_0/X} & \xrightarrow{j_0} & \tilde{X} \\ \pi_0 \downarrow & & \downarrow \pi \\ X_0 & \xrightarrow{i_0} & X. \end{array}$$

Each of these pullbacks has its own Green form  $\mathfrak{g}^\circ(\cdot)$  associated to it, and these satisfy obvious functorial properties, e.g.  $\pi^*\mathfrak{g}^\circ(p) = \mathfrak{g}^\circ(\pi^*p)$ . For any  $t \neq 0$  we have the (particularly simple) logarithmic expansion

$$j_{t*}[\mathfrak{g}^\circ(p)] = \mathfrak{g}^\circ(\pi^*p) \wedge \delta_{\tilde{X}_t},$$

of  $\mathfrak{g}^\circ(p)$  along  $X_0 \subset X$ . Of course one must check that the family of currents on the right hand side is defined at  $t = 0$ , and satisfies the continuity condition of Definition 2.2.4; using the analysis of singularities of  $\mathfrak{g}^\circ(\pi^*p)$  from (3.1.6), this is an easy calculation in local coordinates as in the proof of Lemma 3.3.3. The constant term at  $t = 0$  of this expansion is

$$\mathfrak{g}^\circ(\pi^*p) \wedge \delta_{N_{X_0/X}} = j_{0*}[\mathfrak{g}^\circ(\pi^*p)|_{N_{X_0/X}}] = j_{0*}[\pi_0^*\mathfrak{g}^\circ(p)|_{X_0}],$$

proving the first claim.

The claims about  $\mathfrak{g}^\circ(q)$  and  $\mathfrak{g}^\circ(q_i)$  follow by taking  $t = 0$  in the logarithmic expansions of Lemmas 3.3.3 and 3.3.4, and recalling from Lemma 3.3.1 that the restriction of  $\tilde{q}$  to the fiber  $N_{X_0/X} = \tilde{X}_0$  is  $\sigma_{X_0/X}(q)$ .  $\square$

*Remark 3.3.6.* Using Proposition 3.1.5 and (3.1.8), each section  $\sigma_{X_0/X}(q_i)$  of the hermitian line bundle  $\pi_0^*L_0^\vee$  on  $N_{X_0/X}$  determines a Green function

$$\mathfrak{g}^\circ(\sigma_{X_0/X}(q_i)) = E_1(2\pi h(\sigma_{X_0/X}(q_i)))$$

for the divisor

$$Z(\sigma_{X_0/X}(q_i)) = \sigma_{X_0/X}(Z(q_i)),$$

but Proposition 3.3.5 shows that

$$\mathfrak{g}^\circ(\sigma_{X_0/X}(q_i)) \neq \sigma_{X_0/X}(\mathfrak{g}^\circ(q_i)).$$

This should not cause confusion. The Green function on the left hand side plays no role in our arguments, and will never appear again.

**Proposition 3.3.7.** *The specializations of  $\mathfrak{g}^\circ(s)$ ,  $\mathfrak{g}^\circ(p)$ , and  $\mathfrak{g}^\circ(q)$  to  $N_{X_0/X}$  are related by*

$$\begin{aligned} \sigma_{X_0/X}(\mathfrak{g}^\circ(s)) &= \sigma_{X_0/X}(\mathfrak{g}^\circ(p)) \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q)) \\ &\quad - \partial\sigma_{X_0/X}(A(p; q)) - \bar{\partial}\sigma_{X_0/X}(B(p; q)), \end{aligned}$$

where  $A(p; q)$  and  $B(p; q)$  are the currents of Theorem 3.2.5. Moreover,

$$\begin{aligned} \sigma_{X_0/X}(\mathfrak{g}^\circ(q)) &= \sigma_{X_0/X}(\mathfrak{g}^\circ(q_1)) \star \cdots \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q_\ell)) \\ &\quad - \partial\sigma_{X_0/X}(\mathfrak{a}(q)) - \bar{\partial}\sigma_{X_0/X}(\mathfrak{b}(q)) \end{aligned}$$

where  $\mathfrak{a}(q)$  and  $\mathfrak{b}(q)$  are the currents of (3.2.4). In particular, all currents on  $X$  appearing in these formulas admit logarithmic expansions along  $X_0$ , and the star products in both formulas are defined.

*Proof.* The core of the proof is the following lemma.

**Lemma 3.3.8.** *Suppose  $G \in D_X^{k-1, k-1}$  is any Green current for  $Z(p)$ . If  $G$  admits a logarithmic expansion along  $X_0 \subset X$ , then so does  $G \star \mathfrak{g}^\circ(q)$ , and its specialization to the normal bundle satisfies*

$$\sigma_{X_0/X}(G \star \mathfrak{g}^\circ(q)) = \sigma_{X_0/X}(G) \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q)).$$

*In particular, the star product on the right is defined.*

*Proof.* Abbreviate  $Z = Z(p)$ , and recall from Lemma 3.2.4 that the star product

$$G \star \mathfrak{g}^\circ(q) = \delta_Z \wedge \mathfrak{g}^\circ(q) + G \wedge \omega^\circ(q)$$

is defined. Applying  $j_{t*}$  to both sides results in

$$j_{t*}[G \star \mathfrak{g}^\circ(q)](\varphi) = \int_Z \mathfrak{g}^\circ(q) \wedge j_t^* \varphi + (j_{t*}G)(\pi^* \omega^\circ(q) \wedge \varphi),$$

for any smooth compactly supported form  $\varphi$  on  $\tilde{X}$ , and any  $t \neq 0$ .

Using  $j_t : Z \cong \tilde{Z}_t$  and the equality

$$\mathfrak{g}^\circ(q) = j_t^* \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) + c \cdot \log |\tau| \right)$$

of Lemma 3.3.3, the integral on the right becomes

$$\int_Z \mathfrak{g}^\circ(q) \wedge j_t^* \varphi = \int_{\tilde{Z}_t} \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) + c \cdot \log |\tau| \right) \wedge \varphi.$$

Fixing a logarithmic expansion  $j_{t*}G = \sum_{i \geq 0} G_i(t)(\log |t|)^i$ , we obtain

$$j_{t*}[G \star \mathfrak{g}^\circ(q)] = \sum_{i \geq 0} C_i(t) \cdot (\log |t|)^i$$

in which

$$\begin{aligned} C_0(t) &= \delta_{\tilde{Z}_t} \wedge \left( \frac{a}{h(\tilde{q})^{\ell-1}} + b \log h(\tilde{q}) \right) + G_0(t) \wedge \pi^* \omega^\circ(q) \\ C_1(t) &= c \wedge \delta_{\tilde{Z}_t} + G_1(t) \wedge \pi^* \omega^\circ(q) \\ C_i(t) &= G_i(t) \wedge \pi^* \omega^\circ(q) \quad \text{for } i > 1. \end{aligned}$$

To see that this is a logarithmic expansion of  $G \star \mathfrak{g}^\circ(q)$  one must check that the terms involving  $\delta_{\tilde{Z}_t}$  are well-defined currents (including at  $t = 0$ ) that satisfy the continuity conditions of Definition 2.2.4; this is easily verified in the local coordinates of the proof of Proposition 3.3.2.

The current  $C_0(0)$  is the pushforward via  $j_0 : N_{X_0/X} \rightarrow \tilde{X}$  of

$$\delta_{\sigma_{X_0/X}(Z)} \wedge \sigma_{X_0/X}(\mathfrak{g}^\circ(q)) + \sigma_{X_0/X}(G) \wedge \pi_0^* \omega^\circ(q)$$

which agrees with  $\sigma_{X_0/X}(G) \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q))$  by Proposition 2.2.10.  $\square$

Recall the equality

$$\mathfrak{g}^\circ(s) = \mathfrak{g}^\circ(p) \star \mathfrak{g}^\circ(q) - \partial[A(p; q)] - \bar{\partial}[B(p; q)]$$

of Theorem 3.2.5. The currents  $A(p, q)$  and  $B(p, q)$  admit logarithmic expansions along  $X_0$  by Theorem 2.2.5 and the final claim of Theorem 3.2.5.

The star product admits a logarithmic expansion by Lemma 3.3.8. The Green current on the left admits a logarithmic expansion by Theorem 2.2.5 and (3.1.7), and also because the right hand side does. Specializing both sides to  $N_{X_0/X}$  and using Remark 2.2.9 and Lemma 3.3.8 proves the first claim of Proposition 3.3.7.

For the second claim we use the following lemma.

**Lemma 3.3.9.** *Fix  $1 \leq r < \ell$ , and let  $G \in D_X^{r-1, r-1}$  be any Green current for  $Z(q_1, \dots, q_r)$ . If  $G$  admits a logarithmic expansion along  $X_0 \subset X$ , then so does  $G \star \mathfrak{g}^\circ(q_{r+1})$ , and*

$$\sigma_{X_0/X}(G \star \mathfrak{g}^\circ(q_{r+1})) = \sigma_{X_0/X}(G) \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q_{r+1})).$$

*Proof.* The proof is virtually identical to that of Lemma 3.3.8, using Lemma 3.3.4 in place of Lemma 3.3.3.  $\square$

Applying Lemma 3.3.9 inductively shows that

$$\sigma_{X_0/X}(\mathfrak{g}^\circ(q_1) \star \dots \star \mathfrak{g}^\circ(q_r)) = \sigma_{X_0/X}(\mathfrak{g}^\circ(q_1)) \star \dots \star \sigma_{X_0/X}(\mathfrak{g}^\circ(q_r))$$

and that all star products appearing here are defined. Each term in the iterated star product formula

$$\mathfrak{g}^\circ(q) = \mathfrak{g}^\circ(q_1) \star \dots \star \mathfrak{g}^\circ(q_\ell) - \partial[\mathfrak{a}(q)] - \bar{\partial}[\mathfrak{b}(q)]$$

of (3.2.4) admits a logarithmic expansion along  $X_0 \subset X$ . Thus we may specialize to the normal bundle term-by-term and use Remark 2.2.9 to obtain the desired formula.  $\square$

#### 4. ORTHOGONAL SHIMURA VARIETIES

We now apply the general theory of the previous subsections to the special case in which  $X$  is either the hermitian symmetric domain  $\mathcal{D}$  associated to an orthogonal group over a totally real field, or the complex Shimura variety  $M(\mathbb{C})$  determined by such a group. This allows us to prove our main result: a description of the behavior of special arithmetic cycles on the canonical model  $M$  under pullback via the inclusion  $M_0 \rightarrow M$  of a smaller orthogonal Shimura variety.

**4.1. The hermitian symmetric domain.** Let  $(V, Q)$  be a quadratic space of dimension  $n + 2 \geq 3$  over a totally real number field  $F$ . Assume there is one embedding  $\sigma : F \rightarrow \mathbb{R}$  for which

$$V_\sigma = V \otimes_{F, \sigma} \mathbb{R}$$

has signature  $(n, 2)$ , while  $V_\tau = V \otimes_{F, \tau} \mathbb{R}$  is positive definite for all embeddings  $\tau \neq \sigma$ . Denote by

$$[x, y] = Q(x + y) - Q(x) - Q(y)$$

the associated  $F$ -bilinear form on  $V$ . Extend it  $\mathbb{R}$ -bilinearly to  $V_\sigma$ , and  $\mathbb{C}$ -bilinearly to  $V_\sigma \otimes_{\mathbb{R}} \mathbb{C}$ .

The data  $(V, Q)$  determines a hermitian symmetric domain.

$$\mathcal{D} = \{z \in V_\sigma \otimes_{\mathbb{R}} \mathbb{C} : [z, z] = 0, [z, \bar{z}] < 0\} / \mathbb{C}^\times.$$

Denote by  $L$  the tautological bundle on  $\mathcal{D}$ , whose fiber at a point  $z \in \mathcal{D}$  is the isotropic line  $L_z = \mathbb{C}z \subset V_\sigma \otimes_{\mathbb{R}} \mathbb{C}$ . Endow the line  $L_z$  with the positive definite hermitian form  $h$  determined by

$$(4.1.1) \quad h(z, z) = -\frac{[z, \bar{z}]}{2}.$$

This makes  $L$  into a hermitian line bundle.

The constant vector bundle  $V_{\mathcal{D}} = V_\sigma \otimes_{\mathbb{R}} \mathcal{O}_{\mathcal{D}}$  is endowed with a symmetric bilinear pairing

$$[\cdot, \cdot] : V_{\mathcal{D}} \times V_{\mathcal{D}} \rightarrow \mathcal{O}_{\mathcal{D}}$$

under which  $L \subset V_{\mathcal{D}}$  is isotropic. Hence there is a filtration by local direct summands  $L \subset L^\perp \subset V_{\mathcal{D}}$ , and a canonical isomorphism

$$(4.1.2) \quad V_{\mathcal{D}}/L^\perp \cong L^\vee.$$

Using this last isomorphism, any  $x \in V_\sigma$  determines first a global section of  $V_{\mathcal{D}}$ , and then a global section

$$(4.1.3) \quad s(x) \in H^0(\mathcal{D}, L^\vee)$$

with zero locus the smooth analytic divisor

$$Z(x) = \{z \in \mathcal{D} : [z, x] = 0\}.$$

More generally, any tuple  $x = (x_1, \dots, x_d) \in V_\sigma^d$  determines a tuple  $s(x) = (s(x_1), \dots, s(x_d))$  of sections, and we denote by

$$Z(x) \subset \mathcal{D}$$

the analytic subspace defined by the vanishing of all components. In other words,  $Z(x)$  is those lines  $\mathbb{C}z \subset \mathcal{D}$  such that  $[z, x_i] = 0$  for all  $1 \leq i \leq d$ . This is a complex submanifold, which depends only on  $\text{Span}_{\mathbb{R}}\{x_1, \dots, x_d\} \subset V_\sigma$ . It is nonempty precisely when this subspace is positive definite, in which case it has codimension  $\dim_{\mathbb{R}} \text{Span}_{\mathbb{R}}\{x_1, \dots, x_d\}$ . Recalling the notation and terminology of §3.1, the smoothness and regularity of the tuple  $s(x)$  are both equivalent to the linear independence of the vectors  $x_1, \dots, x_d$ .

Now fix a positive definite  $v \in \text{Sym}_d(\mathbb{R})$  and an  $\alpha \in \text{GL}_d(\mathbb{R})$  with positive determinant such that

$$v = \alpha \cdot {}^t \alpha.$$

If  $x \in V_\sigma^d$  is a tuple with linearly independent components, we may form a new  $d$ -tuple  $x\alpha \in V_\sigma^d$ , and hence a corresponding smooth tuple  $s(x\alpha)$  of sections of  $L^\vee$ . Applying the constructions of §3.1 to this tuple of sections determines forms

$$(4.1.4) \quad \begin{aligned} \mathfrak{g}^\circ(x, v) &= \mathfrak{g}^\circ(s(x\alpha)) \in E_{\mathcal{D} \setminus Z(x)}^{d-1, d-1} \\ \omega^\circ(x, v) &= \omega^\circ(s(x\alpha)) \in E_{\mathcal{D}}^{d, d} \end{aligned}$$

related by the Green equation

$$dd^c[\mathfrak{g}^\circ(x, v)] + \delta_{Z(x)} = [\omega^\circ(x, v)].$$

These forms are independent of  $\alpha$  by Proposition 2.6(d) of [GS19].

**4.2. Arithmetic cycle classes.** The quadratic space  $(V, Q)$  determines a short exact sequence

$$1 \rightarrow \mathbb{G}_{mF} \rightarrow \mathrm{GSpin}(V) \rightarrow \mathrm{SO}(V) \rightarrow 1$$

of reductive groups over  $F$ . From now on we denote by  $G$  either

$$\mathrm{Res}_{F/\mathbb{Q}} \mathrm{GSpin}(V) \quad \text{or} \quad \mathrm{Res}_{F/\mathbb{Q}} \mathrm{SO}(V).$$

For our purposes these two groups are interchangeable. The group  $G(\mathbb{R})$  acts on  $\mathcal{D}$  via the projection

$$G(\mathbb{R}) \rightarrow \prod_{\tau:F \rightarrow \mathbb{R}} \mathrm{SO}(V_\tau) \rightarrow \mathrm{SO}(V_\sigma),$$

and the pair  $(G, \mathcal{D})$  is a Shimura datum. A choice of sufficiently small compact open subgroup  $K \subset G(\mathbb{A}_f)$  determines a smooth quasi-projective variety  $M$  over the reflex field  $F \cong \sigma(F) \subset \mathbb{C}$  with  $\mathbb{C}$ -points

$$M(\mathbb{C}) = G(\mathbb{Q}) \backslash \mathcal{D} \times G(\mathbb{A}_f) / K.$$

It is projective if and only if  $V$  is anisotropic. Any  $g \in G(\mathbb{A}_f)$  determines an open and closed submanifold

$$(4.2.1) \quad (gKg^{-1} \cap G(\mathbb{Q})) \backslash \mathcal{D} \subset M(\mathbb{C}),$$

where the inclusion is  $z \mapsto (z, g)$ .

Let  $V_M = V \otimes_F \mathcal{O}_M$  be the constant vector bundle determined by  $V$ . By the theory of automorphic vector bundles, it has a bilinear pairing

$$[\cdot, \cdot] : V_M \times V_M \rightarrow \mathcal{O}_M$$

and a filtration  $L_M \subset L_M^\perp \subset V_M$  by local direct summands inducing an isomorphism

$$(4.2.2) \quad V_M / L_M^\perp \cong L_M^\vee$$

that agrees with (4.1.2) under the complex uniformization.

Fix an integer  $d$  with  $1 \leq d \leq n + 1$ . The group  $G(\mathbb{A}_f)$  acts on  $\widehat{V} = V \otimes_{\mathbb{Q}} \mathbb{A}_f$ , and we fix a  $K$ -invariant  $\mathbb{Z}$ -valued Schwartz function  $\varphi \in S(\widehat{V}^d)$ . Any  $g \in G(\mathbb{A}_f)$  and  $T \in \mathrm{Sym}_d(F)$  determine an analytic cycle

$$(4.2.3) \quad Z(T, \varphi)(\mathbb{C})_g = \sum_{\substack{x \in V^d \\ T(x)=T}} \varphi(g^{-1}x) \cdot Z(x),$$

on (4.2.1), where we have set

$$T(x) = \left( \frac{[x_i, x_j]}{2} \right) \in \mathrm{Sym}_d(F).$$

Varying  $g$  yields an analytic cycle  $Z(T, \varphi)(\mathbb{C})$  on  $M(\mathbb{C})$ , which, being a union of smaller Shimura varieties constructed in the same way as  $M$ , is the complexification of an algebraic cycle  $Z(T, \varphi)$  of codimension  $\text{rank}(T)$  on the canonical model  $M$ .

Fix a positive definite  $v \in \text{Sym}_d(\mathbb{R})$  and assume  $\det(T) \neq 0$ . As in §4.3 of [GS19], the sums

$$(4.2.4) \quad \begin{aligned} \mathfrak{g}^\circ(T, v, \varphi)_g &= \sum_{\substack{x \in V^d \\ T(x)=T}} \varphi(g^{-1}x) \cdot \mathfrak{g}^\circ(x, v) \in E_{\mathcal{D} \setminus Z(T, \varphi)(\mathbb{C})_g}^{d-1, d-1} \\ \omega^\circ(T, v, \varphi)_g &= \sum_{\substack{x \in V^d \\ T(x)=T}} \varphi(g^{-1}x) \cdot \omega^\circ(x, v) \in E_{\mathcal{D}}^{d, d} \end{aligned}$$

descend to the quotient (4.2.1). By varying  $g$ , we obtain forms

$$\mathfrak{g}^\circ(T, v, \varphi) \in E_{M(\mathbb{C}) \setminus Z(T, \varphi)(\mathbb{C})}^{d-1, d-1} \quad \text{and} \quad \omega^\circ(T, v, \varphi) \in E_{M(\mathbb{C})}^{d, d}$$

related by the Green equation

$$dd^c[\mathfrak{g}^\circ(T, v, \varphi)] + \delta_{Z(T, \varphi)} = [\omega^\circ(T, v, \varphi)],$$

and an arithmetic cycle class

$$(4.2.5) \quad \widehat{Z}(T, v, \varphi) = (Z(T, \varphi), \mathfrak{g}^\circ(T, v, \varphi)) \in \widehat{\text{CH}}^d(M).$$

We would like to extend the definition to include singular  $T$ .

Recall that we have endowed the tautological line bundle  $L$  on  $\mathcal{D}$  with the hermitian metric  $h$  of (4.1.1), and have endowed  $L^\vee$  with the dual metric. These induce metrics on the canonical models  $L_M$  and  $L_M^\vee$ , and so determine arithmetic cycle classes

$$L_M, L_M^\vee \in \widehat{\text{CH}}^1(M)$$

using the arithmetic Chern class map from §III.4.2 of [Sou92]. Of course  $L_M^\vee = -L_M$ . A distinguished role is played by

$$(4.2.6) \quad \widehat{\omega}^{-1} = L_M^\vee + (0, -\log(2\pi e^\gamma)) \in \widehat{\text{CH}}^1(M).$$

In other words, if we endow  $L_M$  with the rescaled metric  $(2\pi e^\gamma)^{-1}h$ , then (4.2.6) is the image of its dual under the arithmetic Chern class map. Abbreviate

$$\Omega = \text{ch}(L_M^\vee) \in E_{M(\mathbb{C})}^{1,1}$$

for the Chern form of the dual of  $(L_M, h)$ , and note that  $\Omega$  is also the Chern form of (4.2.6).

*Remark 4.2.1.* Our  $L_M$  agrees with the  $\mathcal{E}$  in (5.160) of [GS19], but our  $\widehat{\omega}$  differs from theirs by an inverse and a rescaling of metrics.

**Theorem 4.2.2** (Garcia-Sankaran). *Assume that  $V$  is anisotropic. There are arithmetic cycle classes*

$$\widehat{Z}(T, v, \varphi) \in \widehat{\text{CH}}^d(M)$$

indexed by  $T \in \text{Sym}_d(F)$ , positive definite  $v \in \text{Sym}_d(\mathbb{R})$ , and  $K$ -fixed  $\mathbb{Z}$ -valued  $\varphi \in S(\widehat{V})$  satisfying the following properties.

- (1) For fixed  $T$  and  $v$ , the formation of  $\widehat{Z}(T, v, \varphi)$  is linear in  $\varphi$ .
- (2) If  $\det(T) \neq 0$  then  $\widehat{Z}(T, v, \varphi)$  agrees with (4.2.5).
- (3) If  $0_d \in \text{Sym}_d(F)$  denotes the zero matrix, then

$$\widehat{Z}(0_d, v, \varphi) = \varphi(0) \cdot \underbrace{\widehat{\omega}^{-1} \cdots \widehat{\omega}^{-1}}_{d \text{ times}}.$$

- (4) Assume that  $T$  and  $v$  have the form

$$T = \begin{pmatrix} T_0 & \\ & 0_{d-r} \end{pmatrix} \quad \text{and} \quad v = {}^t\theta \cdot \begin{pmatrix} v_0 & \\ & w \end{pmatrix} \cdot \theta$$

with  $T_0 \in \text{Sym}_r(F)$  nonsingular,  $v_0 \in \text{Sym}_r(\mathbb{R})$  and  $w \in \text{Sym}_{d-r}(\mathbb{R})$  of positive determinant, and

$$\theta = \begin{pmatrix} 1_r & * \\ & 1_{d-r} \end{pmatrix} \in \text{GL}_d(\mathbb{R}).$$

If  $\varphi = \varphi^{(r)} \otimes \varphi^{(d-r)} \in S(\widehat{V}^r) \otimes S(\widehat{V}^{d-r})$  then

$$\widehat{Z}(T, v, \varphi) = \widehat{Z}(T_0, v_0, \varphi^{(r)}) \cdot \widehat{Z}(0_{d-r}, w, \varphi^{(d-r)}).$$

- (5) For any  $a \in \text{GL}_d(F)$  we have

$$\widehat{Z}(T, v, \varphi) = \widehat{Z}({}^aT, {}^av, {}^a\varphi)$$

where

$${}^aT = {}^t_aT a, \quad {}^av = \sigma(a^{-1})v\sigma({}^t_a^{-1}), \quad {}^a\varphi(x) = \varphi(xa^{-1}).$$

*Proof.* This is a minor modification of the construction of §5.4 of [GS19].

We have defined the forms (4.2.4) only when  $\det(T) \neq 0$ . If we drop this assumption, the construction of  $\omega^\circ(T, v, \varphi)$  still makes sense word-for-word. The construction of  $\mathfrak{g}^\circ(T, v, \varphi)$  does not, because Proposition 3.1.5 only applies to regular tuples, and the tuple  $s(x\alpha)$  appearing in the definition (4.1.4) is not regular if  $T(x)$  is a singular matrix. Nevertheless, Propositions 4.3 and 4.4 of [GS19] provide the construction of a current  $\mathfrak{g}^\circ(T, v, \varphi)$  on  $M(\mathbb{C})$  of type  $(d-1, d-1)$  satisfying the generalized Green equation

$$dd^c \mathfrak{g}^\circ(T, v, \varphi) + \delta_{Z(T, \varphi)} \wedge \Omega^{d-\text{rank}(T)} = \omega^\circ(T, v, \varphi).$$

We remark that when  $\text{rank}(T) < d$ , the current  $\mathfrak{g}^\circ(T, v, \varphi)$  is not represented by a locally integrable form on  $M(\mathbb{C})$ .

Now let  $\mathfrak{g}$  be any choice of Green current for the cycle  $Z(T, \varphi)$  of codimension  $r = \text{rank}(T)$ . The cycle class

$$\widehat{Z}(T, v, \varphi) = (Z(T, \varphi), \mathfrak{g}) \cdot (\widehat{\omega}^{-1})^{d-r} + (0, \mathfrak{g}^\circ(T, v, \varphi) - \mathfrak{g} \wedge \Omega^{d-r})$$

is easily seen to be independent of  $\mathfrak{g}$ . This is the same definition as (5.158) of [GS19], except that we have used the class (4.2.6) in place of  $L_M^\vee$ , and have used the current  $\mathfrak{g}^\circ(T, v, \varphi)$  instead of the modified version of Definition 4.7 of [GS19].



Properties (1) and (2) are immediate from the definitions. Property (3) follows from  $Z(0_d, \varphi) = \varphi(0) \cdot M$  and  $\mathfrak{g}^\circ(0_d, v, \varphi) = 0$ , as in (4.43) of [GS19]. Property (4) is a consequence of the relations

$$\begin{aligned} Z(T, \varphi) &= \varphi^{(d-r)}(0) \cdot Z(T_0, \varphi^{(r)}) \\ \mathfrak{g}^\circ(T, v, \varphi) &= \varphi^{(d-r)}(0) \cdot \mathfrak{g}^\circ(T_0, v_0, \varphi^{(r)}) \wedge \Omega^{d-r} + \partial A + \bar{\partial} B, \end{aligned}$$

for currents  $A$  and  $B$  on  $M(\mathbb{C})$ , as in Examples 2.14 and 4.8 of [GS19]. Property (5) follows from

$$\begin{aligned} Z(T, \varphi) &= Z({}^a T, {}^a \varphi) \\ \mathfrak{g}^\circ(T, v, \varphi) &= \mathfrak{g}^\circ({}^a T, {}^a v, {}^a \varphi), \end{aligned}$$

as in Remark 4.9 of [GS19].  $\square$

*Remark 4.2.3.* The arithmetic cycle classes of Theorem 4.2.2 are uniquely determined by the properties listed there. The key point is that for any  $T$  and  $v$  one may find an  $a \in \mathrm{GL}_d(F)$  such that the matrices  ${}^a T$  and  ${}^a v$  appearing in (5) have the form described in (4). The classes determined by such matrices are obviously determined by properties (1)-(4).

We now modify the arithmetic cycle classes of Theorem 4.2.2. Given data  $(T, v, \varphi)$  as in that theorem, choose  $a \in \mathrm{GL}_d(F)$  in such a way that

$${}^a T = \begin{pmatrix} T_0 & \\ & 0_{d-\mathrm{rank}(T)} \end{pmatrix} \quad \text{and} \quad {}^a v = {}^t \theta \cdot \begin{pmatrix} v_0 & \\ & w \end{pmatrix} \cdot \theta$$

have the form described in part (4), and define

$$\widehat{C}(T, v, \varphi) = \widehat{Z}({}^a T, {}^a v, {}^a \varphi) + (0, -\log(\det(w)) \cdot \delta_{Z(T, \varphi)} \wedge \Omega^{d-\mathrm{rank}(T)-1}).$$

Note that if  $T$  is not totally positive semi-definite, then  $Z(T, \varphi) = 0$  and the correction term disappears. If  $\det(T) \neq 0$  we understand  $\det(w) = 1$ , so that the correction term again vanishes.

**Proposition 4.2.4.** *The arithmetic cycle class  $\widehat{C}(T, v, \varphi)$  does not depend on the choice of  $a \in \mathrm{GL}_d(F)$  used in its construction. It satisfies all the properties listed in Theorem 4.2.2, except that now*

$$\widehat{C}(0_d, v, \varphi) = \varphi(0) \cdot [\underbrace{\widehat{\omega}^{-1} \cdots \widehat{\omega}^{-1}}_d + (0, -\log(\det(v)) \cdot \Omega^{d-1})].$$

In particular, if  $\varphi = \varphi_1 \otimes \cdots \otimes \varphi_d$  is a pure tensor then

$$\widehat{C}(0_d, v, \varphi) = \widehat{C}(0, v_1, \varphi_1) \cdots \widehat{C}(0, v_d, \varphi_d),$$

where  $v_1, \dots, v_d$  are the eigenvalues of  $v$ , and

$$\widehat{C}(0, v_i, \varphi_i) = \varphi_i(0) \cdot [\widehat{\omega}^{-1} + (0, -\log(v_i))] \in \widehat{\mathrm{CH}}^1(M).$$

*Proof.* For the independence of the choice of  $a$ , a linear algebra exercise shows that choosing a different  $a$  has the effect of multiplying both  $\det(v_0)$

and  $\det(w)$  by nonzero squares in  $\sigma(F)$ . Thus it suffices to show that the arithmetic cycle

$$(4.2.7) \quad (0, -\log \sigma(\xi^2) \cdot \delta_{Z(T, \varphi)} \wedge \Omega^{d-\text{rank}(T)-1}) \in \widehat{\text{CH}}^d(M)$$

is trivial for any  $\xi \in F^\times$ . If we view  $\xi$  as a (constant) rational function on  $Z(T, \varphi)$ , it determines an arithmetic cycle

$$(i_* \text{div}(\xi), i_*[-\log \sigma(\xi^2)]) = (0, -\log \sigma(\xi^2) \wedge \delta_{Z(T, \varphi)}) \in \widehat{Z}^{\text{rank}(T)+1}(M),$$

where  $i : Z(T, \varphi) \rightarrow M$  is the inclusion. As in the discussion leading to Definition 1 in §III.1.1 of [Sou92], this arithmetic cycle is trivial in the arithmetic Chow group. On the other hand, its arithmetic intersection with  $d - \text{rank}(T) - 1$  copies of  $\widehat{\omega}^{-1}$  is (4.2.7), which is therefore also trivial.

The remaining claims follow from Theorem 4.2.2 and the definitions.  $\square$

*Remark 4.2.5.* If there is no  $x \in V^d$  such that  $T(x) = T$ , then

$$\widehat{C}(T, v, \varphi) = \widehat{Z}(T, v, \varphi) = 0.$$

If  $T$  is nonsingular, this is clear from the definitions. The general case can be reduced to the nonsingular case using Remark 4.2.3.

**4.3. The pullback formula.** Now suppose we are given an orthogonal decomposition

$$V = V_0 \oplus W$$

with  $V_0$  and  $W$  of dimensions  $n_0 + 2 \geq 3$  and  $m$ , respectively. Assume moreover that  $W_\tau = W \otimes_{F, \tau} \mathbb{R}$  is positive definite for every  $\tau : F \rightarrow \mathbb{R}$ . The assumptions on  $V$  imposed in §4.1 imply that  $V_{0, \tau} = V_0 \otimes_{F, \tau} \mathbb{R}$  has signature

$$\text{sig}(V_{0, \tau}) = \begin{cases} (n_0, 2) & \text{if } \tau = \sigma \\ (n_0 + 2, 0) & \text{if } \tau \neq \sigma. \end{cases}$$

The quadratic space  $V_0$  therefore has its own Shimura datum  $(G_0, \mathcal{D}_0)$ , and the inclusion  $V_0 \subset V$  induces an injection of Shimura data

$$i_0 : (G_0, \mathcal{D}_0) \rightarrow (G, \mathcal{D})$$

realizing  $\mathcal{D}_0 \subset \mathcal{D}$  as a codimension  $m$  submanifold. Fix a compact open subgroup  $K_0 \subset G_0(\mathbb{A}_f) \cap K$ , and let  $M_0$  be the associated Shimura variety over  $F = \sigma(F)$  with complex points

$$M_0(\mathbb{C}) = G_0(\mathbb{Q}) \backslash \mathcal{D}_0 \times G_0(\mathbb{A}_f) / K_0.$$

The induced map  $i_0 : M_0 \rightarrow M$  is finite and unramified. The Shimura variety  $M_0$  has its own hermitian line bundle  $L_{M_0}$ , related to the one on  $M$  by a canonical isomorphism  $L_{M_0} \cong i_0^* L_M$ .

**Hypothesis 4.3.1.** We assume throughout that the compact open subgroups  $K_0 \subset G_0(\mathbb{A}_f)$  and  $K \subset G(\mathbb{A}_f)$  have been chosen so that

$$i_0 : M_0 \rightarrow M$$

is a closed immersion. This is always possible, by Proposition 1.15 of [Del71].

**Theorem 4.3.2.** *Assume that  $V$  is anisotropic. Fix an integer  $1 \leq d \leq n_0 + 1$ , and a  $K$ -fixed  $\mathbb{Z}$ -valued Schwartz function*

$$\varphi = \sum_j \varphi_{0,j} \otimes \psi_j \in S(\widehat{V}_0^d) \otimes S(\widehat{W}^d) \cong S(\widehat{V}^d),$$

so that §4.2 associates to any  $T \in \text{Sym}_d(F)$  and positive definite  $v \in \text{Sym}_d(\mathbb{R})$  arithmetic cycle classes

$$(4.3.1) \quad \widehat{C}_M(T, v, \varphi) \in \widehat{\text{CH}}^d(M) \quad \text{and} \quad \widehat{C}_{M_0}(T, v, \varphi_0) \in \widehat{\text{CH}}^d(M_0).$$

The specialization to the normal bundle

$$\sigma_{M_0/M} : \widehat{\text{CH}}^d(M) \rightarrow \widehat{\text{CH}}^d(N_{M_0/M})$$

satisfies

$$\sigma_{M_0/M}(\widehat{C}_M(T, v, \varphi)) = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \sum_j \psi_j(y) \cdot \pi_0^* \widehat{C}_{M_0}(T_0, v, \varphi_{0,i}),$$

where  $\pi_0 : N_{M_0/M} \rightarrow M_0$  is the bundle map.

Theorem 4.3.2 will be proved below. First, we record a corollary explaining the precise connection between the classes of (4.3.1).

**Corollary 4.3.3.** *Keeping the notation and assumptions of Theorem 4.3.2, the pullback*

$$i_0^* : \widehat{\text{CH}}^d(M) \rightarrow \widehat{\text{CH}}^d(M_0)$$

satisfies

$$i_0^* \widehat{C}_M(T, v, \varphi) = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \sum_i \psi_i(y) \cdot \widehat{C}_{M_0}(T_0, v, \varphi_{0,i}).$$

*Proof.* This is immediate from Theorems 2.3.1 and 4.3.2, along with the injectivity of  $\pi_0^*$  proved in Proposition 2.3.3.  $\square$

**4.4. Specialization of degenerate cycles.** This subsection is devoted to proving Proposition 4.4.1 below. When we prove Theorem 4.3.2, it will allow us to compute the specializations to  $N_{M_0/M}$  of those arithmetic cycles on  $M$  that intersect  $M_0$  improperly.

The action of  $G_0(\mathbb{R})$  on the pair  $\mathcal{D}_0 \subset \mathcal{D}$  induces an action on  $N_{\mathcal{D}_0/\mathcal{D}}$ , and the normal bundle to  $M_0 \rightarrow M$  has complex points

$$N_{M_0/M}(\mathbb{C}) = G_0(\mathbb{Q}) \backslash N_{\mathcal{D}_0/\mathcal{D}} \times G_0(\mathbb{A}_f) / K_0.$$

In particular, every  $g \in G_0(\mathbb{A}_f)$  determines an open and closed immersion

$$(4.4.1) \quad (gK_0g^{-1} \cap G_0(\mathbb{Q})) \backslash N_{\mathcal{D}_0/\mathcal{D}} \xrightarrow{z \mapsto (z,g)} N_{M_0/M}(\mathbb{C}).$$

**Proposition 4.4.1.** *Fix a tuple  $y = (y_1, \dots, y_d) \in W^d$  with linearly independent components, and a positive definite  $w \in \text{Sym}_d(\mathbb{R})$ . There is a cycle  $Z$  on  $N_{M_0/M}$  of codimension  $d$ , and a Green current  $\mathfrak{g} \in D_{N_{M_0/M}(\mathbb{C})}^{d-1, d-1}$  for it, such that the following properties hold.*

(1) *In the codimension  $d$  arithmetic Chow group of  $N_{M_0/M}$ , we have*

$$(Z, \mathfrak{g}) = \pi_0^* \left( \underbrace{\widehat{\omega}_0^{-1} \cdots \widehat{\omega}_0^{-1}}_d \right) + \pi_0^* (0, -\log(\det(w)) \cdot \Omega_0^{d-1}),$$

where  $\pi_0 : N_{M_0/M} \rightarrow M_0$  is the bundle map,  $\widehat{\omega}_0^{-1}$  is the analogue of (4.2.6) on  $M_0$ , and  $\Omega_0 \in E_{M_0(\mathbb{C})}^{1,1}$  is its Chern form.

(2) *For any  $g \in G_0(\mathbb{A}_f)$ , the restriction of  $(Z, \mathfrak{g})$  to (4.4.1) is equal to the pair obtained by specializing  $(Z(y), \mathfrak{g}^\circ(y, w))$  to  $N_{\mathcal{D}_0/\mathcal{D}}$ .*

Although we suppress it from the notation, the pair  $(Z, \mathfrak{g})$  depends on the the data of  $y$  and  $w$ .

The proof of Proposition 4.4.1 uses the degenerating sections of §3.3 in an essential way. First note that the closed immersion  $\mathcal{D}_0 \subset \mathcal{D}$  admits a presentation of the type considered in §3.3. More precisely, if we define vector bundles  $W_{\mathcal{D}} = W_\sigma \otimes_{\mathbb{R}} \mathcal{O}_{\mathcal{D}}$  and  $N = \underline{\text{Hom}}(W_{\mathcal{D}}, L^\vee)$  on  $\mathcal{D}$ , the composition

$$W_{\mathcal{D}} \rightarrow V_{\mathcal{D}}/L^\perp \xrightarrow{(4.1.2)} L^\vee$$

defines a global section  $u \in H^0(\mathcal{D}, N)$  whose vanishing locus is  $\mathcal{D}_0$ . In particular, (3.3.1) gives us an isomorphism

$$(4.4.2) \quad N_{\mathcal{D}_0/\mathcal{D}} \cong N|_{\mathcal{D}_0} \cong \underline{\text{Hom}}(W_{\mathcal{D}_0}, L_0^\vee).$$

Each  $y_i \in W$  determines a global section of  $W_{\mathcal{D}}$ , and hence a morphism  $y_i : N \rightarrow L^\vee$ . As in (3.3.2), this morphism determines a degenerating section

$$q_i \in H^0(\mathcal{D}, L^\vee).$$

Directly comparing the constructions,  $q_i$  is none other than the section  $s(y_i)$  of (4.1.3), and so

$$(4.4.3) \quad Z(y) = Z(q)$$

is the vanishing locus of the components of  $q = (q_1, \dots, q_d)$ . As in (3.3.3), to the degenerating section  $q_i$  there is an associated section

$$(4.4.4) \quad \sigma_{\mathcal{D}_0/\mathcal{D}}(q_i) \in H^0(N_{\mathcal{D}_0/\mathcal{D}}, \pi_0^* L_0^\vee),$$

and second claim of Proposition 3.3.2 implies

$$\sigma_{\mathcal{D}_0/\mathcal{D}}(Z(y)) = \text{div}(\sigma_{\mathcal{D}_0/\mathcal{D}}(q_1)) \cdots \text{div}(\sigma_{\mathcal{D}_0/\mathcal{D}}(q_d))$$

as analytic cycles on  $N_{\mathcal{D}_0/\mathcal{D}}$ .

The first step in proving Proposition 4.4.1 is to show that the construction of (4.4.4) can be replicated on the canonical model. For this we need an analogue of (4.4.2).

**Lemma 4.4.2.** *Let  $W_{M_0} = W \otimes_F \mathcal{O}_{M_0}$  be the constant vector bundle. There is an isomorphism*

$$N_{M_0/M} \cong \underline{\mathrm{Hom}}(W_{M_0}, L_{M_0}^\vee)$$

*of vector bundles on  $M_0$ , whose restriction to (4.4.1) is (4.4.2).*

*Proof.* Let  $W_M \subset V_M$  be the constant local direct summand determined by  $W \subset V$ . Using (4.2.2), we obtain a morphism

$$u : W_M \rightarrow V_M/L_M \cong L_M^\vee$$

whose vanishing locus is  $M_0$ . Indeed, this can be checked over the complex fiber, where it is already known. Viewing  $u$  as a global section of  $N = \underline{\mathrm{Hom}}(W_M, L_M^\vee)$ , the desired isomorphism is defined as in (3.3.1).  $\square$

We now directly imitate the construction of (4.4.4). Each  $y_i \in W$  determines a global section of  $W_{M_0}$ , and hence (by Lemma 4.4.2), a morphism

$$N_{M_0/M} \rightarrow L_{M_0}^\vee.$$

If we pull this back via the bundle map  $\pi_0 : N_{M_0/M} \rightarrow M_0$ , we obtain a morphism

$$\pi_0^* N_{M_0/M} \rightarrow \pi_0^* L_{M_0}^\vee$$

which can be evaluated at the tautological section of  $\pi_0^* N_{M_0/M}$ , exactly as in the construction of (3.3.3). The result is a section

$$Q_i \in H^0(N_{M_0/M}, \pi_0^* L_{M_0}^\vee)$$

whose restriction to (4.4.1) agrees with the section (4.4.4).

Consider the divisors  $\mathrm{div}(Q_1), \dots, \mathrm{div}(Q_d)$  on  $N_{M_0/M}$ . Their restrictions to (4.4.1) are the divisors of the sections (4.4.4), which satisfy

$$\sigma_{\mathcal{D}_0/\mathcal{D}}(Z(y)) = \mathrm{div}(\sigma_{\mathcal{D}_0/\mathcal{D}}(q_1)) \cdots \mathrm{div}(\sigma_{\mathcal{D}_0/\mathcal{D}}(q_d))$$

by (4.4.3) and Proposition 3.3.2. Thus the codimension  $d$  cycle on  $N_{M_0/M}$  defined by

$$Z = \mathrm{div}(Q_1) \cdots \mathrm{div}(Q_d)$$

has the property that its pullback under (4.4.1) is equal to  $\sigma_{\mathcal{D}_0/\mathcal{D}}(Z(y))$ .

As  $Z$  is defined as a proper intersection of divisors of sections of the hermitian line bundle  $\pi_0^* L_{M_0}^\vee \rightarrow N_{M_0/M}$ , it is easy to construct Green currents for it. Each divisor  $\mathrm{div}(Q_i)$  has a Green current  $-\log(2\pi e^\gamma h(Q_i))$ , and so the iterated star product

$$G = [-\log(2\pi e^\gamma h(Q_1))] \star \cdots \star [-\log(2\pi e^\gamma h(Q_d))] \in D_{N_{M_0/M}(\mathbb{C})}^{d-1, d-1}$$

is a Green current for  $Z$ .

This construction can be generalized. For  $\beta \in \mathrm{GL}_d(\mathbb{R})$ , consider the tuple

$$(Q'_1, \dots, Q'_d) = (Q_1, \dots, Q_d) \cdot \beta \in H^0(N_{M_0/M}(\mathbb{C}), \pi_0^* L_{M_0(\mathbb{C})}^\vee)^d$$

of sections defined over the complex fiber (of course they will not be defined over the reflex field  $F = \sigma(F)$ , unless  $\beta$  is). Because

$$\mathrm{div}(Q'_1) \cdots \mathrm{div}(Q'_d) = Z(\mathbb{C}),$$

the iterated star product

$$G(\beta) = [-\log(2\pi e^\gamma h(Q'_1))] \star \cdots \star [-\log(2\pi e^\gamma h(Q'_d))] \in D_{N_{M_0/M}(\mathbb{C})}^{d-1, d-1}$$

is a Green current for  $Z$ .

**Lemma 4.4.3.** *For any  $\beta \in \mathrm{GL}_d(\mathbb{R})$ , the pullback of*

$$\underbrace{\widehat{\omega}_0^{-1} \cdots \widehat{\omega}_0^{-1}}_{d \text{ times}} + (0, -\log |\beta|^2 \cdot \Omega_0^{d-1}) \in \widehat{\mathrm{CH}}^d(M_0)$$

via the bundle map  $\pi_0 : N_{M_0/M} \rightarrow M_0$  is represented by the arithmetic cycle

$$(Z, G(\beta)) \in \widehat{Z}^d(N_{M_0/M}).$$

*Proof.* By construction,  $(Z, G)$  is the arithmetic intersection of the

$$(\mathrm{div}(Q_i), -\log(h(Q_i))) + (0, -\log(2\pi e^\gamma)) \in \widehat{\mathrm{CH}}^1(N_{M_0/M})$$

as  $i$  varies over  $1 \leq i \leq d$ . Each  $Q_i$  is a section of  $\pi_0^* L_{M_0}^\vee$ , and so, recalling (4.2.6), each of these arithmetic divisors represents

$$\pi_0^* L_{M_0}^\vee + (0, -\log(2\pi e^\gamma)) = \pi_0^* \widehat{\omega}_0^{-1}.$$

Thus

$$(Z, G) = \underbrace{\pi_0^* \widehat{\omega}_0^{-1} \cdots \pi_0^* \widehat{\omega}_0^{-1}}_{d \text{ times}} \in \widehat{\mathrm{CH}}^d(N_{M_0/M}),$$

and the claim is true when  $\beta$  is the identity matrix.

It now suffices to show that for any  $\alpha, \beta \in \mathrm{GL}_d(\mathbb{R})$ , we have

$$(Z, G(\alpha\beta)) = (Z, G(\alpha)) + (0, -\log |\beta|^2 \cdot \Omega_0^{d-1})$$

in the arithmetic Chow group. If  $\beta$  is a permutation matrix, this follows from the usual associativity and commutativity of the star product (modulo currents of the form  $\partial a + \bar{\partial} b$ ). The cases

$$\beta = \begin{pmatrix} 1 & 1 & & \\ 0 & 1 & & \\ & & \ddots & \\ & & & I_{d-2} \end{pmatrix} \quad \text{and} \quad \beta = \begin{pmatrix} \lambda & & & \\ & & & \\ & & & \\ & & & I_{d-1} \end{pmatrix}$$

with  $\lambda \in \mathbb{R}^\times$  follow immediately from the definition of the star product. The general case follows by writing  $\beta$  as a product of such matrices.  $\square$

*Proof of Proposition 4.4.1.* Factor  $w = \beta \cdot {}^t\beta$  with  $\beta \in \mathrm{GL}_d(\mathbb{R})$  of positive determinant, and consider the pair  $(Z, G(\beta))$ . Lemma 4.4.3 shows that its image in the arithmetic Chow group of  $N_{M_0/M}$  represents the pullback of

$$\underbrace{\widehat{\omega}_0^{-1} \cdots \widehat{\omega}_0^{-1}}_d + (0, -\log(\det(w)) \cdot \Omega_0^{d-1}) \in \widehat{\mathrm{CH}}^d(M_0),$$

as desired. We already saw that the restriction of  $Z$  to (4.4.1) is  $\sigma_{\mathcal{D}_0/\mathcal{D}}(Z(y))$ , and so it remains to understand the restriction of  $G(\beta)$ .

Recall that the restriction of  $Q_i$  is the section  $\sigma_{X_0/X}(q_i)$  of (4.4.4). Using this, Proposition 3.3.5 implies that the restriction of  $G(\beta)$  is equal to

$$\begin{aligned} & [-\log(2\pi h(\sigma_{\mathcal{D}_0/\mathcal{D}}(q'_1)))] \star \cdots \star [-\log(2\pi e^\gamma h(\sigma_{\mathcal{D}_0/\mathcal{D}}(q'_d)))] \\ &= \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(q'_1)) \star \cdots \star \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(q'_d)), \end{aligned}$$

where the  $q'_i \in H^0(\mathcal{D}, L^\vee)$  are the components of the tuple  $q' = q\beta$ . By Proposition 3.3.7, this is equal to

$$\sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(q')) + \partial a + \bar{\partial} b$$

for some currents  $a$  and  $b$  on (4.4.1). On the other hand

$$q' = q\beta = s(y\beta),$$

and so  $\mathfrak{g}^\circ(q') = \mathfrak{g}^\circ(y, w)$ . In other words, the restriction of  $G(\beta)$  to (4.4.1) is equal to

$$\sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(y, w)) + \partial a + \bar{\partial} b.$$

Setting  $\mathfrak{g} = G(\beta) - \partial a - \bar{\partial} b$ , the pair  $(Z, \mathfrak{g})$  enjoys the properties stated in Proposition 4.4.1.  $\square$

**4.5. Proof of Theorem 4.3.2.** Keep the notation and assumptions of Theorem 4.3.2. To ease notation, assume  $\varphi = \varphi_0 \otimes \psi \in S(\widehat{V}_0^d) \otimes S(\widehat{W}^d)$ .

Assume for the moment that  $\det(T) \neq 0$ . Using the orthogonal decomposition

$$V = V_0 \oplus W,$$

each  $x \in V^d$  decomposes as  $x = x_0 + y$ , with  $x_0 \in V_0^d$  and  $y \in W^d$  satisfying

$$T(x_0) + T(y) = T(x).$$

Working with one fixed  $g \in G_0(\mathbb{A}_f)$  in (4.2.1), this allows us to decompose (4.2.3) and (4.2.4) as

$$(4.5.1) \quad Z(T, \varphi)(\mathbb{C})_g = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \psi(y) \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1}x_0) Z(x_0 + y)$$

and

$$(4.5.2) \quad \mathfrak{g}^\circ(T, v, \varphi)_g = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \psi(y) \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1}x_0) \mathfrak{g}^\circ(x_0 + y, v).$$

To compute their specializations to  $N_{\mathcal{D}_0/\mathcal{D}}$ , it suffices to do so for the inner summations for a fixed  $T_0$ ,  $v$ , and  $y$ .

This is done by reduction to the following special case. Suppose that for some  $1 \leq r \leq d$  we have

$$T_0 = \begin{pmatrix} S_0 & 0 \\ 0 & 0 \end{pmatrix} \in \text{Sym}_d(F)$$

with  $S_0 \in \text{Sym}_r(F)$  nonsingular, and

$$v = \begin{pmatrix} v_0 & 0 \\ 0 & w \end{pmatrix} \in \text{Sym}_d(\mathbb{R})$$

with  $v_0 \in \text{Sym}_r(\mathbb{R})$  and  $w \in \text{Sym}_{d-r}(\mathbb{R})$ . Let  $y \in W^d$  be any tuple such that

$$\text{rank}(T_0 + T(y)) = d,$$

and write  $y = (y', y'')$  as the concatenation of  $y' \in W^r$  and  $y'' \in W^{d-r}$ .

**Lemma 4.5.1.** *Assume  $V_0$  is anisotropic, and that*

$$\varphi_0 = \varphi_0^{(r)} \otimes \varphi_0^{(d-r)} \in S(\widehat{V}_0^r) \otimes S(\widehat{V}_0^{d-r}).$$

*In the situation above, we have the equalities*

$$\begin{aligned} & \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1}x_0) \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z(x_0 + y)) \\ &= \varphi_0^{(d-r)}(0) \cdot \pi_0^* Z(S_0, \varphi_0, \varphi_0^{(r)})(\mathbb{C})_g \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z(y'')) \end{aligned}$$

and

$$\begin{aligned} & \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1}x_0) \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(x_0 + y, v)) \\ &= \varphi_0^{(d-r)}(0) \cdot \pi_0^* \mathfrak{g}^\circ(S_0, v_0, \varphi_0^{(r)})_g \star \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}^\circ(y'', w)) - \partial A_g - \bar{\partial} B_g \end{aligned}$$

of cycles and currents on

$$(gK_0g^{-1} \cap G_0(\mathbb{Q})) \backslash N_{\mathcal{D}_0/\mathcal{D}} \subset N_{M_0/M}(\mathbb{C}),$$

for some currents  $A_g$  and  $B_g$ . On the right hand side  $\pi_0 : N_{\mathcal{D}_0/\mathcal{D}} \rightarrow \mathcal{D}_0$  is the bundle map, and  $Z(S_0, \varphi_0^{(r)})(\mathbb{C})_g$  and  $\mathfrak{g}^\circ(S_0, v_0, \varphi_0^{(r)})_g$  are the cycle and current on

$$(gK_0g^{-1} \cap G_0(\mathbb{Q})) \backslash \mathcal{D}_0 \subset M_0(\mathbb{C})$$

defined in the same way as (4.2.3) and (4.2.4), but with the Shimura datum  $(G, \mathcal{D})$  replaced by  $(G_0, \mathcal{D}_0)$ .

*Proof.* Given  $x \in V^d$ , abbreviate  $x' \in V^r$  and  $x'' \in V^{d-r}$  for the tuples formed from the first  $r$  and final  $d-r$  components of  $x$ .

For any  $x_0 \in V_0^d$  satisfying  $T(x_0) = T_0$  we have  $T(x'_0) = S_0$  and  $T(x''_0) = 0$ . Hence  $x''_0 = 0$  by our assumption that  $V_0$  is anisotropic, and  $x_0 + y \in V^d$  is the concatenation of  $x'_0 + y' \in V^r$  and  $y'' \in W^{d-r} \subset V^{d-r}$ . As in the discussion surrounding (4.1.3), these tuples determine tuples of sections

$$p = s(x'_0 + y') \in H^0(\mathcal{D}, L^\vee)^r \quad \text{and} \quad q = s(y'') \in H^0(\mathcal{D}, L^\vee)^{d-r},$$

whose concatenation is  $(p, q) = s(x_0 + y)$ . These satisfy the assumptions imposed in §3.3. More precisely:



- (1) The restriction  $p|_{\mathcal{D}_0} = s(x'_0) \in H^0(\mathcal{D}_0, L_0^\vee)^r$  is the tuple of sections formed from  $x'_0 \in V_0^r$ . As  $T(x'_0) = S_0$  is nonsingular, this restriction is again smooth, and

$$Z_{\mathcal{D}_0}(p|_{\mathcal{D}_0}) = Z_{\mathcal{D}_0}(x'_0),$$

where we now use the subscripts  $\mathcal{D}$  and  $\mathcal{D}_0$  to distinguish between cycles on these two hermitian symmetric domains.

- (2) As explained in the proof of Proposition 4.4.1, the components of  $q$  are degenerating sections in the sense of §3.3.

Thus Proposition 3.3.2 applies, and shows that

$$\sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(x_0 + y)) = \pi_0^* Z_{\mathcal{D}_0}(x'_0) \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(y'')).$$

Now sum both sides of this last equality over all  $x_0 \in V_0^d$  with  $T(x_0) = T_0$ . As  $x''_0 = 0$  for every such  $x_0$ , that sum can be replaced by the sum over all  $x'_0 \in V_0^r$  satisfying  $T(x'_0) = S_0$ . The result is

$$\begin{aligned} & \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1}x_0) \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(x_0 + y)) \\ &= \sum_{\substack{x'_0 \in V_0^r \\ T(x'_0) = S_0}} \varphi_0^{(r)}(g^{-1}x'_0) \varphi_0^{(d-r)}(0) \cdot \pi_0^* Z_{\mathcal{D}_0}(x'_0) \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(y'')), \end{aligned}$$

proving the first claim of the proposition.

For the specialization of Green forms, choose the  $\alpha \in \mathrm{GL}_d(\mathbb{R})$  of (4.1.4) in the block diagonal form

$$\alpha = \begin{pmatrix} \alpha_0 & \\ & \beta \end{pmatrix}$$

with  $\alpha_0 \in \mathrm{Sym}_r(\mathbb{R})$  and  $\beta \in \mathrm{Sym}_{d-r}(\mathbb{R})$  of positive determinant. By definition,  $\mathfrak{g}^\circ(x_0 + y, v)$  is the Green current associated to the tuple of sections

$$s(x_0\alpha + y\alpha) \in H^0(\mathcal{D}, L^\vee)^d.$$

Writing this as the concatenation of  $p\alpha_0$  and  $q\beta$ , Propositions 3.3.5 and 3.3.7 imply

$$\begin{aligned} \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}_\mathcal{D}^\circ(x_0 + y, v)) &= \pi_0^* \mathfrak{g}_{\mathcal{D}_0}^\circ(x'_0, v_0) \star \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}_\mathcal{D}^\circ(y'', w)) \\ &\quad - \partial \sigma_{\mathcal{D}_0/\mathcal{D}}(A(p\alpha_0; q\beta)) - \bar{\partial} \sigma_{\mathcal{D}_0/\mathcal{D}}(B(p\alpha_0; q\beta)). \end{aligned}$$

As above, summing both sides over all  $x_0 \in V_0^d$  with  $T(x_0) = T_0$  proves the second claim of the proposition.  $\square$

The proof of Theorem 4.3.2 will now proceed in two steps; first assuming  $\det(T) \neq 0$ , and then without this assumption. As in (4.3.1), we will use subscripts of  $M$  and  $M_0$  to distinguish between the special cycles and Green currents on these two Shimura varieties.

*Proof of Theorem 4.3.2 for nonsingular matrices.* Assume  $\det(T) \neq 0$ , so that

$$\widehat{Z}_M(T, v, \varphi) = (Z_M(T, v, \varphi), \mathfrak{g}_M^\circ(T, v, \varphi))$$

For a given  $T_0 \in \text{Sym}_d(F)$  and  $y \in W^d$  satisfying  $T_0 + T(y) = T$ , abbreviate  $r = \text{rank}(T_0)$ , choose  $a \in \text{GL}_d(F)$  in such a way that the matrices

$${}^a T_0 = {}^t a T_0 a, \quad {}^a v = \sigma(a^{-1}) v \sigma({}^t a^{-1})$$

have the form

$$(4.5.3) \quad {}^a T_0 = \begin{pmatrix} S_0 & \\ & 0_{d-r} \end{pmatrix}, \quad {}^a v = {}^t \theta \cdot \begin{pmatrix} v_0 & \\ & w \end{pmatrix} \cdot \theta$$

of part (4) of Theorem 4.2.2, with  $\det(S_0) \neq 0$ , and decompose

$${}^a \varphi_0 = \sum_i \Phi_i^{(r)} \otimes \Phi_i^{(d-r)} \in S(\widehat{V}_0^r) \otimes S(\widehat{V}_0^{d-r})$$

as a sum of pure tensors.

It follows from  $T(x_0 a) = {}^a T(x_0)$  that

$$\sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1} x_0) Z_{\mathcal{D}}(x_0 + y) = \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = {}^a T_0}} {}^a \varphi_0(g^{-1} x_0) Z_{\mathcal{D}}(x_0 + y a)$$

as cycles on (4.2.1). Specializing both sides to  $N_{\mathcal{D}_0/\mathcal{D}}$  and using Lemma 4.5.1 yields

$$(4.5.4) \quad \begin{aligned} & \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1} x_0) \sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(x_0 + y)) \\ &= \sum_i \Phi_i^{(d-r)}(0) \cdot \pi_0^* Z_{\mathcal{D}_0}(S_0, \Phi_i^{(r)})(\mathbb{C})_g \cdot \sigma_{\mathcal{D}_0/\mathcal{D}}(Z_{\mathcal{D}}(\bar{y})), \end{aligned}$$

where  $\bar{y} = (\bar{y}_1, \dots, \bar{y}_{d-r}) \in W^{d-r}$  consists of the final  $d - r$  components of  $y a$ . The same reasoning shows that

$$(4.5.5) \quad \begin{aligned} & \sum_{\substack{x_0 \in V_0^d \\ T(x_0) = T_0}} \varphi_0(g^{-1} x_0) \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}_{\mathcal{D}}^\circ(x_0 + y, v)) \\ &= \sum_i \Phi_i^{(d-r)}(0) \cdot \pi_0^* \mathfrak{g}_{\mathcal{D}}^\circ(S_0, v_0, \Phi_i^{(r)}) \star \sigma_{\mathcal{D}_0/\mathcal{D}}(\mathfrak{g}_{\mathcal{D}}^\circ(\bar{y}, w)) - \partial A - \bar{\partial} B, \end{aligned}$$

where the currents  $A$  and  $B$  depend on  $T_0$ ,  $y$ , and the choice of  $a$ .

Now sum the equalities (4.5.4) and (4.5.5) over all  $T_0 \in \text{Sym}_d(F)$  and  $y \in W^d$  for which  $T_0 + T(y) = T$ . Combining Proposition 4.4.1 with the decompositions (4.5.1) and (4.5.2) proves the equality

$$\sigma_{M_0/M}(\widehat{Z}_M(T, v, \varphi)) = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \psi(y) \sum_i \pi_0^* \widehat{Z}_{M_0}(S_0, v_0, \Phi_i^{(r)}) \cdot (Z, \mathfrak{g})$$

in the arithmetic Chow group of  $N_{M_0/M}$ , where

$$(Z, \mathfrak{g}) \in \widehat{Z}^d(N_{M_0/M})$$

represents the pullback via  $\pi_0 : N_{M_0/M} \rightarrow M_0$  of the cycle class

$$\widehat{C}_{M_0}(0_{d-r}, w, \Phi_i^{(d-r)}) = \Phi_i^{(d-r)}(0) \cdot \left[ \underbrace{\widehat{\omega}_0^{-1} \cdots \widehat{\omega}_0^{-1}}_{d \text{ times}} + (0, -\log(\det(w)) \cdot \Omega_0^{d-1}) \right]$$

of Proposition 4.2.4. Note that in the inner sum the data  $S_0, v_0, w, \Phi_i, r = \text{rank}(T_0)$ , and  $(Z, \mathfrak{g})$  all depend on  $T_0, y$ , and a choice of  $a \in \text{SL}_d(F)$  as in (4.5.3). We suppress this dependence from the notation, as in the arithmetic Chow group the inner sum simplifies to

$$\begin{aligned} & \sum_i \pi_0^* \widehat{Z}_{M_0}(S_0, v_0, \Phi_i^{(r)}) \cdot (Z, \mathfrak{g}) \\ &= \sum_i \pi_0^* \widehat{C}_{M_0}(S_0, v_0, \Phi_i^{(r)}) \cdot \pi_0^* \widehat{C}_{M_0}(0_{d-r}, w, \Phi_i^{(d-r)}) \\ &= \pi_0^* \widehat{C}_{M_0}(^a T_0, ^a v, ^a \varphi_0) \\ &= \pi_0^* \widehat{C}_{M_0}(T_0, v, \varphi_0) \end{aligned}$$

by Proposition 4.2.4. In other words

$$\sigma_{M_0/M}(\widehat{Z}_M(T, v, \varphi)) = \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \psi(y) \cdot \pi_0^* \widehat{C}_{M_0}(T_0, v, \varphi_0),$$

completing the proof of Theorem 4.3.2 when  $\det(T) \neq 0$ .  $\square$

*Proof of Theorem 4.3.2 for general matrices.* Now let  $T \in \text{Sym}_d(F)$  be arbitrary, and set  $r = \text{rank}(T)$ . Using Remark 4.2.3 and Proposition 4.2.4, one immediately reduces to the case in which

$$T = \begin{pmatrix} S & \\ & 0_{d-r} \end{pmatrix} \quad \text{and} \quad v = {}^t \theta \cdot \begin{pmatrix} v_0 \\ w \end{pmatrix} \cdot \theta$$

as in part (4) of Theorem 4.2.2, with  $S \in \text{Sym}_r(F)$  nonsingular. We may also assume that the factors in  $\varphi = \varphi_0 \otimes \psi$  admit further factorizations

$$\begin{aligned} \varphi_0 &= \varphi_0^{(r)} \otimes \varphi_0^{(d-r)} \in S(\widehat{V}_0^r) \otimes S(\widehat{V}_0^{d-r}) \\ \psi &= \psi^{(r)} \otimes \psi^{(d-r)} \in S(\widehat{W}^r) \otimes S(\widehat{W}^{d-r}), \end{aligned}$$

so that Proposition 4.2.4 implies

$$(4.5.6) \quad \widehat{C}_M(T, v, \varphi) = \widehat{C}_M(S, v_0, \varphi^{(r)}) \cdot \widehat{C}_M(0_{d-r}, w, \varphi^{(d-r)})$$

with  $\varphi^{(r)} = \varphi_0^{(r)} \otimes \psi^{(r)}$ , and similarly with  $r$  replaced by  $d - r$ .

It is clear from the definitions that pullback via  $i_0 : M_0 \rightarrow M$  satisfies

$$i_0^* \widehat{\omega}^{-1} = \widehat{\omega}_0^{-1},$$

and so Proposition 4.2.4 and Theorem 2.3.1 imply

$$\sigma_{M_0/M}(\widehat{C}_M(0_{d-r}, w, \varphi^{(d-r)})) = \psi^{(d-r)}(0) \cdot \pi_0^* \widehat{C}_{M_0}(0_{d-r}, r, \varphi_0^{(d-r)}).$$

We have already proved Theorem 4.3.2 for the nonsingular matrix  $S$ , so

$$\sigma_{M_0/M}(\widehat{C}_M(S, v_0, \varphi^{(r)})) = \sum_{\substack{S_0 \in \text{Sym}_r(F) \\ y \in W^r \\ S_0 + T(y) = S}} \psi^{(r)}(y) \cdot \pi_0^* \widehat{C}_{M_0}(S_0, v_0, \varphi_0^{(r)}).$$

Specialization to the normal bundle commutes with arithmetic intersection (this is immediate from Theorem 2.3.1 and the fact that pullbacks commute with arithmetic intersection), and so the specialization of (4.5.6) is equal to the pullback via  $\pi_0 : N_{M_0/M} \rightarrow M_0$  of

$$(4.5.7) \quad \sum_{\substack{S_0 \in \text{Sym}_r(F) \\ y \in W^r \\ S_0 + T(y) = S}} \psi^{(r)}(y) \cdot \psi^{(d-r)}(0) \cdot \widehat{C}_{M_0}(S_0, v_0, \varphi_0^{(r)}) \cdot \widehat{C}_{M_0}(0_{d-r}, w, \varphi_0^{(d-r)}).$$

To complete the proof, we must show that (4.5.7) is equal to

$$(4.5.8) \quad \sum_{\substack{T_0 \in \text{Sym}_d(F) \\ y \in W^d \\ T_0 + T(y) = T}} \psi(y) \cdot \widehat{C}_{M_0}(T_0, v, \varphi_0).$$

If the  $(T_0, y)$  term in (4.5.8) is nonzero then, by Remark 4.2.5, there is an  $x \in V_0^d$  such that  $T(x) = T_0$ . The tuple  $x + y \in V^d$  then satisfies  $T(x + y) = T$ , and so its  $i^{\text{th}}$  component is isotropic for  $r < i \leq d$ . As we have assumed that  $V$  is anisotropic, we deduce that  $y$  has the form

$$y = (y_1, \dots, y_r, 0, \dots, 0).$$

It then follows from  $T_0 + T(y) = T$  that

$$T_0 = \begin{pmatrix} S_0 & \\ & 0_{d-r} \end{pmatrix}$$

for some  $S_0 \in \text{Sym}_r(F)$ , and we know from Proposition 4.2.4 that

$$\widehat{C}_{M_0}(T_0, v, \varphi_0) = \widehat{C}_{M_0}(S_0, v_0, \varphi_0^{(r)}) \cdot \widehat{C}_{M_0}(0_{d-r}, w, \varphi_0^{(d-r)}).$$

Thus in (4.5.8) we may replace the sum over  $T_0 \in \text{Sym}_d(F)$  with a sum over  $S_0 \in \text{Sym}_r(F)$ , replace the sum over  $y \in W^d$  with a sum over  $y \in W^r$ , and the result is (4.5.7).  $\square$

## REFERENCES

- [AGHMP17] Fabrizio Andreatta, Eyal Z. Goren, Benjamin Howard, and Keerthi Madapusi Pera, *Height pairings on orthogonal Shimura varieties*, *Compos. Math.* **153** (2017), no. 3, 474–534. MR 3705233
- [AM86] Reynir Axelsson and Jón Magnússon, *Complex analytic cones*, *Math. Ann.* **273** (1986), no. 4, 601–627. MR 826461

- [Bis90] Jean-Michel Bismut, *Superconnection currents and complex immersions*, Invent. Math. **99** (1990), no. 1, 59–113. MR 1029391
- [BGS90] J.-M. Bismut, H. Gillet, and C. Soulé, *Bott-Chern currents and complex immersions*, Duke Math. J. **60** (1990), no. 1, 255–284. MR 1047123
- [Bru02] Jan H. Bruinier, *Borchers products on  $O(2, l)$  and Chern classes of Heegner divisors*, Lecture Notes in Mathematics, vol. 1780, Springer-Verlag, Berlin, 2002. MR 1903920
- [BWR15] Jan H. Bruinier and Martin Westerholt-Raum, *Kudla’s modularity conjecture and formal Fourier-Jacobi series*, Forum Math. Pi **3** (2015), e7, 30. MR 3406827
- [BHY15] Jan H. Bruinier, Benjamin Howard, and Tonghai Yang, *Heights of Kudla-Rapoport divisors and derivatives of  $L$ -functions*, Invent. Math. **201** (2015), no. 1, 1–95. MR 3359049
- [Bur94] José Ignacio Burgos, *Green forms and their product*, Duke Math. J. **75** (1994), no. 3, 529–574. MR 1291696
- [Bur97] José Ignacio Burgos, *Arithmetic Chow rings and Deligne-Beilinson cohomology*, J. Algebraic Geom. **6** (1997), no. 2, 335–377. MR 1489119
- [BGKK07] J. I. Burgos Gil, J. Kramer, and U. Kühn, *Cohomological arithmetic Chow rings*, J. Inst. Math. Jussieu **6** (2007), no. 1, 1–172. MR 2285241
- [Del71] Pierre Deligne, *Travaux de Shimura*, Séminaire Bourbaki, 23ème année (1970/71), Exp. No. 389, 1971, pp. 123–165. Lecture Notes in Math., Vol. 244. MR 0498581
- [Dra69] Richard N. Draper, *Intersection theory in analytic geometry*, Math. Ann. **180** (1969), 175–204. MR 247134
- [Fis76] Gerd Fischer, *Complex analytic geometry*, Lecture Notes in Mathematics, Vol. 538, Springer-Verlag, Berlin-New York, 1976. MR 0430286
- [Ful84] William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 2, Springer-Verlag, Berlin, 1984. MR 732620
- [FH21] Jens Funke and Eric Hofmann, *The construction of Green currents and singular theta lifts for unitary groups*, Trans. Amer. Math. Soc. **374** (2021), no. 4, 2909–2947. MR 4223037
- [Gar18] Luis E. Garcia, *Superconnections, theta series, and period domains*, Adv. Math. **329** (2018), 555–589. MR 3783423
- [GS19] Luis E. Garcia and Siddarth Sankaran, *Green forms and the arithmetic Siegel-Weil formula*, Invent. Math. **215** (2019), no. 3, 863–975. MR 3935034
- [Gil81] Henri Gillet, *Riemann-Roch theorems for higher algebraic  $K$ -theory*, Adv. in Math. **40** (1981), no. 3, 203–289. MR 624666
- [GS90] Henri Gillet and Christophe Soulé, *Arithmetic intersection theory*, Inst. Hautes Études Sci. Publ. Math. (1990), no. 72, 93–174 (1991). MR 1087394 (92d:14016)
- [Hu99] Junda Hu, *Deformation to the normal bundle in arithmetic geometry*, ProQuest LLC, Ann Arbor, MI, 1999, Thesis (Ph.D.)—University of Illinois at Chicago. MR 2699571
- [Kin71] James R. King, *The currents defined by analytic varieties*, Acta Math. **127** (1971), no. 3-4, 185–220. MR 393550
- [Kol07] János Kollár, *Lectures on resolution of singularities*, Annals of Mathematics Studies, vol. 166, Princeton University Press, Princeton, NJ, 2007. MR 2289519
- [Kud04] Stephen S. Kudla, *Special cycles and derivatives of Eisenstein series, Heegner points and Rankin  $L$ -series*, Math. Sci. Res. Inst. Publ., vol. 49, Cambridge Univ. Press, Cambridge, 2004, pp. 243–270. MR 2083214

- [Kud21] ———, *Remarks on generating series for special cycles on orthogonal Shimura varieties*, *Algebra Number Theory* **15** (2021), no. 10, 2403–2447. MR 4377855
- [Mat89] Hideyuki Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid. MR 1011461
- [Sou92] Christophe Soulé, *Lectures on Arakelov geometry*, Cambridge Studies in Advanced Mathematics, vol. 33, Cambridge University Press, Cambridge, 1992, With the collaboration of D. Abramovich, J.-F. Burnol and J. Kramer. MR MR1208731 (94e:14031)
- [Wło09] Jarosław Włodarczyk, *Resolution of singularities of analytic spaces*, Proceedings of Gökova Geometry-Topology Conference 2008, Gökova Geometry/Topology Conference (GGT), Gökova, 2009, pp. 31–63. MR 2500573

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, 140 COMMONWEALTH AVE.,  
CHESTNUT HILL, MA 02467, USA  
*Email address:* `howardbe@bc.edu`