

## ON CLIQUE VERSION OF THE RANDIC INDEX

HOSSEIN TEIMOORI FAAL

ABSTRACT. In this paper, we first review the weighted-version of the handshaking lemma based on the idea of a weighted vertex-edge incidence matrix of a given graph  $G$ . Then, we obtain a generalized version of the handshaking lemma based on the concept of the clique value. We also define a generalized version of Randic index. More importantly, we prove an upper bound for the generalized Randic index of a graph  $G$ . We finally conclude the paper with some discussions about possible future works.

## 1. INTRODUCTION AND MOTIVATION

One of the important parameters of a simple, finite graph is the *degree* of a vertex. It simply reflects a topological property of a graph which is the size of an *open neighborhood* of a given vertex. Therefore, one interesting line of research in graph theory is to generalize this local concept and also seeking for its potential applications.

In [1], the author of this paper has suggested an extension this concept to the value of an edge  $e = \{u, v\}$  as the size of the intersection of the open neighborhoods of its end-vertices; that is  $val_G(e) = |N_G(u) \cap N_G(v)|$ . He also has used this idea in [1] to find a new upper bound for the number of edges with respect to the number of triangles in any  $K_4$ -free graph. The well-known *Randič* index  $R(G)$  of a graph  $G$  was introduced in 1975 by *Randič* [2]. More precisely, he has defined it by

$$(1.1) \quad R(G) := \sum_{\{u,v\} \in E(G)} \frac{1}{\sqrt{deg_G(u)deg_G(v)}}.$$

Indeed, in this paper we will call it the *vertex-version* of *Randič* index. It seems that this parameter is very useful in *mathematical chemistry* and has been extensively investigated in the literature (see [3] and the references therein). It is worth to mention the following two classical results in the context of *Randič* index.

**Theorem 1.1** (Bollobas and Erdos [4]). *For any connected graph  $G$  with  $n$  vertices,  $R(G) \geq \sqrt{n-1}$  with equality if and only if  $G \cong K_{1,n-1}$ .*

**Theorem 1.2** (Fajtlowicz [5]). *For a graph  $G$  with  $n$  vertices,  $R(G) \leq \frac{n}{2}$  with equality if and only if each component of  $G$  has at least two vertices and is regular.*

Our main goal in this paper is to obtain an extension of Theorem 1.2, based on the idea of the value of a clique.

## 2. BASIC DEFINITIONS AND NOTATIONS

We assume that our graphs are simple, finite and undirected. For a given graph  $G = (V, E)$  and a vertex  $v \in V(G)$ , the set of vertices adjacent to  $v$  is called the *open neighborhood* of  $v$  in  $G$  and will be denoted by  $N_G(v)$ .

The cardinality of  $N_G(v)$  is called the *degree* of  $v$  and is denoted by  $\deg_G(v)$ . A complete subgraph of  $G$  is called a *clique* of  $G$ . A clique on  $k$  vertices is called a  $k$ -clique. A clique on 3 vertices is called a triangle. We will denote the set of triangles of  $G$  by  $T(G)$ . We denote the set of all  $k$ -cliques of  $G$  by  $\Delta_k(G)$ . The number of  $k$ -cliques of  $G$  is denoted by  $c_k(G)$ . We also recall that the well-known *Cauchy-Schwartz* is the following inequality.

**Lemma 2.1.** [*Geometric-Harmonic Mean Inequality*] For any real sequences  $\{a_k\}_{k \geq 1}$ , we have

$$(2.1) \quad \sqrt[k]{a_1 a_2 \cdots a_k} \geq \frac{k}{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_k}},$$

with equality whenever  $a_1 = a_2 = \cdots = a_k$ .

In what follows, we quickly review the weighted-version of the well-known handshaking lemma and one of its consequences which is known as Mantel's theorem for triangle-free graphs.

The *weighted-version* of the well-known *handshaking lemma* can be read, as follows. From now on, we will denote the set of non-negative real numbers with  $\mathbb{R}^+$ .

**Lemma 2.2** (Weighted Handshaking Lemma [6]). Let  $G = (V, E)$  be a graph and  $f : V(G) \mapsto \mathbb{R}^+$  be a non-negative weight function. Then, we have

$$(2.2) \quad \sum_{v \in V(G)} f(v) \deg_G(v) = \sum_{e=uv \in E(G)} (f(u) + f(v)).$$

In particular, we have

$$(2.3) \quad \sum_{v \in V(G)} \deg_G^2(v) = \sum_{e=uv \in E(G)} (\deg_G(u) + \deg_G(v)).$$

## 3. A VERTEX-VERSION OF RANDIC INDEX

In chemical graph theory literature, the *branching index* of a given graph  $G$  is known as the *Randič* index of the graph  $G$  denoted by  $R(G)$ . Here, we

will denote it by  $R_v(G)$  and call it *vertex-version Randić index* of  $G$ . It is defined, as follows

$$(3.1) \quad R_v(G) = \sum_{e=uv \in E(G)} \frac{1}{\sqrt{\deg_G(u) \deg_G(v)}}.$$

In [6], the following result is proved by a simple argument based on the *weighted version of handshaking lemma* and *geometric-harmonic mean inequality*.

**Theorem 3.1** ([5]). *For a graph  $G$  of order  $n$ ,*

$$R_v(G) \leq \frac{n}{2},$$

*with equality if and only if every component of  $G$  is regular and  $G$  has no isolated vertices.*

For the sake of completeness, here we also provide a short proof of Theorem 3.1 based on the reference [6].

*Proof.* First, we note that by defining the function  $f$  in the equation (2.2) by

$$(f(v) = \begin{cases} \frac{1}{\deg_G(v)} & \text{if } \deg_G(v) > 0, \\ 1 & \text{if } \deg_G(v) = 0, \end{cases}$$

we obtain

$$(3.2) \quad \sum_{uv \in E(G)} \left( \frac{1}{\deg_G(u)} + \frac{1}{\deg_G(v)} \right) = n - n_0,$$

where  $n_0$  denotes the number of isolated vertices in  $G$ . Now considering the *geometric-harmonic inequality* for  $k = 2$ , Lemma 2.1, we have

$$(3.3) \quad \begin{aligned} R_v(G) &= \sum_{e=uv \in E(G)} \frac{1}{\sqrt{\deg_G(u) \deg_G(v)}} \\ &\leq \sum_{uv \in E(G)} \frac{1}{2} \left( \frac{1}{\deg_G(u)} + \frac{1}{\deg_G(v)} \right) \\ &= \frac{n - n_0}{2} \leq \frac{n}{2}, \end{aligned}$$

as required. □

## 4. AN EDGE-VERSION OF RANDIČ INDEX

In this section, we aim to obtain an edge version of the upper bound for the *Randič* index, based on the new concept of the value of an edge.

**Definition 4.1.** Let  $G = (V, E)$  be a graph and  $e = uv$  be an edge of  $G$ . Then, we define the *edge value* of  $e$ , denoted by  $val_G(e)$ , as follows

$$(4.1) \quad val_G(e) = |N_G(e)| = |N_G(u) \cap N_G(v)|.$$

Here,  $N_G(e)$  denotes the set of common neighbors of end vertices of the edge  $e$ .

Next, we generalize the weighted handshaking lemma for values of edges of a given graph.

**Lemma 4.1.** [*Weighted Edge Handshaking Lemma*] Let  $G = (V, E)$  be a graph and  $g : E(G) \mapsto \mathbb{R}^+$  be a non-negative weight function. Then, we have

$$(4.2) \quad \sum_{e \in E(G)} g(e)val_G(e) = \sum_{\delta=e_1e_2e_3 \in T(G)} (g(e_1) + g(e_2) + g(e_3)).$$

In particular, we have

$$(4.3) \quad \sum_{e \in E(G)} val_G^2(e) = \sum_{\delta=e_1e_2e_3 \in T(G)} (val_G(e_1) + val_G(e_2) + val_G(e_3)).$$

As an immediate consequence of the above lemma, we have the following interesting result. Recall that an edge  $e \in E(G)$  is called *isolated* if we have  $val_G(e) = 0$ .

**Corollary 4.2.** For any graph  $G$  with  $m$  edges, we have

$$(4.4) \quad \sum_{\delta=e_1e_2e_3 \in T(G)} \left( \frac{1}{val_G(e_1)} + \frac{1}{val_G(e_2)} + \frac{1}{val_G(e_3)} \right) = m - m_0,$$

in which  $m_0$  is the number of isolated edges.

Next, we give a generalization of the result in [5]. To do so, we first need to give a generalization of the concept of *Randič* index.

**Definition 4.2.** For a given graph  $G = (V, E)$ , the edge-version of the *Randič* index, denoted by  $R_e(G)$ , is defined as

$$(4.5) \quad R_e(G) := \sum_{\delta=e_1e_2e_3 \in T(G)} \frac{1}{\sqrt{val_G(e_1)val_G(e_2)val_G(e_3)}}$$

From now on, we will call a graph *edge-regular* if  $val_G(e)$  is the same for all edges of  $G$ .

**Theorem 4.3.** *For a graph  $G$  of size  $m$ , we have*

$$R_e(G) \leq \frac{m}{3},$$

*with equality if and only if every component of  $G$  is edge-regular and  $G$  has no isolated edges.*

*Proof.* Considering Corollary 4.2 and arithmetic-geometric inequality, we have

$$\begin{aligned} (4.6) \quad R_e(G) &:= \sum_{\delta=e_1e_2e_3 \in T(G)} \frac{1}{\sqrt{\text{val}_G(e_1)\text{val}_G(e_2)\text{val}_G(e_3)}} \\ &\leq \frac{1}{3} \sum_{\delta=e_1e_2e_3 \in T(G)} \left( \frac{1}{\text{val}_G(e_1)} + \frac{1}{\text{val}_G(e_2)} + \frac{1}{\text{val}_G(e_3)} \right) \\ &= \frac{1}{3}(m - m_0) \leq \frac{m}{3}. \end{aligned}$$

□

## 5. THE CLIQUE HANDSHAKING LEMMA

In this section, we attempt to find a more generalized version of *Randič* index which we call it *the generalized Randić* index. To this end, we first need to present an extension of the definition of the degree of vertex to the value of any clique of higher order.

**Definition 5.1.** Let  $G = (V, E)$  be a graph and  $q_k \in \Delta_k(G)$  be a  $k$ -clique in  $G$ . Then, we define the value of the clique  $q_k$  with the vertex set  $V(q_k) = v_{i_1} \cdots v_{i_k}$  denoted by  $\text{val}_G(q_k)$ , as follows

$$(5.1) \quad \text{val}_G(q) = \left| \bigcap_{i=1}^k N_G(v_{j_i}) \right|.$$

We will also call an  $k$ -clique  $q_k$  with  $\text{val}_G(q_k) = 0$ , an *isolated* clique of  $G$ .

Note that any  $k$ -clique  $q_k = v_{i_1} \cdots v_{i_k} \in \Delta_k(G)$  in  $G$  can also be represented (uniquely) by  $q_k = q_{k-1,1} \cdots q_{k-1,k}$  where for each  $i = 1, \dots, k$  the symbol  $q_{k-1,i}$  denotes a  $(k-1)$ -clique subgraph of  $q_k$ . We will use this fact in our next key lemma.

**Lemma 5.1** (Weighted Clique Handshaking Lemma). *Let  $G = (V, E)$  be a graph and let  $h : \Delta_k(G) \mapsto \mathbb{R}^+$  ( $k \geq 2$ ) be a weight function. Then, we have*

$$(5.2) \quad \sum_{q_k \in \Delta_k(G)} h(q_k) \text{val}_G(q_k) = \sum_{q_{k+1}=q_{k,1} \cdots q_{k,k+1} \in \Delta_{k+1}(G)} \left( h(q_{k,1}) + \dots + h(q_{k,k+1}) \right).$$

*In particular, we have*

$$(5.3) \quad \sum_{q_k \in \Delta_k(G)} val_G^2(q_k) = \sum_{q_{k+1}=q_{k,1} \cdots q_{k,k+1} \in \Delta_{k+1}(G)} \left( val_G(q_{k,1}) + \cdots + val_G(q_{k,k+1}) \right).$$

*Proof.* We proceed by defining the weighted *subclique-superclique* matrix  $I_{f,k}(G)$  of order  $k$ , as follows

$$(I_{f,k}(G))_{q_k, q_{k+1}} = \begin{cases} h(q_k) & \text{if } q_k \text{ is a subgraph of } q_{k+1}, \\ 0 & \text{otherwise.} \end{cases}$$

Next, we note that in the matrix  $I_{f,k}(G)$  each row corresponding to the clique  $q_k$  has  $val_G(q_k)$  non-zero entries. Hence, the resulting row-sum equals to  $h(q_k)val_G(q_k)$ . On the other hand, each column corresponding to the clique  $q_{k+1} = q_{k,1} \cdots q_{k,k+1}$  has the column-sum  $h(q_{k,1}) + \cdots + h(q_{k,k+1})$ . Thus, by summing over all rows and columns and equating them we get the desired result.  $\square$

## 6. THE GENERALIZED RANDIČ INDEX

In this last section, we obtain a more generalized version of *Randič* index which we call it the *generalized Randič index*.

Next, we define the *generalized Randič index* of a graph  $G$  or a *clique-version* of the *Randič* index based on the new concept of the *clique value* in graph theory.

**Definition 6.1.** Let  $G = (V, E)$  be a graph. Then, the *generalized Randič index* of  $G$ , denoted by  $R_{cliq}(G)$ , is defined by

$$(6.1) \quad R_{cliq}(G; k) := \sum_{q_{k+1}=q_{k,1} \cdots q_{k,k+1} \in \Delta_{k+1}(G)} \frac{1}{\sqrt{\prod_{j=1}^{k+1} val_G(q_{k,j})}}, \quad (k \geq 1)$$

We first need the following key result.

**Proposition 6.1.** *Let  $G = (V, E)$  be a graph. Then, we have*

$$(6.2) \quad \sum_{q_{k+1}=q_{k,1} \cdots q_{k,k+1} \in \Delta_{k+1}} \left( \frac{1}{val_G(q_{k,1})} + \cdots + \frac{1}{val_G(q_{k,k+1})} \right) = c_k(G) - c_{k,0}(G),$$

in which  $c_{k,0}(G)$  is the number of isolated  $k$ -cliques of  $G$ .

Now, we are at the position to state the main result of this paper which a generalized version of the result in [5]. Form now on, a graph in which all values of  $val_G(q_k)$  are the same for  $k$ -cliques of  $G$  is called *k-clique regular* graph.

**Theorem 6.2.** *Let  $G$  be a graph. Then, we have*

$$(6.3) \quad R_{clq}(G; k) \leq \frac{c_k(G)}{k+1}.$$

*The equality holds if and only if each component of  $G$  is a  $k$ -clique regular graphs.*

*Proof.* The proof is straight forward based on Proposition 6.1 and the arithmetic-geometric mean inequality 2.1. □

#### REFERENCES

- [1] H. Teimoori, *ON CLIQUE VALUES IDENTITIES AND MANTEL-TYPE THEOREMS*, Transactions on Combinatorics, 9 (2020) pp. 139–146.
- [2] M. Randic, *On characterization of molecular branching*, J. Am. Chem. Soc., 97 (1975), pp. 6609–6615.
- [3] X. Li and I. Gutman, *Mathematical Aspects of Randić-Type Molecular Structure Descriptors* University of Kragujevac, Kragujevac, (2006).
- [4] B. Bollobás and P. Erdos, *Graphs of extremal weights*, Ars Combin., 50 (1998), pp. 225–233.
- [5] S. Fajtlowicz, *Written on the Wall, a list of conjectures of Graffiti*, Available from the author.
- [6] B. Wu, *The weighted version of the handshaking lemma with an application*, J. Inequalities and Applications, 1 (2014), pp. 225–233.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE,, ALLAMEH TABATABA'I UNIVERSITY, TEHRAN, IRAN