

REILLY-TYPE INEQUALITIES FOR SUBMANIFOLDS IN CARTAN-HADAMARD MANIFOLDS

HANG CHEN AND XUDONG GUI

ABSTRACT. Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional complete simply-connected Riemannian manifold N , where the sectional curvature of N is bounded above by δ . When $\delta < 0$, inspired by Niu-Xu (arXiv:2106.01912), we give new upper bounds for the first nonzero eigenvalues of the p -Laplacian and the L_T operator, respectively. These generalize Niu-Xu's work for the Laplacian (arXiv:2106.01912) and improve the estimates due to Chen (Nonlinear Anal.,196, 111833, 2020) for the p -Laplacian and Grosjean (Hokkaido Math. J., 33(2), 319–339, 2004) for the L_T operator, respectively. We also obtain several Reilly-type inequalities for the weighted manifolds and some boundary value problems.

1. INTRODUCTION

In geometry of submanifolds, Reilly-type inequality is an important estimate for the first eigenvalue of Laplacian, which gives the upper bound in terms of the mean curvature. It originated from Reilly's work [19] in 1977, where closed (orientable) submanifolds in the Euclidean space were considered and a special case says

$$\lambda_1^\Delta \leq \frac{m}{\text{vol}(M)} \int_M |\mathbf{H}|^2. \quad (1.1)$$

Here m and \mathbf{H} are the dimension and the mean curvature vector of the submanifold respectively, and λ_1^Δ represents the first non-zero eigenvalue of the Laplacian on the submanifold. Later, Heintze [14] considered a more general case that the sectional curvature of the ambient manifold N is bounded above by a constant δ , and showed that

$$\lambda_1^\Delta \leq \begin{cases} \frac{m}{\text{vol}(M)} \int_M (\delta + |\mathbf{H}|^2) & \text{for } \delta \geq 0, \\ m(\delta + \max |\mathbf{H}|^2) & \text{for } \delta < 0, \end{cases} \quad (1.2)$$

$$(1.3)$$

here it is required that N is simply-connected when $\delta \leq 0$ and M lies in a convex ball of N of the radius $\leq \pi/4\sqrt{\delta}$ when $\delta > 0$. Notice that the approach of Heintze cannot give the upper bound involving L^2 -norm the the mean curvature when $\delta < 0$ while there is a restriction on the radius of the convex ball when $\delta > 0$. However, when N is the space form of constant curvature δ , El Soufi and Ilias [11] removed the restrictions and gave a sharp upper bound

$$\lambda_1^\Delta \leq \frac{m}{\text{vol}(M)} \int_M (\delta + |\mathbf{H}|^2) \quad (1.4)$$

by using a conformally mapping method whenever $\delta > 0$ and $\delta < 0$.

2020 *Mathematics Subject Classification.* 58C40, 53C42, 35P15.

Key words and phrases. Reilly-type inequalities, p -Laplacian, L_T operator, eigenvalue estimates, Cartan-Hadamard manifolds.

Chen supported by Natural Science Foundation of Shaanxi Province Grant No. 2020JQ-101.

Very recently, Niu-Xu did nice work which says the estimate (1.3) can be improved to

$$\lambda_1^\Delta \leq \frac{m}{\text{vol}(M)} \int_M (\delta + |\mathbf{H}|^2) \quad (1.5)$$

when $\delta < 0$ and $m \geq 2$ ([18, Theorem 1.1]).

Note that the original Reilly's inequalities in [19] actually involve not only the mean curvature, but also the higher order mean curvatures. More generally, given a symmetric $(1, 1)$ -tensor $T = (T_{ij})$ on M , we can define a normal vector field H_T associated with T by

$$H_T = \sum_{i=1}^m A(Te_i, e_i) = \sum_{\substack{m+1 \leq \alpha \leq n \\ 1 \leq i, j \leq m}} h_{ij}^\alpha T_{ij} e_\alpha,$$

where $\{e_1, \dots, e_n\}$ is a local orthonormal frame of N such that $\{e_1, \dots, e_m\}$ are tangent to M and $\{e_{m+1}, \dots, e_n\}$ are normal to M , and $A = (h_{ij}^\alpha)$ the second fundamental form of M in N . Actually, H_T can be viewed as a natural generalization of the mean curvature and the higher order mean curvature, and has been involved a lot of results ([2, 5, 13]).

To generalize the Reilly-type inequalities, a common approach is considering the other operators instead of the Laplacian. Usually, there are two classes of the elliptic operators as alternatives: one is the p -Laplacian and the other one is the L_T operator. We states them as follows respectively.

1.1. p -Laplacian. The p -Laplacian ($p > 1$) on a compact M is defined by

$$\Delta_p u = \text{div}(|\nabla u|^{p-2} \nabla u).$$

It is a second order quasilinear elliptic operator for $p \neq 2$ and can be viewed as a generalization of the Laplacian (corresponding to $p = 2$). A real number λ is called a Dirichlet (or Neumann) eigenvalue if there exists a non-zero function u satisfying the following equation with Dirichlet boundary condition $u \equiv 0$ on ∂M (or Neumann boundary condition $\nabla_\nu u \equiv 0$ on ∂M):

$$\Delta_p u = -\lambda |u|^{p-2} u \quad \text{on } M,$$

where ν is the outward normal on ∂M . So far, many estimates for the first eigenvalue of Laplacian in Riemannian geometry have been generalized to p -Laplacian (e.g., [4] and the references therein).

The inequality mentioned above have been extended from Laplacian to p -Laplacian. For example, Du-Mao [10] generalized (1.1) in the Euclidean space case; Wei and the first author [6] generalized (1.4) by the conformally mapping method in the space form case.

Recently, the first author generalized (1.2) and (1.3) and proved

Theorem 1.1 (cf. [4, Theorem 1.3]). *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional Riemannian manifold N of the sectional curvature $K_N \leq \delta$. If $\delta \leq 0$, we assume that N is simply-connected; if $\delta > 0$, we assume that M is contained in a convex ball of radius $\leq \pi/4\sqrt{\delta}$; if $\delta < 0$, we assume that M is contained in a convex ball of radius $\leq \frac{1}{2\sqrt{-\delta}} \text{arsinh}(\sqrt{-\delta})$. Let T be a symmetric, positive definite and divergence-free $(1, 1)$ -tensor on M .*

Then the first non-zero eigenvalue $\lambda_{1,p}$ of the p -Laplacian ($p > 1$) on M satisfies:

(1) For $\delta = 0$,

$$\lambda_{1,p} \left(\int_M \text{tr } T \right)^p \leq n^{\frac{|p-2|}{2}} m^{p/2} \text{vol}(M) \left(\int_M |H_T|^{\frac{p}{p-1}} \right)^{p-1}, \quad \text{for } p > 1.$$

In particular, when $T = I$ the identity operator, we have

$$\lambda_{1,p} \leq n^{\frac{|p-2|}{2}} \frac{m^{p/2}}{(\text{vol}(M))^{p-1}} \left(\int_M |\mathbf{H}|^{\frac{p}{p-1}} \right)^{p-1}, \quad \text{for } p > 1.$$

(2) For $\delta > 0$,

$$\lambda_{1,p} \leq (n+1)^{\frac{|p-2|}{2}} m^{p/2} \delta^{p/2} \left(\frac{1}{\text{vol}(M)} \int_M (1 + |\mathbf{H}/\sqrt{\delta}|^2) \right)^{\max\{1, p/2\}};$$

(3) For $\delta < 0$,

$$\lambda_{1,p} \leq n^{\frac{|p-2|}{2}} m (\delta + \max |\mathbf{H}|^2) \quad \text{for } 1 < p \leq 2.$$

The estimate is not good enough in the case of $\delta < 0$ for at least two reasons: (i) there is a restriction on the radius of the convex ball for $1 < p < 2$, but it is not necessary for $p = 2$ (cf. Heintze's upper bound (1.3)); (ii) the estimate is just valid for $1 < p < 2$ and no information is given for $p > 2$.

Inspired by Niu-Xu [18], we extend (1.5) to p -Laplacian and give a new upper bound of $\lambda_{1,p}$ without previous restrictions for $\delta < 0$. Precisely, we prove the following theorem.

Theorem 1.2. *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional complete simply-connected Riemannian manifold N of the sectional curvature $K_N \leq \delta < 0$. Then the first non-zero eigenvalue $\lambda_{1,p}$ of p -Laplacian on M satisfies*

$$\lambda_{1,p} \leq \begin{cases} n^{\frac{|p-2|}{2}} \left(\frac{m}{\text{vol}(M)} \right)^{p/2} \left(\int_M |\mathbf{H}|^2 \right)^{p/2-1} \int_M (\delta + |\mathbf{H}|^2), & \text{for } p \geq 2; \\ n^{\frac{|p-2|}{2}} m^{p/2} \frac{(-\delta)^{p/2-1}}{\text{vol}(M)} \int_M \left(\frac{p}{2} \delta + |\mathbf{H}|^2 \right), & \text{for } 1 < p \leq 2. \end{cases} \quad (1.6)$$

$$\lambda_{1,p} \leq \begin{cases} n^{\frac{|p-2|}{2}} \left(\frac{m}{\text{vol}(M)} \right)^{p/2} \left(\int_M |\mathbf{H}|^2 \right)^{p/2-1} \int_M (\delta + |\mathbf{H}|^2), & \text{for } p \geq 2; \\ n^{\frac{|p-2|}{2}} m^{p/2} \frac{(-\delta)^{p/2-1}}{\text{vol}(M)} \int_M \left(\frac{p}{2} \delta + |\mathbf{H}|^2 \right), & \text{for } 1 < p \leq 2. \end{cases} \quad (1.7)$$

More generally, we can obtain the upper bound involving H_T .

Theorem 1.3. *Assumptions as in Theorem 1.2. Let T be a symmetric, positive definite and divergence-free $(1,1)$ -tensor on M . Suppose the one of the followings holds:*

- (a) $(\text{tr } T)I - 2T \geq 0$, i.e., positive semidefinite;
- (b) M is contained in a convex ball of radius $\leq \frac{1}{2\sqrt{-\delta}} \text{arcosh } \sqrt{2}$.

Then the first non-zero eigenvalue $\lambda_{1,p}$ of p -Laplacian on M satisfies

$$\lambda_{1,p} \leq \begin{cases} n^{\frac{|p-2|}{2}} m^{p/2} Q(T)^{p/2-1} (\delta + Q(T)), & \text{for } p \geq 2; \\ n^{\frac{|p-2|}{2}} m^{p/2} (-\delta)^{p/2-1} \left(\frac{p}{2} \delta + Q(T) \right), & \text{for } 1 < p \leq 2, \end{cases} \quad (1.8)$$

$$\lambda_{1,p} \leq \begin{cases} n^{\frac{|p-2|}{2}} m^{p/2} Q(T)^{p/2-1} (\delta + Q(T)), & \text{for } p \geq 2; \\ n^{\frac{|p-2|}{2}} m^{p/2} (-\delta)^{p/2-1} \left(\frac{p}{2} \delta + Q(T) \right), & \text{for } 1 < p \leq 2, \end{cases} \quad (1.9)$$

where

$$Q(T) = \frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|\mathbf{H}_T|^2}{\text{tr } T}.$$

By taking $p = 2$, we immediately obtain the following

Corollary 1.4. *Assumptions as in Theorem 1.3. We have*

$$\lambda_1^\Delta \leq m \left(\delta + \frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|\mathbf{H}_T|^2}{\text{tr } T} \right). \quad (1.10)$$

Remark 1.5. (1) The condition $(\operatorname{tr} T)I - 2T \geq 0$ is also required in [5, Theorem 1.1].

(2) When $m \geq 2$, $(\operatorname{tr} T)I - 2T \geq 0$ always holds for $T = I$ the identity operator, and then Theorem 1.3 recovers Theorem 1.2; furthermore, this recovers [18, Theorem 1.1] when $p = 2$. However, when $m = 1$, even for the Laplacian operator, $(\operatorname{tr} T)I - 2T \geq 0$ fails, while an analogous estimate involving the curvature of the curve also fails unless “smallness” assumed (see [18] for details).

1.2. L_T operator. Consider a symmetric $(1, 1)$ -tensor $T = (T_{ij})$ on M as previously mentioned. If we additionally assume that T is positive definite and divergence-free, then we can define a self-adjoint elliptic operator L_T associated to T by

$$L_T u = \operatorname{div}(T \nabla u).$$

The eigenvalue equation is given by $L_T u = -\lambda u$. It is also a generalization of the Laplacian since $L_T = \Delta$ when $T = I$.

The L_T operator has been intensively studied. Regarding the Reilly-type inequality, Grosjean proved that

Theorem 1.6 ([13, Theorem 1]). *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional Riemannian manifold N of the sectional curvature $K_N \leq \delta$. If $\delta \leq 0$, we assume that N is simply-connected; if $\delta > 0$, we assume that M is contained in a convex ball of radius $\leq \pi/4\sqrt{\delta}$. Let T be a symmetric, positive definite and divergence-free $(1, 1)$ -tensor on M . Then the first non-zero eigenvalue $\lambda_{1,T}$ of L_T on M satisfies*

$$\lambda_{1,T} \leq \frac{\sup_M |H_T|^2 + \sup_M \delta (\operatorname{tr} T)^2}{\inf_M \operatorname{tr} T}. \quad (1.11)$$

If the equality holds, then M is contained in a geodesic sphere.

When N is the space form with constant curvature δ , the first author and X. Wang [5] showed that the upper bound in (1.11) can be improved by the L_2 -norm of H_T , precisely,

$$\lambda_{1,T} \leq \frac{1}{\operatorname{vol}(M)} \int_M \left(\delta \operatorname{tr} T + \frac{|H_T|^2}{\operatorname{tr} T} \right). \quad (1.12)$$

We remark that in the case, there is no restriction on the radius of M , but it is required that $(\operatorname{tr} T)I - 2T \geq 0$, i.e., positive semidefinite. The sufficient and necessary conditions for the equality holding are also given.

Motivated by Niu-Xu [18] again, we can also improve the estimate (1.11) for $\delta < 0$. Precisely, we prove the following theorem.

Theorem 1.7. *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional complete simply-connected Riemannian manifold N of the sectional curvature $K_N \leq \delta < 0$. Let T be a symmetric, positive definite and divergence-free $(1, 1)$ -tensor on M . Suppose the one of the followings holds:*

- (a) $(\operatorname{tr} T)I - 2T \geq 0$, i.e., positive semidefinite;
- (b) M is contained in a convex ball of radius $\leq \frac{1}{2\sqrt{-\delta}} \operatorname{arcosh} \sqrt{2}$.

Then we have

$$\lambda_{1,T} \leq \frac{\sup_M \operatorname{tr} T}{\int_M \operatorname{tr} T} \int_M \left(\delta \operatorname{tr} T + \frac{|H_T|^2}{\operatorname{tr} T} \right). \quad (1.13)$$

There are other Reilly-type inequalities in various situations, for example, when the ambient Riemannian manifold (N, g_N) is endowed with a density e^{-f} for certain $f \in C^\infty(N)$, one can consider the weighted L_T operator, denoted by $L_{T,f}$; when the submanifold has a non-empty boundary, one can consider the first (non-zero) eigenvalues of the Steklov-type problems (see [7, 9, 15, 16, 20, 21, 23]). We also obtain some new estimates and list them in the last section.

This paper is organized as follows. In Sect. 2, we list some notations and show some lemmas and propositions, some of them are extensions from the mean curvature to H_T . In Sect. 3 and Sect. 4, we give the proofs for the p -Laplacian and L_T , respectively. To estimate the upper bound of the first (non-zero) eigenvalues, a starting point is the Rayleigh-type variational characterization (cf. Eqs. (3.2) and (4.3)). One of key ingredients is to find suitable test functions (Sect. 2.2), and the other one is an estimate of L^2 lower bound of H_T , which extends Niu-Xu's estimate for the mean curvature (Sect. 2.3). In Sect. 5, we give some Reilly-type estimates when the ambient manifold is a weighted manifold.

Acknowledgment: The first author was supported by Natural Science Foundation of Shaanxi Province Grant No. 2020JQ-101.

2. PRELIMINARIES

In this section, we introduce some notations and review some lemmas.

2.1. Some estimates for radial vector fields. Let M be an m -dimensional submanifold of an n -dimensional Riemannian manifold N whose sectional curvature $K_N \leq \delta < 0$. We denote the Levi-Civita connection on M and N by ∇ and ∇^N respectively. For any $q \in N$, we consider its normal coordinates (x_1, \dots, x_n) on N centered at some fixed point $q_0 \in N$.

Let $\text{sh}_\delta(y) = \frac{1}{\sqrt{-\delta}} \sinh(\sqrt{-\delta}y)$, $\text{ch}_\delta(y) = \text{sh}_\delta'(y)$ and $\text{th}_\delta(y) = \text{sh}_\delta(y)/\text{ch}_\delta(y)$. It is easy to check that $\text{ch}_\delta^2 + \delta \text{sh}_\delta^2 = 1$ and $1/\text{ch}_\delta^2 = 1 + \delta \text{th}_\delta^2$.

Consider the vector field $Z = \phi(r)\nabla^N r$, where $r = r(q)$ is the distance function from q_0 to q and $\phi(r)$ be a positive function with $\phi(r) \sim r$ as $r \rightarrow 0$. Then the coordinates of Z in the normal local frame are $\left(\frac{\phi(r)}{r}x_i\right)_{1 \leq i \leq n}$. By using basic properties of the exponential map and the comparison for Jacobi fields, we can prove the following lemma.

Lemma 2.1 (cf. [13, 18]). *Let T be a symmetric, positive definite and divergence-free $(1, 1)$ -tensor on M , and Z^\top be the tangential projection of $Z = \phi(r)\nabla^N r$ to M . Then we have*

(1) $\sum_{i=1}^n \langle T\nabla\left(\frac{\phi(r)}{r}x_i\right), \nabla\left(\frac{\phi(r)}{r}x_i\right) \rangle \leq (\text{tr } T) \frac{\phi^2}{\text{sh}_\delta^2} + \left((\phi')^2 - \frac{\phi^2}{\text{sh}_\delta^2}\right) \langle T\nabla r, \nabla r \rangle$, and equality holds if N has constant sectional curvature δ .

(2) $\text{div}_M(TZ^\top) \geq (\text{tr } T) \frac{\phi}{\text{th}_\delta} + \left(\phi' - \frac{\phi}{\text{th}_\delta}\right) \langle T\nabla r, \nabla r \rangle + \langle Z, H_T \rangle$.

Proof. When $\phi = \text{sh}_\delta$, Items (1) and (2) go back to Lemma 1 and Lemma 2 in [13] respectively, and the proof of this lemma follows from the same steps with similar calculations in [13]. Here we give a quick proof by directly using the results in [13].

Since

$$\nabla\left(\frac{\phi(r)}{r}x_i\right) = \left(\frac{\phi}{r}\right)'x_i\nabla r + \frac{\phi}{r}\nabla x_i, \quad \sum_{i=1}^n (x_i)^2 = r^2, \quad \sum_{i=1}^n x_i\nabla x_i = r\nabla r,$$

we have

$$\begin{aligned}
\sum_{i=1}^n \langle T \nabla \left(\frac{\phi(r)}{r} x_i \right), \nabla \left(\frac{\phi(r)}{r} x_i \right) \rangle &= \sum_{i=1}^n \frac{\phi^2}{r^2} \langle T \nabla x_i, \nabla x_i \rangle + \left(\left(\frac{\phi}{r} \right)' r^2 + 2\phi \left(\frac{\phi}{r} \right)' \right) |\sqrt{T} \nabla r|^2 \\
&= \sum_{i=1}^n \frac{\phi^2}{r^2} \langle T \nabla x_i, \nabla x_i \rangle + \left((\phi')^2 - \frac{\phi^2}{r^2} \right) |\sqrt{T} \nabla r|^2 \\
&= (\phi')^2 |\sqrt{T} \nabla r|^2 - \frac{\phi^2}{\text{sh}_\delta^2} |\sqrt{T} \nabla r|^2 \\
&\quad + \frac{\phi^2}{\text{sh}_\delta^2} \left[\frac{\text{sh}_\delta^2}{r^2} \sum_{i=1}^n \langle T \nabla x_i, \nabla x_i \rangle + \left(1 - \frac{\text{sh}_\delta^2}{r^2} \right) |\sqrt{T} \nabla r|^2 \right] \\
&\leq (\phi')^2 |\sqrt{T} \nabla r|^2 - \frac{\phi^2}{\text{sh}_\delta^2} |\sqrt{T} \nabla r|^2 + \frac{\phi^2}{\text{sh}_\delta^2} (\text{tr } T),
\end{aligned}$$

where we used [13, Eqs. (5) and (6)] in the last inequality.

$$\begin{aligned}
\text{div}_M(TZ^\top) &= \text{div}_M \left(\frac{\phi}{\text{sh}_\delta} T(\text{sh}_\delta \nabla r) \right) \\
&= \frac{\phi}{\text{sh}_\delta} \text{div}_M(T(\text{sh}_\delta \nabla r)) + \left(\frac{\phi}{\text{sh}_\delta} \right)' \langle \nabla r, T(\text{sh}_\delta \nabla r) \rangle \\
&\geq \frac{\phi}{\text{sh}_\delta} \left((\text{tr } T) \text{ch}_\delta + \langle \text{sh}_\delta \nabla^N r, H_T \rangle \right) + \frac{\phi' \text{sh}_\delta - \phi \text{ch}_\delta}{\text{sh}_\delta} \langle \nabla r, T \nabla r \rangle \\
&= (\text{tr } T) \frac{\phi}{\text{th}_\delta} + \left(\phi' - \frac{\phi}{\text{th}_\delta} \right) \langle T \nabla r, \nabla r \rangle + \langle Z, H_T \rangle,
\end{aligned}$$

where we used [13, Lemma 2] in the inequality. \square

Remark 2.2. When $T = I$, Lemma 2.1 recovers [18, Lemma 2.3].

2.2. Test functions. The next lemmas will provide suitable test functions for the upper bound estimates of the first eigenvalue, and the basic idea can be found in [3, 14].

Lemma 2.3. *Under the same assumptions as in Theorem 1.2 or Theorem 1.3, we have for $p > 1$, there exists a point $q_0 \in N$ such that*

$$\int_M \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^{p-2} \frac{\text{th}_\delta(r)}{r} x_i dv_M = 0 \quad (i = 1, \dots, n). \quad (2.1)$$

Here (x_1, \dots, x_n) is the normal coordinate around q_0 , and $r(q) = d(q_0, q)$ is the geodesic distance from q_0 to q on N .

Proof. We can assume M lies in a convex ball $B \subset N$. For each $q \in N$, we define a vector $Y_q \in T_q N$ under the normal coordinate system around q as follows:

$$Y_q := \left(\int_M \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^{p-2} \frac{\text{th}_\delta(r)}{r} x_i dv_M, \dots, \int_M \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^{p-2} \frac{\text{th}_\delta(r)}{r} x_n dv_M \right). \quad (2.2)$$

Hence, we obtain a vector field Y in a neighborhood of B which points at the boundary into the interior of B .

Now suppose $Y(q) \neq 0$ for all $q \in B$, which means $\text{Ind}(Y) = 0$. Since Y is a continuous vector field on B and $Y(q)$ points inside for all $q \in \partial B$, we have (cf. [12, 17])

$$\text{Ind}(Y) + \chi(\partial B) = \chi(B),$$

so $\text{Ind}(Y) = 1 - (1 + (-1)^{n-1}) = (-1)^n \neq 0$, a contradiction. The proof is complete. \square

Remark 2.4. From the proof we see that $q_0 \in B$. If the radius of $B \leq \frac{1}{2\sqrt{-\delta}} \operatorname{arcosh} \sqrt{2}$, then M lies in a ball of radius $\leq \frac{1}{\sqrt{-\delta}} \operatorname{arcosh} \sqrt{2}$ centered at q_0 , which implies $\operatorname{ch}_\delta^2(r) \leq 2$ on M .

2.3. L^2 lower bound of H_T . Recall the following key estimate used to prove (1.5).

Proposition 2.5 ([18, Proposition 2.1]). *Under the same assumptions as in Theorem 1.2, we have*

$$\int_M |\mathbf{H}|^2 \int_M \operatorname{th}_\delta^2(r) \geq (\operatorname{vol}(M))^2. \quad (2.3)$$

We can extend Proposition 2.5 to H_T as follows.

Proposition 2.6. *Under the same assumptions as in Theorem 1.3, we have*

$$\int_M \frac{|H_T|^2}{\operatorname{tr} T} \int_M \operatorname{tr} T \operatorname{th}_\delta^2(r) \geq \left(\int_M \operatorname{tr} T \right)^2. \quad (2.4)$$

Proof. By taking $\phi(r) = \operatorname{th}_\delta(r)$ in Lemma 2.1, we have $Z = \operatorname{th}_\delta(r) \nabla^N r$ and

$$\operatorname{div}_M(TZ^\top) \geq \operatorname{tr} T + \delta |\sqrt{T}Z^\top|^2 + \langle Z^\perp, H_T \rangle. \quad (2.5)$$

Integrating (2.5) over M and using the divergence theorem, the Cauchy-Schwarz inequality and the Hölder inequality, we have

$$\|\sqrt{\operatorname{tr} T}Z^\perp\|_2 \left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2 \geq \int_M -\langle Z^\perp, H_T \rangle \geq \int_M \operatorname{tr} T + \delta \|\sqrt{T}Z^\top\|_2^2, \quad (2.6)$$

where $\|Y\|_2 := (\int_M |Y|^2)^{1/2}$ for some vector field Y , and $^\perp$ represents the normal part of the vector field.

We discuss each case.

Case (a): $(\operatorname{tr} T)I - 2T \geq 0$. Let Λ_T be the largest eigenvalue of T , then $\Lambda_T \leq \frac{1}{2} \operatorname{tr} T$, which implies $|\sqrt{T}Z^\top|^2 \leq \Lambda_T |Z^\top|^2 \leq \frac{1}{2} \operatorname{tr} T |Z^\top|^2$, and it follows from (2.6) that

$$\|\sqrt{\operatorname{tr} T}Z^\perp\|_2 \left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2 \geq \int_M \operatorname{tr} T + \frac{\delta}{2} \|\sqrt{T}Z^\top\|_2^2, \quad (2.7)$$

Denote $X = \sqrt{\operatorname{tr} T}Z$ and note that $\|X^\top\|_2^2 = \|X\|_2^2 - \|X^\perp\|_2^2$, then (2.7) becomes

$$\left(\left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2 + \frac{\delta}{2} \|X^\perp\|_2 \right) \|X^\perp\|_2 \geq \int_M \operatorname{tr} T + \frac{\delta}{2} \|X\|_2^2. \quad (2.8)$$

Now we consider the quadratic function $\phi_h(x) = (h + \frac{\delta}{2}x)x$ with $h = \left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2$. Note that (2.4) is equivalent to

$$\phi_h(\|X\|_2) = \left(h + \frac{\delta}{2} \|X\|_2 \right) \|X\|_2 \geq \int_M \operatorname{tr} T + \frac{\delta}{2} \|X\|_2^2,$$

hence, to prove the lemma, it is sufficient to show that

$$\phi_h(\|X\|_2) \geq \phi_h(\|X^\perp\|_2). \quad (2.9)$$

Next we verify (2.9). Firstly, we have

$$2\|X\|_2^2 = 2 \int_M \operatorname{tr} T \operatorname{th}_\delta^2 = \frac{1}{-\delta} \int_M 2 \operatorname{tr} T \left(1 - \frac{1}{\operatorname{ch}_\delta^2} \right) \leq \frac{2}{-\delta} \int_M \operatorname{tr} T. \quad (2.10)$$

Therefore, we obtain that

$$\begin{aligned}
& (\|X\|_2 + \|X^\perp\|_2)\|X^\perp\|_2 \\
&= \|X\|_2\|X^\perp\|_2 + \|X\|_2^2 - \|X^\top\|_2^2 \leq 2\|X\|_2^2 - \|X^\top\|_2^2 \\
&\stackrel{(2.10)}{\leq} \frac{2}{-\delta} \int_M \operatorname{tr} T - \|X^\top\|_2^2 = \frac{2}{-\delta} \left(\int_M \operatorname{tr} T + \frac{\delta}{2} \|X^\top\|_2^2 \right) \\
&\stackrel{(2.7)}{\leq} \frac{2h}{-\delta} \|X^\perp\|_2.
\end{aligned}$$

Since $\|X^\perp\|_2 \neq 0$, we conclude that $2\|X^\perp\|_2 \leq \|X\|_2 + \|X^\perp\|_2 \leq \frac{2h}{-\delta}$. Hence, (2.9) holds by using the monotonicity of $\phi_h = (h + \frac{\delta}{2}x)x$.

Case (b): $M \subset B(R)$, $R \leq \frac{1}{2\sqrt{-\delta}} \operatorname{arcosh} \sqrt{2}$. We still denote $X = \sqrt{\operatorname{tr} T}Z$, then (2.6) becomes

$$\left(\left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2 + \delta \|X^\perp\|_2 \right) \|X^\perp\|_2 \geq \int_M \operatorname{tr} T + \delta \|X\|_2^2. \quad (2.11)$$

Next we can use the analogous argument as in Case (a) for the quadratic function $\psi_h(x) = (h + \delta x)x$ with $h = \left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2$.

By Remark 2.4 we have $\operatorname{ch}_\delta^2 \leq 2$ on M , then

$$2\|X\|_2^2 = 2 \int_M \operatorname{tr} T \operatorname{th}_\delta^2 = \frac{1}{-\delta} \int_M 2 \operatorname{tr} T \left(1 - \frac{1}{\operatorname{ch}_\delta^2} \right) \leq \frac{1}{-\delta} \int_M \operatorname{tr} T. \quad (2.12)$$

Consequently, $2\|X^\perp\|_2 \leq \|X\|_2 + \|X^\perp\|_2 \leq \frac{h}{-\delta}$ follows from

$$(\|X\|_2 + \|X^\perp\|_2)\|X^\perp\|_2 \leq 2\|X\|_2^2 - \|X^\top\|_2^2 \stackrel{(2.12)}{\leq} \frac{1}{-\delta} \left(\int_M \operatorname{tr} T + \delta \|X^\top\|_2^2 \right) \stackrel{(2.6)}{\leq} \frac{h}{-\delta} \|X^\perp\|_2.$$

Now we derive that

$$\left(\left\| \frac{H_T}{\sqrt{\operatorname{tr} T}} \right\|_2 + \delta \|X\|_2 \right) \|X\|_2 = \psi_h(\|X\|_2) \geq \psi_h(\|X^\perp\|_2) \stackrel{(2.11)}{\geq} \int_M \operatorname{tr} T + \delta \|X\|_2^2,$$

which is equivalent to (2.4).

We complete the proof. \square

3. PROOFS FOR p -LAPLACIAN WITH $\delta < 0$

In this section, we prove Theorems 1.2 and 1.3. Firstly, we give the following estimate for $\lambda_{1,p}$.

Lemma 3.1. *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional Riemannian manifold N . Then we have, for all $p > 1$,*

$$\lambda_{1,p} \int_M \operatorname{th}_\delta^p(r) \leq n^{\frac{p-2}{2}} m^{\frac{p}{2}} \int_M \left(1 + \delta \operatorname{th}_\delta^2(r) \right)^{p/2}. \quad (3.1)$$

Proof. We choose $\frac{\operatorname{th}_\delta(r)}{r}x_i$ in Lemma 2.3 as the test functions. Recall the Rayleigh-type quotient (cf. [22])

$$\lambda_{1,p} = \inf \left\{ \frac{\int_M |\nabla u|^p}{\int_M |u|^p} \mid u \in W^{1,p}(M) \setminus \{0\}, \int_M |u|^{p-2}u = 0 \right\}, \quad (3.2)$$

then we obtain

$$\lambda_{1,p} \int_M \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^p \leq \int_M \left| \nabla \left(\frac{\text{th}_\delta(r)}{r} x_i \right) \right|^p \quad \text{for each } i. \quad (3.3)$$

When $1 < p \leq 2$,

$$\text{th}_\delta^p(r) = (\text{th}_\delta^2(r))^{p/2} = \left(\sum_{i=1}^n \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^2 \right)^{p/2} \leq \sum_{i=1}^n \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^p \quad (3.4)$$

By taking $T = I$, $f(r) = \text{th}_\delta(r)$ in Item (1) of Lemma 2.1 and using the Hölder inequality, we derive that

$$\begin{aligned} n^{1-2/p} \left(\sum_{i=1}^n \left| \nabla \left(\frac{\text{th}_\delta(r)}{r} x_i \right) \right|^p \right)^{\frac{2}{p}} &\leq \sum_{i=1}^n \left| \nabla \left(\frac{\text{th}_\delta(r)}{r} x_i \right) \right|^2 \\ &\leq m \left(\frac{1}{\text{ch}_\delta^2} + \delta \frac{\text{th}_\delta^2}{\text{ch}_\delta^2} |\sqrt{T} \nabla r|^2 \right) \\ &\leq m \left(1 + \delta \text{th}_\delta^2(r) \right), \end{aligned} \quad (3.5)$$

where we used $\delta < 0$ and $1 = \text{ch}_\delta^2 + \delta \text{sh}_\delta^2$. Then (3.1) is from (3.3), (3.4) and (3.5).

When $p \geq 2$, the Hölder inequality gives

$$\text{th}_\delta^2(r) = \sum_{i=1}^n \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^2 \leq n^{1-2/p} \left(\sum_{i=1}^n \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^p \right)^{2/p},$$

so we have

$$\lambda_{1,p} \int_M \text{th}_\delta^p(r) \leq n^{p/2-1} \left(\sum_{i=1}^n \lambda_{1,p} \int_M \left| \frac{\text{th}_\delta(r)}{r} x_i \right|^p \right). \quad (3.6)$$

On the other hand, we have

$$\sum_{i=1}^n \left| \nabla \left(\frac{\text{th}_\delta(r)}{r} x_i \right) \right|^p \leq \left(\sum_{i=1}^n \left| \nabla \left(\frac{\text{th}_\delta(r)}{r} x_i \right) \right|^2 \right)^{p/2} \leq m^{\frac{p}{2}} \left(1 + \delta \text{th}_\delta^2(r) \right)^{p/2}. \quad (3.7)$$

Hence, (3.1) follows from (3.3), (3.6) and (3.7). \square

Now we are ready to prove our theorems. Since Theorem 1.2 is a special case of Theorem 1.3, we only prove the latter.

Proof of Theorem 1.3. We have $\delta \text{th}_\delta^2(r) = -\text{th}^2(\sqrt{-\delta}r) \in (-1, 0]$ since $\delta < 0$. From (2.4) we have

$$\sup_M \text{tr } T \int_M \text{th}_\delta^2 \int_M \frac{|H_T|^2}{\text{tr } T} \geq \int_M \text{tr } T \text{th}_\delta^2 \int_M \frac{|H_T|^2}{\text{tr } T} \geq \left(\int_M \text{tr } T \right)^2. \quad (3.8)$$

Case 1. When $p \geq 2$, we have

$$\begin{aligned} \lambda_{1,p} \int_M \text{th}_\delta^p(r) &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \int_M \left(1 + \delta \text{th}_\delta^2(r) \right)^{p/2} \\ &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \int_M \left(1 + \delta \text{th}_\delta^2(r) \right) \\ &= n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \left(\text{vol}(M) + \delta \int_M \text{th}_\delta^2(r) \right) \\ &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \left(\frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|H_T|^2}{\text{tr } T} + \delta \right) \int_M \text{th}_\delta^2(r), \end{aligned} \quad (3.9)$$

where we used (3.8) in the last inequality. By the Hölder inequality, we have

$$\int_M \text{th}_\delta^2(r) \leq \left(\int_M \text{th}_\delta^p(r) \right)^{2/p} (\text{vol}(M))^{1-2/p},$$

which is equivalent to

$$(\text{vol}(M))^{1-p/2} \left(\int_M \text{th}_\delta^2(r) \right)^{p/2} \leq \int_M \text{th}_\delta^p(r). \quad (3.10)$$

Combining (3.10) with (3.9), we have

$$\begin{aligned} \lambda_{1,p} &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \left(\frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|H_T|^2}{\text{tr } T} + \delta \right) \left(\frac{\text{vol}(M)}{\int_M \text{th}_\delta^2(r)} \right)^{p/2-1} \\ &\leq n^{\frac{|p-2|}{2}} m^{p/2} \left(\frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|H_T|^2}{\text{tr } T} + \delta \right) \left(\frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|H_T|^2}{\text{tr } T} \right)^{p/2-1}, \end{aligned}$$

where we used (3.8) again. This is just (1.8).

Case 2. When $1 < p \leq 2$, we have

$$\begin{aligned} \lambda_{1,p} \int_M \text{th}_\delta^p(r) &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \int_M \left(1 + \delta \text{th}_\delta^2(r) \right)^{p/2} \\ &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \int_M \left(1 + \frac{p}{2} \delta \text{th}_\delta^2(r) \right) \\ &= n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \left(\text{vol}(M) + \frac{p}{2} \delta \int_M \text{th}_\delta^2(r) \right) \\ &\leq n^{\frac{|p-2|}{2}} m^{\frac{p}{2}} \left(\frac{\text{vol}(M) \sup_M \text{tr } T}{\left(\int_M \text{tr } T \right)^2} \int_M \frac{|H_T|^2}{\text{tr } T} + \frac{p}{2} \delta \right) \int_M \text{th}_\delta^2(r), \end{aligned} \quad (3.11)$$

where we used (3.8) in the last inequality. Notice that

$$\text{th}_\delta^p(r) = \text{th}^p(\sqrt{-\delta}r)/(-\delta)^{\frac{p}{2}}$$

and

$$\text{th}_\delta^2(r) = \text{th}^2(\sqrt{-\delta}r)/(-\delta) \leq \text{th}^p(\sqrt{-\delta}r)/(-\delta),$$

so

$$\int_M \text{th}_\delta^2(r) \leq (-\delta)^{\frac{p}{2}-1} \int_M \text{th}_\delta^p(r).$$

Inserting this into (3.11) we derive (1.9).

Hence, we complete the whole proof. \square

4. PROOFS FOR L_T OPERATOR WITH $\delta < 0$

In this section, we prove Theorem 1.7. Actually, we will prove the following stronger result.

Theorem 4.1. *Let M be an $m(\geq 2)$ -dimensional closed orientable submanifold in an n -dimensional complete simply-connected Riemannian manifold N of the sectional curvature $K_N \leq \delta < 0$. Let T, S be a symmetric, positive definite and divergence-free $(1, 1)$ -tensor on M . Suppose the one of the followings holds:*

- (a) $(\text{tr } S)I - 2S \geq 0$, i.e., positive semidefinite;
- (b) M is contained in a convex ball of radius $\leq \frac{1}{2\sqrt{-\delta}} \text{arcosh } \sqrt{2}$.

Then we have either

$$\lambda_{1,T} \leq \sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S} \right) \sup_M (\operatorname{tr} S) \left(\delta + \frac{1}{\int_M \operatorname{tr} S} \int_M \frac{|H_S|^2}{\operatorname{tr} S} \right), \quad (4.1)$$

or

$$\lambda_{1,T} \leq \sup_M (\operatorname{tr} T) \left(\delta + \frac{\operatorname{vol}(M) \sup_M \operatorname{tr} S}{\left(\int_M \operatorname{tr} S \right)^2} \int_M \frac{|H_S|^2}{\operatorname{tr} S} \right). \quad (4.2)$$

Proof. We also have the Rayleigh-type quotient (cf. [13])

$$\lambda_{1,T} = \inf \left\{ \frac{-\int_M u L_T u}{\int_M u^2} \mid u \in W^{1,2}(M) \setminus \{0\}, \int_M u = 0 \right\}. \quad (4.3)$$

By slightly modifying the Lemma 2.3, we can prove that there exists a point $q_0 \in N$ such that

$$\int_M \frac{\operatorname{th}_\delta(r)}{r} x_i dv_M = 0 \quad (i = 1, \dots, n), \quad (4.4)$$

where (x_1, \dots, x_n) is the normal coordinate around q_0 , and $r(q) = d(q_0, q)$ is the geodesic distance from q_0 to q on N . Here we omit the detail.

Now by using Lemma 2.1, we have

$$\begin{aligned} \lambda_{1,T} \int_M \operatorname{th}_\delta^2(r) &\leq - \int_M \left(\frac{\operatorname{th}_\delta(r)}{r} x_i \right) L_T \left(\frac{\operatorname{th}_\delta(r)}{r} x_i \right) \\ &\leq \sum_{i=1}^n \int_M \left\langle T \nabla \left(\frac{\operatorname{th}_\delta(r)}{r} x_i \right), \nabla \left(\frac{\operatorname{th}_\delta(r)}{r} x_i \right) \right\rangle \\ &\leq \int_M (\operatorname{tr} T) \left(1 + \delta \operatorname{th}_\delta^2(r) \right), \end{aligned} \quad (4.5)$$

where we used (2.4).

In order to introduce $\operatorname{tr} S$ and H_S and cancel the term of $\int_M \operatorname{th}_\delta^2(r)$, we have two methods.

Method 1.

$$\begin{aligned} &\int_M (\operatorname{tr} T) \left(1 + \delta \operatorname{th}_\delta^2(r) \right) \\ &\leq \sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S} \right) \int_M (\operatorname{tr} S) \left(1 + \delta \operatorname{th}_\delta^2(r) \right) \\ &= \sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S} \right) \left(\frac{\int_M \operatorname{tr} S}{\int_M \operatorname{tr} S \operatorname{th}_\delta^2(r)} + \delta \right) \int_M \operatorname{tr} S \operatorname{th}_\delta^2(r) \\ &\leq \sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S} \right) \sup_M (\operatorname{tr} S) \left(\frac{1}{\int_M \operatorname{tr} S} \int_M \frac{|H_S|^2}{\operatorname{tr} S} + \delta \right) \int_M \operatorname{th}_\delta^2(r), \end{aligned} \quad (4.6)$$

where we used Proposition 2.6 for the tensor S in the last inequality. Putting (4.6) into (4.5), we obtain (4.1).

Method 2.

$$\begin{aligned}
& \int_M (\operatorname{tr} T) \left(1 + \delta \operatorname{th}_\delta^2(r)\right) \\
& \leq \sup_M (\operatorname{tr} T) \int_M \left(1 + \delta \operatorname{th}_\delta^2(r)\right) \\
& = \sup_M (\operatorname{tr} T) \left(\operatorname{vol}(M) \times 1 + \delta \int_M \operatorname{th}_\delta^2(r)\right) \\
& \leq \sup_M (\operatorname{tr} T) \left(\operatorname{vol}(M) \frac{\int_M \operatorname{tr} S \operatorname{th}_\delta^2(r)}{(\int_M \operatorname{tr} S)^2} \int \frac{|H_S|^2}{\operatorname{tr} S} + \delta \int_M \operatorname{th}_\delta^2(r)\right) \\
& \leq \sup_M (\operatorname{tr} T) \left(\frac{\operatorname{vol}(M) \sup_M \operatorname{tr} S}{(\int_M \operatorname{tr} S)^2} \int \frac{|H_S|^2}{\operatorname{tr} S} + \delta\right) \int_M \operatorname{th}_\delta^2(r) \tag{4.7}
\end{aligned}$$

where we used Proposition 2.6 for the tensor S in the penultimate inequality. Putting (4.7) into (4.5), we obtain (4.2). \square

Remark 4.2. We briefly discuss the upper bounds in Eqs. (4.1) and (4.2).

- (1) When $S = T$, (4.1) becomes (1.13) and is better than (4.2).
- (2) When $\operatorname{tr} S$ is constant, (4.1) is the same as (4.2).
- (3) Generally, $\sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S}\right) \sup_M (\operatorname{tr} S) \geq \sup_M \operatorname{tr} T$, but we cannot compare $\sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S}\right)$ and $\frac{\operatorname{vol}(M) \sup_M \operatorname{tr} T}{\int_M \operatorname{tr} S}$. For example, when $\operatorname{tr} T$ is constant, we have $\sup_M \left(\frac{\operatorname{tr} T}{\operatorname{tr} S}\right) \geq \frac{\operatorname{vol}(M) \sup_M \operatorname{tr} T}{\int_M \operatorname{tr} S}$, so we cannot compare these two upper bounds since $\delta < 0$.
- (4) When $T = I$, (4.1) recovers (1.10).

5. WEIGHTED MANIFOLDS AND BOUNDARY VALUE PROBLEMS

In this section, we consider the weighted manifolds and some boundary value problems and give the corresponding Reilly-type inequalities. Since most of the discussion is similar to the previous section, we will highlight the difference and omit other details.

5.1. Basic settings and lemmas. Firstly we recall some concepts of weighted manifolds. Let (N, \bar{g}) be a Riemannian manifold equipped with the metric \bar{g} . We denote by $dv_{\bar{g}}$ by the usual (non-weighted) volume form on N . For a smooth function f on N , we can consider the *weighted volume form* (or *weighted measure*) $\bar{\mu}_f := e^{-f} dv_{\bar{g}}$. The triplet $(N, \bar{g}, \bar{\mu}_f)$ is called a *weighted manifold*. It is also called a *metric measure space*, or a *manifold with density* in some literature.

Now let M be a submanifold of $(N, \bar{g}, \bar{\mu}_f)$. Then M has an induced metric g and a weighted measure $\mu_f = e^{-f} dv_g$. If M has non-empty boundary ∂M , then we denote by \tilde{g} and $\tilde{\mu}_f = e^{-f} dv_{\tilde{g}}$ on ∂M the induced metric and the weighted measure on ∂M , respectively.

Denote $\dim M = m, \dim N = n$. The corresponding n -volume $\operatorname{vol}_f(N)$, m -volume $\operatorname{vol}_f(M)$ and $(m-1)$ -volume $\operatorname{vol}_f(\partial M)$ (when $\partial M \neq \emptyset$) are respectively given by

$$\operatorname{vol}_f(N) = \int_N \bar{\mu}_f, \quad \operatorname{vol}_f(M) = \int_M \mu_f, \quad \operatorname{vol}_f(\partial M) = \int_{\partial M} \mu_f. \tag{5.1}$$

For a symmetric $(1,1)$ -tensor $T = (T_{ij})$ on M , we can define the weighted divergence (or f -divergence) on M by

$$\operatorname{div}_{f,M} = \operatorname{div}_M - \langle \nabla f, \cdot \rangle. \tag{5.2}$$

Then the weighted L_T operator, denoted by $L_{T,f}$, can be defined by

$$L_{T,f}u := \operatorname{div}_{f,M}(T\nabla u) = \operatorname{div}_M(T\nabla u) - \langle \nabla f, T\nabla u \rangle. \quad (5.3)$$

One can easily obtain the Stokes' formula for the weighted operators with respect to the weighted measure μ_f . When T is divergence-free, $L_{T,f}$ is self-adjoint with respect to the weighted measure μ_f . In particular, the case of $T = I$ has been widely studied, and in this situation, $L_{T,f}$ becomes $\Delta u - \langle \nabla f, \nabla u \rangle$, which is usually called the weighted Laplacian or f -Laplacian.

5.2. The case $\delta = 0$. Let $\operatorname{sn}_\delta(t) = \frac{1}{\sqrt{\delta}} \sin(\sqrt{\delta}t)$, t and $\frac{1}{\sqrt{-\delta}} \sinh(\sqrt{-\delta}t)$ when $\delta > 0$, $\delta = 0$ and $\delta < 0$, respectively. Very recently, Manfio, Roth and Upadhyay [16] studied various extrinsic eigenvalues upper bounds in weighted manifolds and proved the following three results, which are restated by using the notations and settings in Sect. 5.1 as follows¹.

Theorem 5.1 (cf. [16, Theorem 1.1]). *Let M be a closed submanifold immersed in a weighted manifold $(N, \bar{g}, \bar{\mu}_f)$ with sectional curvature $K_N \leq \delta \leq 0$. Let T, S be symmetric, divergence-free and positive definite $(1, 1)$ -tensors on M . Then the first positive eigenvalue of $L_{T,f}$, denoted by λ_1 , satisfies*

$$\lambda_1 \leq \sup_M \left[\delta \operatorname{tr} T + \sup_M \left(\frac{|H_T - T\nabla f|}{\operatorname{tr} S} \right) |H_S - S\nabla f| \right]. \quad (5.4)$$

Theorem 5.2 (cf. [16, Theorem 1.2]). *Let M be a compact submanifold with non-empty boundary ∂M immersed in a weighted manifold $(N, \bar{g}, \bar{\mu}_f)$ with sectional curvature $K_N \leq \delta \leq 0$. Let T, S be symmetric, divergence-free and positive definite $(1, 1)$ -tensors on M and ∂M , respectively, and denote by σ_1 the first eigenvalue of the generalized weighted Steklov problem*

$$(f\text{-Stek}) \begin{cases} L_{T,f}u = 0, & \text{in } M; \\ \frac{\partial u}{\partial \nu_T} = \sigma u, & \text{on } \partial M, \end{cases}$$

where $\frac{\partial u}{\partial \nu_T} = \langle T(\tilde{\nabla}u), \nu \rangle$, $\tilde{\nabla}$ and ν are the Levi-Civita connection and unit normal on ∂M , respectively.

Assume that M is contained in the geodesic ball $B(p, R)$ of radius R , where p is the center of mass of ∂M for the measure $\tilde{\mu}_f$, then

$$\begin{aligned} \sigma_1 &\leq \sup_M \left[\delta \operatorname{tr} T + \sup_M \left(\frac{|H_T - T\nabla f|}{\operatorname{tr} T} \right) |H_T - T\nabla f| \right] \\ &\quad \times \left[\delta + \frac{\sup_{\partial M} |H_S - S(\tilde{\nabla}f)|^2}{\inf_{\partial M} (\operatorname{tr} S)^2} \right] \frac{\operatorname{vol}_f(M)}{\operatorname{vol}_f(\partial M)} \operatorname{sn}_\delta^2(R). \end{aligned} \quad (5.5)$$

Theorem 5.3 (cf. [16, Theorem 1.3]). *Let M be a compact submanifold with non-empty boundary ∂M immersed in a Riemannian manifold (N, \bar{g}) with sectional curvature $K_N \leq \delta \leq 0$. Let S be a symmetric, divergence-free and positive definite $(1, 1)$ -tensors on ∂M , and denote by α_1 the first non-zero eigenvalue of the Steklov-Wentzell problem*

$$(SW) \begin{cases} \Delta u = 0, & \text{in } M; \\ -b\tilde{\Delta}u - \frac{\partial u}{\partial \nu} = \alpha u, & \text{on } \partial M, \end{cases}$$

where b is a given positive constant and $\tilde{\Delta}$ is the Laplacian on ∂M .

¹In fact, the estimates in the case $K_N > 0$ are also obtained, but here we focus on the case $K_N \leq 0$ and assume that N is simply connected.

Assume that M is contained in the geodesic ball $B(p, R)$ of radius R , where p is the center of mass of ∂M , then

$$\begin{aligned} \alpha_1 \leq & \left[m \frac{\text{vol}(M)}{\text{vol}(\partial M)} + b(m-1) - \delta \text{sn}_\delta^2(R) \left(\frac{\text{vol}(M)}{\text{vol}(\partial M)} + b \right) \right] \\ & \times \left(\delta + \frac{\sup_{\partial M} |H_S|^2}{\inf_{\partial M} (\text{tr } S)^2} \right). \end{aligned} \quad (5.6)$$

When $\delta = 0$, we can obtain upper bounds in terms of the integral. Precisely, we have

Theorem 5.4. *Settings and notations as in Theorem 5.1 with $\delta = 0$. Then*

$$\lambda_1 \left(\int_M \text{tr } S \mu_f \right)^2 \leq \int_M \text{tr } T \mu_f \int_M |H_S - S \nabla f|^2 \mu_f. \quad (5.7)$$

Theorem 5.5. *Settings and notations as in Theorem 5.2 with $\delta = 0$. Then*

$$\sigma_1 \left(\int_{\partial M} \text{tr } S \tilde{\mu}_f \right)^2 \leq \int_M \text{tr } T \mu_f \int_{\partial M} |H_S - S \nabla f|^2 \mu_f. \quad (5.8)$$

Theorem 5.6. *Settings and notations as in Theorem 5.3 with $\delta = 0$. Then*

$$\alpha_1 \left(\int_{\partial M} \text{tr } S \right)^2 \leq \left(m \text{vol}(M) + b(m-1) \text{vol}(\partial M) \right) \int_{\partial M} |H_S|^2. \quad (5.9)$$

Proofs of Theorems 5.4, 5.5 and 5.6. We have the variational characterization (cf. [1, 8, 15, 16, 21]):

$$\lambda_1 = \inf \left\{ \frac{\int_M \langle T \nabla u, \nabla u \rangle \mu_f}{\int_M u^2 \mu_f} \mid \int_M u \mu_f = 0 \right\}. \quad (5.10)$$

$$\sigma_1 = \inf \left\{ \frac{\int_M \langle T \nabla u, \nabla u \rangle \mu_f}{\int_{\partial M} u^2 \tilde{\mu}_f} \mid \int_{\partial M} u \tilde{\mu}_f = 0 \right\}. \quad (5.11)$$

$$\alpha_1 = \inf \left\{ \frac{\int_M |\nabla u|^2 + b \int_{\partial M} |\tilde{\nabla} u|^2}{\int_M u^2} \mid \int_{\partial M} u = 0 \right\}. \quad (5.12)$$

When $\delta = 0$, consider $Z = r \nabla^N r$, we have (cf. [16, Lemma 2.1])

$$\text{div}_{f,M}(TZ^\top) \geq \text{tr } T + \langle Z, H_T - T \nabla f \rangle, \quad (5.13)$$

which implies

$$\int_M \text{tr } T \mu_f \leq - \int_M \langle Z, H_T - T \nabla f \rangle \mu_f \leq \int_M |H_T - T \nabla f| r \mu_f \quad (5.14)$$

if M has no boundary.

Similar to the proof of Lemma 2.3, there exists a point $q_0 \in N$ such that

$$\int_M x_i \mu_f = 0 \quad (i = 1, \dots, n), \quad (5.15)$$

or

$$\int_{\partial M} x_i \tilde{\mu}_f = 0 \quad (i = 1, \dots, n), \quad (5.16)$$

By taking $\phi(r) = r$ in Lemma 2.1 (1), we have $\sum_{i=1}^n \langle T \nabla x_i, \nabla x_i \rangle \leq \text{tr } T$. In particular, $\sum_{i=1}^n \langle \nabla x_i, \nabla x_i \rangle \leq m$ for $T = I$. We also have $\sum_{i=1}^n \langle \tilde{\nabla} x_i, \tilde{\nabla} x_i \rangle \leq m-1$ when we consider the boundary ∂M .

Hence, considering (5.15), we have

$$\lambda_1 \int_M r^2 \mu_f = \lambda_1 \int_M \sum_{i=1}^n (x_i)^2 \mu_f \leq \int_M \sum_{i=1}^n \langle T \nabla x_i, \nabla x_i \rangle \mu_f \leq \int_M \operatorname{tr} T \mu_f.$$

Multiplying the both sides by $\int_M |H_S - S \nabla f|^2 \mu_f$, we derive that

$$\begin{aligned} \left(\int_M |H_S - S \nabla f|^2 \mu_f \right) \left(\int_M \operatorname{tr} T \mu_f \right) &\geq \lambda_1 \left(\int_M r^2 \mu_f \right) \left(\int_M |H_S - S \nabla f|^2 \mu_f \right) \\ &\geq \lambda_1 \left(\int_M r |H_S - S \nabla f| \mu_f \right)^2 \\ &\geq \lambda_1 \left(\int_M \operatorname{tr} S \mu_f \right)^2, \end{aligned}$$

where we used (5.14) for the tensor S and the Hölder inequality. This is (5.7).

Considering (5.16), we have

$$\sigma_1 \int_{\partial M} r^2 \tilde{\mu}_f \leq \int_M \sum_{i=1}^n \langle T \nabla x_i, \nabla x_i \rangle \mu_f \leq \int_M \operatorname{tr} T \mu_f.$$

Multiplying the both sides by $\int_{\partial M} |H_S - S \nabla f|^2 \tilde{\mu}_f$, we can obtain (5.8) by using the similar arguments.

Considering (5.16) for $f \equiv 0$, we have

$$\alpha_1 \int_{\partial M} r^2 \leq \int_M \sum_{i=1}^n \langle \nabla x_i, \nabla x_i \rangle + b \int_{\partial M} \sum_{i=1}^n \langle \tilde{\nabla} x_i, \tilde{\nabla} x_i \rangle \leq n \operatorname{vol}(M) + b(n-1) \operatorname{vol}(\partial M).$$

Multiplying the both sides by $\int_{\partial M} |H_S|^2 \tilde{\mu}_f$, we can obtain (5.9) by using the similar arguments. \square

Remark 5.7. Our upper bounds are formally the same forms as the previous results such as in [7, 20, 21]. The main differences are:

(1) In [7, 20, 21], the ambient manifold is the Euclidean space. Here we extend it to a Cartan-Hadamard manifold.

(2) When the ambient manifold is the Euclidean space, one can choose test functions by moving the mass center to the origin and use the Hsiung–Minkowski formulas; when the ambient manifold is a Cartan-Hadamard manifold, we choose test functions by (a modified version of) Lemma 2.3, and use the inequality (5.13).

(3) When the ambient space is a Cartan-Hadamard manifold, giving the sufficient and necessary conditions for the equalities become more difficult.

REFERENCES

- [1] Giles Auchmuty, *Steklov eigenproblems and the representation of solutions of elliptic boundary value problems*, Numer. Funct. Anal. Optim. **25** (2004), no. 3-4, 321–348. MR2072072
- [2] Linfen Cao and Haizhong Li, *r-minimal submanifolds in space forms*, Ann. Global Anal. Geom. **32** (2007), no. 4, 311–341. MR2346221
- [3] Isaac Chavel, *On A. Hurwitz' method in isoperimetric inequalities*, Proc. Amer. Math. Soc. **71** (1978), no. 2, 275–279. MR493885
- [4] Hang Chen, *Extrinsic upper bound of the eigenvalue for p-Laplacian*, Nonlinear Anal. **196** (2020), 111833, 9. MR4074630
- [5] Hang Chen and Xianfeng Wang, *Sharp Reilly-type inequalities for a class of elliptic operators on submanifolds*, Differential Geom. Appl. **63** (2019), 1–29. MR3896192

- [6] Hang Chen and Guofang Wei, *Reilly-type inequalities for p -Laplacian on submanifolds in space forms*, *Nonlinear Anal.* **184** (2019), 210–217. MR3921104
- [7] Qun Chen and Jianghai Shi, *A Reilly inequality for the first non-zero eigenvalue of a class of operators on Riemannian manifold*, *Bull. Braz. Math. Soc. (N.S.)* **49** (2018), no. 3, 481–493. MR3862792
- [8] M. Dambrine, D. Kateb, and J. Lamboley, *An extremal eigenvalue problem for the Wentzell-Laplace operator*, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **33** (2016), no. 2, 409–450. MR3465381
- [9] M. Carmen Domingo-Juan, Vicente Miquel, and Jonathan J. Zhu, *Reilly's type inequality for the Laplacian associated to a density related with shrinkers for MCF*, *J. Differential Equations* **272** (2021), 958–978. MR4166071
- [10] Feng Du and Jing Mao, *Reilly-type inequalities for p -Laplacian on compact Riemannian manifolds*, *Front. Math. China* **10** (2015), no. 3, 583–594. MR3323673
- [11] A. El Soufi and S. Ilias, *Une inégalité du type "Reilly" pour les sous-variétés de l'espace hyperbolique*, *Comment. Math. Helv.* **67** (1992), no. 2, 167–181. MR1161279
- [12] Daniel H. Gottlieb, *A de Moivre like formula for fixed point theory*, *Fixed point theory and its applications* (Berkeley, CA, 1986), 1988, pp. 99–105. MR956481
- [13] Jean-François Grosjean, *Extrinsic upper bounds for the first eigenvalue of elliptic operators*, *Hokkaido Math. J.* **33** (2004), no. 2, 319–339. MR2073001
- [14] Ernst Heintze, *Extrinsic upper bounds for λ_1* , *Math. Ann.* **280** (1988), no. 3, 389–402. MR936318
- [15] Saïd Ilias and Ola Makhoul, *A Reilly inequality for the first Steklov eigenvalue*, *Differential Geom. Appl.* **29** (2011), no. 5, 699–708. MR2831826
- [16] Fernando Manfio, Julien Roth, and Abhitosh Upadhyay, *Extrinsic eigenvalues upper bounds for submanifolds in weighted manifolds* (2021). working paper or preprint.
- [17] Marston Morse, *Singular Points of Vector Fields Under General Boundary Conditions*, *Amer. J. Math.* **51** (1929), no. 2, 165–178. MR1506710
- [18] Yanyan Niu and Shicheng Xu, *Total squared mean curvature of immersed submanifolds in a negatively curved space*, arXiv:2106.01912v1 (2021).
- [19] Robert C. Reilly, *On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space*, *Comment. Math. Helv.* **52** (1977), no. 4, 525–533. MR482597
- [20] Julien Roth, *General Reilly-type inequalities for submanifolds of weighted Euclidean spaces*, *Colloq. Math.* **144** (2016), no. 1, 127–136. MR3485886
- [21] ———, *Reilly-type inequalities for Paneitz and Steklov eigenvalues*, *Potential Anal.* **53** (2020), no. 3, 773–798. MR4140078
- [22] L. Veron, *Some existence and uniqueness results for solution of some quasilinear elliptic equations on compact Riemannian manifolds*, *Differential equations and its applications* (Budapest, 1991), 1991, pp. 317–352. MR1468764
- [23] Rushan Wang, *Reilly inequalities of elliptic operators on closed submanifolds*, *Bull. Aust. Math. Soc.* **80** (2009), no. 2, 335–346. MR2540366

(Hang Chen) SCHOOL OF MATHEMATICS AND STATISTICS, NORTHWESTERN POLYTECHNICAL UNIVERSITY, XI' AN 710129, P. R. CHINA, EMAIL: CHENHANG86@NWPU.EDU.CN

(Xudong Gui) SCHOOL OF MATHEMATICS AND STATISTICS, NORTHWESTERN POLYTECHNICAL UNIVERSITY, XI' AN 710129, P. R. CHINA, EMAIL: 2018302512@MAIL.NWPU.EDU.CN