

Normalization and reduction of the Stark Hamiltonian

Richard Cushman¹

Abstract

This paper details a calculation of the second order normal form of the Stark effect Hamiltonian after regularization, using the Kustaanheimo-Stiefel mapping. After reduction we obtain an integrable two degree of freedom system on $S_h^2 \times S_h^2$, which we reduce again to obtain a one degree of freedom Hamiltonian system.

Most papers on the Stark Hamiltonian discuss only the two degree of freedom version, which was shown to be completely integrable by Lagrange [4], see also [5]. We do not expect the three degree of freedom Stark Hamiltonian to be completely integrable. Our treatment follows that of [3], which does not give any details how the second order normal form was obtained. We use the notation and results of [2].

1 The basic setup

On $T_0\mathbb{R}^3 = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}^3$ with coordinates (x, y) and standard symplectic form $\omega_3 = \sum_{i=1}^3 dx_i \wedge dy_i$ consider the Stark Hamiltonian

$$K(x, y) = \frac{1}{2}\langle y, y \rangle - \frac{1}{|x|} + fx_3. \quad (1)$$

Here $\langle \cdot, \cdot \rangle$ is the Euclidean inner product on \mathbb{R}^3 with associated norm $|\cdot|$. On the negative energy level $-\frac{1}{2}k^2$ with $k > 0$ rescaling time $dt \mapsto \frac{|x|}{k} ds$, we obtain

$$0 = \frac{1}{2k}(|x|\langle y, y \rangle + k^2|x|) - \frac{1}{k} + fx_3\frac{|x|}{k}. \quad (2)$$

In other words, (x, y) lies in the $\frac{1}{k}$ level set of

$$\widehat{K}(x, y) = \frac{1}{2k}|x|(\langle y, y \rangle + k^2|x|) + fx_3\frac{|x|}{k}. \quad (3)$$

We assume that f is small, namely, $f = \varepsilon\beta$. After the symplectic coordinate change $(x, y) \mapsto (\frac{1}{k}x, ky)$ the Hamiltonian \widehat{K} becomes the preregularized Hamiltonian

$$\mathcal{K}(x, y) = \frac{1}{2}|x|(\langle y, y \rangle + |x|) + \varepsilon\beta x_3|x| \quad (4)$$

on the level set $\mathcal{K}^{-1}(1)$.

¹e-mail: r.h.cushman@gmail.com

Let $T_0\mathbb{R}^4 = (\mathbb{R}^4 \setminus \{0\}) \times \mathbb{R}^4$ have coordinates (q, p) and a symplectic form $\omega_4 = \sum_{i=1}^4 dq_i \wedge dp_i$. Pull back \mathcal{K} by the Kustaanheimo-Stiefel mapping

$$\mathcal{KS} : T_0\mathbb{R}^4 \rightarrow T_0\mathbb{R}^3 : (q, p) \mapsto (x, y),$$

where

$$\begin{aligned} x_1 &= 2(q_1q_3 + q_2q_4) = U_2 - K_1 \\ x_2 &= 2(q_1q_4 - q_2q_3) = U_3 - K_2 \\ x_3 &= q_1^2 + q_2^2 - q_3^2 - q_4^2 = U_4 - K_3 \\ y_1 &= (\langle q, q \rangle)^{-1}(q_1p_3 + q_2p_4 + q_3p_1 + q_4p_2) = (H_2 + V_1)^{-1}V_2 \\ y_2 &= (\langle q, q \rangle)^{-1}(q_1p_4 - q_2p_3 - q_3p_2 + q_4p_1) = (H_2 + V_1)^{-1}V_3 \\ y_3 &= (\langle q, q \rangle)^{-1}(q_1p_1 + q_2p_2 + q_3p_3 + q_4p_4) = (H_2 + V_1)^{-1}V_4 \end{aligned}$$

and

$$\begin{aligned} H_2 &= \frac{1}{2}(p_1^2 + p_2^2 + p_3^2 + p_4^2 + q_1^2 + q_2^2 + q_3^2 + q_4^2) \\ \Xi &= q_1p_2 - q_2p_1 + q_3p_4 - q_4p_3, \end{aligned}$$

to get the regularized Stark Hamiltonian

$$\mathcal{H} = H_2 + \varepsilon\beta(U_4V_1 + H_2U_4 - K_3V_1 - H_2K_3) \quad (5)$$

on $\Xi^{-1}(0)$, since $|x| = \langle q, q \rangle = H_1 + V_1$. Here

$$\begin{aligned} K_1 &= -(q_1q_3 + q_2q_4 + p_1p_3 + p_2p_4) \\ K_2 &= -(q_1q_4 - q_2q_3 + p_1p_4 - p_2p_3) \\ K_3 &= \frac{1}{2}(q_3^2 + q_4^2 + p_3^2 + p_4^2 - q_1^2 - q_2^2 - p_1^2 - p_2^2) \\ L_1 &= q_4p_1 - q_3p_2 + q_2p_3 - q_1p_4 \\ L_2 &= q_1p_3 + q_2p_4 - q_3p_1 - q_4p_2 \\ L_3 &= q_3p_4 - q_4p_3 + q_2p_1 - q_1p_2 \\ U_1 &= -(q_1p_1 + q_2p_2 + q_3p_3 + q_4p_4) \\ U_2 &= q_1q_3 + q_2q_4 - p_1p_3 - p_2p_4 \\ U_3 &= q_1q_4 - q_2q_3 + p_2p_3 - p_1p_4 \\ U_4 &= \frac{1}{2}(q_1^2 + q_2^2 - q_3^2 - q_4^2 + p_3^2 + p_4^2 - p_1^2 - p_2^2) \\ V_1 &= \frac{1}{2}(q_1^2 + q_2^2 + q_3^2 + q_4^2 - p_1^2 - p_2^2 - p_3^2 - p_4^2) \\ V_2 &= q_1p_3 + q_2p_4 + q_3p_1 + q_4p_2 \\ V_3 &= q_1p_4 - q_2p_3 + q_4p_1 - q_3p_2 \\ V_4 &= q_1p_1 + q_2p_2 - q_3p_3 - q_4p_4. \end{aligned}$$

generate the algebra of polynomials invariant under the S^1 action φ_s^Ξ given by the flow of X_Ξ on $(T\mathbb{R}^4 = \mathbb{R}^8, \omega_4)$. The Hamiltonian \mathcal{H} (5) is invariant under this S^1 action and thus is a smooth function on the orbit space $\Xi^{-1}(0)/S^1 \subseteq \mathbb{R}^{16}$ with coordinates $(K, L, H, \Xi; U, V)$.

2 The first order normal form on $\Xi^{-1}(0)/S^1$

The harmonic oscillator vector field X_{H_2} on $(T\mathbb{R}^4, \omega_4)$ induces the vector field $Y_{H_2} = \sum_{i=1}^4 (2V_i \frac{\partial}{\partial U_i} - 2U_i \frac{\partial}{\partial V_i})$ on the orbit space $\mathbb{R}^8/S^1 \subseteq \mathbb{R}^{16}$, which leaves $\Xi^{-1}(0)/S^1$ invariant.

We now compute the first order normal form of the Hamiltonian \mathcal{H} (5) on the reduced space $\Xi^{-1}(0)/S^1 \subseteq \mathbb{R}^8/S^1$.

The average of $H_2U_4 - K_3V_1$ over the flow

$$\varphi_t^{Y_{H_2}}(K, L, H_2, \Xi; U, V) = (K, L, H_2, \Xi; U \cos 2t + V \sin 2t, -U \sin 2t + V \cos 2t)$$

of Y_{H_2} is

$$\begin{aligned} \overline{H_2U_4 - K_3V_1} &= \frac{1}{\pi} \int_0^\pi (\varphi_t^{Y_{H_2}})^*(H_2U_4 - K_3V_1) dt \\ &= \frac{1}{\pi} \int_0^\pi (U_4 \cos 2t + V_4 \sin 2t) dt H_2 - \frac{1}{\pi} \int_0^\pi (-U_1 \sin 2t + V_1 \cos 2t) dt K_3 = 0. \end{aligned}$$

The second equality above follows because $L_{X_{H_2}}K_3 = 0$ and the third because $\overline{\cos 2t} = \overline{\sin 2t} = 0$. The average of U_4V_1 over the flow of Y_{H_2} on $\Xi^{-1}(0)/S^1$ is

$$\begin{aligned} \overline{U_4V_1} &= \frac{1}{\pi} \int_0^\pi (\varphi_t^{Y_{H_2}})^*(U_4V_1) dt \\ &= -\frac{1}{2}U_1U_4 \overline{\sin 4t} + U_4V_1 \overline{\cos^2 2t} - U_1V_4 \overline{\sin^2 2t} + \frac{1}{2}V_1V_4 \overline{\sin 4t} \\ &= \frac{1}{2}(U_4V_1 - U_1V_4) = -\frac{1}{2}H_2K_3, \end{aligned}$$

since $\overline{\cos^2 2t} = \overline{\sin^2 2t} = \frac{1}{2}$ and $\overline{\sin 4t} = 0$. The last equality above follows from the explicit description of the orbit space \mathbb{R}^8/S^1 as the semialgebraic variety in \mathbb{R}^{16} with coordinates $(K, L, U, V; H_2, \Xi)$ given by

$$\begin{aligned} \langle U, U \rangle &= U_1^2 + U_2^2 + U_3^2 + U_4^2 = H_2^2 - \Xi^2 \geq 0 \quad H_2 \geq 0 \\ \langle V, V \rangle &= V_1^2 + V_2^2 + V_3^2 + V_4^2 = H_2^2 - \Xi^2 \geq 0 \\ \langle U, V \rangle &= U_1V_1 + U_2V_2 + U_3V_3 + U_4V_4 = 0 \\ U_2V_1 - U_1V_2 &= L_1\Xi - K_1H_2 \\ U_3V_1 - U_1V_3 &= L_2\Xi - K_2H_2 \\ U_4V_1 - U_1V_4 &= L_3\Xi - K_3H_2 \\ U_4V_3 - U_3V_4 &= K_1\Xi - L_1H_2 \\ U_2V_4 - U_4V_2 &= K_2\Xi - L_2H_2 \\ U_3V_2 - U_2V_3 &= K_3\Xi - L_3H_2. \end{aligned} \tag{6}$$

So the average of $U_4V_1 + H_2U_4 - K_3V_1 - H_2K_3$ over the flow of Y_{H_2} is $-\frac{3}{2}H_2K_3$ on $\Xi^{-1}(0)/S^1$. Thus the first order normal form of the regularized Stark Hamiltonian \mathcal{H} (5) on $\Xi^{-1}(0)/S^1$ is

$$\mathcal{H}_{\text{nf}}^{(1)} = H_2 - \frac{3}{2}\beta\varepsilon H_2K_3. \tag{7}$$

3 The second order normal form on $\Xi^{-1}(0)/S^1$

In order to compute the second order normal form of the Hamiltonian \mathcal{H} on $\Xi^{-1}(0)/S^1$, we need to find a function F on \mathbb{R}^{16} such that changing coordinates by the time ε value of the flow of the Hamiltonian vector field Y_F brings the regularized Hamiltonian \mathcal{H} (5) into first order normal form. Choose F so that

$$L_{Y_F}H_2 = \beta\left(-U_4V_1 - \frac{1}{2}H_2K_3 - H_2U_4 + K_3V_1\right). \quad (8)$$

The following calculation shows that this choice does the job.

$$\begin{aligned} (\varphi_\varepsilon^{Y_F})^*\mathcal{H} &= \mathcal{H} + \varepsilon L_{Y_F}\mathcal{H} + \frac{1}{2}\varepsilon^2 L_{Y_F}^2\mathcal{H} + \mathcal{O}(\varepsilon^3) \\ &= H_2 + \varepsilon\beta(U_4V_1 + H_2U_4 - K_3V_1 - H_2K_3) + \varepsilon L_{Y_F}H_2 \\ &\quad + \varepsilon^2\beta L_{Y_F}(U_4V_1 + H_2U_4 - K_3V_1 - H_2K_3) + \frac{1}{2}\varepsilon^2 L_{Y_F}^2H_2 + \mathcal{O}(\varepsilon^3) \\ &= H_2 + \varepsilon\beta(U_4V_1 + H_2U_4 - K_3V_1 - H_2K_3) \\ &\quad + \varepsilon\beta\left(-U_4V_1 - \frac{1}{2}H_2K_3 - H_2U_4 + K_3V_1\right) \\ &\quad + \varepsilon^2[L_{Y_F}(-L_{Y_F}H_2 - \frac{3}{2}\beta H_2K_3) + \frac{1}{2}L_{Y_F}^2H_2] + \mathcal{O}(\varepsilon^3) \\ &= H_2 - \frac{3}{2}\varepsilon\beta H_2K_3 - \frac{1}{2}\varepsilon^2(L_{Y_F}^2H_2 + 3\beta L_{Y_F}(H_2K_3)) + \mathcal{O}(\varepsilon^3). \end{aligned} \quad (9)$$

To determine the function F , we solve equation (8). Write $F = F_1 + F_2$, where $L_{Y_{H_2}}F_1 = \beta(U_4V_1 + \frac{1}{2}H_2K_3)$ and $L_{Y_{H_2}}F_2 = \beta(H_2U_4 - K_3V_1)$. Then

$$\begin{aligned} L_{Y_F}H_2 &= -L_{Y_{H_2}}Y_F = -L_{Y_{H_2}}F_1 - L_{Y_{H_2}}F_2 \\ &= -\beta(U_4V_1 + \frac{1}{2}H_2K_3) - \beta(H_2U_4 - K_3V_1). \end{aligned} \quad (10)$$

Since $L_{Y_{H_2}}V_4 = -2U_4$ and $L_{Y_{H_2}}U_1 = 2V_1$, it follows that

$$F_2 = -\frac{\beta}{2}(H_2V_4 + K_3U_1). \quad (11a)$$

Now

$$F_1 = \frac{\beta}{\pi} \int_0^\pi t(\varphi_t^{Y_{H_2}})^*(U_4V_1 + \frac{1}{2}H_2K_3) dt = \frac{\beta}{\pi} \int_0^\pi t(\varphi_t^{Y_{H_2}})^*(U_4V_1) dt + \frac{\pi\beta}{4}H_2K_3,$$

see [1], and

$$\begin{aligned} \frac{\beta}{\pi} \int_0^\pi t(\varphi_t^{Y_{H_2}})^*(U_4V_1) dt &= \\ &= -\frac{\beta}{2}(U_1U_4) \frac{1}{\pi} \int_0^\pi t \sin 4t dt + \beta(U_4V_1) \frac{1}{\pi} \int_0^\pi t \cos^2 2t dt \\ &\quad - \beta(U_1V_4) \frac{1}{\pi} \int_0^\pi t \sin^2 2t dt + \frac{\beta}{2}(V_1V_4) \frac{1}{\pi} \int_0^\pi t \sin 4t dt \\ &= \frac{\beta}{8}(U_1U_4 - V_1V_4) + \frac{\pi\beta}{4}(U_4V_1 - U_1V_4), \end{aligned}$$

since $\frac{1}{\pi} \int_0^\pi t \sin 4t dt = -\frac{1}{4}$ and $\frac{1}{\pi} \int_0^\pi t \sin^2 2t dt = \frac{1}{\pi} \int_0^\pi t \cos^2 2t dt = \frac{\pi}{4}$. Thus

$$F_1 = \frac{\beta}{8}(U_1U_4 - V_1V_4) + \frac{\pi\beta}{4}(U_4V_1 - U_1V_4 + H_2K_3) = \frac{\beta}{8}(U_1U_4 - V_1V_4)$$

on $\Xi^{-1}(0)/S^1$, see (6). Hence on $\Xi^{-1}(0)/S^1$

$$F = F_1 + F_2 = \frac{\beta}{8}(U_1U_4 - V_1V_4) - \frac{\beta}{2}(H_2V_4 + K_3U_1). \quad (12)$$

We now calculate the average over the flow of Y_{H_2} of

$$-\frac{3}{2}\beta L_{Y_F}(H_2K_3) - \frac{1}{2}L_{Y_F}^2 H_2, \quad (13)$$

which is the ε^2 term in the transformed Hamiltonian $(\varphi_\varepsilon^{Y_F})^*\mathcal{H}$, see (9). This determines the second order normal form of \mathcal{H} on $\Xi^{-1}(0)/S^1$. We begin with the term

$$-\frac{3}{2}\beta L_{Y_F}(H_2K_3) = -\frac{3}{2}\beta[K_3(L_{Y_F}H_2) - H_2(L_{Y_{K_3}}F)].$$

The average of

$$-\frac{3}{2}\beta K_3(L_{Y_F}H_2) = \frac{3}{2}\beta K_3(U_4V_1 + \frac{1}{2}\beta H_2K_3 + H_2U_4 - K_3V_1)$$

vanishes on $\Xi^{-1}(0)/S^1$. The term

$$\begin{aligned} \frac{3}{2}\beta H_2(L_{Y_{K_3}}F) &= \frac{3}{2}\beta^2 H_2 L_{Y_{K_3}} \left(\frac{1}{8}(U_1U_4 - V_1V_4) - \frac{1}{2}(H_2V_4 + K_3U_1) \right) \\ &= \frac{3}{2}\beta^2 H_2 \left[\left(-2L_2 \frac{\partial}{\partial K_1} + 2L_1 \frac{\partial}{\partial K_2} - 2K_2 \frac{\partial}{\partial L_1} + 2K_1 \frac{\partial}{\partial L_2} \right. \right. \\ &\quad \left. \left. - 2U_4 \frac{\partial}{\partial U_1} + 2U_1 \frac{\partial}{\partial U_4} - 2V_4 \frac{\partial}{\partial V_1} + 2V_1 \frac{\partial}{\partial V_4} \right) \right] \left(\frac{1}{8}(U_1U_4 - V_1V_4) \right. \\ &\quad \left. - \frac{1}{2}(H_2V_4 + K_3U_1) \right), \text{ see [2, table 1]} \\ &= \frac{3}{2}\beta^2 H_2 \left[\frac{1}{4}(-U_4^2 + U_1^2 + V_4^2 - V_1^2) - H_2V_1 + K_3U_4 \right]. \end{aligned}$$

Next we calculate $\frac{3}{2}\beta \overline{H_2(L_{Y_{K_3}}F)}$. Since $\overline{H_2V_1} = 0 = \overline{K_3U_4}$ we need only calculate the average of U_1^2 , U_4^2 , V_1^2 and V_4^2 . We get $\overline{U_1^2} = \frac{1}{2}(U_1^2 + V_1^2) = \overline{V_1^2}$ and $\overline{U_4^2} = \frac{1}{2}(U_4^2 + V_4^2) = \overline{V_4^2}$. Thus $\frac{3}{2}\beta \overline{H_2(L_{Y_{K_3}}F)} = 0$. So the average $-\frac{3}{2}\beta \overline{L_{Y_F}(H_2K_3)}$ of the first term in expression (13) vanishes on $\Xi^{-1}(0)/S^1$.

Next we calculate the average of the term $L_{Y_F}^2 H_2$ in expression (13) on $\Xi^{-1}(0)/S^1$. We have

$$\begin{aligned} L_{Y_F}^2 H_2 &= -L_{Y_F}(L_{Y_{H_2}}F) \\ &= -\beta L_{Y_F}(U_4V_1 + \frac{1}{2}H_2K_3 + H_2U_4 - K_3V_1), \text{ using (10)} \\ &= \beta \left[\overbrace{(L_{Y_{U_4}}F)V_1}^I + \overbrace{U_4(L_{Y_{V_1}}F)}^{II} - \frac{1}{2} \overbrace{(L_{Y_F}H_2)K_3}^{III} + \frac{1}{2} \overbrace{H_2(L_{Y_{K_3}}F)}^{IV} \right. \\ &\quad \left. - \overbrace{(L_{Y_F}H_2)U_4}^V + \overbrace{H_2(L_{U_4}F)}^{VI} - \overbrace{(L_{Y_{K_3}}F)V_1}^{VII} - \overbrace{K_3(L_{Y_{V_1}}F)}^{VIII} \right]. \end{aligned}$$

We begin by finding

$$L_{Y_{H_2}}F = \beta \left[2V_1 \frac{\partial}{\partial U_1} + 2V_2 \frac{\partial}{\partial U_2} + 2V_3 \frac{\partial}{\partial U_3} + 2V_4 \frac{\partial}{\partial U_4} - 2U_1 \frac{\partial}{\partial V_1} - 2U_2 \frac{\partial}{\partial V_2} \right]$$

$$\begin{aligned}
& -2U_3 \frac{\partial}{\partial V_3} - 2U_4 \frac{\partial}{\partial V_4} \Big] \left(\frac{1}{8}(U_1 U_4 - V_1 V_4) - \frac{1}{2}(H_2 V_4 + K_3 U_1) \right) \\
& = \beta \left[\frac{1}{2}(V_1 U_4 + U_1 V_4) + H_2 U_4 - K_3 V_1 \right]; \\
L_{Y_{K_3}} F & = \beta \left(\left(-2L_2 \frac{\partial}{\partial K_1} + 2L_1 \frac{\partial}{\partial K_2} - 2K_2 \frac{\partial}{\partial L_1} + 2K_1 \frac{\partial}{\partial L_2} - 2U_4 \frac{\partial}{\partial U_1} + 2U_1 \frac{\partial}{\partial U_4} \right. \right. \\
& \quad \left. \left. - 2V_4 \frac{\partial}{\partial V_1} + 2V_1 \frac{\partial}{\partial V_4} \right) \left(\frac{1}{8}(U_1 U_4 - V_1 V_4) - \frac{1}{2}(H_2 V_4 + K_3 U_1) \right) \right) \\
& = \beta \left[\frac{1}{4}(-U_4^2 + U_1^2 + V_4^2 - V_1^2) - H_2 V_1 + K_3 U_4 \right]; \\
L_{Y_{U_4}} F & = \beta \left[\left(-2U_1 \frac{\partial}{\partial K_1} - 2U_3 \frac{\partial}{\partial L_1} + 2U_3 \frac{\partial}{\partial L_2} - 2V_4 \frac{\partial}{\partial H_2} - 2K_3 \frac{\partial}{\partial U_1} + 2L_2 \frac{\partial}{\partial U_2} \right. \right. \\
& \quad \left. \left. + 2V_3 \frac{\partial}{\partial U_3} - 2H_2 \frac{\partial}{\partial V_4} \right) \left(\frac{1}{8}(U_1 U_4 - V_1 V_4) - \frac{1}{2}(H_2 V_4 + K_3 U_1) \right) \right] \\
& = \beta \left[V_4^2 + K_3^2 - \frac{1}{4}(K_3 U_4 - H_2 V_1) + H_2^2 \right]; \\
L_{Y_{V_1}} F & = \beta \left[\left(-2V_2 \frac{\partial}{\partial K_1} - 2V_1 \frac{\partial}{\partial L_2} + 2V_2 \frac{\partial}{\partial L_2} + 2U_2 \frac{\partial}{\partial H_2} + 2H_2 \frac{\partial}{\partial U_2} + 2K_1 \frac{\partial}{\partial V_2} \right. \right. \\
& \quad \left. \left. + 2K_2 \frac{\partial}{\partial V_3} + 2U_4 \frac{\partial}{\partial V_4} \right) \left(\frac{1}{8}(U_1 U_4 - V_1 V_4) - \frac{1}{2}(H_2 V_4 + K_3 U_1) \right) \right] \\
& = \beta \left(-U_2 V_2 - \frac{1}{4} U_4 V_1 \right).
\end{aligned}$$

So the average of term I on $\Xi^{-1}(0)/S^1$ is

$$\begin{aligned}
\overline{\beta(L_{Y_{U_4}} F)V_1} & = \beta^2 \overline{(V_1 V_4^2 + K_3^2 V_1 - \frac{1}{4} K_3 U_1 V_4 + \frac{1}{4} H_2 V_1^2 + H_2^2 V_2)} \\
& = \beta^2 \left(\frac{1}{8} H_2 K_3^2 + \frac{1}{8} H_2 (U_1^2 + V_1^2) \right), \tag{14a}
\end{aligned}$$

since the average of $V_1 V_4^2$, $K_3^2 V_1$, and $H_2^2 V_1$ are each 0.

Term II is

$$\beta U_4 (L_{Y_{V_1}} F) = -\beta^2 (U_2 U_4 V_2 + \frac{1}{4} U_4^2 V_1).$$

So

$$\overline{\beta U_4 (L_{Y_{V_1}} F)} = 0. \tag{14b}$$

For term III, we have already shown that

$$-\frac{\beta}{2} \overline{(L_{Y_F} H_2) K_3} = 0. \tag{14c}$$

and for term IV we have already shown that

$$\frac{\beta}{2} \overline{H_2 (L_{Y_{K_3}} F)} = 0. \tag{14d}$$

Term V is

$$-\beta (L_{Y_F} H_2) U_4 = \beta^2 (U_4^2 V_1 + \frac{1}{2} H_2 K_3 U_4 + H_2 U_4^2 + K_3 U_4 V_1).$$

So

$$-\beta \overline{(L_{Y_F} H_2) U_4} = \beta^2 \overline{(U_4^2 V_1 + \frac{1}{2} H_2 K_3 U_4 + H_2 U_4^2 + K_3 U_4 V_1)}$$

$$= \frac{\beta^2}{2} H_2 (U_4^2 + V_4^2) + \frac{\beta^2}{2} K_3 (U_4 V_1 - U_1 V_4), \quad (14e)$$

since the average of $U_4^2 V_1$ and $H_2 K_3 U_4$ vanish; while $\overline{U_4^2} = \frac{1}{2}(U_4^2 + V_4^2)$ and $\overline{U_4 V_1} = U_4 V_1 - U_1 V_4$.

Term VI is

$$\beta H_2 (L_{Y_{U_4}} F) = \beta^2 H_2 (V_4^2 + K_3^2 - \frac{1}{4} K_3 U_4 + \frac{1}{4} H_2 V_1 + H_2^2).$$

So

$$\begin{aligned} \beta \overline{H_2 (L_{Y_{U_4}} F)} &= \beta^2 \overline{V_4^2} + \beta^2 H_2 K_3^2 + \beta^2 H_2^3 \\ &= \frac{1}{2} \beta^2 H_2 (U_4^2 + V_4^2) + \beta^2 H_2 K_3^2 + \beta^2 H_2^3, \end{aligned} \quad (14f)$$

since $\overline{K_2 U_4} = 0 = \overline{H_2 V_1}$.

Term VII is

$$-\beta (L_{Y_{K_3}} F) V_1 = \beta^2 (\frac{1}{4} (U_4^2 V_1 - U_1^2 V_1 - V_1 V_4^2 + V_1^3) + H_2 V_1^2 + K_3 U_4 V_1).$$

So

$$\begin{aligned} -\beta \overline{(L_{Y_{K_3}} F) V_1} &= \beta^2 H_2 \overline{V_1^2} + \beta^2 K_3 \overline{U_4 V_1} \\ &= \frac{\beta^2}{2} H_2 (U_1^2 + V_1^2) + \frac{\beta^2}{2} K_3 (U_4 V_1 - U_1 V_4). \end{aligned} \quad (14g)$$

Term VIII is

$$-\beta K_3 (L_{Y_{V_1}} F) = \beta^2 K_3 (U_2 V_2 + \frac{1}{4} U_4 V_1).$$

So

$$-\beta \overline{K_3 (L_{Y_{V_1}} F)} = \beta^2 K_3 \overline{U_2 V_2} + \frac{\beta^2}{4} K_3 \overline{U_4 V_1} = \frac{\beta^2}{4} K_3 (U_4 V_1 - U_1 V_4), \quad (14h)$$

since

$$\begin{aligned} \overline{U_2 V_2} &= \frac{1}{\pi} \int_0^\pi (U_2 \cos 2t + V_2 \sin 2t) (-U_2 \sin 2t + V_2 \cos 2t) dt \\ &= \frac{1}{\pi} \int_0^\pi (-\frac{1}{2} U_2^2 \sin 4t + U_2 V_2 \cos^2 2t - U_2 V_2 \sin^2 2t + \frac{1}{2} V_2^2 \sin 4t) dt \\ &= \frac{1}{2} (U_2 V_2 - U_2 V_2) = 0. \end{aligned}$$

Collecting together the results of all the above term calculations gives

$$\begin{aligned} \beta \overline{L_{Y_F}^2 H_2} &= \beta \overline{(L_{Y_{U_4}} F) V_1} - \beta \overline{(L_{Y_F} H_2) U_4} + \beta \overline{H_2 (L_{Y_{U_4}} F)} - \beta \overline{(L_{Y_{K_3}} F) V_1} - \beta \overline{K_3 (L_{Y_{V_1}} F)} \\ &= \beta^2 \left(\left[\frac{1}{8} H_2 K_3^2 + \frac{1}{8} H_2 (U_1^2 + V_1^2) \right] + \left[\frac{1}{2} H_2 (U_4^2 + V_4^2) + \frac{1}{2} K_3 (U_4 V_1 - U_1 V_4) \right] \right. \\ &\quad \left. + \left[\frac{1}{2} H_2 (U_4^2 + V_4^2) + H_2 K_3^2 + H_2^3 \right] + \left[\frac{1}{2} H_2 (U_1^2 + V_1^2) + \frac{1}{2} K_3 (U_4 V_1 - U_1 V_4) \right] \right. \\ &\quad \left. + \frac{1}{4} K_3 (U_4 V_1 - U_1 V_4) \right) \\ &= \beta^2 \left[\left(\frac{9}{8} H_2 K_3^2 + H_2^3 \right) + \frac{5}{8} H_2 (U_1^2 + V_1^2) + H_2 (U_4^2 + V_4^2) + \frac{5}{4} K_3 (U_4 V_1 - U_1 V_4) \right]. \end{aligned}$$

Thus the second order normal form of the regularized Stark Hamiltonian \mathcal{H} on $\Xi^{-1}(0)/S^1$ is

$$\begin{aligned}\mathcal{H}_{\text{nf}}^{(2)} &= H_2 - \frac{3}{2}\varepsilon\beta H_2 K_3 - \frac{1}{2}\varepsilon^2\beta \overline{L_{Y_F}^2} H_2 \\ &= H_2 - \frac{3}{2}\varepsilon\beta H_2 K_3 - \frac{1}{2}\varepsilon^2\beta^2 \left[\frac{9}{8}H_2 K_3^2 + H_2^3 \right. \\ &\quad \left. + \frac{5}{8}H_2(U_1^2 + V_1^2) + H_2(U_4^2 + V_4^2) + \frac{5}{4}K_3(U_4 V_1 - U_1 V_4) \right].\end{aligned}\quad (15)$$

4 The first order normal form of $\mathcal{H}_{\text{nf}}^{(2)}$ on $T_h S_1^3$

Since $L_{X_{H_2}} \mathcal{H}_{\text{nf}}^{(2)} = 0$ by construction, the second order normal form $\mathcal{H}_{\text{nf}}^{(2)}$ (15) is a smooth function on $(H_2^{-1}(h) \cap \Xi^{-1}(0))/S^1 = T_h S_1^3$, the tangent h -sphere bundle of the unit 3-sphere S_1^3 , given by

$$\begin{aligned}\mathcal{H}_{\text{nf}}^{(2)} &= h - \frac{3}{2}\varepsilon\beta h K_3 - \frac{1}{2}\varepsilon^2\beta^2 \left[\frac{9}{8}K_3^2 + h^3 + \frac{5}{8}(U_1^2 + V_1^2) + h(U_4^2 + V_4^2) \right. \\ &\quad \left. + K_3(U_4 V_1 - U_1 V_4) \right] \\ &= -\frac{3}{2}\varepsilon\beta h K_3 - \frac{1}{2}\varepsilon^2\beta^2 \left[\frac{9}{8}K_3^2 + \frac{5}{8}(U_1^2 + V_1^2) + h(U_4^2 + V_4^2) + K_3(U_4 V_1 - U_1 V_4) \right],\end{aligned}$$

dropping the constant $h - \frac{1}{2}\varepsilon^2\beta^2 h^3$. Rescaling time $dt \mapsto -\frac{3}{2}\varepsilon\beta ds$ gives the Hamiltonian

$$\tilde{\mathcal{H}} = hK_3 + \frac{1}{3}\varepsilon\beta \left[\frac{9}{8}K_3^2 + \frac{5}{8}(U_1^2 + V_1^2) + K_3(U_4 V_1 - U_1 V_4) \right] \quad (16)$$

on $T_h S_1^3$.

We now show that the Hamiltonian $\tilde{\mathcal{H}}$ (16) on $T_h S_1^3$ can be normalized again. On $(T\mathbb{R}^4, \omega_4)$ the Hamiltonian

$$K_3(q, p) = \frac{1}{2}(q_3^2 + q_4^2 + p_3^2 + p_4^2 - q_1^2 - q_2^2 - p_1^2 - p_2^2)$$

gives rise to the Hamiltonian vector field X_{K_3} , whose flow is

$$\begin{aligned}\varphi_t^{X_{K_3}}(q, p) &= \\ &= (q_1 \cos t - p_1 \sin t, q_2 \cos t - p_2 \sin t, q_3 \cos t + p_3 \sin t, q_4 \cos t + p_4 \sin t, \\ &\quad q_1 \sin t + p_1 \cos t, q_2 \sin t + p_2 \cos t, -q_3 \sin t + p_3 \cos t, -q_4 \sin t + p_4 \cos t),\end{aligned}$$

which is periodic of period 2π .

The vector field X_{K_3} on $T\mathbb{R}^4$ induces the vector field

$$\begin{aligned}Y_{K_3} &= -2L_2 \frac{\partial}{\partial K_1} + 2L_1 \frac{\partial}{\partial K_2} - 2K_2 \frac{\partial}{\partial L_1} + 2K_1 \frac{\partial}{\partial L_2} \\ &\quad - 2U_4 \frac{\partial}{\partial U_1} + 2U_1 \frac{\partial}{\partial U_4} - 2V_4 \frac{\partial}{\partial V_1} + 2V_1 \frac{\partial}{\partial V_4},\end{aligned}$$

on $\Xi^{-1}(0)/S^1 \subseteq \mathbb{R}^{16}$ with coordinates $(K, L, H_2, \Xi; U, V)$, whose flow

$$\begin{aligned} \varphi_s^{Y_{K_3}}(K, L, H_2, \Xi; U, V) = & (-L_2 \sin 2s + K_1 \cos 2s, L_1 \sin 2s + K_2 \cos 2s, K_3 \\ & - K_2 \sin 2s + L_1 \cos 2s, K_1 \sin 2s + L_2 \cos 2s, L_3, H_2, \Xi; U_1 \cos 2s - U_4 \sin 2s, \\ & U_2, U_3, U_1 \sin 2s + U_4 \cos 2s, V_1 \cos 2s - V_4 \sin 2s, V_2, V_3, V_1 \sin 2s + V_4 \cos 2s) \end{aligned}$$

is periodic of period π . Since $L_{Y_{K_3}}$ maps the ideal of smooth functions which vanish identically on $\Xi^{-1}(0)/S^1$ into itself, Y_{K_3} is a vector field on $\Xi^{-1}(0)/S^1$. Since $L_{X_{K_3}}H_2 = 0$, it follows that Y_{K_3} induces a vector field on $T_h S_1^3$ with periodic flow. So we can normalize again.

To compute the normal form of the Hamiltonian $\tilde{\mathcal{H}}$ (16) on $T_h S_1^3$ we need only calculate the average of the term

$$T = \frac{9}{8}K_3^2 + \frac{5}{8}(U_1^2 + V_1^2) + h(U_4^2 + V_4^2) + K_3(U_4V_1 - U_1V_4)$$

over the flow $\varphi_s^{Y_{K_3}}$. Since $L_{Y_{K_3}}K_3 = 0$ and

$$L_{Y_{K_3}}(U_4V_1 - U_1V_4) = 2U_1V_1 - 2U_4V_4 + 2U_4V_4 - 2U_1V_1 = 0,$$

we need only calculate $\overline{U_1^2}$, $\overline{U_4^2}$, $\overline{V_1^2}$, and $\overline{V_4^2}$. Now

$$\begin{aligned} \overline{U_1^2} &= \frac{1}{\pi} \int_0^\pi (U_1 \cos 2s - U_4 \sin 2s)^2 ds \\ &= \frac{1}{\pi} \int_0^\pi (U_1^2 \cos^2 2s - U_1U_4 \sin 4s + U_4^2 \sin^2 2s) ds = \frac{1}{2}(U_1^2 + U_4^2). \end{aligned}$$

Similarly, $\overline{V_1^2} = \frac{1}{2}(V_1^2 + V_4^2)$, $\overline{U_4^2} = \frac{1}{2}(U_1^2 + U_4^2)$, and $\overline{V_4^2} = \frac{1}{2}(V_1^2 + V_4^2)$. Thus

$$\overline{T} = \left(\frac{9}{8} - h\right)K_3^2 + \frac{13}{16}h(U_1^2 + V_1^2 + U_4^2 + V_4^2). \quad (17)$$

So the first order normal form of $\tilde{\mathcal{H}}$ (16) on $T_h S_1^3$ is

$$\tilde{\mathcal{H}}_{\text{nf}}^{(1)} = hK_3 + \frac{1}{3}\varepsilon\beta\left[\left(\frac{9}{8} - h\right)K_3^2 + \frac{13}{16}(U_1^2 + V_1^2 + U_4^2 + V_4^2)\right] \quad (18)$$

5 The reduced Hamiltonian $\tilde{\mathcal{H}}_{\text{nf}}^{(1)}$ on $S_h^2 \times S_h^2$

The polynomial $U_1^2 + V_1^2 + U_4^2 + V_4^2$ is invariant under the flows $\varphi_t^{X_{H_2}}$, $\varphi_u^{X_\Xi}$, and thus is a polynomial on the orbit space $T_h S_1^3/S^1 = S_h^2 \times S_h^2$, defined by

$$\begin{aligned} K_1^2 + K_2^2 + K_3^2 + L_1^2 + L_2^2 + L_3^2 &= h \\ K_1L_1 + K_2L_2 + K_3L_3 &= 0. \end{aligned}$$

We now find this polynomial. From the explicit description of \mathbb{R}^8/S^1 in (6) it follows that on $(H_2^{-1}(h) \cap \Xi^{-1}(0))/S^1$

$$\begin{aligned}
U_2V_1 - U_1V_2 &= -hK_1 \\
U_3V_1 - U_1V_3 &= -hK_2 \\
U_4V_1 - U_2V_4 &= -hK_3.
\end{aligned}$$

So on $S_h^2 \times S_h^2$

$$\begin{aligned}
h^2(K_1^2 + K_2^2 + K_3^2) &= (U_2V_1 - U_1V_2)^2 + (U_3V_1 - U_1V_3)^2 + (U_4V_1 - U_1V_4)^2 \\
&= (U_1^2 + U_2^2 + U_3^2 + U_4^2)V_1^2 - U_1^2V_1^2 \\
&\quad - 2(U_1V_1)(U_1V_1 + U_2V_2 + U_3V_3 + U_4V_4) + 2U_1^2V_1^2 \\
&\quad + (V_1^2 + V_2^2 + V_3^2 + V_4^2)U_1^2 - U_1^2V_1^2 \\
&= h^2(V_1^2 + U_1^2),
\end{aligned}$$

since $\langle U, U \rangle = h^2$, $\langle V, V \rangle = h^2$, and $\langle U, V \rangle = 0$. Thus $U_1^2 + V_1^2 = K_1^2 + K_2^2 + K_3^2$. Again from the explicit description of $(H_2^{-1}(h) \cap \Xi^{-1}(0))/S^1$ we have

$$\begin{aligned}
U_4V_3 - U_3V_4 &= -hL_1 \\
U_4V_2 - U_2V_4 &= hL_2 \\
U_4V_1 - U_1V_4 &= -hK_3.
\end{aligned}$$

So on $S_h^2 \times S_h^2$

$$\begin{aligned}
h^2(L_1^2 + L_2^2 + K_3^2) &= (U_4V_3 - U_3V_4)^2 + (U_4V_2 - U_2V_4)^2 + (U_4V_1 - U_1V_4)^2 \\
&= (V_1^2 + V_2^2 + V_3^2 + V_4^2)U_4^2 - U_4^2V_4^2 \\
&\quad - 2(U_4V_4)(U_1V_1 + U_2V_2 + U_3V_3 + U_4V_4) + 2U_4^2V_4^2 \\
&\quad + (U_1^2 + U_2^2 + U_3^2 + U_4^2)V_4^2 - U_4^2V_4^2 \\
&= h^2(U_4^2 + V_4^2).
\end{aligned}$$

Thus $U_4^2 + V_4^2 = L_1^2 + L_2^2 + K_3^2$. Consequently

$$U_1^2 + V_1^2 + U_4^2 + V_4^2 = K_1^2 + K_2^2 + 2K_3^2 + L_1^2 + L_2^2$$

$S_h^2 \times S_h^2$. Hence on $S_h^2 \times S_h^2$

$$\begin{aligned}
\widehat{\mathcal{H}} &= \widetilde{\mathcal{H}}_{\text{nf}}^{(1)} = hK_3 + \frac{1}{3}\varepsilon\beta\left[\frac{1}{16}(18 - 3h)K_3^2 + \frac{13}{16}h(K_1^2 + K_2^2 + K_3^2 + L_1^2 + L_2^2)\right] \\
&= hK_3 + \frac{1}{3}\varepsilon\beta\left[\frac{1}{16}(18 - 3h)K_3^2 + \frac{13}{16}h(h^2 - L_3^2)\right]. \tag{19}
\end{aligned}$$

6 The Hamiltonian system $(\widehat{\mathcal{H}}, S_h^2 \times S_h^2, \{ , \})$

Using the coordinates $(\xi, \eta) = ((K + L)/2, (K - L)/2)$ on $\mathbb{R}^3 \times \mathbb{R}^3$, the space of smooth functions on the reduced space $S_h^2 \times S_h^2$, defined by

$$\xi_1^2 + \xi_2^2 + \xi_3^2 = h^2 \quad \text{and} \quad \eta_1^2 + \eta_2^2 + \eta_3^2 = h^2,$$

has a Poisson structure with bracket relations

$$\{\xi_i, \xi_j\} = \sum_{k=1}^3 \epsilon_{ijk} \xi_k, \quad \{\eta_i, \eta_j\} = - \sum_{k=1}^3 \epsilon_{ijk} \eta_k, \quad \{\xi_i, \eta_j\} = 0.$$

Since $\{K_3, L_3\} = 0$, it follows that $\{K_3, \widehat{\mathcal{H}}\} = 0$. Thus the flow $\varphi_r^{Z_{K_3}}$ of the Hamiltonian vector field Z_{K_3} on $(S_h^2 \times S_h^2, \{, \})$ generates an S^1 symmetry of the Hamiltonian system $(\widehat{\mathcal{H}}, S_h^2 \times S_h^2, \{, \})$. So this system is completely integrable.

We reduce this S^1 symmetry as follows. Consider the vector field Z_{K_3} on $\mathbb{R}^3 \times \mathbb{R}^3$ corresponding to the Hamiltonian $K_3 = \frac{1}{2}(\xi_3 + \eta_3)$. Its integral curves satisfy

$$\begin{aligned} \dot{\xi}_1 &= \{\xi_1, K_3\} = \frac{1}{2}\{\xi_1, \xi_3\} = -\frac{1}{2}\xi_2 \\ \dot{\xi}_2 &= \{\xi_2, K_3\} = \frac{1}{2}\{\xi_2, \xi_3\} = \frac{1}{2}\xi_1 \\ \dot{\xi}_3 &= \{\xi_3, K_3\} = 0 \\ \dot{\eta}_1 &= \{\eta_1, K_3\} = \frac{1}{2}\{\eta_1, \eta_3\} = \frac{1}{2}\eta_2 \\ \dot{\eta}_2 &= \{\eta_2, K_3\} = \frac{1}{2}\{\eta_2, \eta_3\} = -\frac{1}{2}\eta_1 \\ \dot{\eta}_3 &= \{\eta_3, K_3\} = 0. \end{aligned}$$

Thus the flow of Z_{K_3} on $\mathbb{R}^3 \times \mathbb{R}^3$ is

$$\begin{aligned} \varphi_t^{Z_{K_3}}(\xi, \eta) &= (\xi_1 \cos t/2 - \xi_2 \sin t/2, \xi_1 \sin t/2 + \xi_2 \cos t/2, \xi_3, \\ &\quad \eta_1 \cos t/2 + \eta_2 \sin t/2, \eta_1 \sin t/2 - \eta_2 \cos t/2, \eta_3), \end{aligned}$$

which preserves $S_h^2 \times S_h^2$ and is periodic of period 4π .

We now determine the space $(S_h^2 \times S_h^2)/S^1$ of orbits of the vector field Z_{K_3} . We use invariant theory. The algebra of polynomials on $\mathbb{R}^3 \times \mathbb{R}^3$, which are invariant under the S^1 action given by the flow $\varphi_t^{Z_{K_3}}$, is generated by

$$\begin{aligned} \sigma_1 &= \xi_1^2 + \xi_2^2 & \sigma_2 &= \eta_1^2 + \eta_2^2 & \sigma_3 &= \xi_1 \eta_2 - \xi_2 \eta_1 \\ \sigma_4 &= \xi_1 \eta_1 + \xi_2 \eta_2 & \sigma_5 &= \frac{1}{2}(\xi_3 + \eta_3) & \sigma_6 &= \frac{1}{2}(\xi_3 - \eta_3), \end{aligned}$$

which are subject to the relation

$$\begin{aligned} \sigma_3^2 + \sigma_4^2 &= (\xi_1 \eta_2 - \xi_2 \eta_1)^2 + (\xi_1 \eta_1 + \xi_2 \eta_2)^2 \\ &= (\xi_1^2 + \xi_2^2)(\eta_1^2 + \eta_2^2) = \sigma_1 \sigma_2, \quad \sigma_1 \geq 0 \text{ \& } \sigma_2 \geq 0. \end{aligned} \quad (20a)$$

In terms of invariants the defining equations of $S_h^2 \times S_h^2$ become

$$\sigma_1 + (\sigma_5 + \sigma_6)^2 = \xi_1^2 + \xi_2^2 + \xi_3^2 = h^2 \quad (20b)$$

$$\sigma_2 + (\sigma_5 - \sigma_6)^2 = \eta_1^2 + \eta_2^2 + \eta_3^2 = h^2 \quad (20c)$$

Eliminating σ_1 and σ_2 from (20a) using (20b) and (20c) gives

$$\sigma_3^2 + \sigma_4^2 = (h^2 - (\sigma_5 + \sigma_6)^2)(h^2 - (\sigma_5 - \sigma_6)^2), \quad |\sigma_5 + \sigma_6| \leq h \ \& \ |\sigma_5 - \sigma_6| \leq h, \quad (21a)$$

which defines $(S_h^2 \times S_h^2)/S^1$ as a semialgebraic variety in \mathbb{R}^4 with coordinates $(\sigma_3, \sigma_4, \sigma_5, \sigma_6)$. Thus the reduced space $(K_3^{-1}(2k) \cap (S_h^2 \times S_h^2))/S^1$ is defined by (21a) and

$$\sigma_5 = k. \quad (21b)$$

Consequently, $(K_3^{-1} \cap (S_h^2 \times S_h^2))(2\ell)/S^1$ is the semialgebraic variety

$$\begin{aligned} \sigma_3^2 + \sigma_4^2 &= (h^2 - (k + \sigma_6)^2)(h^2 - (k - \sigma_6)^2) \\ &= ((h - k)^2 - \sigma_6^2)((h + k)^2 - \sigma_6^2), \quad |\sigma_6| \leq h - |k| \end{aligned} \quad (22)$$

in \mathbb{R}^3 with coordinates $(\sigma_3, \sigma_4, \sigma_6)$. When $0 < |k| < h$ the reduced space (22) is a smooth 2-sphere. When $|k| = h$ it is a point. When $k = 0$ it is a topological 2-sphere with conical singular points at $(0, 0, \pm h)$. These singular points correspond to the fixed points $h(0, 0, \pm 1, 0, 0, \mp 1)$ of the S^1 action on $S_h^2 \times S_h^2$ generated by the flow of the vector field Z_{K_3} .

By (19) the reduced Hamiltonian on $(K_3^{-1}(2k) \cap (S_h^2 \times S_h^2))/S^1$ is

$$\widehat{\mathcal{H}}_{\text{red}} = -\frac{13}{12}\varepsilon\beta\sigma_6^2, \quad (23)$$

using $L_3 = \xi_3 - \eta_3 = 2\sigma_6$, having dropped the constant $hk + \frac{1}{48}\varepsilon\beta[(18 - 3h)k^2 + 13h^3]$.

References

- [1] Richard Cushman, Normal form for Hamiltonian vector fields with periodic flow, in: “Differential geometric methods in mathematical physics”, ed. Sternberg, Reidel, Dordrecht, 1984, 125–145.
- [2] Richard Cushman, The geometry of the Kustaanheimo-Stiefel mapping, [arXiv:2205.08485](https://arxiv.org/abs/2205.08485).
- [3] R.H. Cushman and D.A. Sadovskii, Monodromy in the hydrogen atom in crossed fields, *Physica D* **142** (2000) 166–196.
- [4] J.L. Lagrange, “Mécanique Analytique”, Courcier, Paris (1788).
- [5] G. Lantoine and R.P. Russell, Complete closed-form solutions of the Stark problem, *Celest. Mech. Dyn. Astr* **109** (2011) 333–366.