

AH CONJECTURE FOR CANTOR MINIMAL DIHEDRAL SYSTEMS

EDUARDO SCARPARO

ABSTRACT. The AH Conjecture relates the low-dimensional homology groups of a groupoid with the abelianization of its topological full group. We show that transformation groupoids of minimal actions of the infinite dihedral group on the Cantor set satisfy this conjecture. The proof uses Kakutani-Rokhlin partitions adapted to such systems.

1. INTRODUCTION

In [Mat16], Matui conjectured (AH conjecture) that, given an effective minimal étale groupoid G with unit space homeomorphic to the Cantor set, there is an exact sequence

$$H_0(G) \otimes \mathbb{Z}_2 \xrightarrow{j} [[G]]_{\text{ab}} \xrightarrow{I} H_1(G) \longrightarrow 0,$$

where $H_*(G)$, for $* = 0, 1$, are homology groups of G , $[[G]]$ is the topological full group, $I: [[G]]_{\text{ab}} \rightarrow H_1(G)$ is the *index map* and $j: H_0(G) \otimes \mathbb{Z}_2 \rightarrow [[G]]_{\text{ab}}$ is a map introduced by Nekrashevych in [Nek19]. This conjecture has been verified for several classes of examples, e.g., principal almost finite groupoids ([Mat16]), transformation groupoids of odometers ([Sca20]), graph groupoids ([Mat15] and [NO21b]) and Katsura-Exel-Pardo groupoids ([NO21]).

In [Mat16], Matui also made a conjecture relating homology groups of a groupoid G and the K-theory of the reduced C^* -algebra of G (HK conjecture). In [Sca20] a counterexample was given to this conjecture involving a certain non-free minimal action of the infinite dihedral group $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set (further counterexamples were later provided in [OS20], [OS22] and [Dee22]).

Minimal actions of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set have been studied in several different contexts. For example, in [BEK93], Bratteli, Evans and Kishimoto showed that, if the action is not free, then the associated crossed product is approximately finite-dimensional. In [Tho10], Thomsen showed that the converse of this statement holds. In [BRY18, Example 3], Baake, Roberts and Yassawi gave an example of an aperiodic minimal subshift whose associated \mathbb{Z} -action cannot be extended to an action of $\mathbb{Z} \rtimes \mathbb{Z}_2$. In [Jia21], Jiang showed that continuously orbit equivalent minimal actions of $\mathbb{Z} \rtimes \mathbb{Z}_2$ are conjugate.

In [OS20], Ortega and the author showed that (transformation groupoids of) Cantor minimal dihedral system satisfy the HK conjecture if and only if the action is free. Furthermore, it was shown that such systems are always almost finite. By Matui's result, it follows that, in the case that the action is free, the AH conjecture holds. But the question of whether non-free actions also satisfy it was left open.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 817597).

In this work, we address this problem. The main tool is a technique of Grigorchuk and Medynets ([GM14]) which, given a minimal homeomorphism on the Cantor set, provides a decomposition of the elements of the topological full group as products of elements adapted to Kakutani-Rokhlin partitions.

2. PRELIMINARIES

2.1. Homology of transformation groupoids. In this section, we provide an ad hoc definition of the low-dimensional homology groups of a transformation groupoid. See, for example, [Mat12, Section 3.1] for a more general perspective.

Let X and Y be locally compact totally disconnected Hausdorff spaces. Given a local homeomorphism $\pi: X \rightarrow Y$, let $\pi_*: C_c(X, \mathbb{Z}) \rightarrow C_c(Y, \mathbb{Z})$ be given by

$$\pi_*(f)(y) := \sum_{x \in \pi^{-1}(y)} f(x),$$

for $f \in C_c(X, \mathbb{Z})$ and $y \in Y$. Notice that, given $U \subset X$ compact-open such that $\pi|_U$ is a homeomorphism, we have that $\pi_*(1_U) = 1_{\pi(U)}$.

Let Γ be a discrete group acting by homeomorphisms on X . As a space, the *transformation groupoid* of the action, denoted by $\Gamma \times X$, is the set $\Gamma \times X$ endowed with the product topology. The *range* and *source* maps $r, s: \Gamma \times X \rightarrow X$ are given by $r(g, x) := gx$ and $s(g, x) := x$, for $(g, x) \in \Gamma \times X$. The product of two elements (h, y) and (g, x) is defined if and only if $y = gx$, in which case $(h, gx)(g, x) := (hg, x)$. Inversion is given by $(g, x)^{-1} := (g^{-1}, gx)$. Let $(\Gamma \times X)^{(2)} := \{(a_1, a_2) \in (\Gamma \times X)^2 : s(a_1) = r(a_2)\}$.

A *bisection* is a compact open subset $U \subset \Gamma \times X$ such that $r|_U$ and $s|_U$ are homeomorphisms. Given U and V bisections, the set

$$(1) \quad UV := \{a_1 a_2 : (a_1, a_2) \in (U \times V) \cap (\Gamma \times X)^{(2)}\}$$

is a bisection as well, and so is U^{-1} .

Let $d_0, d_1, d_2: (\Gamma \times X)^{(2)} \rightarrow \Gamma \times X$ be the local homeomorphisms given by

$$d_i(a_1, a_2) := \begin{cases} a_2 & \text{if } i = 0 \\ a_1 a_2 & \text{if } i = 1 \\ a_1 & \text{if } i = 2. \end{cases}$$

Let $\delta_2: C_c((\Gamma \times X)^{(2)}, \mathbb{Z}) \rightarrow C_c(\Gamma \times X, \mathbb{Z})$ be given by $\delta_2 := d_{0*} - d_{1*} + d_{2*}$ and $\delta_1: C_c(\Gamma \times X, \mathbb{Z}) \rightarrow C_c(X, \mathbb{Z})$ by $\delta_1 := s_* - r_*$. Then $\delta_1 \delta_2 = 0$ and we let $H_0(\Gamma \times X) := \frac{C_c(X, \mathbb{Z})}{\text{Im } \delta_1}$ and $H_1(\Gamma \times X) := \frac{\ker \delta_1}{\text{Im } \delta_2}$.

2.2. Cantor minimal dihedral systems. An action of a group Γ on a compact Hausdorff space X is said to be *topologically free* if, for any $g \in \Gamma \setminus \{e\}$, the set $\text{Fix}_g := \{x \in X : gx = x\}$ has empty interior. It is said to be *minimal* if every point has dense orbit. If φ is a homeomorphism on X , then we say that φ is minimal if the associated \mathbb{Z} -action is minimal.

Recall that the *infinite dihedral group* is the group $\mathbb{Z} \rtimes \mathbb{Z}_2$, where the \mathbb{Z}_2 -action is given by multiplication by -1 . Any action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on a space X is given by a pair (φ, σ) such that $\sigma \varphi \sigma^{-1} = \varphi^{-1}$ and $\sigma^2 = \text{Id}_X$. Alternatively, since $\mathbb{Z} \rtimes \mathbb{Z}_2 \simeq \mathbb{Z}_2 * \mathbb{Z}_2$, any action of the infinite dihedral group on X is given by a pair of homeomorphisms

a and b on X such that $a^2 = b^2 = \text{Id}_X$. Moreover, we can identify σ with a , and $\varphi\sigma$ with b .

Remark 2.1. In [OS20, Proposition 2.8], it was observed that any minimal action (φ, σ) of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set is topologically free. Furthermore, if the action is not free, then φ is minimal, and Fix_σ and $\text{Fix}_{\varphi\sigma}$ are disjoint sets, not both empty.

Notice that σ induces a homomorphism

$$\begin{aligned} \sigma_*: H_0(\mathbb{Z} \times X) &\rightarrow H_0(\mathbb{Z} \times X) \\ [f] &\mapsto [f \circ \sigma], \end{aligned}$$

for $f \in C(X, \mathbb{Z})$.

We recall the following computation of homology groups of Cantor minimal dihedral systems:

Theorem 2.2 ([OS20, Theorem 3.6]). *Let $\alpha := (\varphi, \sigma)$ be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X .*

- (i) *If φ is not minimal, then there exists $Y \subset X$ clopen and φ -invariant such that $X = Y \sqcup \sigma(Y)$ and $\varphi|_Y$ is minimal. Furthermore,*

$$\begin{aligned} H_0((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &\simeq H_0(\mathbb{Z} \times Y), \\ H_1((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &\simeq \mathbb{Z}; \end{aligned}$$

- (ii) *If φ is minimal and α is free, then*

$$\begin{aligned} H_0((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &\simeq \mathbb{Z}_2 \oplus (1 + \sigma_*)H_0(\mathbb{Z} \times X), \\ H_1((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &= 0; \end{aligned}$$

- (iii) *If α is not free, then*

$$\begin{aligned} H_0((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &\simeq (1 + \sigma_*)H_0(\mathbb{Z} \times X), \\ (2) \quad H_1((\mathbb{Z} \rtimes \mathbb{Z}_2) \times X) &\simeq C(\text{Fix}_\sigma \sqcup \text{Fix}_{\varphi\sigma}, \mathbb{Z}_2). \end{aligned}$$

Remark 2.3. (i) We do not know any example of minimal action (φ, σ) of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set for which Fix_σ and $\text{Fix}_{\varphi\sigma}$ are not both finite. For odometers and actions coming from irrational rotations this was shown to be the case in [Sca20, Lemma 3.2] and [BEK93, Corollary 4.4], respectively. Let (X, φ) be a two-sided shift space coming from a primitive substitution. Suppose that, given any word $w_1 \dots w_n$ in the language $\mathcal{L}(X)$, it holds that $w_n \dots w_1 \in \mathcal{L}(X)$ as well. Then the map $\sigma: X \rightarrow X$ defined by $\sigma(x)_n := x_{-n}$ is well-defined and induces an action (φ, σ) of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on X . It follows from [DZ00, Theorem 1.1] that Fix_σ and $\text{Fix}_{\varphi\sigma}$ are both finite.

(ii) Let (φ, σ) be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X such that φ is minimal. For odometers and actions coming from irrational rotations, it was shown in [Sca20, Proposition 3.3] and [BEK93, Corollary 4.4], respectively, that $\sigma_* = \text{Id}_{H_0(\mathbb{Z} \times X)}$. In such cases, $1 + \sigma_*$ becomes simply multiplication by 2. Since $H_0(\mathbb{Z} \times X)$ is always torsion-free, it follows that $(1 + \sigma_*)H_0(\mathbb{Z} \times X) \simeq H_0(\mathbb{Z} \times X)$. It would be interesting to find examples in which $\sigma_* \neq \text{Id}_{H_0(\mathbb{Z} \times X)}$.

We will now present an explicit description of the isomorphism in (2) which will be necessary later.

Let Γ be a group acting on a compact Hausdorff space X . Given $f \in C_c(\Gamma \times X, \mathbb{Z})$ and $w \in \Gamma$, we let $f_w \in C(X, \mathbb{Z})$ be given by $f(x) := f(w, x)$, for $x \in X$. If $g \in \Gamma$ and $x \in X$, then $gf_w(x) := f(g^{-1}x, w)$.

Given $w \in \mathbb{Z}_2 * \mathbb{Z}_2$, let $w_1 \dots w_n$ be the reduced expression of w in the letters a, b associated to the generators of $\mathbb{Z}_2 * \mathbb{Z}_2$. The *length* of w is denoted by $|w|$. Given $1 \leq i \leq j \leq n$, let $w_{(i,j)} := w_{i+1} \dots w_n$ if $i < j$ and e otherwise.

Lemma 2.4. *Let (a, b) be a non-free minimal action of $\mathbb{Z}_2 * \mathbb{Z}_2$ on the Cantor set X . The map $R: C_c((\mathbb{Z}_2 * \mathbb{Z}_2) \times X, \mathbb{Z}) \rightarrow C(\text{Fix}_a \sqcup \text{Fix}_b, \mathbb{Z}_2)$ given by*

$$R(f) := \sum_{w \in \mathbb{Z}_2 * \mathbb{Z}_2} \sum_{i=1}^{|w|} (w_{(i,|w|]} f_w) \mathbf{1}_{\text{Fix}_{w_i}} \pmod{2},$$

is well-defined and vanishes on $\text{Im } \delta_2$.

Proof. We will first show that the image of R consists of continuous functions. Given $A \subset X$ clopen and $w \in \mathbb{Z}_2 * \mathbb{Z}_2$, notice that

$$R(\mathbf{1}_{\{w\} \times A}) = \sum_{i=1}^{|w|} \mathbf{1}_{(w_{(i,|w|]} A) \cap \text{Fix}_{w_i}}$$

(the two sides of this equation are viewed as functions taking values in \mathbb{Z}_2). Since $C_c((\mathbb{Z}_2 * \mathbb{Z}_2) \times X)$ is generated by elements of the form $\mathbf{1}_{\{w\} \times A}$, it follows that R takes values in $C(\text{Fix}_a \sqcup \text{Fix}_b, \mathbb{Z}_2)$.

We will now show that R vanishes on $\text{Im } \delta_2$. Given $u, v \in \mathbb{Z}_2 * \mathbb{Z}_2$, there exist $u', x, v' \in \mathbb{Z}_2 * \mathbb{Z}_2$ such that $u = u'x^{-1}$ and $v = xv'$ are in reduced form as well as $u'v'$. Fix $A \subset X$ clopen and let $U := \{u\} \times vA$ and $V := \{v\} \times A$. Then $UV = \{u'v'\} \times A$ and

$$(3) \quad \delta_2(\mathbf{1}_{(U \times V) \cap ((\mathbb{Z}_2 * \mathbb{Z}_2) \times X)}^{(2)}) = \mathbf{1}_U - \mathbf{1}_{UV} + \mathbf{1}_V$$

Furthermore,

$$\begin{aligned} R(\mathbf{1}_U - \mathbf{1}_{UV} + \mathbf{1}_V) &= \sum_{i=1}^{|u|} \mathbf{1}_{(u_{(i,|u|]} vA) \cap \text{Fix}_{u_i}} - \sum_{i=1}^{|u'v'|} \mathbf{1}_{((u'v')_{(i,|u'v'|]} A) \cap \text{Fix}_{(u'v')_i}} \\ &+ \sum_{i=1}^{|v|} \mathbf{1}_{(v_{(i,|v|]} A) \cap \text{Fix}_{v_i}} = \sum_{i=1}^{|u'|} \mathbf{1}_{(u'_{(i,|u'|]} x^{-1}vA) \cap \text{Fix}_{u'_i}} + \sum_{i=1}^{|x|} \mathbf{1}_{((x^{-1})_{(i,|x|]} vA) \cap \text{Fix}_{(x^{-1})_i}} \\ &\quad - \sum_{i=1}^{|u'|} \mathbf{1}_{(u'_{(i,|u'|]} v'A) \cap \text{Fix}_{u'_i}} - \sum_{i=1}^{|v'|} \mathbf{1}_{(v'_{(i,|v'|]} A) \cap \text{Fix}_{v'_i}} + \sum_{i=1}^{|x|} \mathbf{1}_{(x_{(i,|x|]} v'A) \cap \text{Fix}_{x_i}} \\ &+ \sum_{i=1}^{|v'|} \mathbf{1}_{(v'_{(i,|v'|]} A) \cap \text{Fix}_{v'_i}} = \sum_{i=1}^{|x|} \mathbf{1}_{((x^{-1})_{(i,|x|]} xv'A) \cap \text{Fix}_{(x^{-1})_i}} + \sum_{i=1}^{|x|} \mathbf{1}_{(x_{(i,|x|]} v'A) \cap \text{Fix}_{x_i}} \\ &\quad \stackrel{j:=m-i+1}{=} \sum_{j=1}^{|x|} \mathbf{1}_{(x_1 \dots x_{j-1})^{-1} xv'A) \cap \text{Fix}_{x_j}} + \sum_{i=1}^{|x|} \mathbf{1}_{(x_{(i,|x|]} v'A) \cap \text{Fix}_{x_i}} \\ &= \sum_{j=1}^{|x|} \mathbf{1}_{(x_j x_{(j,|x|]} v'A) \cap \text{Fix}_{x_j}} + \sum_{i=1}^{|x|} \mathbf{1}_{(x_{(i,|x|]} v'A) \cap \text{Fix}_{x_i}} = 0, \end{aligned}$$

where the last equality follows from the fact that, given $B \subset X$ and $g \in \mathbb{Z}_2 * \mathbb{Z}_2$, we have $gB \cap \text{Fix}_g = B \cap \text{Fix}_g$.

Since $\text{Im } \delta_2$ is generated by elements as in (3), the conclusion follows. \square

Theorem 2.5. *Let (a, b) be a non-free minimal action of $\mathbb{Z}_2 * \mathbb{Z}_2$ on the Cantor set X . The map $R: H_1((\mathbb{Z}_2 * \mathbb{Z}_2) \ltimes X) \rightarrow C(\text{Fix}_a \sqcup \text{Fix}_b, \mathbb{Z}_2)$ given by*

$$R([f]) := \sum_{w \in \mathbb{Z}_2 * \mathbb{Z}_2} \sum_{i=1}^{|w|} (w_{(i, |w|]} f_w) 1_{\text{Fix}_{w_i}} \pmod{2},$$

for $f \in \ker \delta_1$, is an isomorphism.

Proof. Surjectivity: Given $U' \subset \text{Fix}_a$ clopen in Fix_a , there exists an open set $U \subset X$ such that $U' = U \cap \text{Fix}_a$. By compactness of U' , we can take U to be clopen. By considering $U \cap a(U)$, we may also assume that $U = a(U)$. Then $R([1_{\{a\} \times U}]) = 1_{U \cap \text{Fix}_a} = 1_{U'}$. By arguing in the same way for b , surjectivity of R follows.

Injectivity: Fix $f \in \ker \delta_1$ such that $R([f]) = 0$, and we will show that $[f] = 0$. Given $A \subset X$ clopen and $g, h \in \mathbb{Z}_2 * \mathbb{Z}_2$, we have by (3) that

$$(4) \quad [1_{\{h\} \times hA} - 1_{\{gh\} \times A} + 1_{\{g\} \times A}] = 0.$$

In particular, $[1_{\{e\} \times A}] = 0$.

By using (4) repeatedly times, we obtain $f_1, f_2 \in C_c((\mathbb{Z}_2 * \mathbb{Z}_2) \ltimes X, \mathbb{Z})$ such that $\text{supp } f_1 \subset \{a\} \times X$, $\text{supp } f_2 \subset \{b\} \times X$ and $[f] = [f_1 + f_2]$. We claim that, in this case, $\delta_1(f_1) = \delta_1(f_2) = 0$. Indeed, we can write $f_1 = \sum_{n \in \mathbb{Z}} n 1_{\{a\} \times A_n}$ and $f_2 = \sum_{n \in \mathbb{Z}} n 1_{\{b\} \times B_n}$ for families of disjoint clopen sets $(A_n)_{n \in \mathbb{Z}}, (B_n)_{n \in \mathbb{Z}}$. We have

$$\begin{aligned} 0 &= \delta_1(f_1 + f_2) = \sum n(1_{A_n} - 1_{aA_n}) + \sum n(1_{B_n} - 1_{bB_n}) \\ &\implies h := \sum n(1_{A_n} - 1_{aA_n}) = \sum n(1_{bB_n} - 1_{B_n}). \end{aligned}$$

Hence, $h \circ a = -h = h \circ b$ and $h \circ (ba) = h$. Since the action is not free, it follows from Remark 2.1 that ba is a minimal homeomorphism. Therefore, h is a constant function. Since the action is not free, we have that either a or b have a fixed point x . In particular, $h(x) = 0$ and $h = 0$. Hence, $\delta_1(f_1) = \delta_1(f_2) = 0$.

Let us now show that $[f_1] = 0$. Since $\delta_1(f_1) = 0$, we have that $\sum n 1_{aA_n} = \sum n 1_{A_n}$. Hence $a(A_n) = A_n$ for every $n \in \mathbb{Z} \setminus \{0\}$. By applying (4), we conclude that for any $D \subset X$ clopen such that $a(D) = D$, it holds that $[2(1_{\{a\} \times D})] = 0$. Hence, there exists $A \subset X$ clopen such that $a(A) = A$ and $[f_1] = [1_{\{a\} \times A}]$. Since $R([f]) = 0$, we obtain that $A \cap \text{Fix}_a = \emptyset$. In particular, there is A' clopen set such that $A = A' \sqcup a(A')$.

Finally, $[f_1] = [1_{\{a\} \times a(A')} + 1_{\{a\} \times A'}] = 0$ by (4) again.

By arguing in the same way for f_2 , we conclude that $[f] = 0$. \square

2.3. Topological full groups. Let α be an action of a group Γ on the Cantor set X . We say that a bisection $U \subset \Gamma \ltimes X$ is *full* if $r(U) = s(U) = X$. In this case, there exist $g_1, \dots, g_n \in \Gamma$ and $A_1, \dots, A_n \subset X$ clopen sets such that $X = \bigsqcup A_i = \bigsqcup g_i A_i$ and $U = \bigsqcup \{g_i\} \times A_i$. Moreover, we can associate to U a homeomorphism θ_U on X given by $\theta_U(x) = g_i x$, for $x \in A_i$. Given full bisections

U and V , it holds that $\theta_{UV} = \theta_U \theta_V$ and $\theta_{U^{-1}} = (\theta_U)^{-1}$. The *topological full group* of α is $[[\alpha]] := \{\theta_U : U \text{ is a full bisection}\}$.

From now on, suppose Γ is countable and α is minimal and topologically free. In this case, the map The *index map* $I: [[\alpha]]_{\text{ab}} \rightarrow H_1(\Gamma \ltimes X)$ is given by $I([\theta_U]) := [1_U]$.

Given $g \in \Gamma$ and $A \subset X$ a clopen set such that $A \cap gA = \emptyset$, let $\tau_{g,A} \in [[\alpha]]$ be the homeomorphism given by

$$(5) \quad \tau_{g,A}(x) := \begin{cases} gx & \text{if } x \in A \\ g^{-1}x & \text{if } x \in gA \\ x & \text{otherwise.} \end{cases}$$

Let $S(\alpha)$ be the subgroup of $[[\alpha]]$ generated by such elements. By [Nek19, Theorem 7.2], there exists a unique map $j: H_0(\Gamma \ltimes X) \rightarrow [[\alpha]]_{\text{ab}}$ which, given $g \in \Gamma$ and $A \subset X$ as above, maps $[1_A] \otimes 1$ to $[\tau_{g,A}]$. It is easy to see that $I \circ j = 0$, and Matui's AH conjecture ([Mat15, Conjecture 2.9]) predicts that the following sequence is exact:

$$H_0(\Gamma \ltimes X) \otimes \mathbb{Z}_2 \xrightarrow{j} [[\alpha]]_{\text{ab}} \xrightarrow{I} H_1(\Gamma \ltimes X) \longrightarrow 0.$$

3. AH CONJECTURE FOR CANTOR MINIMAL DIHEDRAL SYSTEMS

In this section, we show that minimal actions of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set satisfy the AH conjecture. We will use the following result from [BEK93] which provides *Kakutani-Rokhlin partitions* for such systems. All the intervals consist of integers.

Proposition 3.1 ([BEK93]). *Let (φ, σ) be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X such that $\varphi\sigma$ admits a fixed point. Given a partition \mathcal{P} of X and $N \in \mathbb{N}$, there exist clopen sets $Y_1, \dots, Y_K \subset X$ and integers J_1, \dots, J_K such that*

- (i) *For $k \in [1, K]$, $J_k \geq 2N$;*
- (ii) *The sets $\{\varphi^i(Y_k) : k \in [1, K], i \in [0, J_k]\}$ are disjoint with union X . Moreover, this partition refines \mathcal{P} ;*
- (iii) *Let $Y := \bigsqcup_{k=1}^K Y_k$. Then $\sigma(Y) = \varphi^{-1}(Y)$ and, for $i \in [-N, N]$, there exists $P_i \in \mathcal{P}$ such that $\varphi^i(Y) \subset P_i$;*
- (iv) *There exists an involutive bijection $\tau: [1, K] \rightarrow [1, K]$ such that, for $l \in \mathbb{Z}$ and $k \in [1, K]$, we have $J_k = J_{\tau(k)}$ and $\sigma\varphi^l(Y_k) = \varphi^{J_k - l - 1}(Y_{\tau(k)})$.*

Proof. Items (i), (ii) and (iii) are the content of [BEK93, Proposition 1.2]. Let us show (iv). By [BEK93, Remark 1.3], given $k \in [1, K]$, there exists $\tau(k) \in [1, K]$ such that $\sigma(Y_k) = \varphi^{J_{\tau(k)} - 1}(Y_{\tau(k)})$. Furthermore, $J_k = J_{\tau(k)}$ and $\sigma(Y_{\tau(k)}) = \varphi^{J_k - 1}(Y_k)$. In particular, given $l \in \mathbb{Z}$, we have

$$\sigma\varphi^l(Y_k) = \varphi^{-l}\sigma(Y_k) = \varphi^{J_k - l - 1}(Y_{\tau(k)}).$$

□

Given $A \subset X$ clopen and $h \in [[(\varphi, \sigma)]]$ such that $h(A) = A$, let $h_A \in [[(\varphi, \sigma)]]$ be given by $h_A(x) := h(x)$, for $x \in A$, and $h_A(x) := x$ otherwise.

Lemma 3.2. *Let $n \in \mathbb{Z}$ and $U \subset X$ clopen such that $\varphi^n \sigma(U) = U$. If $U \cap \text{Fix}_{\varphi^n \sigma} = \emptyset$, then $(\varphi^n \sigma)_U \in S(\varphi, \sigma)$.*

Proof. Take $V \subset U$ clopen such that $U = V \sqcup \varphi^n \sigma(V)$. Then $(\varphi^n \sigma)_U = \tau_{\varphi^n \sigma, V}$, as in (5). \square

The idea behind the proof of the next result is a technique introduced in [GM14, Theorem 4.7] for factorizing an element of the topological full group of a Cantor minimal \mathbb{Z} -system as a product of elements adapted to a Kakutani-Rokhlin partition of the system.

Theorem 3.3. *Let (φ, σ) be a non-free minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . Given $Q \in [[(\varphi, \sigma)]]$, there exist $n_1, \dots, n_l \in \mathbb{Z}$ and U_1, \dots, U_l disjoint clopen sets such that $\varphi^{n_i} \sigma(U_i) = U_i$ for $1 \leq i \leq l$, and $Q \prod_{i=1}^l (\varphi^{n_i} \sigma)_{U_i} \in [[\varphi]]$.*

Proof. We claim that we can assume that $\varphi \sigma$ has a fixed point. If it does not, then σ does and we can consider the action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ given by $(\psi, \mu) := (\varphi^{-1}, \varphi \sigma)$. For $r \in \mathbb{Z}$, we have $\psi^r \mu = \varphi^{-r+1} \sigma$ and, in particular, $\psi \mu = \sigma$ has a fixed point. Moreover, $[[\psi, \mu]] = [[\varphi, \sigma]]$. Hence we can assume that $\varphi \sigma$ has a fixed point.

Let A_1, \dots, A_n be disjoint clopen sets with union X and $(m_1, p_1), \dots, (m_n, p_n) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ such that, for $1 \leq i \leq n$ and $x \in A_i$, we have $Q(x) = \varphi^{m_i} \sigma^{p_i}(x)$. Denote by \mathcal{P} the partition determined by the sets A_i . Also let $N := \max\{|m_i| : 1 \leq i \leq n\}$.

Adopt the notation of Proposition 3.1. Given $k \in [1, K]$ and $i \in [0, J_k]$, let $f(k, i) := (f_1(k, i), f_2(k, i)) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ be given by $f(k, i) := (m_j, p_j)$ if $\varphi^i(Y_k) \subset A_j$. Notice that, for any $k, l \in [1, K]$ and $i \in [0, N]$, we have $f(k, i) = f(l, i)$ and $f(k, J_k - i - 1) = f(l, J_l - i - 1)$.

Given $k \in [1, K]$, let $S_k := \{i \in [N, J_k - N] : f(k, i) = 1\}$. We will show that $|S_k| = |S_{\tau(k)}|$. Let $T_k := \{f_1(\tau(k), n) - n - 1 : n \in S_{\tau(k)}\}$. Since, given $n \in S_{\tau(k)}$, we have $Q(\varphi^n(Y_{\tau(k)})) = \varphi^{f_1(\tau(k), n) + J_k - n - 1}(Y_k)$ and $0 \leq f_1(\tau(k), n) + J_k - n - 1 < J_k$, it follows that $|T_{\tau(k)}| = |S_{\tau(k)}|$.

Identify $[0, J_k]$ with the group \mathbb{Z}_{J_k} and let $F: \mathbb{Z}_{J_k} \setminus S_k \rightarrow \mathbb{Z}_{J_k}$ be the function given by

$$F(t) := \begin{cases} f_1(k, t) + t & \text{if } f_2(k, t) = 0 \\ f_1(k, t) - t - 1 & \text{if } f_2(k, t) = 1. \end{cases}$$

We will show that $\text{Im } F \supset \mathbb{Z}_{J_k} \setminus T_{\tau(k)}$. Given $m \in \mathbb{Z}_{J_k} \setminus T_k$, let $x \in X$ such that $Q(x) \in \varphi^m(Y_k)$. Take $l \in [1, K]$ and $i \in [0, J_l]$ such that $x \in \varphi^i(Y_l)$.

Case $l = k$: In this case, $i \notin S_k$ and $F(i) = m$.

Case $l = \tau(k)$: Since $m \notin T_k$, we have $i \notin [N, J_k - N]$, hence $f(k, i) = f(\tau(k), i)$ and $F(i) = m$.

Case $l \notin \{k, \tau(k)\}$: Notice that $i \notin [N, J_l - N]$. Suppose first $i \in [0, N]$. If $f_2(l, i) = 0$, then $f_1(l, i) + i < 0$ and $f_1(l, i) + i = m \pmod{J_k}$, hence $F(i) = m$. If $f_2(l, i) = 1$, then $f_1(l, i) + J_l - i - i \geq J_l$ and $f_1(l, i) - i - 1 = m$. In particular, $F(i) = m$.

Finally, assume $i \in [J_l - N, J_l]$. If $f_2(l, i) = 0$, we have $f_1(l, i) + i - J_l = m$ and $F(J_k - J_l + i) = f_1(k, J_k - J_l + i) + J_k - J_l + i = m \pmod{J_k}$. If $f_2(l, i) = 1$, we have $f_1(l, i) + J_l - i - 1 = m \pmod{J_k}$, hence $F(J_k - J_l + i) = f_1(k, J_k - J_l + i) - J_k + J_l - i - 1 = m \pmod{J_k}$.

This finishes the proof that $\text{Im } F \supset \mathbb{Z}_{J_k} \setminus T_{\tau(k)}$. Hence, $|S_k| \leq |T_{\tau(k)}| = |S_{\tau(k)}|$. Since k was arbitrary, we get $|S_k| = |S_{\tau(k)}|$ for any $k \in [1, K]$.

Let $\Lambda_1 := \{i \in [-N, N] : f_2(1, i) = 1\}$. Given $\lambda \in \Lambda_1$, notice that

$$\varphi^{2\lambda+1} \sigma(\varphi^\lambda(Y)) = \varphi^{2\lambda+1} \varphi^{-\lambda-1}(Y) = \varphi^\lambda(Y).$$

Take $\Lambda_2 \subset [1, K]$ minimal such that $[1, K] = \Lambda_2 \cup \tau(\Lambda_2)$. For each $k \in \Lambda_2$, let $\alpha_k: S_k \rightarrow S_{\tau(k)}$ be the order-preserving bijection, and, for $i \in S_k$, let $n_{k,i} := -J_k + i + \alpha_k(i) + 1$ and $U_{k,i} := \varphi^i(Y_k) \cup \varphi^{\alpha_k(i)}(Y_{\tau(k)})$ (so that, if $\tau(k) = k$, then $n_{k,i} = J_k + 2i + 1$ and $U_{k,i} = \varphi^i(Y_k)$). Notice that, for each $k \in \Lambda_2$ and $i \in S_k$, it holds that $\varphi^{n_{k,i}}\sigma(U_{k,i}) = U_{k,i}$.

Finally,

$$Q \prod_{\lambda \in \Lambda_1} (\varphi^{2\lambda+1}\sigma)_{\varphi^\lambda(Y)} \prod_{\substack{k \in \Lambda_2 \\ i \in S_k}} (\varphi^{n_{k,i}}\sigma)_{U_{k,i}} \in [[\varphi]]$$

□

Corollary 3.4. *Let (φ, σ) be a non-free minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . Given $Q \in [[(\varphi, \sigma)]]$, there exist $m_1, \dots, m_l \in \mathbb{Z}$ and V_1, \dots, V_l clopen sets such that $\varphi^{m_i}\sigma(V_i) = V_i$ for $1 \leq i \leq l$, and $Q \prod_{i=1}^l (\varphi^{m_i}\sigma)_{V_i} \in \mathcal{S}(\varphi)$.*

Proof. Recall that we can identify $H_1(\mathbb{Z} \rtimes X)$ with \mathbb{Z} in such a way that the index map $I: [[\varphi]]_{ab} \rightarrow \mathbb{Z}$ satisfies $I([\varphi]) = 1$ (see, e.g., [Nyl20, Proposition 3.3.1]). In particular, by [Mat12, Lemma 6.3] and [Mat16, Theorem 3.3.(4)], given $T \in [[\varphi]]$, there exists $S \in \mathcal{S}(\varphi)$ such that $T = S\varphi^{I(T)}$. Since $\varphi^{I(T)} = ((\varphi\sigma)\sigma)^{I(T)}$, the result follows from Theorem 3.3. □

Theorem 3.5. *Minimal actions of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set satisfy the AH conjecture.*

Proof. By [OS20, Theorem 2.10] and [Mat16, Theorem 3.6], if the action is free, then it satisfies the AH conjecture. Furthermore, surjectivity of the index map in general follows from [Mat12, Theorem 7.5] (alternatively, this can be easily checked by using Theorem 2.5).

Let (φ, σ) be a non-free minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . Given $Q \in [[(\varphi, \sigma)]]$, denote the class of Q in $\frac{[[(\varphi, \sigma)]]_{ab}}{\text{Im } j}$ by $[Q]$. Since $\text{Im } j \subset \ker I$, we have that the index map factors through $\frac{[[(\varphi, \sigma)]]_{ab}}{\text{Im } j}$. Assume that $I([Q]) = 0$, and we will show that $[Q] = 0$.

By Corollary 3.4, there exist clopen sets V_1, \dots, V_l and $m_1, \dots, m_l \in \mathbb{Z}$ such that $\varphi^{m_i}\sigma(V_i) = V_i$ for $1 \leq i \leq l$ and $[Q] = \sum_{i=1}^l [(\varphi^{m_i}\sigma)_{V_i}]$.

Given $A \subset X$ clopen and $n, m \in \mathbb{Z}$ such that $\varphi^m\sigma(A) = A$, notice that $\varphi^{-n}(\varphi^m\sigma)_A\varphi^n = (\varphi^{-2n+m}\sigma)|_{\varphi^{-n}(A)}$. By choosing, for $1 \leq i \leq l$, $n_i \in \mathbb{Z}$ such that $-2n_i + m_i \in \{0, 1\}$, we obtain clopen sets $A_1, \dots, A_k, B_1, \dots, B_k$ such that $\sigma(A_i) = A_i$ and $\varphi\sigma(B_i) = B_i$ for $1 \leq i \leq k$, and $[Q] = \sum_{i=1}^k [(\sigma)_{A_i}] + [(\varphi\sigma)_{B_i}]$.

Let $C_a := \sum_{i=1}^k (\sigma)_{A_i}$ and $C_b := \sum_{i=1}^k (\varphi\sigma)_{B_i}$. Since $I([Q]) = 0$, it follows from Theorem 2.5 that $I(C_a) = I(C_b) = 0$.

By taking intersections and symmetric differences of the sets A_1, \dots, A_k , we obtain W_1, \dots, W_n disjoint clopen sets such that $\sigma(W_j) = W_j$ for $1 \leq j \leq n$ and such that, for $1 \leq i \leq k$, there is $F_i \subset [1, k]$ such that $A_i = \bigsqcup_{j \in F_i} W_j$. In particular, $C_a = \sum_{i=1}^n \sum_{j \in F_i} [(\sigma)_{W_j}]$. Since each $(\sigma)_{W_j}$ is involutive, we conclude that there is $F \subset [1, k]$ such that $C_a = \sum_{j \in F} [(\sigma)_{W_j}]$. Since $0 = I(C_a) = \sum_{j \in F} 1_{W_j \cap \text{Fix}_\sigma}$, it follows that $W_j \cap \text{Fix}_\sigma = \emptyset$ for any $j \in F$. By using Lemma 3.2, we conclude that $C_a = 0$.

By arguing in an analogous way for $\sum_{i=1}^k [(\varphi\sigma)_{B_i}]$, we conclude that $C_b = 0$ and $[Q] = 0$.

□

In [GPS99, Section 5], it was shown that topological full groups of Cantor minimal \mathbb{Z} -systems are *indicable* (i.e., admit a homomorphism onto \mathbb{Z}). The following is an immediate consequence of Theorems 2.2 and 3.5.

Corollary 3.6. *Let (φ, σ) be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set. Then $[[(\varphi, \sigma)]]$ is indicable if and only if φ is not minimal.*

REFERENCES

- [BEK93] O. Bratteli, D. E. Evans, and A. Kishimoto, *Crossed products of totally disconnected spaces by $\mathbb{Z}_2 * \mathbb{Z}_2$* , Ergodic Theory Dynam. Systems **13** (1993), no. 3, 445–484. ↑1, 3, 6
- [BRY18] M. Baake, J. A. G. Roberts, and R. Yassawi, *Reversing and extended symmetries of shift spaces*, Discrete Contin. Dyn. Syst. **38** (2018), no. 2, 835–866. ↑1
- [Dee22] R. J. Deeley, *A counterexample to the HK-conjecture that is principal*, Ergodic Theory and Dynamical Systems (2022), 1–18. ↑1
- [DZ00] D. Damanik and D. Zare, *Palindrome complexity bounds for primitive substitution sequences*, Discrete Math. **222** (2000), no. 1-3, 259–267. ↑3
- [GM14] R. I. Grigorchuk and K. S. Medinets, *On the algebraic properties of topological full groups*, Mat. Sb. **205** (2014), no. 6, 87–108. ↑2, 7
- [GPS99] T. Giordano, I. F. Putnam, and C. F. Skau, *Full groups of Cantor minimal systems*, Israel J. Math. **111** (1999), 285–320. ↑9
- [Jia21] Y. Jiang, *On continuous orbit equivalence rigidity for virtually cyclic group actions*, arXiv:2106.06221 (2021). To appear in Groups, Geometry and Dynamics. ↑1
- [Mat12] H. Matui, *Homology and topological full groups of étale groupoids on totally disconnected spaces*, Proc. Lond. Math. Soc. (3) **104** (2012), no. 1, 27–56. ↑2, 8
- [Mat15] ———, *Topological full groups of one-sided shifts of finite type*, J. Reine Angew. Math. **705** (2015), 35–84. ↑1, 6
- [Mat16] ———, *Étale groupoids arising from products of shifts of finite type*, Adv. Math. **303** (2016), 502–548. ↑1, 8
- [Nek19] V. Nekrashevych, *Simple groups of dynamical origin*, Ergodic Theory Dyn. Syst. **39** (2019), no. 3, 707–732. ↑1, 6
- [NO21a] P. Nyland and E. Ortega, *Katsura-Exel-Pardo groupoids and the AH conjecture*, J. Lond. Math. Soc. (2) **104** (2021), no. 5, 2240–2259. ↑1
- [NO21b] ———, *Matui’s AH conjecture for graph groupoids*, Doc. Math. **26** (2021), 1679–1727. ↑1
- [Nyl20] P. K. Nyland, *Ample groupoids and their topological full groups*, Ph.D. Thesis, NTNU, 2020. Available at <https://hdl.handle.net/11250/2678011>. ↑8
- [OS20] E. Ortega and E. Scarparo, *Almost finiteness and homology of certain non-free actions*, arXiv:2007.02329 (2020). To appear in Groups, Geometry and Dynamics. ↑1, 3, 8
- [OS22] E. Ortega and A. Sanchez, *The homology of the groupoid of the self-similar infinite dihedral group*, Math. Scand. **128** (2022), no. 2. ↑1
- [Sca20] E. Scarparo, *Homology of odometers*, Ergodic Theory Dynam. Systems **40** (2020), no. 9, 2541–2551. ↑1, 3
- [Tho10] K. Thomsen, *The homoclinic and heteroclinic C^* -algebras of a generalized one-dimensional solenoid*, Ergodic Theory Dynam. Systems **30** (2010), no. 1, 263–308. ↑1

EDUARDO SCARPARO, UNIVERSITY OF GLASGOW, UNITED KINGDOM
 Email address: eduardo.scarparo@glasgow.ac.uk