

Dispersing Obnoxious Facilities on Graphs by Rounding Distances

Tim A. Hartmann 

Department of Computer Science, RWTH Aachen University, Germany
hartmann@algo.rwth-aachen.de

Stefan Lendl 

Department of Operations and Information Systems, University of Graz, Austria

Abstract

We continue the study of δ -dispersion, a continuous facility location problem on a graph where all edges have unit length and where the facilities may also be positioned in the interior of the edges. The goal is to position as many facilities as possible subject to the condition that every two facilities have distance at least δ from each other.

Our main technical contribution is an efficient procedure to ‘round-up’ distance δ . It transforms a δ -dispersed set S into a δ^* -dispersed set S^* of same size where distance δ^* is a slightly larger rational $\frac{a}{b}$ with a numerator a upper bounded by the longest (not-induced) path in the input graph.

Based on this rounding procedure and connections to the distance- d independent set problem we derive a number of algorithmic results. When parameterized by treewidth, the problem is in XP. When parameterized by treedepth the problem is FPT and has a matching lower bound on its time complexity under ETH. Moreover, we can also settle the parameterized complexity with the solution size as parameter using our rounding technique: δ -DISPERSION is FPT for every $\delta \leq 2$ and W[1]-hard for every $\delta > 2$.

Further, we show that δ -dispersion is NP-complete for every fixed irrational distance δ , which was left open in a previous work.

2012 ACM Subject Classification Theory of computation \rightarrow Discrete optimization; Mathematics of computing \rightarrow Graph algorithms; Theory of computation \rightarrow Parameterized complexity and exact algorithms; Theory of computation \rightarrow Problems, reductions and completeness

Keywords and phrases facility location, parameterized complexity, packing

Digital Object Identifier 10.4230/LIPIcs...1

1 Introduction

We study the algorithmic behavior of a continuous dispersion problem. Consider an undirected graph G , whose edges have unit length. Let $P(G)$ be the continuum set of points on all the edges and vertices. For two points $p, q \in P(G)$, we denote by $d(p, q)$ the length of a shortest path containing p and q in the underlying metric space. A subset $S \subseteq P(G)$ is δ -dispersed for some positive real number δ , if every distinct points $p, q \in S$ have distance at least $d(p, q) \geq \delta$. Our goal is, for a given graph G and a positive real number δ , to compute a maximum cardinality subset $S \subseteq P(G)$ that is δ -dispersed. We denote by $\delta\text{-disp}(G)$ the maximum size of a δ -dispersed set of G . The decision problem DISPERSION asks for a δ -dispersed set of size at least k , where additionally integer $k \geq 0$ is part of the input. When δ is fixed and not part of the input, we refer to the problem as δ -DISPERSION.

1.1 Known and related results

The area of obnoxious facility location goes back to seminal articles of Goldman & Dearing [6] and Church & Garfinkel [3]. The area includes a wide variety of objectives and models.



© Tim A. Hartmann and Stefan Lendl;
licensed under Creative Commons License CC-BY
Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

For example, purely geometric variants have been studied by Abrevaya & Segal [1], Ben-Moshe, Katz & Segal [2], and Katz, Kedem & Segal [13]. Recently, van Ee studied the approximability of a generalized covering problem in a metric space that also involves dispersion constraints [18]. Another direction is a graph-theoretic model, where every edge of the given graph G is rectifiable and has some individual length. Tamir discusses the complexity and approximability of several optimization problems. For example, when G is a tree, then a δ -dispersed set can be computed in polynomial time [17]. Another task is to place a single obnoxious facility in a network while maximizing, for example, the smallest distance from facility to certain clients, as studied by Segal [15].

In a previous work, the complexity of DISPERSION was studied for every rational distance δ . When δ is a rational number with numerator 1 or 2, the problem is polynomial time solvable, while it is NP-complete for all other rational values of δ [7, 8]. The complexity when δ is irrational was left as an open problem.

A closely related facility location problem is δ -covering. The objective is to place as few locations as possible on $P(G)$ subject to the condition that any point in $P(G)$ is in distance at most δ to a placed location. This problem is polynomial time solvable whenever δ is a unit fraction, while it is NP-hard for all non unit fractions δ [9]. Furthermore, the parameterized complexity with the parameter solution size k is studied. δ -covering is fixed parameter tractable when $\delta < \frac{3}{2}$, while for $\delta \geq \frac{3}{2}$ the problem is W[2]-complete [9]. Tamir [16] showed that for δ -covering only certain distances δ are of interest. For every amount of points p the distance $\max\{\delta^* : |\text{cover}^{\delta^*}(G)| = p\}$ is of the form $\frac{L'}{2^{p'}}$ where $p' \in \{1, \dots, p\}$ and L' is roughly at most twice the length of a non-induced path in G .

1.2 Our contribution

Our main technical contribution is an efficient and constructive rounding procedure. Given a δ -dispersed set S for some distance value $\delta > 0$, it transforms S into a δ^* -dispersed set S^* of equal size with a slightly larger well-behaving distance value $\delta^* \geq \delta$. The new distance δ^* is a rational $\frac{a}{b}$ with small numerator a . More precisely, the numerator is upper bounded by the length of the longest (not-induced) path L , hence upper bounded asymptotically by the number of vertices n of the input graph (see Section 5).

Our second technical contribution relates the optimal solution for distance δ and $\frac{\delta}{\delta+1}$ for $\delta \leq 3$. A δ -dispersed set translates to a $\frac{\delta}{\delta+1}$ -dispersed set by placing one more point on every edge, and vice versa by removing one point (see Section 3).

Further we explore a connection of DISPERSION and an independent set problem (see Section 4). The combination of that connection with our technical contributions yields several algorithmic results for DISPERSION (see Section 6 and Section 7):

- DISPERSION is NP-hard even for chordal graphs of diameter 4.
- DISPERSION is FPT for the graph parameter treedepth $\text{td}(G)$ with a run time matching a lower bound under ETH. We complement this result by showing that δ -DISPERSION is W[1]-hard for the slightly more general graph parameter pathwidth $\text{pw}(G)$, even for the combined parameter $\text{pw}(G) + k$. Similarly, δ -DISPERSION is W[1]-hard for the graph parameter $\text{fvs}(G)$, the minimum size of a feedback vertex set.
- DISPERSION is XP for the parameter treewidth $\text{tw}(G)$, with a running time of $(2L + 2)^{\text{tw}(G)} n^{\mathcal{O}(1)}$, where n is the number of vertices and L is an upper bound on the length of the longest path in G . We complement this result by the more general lower bound of $n^{\mathcal{O}(\text{tw}(G) + \sqrt{k})}$, assuming ETH. It implies the lower bound of $L^{\mathcal{O}(\text{tw}(G) + \sqrt{k})}$ since $L \leq n$. Note that a mere lower bound of $L^{\mathcal{O}(\text{tw}(G) + \sqrt{k})}$ would not exclude an $n^{\mathcal{O}(\text{tw}(G))}$ -algorithm.

In addition, we completely resolve the complexity of δ -dispersion, by showing NP-hardness for irrational δ (see Section 8). We also study the parameterized complexity when parameterized by the solution size k . The problem is W[1]-hard when $\delta > 2$, and FPT otherwise. Thus, there is a sharp threshold at $\delta = 2$ where the complexity jumps from FPT to W[1]-hard (see Section 9).

We mark statements whose proof can be found in the appendix with “(★)”.

2 Preliminaries

We use the word *vertex* in the graph-theoretic sense, while we use the word *point* to denote the elements of the geometric structure $P(G)$. As an input for δ -dispersion, we consider graphs G that are undirected, connected, and without loops and isolated vertices.

For an edge $\{u, v\} \in E(G)$ and a real number $\lambda \in [0, 1]$, let $p(u, v, \lambda) \in P(G)$ be the point on edge $\{u, v\}$ that has distance λ from u . Note that $p(u, v, 0) = u$, $p(u, v, 1) = v$ and $p(u, v, \lambda) = p(v, u, 1 - \lambda)$. Further, we use $d(p, q)$ for the length of a shortest path between points $p, q \in P(G)$.

For a subset of vertices $V' \subseteq V(G)$ or a subset of edges $E' \subseteq E(G)$, we denote by $G[V']$ and $G[E']$ the subgraph induced by V' and E' , respectively. The neighborhood of a vertex u is $N(u) := \{v \in V(G) \mid \{u, v\} \in E(G)\}$. We use n as the number of vertices of G , when G is clear from the context.

For a graph G and integer $c \geq 1$, let the c -subdivision of G be the graph G where every edge is replaced by a path of length c .

► **Lemma 1** ([8]). *Let G be a graph, let $c \geq 1$ be an integer, and let G' be the c -subdivision of G . Then $\delta\text{-disp}(G) = (c\delta)\text{-disp}(G')$.*

For integers a and b , we denote the rational number $\frac{a}{b}$ as b -simple. A set $S \subseteq P(G)$ is b -simple, if for every point $p(u, v, \lambda)$ in S the edge position λ is b -simple.

► **Lemma 2** ([8]). *Let $\delta = \frac{a}{b}$ with integers a and b , and let G be a graph. Then, there exists an optimal δ -dispersed set S^* that is $2b$ -simple.*

For an introduction into parameterized algorithms, we refer to [4]. We study of the complexity of DISPERSION with the natural parameter solution size k , as well as its dependency on structural measures on the input graph. Besides treewidth $\text{tw}(G)$ and pathwidth $\text{pw}(G)$, we also study the parameters ‘feedback vertex set size’ $\text{fvs}(G)$ and treedepth $\text{td}(G)$.

A graph has a feedback vertex set $W \subseteq V(G)$ if G after removing W contains no cycle. The ‘feedback vertex set size’ is the size of a smallest feedback vertex set of G .

The treedepth of a connected graph G can be defined as follows. If G is disconnected, it is the maximum treedepth of its components; If G consists of a single vertex, then $\text{td}(G) = 1$; And else it is one plus the minimum over all $u \in V(G)$ of the treedepth of G without vertex u .

We provide lower bounds for the time-complexity assuming the Exponential Time Hypothesis (ETH): There is no $2^{o(N)}$ -time algorithm for 3-SAT with N variables and $\mathcal{O}(N)$ clauses [10]. For more details on ETH, we refer to [4].

3 Translating δ -Dispersion

There is an intriguing relation of the optimal solution for distance δ and $\frac{\delta}{2\delta+1}$ for the similar problem δ -covering [9]. We may analogously expect that an optimal solution for δ -dispersion translates to an optimal solution for $\frac{\delta}{\delta+1}$ -dispersion; i.e., that an optimal δ -dispersed set

corresponds to an optimal $\frac{\delta}{\delta+1}$ -dispersed set of the same size plus one extra point for every edge.

This is not true for $\delta = 3 + \varepsilon$ for any $\varepsilon > 0$: Consider a triangle, where a $(3 + \varepsilon)$ -dispersed set S contains at most one point p . Since $\frac{\delta}{\delta+1} > \frac{3}{4}$, a $\frac{\delta}{\delta+1}$ -dispersed set however contains at most $3 < |S| + 3$ points.

Causing trouble is a non-trivial closed walk containing p and of length less than δ . The translating lemma may only apply to a variation of dispersion that is sensitive to such walks, a variant which we call *auto-dispersion*. A δ -dispersed set $S \subseteq P(G)$ is δ -*auto-dispersed* if additionally for every point $p \in S$ there is no walk from p to p of length $< \delta$ that is locally-injective and non-trivial. A walk is *locally-injective* if, when interpreted as a continuous mapping $f : [0, 1] \rightarrow P(G)$ from $f(0) = p$ to $f(1) = p$, has for every pre-image $c \in (0, 1)$ a positive range $\varepsilon > 0$ such that f restricted to the interval $(c - \varepsilon, c + \varepsilon)$ is injective.

► **Lemma 3.** *Let G be a graph and $\delta > 0$. Then δ -auto-disp(G) = $\frac{\delta}{\delta+1}$ -auto-disp(G) + $|E(G)|$.*

Fortunately, this translation lemma is still useful for ordinary δ -dispersion. We have δ -auto-disp(G) = δ -disp(G) for $\delta \leq 3$, since there is no such non-trivial closed walk 'without a turn within an edge' of length < 3 . The threshold of 3 is tight according to the above example with graph K_3 .

► **Corollary 4.** *Let G be a graph and $\delta \in (0, 3]$. Then δ -disp(G) = $\frac{\delta}{\delta+1}$ -disp(G) + $|E(G)|$.*

4 Dispersion and Independent Set

To solve DISPERSION we can borrow from algorithmic results from a generalized independent set problem. A classical independent set is a set of *vertices* where each two elements have to be at least 2 apart from each other (when we consider that the edges have unit length). In a 2-dispersed set also each two elements need to be at least 2 apart from each other, though the set contains a set of *points* of the graph.

To generalize the independent set problem, we may ask that the vertices are not 2 apart but some integer d apart from each other. Such a generalization for independent set is called a *distance- d independent set* or *d -scattered set*. They have been studied by Eto et al. [5] and Katsikarelis et al. [12].

Let $\alpha_d(G)$ be the maximum size of a distance d independent set, for a graph G and integer d . We relate δ -dispersion to α_d . We consider the *c -subdivision of a graph G* , denoted as G_c , which is the graph G where every edge is replaced by a path of length c , for some integer $c \geq 1$.

► **Lemma 5.** *Consider integers a, b and a $2b$ -subdivision G_{2b} of a graph G . Then $\frac{a}{b}$ -disp(G) = $\alpha_{2a}(G_{2b})$.*

Proof. Consider the b -subdivision G_b of G . Then G_{2b} is a 2-subdivision of G_b . We know that $\frac{a}{b}$ -disp(G) = $2a$ -disp(G_{2b}) from Lemma 1. Hence it remains to show $2a$ -disp(G_{2b}) = $\alpha_{2a}(G_{2b})$.

Clearly, a distance- $2a$ independent set $I \subseteq V(G)$ is also a $2a$ -dispersed set. For the reverse direction, assume there is a $2a$ -dispersed set I_{2a} of G_{2b} . Then I_{2a} corresponds to an a -dispersed set I of G_b of same size, according to Lemma 1. Since a is integer, we may assume that S contains only half-integral points, hence points with edge position from $\{0, \frac{1}{2}, 1\}$, according to Lemma 2. Let G_{2b} result from G_b by replacing each edge $\{u, v\}$ by a path $uw_{u,v}v$. Then let $I \subseteq V(G_{2b})$ consist of vertex $u \in V(G)$ with a point in S and every $w_{u,v}$ for every point $p(u, v, \frac{1}{2}) \in S$. Then I is a distance $2a$ independent set of G_{2b} of size $|I| = |S|$. ◀

Thus to solve DISPERSION for $\delta = \frac{a}{b}$ we can use algorithms for distance d independent set. For rationals $\frac{a}{b}$ with small values of a and b this possibly leads to efficient algorithms. For example, a distance d independent set on graphs with treewidth $\text{tw}(G)$ (and a given tree decomposition) can be found in time $d^{\text{tw}(G)}n^{\mathcal{O}(1)}$, see [12]. “Simply” subdivide the edges of the input graph sufficiently often, which does not increase the treewidth of the considered graph. To find a $\frac{a}{b}$ -dispersed set in a graph G , we can search for distance $2a$ independent set the $2b$ -subdivision of G .

► **Corollary 6.** *There is an algorithm that, given a rational distance $\frac{a}{b} > 0$ and a graph G , a tree decomposition of width $\text{tw}(G)$, computes a maximum $\frac{a}{b}$ -dispersed set S in time $(2a)^{\text{tw}(G)}(bn)^{\mathcal{O}(1)}$.*

However, in general this constitutes a possibly exponential increase of the input size. While in the input of $\frac{a}{b}$ -dispersion encodes a and b in binary, the subdivided graph essentially encodes b in unary. Further, if δ is irrational, we do not have a suitable subdivision at all.

5 Rounding the Distance

For a given graph G and distance δ , we state a rational $\delta^* \geq \delta$ such that δ -disp(G) = δ^* -disp(G). Our proof is constructive. We give a procedure that efficiently transforms a δ -dispersed set into a δ^* -dispersed set. The guaranteed rational δ^* has a numerator bounded by the longest path in G (or just n as an upper bound thereof). It is independent of the precise structure of the given graph.

To give some intuition: Generally there is some leeway for δ . For example, in a star $K_{1,k}$, $k \geq 1$ for every $\delta \in (1, 2]$ the optimal solution puts a point on every leaf yielding a δ -dispersed set of size k . Hence for instance $\frac{3}{2}$ -disp($K_{1,k}$) = 2-disp($K_{1,k}$). However, for $\delta > 2$ only one point can be placed, such that 2 -disp($K_{1,k}$) \neq $(2 + \varepsilon)$ -disp($K_{1,k}$) for every $\varepsilon > 0$.

So what δ^* can be guaranteed such that δ -disp(G) = δ^* -disp(G)? An illustrative example is a path of length 6. Then $\frac{15}{11}$ -disp(G) = 5 = $\frac{3}{2}$ -disp(G). For $\delta = \frac{15}{11}$ tightly packing 5 points allows to have a space of size $\frac{6}{11}$ at either end of the path, not enough to place another point. However, placing 5 points in distance $\delta = \frac{3}{2}$ allows no leeway; δ is (already) a divisor of 6, the length of the considered path. Distance δ^* relies on L , the length of the longest (not-induced) path in G . We have to take into account that δ might divide any path of length $\leq L$. Our δ^* is the smallest rational $\frac{a^*}{b^*}$ where the numerator $a^* \leq 2L + 2$. In other words, the inverse of δ^* is the next rational number in the Farey sequence of order $2L + 2$.

► **Theorem 7.** *Let $\delta \in \mathbb{R}^+$. Let L be an upper bound on the length of paths in G . Let $\delta^* = \frac{a^*}{b^*} \geq \delta$ minimal for $a^* \leq 2L + 2$ and $b^* \in \mathbb{N}$. Then δ -disp(G) = δ^* -disp(G).*

Clearly, a δ^* -dispersed set S^* , is also δ -dispersed, since $\delta^* \geq \delta$. We have to show the reverse direction. Consider a δ -dispersed set S (of size $|S| \geq 2$) of a connected graph G that is not δ^* -dispersed, hence δ is irrational or is equal to $\frac{a}{b}$ for some co-prime a, b with $a > 2L + 2$.

In the following we develop our rounding procedure that shows the reverse direction. Our presentation aims to be accessible by starting from the core algorithmic idea from which we unravel all involved technical concepts piece by piece. The detailed proofs are placed in the appendix.

5.1 Overview

Our rounding procedure repeatedly applies a pushing algorithm to the current point set S . We show that each such step strictly decreases a polynomially bounded potential $\Phi : P(G) \rightarrow \mathbb{N}$.

► **Theorem 8.** *Suppose that there is an algorithm, that given a δ -dispersed set S with $\delta < \delta^*$ computes an $\varepsilon > 0$ and a $(\delta + \varepsilon)$ -dispersed set S_ε of size $|S_\varepsilon| = |S|$ that satisfies $\Phi(S) > \Phi(S_\varepsilon)$ for some polynomially bounded potential $\Phi : P(G) \rightarrow \mathbb{N}$. Then Theorem 7 follows.*

Proof. Let S be a δ -dispersed set. Apply the assumed algorithm to obtain a $\varepsilon > 0$ and a $(\delta + \varepsilon)$ -dispersed set S_ε of size $|S_\varepsilon| = |S|$. If $\delta = \delta^*$, we reached our goal. Else we apply the assumed algorithm again. Since the potential $\Phi : P(G) \rightarrow \mathbb{N}$ decreases for S_ε compared to S and Φ is polynomially bounded, we have to reach δ^* in polynomial many steps. ◀

In the remainder of this section we will develop such an algorithm. It pushes the points of point set S away from each other such that their pairwise distance increases from ‘at least δ ’ to ‘at least $\delta + \varepsilon$ ’. We choose $\varepsilon \geq 0$ as large as possible limited by some events. Either we already reach $\delta + \varepsilon = \delta^*$, hence we reached our goal, or at least one of three events occurs. We will specify these events in the course of this section. These events mean that one pushing step, i.e., one step for Theorem 8 terminated. All the following preparations for such a pushing step start anew.

We make sure that our potential $\Phi : P(G) \rightarrow \mathbb{N}$ decreases when an event occurs. Each of the three events has a corresponding partial potential $\Phi_1(S)$, $\Phi_2(S)$ and $\Phi_3(S)$. They define the overall potential as $\Phi(S) := \Phi_1(S) + \Phi_2(S) + \Phi_3(S)$. Each part never increases. Whenever event i occurs, $\Phi_i(S)$ strictly decreases.

We denote a pair of points $\{p, q\}$ from our given point set S as δ -critical, if they have distance exactly δ . Hence the critical pairs of points are exactly those that we need to push away from each other. At the same time we make sure that, once $\{p, q\}$ are δ -critical, they never turn uncritical again, i.e., they are $(\delta + \varepsilon)$ -critical in the next step. An uncritical pair of points $\{p, q\}$ might become critical, hence we have to take care of $\{p, q\}$ in future steps. This constitutes our first event. The corresponding partial potential is $\Phi_1(S)$, the number of uncritical pairs of points $\{p, q\}$.

(Event 1) A δ -uncritical pair of points $\{p, q\}$ becomes $(\delta + \varepsilon)$ -critical.


$$\Phi_1(S) := |\{\{p, q\} \in \binom{S}{2} \mid \{p, q\} \text{ are not } \delta\text{-critical}\}| \leq |S|^2$$

5.2 Coordination of Movement

We need to coordinate the movement of all critical pairs of points. To this end, we will fix some set of *root points* R . Our movement will be locally prescribed for sequences of points p_0, p_1, \dots, p_s that originate in $p_0 \in R$ and where each $\{p_0, p_1\}, \dots, \{p_{s-1}, p_s\}$ is critical. The overall movement will be uniquely defined by movement defined for these sequences.

For now, consider such a sequence of points p_0, p_1, p_2, \dots . Our idea is to do not move p_0 , to move p_1 by distance ε away from p_0 , point p_2 by distance 2ε away from p_1 and so on. We have to stop pushing in this way as soon as one of the points, say p_i , becomes half-integral, i.e., p_i is moved onto a vertex or the midpoint of an edge. See Figure 1 for examples. This constitutes the second event.

(Event 2) A non-half-integral $p \in S$ becomes half-integral.



■ **Figure 1** (left) Consider critical points $\{p, q\}$ (points depicted as black dots; vertices as white squares). If we move q away from p by $\varepsilon \geq 0$, their distance increases by ε until q reaches the half-integral point $p(u, v, \frac{1}{2})$. (middle) Let $\{p_i, p_{i-1}\}$ be critical for $i \geq 1$. Consider moving $p_i, i \geq 1$ by $i\varepsilon$ away from p_{i-1} . Once p_2 becomes half-integral, points $\{p_3, p_0\}$ become also critical, hence we cannot continue to move points in the same way. This happens when a point in S becomes half-integral or ... (right) ... a point half-way between two points in S becomes half-integral, as in this example between p_4, p_5 . We say p_4, p_5 witness the pivot $p(u, v, \frac{1}{2})$.

$$\Phi_2(S) := |\{p \in S \mid p \text{ is not half-integral}\}| \leq |S|.$$

The next pushing step will choose p_i as one of the root point R and will move the points away from p_i instead of p_0 . Very similarly, we stop when a point $r \in P(G)$ that is ‘half-way’ between two points p_i, p_{i-1} becomes half-integral. Formally, we denote such a point r as an (S, δ) -pivot, or simply a pivot, if it is half-integral and there is a (critical) pair of points $\{p, q\} \in \binom{S}{2}$, the witnesses, that have equal distances to r , which means $d(p, r) = d(q, r) = \frac{\delta}{2}$. Let $\text{pivots}(S, \delta)$ be the set of (S, δ) -pivots, and let $W(S, \delta) \subseteq \binom{S}{2}$ be the family of pairs of points from S , that witness some (S, δ) -pivot. This leads to the third and final event.

(Event 3) A non-pivot point $r \in P(G)$ becomes a pivot.

$$\Phi_3(S) := |\{r \in P(G) \mid r \text{ is half-integral}\} \setminus \text{pivots}(S, \delta)| \leq |V(G)|^2.$$

Hence a root point R may not only be a point $p \in S$ but also come from the set of pivots. We will later properly define R as a superset of half-integral points $p_i \in S$ and the (S, δ) -pivots.

We use an *auxiliary graph* G_S for the current δ -dispersed set S . Its vertex set is $S \cup \text{pivots}(S, \delta)$. Essentially we make all pairs of critical $\{p, q\}$ adjacent unless they witness a pivot; If they do witness a pivot, we make them adjacent to the pivot:


- For $\{p, q\} \in W(S, \delta)$ and for every pivot $r \in \text{pivots}(S, \delta)$ they witness, add edges $\{p, r\}, \{r, q\}$; and
- for every critical pair of points $\{p, q\} \in \binom{S}{2} \setminus W(S, \delta)$ add edge $\{p, q\}$.

Note that, for every edge $\{r, p\}$ with $r \in \text{pivots}(S, \delta)$, there is at least one other edge $\{r, q\}$ such that p, q witness r as a pivot.

Now we define the sequence of points which serve as the structure to state the movement. A path $P = (p_0, p_1, \dots, p_s)$ in the auxiliary graph G_S of length $s \geq 1$ is a *spine* if p_1, \dots, p_s are not half-integral. Note that any sub-sequence (p_0, \dots, p_i) for $1 \leq i \leq s$ is also a spine.

5.3 Velocities

We assign velocities vel_P to the points p_0, \dots, p_s of a spine P that specify their movement speed. The point p_i for $i \in \{1, \dots, s\}$ is moved by $\text{vel}(p_i)\varepsilon$. Thus setting $\text{vel}(p_1) = 1$ makes the δ -critical $\{p_0, p_1\}$ become $(\delta + \varepsilon)$ -critical, as desired. Setting $\text{vel}(p_i) = i$ for $i \geq 1$, however, can make consecutive points $\{p_{i-1}, p_i\}$ uncritical. Figure 2 provides an example. To see this, fix some shortest p_{i-1}, p_i -path P_i and some shortest p_i, p_{i+1} -path P_{i+1} . The paths P_i and



■ **Figure 2** A spine $P = (p_0, \dots, p_7)$. The shortest path between p_0, p_1 and the shortest path between p_1, p_2 have the trivial intersection of $\{p_1\}$, hence $\text{flip}_P(p_1) = 1$. In turn, the shortest path between p_1, p_2 and the shortest path between p_2, p_3 have a non-trivial intersection, hence $\text{flip}_P(p_1) = -1$. Also $\text{flip}_P(p_3) = \text{flip}_P(p_4) = -1$ while the other values are positive. Consequently $(\text{sgn}_P(p_1), \dots, \text{sgn}_P(p_7)) = (1, 1, -1, 1, -1, -1, -1)$. Thus $(\text{vel}_P(p_1), \dots, \text{vel}_P(p_7)) = (1, 2, 1, 2, 1, 0, -1)$. In particular, $\text{sgn}_P(p_5)$ is negative such that it is moved towards p_4 . In turn, $\text{sgn}_P(p_7)$ and $\text{vel}_P(p_7)$ are negative such that p_7 has a net movement away from p_0 . Under this movement all $\{p_0, p_1\}, \dots, \{p_6, p_7\}$ remain critical.

P_{i+1} can have a trivial intersection of only $\{p_i\}$ or their intersection may contain more than one point. We denote this bit of information as $\text{flip}_P(p_i) \in \{-1, 1\}$. We set $\text{flip}_P(p_i) = 1$ if and only if P_i and P_{i-1} have a trivial intersection. (The definition of flip_P is independent on the exact considered shortest paths and we will define it properly in the next subsection.)

The easy case is when P_i and P_{i+1} have a trivial intersection, i.e., $\text{flip}_P(p_i) = 1$. In this case we increase the velocity of the next point p_{i+1} . The first time we encounter the other case, that $\text{flip}_P(p_i) = -1$, we decrease the velocity of the next point p_{i+1} . Further we move p_{i+1} towards and not away of p_i . Hence we also specify a **sgn** of the velocity that records whether a point p_{i+1} is pushed towards or away from its predecessor p_i . All these changes are relative to whether the movement of predecessor p_i is away from p_{i-1} , i.e., whether $\text{sgn}(p_i)$ is positive. For example, the second time we encounter a point p_j with $\text{flip}_P(p_j) = -1$, point p_{j+1} is again moved away from its predecessor.

This leads to the following definition of vel_P and sgn_P for a spine P . We define its half-integral velocities $\text{vel}_P : \{p_0, \dots, p_s\} \rightarrow \{\frac{z}{2} \mid z \in \mathbb{Z}\}$ depending on signs $\text{sgn}_P : \{p_1, \dots, p_s\} \rightarrow \{-1, 1\}$, which in turn depend on flip_P . We may drop the subscript P , if it is clear from the context. Let $\text{vel}(p_0) = 0$. Let $\text{vel}(p_1) = \frac{1}{2}$, if $p_0 \in \text{pivots}(S, \delta)$, and let $\text{vel}(p_1) = 1$, if $p_0 \in S$. For $i \geq 1$, let

$$\text{vel}(p_{i+1}) := \text{vel}(p_i) + \text{sgn}(p_{i+1}).$$

Thus $\text{sgn} \in \{-1, 1\}$ indicates whether the velocity increases or decreases. The current **sgn** is unchanged unless **flip** is negative. Let $\text{sgn}(p_1) = 1$. For $2 \leq i \leq s$, let

$$\text{sgn}(p_i) := \text{flip}(p_{i-1}) \text{sgn}(p_{i-1}) = \prod_{0 < j < i} \text{flip}(p_j).$$

The movement step of a point p_i in a spine $P = (p_0, \dots, p_i)$ is now as follows. We push the point p_i by the (possibly negative) distance $\text{sgn}_P(p_i) \text{vel}_P(p_i) \varepsilon$ away from its predecessor p_{i-1} . In other words, the point $p_i = p(u, v, \lambda)$ is replaced by the point $p(u, v, \lambda + \text{sgn}_P(p_i) \text{vel}_P(p_i) \varepsilon)$ assuming that vertex u compared to v is in some sense closer to the predecessor point p_{i-1} . We make this notion formal in the next subsection.

5.4 Directions

We formalize the notion the direction of a point p towards another point q . The *direction* $\text{dir}(p \rightarrow q) \in \{u, v\}$ for distinct points $p = p(u, v, \lambda) \in P(G)$ and $q \in P(G)$ is defined as follows:

- For points $p = p(u, v, \lambda_p)$ and $q = p(u, v, \lambda_q)$ on a common edge $\{u, v\} \in E(G)$ with $\lambda_p < \lambda_q$, let $\text{dir}(p \rightarrow q) = v$. Let $\overline{\text{dir}}(p \rightarrow q) = u$.
- For points $p = p(u_p, v_p, \lambda_p)$ and $q = p(u_q, v_q, \lambda_q)$ on distinct edges $\{u_p, v_p\} \neq \{u_q, v_q\}$, let $\text{dir}(p \rightarrow q)$ be the unique vertex of $\{u_p, v_p\}$ that is contained in *every* shortest path between p and q , if such a vertex exists. If $\text{dir}(p \rightarrow q)$ is defined, let $\overline{\text{dir}}(p \rightarrow q)$ be the unique vertex in $\{u_p, v_p\} \setminus \{\text{dir}(p \rightarrow q)\}$.

► **Lemma 9.** (\star) *For distinct points $p, q \in V(G_S)$, $\text{dir}(p \rightarrow q)$ is well-defined, unless p is half-integral.*

Hence we can properly define $\text{flip}(p_i)$ of a point p_i of a spine (p_0, \dots, p_s) with $1 \leq i \leq s-1$, since p_i with $i \geq 1$ is non-half-integral. Let

$$\text{flip}(p_i) := \begin{cases} 1, & \text{dir}(p_i \rightarrow p_{i-1}) \neq \text{dir}(p_i \rightarrow p_{i+1}), \\ -1, & \text{else.} \end{cases}$$

Further, for a non-half-integral point $p = p(u, v, \lambda)$ we have $\{u, v\} = \{\text{dir}(p \rightarrow q), \overline{\text{dir}}(p \rightarrow q)\}$. By symmetry assume that $\overline{\text{dir}}(p \rightarrow q) = u$. We can equivalently specify point p as $p(\overline{\text{dir}}(p \rightarrow q), \text{dir}(p \rightarrow q), \lambda)$. Conveniently, we may write $p = p(\cdot, \overline{\text{dir}}(p \rightarrow q), \lambda)$ since the missing entry is clear from the context. Doing so, the edge position λ measures a part of the length of any shortest p, q -path, specifically the part using the edge of p (assuming q is on another edge).

Now we can also properly define one pushing step for a point p_i of a spine $P = (p_0, \dots, p_s)$ and for $\varepsilon > 0$. Let λ_i be such that $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \lambda_i)$. Then the new point is

$$(p_i)_{P, \varepsilon} := p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \lambda_i + \text{sgn}_P(p_i) \text{vel}_P(p_i) \varepsilon).$$

► **Lemma 10.** (\star) *For a spine $P = (p_0, \dots, p_s)$ and $i \in \{0, \dots, s-1\}$, points $(p_i)_{P, \varepsilon}, (p_{i+1})_{P, \varepsilon}$ are $(\delta + \varepsilon)$ -critical for the maximal $\varepsilon \leq \delta^* - \delta$ that is limited by the Events 1,2,3.*

5.5 Root Points

We formally define the set of root points R . Let R_0 be the set of half-integral points in G_S . There may be some components of the auxiliary graph G_S without a point in R_0 . Let R result from R_0 by adding exactly one point from every component that has no point in R_0 .

We consider only spines $P = (p_0, \dots, p_i)$ where $p_0 \in R$. Clearly every point G_S is part of at least one spine and hence has some movement prescribed. We also claim that the prescribed movement is uniquely defined. In other words, there are no spines $P = (p_0, \dots, p_i)$ and $Q = (q_0, \dots, q_j)$ with $p_0, q_0 \in R$ that terminate at the same point $p_i = q_j$ and contradict in their prescribed movement for $p_i = q_j$.

► **Lemma 11.** (\star) *Let $\delta < \delta^*$. Consider spines $P = (p_0, \dots, p_i)$ and $Q = (q_0, \dots, q_j)$ with $p_0, q_0 \in R$ and $p_i = q_j$. Then (1) $\text{vel}_P(p_i) = \text{vel}_Q(q_j)$; and (2) $\overline{\text{dir}}(p_i \rightarrow p_{i-1}) = \overline{\text{dir}}(q_j \rightarrow q_{j-1})$ if and only if $\text{sgn}_P(p_i) = \text{sgn}_Q(q_j)$.*

Therefore we can uniquely define the moved version of a point p as $p_\varepsilon := (p)_{P, \varepsilon}$ where we may choose P to be an arbitrary spine starting in R and containing p . This defines the set of pushed points $S_\varepsilon := \{p_\varepsilon \mid p \in S\}$.

1:10 Dispersing Obnoxious Facilities on Graphs by Rounding Distances

For the proof of Lemma 11, we use that $\delta^* = \frac{a^*}{b^*} \geq \delta$ is minimal with $a^* \leq 2L + L$. Since spines P, Q meet at the point $p_i = q_j$ their roots p_0, q_0 must be from the same component in the auxiliary graph G_S ; in other words p_0, q_0 are both half-integral or are the same point. Our proof by contradiction considers these two options and whether P, Q reach p_i and q_j from the same vertex relatively to if they agree on the sgn of $p_i = q_j$. An example is that $p_0 = q_0$ and P, Q reach $p_i = q_j$ from the same vertex (formally with $\overline{\text{dir}}(p_i \rightarrow p_{i-1}) \neq \overline{\text{dir}}(q_j \rightarrow q_{j-1})$) while they agree on the sgn (formally $\text{sgn}_P(p_i) = \text{sgn}_Q(q_j)$). Then we can glue P and Q together forming a walk starting in $p_0 = q_0$ and returning to $p_0 = q_0$. This walk then has half-integral length at most $2L + 2$ but is made up hops of length δ . That implies the contradiction that already $\delta = \delta^*$.

5.6 Summery

With the previous observations we can assemble the algorithm for Theorem 8. Our potential counts how many elements can still trigger Events 1,2,3. That is

$$\Phi(S) := \Phi_1(S) + \Phi_2(S) + \Phi_3(S) \leq 2|S|^2 + |V(G)|^2.$$

We define $\varepsilon^* \geq 0$ as the maximal $\varepsilon \leq \delta^* - \delta$ limited by the Events 1,2,3. We claim that $\varepsilon^* \geq 0$ is defined. This is due to that the above events depend on continuous functions in ε , which are the distance of p_ε to its closest half-integral point, and the distance between points p_ε and q_ε for $p, q \in S$.

To show termination, we prove that no such element can trigger its according event more than once. Lemma 10 already implies that a δ -critical pair of points $\{p, q\}$ stays $(\delta + \varepsilon^*)$ -critical. It remains to show the following monotonicities:

- **Lemma 12.** (\star) *Let S be a δ -dispersed set for $\delta < \delta^*$ and ε^* defined as above. Then:*
- (1) S_{ε^*} is a $(\delta + \varepsilon^*)$ -dispersed set of size $|S|$.
 - (2) If $\{p, q\} \in \binom{S}{2}$ is δ -critical, then $\{p_{\varepsilon^*}, q_{\varepsilon^*}\}$ is $(\delta + \varepsilon^*)$ -critical.
 - (3) If $r \in \text{pivots}(S, \delta)$, then $r \in \text{pivots}(S_{\varepsilon^*}, \delta + \varepsilon^*)$.

Now we have all the tools to show Theorem 8. Determine ε^* and the $(\delta + \varepsilon^*)$ -dispersed set S_{ε^*} as defined before. The resulting set S_{ε^*} is a $(\delta + \varepsilon^*)$ -dispersed set of the same size, according to Lemma 12. If $\delta + \varepsilon^* = \delta^*$ then S_{ε^*} is already the desired δ^* -dispersed set. Else one of the Events 1,2,3 occurred. We observe that the potential strictly decreases, that is $\Phi(S_{\varepsilon^*}) < \Phi(S)$. Because of the monotonicities of Lemma 10 and Lemma 12 the partial potentials Φ_1, Φ_2 and Φ_3 do not increase. If Event 1 occurs then Φ_1 strictly decrease. If Event 2 occurs then Φ_2 strictly decrease. If Event 3 occurs then Φ_3 strictly decrease. All in all at least one part strictly decreases and so does Φ . This completes the proof of Theorem 8 and hence of Theorem 7.

6 Algorithmic Implications

Based on the rounding procedure from Section 5, the translation result from Section 3 and connections to distance- d independent set we derive a number of algorithmic results.

- **Theorem 13.** *There is an algorithm that, given distance $\delta \geq 0$, a graph G , a tree decomposition and an upper bound $L \in \mathbb{N}$ on the length of the longest path in G , computes a maximum δ -dispersed set S in time $(2L + 2)^{\text{tw}(G)} n^{\mathcal{O}(1)}$.*

Proof. According to Theorem 7, we may consider the rounded up distance, that is a rational $\frac{a}{b} \geq \delta$ with $a \leq 2L + 2$, instead of δ . Notice that $\frac{a}{b}$ polynomial time computable. As long as $\frac{a}{b} \leq \frac{3}{4}$, we may repeatedly apply Corollary 4 such that eventually we obtain that $\frac{3}{4}b < a \leq 2L + 2$. Let G_{2b} be a $2b$ -subdivision of G . Observe that $\text{tw}(G) = \text{tw}(G_{2b})$ and the number of vertices increases only by a factor of $\mathcal{O}(n^2L)$. According to Lemma 5 $\frac{a}{b}$ -disp(G) = $\alpha_{2a}(G_{2b})$. Thus, to answer the original δ -DISPERSION-instance we may find a maximum distance- $2a$ independent set in G_{2b} , which is possible in time $(2a)^{\text{tw}(G)}n^{\mathcal{O}(1)}$, according to [11]. ◀

This result immediately yields parameterized complexity results for the parameters treedepth and treewidth. Regarding the treewidth, note that n is an upper bound on L . Thus the above algorithm is an XP-algorithm for the parameter treewidth. When a treewidth decomposition is given, DISPERSION can be solved in time $2n^{\text{tw}(G)}n^{\mathcal{O}(1)}$.

▶ **Corollary 14.** *DISPERSION can be solved in time $2n^{\text{tw}(G)}n^{\mathcal{O}(1)}$, assuming a tree decomposition is given.*

Similarly we obtain an FPT algorithm for treedepth $\text{td}(G)$ of the input graph. The treedepth $\text{td}(G)$ implies a bound on L , which is $L \leq 2^{\text{td}(G)}$. Since also $\text{td}(G) \geq \text{tw}(G)$, we obtain an $2^{\mathcal{O}(\text{td}(G)^2)}n^{\mathcal{O}(1)}$ -time algorithm, assuming a treedepth decomposition is given.

▶ **Corollary 15.** *DISPERSION can be solved in time $2^{\mathcal{O}(\text{td}(G)^2)}n^{\mathcal{O}(1)}$, assuming a treedepth decomposition is given.*

7 Parameterized Hardness Results

We complement the positive results by hardness results. These results borrow ideas from hardness-reductions for the similar problem DISTANCE INDEPENDENT SET (DIS), see Section 4.

A natural generalization of treedepth is the maximum diameter of graph G , which is the maximum distance between any vertices $u, v \in V(G)$ (since we only consider connected graphs G). We show NP-hardness for graphs of any diameter ≥ 3 even for chordal graphs by a reduction from INDEPENDENT SET, similarly as NP-hardness for DIS is shown by Eto et al. [5]. Our reduction also shows W[1]-hardness with respect to the solution size k .

▶ **Lemma 16.** (\star) *For every $\delta > 3$, δ -DISPERSION is NP-complete and W[1]-hard with parameter solution size, even for connected chordal graphs of diameter $\leq \lceil \delta \rceil$.*

Another direct generalization of treedepth is pathwidth of the input graph G . We show W[1]-hardness even for the combined parameters pathwidth and solution size $\text{pw}(G) + k$. With the same reduction also W[1]-hardness for the combined parameters ‘feedback vertex set size’ $\text{fvs}(G)$ and solution size k follows. We can essentially use the same reduction as used by Katsikarelis et al. to show W[1]-hardness of DIS when parameterized by $\text{fvs}(G) + k$ by reducing from MULTI-COLORED-INDEPENDENT-SET [11].

▶ **Theorem 17.** (\star) *DISPERSION is W[1]-hard parameterized by $\text{pw}(G) + k$. Further, there is no $n^{o(\sqrt{\text{pw}(G)} + \sqrt{k})}$ -time algorithm unless ETH fails. DISPERSION is W[1]-hard parameterized by $\text{fvs}(G) + k$. Further, there is no $n^{o(\text{fvs}(G) + \sqrt{k})}$ -time algorithm unless ETH fails.*

Since $\text{fvs}(G)$ is a linear upper bound for the treewidth of G , we also obtain: DISPERSION is W[1]-hard parameterized by $\text{tw}(G) + k$. Further, there is no $n^{o(\text{tw}(G) + \sqrt{k})}$ -time algorithm unless ETH fails. Similarly as in [11] we obtain a lower bound for treedepth.

▶ **Theorem 18.** (\star) *Assuming ETH, there is no $2^{o(\text{td}(G)^2)}$ -time algorithm for DISPERSION.*

8 NP-hardness for Irrational Distance

We show NP-hardness of δ -DISPERSION for every irrational distance $\delta > 0$. Thus together with earlier results [8] the complexity for every real $\delta > 0$ is resolved: For rational distance $\delta = \frac{a}{b}$ where $a \in \{1, 2\}$ the problem is polynomial time solvable, while it is NP-complete for every other distance $\delta > 0$.

► **Theorem 19.** *For every irrational $\delta > 0$, δ -DISPERSION is NP-complete.*

The key step is a reduction from INDEPENDENT SET which shows NP-hardness not only for a single distance δ but for the whole interval $\delta \in (2, 3]$.

Construction: Given a graph G and integer $k \in \mathbb{N}$, we construct an input for δ -DISPERSION consisting of a graph G' and integer $k' = k$ as follows. For every vertex $u \in V(G)$ introduce vertices u_1, u_2 and edge $\{u_1, u_2\}$. For every edge $\{u, v\} \in E(G)$ introduce edges $\{u_i, v_j\}$ for every $i, j \in \{1, 2\}$.

► **Lemma 20.** *For every $\delta \in (2, 3]$, δ -DISPERSION is NP-hard and W[1]-hard when parameterized by solution size.*

Proof. Clearly, this construction is polynomial time computable. Further, the reduction is parameter preserving such that W[1]-hardness of INDEPENDENT SET implies W[1]-hardness of DISPERSION, assuming correctness of the reduction.

Hence, it remains to show the correctness, that G has an independent set of size k if and only if G' has a δ -dispersed set of size k .

(\Rightarrow) Let $I \subseteq V(G)$ be an independent set of graph G . We define $S := \{p(u_1, u_2, \frac{1}{2}) \mid u \in I\} \subseteq P(G)$, which has size $|S| = |I|$. We claim that S is δ -dispersed in G' for $\delta \in (2, 3]$. Since any vertices $u, v \in V(G)$ have distance at least 2 in G , their corresponding points $p(u_1, u_2, \frac{1}{2})$ and $p(v_1, v_2, \frac{1}{2})$ have distance at least 3 in $P(G)$. Thus they are δ -dispersed for $\delta \in (2, 3]$.

(\Leftarrow) Let $S \subseteq P(G)$ be a δ -dispersed for some $\delta \in (2, 3]$. We define the ball B_u for $u \in V(G)$ as the points in $P(G)$ with distance at most $\frac{1}{2}$ to u_1 or u_2 , which is $B_u := \{p(u_i, v, \lambda) \mid i \in \{1, 2\}, \{u_i, v\} \in E(G'), \lambda \in [0, \frac{1}{2}]\}$. Every ball B_u for $u \in V(G)$ contains at most one point from S since points $p, q \in B_u$ can be at most $2 < \delta$ apart. Every union $B_u \cup B_v$ for adjacent $\{u, v\} \in E(G)$ contains at most one point from S since points $p, q \in B_u \cup B_v$ can also be at most $2 < \delta$ apart.

Now we define an independent set $I \subseteq V(G)$. Add vertex $u \in V(G)$ for every point $p \in S \cap B_u$ except when $p \in B_u \cap B_v$ for some $v \in P(G)$. If $p \in S \cap B_u \cap B_v$, add either the point u or v to I . Then $|I| = |S|$ since the union of B_u for $u \in V(G)$ is the whole point space $P(G)$. Further, no adjacent vertices u, v are in I since $B_u \cup B_v$ contain at most one point from S . Thus $I \subseteq V(G)$ is an independent set of size $|S|$. ◀

Because $\delta \leq 3$ we may apply Lemma 3 to obtain NP-hardness for δ in the interval $(\frac{2}{2x+1}, \frac{3}{3x+1}]$ for every integer $x \geq 0$. Applying Lemma 1 yields NP-hardness for δ in the interval $(\frac{2c}{2x+1}, \frac{3c}{3x+1}]$ for every integer $c \geq 1$.

Now, consider any irrational distance $\delta > 0$. Consider $F := \{c\delta^{-1} - \lfloor c\delta^{-1} \rfloor \mid c \geq 1\}$, the set of fractional parts of multiples of δ^{-1} . Since δ^{-1} is irrational, F is a dense subset of the interval $[0, 1]$. Let integer $c \geq 1$ be such that $\frac{1}{3} \leq c\delta^{-1} - \lfloor c\delta^{-1} \rfloor < \frac{1}{2}$. Thus there is a non-negative x such that $x + \frac{1}{3} \leq c\delta^{-1} < x + \frac{1}{2}$. This implies that $\frac{2c}{2x+1} < \delta \leq \frac{3c}{3x+1}$ and hence NP-hardness for δ -dispersion. This finishes the proof of Theorem 19.

9 Parameter Solution Size

δ -DISPERSION parameterized by the solution size k is $W[1]$ -hard when $\delta > 2$: When $\delta \in (2, 3]$ Lemma 20 shows $W[1]$ -hardness, while for $\delta > 3$ Lemma 16 implies $W[1]$ -hardness even when the input graph is chordal. It remains to consider $\delta \leq 2$. Observe that for $\delta \leq 2$, every graph G satisfies $\delta\text{-disp}(G) \geq \nu(G)$ [8], where $\nu(G)$ is the maximum matching size of G . Thus, a win-win situation occurs. Determine $\nu(G)$ in polynomial time. If $k \leq \nu(G)$, we may immediately answer ‘yes’. Otherwise $k > \nu(G) \geq \frac{\text{vc}(G)}{2}$, where $\text{vc}(G)$ is the minimum size of a vertex cover in G . The size of a vertex cover upper bounds the treedepth. A treedepth decomposition of size $\text{td}(G)$ is computable in FPT-time [14]. Thus we may apply the FPT algorithm for parameter treedepth from Theorem 13.

► **Theorem 21.** δ -DISPERSION parameterized by solution size k is FPT if $\delta \leq 2$; and $W[1]$ -hard if $\delta > 2$.

References

- 1 Shimon Abavaya and Michael Segal. Maximizing the number of obnoxious facilities to locate within a bounded region. *Comput. Oper. Res.*, 37(1):163–171, 2010. doi:10.1016/j.cor.2009.04.004.
- 2 Boaz Ben-Moshe, Matthew J. Katz, and Michael Segal. Obnoxious facility location: complete service with minimal harm. *International Journal of Computational Geometry and Applications*, 10(6):581–592, 2000.
- 3 Richard L. Church and Robert S. Garfinkel. Locating an obnoxious facility on a network. *Transportation Science*, 12:107–118, 1978.
- 4 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015. doi:10.1007/978-3-319-21275-3.
- 5 Hiroshi Eto, Fengrui Guo, and Eiji Miyano. Distance- d independent set problems for bipartite and chordal graphs. *J. Comb. Optim.*, 27(1):88–99, 2014. doi:10.1007/s10878-012-9594-4.
- 6 Alan J. Goldman and Perino M. Dearing. Concepts of optimal location for partially noxious facilities. *ORSA Bulletin*, 23:B–31, 1975.
- 7 Alexander Grigoriev, Tim A. Hartmann, Stefan Lendl, and Gerhard J. Woeginger. Dispersing obnoxious facilities on a graph. In *36th International Symposium on Theoretical Aspects of Computer Science, STACS 2019, March 13-16, 2019, Berlin, Germany*, pages 33:1–33:11, 2019. doi:10.4230/LIPIcs.STACS.2019.33.
- 8 Alexander Grigoriev, Tim A. Hartmann, Stefan Lendl, and Gerhard J. Woeginger. Dispersing obnoxious facilities on a graph. *Algorithmica*, 83(6):1734–1749, 2021.
- 9 Tim A. Hartmann, Stefan Lendl, and Gerhard J. Woeginger. Continuous facility location on graphs. *Math. Program.*, 192(1):207–227, 2022. doi:10.1007/s10107-021-01646-x.
- 10 Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4):512–530, 2001. doi:10.1006/jcss.2001.1774.
- 11 Ioannis Katsikarelis, Michael Lampis, and Vangelis Th. Paschos. Structurally parameterized d -scattered set. In *Graph-Theoretic Concepts in Computer Science - 44th International Workshop, WG 2018, Cottbus, Germany, June 27-29, 2018, Proceedings*, pages 292–305, 2018. doi:10.1007/978-3-030-00256-5_24.
- 12 Ioannis Katsikarelis, Michael Lampis, and Vangelis Th. Paschos. Structurally parameterized d -scattered set. *Discret. Appl. Math.*, 308:168–186, 2022. doi:10.1016/j.dam.2020.03.052.
- 13 Matthew J. Katz, Klara Kedem, and Michael Segal. Improved algorithms for placing undesirable facilities. *Comput. Oper. Res.*, 29(13):1859–1872, 2002. doi:10.1016/S0305-0548(01)00063-6.

- 14 Felix Reidl, Peter Rossmanith, Fernando Sánchez Villaamil, and Somnath Sikdar. A faster parameterized algorithm for treedepth. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part I*, pages 931–942, 2014. doi:10.1007/978-3-662-43948-7_77.
- 15 Michael Segal. Placing an obnoxious facility in geometric networks. *Nordic Journal of Computing*, 10:224–237, 2003.
- 16 Arie Tamir. On the solution value of the continuous p -center location problem on a graph. *Math. Oper. Res.*, 12(2):340–349, 1987. doi:10.1287/moor.12.2.340.
- 17 Arie Tamir. Obnoxious facility location on graphs. *SIAM J. Discret. Math.*, 4(4):550–567, 1991. doi:10.1137/0404048.
- 18 Martijn van Ee. Approximability of the dispersed $p \rightarrow$ -neighbor k -supplier problem. *Discret. Appl. Math.*, 289:219–229, 2021. doi:10.1016/j.dam.2020.10.007.

A Translating δ -Dispersion

A.1 A Distance Measure

We provide a distance measure for an edge $\{u, v\} \in E(G)$ regarding a point set $S \subseteq P(G)$. This notion will lead us to an alternative definition of auto-dispersion and covering that avoids talking about walks in $P(G)$.

For an edge $\{u, v\} \in E(G)$, let $\text{dr}_S(u, v)$ be the minimum length, over all points $p \in S$, of a u, p -walk in $P(G)$ starting from u in the direction of v .

This notion equals a recursive definition that does not involve walks in $P(G)$. A point $p \in S$ that yields the value $\text{dr}_S(u, v)$, is either already on the current edge $\{u, v\}$ or is found by recursing on another incident edge of v .

► **Lemma 22.** *For every edge $\{u, v\} \in E(G)$,*

$$\text{dr}_S(u, v) = \begin{cases} \min\{\lambda \mid p(u, v, \lambda) \in S\}, & \text{if } S \cap P(G[\{u, v\}]) \neq \emptyset, \\ 1 + \min\{\text{dr}_S(v, w) \mid w \in N(v) \setminus \{u\}\}, & \text{else.} \end{cases}$$

Proof. We prove the statement by induction on $\ell' = \lceil \text{dr}_S(u, v) \rceil$.

If $\ell' = 1$, there is a point $p(u, v, \lambda) \in S$ with edge position $\lambda = \text{dr}_S(u, v)$. Thus, $\{p(u, v, \lambda') \mid \lambda' \in [0, \lambda]\} \subseteq P(G)$ is a u, p -walk starting from u in the direction of v and of minimum length $\text{dr}_S(u, v)$.

Now assume by induction that the statement holds for all $u', v' \in V(G)$ with $\lceil \text{dr}_S(u', v') \rceil < \ell'$. Consider $\lceil \text{dr}_S(u, v) \rceil = \ell'$. Then $\ell' \geq 2$ such that $S \cap P(G[\{u, v\}]) = \emptyset$. Let distance $\text{dr}_S(u, v)$ be realized by a u, p -walk $P \subseteq P(G)$ starting from u in the direction of v . Since P has length > 1 , after traversing edge $\{u, v\}$ it continues along an incident edge $\{v, w\} \in E(G)$ for some neighbor $w \in N(v) \setminus \{u\}$. Then P consists of $P(G[\{u, v\}])$ and a remaining v, p -walk P' starting from v in the direction of w . Thus $\text{dr}_S(v, w) = 1 + \text{dr}_S(v, w)$.

Vice versa, for any neighbor $w' \in N_G(v) \setminus \{u\}$ there is a v, p' -path Q' starting from u in the direction of w' for some point $p' \in S$. Thus $P(G[\{u, v\}])$ combined with Q' is a u, p' -walk $Q \subseteq P(G)$. By the minimality of P , walk Q' has length at least that of P' . Thus w minimizes $\min_{w' \in N(v) \setminus \{u\}} \text{dr}_S(v, w')$, which implies the statement. ◀

For convenience, let $\ell(p, u) > 0$ be minimum length of a non-trivial p, q -walk containing u for any point $q \in S$ (possibly $p = q$).

► **Lemma 23.** *Let $p = p(u, v, \lambda) \in S$ with $\text{dr}_S(u, v) = \lambda \in (0, 1]$. Then, there exists a neighbor $w \in N_G(u)$ such that $\text{dr}_S(u, v) + \text{dr}_S(u, w) = \ell(p, u)$.*

Proof. Consider a point $p = p(u, v, \lambda) \in S$ with $\text{dr}_S(u, v) = \lambda \in (0, 1)$.

(\geq) Let $w \in N_G(u) \setminus \{v\}$ be arbitrary. Let P' realize $\text{dr}_S(u, w)$ for some neighbor, which means that P' is u, q -walk starting from u in the direction of w be of length $\text{dr}_S(u, w)$, for a point $q \in S$. Combine P' and $\{p(u, v, \lambda) \mid \lambda' \in [0, \lambda]\}$ to a non-trivial p, q -walk P containing u of length $\text{dr}_S(u, v) + \text{dr}_S(u, w)$. Thus $\text{dr}_S(u, v) + \text{dr}_S(u, w) \geq \ell(p, u)$ for every $w \in N_G(u) \setminus \{v\}$.

(\leq) Let P realize $\ell(p, u)$, which means that P is a non-trivial p, q -walk containing u of length $\ell(p, u)$ for some point $q \in S$. Let neighbor $w \in N_G(u) \setminus \{v\}$ be such that $\{u, w\}$ is the first edge after $\{v, u\}$ visited by P . Let P' be the subwalk P starting from u and ending in q of length $\ell(p, u) - \text{dr}_S(u, v)$. Since P' is a u, q -walk starting from u in the direction of w , it has length $\geq \text{dr}_S(u, w)$. Thus $\text{dr}_S(u, v) + \text{dr}_S(u, w) \leq \ell(p, u)$. ◀

A.2 Every Edge Contains at Least One Point

As a preliminary step, we need the following result.

► **Lemma 24.** *Let G be a graph and $\delta < 1$. There exists a maximum δ -dispersed set $S \subseteq P(G)$ such that every edge $\{u, v\} \in E(G)$ contains a point $p(u, v, \lambda) \in S$ with $\lambda \in (0, 1)$.*

Proof. Consider a maximum δ -dispersed set S . Then $|S| \geq |E(G)|$, since placing a point on every mid-point of each edge is a δ -dispersed set. For $i \geq 0$, let $E_i \subseteq E(G)$ be the set of edges $\{u, v\}$ that contain exactly i points from S , which is $|S \cap \{p(u, v, \lambda) \mid 0 \leq \lambda \leq 1\}| = i$. Let $E_{\geq i}$ be the union of E_j for $j \geq i$. Let an edge $\{u, v\} \in E(G)$ be *happy* if there is a point $p(u, v, \lambda) \in S$ with $\lambda \in (0, 1)$.

Consider an unhappy edge $\{u, v\}$. We will modify S such that $\{u, v\}$ becomes happy, all happy edges remain happy and S remains equal-sized and δ -dispersed: Either there will be enough room such that $p(u, v, \frac{1}{2})$ may be added, or we will find a sequence of point movements that also adds a point on edge $\{u, v\}$.

If already $S \cup p(u, v, \frac{1}{2})$ is δ -dispersed, it contradicts that S is maximum. Hence $\delta > \frac{1}{2}$, and there is at least one point $p_1 \in S$ in distance $< \delta - \frac{1}{2} \leq \frac{\delta}{2}$ to $\{u, v\}$. Let point $p_1 = p(w_1, w_2, \lambda_1) \in S$ with $\lambda_1 < \frac{\delta}{2}$ have minimal distance to $p(u, v, \frac{1}{2})$. (We even know that $\lambda_1 < \delta - \frac{1}{2}$, but this is not essential for the proof). By symmetry, we may assume that $w_1 = v$. Further, let $w_0 := u$.

Based on vertices w_0, w_1, w_2 and point p_1 we define an *exchange sequence* w_0, w_1, \dots, w_{s+1} with points $p(w_i, w_{i+1}, \lambda_i) \in S$ for $1 \leq i \leq s$ as follows. Assume, we have w_0, w_1, \dots, w_i defined for an $i \geq 2$. If $\{w_{i-1}, w_i\} \in E_{\geq 2}$, stop and set $s := i - 1$. Else $\{w_{i-1}, w_i\} \in E_1$. Let E_{w_i} be the subset of edges $\{w_i, w_{i+1}\} \in E_{\geq 1} \setminus \{\{w_{i-1}, w_i\}\}$ that are incident to w_i . If E_{w_i} is empty, stop and set $s := i - 1$. Otherwise, let $p_i = p(w_i, w_{i+1}, \lambda_i) \in S$ be a point on an edge $\{w_i, w_{i+1}\} \in E_{w_i}$ where p_i has minimal distance to vertex w_i . If $\lambda_i > \lambda_{i-1}$ or $w_{i+1} \in \{w_1, \dots, w_i\}$, stop and set $s := i$. Else, continue with increased i .

We denote an exchange sequence $P = w_0, w_1, \dots, w_{s+1}$ as *non-increasing* if $\lambda_s \leq \dots \leq \lambda_1 < \frac{\delta}{2}$. We claim that we can modify S such that every exchange sequence is non-increasing. We achieve this goal by moving a point closer to the midpoint of its edge.

Consider an exchange sequence $P = w_0, w_1, \dots, w_{s+1}$ that is not non-increasing. By construction, we have that $\lambda_s > \lambda_{s-1} \leq \dots \leq \lambda_1 < \frac{\delta}{2}$ for $s \geq 2$. We replace the second to last point $p_{s-1} = p(w_{s-1}, w_s, \lambda_{s-1})$ by a new point $p'_{s-1} = p(w_{s-1}, w_s, \min\{\frac{\delta}{2}, \lambda_s\})$. Observe that S under this modification remains equal-sized, δ -dispersed, and no happy edge becomes unhappy. After this modification, if $s \geq 3$, the shortened sequence w_0, w_1, \dots, w_s is an exchange sequence that terminates with a point $p'_{s-1} = p(w_{s-1}, w_s, \lambda'_{s-1})$. Possibly $\lambda'_{s-1} > \lambda_{s-2}$, in which case this exchange sequence is also not non-increasing. We apply this modification iteratively, until every exchange sequence starting with $w_0 = u, w_1 = v$ is non-increasing.

We claim that this process terminates. If at one step $p'_{s-1} = p(u, v, \frac{\delta}{2})$, then it cannot be the second to last point of an exchange sequence again, and hence will not be moved again. Else p'_{s-1} compared to p_{s-1} is moved by $\lambda_\varepsilon := \lambda_s - \lambda_{s-1}$ closer to $\frac{\delta}{2}$. Let

$$\Lambda = \{\lambda \in (0, 1) \mid p(u, v, \lambda) \in S, \{u, v\} \in E(G)\}$$

be fixed before starting this modification process. Throughout this process it holds that $\lambda_\varepsilon \geq (\prod_{\lambda \in \Lambda} \lambda)^{-1}$, since $\lambda_\varepsilon > 0$ and $\lambda_\varepsilon = \sum_{\lambda \in \Lambda} m_\lambda \lambda$ with $m_\lambda \in \mathbb{Z}$ for each $\lambda \in \Lambda$. Thus after finitely many moves of p_{s-1} , it reaches edges position $\frac{\delta}{2}$.

If now no exchange sequence starting with $w_0 = u, w_1 = v$ exists, apply the process to $w_0 = v, w_1 = u$. If also in this case, no exchange sequence starting with $w_0 = v, w_1 = u$ exists, there is no point of S in distance δ to $p(u, v, \frac{1}{2})$. A contradiction to S being maximum.

Now, every exchange sequence P starting with $w_0 = u, w_1 = v$ is non-increasing, in other words, $\lambda_s \leq \dots \leq \lambda_1 < \frac{\delta}{2} < \frac{1}{2}$, and such an exchange sequence exists. Then, we may *apply* such an exchange sequence w_0, w_1, \dots, w_s to S as follows. For $1 \leq i \leq s$, replace every point $p_i = p(w_i, w_{i+1}, \lambda_i) \in S$ with the point $p'_i = p(w_{i-1}, w_i, 1 - \lambda_i)$, which is the point ‘mirrored’ at w_i . It is easy to see that S remains equal-sized and δ -dispersed under this modification.

Now consider an exchange sequence $P = w_0, w_1, \dots, w_{s+1}$. We distinguish the following cases.

- If P terminates because $\{w_s, w_{s+1}\} \in E_{\geq 2}$, apply P to S resulting in $S' \subseteq P(G)$. Then S under this modification has $\{w_0, w_1\}$ newly happy and every happy edge remains happy, beside possibly $\{w_s, w_{s+1}\}$. If $\{w_s, w_{s+1}\}$ became unhappy, it contained only the point $p_s \in S$ and point $q = p(w_s, w_{s+1}, 1) \in S$ on the vertex w_{s+1} . Replace q by point $p(w_s, w_{s+1}, \delta)$. Then, S remains equal-sized and δ -dispersed and edge $\{w_s, w_{s+1}\}$ is happy.
- If P terminates because $E_{w_{s+1}}$ is empty, then w_{s+1} is only incident to edge $\{w_s, w_{s+1}\}$ and edges from E_0 . Apply P to S resulting in $S' \subseteq P(G)$. Then S under this modification is δ -dispersed, $\{w_0, w_1\}$ is newly happy and every happy edge remains happy, except for $\{w_s, w_{s+1}\}$ which becomes unhappy. There is no point of S in distance δ to $p(w_s, w_{s+1}, \delta)$. A contradiction to the maximality of S .
- If P terminates because $w_{s+1} = w_k \in \{w_1, \dots, w_s\}$, replace every point p_i by a point $p'_i = p(w_i, w_{i+1}, \frac{\delta}{2})$ for $1 \leq i \leq s$, resulting in a set S' . Clearly, $|S'| = |S|$ and every edge for S is also happy for S' .

We claim that S' is δ -dispersed. The points p'_i, p'_{i+1} for $i \in \{1, \dots, s\}$ have distance 1. Further, point p'_s has distance 1 to p_k and distance $2 - \frac{\delta}{2}$ to p_{k-1} (if $k \geq 2$). It follows that p_1, \dots, p_s are δ -dispersed. Assume, for the sake of contradiction, that there is a point $q \in S' \setminus \{p'_1, \dots, p'_s\}$ that has a shortest q, p'_i -path for some $i \in \{1, \dots, s\}$. Since point p'_i compared to p_i is closer to w_{i+1} , path P uses vertex w_{i+1} . Thus, q is at position $q = p(w_{i+1}, w', \lambda)$ for some neighbor $w' \in N_G(w_{i+1})$ and $\lambda < \frac{\delta}{2}$. Because vertex w_{i+1} has at most one neighbor w' that contains a point in distance $< \frac{\delta}{2}$ to w_{i+1} , it must be that $w' = w_{i+2}$ and $i < s$. A contradiction to that S' contains only point p'_{i+1} on edge $\{w_{i+1}, w_{i+2}\}$.

After this modification, the closest point in S' to vertex u is p' with distance $\frac{\delta}{2}$. Consider exchange paths starting with $w_0 = v$ and $w_1 = u$ instead and apply above modifications. Possibly, the closest point in the updated set of points has distance $\frac{\delta}{2}$, such that a point $p(u, v, \frac{1}{2})$ can be added, which makes also edge $\{u, v\}$ happy. Otherwise, some construction made edge $\{u, v\}$ happy.

By repeatedly applying this procedure for every unhappy edge $\{u, v\}$, eventually we obtain a δ -dispersed set of size $|S|$ without unhappy edges. ◀

► **Corollary 25.** *Let $\delta < 1$. Then there exists a δ -auto-dispersed set S such that $|S| = \delta$ -auto-disp(G) and for every edge $\{u, v\} \in E(G)$ there is a point $p = p(u, v, \lambda) \in S$ with $\lambda \in (0, 1)$.*

A.3 An Alternative Definition

We obtain the following equivalence to auto-dispersion. A set of points $S \subseteq P(G)$ is *edge internally δ -dispersed* if every edge by itself does not falsify that S is δ -dispersed, formally that every distinct points $p, q \in P(G[\{u, v\}])$ for every edges $\{u, v\} \in E(G)$ are δ -dispersed.

► **Lemma 26.** *Let G be a graph and $\delta > 0$. A subset $S \subseteq P(G)$ is δ -auto-dispersed if and only if*

1. (A1) S is edge internally δ -dispersed,
2. (A2) $\text{dr}_S(u, v) + \text{dr}_S(u, w) \geq \delta$ for every distinct adjacent edges $\{u, v\}, \{u, w\} \in E(G)$ where vertex u is not in S .

Proof. (\Rightarrow) Let $S \subseteq P(G)$ be δ -auto-dispersed. Then clearly S is edge internally δ -dispersed. Now assume, for the sake of contradiction, that (A2) is violated, which means that there are distinct adjacent edges $\{u, v\}, \{u, w\} \in E(G)$ where vertex u is not in S and $\text{dr}_S(u, v) + \text{dr}_S(u, w) = \ell < \delta$.

If $\text{dr}_S(u, v) \leq 1$ there is a point $p \in S$ on edge $p(u, v, \lambda)$ with $\lambda > 0$ since u is not in S . Then Lemma 23 implies that $\ell(u, p) = \text{dr}_S(u, v) + \text{dr}_S(u, w) < \delta$, hence that there is non-trivial p, q -walk containing u of length $< \delta$ for some point $q \in S$. A contradiction to that S is δ -dispersed. Symmetrically, assuming $\text{dr}_S(u, w) \leq 1$ leads to a contradiction.

Thus, it remains to consider that $1 < \text{dr}_S(u, v) \leq \text{dr}_S(u, w)$, up to symmetry. Then $\text{dr}_S(u, v) = \min_{v' \in N_G(v)} 1 + \text{dr}_S(v, v')$ according to Lemma 22. Thus, $\text{dr}_S(v, v') = \text{dr}_S(u, v) - 1$. Consider the pair of edges $(v, v'), (v, u)$ instead, which has at least $\text{dr}_S(v, v')$ smaller than $\text{dr}_S(u, v), \text{dr}_S(u, w)$. We repeat this argument until $\text{dr}_S(v, v') < 1$. Then again the previous case applies. Therefore, also condition (A2) is satisfied.

(\Leftarrow) Let $S \subseteq P(G)$ be not δ -dispersed. Then there are points $p, q \in S$ (possibly $p = q$) with a non-trivial p, q -walk in $P(G)$. If p, q are distinct and from the same edge $\{u, v\} \in E(G)$, hence $p, q \in P(G[\{u, v\}])$, then S is not edge internally δ -dispersed.

Else, we may assume that $p = p(u, v, \lambda)$ for some edge $\{u, v\} \in E(G)$ for $\lambda \in (0, 1]$ and a shortest non-trivial p, q -walk contains u . In particular $\lambda = \text{dr}_S(u, v)$ since otherwise we may consider $p', q' \in S$ with smaller distance than p, q . Then $\ell(p, u) \leq \ell < \delta$ by definition. According to Lemma 23, $\text{dr}_S(u, v) + \text{dr}_S(u, w) = \ell < \delta$ for some neighbor $w \in N_G(u)$, which violates condition (A2). ◀

A.4 Constructive Part

Now, we are ready to prove the actual translation lemma.

► **Lemma 27** ((Lemma 3 repeated)). *Let G be a graph and $\delta > 0$. Then δ -auto-disp(G) = $\frac{\delta}{\delta+1}$ -auto-disp(G) + $|E(G)|$.*

Proof. (\leq) First, we show that δ -auto-disp(G) $\leq \frac{\delta}{\delta+1}$ -auto-disp(G) + $|E(G)|$. Consider a δ -dispersed set $S \subseteq P(G)$ of size $|S| = \delta$ -disp(G). Let $\rho = (\delta + 1)^{-1}$, the conversion ratio. We construct a $(\delta\rho)$ -dispersed set S' of size $|S'| = |S| + |E(G)|$. A key observation is that

$$\rho + \delta\rho = 1.$$

We construct S' by considering each edge $\{u, v\} \in E(G)$ separately. Set S' compared to S will contain one more point in $P(G[\{u, v\}])$ for every edge $\{u, v\} \in E(G)$. Any point $p \in S$ on a vertex will also be in S' . Thus $|S'| = |S| + |E(G)|$.

- Consider that edge $\{u, v\} \in E(G)$ contains $m \geq 1$ points from S , which is $|S \cap P(G[\{u, v\}])| = m$. Let $p = p(u, v, \lambda)$ be the point among $S \cap P(G[\{u, v\}])$ with minimum distance to u , hence with distance λ . Analogously, let $q = p(u, v, 1 - \mu)$ be the point among $S \cap P(G[\{u, v\}])$ with minimal distance to v , hence with distance μ , possibly $p = q$. Then $\text{dr}_S(u, v) = \lambda$ and $\text{dr}_S(v, u) = \mu$. There are $m - 1 \geq 0$ further points between p and q on edge $\{u, v\}$, such that p and q have distance $1 - (\lambda + \mu) \geq m\delta$. In other words $\lambda + \mu \leq 1 - m\delta$.

Add points $p' = p(u, v, \lambda\rho)$ and $q' = p(u, v, 1 - \mu\rho)$ to S' , which are distinct even in when $p = q$. Note that if p is positioned on a vertex, also p' is; analogously for q and q' . Thus, the distance between the new points p' and q' is

$$1 - (\lambda + \mu)\rho \geq 1 - (1 - m\delta)\rho = 1 - \rho + m\delta\rho = (m + 1)\delta\rho.$$

Hence, we may add m further points between p' and q' on edge $\{u, v\}$ to set S' in such a way that the now $m + 2$ points in $S' \cap P(G[\{u, v\}])$ have pairwise distance at least $\delta\rho$.

- It remains to consider that edge $\{u, v\} \in E(G)$ contains no point from S , which is $S \cap P(G[\{u, v\}]) = \emptyset$. Fix a neighbor $w_u \in N(u)$ with minimum distance $\text{dr}_S(u, w_u)$. Analogously, fix a neighbor $w_v \in N(v)$ with minimum distance $\text{dr}_S(v, w_v)$. By symmetry, we may assume the inequality $\text{dr}_S(u, w_u) \leq \text{dr}_S(v, w_v)$. Then $w_u \neq v$ since then $\text{dr}_S(u, w_u) = 1 + \text{dr}_S(v, w_v)$. Then, add a new point $p(u, v, \lambda')$ at edge position $\lambda' = \max\{\frac{\delta\rho}{2}, \delta\rho - \text{dr}_S(u, w_u)\rho\}$ to S' .

We show that S' is $(\delta\rho)$ -auto-dispersed by proving the conditions of Lemma 26. Set S' is edge internally $(\delta\rho)$ -dispersed since for every edge we have added points that have pairwise distance at least δ . It remains to show condition (A2), which is that $\text{dr}_{S'}(u, v) + \text{dr}_{S'}(u, w) \geq \delta$ for every distinct adjacent edges $\{u, v\}, \{u, w\} \in E(G)$ where vertex u is not in S .

We partition the directed edges (u, v) for $\{u, v\} \in E(G)$ as follows:

- A directed edge (u, v) is *positive* if it satisfies $\text{dr}_{S'}(u, v) \geq \text{dr}_S(u, v)\rho$. This is the case when $S \cap G[\{u, v\}] \neq \emptyset$, by construction. Further, we claim that (u, v) is also positive if $S \cap G[\{u, v\}] = \emptyset$ and point $p(u, v, \lambda')$ with $\lambda' = 1 - (\delta\rho - \text{dr}_S(v, w_v)\rho)$ is added to S' , where $w_v \in N_G(v)$ is a neighbor with minimum $\text{dr}_S(v, w_v)$, which has $w_v \neq u$. Then $\text{dr}_S(u, v) = 1 + \text{dr}_S(v, w_v)$. It follows that

$$\text{dr}_{S'}(u, v) = 1 - \delta\rho + \text{dr}_S(v, w_v)\rho = 1 - \delta\rho + \text{dr}_S(u, v)\rho - \rho = \text{dr}_S(u, v)\rho.$$

- A not positive directed edge (u, v) is *neutral* if $S \cap G[\{u, v\}] = \emptyset$ and point $p(u, v, \lambda')$ with $\lambda' = 1 - \frac{\delta\rho}{2}$ is added to S' . Then, since it is not positive, $\text{dr}_S(v, w_v)\rho > \frac{\delta\rho}{2}$ where $w_v \in N_G(u)$ is a neighbor with minimum $\text{dr}_S(v, w_v)$. Also $w_v \neq u$.
- A directed edge (u, v) is *negative* if $S \cap G[\{u, v\}] = \emptyset$ and point $p(u, v, \lambda')$ with $\lambda' = \max\{\frac{\delta\rho}{2}, \delta\rho - \text{dr}_S(u, w_u)\rho\}$ is added to S' , in which case $w_v \neq u$.

Now, we observe that distinct adjacent edges $\{u, v\}, \{u, w\} \in E(G)$ satisfy condition (A2), that is $\text{dr}_{S'}(u, v) + \text{dr}_{S'}(u, w) \geq 2(\frac{\delta\rho}{2}) \geq \delta\rho$. We distinguish which type of (u, v) and (u, w) have.

- If $(u, v), (u, w)$ are positive, $\text{dr}_{S'}(u, v) + \text{dr}_{S'}(u, w) \geq (\text{dr}_S(u, v) + \text{dr}_S(u, w))\rho \geq \delta\rho = \delta\rho$. Here we used that Lemma 26 applies to S and δ , hence that $\text{dr}_S(u, v) + \text{dr}_S(u, w) \geq \delta$.
- If $(u, v), (u, w)$ are both not positive, then $\text{dr}_{S'}(u, v) + \text{dr}_{S'}(u, w) \geq 2\frac{\delta\rho}{2} \geq \delta\rho$, which follows from the fact that $\delta\rho < 1$.
- Consider that (u, v) is neutral and (u, w) is positive. By construction, we have $\text{dr}_S(u, w) \geq \text{dr}_S(u, w_u) \geq \text{dr}_S(v, w_v)$. Then

$$\begin{aligned} \text{dr}_{S'}(u, w) + \text{dr}_{S'}(u, v) &\geq \text{dr}_S(u, w)\rho + 1 - \frac{\delta\rho}{2} \\ &\geq \text{dr}_S(v, w_v)\rho + 1 - \frac{\delta\rho}{2} \\ &\geq 1 \\ &> \delta\rho. \end{aligned}$$

- It remains to consider that (u, v) is negative and (u, w) is positive. Observe that $\text{dr}_{S'}(u, w) + \text{dr}_{S'}(u, v) \geq \text{dr}_{S'}(u, w_u) + \text{dr}_{S'}(u, v)$. If (u, w_u) is neutral or negative, we

obtain that $\text{dr}_{S'}(u, w_u) + \text{dr}_{S'}(u, v) \geq \delta\rho$, by the previous cases, as desired. Thus, consider that (u, w_u) is positive. Then

$$\begin{aligned} \text{dr}_{S'}(u, w_u) + \text{dr}_{S'}(u, v) &\geq \text{dr}_{S'}(u, w_u) + \delta\rho - \text{dr}_S(u, w_u)\rho \\ &\geq \text{dr}_{S'}(u, w_u) + \delta\rho - \text{dr}_{S'}(u, w_u) = \delta\rho. \end{aligned}$$

Therefore, Lemma 26 applies to S' and $\delta\rho$, which shows that S' is $\delta\rho$ -auto-dispersed.

(\geq) Now, we show that δ -auto-disp(G) $\geq \frac{\delta}{\delta+1}$ -auto-disp(G) + $|E(G)|$. Let $\rho^{-1} = (\delta + 1)$, the conversion ratio. Consider a $(\delta\rho)$ -dispersed set $S' \subseteq P(G)$ of size $|S'| = \delta$ -auto-disp(G). We construct a δ -covering set $S \subseteq P(G)$ of size $|S| = |S'| - |E(G)|$. A key observation is that

$$\rho^{-1} + \delta\rho = 1.$$

Similarly to before, we construct S by considering each edge $\{u, v\} \in E(G)$ separately. We may assume that every edge contains at least one point, according to Corollary 25. Set S compared to S' will contain one less point $p(u, v, \lambda)$ with $\lambda \in (0, 1)$ for every edge $\{u, v\}$. Any point $p \in S'$ on a vertex will also be in S . Thus $|S| = |S'| - |E(G)|$.

■ Consider that edge $\{u, v\}$ contains exactly one point from S' , which is $S' \cap P(G[\{u, v\}]) = \emptyset$. Then add no point for edge $\{u, v\}$ to S .

■ Consider that edge $\{u, v\}$ contains $m+1 \geq 2$ points from S' , which is $|S' \cap P(G[\{u, v\}])| = m+1$, for $m \geq 1$. Let $p' = p(u, v, \lambda)$ be the point among $S' \cap P(G[\{u, v\}])$ with minimal distance to vertex u , hence with distance λ . Analogously, let $q' = p(u, v, 1 - \mu)$ be the point among $S' \cap P(G[\{u, v\}])$ with minimal distance to v , hence with distance μ . Then $\text{dr}_{S'}(u, v) = \lambda$ and $\text{dr}_{S'}(v, u) = \mu$. We have that $p' \neq q'$.

Add points $p = p(u, v, \lambda\rho^{-1})$ and $q = p(u, v, 1 - \mu\rho^{-1})$ to S . If p' is on a vertex, also the new point p is on the same vertex; analogously for q' and q .

Observe that the distance between the old points p' and q' is $1 - (\lambda + \mu) \geq \delta\rho m$ since points $S' \cap G[\{u, v\}]$ have pairwise distance at least δ . Thus, the distance between the new points p and q is

$$1 - (\lambda + \mu)\rho^{-1} = (1 - (\lambda + \mu))\rho^{-1} + (1 - \rho^{-1}) \geq \delta\rho m\rho^{-1} - \delta = \delta(m - 1).$$

If $m = 1$, then $1 - (\mu + \lambda) \geq \delta\rho$, and hence the distance between p and q is 0. In other words, $p = q$ such that S contains one point less on edge $\{u, v\}$. Else, if $m \geq 2$, we may add $m - 2 \geq 0$ further points between p and q such that the m points $S \cap P(G[\{u, v\}])$ have pairwise distance at least δ . Then, S contains one more point on edged $\{u, v\}$ than S' .

We prove that S is δ -auto-dispersed by showing the conditions of Lemma 26. Set S is edge internally δ -dispersed since for every edge we have added points that have pairwise distance at least δ . For condition (A2) it suffices to show that $\text{dr}_S(u, v) \geq \text{dr}_{S'}(u, v)\rho^{-1}$ for every pair (u, v) where $\{u, v\} \in E(G)$. Then for every distinct adjacent edges $\{u, v\}, \{u, w\} \in E(G)$ we have $\text{dr}_S(u, v) + \text{dr}_S(u, w) = (\text{dr}_S(u, v) + \text{dr}_S(u, w))\rho^{-1} \geq \delta\rho\rho^{-1} = \delta$ since Lemma 26 applies to the $(\delta\rho)$ -auto-dispersed set S' .

Hence, assume, for the sake of contradiction, that there are (u, v) with $\{u, v\} \in E(G)$ where $\text{dr}_S(u, v) < \text{dr}_{S'}(u, v)\rho^{-1}$. Among all such pairs (u, v) where $\text{dr}_S(u, v) < \text{dr}_{S'}(u, v)\rho^{-1}$, let (u, v) be where $\text{dr}_S(u, v)$ is minimum.

Consider that $\text{dr}_S(u, v) \leq 1$. Then S contains a point $p = p(u, v, \text{dr}_S(u, v))$. By construction, S' contains point $p' = p(u, v, \lambda')$ with $\lambda' = \text{dr}_S(u, v)\rho = \text{dr}_{S'}(u, v)$. Thus, actually, $\text{dr}_S(u, v) \leq \text{dr}_{S'}(u, v)\rho^{-1}$.

Otherwise, we have $\text{dr}_S(u, v) > 1$. Then $\text{dr}_S(u, v) = 1 + \text{dr}_S(v, w)$ for some neighbor $w \in N_G(v) \setminus \{u\}$. Because $\text{dr}_S(u, v) > \text{dr}_S(v, w)$, the assumption $\text{dr}_S(v, w) \geq \text{dr}_{S'}(v, w)\rho^{-1}$ holds. We will follow that $\text{dr}_S(u, v) \geq \text{dr}_S(u, v)\rho^{-1}$. Again, we use that Lemma 26 applies to S' , such that $\text{dr}_{S'}(v, u) \geq \delta\rho - \text{dr}_{S'}(v, w)$. By construction, set $S' \cap P(G[\{u, v\}])$ contains only one point. Thus $\text{dr}_{S'}(v, u) = 1 - \text{dr}_{S'}(u, v)$. It follows that

$$\begin{aligned} \text{dr}_S(u, v) &= 1 + \text{dr}_S(v, w) \\ &\geq (1 - \delta\rho + \text{dr}_{S'}(v, w))\rho^{-1} \\ &\geq (1 - \text{dr}_{S'}(v, u))\rho^{-1} \\ &= \text{dr}_S(u, v)\rho^{-1}. \end{aligned}$$

Hence, in both cases, we obtain the contradiction that $\text{dr}_S(u, v) \geq \text{dr}_S(u, v)\rho^{-1}$. Thus, by Lemma 26, S is δ -auto-dispersed. \blacktriangleleft

B Rounding the Distance

B.1 Proof of Lemma 9

We rely on the *direction* $\text{dir}(p \rightarrow q) \in \{u, v\}$ for distinct points $p, q \in P(G)$ defined as follows:

- For points $p = p(u, v, \lambda_p)$ and $q = p(u, v, \lambda_q)$ on a common edge $\{u, v\}$ with $\lambda_p < \lambda_q$, let $\text{dir}(p \rightarrow q) = v$. Let $\overline{\text{dir}}(p \rightarrow q) = u$.
- For points $p = p(u_p, v_p, \lambda_p)$ and $q = p(u_q, v_q, \lambda_q)$ on distinct edges $\{u_p, v_p\} \neq \{u_q, v_q\}$, let $\text{dir}(p \rightarrow q)$ be the unique vertex of $\{u_p, v_p\}$ that is contained in *every* shortest path between p and q , if such vertex exists. If $\text{dir}(p \rightarrow q)$ is defined, let $\overline{\text{dir}}(p \rightarrow q)$ be the unique vertex in $\{u_p, v_p\} \setminus \{\text{dir}(p \rightarrow q)\}$.

► **Lemma 28** ((Lemma 9 restated)). *For distinct points $p, q \in V(G_S)$, $\text{dir}(p \rightarrow q)$ is well-defined, unless p is half-integral.*

Proof. Let $p = p(u_p, v_p, \lambda_p)$ and $q = p(u_q, v_q, \lambda_q)$. If $\{u_p, v_p\} = \{u_q, v_q\}$, then $\text{dir}(p \rightarrow q)$ is defined for p, q . It remains to consider that $\{u_p, v_p\} \neq \{u_q, v_q\}$. Assume, for the sake of contradiction, that there are shortest paths $P_u, P_v \in P(G)$ which contain u_p respectively v_p . Let $\ell_u \in \mathbb{N}$ be the length of the path P_u without edges $\{u_p, v_p\}$ and $\{u_q, v_q\}$. Analogously let $\ell_v \in \mathbb{N}$ be the length of the path P_v without edges $\{u_p, v_p\}$ and $\{u_q, v_q\}$.

If P_u and P_v both contain u_q or both contain v_q , then their length may only differ by $|\lambda_p - (1 - \lambda_p)| < 1$. Since P_u and P_v have equal length, $\lambda_p = \frac{1}{2}$ in contradiction that p is not half-integral. Thus, up to symmetry, P_u contains u_q and P_v contains v_q .

First, assume that $q \in S \setminus \text{pivots}(S, \delta)$, such that p, q have distance δ . Then the distance between p and q also computes as $\lambda_p + \ell_u + \lambda_q = (1 - \lambda_p) + \ell_v + (1 - \lambda_q) = \delta$. In other words $2 + \ell_u + \ell_v = 2\delta$, which implies that δ is half-integral. Contradiction to Lemma 32.

Now, assume that $q \in \text{pivots}(S, \delta)$. Then $\lambda_q \in \{0, \frac{1}{2}, 1\}$. The length of path P_v is $\lambda_p + x$ for some $x \in \{\frac{z}{2} \mid z \in \mathbb{Z}\}$. Similarly the length of path P_u is $(1 - \lambda_p) + y$ for some $y \in \{\frac{z}{2} \mid z \in \mathbb{Z}\}$. Since λ_p is not a half integral, paths P_u and P_v have different length. Contradiction. \blacktriangleleft

B.2 Proof of Lemma 10

We work towards proving Lemma 10. We observe the following as an underlying principle.

► **Lemma 29.** *Consider points $p = p(u_p, v_p, \lambda_p)$, $q = p(u_q, v_q, \lambda_q)$ in distance $\delta' > 0$ and a shortest p, q -path that does not visit v_p, v_q . Let $x, y \in \mathbb{R}$, $x + y = 1$ and $\varepsilon \geq 0$. For $\varepsilon' \in [0, \varepsilon)$,*

let no point $p(u_p, v_p, \lambda_p + x\varepsilon')$, $p(u_q, v_q, \lambda_q + y\varepsilon')$ be half-integral nor $\delta' + \varepsilon'$ be half-integral. Then points $p(u_p, v_p, \lambda_p + x\varepsilon)$ and $p(u_q, v_q, \lambda_q + y\varepsilon)$ have distance $\delta' + \varepsilon$.

Proof. Let $p_\varepsilon := p(u_p, v_p, \lambda_p + x\varepsilon)$ and $q_\varepsilon := p(u_q, v_q, \lambda_q + y\varepsilon)$.

The simple case is that points p, q are from a common edge $\{u, v\}$. Then any shortest path between $p_\varepsilon, q_\varepsilon$ uses edge $\{u, v\}$ such that points $p_\varepsilon, q_\varepsilon$ have distance $\delta' + x\varepsilon + y\varepsilon = \delta' + \varepsilon$, as desired.

The interesting case is that p, q are from distinct edges. Then, there is a shortest path $P \subseteq P(G)$ of length δ' between the points p and q that uses vertices u_p and u_q . Let $Q \subseteq P$ be the subpath of P between the vertices u_p and u_q , which has some integer length $\ell \in \mathbb{N}_0$. Hence P has length $\delta' = \ell + \lambda_p + \lambda_q$.

Let $P' \subseteq P(G)$ be the path between the new vertices p_ε and q_ε consisting of Q and $\{p(u_p, v_p, \lambda) \mid 0 \leq \lambda \leq \lambda_p + x\varepsilon\}$ and $\{p(u_q, v_q, \lambda) \mid 0 \leq \lambda \leq \lambda_q + y\varepsilon\}$. Path P' has minimum length of $p_\varepsilon, q_\varepsilon$ -paths that visit vertices u_p and u_q . Further, P' has length $\ell + \lambda_p + \lambda_q + \varepsilon = \delta' + \varepsilon$.

Assume, for the sake of contradiction, that there is a path between p_ε and q_ε of length $< \delta' + \varepsilon$. Then this path may not visit u_p and u_q . Further, there is an $\varepsilon'' \in [0, \varepsilon)$ such that there is a shortest path P'' between $p_{\varepsilon''}$ and $q_{\varepsilon''}$ using not both of u_p and u_q and of length exactly $\delta' + \varepsilon''$. To see this, note that the distance of a shortest path between p_ε and q_ε that does not use both vertices u_p and u_q is a continuous function in ε . By symmetry assume that P'' does not use v_p .

- First, consider that P'' uses vertices v_p and u_q . Consider that $d(u_p, u_q) = d(v_p, u_q)$. Let path P^* , analogously to P' , consist of Q , $\{p(u_p, v_p, \lambda) \mid 0 \leq \lambda \leq \lambda_p + x\varepsilon''\}$ and $\{p(u_q, v_q, \lambda) \mid 0 \leq \lambda \leq \lambda_q + y\varepsilon''\}$. Then each of P'' and P^* has length $\delta' + \varepsilon''$. Since $d(u_p, u_q) = d(v_p, u_q)$, point $p_{\varepsilon''}$ has to have equal distance to u_p and to u_q . Thus, $p_{\varepsilon''}$ is the half-integral point $p(u_p, v_p, \frac{1}{2})$. A contradiction to that no point $p(u_p, v_p, \lambda_p + x\varepsilon')$ for $0 \leq \varepsilon' < \varepsilon''$ is half-integral.

If $d(u_p, u_q) \neq d(v_p, u_q)$, the distances $d(u_p, u_q)$ and $d(v_p, u_q)$ differ by at least one. Then by similar arguments it follows that $p_{\varepsilon''} = p(u_p, v_p, 0)$ or $p_{\varepsilon''} = p(u_p, v_p, 1)$, hence is half-integral. Again a contradiction to that no point $p(u_p, v_p, \lambda_p + x\varepsilon')$ for $0 \leq \varepsilon' < \varepsilon''$ is half-integral.

- Now, consider that path P'' uses vertices v_p and v_q . Assume that P and P'' intersect in an inner point. Then, they in particular intersect in a point that is at a vertex $u \notin \{u_p, v_p\}$. Analogously to the previous case, this implies that $p_{\varepsilon''}$ is half-integral. Thus P and P'' must have inner-points disjoint. Then $P \cup P'' \subseteq P(G)$ forms a cycle of length $2(\delta' + \varepsilon'')$ which is an integer. Hence $\delta' + \varepsilon''$ is half-integral. A contradiction to that no $\delta' + \varepsilon'$ for $\varepsilon' < \varepsilon$ is half-integral.

Thus, P' is indeed a shortest path between p_ε and q_ε , which has length $\delta' + \varepsilon$, as desired. ◀

To properly prove Lemma 10, we recall the technical framework.

An (S, δ) -pivot, or simply a pivot, is a half-integral point $r \in P(G)$ where two points $p, q \in S$, the witnesses, have distances $d(p, r) = d(r, q) = \frac{\delta}{2}$. Since S is δ -dispersed, there is a shortest p, q -path of length δ containing r . Let $\text{pivots}(S, \delta)$ be the set of (S, δ) -pivots, and let $W(S, \delta) \subseteq \binom{S}{2}$ be the family of pairs of points from S , that witness some (S, δ) -pivot.

We construct an auxiliary graph G_S on vertex set $S \cup \text{pivots}(S, \delta)$:

- For $\{p, q\} \in W(S, \delta)$ and for every point $r \in \text{pivots}(S, \delta)$ they witness, add edges $\{p, r\}, \{r, q\}$; and
- for every pair of points $\{p, q\} \in \binom{S}{2} \setminus W(S, \delta)$ with distance $d(p, q) = \delta$, add edge $\{p, q\}$. Note that, for every edge $\{r, p_1\}$ with $r \in \text{pivots}(S, \delta)$, there is at least one other edge $\{r, p_2\}$ such that p_1, p_2 witness pivot r .

A path $P = (p_0, p_1, \dots, p_s)$, $s \geq 1$ in graph G_S is a *spine* if points p_1, \dots, p_s are not half-integral. Note that any sub-sequence (p_0, \dots, p_i) , $1 \leq i \leq s$ is also a spine.

Consider a spine $P = (p_0, \dots, p_s)$. We define its half-integral *velocities* $\text{vel}_P : \{p_0, \dots, p_s\} \rightarrow \{\frac{z}{2} \mid z \in \mathbb{Z}\}$ depending on *signs* $\text{sgn}_P : \{p_1, \dots, p_s\} \rightarrow \{-1, 1\}$, which in turn depend on *flip* $_P : \{p_0, \dots, p_s\} \rightarrow \{-1, 1\}$. We may drop the subscript P , if it is clear from the context. Let $\text{vel}(p_0) = 0$. Let $\text{vel}(p_1) = \frac{1}{2}$, if $p_0 \in \text{pivots}(S, \delta)$, and let $\text{vel}(p_1) = 1$, if $p_0 \in S$. For $i \geq 1$, let

$$\text{vel}(p_{i+1}) := \text{vel}(p_i) + \text{sgn}(p_{i+1}).$$

Thus, $\text{sgn} \in \{-1, 1\}$ indicates whether the velocity increases or decreases. The current sgn is unchanged unless flip is negative. Let $\text{sgn}(p_1) = 1$. For $2 \leq i \leq s$, let

$$\text{sgn}(p_i) := \text{flip}(p_{i-1}) \text{sgn}(p_{i-1}) = \prod_{0 < j < i} \text{flip}(p_j),$$

where for $1 \leq i \leq s-1$, $\text{flip}(p_i) \in \{1, -1\}$, and $\text{flip}(p_i) = 1$ if and only if $\text{dir}(p_i \rightarrow p_{i-1}) \neq \text{dir}(p_i \rightarrow p_{i+1})$.

$$\text{flip}(p_i) := \begin{cases} 1, & \text{dir}(p_i \rightarrow p_{i-1}) \neq \text{dir}(p_i \rightarrow p_{i+1}), \\ -1, & \text{else.} \end{cases}$$

Value $\text{flip}(p_i)$ describes whether a shortest path between p_{i-1} and p_i and a shortest path between p_i and p_{i+1} overlap. If $\text{flip}(p_i) = 1$, they do not overlap. If $\text{flip}(p_i) = -1$, the paths overlap; and this will cause a flip for the sign, i.e., whether the speed is increasing or decreasing.

Let $\varepsilon \geq 0$ and consider a spine $P = (p_0, \dots, p_i)$. We move all points p_i for $1 \leq i \leq s$ by $\text{sgn}_P(p_i) \text{vel}_P(p_i) \varepsilon$ away from their predecessor p_{i-1} . Specifically, when λ_i is such that $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \lambda_i)$, we define the moved point as

$$(p_i)_{P,\varepsilon} := p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \lambda_i + \text{sgn}_P(p_i) \text{vel}_P(p_i) \varepsilon).$$

Doing so, points $(p_i)_{P,\varepsilon}, (p_{i+1})_{P,\varepsilon}$ maintain a distance of $\delta + \varepsilon$, under the condition that we do not encounter a half-integral point in the process and that $\delta + \varepsilon$ itself does not become half-integral.

► **Lemma 30.** *Let $P = (p_0, \dots, p_i)$ be a spine. For $\varepsilon' \in (0, \varepsilon)$, let neither $(p_{i-1})_{P,\varepsilon'}$ nor $(p_i)_{P,\varepsilon'}$ nor $\delta' + \varepsilon'$ be half-integral. Then points $\{(p_{i-1})_{P,\varepsilon}, (p_i)_{P,\varepsilon}\}$ are $(\delta + \varepsilon)$ -critical.*

Proof. For simplicity, let us write $(p_{i'})_\varepsilon$ instead of $(p_{i'})_{P,\varepsilon}$ for $0 \leq i' \leq i$.

If $i = 1$, then $\text{vel}(p_0) = 0$, and $p_0 = (p_0)_\varepsilon$ by definition. Further $\text{vel}(p_1) = 1$, since $p_0, p_1 \in S$. Also $\text{sgn}(p_1) = 1$. Let $p_1 = p(\cdot, \overline{\text{dir}}(p_1 \rightarrow p_0), \mu)$. Then $(p_1)_\varepsilon = p(\cdot, \overline{\text{dir}}(p_1 \rightarrow p_0), \mu + \text{sgn}(p_1) \text{vel}(p_1) \varepsilon) = p(\cdot, \overline{\text{dir}}(p_1 \rightarrow p_0), \mu + \varepsilon)$. There is a shortest path between p_0, p_1 that uses $\text{dir}(p_1 \rightarrow p_0)$. Hence Lemma 29 applies to old points p_0, p_1 and new points $(p_0)_\varepsilon, (p_1)_\varepsilon$ with $x = 0$ and $y = 1$. It follows that $d(p_\varepsilon, q_\varepsilon) = \delta + \varepsilon$, as desired.

Now consider that $i > 1$, and hence point p_{i-1} is not half-integral. Then there is an edge position $\lambda \in (-\frac{1}{2}, \frac{1}{2})$ such that

$$\begin{aligned} p_{i-1} &= p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2}), \frac{1}{2} + \lambda) \\ &= p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_i), \frac{1}{2} - \text{flip}(p_{i-1}) \lambda) \end{aligned}$$

For the new point $(p_{i-1})_\varepsilon$ there is some edge position $\lambda^* = \lambda + \text{sgn}(p_{i-1}) \text{vel}(p_{i-1})\varepsilon \in [-\frac{1}{2}, \frac{1}{2}]$ such that $(p_{i-1})_\varepsilon = p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2}), \frac{1}{2} + \lambda^*)$. Indeed, the new edge position may be integral. Still $\overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2})$ is defined, since it relies on the old point p_{i-1} . By Lemma 36,

$$\begin{aligned} (p_{i-1})_\varepsilon &= p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_i), \frac{1}{2} - \text{flip}(p_{i-1})\lambda^*) \\ &= p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_i), \frac{1}{2} - \text{flip}(p_{i-1})\lambda - \text{flip}(p_{i-1}) \text{sgn}(p_{i-1}) \text{vel}_S(p_{i-1})\varepsilon) \\ &= p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_i), \frac{1}{2} - \text{flip}(p_{i-1})\lambda - \text{sgn}(p_i) \text{vel}_S(p_{i-1})\varepsilon). \end{aligned}$$

Let edge position $\mu \in (-\frac{1}{2}, \frac{1}{2})$ be such that $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \mu)$. Then

$$\begin{aligned} (p_i)_\varepsilon &= p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \mu + \text{sgn}(p_i) \text{vel}(p_i)\varepsilon) \\ &= p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \mu + \text{sgn}(p_i)(\text{vel}(p_{i-1}) + \text{sgn}(p_i))\varepsilon) \\ &= p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \mu + (\text{sgn}(p_i) \text{vel}(p_{i-1}) + 1)\varepsilon). \end{aligned}$$

There is a shortest path between p_{i-1}, p_i that does not use $\overline{\text{dir}}(p_i \rightarrow p_{i-1})$ and $\overline{\text{dir}}(p_{i-1} \rightarrow p_i)$. Thus Lemma 29 applies to points $(p_{i-1})_\varepsilon, (p_i)_\varepsilon$ with $x = -\text{sgn}(p_i) \text{vel}_S(p_{i-1})$ and $y = \text{sgn}(p_i) \text{vel}_S(p_{i-1}) + 1$. It follows that $d(p_\varepsilon, q_\varepsilon) = \delta + \varepsilon$, as desired. \blacktriangleleft

► **Lemma 31.** *Let $P \subseteq P(G)$ be a path of points of length $\ell \in \mathbb{R}^+$. Then G contains a path of length at least $\lfloor \ell - 1 \rfloor$.*

Proof. Let $\{u_1, u_2\}, \{u_2, u_3\}, \dots, \{u_s, u_{s+1}\}$ be the list of edges visited by *non-integral* points when we follow the path P of length ℓ from start to finish. Beside possibly $\{u_1, u_2\} = \{u_s, u_{s+1}\}$, no edge may be repeated. Hence also $\lceil \ell \rceil \geq s$. Particularly none of the vertices u_2, u_3, \dots, u_s occurs twice in P . Thus u_2, u_3, \dots, u_s is a path in G consisting of at least $\lfloor \ell - 1 \rfloor$ edges. \blacktriangleleft

► **Lemma 32.** *δ is not half-integral.*

Proof. Assume, for the sake of contradiction, that δ is half-integral, hence that $\delta = a$ or $\delta = \frac{a}{2}$ for some $a \in \mathbb{N}$. Recall that $a > 2L + 2$. Then $\delta \geq \frac{a}{2} > L$. Recall that $|S| > 2$. Let $P \subseteq P(G)$ be a shortest path between distinct points $p, q \in S$, which is of length $\delta > L$. Then G contains a path of length at least L , as seen in Lemma 31. Contradiction. \blacktriangleleft

Recall the definition of the root points R . Let R_0 be the set of half-integral points in G_S . There may be some components of the auxiliary graph G_S without a point in R_0 . Let R result from R_0 by adding exactly one point from every component that has no point in R_0 . Now, when we restrict spines to those that begin with a point from R , the implied movement of points is uniquely determined.

For each point p , the new point is p_ε , and the resulting set is $S_\varepsilon := \{p_\varepsilon \mid p \in S\}$. For a point $p_0 \in R$, simply $(p_0)_\varepsilon := p_0$. For other points $p_i \in S \setminus R$, hence that are the i -th point, for some $i \geq 1$, of some spine $P = (p_0, \dots, p_i)$ with start $p_0 \in R$, point $(p_i)_\varepsilon$ is defined as for the spine P , formally $(p_i)_\varepsilon := (p_i)_{P, \varepsilon}$. By Lemma 11, the definition of p_ε is independent of the considered spine.

We further recall ε^* and its defining events. Let $\varepsilon^* \geq 0$ be minimum such that $\delta + \varepsilon^* = \delta^*$ or one of the following events occurs:

- (Event 1) For a δ -uncritical pair of points $\{p, q\} \in \binom{S}{2}$, now $\{p_{\varepsilon^*}, q_{\varepsilon^*}\}$ $(\delta + \varepsilon)$ -critical.
- (Event 2) a non-half-integral $p \in S$, now p_{ε^*} half-integral, or
- (Event 3) for $r \in P(G) \setminus \text{pivots}(S, \delta)$, now $r \in \text{pivots}(S_{\varepsilon^*}, \delta + \varepsilon^*)$.

We claim that the minimum $\varepsilon^* \geq 0$ is defined. This is due to that the above events depend on continuous functions in ε , which are the distance of p_ε to its closest half-integral point, and the distance between points p_ε and q_ε for $p, q \in S$.

► **Lemma 33** ((Lemma 10 restated)). *For a spine $P = (p_0, \dots, p_s)$ and $i \in \{0, \dots, s-1\}$, points $(p_i)_{P, \varepsilon}, (p_{i+1})_{P, \varepsilon}$ are $(\delta + \varepsilon)$ -critical for the maximal $\varepsilon \leq \delta^* - \delta$ that is limited by the Events 1, 2, 3.*

Proof. This is an immediate consequence from Lemma 30. Because of Lemma 32 $\delta + \varepsilon'$ is not half-integral for $\varepsilon' \in (0, \varepsilon)$ is $\varepsilon' \in$. That event 2 did not occur before implies that neither $(p_i)_{P, \varepsilon'}$ nor $(p_{i+1})_{P, \varepsilon'}$ is half-integral for $\varepsilon' \in (0, \varepsilon)$ is $\varepsilon' \in$. ◀

B.3 Proof of Lemma 12

We delay the (lengthy and technical) proof of Lemma 11 to the next subsection. Here we prove the monotonicity result, Lemma 12.

► **Lemma 34** ((Lemma 12 restated)). *Let S be a δ -dispersed set for $\delta < \delta^*$ and ε^* defined as above. Then:*

- (1) S_{ε^*} is a $(\delta + \varepsilon^*)$ -dispersed set of size $|S|$.
- (2) If $\{p, q\} \in \binom{S}{2}$ is δ -critical, then $\{p_{\varepsilon^*}, q_{\varepsilon^*}\}$ is $(\delta + \varepsilon^*)$ -critical.
- (3) If $r \in \text{pivots}(S, \delta)$, then $r \in \text{pivots}(S_{\varepsilon^*}, \delta + \varepsilon^*)$.

Proof. To show claims (1), (2), (3), we prove the following more convenient version:

- (1') For every $\{p, q\} \in \binom{S}{2} \setminus (E(G_S) \cup W(S, \delta))$ we have $d(p_{\varepsilon^*}, q_{\varepsilon^*}) \geq \delta + \varepsilon^*$.
- (3') For every $\{p, q\} \in \binom{S}{2} \cap W(S, \delta)$ that witness some pivot $r \in \text{pivots}(S, \delta)$, we have $\{p_{\varepsilon^*}, q_{\varepsilon^*}\} \in W(S_{\varepsilon^*}, \delta + \varepsilon^*)$ and $r \in \text{pivots}(S_{\varepsilon^*}, \delta + \varepsilon^*)$. Moreover, for every pivot $r \in \text{pivots}(S, \delta)$ with edge $\{r, p\} \in E(G_S)$, we have $\{r, p_{\varepsilon^*}\} \in E(G_{S_{\varepsilon^*}})$.
- (2') For every $\{p, q\} \in \binom{S}{2} \cap E(G_S)$, we have $d(p_{\varepsilon^*}, q_{\varepsilon^*}) = \delta + \varepsilon^*$.

Here (1'), (3'), (2') is a case-distinction for (1). Statement (3) follows from (3') since any pivot r has some witnesses p, q . Claim (2) for edges in G_S involving a pivot is shown by (3'). Claim (2) for the remaining edges $\{p, q\} \in \binom{S}{2} \cap E(G_S)$ follows from (2'). So it remains to show the above claims.

- (1') Assume, for the sake of contradiction, that there are points $\{p, q\} \in \binom{S}{2} \setminus (E(G_S) \cup W(S, \delta))$ where $d(p_{\varepsilon^*}, q_{\varepsilon^*}) < \delta + \varepsilon^*$. Then there is an $\varepsilon' \in [0, \varepsilon^*)$ such that points $p_{\varepsilon'}, q_{\varepsilon'}$ have distance exactly $\delta + \varepsilon'$. Then event Event 1 should have triggered for ε' and points $p_{\varepsilon'}, q_{\varepsilon'}$. A contradiction to the minimality of ε^* .
- (3') Consider points $\{p, q\} \in \binom{S}{2} \cap W(S, \delta)$ that witness some pivot $r \in \text{pivots}(S, \delta)$. Especially p, q may not be half-integral due to Lemma 32. Hence there are edge positions $\lambda_p, \lambda_q \in (0, 1)$ such that $p = p(\cdot, \overline{\text{dir}}(p \rightarrow r), \lambda_p)$ and $q = p(\cdot, \overline{\text{dir}}(q \rightarrow r), \lambda_q)$. Further, the two-element sequences $P = (r, p)$ and $Q = (r, q)$ are spines. By construction, we have that $\text{vel}_P(p) = \text{vel}_Q(q) = \frac{1}{2}$. It follows that the new points are at $p_{\varepsilon^*} = p(\cdot, \overline{\text{dir}}(p \rightarrow r), \lambda_p + \frac{1}{2}\varepsilon^*)$ and $q_{\varepsilon^*} = p(\cdot, \overline{\text{dir}}(q \rightarrow r), \lambda_q + \frac{1}{2}\varepsilon^*)$. Then Lemma 29 applies to points $p_{\varepsilon^*}, q_{\varepsilon^*}$ with $\delta' = \delta$, $\varepsilon' = \varepsilon^*$ and with $x + y = \frac{1}{2} + \frac{1}{2} = 1$, since $\delta + \varepsilon^* \leq \delta^*$ and Event 2 has not occurred. It follows that $d(p_{\varepsilon^*}, q_{\varepsilon^*}) = \delta + \varepsilon^*$, as desired. Further points $p_{\varepsilon^*}, q_{\varepsilon^*}$ still witness r as a $(\delta + \varepsilon^*)$ -pivot. In other words $r \in \text{pivots}(S_{\varepsilon^*}, \delta + \varepsilon^*)$.

Finally, consider pivot $r \in \text{pivots}(S, \delta)$ of S with an edge $\{r, p\} \in E(G_S)$. Then there is $q \in S$ such that p, q witness pivot r . As just observed, then $d(p_{\varepsilon^*}, r) = \frac{\delta + \varepsilon^*}{2}$ and hence $\{p_{\varepsilon^*}, r\} \in E(G_{S_{\varepsilon^*}})$.

- (2') Consider points $\{p, q\} \in \binom{S}{2} \cap E(G_S)$. We will show that $d(p_{\varepsilon^*}, q_{\varepsilon^*}) = \delta + \varepsilon^*$. There is a spine $P = (p_0, \dots, p_i)$ with $p_0 \in R$ and $\{p_{i-1}, p_i\} = \{p, q\}$. By symmetry, we may assume that $p_{i-1} = p$ and $p_i = q$. Since event Event 2 did not occur for $\varepsilon' \in (0, \varepsilon)$ Lemma 30 applies. It follows that $d(p_{\varepsilon^*}, q_{\varepsilon^*}) = \delta + \varepsilon^*$.

This finishes the proof. \blacktriangleleft

B.4 Proof of Lemma 11

Now we work towards proving Lemma 11.

- **Lemma 35** ((Lemma 11 restated)). *Let $\delta < \delta^*$. Consider spines $P = (p_0, \dots, p_i)$ and $Q = (q_0, \dots, q_j)$ with $p_0, q_0 \in R$ and $p_i = q_j$. Then (1) $\text{vel}_P(p_i) = \text{vel}_Q(q_j)$; and (2) $\overline{\text{dir}}(p_i \rightarrow p_{i-1}) = \overline{\text{dir}}(q_j \rightarrow q_{j-1})$ if and only if $\text{sgn}_P(p_i) = \text{sgn}_Q(q_j)$.*

We begin by explicitly stating the edge position of a point p_i of a spine $P = (p_0, \dots, p_i)$ depending on $\text{vel}(p_i)$, $\text{sgn}(p_i)$ and the edge position λ_0 of p_0 . Then our final proof is by contradiction. It leads the assumption that different spines that end in the same point do not agree on vel and sgn , such that there must be a path G longer than L , a contradiction.

Consider a point p_i that is part of a spine $P = (p_0, \dots, p_i)$. We will state its edge position μ_i depending on its velocity $\text{vel}(p_i)$, its sign $\text{sgn}(p_i)$ and the edge position λ_0 of p_0 . We orientate the edge position μ_i according to $\overline{\text{dir}}(p_i \rightarrow p_{i-1})$, its direction to the predecessor point p_{i-1} . That is, let μ_i be such that $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \mu_i)$; hence μ_i is a part of the length of any shortest path between p_i and p_{i-1} , specifically the part using the edge of p_i (assuming q is on another edge).

We give some intuition: Assume that we already have some expression for the edge position μ_{i-1} of the previous point p_{i-1} , hence we know the value $\mu_{i-1} \in (0, 1)$ such that $p_{i-1} = p(\cdot, \overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2}), \mu_{i-1})$. Consider that a shortest p_{i-1}, p_i -path starts from p_{i-1} in the direction of vertex $\overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2})$, in other words $\overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2}) \neq \overline{\text{dir}}(p_{i-1} \rightarrow p_i)$ and by definition $\text{flip}(p_{i-1}) = 1$. Then it is not difficult to follow that $\mu_i = \text{fp}(\mu_{i-1} + \delta)$ where $\text{fp}(y) = y - \lfloor y \rfloor$ is the *fractional part* of $y \in \mathbb{R}$. If however $\text{flip}(p_{i-1}) = -1$, hence $\overline{\text{dir}}(p_{i-1} \rightarrow p_{i-2}) = \overline{\text{dir}}(p_{i-1} \rightarrow p_i)$, then point p_i has the same edge position as point p_{i-2} but ‘flipped’. More precisely, when $\mu_{i-1} \in (0, 1)$ is such that $p_{i-2} = p(\cdot, \overline{\text{dir}}(p_{i-2} \rightarrow p_{i-3}), \mu_{i-2})$, we have $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), 1 - \mu_{i-2})$.

To handle this flip of the edge position more conveniently, let us specify the edge positions relatively to the middle $\frac{1}{2}$. That is, we are interested in the (shifted) edge position $\lambda_i \in (-\frac{1}{2}, \frac{1}{2})$ such that $p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \lambda_i)$ for $i \geq 1$. It follows that:

- **Lemma 36.** *For a spine $P = (p_0, \dots, p_{i+1})$, $i \geq 1$, there is a $\lambda \in (-\frac{1}{2}, \frac{1}{2})$ such that $p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \lambda) = p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i+1}), \frac{1}{2} - \text{flip}(p_{i-1})\lambda)$.*

As noted earlier, if the predecessor point p_{i-1} has $\text{flip}(p_{i-1}) = -1$, the sign of the edge position of p_i should flip. This sign change is tracked by sgn . In turn $\text{vel}(p_i)$ sums up the values of sgn of points p_1, \dots, p_i , and hence allows to state the edge position relatively to that of p_0 . For $\lambda \in (-\frac{1}{2}, \frac{1}{2})$ and $x \in \{\frac{z}{2} \mid z \in \mathbb{Z}\}$, let

$$\text{pos}_\lambda(x) := \text{fp}\left(\frac{1}{2} + \lambda + x\delta\right) - \frac{1}{2}.$$

Consider that λ is such that $p_0 = p(\cdot, \overline{\text{dir}}(p_0 \rightarrow p_1), \frac{1}{2} + \lambda)$, assuming for now that p_0 is non-half-integral. Then $\text{pos}_\lambda(\text{vel}(p_i))$ is the distance of the edge position of p_i to the middle $\frac{1}{2}$, in other words, the edge position of p_i up to the sign.

Later we rely on the following observation:

► **Lemma 37.** $\text{pos}_\lambda(x + y) = \text{pos}_{\text{pos}_\lambda(x)}(y)$, for every $\lambda \in [-\frac{1}{2}, \frac{1}{2})$ and $x, y \in \{\frac{z}{2} \mid z \in \mathbb{Z}\}$.

Further, let us state the edge position even for half-integral points, as for example it may be the case for p_0 . To do so, we extend the definition of $\text{dir}(q \rightarrow p)$ to also apply for half-integral p : For points $p, q \in P(G)$, let $\text{dir}^*(p \rightarrow q)$ be equal to $\text{dir}(q \rightarrow p)$ when $\text{dir}(q \rightarrow p)$ is defined. Else, p must be half-integral as observed in Lemma 9. If p is such that $p = (u, v, \frac{1}{2})$, fix $\text{dir}^*(p \rightarrow q)$ to be u or v arbitrarily, and let $\overline{\text{dir}}^*(q \rightarrow p)$ be the unique vertex in $\{u, v\} \setminus \{\text{dir}^*(q \rightarrow p)\}$. Else, if p is such that $p = p(u, v, 0)$ for some edge $\{u, v\} \in E(G)$, let $\text{dir}^*(p \rightarrow q) = u$ and let $\overline{\text{dir}}^*(p \rightarrow q) = v'$ where v' is an arbitrary neighbor of v in G . As a result, we may for example define $\lambda_0 \in [-\frac{1}{2}, \frac{1}{2})$ such that $p_0 = p(\cdot, \overline{\text{dir}}^*(p \rightarrow q), \frac{1}{2} + \lambda_0)$. Here again the missing entry is irrelevant for λ_0 .

Moreover, let us also extend the definition of a spine to allow a final half-integral point. A path $P = (p_0, \dots, p_s)$, $s \geq 1$ in graph G_S is a *generalized spine* if (p_0, \dots, p_{s-1}) is a spine. Values vel , sgn and flip are defined analogously as we did for spines (still based on $\overline{\text{dir}}$). Also vel and sgn are defined for p_s . Only $\text{flip}(p_s)$ is undefined which however is not used.

► **Lemma 38.** Let $P = (p_0, \dots, p_s)$ be a generalized spine. Let $\lambda \in [-\frac{1}{2}, \frac{1}{2})$ be such that $p_0 = p(\cdot, \overline{\text{dir}}^*(p_0 \rightarrow p_1), \frac{1}{2} + \lambda)$. Then, for $i \in \{1, \dots, s\}$,

$$p_i = p(\cdot, \overline{\text{dir}}^*(p_i \rightarrow p_{i-1}), \frac{1}{2} + \text{sgn}(p_i) \cdot \text{pos}_\lambda(\text{vel}(p_i))).$$

Proof. Before we prove the actual statement, let us consider how the edge positions of some non-half-integral points p, q in distance $\delta' \in \{\frac{\delta}{2}, \delta\}$ relate. Let $\mu_p \in [0, 1)$ be such that $p = p(\cdot, \text{dir}^*(p \rightarrow q), \mu_p)$, and let $\mu_q \in [0, 1)$ be such that $q = p(\cdot, \overline{\text{dir}}^*(q \rightarrow p), \mu_q)$. If p, q are on the same edge, then $\mu_q = \mu_p + \delta'$. If p, q are not on the same edge, there is a shortest path P between p and q that uses vertices $\text{dir}^*(p \rightarrow q)$ and $\text{dir}^*(q \rightarrow p)$. Let integer $\ell \geq 0$ be the length of path P restricted to the path between $\text{dir}^*(p \rightarrow q)$ and $\text{dir}^*(q \rightarrow p)$. Then $\delta' = (1 - \mu_p) + \ell + \mu_q$. Hence $\mu_p + \delta' = \ell + \mu_q + 1$, which implies $\lfloor \mu_p + \delta' \rfloor = \ell + 1$, since $\ell, 1$ are integer and $\lfloor \mu_q \rfloor = 0$. Then

$$\mu_q = \mu_p + \delta' - \lfloor \mu_p + \delta' \rfloor. \quad (1)$$

This also applies when p, q are on the same edge, where we followed $\mu_q = \mu_p + \delta' \in [0, 1)$. Let us state the edge positions relatively to the middle $\frac{1}{2}$. Let $\lambda_p \in [-\frac{1}{2}, \frac{1}{2})$ be such that $p = p(\cdot, \text{dir}^*(p \rightarrow q), \frac{1}{2} + \lambda_p)$, and let $\lambda_q \in [-\frac{1}{2}, \frac{1}{2})$ be such that $q = p(\cdot, \overline{\text{dir}}^*(q \rightarrow p), \frac{1}{2} + \lambda_q)$. That means $\frac{1}{2} + \lambda_p = \mu_p$ and $\frac{1}{2} + \lambda_q = \mu_q$. It follows that

$$\lambda_q = \lambda_p + \delta' - \lfloor \frac{1}{2} + \lambda_p + \delta' \rfloor = \text{pos}_{\lambda_p}(\frac{\delta'}{\delta}). \quad (2)$$

Now, we are ready to prove the statement that

$$p_i = p(\cdot, \overline{\text{dir}}^*(p_i \rightarrow p_{i-1}), \frac{1}{2} + \text{sgn}(p_i) \cdot \text{pos}_\lambda(\text{vel}(p_i)))$$

by induction on $i \in \{1, \dots, s\}$.

Induction base, $i = 1$: Let δ' be the distance between p_0, p_1 which is either δ or $\frac{\delta}{2}$. Then p_1 has velocity $\text{vel}(p_1) = \frac{\delta'}{\delta}$. Note that Equation 2 applies for points $p_0 = p, p_1 = q$ and $\lambda_p = \lambda$. It follows that $p_1 = p(\cdot, \overline{\text{dir}}^*(p_1 \rightarrow p_0), \frac{1}{2} + \text{pos}_\lambda(\text{vel}(p_1)))$. Since $\text{sgn}(p_1) = 1$, we are done.

Induction step, $i \rightsquigarrow i + 1$, $i \geq 1$: Note that point p_i is non-half-integral. Let $z := \text{pos}_\lambda(\text{vel}(p_i))$. By the induction hypothesis,

$$\begin{aligned}
 p_i &= p(\cdot, \overline{\text{dir}}^*(p_i \rightarrow p_{i-1}), \frac{1}{2} + \text{sgn}(p_i)z) \\
 &= p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \text{sgn}(p_i)z), && | p_i \text{ is non-half-integral} \\
 &= p(\cdot, \text{dir}(p_i \rightarrow p_{i+1}), \frac{1}{2} + \text{flip}(p_i) \text{sgn}(p_i)z), && | \text{Lemma 36} \\
 &= p(\cdot, \text{dir}^*(p_i \rightarrow p_{i+1}), \frac{1}{2} + \text{flip}(p_i) \text{sgn}(p_i)z) \\
 &= p(\cdot, \text{dir}^*(p_i \rightarrow p_{i+1}), \frac{1}{2} + \text{sgn}(p_{i+1})z).
 \end{aligned}$$

Hence, the edge position of point p_i is $\lambda_i = \text{sgn}(p_{i+1})z$ when defined as above. Let value $\lambda_{i+1} \in [-\frac{1}{2}, \frac{1}{2})$ be such that $p_{i+1} = p(\cdot, \overline{\text{dir}}^*(p_{i+1} \rightarrow p_i), \frac{1}{2} + \lambda_{i+1})$. Then Equation 2 applies for points $p = p_i$ and $q = p_{i+1}$ with edge positions λ_i and λ_{i+1} , respectively, and distance between them of $\delta' = \delta$. We define $\lambda'_{i+1} := \text{sgn}(p_{i+1})\lambda_{i+1}$. Thus we have to show that $\lambda'_{i+1} = \text{pos}_\lambda(\text{vel}(p_{i+1}))$. We have that $\lambda'_{i+1} = \text{sgn}(p_{i+1}) \cdot \text{pos}_{\lambda_i}(1)$, according to Equation 2.

First, consider that $\text{sgn}(p_{i+1}) = 1$, and hence $\text{vel}(p_{i+1}) = \text{vel}(p_i) + 1$. Then

$$\lambda'_{i+1} = \text{pos}_{\lambda_i}(1) = \text{pos}_{\text{pos}_\lambda(\text{vel}(p_i))}(1) \stackrel{\text{Obs. 37}}{=} \text{pos}_\lambda(\text{vel}(p_i) + 1) = \text{pos}_\lambda(\text{vel}(p_{i+1})),$$

as desired. Now, consider that $\text{sgn}(p_{i+1}) = -1$, and hence $\text{vel}(p_{i+1}) = \text{vel}(p_i) - 1$. We use the fact that $\lfloor -x \rfloor = \lfloor -x + 1 \rfloor$ for non-integral x . Then

$$\begin{aligned}
 \lambda'_{i+1} &= -\text{pos}_{\lambda_i}(1) \\
 &= -(\lambda_i + \delta - \lfloor \frac{1}{2} + \lambda_i + \delta \rfloor) \\
 &= z - \delta + \lfloor \frac{1}{2} - z + \delta \rfloor \\
 &= z - \delta - \lfloor \frac{1}{2} + z - \delta \rfloor \\
 &= \text{pos}_z(-1) \\
 &= \text{pos}_\lambda(\text{vel}(p_i) - 1) && | \text{Lemma 37}
 \end{aligned}$$

as desired. ◀

Now, we show that a generalized spine (p_0, p_1, \dots, p_s) implies a path of length $\lfloor |\text{vel}(p_s)|\delta \rfloor$ in G . Roughly speaking, we can combine shortest paths between the pairs $(p_0, p_1), (p_1, p_2), \dots$ to obtain a long path in $P(G)$, which in turn implies a long path in G .

► **Lemma 39.** *Let $P = (p_0, p_1, \dots, p_s)$ be a generalized spine. Then G contains a path of length $\lfloor |\text{vel}(p_s)|\delta - 1 \rfloor$.*

Proof. Before we begin, let us rule out the special case that $\text{vel}(p_{i-1}) = \frac{1}{2}$ and $\text{vel}(p_i) = -\frac{1}{2}$, $i \in \{1, \dots, s\}$, which would imply that p_{i-1} and p_i witness a pivot.

▷ **Claim 40.** For no $i \in \{1, \dots, s\}$ we have $\text{vel}(p_{i-1}) = \frac{1}{2}$ and $\text{vel}(p_i) = -\frac{1}{2}$.

Proof. Clearly we have $i \geq 2$ since $\text{vel}(p_0) = 0$. There is a position $\lambda \in (-\frac{1}{2}, \frac{1}{2})$ such that $p_{i-1} = p(\cdot, \overline{\text{dir}}^*(p_{i-1} \rightarrow p_{i-2}), \frac{1}{2} + \lambda)$ and $p_i = p(\cdot, \overline{\text{dir}}^*(p_i \rightarrow p_{i-1}), \frac{1}{2} - \lambda)$, according to Lemma 38. First assume that $\text{flip}(p_{i-1}) = 1$ hence that $\text{sgn}(p_{i-1}) = \text{sgn}(p_i)$. Then we know that $p_{i-1} = p(\cdot, \overline{\text{dir}}^*(p_{i-1} \rightarrow p_i), \frac{1}{2} - \lambda)$, as seen in Lemma 36. It follows that points p_{i-1}, p_i have integer distance $\frac{1}{2} + \lambda + \frac{1}{2} - \lambda + \ell = 1 + \ell$ for some $\ell \in \mathbb{N}$, hence that δ is integer, a contradiction to Lemma 32.

Thus, it remains to consider that $\text{flip}(p_{i-1}) = -1$ and hence that $\text{sgn}(p_{i-1}) \neq \text{sgn}(p_i)$. Then we know that $p_{i-1} = p(\cdot, \overline{\text{dir}}^*(p_{i-1} \rightarrow p_i), \frac{1}{2} + \lambda)$, as seen in Lemma 36. That is,

points p_{i-1}, p_i have some equal distance to a half-integral point $r \in P(G)$, which is midway on some path between the vertices $\text{dir}^*(p_{i-1} \rightarrow p_i)$ and $\text{dir}^*(p_i \rightarrow p_{i-1})$ (even when p_{i-1} and p_i are on the same edge). Then p_i, p_{i-1} witness the pivot r , in contradiction to that $\{p_i, p_{i-1}\} \in E(G_S)$. \triangleleft

Further, we claim that it is safe to assume that $\text{vel}(p_s)$ is positive and has maximum velocity among all points in P . Clearly if $\text{vel}(p_s) = 0$ there is nothing to prove for the lemma. Now consider that $\text{vel}(p_s) < 0$. Then there is an index $i \in \{1, \dots, s-1\}$ where $\text{vel}(p_i) \in \{\frac{1}{2}, 0\}$ and $\text{vel}(p_j) < 0$ for every $j \in \{i+1, \dots, s\}$. Particularly, $\text{vel}(p_i) = 0$ since otherwise it contradicts Claim 40. Consider the spine $P' = (p_i, \dots, p_s)$. Observe that $\text{sgn}_P(p_i) = -1 = -\text{sgn}_{P'}(p_i)$ and $\text{vel}_P(p_i) = 0 = \text{vel}_{P'}(p_i)$. Further, $\text{flip}_P(p_j) = \text{flip}_{P'}(p_j)$ for $j \in \{i+1, \dots, s\}$. Inductively, it follows that $\text{sgn}_{P'}(p_j) = -\text{sgn}_P(p_j)$ and hence $\text{vel}_{P'}(p_j) = -\text{vel}_P(p_j)$, for $j \in \{1, \dots, s\}$. Thus, if $\text{vel}_P(p_s) < 0$, let us consider the spine P' with non-negative $\text{vel}_{P'}(p_s) = -\text{vel}_P(p_s)$ instead of P . Hence, it is safe to only consider the case that $\text{vel}(p_s) > 0$. Now, if $\text{vel}(p_i) \geq \text{vel}(p_s)$, for some $i < s$, consider the sub-spine (p_0, \dots, p_i) instead, which has the same flip , sgn and vel values. Since $\text{vel}(p_i) \geq \text{vel}(p_s) > 0$ we have that $i \geq 1$. Hence, in the following, we may then assume that $z = \text{vel}(p_s)$ is positive and has maximum velocity among the points of P .

Let us define shortest paths $P_i \subseteq P(G)$ between p_{i-1} and p_i for $i \in \{1, \dots, s\}$. For convenience, let $P_i := \emptyset$ for $i \in \mathbb{N} \setminus \{1, \dots, s\}$. For $i \in \{1, \dots, s\}$, let P_i be some shortest path between p_{i-1} and p_i with maximum intersection with the previous path P_{i-1} . Possibly their intersection is only $\{p_{i-1}\}$, in case that $\text{flip}(p_{i-1}) = 1$. Hence, the intersection of paths P_{i-1} and P_i forms a path starting at p_{i-1} . It follows that $P_i \setminus P_{i-1}$ is a path.

For $j \geq 1$, let $Q_j \subseteq P(G)$ be the set of points resulting from the union of $P_i \setminus (P_{i-1} \cup P_{i+1})$ for $i \leq j$, i.e., $Q_j = \bigcup_{i \leq j} (P_i \setminus (P_{i-1} \cup P_{i+1}))$. We will show that Q_s is a path of length at least $z\delta$. Then it follows that G contain a path of length at least $z\delta$, according to Lemma 31. First, let us study the intersection of the described paths.

\triangleright **Claim 41.** For $i, j \in \{1, \dots, s\}$ and $i < j$ we have

- (A) the interior of P_i and P_j are disjoint, unless $j = i+1$ and $\text{flip}(p_i) = -1$; and
- (B) path $P_i \setminus P_{i-1}$ has length $\geq \frac{\delta}{2}$. Symmetrically, $P_i \setminus P_{i+1}$ has length $\geq \frac{\delta}{2}$.

Above (A), (B) and that p_0, \dots, p_s are pairwise distinct imply that Q_s is a path in $P(G)$.

Proof. (A) Let I be the intersection of interior points of P_i and the interior points of P_j . Assume, for the sake of contradiction, that $I \neq \emptyset$.

Consider that $j = i+1$ and $\text{flip}(p_i) = 1$, meaning $\text{dir}(p_i \rightarrow p_{i-1}) \neq \text{dir}(p_i \rightarrow p_{i+1})$. Let $p_i = p(u, v, \lambda)$ for some $\lambda \in (0, 1)$. By the definition of $\text{dir}(p_i \rightarrow p_{i-1})$ and $\text{dir}(p_i \rightarrow p_{i+1})$, the paths P_i and P_{i+1} and $\{p(u, v, \lambda') \mid \lambda' \in (0, 1)\}$ intersect only in p_i . Then there exists a point $p \in I$ with minimum distance to p_i . Let P_i^* be the subpath of P_i from p_i to p . Analogously, P_{i+1}^* be the subpath of P_{i+1} from p_i to p . Then, P_i^* is a shortest p, p_i -path. Analogously, P_{i+1}^* is a shortest p, p_{i+1} -path. We claim that $P_i^* \cup P_{i+1}^*$ forms a cycle. If not, there is a point $p' \in (P_i^* \cap P_{i+1}^*) \setminus \{p, p_i\}$, where $d(p', p_i) < d(p, p_i)$. Then $p' \in I$ has smaller distance to p_i than p , in contradiction to the definition of p . Further, we claim that p is at a vertex. Assuming otherwise, p is at the position $p = p(u', v', \lambda')$ for some $\lambda' \in (0, 1)$. Then $p(u', v', 0) \in I$ or $p(u', v', 1) \in I$ and hence p cannot have minimum distance to p_i .

Since p is integral, paths P_i^*, P_{i+1}^* have equal length and $P_i^* \cup P_{i+1}^*$ forms a cycle, point p_i must also be half-integral. A contradiction to the definition of a generalized spine.

It remains to consider that $j \neq i+1$. We distinguish between possible lengths of paths P_i, P_j , which must be δ or $\frac{\delta}{2}$. Fix point $p \in I$ arbitrarily. All cases lead to a contradiction:

- Case, paths P_i and P_j both have length $\frac{\delta}{2}$. Then $p_i, p_{j-1} \in S$ such that they have distance δ . However, $d(p_i, p_{j-1}) \leq d(p_i, p) + d(p, p_{j-1}) < \frac{\delta}{2} + \frac{\delta}{2}$. Contradiction.
- Case, paths P_i and P_j have length δ . Then $p_i, p_{j-1}, p_{j-1}, p_j \in S$ such that they pairwise have distance at least δ . Further $d(p_{i-1}, p_i) = \delta$ and $d(p_{j-1}, p_j) = \delta$. It follows that the distance of respectively $p_i, p_{j-1}, p_{j-1}, p_j$ to p is exactly $\frac{\delta}{2}$. Then however, p_{i-1}, p_i witness the pivot p , and hence they may not be part of a spine. Contradiction.
- $\{P_i, P_j\} = \{P', P''\}$ where P' has length δ and P'' has length $\frac{\delta}{2}$. Path P' connects some points $p', p'' \in S$. Then $d(p', p) \leq \frac{\delta}{2}$. Path P'' connects a pivot and some point $q \in S$. Then $d(q, p) < \frac{\delta}{2}$. Thus, $d(q, p') \leq d(q, p) + d(p, p') < \delta$, in contradiction to that S is δ -dispersed.

(B) We show that $P_i \setminus P_{i-1}$ has length $\geq \frac{\delta}{2}$, for $i \in \{1, \dots, s\}$. Then that $P_i \setminus P_{i+1}$ has length $\geq \frac{\delta}{2}$, for $i \in \{0, \dots, s-1\}$, follows from considering the reversed generalized spine p_s, p_{s-1}, \dots, p_0 .

If $\text{flip}(p_{i-1}) = 1$, then (A) applies and we are done. Thus consider $\text{flip}(p_{i-1}) = -1$.

If p_{i-2} exists and $p_{i-2} \in \text{pivots}(S, \delta)$, then $p_{i-2} = p_0$ is a pivot. Hence $\text{vel}(p_1) = \frac{1}{2}$. Having $\text{flip}(p_1) = -1$ implies that $\text{vel}(p_2) = -\frac{1}{2}$. Then points p_1, p_2 contradict Claim 40.

If p_i exists and $p_i \in \text{pivots}(S, \delta)$, then p_i is a pivot and hence $p_i = p_s$. Having that $\text{flip}(p_{i-1}) = -1$ it follows that $\text{vel}(p_{i-1}) > \text{vel}(p_i) = z$, which we excluded earlier.

If p_{i-1} exists and $p_{i-1} \in \text{pivots}(S, \delta)$, then p_{i-1} is a pivot and hence $p_0 = p_{i-1}$. Especially p_{i-2} does not exist, such that (B) follows immediately for $P_i \setminus P_{i-1}$.

Thus it remains to consider that points p_{i-2}, p_{i-1}, p_i , if they exist, are in S . If paths P_i and P_{i-1} have inner points disjoint, we are done. Hence consider that they intersect in a point $p \notin \{p_{i-2}, p_{i-1}, p_i\}$. Since $p_{i-2} \neq p_i$, there is a well-defined intersection point p that maximizes the distance to p_{i-1} . If $d(p, p_{i-2}) \geq \frac{\delta}{2}$, then also $P_i \setminus P_{i-1}$ has length $\geq \frac{\delta}{2}$. Thus consider that $d(p, p_{i-2}) < \frac{\delta}{2}$. Then $d(p, p_{i-1}) > \frac{\delta}{2}$. Further $d(p, p_{i+1}) < \frac{\delta}{2}$ since $p \in P_i$. Then we have the contradiction $d(p_{i-2}, p_i) \leq d(p_{i-2}, p) + d(p, p_i) < \delta$. \triangleleft

Let index i_x describe the first point that reaches a velocity $x \in \{1, \dots, z\}$, which is $i_x := \min\{i_x \mid \text{vel}(p_{i_x}) \geq x\}$. We prove by induction on $x \in \{1, \dots, \text{vel}(p_s) - 1\}$ that the path $Q_{i_{x+1}-1}$ has length $x\delta$. (We use that $x < \text{vel}(p_s)$, implies that the point of index i_{x+1} exists.) For $x = \text{vel}(p_s)$ we have $i_x = s$ and we will follow that $Q_s = s\delta$. Recall that the difference in the velocity of consecutive points is always ± 1 with exceptions at the very beginning and at the very end.

Induction base, $x = 1$: Then $i_x = 1$ and $Q_1 = P_1$ has length $\text{vel}(p_1) \in \{\frac{1}{2}, 1\}$, as desired.

Induction step, $x - 1 \rightsquigarrow x$: By the induction hypothesis, path Q_{i_x-1} has length at least $\delta(x - 1)$. By the definition of i_x , the velocity increased twice before reaching i_x , in other words $\text{vel}(p_{i_x-1}) = x - 1$ and $\text{vel}(p_{i_x-2}) = x - 2$. That means $\text{flip}(p_{i_x-1}) = 1$. Thus the path Q_{i_x-1} ending at p_{i_x-1} has inner points disjoint from P_j for $j \geq i_x$, according to Claim 41. If $i_x = s$, then path Q_s consists of the disjoint parts Q_{s-1} and P_s , and hence has length $(\text{vel}(p_{s-1}) + (\text{vel}(p_s) - \text{vel}(p_{s-1})))\delta = \text{vel}(p_s)\delta$, as desired. Else, if $i_x < s$, consider the point with index $i_{x+1} - 1$. It remains to show that $Q_{i_{x+1}-1} \setminus Q_{i_x-1}$ has length at least δ .

If $i_{x+1} - 1 = i_x$, then $Q_{i_{x+1}-1} \setminus Q_{i_x-1}$ consists only of P_{i_x} which has length δ , as desired.

Else, consider that $i_{x+1} - 1 \neq i_x$. Then $P_{i_x} \setminus \bigcup_{j \neq i_x} P_j$ has length $\frac{\delta}{2}$, according to Claim 41. Again the velocity increased twice before reaching i_{x+1} , and hence $\text{flip}(p_{i_{x+1}-1}) = 1$. That means $P_{i_{x+1}-1} \setminus \bigcup_{j \neq i_{x+1}-1} P_j$ has length $\frac{\delta}{2}$. Hence $Q_{i_{x+1}-1} \setminus Q_{i_x-1}$ has length at least $2\frac{\delta}{2} = \delta$, as desired.

As discussed within the induction step, for $i_x = s$ we follow that Q_s has length $z\delta$. It follows that G has a path of length $\lfloor z\delta - 1 \rfloor$ as a non-induced subgraph, according to Lemma 31. \blacktriangleleft

► **Corollary 42.** *Let $P = (p_0, p_1, \dots, p_s)$ be a generalized spine. Let $\delta = \frac{a}{b}$ be a rational number. Then $|\text{vel}(p_s)| < \frac{b}{2}$.*

Proof. Assume, for the sake of contradiction, that $|\text{vel}(p_s)| \geq \frac{b}{2}$. Then by Lemma 39 there is a path in G of length $\lfloor |\text{vel}(p_s)|\delta - 1 \rfloor \geq \lfloor \frac{b}{2} \cdot \frac{a}{b} - 1 \rfloor = \lfloor \frac{a}{2} - 1 \rfloor$. Hence $a \leq 2L + 2$. A contradiction to that $\frac{a^*}{b^*} > \delta$ where $\frac{a^*}{b^*}$ is smallest possible rational $\frac{a^*}{b^*} \geq \delta$ for co-prime a^*, b^* and $a^* < 2L + 2$. \blacktriangleleft

Finally, we are able to prove Lemma 11.

► **Lemma 43** ((Lemma 11 restated)). *Let $\delta < \delta^*$. Consider spines $P = (p_0, \dots, p_i)$ and $Q = (q_0, \dots, q_j)$ with $p_0, q_0 \in R$ and $p_i = q_j$. Then (1) $\text{vel}_P(p_i) = \text{vel}_Q(q_j)$; and (2) $\overline{\text{dir}}(p_i \rightarrow p_{i-1}) = \overline{\text{dir}}(q_j \rightarrow q_{j-1})$ if and only if $\text{sgn}_P(p_i) = \text{sgn}_Q(q_j)$.*

Proof. By the definition of the root points R , points p_0, q_0 are either both half-integral or $p_0 = q_0$, since p_0, q_0 must be from the same connected component of G_S . First, let us rule out that p_0 is integral and q_0 is at the middle position of an edge (and analogously the symmetric case). Consider the walk $p_0, p_1, \dots, p_{i-1}, q_j, q_{j-1}, \dots, q_0$ in G_S . Remove any loop from this walk to obtain a path $P^* = (p_0^*, p_1^*, \dots, p_k^*)$ with $p_0^* = p_0 \neq p_k^* = q_0$. Then P^* is a generalized spine. Lemma 38 applies with $\lambda = -\frac{1}{2}$. Therefore $\text{sgn}(p_i) \cdot \text{pos}_\lambda(\text{vel}(p_i))$ is the edge position of q_0 (relatively to the middle $\frac{1}{2}$) which must be 0. Hence $\text{pos}_\lambda(\text{vel}(p_i)) = \text{fp}(\text{vel}(q_0)\delta) - \frac{1}{2} = 0$. Then δ must be a rational $\frac{a}{b}$ with co-prime $a, b \in \mathbb{N}$ and $|\text{vel}_{P^*}(q_0)|$ is a multiple of $\frac{b}{2}$, which contradicts Corollary 42.

Thus it remains to consider that p_0, q_0 are equal, are both integral or both are at the middle of an edge. Let $\lambda_0 \in [-\frac{1}{2}, \frac{1}{2})$ be such that $p_0 = p(\cdot, \overline{\text{dir}}^*(p_0 \rightarrow p_1), \frac{1}{2} + \lambda_0)$. Analogously, let $\lambda'_0 \in [-\frac{1}{2}, \frac{1}{2})$ be such that $p_0 = p(\cdot, \overline{\text{dir}}^*(q_0 \rightarrow q_1), \frac{1}{2} + \lambda'_0)$. We observe that there is a value $\beta_0 \in \{-1, 1\}$ such that $\lambda'_0 = \beta_0 \lambda_0$. If $\lambda'_0 = \lambda_0 = 0$, we fix β_0 to be 1.

Since $p_i = q_j$, their edge positions defined relatively to their spine P and Q are equal up to the sign. Also recall that $p_i = q_j$ is not half-integral. That means there is an edge position $\lambda \in (-\frac{1}{2}, \frac{1}{2}) \setminus \{0\}$ and a value $\beta' \in \{-1, 1\}$ such that

$$p_i = p(\cdot, \overline{\text{dir}}(p_i \rightarrow p_{i-1}), \frac{1}{2} + \lambda) = p(\cdot, \overline{\text{dir}}(q_j \rightarrow q_{j-1}), \frac{1}{2} + \beta'\lambda) = q_j.$$

Figuratively speaking, β' indicates whether spines P, Q reach $p_i = q_j$ from the same direction. We observe that $\overline{\text{dir}}(p_i \rightarrow p_{i-1}) = \overline{\text{dir}}(q_j \rightarrow q_{j-1})$ if and only if $\beta' = 1$. Hence for the claim (2) we have to show that $\beta := \beta' \text{sgn}_P(p_i) \text{sgn}_Q(q_j) = 1$.

Let $x = \text{vel}_P(p_i)$ and $y = \text{vel}_Q(q_j)$. For the claim (1) we have to show that $x = y$. By Lemma 38 we have

$$\lambda = \text{sgn}_P(p_i) \text{pos}_{\lambda_0}(x) \quad \text{and} \quad \beta'\lambda = \text{sgn}_Q(q_j) \text{pos}_{\beta_0 \lambda_0}(y).$$

Then $\text{pos}_{\beta_0 \lambda_0}(y) = \beta \text{pos}_{\lambda_0}(x)$ for above defined $\beta \in \{-1, 1\}$. Hence $\text{pos}_{\lambda_0}(x) = \beta \text{pos}_{\beta_0 \lambda_0}(y)$, which is equivalent to

$$\text{fp}\left(\frac{1}{2} + \lambda_0 + x\delta\right) - \frac{1}{2} = \beta \left(\text{fp}\left(\frac{1}{2} + \beta_0 \lambda_0 + y\delta\right) - \frac{1}{2}\right).$$

We distinguish the cases whether $\beta_0, \beta \in \{-1, 1\}$.

- Case $\beta_0 = \beta = 1$: Then claim (2) holds. Further $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) = \text{fp}(\frac{1}{2} + \lambda_0 + y\delta)$, which implies that $\text{fp}(x\delta) = \text{fp}(y\delta)$. Assume, for the sake of contradiction, claim (1) is false, hence that $x \neq y$. Then δ is a rational $\frac{a}{b}$ with co-prime $a, b \in \mathbb{N}$. By symmetry, we may assume that $|x| < |y|$. Also x, y are both either non-negative or non-positive. Then $(y - x)\delta \in \mathbb{N}$, and hence $y - x$ is a multiple of b . Then $|y| \geq \frac{b}{2}$ in contradiction to Corollary 42.

All other cases lead to a contradiction, regardless whether $x \neq y$.

- Case $\beta_0 = \beta = -1$: Then $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) - \frac{1}{2} = -\text{fp}(\frac{1}{2} - \lambda_0 + y\delta) + \frac{1}{2}$. Thus $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) + \text{fp}(\frac{1}{2} - \lambda_0 + y\delta) = 0$. Hence $(\frac{1}{2} + \lambda_0 + x\delta) + (\frac{1}{2} - \lambda_0 + y\delta) = (x + y)\delta \in \mathbb{Z}$. This implies that δ is a rational $\frac{a}{b}$ with co-prime $a, b \in \mathbb{N}$, and that $|x + y|$ is a multiple of b . For either $z \in \{x, y\}$ we have $|z| \geq \frac{|x+y|}{2}$. By symmetry, assume that $|x| \geq \frac{|x+y|}{2}$. Thus $|x|$ is larger than a multiple of $\frac{b}{2}$, in contradiction to Corollary 42.
- Case $\beta = 1$ and $\beta_0 = -1$: Then $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) = \text{fp}(\frac{1}{2} - \lambda_0 + y\delta)$. Thus $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) - \text{fp}(\frac{1}{2} - \lambda_0 + y\delta) = 0$. That means $(\frac{1}{2} + \lambda_0 + x\delta) - (\frac{1}{2} - \lambda_0 + y\delta) \in \mathbb{Z}$ and hence $2\lambda_0 + (x - y)\delta \in \mathbb{Z}$. Let $z := \frac{x-y}{2}$. Then $\text{pos}_{\lambda_0}(z) = \text{fp}(\frac{1}{2}(1 + 2\lambda_0 + 2z\delta)) - \frac{1}{2} \in \{0, -\frac{1}{2}\}$, i.e., it is half-integral. Hence no point $p \in \{p_1, \dots, p_i\}$ has velocity $\text{pos}_{\lambda_0}(\text{vel}_P(p)) = z$, since p is not half-integral. Thus $z \notin \{x, y\}$. Further $z \neq 0$; Indeed assuming that $z = 0$, implies that $2\lambda_0 + 0\delta \in \mathbb{Z}$ and hence that λ_0 is half-integral. In that case, however, $\beta_0 = 1$, and hence this is covered by other cases. Now, by symmetry, we may assume that $z > 0$. Either z occurs as a potential $\text{vel}_P(p_{i'})$ or $z - \frac{1}{2}, z + \frac{1}{2}$ occur as potentials $\text{vel}_P(p_{i'}), \text{vel}_P(p_{i+1'})$ for some $j \in \{1, \dots, i-1\}$. The former implies that $p_{i'}$ is half-integral. The latter implies that $p_{i-1}, p_{i'}$ witness a pivot half-way on a shortest path between them. Both cases contradict the definition of the spine P .
- Case $\beta = -1$ and $\beta_0 = 1$: Then $\text{fp}(\frac{1}{2} + \lambda_0 + x\delta) + \text{fp}(\frac{1}{2} + \lambda_0 + y\delta) = 1$. That means $\frac{1}{2} + \lambda_0 + x\delta + \frac{1}{2} + \lambda_0 + y\delta \in \mathbb{Z}$, and hence $2\lambda_0 + (x + y)\delta \in \mathbb{Z}$. Let $z = \frac{x+y}{2}$. Then $\text{pos}_{\lambda_0}(z) = \text{fp}(\frac{1}{2}(1 + 2\lambda_0 + (x + y)\delta)) \in \{0, -\frac{1}{2}\}$, i.e., it is half-integral. Hence no point $p \in \{p_1, \dots, p_i\}$ has velocity $\text{pos}_{\lambda_0}(\text{vel}_P(p)) = z$, since p is not half-integral. Note that $z \notin \{0, x, y\}$ since $x, y \geq 1$. Then, analogously to the previous case it follows either contradiction that an inner point of P, Q is half-integral or two points witness a pivot.

To summarize, we always have $\beta = \beta_0 = 1$ and there claim (1) and (2) hold. ◀

C Hardness Results

C.1 Diameter

► **Lemma 44** ((Lemma 16 restated)). *For every $\delta > 3$, δ -DISPERSION is NP-complete and $W[1]$ -hard with parameter solution size, even for connected chordal graphs of diameter $\leq \lceil \delta \rceil$.*

Construction: Let $\delta > 3$ be arbitrary. Given a INDEPENDENT SET-Instance consisting of a graph G and integer k , we construct a δ -DISPERSION-instance consisting of a graph G' and integer $k' = k$. For every edge $\{u, v\} \in E(G)$, add a vertex $w_{\{u,v\}}$ to $V(G')$. Let $w_{\{u,v\}}$ for $\{u, v\} \in E(G)$ form a clique. For every vertex $u \in V(G)$ add a vertex u' to $V(G')$ and make it adjacent to $w_{\{u,v\}}$ for every $v \in N_G(u)$. For every $u \in V(G)$, add a path of length $\lceil \frac{\delta}{2} \rceil - 2 \geq 0$ starting at u' and terminating at a vertex which we call u_1 . Finally, if $\lceil \delta \rceil$ is even, add a true twin u_2 to u_1 for every $u \in V(G)$, that is u_2 is adjacent to $\{u_1\} \cup N_{G'}(u_1)$.

The diameter of G' is at most the distance between u_1 and v_1 for non-adjacent $u, v \in V(G)$, hence at most $2(\lceil \frac{\delta}{2} \rceil - 2) + 3 \leq \delta$. Observe that G' is connected and chordal. Further, the construction is possible in polynomial time. Moreover, INDEPENDENT SET is $W[1]$ -hard and the reduction preserves the parameter. Hence, in order to show NP-hardness and $W[1]$ -hardness, it remains to show the correctness:

► **Lemma 45.** *Graph G has an independent set of size k if and only if G' has a δ -dispersed set of size k .*

Proof. (\Rightarrow) Let $I \subseteq V(G)$ be an independent set of G . If $\lceil \delta \rceil$ is odd, then let $S \subseteq P(G)$ be the point set consisting of points p_u at the vertices $u \in I$. Then, every $p_u, p_v \in S$ have distance $2(\lceil \frac{\delta}{2} \rceil - 2) + 3 \leq \delta$, since $\{u, v\} \notin E(G)$. Else, when $\lceil \delta \rceil$ is even, let $S := \{p(u_1, u_2, \frac{1}{2}) \mid u \in I\}$. Then, every $p(u_1, u_2, \frac{1}{2}), p(v_1, v_2, \frac{1}{2})$ have distance $2(\lceil \frac{\delta}{2} \rceil - 2) + 4 \leq \delta$, again since $\{u, v\} \notin E(G)$. Thus, in both cases, S is δ -dispersed.

(\Leftarrow) Let $S \subseteq P(G')$ be a δ -dispersed set of G' . Let $C \subseteq P(G')$ be the set of points located at the central clique, that is the set of points $p(u, v, \lambda)$ with $u, v \in V(G)$ and $\lambda \in [0, 1]$. Consider that S contains a point $p \in C$, hence a point $p = p(u, v, \lambda)$ with $u, v \in V(G)$ and $\lambda \in [0, 1]$. Then p has distance to any vertex u' for $u \in V(G)$ of at most $\frac{3}{2}$. Thus, p has distance to any other point in $P(G')$ of at most $\frac{3}{2} + \lceil \frac{\delta}{2} \rceil - 2 < \delta$. That means, $|S| = 1$ and any vertex in G forms an independent set of the same size. Hence it remains to consider that $S \cap C = \emptyset$.

Let ball $B_u \subseteq P(G)$, for $u \in V(G)$, consist of every point in distance $\lceil \frac{\delta}{2} \rceil - 1 \geq 1$ to u_1 or u_2 (if u_2 exists). Then the balls B_u for $u \in V(G)$ and C together make up the whole set of points $P(G)$. Clearly, every ball B_u , for $u \in V(G)$, contains at most one point from S . Then $I := \{u \in V(G) \mid S \cap B_u \neq \emptyset\}$ has size $|S|$. We claim that I is an independent set. First consider that $\delta > 4$. Consider an edge $\{u, v\} \in E(G)$. Vertices u' and v' have the common neighbor $w_{u,v}$, hence have distance 2. Any point $p \in B_u$, for $u \in V(G)$, has distance at most $\lceil \frac{\delta}{2} \rceil - \frac{3}{2}$ to u' . Hence, any distinct points $p, q \in B_u \cup B_v$ have distance at most $2(\lceil \frac{\delta}{2} \rceil - \frac{3}{2}) + 1 < \delta$. Therefore, for adjacent $u, v \in V(G)$ either $S \cap B_u \neq \emptyset$ or $S \cap B_v \neq \emptyset$, such that $|I \cap \{u, v\}| = 1$, hence that I is an independent set of G .

A remaining special case is when $\delta \leq 4$. Then u', v' have a true twins u_2 and v_2 respectively, the distance of any u', u_2 to any v', v_2 is at most 2. Again, any point $p \in B_u$ has distance at most $\lceil \frac{\delta}{2} \rceil - \frac{3}{2}$ to either $u' = u_1$ or its true twin u_2 . Thus, analogously follows that I is an independent set of G . ◀

C.2 Pathwidth and Feedback Vertex Set Size

► **Theorem 46** ((Theorem 17 restated)). *DISPERSION is $W[1]$ -hard parameterized by $\text{pw}(G) + k$. Further, there is no $n^{o(\sqrt{\text{pw}(G)} + \sqrt{k})}$ -time algorithm unless ETH fails. DISPERSION is $W[1]$ -hard parameterized by $\text{fvs}(G) + k$. Further, there is no $n^{o(\text{fvs}(G) + \sqrt{k})}$ -time algorithm unless ETH fails.*

In MULTI-COLORED-INDEPENDENT-SET (MCIS) we are given a graph G with a partition of $V(G)$ into k independent sets, called *color classes*, V_1, \dots, V_k , each of size n . The task is to find a *multi-colored independent set*, an independent set that contains exactly one vertex from each color class.

Construction (as in [11]) Let the given MCIS consist of graph G where $V(G)$ partitions into k color classes V_1, \dots, V_k , each of size n . We construct a DISPERSION instance consisting of a graph G' , distance $\delta := 6n$ that asks for a set of size at least $k' := k^2$.

- For every color class V_i , $1 \leq i \leq k$: Enumerate the vertices in V_i as v_1^i, \dots, v_n^i . Create a set $P_i \subseteq V(G')$ of n vertices p_1^i, \dots, p_n^i ; Add vertex a_i and connect it to each p_ℓ^i by a path of length $n + \ell$ for $1 \leq \ell \leq n$; Similarly add a vertex b_i and connect it to each p_ℓ^i by a path of length $n + \ell$ for $1 \leq \ell \leq n$.
- For every non-edge $e = \{v_{j_1}, v_{j_2}\} \in V_{i_1} \times V_{i_2} \setminus E(G)$ (where $i_1 \neq i_2$) add a vertex u_e . We connect u_e to a_{i_1}, b_{i_1} and to a_{i_2}, b_{i_2} by paths of lengths that encode the indices j_1 and j_2 : Add a path between u_e and a_{i_1} of length $5n - j_1$ and add a path between u_e and b_{i_1} of length $4n + j_1$; Similarly between u_e and a_{i_2} of length $5n - j_2$ and add a path between u_e and b_{i_2} of length $4n + j_2$.
- For every distinct indices $i_1, i_2 \in \{1, \dots, k\}$ introduce a path of length $6n - 1$ with start vertex g_{i_1, i_2} and end vertex g'_{i_1, i_2} . Make g_{i_1, i_2} adjacent to every u_e with $e \in V_{i_1} \times V_{i_2}$.

A multi-colored independent set $I \subseteq V(G)$ translates to a $6n$ -dispersed set of size k^2 by selecting the corresponding p_j^i vertices and for every pair of selected $p_{j_1}^{i_1}, p_{j_2}^{i_2}$, for $i_1 \neq i_2$, selecting g'_{i_1, i_2} and u_e where $e = \{v_{j_1}^{i_1}, v_{j_2}^{i_2}\}$. Similarly, a maximum $6n$ -dispersed is forced to select vertices corresponding to an independent set and non-edges between those in the original graph G . The details of the correctness easily follow from the reduction of that of Katsikarelis et al. [11].

The resulting graph G' has a feedback vertex set of size $\mathcal{O}(k)$ which consists of a_i, b_i for every $1 \leq i \leq k$. Also G' has pathwidth $\mathcal{O}(k^2)$: Removing all a_i, b_i vertices and all g_{i_1, i_2} vertices results in a forest of stars which has a constant width pathwidth decomposition. By adding the $\mathcal{O}(k^2)$ removed vertices to every bag, we obtain a pathwidth decomposition of G' and of width $\mathcal{O}(k^2)$.

The bounds on the feedback vertex set size and pathwidth imply lower bounds assuming the Exponential Time Hypothesis (ETH). MCIS cannot be solved in time $n^{o(k)}$ assuming ETH is true [4]. However, assuming an $n^{o(\sqrt{\text{pw}(G)} + \sqrt{k})}$ -time algorithm for DISPERSION implies an $n^{o(k)}$ -time algorithm for MCIS. Similarly, an $n^{o(\text{fvs}(G) + \sqrt{k})}$ -time algorithm for DISPERSION implies an $n^{o(k)}$ -time algorithm for MCIS.

C.3 Treedepth

► **Theorem 47** ((Theorem 18 repeated)). *Assuming ETH, there is no $2^{o(\text{td}(G)^2)}$ -time algorithm for DISPERSION.*

To show a lower bound regarding the treedepth, a start is to use same reduction as for parameter pathwidth. However, this alone provides only a lower bound of $n^{\mathcal{O}(k + \log \ell)}$ where ℓ is the number of colors of the original MCIS-instance. To avoid the dependency on n we further use a reduction from 3-SAT to MCIS that outputs instance with relatively small ℓ (similar as in [11]).

First, consider the treedepth of G' of the resulting DISPERSION-instance when the original MCIS-instance consists of a graph G with k color classes each containing ℓ vertices. The treedepth of G' is at most $\mathcal{O}(k + \log \ell)$, as observed in [11]: To see this, consider the components of graph G' after removing the $2k$ vertices a_i, b_i for $1 \leq i \leq k$. The treedepth of G' is then at most $2k$ plus the maximum treedepth of the remaining components. The remaining components are a tree for each vertex g_{i_1, i_2} and a path of length 3ℓ for each vertex p_j^i . A path of length 3ℓ has treedepth $\mathcal{O}(\log \ell)$. For a tree containing some g_{i_1, i_2} , note that after removing g_{i_1, i_2} only paths of length $< 6\ell$ remain. Hence, each remaining tree has treedepth $\mathcal{O}(\log \ell)$. In total G' has treedepth $\mathcal{O}(k + \log \ell)$.

Now, we give a (non-polynomial time) reduction from 3-SAT to MCIS. There, a 3-SAT-instance φ with N variables and $\mathcal{O}(N)$ clauses reduces to a MCIS-instance with \sqrt{N} color classes each containing $c^{\sqrt{N}}$ vertices, for some constant c . Further, the run-time is $\mathcal{O}(\sqrt{N} \cdot c^{2\sqrt{N}})$.

Construction (similar to [11]¹): Group the clauses of φ into $\ell = \mathcal{O}(\sqrt{N})$ groups F_1, \dots, F_ℓ each of size $\mathcal{O}(\sqrt{N})$. Thus, every group F_i consists of $\mathcal{O}(\sqrt{N})$ variables, which have $2^{\mathcal{O}(\sqrt{N})}$ truth assignments. Let c be such that $2^{\mathcal{O}(\sqrt{N})} = c^{\sqrt{N}}$. Let $\varphi_1^i, \dots, \varphi_s^i$ for $s \leq c^{\sqrt{N}}$ be an enumeration of truth assignments to the variables of F_i that satisfy the clauses of F_i , for each $i \in \ell$. For each truth assignments φ_j^i introduce a corresponding vertex u_j^i of color i . Add an edge between u_j^i and $u_{j'}^{i'}$ for $i \neq i'$, if the truth assignments φ_j^i and $\varphi_{j'}^{i'}$ are contradicting each other in variables they have in common.

It easily follows that φ is satisfiable if and only if the constructed graph G' has a multi-colored independent set of size ℓ . The run-time is $\mathcal{O}(\sqrt{N} \cdot c^{\sqrt{N}} + c^{2\sqrt{N}})$. For more details, we refer to [11].

By combining these two reductions, 3-SAT reduces to DISPERSION with treedepth $\mathcal{O}(k + \log \ell) = \mathcal{O}(\sqrt{N} + \log c^{\sqrt{N}}) = \mathcal{O}(\sqrt{N})$. Assuming a $2^{o(\text{td}(G)^2)}$ -time algorithm for DISPERSION implies a $2^{o(N)}$ -time algorithm for 3-SAT with N variables and $\mathcal{O}(N)$ clauses, contradicting ETH [10]. This finishes the proof of Theorem 47.

¹ The reduction there directly reduces to DISTANCE INDEPENDENT SET. The idea is essentially to use the following reduction to MCIS as the first step.