

ASYMPTOTIC GEOMETRY OF THE MODULI SPACE OF RANK TWO IRREGULAR HIGGS BUNDLES OVER THE PROJECTIVE LINE

GAO CHEN AND NIANZI LI

ABSTRACT. We study the asymptotic behavior of Hitchin's hyperkähler metric on the moduli space of rank two irregular Higgs bundles over $\mathbb{C}P^1$, with all tame singularities (if any) assumed to be strongly parabolic. When the moduli space is four-dimensional, we show that Hitchin's metric is asymptotic to an ALG/ALG* model metric.

1. INTRODUCTION

Many conjectures on the moduli of Higgs bundles require a deeper understanding of the ends of the moduli space, among which the most well-known are the GMN (Gaiotto-Moore-Neitzke) conjecture [GMN10] on the asymptotic metric and Hausel's conjecture [Hau01] on the vanishing of the L^2 -cohomology. A consequence of the GMN conjecture is that $g_{L^2} - g_{sf} = O(e^{-ct})$ along a generic ray parametrized by $t \in \mathbb{R}^+$, where g_{L^2} is the natural complete hyperkähler metric on the moduli space, and g_{sf} is the semiflat metric [Fre99] arising from the algebraic integrability. With analytic techniques, many progresses have been made towards proving this result. The pioneering work [MSWW19] used fiducial solutions and limiting configurations in [MSWW16] to establish the polynomial decay of the above difference. In [DN19], Dumas and Neitzke used the local biholomorphic flow to show that the decay is exponential on the Hitchin section. These techniques were combined in [Fre18, Fre20] to prove the exponential decay for $SL(n, \mathbb{C})$ -Higgs bundles. Recently in [FMSW22], the authors extended this result to the moduli space of parabolic $SL(2, \mathbb{C})$ -Higgs bundles, where the Higgs fields admit simple poles, and they showed that the moduli space of $SL(2, \mathbb{C})$ -Higgs bundles with four strongly parabolic points over $\mathbb{C}P^1$ is an ALG gravitational instanton.

The goal of this paper is to extend the result of [FMSW22] to the moduli space of irregular (wild) Higgs bundles, which means the Higgs fields can have higher order poles. Irregular Higgs bundles have received an increasing interest in recent years, because of their appearances in isomonodromic deformations [Boa01], wild character varieties [Boa14], and the geometric Langlands program [Wit08]. In [BB04], Biquard and Boalch established the wild non-abelian Hodge correspondence, which exhibits the moduli space of irregular Higgs bundles as a complete hyperkähler manifold. When the moduli space is four-dimensional, it is expected to be a gravitational instanton of ALG type [Boa12]. In this paper we consider the moduli space \mathcal{M} of rank two irregular Higgs bundles over $\mathbb{C}P^1$. Using the methods of [FMSW22] and the analytic results of [BB04], we obtain the following main result.

Theorem 1.1. *Fix a generic curve $[(\bar{\partial}_E, \Phi_t)]$ in \mathcal{M} , and an infinitesimal deformation $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]}\mathcal{M}$. As $t \rightarrow \infty$, there exist positive constants c, σ such that*

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2 - \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{sf}}^2 = O(e^{-ct^\sigma}).$$

The first named author is supported by the Project of Stable Support for Youth Team in Basic Research Field, Chinese Academy of Sciences, YSBR-001.

The precise definition of the curve $[(\bar{\partial}_E, \Phi_t)]$ will be given in Section 3. Later in Section 6.3, we will specialize to the case when the moduli space is (real) four dimensional, with this result, we are able to show that g_{L^2} is polynomially close to an ALG/ALG* model metric g_{model} . Moreover, the constants in the difference $g_{L^2} - g_{\text{model}}$ can be shown to be independent of the choice of the curve $[(\bar{\partial}_E, \Phi_t)]$. The correspondences between the singularity configurations and the types of the model metrics are listed below (see also [Boa18, p. 448]). The tilde ($\tilde{}$) on the pole indicates a twisted irregular type (defined in Section 2).

Kodaira type	II^*	III^*	IV^*
Dynkin diagram	A_0	A_1	A_2
β	$\frac{5}{6}$	$\frac{3}{4}$	$\frac{2}{3}$
τ	$e^{2\pi i/3}$	i	$e^{2\pi i/3}$
D	$4 \cdot \{\tilde{0}\}$	$4 \cdot \{0\}$ or $3 \cdot \{\tilde{0}\} + \{\infty\}$	$3 \cdot \{0\} + \{\infty\}$

TABLE 1. ALG

In the previous table, (β, τ) means that \mathcal{M} is asymptotic to the standard ALG model $(X, G_{\beta, \tau})$ of type (β, τ) . Here X is obtained by identifying two boundary components of

$$\{u \in \mathbb{C} \mid \text{Arg}(u) \in [0, 2\pi\beta] \text{ and } |u| \geq R\} \times \mathbb{C}_v / (\mathbb{Z} \oplus \mathbb{Z}\tau)$$

via the gluing map $(u, v) \sim (e^{2\pi i\beta}u, e^{-2\pi i\beta}v)$. $G_{\beta, \tau}$ is a flat hyperkähler metric on X such that $\omega^1 = i/2(du \wedge d\bar{u} + dv \wedge d\bar{v})$ and $\omega^2 + i\omega^3 = du \wedge dv$. Kodaira type means that \mathcal{M} is biholomorphic to a rational elliptic surface minus a fiber with the given Kodaira type [CC21]. Dynkin diagram means that $H^2(\mathcal{M})$ is generated by the given extended Dynkin diagram. This makes sense because ALG gravitational instantons with the same β are diffeomorphic to each other [CV21].

Kodaira type	I_4^*	I_3^*	I_2^*	I_1^*
Dynkin diagram	D_0	D_1	D_2	$D_3 = A_3$
D	$2 \cdot \{\tilde{0}\} + 2 \cdot \{\tilde{\infty}\}$	$2 \cdot \{0\} + 2 \cdot \{\tilde{\infty}\}$	$2 \cdot \{0\} + 2 \cdot \{\infty\}$ or $\{0\} + \{1\} + 2 \cdot \{\tilde{\infty}\}$	$\{0\} + \{1\}$ $+ 2 \cdot \{\infty\}$

TABLE 2. ALG*

In the previous table, Kodaira type means that \mathcal{M} is biholomorphic to a rational elliptic surface minus a fiber with the given Kodaira type [CV21]. Dynkin diagram means that $H^2(\mathcal{M})$ is generated by the given extended Dynkin diagram. This makes sense because ALG* gravitational instantons with the same Kodaira type at infinity are diffeomorphic to each other [CV21].

To prove Theorem 1.1, we wish to use a similar strategy as that in [FMSW22]:

- Construct the harmonic metric h_t for $(\bar{\partial}_E, t\Phi)$ by a gluing method, with the following building blocks.
 - Limiting metric h_∞ , which is singular at the points where the spectral cover is ramified.
 - Fiducial solutions h_t^{model} in some fixed disks around the ramification points.
 - Desingularize h_∞ by h_t^{model} to obtain h_t^{app} , and perturb h_t^{app} to h_t .

- Interpolate g_{app} between g_{L^2} and g_{sf} , compare tangent vectors in Coulomb gauges using L^2 descriptions of these metrics.

However, there are two major new challenges in our case. The first is the lack of a natural \mathbb{C}^* action on \mathcal{M} since the irregular types are fixed. Fortunately, irregular Higgs bundles over $\mathbb{C}P^1$ can be explicitly described, and we can find some curve $[(\bar{\partial}_E, \Phi_t)]$ in \mathcal{M} tending to infinity in place of the ray $[(\bar{\partial}_E, t\Phi)]$ considered in [FMSW22]. The second is that the analysis of the linearized operator requires modifications because of the presence of irregular singularities. We use the analytic tools from [BB04] to handle these singularities.

This paper is organized as follows. In Section 2, we give some definitions relating to irregular Higgs bundles, and describe the Hitchin base and generic Hitchin fibers explicitly. In Section 3, we introduce the building blocks, the fiducial solutions and the limiting metric, and construct an approximate solution by gluing these together. Note that in our case the zeros of $\det \Phi_t$ are moving with t , so the gluing regions also depend on t . The decaying rates of the error terms in different regions depend on some local parameters, which are called *local masses* by analogy with those appearing in [Fos17]. We analyze the linearized Hitchin equation in Section 4, and nonlinear terms in Section 5, then a genuine solution to the Hitchin equation is obtained. Finally in Section 6, we prove Theorem 1.1, and analyze the semiflat metrics in detail for the four-dimensional moduli spaces.

2. PRELIMINARIES

Let E be a complex rank two vector bundle over $C = \mathbb{C}P^1$, S be a set of points on C . Cover C by the usual coordinate charts $U = \mathbb{C}_z$, $V = \mathbb{C}_w$ with $w = 1/z$, and such that $\{w = 0\} \notin S$. For $x \in S$, we also denote its z -coordinate by $x \in \mathbb{C}$. Furthermore, we fix the *parabolic structure* at each $x \in S$, which consists of a one-dimensional subspace $L_x \subset E_x$ (a full flag) and the associated parabolic weights $1/2 < \alpha_{x,2} < 1$, $\alpha_{x,1} = 1 - \alpha_{x,2}$.

Recall that an $\text{SL}(2, \mathbb{C})$ -irregular Higgs bundle [BB04] is a pair $(\bar{\partial}_E, \Phi)$, where $\bar{\partial}_E$ is a holomorphic structure on E , and the Higgs field $\Phi \in H^0(C, \mathfrak{sl}(E) \otimes K(D))$, for the divisor $D = \sum_{x \in S} m_x \cdot \{x\}$ with $m_x \in \mathbb{Z}^+$. Write $S = I \cup T$, where $m_x > 1$ for $x \in I$ and $m_x = 1$ for $x \in T$. Then Φ is a meromorphic section of $\Omega^{1,0}(\mathfrak{sl}(E))$ with singularities at points in S , which are called *irregular singularities* when they belong to I , otherwise called *tame singularities*. We require that Φ has a fixed *irregular type* at each $x \in I$, meaning that there exists a holomorphic trivialization of $(E, \bar{\partial}_E)$ over a neighborhood U_x of x such that

$$\Phi = \left(\frac{\phi_{x,m_x}}{z_x^{m_x}} + \cdots + \frac{\phi_{x,1}}{z_x} + \text{holomorphic terms} \right) dz_x,$$

where $z_x = z - x$, and $\phi_{x,m_x}, \dots, \phi_{x,1} \in \mathfrak{sl}(2, \mathbb{C})$ are given. Moreover, Φ is adapted to the parabolic structure, i.e., $\phi_{x,m_x}(L_x) \subset L_x$. We assume that Φ is *strongly parabolic* at each $x \in T$ (see Remark 3.6), meaning that $\phi_{x,1}$, the residue of Φ at x , is nilpotent and not fixed.

Definition 2.1. An irregular Higgs bundle $(\bar{\partial}_E, \Phi)$ is *stable* if for every rank one Φ -invariant holomorphic subbundle F of $(E, \bar{\partial}_E)$,

$$\text{pdeg } F < \frac{1}{2} \text{pdeg } E,$$

where $\text{pdeg } E = \deg E + |S|$, and $\text{pdeg } F = \deg F + \sum_{x \in S} \alpha_{x,F}$, $\alpha_{x,F} = \alpha_{x,2}$ if $F_x = L_x$, otherwise $\alpha_{x,F} = \alpha_{x,1}$.

Definition 2.2. The *moduli space* of irregular Higgs bundles is

$$\mathcal{M} = \{(\bar{\partial}_E, \Phi) \text{ stable and compatible with the parabolic structure and the irregular type at each } x \in S\} / \mathcal{G}_C,$$

where $\mathcal{G}_C = \Gamma(\text{ParEnd}(E) \cap SL(E))$, which consists of sections of $SL(E)$ preserving L_x for each $x \in S$. Alternatively, $\mathcal{M} = \{(\bar{\partial}_E, \Phi, L)\} / \Gamma(SL(E))$, where $L = \{L_x\}_{x \in S}$, $L_x \in \mathbb{C}\mathbb{P}_x^1$ which parametrizes one-dimensional subspaces of E_x . The gauge group acts as

$$g \cdot (\bar{\partial}_E, \Phi, L) = (g^{-1} \circ \bar{\partial}_E \circ g, g^{-1} \Phi g, g^{-1} \cdot L).$$

By [Ina16, Th. 3.1], $\dim_{\mathbb{C}} \mathcal{M} = 2(N - 3)$, where $N = \deg D$. We assume that $N \geq 4$.

We further decompose $I = I_u \cup I_t$. For $x \in I_u$, the leading term ϕ_{x, m_x} is diagonalizable with opposite nonzero eigenvalues $\pm \rho_{x, m_x}$ (x is called an *untwisted singularity*), while for $x \in I_t$, ϕ_{x, m_x} is nilpotent (x is called a *twisted singularity*). In the triple $(\bar{\partial}_E, \Phi, L)$, as Φ is compatible with L , we can determine L_x from Φ as the kernel of ϕ_{x, m_x} for $x \in I_t \cup T$, and as one of the eigenspace of ϕ_{x, m_x} for $x \in I_u$. There are $2^{|I_u|}$ choices, depending on whether L_x corresponds to ρ_{x, m_x} or $-\rho_{x, m_x}$, where $0 \leq \text{Arg}(\rho_{x, m_x}) < \pi$. These $2^{|I_u|}$ components of \mathcal{M} are equivalent, so we only need to consider the one where L_x corresponds to ρ_{x, m_x} , and still denote this component by \mathcal{M} . We also suppress $[(\bar{\partial}_E, \Phi, L_\Phi)] \in \mathcal{M}$ as $[(\bar{\partial}_E, \Phi)]$.

Definition 2.3. The *Hitchin fibration (map)* is defined as

$$H : \mathcal{M} \rightarrow \mathcal{B}, [(\bar{\partial}_E, \Phi)] \mapsto \det \Phi.$$

Here $\mathcal{B} \subset H^0(C, K(D)^2)$ consists of quadratic differentials ν which has the expansion

$$\nu = \left(\frac{\mu_{x, 2m_x}}{z_x^{2m_x}} + \cdots + \frac{\mu_{x, m_x+1}}{z_x^{m_x+1}} + \text{higher order terms} \right) dz_x^2,$$

near each $x \in I$. $\mu_{x, j}$'s are complex numbers fixed by the irregular type at x , and clearly $\mu_{x, 2m_x} \neq 0$ for $x \in I_u$, while $\mu_{x, 2m_x} = 0$ for $x \in I_t$. Moreover, we assume that $\mu_{x, 2m_x-1} \neq 0$ for $x \in I_t$ (when it is zero, by choosing a different extension of the holomorphic structure over x , this case can be reduced to the nonzero case or the case where the leading coefficient is diagonalizable [Wit08, p. 53]).

Lemma 2.4. *The Hitchin base \mathcal{B} is a dimension $N - 3$ affine subspace of $H^0(C, K(D)^2)$, which can be explicitly written as*

$$\mathcal{B} = \left\{ \left(\sum_{x \in I_{\geq 3}} \sum_{a=m_x+1}^{2m_x} \frac{\mu_{x, a}}{(z-x)^a} + \sum_{x \in I_{=2}} \left(\frac{\mu_{x, 4}}{(z-x)^4} + \frac{\mu_{x, 3}(x-y_x)}{(z-x)^3(z-y_x)} \right) + \frac{\sum_{b=0}^{N-4} \nu_b z^b}{\prod_{x \in S} (z-x)^{m_x}} \right) dz^2 \mid \nu_0, \dots, \nu_{N-4} \in \mathbb{C} \right\}, \quad (1)$$

where $I_{\geq 3}$ consists of poles with order ≥ 3 , and $I_{=2}$ consists of order two poles. If $I_{=2} \neq \emptyset$, then we can choose $y_x \in S \setminus \{x\} \neq \emptyset$ since $N \geq 4$. Choosing a different y_x for $x \in I_{=2}$ amounts to shifting $(\nu_0, \dots, \nu_{N-4})$ by an element of \mathbb{C}^{N-3} .

Proof. By considering the difference of two elements in \mathcal{B} , one can see that \mathcal{B} is an affine space modeled on $H^0(C, K^2 \otimes \mathcal{O}(D))$, which has dimension $N - 3$. Let the right hand side of

(1) be \mathcal{B}_1 , then $\mathcal{B}_1 \subset \mathcal{B}$. Since $\prod_{x \in S} (z-x)^{-m_x} z^i$ ($i = 0, \dots, N-4$) are linearly independent, we have $\dim \mathcal{B}_1 = N-3$ and $\mathcal{B}_1 = \mathcal{B}$. For $y_{x,1}, y_{x,2} \in S \setminus \{x\}$,

$$\frac{\mu_{x,3}(x-y_{x,1})}{(z-x)^3(z-y_{x,1})} - \frac{\mu_{x,3}(x-y_{x,2})}{(z-x)^3(z-y_{x,2})} = \frac{\mu_{x,3}(y_{x,2}-y_{x,1})}{(z-x)^2(z-y_{x,1})(z-y_{x,2})},$$

the last statement follows. \square

Remark 2.5. Let $\widetilde{\mathcal{M}}$ be a moduli space of $\mathrm{GL}(2, \mathbb{C})$ -irregular Higgs bundles, defined in the same way as for \mathcal{M} except that Φ belongs to $H^0(C, \mathfrak{gl}(E) \otimes K(D))$. Then for two different Higgs fields Φ_1, Φ_2 in $\widetilde{\mathcal{M}}$, since the irregular types are fixed, we have $\mathrm{tr}(\Phi_1) - \mathrm{tr}(\Phi_2) \in H^0(C, K)$ which is trivial. This means that Higgs fields in $\widetilde{\mathcal{M}}$ have constant traces, and there is no loss in considering $\mathrm{SL}(2, \mathbb{C})$ -irregular Higgs bundles over C .

Let $\pi : K(D) \rightarrow C$ be the projection, and λ be the tautological section of $\pi^*K(D)$. Each point ν in \mathcal{B} defines a section of $(\pi^*K(D))^2$ of the form $\lambda^2 + \pi^*\nu$ whose zero locus S_ν is called the *spectral curve* determined by ν . From now on we assume that S_ν is smooth, which is true for generic ν by Bertini's theorem. $\pi : S_\nu \rightarrow C$ is a two-fold ramified cover with ramification divisor R_ν . By the Riemann-Hurwitz formula, the genus of S_ν is $N-3$. For Φ with $\det \Phi = \nu$, one can associate the *spectral line bundle* \mathcal{L}_Φ over S_ν such that $\mathcal{L}_\Phi(-R_\nu) \subset \pi^*E$ and $\pi_*\mathcal{L}_\Phi = E$. Away from the support of R_ν , the fiber $\mathcal{L}_{\Phi,q}$ is the eigenspace in $E_{\pi(q)}$ of Φ with the eigenvalue q . By the Grothendieck–Riemann–Roch theorem [LM10, p. 96],

$$\deg \mathcal{L}_\Phi = \deg E + N - 2. \quad (2)$$

Assume for simplicity that $\mathrm{pdeg} E = 0$ (if $\mathrm{pdeg} E = 2d$ for some $d \in \mathbb{Z}$, one can replace E by $E \otimes \mathcal{O}(-d)$ [FN21, p. 624]). Then $\deg E = -|S|$ and $\mathcal{E} = (E, \bar{\partial}_E) \cong \mathcal{O}(m) \oplus \mathcal{O}(-|S|-m)$, $m \geq -|S|/2$, by the Birkhoff–Grothendieck theorem. Now

$$\mathrm{End} \mathcal{E} \cong \begin{pmatrix} \mathcal{O} & \mathcal{O}(2m+|S|) \\ \mathcal{O}(-|S|-2m) & \mathcal{O} \end{pmatrix}.$$

In a trivialization of $\mathrm{End} \mathcal{E}$ over U , we have

$$\Phi = \frac{dz}{\prod_{x \in S} (z-x)^{m_x}} \begin{pmatrix} a(z) & b(z) \\ c(z) & -a(z) \end{pmatrix}, \quad \text{where } a(z), b(z), c(z) \in \mathbb{C}[z]. \quad (3)$$

Lemma 2.6. *Every Φ in (3) with $\det \Phi \in \mathcal{B}$ of (1) is compatible with some fixed irregular data at points in I , i.e.,*

(i) *there exists a local holomorphic frame around $x \in I_u$ in which*

$$\Phi = \sum_{j=1}^{m_x} \rho_{x,j} z_x^{-j} \sigma_3 dz_x + \text{holomorphic terms}, \quad (4)$$

where σ_3 is the Pauli matrix $\mathrm{diag}(1, -1)$ and $\rho_{x,j}$'s are determined by $\mu_{x,a}$'s in (1);

(ii) *there exists a local holomorphic frame around $x \in I_t$ in which*

$$\Phi = \begin{pmatrix} 0 & -\sum_{j=1}^{m_x-1} \mu_{x,m_x+j} z_x^{-j} \\ z_x^{-m_x} & 0 \end{pmatrix} dz_x + \text{holomorphic terms}. \quad (5)$$

Proof. (i) This is essentially [BB04, Lem. 1.1]. By (1) and (3), in a neighborhood of x , we have the expansion

$$\det \Phi(z_x) = \left(\sum_{j=m_x+1}^{2m_x} \mu_{x,j} z_x^{-j} + \text{higher order terms} \right) dz_x^2, \quad (6)$$

$$\Phi(z_x) = \left(\sum_{j=1}^{m_x} \phi_{x,j} z_x^{-j} + \text{holomorphic terms} \right) dz_x, \quad (7)$$

$\mu_{x,2m_x} = \det \phi_{x,m_x} \neq 0$, so we can find a frame where $\phi_{x,m_x} = \rho_{x,m_x} \sigma_3$ with $\rho_{x,m_x} = \mu_{x,2m_x}^{1/2}$. Then we use gauge transformations of the form $g = 1 - g_j z_x^j$ ($j = 1, \dots, m_x - 1$) to successively cancel the off-diagonal terms of ϕ_{x,m_x-j} . In each stage, if necessary, we shrink the neighborhood of 0 so that g is invertible. For example, for $g = 1 - g_1 z$,

$$g^{-1} \Phi g = (\phi_{x,m_x} z^{-m_x} + (\phi_{x,m_x-1} - [\phi_{x,m_x}, g_1]) z^{-m_x+1} + O(z^{-m_x+2})) dz_x,$$

and $\phi_{x,m_x-1} - [\phi_{x,m_x}, g_1]$ becomes diagonal if we choose

$$g_1 = \begin{pmatrix} 0 & \phi_{x,m_x-1,12}/2 \\ -\phi_{x,m_x-1,21}/2 & 0 \end{pmatrix}, \text{ where } \phi_{x,m_x-1,ij} \text{ is the } (i,j) \text{ - entry of } \phi_{x,m_x-1}.$$

Therefore, in some local holomorphic frame around x , Φ has the form in (4). Now $\rho_{x,m_x} = \mu_{x,2m_x}^{1/2} \neq 0$, and $\rho_{x,j}$ ($j = m_x - 1, \dots, 1$) can be recursively determined as

$$\rho_{x,j} = (-2\rho_{x,m_x})^{-1} \left(\mu_{x,j+m_x} + \sum_{k=1}^{m_x-j-1} \rho_{x,j+k} \rho_{x,m_x-k} \right).$$

(ii) The proof is similar to (i). Around x , we can expand $\det \Phi$ and Φ as in (6), (7). Since $\mu_{x,2m_x} = 0$ and $\mu_{x,2m_x-1} \neq 0$, then $\phi_{x,m_x} \neq 0$ and is nilpotent, so we can find a local frame in which $\phi_{x,m_x} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Then we use gauge transformations of the form $g = 1 - g_j z_x^j$ ($j = 1, \dots, m_x - 1$) to successively cancel the lower-triangular part of ϕ_{x,m_x-j} . For example if we choose

$$g_1 = \begin{pmatrix} \phi_{x,m_x-1,21} & \phi_{x,m_x-1,22} \\ 0 & 0 \end{pmatrix}$$

in $g = 1 - g_1 z$, then $\phi_{x,m_x-1} - [\phi_{x,m_x}, g_1]$ becomes strictly upper-triangular. Therefore, in some local holomorphic frame around x , Φ has the form in (5). \square

In V , we can write Φ as

$$\Phi = \frac{-w^{N-2} dw}{\prod_{x \in S} (1 - xw)^{m_x}} \begin{pmatrix} a(w^{-1}) & w^{2m+|S|} b(w^{-1}) \\ w^{-2m-|S|} c(w^{-1}) & -a(w^{-1}) \end{pmatrix}.$$

Φ is regular at ∞ , so $\deg a \leq N - 2$, $\deg b \leq 2m + |S| + N - 2$, $\deg c \leq -2m - |S| + N - 2$. Since $\deg c \geq 0$, and recall that $m \geq -|S|/2$, we have

$$-|S| \leq 2m \leq -|S| + N - 2,$$

which yields a stratification of \mathcal{M} :

$$\mathcal{M} = \bigsqcup_{m=\lceil -|S|/2 \rceil}^{\lfloor (-|S|+N-2)/2 \rfloor} \mathcal{M}_m, \quad (8)$$

where $[(\bar{\partial}_E, \Phi)] \in \mathcal{M}_m$ if $(E, \bar{\partial}_E) \cong \mathcal{O}(m) \oplus \mathcal{O}(-|S| - m)$. Let H_m be the restriction of H to \mathcal{M}_m . Note that

$$-a(z)^2 - b(z)c(z) = \prod_{x \in S} (z - x)^{2m_x} \nu(z) := \tilde{\nu}(z) \text{ where } \det \Phi = \nu = \nu(z) dz^2, \quad (9)$$

then $b(z) = (-a(z)^2 - \tilde{\nu}(z))/c(z)$. The spectral cover $S_\nu \rightarrow C$ ramifies exactly at the $2N - 4$ zeros (counted with multiplicity) of $\tilde{\nu}(z)$. We say that $\nu \in \mathcal{B}'$, the *regular locus*, if $\tilde{\nu}(z)$ only has simple zeros. When $\nu = \det \Phi \in \mathcal{B}'$, the Higgs bundle is stable [MSWW16, p. 7] and S_ν is smooth [FMSW22, p. 10]. Fix $\nu \in \mathcal{B}'$, and next we study the fiber $H_m^{-1}(\nu)$. Parabolic Higgs bundles have been studied in [FMSW22], so from now on we suppose that $I \neq \emptyset$. If $I_u \neq \emptyset$, by a coordinate change when necessary, we may assume that $0 \in I_u$, or else we assume that $0 \in I_t$.

If $m = -|S|/2$, then $|S|$ is even, the holomorphic gauge transformations are of the form

$$g = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}, \quad g_{11}, g_{12}, g_{21}, g_{22} \in \mathbb{C}, \det g = 1.$$

Let $a_i, b_i, c_i, \tilde{\nu}_i$ be the coefficients of z^i in $a(z), b(z), c(z), \tilde{\nu}(z)$, and make the generic assumption that $\tilde{\nu}_{2N-4}, \tilde{\nu}_{2N-5} \neq 0$. By some gauge transformation g above, we can make $b_0 = c_0 = 0$ when $0 \in I_u$, or $a_0 = c_0 = 0$ when $0 \in I_t$. Now $c(z) \neq 0$, otherwise $\tilde{\nu}(z) = -a(z)^2$, contradicting that $\tilde{\nu}(z)$ only has simple zeros. If $\deg c = N - 2$, using the diagonal gauge transformation $g = \text{diag}(c_{N-2}^{-1/2}, c_{N-2}^{1/2})$ we can make $c(z)$ monic. Then we apply

$$g = \begin{pmatrix} 1 & a_{N-2} \\ 0 & 1 \end{pmatrix}$$

to make $a_{N-2} = 0$. There is no gauge freedom left. The remaining $N - 2$ (if $0 \in I_u$, or $N - 3$ if $0 \in I_t$) coefficients of $a(z)$ can be determined (up to finitely many choices) by c_1, c_2, \dots, c_{N-3} and the relation (9). For example, if $c(z)$ has distinct roots $x_1 = 0, \dots, x_{N-2}$, then a_0, a_1, \dots, a_{N-3} satisfy $N - 2$ linear equations

$$a(x_i) = y_i, \quad i = 1, \dots, N - 2, \text{ where } y_i = \pm(-\tilde{\nu}(x_i))^{1/2}.$$

If $\deg c < N - 2$, then $a_{N-2}^2 = -\tilde{\nu}_{2N-4}$ and $\deg c = N - 3$. In fact, if $a_{N-2}^2 \neq -\tilde{\nu}_{2N-4}$, then $\deg(-a(z)^2 - \tilde{\nu}(z)) = 2N - 4 > 2N - 5 \geq \deg(b(z)c(z))$, a contradiction. Then we have $\deg(-a(z)^2 - \tilde{\nu}(z)) = 2N - 5$, and $\deg c = 2N - 5 - \deg b \geq N - 3$, and $\deg c = N - 3$. Applying gauge transformations as above we can make $c(z)$ monic and $a_{N-3} = 0$. As before, the remaining coefficients of $a(z)$ are determined by c_1, \dots, c_{N-4} . Therefore

$$\dim(\mathcal{M}_{-|S|/2}) = N - 3.$$

If $m > -|S|/2$, the holomorphic gauge transformations are of the form

$$g = \begin{pmatrix} g_{11} & g_{12}(z) \\ 0 & g_{11}^{-1} \end{pmatrix}, \quad g_{11} \in \mathbb{C}, g_{12}(z) \in \mathbb{C}[z], \deg g_{12}(z) \leq 2m + |S|.$$

If $\deg c = -2m - |S| + N - 2$, we may assume that $c(z)$ is monic after applying a diagonal gauge transformation as above. Then we successively apply

$$g = \begin{pmatrix} 1 & a_{N-2-i} z^{2m+|S|-i} \\ 0 & 1 \end{pmatrix}, \quad i = 0, 1, \dots, 2m + |S|,$$

to make $a_{N-2} = a_{N-3} = \cdots = a_{N-2-2m-|S|} = 0$. There is no gauge freedom left. The remaining $N - 2 - 2m - |S|$ coefficients of $a(z)$ are determined by $c(z)$ and (9). If $\deg c < -2m - |S| + N - 2$, then $a_{N-2}^2 = -\tilde{\nu}_{2N-4}$ and $\deg c = -2m - |S| + N - 3$. Again we can find a representative in each gauge orbit with $c(z)$ monic and $a_{N-3} = \cdots = a_{N-3-2m-|S|} = 0$. The remaining coefficients of $a(z)$ are determined by $c(z)$. Therefore

$$\dim(\mathcal{M}_m) = -2m - |S| + N - 2.$$

3. APPROXIMATE SOLUTIONS

We consider the case $I_u \neq \emptyset$, then $0 \in I_u$ by assumption. Fix $\nu \in \mathcal{B}'$ satisfying the generic conditions $\tilde{\nu}_{2N-4}, \tilde{\nu}_{2N-5} \neq 0$, and $\nu_{N-4} \neq 0$. By (1), we can represent ν by $(\nu_0, \dots, \nu_{N-4}) \in \mathbb{C}^{N-3}$. Consider a curve $\nu_t = (f_0(t)\nu_0, \dots, f_{N-4}(t)\nu_{N-4})$ in \mathcal{B} with $\nu_{t=1} = \nu$ and its norm tending to ∞ , where $f_i(t)$ is positive and nondecreasing in t . For simplicity, let $f_{N-4}(t) = t$ and $f_i(t) = o(t^{(m_0+i)/(N-4+m_0)})$ for $i = 0, \dots, N-5$, so that the following holds.

Lemma 3.1. *For t sufficiently large, $\tilde{\nu}_t(z)$ has $2N - 4$ distinct roots, with leading terms given by*

$$z_{0,j}(t) \sim \left(-\frac{\mu_{0,2m_0}}{\nu_{N-4}} \prod_{y \in S \setminus \{0\}} (-y)^{m_y} \right)^{\frac{1}{m_0+N-4}} e^{\frac{2j\pi i}{m_0+N-4}} t^{-\frac{1}{m_0+N-4}}, \quad j = 0, \dots, m_0 + N - 5;$$

$$z_{x,j}(t) \sim x + \left(-\frac{\mu_{x,2m_x}}{\nu_{N-4}} x^{4-N} \prod_{y \in S \setminus \{x\}} (x-y)^{m_y} \right)^{\frac{1}{m_x}} e^{\frac{2j\pi i}{m_x}} t^{-\frac{1}{m_x}}, \quad j = 0, \dots, m_x - 1,$$

for $x \in I_u \setminus \{0\}$;

$$z_{x,j}(t) \sim x + \left(-\frac{\mu_{x,2m_x-1}}{\nu_{N-4}} x^{4-N} \prod_{y \in S \setminus \{x\}} (x-y)^{m_y} \right)^{\frac{1}{m_x-1}} e^{\frac{2j\pi i}{m_x-1}} t^{-\frac{1}{m_x-1}}, \quad j = 0, \dots, m_x - 2,$$

for $x \in I_t$; and $z_{x,m_x-1}(t) = x$, for $x \in I_t \cup T$.

Here the notation $A \sim B$ means $A = B + o(B)$. Therefore $\nu_t \in \mathcal{B}'$ for t large enough.

Proof. By (1), we can write

$$\tilde{\nu}_t(z) = \left(\sum_{b=0}^{N-4} f_b(t)\nu_b z^b \right) z^{m_0} \prod_{y \in S \setminus \{0\}} (z-y)^{m_y} + \mu_{0,2m_0} \prod_{y \in S \setminus \{0\}} (z-y)^{2m_y} + z g_0(z),$$

where $g_0(z)$ is some polynomial independent of t . By Rouché's theorem, there is a constant κ such that in the disk $B_{\kappa t^{-1/(m_0+N-4)}}(0)$, $\tilde{\nu}_t(z)$ has the same number of zeros as that for

$$\tilde{\nu}_t^{(0)}(z) := \nu_{N-4} t z^{m_0+N-4} \prod_{y \in S \setminus \{0\}} (-y)^{m_y} + \mu_{0,2m_0} \prod_{y \in S \setminus \{0\}} (-y)^{2m_y},$$

for t large enough. Choose κ , so that $\tilde{\nu}_t^{(0)}$ has $m_0 + N - 4$ distinct roots in the disk. Using the scaling $z \mapsto t^{1/(m_0+N-4)} z$, we see that the roots $z_{0,j}(t)$ ($j = 0, \dots, m_0 + N - 5$) of $\tilde{\nu}_t(z)$ in the disk are asymptotic to those of $\tilde{\nu}_t^{(0)}(z)$, which are given in the statement of the lemma. The analysis near $x \in I \setminus \{0\}$ is similar, using the coordinate z_x in place of z . \square

The hyperkähler metric on \mathcal{M} is defined through the nonabelian Hodge correspondence, which realizes \mathcal{M} as a hyperkähler quotient, i.e., unitary gauge equivalence classes of solutions to Hitchin's equations. Thus, we need to find the solution of the following *Hitchin's equation* for $[(\bar{\partial}_E, \Phi)] \in \mathcal{M}$:

$$F_h + [\Phi, \Phi^{*h}] = 0,$$

where F_h is the curvature of the Chern connection $D(\bar{\partial}_E, h)$. h is called a *harmonic metric* if it satisfies the above equation and adapted to the parabolic structure, which means for each x the filtration of the fiber E_x given by $E_{x,\beta} := \{s(0) : s \text{ holomorphic, } |s(z)|_h = O(|z|^\beta)\}$ coincides with that of the parabolic structure. When $I_t = \emptyset$, such a metric h always exists on a stable Higgs bundle by [BB04]. However, if $I_t \neq \emptyset$, the existence of h will impose further constraints on the parabolic weights $\alpha_{x,1}, \alpha_{x,2}$ for $x \in I_t$.

Lemma 3.2. *Let (E, Φ) be an $\text{SL}(2, \mathbb{C})$ -Higgs bundle over a Riemann surface C , where Φ has an order $n \geq 2$ pole at x , with nilpotent leading coefficient ϕ_{-n} . The kernel L of ϕ_{-n} determines a filtration $0 \subset L \subset E_x$ and let the associated parabolic weights be $1 > \alpha_2 > \alpha_1 > 0$, $\alpha_1 + \alpha_2 = 1$. Suppose $\text{pdeg } E = 0$. If h is a harmonic metric adapted to the parabolic structure, then $\alpha_1 = 1/4, \alpha_2 = 3/4$.*

Proof. As in [MSWW16, Lem. 3.6], we first find a standard form of Φ near x . Since ϕ_{-n} is nilpotent, $\det \Phi$ has an order $2n - 1$ pole at p_0 . By [Str84, p. 28], one can find a local coordinate z centered at p_0 , such that $\det \Phi = -z^{1-2n} dz^2$. Again by the nilpotency of the leading term, in some local holomorphic frame

$$z^n \phi(z) = \begin{pmatrix} a(z) & b(z) \\ c(z) & -a(z) \end{pmatrix}, \quad z^n \phi(z)|_{z=0} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \text{where } \Phi(z) = \phi(z) dz.$$

Now $\sqrt{c(z)}$ is well-defined near 0 since $c(0) = 1$, and

$$g^{-1} \Phi g = \begin{pmatrix} 0 & \frac{1}{z^{n-1}} \\ \frac{1}{z^n} & 0 \end{pmatrix} dz, \quad \text{where } g = \frac{1}{\sqrt{c(z)}} \begin{pmatrix} 1 & a(z) \\ 0 & c(z) \end{pmatrix}.$$

In the local holomorphic frame determined by g , we have

$$h = \begin{pmatrix} \frac{h_{11}(z)}{h_{12}(z)} & \frac{h_{12}(z)}{h_{22}(z)} \end{pmatrix}, \quad h_{11}(z) \sim c_1 r^{2\alpha_1}, h_{22}(z) \sim c_2 r^{2\alpha_2}, \det h = Q(z) r^2, h_{12}(z) = O(r),$$

where c_1, c_2 are positive constants, $r = |z|$, $Q(z)$ is determined by $h_{\det E}$, the metric on $\det E$ induced by h , $Q(0) > 0$. The $(1, 1)$ -entry of F_h is $O(r^{-2})$ while that of $[\Phi, \Phi^{*h}]$ is $O(r^{4\alpha_1-2n}) + O(r^{4\alpha_2-2-2n})$ (the leading terms in these $O(\cdot)$'s are nonzero), which grows faster than r^{-2} near 0 if the sum of the leading terms is nonzero. Therefore $F_h + [\Phi, \Phi^{*h}] = 0$ implies that $4\alpha_1 - 2n = 4\alpha_2 - 2 - 2n$, and $\alpha_1 = 1/4, \alpha_2 = 3/4$. \square

Henceforth, we assume that $\alpha_{x,1} = 1/4, \alpha_{x,2} = 3/4$, for $x \in I_t$. To get a more explicit description of the hyperkähler metric near the ends of \mathcal{M} , we need more information on h , which can be obtained via gluing constructions as in [FMSW22], and we also follow this strategy. Fix $[(\bar{\partial}_E, \Phi)] \in H^{-1}(\nu)$, and choose $[(\bar{\partial}_E, \Phi_t)] \in H^{-1}(\nu_t)$, such that $\Phi_t(z)$ and $\Phi(z)$ have the same $c(z)$ when written as in (3) (by the previous section, there are only finitely many gauge inequivalent choices of Φ_t). Note that if $[(\bar{\partial}_E, \Phi)]$ belongs to \mathcal{M}_m of (8), then the whole curve $[(\bar{\partial}_E, \Phi_t)]$ remains in \mathcal{M}_m as t varies.

Now we aim to find a solution h_t of $F_{h_t} + [\Phi_t, \Phi_t^{*h_t}] = 0$ for large t . First we construct an approximate solution h_t^{app} , and later we will deform it into a genuine solution. The construction of h_t^{app} is similar to that in [FMSW22, Sec. 3]. Since we are considering $\text{SL}(2, \mathbb{C})$ Higgs bundles, the data on $\det E \simeq \mathcal{O}(-|S|)$ is fixed. In other words, we fix the holomorphic structure induced from $\bar{\partial}_E$ and the harmonic metric $h_{\det E} = \prod_{x \in S} |z - x|^2$ adapted to the induced parabolic weight $\alpha_{x,1} + \alpha_{x,2} = 1$ at x (in V , $h_{\det E} = \prod_{x \in S} |1 - xw|^2$). Let (S_t, \mathcal{L}_t) be the spectral data associated with $(\bar{\partial}_E, \Phi_t)$. The spectral cover $\pi : S_t \rightarrow C$ ramifies at Z_t , the zero locus of $\tilde{\nu}_t(z)$. By (2), $\deg \mathcal{L}_t = -|S| + N - 2$. Now \mathcal{L}_t has rank one, a parabolic structure at a point is equivalent to a parabolic weight since there is only a trivial filtration. For each of the two points in $\pi^{-1}(x)$, $x \in I_u$, let the weight be $\alpha_{x,1}$ if it corresponds to the eigenvalue ρ_{x,m_x} of ϕ_{x,m_x} and otherwise be $\alpha_{x,2}$. For each point in $\pi^{-1}(Z_t \setminus (I_t \cup T))$, let the weight be $-1/2$. For each point $\pi^{-1}(x)$, $x \in I_t \cup T$, let the weight be $\alpha_{x,1} + \alpha_{x,2} - 1/2 = 1/2$. Endowed with this parabolic structure we have

$$\text{pdeg } \mathcal{L}_t = -|S| + N - 2 + |I_u| + |I_t|/2 + |T|/2 - (2N - 4 - |I_t| - |T|)/2 = 0 = \text{pdeg } E.$$

By [Sim90], there is a unique (up to scaling) Hermitian-Einstein metric $h_{\mathcal{L}_t}$ on \mathcal{L}_t adapted to the parabolic structure, which is flat in our setting. Finally, let $h_{t,\infty}$ be the metric on $E|_{C \setminus (Z_t \cup S)}$ defined as the orthogonal pushforward of $h_{\mathcal{L}_t}$. Then $h_{t,\infty}$ solves the decoupled Hitchin's equations:

$$F_{h_{t,\infty}} = 0, \quad [\Phi_t, \Phi_t^{*h_{t,\infty}}] = 0.$$

The metric on $\det E|_{C \setminus (Z_t \cup S)}$ induced by $h_{t,\infty}$ can be extended to $C \setminus \{S\}$ and is a harmonic metric. By uniqueness, we may assume that this metric is $h_{\det E}$ defined above, by rescaling $h_{\mathcal{L}_t}$ if necessary. Near $x \in Z_t \cup S$, Φ_t and $h_{t,\infty}$ can be explicitly described using the following normal forms.

Proposition 3.3. *There exist constants $1 > \kappa, \kappa_1, \kappa_2 > 0$, such that for t large enough, the followings hold.*

- (i) *There is a holomorphic coordinate $\zeta_{x,j,t}$ centered at $z_{x,j}(t)$ for $x \in I$, $j = 0, \dots, j(x) - 1$ where*

$$j(x) = \begin{cases} m_0 + N - 4, & \text{if } x = 0, \\ m_x, & \text{if } x \in I_u \setminus \{0\}, \\ m_x - 1, & \text{if } x \in I_t, \end{cases}$$

and a local holomorphic frame of $(E, \bar{\partial}_E)$ over $\tilde{B}_{x,j,t} := \{|\zeta_{x,j,t}| < \kappa t^{-1/j(x)}\}$ in which

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \lambda_{x,j}(t) \begin{pmatrix} 0 & 1 \\ \zeta_{x,j,t} & 0 \end{pmatrix} d\zeta_{x,j,t}, \quad h_{t,\infty} = \begin{pmatrix} |\zeta_{x,j,t}|^{1/2} & 0 \\ 0 & |\zeta_{x,j,t}|^{-1/2} \end{pmatrix}. \quad (10)$$

Here the local mass $\lambda_{x,j}(t) = |\nu_{x,j,t}(z_{x,j}(t))|^{1/2}$, where $\nu_{x,j,t}(z) = (z - z_{x,j}(t))^{-1} \nu_t(z)$.

- (ii) *There is a holomorphic coordinate $\zeta_{x,t}$ centered at x for $x \in T$, and a local holomorphic frame of $(E, \bar{\partial}_E)$ over $\tilde{B}_{x,t} := \{|\zeta_{x,t}| < \kappa\}$ in which*

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \lambda_x(t) \begin{pmatrix} 0 & 1 \\ \frac{1}{\zeta_{x,t}} & 0 \end{pmatrix} d\zeta_{x,t}, \quad h_{t,\infty} = \begin{pmatrix} |\zeta_{x,t}|^{1/2} & 0 \\ 0 & |\zeta_{x,t}|^{3/2} \end{pmatrix}. \quad (11)$$

Here $\lambda_x(t) = |\nu_{x,t}(x)|^{1/2}$, and $\nu_{x,t}(z) = (z - x)\nu_t(z)$.

(iii) There is a holomorphic coordinate $\zeta_{x,t}$ centered at x for $x \in I_t$, and a local holomorphic frame of $(E, \bar{\partial}_E)$ over $\tilde{B}_{x,t} := \{|\zeta_{x,t}| < \kappa t^{-1/(m_x-1)}\}$ in which

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \lambda_x(t) \begin{pmatrix} 0 & \frac{1}{\zeta_{x,t}^{m_x-1}} \\ \frac{1}{\zeta_{x,t}^{m_x}} & 0 \end{pmatrix} d\zeta_{x,t}, \quad h_{t,\infty} = \begin{pmatrix} |\zeta_{x,t}|^{1/2} & 0 \\ 0 & |\zeta_{x,t}|^{3/2} \end{pmatrix}. \quad (12)$$

Here $\lambda_x(t) = |\nu_{x,t}(x)|^{1/2}$, and $\nu_{x,t}(z) = (z-x)^{2m_x-1} \nu_t(z)$.

(iv) For $x \in I_u$, there is a local holomorphic frame of $(E, \bar{\partial}_E)$ over $\tilde{B}_{x,t} := \{|z-x| < \kappa t^{-1/j(x)}\}$ in which

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \frac{1}{z_x^{m_x}} (z_x^{2m_x} \nu_t(z))^{1/2} \sigma_3 dz, \quad h_{t,\infty} = \begin{pmatrix} |z_x|^{2\alpha_{x,1}} & 0 \\ 0 & |z_x|^{2\alpha_{x,2}} \end{pmatrix} \quad (13)$$

The disks $\tilde{B}_{x(j),t}$ are disjoint, and $B_{x(j),t,\kappa_1} \subset \tilde{B}_{x(j),t} \subset B_{x(j),t,\kappa_2}$, where B_\bullet are disks defined using the coordinate z , for example, $B_{x,j,t,\kappa_1} = \{|z - z_{x,j}(t)| < \kappa_1 t^{-1/j(x)}\}$, for $x \in I, j = 0, \dots, j(x) - 1$.

Proof. By [Str84, Th. 6.1], any quadratic differential having a critical point of odd order can be written in a standard form using some local holomorphic coordinate around that point. This can be applied to (i)-(iii), where the quadratic differential $\det \Phi_t(z)$ has a critical point of order 1, -1 , $-(2m_x - 1)$ respectively.

(i) By a straightforward adaptation of the proof in [Str84], one can find a biholomorphic map $\sigma_{x,j,t}(z - z_{x,j}(t)) = \zeta_{x,j,t}$ from some disk B_{x,j,t,κ_2} to its image, such that $\det \Phi_t = -\lambda_{x,j}(t)^2 \zeta_{x,j,t} d\zeta_{x,j,t}^2$. $\sigma_{x,j,t}(0) = 0$, and moreover for t large enough, $|\sigma'_{x,j,t}|$ is bounded above and below by some positive constants. Therefore one can find constants κ, κ_1 so that $\tilde{B}_{x,j,t}$ lies inside the image $\sigma_{x,j,t}(B_{x,j,t,\kappa_2})$ and its preimage contains a smaller disk B_{x,j,t,κ_1} , for all sufficiently large t . As in [MSWW16, Lem. 3.6], we will write Φ_t in a standard form using local holomorphic gauge transformations, well defined over $\tilde{B}_{x,j,t}$. Recall that $c(z) \neq 0$ in Φ_t is fixed, for t large, $c(z)$ is nonzero on $\tilde{B}_{x,j,t}$. So we can write Φ_t in the coordinate $\zeta_{x,j,t}$ as $\Phi_t = \phi_{x,j,t}(\zeta_{x,j,t}) d\zeta_{x,j,t}$, with $\phi_{x,j,t,21}(\zeta_{x,j,t})$ nonvanishing on $\tilde{B}_{x,j,t}$, $\phi_{x,j,t,ij}(\zeta_{x,j,t})$ being the (i,j) -entry of $\phi_{x,j,t}(\zeta_{x,j,t})$. By a constant gauge transformation

$$\begin{pmatrix} \phi_{x,j,t,11}(0) & 1 \\ \phi_{x,j,t,21}(0) & 0 \end{pmatrix},$$

we can make $\Phi_t = \phi_{x,j,t}^{(1)}(\zeta_{x,j,t}) d\zeta_{x,j,t}$, with

$$\phi_{x,j,t}^{(1)}(0) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \text{and } \phi_{x,j,t,12}^{(1)} = \phi_{x,j,t,21}(\zeta_{x,j,t}) / \phi_{x,j,t,21}(0),$$

Now for κ suitably small, $|\phi_{x,j,t,12}^{(1)} - 1| < 1/2$ and $\sqrt{\phi_{x,j,t,12}^{(1)}}$ is well defined on $\tilde{B}_{x,j,t}$. Take the gauge transformation

$$\frac{1}{\sqrt{\phi_{x,j,t,12}^{(1)}}} \begin{pmatrix} \phi_{x,j,t,12}^{(1)} & 0 \\ -\phi_{x,j,t,11}^{(1)} & 1 \end{pmatrix}, \quad \text{such that } \Phi_t = \begin{pmatrix} 0 & 1 \\ \lambda_{x,j}(t)^2 \zeta_{x,j,t} & 0 \end{pmatrix} d\zeta_{x,j,t}.$$

Use the constant gauge transformation

$$\begin{pmatrix} 1 & 0 \\ 0 & \lambda_{x,j}(t) \end{pmatrix}, \text{ and then } \Phi_t = \lambda_{x,j}(t) \begin{pmatrix} 0 & 1 \\ \zeta_{x,j,t} & 0 \end{pmatrix} d\zeta_{x,j,t}.$$

Finally, as in [Fre18, Prop. 3.5], $h_{t,\infty}$ has the form given above, after applying a holomorphic gauge transformation on $\tilde{B}_{x,j,t}$ preserving Φ_t .

(ii) is similar to [FMSW22, Prop. 3.5]. (iii) is similar to Lemma 3.2 and (ii). (iv) is similar to [FMSW22, Prop. 3.7], noting that $(z_x^{2m_x} \nu_t(z))^{1/2}$ is well defined on $\tilde{B}_{x,t}$ for κ suitably small. We can also choose κ so that all the disks above are disjoint. \square

Near the points in (iii) and (iv) above, $h_{t,\infty}$ is already adapted to the parabolic structure. Near the points in (i), $h_{t,\infty}$ is singular, while near the points in (ii), $h_{t,\infty}$ is not adapted to the parabolic weights. So we need to desingularize $h_{t,\infty}$ using the following *fiducial solutions* h_t^{model} [FMSW22, Prop.s 3.8 & 3.9], and obtain the approximate metric h_t^{app} .

Definition 3.4. (i) For $x \in I$, in the above holomorphic frame over $\tilde{B}_{x,j,t}$, define

$$h_t^{\text{model}} = \begin{pmatrix} |\zeta_{x,j,t}|^{1/2} e^{l_{x,j,t}(|\zeta_{x,j,t}|)} & 0 \\ 0 & |\zeta_{x,j,t}|^{-1/2} e^{-l_{x,j,t}(|\zeta_{x,j,t}|)} \end{pmatrix},$$

where $l_{x,j,t}(r) = \psi_1(8\lambda_{x,j}(t)r^{3/2}/3)$, and ψ_1 satisfies the ODE

$$\left(\frac{d^2}{d\rho^2} + \frac{1}{\rho} \frac{d}{d\rho} \right) \psi_1(\rho) = \frac{1}{2} \sinh(2\psi_1(\rho)), \quad (14)$$

with asymptotics

$$\psi_1(\rho) \sim \frac{1}{\pi} K_0(\rho) \text{ as } \rho \rightarrow \infty, \quad \psi_1(\rho) \sim -\frac{1}{3} \log \rho \text{ as } \rho \rightarrow 0.$$

Here K_0 is the modified Bessel function of the second kind, $K_0(\rho) \sim \rho^{-1/2} e^{-\rho}$ as $\rho \rightarrow \infty$. Then we define

$$h_t^{\text{app}} = \begin{pmatrix} |\zeta_{x,j,t}|^{1/2} e^{\chi(|\zeta_{x,j,t}| \kappa^{-1} t^{1/j(x)}) l_{x,j,t}(|\zeta_{x,j,t}|)} & 0 \\ 0 & |\zeta_{x,j,t}|^{-1/2} e^{-\chi(|\zeta_{x,j,t}| \kappa^{-1} t^{1/j(x)}) l_{x,j,t}(|\zeta_{x,j,t}|)} \end{pmatrix}, \quad (15)$$

where χ is a nonincreasing smooth cutoff function with $\chi(r) = 1$ for $r \leq 1/2$ and $\chi(r) = 0$ for $r \geq 1$.

(ii) For $x \in T$, in the above holomorphic frame over $\tilde{B}_{x,t}$, define

$$h_t^{\text{model}} = \begin{pmatrix} |\zeta_{x,t}|^{1/2} e^{m_{x,t}(|\zeta_{x,t}|)} & 0 \\ 0 & |\zeta_{x,t}|^{3/2} e^{-m_{x,t}(|\zeta_{x,t}|)} \end{pmatrix},$$

where $m_{x,t}(r) = \psi_{2,x}(8\lambda_x(t)r^{1/2})$, and $\psi_{2,x}$ satisfies (14) with asymptotics

$$\psi_{2,x}(\rho) \sim \frac{1}{\pi} K_0(\rho) \text{ as } \rho \rightarrow \infty, \quad \psi_{2,x}(\rho) \sim (1 + 2\alpha_{x,1} - 2\alpha_{x,2}) \log \rho \text{ as } \rho \rightarrow 0.$$

Then we define

$$h_t^{\text{app}} = \begin{pmatrix} |\zeta_{x,t}|^{1/2} e^{\chi(|\zeta_{x,t}| \kappa^{-1}) m_{x,t}(|\zeta_{x,t}|)} & 0 \\ 0 & |\zeta_{x,t}|^{3/2} e^{-\chi(|\zeta_{x,t}| \kappa^{-1}) m_{x,t}(|\zeta_{x,t}|)} \end{pmatrix}. \quad (16)$$

It is proved in [MTW77] that there exist unique functions $\psi_1, \psi_{2,x}$ solving (14) with the prescribed asymptotics. Finally, set $h_t^{\text{app}} = h_{t,\infty}$ on the complement of the above disks.

It is straightforward to check that h_t^{model} satisfies Hitchin's equation over $\tilde{B}_{x(j),t}$, smooth near $z_{x,j}(t) \in Z_t \setminus (I_t \cup T)$ and adapted to the parabolic structure at $x \in T$. h_t^{app} approximately solves Hitchin's equation in the following sense.

Proposition 3.5. *There exists positive constants t_0, c, c' , such that for any $t \geq t_0$,*

$$\left\| F_{h_t^{\text{app}}} + \left[\Phi_t, \Phi_t^{*h_t^{\text{app}}} \right] \right\|_{L^2} \leq ce^{-c't^\sigma}, \quad (17)$$

where the L^2 -norm is taken with respect to h_t^{app} and a fixed Riemannian metric on C , and

$$\sigma = \min \left(\frac{m_0 - 1}{m_0 + N - 4}, \min_{x \in I_u \setminus \{0\}} \frac{m_x - 1}{m_x}, \min_{x \in I_t} \frac{2m_x - 3}{2(m_x - 1)}, \frac{1}{2} \delta_{T \neq \emptyset} \right),$$

$\delta_{T \neq \emptyset} = 1$ if $T \neq \emptyset$, otherwise it equals to 2.

Proof. Since h_t^{app} coincides with h_t^{model} and $h_{t,\infty}$ away from some annuli, solving Hitchin's equation, we only need to estimate the error in these regions. First, we consider $\text{Ann}_{x,j} = \{ \kappa t^{-1/j(x)}/2 < |\zeta_{x,j,t}| < \kappa t^{-1/j(x)} \}$. Denote $|\zeta_{x,j,t}|$ by r for simplicity.

$$\begin{aligned} F_{h_t^{\text{app}}} &= -\frac{1}{4} (\Delta l_{x,j,t}(r) \chi(r \kappa^{-1} t^{1/j(x)})) \sigma_3 d\zeta_{x,j,t} d\bar{\zeta}_{x,j,t} \\ &= -\frac{1}{4} \left(l''_{x,j,t}(r) \chi(r \kappa^{-1} t^{1/j(x)}) + 2\kappa^{-1} t^{1/j(x)} l'_{x,j,t}(r) \chi'(r \kappa^{-1} t^{1/j(x)}) \right. \\ &\quad \left. + \kappa^{-2} t^{2/j(x)} l_{x,j,t}(r) \chi''(r \kappa^{-1} t^{1/j(x)}) \right. \\ &\quad \left. + \frac{1}{r} l'_{x,j,t}(r) \chi(r \kappa^{-1} t^{1/j(x)}) + \frac{1}{r} \kappa^{-1} t^{1/j(x)} l_{x,j,t}(r) \chi'(r \kappa^{-1} t^{1/j(x)}) \right) \sigma_3 d\zeta_{x,j,t} d\bar{\zeta}_{x,j,t}. \end{aligned}$$

On the other hand,

$$\left[\Phi, \Phi^{*h_t^{\text{app}}} \right] = 2\lambda_{x,j}(t)^2 r \sinh(2l_{x,j,t}(r) \chi(r \kappa^{-1} t^{1/j(x)})) \sigma_3 d\zeta_{x,j,t} d\bar{\zeta}_{x,j,t}.$$

By (14) and the definition of $l_{x,j,t}$, we have

$$l''_{x,j,t}(r) + r^{-1} l'_{x,j,t}(r) = 8\lambda_{x,j}(t)^2 r \sinh(2l_{x,j,t}), \quad l_{x,j,t} = O(e^{-c'\lambda_{x,j}(t)t^{-3/2j(x)}}).$$

We can write $F_{h_t^{\text{app}}} + \left[\Phi_t, \Phi_t^{*h_t^{\text{app}}} \right] = E_{x,j,t} \sigma_3 d\zeta_{x,j,t} d\bar{\zeta}_{x,j,t}$, the L^2 norm of which is bounded by $ct^{-1/j(x)} \sup |E_{x,j,t}|$. Here and below, $c, c' > 0$ are generic constants which may vary, but does not depend on t . By Lemma 3.1, as $t \rightarrow \infty$,

$$\lambda_{x,j}(t) \sim t^{1/2} \prod_{y \in S} |z_{x,j}(t) - y|^{-m_x} \prod_{(y,k) \neq (x,j)} |z_{x,j}(t) - z_{y,k}(t)|^{1/2} \sim \begin{cases} c't^{\frac{2m_x+1}{2j(x)}} & \text{if } x \in I_u, \\ c't^{\frac{m_x}{j(x)}} & \text{if } x \in I_t. \end{cases}$$

Then for $x \in I_u$,

$$\begin{aligned} |E_{x,j,t}| &\leq c (\lambda_{x,j}(t)^2 r (|\sinh(2l_{x,j,t}) - \sinh(2l_{x,j,t}\chi)| + |\chi - 1| |\sinh(2l_{x,j,t})|) \\ &\quad + t^{1/j(x)} |l'_{x,j,t}| + t^{2/j(x)} |l_{x,j,t}|) \\ &\leq c (t^{2m_x/j(x)} \cosh(2l_{x,j,t}) |l_{x,j,t}| + t^{1/j(x)} |l'_{x,j,t}| + t^{2/j(x)} |l_{x,j,t}|) \\ &\leq c (t^{2m_x/j(x)} |l_{x,j,t}| + t^{2/j(x)} |l'_{x,j,t}|) \leq ce^{-c't^{(m_x-1)/j(x)}}. \end{aligned}$$

For $x \in I_t$, $|E_{x,j,t}| \leq ce^{-c't(2m_x-3)/2j(x)}$. Let $\text{Ann}_x = \{\kappa/2 < |\zeta_{x,t}| < \kappa\}$ for $x \in T$, and denote $r = |\zeta_{x,t}|$. Similar as before,

$$\begin{aligned} F_{h_t^{\text{app}}} &= -\frac{1}{4} \left(m''_{x,t}(r) \chi(r\kappa^{-1}) + 2\kappa^{-1} m'_{x,t}(r) \chi'(r\kappa^{-1}) + \kappa^{-2} m_{x,t}(r) \chi''(r\kappa^{-1}) \right. \\ &\quad \left. + \frac{1}{r} m'_{x,t}(r) \chi(r\kappa^{-1}) + \frac{1}{r} \kappa^{-1} m_{x,t}(r) \chi'(r\kappa^{-1}) \right) \sigma_3 d\zeta_{x,t} d\bar{\zeta}_{x,t}, \\ \left[\Phi, \Phi^{*h_t^{\text{app}}} \right] &= 2\lambda_x(t)^2 r^{-1} \sinh(2m_{x,t}(r) \chi(r\kappa^{-1})) \sigma_3 d\zeta_{x,t} d\bar{\zeta}_{x,t}. \end{aligned}$$

Now we have

$$m''_{x,t}(r) + r^{-1} m'_{x,t}(r) = 8\lambda_x(t)^2 r^{-1} \sinh(2m_{x,t}), \quad m_{x,j,t} = O(e^{-c'\lambda_x(t)}),$$

where $\lambda_x(t) \sim c't^{1/2}$. For $E_{x,t}$ defined similarly as above, we have the estimate

$$|E_{x,t}| \leq ce^{-c't^{1/2}}.$$

Combining all the estimates above, the result follows. \square

Remark 3.6. If we allow tame untwisted singularities (also called weakly parabolic), then $\lambda_{x,j}(t) \sim c't^{3/2}$, and $l_{x,j,t} = O(e^{-c't^{3/2}t^{-3/2}}) = O(1)$ which does not decay to 0 as $t \rightarrow \infty$. Thus, the error cannot be arbitrarily small when tame untwisted poles exist.

4. LINEAR ANALYSIS

We wish to obtain the solution h_t to Hitchin's equation by perturbing h_t^{app} . This is equivalent to finding an $\text{SL}(E)$ -valued h_t^{app} -Hermitian section H_t such that $h_t(v, w) = h_t^{\text{app}}(H_t v, w)$. Let $e^{\gamma_t} = H_t^{-1/2}$, then $e^{\gamma_t} \cdot h_t(v, w) = h_t(H_t^{-1/2} v, H_t^{-1/2} w) = h_t^{\text{app}}(v, w)$. By the gauge invariance of Hitchin's equations, $(\bar{\partial}_E, \Phi_t, h_t)$ is a solution is equivalent to that

$$e^{\gamma_t} \cdot (\bar{\partial}_E, \Phi_t, h_t) = (e^{-\gamma_t} \bar{\partial}_E e^{\gamma_t}, e^{-\gamma_t} \Phi_t e^{\gamma_t}, h_t^{\text{app}})$$

is a solution. Therefore, we need to find $\gamma_t \in \Omega^0(i\mathfrak{su}(E))$ (the bundle is defined over $C \setminus S$ using E and h_t^{app}), such that $F_t(\gamma_t) = 0$, where

$$F_t(\gamma) := F_{A_t^{\text{exp}(\gamma)}} + [e^{-\gamma} \Phi_t e^{\gamma}, e^{\gamma} \Phi_t^{*h_t^{\text{app}}} e^{-\gamma}],$$

A_t is the Chern connection $D(\bar{\partial}_E, h_t^{\text{app}})$ and $A_t^{\text{exp}(\gamma)} = D(e^{-\gamma} \bar{\partial}_E e^{\gamma}, h_t^{\text{app}})$. Let $d_{A_t} = \partial_{A_t} + \bar{\partial}_E$ be the exterior derivative associated with A_t , ∇_{A_t} be the covariant derivative. We have

$$F_{A_t^{\text{exp}(\gamma)}} = e^{-\gamma} (F_{A_t} + \bar{\partial}_{A_t} (e^{2\gamma} \partial_{A_t} (e^{-2\gamma}))) e^{\gamma}.$$

Linearize F_t at $\gamma = 0$ we get

$$\begin{aligned} DF_t(\gamma) &= [F_{A_t}, \gamma] - 2\bar{\partial}_{A_t} \partial_{A_t} \gamma + [[\Phi_t, \gamma], \Phi_t^{*h_t^{\text{app}}}] + [\Phi_t, [\gamma, \Phi_t^{*h_t^{\text{app}}}] \\ &= i \star \Delta_{A_t} \gamma + M_{\Phi_t} \gamma, \end{aligned}$$

where $\Delta_{A_t} = d_{A_t}^* d_{A_t}$ on $\Omega^0(i\mathfrak{su}(E))$, $M_{\Phi_t} \gamma = [\Phi_t^{*h_t^{\text{app}}}, [\Phi_t, \gamma]] - [\Phi_t, [\Phi_t^{*h_t^{\text{app}}}, \gamma]]$, and we used the identity $2\bar{\partial}_A \partial_A = F_A - i \star \Delta_A$. By composing $-i \star : \Omega^2(\mathfrak{su}(E)) \rightarrow \Omega^0(i\mathfrak{su}(E))$ we get the operator

$$L_t(\gamma) = \Delta_{A_t} \gamma - i \star M_{\Phi_t} \gamma. \quad (18)$$

Consider the irregular connection defined by $D_t^\gamma = d_{A_t^{\exp(\gamma)}} + e^{-\gamma}\Phi_t e^\gamma + e^\gamma\Phi_t^* e^{-\gamma}$, here and below we drop the subscript of $*_{h_t^{\text{app}}}$. Then its curvature $(D_t^\gamma)^2 = F_t(\gamma)$, since $\bar{\partial}_E\Phi_t = \partial_{A_t}\Phi_t^* = 0$. So we can also regard the problem as finding γ so that this connection is flat. When $\gamma = 0$, write the operator as D_t , which is the sum of the unitary part d_{A_t} and the self-adjoint part $\Psi_t := \Phi_t + \Phi_t^*$.

Lemma 4.1. (*Weitzenböck formulas*) *Suppose $\gamma \in \Omega^0(\text{isu}(E))$, $w \in \Omega^p(\text{End } E)$, then*

- (i) $(D_t^*D_t + D_tD_t^*)w = \nabla_{A_t}^*\nabla_{A_t}w + (\Psi_t \otimes)^*\Psi_t \otimes w + \mathcal{F}(w) + \mathcal{R}(w)$, where \mathcal{F}, \mathcal{R} are curvature operators, $\Psi_t \otimes \cdot$ is a combination of tensor product on the manifold part and Lie bracket on the bundle part. More explicitly, let e_1, e_2 be a local orthonormal frame of TC and e^1, e^2 be the dual frame, write the curvature D_t^2 as $F_{t,12}e^1 \wedge e^2$, then $\mathcal{F}(w) = [F_{t,12}, (e^1 \wedge i_{e_2} - e^2 \wedge i_{e_1})w]$, and

$$\begin{aligned} \mathcal{R}(w)(X_1, \dots, X_p) &= \sum_{j=1}^p w(X_1, \dots, \text{Ric}(X_j), \dots, X_p) \\ &\quad + \sum_{a=1}^2 \sum_{j < k} (-1)^{j+k+1} w(e_a, R(X_j, X_k)e_a, X_1, \dots, \hat{X}_j, \dots, \hat{X}_k, \dots, X_p). \end{aligned}$$

- (ii) $D_t^*D_t\gamma = d_{A_t}^*d_{A_t}\gamma + ([\Psi_t, \cdot])^*[\Psi_t, \gamma] = L_t\gamma$.

- (iii) $\text{Re}(\langle d_{A_t}\gamma, [\Psi_t, \gamma] \rangle) = 0$, so the following pointwise equality holds

$$\langle D_t\gamma, D_t\gamma \rangle = |D_t\gamma|^2 = |d_{A_t}\gamma|^2 + |[\Psi_t, \gamma]|^2.$$

Proof. (i) This follows from [Biq97, Theorem 5.4], since $d_{A_t}^*\Psi_t = 0$.

(ii) The first equality is a special case of (i). The adjoint of $[\Phi_t, \cdot]$ is $-\bar{*}[\Phi_t, \bar{*}]$, where $\bar{*}$ is the conjugate linear operator combining $\bar{\star}$ and the Hermitian adjoint. Actually, for a fixed holomorphic coordinate z where $\Phi = \varphi dz$ (subscript t omitted),

$$\begin{aligned} \langle [\Phi, a], b dz \rangle &= \langle [\varphi, a] dz, b dz \rangle = 2\langle [\varphi, a], b \rangle, \\ \langle a, -\bar{*}[\Phi, \bar{*}b dz] \rangle &= \langle a, -\bar{*}[\Phi, ib^* d\bar{z}] \rangle = \langle a, -[b, \varphi^*] \bar{\star}(i dz d\bar{z}) \rangle = 2\langle a, [\varphi^*, b] \rangle = 2\langle [\varphi, a], b \rangle. \end{aligned}$$

Similarly, the adjoint of $[\Phi_t^*, \cdot]$ is $-\bar{*}[\Phi_t^*, \bar{*}]$. Then (recall that $\bar{d}z = i d\bar{z}$, $\bar{d}\bar{z} = -i dz$)

$$\begin{aligned} ([\Psi_t, \cdot])^*[\Psi_t, \gamma] &= -\bar{*}[\Phi_t, \bar{*}[\Phi_t, \gamma]] - \bar{*}[\Phi_t^*, \bar{*}[\Phi_t^*, \gamma]] \\ &= -\bar{*}[\Phi_t, i[\Phi_t^*, \gamma]] + \bar{*}[\Phi_t^*, i[\Phi_t, \gamma]] \\ &= -i \star [\Phi_t^*, [\Phi_t, \gamma]] + i \star [\Phi_t, [\Phi_t^*, \gamma]] = -i \star M_{\Phi_t}\gamma. \end{aligned}$$

(iii) Write $d_{A_t}\gamma = \gamma_z dz + \gamma_{\bar{z}} d\bar{z}$. $\gamma_{\bar{z}} = \gamma_z^*$, since $\gamma = \gamma^*$ and d_{A_t} is a Hermitian connection. As above, write $\Phi = \varphi dz$, then

$$\begin{aligned} \text{Re}(\langle d_{A_t}\gamma, [\Phi + \Phi^*, \gamma] \rangle) &= 2\text{Re}(\text{tr}(\gamma_z[\varphi, \gamma]^*) + \text{tr}(\gamma_{\bar{z}}[\varphi^*, \gamma]^*)) \\ &= 2\text{Re}(\text{tr}(\gamma_z[\varphi, \gamma]^*) + \text{tr}(\gamma_z^*[\gamma, \varphi])) \\ &= 2\text{Re}\left(\text{tr}(\gamma_z[\varphi, \gamma]^*) - \overline{\text{tr}(\gamma_z[\varphi, \gamma]^*)}\right) = 0. \end{aligned}$$

□

Choose a local holomorphic trivialization as in Proposition 3.3 (iii). Consider the map $\sigma_{x,t} : D_{x,t} \rightarrow \tilde{B}_{x,t}$, $\xi_{x,t} \rightarrow \xi_{x,t}^2 = \zeta_{x,t}$, where $D_{x,t} = \{|\xi_{x,t}| < \kappa^{1/2}t^{-1/2(m_x-1)}\}$. Then for the

pullbacks $\tilde{E} = \sigma_{x,t}^* E$, $\tilde{\Phi}_t = \sigma_{x,t}^* \Phi_t$, $\tilde{h}_t^{\text{app}} = \sigma_{x,t}^* h_t^{\text{app}}$, we have

$$\tilde{\partial}_E = \bar{\partial}, \quad \tilde{\Phi}_t = 2\lambda_x(t) \begin{pmatrix} 0 & \frac{1}{\xi_{x,t}^{2m_x-3}} \\ \frac{1}{\xi_{x,t}^{2m_x-1}} & 0 \end{pmatrix} d\xi_{x,t}, \quad \tilde{h}_t^{\text{app}} = \begin{pmatrix} |\xi_{x,t}| & 0 \\ 0 & |\xi_{x,t}|^3 \end{pmatrix}.$$

After applying the gauge transformation $\begin{pmatrix} \xi_{x,t} & -\xi_{x,t} \\ 1 & 1 \end{pmatrix}$ (noninvertible at 0),

$$\tilde{\Phi}_t = 2\lambda_x(t) \begin{pmatrix} \frac{1}{\xi_{x,t}^{2(m_x-1)}} & 0 \\ 0 & \frac{1}{\xi_{x,t}^{2(m_x-1)}} \end{pmatrix} d\xi_{x,t}, \quad \tilde{h}_t^{\text{app}} = 2 \begin{pmatrix} |\xi_{x,t}|^3 & 0 \\ 0 & |\xi_{x,t}|^3 \end{pmatrix}. \quad (19)$$

For $x \in T$, denote $\hat{B}_{x,t} := \{|\zeta_{x,t}| < \lambda_x(t)^{-2}\}$. Choose $t_0 \geq 1$ so that when $t \geq t_0$, we have $\lambda_x(t)^{-2} < \kappa/2$, and then $h_t^{\text{app}} = h_t^{\text{model}}$ over $\hat{B}_{x,t}$. From now on, we assume that $t \geq t_0$.

Definition 4.2. Let r_t be a smooth weight function on C such that

- (i) $r_t = 1$ in $C \setminus (\bigcup_{x \in I} \tilde{B}_{x,t} \cup \bigcup_{x \in T} \hat{B}_{x,t})$.
- (ii) In $\tilde{B}_{x,t}$, $x \in S$, r_t is nondecreasing in $|\zeta_{x,t}|$ (or $|z_x|$ for $x \in I_u$).
- (iii)

$$r_t = \begin{cases} |t^{1/j(x)} z_x| & \text{if } x \in I_u, |z_x| \leq \kappa t^{-1/j(x)}/2, \\ |t^{1/(m_x-1)} \zeta_{x,t}|^{1/2} & \text{if } x \in I_t, |\zeta_{x,t}| \leq \kappa t^{-1/(m_x-1)}/2, \\ |\lambda_x(t)^2 \zeta_{x,t}| & \text{if } x \in T, |\zeta_{x,t}| \leq \lambda_x(t)^{-2}/2. \end{cases}$$

Denote $B_x = \{|z_x| < \kappa_2\}$, where κ_2 is from Proposition 3.3, and we can assume that κ_2 is small enough so that the disks B_x are disjoint.

Definition 4.3. Let $g_t = f_t g_C$ be the metric on C conformal to the round metric g_C (Hitchin's equations are conformally invariant, we consider L_t above with \star and Δ_{A_t} defined using g_t), which satisfies

- (i) $f_t = 1$ in $C \setminus (\bigcup_{x \in S} B_x)$, independent of t in $C \setminus (\bigcup_{x \in I_t \cup T} B_x)$.
- (ii) g_t induces the volume form $(i/8|\zeta_{x,t}|) d\zeta_{x,t} d\bar{\zeta}_{x,t}$ when restricted to $\tilde{B}_{x,t}$ for $x \in I_t$, its pullback under $\sigma_{x,t}$ is $(i/2) d\xi_{x,t} d\bar{\xi}_{x,t}$, the Euclidean volume form on $D_{x,t}$.
- (iii) g_t is Euclidean in B_x for $x \in I_u$.
- (iv) g_t is Euclidean in $\tilde{B}_{x,t}$ for $x \in T$.
- (v) $|\nabla_{g_C}^j \log f_t|_{g_C} \leq c$ in B_x for $x \in I_u \cup T$, $j = 0, 1, 2$, and $c^{-1}|z_x|^{-1} \leq f_t \leq c|z_x|^{-1}$, $|\nabla_{g_C}^j \log f_t|_{g_C} \leq ct^{j/(m_x-1)}$ in $B_x \setminus \tilde{B}_{x,t}$ for $x \in I_t$, $j = 1, 2$.

Definition 4.4. For smooth sections $w \in \Omega^p(\text{End}E)$, define

$$\|w\|_{L_\delta^p} = \|r_t^{-\delta-2/p} w\|_{L^p}, \quad \|w\|_{L_\delta^{k,p}}^p = \|w\|_{L_\delta^{k-1,p}}^p + \|\nabla_{A_t} w\|_{L_\delta^{k-1,p}}^p + \|\Psi_t \otimes w\|_{L_\delta^{k-1,p}}^p \quad (20)$$

inductively for $k = 1, 2, \dots$. Here the L^p norms are taken with respect to g_t and h_t^{app} . Let the spaces L_δ^p and $L_\delta^{k,p}$ be the completions of smooth sections over $C \setminus S$ with respect to these norms. We also introduce spaces with additional weighted conditions,

$$\hat{L}_\delta^{k,p} = \left\{ f \in L_\delta^{k,p}, r_t^{j-k} \nabla_{A_t}^j f \in L_\delta^p \text{ for } 0 \leq j \leq k \right\}.$$

By using cutoff functions of the form $1 - \chi(nr_t)$, one can see that $C_0^\infty(C \setminus S)$ is dense in $\hat{L}_\delta^{k,p}$.

In $\tilde{B}_{x,t}$ for $x \in I_u$, we can decompose $\text{End } E$ as $\text{End}_D E \oplus \text{End}_T E$ in the holomorphic frame above, where End_D is the diagonal part. Then $\text{End}_D E = \ker \text{ad}(\Phi_t)$, $\text{End}_T E = (\ker \text{ad}(\Phi_t))^\perp$. Φ_t acts on $u \in \text{End}_T E$ as

$$[\Phi_t, u] = 2 \frac{\sqrt{-z_x^{2m_x} \nu_t(z)}}{z_x^{m_x}} \begin{pmatrix} 0 & u_{12} \\ -u_{21} & 0 \end{pmatrix} dz \Rightarrow |[\Phi_t, u]| = 2 \frac{|-z_x^{2m_x} \nu_t(z)|^{1/2}}{|z_x|^{m_x}} |u|,$$

and then

$$c^{-1} t^{m_x/j(x)} |u| r_t^{-m_x} \leq |[\Phi_t, u]| \leq c t^{m_x/j(x)} |u| r_t^{-m_x} \text{ in } \tilde{B}_{x,t}. \quad (21)$$

Therefore, over $\tilde{B}_{x,t}$, the spaces defined above are equivalent (may not be uniformly in t) to the spaces defined in [BB04], using the irregular Hermitian connection $A_t + \Phi_t - \Phi_t^*$ and the decomposition of $\text{End } E$ into components $\text{End}(E)_0 = \text{End}_D E$, $\text{End}(E)_{m_x} = \text{End}_T E$, from which we collect the following facts over $\tilde{B}_{x,t}$.

Lemma 4.5. (i) For functions, we have $\hat{L}_{\delta-1}^{1,p} \hookrightarrow L_\delta^q$ if $1/2 \geq 1/p - 1/q$ (with strict inequality for $q = \infty$). In particular, for $p > 2$ we have $\hat{L}_{\delta-1}^{1,p} \hookrightarrow C_\delta^0$.

(ii) If $\delta < 0$ and the function f vanishes on $\partial \tilde{B}_{x,t}$, or if $\delta > 0$ and f vanishes near x , then

$$\|\partial f / \partial r\|_{L_{\delta-1}^p} \geq c^{-1} t^{1/j(x)} \|f\|_{L_\delta^p},$$

where $r = |z_x|$. This implies that $\hat{L}_{\delta-1}^{1,p} = L_{\delta-1}^{1,p}$ for $\delta < 0$ and $p \in [1, \infty)$.

(iii) If $u \in L_{-2+\delta}^{2,2}$ is a section of $\text{End}_D E$, then u is continuous and $u - u(x) \in \hat{L}_{-2+\delta}^{2,2} \subset C_\delta^0 \cap L_\delta^p$.

(iv) For $\delta' < \delta$, $p > 2$, there are compact Sobolev embeddings for sections of $\text{End}_T E$:

$$L_\delta^{1,2} \hookrightarrow L_{\delta'+1+(2m_x-2)/p}^p, \quad L_\delta^{1,p} \hookrightarrow C_{\delta'+m_x-2(m_x-1)/p}^0, \quad L_\delta^{2,2} \hookrightarrow C_{\delta'+m_x+1}^0.$$

Now we consider the Dirichlet problem on $\tilde{B}_{x,t}$ and derive some a priori estimates. The proofs of the following two results are adapted from [BB04, Cor. 4.2, Lem. 4.4], with the t -dependence of the operator L_t taken into account.

Lemma 4.6. Suppose $w \in L_{-2+\delta}^{1,2}(\Omega^1(\text{End}_T E))$, which vanishes on $\partial \tilde{B}_{x,t}$, then

$$\|D_t w\|_{L_{-2+\delta}^2} + \|D_t^* w\|_{L_{-2+\delta}^2} \geq c^{-1} (\|\nabla_{A_t} w\|_{L_{-2+\delta}^2} + \|\Psi_t \otimes w\|_{L_{-2+\delta}^2}),$$

for all weights δ , when t is sufficiently large.

Proof. By Lemma 4.1 (i), noting that the curvature terms vanish,

$$\|(D_t + D_t^*)(r_t^{1-\delta} w)\|_{L^2}^2 = \|\nabla_{A_t}(r_t^{1-\delta} w)\|_{L^2}^2 + \|r_t^{1-\delta} \Psi_t \otimes w\|_{L^2}^2.$$

The commutator can be controlled as

$$|[D_t + D_t^*, r_t^{1-\delta}]w| + |[\nabla_{A_t}, r_t^{1-\delta}]w| \leq c t^{1/j(x)} |r_t^{-\delta} w|.$$

Then we have

$$\|r_t^{1-\delta} (D_t + D_t^*) w\|_{L^2} \geq c^{-1} (\|r_t^{1-\delta} \nabla_{A_t} w\|_{L^2} + \|r_t^{1-\delta} \Psi_t \otimes w\|_{L^2}) - c t^{1/j(x)} \|r_t^{-\delta} w\|_{L^2}.$$

On the other hand,

$$\|r_t^{-\delta} w\|_{L^2} \leq \|r_t^{1-\delta} r_t^{-m_x} w\|_{L^2} \leq c t^{-m_x/j(x)} \|r_t^{1-\delta} \Psi_t \otimes w\|_{L^2}.$$

Thus

$$\begin{aligned} \|r_t^{1-\delta}(D_t + D_t^*)w\|_{L^2} &\geq (c^{-1} - ct^{-(m_x-1)/j(x)}) (\|r_t^{1-\delta}\nabla_{A_t}w\|_{L^2} + \|r_t^{1-\delta}\Psi_t \otimes w\|_{L^2}) \\ &\geq (c^{-1}/2) (\|r_t^{1-\delta}\nabla_{A_t}w\|_{L^2} + \|r_t^{1-\delta}\Psi_t \otimes w\|_{L^2}), \end{aligned}$$

when t is large enough. \square

Proposition 4.7. *On $\tilde{B}_{x,t}$ for $x \in I_u$, $L_t : L_{-2+\delta}^{2,2}(i\mathfrak{su}(E)) \rightarrow L_{-2+\delta}^2(i\mathfrak{su}(E))$ is an isomorphism with zero Dirichlet boundary condition for small weights $\delta > 0$. When restricted to the off-diagonal part, the same holds for any δ . Moreover, the norm of L_t^{-1} is uniformly bounded in t .*

Proof. Observe that the decomposition $\text{End } E = \text{End}_D E \oplus \text{End}_T E$ is preserved by L_t , we can consider the two components separately. For the off-diagonal part, we find a solution of $L_t u = v$ by minimizing the functional (recall Lemma 4.1 (iii))

$$S(u) = \frac{1}{2} \int_{\tilde{B}_{x,t}} |D_t u|^2 \, \text{dvol} - \langle u, v \rangle_{L^2} = \frac{1}{2} \|d_{A_t} u\|_{L^2}^2 + \|[\Phi_t, u]\|_{L^2}^2 - \langle u, v \rangle_{L^2}$$

among $u \in L_{-1}^{1,2}(i\mathfrak{su}_T(E))$ vanishing on the boundary, where $v \in L_{-m_x-1}^2(i\mathfrak{su}_T(E))$. $S(u)$ is continuous:

$$S(u) \leq \|\nabla_{A_t} u\|_{L^2}^2 + \|\Psi_t \otimes u\|_{L^2}^2 + \|r_t^{-m_x} u\|_{L^2} \|r_t^{m_x} v\|_{L^2} \leq \|u\|_{L_{-1}^{1,2}}^2 + c \|u\|_{L_{-1}^{1,2}} \|v\|_{L_{-m_x-1}^2},$$

where we used (21) in the last inequality. On the other hand,

$$\begin{aligned} S(u) &= \frac{1}{2} (\|d_{A_t} u\|_{L^2}^2 + 2\|[\Phi_t, u]\|_{L^2}^2) - \langle u, v \rangle_{L^2} \\ &\geq c^{-1} \left(\|\nabla_{A_t} u\|_{L^2}^2 + \|\Psi_t \otimes u\|_{L^2}^2 + t^{2m_x/j(x)} \|r_t^{-m_x} u\|_{L^2}^2 \right) - \|r_t^{-m_x} u\|_{L^2} \|v\|_{L_{-m_x-1}^2} \\ &\geq c^{-1} \|u\|_{L_{-1}^{1,2}}^2 - c \|u\|_{L_{-1}^{1,2}} \|v\|_{L_{-m_x-1}^2}. \end{aligned} \tag{22}$$

Therefore $S(u)$ is coercive. If $d_{A_t} u = 0$, then $|u|^2$ is constant and $u = 0$ since it vanishes on the boundary. This implies that S is strictly convex, and a unique minimum of S can be found, being a weak solution of the equation of the form

$$\Delta_{A_t} (r_t^{m_x} u) = r_t^{m_x} v + P_0 (r_t^{-m_x} u) + P_1 (\nabla_{A_t} u),$$

where $P_0, P_1 : L^2 \rightarrow L^2$ are bounded. By elliptic regularity, $r_t^{m_x} \nabla_{A_t}^2 u \in L^2$, and then $u \in L_{-m_x-1}^{2,2}$. With the Dirichlet boundary condition, we have obtained an isomorphism $L_t : L_{-m_x-1}^{2,2} \rightarrow L_{-m_x-1}^2$. The solution satisfies $\langle D_t u, D_t f \rangle_{L^2} = \langle v, f \rangle_{L^2}$ for any $f \in L_{-1}^{1,2}$ vanishing on the boundary. Choose $f = u$ (the first inequality follows by setting $v = 0$ in (22)),

$$\begin{aligned} \|u\|_{L_{-1}^{1,2}}^2 &\leq c \|D_t u\|_{L^2}^2 = c \langle v, u \rangle_{L^2} \leq ct^{-m_x/j(x)} \|\Psi_t \otimes u\|_{L^2} \|r_t^{m_x} v\|_{L^2} \\ &\leq ct^{-m_x/j(x)} \|u\|_{L_{-1}^{1,2}} \|v\|_{L_{-m_x-1}^2} \\ &\Rightarrow \|u\|_{L_{-1}^{1,2}} \leq ct^{-m_x/j(x)} \|v\|_{L_{-m_x-1}^2}. \end{aligned} \tag{23}$$

For a one form w with coefficient in $i\mathfrak{su}(E)$, by Lemma 4.6 we have

$$\|\nabla_{A_t} w\|_{L_{-m_x-1}^2} \leq c \left(\|D_t w\|_{L_{-m_x-1}^2} + \|D_t^* w\|_{L_{-m_x-1}^2} \right).$$

Then for $w = \nabla_{A_t} u$,

$$\begin{aligned} D_t w &= (d_{A_t} + [\Psi_t, \cdot])(d_{A_t} u) = [\Psi_t, d_{A_t} u], \\ D_t^* w &= D_t^* D_t u - D_t^* [\Psi_t, u] = D_t^* D_t u + *[\Psi_t, d_{A_t} u] - [\Psi_t, \cdot]^* [\Psi_t, u], \\ \|r_t^{m_x} (\Psi_t \otimes \cdot)^2 u\|_{L^2} &\leq t^{m_x/j(x)} \|\Psi_t \otimes u\|_{L^2} \leq t^{m_x/j(x)} \|u\|_{L_{-1}^{1,2}}, \\ \|r_t^{m_x} \Psi_t \otimes \nabla_{A_t} u\|_{L^2} &\leq t^{m_x/j(x)} \|\nabla_{A_t} u\|_{L^2} \leq t^{m_x/j(x)} \|u\|_{L_{-1}^{1,2}}, \\ \|r_t^{m_x} \nabla_{A_t}^2 u\|_{L^2} &\leq c \left(t^{m_x/j(x)} \|u\|_{L_{-1}^{1,2}} + \|D_t^* D_t u\|_{L^2_{-m_x-1}} \right) \leq c \|v\|_{L^2_{-m_x-1}}, \end{aligned}$$

by (23). Similarly, with $w = [\Psi_t, u]$ one deduces that $\|r_t^{m_x} \nabla_{A_t} [\Psi_t, u]\|_{L^2} \leq c \|v\|_{L^2_{-m_x-1}}$. Hence

$$\begin{aligned} \|u\|_{L_{-m_x-1}^{2,2}} &\leq c \left(\|u\|_{L_{-1}^{1,2}} + \|r_t^{m_x} \nabla_{A_t}^2 u\|_{L^2} + \|r_t^{m_x} \nabla_{A_t} [\Psi_t, u]\|_{L^2} \right. \\ &\quad \left. + \|r_t^{m_x} (\Psi_t \otimes \cdot)^2 u\|_{L^2} + \|r_t^{m_x} \Psi_t \otimes \nabla_{A_t} u\|_{L^2} \right) \leq c \|v\|_{L^2_{-m_x-1}}. \end{aligned}$$

Next we prove that the isomorphism extends to all weights, by showing that the inverse is continuous in other weighted spaces. For any weight δ , it suffices to prove

$$\|r_t^\delta D_t^* D_t u\|_{L^2_{-m_x-1}} = \|D_t^* D_t u\|_{L^2_{-m_x-1-\delta}} \geq c^{-1} \|u\|_{L_{-m_x-1-\delta}^{2,2}},$$

using the established a priori estimates

$$\|r_t^\delta u\|_{L_{-m_x-1}^{2,2}} \leq c \|D_t^* D_t r_t^\delta u\|_{L^2_{-m_x-1}}. \quad (24)$$

We have

$$\begin{aligned} \| [D_t^* D_t, r_t^\delta] u \|_{L^2_{-m_x-1}} &\leq c t^{2/j(x)} \|r_t^{\delta-2} u\|_{L^2_{-m_x-1}} \\ &\quad + c t^{1/j(x)} \left(\|r_t^{\delta-1} \nabla_{A_t} u\|_{L^2_{-m_x-1}} + \|r_t^{\delta-1} [\Phi_t, u]\|_{L^2_{-m_x-1}} \right) \\ &\leq c t^{2/j(x)} \|r_t^\delta r_t^{-2m_x} u\|_{L^2_{-m_x-1}} \\ &\quad + c t^{1/j(x)} \left(\|r_t^\delta r_t^{-m_x} \nabla_{A_t} u\|_{L^2_{-m_x-1}} + \|r_t^\delta r_t^{-m_x} [\Phi_t, u]\|_{L^2_{-m_x-1}} \right) \\ &\leq c t^{(1-m_x)/j(x)} \left(\|r_t^\delta \Psi_t \otimes \nabla_{A_t} u\|_{L^2_{-m_x-1}} + \|r_t^\delta (\Psi_t \otimes \cdot)^2 u\|_{L^2_{-m_x-1}} \right) \\ &\leq c t^{(1-m_x)/j(x)} \|u\|_{L_{-m_x-1-\delta}^{2,2}}. \end{aligned}$$

Similarly, noting that $\|[\nabla_{A_t}, r_t^\delta] w\| \leq c t^{(1-m_x)/j(x)} |r_t^\delta \Psi_t \otimes w|$,

$$\begin{aligned} \|u\|_{L_{-m_x-1-\delta}^{2,2}} &\leq \|r_t^\delta u\|_{L_{-m_x-1}^{2,2}} + \|[\nabla_{A_t}^2, r_t^\delta] u\|_{L^2_{-m_x-1}} + \|[\nabla_{A_t}, r_t^\delta] u\|_{L^2_{-m_x-1}} \\ &\quad + \|\Psi_t \otimes [\nabla_{A_t}, r_t^\delta] u\|_{L^2_{-m_x-1}} + \|[\nabla_{A_t}, r_t^\delta] \Psi_t \otimes u\|_{L^2_{-m_x-1}} \\ &\leq \|r_t^\delta u\|_{L_{-m_x-1}^{2,2}} + c t^{(1-m_x)/j(x)} \|u\|_{L_{-m_x-1-\delta}^{2,2}}. \end{aligned}$$

Finally by (24),

$$\begin{aligned} \|r_t^\delta D_t^* D_t u\|_{L^2_{-m_x-1}} &\geq \|D_t^* D_t r_t^\delta u\|_{L^2_{-m_x-1}} - \| [D_t^* D_t, r_t^\delta] u \|_{L^2_{-m_x-1}} \\ &\geq c^{-1} \|r_t^\delta u\|_{L_{-m_x-1}^{2,2}} - \| [D_t^* D_t, r_t^\delta] u \|_{L^2_{-m_x-1}} \\ &\geq (c^{-1} - c t^{(1-m_x)/j(x)}) \|u\|_{L_{-m_x-1-\delta}^{2,2}} \geq (c^{-1}/2) \|u\|_{L_{-m_x-1-\delta}^{2,2}}, \end{aligned}$$

for t large. We have proved that for the off-diagonal part, $L_t : L_{-2+\delta}^{2,2} \rightarrow L_{-2+\delta}^2$ is an isomorphism for all weights δ with the norm of the inverse uniformly bounded in t .

For the diagonal part, the equation becomes the Laplace equation $\Delta u = v$ with the Dirichlet boundary condition. Let $\tau : \tilde{B}_{x,t} \rightarrow B_0(\kappa) := \{|\tilde{z}| < \kappa\}$, $z_x \mapsto t^{1/j(x)} z_x := \tilde{z}$ be the rescaling map, and $\tilde{u} = (\tau^{-1})^* u$, $\tilde{v} = (\tau^{-1})^* v$. Then $\Delta u = v$ is equivalent to $t^{2/j(x)} \Delta \tilde{u} = \tilde{v}$ on $B_0(\kappa)$, where $\Delta : L_{-2+\delta}^{2,2} \rightarrow L_{-2+\delta}^2$ is an isomorphism by classical elliptic theory on cylinders (see [BB04]), and the weighted spaces on $B_0(\kappa)$ are defined similarly as in (20), using the Euclidean distance from 0 as the weight function, which is equivalent to $(\tau^{-1})^* r_t$ uniformly in t . Then $\|\tilde{v}\|_{L_{-2+\delta}^2} \geq c^{-1} \|t^{2/j(x)} \tilde{u}\|_{L_{-2+\delta}^{2,2}}$, which implies that $\|v\|_{L_{-2+\delta}^2} \geq c^{-1} \|u\|_{L_{-2+\delta}^{2,2}}$. \square

Next we study the behavior of L_t around $x \in I_t$. Define $L_\delta^{k,p}(D_{x,t})$, $\hat{L}_\delta^{k,p}(D_{x,t})$ as in (20) using $\tilde{r}_t = \sigma_{x,t}^* r_t$, $\tilde{\Phi}_t$, \tilde{h}_t^{app} , the Euclidean metric on $D_{x,t}$ (which is $\sigma_{x,t}^* g_t$), and $\tilde{A}_t = D(\tilde{\partial}_E, \tilde{h}_t^{\text{app}})$. For $w \in \Omega^q(\text{End } E)$ we have

$$\|\sigma_{x,t}^* w\|_{L_\delta^{k,p}(D_{x,t})} = 2 \|w\|_{L_\delta^{k,p}(\tilde{B}_{x,t})}. \quad (25)$$

Let $\tilde{L}_t(\gamma) = \Delta_{\tilde{A}_t} \gamma - i \star M_{\tilde{\Phi}_t} \gamma$, then $\tilde{L}_t(\sigma^* \gamma) = \sigma^*(L_t \gamma)$. The previous analysis near $x \in I_u$ yields the following.

Corollary 4.8. *On $\tilde{B}_{x,t}$ for $x \in I_t$, $L_t : L_{-2+\delta}^{2,2}(\text{isu}(E)) \rightarrow L_{-2+\delta}^2(\text{isu}(E))$ is an isomorphism with zero Dirichlet condition on the boundary for small weights $\delta > 0$. Moreover, the norm of L_t^{-1} is uniformly bounded in t .*

Proof. Let $v \in L_{-2+\delta}^2(\text{isu}(E))(\tilde{B}_{x,t})$ then $\tilde{v} = \sigma_{x,t}^* v \in L_{-2+\delta}^2(\text{isu}(E))(D_{x,t})$, and by Proposition 4.7 with m_x and $j(x)$ both replaced by $2(m_x - 1)$, there exists $\tilde{u} \in L_{-2+\delta}^{2,2}(\text{isu}(E))(D_{x,t})$ such that $\tilde{L}_t \tilde{u} = \tilde{v}$ and $\tilde{u} = 0$ on $\partial D_{x,t}$. Note that \tilde{L}_t is \mathbb{Z}_2 -equivariant, meaning that $\iota^*(\tilde{L}_t(\tilde{u})) = \tilde{L}_t(\iota^* \tilde{u})$ for the involution $\iota : \xi_{x,t} \mapsto -\xi_{x,t}$. Then $\tilde{L}_t(\iota^* \tilde{u} - \tilde{u}) = \iota^* \tilde{v} - \tilde{v} = 0$, and $\iota^* \tilde{u} = \tilde{u}$ since \tilde{L}_t is injective. \tilde{u} descends to $\tilde{B}_{x,t}$ to give $u \in L_{-2+\delta}^{2,2}(\text{isu}(E))(\tilde{B}_{x,t})$ with $L_t u = v$ and $u = 0$ on $\partial \tilde{B}_{x,t}$. If $L_t u = 0$, then $\tilde{L}_t(\sigma_{x,t}^* u) = 0$ and $\sigma_{x,t}^* u = 0$ which implies that $u = 0$. Therefore L_t is an isomorphism. The uniform boundedness of L_t^{-1} follows from that of \tilde{L}_t^{-1} and (25). \square

The local behavior of L_t around $x \in T$ has been studied in [FMSW22], and we have the following result.

Proposition 4.9. *On $\hat{B}_{x,t}$, for $x \in T$, $L_t : L_{-2+\delta}^{2,2}(\text{isu}(E)) \rightarrow L_{-2+\delta}^2(\text{isu}(E))$ is an isomorphism with zero Dirichlet condition on the boundary for small weights $\delta > 0$. Moreover, the norm of L_t^{-1} is uniformly bounded in t .*

Proof. Consider $\tau : \hat{B}_{x,t} \rightarrow B_0(1) := \{|\tilde{z}| < 1\}$, $\zeta_{x,t} \mapsto \lambda_x(t)^2 \zeta_{x,t} := \tilde{z}$. Let $\tilde{\Phi}_t = (\tau^{-1})^* \Phi_t$, and $\tilde{h}_t^{\text{app}} = (\tau^{-1})^* \hat{h}_t^{\text{app}}$. Then by (11) and (16),

$$\tilde{\Phi}_t = \begin{pmatrix} 0 & \frac{1}{\lambda_x(t)} \\ \frac{\lambda_x(t)}{\tilde{z}} & 0 \end{pmatrix} d\tilde{z}, \quad \tilde{h}_t^{\text{app}} = \begin{pmatrix} \frac{1}{\lambda_x(t)} \tilde{r}^{1/2} e^{\psi_{2,x}(8\tilde{r}^{1/2})} & 0 \\ 0 & \frac{1}{\lambda_x(t)^3} \tilde{r}^{3/2} e^{-\psi_{2,x}(8\tilde{r}^{1/2})} \end{pmatrix},$$

where $\tilde{r} = |\tilde{z}|$. By the gauge transformation $g = \begin{pmatrix} \lambda_x(t)^{-1/2} & 0 \\ 0 & \lambda_x(t)^{1/2} \end{pmatrix}$, we have

$$\tilde{\Phi}_t = \begin{pmatrix} 0 & 1 \\ \frac{1}{\tilde{z}} & 0 \end{pmatrix} d\tilde{z}, \quad \tilde{h}_t^{\text{app}} = \frac{1}{\lambda_x(t)^2} \begin{pmatrix} \tilde{r}^{1/2} e^{\psi_{2,x}(8\tilde{r}^{1/2})} & 0 \\ 0 & \tilde{r}^{3/2} e^{-\psi_{2,x}(8\tilde{r}^{1/2})} \end{pmatrix}.$$

Then in this trivialization, $\tilde{\Phi}_t$ and $\lambda_x(t)^2 \tilde{h}_t^{\text{app}}$ are independent of t , which together with the Euclidean metric g_e on $B_0(1)$ define an operator \tilde{L} as in (18). The indicial root analysis as in [FMSW22, Sec. 4.2, Lem. 5.8] implies that $\tilde{L} : L_{-2+\delta}^{2,2}(\text{isu}(E)) \rightarrow L_{-2+\delta}^2(\text{isu}(E))$ (these spaces are defined on $B_0(1)$ as in (20) using g_e , $\tilde{\Phi}_t$, $\lambda_x(t)^2 \tilde{h}_t^{\text{app}}$) is an isomorphism for $\delta > 0$ small. Therefore L_t is also an isomorphism, and the uniform boundedness of L_t^{-1} follows in the same way as in Proposition 4.7 for the diagonal part. \square

Let $U_{\text{ext}} := C \setminus (\bigcup_{x \in I} \frac{1}{2} \tilde{B}_{x,t} \cup \bigcup_{x \in T} \frac{1}{2} \hat{B}_{x,t})$, where $\frac{1}{2} \tilde{B}_{x,t} = \{|\zeta_{x,t}| < \kappa t^{-1/(m_x-1)}/2\}$ for $x \in I$, $\frac{1}{2} \hat{B}_{x,t} = \{|z_x| < \kappa t^{-1/j(x)}/2\}$ for $x \in I_u$, and $\frac{1}{2} \hat{B}_{x,t} = \{|\zeta_{x,t}| < \lambda_x(t)^{-2}/2\}$ for $x \in T$. In U_{ext} , $c^{-1} \leq r_t \leq 1$, which is immaterial in the analysis, so we omit the subscript of $L_{-2+\delta}^{k,2}$.

Proposition 4.10. *On U_{ext} , $L_t : L^{2,2}(\text{isu}(E)) \rightarrow L^2(\text{isu}(E))$ is an isomorphism when imposing the zero Dirichlet boundary condition. The norm of L_t^{-1} is bounded by ct^4 .*

Proof. Invertibility is standard (see [MSWW16, Sec. 5.2]). Let $u \in L^{2,2}(\text{isu}(E))$ be vanishing on ∂U_{ext} , then

$$\|\nabla_{A_t} u\|_{L^2}^2 + \|[\Psi_t, u]\|_{L^2}^2 = \langle D_t^* D_t u, u \rangle_{L^2} \leq \|L_t u\|_{L^2} \|u\|_{L^2}.$$

Let $\check{U}_{\text{ext}} = \{|w| < 2\kappa^{-1} t^{1/(m_0+N-4)}\}$, where $w = 1/z$. Then \check{U}_{ext} is a disk centered at ∞ and contains U_{ext} . Let L_e^2 be the L^2 norm on \check{U}_{ext} defined using the Euclidean metric g_e on this disk. By Kato's inequality and Poincaré inequality, $\|u\|_{L_e^2} \leq ct^{1/(m_0+N-4)} \|\nabla_{A_t} u\|_{L_e^2} \leq ct^{1/2} \|\nabla_{A_t} u\|_{L_e^2}$. By the Definition 4.3 of g_t , we have $c^{-1} t^{-1} g_t \leq g_e \leq ct^{4/(m_0+N-4)} g_t \leq ct^2 g_t$ on U_{ext} . Therefore,

$$\|u\|_{L^2}^2 \leq ct \|u\|_{L_e^2}^2 \leq ct^2 \|\nabla_{A_t} u\|_{L_e^2}^2 = ct^2 \|\nabla_{A_t} u\|_{L^2}^2,$$

where the last equality follows from the conformal invariance of the L^2 norm of a 1-form. Then we have

$$\begin{aligned} \|u\|_{L^2}^2 &\leq ct^2 \|L_t u\|_{L^2} \|u\|_{L^2} \Rightarrow \|u\|_{L^2} \leq ct^2 \|L_t u\|_{L^2}, \\ \|\nabla_{A_t} u\|_{L^2} &\leq ct \|L_t u\|_{L^2}, \quad \|\Psi_t \otimes u\|_{L^2} \leq ct \|L_t u\|_{L^2}. \end{aligned}$$

We deduce that

$$\begin{aligned} \|(\Psi_t \otimes \cdot)^2 u\|_{L^2} &\leq c \sup |\Phi_t| \|\Psi_t \otimes u\|_{L^2} \leq ct \sup |\Phi_t| \|L_t u\|_{L^2} \\ \|\Psi_t \otimes \nabla_{A_t} u\|_{L^2} &\leq c \sup |\Phi_t| \|\nabla_{A_t} u\|_{L^2} \leq ct \sup |\Phi_t| \|L_t u\|_{L^2}. \end{aligned}$$

By Lemma 4.1 (i), and the conformal change formula for \mathcal{R} ,

$$\begin{aligned} \|\nabla_{A_t}^2 u\|_{L^2} &\leq c (\|D_t \nabla_{A_t} u\|_{L^2} + \|D_t^* \nabla_{A_t} u\|_{L^2} + t^3 \|\nabla_{A_t} u\|_{L^2} + \|\mathcal{F}(\nabla_{A_t} u)\|_{L^2}) \\ &\leq c (t \sup |\Phi_t| + t^4 + t \max \sup |E_{x(j),t}|) \|L_t u\|_{L^2}, \end{aligned}$$

where the error terms $E_{x(j),t}$ decay exponentially in t . In $U_{\text{ext}} \setminus (\bigcup_{x(j)} \tilde{B}_{x(j),t})$, Φ_t can be diagonalized as $\sqrt{-\nu_t(z)} \sigma_3 dz$, and h_t^{app} is also diagonal so that (recall that the norm $|dz|$ is taken with respect to g_t)

$$|\Phi_t|^2 = 2|\nu_t(z)| |dz|^2 \leq ct^2.$$

In $\tilde{B}_{x,j,t}$, Φ_t and h_t^{app} are given by (10) and (15) in some holomorphic frame, so we have

$$\begin{aligned} |\Phi_t|^2 &= 2\lambda_{x,j}(t)^2 r \cosh(2l_{x,j,t}(r) \chi(r \kappa^{-1} t^{1/j(x)})) |d\zeta_{x,j,t}|^2 \\ &\leq c\lambda_{x,j}(t)^{4/3} \rho^{2/3} \cosh(2\psi_1(\rho) \chi) |d\zeta_{x,j,t}|^2 \leq ct^2, \end{aligned}$$

where $r = |\zeta_{x,j,t}|$, $\rho = 8\lambda_{x,j}(t)r^{3/2}/3$. Similarly, in $\tilde{B}_{x,t} \cap U_{\text{ext}}$, Φ_t and h_t^{app} are given by (11) and (16) in some holomorphic frame, then

$$\begin{aligned} |\Phi_t|^2 &= 2\lambda_x(t)^2 r \cosh(2m_{x,t}(r)\chi(r\kappa^{-1})) |d\zeta_{x,t}|^2 \\ &\leq c\rho^2 \cosh(2\psi_{2,x}(\rho)\chi) |d\zeta_{x,t}|^2 \leq ct, \end{aligned}$$

where $r = |\zeta_{x,t}|$, $\rho = 8\lambda_x(t)r^{1/2}$. Therefore, $\|\nabla_{A_t}^2 u\|_{L^2} \leq ct^4 \|L_t u\|_{L^2}$. Similarly, applying the Weitzenböck formula for $w = [\Psi_t, u]$ gives $\|\nabla_{A_t}[\Psi_t, u]\|_{L^2} \leq ct^4 \|L_t u\|_{L^2}$. Combining all the estimates above, $\|u\|_{L^{2,2}(U_{\text{ext}})} \leq ct^4 \|L_t u\|_{L^2}$. \square

Finally, we consider the global behavior of L_t .

Lemma 4.11. $L_t : L_{-2+\delta}^{2,2}(\text{isu}(E)) \rightarrow L_{-2+\delta}^2(\text{isu}(E))$ is an isomorphism for small $\delta > 0$.

Proof. By [BB04, Lemma 5.1], L_t is Fredholm of index zero (the proof still work for $x \in I_t$ by considering the local lifted problem as in Corollary 4.8 and for $x \in T$ by [FMSW22, Lem. 5.8]). For $u \in \ker L_t$, near $x \in I_u \cup T$ we have $|D_t u| = O(r^{\delta-1})$ ($r = |\zeta_{x,t}|$ for $x \in T$, and $r = |z_x|$ for $x \in I_u$) by the proof of [BB04, Lem. 4.6]. The boundary term in the integration by parts is $\lim_{\epsilon \rightarrow 0^+} \int_{r=\epsilon} \langle \partial_r u, u \rangle r d\theta$ [FMSW22, Lem. 5.5] which vanishes for $x \in I_u \cup T$ since $\langle [\Phi_t, u], u \rangle = 0$ and $\langle \partial_r u, u \rangle = O(r^{\delta-1})$, and similarly vanishes for $x \in I_t$ by considering the double lifting. Therefore

$$\langle L_t u, u \rangle_{L^2} = \|d_{A_t} u\|_{L^2}^2 + \|[\Psi_t, u]\|_{L^2}^2. \quad (26)$$

Now we have $d_{A_t} u = [\Phi_t, u] = 0$, which implies that $u \equiv 0$ as in the proof of [FMSW22, Lem. 5.6] since $Z_t \setminus (I_t \cup T) \neq \emptyset$. L_t has trivial kernel, and then it is an isomorphism. \square

Lemma 4.12. Let $u \in L_{-2+\delta}^{2,2}(\text{isu}(E))$, then for t large enough,

$$\|u\|_{L_{-\delta}^2}^2 \leq ct^2 (\|\nabla_{A_t} u\|_{L^2}^2 + \|[\Phi_t, u]\|_{L^2}^2). \quad (27)$$

Proof. The idea is to use $[\Phi_t, u]$ to control u around a zero of $\tilde{v}_t(z)$ near each $x \in I$ (we choose $z_{x,0}(t)$, which has asymptotics given in Lemma 3.1), and use $\nabla_{A_t} u$ to control u elsewhere. The inequality is obtained by gluing these estimates together.

For t large, $x \in I_u$, define $W_x := \{|\zeta_{x,0,t}| < t^{-(2m_x+1)/3j(x)}\} \subset \frac{1}{2}\tilde{B}_{x,0,t}$, a disk centered at $z_{x,0}(t)$. Then $\chi(r\kappa^{-1}t^{1/j(x)}) = 1$ on W_x , where $r = |\zeta_{x,0,t}|$, and

$$\rho = 8\lambda_{x,0}(t)r^{3/2}/3 \leq c \text{ on } W_x.$$

Recall that Φ_t is given by (10) in some holomorphic frame over W_x . Since u is a section of $\text{isu}(E)$, it has form

$$u = \begin{pmatrix} p & \bar{s}r^{-1}e^{-2l_{x,0,t}(r)} \\ s & -p \end{pmatrix},$$

where p is real valued. Φ_t acts on u as

$$[\Phi_t, u] = \lambda_{x,0}(t) \begin{pmatrix} s - \bar{s}\zeta_{x,0,t}r^{-1}e^{-2l_{x,0,t}(r)} & -2p \\ 2\zeta_{x,0,t}p & -s + \bar{s}\zeta_{x,0,t}r^{-1}e^{-2l_{x,0,t}(r)} \end{pmatrix} d\zeta_{x,0,t}.$$

Then on W_x we have

$$\begin{aligned}
|u|^2 &= 2p^2 + 2|s|^2 r^{-1} e^{-2l_{x,0,t}(r)} \leq 2p^2 + c|s|^2 t^{(2m_x+1)/3j(x)} \rho^{-2/3} e^{-2\psi_1(\rho)} \\
&\leq 2p^2 + ct^{(2m_x+1)/3j(x)} |s|^2, \\
\|\Phi_t, u\|^2 &= \lambda_{x,0}(t)^2 \left(8p^2 r \cosh(2l_{x,0,t}(r)) + 2|s - \bar{s}\zeta_{x,0,t} r^{-1} e^{-2l_{x,0,t}(r)}|^2 \right) \\
&\geq c^{-1} t^{(2m_x+1)/j(x)} \left(p^2 t^{-(2m_x+1)/3j(x)} \rho^{2/3} \cosh(2\psi_1(\rho)) + |s|^2 (1 - e^{-2\psi_1(\rho)}) \right) \\
&\geq c^{-1} t^{(4m_x+2)/3j(x)} |u|^2.
\end{aligned} \tag{28}$$

Consider $\chi_x(r) := \chi(t^{(2m_x+1)/3j(x)} r)$, then $\chi_x = 0$ in $C \setminus W_x$ and $\chi_x = 1$ in $\hat{W}_x := \frac{1}{2}W_x$. Let $r_{t,x}$ be the function obtained by extending $r_t|_{\hat{B}_{x,t}}$ by 1, and $\hat{g}_t = \hat{f}_t g_C$ where \hat{f}_t is the extension of $f_t|_{C \setminus \cup_{y \in I_t} B_y}$ by 1. We claim that for all $v \in L_{-2+\delta}^{2,2}$ with v vanishing in \hat{W}_x ,

$$\|v\|_{L_{-\delta,x}^2}^2 \leq ct^{(6+(4m_x-4)\delta)/3j(x)} \|\nabla_{A_t} v\|_{L_{-\delta-1,x}^2}^2, \tag{29}$$

where the subscripts indicate the norms are defined using $r_{t,x}, \hat{g}_t$ instead of r_t, g_t . By Kato's inequality, we may assume that v is a function, and prove the inequality with $\nabla_{A_t} v$ replaced by dv . First consider the biholomorphism $\tau_x : C \rightarrow C_1 = \mathbb{C}P^1$, defined by $z_x \mapsto c_x t^{1/j(x)} z_x$, where c_x is the constant satisfying $\lim_{t \rightarrow \infty} \tau_x(z_{x,0}(t)) = x + 1$. Define weighted spaces on C_1 using the round metric on C_1 and the weight function r_x given by the geodesic distance from x , then r_x is uniformly equivalent to $(\tau_x^{-1})^* r_{t,x}$, and $c^{-1} t^{-2/j(x)} (\tau_x^{-1})^* \hat{g}_t \leq g_{C_1}$. Then for $v_1 = (\tau_x^{-1})^* v$ we have

$$\begin{aligned}
\|v\|_{L_{-\delta,x}^2}^2 &= \int_C r_{t,x}^{2\delta-2} |v|^2 d\text{vol}_{\hat{g}_t} \leq c \int_{C_1} r_x^{2\delta-2} |v_1|^2 (\tau_x^{-1})^*(d\text{vol}_{\hat{g}_t}) \\
&\leq ct^{2/j(x)} \int_{C_1} r_x^{2\delta-2} |v_1|^2 d\text{vol}_{C_1} = ct^{2/j(x)} \|v_1\|_{L_{-\delta}^2(C_1)}^2 \\
\|dv_1\|_{L_{-\delta-1}^2(C_1)}^2 &= \int_{C_1} r_x^{2\delta} |dv_1|_{g_{C_1}}^2 d\text{vol}_{C_1} \leq c \int_C r_{t,x}^{2\delta} |dv|_{\hat{g}_t}^2 d\text{vol}_{\hat{g}_t} = c \|dv\|_{L_{-\delta-1,x}^2}^2.
\end{aligned}$$

(29) will follow from

$$\|v_1\|_{L_{-\delta}^2(C_1)}^2 \leq ct^{(4m_x-4)\delta/3j(x)} \|dv_1\|_{L_{-\delta-1}^2(C_1)}^2. \tag{30}$$

Suppose C_1 is embedded in \mathbb{R}^3 so that $x = (0, 1, 0)$ and $x + 1 = (0, 0, 1)$. Consider the stereographic projection $\hat{\tau}_x : C_1 \setminus \{x + 1\} \rightarrow \mathbb{R}^2$, which sends $x + 1$ to ∞ and x to $(0, 1)$. Note that as $t \rightarrow \infty$, $\tau_x(z_{x,0}(t)) \rightarrow x + 1$ and the diameter of $\tau_x(\hat{W}_x)$ is $O(t^{-(2m_x-2)/3j(x)})$, then for some constant κ_x , we have $\hat{\tau}_x^{-1}(\mathbb{R}^2 \setminus C_2) \subset \tau_x(\hat{W}_x)$, where $C_2 := B_{(0,1)}(\kappa_x t^{(2m_x-2)/3j(x)})$. Define weighted spaces on C_2 using the Euclidean metric and the weight function \hat{r} given by the Euclidean distance from $(0, 1)$. Then $v_2 = (\hat{\tau}_x^{-1})^* v_1$ vanishes on ∂C_2 , $c^{-1} (\hat{\tau}_x^{-1})^* r_x \leq \hat{r} \leq ct^{(2m_x-2)/3j(x)} (\hat{\tau}_x^{-1})^* r_x$, and $c^{-1} (\hat{\tau}_x^{-1})^* g_{C_1} \leq g_{C_2}$ on C_2 . We deduce that

$$\begin{aligned}
\|v_1\|_{L_{-\delta}^2(C_1)}^2 &= \int_{C_1} r_x^{2\delta-2} |v_1|^2 d\text{vol}_{C_1} \leq c \int_{C_2} \hat{r}^{2\delta-2} |v_2|^2 d\text{vol}_{C_2} = c \|v_2\|_{L_{-\delta}^2(C_2)}^2 \\
\|dv_2\|_{L_{-\delta-1}^2(C_2)}^2 &= \int_{C_2} \hat{r}^{2\delta} |dv_2|_{g_{C_2}}^2 d\text{vol}_{C_2} \leq ct^{(4m_x-4)\delta/3j(x)} \|dv_1\|_{L_{-\delta-1}^2(C_1)}^2.
\end{aligned}$$

An analogue of Lemma 4.5 (ii) implies that

$$\|v_2\|_{L^2_{-\delta}(C_2)}^2 \leq c \|dv_2\|_{L^2_{-\delta-1}(C_2)}^2,$$

since v_2 vanishes on ∂C_2 . Therefore

$$\|v_1\|_{L^2_{-\delta}(C_1)}^2 \leq c \|v_2\|_{L^2_{-\delta}(C_2)}^2 \leq c \|dv_2\|_{L^2_{-\delta-1}(C_2)}^2 \leq ct^{(4m_x-4)\delta/3j(x)} \|dv_1\|_{L^2_{-\delta-1}(C_1)}^2,$$

verifying (30), and (29) holds.

For t large, $x \in I_t$, $W_x := \{|\zeta_{x,0,t}| < t^{-2m_x/3j(x)}\} \subset \frac{1}{2}\tilde{B}_{x,0,t}$, so that $\chi(r\kappa^{-1}t^{1/j(x)}) = 1$ and $\rho \leq c$ on W_x , where $r = |\zeta_{x,0,t}|$ and $\rho = 8\lambda_{x,0}(t)r^{3/2}/3$. Similar to (28),

$$|[\Phi_t, u]|^2 \geq c^{-1}t^{4m_x/3j(x)}|u|^2. \quad (31)$$

Consider $\chi_x(r) := \chi(t^{2m_x/3j(x)}r)$, then $\chi_x = 0$ in $C \setminus W_x$ and $\hat{\chi} = 1$ in $\hat{W}_x := \frac{1}{2}W_x$. Let $r_{t,x}$ be the function on C which is the extension of $r_t|_{\tilde{B}_{x,t}}$ by 1, and $g_{t,x} = f_{t,x}g_C$ where $f_{t,x}$ is the extension of $f_t|_{C \setminus \cup_{y \in I_t \setminus \{x\}} B_y}$ by 1. We claim that for all $v \in L^2_{-2+\delta}$ with v vanishing in \hat{W}_x ,

$$\|v\|_{L^2_{-\delta,x}}^2 \leq ct^{(6+(2m_x-3)\delta)/3j(x)} \|\nabla_{A_t} v\|_{L^2_{-\delta-1,x}}^2, \quad (32)$$

where the subscripts indicate the norms are defined using $r_{t,x}, g_{t,x}$ instead of r_t, g_t . As above we prove the inequality for v being a function. Consider the biholomorphism $\tau_x : C \rightarrow C_1$, defined by $z_x \mapsto c_x t^{1/j(x)} z_x$, where c_x is the constant chosen so that $\lim_{t \rightarrow \infty} \tau_x(z_{x,0}(t)) = x + 1$. Define weighted spaces on C_1 using the singular metric $g_{C_1} = r_x^{-1}g_{\text{FS}(C_1)}$ where $g_{\text{FS}(C_1)}$ is the Fubini-Study metric on C_1 , and the weight function $r_x^{1/2}$ where r_x is the geodesic distance from x . Note that $r_x^{1/2}$ is uniformly equivalent to $(\tau_x^{-1})^*r_{t,x}$, and $c^{-1}t^{-2/j(x)}(\tau_x^{-1})^*g_{t,x} \leq g_{C_1}$. Then for $v_1 = (\tau^{-1})^*v$ we have

$$\|v\|_{L^2_{-\delta,x}}^2 \leq ct^{2/j(x)} \|v_1\|_{L^2_{-\delta}(C_1)}^2, \quad \|dv_1\|_{L^2_{-\delta-1}(C_1)}^2 \leq c \|dv\|_{L^2_{-\delta-1,x}}^2.$$

(32) follows once we prove

$$\|v_1\|_{L^2_{-\delta}(C_1)}^2 \leq ct^{(2m_x-3)\delta/3j(x)} \|dv_1\|_{L^2_{-\delta-1}(C_1)}^2. \quad (33)$$

Suppose C_1 is embedded in \mathbb{R}^3 as above and consider the stereographic projection $\hat{\tau}_x : C_1 \setminus \{x+1\} \rightarrow \mathbb{R}^2$, which sends $x+1$ to ∞ and x to $(0,1)$. Note that the diameter of $\tau_x(\hat{W}_x)$ is $O(t^{-(2m_x-3)/3j(x)})$, then for some constant κ_x , we have $\hat{\tau}_x^{-1}(\mathbb{R}^2 \setminus C_2) \subset \tau_x(W_2)$, where $C_2 := B_{(0,1)}(\kappa_x t^{(2m_x-3)/3j(x)})$. Define weighted spaces on C_2 using the singular metric $g_{C_2} = \hat{r}^{-1}g_{e(C_2)}$ where $g_{e(C_2)}$ is the Euclidean metric, and the weight function $\hat{r}^{1/2}$ where \hat{r} is the Euclidean distance from $(0,1)$. Then $v_2 = (\hat{\tau}_x^{-1})^*v_1$ vanishes on ∂C_2 , $c^{-1}(\hat{\tau}_x^{-1})^*r_x \leq \hat{r} \leq ct^{(2m_x-3)/3j(x)}(\hat{\tau}_x^{-1})^*r_x$, and $c^{-1}(\hat{\tau}_x^{-1})^*g_{C_1} \leq g_{C_2}$ on C_2 . We have

$$\|v_1\|_{L^2_{-\delta}(C_1)}^2 \leq c \|v_2\|_{L^2_{-\delta}(C_2)}^2, \quad \|dv_2\|_{L^2_{-\delta-1}(C_2)}^2 \leq ct^{(2m_x-3)\delta/3j(x)} \|dv_1\|_{L^2_{-\delta-1}(C_1)}^2.$$

Let $\tilde{L}_\delta^{k,p}(C_2)$ be defined using the Euclidean metric $g_{e(C_2)}$ and the weight function \hat{r} . An analogue of Lemma 4.5 (ii) implies that

$$\|v_2\|_{\tilde{L}^2_{-\delta/2}(C_2)}^2 \leq c \|dv_2\|_{\tilde{L}^2_{-\delta/2-1}(C_2)}^2,$$

since v_2 vanishes on ∂C_2 . This is equivalent to $\|v_2\|_{L^2_{-\delta}(C_2)}^2 \leq c \|dv_2\|_{L^2_{-\delta-1}(C_2)}^2$. Therefore (33) and (32) are verified.

For $x \in T$, let $\hat{\chi}_x(r) := \chi(\lambda_x(t)^2 r)$, where $r = |\zeta_{x,t}|$. In $\hat{B}_{x,t}$, similar to (28), we have

$$|[\Phi_t, u]|^2 \geq c^{-1} t^2 |u|^2. \quad (34)$$

Let v be vanishing on $\partial \hat{B}_{x,t}$, then as before,

$$\|v\|_{L^2_{-\delta}(\hat{B}_{x,t})}^2 \leq ct^{-2} \|\nabla_{A_t} u\|_{L^2_{-\delta-1}(\hat{B}_{x,t})}^2. \quad (35)$$

Applying (29), (32) to $v = (1 - \chi_x)u$ and (35) to $v = \hat{\chi}_x u$, and using (28), (31), (34),

$$\begin{aligned} \|u\|_{L^2_{-\delta}}^2 &\leq c \sum_{x \in I} \|u\|_{L^2_{-\delta,x}}^2 + c \sum_{x \in T} \|\hat{\chi}_x u\|_{L^2_{-\delta}(\hat{B}_{x,t})}^2 \\ &\leq c \sum_{x \in I} \|\chi_x u\|_{L^2(W_x)}^2 + c \sum_{x \in I} \|(1 - \chi_x)u\|_{L^2_{-\delta,x}}^2 + c \sum_{x \in T} \|\hat{\chi}_x u\|_{L^2_{-\delta}(\hat{B}_{x,t})}^2 \\ &\leq c \left(\sum_{x \in I_u} t^{-(4m_x+2)/3j(x)} + \sum_{x \in I_t} t^{-4m_x/3j(x)} \right) \|[\Phi_t, u]\|_{L^2}^2 \\ &\quad + c \sum_{x \in I_u} t^{(6+(4m_x-4)\delta)/3j(x)} \|\nabla_{A_t}(1 - \chi_x)u\|_{L^2_{-\delta-1,x}}^2 \\ &\quad + c \sum_{x \in I_t} t^{(6+(2m_x-3)\delta)/3j(x)} \|\nabla_{A_t}(1 - \chi_x)u\|_{L^2_{-\delta-1,x}}^2 + ct^{-2} \sum_{x \in T} \|\nabla_{A_t} \hat{\chi}_x u\|_{L^2_{-\delta-1,x}}^2 \\ &\leq ct^2 (\|\nabla_{A_t} u\|_{L^2}^2 + \|[\Phi_t, u]\|_{L^2}^2), \end{aligned}$$

where in the last inequality we used $\delta < 1$, $(4m_x + 2)/3j(x) < 4/3 + 2/3 = 2$, and the L^2 norm is uniformly stronger than various $L^2_{-\delta-1}$ norms. \square

Proposition 4.13. *The norm of $L_t^{-1} : L^2_{-2+\delta}(i\mathfrak{su}(E)) \rightarrow L^{2,2}_{-2+\delta}(i\mathfrak{su}(E))$ is bounded by ct^8 .*

Proof. Let $v \in C_0^\infty(C \setminus S)$, then $u = L_t^{-1}(v) \in L^2_{-2+\delta}$ and the boundary terms leading to the integration by parts vanish as above, so (26) holds. By the density of $C_0^\infty(C \setminus S)$ in $L^2_{-2+\delta}$ and the continuity of $L_t^{-1}, d_{A_t}, [\Psi_t, \cdot]$, (26) holds for all $u \in L^2_{-2+\delta}$. Using (26) and (27),

$$\begin{aligned} \|u\|_{L^2_{-\delta}}^2 &\leq ct^2 (\|\nabla_{A_t} u\|_{L^2}^2 + \|[\Phi_t, u]\|_{L^2}^2) \leq ct^2 \langle L_t u, u \rangle_{L^2} \\ &\leq ct^2 \|u\|_{L^2_{-\delta}} \|L_t u\|_{L^2_{-2+\delta}}, \\ \|u\|_{L^2_{-2+\delta}} &\leq \|u\|_{L^2_{-\delta}} \leq ct^2 \|L_t u\|_{L^2_{-2+\delta}}, \\ \|\nabla_{A_t} u\|_{L^2_{-2+\delta}} &\leq \|\nabla_{A_t} u\|_{L^2} \leq ct \|L_t u\|_{L^2_{-2+\delta}}, \\ \|[\Phi_t, u]\|_{L^2} &\leq ct \|L_t u\|_{L^2_{-2+\delta}}. \end{aligned}$$

For $x \in I$, let $\hat{\chi}_x$ be the function supported in $\tilde{B}_{x,t}$ where it satisfies $\hat{\chi}_x(r) = \chi(\kappa^{-1} t^{1/j(x)} r)$ ($r = |z_x|$ for $x \in I_u$ and $r = |\zeta_{x,t}|$ for $x \in I_t$). Recall that we have defined $\hat{\chi}_x$ for $x \in T$ in a

similar way. Using Propositions 4.7, 4.8, 4.9, 4.10 we have

$$\|u\|_{L_{-2+\delta}^{2,2}} \leq \sum_{x \in S} \|\hat{\chi}_x u\|_{L_{-2+\delta}^{2,2}} + \|(1 - \sum_{x \in S} \hat{\chi}_x)u\|_{L^{2,2}} \quad (36)$$

$$\begin{aligned} &\leq c \sum_{x \in S} \|L_t \hat{\chi}_x u\|_{L_{-2+\delta}^2} + ct^4 \|L_t(1 - \sum_{x \in S} \hat{\chi}_x)u\|_{L^2} \\ &\leq ct^4 \left(\|L_t u\|_{L_{-2+\delta}^2} + t^2 \|u\|_{L_{-\delta}^2} + t \|\nabla_{A_t} u\|_{L^2} + t \|[\Phi_t, u]\|_{L^2} \right) \\ &\leq ct^8 \|L_t u\|_{L_{-2+\delta}^2}. \end{aligned} \quad (37)$$

□

5. DEFORMING TO A SOLUTION

Now we find a solution to Hitchin's equation by deforming the approximate solution (A_t, Φ_t) via a complex gauge transformation $\exp(\gamma)$ for $\gamma \in \Omega^0(i\mathfrak{su}(E))$, which satisfies

$$F_t(\gamma) = F_{A_t^{\exp(\gamma)}} + [e^{-\gamma} \Phi_t e^\gamma, e^\gamma \Phi_t^* e^{-\gamma}] =: F_{A_t} + [\Phi_t, \Phi_t^*] + L_t \gamma + Q_t \gamma = 0,$$

where by abuse of notation we still denote by L_t the linear operator in the previous section composed with the isomorphism $i\star$. We will analyze the nonlinear term Q_t in a similar way as that in [MSWW16]. Denote

$$\begin{aligned} A_t^{\exp \gamma} &= A_t + (\bar{\partial}_{A_t} - \partial_{A_t})\gamma + R_{A_t}(\gamma), \\ \Phi_t^{\exp \gamma} &= \Phi_t + [\Phi_t, \gamma] + R_{\Phi_t}(\gamma), \\ R_{A_t}(\gamma) &= e^{-\gamma}(\bar{\partial}_{A_t} e^\gamma) - (\partial_{A_t} e^\gamma) e^{-\gamma} - (\bar{\partial}_{A_t} - \partial_{A_t})\gamma, \\ R_{\Phi_t}(\gamma) &= e^{-\gamma} \Phi_t e^\gamma - [\Phi_t, \gamma] - \Phi_t. \end{aligned}$$

Then we have

$$\begin{aligned} Q_t(\gamma) &= d_{A_t}(R_{A_t}(\gamma)) + [R_{\Phi_t}(\gamma), \Phi_t^*] + [\Phi_t, R_{\Phi_t}(\gamma)^*] \\ &\quad + \frac{1}{2}[(\bar{\partial}_{A_t} - \partial_{A_t})\gamma + R_{A_t}(\gamma), (\bar{\partial}_{A_t} - \partial_{A_t})\gamma + R_{A_t}(\gamma)] \\ &\quad + [[\Phi_t, \gamma] + R_{\Phi_t}(\gamma), ([\Phi_t, \gamma] + R_{\Phi_t}(\gamma))^*]. \end{aligned} \quad (38)$$

For $u \in \Omega^p(\text{End } E)$, define

$$\|u\|_{\tilde{L}_\delta^{k,q}} := \left(\sum_{j=0}^k \|\nabla_{A_t}^j u\|_{L_\delta^q}^q \right)^{1/q}.$$

Clearly $\|u\|_{\tilde{L}_\delta^{k,q}} \leq \|u\|_{L_\delta^{k,q}}$.

Lemma 5.1. $\tilde{L}_{-2+\delta}^{2,2}(\text{End } E)$ is an algebra, and $\tilde{L}_{-2+\delta}^{1,2}(\text{End } E)$ is a module over this algebra. Moreover,

$$\|uv\|_{\tilde{L}_{-2+\delta}^{2,2}} \leq ct^5 \|u\|_{\tilde{L}_{-2+\delta}^{2,2}} \|v\|_{\tilde{L}_{-2+\delta}^{2,2}}, \quad (39)$$

$$\|uv\|_{\tilde{L}_{-2+\delta}^{1,2}} \leq ct^{9/2} \|u\|_{\tilde{L}_{-2+\delta}^{2,2}} \|v\|_{\tilde{L}_{-2+\delta}^{1,2}}, \quad (40)$$

$$\|u\|_{L^\infty} \leq ct^3 \|u\|_{\tilde{L}_{-2+\delta}^{2,2}}. \quad (41)$$

Proof. Let τ_x and $\hat{\chi}_x$ be defined as in the previous section, and denote $u_x := \hat{\chi}_x u$, $u_{\text{ext}} := (1 - \sum_{x \in S} \hat{\chi}_x)u$. For $x \in I_u$, τ_x maps $\tilde{B}_{x,t}$ to $B_0(\kappa|c_x|) := \{|z| < \kappa|c_x|\}$. Denote the Euclidean metric on this disk by g_e , then $c^{-1}t^{2/j(x)}(\tau_x^{-1})^*g_t \leq g_e \leq ct^{2/j(x)}(\tau_x^{-1})^*g_t$. Define the Sobolev spaces on $B_0(\kappa|c_x|)$ using the weight function $r = |z|$ and the metric g_e . For w a p -form with values in $\text{End}E$, using the Sobolev embedding $L_{-2+\delta}^{1,2} \hookrightarrow L_{-1+\delta}^q$ on $B_0(\kappa|c_x|)$ for any fixed $q \in [2, \infty)$,

$$\begin{aligned} \|w_x\|_{L_{-1+\delta}^q} &\leq ct^{-2/qj(x)} \|(\tau_x^{-1})^*|w_x|\|_{L_{-1+\delta}^q(B_0(\kappa|c_x|))} \\ &\leq ct^{-2/qj(x)} \left(\|(\tau_x^{-1})^*|w_x|\|_{L_{-2+\delta}^2(B_0(\kappa|c_x|))} + \|(\tau_x^{-1})^*d|w_x|\|_{L_{-2+\delta}^2(B_0(\kappa|c_x|))} \right) \\ &\leq ct^{-2/qj(x)} \left(t^{1/j(x)} \|w_x\|_{L_{-2+\delta}^2} + \|\nabla_{A_t} w_x\|_{L_{-2+\delta}^2} \right) \leq ct^{(q-2)/qj(x)} \|w_x\|_{\tilde{L}_{-2+\delta}^{1,2}}. \end{aligned}$$

Similarly for $x \in T$, we have $\|w_x\|_{L_{-1+\delta}^q} \leq ct^{(q-2)/q} \|w_x\|_{\tilde{L}_{-2+\delta}^{1,2}}$. For $x \in I_t$, let $\check{L}_\delta^{k,m}(B_0(\kappa|c_x|))$ be defined using $r^{-1}g_e$ and the weight function $r^{1/2}$, then \check{L}_δ^q is equivalent to $L_{\delta/2}$ for functions on $B_0(\kappa|c_x|)$, and we have

$$\begin{aligned} \|w_x\|_{L_{-1+\delta}^q} &\leq ct^{-2/qj(x)} \|(\tau_x^{-1})^*|w_x|\|_{\check{L}_{-1+\delta}^q(B_0(\kappa|c_x|))} \leq ct^{-2/qj(x)} \|(\tau_x^{-1})^*|w_x|\|_{L_{(-1+\delta)/2}^q(B_0(\kappa|c_x|))} \\ &\leq ct^{-2/qj(x)} \left(\|(\tau_x^{-1})^*|w_x|\|_{L_{(-1+\delta)/2-1}^2(B_0(\kappa|c_x|))} + \|(\tau_x^{-1})^*d|w_x|\|_{L_{(-1+\delta)/2-1}^2(B_0(\kappa|c_x|))} \right) \\ &\leq ct^{-2/qj(x)} \left(\|(\tau_x^{-1})^*|w_x|\|_{\check{L}_{-2+\delta}^2(B_0(\kappa|c_x|))} + \|(\tau_x^{-1})^*d|w_x|\|_{\check{L}_{-2+\delta}^2(B_0(\kappa|c_x|))} \right) \\ &\leq ct^{-2/qj(x)} \left(t^{1/j(x)} \|w_x\|_{L_{-2+\delta}^2} + \|\nabla_{A_t} w_x\|_{L_{-2+\delta}^2} \right) \leq ct^{(q-2)/qj(x)} \|w_x\|_{\tilde{L}_{-2+\delta}^{1,2}}. \end{aligned}$$

On U_{ext} , $c^{-1}g_C \leq g_t \leq ctg_C$, and r_t is uniformly bounded, then

$$\begin{aligned} \|w_{\text{ext}}\|_{L^q} &\leq ct^{1/q} \|w_{\text{ext}}\|_{L^q(g_C)} \leq ct^{1/q} \left(\|w_{\text{ext}}\|_{L^2(g_C)} + \|\nabla_{A_t} w_{\text{ext}}\|_{L^2(g_C)} \right) \\ &\leq ct^{1/q} \left(t^{\max(p-1,0)/2} \|w_{\text{ext}}\|_{L^2} + t^{p/2} \|\nabla_{A_t} w_{\text{ext}}\|_{L^2} \right). \end{aligned} \quad (42)$$

Combining the above inequalities,

$$\|w\|_{L_{-1+\delta}^q} \leq \sum_{x \in S} \|w_x\|_{L_{-1+\delta}^q} + \|w_{\text{ext}}\|_{L_{-1+\delta}^q} \leq ct^{2+p/2} \|w\|_{\tilde{L}_{-2+\delta}^{1,2}}.$$

For $w_i \in \Omega^{p_i}(\text{End}E)$ ($i = 1, 2$),

$$\begin{aligned} \|w_1 w_2\|_{L_{-2+\delta}^2} &= \|r_t^{1-\delta} w_1 w_2\|_{L^2} \leq \|r_t^{1-2\delta} w_1 w_2\|_{L^2} \leq \|r_t^{1/2-\delta} w_1\|_{L^4} \|r_t^{1/2-\delta} w_2\|_{L^4} \\ &= \|w_1\|_{L_{-1+\delta}^4} \|w_2\|_{L_{-1+\delta}^4} \leq ct^{4+(p_1+p_2)/2} \|w_1\|_{\tilde{L}_{-2+\delta}^{1,2}} \|w_2\|_{\tilde{L}_{-2+\delta}^{1,2}}. \end{aligned} \quad (43)$$

For $2 < q < 2/(1-\delta)$, $x \in I_u$, using $L_{-1+\delta}^q \subset L^q$, $L^{1,q} \subset L^\infty$ on $B_0(\kappa|c_x|)$,

$$\begin{aligned} \|u_x\|_{L^\infty} &= \|(\tau_x^{-1})^*|u_x|\|_{L^\infty(B_0(\kappa|c_x|))} \\ &\leq c \left(\|(\tau_x^{-1})^*|u_x|\|_{L_{-1+\delta}^q(B_0(\kappa|c_x|))} + \|(\tau_x^{-1})^*d|u_x|\|_{L_{-1+\delta}^q(B_0(\kappa|c_x|))} \right) \\ &\leq c \left(t^{2/qj(x)} \|u_x\|_{L_{-1+\delta}^q} + t^{(2-q)/qj(x)} \|\nabla_{A_t} u_x\|_{L_{-1+\delta}^q} \right) \\ &\leq c \left(t^{2/qj(x)} t^{(q-2)/qj(x)} \|u_x\|_{\tilde{L}_{-2+\delta}^{1,2}} + \|\nabla_{A_t} u_x\|_{\tilde{L}_{-2+\delta}^{1,2}} \right) \leq ct^{1/j(x)} \|u\|_{\tilde{L}_{-2+\delta}^{2,2}}. \end{aligned}$$

Similarly, we have $\|u_x\|_{L^\infty} \leq ct\|u_x\|_{\tilde{L}_{-2+\delta}^{2,2}}$ for $x \in T$, and $\|u_x\|_{L^\infty} \leq ct^{1/j(x)}\|u_x\|_{\tilde{L}_{-2+\delta}^{2,2}}$ for $x \in I_t$. On U_{ext} , by (42),

$$\|u_{\text{ext}}\|_{L^\infty} \leq c \left(\|u_{\text{ext}}\|_{L^q(g_C)} + \|\nabla_{A_t} u_{\text{ext}}\|_{L^q(g_C)} \right) \leq ct^{1/2} \|u_{\text{ext}}\|_{\tilde{L}_{-2+\delta}^{2,2}}.$$

Therefore,

$$\|u\|_{L^\infty} \leq \sum_{x \in S} \|u_x\|_{L^\infty} + \|u_{\text{ext}}\|_{L^\infty} \leq ct^3 \|u\|_{\tilde{L}_{-2+\delta}^{2,2}}.$$

Then by (43),

$$\begin{aligned} \|\nabla_{A_t}(uv)\|_{L_{-2+\delta}^2} &\leq \|(\nabla_{A_t} u)v\|_{L_{-2+\delta}^2} + \|u\nabla_{A_t} v\|_{L_{-2+\delta}^2} \\ &\leq ct^{9/2} \|u\|_{\tilde{L}_{-2+\delta}^{2,2}} \|v\|_{\tilde{L}_{-2+\delta}^{1,2}} + \|u\|_{L^\infty} \|v\|_{\tilde{L}_{-2+\delta}^{1,2}} \leq ct^{9/2} \|u\|_{\tilde{L}_{-2+\delta}^{2,2}} \|v\|_{\tilde{L}_{-2+\delta}^{1,2}}. \end{aligned}$$

Finally we estimate

$$\begin{aligned} \|\nabla_{A_t}^2 uv\|_{L_{-2+\delta}^2} &\leq \|u\|_{L^\infty} \|\nabla_{A_t}^2 v\|_{L_{-2+\delta}^2} + ct^5 \|\nabla_{A_t} u\|_{\tilde{L}_{-2+\delta}^{1,2}} \|\nabla_{A_t} v\|_{\tilde{L}_{-2+\delta}^{1,2}} + \|v\|_{L^\infty} \|\nabla_{A_t}^2 u\|_{L_{-2+\delta}^2} \\ &\leq ct^5 \|u\|_{\tilde{L}_{-2+\delta}^{2,2}} \|v\|_{\tilde{L}_{-2+\delta}^{2,2}}. \end{aligned}$$

□

Theorem 5.2. *There exists t_1, c, c' such that for every $t \geq t_1$, there exists a unique h_t^{app} -Hermitian $\gamma_t \in L_{-2+\delta}^{2,2}(\text{isu}(E))$ with $\|\gamma_t\|_{L_{-2+\delta}^{2,2}} \leq c \exp(-c't^\sigma)$, such that $F_t(\gamma_t) = 0$. Hence h_t defined by $h_t(v, w) = h_t^{\text{app}}(e^{-\gamma_t} v, e^{-\gamma_t} w)$ is a harmonic metric associated to $(\bar{\partial}_E, \Phi_t)$ and is compatible with the parabolic structure at each $x \in S$.*

Proof. Let $0 < \rho \leq 1$ and $\gamma_1, \gamma_2 \in B_\rho := \{\gamma \in L_{-2+\delta}^{2,2}(\text{isu}(E)) : \|\gamma\|_{L_{-2+\delta}^{2,2}} < \rho\}$. We estimate each of the five terms in the difference $Q_t(\gamma_1) - Q_t(\gamma_2)$, with Q_t given by (38). First,

$$\begin{aligned} \|R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} &\leq \|(e^{-\gamma_1}(\bar{\partial}_{A_t} e^{\gamma_1}) - e^{-\gamma_2}(\bar{\partial}_{A_t} e^{\gamma_2}) - \bar{\partial}_{A_t}(\gamma_1 - \gamma_2))\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\quad + \|((\partial_{A_t} e^{\gamma_1})e^{-\gamma_1} - (\partial_{A_t} e^{\gamma_2})e^{-\gamma_2} + \partial_{A_t}(\gamma_1 - \gamma_2))\|_{\tilde{L}_{-2+\delta}^{1,2}}. \end{aligned} \quad (44)$$

The first term is bounded by

$$\|(e^{-\gamma_1} - e^{-\gamma_2})\bar{\partial}_{A_t} e^{\gamma_1}\|_{\tilde{L}_{-2+\delta}^{1,2}} + \|e^{-\gamma_2}(\bar{\partial}_{A_t}(e^{\gamma_1} - e^{\gamma_2})) - \bar{\partial}_{A_t}(\gamma_1 - \gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} := I + II.$$

Write $e^\gamma = 1 + \gamma + S(\gamma)$, then

$$\begin{aligned} I &\leq ct^{9/2} \|e^{-\gamma_1} - e^{-\gamma_2}\|_{\tilde{L}_{-2+\delta}^{2,2}} \|\bar{\partial}_{A_t} e^{\gamma_1}\|_{\tilde{L}_{-2+\delta}^{1,2}} \leq ct^{9/2} \|e^{-\gamma_1} - e^{-\gamma_2}\|_{\tilde{L}_{-2+\delta}^{2,2}} \|\gamma_1 + S(\gamma_1)\|_{\tilde{L}_{-2+\delta}^{2,2}} \\ &\leq ct^{9/2} \|\gamma_1 - \gamma_2\|_{\tilde{L}_{-2+\delta}^{2,2}} \left(\sum_{j=0}^{\infty} \frac{(\rho ct^5)^j}{j!} \right) \rho \left(\sum_{j=0}^{\infty} \frac{(\rho ct^5)^j}{(j+1)!} \right) \\ &\leq ct^{9/2} \rho \|\gamma_1 - \gamma_2\|_{\tilde{L}_{-2+\delta}^{2,2}}, \end{aligned}$$

for $\rho < t^{-5}$. Similarly,

$$\begin{aligned} II &= \|(1 - \gamma_2 + S(-\gamma_2))(\bar{\partial}_{A_t}(\gamma_1 - \gamma_2 + S(\gamma_1) - S(\gamma_2))) - \bar{\partial}_{A_t}(\gamma_1 - \gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\leq \|\bar{\partial}_{A_t}(S(\gamma_1) - S(\gamma_2))\|_{\tilde{L}_{-2+\delta}^{1,2}} + \|(-\gamma_2 + S(-\gamma_2))\bar{\partial}_{A_t}(\gamma_1 - \gamma_2 + S(\gamma_1) - S(\gamma_2))\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\leq \|S(\gamma_1) - S(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{2,2}} + ct^{9/2} \|-\gamma_2 + S(-\gamma_2)\|_{\tilde{L}_{-2+\delta}^{2,2}} \|\gamma_1 - \gamma_2 + S(\gamma_1) - S(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{2,2}} \\ &\leq ct^{9/2} \rho \|\gamma_1 - \gamma_2\|_{\tilde{L}_{-2+\delta}^{2,2}}. \end{aligned}$$

The estimate for the other term in (44) is analogous, and we obtain

$$\begin{aligned} \|R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} &\leq ct^{9/2}\rho\|\gamma_1 - \gamma_2\|_{\tilde{L}_{-2+\delta}^{2,2}} \leq ct^{9/2}\rho\|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}, \\ \|d_{A_t}(R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2))\|_{L_{-2+\delta}^2} &\leq ct^{9/2}\rho\|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}. \end{aligned} \quad (45)$$

Let $\beta_1, \dots, \beta_n \in L_{-2+\delta}^{2,2}$, where $n \geq 2$. By the Jacobi identity, $[[\Phi_t^*, [\beta_1, \dots, [\beta_n, \Phi_t] \dots]]]$ is bounded by

$$\begin{aligned} &|[[\Phi_t^*, \beta_1], [\beta_2, \dots, [\Phi_t, \beta_n] \dots]]| + |[\beta_1, [[\Phi_t^*, \beta_2], [\beta_3, \dots, [\Phi_t, \beta_n] \dots]]]| \\ &+ \dots + |[\beta_1, \dots, [[\Phi_t^*, \beta_{n-1}], [\Phi_t, \beta_n]] \dots]| + |[\beta_1, \dots, [\beta_{n-1}, [\Phi_t^*, [\Phi_t, \beta_n]]] \dots]| \\ &\leq 2^n \left(\sum_{j=1}^{n-1} \left(|[\Phi_t^*, \beta_j]| |[\Phi_t, \beta_n]| \prod_{m \neq j, n} \|\beta_m\|_{L^\infty} \right) + |[\Phi_t^*, [\Phi_t, \beta_n]]| \prod_{m=1}^{n-1} \|\beta_m\|_{L^\infty} \right). \end{aligned}$$

Integrate both sides, then $[[\Phi_t^*, [\beta_1, \dots, [\beta_n, \Phi_t] \dots]]]$ is bounded by

$$\begin{aligned} &2^n \left(\sum_{j=1}^{n-1} ct^5 \|[\Phi_t^*, \beta_j]\|_{\tilde{L}_{-2+\delta}^{1,2}} \|[\Phi_t, \beta_n]\|_{\tilde{L}_{-2+\delta}^{1,2}} (ct^3)^{n-2} \prod_{m \neq j, n} \|\beta_m\|_{\tilde{L}_{-2+\delta}^{2,2}} \right. \\ &\quad \left. + \|[\Phi_t^*, [\Phi_t, \beta_n]]\|_{L_{-2+\delta}^2} (ct^3)^{n-1} \prod_{m=1}^{n-1} \|\beta_m\|_{\tilde{L}_{-2+\delta}^{2,2}} \right) \\ &\leq (ct^{9/2})^{n-1} t^{1/2} \prod_{m=1}^n \|\beta_m\|_{L_{-2+\delta}^{2,2}}. \end{aligned}$$

Then for $\gamma_1, \gamma_2 \in B_\rho$,

$$\begin{aligned} \|[\Phi_t^*, (\text{ad}_{\gamma_1})^n \Phi_t - (\text{ad}_{\gamma_2})^n \Phi_t]\|_{L_{-2+\delta}^2} &\leq \|[\Phi_t^*, [\gamma_1 - \gamma_2, [\gamma_1, \dots, [\gamma_1, \Phi_t] \dots]]]\|_{L_{-2+\delta}^2} \\ &\quad + \|[\Phi_t^*, [\gamma_2, [\gamma_1 - \gamma_2, [\gamma_1, \dots, [\gamma_1, \Phi_t] \dots]]]\|_{L_{-2+\delta}^2} \\ &\quad + \dots + \|[\Phi_t^*, [\gamma_2, [\gamma_2, \dots, [\gamma_1 - \gamma_2, \Phi_t] \dots]]]\|_{L_{-2+\delta}^2} \\ &\leq n(ct^{9/2}\rho)^{n-1} t^{1/2} \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}. \end{aligned}$$

Then we compute that

$$\begin{aligned} \| [R_{\Phi_t}(\gamma_1), \Phi_t^*] - [R_{\Phi_t}(\gamma_2), \Phi_t^*] \|_{L_{-2+\delta}^2} &= \| [\Phi_t^*, \sum_{j=2}^{\infty} \frac{1}{j!} (\text{ad}_{-\gamma_1})^j \Phi_t - (\text{ad}_{-\gamma_2})^j \Phi_t] \|_{L_{-2+\delta}^2} \\ &\leq \sum_{j=2}^{\infty} \frac{1}{j!} j (ct^{9/2}\rho)^{j-1} t^{1/2} \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}} \\ &\leq ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}. \end{aligned}$$

Similarly,

$$\| [\Phi_t, R_{\Phi_t}(\gamma_1)^*] - [\Phi_t, R_{\Phi_t}(\gamma_2)^*] \|_{L_{-2+\delta}^2} \leq ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}.$$

The $L_{-2+\delta}^2$ norm of the fourth term in $Q_t(\gamma_1) - Q_t(\gamma_2)$ is bounded by

$$\begin{aligned} &\| [(\bar{\partial}_{A_t} - \partial_{A_t})(\gamma_1 - \gamma_2) + R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2), (\bar{\partial}_{A_t} - \partial_{A_t})\gamma_1 + R_{A_t}(\gamma_1)] \|_{L_{-2+\delta}^2} \\ &+ \| [(\bar{\partial}_{A_t} - \partial_{A_t})\gamma_2 + R_{A_t}(\gamma_2), (\bar{\partial}_{A_t} - \partial_{A_t})(\gamma_1 - \gamma_2) + R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2)] \|_{L_{-2+\delta}^2} := I' + II'. \end{aligned}$$

By (43), (45),

$$\begin{aligned} I' &\leq ct^5 \|(\bar{\partial}_{A_t} - \partial_{A_t})(\gamma_1 - \gamma_2) + R_{A_t}(\gamma_1) - R_{A_t}(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} \|(\bar{\partial}_{A_t} - \partial_{A_t})\gamma_1 + R_{A_t}(\gamma_1)\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\leq ct^5 \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}} (1 + ct^{9/2}\rho)^2 \rho \leq ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}. \end{aligned}$$

Similarly, II' is also bounded by $ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}$. Consider $\beta_1, \dots, \beta_n \in L_{-2+\delta}^{2,2}$, then

$$\begin{aligned} \|[\beta_1, [\beta_2, \dots, [\beta_n, \Phi_t] \dots]]\|_{\tilde{L}_{-2+\delta}^{1,2}} &\leq (ct^{9/2})^{n-1} \prod_{m=1}^{n-1} \|\beta_m\|_{L_{-2+\delta}^{2,2}} \|[\Phi_t, \beta_n]\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\leq (ct^{9/2})^{n-1} \prod_{m=1}^n \|\beta_m\|_{L_{-2+\delta}^{2,2}}. \end{aligned}$$

This implies that

$$\|R_{\Phi_t}(\gamma_1) - R_{\Phi_t}(\gamma_2)\|_{\tilde{L}_{-2+\delta}^{1,2}} \leq \sum_{j=2}^{\infty} \frac{1}{j!} j (ct^{9/2}\rho)^{j-1} \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}} \leq ct^{9/2}\rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}.$$

The $L_{-2+\delta}^2$ norm of the last term in $Q_t(\gamma_1) - Q_t(\gamma_2)$ is bounded by

$$\begin{aligned} &\|[[\Phi_t, \gamma_1 - \gamma_2] + R_{\Phi_t}(\gamma_1) - R_{\Phi_t}(\gamma_2), ([\Phi_t, \gamma_1] + R_{\Phi_t}(\gamma_1))^*]\|_{L_{-2+\delta}^2} \\ &+ \|[[\Phi_t, \gamma_2] + R_{\Phi_t}(\gamma_2), ([\Phi_t, \gamma_1 - \gamma_2] + R_{\Phi_t}(\gamma_1) - R_{\Phi_t}(\gamma_2))^*]\|_{L_{-2+\delta}^2} := I'' + II'', \end{aligned}$$

where I'' can be estimated as

$$\begin{aligned} I'' &\leq ct^5 \|[[\Phi_t, \gamma_1 - \gamma_2] + R_{\Phi_t}(\gamma_1) - R_{\Phi_t}(\gamma_2)]\|_{\tilde{L}_{-2+\delta}^{1,2}} \|[\Phi_t, \gamma_1] + R_{\Phi_t}(\gamma_1)\|_{\tilde{L}_{-2+\delta}^{1,2}} \\ &\leq ct^5 \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}} (1 + ct^{9/2}\rho)^2 \rho \leq ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}, \end{aligned}$$

similar for II'' . Therefore

$$\|Q_t(\gamma_1) - Q_t(\gamma_2)\|_{L_{-2+\delta}^2} \leq ct^5 \rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}. \quad (46)$$

Now $F_t(\gamma) = 0$ is equivalent to

$$\gamma = -L_t^{-1}(F_{A_t} + [\Phi_t, \Phi_t^*] + Q_t(\gamma)) := T_t(\gamma).$$

By (37), (46), $\|T_t(\gamma_1) - T_t(\gamma_2)\|_{L_{-2+\delta}^{2,2}} \leq ct^{13}\rho \|\gamma_1 - \gamma_2\|_{L_{-2+\delta}^{2,2}}$. Furthermore, by (17) (which is still valid when the norm is taken with respect to g_t),

$$\|T_t(\gamma)\|_{L_{-2+\delta}^{2,2}} \leq ct^8 e^{-c't^\sigma} + ct^{13}\rho^2,$$

for $\gamma \in B_\rho$. Let $\rho_t = e^{(-c'/2+\epsilon)t^\sigma}$, where $\epsilon < c'/2$ is small. Then for t large, $\rho_t < t^{-5}$, T_t maps B_{ρ_t} to B_{ρ_t} , and is a contraction mapping, hence there is a unique fixed point γ_t in B_{ρ_t} . In other words, the metric h_t is harmonic, which is also compatible with the parabolic structure as h_t^{app} is and $\gamma_t \in L_{-2+\delta}^{2,2}$. \square

6. ASYMPTOTIC METRIC

In this section, we study the asymptotic geometry of \mathcal{M} using the framework provided by [Fre20, FMSW22].

Elements in \mathcal{M} are gauge equivalence classes of pairs $(\bar{\partial}_E, \Phi)$, with infinitesimal deformations denoted by $(\dot{\eta}, \dot{\Phi})$. Linearize $\bar{\partial}_E \Phi = 0$ and we get

$$\bar{\partial}_E \dot{\Phi} + [\dot{\eta}, \Phi] = 0. \quad (47)$$

For $[(\bar{\partial}_E, \Phi)] \in \mathcal{M}$, let h be the unique harmonic metric (if exists) which induces $h_{\det E}$, and denote its infinitesimal deformation by \dot{h} . The infinitesimal gauge transformation $\dot{\gamma}$ acts as

$$(\dot{\eta}, \dot{\Phi}, \dot{h}) \mapsto (\dot{\eta} + \bar{\partial}_E \dot{\gamma}, \dot{\Phi} + [\Phi, \dot{\gamma}], \dot{h} + \dot{\gamma}).$$

The linearized Hitchin equation and the Coulomb gauge condition can be combined to form a single equation [Fre20, Prop. 2.2]

$$\partial_E^h \bar{\partial}_E \dot{h} - \partial_E^h \dot{\eta} - [\Phi^{*h}, \dot{\Phi} + [\dot{h}, \Phi]] = 0. \quad (48)$$

Recall that $2\partial_E^h \bar{\partial}_E = F_{A_h} + i \star \Delta_{A_h} = -[\Phi, \Phi^{*h}] + i \star \Delta_{A_h}$, the equation can be written as

$$-\frac{1}{2}[[\Phi, \Phi^{*h}], \dot{h}] + \frac{1}{2}i \star \Delta_{A_h} \dot{h} - [\Phi^{*h}, [\dot{h}, \Phi]] = \partial_E^h \dot{\eta} + [\Phi^{*h}, \dot{\Phi}].$$

Composing with $-2i \star$ and simplifying, we obtain

$$L_\Phi \dot{h} := \Delta_{A_h} \dot{h} - i \star ([\Phi^{*h}, [\Phi, \dot{h}]] - [\Phi, [\Phi^{*h}, \dot{h}]]) = -2i \star (\partial_E^h \dot{\eta} + [\Phi^{*h}, \dot{\Phi}]).$$

Note that L_Φ is exactly L_t in (18) with $(\bar{\partial}_E, \Phi_t, h_t^{\text{app}})$ replaced by $(\bar{\partial}_E, \Phi, h)$. Now we consider the solvability of (48). Let $(\bar{\partial}_E, \Phi_t)$ and $L_\delta^{k,p}$ (depending on g_t, Φ_t , and h_t^{app}) be defined as before, ∇_t be the Chern connection determined by $\bar{\partial}_E$ and h_t .

Lemma 6.1. *Fix a pair $[(\bar{\partial}_E, \Phi_t)] \in \mathcal{M}$, $t \geq t_1$ so that the harmonic metric h_t exists. Then $L_{\Phi_t} : L_{-2+\delta}^{2,2}(\mathfrak{sl}(E)) \rightarrow L_{-2+\delta}^2(\mathfrak{sl}(E))$ is an isomorphism.*

Proof. That L_t is Fredholm of index zero follows from Lemma 4.11, and the fact that L_t preserves the decomposition $\mathfrak{sl}(E) = \mathfrak{su}(E) \oplus i\mathfrak{su}(E)$. Since $\gamma_t \in L_{-2+\delta}^{2,2}$, L_{Φ_t} differs from L_t by a compact perturbation, it is also Fredholm of index zero. If $u \in \ker L_{\Phi_t}$, then as before the integration by parts yields $\nabla_t u = [\Phi_t, u] = 0$. By the proof of [FMSW22, Prop. 7.4], we have $u = 0$. Therefore L_{Φ_t} is an isomorphism. \square

Lemma 6.2. *Let $[(\bar{\partial}_E, \Phi)] \in \mathcal{M}$ and $(\dot{\eta}_1, \dot{\Phi}_1)$ be an infinitesimal variation. There exists $(\dot{\eta}_2, \dot{\Phi}_2)$ infinitesimally gauge equivalent to $(\dot{\eta}_1, \dot{\Phi}_1)$ such that the followings hold.*

(i) *In a neighborhood of $x \in T$ and a holomorphic frame where*

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi = \begin{pmatrix} 0 & 1 \\ \zeta_x^{-1} & 0 \end{pmatrix} d\zeta_x,$$

we have

$$\dot{\eta}_2 = 0, \quad \dot{\Phi}_2 = \begin{pmatrix} 0 & 0 \\ \dot{P} & 0 \end{pmatrix} d\zeta_x, \quad \dot{P} \text{ holomorphic.}$$

(ii) *In a neighborhood of $x \in I_u$ and a holomorphic frame where $\bar{\partial}_E = \bar{\partial}$, $\Phi = f z_x^{-2m_x} \sigma_3 dz_x$ with f being holomorphic and nowhere vanishing, we have $\dot{\eta}_2 = 0$, $\dot{\Phi}_2 = \dot{P} \sigma_3 dz$, \dot{P} is holomorphic.*

(iii) In a neighborhood of $x \in I_t$ and a holomorphic frame where

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi = \begin{pmatrix} 0 & \frac{1}{\zeta_x^{m_x-1}} \\ \frac{1}{\zeta_x^{m_x}} & 0 \end{pmatrix} d\zeta_x,$$

we have

$$\dot{\eta}_2 = 0, \quad \dot{\Phi}_2 = \begin{pmatrix} 0 & \dot{P} \\ \frac{\dot{P}}{\zeta_x} & 0 \end{pmatrix} d\zeta_x, \quad \dot{P} \text{ is holomorphic.}$$

Proof. (i) is just [FMSW22, Lem. 7.3].

(ii) By the Poincaré lemma we may assume that $\dot{\eta}_1 = 0$ in the disk. The leading term of $\dot{\Phi}_1$ preserves the flag at x , then we can write $\dot{\Phi}_1$ as

$$\dot{\Phi}_1 = \frac{1}{z_x^{2m_x}} \begin{pmatrix} \dot{P}_1 & \dot{P}_2 z_x \\ \dot{P}_3 & -\dot{P}_1 \end{pmatrix}.$$

where $\dot{P}_1, \dot{P}_2, \dot{P}_3$ are holomorphic by (47). Then $\dot{\Phi}_2 = \dot{\Phi}_1 + [\Phi, \dot{\gamma}] = \dot{P}_1 z_x^{-2m_x} \sigma_3 dz$ if we choose

$$\dot{\gamma} = \begin{pmatrix} 0 & -\frac{\dot{P}_2 z_x}{2f} \\ \frac{\dot{P}_3}{2f} & 0 \end{pmatrix},$$

which satisfies $\bar{\partial}\dot{\gamma} = 0$ and preserves the flag. Higgs fields in \mathcal{M} have fixed polar parts, implying that $\dot{P}_1 z_x^{-2m_x} = \dot{P}$ for some holomorphic \dot{P} .

(iii) As above, assume that $\dot{\eta}_1 = 0$. The leading term of $\dot{\Phi}_1$ preserves the flag at x and the leading term of the Higgs field is nilpotent, then we can write $\dot{\Phi}_1$ as

$$\dot{\Phi}_1 = \frac{1}{\zeta_x^{2m_x-1}} \begin{pmatrix} \dot{P}_1 & \dot{P}_2 \\ \frac{\dot{P}_3}{\zeta_x} & -\dot{P}_1 \end{pmatrix}.$$

where $\dot{P}_1, \dot{P}_2, \dot{P}_3$ are holomorphic by (47). We choose

$$\dot{\gamma} = \begin{pmatrix} \frac{\dot{P}_2 - \dot{P}_3}{4} & \zeta_x \dot{P}_1 \\ 0 & -\frac{\dot{P}_2 - \dot{P}_3}{4} \end{pmatrix},$$

which satisfies $\bar{\partial}\dot{\gamma} = 0$ and preserves the flag. Then

$$\dot{\Phi}_2 = \dot{\Phi}_1 + [\Phi, \dot{\gamma}] = \begin{pmatrix} 0 & \frac{\dot{P}_2 + \dot{P}_3}{2\zeta_x^{2m_x-1}} \\ \frac{\dot{P}_2 + \dot{P}_3}{2\zeta_x^{2m_x}} & 0 \end{pmatrix} d\zeta_x.$$

Higgs fields in \mathcal{M} have fixed polar parts, implying that $(\dot{P}_2 + \dot{P}_3)/(2\zeta_x^{2m_x-1}) = \dot{P}$ for \dot{P} holomorphic. \square

Suppose $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$, by the above lemma we may assume that $\dot{\eta} = 0$ and $\dot{\Phi}$ has the above forms near every $x \in S$. Then $\partial_E^{h_t} \dot{\eta} + [\Phi_t^{*h_t}, \dot{\Phi}] \in L_{-2+\delta}^2$, as $\partial_E^{h_t} \dot{\eta} = 0$ near $x \in S$, $[\Phi_t^{*h_t, \text{app}}, \dot{\Phi}] \in L_{-2+\delta}^2$ for δ small and $\gamma_t \in L_{-2+\delta}^{2,2}$. By Lemma 6.1, a unique $\dot{h} \in L_{-2+\delta}^{2,2}$ solving (48) can be found. Now the hyperkähler metric is given by [FMSW22, (7.5)]

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2 = 2 \int_C |\dot{\eta} - \bar{\partial}_E \dot{h}|_{h_t}^2 + |\dot{\Phi} + [\dot{h}, \Phi_t]|_{h_t}^2 = 2 \int_C \langle \dot{\eta} - \bar{\partial}_E \dot{h}, \dot{\eta} \rangle_{h_t} + \langle \dot{\Phi} + [\dot{h}, \Phi_t], \dot{\Phi} \rangle_{h_t}, \quad (49)$$

where the second equality follows from (48) and the integration by parts which is justified as in the proof of Lemma 4.11 and [FMSW22, p. 43].

For the pair $[(\bar{\partial}_E, \Phi_t)] \in \mathcal{M}$, one can associate metrics h_t^{app} and $h_{t,\infty}$, with infinitesimal deformations denoted by \dot{h}_{app} and \dot{h}_∞ . The metric g_{app} on \mathcal{M} is defined by

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{\text{app}}}^2 = 2 \int_C \langle \dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}, \dot{\eta} \rangle_{h_t^{\text{app}}} + \langle \dot{\Phi} + [\dot{h}_{\text{app}}, \Phi_t], \dot{\Phi} \rangle_{h_t^{\text{app}}}, \quad (50)$$

where \dot{h}_{app} satisfies

$$\partial_E^{h_t^{\text{app}}} \bar{\partial}_E \dot{h}_{\text{app}} - \partial_E^{h_t^{\text{app}}} \dot{\eta} - [\Phi_t^{*h_t^{\text{app}}}, \dot{\Phi} + [\dot{h}_{\text{app}}, \Phi_t]] = 0. \quad (51)$$

Similarly, the metric g_∞ is

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_\infty}^2 = 2 \int_C \langle \dot{\eta} - \bar{\partial}_E \dot{h}_\infty, \dot{\eta} \rangle_{h_{t,\infty}} + \langle \dot{\Phi} + [\dot{h}_\infty, \Phi_t], \dot{\Phi} \rangle_{h_{t,\infty}}, \quad (52)$$

where \dot{h}_∞ satisfies the decoupled equations

$$\partial_E^{h_{t,\infty}} \bar{\partial}_E \dot{h}_\infty - \partial_E^{h_{t,\infty}} \dot{\eta} = 0, \quad [\Phi_t^{*h_{t,\infty}}, \dot{\Phi} + [\dot{h}_\infty, \Phi_t]] = 0. \quad (53)$$

Recall that for each $[(\bar{\partial}_E, \Phi_t)] \in \mathcal{M}$, we can view the associated spectral data as a pair $(\bar{\partial}_{L,t}, \tau_t)$, where L is a complex line bundle of degree $-|S| + N - 2$ over a surface S_t with genus $N - 3$, τ_t is the tautological eigenvalue of $\pi^* \Phi_t$ which embeds S_t in $\pi^* K(D)$. The tangent map induces an isomorphism

$$T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M} \rightarrow H^0(S_t, K_{S_t})_{\text{odd}}^* \oplus H^0(S_t, K_{S_t})_{\text{odd}}, \quad [(\dot{\eta}, \dot{\Phi})] \mapsto (\dot{\xi}, \dot{\tau}),$$

where $H^0(S_t, K_{S_t})_{\text{odd}}$ consists of elements of $H^0(S_t, K_{S_t})$ which are odd under the exchange of two sheets of $\pi : S_t \rightarrow C$, and $\dot{\xi}, \dot{\tau}$ are the infinitesimal deformations of $\bar{\partial}_{L,t}$ and τ . The metric $h_{t,\infty}$ is the orthogonal pushforward of the harmonic metric $h_{\mathcal{L}_t}$ on the spectral line bundle $\mathcal{L}_t = (L, \bar{\partial}_{L,t}) \rightarrow S_t$, and the infinitesimal deformation \dot{h}_L of $h_{\mathcal{L}_t}$ is determined by

$$\partial_L^{h_{\mathcal{L}_t}} \bar{\partial}_{L,t} \dot{h}_L - \partial_L^{h_{\mathcal{L}_t}} \dot{\xi} = 0.$$

Lemma 6.3 ([Fre20, Prop. 2.14]). *The semiflat metric is characterized by three properties:*

- (1) *On horizontal deformations, the semiflat metric is $\int_{S_t} 2|\dot{\tau}|^2$.*
- (2) *On vertical deformations, the semiflat metric is $\int_{S_t} 2|\dot{\xi} - \bar{\partial}_L \dot{h}_L|^2$.*
- (3) *Horizontal and vertical deformations are orthogonal.*

Lemma 6.4 ([FMSW22, Cor. 7.9]). *The semiflat metric g_{sf} on \mathcal{M} coincides with g_∞ .*

6.1. Comparing g_{L^2} and g_{app} . The difference $\frac{1}{2} \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{\text{app}}}^2 - \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2$ is bounded by

$$\left| \int_C |\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}|_{h_t^{\text{app}}}^2 - |\dot{\eta} - \bar{\partial}_E \dot{h}|_{h_t}^2 \right| + \left| \int_C |\dot{\Phi} + [\dot{h}_{\text{app}}, \Phi_t]|_{h_t^{\text{app}}}^2 - |\dot{\Phi} + [\dot{h}, \Phi_t]|_{h_t}^2 \right| := I + II.$$

We first estimate I ,

$$\begin{aligned} I &\leq \int_C |\langle (\text{Id} - e^{-\gamma t})(\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}), \dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}} \rangle_{h_t^{\text{app}}}| \\ &\quad + |\langle e^{-\gamma t}(\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}), (\text{Id} - e^{-\gamma t})(\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}) \rangle_{h_t^{\text{app}}}| \\ &\quad + 2|\langle e^{-\gamma t} \bar{\partial}_E(\dot{h}_{\text{app}} - \dot{h}), e^{-\gamma t}(\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}) \rangle_{h_t^{\text{app}}}| + |e^{-\gamma t} \bar{\partial}_E(\dot{h}_{\text{app}} - \dot{h})|_{h_t^{\text{app}}}^2. \end{aligned} \quad (54)$$

Since the $L_{-2+\delta}^{2,2}$ norm of γ_t is exponentially decaying, the first two terms are bounded by

$$ce^{-c't^\sigma} \|\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}\|_{L^2}^2.$$

Here and following, unless specified, the norms are taken with respect to h_t^{app} as in (20). It remains to estimate $\|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2}$, where $\dot{\mathbf{v}} = \dot{h}_{\text{app}} - \dot{h}$. By (48), (51), we have

$$\begin{aligned} \partial_E^{h_t^{\text{app}}} \bar{\partial}_E \dot{\mathbf{v}} - [\Phi_t^{*h_t^{\text{app}}}, [\dot{\mathbf{v}}, \Phi_t]] + T_t^1(\dot{\eta} - \bar{\partial}_E \dot{h}) + T_t^2(\dot{\Phi} + [\dot{h}, \Phi_t]) &= 0, \text{ where} \\ T_t^1 = \partial_E^{h_t} - \partial_E^{h_t^{\text{app}}} &= -2[\partial_E^{h_t^{\text{app}}} \gamma_t, \cdot] + R_t^1, \quad T_t^2 = [\Phi_t^{*h_t} - \Phi_t^{*h_t^{\text{app}}}, \cdot] = -2[[\gamma_t, \Phi_t^{*h_t^{\text{app}}}], \cdot] + R_t^2, \end{aligned}$$

R_t^1, R_t^2 are nonlinear in γ_t . As before, by composing with $-2i\star$, the equation becomes

$$L_t \dot{\mathbf{v}} = 2i\star (T_t^1(\dot{\eta} - \bar{\partial}_E \dot{h}) + T_t^2(\dot{\Phi} + [\dot{h}, \Phi_t]) + E_t \dot{\mathbf{v}}) := \dot{\mathbf{w}},$$

where $E_t = [F_{h_t^{\text{app}}} + [\Phi_t, \Phi_t^{*h_t^{\text{app}}}], \cdot]$. Note that $\nabla_{A_t} \gamma_t, [\Psi_t, \gamma_t] \in L_{-2+\delta}^{1,2}$ and $L_{-2+\delta}^{1,2} \hookrightarrow L_{-1+\delta}^p$ for any fixed $p \in [2, \infty)$, $L_{-1+\delta}^p \cdot L^2 \subset L_{-2+\delta}^{2p/(p+2)}$, similar to the proof of Theorem 5.2, we have

$$\|\star T_t^1(\dot{\eta} - \bar{\partial}_E \dot{h})\|_{L_{-2+\delta}^{p_1}} + \|\star T_t^2(\dot{\Phi} + [\dot{h}, \Phi_t])\|_{L_{-2+\delta}^{p_1}} \leq ce^{-c't^\sigma} \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}},$$

where $p_1 = 2p/(p+2) < 2$. By the construction of h_t^{app} , E_t is supported on a union of annuli and the C^0 norm is exponentially decaying, so that

$$\|E_t \dot{\mathbf{v}}\|_{L_{-2+\delta}^{p_1}} \leq ce^{-c't^\sigma} \|\dot{\mathbf{v}}\|_{L_{-2+\delta}^{p_1}}.$$

Therefore,

$$\|\dot{\mathbf{w}}\|_{L_{-2+\delta}^{p_1}} \leq ce^{-c't^\sigma} (\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}} + \|\dot{\mathbf{v}}\|_{L_{-2+\delta}^{p_1}}). \quad (55)$$

By (26),

$$\|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2}^2 + \|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2}^2 = \langle \dot{\mathbf{w}}, \dot{\mathbf{v}} \rangle_{L^2} \leq \|\dot{\mathbf{w}}\|_{L_{-2+\delta}^{p_1}} \|\dot{\mathbf{v}}\|_{L_{-1-\delta}^{p_1^*}}, \quad (56)$$

where $1/p_1 + 1/p_1^* = 1$. Let u be a function on the unit disk vanishing on the boundary, then we have the Sobolev inequality $\|u\|_{L_{-1}^{p_1^*}} \leq c\|\nabla u\|_{L^2}$. On the other hand, by Lemma 4.5, $L_{-\delta-1}^{1,2} = \hat{L}_{-\delta-1}^{1,2} \hookrightarrow L_{-\delta}^{p_1^*}$, then

$$\|u\|_{L_{-\delta}^{p_1^*}} \leq c(\|\nabla u\|_{L_{-\delta-1}^2} + \|u\|_{L_{-\delta-1}^2}) \leq c(\|\nabla u\|_{L_{-\delta-1}^2} + \|\nabla u\|_{L_{-\delta-2}^2}) \leq c\|\nabla u\|_{L^2}.$$

These estimates can be translated to C as in the proof of (27), and we obtain

$$\|\dot{\mathbf{v}}\|_{L_{-\delta}^{p_1^*}} \leq ct^M (\|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2} + \|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2}),$$

for some constant M . Combining with (55), (56), we deduce that

$$\begin{aligned} \|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2} + \|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2} &\leq ct^M \|\dot{\mathbf{w}}\|_{L_{-2+\delta}^{p_1}} \leq ce^{-c't^\sigma} (\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}} + \|\dot{\mathbf{v}}\|_{L_{-2+\delta}^{p_1}}) \\ &\leq ce^{-c't^\sigma} (\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}} + \|\dot{\mathbf{v}}\|_{L_{-\delta}^{p_1^*}}) \\ &\leq ce^{-c't^\sigma} (\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}} + \|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2} + \|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2}), \\ \|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2} + \|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2} &\leq ce^{-c't^\sigma} \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}, \end{aligned}$$

for t large. By (54), we conclude that

$$I \leq ce^{-c't^\sigma} (\|\dot{\eta} - \bar{\partial}_E \dot{h}_{\text{app}}\|_{L^2}^2 + \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2).$$

Similarly, with the roll of $\|\nabla_{A_t} \dot{\mathbf{v}}\|_{L^2}$ replaced by $\|[\Psi_t, \dot{\mathbf{v}}]\|_{L^2}$, we have

$$II \leqslant ce^{-c't^\sigma} (\|\dot{\Phi} + [\dot{h}_{\text{app}}, \Phi_t]\|_{L^2}^2 + \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2).$$

Therefore $|\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{\text{app}}}^2 - \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2| \leqslant ce^{-c't^\sigma} \|[(\dot{\eta}, \dot{\Phi})]\|_{g_{L^2}}^2$, meaning that along the curve $[(\bar{\partial}_E, \Phi_t)]$ in \mathcal{M} , g_{L^2} and g_{app} are exponentially close in t .

6.2. Comparing g_{app} and g_{sf} . By the construction, h_t^{app} and $h_{t,\infty}$ coincide on the complement of the disks $\tilde{B}_{x,j,t}$ ($x \in I$) and $\tilde{B}_{x,t}$ ($x \in T$), and so do the infinitesimal deformations \dot{h}_{app} and \dot{h}_∞ . Accordingly, the difference $g_{\text{app}} - g_{\text{sf}}$ localizes on these disks.

On $\tilde{B}_{x,j,t}$, in a local holomorphic coordinate $\zeta_{x,j,t}$ and a holomorphic frame we have

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \lambda_{x,j}(t) \begin{pmatrix} 0 & 1 \\ \zeta_{x,j,t} & 0 \end{pmatrix} d\zeta_{x,j,t}, \quad h_{t,\infty} = \begin{pmatrix} |\zeta_{x,j,t}|^{1/2} & 0 \\ 0 & |\zeta_{x,j,t}|^{-1/2} \end{pmatrix},$$

and h_t^{app} is given by (15). Let $\hat{\zeta}_{x,j,t} = \kappa^{-1} t^{1/j(x)} \zeta_{x,j,t}$ and $\hat{r}_{x,j,t} = |\hat{\zeta}_{x,j,t}|$, then the disk $\tilde{B}_{x,j,t} = \{\hat{r}_{x,j,t} < 1\}$. Applying the local holomorphic gauge transformation

$$g = \begin{pmatrix} \kappa^{-1/4} t^{1/4j(x)} & 0 \\ 0 & \kappa^{1/4} t^{-1/4j(x)} \end{pmatrix}, \quad \text{then } g^{-1} \Phi_t g = t_{x,j} \begin{pmatrix} 0 & 1 \\ \hat{\zeta}_{x,j,t} & 0 \end{pmatrix} d\hat{\zeta}_{x,j,t},$$

where $t_{x,j} = \lambda_{x,j}(t) \kappa^{3/2} t^{-3/2j(x)}$.

$$g^* h_{t,\infty} g = \begin{pmatrix} \hat{r}_{x,j,t}^{1/2} & 0 \\ 0 & \hat{r}_{x,j,t}^{-1/2} \end{pmatrix}, \quad g^* h_t^{\text{app}} g = \begin{pmatrix} \hat{r}_{x,j,t}^{1/2} e^{l_{t,x,j}(\hat{r}_{x,j,t}) \chi(\hat{r}_{x,j,t})} & 0 \\ 0 & \hat{r}_{x,j,t}^{-1/2} e^{-l_{t,x,j}(\hat{r}_{x,j,t}) \chi(\hat{r}_{x,j,t})} \end{pmatrix},$$

where $l_{t,x,j}(\hat{r}_{x,j,t}) = \psi_1(8t_{x,j} \hat{r}_{x,j,t}^{3/2}/3)$. In this local holomorphic frame, the following result holds.

Proposition 6.5 ([Fre20, Th. 3.1]). *Let $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$, then on the disk $\{\hat{r}_{x,j,t} < 1\}$ there is a unique representative in the equivalence class where*

$$\dot{\eta} = 0, \quad \dot{\Phi} = t_{x,j} \begin{pmatrix} 0 & 0 \\ \dot{P} & 0 \end{pmatrix} d\hat{\zeta}_{x,j,t}, \quad \dot{h}_\infty = \frac{\dot{P}}{4\hat{\zeta}_{x,j,t}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{for } \bar{\partial} \dot{P} = 0.$$

These infinitesimal deformations satisfy (53), and \dot{h}_{app} solving (51) is diagonal. For any $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$, there exists a constant $c > 0$ such that

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{\text{app}}(\tilde{B}_{x,j,t})}^2 - \|[(\dot{\eta}, \dot{\Phi})]\|_{g_\infty(\tilde{B}_{x,j,t})}^2 = O(e^{-ct_{x,j}}).$$

On $\tilde{B}_{x,t}$ ($x \in T$), in a local holomorphic coordinate $\zeta_{x,t}$ and holomorphic frame we have

$$\bar{\partial}_E = \bar{\partial}, \quad \Phi_t = \lambda_x(t) \begin{pmatrix} 0 & 1 \\ \frac{1}{\zeta_{x,t}} & 0 \end{pmatrix} d\zeta_{x,t}, \quad h_{t,\infty} = \begin{pmatrix} |\zeta_{x,t}|^{1/2} & 0 \\ 0 & |\zeta_{x,t}|^{3/2} \end{pmatrix},$$

and h_t^{app} is given by (16). Let $\hat{\zeta}_{x,t} = \kappa^{-1} \zeta_{x,t}$ and $\hat{r}_{x,t} = |\hat{\zeta}_{x,t}|$, then the disk $\tilde{B}_{x,t} = \{\hat{r}_{x,t} < 1\}$. Applying the local holomorphic gauge transformation

$$g = \kappa^{-1/2} \begin{pmatrix} \kappa^{1/4} & 0 \\ 0 & \kappa^{-1/4} \end{pmatrix}, \quad \text{then } g^{-1} \Phi_t g = t_x \begin{pmatrix} 0 & 1 \\ \hat{\zeta}_{x,t} & 0 \end{pmatrix} d\hat{\zeta}_{x,t},$$

where $t_x = \lambda_x(t)\kappa^{1/2}$.

$$g^* h_{t,\infty} g = \begin{pmatrix} \hat{r}_{x,t}^{1/2} & 0 \\ 0 & \hat{r}_{x,t}^{3/2} \end{pmatrix}, \quad g^* h_t^{\text{app}} g = \begin{pmatrix} \hat{r}_{x,t}^{1/2} e^{m_{t_x}(\hat{r}_{x,t})\chi(\hat{r}_{x,t})} & 0 \\ 0 & \hat{r}_{x,t}^{3/2} e^{-m_{t_x}(\hat{r}_{x,t})\chi(\hat{r}_{x,t})} \end{pmatrix},$$

where $m_{t_x}(\hat{r}_{x,t}) = \psi_{2,x}(8t_x \hat{r}_{x,t}^{1/2})$. In this local holomorphic frame, the following result holds.

Proposition 6.6 ([FMSW22, Lem. 7.10, Prop. 7.11]). *Let $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$, then on the disk $\{\hat{r}_{x,t} < 1\}$ there is a unique representative in the equivalence class where*

$$\dot{\eta} = 0, \quad \dot{\Phi} = t_x \begin{pmatrix} 0 & 0 \\ \dot{P} & 0 \end{pmatrix} d\hat{\zeta}_{x,t}, \quad \dot{h}_\infty = \frac{\dot{P}}{4} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{for } \bar{\partial}\dot{P} = 0.$$

These infinitesimal deformations satisfy (53). For any $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$, there exists a constant $c > 0$ such that

$$\|[(\dot{\eta}, \dot{\Phi})]\|_{g_{\text{app}}(\tilde{B}_{x,t})}^2 - \|[(\dot{\eta}, \dot{\Phi})]\|_{g_\infty(\tilde{B}_{x,t})}^2 = O(e^{-ct_x}).$$

The above two propositions immediately imply $g_{\text{app}} - g_{\text{sf}} = O(e^{-ct^\sigma})$ along the curve $[(\bar{\partial}_E, \Phi_t)]$ in \mathcal{M} . Hence Theorem 1.1 is established.

Remark 6.7. $g_{L^2} - g_{\text{app}}$ and $g_{\text{app}} - g_{\text{sf}}$ are still exponentially decaying when $I_u = \emptyset$, only slight modifications of the previous proofs are needed.

6.3. Comparing g_{sf} and g_{model} . In this subsection we assume that $\dim_{\mathbb{C}} \mathcal{M} = 2$, or equivalently $N = \deg D = 4$. The Hitchin base now has complex dimension one, parametrized by $t \in \mathbb{C}$. The constants in this subsection will not depend on $\text{Arg}(t)$ and the choice of the Higgs bundle in the Hitchin fiber $H^{-1}(t)$.

6.3.1. An untwisted order four pole. The Hitchin base is given by

$$\mathcal{B} = \left\{ \left(\sum_{k=5}^8 \frac{\mu_k}{z^k} + \frac{t}{z^4} \right) dz^2 \mid t \in \mathbb{C} \right\}, \quad (57)$$

where μ_5, \dots, μ_8 are constants determined by the irregular type at 0, and $\mu_8 \neq 0$. We may assume that $\mu_8 = -1$, using a rescaling of z if necessary. Fix

$$\nu = \nu(z) dz^2 = \left(\sum_{k=5}^8 \frac{\mu_k}{z^k} + \frac{t}{z^4} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z^4} dz^2 \in T_\nu \mathcal{B}'.$$

Then the special Kähler metric g_{SK} , which is the restriction of g_{sf} on \mathcal{B}' , is given by [FMSW22, Prop. 8.3]

$$g_{\text{SK}}(\dot{\nu}, \dot{\nu}) = \int_{\mathbb{C}} \frac{|\dot{\nu}|^2}{|\nu|} d\text{vol}_{\mathbb{C}} = \frac{|\dot{t}|^2}{|t|} \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (z - z_k(t))|} i dz d\bar{z},$$

where $z_k(t) = t^{-1/4} e^{i(k-1)\pi/2} + t^{-1/2} (-1)^k \mu_7/4 + t^{-3/4} e^{-i(k-1)\pi/2} (-\mu_6/4 - \mu_7^2/32) + O(|t|^{-1})$ ($k = 1, \dots, 4$) are the four roots of $z^8 \nu(z) := \tilde{\nu}(z)$. Let $z = t^{-1/4} \xi$ and $z_k = t^{-1/4} \xi_k$, then the integral becomes

$$|t|^{1/2} \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (\xi - \xi_k)|} i d\xi d\bar{\xi}.$$

We define a function

$$F(\xi_1, \dots, \xi_4, \bar{\xi}_1, \dots, \bar{\xi}_4) := \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (\xi - \xi_k)|} i d\xi d\bar{\xi}.$$

Then we can prove that

$$\frac{\partial F}{\partial \xi_k} = \lim_{\epsilon \rightarrow 0} \int_{|\xi - \xi_k| > \epsilon} \frac{\bar{\xi} - \bar{\xi}_k}{2 |\xi - \xi_k|^3} \frac{1}{|\prod_{l \neq k} (\xi - \xi_l)|} i d\xi d\bar{\xi}, \quad \frac{\partial F}{\partial \bar{\xi}_k} = \overline{\frac{\partial F}{\partial \xi_k}}.$$

Then, for $l \neq k$,

$$\begin{aligned} \frac{\partial^2 F}{\partial \xi_k \partial \xi_l} &= \lim_{\epsilon \rightarrow 0} \int_{|\xi - \xi_k| > \epsilon, |\xi - \xi_l| > \epsilon} \frac{\bar{\xi} - \bar{\xi}_k}{2 |\xi - \xi_k|^3} \frac{\bar{\xi} - \bar{\xi}_l}{2 |\xi - \xi_l|^3} \frac{1}{|\prod_{m \neq k, l} (\xi - \xi_m)|} i d\xi d\bar{\xi}, \\ \frac{\partial^2 F}{\partial \xi_k \partial \bar{\xi}_l} &= \lim_{\epsilon \rightarrow 0} \int_{|\xi - \xi_k| > \epsilon, |\xi - \xi_l| > \epsilon} \frac{\bar{\xi} - \bar{\xi}_k}{2 |\xi - \xi_k|^3} \frac{\xi - \xi_l}{2 |\xi - \xi_l|^3} \frac{1}{|\prod_{m \neq k, l} (\xi - \xi_m)|} i d\xi d\bar{\xi}, \\ \frac{\partial^2 F}{\partial \bar{\xi}_k \partial \bar{\xi}_l} &= \lim_{\epsilon \rightarrow 0} \int_{|\xi - \xi_k| > \epsilon, |\xi - \xi_l| > \epsilon} \frac{\xi - \xi_k}{2 |\xi - \xi_k|^3} \frac{\xi - \xi_l}{2 |\xi - \xi_l|^3} \frac{1}{|\prod_{m \neq k, l} (\xi - \xi_m)|} i d\xi d\bar{\xi}. \end{aligned}$$

For any $k = 1, 2, 3, 4$,

$$\frac{\partial^2 F}{\partial \bar{\xi}_k \partial \xi_k} = - \sum_{l \neq k} \frac{\partial^2 F}{\partial \xi_k \partial \xi_l}, \quad \frac{\partial^2 F}{\partial \xi_k \partial \bar{\xi}_k} = - \sum_{l \neq k} \frac{\partial^2 F}{\partial \xi_k \partial \bar{\xi}_l}, \quad \frac{\partial^2 F}{\partial \bar{\xi}_k \partial \bar{\xi}_k} = - \sum_{l \neq k} \frac{\partial^2 F}{\partial \bar{\xi}_k \partial \bar{\xi}_l}$$

because we can shift ξ by any variation of ξ_k in $\frac{\partial F}{\partial \xi_k}$ or $\frac{\partial F}{\partial \bar{\xi}_k}$.

Now we use these to calculate

$$\begin{aligned} g_{sK}(\dot{\nu}, \dot{\nu}) &= |t|^{1/2} \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (\xi - \xi_k)|} i d\xi d\bar{\xi} = |t|^{1/2} F(\xi_k, \bar{\xi}_k) \\ &= |t|^{1/2} \left(F(e^{i(k-1)\pi/2}, e^{-i(k-1)\pi/2}) + \sum_{k=1}^4 2\operatorname{Re} \left(\frac{\partial F}{\partial \xi_k} \frac{(-1)^k \mu_7 t^{-1/4}}{4} \right) \right. \\ &\quad + \sum_{k=1}^4 2\operatorname{Re} \left(\frac{\partial F}{\partial \bar{\xi}_k} (t^{-1/2} e^{-i(k-1)\pi/2}) (-\mu_6/4 - \mu_7^2/32) \right) \\ &\quad + \operatorname{Re} \left(\sum_{k=1}^4 \sum_{l=1}^4 \frac{\partial^2 F}{\partial \xi_k \partial \xi_l} (-1)^{k+l} \frac{\mu_7^2 t^{-1/2}}{16} \right) \\ &\quad \left. + \operatorname{Re} \left(\sum_{k=1}^4 \sum_{l=1}^4 \frac{\partial^2 F}{\partial \bar{\xi}_k \partial \bar{\xi}_l} \right) (-1)^{k+l} \frac{|\mu_7|^2 |t|^{-1/2}}{16} + O(|t|^{-3/4}) \right). \end{aligned}$$

The terms involving $\frac{\partial F}{\partial \xi_k}$ cancel with each other. So in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(C_0 + \operatorname{Re}(C_1 \mu_7^2 t^{-1/2}) + C_2 |\mu_7|^2 |t|^{-1/2} + O(r^{-3/4})) \frac{dr^2 + r^2 d\theta^2}{\sqrt{r}},$$

which is asymptotic to a conic metric with cone angle $3\pi/2$.

Next we consider g_{sf} along the fiber direction. By the discussion in Section 2, the Hitchin fiber $H^{-1}(t)$ can be written as

$$H^{-1}(t) = \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{dz}{z^4} \begin{pmatrix} \pm(-t)^{1/2}z^2 & 1 - \mu_7 z - \mu_6 z^2 - \mu_5 z^3 \\ 1 & \mp(-t)^{1/2}z^2 \end{pmatrix} \right. \right\} \\ \cup \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{dz}{z^4} \begin{pmatrix} a_0 & -tz^3 + (-\mu_5 + tc_0)z^2 + (-\mu_6 + \mu_5 c_0 - tc_0^2)z - \\ \mu_7 + \mu_6 c_0 - \mu_5 c_0^2 + tc_0^3 & \\ c_0 + z & -a_0 \end{pmatrix}, a_0^2 = -\tilde{\nu}(-c_0) \right. \right\}, \quad (58)$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O} \oplus \mathcal{O}(-1)$. Therefore $H^{-1}(t)$ is the closure of the second set, which is the elliptic curve

$$a_0^2 = -\tilde{\nu}(-c_0) = -t \prod_{k=1}^4 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$. Here λ is the modular lambda function, and

$$l(t) = \frac{(z_4(t) - z_1(t))(z_3(t) - z_2(t))}{(z_3(t) - z_1(t))(z_4(t) - z_2(t))} = \frac{1}{2} + \frac{3i\mu_7^2}{32} t^{-1/2} + O(|t|^{-3/4}), \\ \tau(t) = i + \frac{3\pi\mu_7^2}{32K(\sqrt{1/2})^2} t^{-1/2} + O(|t|^{-3/4}),$$

where $K(x) = \int_0^{\pi/2} (1 - x^2 \sin^2 \theta)^{-1/2} d\theta$ is the complete elliptic integral of the first kind. The semiflat metric induces a constant multiple of the Euclidean metric on $H^{-1}(t)$:

Proposition 6.8 ([FMSW22, Proposition 8.4]). *The area of $H^{-1}(t)$ in the semiflat metric is $4\pi^2$, so $H^{-1}(t) \cong \mathbb{C} / c_t(\mathbb{Z} \oplus \tau(t)\mathbb{Z})$ with Euclidean metric $dx^2 + dy^2$ and $c_t = 2\pi/\sqrt{\text{Im } \tau(t)}$.*

Let $T_{\hat{\tau}}^2 = \mathbb{C} / \hat{c}(\mathbb{Z} \oplus \hat{\tau}\mathbb{Z})$, where $\hat{\tau} = i$, $\hat{c} = 2\pi/\sqrt{\text{Im } \hat{\tau}}$. Define $\mu_t : T_{\hat{\tau}}^2 \rightarrow H^{-1}(t)$ as

$$x + iy \mapsto \frac{c_t}{\hat{c}} \left(x + \frac{\text{Re}(\tau(t) - \hat{\tau})}{\text{Im } \hat{\tau}} y + i \frac{\text{Im } \tau(t)}{\text{Im } \hat{\tau}} y \right) = \frac{c_t}{\hat{c}} (x + y \text{Re } \tau(t) + iy \text{Im } \tau(t)).$$

Then we have

$$\mu_t^*(g_{\text{sf}}|_{H^{-1}(t)}) = \frac{c_t^2}{\hat{c}^2} ((dx + \text{Re } \tau(t) dy)^2 + (\text{Im } \tau(t))^2 dy^2) \\ = dx^2 + dy^2 + \text{Im} \left(\frac{3\pi\mu_7^2}{32K(\sqrt{1/2})^2} t^{-1/2} \right) (-dx^2 + dy^2) \\ + 2\text{Re} \left(\frac{3\pi\mu_7^2}{32K(\sqrt{1/2})^2} t^{-1/2} \right) dx \otimes dy + O(|t|^{-3/4}).$$

6.3.2. *A twisted order four pole.* In this case,

$$\mathcal{B} = \left\{ \left(\sum_{k=5}^7 \frac{\mu_k}{z^k} + \frac{t}{z^4} \right) dz^2 \left| t \in \mathbb{C} \right. \right\},$$

where μ_5, μ_6, μ_7 are fixed and $\mu_7 \neq 0$, we may assume that $\mu_7 = -1$. Fix

$$\nu = \left(\sum_{k=5}^7 \frac{\mu_k}{z^k} + \frac{t}{z^4} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z^4} dz^2 \in T_{\nu} \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$g_{\text{sK}}(\dot{\nu}, \dot{\nu}) = \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = \frac{|t|^2}{|t|} \int_C \frac{1}{|\prod_{k=1}^3 z(z - z_k(t))|} i \text{d}z \text{d}\bar{z},$$

where $z_k(t) = t^{-1/3} e^{2(k-1)\pi i/3} - \mu_6 e^{-2(k-1)\pi i/3} t^{-2/3}/3 - \mu_5 t^{-1}/3 + O(|t|^{-4/3})$ ($k = 1, 2, 3$) are the three roots of $z^7 \nu(z) := \tilde{\nu}(z)$. Let $z = t^{-1/3} \xi$, the integral becomes

$$|t|^{2/3} \int_{\mathbb{C}} \frac{1}{|\xi \prod_{k=1}^3 (\xi - t^{1/3} z_k(t))|} i \text{d}\xi \text{d}\bar{\xi}.$$

As before, the linear terms cancel with each other. Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(C_0 + \text{Re}(C_1 \mu_5 t^{-2/3}) + \text{Re}(C_2 \mu_6^2 t^{-2/3}) + C_3 |\mu_6|^2 |t|^{-2/3} + O(r^{-1})) \frac{\text{d}r^2 + r^2 \text{d}\theta^2}{r^{1/3}},$$

which is asymptotic to a conic metric with cone angle $5\pi/3$.

The fiber $H^{-1}(t)$ can be written as

$$H^{-1}(t) = \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{\text{d}z}{z^4} \begin{pmatrix} \pm(-t)^{1/2} z^2 & z - \mu_6 z^2 - \mu_5 z^3 \\ 1 & \mp(-t)^{1/2} z^2 \end{pmatrix} \right. \right\} \\ \cup \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{\text{d}z}{z^4} \begin{pmatrix} a_0 & -tz^3 + (-\mu_5 + tc_0)z^2 + (-\mu_6 + \mu_5 c_0 - tc_0^2)z + \\ c_0 + z & 1 + \mu_6 c_0 - \mu_5 c_0^2 + tc_0^3 \\ & -a_0 \end{pmatrix}, a_0^2 = c_0 \tilde{\nu}(-c_0) \right. \right\},$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O} \oplus \mathcal{O}(-1)$. Therefore $H^{-1}(t)$ is the closure of the second set, which is the elliptic curve

$$a_0^2 = c_0 \tilde{\nu}(-c_0) = tc_0 \prod_{k=1}^3 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{z_2(t)(z_3(t) - z_1(t))}{z_3(t)(z_2(t) - z_1(t))} = e^{-\pi i/3} + \frac{e^{\pi i/6}}{3\sqrt{3}} (3\mu_5 + \mu_6^2) t^{-2/3} + O(|t|^{-4/3}), \\ \tau(t) = e^{2\pi i/3} + \frac{e^{-\pi i/3} \pi (3\mu_5 + \mu_6^2)}{12\sqrt{3} K(e^{-\pi i/6})^2} t^{-2/3} + O(|t|^{-4/3}).$$

Now $H^{-1}(t) \cong \mathbb{C} / c_t (\mathbb{Z} \oplus \tau(t)\mathbb{Z})$. Let $T_{\hat{\tau}}^2 = \mathbb{C} / \hat{c} (\mathbb{Z} \oplus \hat{\tau}\mathbb{Z})$, where $\hat{\tau} = e^{2\pi i/3}$, $\hat{c} = 2\pi / \sqrt{\text{Im } \hat{\tau}}$. Define $\mu_t : T_{\hat{\tau}}^2 \rightarrow H^{-1}(t)$ as

$$x + iy \mapsto \frac{c_t}{\hat{c}} \left(x + \frac{\text{Re}(\tau(t) - \hat{\tau})}{\text{Im } \hat{\tau}} y + i \frac{\text{Im } \tau(t)}{\text{Im } \hat{\tau}} y \right) \\ = \frac{c_t}{\hat{c}} (x + 2y(\text{Re } \tau(t) + 1/2)/\sqrt{3} + 2iy \text{Im } \tau(t)/\sqrt{3}).$$

Then we have

$$\begin{aligned}
\mu_t^*(g_{\text{sf}}|_{H^{-1}(t)}) &= \frac{c_t^2}{\hat{c}^2} \left((dx + 2(\operatorname{Re} \tau(t) + 1/2)/\sqrt{3} dy)^2 + (2\operatorname{Im} \tau(t)/\sqrt{3})^2 dy^2 \right) \\
&= dx^2 + dy^2 + \frac{2}{\sqrt{3}} \operatorname{Im} \left(\frac{e^{-\pi i/3} \pi (3\mu_5 + \mu_6^2)}{12\sqrt{3}K(e^{-\pi i/6})^2} t^{-2/3} \right) (-dx^2 + dy^2) \\
&\quad + \frac{4}{\sqrt{3}} \operatorname{Re} \left(\frac{e^{-\pi i/3} \pi (3\mu_5 + \mu_6^2)}{12\sqrt{3}K(e^{-\pi i/6})^2} t^{-2/3} \right) dx \otimes dy + O(|t|^{-4/3}).
\end{aligned}$$

6.3.3. *An untwisted order three pole and a simple pole.* Suppose these poles are located at 0 and ∞ respectively, then

$$\begin{aligned}
\mathcal{B} &= \left\{ \left(\sum_{k=4}^6 \frac{\mu_k}{z^k} + \frac{t}{z^3} + \frac{\mu_2}{z^2} \right) dz^2 \mid t \in \mathbb{C} \right\} \\
&= \left\{ \frac{1}{w^2} (\mu_6 w^4 + \mu_5 w^3 + \mu_4 w^2 + tw + \mu_2) dw^2 \mid t \in \mathbb{C} \right\}
\end{aligned}$$

where $w = 1/z$, $\mu_2, \mu_4, \mu_5, \mu_6$ are fixed and $\mu_6 \neq 0$, we may assume that $\mu_6 = -1$. Here we do not require that the simple pole is twisted, if it is, then $\mu_2 = 0$. Fix

$$\nu = \frac{1}{w^2} (-w^4 + \mu_5 w^3 + \mu_4 w^2 + tw + \mu_2) dw^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{w} dw^2 \in T_\nu \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$g_{\text{sK}}(\dot{\nu}, \dot{\nu}) = \int_C \frac{|\dot{\nu}|^2}{|\nu|} d\operatorname{vol}_C = |\dot{t}|^2 \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (w - w_k(t))|} i dw d\bar{w},$$

where $w_k(t) = t^{1/3} e^{2(k-1)\pi i/3} + 1 + O(|t|^{-1/3})$ ($k = 1, 2, 3$), $w_4(t) = \mu_2(-t^{-1} - \mu_2 \mu_4 t^{-3} + O(|t|^{-5}))$ are the four roots of $w^2 \nu(w) := \tilde{\nu}(w)$. Let $w = t^{1/3} \eta$, the integral becomes

$$\begin{aligned}
&|t|^{-2/3} \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^4 (\eta - t^{-1/3} w_k(t))|} i d\eta d\bar{\eta} \\
&= |t|^{-2/3} \left(\int_{\mathbb{C}} \frac{1}{|\eta| |\prod_{k=1}^3 (\eta - e^{i(k-1)\pi/2})|} i d\eta d\bar{\eta} + O(|t|^{-1/3}) \right) \\
&= |t|^{-2/3} (C_0 + \operatorname{Re}(C_1 t^{-1/3}) + O(|t|^{-2/3}))
\end{aligned}$$

Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(C_0 + \operatorname{Re}(C_1 t^{-1/3}) + O(r^{-2/3})) \frac{dr^2 + r^2 d\theta^2}{r^{2/3}},$$

which is asymptotic to a conic metric with cone angle $4\pi/3$.

The fiber $H^{-1}(t)$ can be written as

$$H^{-1}(t) = \left\{ \begin{aligned} & [(\bar{\partial}_E^1, \Phi)] \Big| \Phi = \frac{dw}{w} \begin{pmatrix} 0 & w^4 - \mu_5 w^3 - \mu_4 w^2 - tw - \mu_2 \\ 1 & 0 \end{pmatrix} \\ \cup & [(\bar{\partial}_E^0, \Phi)] \Big| \Phi = \frac{dw}{w} \begin{pmatrix} -\mu_5 w/2 + w^2 & -(\mu_5^2/4 + \mu_4)w^2 - tw - \mu_2 \\ 1 & \mu_5 w/2 - w^2 \end{pmatrix} \\ \cup & [(\bar{\partial}_E^0, \Phi)] \Big| \Phi = \frac{dw}{w} \begin{pmatrix} a_0 + w^2 & -\mu_5 w^2 + (c_0 \mu_5 - \mu_4 - 2a_0)w^- \\ c_0 + w & c_0^2 \mu_5 + c_0 \mu_4 + 2a_0 c_0 - t \\ & -a_0 - w^2 \end{pmatrix}, (a_0 + c_0^2)^2 = -\tilde{\nu}(-c_0) \end{aligned} \right\},$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, $(E, \bar{\partial}_E^1) \cong \mathcal{O} \oplus \mathcal{O}(-2)$. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$(a_0 + c_0^2)^2 = -\tilde{\nu}(-c_0) = \prod_{k=1}^4 (-c_0 - w_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{(w_1(t) - w_2(t))(w_3(t) - w_4(t))}{(w_3(t) - w_2(t))(w_1(t) - w_4(t))} = e^{-\pi i/3} + \sqrt{3}it^{-1/3} + O(|t|^{-2/3}),$$

$$\tau(t) = e^{2\pi i/3} + \frac{\sqrt{3}\pi}{4K(e^{-\pi/6})^2} t^{-1/3} + O(|t|^{-2/3}).$$

Define $\mu_t : T_{\tilde{\tau}}^2 \rightarrow H^{-1}(t)$ as above, then we have

$$\begin{aligned} \mu_t^*(g_{\text{sf}}|_{H^{-1}(t)}) &= dx^2 + dy^2 + \frac{2}{\sqrt{3}} \text{Im} \left(\frac{\sqrt{3}\pi}{4K(e^{-\pi/6})^2} t^{-1/3} \right) (-dx^2 + dy^2) \\ &+ \frac{4}{\sqrt{3}} \text{Re} \left(\frac{\sqrt{3}\pi}{4K(e^{-\pi/6})^2} t^{-1/3} \right) dx \otimes dy + O(|t|^{-2/3}). \end{aligned}$$

6.3.4. *A twisted order three pole and a simple pole.* Suppose these poles are located at 0 and ∞ respectively, then

$$\begin{aligned} \mathcal{B} &= \left\{ \left(\frac{\mu_5}{z^5} + \frac{\mu_4}{z^4} + \frac{t}{z^3} + \frac{\mu_2}{z^2} \right) dz^2 \Big| t \in \mathbb{C} \right\} \\ &= \left\{ \frac{1}{w^2} (\mu_5 w^3 + \mu_4 w^2 + tw + \mu_2) dw^2 \Big| t \in \mathbb{C} \right\} \end{aligned}$$

where $w = 1/z$, μ_2, μ_4, μ_5 are fixed and $\mu_5 \neq 0$, we may assume that $\mu_5 = -1$. Fix

$$\nu = \frac{1}{w^2} (-w^3 + \mu_4 w^2 + tw + \mu_2) dw^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{w} dw^2 \in T_{\nu} \mathcal{B}'.$$

Then g_{SK} on \mathcal{B}' is

$$g_{\text{SK}}(\dot{\nu}, \dot{\nu}) = \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = |\dot{t}|^2 \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^3 (w - w_k(t))|} i dw d\bar{w},$$

where $w_k(t) = (-1)^{k-1}(t^{1/2} + \mu_4^2 t^{-1/2}/8) + \mu_4/2 + O(|t|^{-1})$ ($k = 1, 2$), $w_3(t) = \mu_2(-t^{-1} - \mu_2 \mu_4 t^{-3} + O(|t|^{-4}))$ are the three roots of $w^2 \nu(w) := \tilde{\nu}(w)$. Let $w = t^{1/2} \eta$, the integral

becomes

$$\begin{aligned}
& |t|^{-1/2} \int_{\mathbb{C}} \frac{1}{|\prod_{k=1}^3 (\eta - t^{-1/2} w_k(t))|} i d\eta d\bar{\eta} \\
&= |t|^{-1/2} \left(\int_{\mathbb{C}} \frac{1}{|\eta(\eta-1)(\eta+1)|} i d\eta d\bar{\eta} + O(|t|^{-1/2}) \right) \\
&= |t|^{-1/2} (C_0 + \operatorname{Re}(C_1 \mu_4 t^{-1/2}) + O(|t|^{-1}))
\end{aligned}$$

Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(C_0 + \operatorname{Re}(C_1 \mu_4 t^{-1/2}) + O(r^{-1})) \frac{dr^2 + r^2 d\theta^2}{r^{1/2}},$$

which is asymptotic to a conic metric with cone angle $3\pi/2$.

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned}
H^{-1}(t) = & \left\{ [(\bar{\partial}_E^1, \Phi)] \left| \Phi = \frac{dw}{w} \begin{pmatrix} 0 & w^3 - \mu_4 w^2 - tw - \mu_2 \\ 1 & 0 \end{pmatrix} \right. \right\} \\
& \cup \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{dw}{w} \begin{pmatrix} a_0 & w^2 - (\mu_4 + c_0)w + \mu_4 c_0 + c_0^2 - t \\ c_0 + w & -a_0 \end{pmatrix}, a_0^2 = -\tilde{\nu}(-c_0) \right. \right\},
\end{aligned}$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, $(E, \bar{\partial}_E^1) \cong \mathcal{O} \oplus \mathcal{O}(-2)$. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$a_0^2 = -\tilde{\nu}(-c_0) = \prod_{k=1}^3 (-c_0 - w_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$\begin{aligned}
l(t) &= \frac{w_3(t) - w_2(t)}{w_1(t) - w_2(t)} = \frac{1}{2} - \mu_4 t^{-1/2}/4 + O(|t|^{-1}), \\
\tau(t) &= i + \frac{\mu_4 \pi i}{4K(\sqrt{1/2})^2} t^{-1/2} + O(|t|^{-1}).
\end{aligned}$$

Define $\mu_t : T_{\hat{\tau}}^2 \rightarrow H^{-1}(t)$ as above, then we have

$$\begin{aligned}
\mu_t^*(g_{\text{sf}}|_{H^{-1}(t)}) &= dx^2 + dy^2 + \operatorname{Im} \left(\frac{\mu_4 \pi i}{4K(\sqrt{1/2})^2} t^{-1/2} \right) (-dx^2 + dy^2) \\
&\quad + 2\operatorname{Re} \left(\frac{\mu_4 \pi i}{4K(\sqrt{1/2})^2} t^{-1/2} \right) dx \otimes dy + O(|t|^{-1}).
\end{aligned}$$

6.3.5. *Two untwisted order two poles.* Suppose these poles are located at 0 and ∞ respectively, then

$$\mathcal{B} = \left\{ \left(\frac{\mu_4}{z^4} + \frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} + \mu_0 \right) dz^2 \left| t \in \mathbb{C} \right. \right\},$$

where $\mu_0, \mu_1, \mu_3, \mu_4$ are fixed and $\mu_0, \mu_4 \neq 0$, we may assume that $\mu_4 = -1$. Fix

$$\nu = \left(\frac{-1}{z^4} + \frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} + \mu_0 \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z^2} dz^2 \in T_{\nu} \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$\begin{aligned} g_{\text{sK}}(\dot{\nu}, \dot{\nu}) &= \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = |\dot{t}|^2 \int_C \frac{1}{|\mu_0 \prod_{k=1}^4 (z - z_k(t))|} i \, dz d\bar{z} \\ &= \frac{|\dot{t}|^2}{|\mu_0|} \int_{|z| \leq 1} \frac{1}{|\prod_{k=1}^4 (z - z_k(t))|} i \, dz d\bar{z} + |\dot{t}|^2 \int_{|w| \leq 1} \frac{1}{|\prod_{k=1}^4 (w - z_k(t)^{-1})|} i \, dw d\bar{w}, \end{aligned}$$

where $z_k(t) = (-1)^{k-1} t^{-1/2} - \mu_3 t^{-1}/2 + O(|t|^{-3/2})$ ($k = 1, 2$), $z_k(t) = (-1)^{k-1} (-\mu_0)^{-1/2} t^{1/2} - \mu_1/2\mu_0 + O(|t|^{-1/2})$ ($k = 3, 4$) are the four roots of $z^4 \nu(z) =: \tilde{\nu}(z)$. In $\{|z| \leq 1\}$, $|z - z_k(t)| = |\mu_0^{-1/2} t^{1/2}| (1 + O(|t|^{-1/2}))$ for $k = 3, 4$, then the first integral is

$$\begin{aligned} & \int_{|z| \leq 1} \frac{|\mu_0 t^{-1}|}{|z - z_1(t)| |z - z_2(t)|} (1 + O(|t|^{-1/2})) i \, dz d\bar{z} \quad (\text{let } \xi = t^{1/2} z) \\ &= \int_{|\xi| \leq |t^{1/2}|} \frac{1}{|\xi - t^{1/2} z_1(t)| |\xi - t^{1/2} z_2(t)|} i \, d\xi d\bar{\xi} \cdot |\mu_0 t^{-1}| (1 + O(|t|^{-1/2})) \\ &= \int_{|\xi| \leq |t^{1/2}|} \frac{1}{|\xi - 1| |\xi + 1|} i \, d\xi d\bar{\xi} \cdot |\mu_0 t^{-1}| (1 + O(|t|^{-1/2})) \\ &= 2\pi |\mu_0| |t|^{-1} \log |t| + O(|t|^{-1}). \end{aligned}$$

Similarly, the second integral is

$$2\pi |t|^{-1} \log |t| + O(|t|^{-1}).$$

Therefore in polar coordinates, as $|a| = r \rightarrow \infty$, the special Kähler metric is

$$(4\pi \log r + O(1)) \frac{dr^2 + r^2 d\theta^2}{r}.$$

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned} H^{-1}(t) &= \left\{ [(\bar{\partial}_E^1, \Phi)] \left| \Phi = \frac{dz}{z^2} \begin{pmatrix} 0 & 1 - \mu_3 z - tz^2 - \mu_1 z^3 - \mu_0 z^4 \\ 1 & 0 \end{pmatrix} \right. \right\} \\ &\cup \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{dz}{z^2} \begin{pmatrix} -\frac{\mu_1 z}{2(-\mu_0)^{1/2}} + (-\mu_0)^{1/2} z^2 & -(t + \mu_1^2/4\mu_0) z^2 - \mu_3 z + 1 \\ 1 & \frac{\mu_1 z}{2(-\mu_0)^{1/2}} - (-\mu_0)^{1/2} z^2 \end{pmatrix} \right. \right\} \\ &\cup \left\{ [(\bar{\partial}_E^0, \Phi)] \left| \Phi = \frac{dz}{z^2} \begin{pmatrix} a_0 + (-\mu_0)^{1/2} z^2 & -\mu_1 z^2 + (\mu_1 c_0 - 2a_0 (-\mu_0)^{1/2} - t) z - \\ & \mu_1 c_0^2 + 2a_0 c_0 (-\mu_0)^{1/2} + t c_0 - \mu_3 \\ c_0 + z & -a_0 - (-\mu_0)^{1/2} z^2 \end{pmatrix}, \right. \right. \\ &\quad \left. \left. (a_0 + (-\mu_0)^{1/2} c_0^2)^2 = -\tilde{\nu}(-c_0) \right\}, \end{aligned}$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, $(E, \bar{\partial}_E^1) \cong \mathcal{O} \oplus \mathcal{O}(-2)$. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$(a_0 + (-\mu_0)^{1/2} c_0^2)^2 = -\tilde{\nu}(-c_0) = -\mu_0 \prod_{k=1}^4 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$\begin{aligned} l(t) &= \frac{(z_1(t) - z_2(t))(z_3(t) - z_4(t))}{(z_3(t) - z_2(t))(z_1(t) - z_4(t))} \\ &= 4(-\mu_0)^{1/2}t^{-1} + (8\mu_0 + (-\mu_0)^{-1/2}(\mu_1^2 - \mu_0\mu_3^2)/2)t^{-2} + O(|t|^{-3}), \\ \tau(t) &= \frac{i}{\pi} \log(4(-\mu_0)^{-1/2}t) + \frac{i}{\pi}(\mu_0^{-1}(\mu_1^2 - \mu_0\mu_3^2)/8)t^{-1} + O(|t|^{-2}). \end{aligned}$$

$H^{-1}(t) \cong \mathbb{C} / c_t(\mathbb{Z} \oplus \tau(t)\mathbb{Z})$ with Euclidean metric $dx^2 + dy^2$ and $c_t = 2\pi/\sqrt{\text{Im } \tau(t)}$.

6.3.6. *Two order two poles with one twisted.* Suppose the pole at 0 is untwisted and the pole at ∞ is twisted, then

$$\mathcal{B} = \left\{ \left(\frac{\mu_4}{z^4} + \frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} \right) dz^2 \middle| t \in \mathbb{C} \right\},$$

where μ_1, μ_3, μ_4 are fixed and $\mu_1, \mu_4 \neq 0$, we may assume that $\mu_4 = -1$. Fix

$$\nu = \left(\frac{\mu_4}{z^4} + \frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z^2} dz^2 \in T_\nu \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$\begin{aligned} g_{\text{sK}}(\dot{\nu}, \dot{\nu}) &= \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = \frac{|\dot{t}|^2}{|\mu_1|} \int_C \frac{1}{|\prod_{k=1}^3 (z - z_k(t))|} i dz d\bar{z} \\ &= \frac{|\dot{t}|^2}{|\mu_1|} \int_{|z| \leq 1} \frac{1}{|\prod_{k=1}^3 (z - z_k(t))|} i dz d\bar{z} + |\dot{t}|^2 \int_{|w| \leq 1} \frac{1}{|w| |\prod_{k=1}^3 (w - z_k(t)^{-1})|} i dw d\bar{w}, \end{aligned}$$

where $z_k(t) = (-1)^{k-1}t^{-1/2} - \mu_3 t^{-1}/2 + O(|t|^{-3/2})$ ($k = 1, 2$), $z_3(t) = -t/\mu_1 + \mu_3 t^{-1} + O(|t|^{-2})$ are the three roots of $z^4 \nu(z) =: \tilde{\nu}(z)$. In $\{|z| \leq 1\}$, $|z - z_3(t)| = |t/\mu_1|(1 + O(|t|^{-1}))$, then the first integral is

$$\begin{aligned} &\int_{|z| \leq 1} \frac{|\mu_1 t^{-1}|}{|z - z_1(t)||z - z_2(t)|} (1 + O(|t|^{-1})) i dz d\bar{z} \\ &= 2\pi |\mu_1| |t|^{-1} \log |t| + O(|t|^{-1}), \end{aligned}$$

as before. Similarly, the second integral is $4\pi |t|^{-1} \log |t| + O(|t|^{-1})$. Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(6\pi \log r + O(1)) \frac{dr^2 + r^2 d\theta^2}{r}.$$

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned} H^{-1}(t) &= \left\{ [(\bar{\partial}_E^1, \Phi)] \middle| \Phi = \frac{dz}{z^2} \begin{pmatrix} 0 & 1 - \mu_3 z - tz^2 - \mu_1 z^3 \\ 1 & 0 \end{pmatrix} \right\} \\ &\cup \left\{ [(\bar{\partial}_E^0, \Phi)] \middle| \Phi = \frac{dz}{z^2} \begin{pmatrix} a_0 & -\mu_1 z^2 + (\mu_1 c_0 - t)z \\ c_0 + z & \mu_3 + tc_0 - \mu_1 c_0^2 \\ & -a_0 \end{pmatrix}, a_0^2 = -\tilde{\nu}(-c_0) \right\}, \end{aligned}$$

where $\bar{\partial}_E^0, \bar{\partial}_E^1$ are holomorphic structures as before. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$a_0^2 = -\tilde{\nu}(-c_0) = -\mu_1 \prod_{k=1}^3 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{z_1(t) - z_2(t)}{z_1(t) - z_3(t)} = 2\mu_1 t^{-3/2} + \mu_1 \mu_3^2 t^{-5/2}/4 + O(|t|^{-3}),$$

$$\tau(t) = \frac{i}{\pi} (\log(8\mu_1^{-1} t^{3/2}) - \mu_3^2 t^{-1}/8 - \mu_1 t^{-3/2}) + O(|t|^{-2}).$$

6.3.7. *Two twisted order two poles.* Suppose the poles are 0 and ∞ , then

$$\mathcal{B} = \left\{ \left(\frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} \right) dz^2 \middle| t \in \mathbb{C} \right\},$$

where $\mu_1, \mu_3 \neq 0$ are fixed, we may assume that $\mu_3 = -1$. Fix

$$\nu = \left(\frac{\mu_3}{z^3} + \frac{t}{z^2} + \frac{\mu_1}{z} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z^2} dz^2 \in T_\nu \mathcal{B}'.$$

By a similar computation as before, g_{sK} on \mathcal{B}' is

$$\begin{aligned} g_{\text{sK}}(\dot{\nu}, \dot{\nu}) &= \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = \frac{|\dot{t}|^2}{|\mu_1|} \int_{\mathbb{C}} \frac{1}{|z(z - z_1(t))(z - z_2(t))|} i dz d\bar{z} \\ &= \frac{|\dot{t}|^2}{|\mu_1|} \int_{|z| \leq 1} \frac{1}{|z||z - z_1(t)||z - z_2(t)|} i dz d\bar{z} \\ &\quad + |\dot{t}|^2 \int_{|w| \leq 1} \frac{1}{|w||w - z_1(t)^{-1}||w - z_2(t)^{-1}|} i dw d\bar{w} \\ &= |\dot{t}|^2 (8\pi |t|^{-1} \log |t| + O(|t|^{-1})). \end{aligned}$$

Here $z_1(t) = t^{-1} - \mu_1 t^{-3} + O(|t|^{-5})$, $z_2(t) = -t/\mu_1 - t^{-1} + O(|t|^{-3})$ are the two roots of $z^3 \nu(z) := \tilde{\nu}(z)$. Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(8\pi \log r + O(1)) \frac{dr^2 + r^2 d\theta^2}{r}.$$

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned} H^{-1}(t) &= \left\{ [(\bar{\partial}_E^1, \Phi)] \middle| \Phi = \frac{dz}{z^2} \begin{pmatrix} 0 & z(1 - tz + \mu_1 z^2) \\ 1 & 0 \end{pmatrix} \right\} \\ \cup &\left\{ [(\bar{\partial}_E^0, \Phi)] \middle| \Phi = \frac{dz}{z^2} \begin{pmatrix} a_0 & -\mu_1 z^2 + (\mu_1 c_0 - a)z + \\ & 1 + tc_0 - \mu_1 c_0^2 \\ c_0 + z & -a_0 \end{pmatrix}, a_0^2 = c_0 \tilde{\nu}(-c_0) \right\}, \end{aligned}$$

where $\bar{\partial}_E^0, \bar{\partial}_E^1$ are as above. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$a_0^2 = c_0 \tilde{\nu}(-c_0) = \mu_1 c_0 (c_0 + z_1(t))(c_0 + z_2(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{z_1(t) - 0}{z_1(t) - z_2(t)} = \mu_1 t^{-2} - 3\mu_1^2 t^{-4} + O(|t|^{-6}),$$

$$\tau(t) = \frac{i}{\pi} (\log(16\mu_1^{-1} t^2) + 5\mu_1 t^{-2}/2) + O(|t|^{-4}).$$

6.3.8. *An untwisted order two pole and two simple poles.* Suppose the two simple poles are 0 and 1, the order two pole is ∞ , then

$$\mathcal{B} = \left\{ \left(\frac{\mu_0}{z^2} + \frac{\mu_1}{(z-1)^2} + \frac{\mu_3}{z} + \mu_4 + \frac{t}{z(z-1)} \right) dz^2 \mid t \in \mathbb{C} \right\},$$

where $\mu_0, \mu_1, \mu_3, \mu_4$ are fixed, $\mu_4 \neq 0$. Fix

$$\nu = \left(\frac{\mu_0}{z^2} + \frac{\mu_1}{(z-1)^2} + \frac{\mu_3}{z} + \mu_4 + \frac{t}{z(z-1)} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z(z-1)} dz^2 \in T_\nu \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$\begin{aligned} g_{\text{sK}}(\dot{\nu}, \dot{\nu}) &= \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = |\dot{t}|^2 \int_{\mathbb{C}} \frac{1}{|\mu_4 \prod_{k=1}^4 (z - z_k(t))|} i dz d\bar{z} \\ &= \frac{|\dot{t}|^2}{|\mu_4|} \int_{|z| \leq 2} + \int_{2 \leq |z| \leq |\mu_4|^{-1/2} |t|^{1/2}/2} \frac{1}{|\prod_{k=1}^4 (z - z_k(t))|} i dz d\bar{z} \\ &\quad + \frac{|\dot{t}|^2}{|\mu_0|} \int_{|w| \leq 2|\mu_4|^{1/2} |a|^{-1/2}} \frac{1}{|\prod_{k=1}^4 (w - z_k(t)^{-1})|} i dw d\bar{w} \\ &= |\dot{t}|^2 (O(1/|t|) + 2\pi |t|^{-1} \log |t| + O(1/|t|)) = 2\pi |\dot{t}|^2 |t|^{-1} (\log |t| + O(1)). \end{aligned}$$

Here $z_1(t) = \mu_0 t^{-1} + \mu_0(\mu_3 - \mu_0)t^{-2} + O(|t|^{-3})$, $z_2(t) = 1 - \mu_1 t^{-1} + \mu_1^2 t^{-2} + O(|t|^{-3})$, $z_k(t) = (-1)^{k-1} (-\mu_4)^{-1/2} t^{1/2} + (1 - \mu_3/\mu_4)/2 + O(|t|^{-1/2})$ ($k = 3, 4$) are the four roots of $z^2(z-1)^2\nu(z) := \tilde{\nu}(z)$. Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(2\pi \log r + O(1)) \frac{dr^2 + r^2 d\theta^2}{r}.$$

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned} H^{-1}(t) &= \left\{ [(\bar{\partial}_E^0, \Phi)] \mid \Phi = \frac{dz}{z(z-1)} \begin{pmatrix} \pm(-\mu_4)^{1/2} z^2 & \mu_4 z^4 - \tilde{\nu}(z) \\ 1 & \mp(-\mu_4)^{1/2} z^2 \end{pmatrix} \right\} \\ &\cup \left\{ [(\bar{\partial}_E^0, \Phi)] \mid \Phi = \frac{dz}{z(z-1)} \begin{pmatrix} a_0 & \begin{matrix} -\mu_4 z^3 + (\mu_4 c_0 + 2\mu_4 - \mu_3) z^2 + \\ (2\mu_3 - \mu_4 - \mu_1 - t - \mu_4 c_0^2 - 2\mu_4 c_0 + \mu_3 c_0) z + \\ 2\mu_0 + t - \mu_3 + (\mu_4 - 2\mu_3 + \mu_1 + t) c_0 + \\ (2\mu_4 - \mu_3) c_0^2 + \mu_4 c_0^3 \end{matrix} \\ c_0 + z & -a_0 \end{pmatrix}, a_0^2 = -\tilde{\nu}(-c_0) \right\}, \end{aligned}$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O}(-1) \oplus \mathcal{O}(-2)$. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$a_0^2 = -\tilde{\nu}(-c_0) = -\mu_4 \prod_{k=1}^4 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{(z_1(t) - z_2(t))(z_3(t) - z_4(t))}{(z_3(t) - z_2(t))(z_1(t) - z_4(t))} = -2(-\mu_4)^{1/2}t^{-1/2} + 2\mu_4t^{-1} + O(|t|^{-3/2}),$$

$$\tau(t) = \frac{i}{\pi} \log(-8(-\mu_4)^{-1/2}t^{1/2}) + O(|t|^{-1}).$$

6.3.9. *A twisted order two pole and two simple poles.* Let $0, 1$ be the two simple poles and ∞ be the order two pole, then

$$\mathcal{B} = \left\{ \left(\frac{\mu_0}{z^2} + \frac{\mu_1}{(z-1)^2} + \frac{\mu_3}{z} + \frac{t}{z(z-1)} \right) dz^2 \mid t \in \mathbb{C} \right\},$$

where μ_0, μ_1, μ_3 are fixed, $\mu_3 \neq 0$. Fix

$$\nu = \left(\frac{\mu_0}{z^2} + \frac{\mu_1}{(z-1)^2} + \frac{\mu_3}{z} + \frac{t}{z(z-1)} \right) dz^2 \in \mathcal{B}', \quad \dot{\nu} = \frac{\dot{t}}{z(z-1)} dz^2 \in T_\nu \mathcal{B}'.$$

Then g_{sK} on \mathcal{B}' is

$$\begin{aligned} g_{\text{sK}}(\dot{\nu}, \dot{\nu}) &= \int_C \frac{|\dot{\nu}|^2}{|\nu|} \text{dvol}_C = |\dot{t}|^2 \int_{\mathbb{C}} \frac{1}{|\mu_3 \prod_{k=1}^3 (z - z_k(t))|} i dz d\bar{z} \\ &= \frac{|\dot{t}|^2}{|\mu_3|} \int_{|z| \leq 2} + \int_{2 \leq |z| \leq |t|/(2|\mu_3|)} \frac{1}{|\prod_{k=1}^3 (z - z_k(a))|} i dz d\bar{z} \\ &+ \frac{|\dot{t}|^2}{|\mu_0|} \int_{|w| \leq 2|\mu_3|/|t|} \frac{1}{|w| |\prod_{k=1}^3 (w - z_k(t)^{-1})|} i dw d\bar{w} \\ &= |\dot{t}|^2 (O(1/|t|) + 4\pi |t|^{-1} \log |t| + O(1/|t|)) = 4\pi |\dot{t}|^2 |t|^{-1} (\log |t| + O(1)). \end{aligned}$$

Here $z_1(t) = \mu_0 t^{-1} + \mu_0(\mu_3 - \mu_0)t^{-2} + O(|t|^{-3})$, $z_2(t) = 1 - \mu_1 t^{-1} + \mu_1^2 t^{-2} + O(|t|^{-3})$, $z_3(t) = -t/\mu_3 + (1 - (\mu_0 + \mu_1)/\mu_3) + O(|t|^{-1})$ are the three roots of $z^2(z-1)^2\nu(z) := \tilde{\nu}(z)$. Therefore in polar coordinates, as $|t| = r \rightarrow \infty$, the special Kähler metric is

$$(4\pi \log r + O(1)) \frac{dr^2 + r^2 d\theta^2}{r}.$$

The fiber $H^{-1}(t)$ can be written as

$$\begin{aligned} H^{-1}(t) &= \left\{ [(\bar{\partial}_E^0, \Phi)] \mid \Phi = \frac{dz}{z(z-1)} \begin{pmatrix} 0 & -\tilde{\nu}(z) \\ 1 & 0 \end{pmatrix} \right\} \\ &\cup \left\{ [(\bar{\partial}_E^0, \Phi)] \mid \Phi = \frac{dz}{z(z-1)} \begin{pmatrix} a_0 & -\mu_3 z^2 + (\mu_3 c_0 + 2\mu_3 - \mu_0 - \mu_1 - t)z + \\ & t + 2\mu_0 - \mu_3 + \\ & (t + \mu_0 + \mu_1 - 2\mu_3)c_0 - \mu_3 c_0^2 \\ c_0 + z & -a_0 \end{pmatrix}, a_0^2 = -\tilde{\nu}(-c_0) \right\}, \end{aligned}$$

where $(E, \bar{\partial}_E^0) \cong \mathcal{O}(-1) \oplus \mathcal{O}(-2)$. $H^{-1}(t)$ is the closure of the last set, which is the elliptic curve

$$a_0^2 = -\tilde{\nu}(-c_0) = -\mu_3 \prod_{k=1}^3 (-c_0 - z_k(t)), \quad (a_0, c_0) \in \mathbb{C}^2,$$

with modulus $\tau(t) = \lambda^{-1}(l(t))$, where

$$l(t) = \frac{(z_1(t) - z_2(t))}{(z_1(t) - z_3(t))} = -\mu_3 t^{-1} - \mu_3 t^{-2} + O(|t|^{-3}),$$

$$\tau(t) = \frac{i}{\pi} (\log(-16\mu_3^{-1}t) + (\mu_3/2 - 1)t^{-1}) + O(|t|^{-2}).$$

6.4. The Uniformness. In Theorem 1.1, the constants in the decay rate of $g_{L^2} - g_{\text{sf}}$ depend on the choice of the curve. Now we aim to find constants uniform in these choices for a four-dimensional moduli space \mathcal{M} . For simplicity, here we only consider the moduli space in Section 6.3.1.

As in (57), $t \in \mathbb{C}$ parametrizes the quadratic differential ν_t in the Hitchin base, and is no longer a positive real number as in the previous sections. Let $(\bar{\partial}_E, \Phi_t)$ be any Higgs bundle in the Hitchin fiber $H^{-1}(t)$. By Lemma 3.1, the four roots of $\tilde{\nu}_t(z)$ have asymptotics

$$z_k(t) = t^{-1/4} e^{i(k-1)\pi/2} + O(|t|^{-1/2}),$$

as $|t| \rightarrow \infty$. The local mass at $z_k(t)$ ($k = 1, 2, 3, 4$) is

$$\lambda_k(t) = \left| t \prod_{j \neq k} (z_k(t) - z_j(t)) z_k(t)^{-8} \right|^{1/2} = 4t^{9/4} (1 + O(|t|^{-1/4})).$$

Here and below, the constants are independent of $\text{Arg}(t)$ and the choice of $(\bar{\partial}_E, \Phi_t)$ in $H^{-1}(t)$. Proposition 3.3 still hold on disks $\tilde{B}_{k,t} := \{ |\zeta_{k,t}| < \kappa |t|^{-1/4} \}$, where $\zeta_{k,t}$ is the local holomorphic coordinate centered at $z_k(t)$ such that $q_t = -\lambda_k(t)^2 \zeta_{k,t} d\zeta_{k,t}^2$, and κ is independent of the above choices. One only need to check that the gauge transformations in the proof is well-defined on the whole disk $\tilde{B}_{k,t}$ for suitable κ . This is clear for Φ_t in the first subset of (58). For Φ_t in the second subset, now a_0, c_0 may depend on t . Suppose $|c_0 + z_k(t)| \geq |t|^{-1/4}/2$, the first gauge transformation is $\begin{pmatrix} a_0 & 1 \\ z_k(t) + c_0 & 0 \end{pmatrix}$, which is constant and invertible over $\tilde{B}_{k,t}$. The second gauge transformation is well-defined since

$$\left| \frac{z + c_0}{z_k(t) + c_0} - 1 \right| = \left| \frac{z - z_k(t)}{z_k(t)} \right| \left| \frac{z_k(t)}{c_0 + z_k(t)} \right| \leq c\kappa \leq 1/2,$$

for κ small. The rest of the proof remains the same. If $|c_0 + z_k(t)| \leq |t|^{-1/4}/2$ for some k , then $|c_0 + z_j(t)| \geq |t|^{-1/4}/2$ for $j \neq k$ and $|t|$ large, and the previous discussion applies for the disks $\tilde{B}_{j,t}$. On $\tilde{B}_{k,t}$,

$$\Phi_t = \frac{dz}{z^4} \begin{pmatrix} a_0 & -tz^3 + (-\mu_5 + tc_0)z^2 + (-\mu_6 + \mu_5 c_0 - tc_0^2)z - \\ & \mu_7 + \mu_6 c_0 - \mu_5 c_0^2 + tc_0^3 \\ c_0 + z & -a_0 \end{pmatrix} := \begin{pmatrix} a_t(\zeta_{k,t}) & b_t(\zeta_{k,t}) \\ c_t(\zeta_{k,t}) & -a_t(\zeta_{k,t}) \end{pmatrix} d\zeta_{k,t},$$

where

$$b_t(\zeta_{k,t}) = \frac{dz}{d\zeta_{k,t}}(\zeta_{k,t}) \frac{-t}{z^4} (z^3 - c_0 z^2 + c_0^2 z - c_0^3) (1 + O(|t|^{-1/4}))$$

$$= \frac{dz}{d\zeta_{k,t}}(\zeta_{k,t}) \frac{-t}{z^4} (z - c_0)(z^2 + c_0^2) (1 + O(|t|^{-1/4})).$$

Then $\sqrt{b_t(\zeta_{k,t})}$ is well-defined on $\tilde{B}_{k,t}$ by the assumption that $|c_0 + z_k(t)| \leq |t|^{-1/4}/2$, and we have

$$g^{-1}\Phi_t g = \begin{pmatrix} 0 & 1 \\ \lambda_k(t)^2 \zeta_{k,t} & 0 \end{pmatrix} d\zeta_{k,t}, \text{ for } g = \frac{1}{\sqrt{b_t(\zeta_{k,t})}} \begin{pmatrix} b_t(\zeta_{k,t}) & 0 \\ -a_t(\zeta_{k,t}) & 1 \end{pmatrix}.$$

The remaining proof is the same. The approximate metric h_t^{app} is constructed as in Definition 3.4, and the proof of the error estimate in (17) works uniformly. Then there are uniform constants t_0, c, c' , such that for any $|t| \geq t_0$,

$$\left\| F_{h_t^{\text{app}}} + \left[\Phi_t, \Phi_t^{*h_t^{\text{app}}} \right] \right\|_{L^2} \leq ce^{-c'|t|^{3/4}}.$$

The computations in Sections 4, 5, 6 only use the expression of $q_t = \det \Phi_t$ and the local forms of Φ_t provided by Proposition 3.3, so the estimates remain valid uniformly for the choice of Φ_t in $H^{-1}(t)$. Moreover, the constants can also be chosen to be independent of $\text{Arg}(t)$. Therefore, we find uniform constants t_0, c, c' so that for $[(\dot{\eta}, \dot{\Phi})] \in T_{[(\bar{\partial}_E, \Phi_t)]} \mathcal{M}$ and $|t| \geq t_0$,

$$\left| \left\| [(\dot{\eta}, \dot{\Phi})] \right\|_{g_{L^2}} / \left\| [(\dot{\eta}, \dot{\Phi})] \right\|_{g_{\text{sf}}} - 1 \right| \leq ce^{-c'|t|^{3/4}}.$$

REFERENCES

- [BB04] Olivier Biquard and Philip Boalch, *Wild non-abelian Hodge theory on curves*, Compos. Math. **140** (2004), no. 1, 179–204. MR 2004129
- [Biq97] Olivier Biquard, *Fibrés de Higgs et connexions intégrables: le cas logarithmique (diviseur lisse)*, Ann. Sci. École Norm. Sup. (4) **30** (1997), no. 1, 41–96. MR 1422313
- [Boa01] Philip Boalch, *Symplectic manifolds and isomonodromic deformations*, Adv. Math. **163** (2001), no. 2, 137–205. MR 1864833
- [Boa12] Philip Boalch, *Hyperkahler manifolds and nonabelian Hodge theory of (irregular) curves*, arXiv e-prints (2012), arXiv:1203.6607.
- [Boa14] P. P. Boalch, *Geometry and braiding of Stokes data; fission and wild character varieties*, Ann. of Math. (2) **179** (2014), no. 1, 301–365. MR 3126570
- [Boa18] Philip Boalch, *Wild character varieties, meromorphic Hitchin systems and Dynkin diagrams*, Geometry and physics. Vol. II, Oxford Univ. Press, Oxford, 2018, pp. 433–454. MR 3931781
- [CC21] Gao Chen and Xiuxiong Chen, *Gravitational instantons with faster than quadratic curvature decay (III)*, Math. Ann. **380** (2021), no. 1-2, 687–717. MR 4263695
- [CV21] Gao Chen and Jeff Viaclovsky, *Gravitational instantons with quadratic volume growth*, arXiv e-prints (2021), arXiv:2110.06498.
- [DN19] David Dumas and Andrew Neitzke, *Asymptotics of Hitchin’s metric on the Hitchin section*, Comm. Math. Phys. **367** (2019), no. 1, 127–150. MR 3933406
- [FMSW22] Laura Fredrickson, Rafe Mazzeo, Jan Swoboda, and Hartmut Weiss, *Asymptotic geometry of the moduli space of parabolic $SL(2, \mathbb{C})$ -higgs bundles*, Journal of the London Mathematical Society (2022).
- [FN21] Laura Fredrickson and Andrew Neitzke, *Moduli of wild Higgs bundles on $\mathbb{C}P^1$ with \mathbb{C}^\times -actions*, Math. Proc. Cambridge Philos. Soc. **171** (2021), no. 3, 623–656. MR 4324961
- [Fos17] Lorenzo Foscolo, *A gluing construction for periodic monopoles*, Int. Math. Res. Not. IMRN (2017), no. 24, 7504–7550. MR 3801425
- [Fre99] Daniel S. Freed, *Special Kähler manifolds*, Comm. Math. Phys. **203** (1999), no. 1, 31–52. MR 1695113
- [Fre18] Laura Fredrickson, *Generic Ends of the Moduli Space of $SL(n, \mathbb{C})$ -Higgs Bundles*, arXiv e-prints (2018), arXiv:1810.01556.
- [Fre20] Laura Fredrickson, *Exponential decay for the asymptotic geometry of the Hitchin metric*, Comm. Math. Phys. **375** (2020), no. 2, 1393–1426. MR 4083884

- [GMN10] Davide Gaiotto, Gregory W. Moore, and Andrew Neitzke, *Four-dimensional wall-crossing via three-dimensional field theory*, *Comm. Math. Phys.* **299** (2010), no. 1, 163–224. MR 2672801
- [Hau01] Tamas Hausel, *Geometry of the moduli space of Higgs bundles*, arXiv e-prints (2001), arXiv:0107040.
- [Ina16] Michi-aki Inaba, *Moduli space of irregular singular parabolic connections of generic ramified type on a smooth projective curve*, arXiv e-prints (2016), arXiv:1606.02369.
- [LM10] Marina Logares and Johan Martens, *Moduli of parabolic Higgs bundles and Atiyah algebroids*, *J. Reine Angew. Math.* **649** (2010), 89–116. MR 2746468
- [MSWW16] Rafe Mazzeo, Jan Swoboda, Hartmut Weiss, and Frederik Witt, *Ends of the moduli space of Higgs bundles*, *Duke Math. J.* **165** (2016), no. 12, 2227–2271. MR 3544281
- [MSWW19] ———, *Asymptotic geometry of the Hitchin metric*, *Comm. Math. Phys.* **367** (2019), no. 1, 151–191. MR 3933407
- [MTW77] Barry M. McCoy, Craig A. Tracy, and Tai Tsun Wu, *Painlevé functions of the third kind*, *J. Mathematical Phys.* **18** (1977), no. 5, 1058–1092. MR 473322
- [Sim90] Carlos T. Simpson, *Harmonic bundles on noncompact curves*, *J. Amer. Math. Soc.* **3** (1990), no. 3, 713–770. MR 1040197
- [Str84] Kurt Strebel, *Quadratic differentials*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, vol. 5, Springer-Verlag, Berlin, 1984. MR 743423
- [Wit08] Edward Witten, *Gauge theory and wild ramification*, *Anal. Appl. (Singap.)* **6** (2008), no. 4, 429–501. MR 2459120

INSTITUTE OF GEOMETRY AND PHYSICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA,
SHANGHAI, CHINA, 201315

Email address: chengao1@ustc.edu.cn

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI 53706, USA

Email address: nli62@wisc.edu