

ON CODIMENSION ONE HOLOMORPHIC DISTRIBUTIONS ON COMPACT TORIC ORBIFOLDS

MIGUEL RODRÍGUEZ PEÑA

ABSTRACT. The number of singularities, counted with multiplicity, of a generic codimension one holomorphic distribution on a compact toric orbifold is determined. As a consequence, we give the classification of regular distributions on rational normal scrolls and weighted projective spaces. Under certain conditions, we also prove that the singular set of a codimension one holomorphic foliation on a toric orbifold admits at least one irreducible component of codimension two, and we give a Darboux-Jouanolou type integrability theorem for codimension one holomorphic foliations. We illustrate our results with several examples.

1. INTRODUCTION

In this paper, we are interested in the study of codimension one holomorphic distributions on compact toric orbifolds, i.e. on complete simplicial toric varieties, for this we will consider homogeneous coordinates [13]. In Section 2, we start with some preliminaries on toric varieties and orbifolds. In Section 3, after defining codimension one holomorphic distributions and foliations, we present two propositions in analogy with results well-known on the complex projective spaces: in Proposition 3.1, we give a condition for the singular set of a codimension one holomorphic foliation admits at least a codimension two irreducible component (in the case of the complex projective space, this result is due to Jouanolou [21]), and in Proposition 3.2, we give a Darboux-Jouanolou type integrability theorem for codimension one holomorphic foliations, see also [8]. Later, we present two Bott-type formulas for the number of singularities counted with multiplicity of a generic distribution, i.e. when the distribution has at most isolated singularities; see for instance Proposition 3.3 and Proposition 3.5. To end this section we present some examples where our results are applied. Finally, in Section 4, we give some applications; we impose conditions for a distribution to be regular, then we present some examples and applications, among them we give a classification of regular distributions on rational normal scrolls (see Theorem 4.1 and Corollary 4.4), and on weighted projective spaces (see Corollary 4.5).

2. PRELIMINARIES

In this section we recall some basic definitions and results about simplicial complete toric varieties. For more details about toric varieties see [5], [12], [13], [17] and [24].

Let N be a free \mathbb{Z} -module of rank n and $M = \text{Hom}(N, \mathbb{Z})$ be its dual. A subset $\sigma \subset N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R} \simeq \mathbb{R}^n$ is called a strongly convex rational polyhedral cone if there exists a finite number of

elements v_1, \dots, v_k in the lattice N such that

$$\sigma = \{a_1 v_1 + \dots + a_k v_k : a_i \in \mathbb{R}, a_i \geq 0\},$$

and σ does not contain any straight line going through the origin, i.e. $\sigma \cap (-\sigma) = \{0\}$. A cone σ is called simplicial if its generators can be chosen to be linearly independent over \mathbb{R} . The dimension of a cone σ is, by definition, the dimension of a minimal subspace of \mathbb{R}^n containing σ .

Set $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ and $\langle \cdot, \cdot \rangle : M_{\mathbb{R}} \times N_{\mathbb{R}} \rightarrow \mathbb{R}$ the dual pairing. To each cone σ we associate the dual cone $\check{\sigma}$

$$\check{\sigma} = \{m \in M_{\mathbb{R}} : \langle m, v \rangle \geq 0 \ \forall v \in \sigma\},$$

which is a rational polyhedral cone in $M_{\mathbb{R}}$. It follows from Gordan's Lemma that $\check{\sigma} \cap M$ is also a finitely generated semigroup. A subset τ of σ is called a face and is denoted $\tau \prec \sigma$, if

$$\tau = \sigma \cap \{m\}^{\perp} = \{v \in \sigma : \langle m, v \rangle = 0\},$$

for some $m \in \check{\sigma}$. A cone is a face of itself, other faces are called proper faces.

Definition 2.1. A non-empty collection $\Delta = \{\sigma_1, \dots, \sigma_s\}$ of strongly convex rational polyhedral cones in $N_{\mathbb{R}} \simeq \mathbb{R}^n$ is called a fan if it satisfies:

- (i) if $\sigma \in \Delta$ and $\tau \prec \sigma$, then $\tau \in \Delta$,
- (ii) if $\sigma_i, \sigma_j \in \Delta$, then $\sigma_i \cap \sigma_j \prec \sigma_i$ and $\sigma_i \cap \sigma_j \prec \sigma_j$.

The fan Δ is called complete if $N_{\mathbb{R}} = \sigma_1 \cup \dots \cup \sigma_s$. The dimension of a fan is the maximal dimension of its cones. An n -dimensional complete fan is called simplicial if all its n -dimensional cones are simplicial. An affine n -dimensional toric variety corresponding to σ is the variety

$$\mathcal{U}_{\sigma} = \text{Spec } \mathbb{C}[\check{\sigma} \cap M].$$

If a cone τ is a face of σ then $\check{\tau} \cap M$ is a subsemigroup of $\check{\sigma} \cap M$, hence \mathcal{U}_{τ} is embedded into \mathcal{U}_{σ} as an open subset. The affine varieties corresponding to all cones of the fan Δ are glued together according to this rule into the toric variety \mathbb{P}_{Δ} associated with Δ . It is possible to show that a toric variety \mathbb{P}_{Δ} contains a complex torus $\mathbb{T}^n = (\mathbb{C}^*)^n$ as a Zariski open subset such that the action of \mathbb{T}^n on itself extends to an action of \mathbb{T}^n on \mathbb{P}_{Δ} . A toric variety \mathbb{P}_{Δ} determined by a complete simplicial fan Δ is a compact complex orbifold, i.e. a compact complex variety with at most quotient singularities. Note that \mathbb{T}^n , \mathbb{C}^n and \mathbb{P}^n are toric varieties.

2.1. The homogeneous coordinate ring. Let \mathbb{P}_{Δ} be the toric variety determined by a fan Δ in $N \simeq \mathbb{Z}^n$. The one-dimensional cones of Δ form the set $\Delta(1)$, and given $\rho \in \Delta(1)$, we set n_{ρ} the unique generator of $\rho \cap N$. If σ is any cone in Δ , then $\sigma(1) = \{\rho \in \Delta(1) : \rho \prec \sigma\}$ is the set of one-dimensional faces of σ . We will assume that $\Delta(1)$ spans $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R} \simeq \mathbb{R}^n$.

Each $\rho \in \Delta(1)$ corresponds to an irreducible \mathbb{T} -invariant Weil divisor D_{ρ} in \mathbb{P}_{Δ} , where $\mathbb{T} = N \otimes_{\mathbb{Z}} \mathbb{C}^* \simeq \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}^*)$ is the torus acting on \mathbb{P}_{Δ} . The \mathbb{T} -invariant Weil divisors on \mathbb{P}_{Δ} form a free abelian group of rank $|\Delta(1)|$, that will be denoted $\mathbb{Z}^{\Delta(1)}$. Thus an element $D \in \mathbb{Z}^{\Delta(1)}$ is a sum $D = \sum_{\rho} a_{\rho} D_{\rho}$. The \mathbb{T} -invariant Cartier divisors form a subgroup $\text{Div}_{\mathbb{T}}(\mathbb{P}_{\Delta}) \subset \mathbb{Z}^{\Delta(1)}$. The Weil divisor class group of \mathbb{P}_{Δ} is denoted by $\mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$.

Each $m \in M$ gives a character $\chi^m : \mathbb{T} \rightarrow \mathbb{C}^*$, hence χ^m extends to a rational function on \mathbb{P}_{Δ} . As is well-known, χ^m gives the \mathbb{T} -invariant Cartier divisor $\text{div}(\chi^m) = -\sum_{\rho} \langle m, n_{\rho} \rangle D_{\rho}$. We will

consider the map

$$\begin{aligned} M &\longrightarrow \mathbb{Z}^{\Delta(1)} \\ m &\longmapsto D_m = \sum_{\rho} \langle m, n_{\rho} \rangle D_{\rho}. \end{aligned}$$

This map is injective since $\Delta(1)$ spans $N_{\mathbb{R}}$. By [17], we have a commutative diagram

$$(1) \quad \begin{array}{ccccccc} 0 & \rightarrow & M & \rightarrow & \text{Div}_{\mathbb{T}}(\mathbb{P}_{\Delta}) & \rightarrow & \text{Pic}(\mathbb{P}_{\Delta}) & \rightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & M & \rightarrow & \mathbb{Z}^{\Delta(1)} & \rightarrow & \mathcal{A}_{n-1}(\mathbb{P}_{\Delta}) & \rightarrow & 0 \end{array}$$

For each $\rho \in \Delta(1)$, introduce a variable z_{ρ} , and consider the polynomial ring

$$S = \mathbb{C}[z_{\rho}] = \mathbb{C}[z_{\rho} : \rho \in \Delta(1)].$$

Note that a monomial $\prod_{\rho} z_{\rho}^{a_{\rho}}$ determines a divisor $D = \sum_{\rho} a_{\rho} D_{\rho}$ and to emphasize this relationship, we will write the monomial as z^D . We define the degree of a monomial $z^D \in S$ as $\deg(z^D) = [D] \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$. Using the commutative diagram (1), it follows that two monomials $\prod_{\rho} z_{\rho}^{a_{\rho}}$ and $\prod_{\rho} z_{\rho}^{b_{\rho}}$ in S have the same degree if and only if there is some $m \in M$ such that $a_{\rho} = \langle m, n_{\rho} \rangle + b_{\rho}$ for every ρ . Then

$$S = \bigoplus_{\alpha \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})} S_{\alpha},$$

where $S_{\alpha} = \bigoplus_{\deg(z^D) = \alpha} \mathbb{C} \cdot z^D$. Note also that $S_{\alpha} \cdot S_{\beta} = S_{\alpha+\beta}$. The polynomial ring S is called homogeneous coordinate ring of the toric variety \mathbb{P}_{Δ} .

Denote by $\mathcal{O}_{\mathbb{P}_{\Delta}}$ the structure sheaf of \mathbb{P}_{Δ} . Let $\mathcal{O}_{\mathbb{P}_{\Delta}}(D)$ be the coherent sheaf on \mathbb{P}_{Δ} determined by a Weil divisor D . If $\alpha = [D] \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$, then it follows from [13] that

$$S_{\alpha} \simeq H^0(\mathbb{P}_{\Delta}, \mathcal{O}_{\mathbb{P}_{\Delta}}(D)),$$

moreover, if $\alpha = [D_1]$ and $\beta = [D_2]$, there is a commutative diagram

$$\begin{array}{ccc} S_{\alpha} \otimes S_{\beta} & \longrightarrow & S_{\alpha+\beta} \\ \downarrow & & \downarrow \\ H^0(\mathbb{P}_{\Delta}, \mathcal{O}_{\mathbb{P}_{\Delta}}(D_1)) \otimes H^0(\mathbb{P}_{\Delta}, \mathcal{O}_{\mathbb{P}_{\Delta}}(D_2)) & \longrightarrow & H^0(\mathbb{P}_{\Delta}, \mathcal{O}_{\mathbb{P}_{\Delta}}(D_1 + D_2)) \end{array}$$

where the top arrow is the polynomial multiplication. If \mathbb{P}_{Δ} is a complete toric variety, then it follows from [13] that

- (i) S_{α} is finite dimensional for every α , and in particular, $S_0 = \mathbb{C}$.
- (ii) If $\alpha = [D]$ for an effective divisor $D = \sum_{\rho} a_{\rho} D_{\rho}$, then $\dim_{\mathbb{C}}(S_{\alpha}) = \#(P_D \cap M)$, where $P_D = \{m \in M_{\mathbb{R}} : \langle m, n_{\rho} \rangle \geq -a_{\rho} \text{ for all } \rho\}$.

2.2. The toric homogeneous coordinates. Given a toric variety \mathbb{P}_{Δ} , the Weil divisor class group $\mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$ is a finitely generated abelian group of rank $r = k - n$, where $k = |\Delta(1)|$. If we apply $\text{Hom}_{\mathbb{Z}}(-, \mathbb{C}^*)$ to the bottom exact sequence of (1), then we get the exact sequence

$$1 \longrightarrow G \longrightarrow (\mathbb{C}^*)^{\Delta(1)} \longrightarrow \mathbb{T} \longrightarrow 1,$$

where $G = \text{Hom}_{\mathbb{Z}}(\mathcal{A}_{n-1}(\mathbb{P}_{\Delta}), \mathbb{C}^*)$. Since $(\mathbb{C}^*)^{\Delta(1)}$ acts naturally on $\mathbb{C}^{\Delta(1)}$, the subgroup $G \subset (\mathbb{C}^*)^{\Delta(1)}$ acts on $\mathbb{C}^{\Delta(1)}$ by

$$g \cdot t = \left(g([D_{\rho}]) t_{\rho} \right),$$

for $g : \mathcal{A}_{n-1}(\mathbb{P}_{\Delta}) \rightarrow \mathbb{C}^*$ in G , and $t = (t_{\rho}) \in \mathbb{C}^{\Delta(1)}$. The explicit equations for G as a subgroup of the torus $(\mathbb{C}^*)^{\Delta(1)}$ is given by

$$G = \left\{ (t_{\rho}) \in (\mathbb{C}^*)^{\Delta(1)} \left| \prod_{\rho} t_{\rho}^{\langle m_i, n_{\rho} \rangle} = 1, 1 \leq i \leq n \right. \right\},$$

where m_1, \dots, m_n is a basis of M ; see [12].

For each cone $\sigma \in \Delta$, define the monomial

$$z^{\hat{\sigma}} = \prod_{\rho \notin \sigma(1)} z_{\rho},$$

which is the product of the variables corresponding to rays not in σ . Then define

$$\mathcal{Z} = V(\{z^{\hat{\sigma}} : \sigma \in \Delta\}) \subset \mathbb{C}^{\Delta(1)},$$

we have that $\mathcal{Z} \subset \mathbb{C}^{\Delta(1)}$ has codimension at least two, and $\mathcal{Z} = \{0\}$ when $r = 1$; see [3].

Theorem 2.1. [13] *Let \mathbb{P}_{Δ} be a n -dimensional toric variety such that $\Delta(1)$ spans $N_{\mathbb{R}}$. Then*

- (i) \mathbb{P}_{Δ} is naturally isomorphic to the categorical quotient $(\mathbb{C}^{\Delta(1)} - \mathcal{Z})/G$.
- (ii) \mathbb{P}_{Δ} is the geometric quotient $(\mathbb{C}^{\Delta(1)} - \mathcal{Z})/G$ if and only if \mathbb{P}_{Δ} is an orbifold; see Section 2.5.

We shall consider the following subfield of $\mathbb{C}(z_{\rho}) = \text{Frac}(\mathbb{C}[z_{\rho}])$ given by

$$K(\mathbb{P}_{\Delta}) = \left\{ \frac{P}{Q} \in \mathbb{C}(z_{\rho}) : P \in S_{\alpha}, Q \in S_{\alpha} \right\}.$$

The field of rational functions $K(\mathbb{P}_{\Delta})$ on \mathbb{P}_{Δ} , is the subfield of $\mathbb{C}(z_{\rho})$ such that $\deg(P) = \deg(Q)$. It follows that the polynomials $P, Q \in S_{\alpha}$ define a rational function $\frac{P}{Q} : \mathbb{P}_{\Delta} \dashrightarrow \mathbb{P}^1$.

2.3. Quasi-homogeneous hypersurfaces. Assume that \mathbb{P}_{Δ} is a complex orbifold. An element $\alpha \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$ gives the character $\chi^{\alpha} : G \rightarrow \mathbb{C}^*$ that sends $g \in G = \text{Hom}_{\mathbb{Z}}(\mathcal{A}_{n-1}(\mathbb{P}_{\Delta}), \mathbb{C}^*)$ to $g(\alpha) \in \mathbb{C}^*$. The action of G on $\mathbb{C}^{\Delta(1)}$ induces an action on S with the property that given $f \in S$, we have

$$f \in S_{\alpha} \Leftrightarrow f(g \cdot z) = \chi^{\alpha}(g) f(z), \quad \forall g \in G, z \in \mathbb{C}^{\Delta(1)}.$$

We say that $f \in S_{\alpha}$ is quasi-homogeneous of degree α . It follows that the equation $V = \{f = 0\}$ is a well-defined hypersurface in \mathbb{P}_{Δ} . We say that $V = \{f = 0\}$ is a quasi-homogeneous hypersurface of degree α .

Suppose there is a complex number a_{ρ} for each $\rho \in \Delta(1)$ with the property that $\sum_{\rho} a_{\rho} n_{\rho} = 0$ in $N_{\mathbb{C}}$. Then, for any class $\alpha \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$, there is a constant $\theta(\alpha)$ with the property that for any quasi-homogeneous polynomial $f \in S$ of degree α , we have

$$(2) \quad i_R(df) = \theta(\alpha) \cdot f,$$

where $R = \sum_{\rho} a_{\rho} z_{\rho} \frac{\partial}{\partial z_{\rho}}$. The identity (2) is called the Euler formula determined by R . Moreover, considering the $r = k - n$ linearly independent over \mathbb{Z} relations between the n_{ρ} , we have r vector

fields R_i tangent to the orbits of G and $\text{Lie}(G) = \langle R_1, \dots, R_r \rangle$; for more details see [3]. We will call these vector fields $R_i, i = 1, \dots, r$, the radial vector fields on \mathbb{P}_Δ .

2.4. Examples.

- (1) **Weighted projective spaces.** [12] Let $\omega_0, \dots, \omega_n$ be positive integers with $\gcd(\omega_0, \dots, \omega_n) = 1$. Set $\omega = (\omega_0, \dots, \omega_n)$. Consider the \mathbb{C}^* action on \mathbb{C}^{n+1} given by

$$t \cdot (z_0, \dots, z_n) = (t^{\omega_0} z_0, \dots, t^{\omega_n} z_n).$$

Then, the weighted projective spaces is defined by

$$\mathbb{P}(\omega) = (\mathbb{C}^{n+1} - \{0\}) / \mathbb{C}^*.$$

If $\omega_0 = \dots = \omega_n = 1$, then $\mathbb{P}(\omega) = \mathbb{P}^n$. When $\omega_0, \dots, \omega_n$ are pairwise coprime, we say that $\mathbb{P}(\omega)$ is a well formed weighted projective space and we have

$$\text{Sing}(\mathbb{P}(\omega)) = \{\bar{e}_i \mid \omega_i > 1\}.$$

Moreover, we have $\mathcal{A}_{n-1}(\mathbb{P}(\omega)) \simeq \mathbb{Z}$ and $\deg(z_i) = \omega_i$ for all $0 \leq i \leq n+1$. Consequently the homogeneous coordinate ring of $\mathbb{P}(\omega)$ is given by $S = \bigoplus_{\alpha \geq 0} S_\alpha$, where

$$S_\alpha = \bigoplus_{p_0 \omega_0 + \dots + p_n \omega_n = \alpha} \mathbb{C} \cdot z_0^{p_0} \dots z_n^{p_n}.$$

- (2) **Multiprojective spaces.** [12] Consider $\mathbb{C}^{n+1} \times \mathbb{C}^{m+1}$ in coordinates $(z_1, z_2) = (z_{1,0}, \dots, z_{1,n}, z_{2,0}, \dots, z_{2,m})$. Now, consider the $(\mathbb{C}^*)^2$ action on $\mathbb{C}^{n+1} \times \mathbb{C}^{m+1}$ given by

$$(\mu, \lambda) \cdot (z_1, z_2) = (\mu z_{1,0}, \dots, \mu z_{1,n}, \lambda z_{2,0}, \dots, \lambda z_{2,m}).$$

Then, the multiprojective spaces is defined by

$$\mathbb{P}^n \times \mathbb{P}^m = (\mathbb{C}^{n+1} \times \mathbb{C}^{m+1} - \mathcal{Z}) / (\mathbb{C}^*)^2,$$

where $\mathcal{Z} = (\{0\} \times \mathbb{C}^{m+1}) \cup (\mathbb{C}^{n+1} \times \{0\})$.

Moreover, we have $\mathcal{A}_{n+m-1}(\mathbb{P}^n \times \mathbb{P}^m) \simeq \mathbb{Z}^2$ and $\deg(z_{1,i}) = (1, 0)$ for all $0 \leq i \leq n$, and $\deg(z_{2,j}) = (0, 1)$ for all $0 \leq j \leq m$. Consequently the homogeneous coordinate ring of $\mathbb{P}^n \times \mathbb{P}^m$ is given by $S = \bigoplus_{\alpha, \beta \geq 0} S_{(\alpha, \beta)}$, where

$$S_{(\alpha, \beta)} = \bigoplus_{p_0 + \dots + p_n = \alpha; q_0 + \dots + q_m = \beta} \mathbb{C} \cdot z_{1,0}^{p_0} \dots z_{1,n}^{p_n} z_{2,0}^{q_0} \dots z_{2,m}^{q_m},$$

is the ring of bihomogeneous polynomials of bidegree (α, β) . More generally we can define $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ (in general, a finite product of toric varieties is again a toric variety).

- (3) **Hirzebruch surfaces.** Let $r \geq 0$ be an integer. The Hirzebruch surface \mathcal{H}_r is a rational ruled surface defined in $\mathbb{P}^1 \times \mathbb{P}^2$ by

$$\{((\lambda_0 : \lambda_1), (\mu_0 : \mu_1 : \mu_2)) \mid \lambda_0^r \mu_0 = \lambda_1^r \mu_1\}.$$

In particular $\mathcal{H}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{H}_1 \simeq \text{Bl}_p(\mathbb{P}^2)$ (Blow-up of \mathbb{P}^2 at a point p). As a toric variety, $\mathbb{P}_\Delta = \mathcal{H}_r$ has the ray generators $u_1 = -e_1 + r e_2$, $u_2 = e_2$, $u_3 = e_1$ and $u_4 = -e_2$ in

$N_{\mathbb{R}} = \mathbb{R}^2$; see [12].

Consider the $(\mathbb{C}^*)^2$ action on $\mathbb{C}^2 \times \mathbb{C}^2$ given as follows

$$(\lambda, \mu)(z_{1,1}, z_{1,2}, z_{2,1}, z_{2,2}) = (\lambda z_{1,1}, \mu z_{1,2}, \lambda z_{2,1}, \mu \lambda^r z_{2,2}).$$

The quotient representation of the Hirzebruch surface is given by

$$\mathcal{H}_r = (\mathbb{C}^2 \times \mathbb{C}^2 - \mathcal{Z}) / (\mathbb{C}^*)^2,$$

where $\mathcal{Z} = Z(z_{1,1}, z_{2,1}) \cup Z(z_{1,2}, z_{2,2})$.

Moreover, we have $\mathcal{A}_1(\mathcal{H}_r) \simeq \mathbb{Z}^2$ and the homogeneous coordinate ring associated to \mathcal{H}_r is given by $S = \bigoplus_{\alpha \geq 0, \beta \geq 0} S_{(\alpha, \beta)}$, where

$$S_{(\alpha, \beta)} = \bigoplus_{\alpha = p_1 + q_1 + r q_2; \beta = p_2 + q_2} \mathbb{C} \cdot z_{1,1}^{p_1} z_{1,2}^{p_2} z_{2,1}^{q_1} z_{2,2}^{q_2}.$$

In particular $\deg(z_{1,1}) = (1, 0)$, $\deg(z_{1,2}) = (0, 1)$, $\deg(z_{2,1}) = (1, 0)$ and $\deg(z_{2,2}) = (r, 1)$.

- (4) **Rational normal scrolls.** Let a_1, \dots, a_n be integers. Consider the $(\mathbb{C}^*)^2$ action on $\mathbb{C}^2 \times \mathbb{C}^n$ given by

$$(\lambda, \mu)(z_{1,1}, z_{1,2}, z_{2,1}, \dots, z_{2,n}) = (\lambda z_{1,1}, \lambda z_{1,2}, \mu \lambda^{-a_1} z_{2,1}, \dots, \mu \lambda^{-a_n} z_{2,n}).$$

Then, a n -dimensional rational normal scroll $\mathbb{F}(a_1, \dots, a_n)$ is a projective toric manifold defined by

$$\mathbb{F}(a_1, \dots, a_n) = (\mathbb{C}^2 \times \mathbb{C}^n - \mathcal{Z}) / (\mathbb{C}^*)^2,$$

where $\mathcal{Z} = (\{0\} \times \mathbb{C}^n) \cup (\mathbb{C}^2 \times \{0\})$.

Moreover, we have $\mathcal{A}_{n-1}(\mathbb{F}(a_1, \dots, a_n)) \simeq \mathbb{Z}^2$ and the homogeneous coordinate ring associated to $\mathbb{F}(a_1, \dots, a_n)$ is given by $S = \bigoplus_{\alpha \in \mathbb{Z}, \beta \geq 0} S_{(\alpha, \beta)}$, where

$$S_{(\alpha, \beta)} = \bigoplus_{\alpha = p_1 + p_2 - \sum_i q_i a_i; \beta = \sum_i q_i} \mathbb{C} \cdot z_{1,1}^{p_1} z_{1,2}^{p_2} z_{2,1}^{q_1} \dots z_{2,n}^{q_n}.$$

In particular $\deg(z_{1,1}) = \deg(z_{1,2}) = (1, 0)$ and $\deg(z_{2,i}) = (-a_i, 1)$.

Consider $1 \leq a_1 \leq a_2 \leq \dots \leq a_n$ integers, it is possible to show that $\mathbb{F}(a_1, \dots, a_n) \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \dots \oplus \mathcal{O}_{\mathbb{P}^1}(a_n))$, and

$$\mathbb{F}(a_1, \dots, a_n) = \mathbb{F}(b_1, \dots, b_n) \iff \{a_1, \dots, a_n\} = \{b_1 + c, \dots, b_n + c\},$$

for some $c \in \mathbb{Z}$. For more details see [12, 27].

2.5. Orbifolds. A n -dimensional orbifold X is a complex space endowed with the following property: each point $x \in X$ possesses a open neighborhood, which is the quotient $U_x = \tilde{U}/G_x$, where \tilde{U} is a complex manifold, of dimension n , and G_x is a properly discontinuous finite group of automorphisms of \tilde{U} , so that locally we have a quotient map $\pi_x : (\tilde{U}, \tilde{x}) \rightarrow (U_x, x)$. Since the possible singularities appearing in a orbifold X are quotient singularities, we have that X is reduced, normal, Cohen-Macaulay and with only rational singularities.

Satake's fundamental idea was to use local smoothing coverings to extend the definition of usual known objects to orbifolds. For instance, smooth differential k -forms on X are C^∞ -differential k -forms ω on $X_{\text{reg}} = X \setminus \text{Sing}(X)$ such that the pull-back $\pi_x^* \omega$ extends to a C^∞ -differential k -form

on every local smoothing covering $\pi_x : (\tilde{U}, \tilde{x}) \rightarrow (X, x)$ of X . Hence, if ω is a smooth $2n$ -form on X with compact support $\text{Supp}(\omega) \subset (X, x)$, then by definition

$$\int_X^{orb} \omega = \frac{1}{\#G_x} \int_{\tilde{U}} \pi^* \omega.$$

If now the $2n$ -form ω has compact support, we use a partition of unity $\{\rho_\alpha, U_\alpha\}_{\alpha \in \Lambda}$, where $\pi_\alpha : (\tilde{U}_\alpha, \tilde{x}_\alpha) \rightarrow (U_\alpha, x_\alpha)$ is a local smoothing covering and $\sum \rho_\alpha(x) = 1$ for all $x \in \text{Supp}(\omega)$, and set

$$\int_X^{orb} \omega = \sum_\alpha \int_X^{orb} \rho_\alpha \omega.$$

Analogous to the manifold case, we have the sheaf of C^∞ -differential k -forms and exterior differentiation. Stokes' formula, Poincaré's duality and Rham's theorem also hold on orbifolds. For a deeper discussion of orbifolds we refer the reader to [1, 4, 19].

3. CODIMENSION ONE HOLOMORPHIC DISTRIBUTIONS

Let \mathbb{P}_Δ be a complete simplicial toric variety of dimension n where $\Delta(1)$ spans $N_{\mathbb{R}}$. We know that $|\Delta(1)| = n + r$, where r is the rank of the finitely generated abelian group $\mathcal{A}_{n-1}(\mathbb{P}_\Delta)$. We will denote $\Delta(1) = \{\rho_1, \dots, \rho_{n+r}\}$, $S = \mathbb{C}[z_1, \dots, z_{n+r}]$, and $D_i = D_{\rho_i}$ for all $i = 1, \dots, n + r$.

There exists an exact sequence known as the generalized Euler's sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_\Delta}^{\oplus r} \rightarrow \bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_\Delta}(D_i) \rightarrow \mathcal{T}\mathbb{P}_\Delta \rightarrow 0,$$

where $\mathcal{T}\mathbb{P}_\Delta = \mathcal{H}om(\Omega_{\mathbb{P}_\Delta}^1, \mathcal{O}_{\mathbb{P}_\Delta})$ is the so-called Zariski tangent sheaf of \mathbb{P}_Δ . Let $i : \mathbb{P}_{\Delta \text{reg}} \rightarrow \mathbb{P}_\Delta$ be the inclusion of the regular part $\mathbb{P}_{\Delta \text{reg}} = \mathbb{P}_\Delta - \text{Sing}(\mathbb{P}_\Delta)$. Since \mathbb{P}_Δ is a complex orbifold then $\mathcal{T}\mathbb{P}_\Delta \simeq i_* \mathcal{T}\mathbb{P}_{\Delta \text{reg}}$, where $\mathcal{T}\mathbb{P}_{\Delta \text{reg}}$ is the tangent sheaf of $\mathbb{P}_{\Delta \text{reg}}$; see [11, Appendix A.2].

Let $\mathcal{O}_{\mathbb{P}_\Delta}(d) = \mathcal{O}_{\mathbb{P}_\Delta}(d_1, \dots, d_{n+r}) = \mathcal{O}_{\mathbb{P}_\Delta}(\sum_{i=1}^{n+r} d_i D_i)$, where $\sum_{i=1}^{n+r} d_i D_i$ is a Cartier divisor. Set $\Omega_{\mathbb{P}_\Delta}^1(d) = \Omega_{\mathbb{P}_\Delta}^1 \otimes \mathcal{O}_{\mathbb{P}_\Delta}(d)$. Tensorizing the dual Euler's sequence by $\mathcal{O}_{\mathbb{P}_\Delta}(d)$ we get

$$0 \rightarrow \Omega_{\mathbb{P}_\Delta}^1(d) \rightarrow \bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_\Delta}(d_1, \dots, d_i - 1, \dots, d_{n+r}) \rightarrow Cl(\mathbb{P}_\Delta) \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{P}_\Delta}(d)^{\oplus r} \rightarrow 0.$$

For more details, see for instance [3, 12].

Definition 3.1. *A codimension one holomorphic distribution \mathcal{D} on \mathbb{P}_Δ of degree $d = \sum_{i=1}^{n+r} d_i [D_i] \in \mathcal{A}_{n-1}(\mathbb{P}_\Delta)$ is a global section ω of $\Omega_{\mathbb{P}_\Delta}^1(d)$. For simplicity of notation we say that \mathcal{D} has degree $d = (d_1, \dots, d_{n+r})$. A codimension one holomorphic foliation \mathcal{F} on \mathbb{P}_Δ is a distribution that satisfies the Frobenius integrability condition $\omega \wedge d\omega = 0$. We will consider codimension one holomorphic foliations whose singular scheme has codimension greater than 1.*

If $H^1(\mathbb{P}_\Delta, \Omega_{\mathbb{P}_\Delta}^1(d)) = 0$; see for instance the Bott-Steenbrink-Danilov vanishing theorem [12, Theorem 9.3.1], then we have that a codimension one holomorphic distribution \mathcal{D} on \mathbb{P}_Δ of degree d is given by a polynomial form in homogeneous coordinates of the form

$$\omega = \sum_{i=1}^{n+r} P_i dz_i,$$

where P_i is a quasi-homogeneous polynomial of degree $(d_1, \dots, d_i-1, \dots, d_{n+r})$ for all $i = 1, \dots, n+r$, such that $i_R \omega = 0$ for all $R \in \text{Lie}(G)$. Note that ω is a quasi-homogeneous form of degree d . We define the singular set of \mathcal{D} by

$$\text{Sing}(\mathcal{D}) = \pi(\{p \in \mathbb{C}^{n+r} : \omega(p) = 0\}),$$

where $\pi : \mathbb{C}^{n+r} - \mathcal{Z} \rightarrow \mathbb{P}_\Delta$ is the canonical projection.

Let $V = \{f = 0\}$ be a quasi-homogeneous hypersurface. We recall that V is invariant by \mathcal{D} if and only if there is a holomorphic 2-form Θ in homogeneous coordinates, such that $\omega \wedge df = f \cdot \Theta$. We also recall that a rational first integral of \mathcal{D} is a non-constant rational function f such that $\omega \wedge df = 0$.

It is worth mentioning that there are some works about one-dimensional holomorphic foliations on compact toric orbifolds, see for instance [8, 28]. In [2] the authors studied codimension one Fano distributions on Fano manifolds with Picard number one.

3.1. Examples.

- (1) **Weighted projective spaces.** The Euler's sequence on $\mathbb{P}(\omega)$ is an exact sequence of orbibundles

$$0 \longrightarrow \underline{\mathbb{C}} \longrightarrow \bigoplus_{i=0}^n \mathcal{O}_{\mathbb{P}(\omega)}(\omega_i) \longrightarrow T\mathbb{P}(\omega) \longrightarrow 0,$$

where $\underline{\mathbb{C}}$ is the trivial line orbibundle on $\mathbb{P}(\omega)$; see [22]. The radial vector field is given by

$$R = \omega_0 z_0 \frac{\partial}{\partial z_0} + \dots + \omega_n z_n \frac{\partial}{\partial z_n}.$$

- (2) **Multiprojective spaces.** The Euler's sequence over \mathbb{P}^n

$$0 \longrightarrow \underline{\mathbb{C}} \longrightarrow \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n+1} \longrightarrow T\mathbb{P}^n \longrightarrow 0,$$

gives, by direct summation, the exact sequence

$$0 \longrightarrow \underline{\mathbb{C}}^2 \longrightarrow \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^m}(1, 0)^{\oplus n+1} \oplus \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^m}(0, 1)^{\oplus m+1} \longrightarrow T(\mathbb{P}^n \times \mathbb{P}^m) \longrightarrow 0.$$

Here, the radial vector fields are given by

$$R_1 = z_{1,0} \frac{\partial}{\partial z_{1,0}} + \dots + z_{1,n} \frac{\partial}{\partial z_{1,n}}, \text{ and } R_2 = z_{2,0} \frac{\partial}{\partial z_{2,0}} + \dots + z_{2,m} \frac{\partial}{\partial z_{2,m}}.$$

- (3) **Hirzebruch surface.** The Euler's sequence on \mathcal{H}_r is

$$0 \rightarrow \mathcal{O}_{\mathcal{H}_r}^{\oplus 2} \rightarrow \mathcal{O}_{\mathcal{H}_r}(1, 0) \oplus \mathcal{O}_{\mathcal{H}_r}(0, 1) \oplus \mathcal{O}_{\mathcal{H}_r}(1, 0) \oplus \mathcal{O}_{\mathcal{H}_r}(r, 1) \rightarrow \mathcal{T}\mathcal{H}_r \rightarrow 0.$$

Here, the radial vector fields are given by

$$R_1 = z_{1,1} \frac{\partial}{\partial z_{1,1}} + z_{2,1} \frac{\partial}{\partial z_{2,1}} + r z_{2,2} \frac{\partial}{\partial z_{2,2}}, \text{ and } R_2 = z_{1,2} \frac{\partial}{\partial z_{1,2}} + z_{2,2} \frac{\partial}{\partial z_{2,2}}.$$

(4) **Rational normal scrolls.** The Euler's sequence on $\mathbb{F}(a) := \mathbb{F}(a_1, \dots, a_n)$ is

$$0 \rightarrow \mathcal{O}_{\mathbb{F}(a)}^{\oplus 2} \rightarrow \mathcal{O}_{\mathbb{F}(a)}(1, 0)^{\oplus 2} \oplus \bigoplus_{i=1}^n \mathcal{O}_{\mathbb{F}(a)}(-a_i, 1) \rightarrow \mathcal{T}\mathbb{F}(a) \rightarrow 0.$$

Here, the radial vector fields are given by

$$R_1 = z_{1,1} \frac{\partial}{\partial z_{1,1}} + z_{1,2} \frac{\partial}{\partial z_{1,2}} - \sum_{i=1}^n a_i z_{2,i} \frac{\partial}{\partial z_{2,i}}, \text{ and } R_2 = \sum_{i=1}^n z_{2,i} \frac{\partial}{\partial z_{2,i}}.$$

The following proposition is due to Jouanolou [21, Proposition 2.6], in the case of the complex projective space

Proposition 3.1. *Let \mathbb{P}_Δ be a complete simplicial toric variety of dimension n , with homogeneous coordinates z_1, \dots, z_{n+r} . Let $\omega \in \Omega_{\mathbb{P}_\Delta}^1(d)$ be nonzero integrable, and suppose there exists a radial vector field $R = \sum_i a_i z_i \frac{\partial}{\partial z_i}$ on \mathbb{P}_Δ such that $\sum_i a_i d_i \neq 0$. Then, the singular set of ω must contain at least a codimension two irreducible component.*

Proof. Let us first prove the following identity

$$i_R(d\omega) = (\sum_i a_i d_i)\omega.$$

Let $R_t(z) = (e^{a_1 t} z_1, \dots, e^{a_{n+r} t} z_{n+r})$ be the flow of R ; $R_0(z) = z$ and $R'_0(z) = R(z)$. Set $\omega = \sum_i P_i(z) dz_i$. Then

$$\begin{aligned} \mathcal{L}_R(\omega) &= \left. \frac{d}{dt} R_t^*(\omega) \right|_{t=0} = \left. \frac{d}{dt} \sum_i P_i(e^{a_1 t} z_1, \dots, e^{a_{n+r} t} z_{n+r}) d(e^{a_i t} z_i) \right|_{t=0} \\ &= \left. \frac{d}{dt} \left(e^{(\sum_i a_i d_i)t} \right) \right|_{t=0} \cdot \sum_i P_i(z) dz_i = (\sum_i a_i d_i)\omega, \end{aligned}$$

and consequently we have $(\sum_i a_i d_i)\omega = \mathcal{L}_R(\omega) = i_R(d\omega) + d(i_R\omega) = i_R(d\omega)$. Now, the proposition follows from Saito's division Lemma [29, 21]. In fact, suppose that $\text{codim Sing}(\omega) \geq 3$, the condition $\omega \wedge d\omega = 0$ implies that $d\omega = \alpha \wedge \omega$ for some polynomial 1-form α , and so $d\omega = 0$. Therefore we have $(\sum_i a_i d_i)\omega = i_R(d\omega) = 0$, an absurd. \square

Let us prove the following result due to Darboux, in the case of the complex projective space

Proposition 3.2. *Let \mathbb{P}_Δ be a complete simplicial toric variety of dimension n , with homogeneous coordinates z_1, \dots, z_{n+r} and Picard number $r = \text{rank } \mathcal{A}_{n-1}(\mathbb{P}_\Delta)$. Let \mathcal{F} be a codimension one holomorphic foliation of degree d on \mathbb{P}_Δ . Denote $h_\alpha = \dim_{\mathbb{C}} H^0(\mathbb{P}_\Delta, \mathcal{O}_{\mathbb{P}_\Delta}(\alpha)) = \dim_{\mathbb{C}} \mathcal{S}(\Delta)_\alpha$. If \mathcal{F} admits*

$$\mathcal{N}_\Delta = 2 + \sum_{1 \leq i < j \leq n+r} h_{d - \deg(z_i) - \deg(z_j)}$$

invariant irreducible quasi-homogeneous hypersurfaces, then \mathcal{F} admits a rational first integral.

In particular, \mathcal{F} admits a rational first integral if and only if it admits an infinite number of invariants irreducible quasi-homogeneous hypersurfaces.

Proof. The proof is adapted from [23, Theorem 1.3]. Set $N = \mathcal{N}_\Delta$. Let f_1, \dots, f_N be the \mathcal{F} -invariant irreducible quasi-homogeneous hypersurfaces. Let ω be a polynomial 1-form that defines \mathcal{F} in homogeneous coordinates. The invariance condition of f_i implies that

$$(3) \quad \frac{df_i}{f_i} \wedge \omega = \Theta_i, \quad i = 1, \dots, N,$$

where the Θ_i are quasi-homogeneous 2-forms of degree d on \mathbb{P}_Δ . Since the vector space of quasi-homogeneous 2-forms of degree d on \mathbb{P}_Δ has finite dimension equal to

$$\sum_{1 \leq i < j \leq n+r} h_{d-\deg(z_i)-\deg(z_j)},$$

the set $\{\Theta_1, \dots, \Theta_N\}$ is linearly dependent. Then there exist complex numbers $\alpha_1, \dots, \alpha_N$ not all equal to zero, such that $\sum_{i=1}^N \alpha_i \cdot \Theta_i = 0$. By (3) it follows that $\eta_1 \wedge \omega = 0$, where

$$\eta_1 = \sum_{i=1}^N \alpha_i \frac{df_i}{f_i}.$$

In particular, the 1-form $\omega_1 = f_1 \dots f_N \cdot \eta_1$ is holomorphic and such that $\omega_1 \wedge \omega = 0$. The condition $\text{codim Sing}(\omega) \geq 3$ implies there exists $p \in \mathbb{C}[z_1, \dots, z_{n+r}]$ such that $\omega_1 = p \cdot \omega$, so we can write $\eta_1 = h_1 \cdot \omega$, where $h_1 = \frac{p}{f_1 \dots f_N}$. By a similar argument, considering the indices $2, \dots, N+1$, we can build a 1-form

$$\eta_2 = \sum_{i=2}^{N+1} \beta_i \frac{df_i}{f_i},$$

such that the vectors $(\alpha_1, \dots, \alpha_N, 0)$ and $(0, \beta_2, \dots, \beta_{N+1})$ are linearly independent, and there exists a meromorphic function h_2 such that $\eta_2 = h_2 \cdot \omega$. The relation $\eta_1 = h_1 \cdot \omega$ and $\eta_2 = h_2 \cdot \omega$ implies that $\eta_2 = \frac{h_2}{h_1} \cdot \eta_1$. From this we obtain $d\left(\frac{h_2}{h_1}\right) \wedge \eta_1 = d\eta_2 = 0$, hence that $d\left(\frac{h_2}{h_1}\right) \wedge \omega = 0$, and finally that $d\left(\frac{h_2}{h_1}\right) = h_3 \cdot \omega$, where h_3 is a meromorphic function.

Let us verify now that $\frac{h_2}{h_1}$ is not constant. Suppose that $\frac{h_2}{h_1} = c \in \mathbb{C}$, then $\eta_2 = h_2 \cdot \omega = c \cdot \eta_1$, which is absurd. From this we conclude that $\frac{h_2}{h_1}$ is a rational first integral of \mathcal{D} . \square

Remark 3.1. Consider the multiprojective space $\mathbb{P}_\Delta = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_r}$. Set $\alpha = (m_1, \dots, m_r) \in \mathcal{A}_{n-1}(\mathbb{P}_\Delta)$. Then

$$h_\alpha = \prod_{i=1}^r \binom{n_i + m_i}{n_i}.$$

Similarly, consider the n -dimensional rational normal scroll $\mathbb{P}_\Delta = \mathbb{F}(a_1, \dots, a_n)$ and $\alpha = (m_1, m_2) \in \mathcal{A}_{n-1}(\mathbb{P}_\Delta)$. Then

$$h_\alpha = \left(\sum_{i=1}^n a_i \right) \binom{m_1 + n - 1}{n} + (m_2 + 1) \binom{m_1 + n - 1}{n - 1},$$

see [27].

Example 3.1. Let $\mathbb{P}(\omega)$ be an n -dimensional weighted projective space. Let $P_S(t)$ be the Poincaré series of the graded algebra $S(\omega)$ of $\mathbb{P}(\omega)$ given by

$$P_S(t) = \sum_{k=0}^{\infty} h_k t^k = \prod_{k=0}^n (1 - t^{\omega_k})^{-1},$$

where $h_k = \dim_{\mathbb{C}} S(\omega)_k$; see [15]. Let \mathcal{F} be a codimension one holomorphic foliation on $\mathbb{P}(\omega)$ of degree d . Then, if \mathcal{F} admits

$$\mathcal{N}(\omega) = 2 + \sum_{1 \leq i < j \leq n+1} h_{d-\omega_i-\omega_j}$$

invariant irreducible quasi-homogeneous hypersurfaces, then \mathcal{F} admits a rational first integral.

In particular, if $\omega = (1, \dots, 1)$ and \mathcal{F} is a codimension one holomorphic foliation on \mathbb{P}^n of degree $d + 2$, then we have

$$h_{d+2-\omega_i-\omega_j} = h_d = \frac{1}{d!} P_S^{(d)}(0) = \binom{n+d}{n}, \quad \text{and} \quad \mathcal{N}(\omega) = \binom{n+d}{n} \binom{n+1}{2} + 2.$$

Consequently we recover the Darboux-Jouanolou type integrability theorem for codimension one holomorphic foliations in the projective space. Similarly, for a Darboux-Jouanolou-Ghys type integrability theorem for one-dimensional foliations on toric varieties, see [8].

3.2. Codimension one distribution with isolated singularities. Let $k = (k_1, \dots, k_n) \in \mathbb{N}^n$ and $m \in \mathbb{N}$. Set $|k| = \sum k_i$ and $\binom{m}{k} = \frac{m!}{k_1! \dots k_n!}$.

Proposition 3.3. *Let \mathbb{P}_{Δ} be an n -dimensional compact toric manifold. Set $h_i = [D_i] \in \mathcal{A}_{n-1}(\mathbb{P}_{\Delta})$. Let \mathcal{D} be a codimension one distribution of degree $d = \sum_i d_i h_i$ with isolated zeros on \mathbb{P}_{Δ} . Then, the singular scheme of \mathcal{D} consists of*

$$\# \text{Sing}(\mathcal{D}) = \sum_{j=0}^n \left\{ (-1)^j \sum_{|k|=n-j} \binom{n-j}{k} \int_{\mathbb{P}_{\Delta}} \mathcal{C}_j(h) \cdot \prod_i (d_i h_i)^{k_i} \right\}$$

points counted with multiplicity, where $\mathcal{C}_j(h)$ is the j th elementary symmetric function of the variables h_1, \dots, h_{n+r} .

Proof. Using the generalized Euler's sequence, we have

$$c(\mathcal{T}\mathbb{P}_{\Delta}) = c\left(\mathcal{O}_{\mathbb{P}_{\Delta}}^{\oplus r}\right) c(\mathcal{T}\mathbb{P}_{\Delta}) = c\left(\bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_{\Delta}}(D_i)\right),$$

where c denotes the total Chern classes. Note that $h_i = c_1(\mathcal{O}_{\mathbb{P}_\Delta}(D_i)) = [D_i]$. Then

$$\begin{aligned} c_n(\mathcal{T}\mathbb{P}_\Delta \otimes \mathcal{O}_{\mathbb{P}_\Delta}(d)) &= \sum_{j=0}^n c_j(\mathcal{T}\mathbb{P}_\Delta) c_1(\mathcal{O}_{\mathbb{P}_\Delta}(d))^{n-j} \\ &= \sum_{j=0}^n c_j \left(\bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_\Delta}(D_i) \right) \left(\sum_{i=1}^{n+r} c_1(\mathcal{O}_{\mathbb{P}_\Delta}(d_i D_i)) \right)^{n-j} \\ &= \sum_{j=0}^n c_j \left(\bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_\Delta}(D_i) \right) \left(\sum_{i=1}^{n+r} d_i h_i \right)^{n-j}. \end{aligned}$$

On the other hand, we have

$$c \left(\bigoplus_{i=1}^{n+r} \mathcal{O}_{\mathbb{P}_\Delta}(D_i) \right) = \prod_{i=1}^{n+r} c(\mathcal{O}_{\mathbb{P}_\Delta}(D_i)) = \prod_{i=1}^{n+r} (1 + c_1(\mathcal{O}_{\mathbb{P}_\Delta}(D_i))) = \prod_{i=1}^{n+r} (1 + h_i) = \sum_{j=0}^{n+r} \mathbf{C}_j(h).$$

Then

$$c_n(\mathcal{T}\mathbb{P}_\Delta \otimes \mathcal{O}_{\mathbb{P}_\Delta}(d)) = \sum_{j=0}^n \mathbf{C}_j(h) \left(\sum_{i=1}^{n+r} d_i h_i \right)^{n-j} = \sum_{j=0}^n \left\{ \sum_{|k|=n-j} \binom{n-j}{k} \mathbf{C}_j(h) \prod_i (d_i h_i)^{k_i} \right\},$$

hence

$$c_n(\Omega_{\mathbb{P}_\Delta}^1(d)) = (-1)^n c_n(\mathcal{T}\mathbb{P}_\Delta \otimes \mathcal{O}_{\mathbb{P}_\Delta}(-d)) = \sum_{j=0}^n \left\{ (-1)^j \sum_{|k|=n-j} \binom{n-j}{k} \mathbf{C}_j(h) \prod_i (d_i h_i)^{k_i} \right\},$$

so we get

$$\# \text{Sing}(\mathcal{D}) = \int_{\mathbb{P}_\Delta} c_n(\Omega_{\mathbb{P}_\Delta}^1(d)) = \sum_{j=0}^n \left\{ (-1)^j \sum_{|k|=n-j} \binom{n-j}{k} \int_{\mathbb{P}_\Delta} \mathbf{C}_j(h) \cdot \prod_i (d_i h_i)^{k_i} \right\}.$$

□

Example 3.2. Consider the multiprojective space $\mathbb{P}_\Delta = \mathbb{P}^n \times \mathbb{P}^m$. Here $h_1 = h_{10} = \dots = h_{1n}$ and $h_2 = h_{20} = \dots = h_{2m}$. Let \mathcal{D} be a codimension one distribution of degree $d = (d_1, d_2) = d_1 h_1 + d_2 h_2$ with isolated zeros on $\mathbb{P}^n \times \mathbb{P}^m$. Then

$$\begin{aligned} \# \text{Sing}(\mathcal{D}) &= \sum_{j=0}^{n+m} \left\{ (-1)^j \sum_{|k|=n+m-j} \binom{n+m-j}{k_1, k_2} \int_{\mathbb{P}_\Delta} \mathbf{C}_j(h) \cdot \prod_{i=1}^2 (d_i h_i)^{k_i} \right\} \\ &= (-1)^{n+m} \sum_{\substack{0 \leq k_1 \leq n, \\ 0 \leq k_2 \leq m}} (-1)^{|k|} \binom{|k|}{k_1, k_2} d_1^{k_1} d_2^{k_2} \int_{\mathbb{P}_\Delta} \mathbf{C}_{n+m-|k|}(h) \cdot h_1^{k_1} h_2^{k_2} \\ &= (-1)^{n+m} \sum_{\substack{0 \leq k_1 \leq n, \\ 0 \leq k_2 \leq m}} (-1)^{k_1+k_2} \binom{k_1+k_2}{k_1, k_2} \binom{n+1}{n-k_1} \binom{m+1}{m-k_2} d_1^{k_1} d_2^{k_2}. \end{aligned}$$

Similarly, for generic one-dimensional foliations, a formula for the number of singularities in multi-projective spaces is given in [10].

Let us consider an example in blow-ups:

Example 3.3. It is well-known that the blow-ups X_r of \mathbb{P}^2 in $0 \leq r \leq 3$ points in general position are toric del Pezzo surfaces; see for instance [25].

Let us consider X_3 ; it is called the del Pezzo surface of degree six. The morphism $\pi : X_3 \rightarrow \mathbb{P}^2$ induces an injective group homomorphism $\pi^* : \text{Pic}(\mathbb{P}^2) \rightarrow \text{Pic}(X_3)$. We write $H = \pi^*L$ where L is a line in \mathbb{P}^2 , and let us denote by E_1, E_2 and E_3 the exceptional curves. Then $\text{Pic}(X_3) = \mathbb{Z}H \oplus \mathbb{Z}(-E_2) \oplus \mathbb{Z}(-E_1) \oplus \mathbb{Z}(-E_3) \simeq \mathbb{Z}^4$, where

$$H^2 = 1, E_i \cdot E_j = -\delta_{ij} \text{ and } H \cdot E_i = 0.$$

Now, let x, y, z, s, t, u be coordinate functions on \mathbb{C}^6 . As a toric variety we have

$$X_3 = (\mathbb{C}^6 - \mathcal{Z}) / (\mathbb{C}^*)^4,$$

where $\mathcal{Z} = Z(x, t) \cup Z(y, s) \cup Z(z, u) \cup Z(x, y) \cup Z(y, z) \cup Z(z, x) \cup Z(s, t) \cup Z(u, t) \cup Z(s, u)$.

Let us denote by L_1, L_2 and L_3 the strict transforms of the lines in \mathbb{P}^2 spanned by the points that are blown-up. Then we have

$$H = L_1 + E_2 + E_3 = L_2 + E_1 + E_3 = L_3 + E_1 + E_2,$$

and $h_1 = \deg(x) = L_1, h_2 = \deg(y) = L_2, h_3 = \deg(z) = L_3, h_4 = \deg(s) = E_2, h_5 = \deg(t) = E_1$ and $h_6 = \deg(u) = E_3$; for more details see [26].

Now, let \mathcal{F} be a generic holomorphic foliation of degree $d = (d_0, d_1, d_2, d_3) = d_0H - d_1E_2 - d_2E_1 - d_3E_3$ on X_3 . Since $\mathcal{C}_0(h) = 1, \mathcal{C}_1(h) = 3H - E_1 - E_2 - E_3$ and $\mathcal{C}_2(h) = 6$, we have

$$\#\text{Sing}(\mathcal{F}) = \sum_{j=0}^2 (-1)^j d^{2-j} \mathcal{C}_j(h) = d_0(d_0 - 3) + d_1(1 - d_1) + d_2(1 - d_2) + d_3(1 - d_3) + 6.$$

It is of interest to know that blow-ups of \mathbb{P}^n in $0 \leq r \leq n + 1$ points in general position are toric varieties, and that for $r > n + 1$ these are not toric anymore. Moreover, blow-ups of \mathbb{P}^3 along $k = 1, 2$ lines in general position are toric threefolds; see [16].

Definition 3.2. Let \mathbb{P}_Δ be an n -dimensional compact toric orbifold with at most isolated singularities. Let \mathcal{D} be a codimension one distribution with isolated zeros and let ω be a holomorphic section of $\Omega_{\mathbb{P}_\Delta}^1(d)$ that induces \mathcal{D} . The index of \mathcal{D} at a zero p is defined by

$$\mathcal{I}_p^{\text{orb}}(\mathcal{D}) = \frac{1}{\#G_p} \text{Res}_{\tilde{p}} \left\{ \frac{\text{Det}(J\tilde{\omega}) d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\omega}_1 \cdots \tilde{\omega}_n} \right\},$$

where $\pi_p : (\tilde{U}, \tilde{p}) \rightarrow (U, p)$ denotes the local projection of \mathbb{P}_Δ at p : $\tilde{\omega} = \pi_p^* \omega, J\tilde{\omega} = \left(\frac{\partial \tilde{\omega}_i}{\partial \tilde{z}_j} \right)_{1 \leq i, j \leq n}$,

$\text{Res}_{\tilde{p}} \left\{ \frac{\text{Det}(J\tilde{\omega}) d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\omega}_1 \cdots \tilde{\omega}_n} \right\}$ is Grothendieck's point residue and $(\tilde{z}_1, \dots, \tilde{z}_n)$ is a germ of coordinate system on (\tilde{U}, \tilde{p}) .

This index was introduced by Satake [19] for C^∞ -vector fields in the orbifold case, see also [7, 9]. The Grothendieck point residue embodies the Poincaré-Hopf index, the Milnor number and the intersection number of n divisors in \mathbb{C}^n which intersect properly, and has many uses in deep results such as the Baum-Bott theorem, which is a generalization of both the Poincaré-Hopf theorem and the Gauss-Bonnet theorem in the complex realm, see [18, 30, 31].

Proposition 3.4. *Let $\mathbb{P}(\omega)$ be a well formed weighted projective space. Let \mathcal{D} be a codimension one distribution of degree d with isolated zeros on $\mathbb{P}(\omega)$. Then*

$$(4) \quad \sum_{p \in \text{Sing}(\mathcal{D})} \mathcal{I}_p^{orb}(\mathcal{D}) = \frac{1}{\omega_0 \cdots \omega_n} \sum_{j=0}^n (-1)^j \mathcal{C}_j(\omega) d^{n-j}.$$

Proof. Consider the natural map $\varphi : \mathbb{P}^n \rightarrow \mathbb{P}(\omega)$, $\varphi([z_0, \dots, z_n]) = [z_0^{\omega_0}, \dots, z_n^{\omega_n}]_\omega$ of degree $\deg \varphi = \omega_0 \cdots \omega_n$. It was shown in [22] that there is a line orbifold bundle $\mathcal{O}_\omega(1)$ on $\mathbb{P}(\omega)$, unique up to isomorphism, such that $\varphi^* \mathcal{O}_\omega(1) = \mathcal{O}(1)$. Then

$$\int_{\mathbb{P}(\omega)}^{orb} c_1(\mathcal{O}_\omega(1))^n = \frac{1}{\deg \varphi} \int_{\mathbb{P}^n} \varphi^* c_1(\mathcal{O}_\omega(1))^n = \frac{1}{\deg \varphi} \int_{\mathbb{P}^n} c_1(\mathcal{O}(1))^n = \frac{1}{\omega_0 \cdots \omega_n}.$$

Set $H = c_1(\mathcal{O}_\omega(1))$, $d = dH$ and $h_i = \deg(z_i) = \omega_i H$. Note that Proposition 3.3 is still valid here, for more details see [9, 18, 20, 21]. In this case we have

$$\begin{aligned} \sum_{p \in \text{Sing}(\mathcal{D})} \mathcal{I}_p^{orb}(\mathcal{D}) &= \sum_{j=0}^n \left\{ (-1)^j d^{n-j} \int_{\mathbb{P}(\omega)}^{orb} \mathcal{C}_j(\omega_1 H, \dots, \omega_n H) \cdot H^{n-j} \right\} \\ &= \sum_{j=0}^n \left\{ (-1)^j d^{n-j} \mathcal{C}_j(\omega) \int_{\mathbb{P}(\omega)}^{orb} H^n \right\} \\ &= \frac{1}{\omega_0 \cdots \omega_n} \sum_{j=0}^n (-1)^j \mathcal{C}_j(\omega) d^{n-j}. \end{aligned}$$

□

Example 3.4. We present three examples that give a numerical illustration of our proposition above

- (i) Consider the codimension one holomorphic foliation \mathcal{F} of degree $d = 2m$ on $\mathbb{P}(1, 1, m)$, defined by $w = mz_2 z_0^{m-1} dz_0 + mz_2 z_1^{m-1} dz_1 - (z_0^m + z_1^m) dz_2$. Set $\mu_m = \{\alpha \in \mathbb{C}^* : \alpha^m = 1\}$. Then

$$\text{Sing}(\mathcal{F}) = \{ [0 : 0 : 1]_\omega, [1 : i\alpha : 0]_\omega : \alpha \in \mu_m \}.$$

In coordinates $(U_0, \mu_m) : z_0 \neq 0$, we have $w = mz_2 z_1^{m-1} dz_1 - (1 + z_1^m) dz_2$ and $\mathcal{I}_{(i\alpha, 0)}^{orb} = 1$ for all $\alpha \in \mu_m$. In coordinates $(U_2, \{+1\}) : z_2 \neq 0$, we have $w = mz_0^{m-1} dz_0 + mz_1^{m-1} dz_1$ and $\mathcal{I}_{(0,0)}^{orb} = \frac{(m-1)^2}{m}$. Therefore, we get

$$\sum_{p \in \text{Sing}(\mathcal{F})} \mathcal{I}_p^{orb}(\mathcal{F}) = m + \frac{(m-1)^2}{m} = \frac{1}{1 \cdot 1 \cdot m} \sum_{j=0}^2 (-1)^j \mathcal{C}_j(1, 1, m) (2m)^{2-j}.$$

- (ii) Consider the codimension one holomorphic distribution \mathcal{D} of degree $d = \omega_0 + \omega_1 = \omega_2 + m\omega_3$ on $\mathbb{P}(\omega_0, \omega_1, \omega_2, \omega_3)$, defined by $w = -\omega_1 z_1 dz_0 + \omega_0 z_0 dz_1 - \omega_3 z_3^m dz_2 + \omega_2 z_2 z_3^{m-1} dz_3$. Then

$$\text{Sing}(\mathcal{D}) = \{ [0 : 0 : 1 : 0]_\omega \}.$$

In coordinates $(U_2, \mu_{\omega_2}) : z_2 \neq 0$, we have $w = -\omega_1 z_1 dz_0 + \omega_0 z_0 dz_1 + \omega_2 z_3^{m-1} dz_3$ and $\mathcal{I}_{(0,0,0)}^{orb} = \frac{m-1}{\omega_2}$. On the other hand, writing $k = \omega_2 + \omega_3$, we obtain $\mathcal{C}_1(\omega) = d + k$, $\mathcal{C}_2(\omega) = \omega_0 \omega_1 + \omega_2 \omega_3 + dk$ and $\mathcal{C}_3(\omega) = \omega_0 \omega_1 k + \omega_2 \omega_3 d$. Therefore, we have

$$\frac{1}{\omega_0 \omega_1 \omega_2 \omega_3} \sum_{j=0}^3 (-1)^j \mathcal{C}_j(\omega) d^{3-j} = \frac{\omega_0 \omega_1 (d - k)}{\omega_0 \omega_1 \omega_2 \omega_3} = \frac{m-1}{\omega_2} = \sum_{p \in \text{Sing}(\mathcal{D})} \mathcal{I}_p^{orb}(\mathcal{D}).$$

- (iii) Let \mathcal{D} be a distribution of degree d on $\mathbb{P}(1, 1, 1, k)$, $k > 1$. Suppose \mathcal{D} has only isolated singularities. If $k \nmid d^3 - 2$, then \mathcal{D} has a singularity at $\text{Sing}(\mathbb{P}(1, 1, 1, k)) = \{\bar{e}_3\}$. In fact, suppose \bar{e}_3 is non-singular for \mathcal{D} . Then the right-hand side of (4) is an integer and this happens only if $k \mid d^3 - 2$.

Conditions for the existence of such natural maps $\varphi : \mathbb{P}^n \rightarrow \mathbb{P}_\Delta$ as in Proposition 3.4 are given in [14]. More generally, it is easy to understand that the same arguments from the previous proposition can be applied here to obtain:

Proposition 3.5. *Let \mathbb{P}_Δ be an n -dimensional compact toric orbifold with only isolated singularities. Let \mathcal{D} be a codimension one distribution of degree d with isolated zeros. Suppose there is a natural map $\varphi : \mathbb{P}^n \rightarrow \mathbb{P}_\Delta$ of finite degree such that $\varphi^* \mathcal{O}_{\mathbb{P}_\Delta}([D_i]) = \mathcal{O}_{\mathbb{P}^n}(m_i)$. Set $\varphi^* \mathcal{O}_{\mathbb{P}_\Delta}(d) = \mathcal{O}_{\mathbb{P}^n}(k)$. Then*

$$\sum_{p \in \text{Sing}(\mathcal{D})} \mathcal{I}_p^{orb}(\mathcal{D}) = \frac{1}{\deg \varphi} \sum_{j=0}^n (-1)^j \mathcal{C}_j(m) k^{n-j},$$

where $\mathcal{C}_j(m)$ is the j th elementary symmetric function of the variables m_1, \dots, m_{n+r} .

4. APPLICATIONS

We will say that a codimension one distribution \mathcal{D} on \mathbb{P}_Δ is *generic* if it has at most isolated singularities, and that \mathcal{D} is *regular* if $\text{Sing}(\mathcal{D}) = \emptyset$. In the remainder of this section we will assume that all distributions are generic and we will use homogeneous coordinates.

Corollary 4.1. *Let \mathbb{P}_Δ be an n -dimensional compact toric manifold. Set $h_i = [D_i] \in \mathcal{A}_{n-1}(\mathbb{P}_\Delta)$. Let \mathcal{D} be a codimension one distribution of degree d . Then, \mathcal{D} is singular if $\gcd\{d_i\} \nmid \mathcal{C}_n(h)$.*

Proof. By Proposition 3.3, if $\text{Sing}(\mathcal{D}) = \emptyset$, then we can write

$$0 = \sum_{j=0}^n \left\{ (-1)^j \sum_{|k|=n-j} \binom{n-j}{k} \int_{\mathbb{P}_\Delta} \mathcal{C}_j(h) \cdot \prod_i (d_i h_i)^{k_i} \right\} = (-1)^n \mathcal{C}_n(h) + \sum_i a_i d_i,$$

for some $a_i \in \mathbb{Z}$. Then $\gcd\{d_i\} \mid \mathcal{C}_n(h)$. □

Similarly, the following corollary can be proved from Proposition 3.5 and Corollary 4.1:

Corollary 4.2. *Let \mathbb{P}_Δ be an n -dimensional compact toric orbifold with at most isolated singularities. Let \mathcal{D} be a codimension one distribution of degree d . Suppose there is a natural map $\varphi : \mathbb{P}^n \rightarrow \mathbb{P}_\Delta$ of finite degree such that $\varphi^* \mathcal{O}_{\mathbb{P}_\Delta}([D_i]) = \mathcal{O}_{\mathbb{P}^n}(m_i)$. Set $\varphi^* \mathcal{O}_{\mathbb{P}_\Delta}(d) = \mathcal{O}_{\mathbb{P}^n}(k)$, $k \neq 0$. Then, \mathcal{D} is regular if and only if*

$$\prod_{i=1}^{n+r} (k - m_i) = (-1)^n \sum_{i=1}^r (-1)^i \mathcal{C}_{n+i}(m) k^{r-i}.$$

Moreover, \mathcal{D} is singular if $\gcd\{d_i\} \nmid \mathcal{C}_n(m)$.

Proof. By Proposition 3.5 we have, \mathcal{D} is regular if and only if

$$\sum_{j=0}^n (-1)^j \mathcal{C}_j(m) k^{n-j} = 0.$$

Then, \mathcal{D} is regular if and only if

$$\begin{aligned} \prod_{i=1}^{n+r} (k - m_i) &= \sum_{j=0}^{n+r} (-1)^j \mathcal{C}_j(m) k^{n+r-j} \\ &= k^r \sum_{j=0}^n (-1)^j \mathcal{C}_j(m) k^{n-j} + \sum_{j=n+1}^{n+r} (-1)^j \mathcal{C}_j(m) k^{n+r-j} \\ &= (-1)^n \sum_{i=1}^r (-1)^i \mathcal{C}_{n+i}(m) k^{r-i}. \end{aligned}$$

□

Example 4.1. Consider the multiprojective space $\mathbb{P}_\Delta = \mathbb{P}^n \times \mathbb{P}^m$. Let \mathcal{D} be a codimension one distribution of degree $d = (d_1, d_2)$ on $\mathbb{P}^n \times \mathbb{P}^m$. Then, \mathcal{D} is singular if $\gcd(d_1, d_2) \nmid (n+1)(m+1)$, see Example 3.2. Similarly, it is easy to see that on $\mathbb{P}_\Delta = \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$, a codimension one distribution of degree $d = (d_1, \dots, d_k)$ is singular if $\gcd\{d_i\} \nmid \prod_{i=1}^k (n_i + 1)$.

Now let us consider low-dimensional multiprojective spaces:

Corollary 4.3. *Let \mathcal{D} be a codimension one distribution on \mathbb{P}_Δ .*

- (1) *If $\mathbb{P}_\Delta = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and \mathcal{D} regular, we have $\deg(\mathcal{D}) \in \{(2, 0, 0), (0, 2, 0), (0, 0, 2)\}$. Then, in homogeneous coordinates, \mathcal{D} is induced by*

$$\omega = z_{i1} dz_{i0} - z_{i0} dz_{i1}, \text{ for some } i = 1, 2, 3.$$

In particular \mathcal{D} is integrable and admits a rational first integral.

- (2) *If $\mathbb{P}_\Delta = \mathbb{P}^2 \times \mathbb{P}^1$ and \mathcal{D} regular, we have $\deg(\mathcal{D}) = (0, 2)$. Then, in homogeneous coordinates, \mathcal{D} is induced by*

$$\omega = z_{21} dz_{20} - z_{20} dz_{21}.$$

In particular \mathcal{D} is integrable and admits a rational first integral.

- (3) *If $\mathbb{P}_\Delta = \mathbb{P}^2 \times \mathbb{P}^2$, we have $\text{Sing}(\mathcal{D}) \neq \emptyset$, that is, there are no regular distributions.*

Proof. (1) Consider $\mathbb{P}_\Delta = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Here $h_1 = h_{10} = h_{11}$, $h_2 = h_{20} = h_{21}$ and $h_3 = h_{30} = h_{31}$, with $h_1^2 = h_2^2 = 0$ and $h_1 \cdot h_2 = 1$. Let \mathcal{D} be a codimension one regular distribution of degree $d = (d_1, d_2, d_3) = d_1 h_1 + d_2 h_2 + d_3 h_3$ on \mathbb{P}_Δ . By Proposition 3.3 we have

$$\sum_{j=0}^3 (-1)^j \mathbf{C}_j(h) (d_1 h_1 + d_2 h_2 + d_3 h_3)^{3-j} = 0.$$

Substituting $\mathbf{C}_0(h) = 1$, $\mathbf{C}_1(h) = 2(h_1 + h_2 + h_3)$, $\mathbf{C}_2(h) = 4(h_1 h_2 + h_1 h_3 + h_2 h_3)$ and $\mathbf{C}_3(h) = 8 h_1 h_2 h_3$ we obtain

$$3 d_1 d_2 d_3 - 2(d_1 d_2 + d_1 d_3 + d_2 d_3) + 2(d_1 + d_2 + d_3) - 4 = 0.$$

Since at least one $d_i \geq 1$, the integer solutions are given by $d = (1, 1, 2)$, $d = (1, -1, -2)$ and $d = (2, 0, 0)$ (and permutation of d_i). Using homogeneous coordinates, \mathcal{D} is induced by a polynomial form

$$\omega = \sum_{i=1}^3 (P_{i0} dz_{i0} + P_{i1} dz_{i1}),$$

where $\deg(P_{10}) = \deg(P_{11}) = (d_1 - 1, d_2, d_3)$, $\deg(P_{20}) = \deg(P_{21}) = (d_1, d_2 - 1, d_3)$ and $\deg(P_{30}) = \deg(P_{31}) = (d_1, d_2, d_3 - 1)$. Also, $i_{R_i} \omega = 0$ for $R_i = z_{i0} \frac{\partial}{\partial z_{i0}} + z_{i1} \frac{\partial}{\partial z_{i1}}$, $i = 1, 2, 3$. So, it is easy to check that $d = (2, 0, 0)$ (and permutation of d_i), and so \mathcal{D} is induced by

$$\omega = z_{i1} dz_{i0} - z_{i0} dz_{i1}, \text{ for some } i = 1, 2, 3.$$

(2) Consider $\mathbb{P}_\Delta = \mathbb{P}^2 \times \mathbb{P}^1$. Here $h_1 = h_{10} = h_{11} = h_{12}$ and $h_2 = h_{20} = h_{21}$, with $h_1^3 = h_2^2 = 0$ and $h_1^2 \cdot h_2 = 1$. Let \mathcal{D} be a codimension one regular distribution of degree $d = (d_1, d_2) = d_1 h_1 + d_2 h_2$ on \mathbb{P}_Δ . By Proposition 3.3 we have

$$\sum_{j=0}^3 (-1)^j \mathbf{C}_j(h) (d_1 h_1 + d_2 h_2)^{3-j} = 0.$$

Substituting $\mathbf{C}_0(h) = 1$, $\mathbf{C}_1(h) = 3h_1 + 2h_2$, $\mathbf{C}_2(h) = 3h_1^2 + 6h_1 h_2$ and $\mathbf{C}_3(h) = 6 h_1^2 h_2$ we obtain

$$d_2 = \frac{2(d_1^2 - 3d_1 + 3)}{3(d_1 - 1)^2}.$$

Thus, it is easy to check that $d = (0, 2)$, and so \mathcal{D} is induced by

$$\omega = z_{21} dz_{20} - z_{20} dz_{21}.$$

(3) Consider $\mathbb{P}_\Delta = \mathbb{P}^2 \times \mathbb{P}^2$. Here $h_1 = h_{10} = h_{11} = h_{12}$ and $h_2 = h_{20} = h_{21} = h_{22}$, with $h_1^3 = h_2^3 = 0$ and $h_1^2 \cdot h_2^2 = 1$. Let \mathcal{D} be a codimension one distribution of degree $d = (d_1, d_2) = d_1 h_1 + d_2 h_2$ on \mathbb{P}_Δ . By Proposition 3.3 we have

$$\# \text{Sing}(\mathcal{D}) = \sum_{j=0}^4 (-1)^j \mathbf{C}_j(h) (d_1 h_1 + d_2 h_2)^{4-j}.$$

Substituting $\mathbf{C}_0(h) = 1$, $\mathbf{C}_1(h) = 3h_1 + 3h_2$, $\mathbf{C}_2(h) = 3h_1^2 + 9h_1 h_2 + 3h_2^2$, $\mathbf{C}_3(h) = 9h_1^2 h_2 + 9h_1 h_2^2$ and $\mathbf{C}_4(h) = 9h_1^2 h_2^2$ we obtain

$$\# \text{Sing}(\mathcal{D}) = 2d_1^2 d_2^2 - 3d_1 d_2^2 - 3d_1^2 d_2 + d_2^2 + 6d_1 d_2 + d_1^2 - 3d_2 - 3d_1 + 3.$$

Hence, it is easy to check that $\#\text{Sing}(\mathcal{D}) \neq 0$ for all $d \in \mathbb{Z} \times \mathbb{Z}$. \square

Example 4.2. Consider the Hirzebruch surface \mathcal{H}_r , $r \geq 0$. Here $\text{Pic}(\mathcal{H}_r) = \mathbb{Z}h_1 \oplus \mathbb{Z}h_2$, where $h_1^2 = 0$, $h_1 \cdot h_2 = 1$ and $h_2^2 = -r$. Moreover $h_3 = h_1$ and $h_4 = rh_1 + h_2$; see [12]. In particular $\mathcal{C}_2(h) = \sum_{1 \leq i < j \leq 4} h_i \cdot h_j = 4$. Let \mathcal{F} be a holomorphic foliation of degree $d = (d_1, d_2)$ on \mathcal{H}_r . Then, \mathcal{F} is singular if $\gcd(d_1, d_2) \nmid 4$. Noting that $\mathcal{H}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$.

In [6, pp. 37], M. Brunella shows that if \mathcal{F} is a regular foliation on \mathcal{H}_r , then \mathcal{F} is a \mathbb{P}^1 -fibration over \mathbb{P}^1 . Let us give a different proof of this fact. By using homogeneous coordinates, we will give normal forms for these foliations:

Corollary 4.4. *Let \mathcal{F} be a holomorphic foliation of degree $d = (d_1, d_2)$ on \mathcal{H}_r . Then, \mathcal{F} is regular if and only if*

- (1) $d_1 = 0$ and $d_2 = 2$, for $r = 0$. Then, in homogeneous coordinates, \mathcal{F} is induced by

$$\omega = z_{2,2}dz_{1,2} - z_{1,2}dz_{2,2},$$

or

- (2) $d_1 = 2$ and $d_2 = 0$, for $r \geq 0$. Then, in homogeneous coordinates, \mathcal{F} is induced by

$$\omega = z_{2,1}dz_{1,1} - z_{1,1}dz_{2,1}.$$

In particular \mathcal{F} is a \mathbb{P}^1 -fibration over \mathbb{P}^1 ; \mathcal{F} is given by the canonical projection $\mathcal{H}_r \rightarrow \mathbb{P}^1$. Moreover, there are no holomorphic foliations with a unique singularity (with multiplicity 1).

Proof. Set $d = d_1h_1 + d_2h_2$. By Proposition 3.3 we have, \mathcal{F} is regular if and only if

$$\begin{aligned} 0 &= \sum_{j=0}^2 (-1)^j d^{2-j} \mathcal{C}_j(h) \\ &= d^2 - d\mathcal{C}_1(h) + \mathcal{C}_2(h) \\ &= 2d_1d_2 - d_2^2r - (d_1h_1 + d_2h_2) \left((2+r)h_1 + 2h_2 \right) + 4 \\ &= 2d_1d_2 - 2d_1 - 2d_2 + 4 - d_2(d_2 - 1)r \\ &= 2(d_1 - 1)(d_2 - 1) + 2 - d_2(d_2 - 1)r. \end{aligned}$$

Then, \mathcal{F} is regular if and only if $(d_2 - 1)(d_2r - 2(d_1 - 1)) = 2$. Then $d_2 = 0$, $d_2 = 2$ or $d_2 = 3$. If $d_2 = 0$, then $d_1 = 2$. If $d_2 = 2$, then $d_1 = r$. If $d_2 = 3$, then we have $2d_1 - 3r = 1$. Now, let us show $d_1 = r > 0$ and $d_2 = 2$ is not possible. Suppose $d_1 = r > 0$ and $d_2 = 2$, then \mathcal{F} is induced by a polynomial form $\omega = \sum P_{i,j}dz_{i,j}$, where $\deg(P_{1,1}) = \deg(P_{2,1}) = (d_1 - 1, d_2) = (r - 1, 2)$, $\deg(P_{1,2}) = (d_1, d_2 - 1) = (r, 1)$ and $\deg(P_{2,2}) = (d_1 - r, d_2 - 1) = (0, 1)$. Then, the polynomials $P_{i,j}$ are of the form: $P_{1,1} = Az_{1,2}^2$, $P_{2,1} = Bz_{1,2}^2$, $P_{1,2} = Cz_{1,2} + \alpha z_{2,2}$ and $P_{2,2} = \beta z_{1,2}$, for certain polynomials $A, B, C \in \mathbb{C}[z_{1,1}, z_{2,1}]$ and $\alpha, \beta \in \mathbb{C}$. Since $i_{R_2}\omega = 0$, we have

$$0 = z_{1,2}P_{1,2} + z_{2,2}P_{2,2} = Cz_{1,2}^2 + \alpha z_{1,2}z_{2,2} + \beta z_{2,2}z_{1,2},$$

which implies that $\alpha + \beta = 0$ and $C = 0$. On the other hand, since $i_{R_1}\omega = 0$, we have

$$0 = z_{1,1}P_{1,1} + z_{2,1}P_{2,1} + rz_{2,2}P_{2,2} = z_{1,1}Az_{1,2}^2 + z_{2,1}Bz_{1,2}^2 + r\beta z_{2,2}z_{1,2},$$

which implies that $\beta = 0$. Therefore \mathcal{F} is induced by

$$\omega = P_{1,1}dz_{1,1} + P_{2,1}dz_{2,1} = z_{1,2}^2(Adz_{1,1} + Bdz_{2,1}),$$

that contradicts the fact that \mathcal{F} is regular. Now, let us show $d_2 = 3$ and $2d_1 - 3r = 1$ is not possible. Suppose $d_2 = 3$ and $2d_1 - 3r = 1$, then \mathcal{F} is induced by a polynomial form $\omega = \sum P_{i,j}dz_{i,j}$, where $\deg(P_{1,1}) = \deg(P_{2,1}) = (d_1 - 1, d_2) = (\frac{3r-1}{2}, 3)$, $\deg(P_{1,2}) = (d_1, d_2 - 1) = (\frac{3r+1}{2}, 2)$ and $\deg(P_{2,2}) = (d_1 - r, d_2 - 1) = (\frac{r+1}{2}, 2)$. Then, the polynomials $P_{i,j}$ are of the form: $P_{1,1} = A_1z_{1,2}^3 + A_2z_{1,2}^2z_{2,2}$, $P_{2,1} = B_1z_{1,2}^3 + B_2z_{1,2}^2z_{2,2}$, $P_{1,2} = C_1z_{1,2}^2 + C_2z_{1,2}z_{2,2} + \alpha z_{2,2}^2$ and $P_{2,2} = Dz_{1,2}^2 + \beta z_{1,2}z_{2,2}$, for certain polynomials $A_1, A_2, B_1, B_2, C_1, C_2, D \in \mathbb{C}[z_{1,1}, z_{2,1}]$ and $\alpha, \beta \in \mathbb{C}$, with $\alpha = \beta = 0$ if $r \neq 1$. Since $i_{R_1}\omega = 0$, we have $z_{1,1}P_{1,1} + z_{2,1}P_{2,1} + rz_{2,2}P_{2,2} = 0$, which implies that $\beta = 0$. On the other hand, since $i_{R_2}\omega = 0$, we have $z_{1,2}P_{1,2} + z_{2,2}P_{2,2} = 0$, which implies that $\alpha + \beta = 0$. Therefore \mathcal{F} is induced by

$$\omega = z_{1,2}\omega_0,$$

that contradicts the fact that \mathcal{F} is regular.

Let us see the case $d_1 = r = 0$ and $d_2 = 2$. Then \mathcal{F} is induced by a polynomial form $\omega = \sum P_{i,j}dz_{i,j}$ with $\deg(P_{1,1}) = \deg(P_{2,1}) = (d_1 - 1, d_2) = (-1, 2)$, $\deg(P_{1,2}) = (d_1, d_2 - 1) = (0, 1)$ and $\deg(P_{2,2}) = (d_1 - r, d_2 - 1) = (0, 1)$. Then $\omega = (a_{1,2}z_{1,2} + a_{2,2}z_{2,2})dz_{1,2} + (b_{1,2}z_{1,2} + b_{2,2}z_{2,2})dz_{2,2}$. Since $i_{R_2}\omega = 0$, we have $a_{1,2} = b_{2,2} = 0$ and $a_{2,2} + b_{1,2} = 0$. Therefore \mathcal{F} is induced by

$$\omega = k(z_{2,2}dz_{1,2} - z_{1,2}dz_{2,2}), \quad k \in \mathbb{C}^*.$$

Similarly, if $d_1 = 2$ and $d_2 = 0$, then \mathcal{F} is induced by a polynomial form $\omega = \sum P_{i,j}dz_{i,j}$ with $\deg(P_{1,1}) = \deg(P_{2,1}) = (d_1 - 1, d_2) = (1, 0)$, $\deg(P_{1,2}) = (d_1, d_2 - 1) = (2, -1)$ and $\deg(P_{2,2}) = (d_1 - r, d_2 - 1) = (2 - r, -1)$. Then $\omega = (a_{1,1}z_{1,1} + a_{2,1}z_{2,1})dz_{1,1} + (b_{1,1}z_{1,1} + b_{2,1}z_{2,1})dz_{2,1}$. Since $i_{R_1}\omega = 0$, we have $a_{1,1} = b_{2,1} = 0$ and $a_{2,1} + b_{1,1} = 0$. Therefore \mathcal{F} is induced by

$$\omega = k(z_{2,1}dz_{1,1} - z_{1,1}dz_{2,1}), \quad k \in \mathbb{C}^*.$$

Finally, \mathcal{F} has a unique singularity if and only if $(d_2 - 1)(d_2r - 2(d_1 - 1)) = -1$. Now, it is easy to see that this last equation has no integer solutions. \square

Example 4.3. Consider the n -dimensional rational normal scroll $\mathbb{P}_\Delta = \mathbb{F}(a) = \mathbb{F}(a_1, \dots, a_n)$, $a_1, \dots, a_n \in \mathbb{Z}$. Here $\text{Pic}(\mathbb{F}(a)) = \mathbb{Z}L \oplus \mathbb{Z}M$, subject to the relations $L^2 = 0$, $M^n = |a| = \sum_{i=1}^n a_i$ and $M^{n-1} \cdot L = 1$. Moreover $h_1 = h_2 = L$ and $h'_i = h_{i+2} = -a_iL + M$ for all $i = 1, \dots, n$; see [27]. In particular we have

$$\begin{aligned} \mathbf{C}_n(h) &= \mathbf{C}_n(h_1, h_2, h'_1, \dots, h'_n) \\ &= h'_1 \cdots h'_n + 2 \sum_{i_1 < \cdots < i_{n-1}} L \cdot h'_{i_1} \cdots h'_{i_{n-1}} \\ &= M^n - \left(\sum_{i=1}^n a_i \right) M^{n-1} \cdot L + 2 \sum_{i_1 < \cdots < i_{n-1}} L \cdot M^{n-1} \\ &= 2n. \end{aligned}$$

Let \mathcal{D} be a codimension one distribution of degree $d = (d_1, d_2)$ on $\mathbb{F}(a)$. Then, \mathcal{D} is singular if $\gcd(d_1, d_2) \nmid 2n$. Noting that $\mathbb{F}(-a, 0) \simeq \mathcal{H}_a$. More generally we prove the following theorem:

Theorem 4.1. *Let \mathcal{D} be a codimension one distribution of degree $d = (d_1, d_2)$ on $\mathbb{F}(a) = \mathbb{F}(a_1, \dots, a_n)$, $n > 2$. Then, \mathcal{D} is regular if and only if $d_1 = 2$ and $d_2 = 0$. Then, in homogeneous coordinates, \mathcal{D} is induced by*

$$\omega = z_{1,2}dz_{1,1} - z_{1,1}dz_{1,2}.$$

In particular \mathcal{D} is integrable and admits a rational first integral. Moreover \mathcal{D} is the natural fibration given by $\mathbb{F}(a) \rightarrow \mathbb{P}^1$.

Proof. Set $d = d_1L + d_2M$ and put $|a| = \sum a_i$. Then, \mathcal{D} is regular if and only if

$$(5) \quad \sum_{i=0}^n (-1)^i d^{n-i} \mathbf{C}_i(h) = 0.$$

On the other hand, for $m > 0$ we have

$$d^m = \sum_{i=0}^m \binom{m}{i} (d_1L)^{m-i} (d_2M)^i = md_1d_2^{m-1}LM^{m-1} + d_2^mM^m,$$

and

$$\begin{aligned} \mathbf{C}_m(h) &= 2 \sum_{i_1 < \dots < i_{m-1}} L \prod_{k=1}^{m-1} (-a_{i_k}L + M) + \sum_{i_1 < \dots < i_m} \prod_{k=1}^m (-a_{i_k}L + M) \\ &= 2 \sum_{i_1 < \dots < i_{m-1}} LM^{m-1} + \sum_{i_1 < \dots < i_m} \left(-(\sum_{k=1}^m a_k) LM^{m-1} + M^m \right) \\ &= 2 \binom{n}{m-1} LM^{m-1} - \left(\sum_{i_1 < \dots < i_m} (\sum_{k=1}^m a_k) \right) LM^{m-1} + \binom{n}{m} M^m \\ &= 2 \binom{n}{m-1} LM^{m-1} - |a| \binom{n-1}{m-1} LM^{m-1} + \binom{n}{m} M^m \\ &= \left(2 \binom{n}{m-1} - |a| \binom{n-1}{m-1} \right) LM^{m-1} + \binom{n}{m} M^m. \end{aligned}$$

Then, for $0 < i < n$ we have

$$d^{n-i} \mathbf{C}_i(h) = (n-i) \binom{n}{i} d_1 d_2^{n-i-1} + |a| \binom{n}{i} d_2^{n-i} + \left(2 \binom{n}{i-1} - |a| \binom{n-1}{i-1} \right) d_2^{n-i}.$$

Substituting into (5) we obtain

$$\begin{aligned}
 0 &= (nd_1d_2^{n-1} + |a|d_2^n) + \sum_{i=1}^{n-1} (-1)^i d^{n-i} \mathbf{C}_i(h) + (-1)^n \mathbf{C}_n(h) \\
 &= nd_1d_2^{n-1} + |a|d_2^n + \sum_{i=1}^{n-1} (-1)^i (n-i) \binom{n}{i} d_1d_2^{n-i-1} + |a| \sum_{i=1}^{n-1} (-1)^i \binom{n}{i} d_2^{n-i} \\
 &\quad + \sum_{i=1}^{n-1} (-1)^i \left(2 \binom{n}{i-1} - |a| \binom{n-1}{i-1} \right) d_2^{n-i} + (-1)^n 2n \\
 &= d_1 \sum_{i=0}^{n-1} (-1)^i (n-i) \binom{n}{i} d_2^{n-i-1} - 2 \sum_{i=1}^{n-1} (-1)^{i-1} \binom{n}{i-1} d_2^{n-i} + (-1)^n 2n \\
 &\quad + |a| \left(d_2^n + \sum_{i=1}^{n-1} (-1)^i \left(\binom{n}{i} - \binom{n-1}{i-1} \right) d_2^{n-i} \right) \\
 &= nd_1 \sum_{i=0}^{n-1} (-1)^i \binom{n-1}{i} d_2^{n-1-i} - 2 \left(\sum_{i=1}^{n-1} (-1)^{i-1} \binom{n}{i-1} d_2^{n-i} + (-1)^n (1-n) \right) + 2(-1)^n \\
 &\quad + |a| d_2 \left(d_2^{n-1} + \sum_{i=1}^{n-1} (-1)^i \binom{n-1}{i} d_2^{n-1-i} \right) \\
 &= nd_1(d_2 - 1)^{n-1} - 2P(d_2) + 2(-1)^n + |a| d_2 (d_2 - 1)^{n-1},
 \end{aligned}$$

where $P(t) = \sum_{i=1}^{n-1} (-1)^{i-1} \binom{n}{i-1} t^{n-i} + (-1)^n (1-n) = t \sum_{i=0}^{n-2} (-1)^i \binom{n}{i} t^{n-2-i} + (-1)^n (1-n)$. Note that $P(1) = \sum_{i=0}^{n-2} (-1)^i \binom{n}{i} + (-1)^n (1-n) = \sum_{i=0}^n (-1)^i \binom{n}{i} = 0$, so there exists a polynomial $Q(t) \in \mathbb{Z}[t]$ such that $P(t) = (t-1)Q(t)$. Then we have, \mathcal{D} is regular if and only if

$$(6) \quad (d_2 - 1) \left((nd_1 + |a|d_2)(d_2 - 1)^{n-2} - 2Q(d_2) \right) = 2(-1)^{n+1}.$$

Therefore $d_2 \in \{-1, 0, 2, 3\}$. Note that $d_2 \neq -1, 3$ because $n > 2$ in (6). If $d_2 = 0$, substituting into (6) we have

$$d_1 = \frac{2(-1)^n + 2Q(0)}{n(-1)^{n-2}} = \frac{2(-1)^n - 2(-1)^n(1-n)}{n(-1)^{n-2}} = 2.$$

Finally, if $d_2 = 2$, substituting into (6) we have

$$\begin{aligned}
d_1 &= \frac{2P(2) + 2(-1)^{n+1} - 2|a|}{n} \\
&= \frac{2}{n}(P(2) - (-1)^n - |a|) \\
&= \frac{2}{n} \left(2 \sum_{i=0}^{n-2} (-1)^i \binom{n}{i} 2^{n-2-i} - n(-1)^n - |a| \right). \\
&= \frac{1}{n} \left(\sum_{i=0}^{n-2} (-1)^i \binom{n}{i} 2^{n-i} - 2n(-1)^n - 2|a| \right). \\
&= \frac{1}{n} \left(\sum_{i=0}^n (-1)^i \binom{n}{i} 2^{n-i} - (-1)^{n-1} \binom{n}{n-1} 2^1 - (-1)^n \binom{n}{n} 2^0 - 2n(-1)^n - 2|a| \right) \\
&= \frac{1}{n} \left((2-1)^n + 2n(-1)^n - (-1)^n - 2n(-1)^n - 2|a| \right) \\
&= \frac{1}{n} (1 + (-1)^{n+1} - 2|a|).
\end{aligned}$$

Suppose $d_2 = 2$ and $d_1 = \frac{1}{n}(1 + (-1)^{n+1} - 2|a|)$. Then \mathcal{D} is induced by a polynomial form $\omega = \sum P_{i,j} dz_{i,j}$, where $\deg(P_{1,1}) = \deg(P_{1,2}) = (d_1 - 1, d_2) = (d_1 - 1, 2)$ and $\deg(P_{2,i}) = (d_1 + a_i, d_2 - 1) = (d_1 + a_i, 1)$. Then, the polynomials $P_{i,j}$ are of the form:

$$P_{1,1} = \sum_{q_i+q_j=2} A_{ij} z_{2,i}^{q_i} z_{2,j}^{q_j}, \quad P_{1,2} = \sum_{q_i+q_j=2} B_{ij} z_{2,i}^{q_i} z_{2,j}^{q_j}, \quad \text{and} \quad P_{2,i} = \sum_{i=1}^n C_i z_{2,i},$$

for certain polynomials $A_{ij}, B_{ij}, C_i \in \mathbb{C}[z_{1,1}, z_{1,2}]$. Since $i_{R_2}\omega = 0$, we have $0 = \sum_{i=1}^n z_{2,i} P_{2,i} = \sum_{i=1}^n C_i z_{2,i}^2$, which implies that $C_i = 0$. On the other hand, since $i_{R_1}\omega = 0$, we have $z_{1,1}P_{1,1} + z_{1,2}P_{1,2} = 0$, which implies that $P_{1,1} = z_{1,2}Q$ and $P_{1,2} = -z_{1,1}Q$, for some non-constant polynomial Q . Therefore \mathcal{D} is induced by

$$\omega = Q\omega_0,$$

that contradicts the fact that \mathcal{D} is regular.

Let us see the case $d_1 = 2$ and $d_2 = 0$. Then \mathcal{D} is induced by a polynomial form $\omega = \sum P_{i,j} dz_{i,j}$ with $\deg(P_{1,1}) = \deg(P_{1,2}) = (1, 0)$ and $\deg(P_{2,i}) = (2 + a_i, -1)$ for all $i = 0, \dots, n$. Then $\omega = (a_{1,1}z_{1,1} + a_{1,2}z_{1,2}) dz_{1,1} + (b_{1,1}z_{1,1} + b_{1,2}z_{1,2}) dz_{1,2}$. Since $i_{R_1}\omega = 0$, we have $a_{1,1} = b_{1,2} = 0$ and $a_{1,2} + b_{1,1} = 0$. Therefore \mathcal{D} is induced by

$$\omega = k(z_{1,2}dz_{1,1} - z_{1,1}dz_{1,2}), \quad k \in \mathbb{C}^*.$$

Note that $\text{Sing}(\omega) = Z(z_{1,1}, z_{1,2}) \subset \mathcal{Z}$. □

The following corollary characterize all regular distributions on a well formed weighted projective space:

Corollary 4.5. *Let $\mathbb{P}(\omega)$ be a well formed n -dimensional weighted projective space. Let \mathcal{D} be a codimension one distribution of degree d on $\mathbb{P}(\omega)$. Then \mathcal{D} is regular if and only if n is odd and*

$$\prod_{i=0}^n \omega_i = \prod_{i=0}^n (d - \omega_i).$$

Proof. By Corollary 4.2 ($r = 1$) we have, \mathcal{D} is regular if and only if

$$\prod_{i=0}^n (d - \omega_i) = (-1)^{n+1} \mathbf{c}_{n+1}(\omega).$$

Suppose n is even, then there exists i_0 such that $d - \omega_{i_0} < 0$. This is a contradiction since in homogeneous coordinates, \mathcal{D} is induced by a polynomial form $\Omega = \sum_i P_i dz_i$ with $\deg(P_i) = d - \omega_i$ for all $i = 0, \dots, n$, and so n is odd. \square

Finally, let us use Corollary 4.5 to build a family of examples

Example 4.4. Let $\omega_0, \omega_1, \dots, \omega_{2m+1}$ be positive integers, pairwise coprime and d such that $d = \omega_i + \omega_{2m+1-i}$ for all $i = 0, \dots, m$. Let us consider the $(2m + 1)$ -dimensional well formed weighted projective space $\mathbb{P}(\omega) = \mathbb{P}(\omega_0, \omega_1, \dots, \omega_{2m+1})$. In homogeneous coordinates, let Ω be a form

$$\Omega = \sum_{i=0}^m (\omega_{2m+1-i} z_{2m+1-i} dz_i - \omega_i z_i dz_{2m+1-i}).$$

Note that $i_R \Omega = 0$, where $R = \sum_i \omega_i z_i \frac{\partial}{\partial z_i}$. Then \mathcal{D} is a regular distribution of degree d .

ACKNOWLEDGMENTS. I would like to be thankful to Mauricio Corrêa for interesting conversations during the preparation of this paper. I also thank Alan Muniz and Arturo Fernández for useful suggestions which improved the readability of this paper.

REFERENCES

- [1] Adem A., Leida J. and Ruan Y.: *Orbifolds and String Topology*. Cambridge University Press, Cambridge, UK, ISBN 0-511-28288-5 (2007).
- [2] Araujo C., Corrêa M. and Massarenti A.: *Codimension one Fano distributions on Fano manifolds*. Communications in Contemporary Mathematics. Vol. 20, No. 05 (2017).
- [3] Batyrev V. and Cox D. A.: *On the Hodge Structure of Projective Hypersurfaces in Toric Varieties*. Duke Mathematical Journal. Vol. 75, No. 2, 293-338 (1994).
- [4] Blache R. *Chern classes and Hirzebruch-Riemann-Roch theorem for coherent sheaves on complex-projective orbifolds with isolated singularities*. Mathematische Zeitschrift 222: 7-57 (1996).
- [5] Brasselet J. P.: *Introduction to Toric Varieties*. IMPA, Rio de Janeiro (2008).
- [6] Brunella M.: *Birational geometry of foliations* IMPA Monographs, Springer (2015).
- [7] Ding W. and Tian G.: *Kähler-Einstein metrics and the generalized Futaki invariant*. Invent. Math. 110, 315-335 (1992).
- [8] Corrêa Jr M.: *Darboux-Jouanolou-Ghys integrability for one-dimensional foliations on toric varieties*. Bulletin des Sciences Mathématiques. Vol. 134, No. 7, 693-704 (2010).
- [9] Corrêa Jr M., Rodríguez A. M. and Soares M. G.: *A Bott type residue formula on complex orbifolds*. Int. Math. Res. Notices (2015).
- [10] Corrêa Jr M. and Soares M. G.: *A Poincaré type inequality for one-dimensional multiprojective foliations*. Bulletin of the Brazilian Mathematical Society, New Series. 42 (3), 485-503 (2011).

- [11] Cox D. A. and Katz S.: *Mirror Symmetry and Algebraic Geometry*. Mathematical Surveys and Monographs. American Mathematical Society. Vol. 68 (1999).
- [12] Cox D. A., Little J. and Schenck H.: *Toric Varieties*. Graduate Studies in Mathematics. American Mathematical Society. Vol. 124 (2011).
- [13] Cox D. A.: *The Homogeneous Coordinate Ring of a Toric Variety*. J. Algebraic Geometry 4, 17-50 (1995).
- [14] Cox D. A.: *The functor of a smooth toric variety*. Tohoku Mathematical Journal, Second Series. Vol. 47, 251-262 (1995).
- [15] Dolgachev I.: *Weighted projective varieties*. Lecture Notes in Mathematics. Springer Verlag. Vol. 956 (1982).
- [16] Dumitrescu O., Postinghel E. and Urbinati S.: *Cones of effective divisors on the blown-up \mathbb{P}^3 in general lines*. Rend. Circ. Mat. Palermo, II. Ser 66, 205-216 (2017).
- [17] Fulton W.: *Introduction to Toric Varieties*. Princeton University Press (1993).
- [18] Griffiths P. and Harris J.: *Principles of Algebraic Geometry*. John Wiley & Sons, New York (1978).
- [19] Ichiro Satake: *The Gauss-Bonnet theorem for V-manifolds*. J. Math. Soc. Japan. Vol. 9: 464-492 (1957).
- [20] Izawa T.: *Residues of codimension one singular holomorphic distributions*. Bull. Braz. Math. Soc., New Series (2008).
- [21] Jouanolou J. P.: *Équations de Pfaff algébriques*. Lecture Notes in Math., Springer. Vol. 708 (1979).
- [22] Mann E.: *Cohomologie quantique orbifolde des espaces projectifs à poids*. J. Algebraic Geometry 17, 137-166 (2008).
- [23] Neto A. L.: *Componentes irredutíveis dos espaços de folheações*. Publicações Matemáticas do IMPA (2007).
- [24] Oda T.: *Convex bodies and algebraic geometry*. Springer-Verlag (1985).
- [25] Ottem J. C.: *Cox rings of projective varieties* (2009).
<https://www.duo.uio.no/bitstream/handle/10852/10769/1/Master.pdf>.
- [26] Smith Paul: *A non-commutative homogeneous coordinate ring for the degree six del Pezzo surface*. Journal of Algebra. Vol 354, 95-109 (2012).
- [27] Reid M.: *Chapters on algebraic surfaces*. AMS (1996).
- [28] Rodríguez A. M.: *The Poincaré problem for foliations on compact toric orbifolds*. Geometriae Dedicata 215, 333-353 (2021).
- [29] Saito K.: *On a generalization of de-Rham lemma*. Ann. Inst. Fourier 26, no. 2, pp. 165-170 (1976).
- [30] Soares M. G.: *Lectures on point residues*. Monografias IMCA (2002).
- [31] Suwa T.: *Indices of vector fields and residues of singular holomorphic foliations*. Actualités Mathématiques, Hermann, Paris (1998).

A. M. RODRÍGUEZ, DEMAT - UFSJ, DEPARTAMENTO DE MATEMÁTICA E ESTADÍSTICA, SÃO JOÃO DEL-REI MG, BRAZIL.

Email address: miguel.rodriguez.mat@ufs.br