

Consensus for the Hegselmann-Krause model with time variable time delays

ELISA CONTINELLI & CRISTINA PIGNOTTI
Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica
Università degli Studi di L'Aquila
Via Vetoio, Loc. Coppito, 67010 L'Aquila Italy

June 27, 2022

Abstract

In this paper, we analyze a Hegselmann-Krause opinion formation model with time-variable time delay and prove that, if the influence function is always positive, then there is exponential convergence to consensus without requiring any smallness assumptions on the time delay function. The analysis is then extended to a model with distributed time delay.

1 Introduction

Multiagent systems attracted, in recent years, the attention of many researchers in several scientific disciplines, such as biology [5, 13], economics [24], robotics [4, 21], control theory [1, 3, 33, 28, 27], social sciences [2, 31, 8]. In particular, we mention the celebrated Hegselmann-Krause opinion formation model [20] (see [6, 7] for the related PDE model) and its second-order version, i.e. the Cucker-Smale model [13] introduced to describe flocking phenomena. A typical feature is the emergence of a collective behavior, namely, under quite general assumptions, solutions to such systems converge to consensus or, in the case of the CS-model, to flocking. It is also natural to include in such models time delays, taking into account the times necessary for each agent to receive information from other agents or reaction times.

In this paper, we deal with an opinion formation model with time-variable time delay. Multiagent systems with time delays have already been studied by some authors. Flocking results for the CS-model with delay have been proved [22, 9, 10, 12, 30, 19] in different settings, under a smallness assumption on the time delay size. We mention also [14] for the analysis of a thermomechanical CS-model with delay.

Concerning the Hegselmann-Krause model for opinion formation, convergence to consensus results have been proved in presence of small time delays in [11, 25, 17]. More recently, in [18], a consensus result is proved, in the case of a constant time delay, without requiring any upper bound on the time delay. We mention also [23] for a consensus result without any smallness assumptions on the time delay size but in the particular case of constant interaction coefficients. Finally, a flocking result has been recently obtained by [32] for a CS-model with constant time delay without any restrictions on the time delay size, applying a step by step procedure. We mention also [29] for a flocking result without smallness assumption on the time delay related to

a CS-model with leadership. Here, we extend the argument of [32] to the Hegselmann-Krause opinion formation model in the case of a time variable time delay. We then improve previous convergence to consensus results by removing the smallness assumption on the time variable time delay. We are also able to consider a more general influence function, without monotonicity assumptions.

Consider a finite set of $N \in \mathbb{N}$ particles, with $N \geq 2$. Let $x_i(t) \in \mathbb{R}^d$ be the opinion of the i -th particle at time t . We shall denote with $|\cdot|$ and $\langle \cdot, \cdot \rangle$ the usual norm and scalar product on \mathbb{R}^d , respectively. The interactions between the elements of the system are described by the following Hegselmann-Krause type model with variable time delays

$$\frac{d}{dt}x_i(t) = \sum_{j:j \neq i} a_{ij}(t)(x_j(t - \tau(t)) - x_i(t)), \quad t > 0, \forall i = 1, \dots, N, \quad (1.1)$$

with weights a_{ij} of the form

$$a_{ij}(t) := \frac{1}{N-1} \psi(x_i(t), x_j(t - \tau(t))), \quad \forall t > 0, \forall i, j = 1, \dots, N, \quad (1.2)$$

where $\psi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a positive function, and the time delay $\tau : [0, \infty) \rightarrow [0, \infty)$ is a continuous function satisfying

$$0 \leq \tau(t) \leq \bar{\tau}, \quad \forall t \geq 0, \quad (1.3)$$

for some positive constant $\bar{\tau}$. The initial conditions

$$x_i(s) = x_i^0(s), \quad \forall s \in [-\bar{\tau}, 0], \forall i = 1, \dots, N, \quad (1.4)$$

are assumed to be continuous functions.

The influence function ψ is assumed to be continuous. Moreover, we assume that it is bounded and we denote

$$K := \|\psi\|_\infty.$$

For existence results for the above model, we refer e.g. to [15, 16]. Here, we will concentrate on the asymptotic behavior of solutions.

For each $t \geq -\bar{\tau}$, we define the diameter $d(\cdot)$ as

$$d(t) := \max_{i,j=1,\dots,N} |x_i(t) - x_j(t)|.$$

Definition 1.1. We say that a solution $\{x_i\}_{i=1,\dots,N}$ to system (1.1) converges to *consensus* if

$$\lim_{t \rightarrow +\infty} d(t) = 0.$$

We will prove the following convergence to consensus result.

Theorem 1.1. *Assume that $\psi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a positive, bounded, continuous function. Moreover, let $x_i^0 : [-\bar{\tau}, 0] \rightarrow \mathbb{R}^d$ be a continuous function, for any $i = 1, \dots, N$. Then, every solution $\{x_i\}_{i=1,\dots,N}$ to (1.1) under the initial conditions (1.4) satisfies the exponential decay estimate*

$$d(t) \leq \left(\max_{i,j=1,\dots,N} \max_{r,s \in [-\bar{\tau}, 0]} |x_i(r) - x_j(s)| \right) e^{-\gamma(t-2\bar{\tau})}, \quad \forall t \geq 0, \quad (1.5)$$

for a suitable positive constant γ independent of N .

We will also consider the continuity type equation obtained as mean-field limit of the particle model when the number N of the agents tends to infinity. Indeed, since the constants appearing in the exponential consensus estimate for the discrete model are independent of the number of the agents, we can extend the consensus result to the related PDE model.

Moreover, we extend the results obtained for the Hegselmann-Krause model with a pointwise time delay to a model with distributed time delay, namely each agent is influenced by other agents' opinions in a certain time interval (cf. [12, 25]). In particular, we consider the system

$$\frac{d}{dt}x_i(t) = \frac{1}{h(t)} \sum_{j:j \neq i} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s)a_{ij}(t;s)(x_j(s) - x_i(t))ds, \quad t > 0, \quad \forall i = 1, \dots, N, \quad (1.6)$$

where the time delays $\tau_1 : [0, \infty) \rightarrow [0, \infty)$, $\tau_2 : [0, \infty) \rightarrow [0, \infty)$ are continuous functions satisfying

$$0 \leq \tau_1(t) < \tau_2(t) \leq \bar{\tau}, \quad \forall t \geq 0, \quad (1.7)$$

for some positive constant $\bar{\tau}$.

The communication rates $a_{ij}(t;s)$ are of the form

$$a_{ij}(t;s) := \frac{1}{N-1} \psi(x_i(t), x_j(s)), \quad \forall t \geq 0, \quad \forall i, j = 1, \dots, N, \quad (1.8)$$

where $\psi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a positive function.

Moreover, $\alpha : [0, \bar{\tau}] \rightarrow (0, +\infty)$ is a continuous weight function and

$$h(t) := \int_{\tau_1(t)}^{\tau_2(t)} \alpha(s)ds, \quad \forall t \geq 0. \quad (1.9)$$

Note that, since we assume $\tau_1(t) < \tau_2(t)$ and $\alpha(t) > 0$, $\forall t \geq 0$, then the function $h(t)$ is always positive.

As before, the initial conditions

$$x_i(s) = x_i^0(s), \quad \forall s \in [-\bar{\tau}, 0], \quad \forall i = 1, \dots, N,$$

are assumed to be continuous functions. Moreover, the influence function ψ is assumed to be continuous and bounded, and let us denote $K := \|\psi\|_\infty$.

Also in this case, we obtain an exponential consensus estimate without any restrictions on the time delays sizes. This extends and improves the analysis in [25] where a consensus estimate has been obtained, in the case $\tau_1(t) \equiv 0$, $\forall t \geq 0$, subject to a smallness assumption on the time delay size. Moreover, here, as for the pointwise time delay case, we do not require any monotonicity properties on the influence function ψ that is assumed only continuous and bounded.

Even in the distributed case, since the constants in the consensus estimate are independent of the number of the agents, one can extend the consensus theorem to the related PDE model.

The rest of the paper is organized as follows. In Sect. 2 we give some preliminary results based on continuity arguments and Gronwall's inequality. In Sect. 3 we prove our consensus result for the particle model with pointwise time variable time delay, while in Sect. 4 we formulate its extension to the related continuity type equation. Finally, in Sect. 5 we analyze the HK-model with distributed time delay (1.6).

2 Preliminaries

Let $\{x_i\}_{i=1,\dots,N}$ be solution to (1.1) under the initial conditions (1.4). In this section we present some auxiliary lemmas. We assume that the hypotheses of Theorem 1.1 are satisfied.

The following arguments generalize and extend the ones developed in [32] in the case of a Cucker-Smale model with constant time delay.

Lemma 2.1. *For each $v \in \mathbb{R}^d$ and $T \geq 0$, we have that*

$$\min_{j=1,\dots,N} \min_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle \leq \langle x_i(t), v \rangle \leq \max_{j=1,\dots,N} \max_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle, \quad (2.1)$$

for all $t \geq T - \bar{\tau}$ and $i = 1, \dots, N$.

Proof. First of all, we note that the inequalities in (2.1) are satisfied for every $t \in [T - \bar{\tau}, T]$. Now, let $T \geq 0$. Given a vector $v \in \mathbb{R}^d$, we set

$$M_T = \max_{j=1,\dots,N} \max_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle.$$

For all $\epsilon > 0$, let us define

$$K^\epsilon := \left\{ t > T : \max_{i=1,\dots,N} \langle x_i(s), v \rangle < M_T + \epsilon, \forall s \in [T, t] \right\}.$$

By continuity, we have that $K^\epsilon \neq \emptyset$. Thus, denoted with

$$S^\epsilon := \sup K^\epsilon,$$

it holds that $S^\epsilon > T$.

We claim that $S^\epsilon = +\infty$. Indeed, suppose by contradiction that $S^\epsilon < +\infty$. Note that by definition of S^ϵ it turns out that

$$\max_{i=1,\dots,N} \langle x_i(t), v \rangle < M_T + \epsilon, \quad \forall t \in (T, S^\epsilon), \quad (2.2)$$

and

$$\lim_{t \rightarrow S^{\epsilon-}} \max_{i=1,\dots,N} \langle x_i(t), v \rangle = M_T + \epsilon. \quad (2.3)$$

For all $i = 1, \dots, N$ and $t \in (T, S^\epsilon)$, we compute

$$\frac{d}{dt} \langle x_i(t), v \rangle = \frac{1}{N-1} \sum_{j:j \neq i} \psi(x_i(t), x_j(t-\tau(t))) \langle x_j(t-\tau(t)) - x_i(t), v \rangle.$$

Notice that, being $t \in (T, S^\epsilon)$, then $t - \tau(t) \in (T - \bar{\tau}, S^\epsilon)$ and

$$\langle x_j(t - \tau(t)), v \rangle < M_T + \epsilon, \quad \forall j = 1, \dots, N. \quad (2.4)$$

Moreover, (2.2) implies that

$$\langle x_i(t), v \rangle < M_T + \epsilon,$$

so that

$$M_T + \epsilon - \langle x_i(t), v \rangle \geq 0.$$

Combining this last fact with (2.4), we can write

$$\begin{aligned} \frac{d}{dt} \langle x_i(t), v \rangle &\leq \frac{1}{N-1} \sum_{j:j \neq i} \psi(x_i(t), x_j(t - \tau(t))) (M_T + \epsilon - \langle x_i(t), v \rangle) \\ &\leq K(M_T + \epsilon - \langle x_i(t), v \rangle), \quad \forall t \in (T, S^\epsilon). \end{aligned}$$

Then, from Gronwall's inequality we get

$$\begin{aligned} \langle x_i(t), v \rangle &\leq e^{-K(t-T)} \langle x_i(T), v \rangle + K(M_T + \epsilon) \int_T^t e^{-K(t-s)} ds \\ &= e^{-K(t-T)} \langle x_i(T), v \rangle + (M_T + \epsilon) e^{-Kt} (e^{Kt} - e^{KT}) \\ &= e^{-K(t-T)} \langle x_i(T), v \rangle + (M_T + \epsilon) (1 - e^{-K(t-T)}) \\ &\leq e^{-K(t-T)} M_T + M_T + \epsilon - M_T e^{-K(t-T)} - \epsilon e^{-K(t-T)} \\ &= M_T + \epsilon - \epsilon e^{-K(t-T)} \\ &= M_T + \epsilon - \epsilon e^{-K(S^\epsilon - T)}, \end{aligned}$$

for all $t \in (T, S^\epsilon)$. We have so proved that, $\forall i = 1, \dots, N$,

$$\langle x_i(t), v \rangle \leq M_T + \epsilon - \epsilon e^{-K(S^\epsilon - T)}, \quad \forall t \in (T, S^\epsilon).$$

Thus, we get

$$\max_{i=1, \dots, N} \langle x_i(t), v \rangle \leq M_T + \epsilon - \epsilon e^{-K(S^\epsilon - T)}, \quad \forall t \in (T, S^\epsilon). \quad (2.5)$$

Letting $t \rightarrow S^{\epsilon-}$ in (2.5), from (2.3) we have that

$$M_T + \epsilon \leq M_T + \epsilon - \epsilon e^{-K(S^\epsilon - T)} < M_T + \epsilon,$$

which is a contradiction. Thus, $S^\epsilon = +\infty$, which means that

$$\max_{i=1, \dots, N} \langle x_i(t), v \rangle < M_T + \epsilon, \quad \forall t > T.$$

From the arbitrariness of ϵ we can conclude that

$$\max_{i=1, \dots, N} \langle x_i(t), v \rangle \leq M_T, \quad \forall t > T,$$

from which

$$\langle x_i(t), v \rangle \leq M_T, \quad \forall t > T, \forall i = 1, \dots, N,$$

which proves the second inequality in (2.1). Now, to prove the other inequality, let $v \in \mathbb{R}^d$ and define

$$m_T = \min_{j=1, \dots, N} \min_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle.$$

Then, for all $i = 1, \dots, N$ and $t > T$, by applying the second inequality in (2.1) to the vector $-v \in \mathbb{R}^d$ we get

$$-\langle x_j(s), v \rangle = \langle x_i(t), -v \rangle \leq \max_{j=1, \dots, N} \max_{s \in [T-\bar{\tau}, T]} \langle x_j(s), -v \rangle$$

$$= - \min_{j=1,\dots,N} \min_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle = -m_T,$$

from which

$$\langle x_j(s), v \rangle \geq m_T.$$

Thus, also the first inequality in (2.1) is fulfilled. \square

We now introduce some notation.

Definition 2.1. We define

$$D_0 = \max_{i,j=1,\dots,N} \max_{s,t \in [-\bar{\tau}, 0]} |x_i(s) - x_j(t)|,$$

and in general, $\forall n \in \mathbb{N}$,

$$D_n := \max_{i,j=1,\dots,N} \max_{s,t \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} |x_i(s) - x_j(t)|.$$

Note that inequality (1.5) can be written as

$$d(t) \leq e^{-\gamma(t-2\bar{\tau})} D_0, \quad \forall t \geq 0.$$

Let us denote with $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Lemma 2.2. For each $n \in \mathbb{N}_0$ and $i, j = 1, \dots, N$, we get

$$|x_i(s) - x_j(t)| \leq D_n, \quad \forall s, t \geq n\bar{\tau} - \bar{\tau}. \quad (2.6)$$

Proof. Fix $n \in \mathbb{N}_0$ and $i, j = 1, \dots, N$. Given $s, t \geq n\bar{\tau} - \bar{\tau}$, if $|x_i(s) - x_j(t)| = 0$ then of course $D_n \geq 0 = |x_i(s) - x_j(t)|$. Thus, we can assume $|x_i(s) - x_j(t)| > 0$ and we set

$$v = \frac{x_i(s) - x_j(t)}{|x_i(s) - x_j(t)|}.$$

It turns out that v is a unit vector and, by using (2.1) with $T = n\bar{\tau}$ and the Cauchy-Schwarz inequality, we can write

$$\begin{aligned} |x_i(s) - x_j(t)| &= \langle x_i(s) - x_j(t), v \rangle = \langle x_i(s), v \rangle - \langle x_j(t), v \rangle \\ &\leq \max_{l=1,\dots,N} \max_{r \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} \langle x_l(r), v \rangle - \min_{l=1,\dots,N} \min_{r \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} \langle x_l(r), v \rangle \\ &\leq \max_{l,k=1,\dots,N} \max_{r,\sigma \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} \langle x_l(r) - x_k(\sigma), v \rangle \\ &\leq \max_{l,k=1,\dots,N} \max_{r,\sigma \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} |x_l(r) - x_k(\sigma)| |v| = D_n, \end{aligned}$$

which proves (2.6). \square

Remark 2.3. Let us note that from (2.6), in particular, it follows that

$$|x_i(s) - x_j(t)| \leq D_0, \quad \forall s, t \geq -\bar{\tau}. \quad (2.7)$$

Moreover, it holds

$$D_{n+1} \leq D_n, \quad \forall n \in \mathbb{N}_0. \quad (2.8)$$

With an analogous argument, one can find a bound on $|x_i(t)|$, uniform with respect to t and $i = 1, \dots, N$. Indeed, we have the following lemma.

Lemma 2.4. *For every $i = 1, \dots, N$, we have that*

$$|x_i(t)| \leq M^0, \quad \forall t \geq -\bar{\tau}, \quad (2.9)$$

where

$$M^0 := \max_{i=1, \dots, N} \max_{s \in [-\bar{\tau}, 0]} |x_i(s)|.$$

Proof. Given $i = 1, \dots, N$ and $t \geq -\bar{\tau}$, if $|x_i(t)| = 0$ then trivially $M^0 \geq 0 = |x_i(t)|$. On the contrary, if $|x_i(t)| > 0$, we define

$$v = \frac{x_i(t)}{|x_i(t)|},$$

which is a unit vector for which we can write

$$|x_i(t)| = \langle x_i(t), v \rangle.$$

Then, by applying (2.1) for $T = 0$ and by using the Cauchy-Schwarz inequality we get

$$\begin{aligned} |x_i(t)| &\leq \max_{j=1, \dots, N} \max_{s \in [-\bar{\tau}, 0]} \langle x_j(s), v \rangle \leq \max_{j=1, \dots, N} \max_{s \in [-\bar{\tau}, 0]} |x_j(s)| |v| \\ &= \max_{j=1, \dots, N} \max_{s \in [-\bar{\tau}, 0]} |x_j(s)| = M^0, \end{aligned}$$

which proves (2.9). \square

Remark 2.5. From the estimate (2.9), since the influence function ψ is continuous, we deduce that

$$\psi(x_i(t), x_j(t - \tau(t))) \geq \psi_0 := \min_{|y|, |z| \leq M^0} \psi(y, z) > 0, \quad (2.10)$$

for all $t \geq 0$, for all $i, j = 1, \dots, N$.

Lemma 2.6. *For all $i, j = 1, \dots, N$, unit vector $v \in \mathbb{R}^d$ and $n \in \mathbb{N}_0$ we have that*

$$\langle x_i(t) - x_j(t), v \rangle \leq e^{-K(t-t_0)} \langle x_i(t_0) - x_j(t_0), v \rangle + (1 - e^{-K(t-t_0)}) D_n, \quad (2.11)$$

for all $t \geq t_0 \geq n\bar{\tau}$.

Moreover, for all $n \in \mathbb{N}_0$, we get

$$D_{n+1} \leq e^{-K\bar{\tau}} d(n\bar{\tau}) + (1 - e^{-K\bar{\tau}}) D_n. \quad (2.12)$$

Proof. Fix $n \in \mathbb{N}_0$ and $v \in \mathbb{R}^d$ such that $|v| = 1$. We set

$$M_n = \max_{i=1, \dots, N} \max_{t \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} \langle x_i(t), v \rangle,$$

$$m_n = \min_{i=1, \dots, N} \min_{t \in [n\bar{\tau}-\bar{\tau}, n\bar{\tau}]} \langle x_i(t), v \rangle.$$

Then, it is easy to see that $M_n - m_n \leq D_n$. Now, for all $i = 1, \dots, N$ and $t \geq t_0 \geq n\bar{\tau}$ we have that

$$\begin{aligned}
\frac{d}{dt}\langle x_i(t), v \rangle &= \sum_{j:j \neq i} a_{ij}(t) \langle x_j(t - \tau(t)) - x_i(t), v \rangle \\
&= \frac{1}{N-1} \sum_{j:j \neq i} \psi(x_i(t), x_j(t - \tau(t))) (\langle x_j(t - \tau(t)), v \rangle - \langle x_i(t), v \rangle) \\
&\leq \frac{1}{N-1} \sum_{j:j \neq i} \psi(x_i(t), x_j(t - \tau(t))) (M_n - \langle x_i(t), v \rangle).
\end{aligned}$$

Note that, being $t \geq n\bar{\tau}$, $\langle x_i(t), v \rangle \leq M_n$ from (2.1). Therefore, we have that $M_n - \langle x_i(t), v \rangle \geq 0$ and we can write

$$\frac{d}{dt}\langle x_i(t), v \rangle \leq \frac{1}{N-1} K \sum_{j:j \neq i} (M_n - \langle x_i(t), v \rangle) = K(M_n - \langle x_i(t), v \rangle).$$

Thus, from the Gronwall's inequality it comes that

$$\begin{aligned}
\langle x_i(t), v \rangle &\leq e^{-K(t-t_0)} \langle x_i(t_0), v \rangle + \int_{t_0}^t K M_n e^{-K(t-t_0)+K(s-t_0)} ds \\
&= e^{-K(t-t_0)} \langle x_i(t_0), v \rangle + e^{-K(t-t_0)} M_n (e^{K(t-t_0)} - 1),
\end{aligned}$$

that is

$$\langle x_i(t), v \rangle \leq e^{-K(t-t_0)} \langle x_i(t_0), v \rangle + (1 - e^{-K(t-t_0)}) M_n. \quad (2.13)$$

On the other hand, for all $i = 1, \dots, N$ and $t \geq t_0 \geq n\bar{\tau}$ it holds that

$$\begin{aligned}
\frac{d}{dt}\langle x_i(t), v \rangle &= \frac{1}{N-1} \sum_{i:j \neq i} \psi(x_i(t), x_j(t - \tau(t))) (\langle x_j(t - \tau(t)), v \rangle - \langle x_i(t), v \rangle) \\
&\geq \frac{1}{N-1} \sum_{j:j \neq i} \psi(x_i(t), x_j(t - \tau(t))) (m_n - \langle x_i(t), v \rangle).
\end{aligned}$$

Note that from (2.1) $\langle x_i(t), v \rangle \geq m_n$ since $t \geq n\bar{\tau}$. Thus, $m_n - \langle x_i(t), v \rangle \leq 0$ and, by recalling that ψ is bounded, we get

$$\frac{d}{dt}\langle x_i(t), v \rangle \geq K(m_n - \langle x_i(t), v \rangle).$$

Hence, by using the Gronwall's inequality it turns out that

$$\langle x_i(t), v \rangle \geq e^{-K(t-t_0)} \langle x_i(t_0), v \rangle + (1 - e^{-K(t-t_0)}) m_n. \quad (2.14)$$

Therefore, for all $i, j = 1, \dots, N$ and $t \geq t_0 \geq n\bar{\tau}$, by using (2.13) and (2.14) and by recalling that $M_n - m_n \leq D_n$, we finally get

$$\begin{aligned}
\langle x_i(t) - x_j(t), v \rangle &= \langle x_i(t), v \rangle - \langle x_j(t), v \rangle \\
&\leq e^{-K(t-t_0)} \langle x_i(t_0), v \rangle + (1 - e^{-K(t-t_0)}) M_n \\
&\quad - e^{-K(t-t_0)} \langle x_j(t_0), v \rangle - (1 - e^{-K(t-t_0)}) m_n \\
&= e^{-K(t-t_0)} \langle x_i(t_0) - x_j(t_0), v \rangle + (1 - e^{-K(t-t_0)}) (M_n - m_n) \\
&\leq e^{-K(t-t_0)} \langle x_i(t_0) - x_j(t_0), v \rangle + (1 - e^{-K(t-t_0)}) D_n,
\end{aligned}$$

i.e. (2.11) holds true.

Now we prove (2.12). Given $n \in \mathbb{N}_0$, let $i, j = 1, \dots, N$ and $s, t \in [n\bar{\tau}, n\bar{\tau} + \bar{\tau}]$ be such that $D_{n+1} = |x_i(s) - x_j(t)|$. Note that, if $|x_i(s) - x_j(t)| = 0$, then obviously

$$0 = D_{n+1} \leq e^{-K\bar{\tau}}d(n\bar{\tau}) + (1 - e^{-K\bar{\tau}})D_n.$$

So we can assume $|x_i(s) - x_j(t)| > 0$. Let us define the unit vector

$$v = \frac{x_i(s) - x_j(t)}{|x_i(s) - x_j(t)|}.$$

Hence, we can write

$$D_{n+1} = \langle x_i(s) - x_j(t), v \rangle = \langle x_i(s), v \rangle - \langle x_j(t), v \rangle.$$

Now, by using (2.13) with $t_0 = n\bar{\tau}$, we have that

$$\begin{aligned} \langle x_i(s), v \rangle &\leq e^{-K(s-n\bar{\tau})} \langle x_i(n\bar{\tau}), v \rangle + (1 - e^{-K(s-n\bar{\tau})})M_n \\ &= e^{-K(s-n\bar{\tau})} (\langle x_i(n\bar{\tau}), v \rangle - M_n) + M_n. \end{aligned}$$

Thus, since $s \leq n\bar{\tau} + \bar{\tau}$ and $\langle x_i(n\bar{\tau}), v \rangle - M_n \leq 0$ from (2.1), we get

$$\begin{aligned} \langle x_i(s), v \rangle &\leq e^{-K\bar{\tau}} (\langle x_i(n\bar{\tau}), v \rangle - M_n) + M_n \\ &\leq e^{-K\bar{\tau}} \langle x_i(n\bar{\tau}), v \rangle + (1 - e^{-K\bar{\tau}})M_n. \end{aligned} \tag{2.15}$$

Similarly, by taking into account of (2.1) and (2.14), we have that

$$\langle x_j(t), v \rangle \geq e^{-K\bar{\tau}} \langle x_j(n\bar{\tau}), v \rangle + (1 - e^{-K\bar{\tau}})m_n. \tag{2.16}$$

Therefore, combining (2.15) and (2.16), we can write

$$\begin{aligned} D_{n+1} &\leq e^{-K\bar{\tau}} \langle x_i(n\bar{\tau}), v \rangle + (1 - e^{-K\bar{\tau}})M_n - e^{-K\bar{\tau}} \langle x_j(n\bar{\tau}), v \rangle - (1 - e^{-K\bar{\tau}})m_n \\ &= e^{-K\bar{\tau}} \langle x_i(n\bar{\tau}) - x_j(n\bar{\tau}), v \rangle + (1 - e^{-K\bar{\tau}})(M_n - m_n). \end{aligned}$$

Then, by recalling that $M_n - m_n \leq D_n$ and by using the Cauchy-Schwarz inequality, we can conclude that

$$\begin{aligned} D_{n+1} &\leq e^{-K\bar{\tau}} |x_i(n\bar{\tau}) - x_j(n\bar{\tau})| |v| + (1 - e^{-K\bar{\tau}})D_n \\ &\leq e^{-K\bar{\tau}} d(n\bar{\tau}) + (1 - e^{-K\bar{\tau}})D_n. \end{aligned}$$

□

Lemma 2.7. *There exists a constant $C \in (0, 1)$, independent of $N \in \mathbb{N}$, such that*

$$d(n\bar{\tau}) \leq CD_{n-2}, \tag{2.17}$$

for all $n \geq 2$.

Proof. Trivially, if $d(n\bar{\tau}) = 0$, then of course inequality (2.17) holds for any constant $C \in (0, 1)$. So, suppose $d(n\bar{\tau}) > 0$. Let $i, j = 1, \dots, N$ be such that $d(n\bar{\tau}) = |x_i(n\bar{\tau}) - x_j(n\bar{\tau})|$. We set

$$v = \frac{x_i(n\bar{\tau}) - x_j(n\bar{\tau})}{|x_i(n\bar{\tau}) - x_j(n\bar{\tau})|}.$$

Then, v is a unit vector for which we can write

$$d(n\bar{\tau}) = \langle x_i(n\bar{\tau}) - x_j(n\bar{\tau}), v \rangle.$$

Let us define

$$\begin{aligned} M_{n-1} &= \max_{l=1, \dots, N} \max_{s \in [n\bar{\tau}-2\bar{\tau}, n\bar{\tau}-\bar{\tau}]} \langle x_l(s), v \rangle, \\ m_{n-1} &= \min_{l=1, \dots, N} \min_{s \in [n\bar{\tau}-2\bar{\tau}, n\bar{\tau}-\bar{\tau}]} \langle x_l(s), v \rangle. \end{aligned}$$

Then $M_{n-1} - m_{n-1} \leq D_{n-1}$. Now, we distinguish two different situations.

Case I. Assume that there exists $t_0 \in [n\bar{\tau} - 2\bar{\tau}, n\bar{\tau}]$ such that

$$\langle x_i(t_0) - x_j(t_0), v \rangle < 0.$$

Then from (2.11) with $n\bar{\tau} \geq t_0 \geq n\bar{\tau} - 2\bar{\tau}$ we have that

$$\begin{aligned} d(n\bar{\tau}) &\leq e^{-K(n\bar{\tau}-t_0)} \langle x_i(t_0) - x_j(t_0), v \rangle + (1 - e^{-K(n\bar{\tau}-t_0)}) D_{n-2} \\ &\leq (1 - e^{-K(n\bar{\tau}-t_0)}) D_{n-2} \leq (1 - e^{-2K\bar{\tau}}) D_{n-2}. \end{aligned}$$

Case II. Suppose that

$$\langle x_i(t) - x_j(t), v \rangle \geq 0, \quad \forall t \in [n\bar{\tau} - 2\bar{\tau}, n\bar{\tau}]. \quad (2.18)$$

Then, for every $t \in [n\bar{\tau} - \bar{\tau}, n\bar{\tau}]$ we have that

$$\begin{aligned} \frac{d}{dt} \langle x_i(t) - x_j(t), v \rangle &= \frac{1}{N-1} \sum_{l:l \neq i} \psi(x_i(t), x_l(t - \tau(t))) \langle x_l(t - \tau(t)) - x_i(t), v \rangle \\ &\quad - \frac{1}{N-1} \sum_{l:l \neq j} \psi(x_i(t), x_l(t - \tau(t))) \langle x_l(t - \tau(t)) - x_j(t), v \rangle \\ &= \frac{1}{N-1} \sum_{l:l \neq i} \psi(x_i(t), x_l(t - \tau(t))) (\langle x_l(t - \tau(t)), v \rangle - M_{n-1} + M_{n-1} - \langle x_i(t), v \rangle) \\ &\quad + \frac{1}{N-1} \sum_{l:l \neq j} \psi(x_i(t), x_l(t - \tau(t))) (\langle x_j(t), v \rangle - m_{n-1} + m_{n-1} - \langle x_l(t - \tau(t)), v \rangle) \\ &:= S_1 + S_2. \end{aligned}$$

Now, being $t \in [n\bar{\tau} - \bar{\tau}, n\bar{\tau}]$, it holds that $t - \tau(t) \in [n\bar{\tau} - 2\bar{\tau}, n\bar{\tau}]$. Therefore, both $t, t - \tau(t) \geq n\bar{\tau} - 2\bar{\tau}$ and from (2.1) we have that

$$m_{n-1} \leq \langle x_k(t), v \rangle \leq M_{n-1}, \quad m_{n-1} \leq \langle x_k(t - \tau(t)), v \rangle \leq M_{n-1}, \quad \forall k = 1, \dots, N. \quad (2.19)$$

Therefore, using (2.9), we get

$$\begin{aligned}
S_1 &= \frac{1}{N-1} \sum_{l:l \neq i} \psi(x_i(t), x_l(t - \tau(t))) (\langle x_l(t - \tau(t)), v \rangle - M_{n-1}) \\
&\quad + \frac{1}{N-1} \sum_{l:l \neq i} \psi(x_i(t), x_l(t - \tau(t))) (M_{n-1} - \langle x_i(t), v \rangle) \\
&\leq \frac{1}{N-1} \psi_0 \sum_{l:l \neq i} (\langle x_l(t - \tau(t)), v \rangle - M_{n-1}) + K(M_{n-1} - \langle x_i(t), v \rangle),
\end{aligned}$$

and

$$\begin{aligned}
S_2 &= \frac{1}{N-1} \sum_{l:l \neq j} \psi(x_i(t), x_l(t - \tau(t))) (\langle x_j(t), v \rangle - m_{n-1}) \\
&\quad + \frac{1}{N-1} \sum_{l:l \neq j} \psi(x_i(t), x_l(t - \tau(t))) (m_{n-1} - \langle x_l(t - \tau(t)), v \rangle) \\
&\leq K(\langle x_j(t), v \rangle - m_{n-1}) + \frac{1}{N-1} \psi_0 \sum_{l:l \neq j} (m_{n-1} - \langle x_l(t - \tau(t)), v \rangle).
\end{aligned}$$

Combining this last fact with (2.19) it comes that

$$\begin{aligned}
\frac{d}{dt} \langle x_i(t) - x_j(t), v \rangle &\leq K(M_{n-1} - m_{n-1} - \langle x_i(t) - x_j(t), v \rangle) \\
&\quad + \frac{1}{N-1} \psi_0 \sum_{l:l \neq i, j} (\langle x_l(t - \tau(t)), v \rangle - M_{n-1} + m_{n-1} - \langle x_l(t - \tau(t)), v \rangle) \\
&\quad + \frac{1}{N-1} \psi_0 (\langle x_j(t - \tau(t)), v \rangle - M_{n-1} + m_{n-1} - \langle x_i(t - \tau(t)), v \rangle) \\
&= K(M_{n-1} - m_{n-1}) - K \langle x_i(t) - x_j(t), v \rangle + \frac{N-2}{N-1} \psi_0 (-M_{n-1} + m_{n-1}) \\
&\quad + \frac{1}{N-1} \psi_0 (\langle x_j(t - \tau(t)), v \rangle - M_{n-1} + m_{n-1} - \langle x_i(t - \tau(t)), v \rangle).
\end{aligned}$$

Therefore, since from (2.18) $\langle x_i(t - \tau(t)) - x_j(t - \tau(t)), v \rangle \geq 0$, we get

$$\begin{aligned}
\frac{d}{dt} \langle x_i(t) - x_j(t), v \rangle &\leq K(M_{n-1} - m_{n-1}) - K \langle x_i(t) - x_j(t), v \rangle \\
&\quad + \frac{N-2}{N-1} \psi_0 (-M_{n-1} + m_{n-1}) + \frac{1}{N-1} \psi_0 (-M_{n-1} + m_{n-1}) \\
&\quad - \frac{1}{N-1} \psi_0 \langle x_i(t - \tau(t)) - x_j(t - \tau(t)), v \rangle \\
&\leq K(M_{n-1} - m_{n-1}) - K \langle x_i(t) - x_j(t), v \rangle + \psi_0 (-M_{n-1} + m_{n-1}) \\
&= (K - \psi_0) (M_{n-1} - m_{n-1}) - K \langle x_i(t) - x_j(t), v \rangle.
\end{aligned}$$

Hence, from Gronwall's inequality it comes that

$$\begin{aligned}
\langle x_i(t) - x_j(t), v \rangle &\leq e^{-K(t-n\bar{\tau}+\bar{\tau})} \langle x_i(n\bar{\tau} - \bar{\tau}) - x_j(n\bar{\tau} - \bar{\tau}), v \rangle \\
&\quad + (K - \psi_0) (M_{n-1} - m_{n-1}) \int_{n\bar{\tau} - \bar{\tau}}^t e^{-K(t-s)} ds,
\end{aligned}$$

for all $t \in [n\bar{\tau} - \bar{\tau}, n\bar{\tau}]$. In particular, for $t = n\bar{\tau}$ it comes that

$$\begin{aligned} d(n\bar{\tau}) &\leq e^{-K\bar{\tau}} \langle x_i(n\bar{\tau} - \bar{\tau}) - x_j(n\bar{\tau} - \bar{\tau}), v \rangle + \frac{K - \psi_0}{K} (M_{n-1} - m_{n-1}) (1 - e^{-K\bar{\tau}}) \\ &\leq e^{-K\bar{\tau}} |x_i(n\bar{\tau} - \bar{\tau}) - x_j(n\bar{\tau} - \bar{\tau})| |v| + \frac{K - \psi_0}{K} (M_{n-1} - m_{n-1}) (1 - e^{-K\bar{\tau}}) \\ &\leq e^{-K\bar{\tau}} d(n\bar{\tau} - \bar{\tau}) + \frac{K - \psi_0}{K} (M_{n-1} - m_{n-1}) (1 - e^{-K\bar{\tau}}) \end{aligned}$$

Then, by recalling that $M_{n-1} - m_{n-1} \leq D_{n-1}$ we get

$$\begin{aligned} d(n\bar{\tau}) &\leq e^{-K\bar{\tau}} d(n\bar{\tau} - \bar{\tau}) + \frac{K - \psi_0}{K} D_{n-1} (1 - e^{-K\bar{\tau}}) \\ &\leq e^{-K\bar{\tau}} d(n\bar{\tau} - \bar{\tau}) + \frac{K - \psi_0}{K} D_{n-1} (1 - e^{-K\bar{\tau}}). \end{aligned}$$

Finally, by using (2.6) and (2.8) we have that that

$$\begin{aligned} d(n\bar{\tau}) &\leq e^{-K\bar{\tau}} D_n + \frac{K - \psi_0}{K} D_{n-1} (1 - e^{-K\bar{\tau}}) \\ &\leq e^{-K\bar{\tau}} D_{n-2} + \frac{K - \psi_0}{K} D_{n-2} (1 - e^{-K\bar{\tau}}) \\ &= \left[1 - \frac{\psi_0}{K} (1 - e^{-K\bar{\tau}}) \right] D_{n-2}. \end{aligned} \tag{2.20}$$

Now, we set

$$C = \max \left\{ 1 - e^{-2K\bar{\tau}}, 1 - \frac{\psi_0}{K} (1 - e^{-K\bar{\tau}}) \right\} \in (0, 1). \tag{2.21}$$

Then, taking into account (2.20), we can conclude that C is the constant for which inequality (2.17) holds. \square

3 Proof of Theorem 1.1

Proof. Let $(x_i)_{i=1, \dots, N}$ be solution to (1.1), (1.4). We claim that

$$D_{n+1} \leq \tilde{C} D_{n-2}, \quad \forall n \geq 2, \tag{3.1}$$

for some constant $\tilde{C} \in (0, 1)$. Indeed, given $n \geq 2$, from (2.8), (2.12) and (2.17) we have that

$$\begin{aligned} D_{n+1} &\leq e^{-K\bar{\tau}} d(n\bar{\tau}) + (1 - e^{-K\bar{\tau}}) D_n \\ &\leq e^{-K\bar{\tau}} C D_{n-2} + (1 - e^{-K\bar{\tau}}) D_n \\ &\leq e^{-K\bar{\tau}} C D_{n-2} + (1 - e^{-K\bar{\tau}}) D_{n-2} \\ &\leq (1 - e^{-K\bar{\tau}} (1 - C)) D_{n-2}, \end{aligned}$$

where the constant C is defined in (2.21). So, setting

$$\tilde{C} = 1 - e^{-K\bar{\tau}} (1 - C),$$

we can conclude that $\tilde{C} \in (0, 1)$ is the constant for which (3.1) holds true. This implies that

$$D_{3n} \leq \tilde{C}^n D_0, \quad \forall n \geq 1. \quad (3.2)$$

Indeed, by induction, if $n = 1$ we know from (3.1) that

$$D_3 \leq \tilde{C} D_0.$$

So, assume that (3.2) holds for $n > 1$ and we prove it for $n + 1$. By using again (3.1) and from the induction hypothesis it comes that

$$D_{3(n+1)} \leq \tilde{C} D_{3n} \leq \tilde{C} \tilde{C}^n D_0 = \tilde{C}^{n+1} D_0,$$

i.e. (3.2) is fulfilled.

Notice that, from (3.2), we have that

$$D_{3n} \leq \left(\sqrt[3]{\tilde{C}} \right)^{3n} D_0 = e^{3n \ln \left(\sqrt[3]{\tilde{C}} \right)} D_0 = e^{3n\bar{\tau} \ln(\tilde{C}) \frac{1}{3\bar{\tau}}} D_0, \quad \forall n \in \mathbb{N}.$$

Therefore, if we set

$$\gamma = \frac{1}{3\bar{\tau}} \ln \left(\frac{1}{\tilde{C}} \right),$$

we can write

$$D_{3n} \leq e^{-3n\gamma\bar{\tau}} D_0, \quad \forall n \in \mathbb{N}_0. \quad (3.3)$$

Now, fix $i, j = 1, \dots, N$ and $t \geq 0$. Then, $t \in [3n\bar{\tau} - \bar{\tau}, 3n\bar{\tau} + 2\bar{\tau}]$, for some $n \in \mathbb{N}_0$. Therefore, by using (2.6) and (3.3), it turns out that

$$|x_i(t) - x_j(t)| \leq D_{3n} \leq e^{-3n\gamma\bar{\tau}} D_0.$$

Thus, being $t \leq 3n\bar{\tau} + 2\bar{\tau}$, we get

$$|x_i(t) - x_j(t)| \leq e^{-\gamma t} e^{2\gamma\bar{\tau}} D_0.$$

Then,

$$d(t) \leq e^{-\gamma(t-2\bar{\tau})} D_0, \quad \forall t \geq 0$$

and (1.5) is proved. \square

Remark 3.1. Let us note that Theorem 1.1 holds true, in particular, for weights a_{ij} of the form

$$a_{ij}(t) := \frac{1}{N-1} \tilde{\psi}(|x_i(t) - x_j(t - \tau(t))|), \quad \forall t > 0, \forall i, j = 1, \dots, N,$$

where $\tilde{\psi} : \mathbb{R} \rightarrow \mathbb{R}$ is a positive, bounded and continuous function. In this case, we can estimate from below $\tilde{\psi}$ in terms of the distance between the initial opinions, namely

$$\tilde{\psi}(|x_i(t) - x_j(t - \tau(t))|) \geq \tilde{\psi}_0 := \min_{s \in [0, D_0]} \tilde{\psi}(s) > 0, \quad \forall t \geq 0,$$

where D_0 is as in Definition 2.1. Then, the proof of Theorem 1.1 follows with the same arguments we have employed in the general case of weights of the type (1.2).

4 The continuum HK-model with pointwise time delay

In this section, we consider the continuum model obtained as mean-field limit of the particle system when $N \rightarrow \infty$. Let $\mathcal{M}(\mathbb{R}^d)$ be the set of probability measures on the space \mathbb{R}^d . Then, the continuum model associated to the particle system (1.1) is given by

$$\begin{aligned} \partial_t \mu_t + \operatorname{div}(F[\mu_{t-\tau(t)}]\mu_t) &= 0, \quad t > 0, \\ \mu_s &= g_s, \quad x \in \mathbb{R}^d, \quad s \in [-\bar{\tau}, 0], \end{aligned} \quad (4.1)$$

where the velocity field F is defined as

$$F[\mu_{t-\tau(t)}](x) = \int_{\mathbb{R}^d} \psi(x, y)(y - x) d\mu_{t-\tau(t)}(y), \quad (4.2)$$

and $g_s \in \mathcal{C}([-\bar{\tau}, 0]; \mathcal{M}(\mathbb{R}^d))$. For the continuum model, as in [11], we assume that the delay function $\tau(\cdot)$ is bounded from below, namely there exists a strictly positive constant $\tau^* > 0$ such that

$$\tau(t) \geq \tau^*, \quad \forall t \geq 0.$$

Moreover, we assume that the potential $\psi(\cdot, \cdot)$ in (4.2) is also Lipschitz continuous, namely for any $(x, y), (x', y') \in \mathbb{R}^{2d}$ there exists $L > 0$ such that

$$|\psi(x, y) - \psi(x', y')| \leq L(|y - y'| + |x - x'|).$$

Definition 4.1. Let $T > 0$. We say that $\mu_t \in \mathcal{C}([0, T]; \mathcal{M}(\mathbb{R}^d))$ is a measure-valued solution to (4.1) on the time interval $[0, T]$ if for all $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d \times [0, T])$ we have:

$$\int_0^T \int_{\mathbb{R}^d} (\partial_t \varphi + F[\mu_{t-\tau(t)}](x) \cdot \nabla_x \varphi) d\mu_t(x) dt + \int_{\mathbb{R}^d} \varphi(x, 0) dg_0(x) = 0. \quad (4.3)$$

Before stating the consensus result for solutions to model (4.1), we first recall some basic tools on probability spaces and measures.

Definition 4.2. Let $\mu, \nu \in \mathcal{M}(\mathbb{R}^d)$ be two probability measures on \mathbb{R}^d . We define the 1-Wasserstein distance between μ and ν as

$$d_1(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y| d\pi(x, y),$$

where $\Pi(\mu, \nu)$ is the space of all couplings for μ and ν , namely all those probability measures on \mathbb{R}^{2d} having as marginals μ and ν :

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \varphi(x) d\pi(x, y) = \int_{\mathbb{R}^d} \varphi(x) d\mu(x), \quad \int_{\mathbb{R}^d \times \mathbb{R}^d} \varphi(y) d\pi(x, y) = \int_{\mathbb{R}^d} \varphi(y) d\nu(y),$$

for all $\varphi \in \mathcal{C}_b(\mathbb{R}^d)$.

Let us introduce the space \mathcal{P}_1 of all probability measures with finite first-order moment. It is well-known that $(\mathcal{P}_1(\mathbb{R}^d), d_1(\cdot, \cdot))$ is a complete metric space.

Now, we define the position diameter for a compactly supported measure $g \in \mathcal{P}_1(\mathbb{R}^d)$ as follows:

$$d_X[g] := \operatorname{diam}(\operatorname{supp} g).$$

Since the consensus result for the particle model (1.1) holds without any upper bounds on the time delay $\tau(\cdot)$, one can improve the consensus theorem for the PDE model (4.1) obtained in [11] removing the smallness assumption on the time delay $\tau(t)$. We omit the proof since, once we have the result for the particle system (1.1), the consensus estimate for the continuum model is obtained with arguments analogous to the ones in [11] (see also [26]).

Theorem 4.1. *Let $\mu_t \in \mathcal{C}([0, T]; \mathcal{P}_1(\mathbb{R}^d))$ be a measure-valued solution to (4.1) with compactly supported initial datum $g_s \in \mathcal{C}([-\bar{\tau}, 0]; \mathcal{P}_1(\mathbb{R}^d))$ and let F as in (4.2). Then, there exists a constant $C > 0$ such that*

$$d_X(\mu_t) \leq \left(\max_{s \in [-\bar{\tau}, 0]} d_X(g_s) \right) e^{-Ct},$$

for all $t \geq 0$.

5 Distributed time delay

Here, we study the asymptotic behavior of solutions to system (1.6). As before, one can prove the following crucial lemma.

Lemma 5.1. *Let $\{x_i\}_{i=1, \dots, N}$ be a solution to system (1.6) with continuous initial conditions. Then, for each $v \in \mathbb{R}^d$ and $T \geq 0$, we have that*

$$\min_{j=1, \dots, N} \min_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle \leq \langle x_i(t), v \rangle \leq \max_{j=1, \dots, N} \max_{s \in [T-\bar{\tau}, T]} \langle x_j(s), v \rangle, \quad (5.1)$$

for all $t \geq T - \bar{\tau}$ and $i = 1, \dots, N$.

Proof. First of all, we note that the inequalities in the statement are satisfied for every $t \in [T - \bar{\tau}, T]$.

Now, fix $T \geq 0$, a vector $v \in \mathbb{R}^d$ and a positive constant ϵ . Define the constant M_T and the set K^ϵ as in the proof of Lemma 2.1. Then, denoted as before $S^\epsilon := \sup K^\epsilon$, it holds that $S^\epsilon > T$. We claim that $S^\epsilon = +\infty$. Indeed, suppose by contradiction that $S^\epsilon < +\infty$. Note that by definition of S^ϵ it turns out that

$$\max_{i=1, \dots, N} \langle x_i(t), v \rangle < M_T + \epsilon, \quad \forall t \in (T, S^\epsilon), \quad (5.2)$$

and

$$\lim_{t \rightarrow S^{\epsilon-}} \max_{i=1, \dots, N} \langle x_i(t), v \rangle = M_T + \epsilon. \quad (5.3)$$

For all $i = 1, \dots, N$ and $t \in (T, S^\epsilon)$, we compute

$$\begin{aligned} \frac{d}{dt} \langle x_i(t), v \rangle &= \frac{1}{h(t)} \sum_{j: j \neq i} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) a_{ij}(t; s) \langle x_j(s) - x_i(t), v \rangle ds \\ &= \frac{1}{N-1} \frac{1}{h(t)} \sum_{j: j \neq i} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) \psi(x_i(t), x_j(s)) (\langle x_j(s), v \rangle - \langle x_i(t), v \rangle) ds. \end{aligned}$$

Notice that, being $t \in (T, S^\epsilon)$, then $t - \tau_2(t), t - \tau_1(t) \in (T - \bar{\tau}, S^\epsilon)$ and

$$\langle x_j(s), v \rangle < M_T + \epsilon, \quad \forall s \in [t - \tau_2(t), t - \tau_1(t)], \forall j = 1, \dots, N. \quad (5.4)$$

Moreover, (5.2) implies that

$$\langle x_i(t), v \rangle < M_T + \epsilon,$$

so that

$$M_T + \epsilon - \langle x_i(t), v \rangle \geq 0.$$

Combining this last fact with (5.4) and by recalling of (1.9), we can write

$$\begin{aligned} \frac{d}{dt} \langle x_i(t), v \rangle &\leq \frac{1}{N-1} \frac{1}{h(t)} \sum_{j:j \neq i} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) \psi(x_i(t), x_j(s)) (M_T + \epsilon - \langle x_i(t), v \rangle) ds \\ &\leq \frac{K}{N-1} \frac{1}{h(t)} (M_T + \epsilon - \langle x_i(t), v \rangle) \sum_{j:j \neq i} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) ds \\ &= K \frac{1}{h(t)} (M_T + \epsilon - \langle x_i(t), v \rangle) \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) ds \\ &= K (M_T + \epsilon - \langle x_i(t), v \rangle), \end{aligned}$$

for all $t \in (T, S^\epsilon)$. Then, the Gronwall's Lemma allows us to conclude the proof of the second inequality arguing analogously to the proof of Lemma 2.1. Also the proof of the first inequality is obtained similarly with respect to the pointwise time delay case. We omit the details. \square

As before, one can define the quantities D_n , $n \in \mathbb{N}_0$, and prove the analogous, for solutions to the model with distributed time delay (1.6), of the lemmas in Section 2. Then, the following exponential convergence to consensus holds.

Theorem 5.2. *Assume that $\psi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a positive, bounded, continuous function. Then, every solution $\{x_i\}_{i=1, \dots, N}$ to (1.6), with continuous initial conditions $x_i^0 : [-\bar{\tau}, 0] \rightarrow \mathbb{R}^d$, satisfies the exponential decay estimate*

$$d(t) \leq \left(\max_{i,j=1, \dots, N} \max_{r,s \in [-\bar{\tau}, 0]} |x_i(r) - x_j(s)| \right) e^{-\gamma(t-2\bar{\tau})}, \quad \forall t \geq 0,$$

for a suitable positive constant γ independent of N .

Remark 5.3. Let us note that Theorem 5.2 holds true, in particular, for weights a_{ij} of the form

$$a_{ij}(t; s) := \frac{1}{N-1} \tilde{\psi}(|x_i(t) - x_j(s)|), \quad \forall t > 0, \forall i, j = 1, \dots, N,$$

where $\tilde{\psi} : \mathbb{R} \rightarrow \mathbb{R}$ is a positive, bounded and continuous function. In this case, one can bound from below the influence function $\tilde{\psi}$ as explained in Remark 3.1.

The related PDE model is now:

$$\begin{aligned} \partial_t \mu_t + \operatorname{div} \left(\frac{1}{h(t)} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) F[\mu_s] ds \mu_t \right) &= 0, \quad t > 0, \\ \mu_s = g_s, \quad x \in \mathbb{R}^d, \quad s \in [-\bar{\tau}, 0], \end{aligned} \tag{5.5}$$

where the velocity field F is given by

$$F[\mu_s](x) = \int_{\mathbb{R}^d} \psi(x, y)(y - x) d\mu_s(y), \quad (5.6)$$

and $g_s \in \mathcal{C}([-\bar{\tau}, 0]; \mathcal{M}(\mathbb{R}^d))$. For the continuum model, as in [25], we assume that the delay functions are bounded from below, namely there exists a strictly positive constant $\tau^* > 0$ such that

$$\tau_2(t) > \tau_1(t) \geq \tau^*, \quad \forall t \geq 0.$$

Moreover, as before, we assume that the potential $\psi(\cdot, \cdot)$ in (5.6) is also Lipschitz continuous with respect to the two arguments.

Definition 5.1. Let $T > 0$. We say that $\mu_t \in \mathcal{C}([0, T]; \mathcal{M}(\mathbb{R}^d))$ is a measure-valued solution to (5.5) on the time interval $[0, T]$ if for all $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d \times [0, T])$ we have:

$$\int_0^T \int_{\mathbb{R}^d} \left(\partial_t \varphi + \frac{1}{h(t)} \int_{t-\tau_2(t)}^{t-\tau_1(t)} \alpha(t-s) F[\mu_s](x) ds \cdot \nabla_x \varphi \right) d\mu_t(x) dt + \int_{\mathbb{R}^d} \varphi(x, 0) dg_0(x) = 0.$$

Since the consensus result for the particle model (1.6) holds without any upper bounds on the time delays $\tau_1(\cdot), \tau_2(\cdot)$, one can improve the consensus theorem for the PDE model (5.5) of [25]. Indeed, in [25], where the author concentrates in the case $\tau_1(t) \equiv 0$, the consensus estimate is obtained under a smallness condition on the time delay. The proof is analogous, then we omit it.

Theorem 5.4. Let $\mu_t \in \mathcal{C}([0, T]; \mathcal{P}_1(\mathbb{R}^d))$ be a measure-valued solution to (5.5) with compactly supported initial datum $g_s \in \mathcal{C}([-\bar{\tau}, 0]; \mathcal{P}_1(\mathbb{R}^d))$ and let F as in (5.6). Then, there exists a constant $C > 0$ such that

$$d_X(\mu_t) \leq \left(\max_{s \in [-\bar{\tau}, 0]} d_X(g_s) \right) e^{-Ct},$$

for all $t \geq 0$.

References

- [1] A. Aydogdu, M. Caponigro, S. McQuade, B. Piccoli, N. Pouradier Duteil, F. Rossi and E. Trélat, *Interaction network, state space and control in social dynamics*. Active particles. Vol. 1. Advances in theory, models, and applications, 99–140. Birkhäuser/Springer, Cham, 2017.
- [2] N. Bellomo, M. A. Herrero and A. Tosin, On the dynamics of social conflict: Looking for the Black Swan, *Kinet. Relat. Models*, 6, (2013), 459–479.
- [3] A. Borzi and S. Wongkaew. Modeling and control through leadership of a refined flocking system. *Math. Models Methods Appl. Sci.*, 25, (2015), 255–282.
- [4] F. Bullo, J. Cortés and S. Martínez, *Distributed control of robotic networks: a mathematical approach to motion coordination algorithms*. Princeton series in applied mathematics. Princeton University Press, Princeton, 2009.
- [5] S. Camazine, J. L. Deneubourg, N.R. Franks, J. Sneyd, G. Theraulaz and E. Bonabeau, *Self-Organization in Biological Systems*, Princeton University Press, Princeton, NJ, 2001.

- [6] C. Canuto, F. Fagnani and P. Tilli, A Eulerian approach to the analysis of rendez-vous algorithms, *IFAC Proceedings Volumes*, 41, (2008), 9039–9044.
- [7] C. Canuto, F. Fagnani and P. Tilli, An Eulerian approach to the analysis of Krause’s consensus models, *SIAM J. Control Optim.*, 50, (2012), 243–265.
- [8] F. Ceragioli and P. Frasca, Continuous and discontinuous opinion dynamics with bounded confidence, *Nonlinear Anal. Real World Appl.*, 13, (2012), 1239–1251.
- [9] Y.-P. Choi and J. Haskovec, Cucker-Smale model with normalized communication weights and time delay, *Kinet. Relat. Models*, 10, (2017), 1011–1033.
- [10] Y.-P. Choi and Z. Li, Emergent behavior of Cucker-Smale flocking particles with heterogeneous time delays, *Appl. Math. Lett.*, 86, (2018), 49–56.
- [11] Y.-P. Choi, A. Paolucci and C. Pignotti, Consensus of the Hegselmann-Krause opinion formation model with time delay, *Math. Methods Appl. Sci.*, 44, (2021), 4560–4579.
- [12] Y.-P. Choi and C. Pignotti, Emergent behavior of Cucker-Smale model with normalized weights and distributed time delays, *Netw. Heterog. Media*, 14, (2019), 789–804.
- [13] F. Cucker and S. Smale, Emergent behaviour in flocks, *IEEE Transactions on Automatic Control*, 52, (2007), 852–862.
- [14] J.-G. Dong, S.-Y. Ha, K. Doheon and K. Jeongho, Time-delay effect on the flocking in an ensemble of thermomechanical Cucker-Smale particles, *J. Differential Equations*, 266, (2019), 2373–2407.
- [15] A. Halanay, *Differential Equations: Stability, Oscillations, Time Lags*, Academic Press, New York and London, 1966.
- [16] J. K. Hale and S.M. V. Lunel, *Introduction to functional differential equations*, Applied Mathematical Sciences, 99, Springer, 1993.
- [17] J. Haskovec, A simple proof of asymptotic consensus in the Hegselmann-Krause and Cucker-Smale models with normalization and delay, *SIAM J. Appl. Dyn. Syst.*, 20, (2021), 130–148.
- [18] J. Haskovec, Direct proof of unconditional asymptotic consensus in the Hegselmann-Krause model with transmission-type delay, *Bull. Lond. Math. Soc.*, 53, (2021), pp. 1312–1323.
- [19] J. Haskovec and I. Markou, Asymptotic flocking in the Cucker-Smale model with reaction-type delays in the non-oscillatory regime, *Kinet. Relat. Models*, 13, (2020), 795–813.
- [20] R. Hegselmann and U. Krause, Opinion dynamics and bounded confidence models, analysis, and simulation, *J. Artif. Soc. Soc. Simul.*, 5, (2002), 1–24.
- [21] A. Jadbabaie, J. Lin and A. S. Morse, Coordination of groups of mobile autonomous agents using nearest neighbor rules, *IEEE Trans. Automat. Control*, 48, (2003), 988–1001.
- [22] Y. Liu and J. Wu, Flocking and asymptotic velocity of the Cucker-Smale model with processing delay, *J. Math. Anal. Appl.*, 415, (2014), 53–61.

- [23] J. Lu, D. W. C. Ho and J. Kurths, Consensus over directed static networks with arbitrary finite communications delays, *Phys. Rev. E*, 80, (2009), 066121, 7 pp.
- [24] G. A. Marsan, N. Bellomo and M. Egidi, Towards a mathematical theory of complex socio-economical systems by functional subsystems representation, *Kinet. Relat. Models*, 1, (2008), 249–278.
- [25] A. Paolucci, Convergence to consensus for a Hegselmann-Krause-type model with distributed time delay, *Minimax Theory Appl.*, 6, (2021), 379–394.
- [26] A. Paolucci and C. Pignotti, A note on the Hegselmann-Krause opinion formation model with time variable time delay, *preprint 2022*.
- [27] B. Piccoli, N. Pouradier Duteil and E. Trélat, Sparse control of Hegselmann-Krause models: Black hole and declustering, *SIAM J. Control Optim.*, 57, (2019), 2628–2659.
- [28] B. Piccoli, F. Rossi and E. Trélat. Control to flocking of the kinetic Cucker-Smale model. *SIAM J. Math. Anal.*, 47, (2015), 4685–4719.
- [29] C. Pignotti and I. Reche Vallejo, Flocking estimates for the Cucker-Smale model with time lag and hierarchical leadership, *J. Math. Anal. Appl.*, 464, (2018), 1313–1332.
- [30] C. Pignotti and E. Trélat, Convergence to consensus of the general finite-dimensional Cucker-Smale model with time-varying delays, *Commun. Math. Sci.*, 16, (2018), 2053–2076.
- [31] S. Y. Pilyugin and M. C. Campi, Opinion formation in voting processes under bounded confidence, *Netw. Heterog. Media*, 14, (2019), 617–632.
- [32] M. Rodriguez Cartabia, Cucker-Smale model with time delay, *Discrete Contin. Dynam. Systems*, 42, (2022), 2409–2432.
- [33] S. Wongkaew, M. Caponigro and A. Borzi, On the control through leadership of the Hegselmann-Krause opinion formation model. *Math. Models Methods Appl. Sci.*, 25, (2015), 565–585.