

CANAL HYPERSURFACES GENERATED BY NON-NULL CURVES IN LORENTZ-MINKOWSKI 4-SPACE

AHMET KAZAN¹, MUSTAFA ALTIN² AND DAE WON YOON³

ABSTRACT. In the present paper, firstly we obtain the general expression of the canal hypersurfaces which are formed as the envelope of a family of pseudo hyperspheres, pseudo hyperbolic hyperspheres and null hypercones in E_1^4 and give their some geometric invariants such as unit normal vector fields, Gaussian curvatures, mean curvatures and principal curvatures. Also we give some results about their flatness and minimality conditions and Weingarten canal hypersurfaces. Also, we obtain these characterizations for tubular hypersurfaces in E_1^4 by taking constant radius function and finally we construct some examples and visualize them with the aid of Mathematica.

1. GENERAL INFORMATION AND BASIC CONCEPTS

Canal surfaces, which are the class of surfaces formed by sweeping a sphere, have been investigated by Monge in 1850. Thus, a canal surface can be seen as the envelope of a moving sphere with varying radius, defined by the trajectory $\beta(s)$ of its centers and a radius function $r(s)$. If the radius function is constant, then the canal surface is called a tubular surface or pipe surface. Canal surfaces (especially tubular surfaces) have been applied to many fields, such as the solid and the surface modeling for CAD/CAM, construction of blending surfaces, shape re-construction and they are useful to represent various objects such as pipe, hose, rope or intestine of a body [16], [27]. In this context, canal and tubular (hyper)surfaces have been studied by many mathematicians in different three, four or higher dimensional spaces (see [1], [5], [7], [8], [9], [11], [12], [13], [14], [15], [18], [19], [21], [24], [25], [26], [27], [29], [30] and etc.).

Since the extrinsic differential geometry of submanifolds in Lorentz-Minkowski 4-space E_1^4 is of special interest in Relativity Theory, lots of studies about curves and (hyper)surfaces have been done in this space and this motivated us to construct the canal hypersurfaces in E_1^4 . Now, let us recall some basic concepts for curves and hypersurfaces in E_1^4 .

If $\vec{x} = (x_1, x_2, x_3, x_4)$, $\vec{y} = (y_1, y_2, y_3, y_4)$ and $\vec{z} = (z_1, z_2, z_3, z_4)$ are three vectors in E_1^4 , then the inner product and vector product are defined by

$$\langle \vec{x}, \vec{y} \rangle = -x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4 \quad (1.1)$$

and

$$\vec{x} \times \vec{y} \times \vec{z} = \det \begin{bmatrix} -e_1 & e_2 & e_3 & e_4 \\ x_1 & x_2 & x_3 & x_4 \\ y_1 & y_2 & y_3 & y_4 \\ z_1 & z_2 & z_3 & z_4 \end{bmatrix}, \quad (1.2)$$

respectively.

A vector $\vec{x} \in E_1^4 - \{0\}$ is called spacelike if $\langle \vec{x}, \vec{x} \rangle > 0$; timelike if $\langle \vec{x}, \vec{x} \rangle < 0$ and lightlike (null) if $\langle \vec{x}, \vec{x} \rangle = 0$. In particular, the vector $\vec{x} = 0$ is spacelike. Also, the norm of the vector \vec{x} is $\|\vec{x}\| = \sqrt{|\langle \vec{x}, \vec{x} \rangle|}$. A curve $\beta(s)$ in E_1^4 is spacelike, timelike or lightlike (null), if all its

2010 *Mathematics Subject Classification.* 14J70, 53A07, 53A10.

Key words and phrases. Canal hypersurface, Tubular hypersurface, Lorentz-Minkowski 4-space, Weingarten Hypersurface.

velocity vectors $\beta'(s)$ are spacelike, timelike or lightlike, respectively and a non-null (i.e. timelike or spacelike) curve has unit speed if $\langle \beta', \beta' \rangle = \mp 1$ [22].

If $\{F_1, F_2, F_3, F_4\}$ is the moving Frenet frame along the timelike or spacelike curve $\beta(s)$ in E_1^4 , where we'll call F_1, F_2, F_3 and F_4 are unit tangent vector field, principal normal vector field, binormal vector field and trinormal vector field, respectively, then the Frenet equations can be given according to the causal characters of non-null Frenet vector fields F_1, F_2, F_3 and F_4 as follows [28]:

If the curve $\beta(s)$ is timelike, i.e. $\langle F_1, F_1 \rangle = -1, \langle F_2, F_2 \rangle = \langle F_3, F_3 \rangle = \langle F_4, F_4 \rangle = 1$, then

$$\begin{bmatrix} F_1' \\ F_2' \\ F_3' \\ F_4' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ k_1 & 0 & k_2 & 0 \\ 0 & -k_2 & 0 & k_3 \\ 0 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \end{bmatrix}; \quad (1.3)$$

if the curve $\beta(s)$ is spacelike with timelike principal normal vector field F_2 , i.e. $\langle F_2, F_2 \rangle = -1, \langle F_1, F_1 \rangle = \langle F_3, F_3 \rangle = \langle F_4, F_4 \rangle = 1$, then

$$\begin{bmatrix} F_1' \\ F_2' \\ F_3' \\ F_4' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ k_1 & 0 & k_2 & 0 \\ 0 & k_2 & 0 & k_3 \\ 0 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \end{bmatrix}; \quad (1.4)$$

if the curve $\beta(s)$ is spacelike with timelike binormal vector field F_3 , i.e. $\langle F_3, F_3 \rangle = -1, \langle F_1, F_1 \rangle = \langle F_2, F_2 \rangle = \langle F_4, F_4 \rangle = 1$, then

$$\begin{bmatrix} F_1' \\ F_2' \\ F_3' \\ F_4' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & k_2 & 0 & k_3 \\ 0 & 0 & k_3 & 0 \end{bmatrix} \begin{bmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \end{bmatrix}; \quad (1.5)$$

if the curve $\beta(s)$ is spacelike with timelike trinormal vector field F_4 , i.e. $\langle F_4, F_4 \rangle = -1, \langle F_1, F_1 \rangle = \langle F_2, F_2 \rangle = \langle F_3, F_3 \rangle = 1$, then

$$\begin{bmatrix} F_1' \\ F_2' \\ F_3' \\ F_4' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & -k_2 & 0 & k_3 \\ 0 & 0 & k_3 & 0 \end{bmatrix} \begin{bmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \end{bmatrix}, \quad (1.6)$$

where k_1, k_2, k_3 are the first, second and third curvatures of the non-null curve $\beta(s)$.

Furthermore, if p is a fixed point in E_1^4 and r is a positive constant, then the pseudo-Riemannian hypersphere is defined by

$$S_1^3(p, r) = \{x \in E_1^4 : \langle x - p, x - p \rangle = r^2\},$$

the pseudo-Riemannian hyperbolic space is defined by

$$H_0^3(p, r) = \{x \in E_1^4 : \langle x - p, x - p \rangle = -r^2\},$$

the pseudo-Riemannian null hypercone is defined by

$$Q_1^3 = \{x \in E_1^4 : \langle x - p, x - p \rangle = 0\}.$$

In the present study, we construct the canal hypersurfaces in E_1^4 as the envelope of a family of pseudo hyperspheres, pseudo hyperbolic hyperspheres or null hypercones whose centers lie on a non-null space curve.

On the other hand, the differential geometry of different types of (hyper)surfaces in 4-dimensional spaces has been a popular topic for geometers, recently ([2], [3], [4], [6], [17], [20] and etc). In this context, let Γ be a hypersurface in E_1^4 given by

$$\begin{aligned} \Gamma : U \subset E^3 &\longrightarrow E_1^4 \\ (u_1, u_2, u_3) &\longrightarrow \Gamma(u_1, u_2, u_3) = (\Gamma_1(u_1, u_2, u_3), \Gamma_2(u_1, u_2, u_3), \Gamma_3(u_1, u_2, u_3), \Gamma_4(u_1, u_2, u_3)). \end{aligned} \quad (1.7)$$

Then the Gauss map (i.e., the unit normal vector field), the matrix forms of the first and second fundamental forms are

$$N_\Gamma = \frac{\Gamma_{u_1} \times \Gamma_{u_2} \times \Gamma_{u_3}}{\|\Gamma_{u_1} \times \Gamma_{u_2} \times \Gamma_{u_3}\|}, \quad (1.8)$$

$$[g_{ij}] = \begin{bmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{bmatrix} \quad (1.9)$$

and

$$[h_{ij}] = \begin{bmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{bmatrix}, \quad (1.10)$$

respectively. Here $g_{ij} = \langle \Gamma_{u_i}, \Gamma_{u_j} \rangle$, $h_{ij} = \langle \Gamma_{u_i u_j}, N_\Gamma \rangle$, $\Gamma_{u_i} = \frac{\partial \Gamma}{\partial u_i}$, $\Gamma_{u_i u_j} = \frac{\partial^2 \Gamma}{\partial u_i \partial u_j}$, $i, j \in \{1, 2, 3\}$.

Also, the matrix of shape operator of the hypersurface (1.7) is

$$S = [a_{ij}] = [g^{ij}] \cdot [h_{ij}], \quad (1.11)$$

where $[g^{ij}]$ is the inverse matrix of $[g_{ij}]$.

With the aid of (1.9)-(1.11), the Gaussian curvature and mean curvature of a hypersurface in E_1^4 are given by

$$K = \varepsilon \frac{\det[h_{ij}]}{\det[g_{ij}]} \quad (1.12)$$

and

$$3\varepsilon H = \text{tr}(S), \quad (1.13)$$

respectively. Here, $\varepsilon = \langle N_\Gamma, N_\Gamma \rangle$. We say that a hypersurface is flat or minimal, if it has zero Gaussian or zero mean curvature, respectively. For more details about hypersurfaces in E_1^4 , we refer to [10], [23] and etc.

In the second section of this paper, we obtain the general expression of the canal hypersurfaces which are formed as the envelope of a family of pseudo hyperspheres, pseudo hyperbolic hyperspheres and null hypercones in E_1^4 and give some characterizations for them. In the third section, we obtain these results for tubular hypersurfaces by taking constant radius function and in the last section, we construct some examples and visualize them with the aid of Mathematica.

2. CANAL HYPERSURFACES GENERATED BY NON-NULL CURVES IN E_1^4

In this section, firstly we construct the canal hypersurfaces which are formed as the envelope of a family of pseudo hyperspheres, pseudo hyperbolic hyperspheres and null hypercones in E_1^4 . After that, we obtain some important geometric invariants such as unit normal vector fields, Gaussian curvatures, mean curvatures and principal curvatures of these canal hypersurfaces in general form. Also, we give some results for flat, minimal and Weingarten canal hypersurfaces in E_1^4 .

2.1. CONSTRUCTION OF CANAL HYPERSURFACES.

Here, we prove a theorem which gives us the general parametric expressions of 11 different types of canal hypersurfaces obtained by pseudo hyperspheres, pseudo hyperbolic hyperspheres and null hypercones in E_1^4 . Also, we write the parametric expressions of these canal hypersurfaces explicitly.

Theorem 1. *The canal hypersurfaces which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by spacelike or timelike center curves with non-null Frenet vector fields can be parametrized by*

$$\mathfrak{C}^{\{j;\lambda\}}(s, t, w) = \beta(s) - \lambda \varepsilon_1 r(s) r'(s) F_1(s) \mp r(s) \sqrt{r'(s)^2 - \lambda \varepsilon_1} \left(\sum_{i=2}^4 \mathfrak{a}_i(s, t, w) F_i(s) \right), \quad (2.1)$$

where

$$i) \ g(F_j, F_j) = -1 = \varepsilon_j \quad \text{and for } i \neq j, \ \varepsilon_i = 1, \ i, j \in \{1, 2, 3, 4\};$$

$$ii) \ \begin{cases} j = 1 \Rightarrow \mathfrak{a}_2(s, t, w) = \cos t \cos w, \ \mathfrak{a}_3(s, t, w) = \sin t \cos w, \ \mathfrak{a}_4(s, t, w) = \sin w, \\ j = 2, 3, 4 \Rightarrow \begin{cases} \mathfrak{a}_j(s, t, w) = \cosh t \cosh w, \ \mathfrak{a}_{j+1}(s, t, w) = \sinh w, \ \mathfrak{a}_{j+2}(s, t, w) = \sinh t \cosh w, \\ \mathfrak{a}_5(s, t, w) = \mathfrak{a}_2(s, t, w), \ \mathfrak{a}_6(s, t, w) = \mathfrak{a}_3(s, t, w); \end{cases} \end{cases}$$

also, we suppose $r'(s)^2 > \lambda \varepsilon_1$ and if the canal hypersurface is foliated by pseudo hyperspheres or pseudo hyperbolic hyperspheres, then $\lambda = 1$ or $\lambda = -1$, respectively.

Furthermore, the canal hypersurfaces which are formed as the envelope of a family of null hypercones can be parametrized by

$$\mathfrak{C}^{\{j;0\}}(s, t, w) = \beta(s) \mp \left(\sum_{i=2}^4 a_i(s, t, w) F_i(s) \right), \quad j = 2, 3, 4 \quad (2.2)$$

where $a_i(s, t, w)$ are arbitrary functions satisfying

$$\sum_{i=2}^4 \varepsilon_i a_i^2(s, t, w) = 0.$$

It is obvious from this expression that, the canal hypersurface $\mathfrak{C}^{\{1;0\}}$ cannot be defined.

Here, the center curve which generates the canal hypersurface is timelike or spacelike if $j = 1$ or $j = 2, 3, 4$, respectively.

Proof. Let $\beta : I \subseteq \mathbb{R} \rightarrow E_1^4$ be a spacelike or timelike center curve parametrized by arc-length with non-zero curvature and $g(F_j, F_j) = \varepsilon_j = -1$, where $F_1(s), F_2(s), F_3(s), F_4(s)$ are unit tangent, principal normal, binormal and trinormal vectors of $\beta(s)$, respectively. Then, the parametrization of the envelope of pseudo hyperspheres ($\lambda = 1$) (resp. pseudo hyperbolic hyperspheres ($\lambda = -1$)) or null hypercones ($\lambda = 0$)) defining the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}$ in E_1^4 can be given by

$$\mathfrak{C}^{\{j;\lambda\}}(s, t, w) - \beta(s) = a_1(s, t, w) F_1(s) + a_2(s, t, w) F_2(s) + a_3(s, t, w) F_3(s) + a_4(s, t, w) F_4(s), \quad (2.3)$$

where $a_i(s, t, w)$ are differentiable functions of s, t, w on the interval I . Furthermore, since $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ lies on the pseudo hyperspheres (resp. pseudo hyperbolic hyperspheres or null hypercones), we have

$$g(\mathfrak{C}^{\{j;\lambda\}}(s, t, w) - \beta(s), \mathfrak{C}^{\{j;\lambda\}}(s, t, w) - \beta(s)) = \lambda r^2(s) \quad (2.4)$$

which leads to from (2.3) that

$$\varepsilon_1 a_1^2 + \varepsilon_2 a_2^2 + \varepsilon_3 a_3^2 + \varepsilon_4 a_4^2 = \lambda r^2 \quad (2.5)$$

and

$$\varepsilon_1 a_1 a_{1_s} + \varepsilon_2 a_2 a_{2_s} + \varepsilon_3 a_3 a_{3_s} + \varepsilon_4 a_4 a_{4_s} = \lambda r r_s, \quad (2.6)$$

where $r(s)$ is the radius function; $r = r(s)$, $r_s = \frac{dr(s)}{ds}$, $a_i = a_i(s, t, w)$, $a_{i_s} = \frac{\partial a_i(s, t, w)}{\partial s}$.

Also from (1.3)-(1.6), the non-null Frenet vectors $F_i(s)$, $i \in \{1, 2, 3, 4\}$, of the spacelike or timelike curve $\beta(s)$ satisfy the relations

$$\left. \begin{aligned} F_1'(s) &= k_1(s)F_2(s), \\ F_2'(s) &= \varepsilon_3\varepsilon_4k_1(s)F_1(s) + k_2(s)F_3(s), \\ F_3'(s) &= \varepsilon_1\varepsilon_4k_2(s)F_2(s) + k_3(s)F_4(s), \\ F_4'(s) &= \varepsilon_1\varepsilon_2k_3(s)F_3(s). \end{aligned} \right\} \quad (2.7)$$

Here, $\beta(s)$ is a timelike curve if $\varepsilon_1 = -1$. Also, $\beta(s)$ is a spacelike curve with timelike principal normal F_2 or timelike binormal F_3 or timelike trinormal F_4 if $\varepsilon_2 = -1$ or $\varepsilon_3 = -1$ or $\varepsilon_4 = -1$, respectively, where $g(F_i, F_i) = \varepsilon_i$, $i \in \{1, 2, 3, 4\}$.

So, differentiating (2.3) with respect to s and using the Frenet formula (2.7), we get

$$\mathfrak{C}_s^{\{j;\lambda\}} = (1 + \varepsilon_3\varepsilon_4a_2k_1 + a_{1_s})F_1 + (a_1k_1 + \varepsilon_1\varepsilon_4a_3k_2 + a_{2_s})F_2 + (a_2k_2 + \varepsilon_1\varepsilon_2a_4k_3 + a_{3_s})F_3 + (a_3k_3 + a_{4_s})F_4, \quad (2.8)$$

where $\mathfrak{C}_s^{\{j;\lambda\}} = \frac{\partial \mathfrak{C}_s^{\{j;\lambda\}}(s, t, w)}{\partial s}$. Furthermore, $\mathfrak{C}^{\{j;\lambda\}}(s, t, w) - \beta(s)$ is a normal vector to the canal hypersurfaces, which implies that

$$g(\mathfrak{C}^{\{j;\lambda\}}(s, t, w) - \beta(s), \mathfrak{C}_s^{\{j;\lambda\}}(s, t, w)) = 0 \quad (2.9)$$

and so, from (2.3), (2.8) and (2.9) we have

$$\begin{aligned} \varepsilon_1a_1(1 + \varepsilon_3\varepsilon_4a_2k_1 + a_{1_s}) + \varepsilon_2a_2(a_1k_1 + \varepsilon_1\varepsilon_4a_3k_2 + a_{2_s}) \\ + \varepsilon_3a_3(a_2k_2 + \varepsilon_1\varepsilon_2a_4k_3 + a_{3_s}) + \varepsilon_4a_4(a_3k_3 + a_{4_s}) = 0. \end{aligned} \quad (2.10)$$

Using (2.6) in (2.10), we get

$$\varepsilon_1a_1 + \varepsilon_1a_1a_{1_s} + \varepsilon_2a_2a_{2_s} + \varepsilon_3a_3a_{3_s} + \varepsilon_4a_4a_{4_s} = 0 \quad (2.11)$$

and thus, from (2.6) and the definition of ε_i , we obtain

$$a_1 = -\varepsilon_1\lambda rr_s. \quad (2.12)$$

Hence, using (2.12) in (2.5), we reach that

$$\varepsilon_1(\varepsilon_2a_2^2 + \varepsilon_3a_3^2 + \varepsilon_4a_4^2) = -r^2(-\varepsilon_1\lambda + \lambda^2r_s^2). \quad (2.13)$$

Here from (2.13);

- i:** if $\lambda = 1$ or $\lambda = -1$, then the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}$ which is formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by spacelike or timelike center curves can be parametrized by (2.1);
- ii:** if $\lambda = 0$, then the canal hypersurfaces $\mathfrak{C}^{\{j;0\}}$ which is formed as the envelope of a family of null hypercones in E_1^4 generated by spacelike or timelike center curves can be parametrized by (2.2)

and this completes the proof. \square

Thus, the explicit parametric expressions of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ and $\mathfrak{C}^{\{j;0\}}(s, t, w)$ from Theorem 1 are as follows:

$$\left. \begin{aligned}
\mathfrak{C}^{\{1;1\}} &= \beta + rr' F_1 \mp r \sqrt{r'^2 + 1} (\cos t \cos w F_2 + \sin t \cos w F_3 + \sin w F_4), \\
\mathfrak{C}^{\{1;-1\}} &= \beta - rr' F_1 \mp r \sqrt{r'^2 - 1} (\cos t \cos w F_2 + \sin t \cos w F_3 + \sin w F_4), \\
\mathfrak{C}^{\{2;1\}} &= \beta - rr' F_1 \mp r \sqrt{r'^2 - 1} (\cosh t \cosh w F_2 + \sinh w F_3 + \sinh t \cosh w F_4), \\
\mathfrak{C}^{\{2;-1\}} &= \beta + rr' F_1 \mp r \sqrt{r'^2 + 1} (\cosh t \cosh w F_2 + \sinh w F_3 + \sinh t \cosh w F_4), \\
\mathfrak{C}^{\{3;1\}} &= \beta - rr' F_1 \mp r \sqrt{r'^2 - 1} (\sinh t \cosh w F_2 + \cosh t \cosh w F_3 + \sinh w F_4), \\
\mathfrak{C}^{\{3;-1\}} &= \beta + rr' F_1 \mp r \sqrt{r'^2 + 1} (\sinh t \cosh w F_2 + \cosh t \cosh w F_3 + \sinh w F_4), \\
\mathfrak{C}^{\{4;1\}} &= \beta - rr' F_1 \mp r \sqrt{r'^2 - 1} (\sinh w F_2 + \sinh t \cosh w F_3 + \cosh t \cosh w F_4), \\
\mathfrak{C}^{\{4;-1\}} &= \beta + rr' F_1 \mp r \sqrt{r'^2 + 1} (\sinh w F_2 + \sinh t \cosh w F_3 + \cosh t \cosh w F_4)
\end{aligned} \right\} \quad (2.14)$$

and

$$\left. \begin{aligned}
\mathfrak{C}^{\{2;0\}} &= \beta \mp \sqrt{a_3^2 + a_4^2} F_2 + a_3 F_3 + a_4 F_4, \\
\mathfrak{C}^{\{3;0\}} &= \beta + a_2 F_2 \mp \sqrt{a_2^2 + a_4^2} F_3 + a_4 F_4, \\
\mathfrak{C}^{\{4;0\}} &= \beta + a_2 F_2 + a_3 F_3 \mp \sqrt{a_2^2 + a_3^2} F_4.
\end{aligned} \right\} \quad (2.15)$$

Remark 1. As one can see in Theorem 1, we deal with the canal hypersurfaces satisfying $r'(s)^2 > \lambda \varepsilon_1$. If we assume that $r'(s)^2 < \lambda \varepsilon_1$, then the canal hypersurfaces $\mathfrak{C}^{\{2;1\}}$, $\mathfrak{C}^{\{3;1\}}$ and $\mathfrak{C}^{\{4;1\}}$ can be rewritten as

$$\left. \begin{aligned}
\mathfrak{C}^{\{2;1\}} &= \beta - rr' F_1 \mp r \sqrt{1 - r'^2} (\cosh t \sinh w F_2 + \cosh w F_3 + \sinh t \sinh w F_4), \\
\mathfrak{C}^{\{3;1\}} &= \beta - rr' F_1 \mp r \sqrt{1 - r'^2} (\sinh t \sinh w F_2 + \cosh t \sinh w F_3 + \cosh w F_4), \\
\mathfrak{C}^{\{4;1\}} &= \beta - rr' F_1 \mp r \sqrt{1 - r'^2} (\cosh w F_2 + \sinh t \sinh w F_3 + \cosh t \sinh w F_4).
\end{aligned} \right\} \quad (2.16)$$

2.2. CURVATURES OF CANAL HYPERSURFACES.

At the beginning of this subsection, we must note that we'll obtain the following results by taking $r'(s)^2 > \lambda \varepsilon_1$ and " \mp " which is in (2.1) as "+". Similar characterizations can be obtained by taking $r'(s)^2 > \lambda \varepsilon_1$ and " \mp " as "-", (or $r'(s)^2 < \lambda \varepsilon_1$ and "+" or "-").

Here, we'll obtain some important geometric invariants of these canal hypersurfaces by proving the following theorem:

Theorem 2. The unit normal vector fields $N^{\{j;\lambda\}}$, Gaussian curvatures $K^{\{j;\lambda\}}$, mean curvatures $H^{\{j;\lambda\}}$ and principle curvatures $\mu_i^{\{j;\lambda\}}$, $i \in \{1, 2, 3\}$, of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given

by (2.1) in E_1^4 , are

$$N^{\{j;\lambda\}} = -\varepsilon_3\varepsilon_4\lambda^j \left(-\lambda\varepsilon_1 r' F_1(s) + \sqrt{r'^2 - \lambda\varepsilon_1} \left(\sum_{i=2}^4 \mathbf{a}_i(s, t, w) F_i(s) \right) \right), \quad (2.17)$$

$$K^{\{j;\lambda\}} = \frac{\varepsilon_3\varepsilon_4\lambda^j \left(r k_1^2 f_j^2 (r'^2 - \lambda\varepsilon_1) + r'' (r'^2 - \lambda\varepsilon_1 + r r'') + \varepsilon_2 \lambda k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} (r'^2 - \lambda\varepsilon_1 + 2r r'') \right)}{r^2 (r'^2 - \lambda\varepsilon_1 + \varepsilon_2 \lambda r k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} + r r'')^2}, \quad (2.18)$$

$$H^{\{j;\lambda\}} = \frac{\varepsilon_3\varepsilon_4\lambda^j}{3} \left(\frac{2}{r} + \frac{(r k_1^2 f_j^2 (r'^2 - \lambda\varepsilon_1) + r'' (r'^2 - \lambda\varepsilon_1 + r r'') + \varepsilon_2 \lambda k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} (r'^2 - \lambda\varepsilon_1 + 2r r''))}{(r'^2 - \lambda\varepsilon_1 + \varepsilon_2 \lambda r k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} + r r'')^2} \right), \quad (2.19)$$

$$\left. \begin{aligned} \mu_1^{\{j;\lambda\}} &= \mu_2^{\{j;\lambda\}} = \frac{\varepsilon_3\varepsilon_4\lambda^j}{r}, \\ \mu_3^{\{j;\lambda\}} &= \frac{\varepsilon_3\varepsilon_4\lambda^j \left(r k_1^2 f_j^2 (r'^2 - \lambda\varepsilon_1) + r'' (r'^2 - \lambda\varepsilon_1 + r r'') + \varepsilon_2 \lambda k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} (r'^2 - \lambda\varepsilon_1 + 2r r'') \right)}{(r'^2 - \lambda\varepsilon_1 + \varepsilon_2 \lambda r k_1 f_j \sqrt{r'^2 - \lambda\varepsilon_1} + r r'')^2}, \end{aligned} \right\} \quad (2.20)$$

where

$$f_j = \begin{cases} \cos t \cos w; & \text{for } j = 1, \\ \cosh t \cosh w; & \text{for } j = 2, \\ \sinh t \cosh w; & \text{for } j = 3, \\ \sinh w; & \text{for } j = 4. \end{cases} \quad (2.21)$$

Proof. Firstly, with the aid of the first derivatives of (2.14) according to s , t and w , we obtain the normals of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ from (1.8) as

$$\left. \begin{aligned} N^{\{1;1\}} &= -r'(s)F_1(s) - \sqrt{r'(s)^2 + 1} (\cos t \cos w F_2 + \sin t \cos w F_3 + \sin w F_4), \\ N^{\{1;-1\}} &= -r'(s)F_1(s) + \sqrt{r'(s)^2 - 1} (\cos t \cos w F_2 + \sin t \cos w F_3 + \sin w F_4), \\ N^{\{2;1\}} &= r'(s)F_1(s) - \sqrt{r'(s)^2 - 1} (\cosh t \cosh w F_2 + \sinh w F_3 + \sinh t \cosh w F_4), \\ N^{\{2;-1\}} &= -r'(s)F_1(s) + \sqrt{r'(s)^2 + 1} (\cosh t \cosh w F_2 + \sinh w F_3 + \sinh t \cosh w F_4), \\ N^{\{3;1\}} &= -r'(s)F_1(s) + \sqrt{r'(s)^2 - 1} (\sinh t \cosh w F_2 + \cosh t \cosh w F_3 + \sinh w F_4), \\ N^{\{3;-1\}} &= -r'(s)F_1(s) - \sqrt{r'(s)^2 + 1} (\sinh t \cosh w F_2 + \cosh t \cosh w F_3 + \sinh w F_4), \\ N^{\{4;1\}} &= -r'(s)F_1(s) + \sqrt{r'(s)^2 - 1} (\sinh w F_2 + \sinh t \cosh w F_3 + \cosh t \cosh w F_4), \\ N^{\{4;-1\}} &= r'(s)F_1(s) + \sqrt{r'(s)^2 + 1} (\sinh w F_2 + \sinh t \cosh w F_3 + \cosh t \cosh w F_4) \end{aligned} \right\} \quad (2.22)$$

and so, we can write (2.17).

Now, the coefficients of the first and second fundamental forms of the canal hypersurfaces $\mathfrak{C}^{\{1;\lambda\}}$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic

hyperspheres in E_1^4 generated by the timelike center curve are

$$\left. \begin{aligned} g_{11}^{\{1;\lambda\}} &= \frac{r'^2 + \lambda}{4} \left(r^2 \left(\begin{aligned} &4k_2^2 \cos^2 w - 4k_2 k_3 \cos t \sin 2w \\ &-(2 \cos 2t \cos^2 w + \cos 2w - 3) k_3^2 \end{aligned} \right) - 4\lambda \right) \\ &\quad - \lambda r^2 k_1^2 (\cos^2 t \cos^2 w + (\lambda \cos^2 t \cos^2 w - \lambda) r'^2) - 2\lambda r r'' - \frac{\lambda r^2 r''^2}{\lambda + r'^2} \\ &\quad - \frac{2\lambda k_1 r \cos w}{\sqrt{\lambda + r'^2}} ((\lambda k_2 r r' \cos t \sin t) (r'^2 + \lambda) + \lambda r r'' \cos t), \\ g_{12}^{\{1;\lambda\}} &= g_{21}^{\{1;\lambda\}} = r^2 (k_2 (r'^2 + \lambda) \cos w - \lambda k_1 r' \sqrt{r'^2 + \lambda} \sin t - k_3 (r'^2 + \lambda) \cos t \sin w) \cos w, \\ g_{13}^{\{1;\lambda\}} &= g_{31}^{\{1;\lambda\}} = r^2 (k_3 (r'^2 + \lambda) \sin t - \lambda k_1 r' \sqrt{r'^2 + \lambda} \cos t \sin w), \\ g_{22}^{\{1;\lambda\}} &= (r'^2 + \lambda) r^2 \cos^2 w, \quad g_{23}^{\{1;\lambda\}} = g_{32}^{\{1;\lambda\}} = 0, \quad g_{33}^{\{1;\lambda\}} = (r'^2 + \lambda) r^2; \end{aligned} \right\} \quad (2.23)$$

$$\left. \begin{aligned} h_{11}^{\{1;\lambda\}} &= \frac{\lambda r (r'^2 + \lambda)}{4} (4k_2^2 \cos^2 w - (\cos(2t) + 2 \cos^2 t \cos(2w) - 3) k_3^2 - 4k_2 k_3 \cos t \sin(2w)) \\ &\quad - \lambda k_1^2 r (\lambda \cos^2 t \cos^2 w + (\cos^2 t \cos^2 w - 1) r'^2) - r'' - \frac{r r''^2}{\lambda + r'^2} \\ &\quad - \frac{\lambda k_1 \cos w}{\sqrt{\lambda + r'^2}} ((\cos t + 2\lambda k_2 r r' \sin t) (r'^2 + \lambda) + 2r r'' \cos t), \\ h_{12}^{\{1;\lambda\}} &= h_{21}^{\{1;\lambda\}} = \lambda r (\cos w k_2 (r'^2 + \lambda) - \lambda k_1 \sin t r' \sqrt{r'^2 + \lambda} - \cos t k_3 \sin w (r'^2 + \lambda)) \cos w, \\ h_{13}^{\{1;\lambda\}} &= h_{31}^{\{1;\lambda\}} = \lambda r (k_3 (r'^2 + \lambda) \sin t - \lambda k_1 r' \sqrt{r'^2 + \lambda} \cos t \sin w), \\ h_{22}^{\{1;\lambda\}} &= \lambda r (r'^2 + \lambda) \cos^2 w, \quad h_{23}^{\{1;\lambda\}} = h_{32}^{\{1;\lambda\}} = 0, \quad h_{33}^{\{1;\lambda\}} = \lambda r (r'^2 + \lambda); \end{aligned} \right\} \quad (2.24)$$

the coefficients of the first and second fundamental forms of the canal hypersurfaces $\mathfrak{C}^{\{2;\lambda\}}$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by the spacelike center curve with the timelike principal normal vector are

$$\left. \begin{aligned} g_{11}^{\{2;\lambda\}} &= \frac{1}{4} \left(4 - 4\lambda r'^2 + r^2 (r'^2 - \lambda) \left(\begin{aligned} &(\cosh 2t + 2 \cosh^2 t \cosh 2w - 3) k_3^2 \\ &+ (\cosh 2t + 2 \cosh 2w \sinh^2 t + 3) k_2^2 \\ &- 4k_2 k_3 \cosh^2 w \sinh 2t \end{aligned} \right) \right) \\ &\quad + r^2 k_1^2 ((\cosh^2 t \cosh^2 w - 1) r'^2 - \lambda \cosh^2 t \cosh^2 w) - 2\lambda r r'' - \frac{\lambda r^2 r''^2}{r'^2 - \lambda} \\ &\quad + \frac{2k_1 r}{\sqrt{r'^2 - \lambda}} ((\cosh t \cosh w + \lambda k_2 r r' \sinh w) (r'^2 - \lambda) + r r'' \cosh t \cosh w), \\ g_{12}^{\{2;\lambda\}} &= g_{21}^{\{2;\lambda\}} = r^2 \left(\begin{aligned} &\lambda (k_1 r' \sqrt{r'^2 - \lambda} - k_2 \lambda (r'^2 - \lambda) \sinh w) \sinh t \\ &+ k_3 (r'^2 - \lambda) \cosh t \sinh w \end{aligned} \right) \cosh w, \\ g_{13}^{\{2;\lambda\}} &= g_{31}^{\{2;\lambda\}} = r^2 (\lambda k_1 r' \sqrt{r'^2 - \lambda} \cosh t \sinh w + k_2 (r'^2 - \lambda) \cosh t + k_3 (r'^2 - \lambda) \sinh t), \\ g_{22}^{\{2;\lambda\}} &= (r'^2 - \lambda) r^2 \cosh^2 w, \quad g_{23}^{\{2;\lambda\}} = g_{32}^{\{2;\lambda\}} = 0, \quad g_{33}^{\{2;\lambda\}} = r^2 (r'^2 - \lambda); \end{aligned} \right\} \quad (2.25)$$

$$\left. \begin{aligned} h_{11}^{\{2;\lambda\}} &= \frac{1}{4} \left(\begin{aligned} &r (r'^2 - \lambda) \left(\begin{aligned} &(\cosh(2t) + 2 \cosh^2 t \cosh(2w) - 3) k_3^2 \\ &- 4k_2 k_3 \cosh^2 w \sinh(2t) \\ &+ k_2^2 (3 + \cosh(2t) + 2 \sinh^2 t \cosh(2w)) \end{aligned} \right) - 4\lambda r'' \\ &+ k_1^2 r (r'^2 (\cosh(2t) + 2 \cosh^2 t \cosh(2w) - 3) - 4\lambda \cosh^2 t \cosh^2 w) \\ &+ \frac{4k_1}{\sqrt{r'^2 - \lambda}} \left(\begin{aligned} &(\cosh t \cosh w + 2\lambda k_2 r r' \sinh w) (r'^2 - \lambda) \\ &+ 2r r'' \cosh t \cosh w \end{aligned} \right) - \frac{4\lambda r r''^2}{r'^2 - \lambda} \end{aligned} \right), \\ h_{12}^{\{2;\lambda\}} &= h_{21}^{\{2;\lambda\}} = r (\lambda k_1 r' \sqrt{r'^2 - \lambda} \sinh t + (r'^2 - \lambda) (k_3 \cosh t - k_2 \sinh t) \sinh w) \cosh w, \\ h_{13}^{\{2;\lambda\}} &= h_{31}^{\{2;\lambda\}} = r (\lambda k_1 r' \sqrt{r'^2 - \lambda} \cosh t \sinh w + (r'^2 - \lambda) (k_2 \cosh t - k_3 \sinh t)), \\ h_{22}^{\{2;\lambda\}} &= r (r'^2 - \lambda) \cosh^2 w, \quad h_{23}^{\{2;\lambda\}} = h_{32}^{\{2;\lambda\}} = 0, \quad h_{33}^{\{2;\lambda\}} = r (r'^2 - \lambda); \end{aligned} \right\} \quad (2.26)$$

the coefficients of the first and second fundamental forms of the canal hypersurfaces $\mathfrak{C}^{\{3;\lambda\}}$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres

in E_1^4 generated by the spacelike center curve with the timelike binormal vector are

$$\left. \begin{aligned} g_{11}^{\{3;\lambda\}} &= \frac{r'^2 - \lambda}{4} \left(r^2 \left(\begin{aligned} &k_3^2 (3 + \cosh 2t + 2 \cosh 2w \sinh^2 t) \\ &+ 4k_2^2 \cosh^2 w - 4k_2 k_3 \sinh t \sinh 2w \end{aligned} \right) - 4\lambda \right) \\ &\quad + r^2 k_1^2 (r'^2 (1 + \sinh^2 t \cosh^2 w) - \lambda \sinh^2 t \cosh^2 w) - 2\lambda r r'' - \frac{\lambda r^2 r'^2}{r'^2 - \lambda} \\ &\quad - \frac{2k_1 r \cosh w}{\sqrt{r'^2 - \lambda}} ((r'^2 - \lambda) (\lambda k_2 r r' \cosh t \sinh t) + r r'' \sinh t), \\ g_{12}^{\{3;\lambda\}} &= g_{21}^{\{3;\lambda\}} = r^2 \left(\begin{aligned} &k_2 (r'^2 - \lambda) \cosh w - \lambda k_1 r' \sqrt{r'^2 - \lambda} \cosh t \\ &- k_3 (r'^2 - \lambda) \sinh t \sinh w \end{aligned} \right) \cosh^2 w, \\ g_{13}^{\{3;\lambda\}} &= g_{31}^{\{3;\lambda\}} = r^2 (k_3 (r'^2 - \lambda) \cosh t - \lambda k_1 r' \sqrt{r'^2 - \lambda} \sinh t \sinh w), \\ g_{22}^{\{3;\lambda\}} &= (r'^2 - \lambda) r^2 \cosh^2 w, \quad g_{23}^{\{3;\lambda\}} = g_{32}^{\{3;\lambda\}} = 0, \quad g_{33}^{\{3;\lambda\}} = (r'^2 - \lambda) r^2; \end{aligned} \right\} \quad (2.27)$$

$$\left. \begin{aligned} h_{11}^{\{3;\lambda\}} &= 2k_1 k_2 r r' \sqrt{r'^2 - \lambda} \cosh t \cosh w - \lambda k_2^2 r (r'^2 - \lambda) \cosh^2 w \\ &\quad - \frac{\lambda k_3^2 r (r'^2 - \lambda)}{4} (3 + \cosh(2t) + 2 \sinh^2 t \cosh(2w)) + \lambda k_2 k_3 r (r'^2 - \lambda) \sinh t \sinh(2w) \\ &\quad + \frac{k_1^2 r}{4} (4 \cosh^2 w \sinh^2 t - \lambda r'^2 (3 + \cosh(2t) + 2 \sinh^2 t \cosh(2w))) + r'' + \frac{r r'^2}{-\lambda + r'^2} \\ &\quad + \frac{\lambda}{\sqrt{r'^2 - \lambda}} (k_1 (r'^2 - \lambda + 2r r'') \cosh w \sinh t), \\ h_{12}^{\{3;\lambda\}} &= h_{21}^{\{3;\lambda\}} = r (k_1 r' \sqrt{r'^2 - \lambda} \cosh t - \lambda (r'^2 - \lambda) (k_2 \cosh w + k_3 \sinh t \sinh w)) \cosh w, \\ h_{13}^{\{3;\lambda\}} &= h_{31}^{\{3;\lambda\}} = r (k_1 r' \sqrt{r'^2 - \lambda} \sinh t \sinh w - \lambda k_3 (r'^2 - \lambda) \cosh t), \\ h_{22}^{\{3;\lambda\}} &= -\lambda r (r'^2 - \lambda) \cosh^2 w, \quad h_{23}^{\{3;\lambda\}} = h_{32}^{\{3;\lambda\}} = 0, \quad h_{33}^{\{3;\lambda\}} = -\lambda r (r'^2 - \lambda) \end{aligned} \right\} \quad (2.28)$$

and the coefficients of the first and second fundamental forms of the canal hypersurfaces $\mathfrak{C}^{\{4;\lambda\}}$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by the spacelike center curve with the timelike trinormal vector are

$$\left. \begin{aligned} g_{11}^{\{4;\lambda\}} &= \frac{-\lambda + r'^2}{4} \left(r^2 \left(\begin{aligned} &k_2^2 (2 \cosh 2t \cosh^2 w + \cosh 2w - 3) \\ &+ 4k_3^2 \cosh^2 w + 4k_2 k_3 \cosh t \sinh 2w \end{aligned} \right) - 4\lambda \right) \\ &\quad + r^2 k_1^2 (r'^2 \cosh^2 w - \lambda \sinh^2 w) - 2\lambda r r'' - \frac{\lambda r^2 r'^2}{-\lambda + r'^2} \\ &\quad + \frac{2\lambda k_1 r}{\sqrt{r'^2 - \lambda}} (k_2 r r' (r'^2 - \lambda) \cosh w \sinh t - \lambda (r'^2 - \lambda + r r'') \sinh w), \\ g_{12}^{\{4;\lambda\}} &= g_{21}^{\{4;\lambda\}} = r^2 (r'^2 - \lambda) (k_2 \cosh t \sinh w + k_3 \cosh w) \cosh w, \\ g_{13}^{\{4;\lambda\}} &= g_{31}^{\{4;\lambda\}} = -r^2 (\lambda k_1 r' \sqrt{r'^2 - \lambda} \cosh w + k_2 (r'^2 - \lambda) \sinh t), \\ g_{22}^{\{4;\lambda\}} &= r^2 (r'^2 - \lambda) \cosh^2 w, \quad g_{23}^{\{4;\lambda\}} = g_{32}^{\{4;\lambda\}} = 0, \quad g_{33}^{\{4;\lambda\}} = r^2 (r'^2 - \lambda) \end{aligned} \right\} \quad (2.29)$$

$$\left. \begin{aligned} h_{11}^{\{4;\lambda\}} &= \frac{-r(r'^2 - \lambda)}{4} (k_2^2 (\cosh(2t) + 2 \cosh^2 t \cosh(2w) - 3) + 4k_3 (k_3 \cosh^2 w + k_2 \cosh t \sinh(2w))) \\ &\quad + \lambda k_1^2 r (\sinh^2 w - \lambda r'^2 \cosh^2 w) + \lambda r'' + \frac{\lambda r r'^2}{r'^2 - \lambda} \\ &\quad + \frac{k_1}{\sqrt{r'^2 - \lambda}} ((\sinh w - 2\lambda k_2 r r' \sinh t \cosh w) (r'^2 - \lambda) + 2r r'' \sinh w), \\ h_{12}^{\{4;\lambda\}} &= h_{21}^{\{4;\lambda\}} = -r (k_3 \cosh w + k_2 \cosh t \sinh w) (r'^2 - \lambda) \cosh w, \\ h_{13}^{\{4;\lambda\}} &= h_{31}^{\{4;\lambda\}} = r (\lambda k_1 r' \sqrt{r'^2 - \lambda} \cosh w + k_2 (r'^2 - \lambda) \sinh t), \\ h_{22}^{\{4;\lambda\}} &= -r (r'^2 - \lambda) \cosh^2 w, \quad h_{23}^{\{4;\lambda\}} = h_{32}^{\{4;\lambda\}} = 0, \quad h_{33}^{\{4;\lambda\}} = -r (r'^2 - \lambda) \end{aligned} \right\} \quad (2.30)$$

and these imply that

$$\det g_{ij}^{\{j;\lambda\}} = -\lambda A^2 r^4 (r'^2 - \lambda \varepsilon_1) \left(r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda k_1 f_j r \sqrt{r'^2 - \lambda \varepsilon_1 + r r''} \right)^2, \quad (2.31)$$

$$\det h_{ij}^{\{j;\lambda\}} = -\lambda A^2 r^2 (r'^2 - \lambda \varepsilon_1) \left(\begin{aligned} &\varepsilon_3 \varepsilon_4 \lambda^j f_j^2 k_1^2 r (r'^2 - \lambda \varepsilon_1) + \varepsilon_3 \varepsilon_4 \lambda^j r'' (r'^2 - \lambda \varepsilon_1 + r r'') \\ &+ \varepsilon_2 \varepsilon_3 \varepsilon_4 \lambda^{j+1} f_j k_1 \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + 2r r'') \end{aligned} \right), \quad (2.32)$$

where $A = \cos w$, for $j = 1$ and $A = \cosh w$, for $j = 2, 3, 4$. Thus, from (1.12), (2.31) and (2.32), the Gaussian curvatures are obtained as (2.18).

Here, obtaining the inverse of the first fundamental forms and using these and second fundamental forms in (1.11), we obtain the components of the shape operators of canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}$ as

$$\left. \begin{aligned} S_{11}^{\{1;\lambda\}} &= \frac{\lambda k_1^2 r (r'^2 + \lambda) \cos^2 t \cos^2 w + \lambda r'' (r'^2 + \lambda + r r'') k_1 \sqrt{r'^2 + \lambda} (r'^2 + \lambda + 2r r'') \cos t \cos w}{(r'^2 + \lambda + r (\lambda k_1 \sqrt{r'^2 + \lambda} \cos t \cos w + r''))^2}, \\ S_{21}^{\{1;\lambda\}} &= \frac{-\lambda \sqrt{r'^2 + \lambda} (k_1 r' \sin t \sec w - \lambda k_2 \sqrt{r'^2 + \lambda} + \lambda k_3 \sqrt{r'^2 + \lambda} \cos t \tan w) \sec w}{r (k_1 r \sqrt{r'^2 + \lambda} \cos t + \lambda (r'^2 + \lambda + r r'') \sec w)}, \\ S_{31}^{\{1;\lambda\}} &= \frac{\lambda (\lambda k_3 (r'^2 + \lambda) \sin t - k_1 r' \sqrt{r'^2 + \lambda} \cos t \sin w)}{r (1 + \lambda r'^2 + r (k_1 \sqrt{\lambda + r'^2} \cos t \cos w + \lambda r''))}, \\ S_{22}^{\{1;\lambda\}} &= S_{33}^{\{1;\lambda\}} = \frac{\lambda}{r}, \\ S_{12}^{\{1;\lambda\}} &= S_{13}^{\{1;\lambda\}} = S_{23}^{\{1;\lambda\}} = S_{32}^{\{1;\lambda\}} = 0; \end{aligned} \right\} \quad (2.33)$$

$$\left. \begin{aligned} S_{11}^{\{2;\lambda\}} &= \frac{k_1^2 r (r'^2 - \lambda) \cosh^2 t \cosh^2 w + r'' (r'^2 - \lambda + r r'') - \lambda k_1 \sqrt{r'^2 - \lambda} (r'^2 - \lambda + 2r r'') \cosh t \cosh w}{(r'^2 - \lambda + r (-\lambda k_1 \sqrt{r'^2 - \lambda} \cosh t \cosh w + r''))^2}, \\ S_{21}^{\{2;\lambda\}} &= \frac{(\lambda k_1 r' \sqrt{r'^2 - \lambda} \sinh t \sec h w + (k_3 \cosh t - k_2 \sinh t) \sqrt{r'^2 - \lambda} \tan w) \sec h w}{r ((r'^2 - \lambda + r r'') \sec h w - \lambda k_1 r \sqrt{r'^2 - \lambda} \cosh t)}, \\ S_{31}^{\{2;\lambda\}} &= \frac{-k_1 r' \sqrt{r'^2 - \lambda} \cosh t \sinh w - \lambda k_2 (r'^2 - \lambda) \cosh t + \lambda k_3 (r'^2 - \lambda) \sinh t}{r (1 - \lambda r'^2 + r (k_1 \sqrt{r'^2 - \lambda} \cosh t \cosh w - \lambda r''))}, \\ S_{22}^{\{2;\lambda\}} &= S_{33}^{\{2;\lambda\}} = \frac{1}{r}, \\ S_{12}^{\{2;\lambda\}} &= S_{13}^{\{2;\lambda\}} = S_{23}^{\{2;\lambda\}} = S_{32}^{\{2;\lambda\}} = 0; \end{aligned} \right\} \quad (2.34)$$

$$\left. \begin{aligned} S_{11}^{\{3;\lambda\}} &= -\lambda \frac{k_1^2 r (r'^2 - \lambda) \sinh^2 t \cosh^2 w + r'' (r'^2 - \lambda + r r'') + \lambda k_1 \sqrt{r'^2 - \lambda} (r'^2 - \lambda + 2r r'') \sinh t \cosh w}{(r'^2 - \lambda + r (\lambda k_1 \sqrt{r'^2 - \lambda} \sinh t \cosh w + r''))^2}, \\ S_{21}^{\{3;\lambda\}} &= \frac{\sec h w (k_1 r' \sqrt{r'^2 - \lambda} \cosh t \sec h w - \lambda (k_2 - k_3 \sinh t \tanh w) (r'^2 - \lambda))}{r (\lambda k_1 r \sqrt{r'^2 - \lambda} \sinh t + (-\lambda + r'^2 + r r'') \sec h w)}, \\ S_{31}^{\{3;\lambda\}} &= \frac{-k_1 r' \sqrt{r'^2 - \lambda} \sinh t \sinh w - \lambda k_3 (r'^2 - \lambda) \cosh t}{r (r'^2 - \lambda + r (\lambda k_1 \sqrt{-\lambda + r'^2} \sinh t \cosh w + r''))}, \\ S_{22}^{\{3;\lambda\}} &= S_{33}^{\{3;\lambda\}} = -\frac{\lambda}{r}, \\ S_{12}^{\{3;\lambda\}} &= S_{13}^{\{3;\lambda\}} = S_{23}^{\{3;\lambda\}} = S_{32}^{\{3;\lambda\}} = 0; \end{aligned} \right\} \quad (2.35)$$

$$\left. \begin{aligned} S_{11}^{\{4;\lambda\}} &= \frac{-k_1^2 r (r'^2 - \lambda) \sinh^2 w - r'' (r'^2 - \lambda + r r'') - \lambda k_1 \sqrt{r'^2 - \lambda} (r'^2 - \lambda + 2r r'') \sinh w}{(r'^2 - \lambda + r (\lambda k_1 \sqrt{-\lambda + r'^2} \sinh w + r''))^2}, \\ S_{21}^{\{4;\lambda\}} &= -\lambda \frac{(k_3 + k_2 \tanh w \cosh t) (r'^2 - \lambda)}{r (\lambda r'^2 - 1 + r (k_1 \sqrt{r'^2 - \lambda} \sinh w + \lambda r''))}, \\ S_{31}^{\{4;\lambda\}} &= \frac{-k_1 r' \sqrt{r'^2 - \lambda} \cosh w + \lambda k_2 (r'^2 - \lambda) \sinh t}{r (\lambda r'^2 - 1 + r (k_1 \sqrt{r'^2 - \lambda} \sinh w + \lambda r''))}, \\ S_{22}^{\{4;\lambda\}} &= S_{33}^{\{4;\lambda\}} = -\frac{1}{r}, \\ S_{12}^{\{4;\lambda\}} &= S_{13}^{\{4;\lambda\}} = S_{23}^{\{4;\lambda\}} = S_{32}^{\{4;\lambda\}} = 0. \end{aligned} \right\} \quad (2.36)$$

From (1.13) and (2.33)-(2.36), we obtain the mean curvatures of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}$ as (2.19).

Finally, from (2.33)-(2.36), we get

$$\det(S - \mu I) = \left(\mu - \frac{\varepsilon_3 \varepsilon_4 \lambda^j}{r} \right)^2 \left(\frac{\left(\begin{aligned} &\varepsilon_3 \varepsilon_4 \lambda^j (r k_1^2 f_j^2 (r'^2 - \lambda \varepsilon_1) + r'' (r'^2 - \lambda \varepsilon_1 + r r'')) \\ &+ \varepsilon_2 \varepsilon_3 \varepsilon_4 \lambda^{j+1} k_1 f_j \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + 2r r'') \end{aligned} \right)}{(r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda r k_1 f_j \sqrt{r'^2 - \lambda \varepsilon_1} + r r'')^2} - \mu \right) \quad (2.37)$$

and solving the equation of $\det(S - \mu I) = 0$ we have (2.20). So, the proof is completed. \square

Here, from (2.22) or (2.31), we can state the following proposition:

Proposition 1. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ in E_1^4 are spacelike or timelike if $\lambda = -1$ or $\lambda = 1$, respectively.*

2.3. SOME GEOMETRIC CHARACTERIZATIONS FOR CANAL HYPERSURFACES.

From (2.18) and (2.19), we can obtain the following important relation between the Gaussian and mean curvatures of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}$:

Theorem 3. *The Gaussian and mean curvatures of the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , satisfy*

$$3H^{\{j;\lambda\}}r - K^{\{j;\lambda\}}r^3 - 2\varepsilon_3\varepsilon_4\lambda^j = 0. \quad (2.38)$$

Theorem 4. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , are flat if and only if $k_1 = 0$ and $r(s) = as + b$, $a, b \in \mathbb{R}$, $a \neq \mp 1$.*

Proof. Firstly, let us suppose that the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ in E_1^4 are flat. Then from (2.18) we get

$$rk_1^2(r'^2 - \lambda\varepsilon_1) f_j^2 + \varepsilon_2\lambda^j k_1 \sqrt{r'^2 - \lambda\varepsilon_1} (r'^2 - \lambda\varepsilon_1 + 2rr'') f_j + r''(r'^2 - \lambda\varepsilon_1 + rr'') = 0. \quad (2.39)$$

Since $\{f_j, f_j^2, 1\}$ are linearly independent, we have

$$rk_1^2(r'^2 - \lambda\varepsilon_1) = \varepsilon_2\lambda^j k_1 \sqrt{r'^2 - \lambda\varepsilon_1} (r'^2 - \lambda\varepsilon_1 + 2rr'') = r''(r'^2 - \lambda\varepsilon_1 + rr'') = 0. \quad (2.40)$$

From the (2.40), we get $k_1 = 0$ and $r''(r'^2 - \lambda\varepsilon_1 + rr'') = 0$. When $k_1 = 0$, from (2.18) we have $r'^2 - \lambda\varepsilon_1 + rr'' \neq 0$ and so it must be $r'' = 0$.

Conversely, if $k_1 = 0$ and $r(s) = as + b$, $a, b \in \mathbb{R}$, $a \neq \mp 1$, from (2.18) we get $K = 0$ and this completes the proof. \square

Theorem 5. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , are minimal if and only if $k_1 = 0$ and the radius $r(s)$ satisfies $\int \frac{dr}{\sqrt{\varepsilon_1\lambda + (\frac{c_1}{r})^{\frac{4}{3}}}} = \pm s + c_2$, $c_1, c_2 \in \mathbb{R}$.*

Proof. Firstly, let us suppose that the canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$ in E_1^4 are minimal. Then from (2.19) we get

$$\begin{aligned} & -3r^2k_1^2(\varepsilon_1\lambda - r'^2) f_j^2 - \varepsilon_2\lambda k_1 r \sqrt{r'^2 - \lambda\varepsilon_1} (5\varepsilon_1\lambda - 5r'^2 - 6rr'') f_j \\ & + 2 + 2r'^4 - 5\varepsilon_1\lambda rr'' + 3r^2r''^2 + r'^2(-4\varepsilon_1\lambda + 5rr'') = 0. \end{aligned} \quad (2.41)$$

Since $\{f_j, f_j^2, 1\}$ are linearly independent, we have

$$\left. \begin{aligned} & -3r^2k_1^2(\varepsilon_1\lambda - r'^2) = 0, \\ & \varepsilon_2\lambda k_1 r \sqrt{r'^2 - \lambda\varepsilon_1} (5\varepsilon_1\lambda - 5r'^2 - 6rr'') = 0, \\ & 2 + 2r'^4 - 5\varepsilon_1\lambda rr'' + 3r^2r''^2 + r'^2(-4\varepsilon_1\lambda + 5rr'') = 0. \end{aligned} \right\} \quad (2.42)$$

From (2.42), we get $k_1 = 0$ and $2 + 2r'^4 - 5\varepsilon_1\lambda rr'' + 3r^2r''^2 + r'^2(-4\varepsilon_1\lambda + 5rr'') = 0$; i.e.

$$(2\varepsilon_1\lambda - 2r'^2 - 3rr'')(\varepsilon_1\lambda - r'^2 - rr'') = 0. \quad (2.43)$$

When $k_1 = 0$, from (2.19) we have $r'^2 - \lambda\varepsilon_1 + rr'' \neq 0$ and so it must be

$$-2(r'^2 - \varepsilon_1\lambda) - 3rr'' = 0. \quad (2.44)$$

Now, let us solve the equation (2.44).

If we take $r'(s) = h(s)$, we get

$$r'' = h' = \frac{dh}{dr} \frac{dr}{ds} = \frac{dh}{dr} h. \quad (2.45)$$

Using (2.45) in (2.44), we have

$$3r \frac{dh}{dr} h + 2h^2 - 2\varepsilon_1 \lambda = 0. \quad (2.46)$$

From (2.44), $r'(s) = h(s) \neq 0$ and so we reach that

$$\frac{3h}{2(\varepsilon_1 \lambda - h^2)} dh = \frac{dr}{r}. \quad (2.47)$$

By integrating (2.47), we have

$$h = \pm \sqrt{\varepsilon_1 \lambda + \left(\frac{c_1}{r}\right)^{\frac{4}{3}}}, \quad (2.48)$$

where c_1 is constant. Since $r' = \frac{dr}{ds} = h$, from (2.48) we get

$$\int \frac{dr}{\sqrt{\varepsilon_1 \lambda + \left(\frac{c_1}{r}\right)^{\frac{4}{3}}}} = \pm \int ds. \quad (2.49)$$

Conversely, if $k_1 = 0$ and $r(s)$ satisfies $\int \frac{dr}{\sqrt{\varepsilon_1 \lambda + \left(\frac{c_1}{r}\right)^{\frac{4}{3}}}} = \pm s + c_2$, $c_1, c_2 \in \mathbb{R}$, then we have $H = 0$ and this completes the proof. \square

Now, if

$$H_s K_t - H_t K_s = 0, \quad H_s K_w - H_w K_s = 0, \quad H_t K_w - H_w K_t = 0, \quad (2.50)$$

hold on a hypersurface, then we call the hypersurface as $(H, K)_{st}$ -Weingarten, $(H, K)_{sw}$ -Weingarten, $(H, K)_{tw}$ -Weingarten hypersurface, respectively, where $H_s = \frac{\partial H}{\partial s}$ and so on. So, from (2.18) and (2.19) we have

Theorem 6. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , are $(H, K)_{tw}$ -Weingarten.*

Proof. From (2.18) and (2.19), we get $H_t^{\{j;\lambda\}} K_w^{\{j;\lambda\}} - H_w^{\{j;\lambda\}} K_t^{\{j;\lambda\}} = 0$ and this completes the proof. \square

Theorem 7. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , are $(H, K)_{sw}$ -Weingarten if and only if $k_1 = 0$ or $r(s) = \text{constant}$.*

Proof. From (2.18) and (2.19), we have

$$\begin{aligned} & H_s^{\{j;\lambda\}} K_w^{\{j;\lambda\}} - H_w^{\{j;\lambda\}} K_s^{\{j;\lambda\}} \\ &= - \frac{\left(\begin{array}{l} 2\varepsilon_3^2 \varepsilon_4^2 \lambda^{2j} k_1 r' (r'^2 - \lambda \varepsilon_1)^2 \\ \left((\varepsilon_2^2 \lambda^2 - 1) f_j^2 k_1^2 r^2 + r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j + r r'' \right) \\ \left(\varepsilon_2 \lambda \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + r r'') \right. \\ \left. + k_1 r ((2 - \varepsilon_2^2 \lambda^2) (r'^2 - \lambda \varepsilon_1) - 2 (\varepsilon_2^2 \lambda^2 - 1) r r'') f_j \right) f_{jw} \end{array} \right)}{3r^4 (r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j + r r'')^5}. \end{aligned} \quad (2.51)$$

If $H_s^{\{j;\lambda\}} K_w^{\{j;\lambda\}} - H_w^{\{j;\lambda\}} K_s^{\{j;\lambda\}} = 0$, then from (2.51) we have

$$2k_1 r' (r'^2 - \lambda \varepsilon_1)^2 \left(\begin{array}{l} r'^2 - \lambda \varepsilon_1 + r r'' \\ + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j \end{array} \right) \left(\begin{array}{l} \varepsilon_2 \lambda \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + r r'') \\ + k_1 r (r'^2 - \lambda \varepsilon_1) f_j \end{array} \right) f_{jw} = 0 \quad (2.52)$$

and so,

$$k_1 r' \begin{pmatrix} r'^2 - \lambda \varepsilon_1 + r r'' \\ + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j \end{pmatrix} \begin{pmatrix} \varepsilon_2 \lambda \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + r r'') \\ + k_1 r (r'^2 - \lambda \varepsilon_1) f_j \end{pmatrix} = 0. \quad (2.53)$$

From (2.51), the second and third component of (2.53) cannot be zero and so it must be

$$k_1 r' = 0. \quad (2.54)$$

This completes the proof. \square

Theorem 8. *The canal hypersurfaces $\mathfrak{C}^{\{j;\lambda\}}(s, t, w)$, given by (2.1) in E_1^4 , are $(H, K)_{st}$ -Weingarten and the canal hypersurfaces $\mathfrak{C}^{\{1;\lambda\}}(s, t, w)$, $\mathfrak{C}^{\{2;\lambda\}}(s, t, w)$, $\mathfrak{C}^{\{3;\lambda\}}(s, t, w)$ are $(H, K)_{st}$ -Weingarten if and only if $k_1 = 0$ or $r(s) = \text{constant}$.*

Proof. From (2.18) and (2.19), we have

$$\begin{aligned} & H_s^{\{j;\lambda\}} K_t^{\{j;\lambda\}} - H_t^{\{j;\lambda\}} K_s^{\{j;\lambda\}} \\ &= - \frac{\begin{pmatrix} 2\varepsilon_3^2 \varepsilon_4^2 \lambda^{2j} k_1 r' (r'^2 - \lambda \varepsilon_1)^2 \\ \left((\varepsilon_2^2 \lambda^2 - 1) k_1^2 r^2 f_j^2 + r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j + r r'' \right) \\ \left(\varepsilon_2 \lambda \sqrt{r'^2 - \lambda \varepsilon_1} (r'^2 - \lambda \varepsilon_1 + r r'') \right. \\ \left. + k_1 r ((2 - \varepsilon_2^2 \lambda^2) (r'^2 - \lambda \varepsilon_1) - 2 (\varepsilon_2^2 \lambda^2 - 1) r r'') f_j \right) f_{jt} \end{pmatrix}}{3r^4 (r'^2 - \lambda \varepsilon_1 + \varepsilon_2 \lambda k_1 r \sqrt{r'^2 - \lambda \varepsilon_1} f_j + r r'')^5}. \end{aligned} \quad (2.55)$$

Using (2.21), for $j = 1, 2, 3$ the proof can be seen with similar method in Theorem 7. Also, for $j = 4$ we have $f_{4t} = 0$ and so $H_s^{\{4;\lambda\}} K_t^{\{4;\lambda\}} - H_t^{\{4;\lambda\}} K_s^{\{4;\lambda\}} = 0$ and this completes the proof. \square

3. TUBULAR HYPERSURFACES GENERATED BY NON-NULL CURVES IN E_1^4

In this section, we give the above results, which are given for the canal hypersurfaces in E_1^4 , for tubular hypersurfaces.

Theorem 9. *The tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by spacelike or timelike center curves can be parametrized by*

$$\left. \begin{aligned} \mathfrak{T}^{\{1;1\}}(s, t, w) &= \beta(s) \mp r(s) (\cos t \cos w F_2(s) + \sin t \cos w F_3 + \sin w F_4), \\ \mathfrak{T}^{\{2;1\}}(s, t, w) &= \beta(s) \mp r(s) (\cosh t \sinh w F_2(s) + \cosh w F_3 + \sinh t \sinh w F_4), \\ \mathfrak{T}^{\{2;-1\}}(s, t, w) &= \beta(s) \mp r(s) (\cosh t \cosh w F_2(s) + \sinh w F_3 + \sinh t \cosh w F_4), \\ \mathfrak{T}^{\{3;1\}}(s, t, w) &= \beta(s) \mp r(s) (\sinh t \sinh w F_2(s) + \cosh t \sinh w F_3 + \cosh w F_4), \\ \mathfrak{T}^{\{3;-1\}}(s, t, w) &= \beta(s) \mp r(s) (\sinh t \cosh w F_2(s) + \cosh t \cosh w F_3 + \sinh w F_4), \\ \mathfrak{T}^{\{4;1\}}(s, t, w) &= \beta(s) \mp r(s) (\cosh w F_2(s) + \sinh t \sinh w F_3 + \cosh t \sinh w F_4), \\ \mathfrak{T}^{\{4;-1\}}(s, t, w) &= \beta(s) \mp r(s) (\sinh w F_2(s) + \sinh t \cosh w F_3 + \cosh t \cosh w F_4). \end{aligned} \right\} \quad (3.1)$$

Also, there is no tubular hypersurface $\mathfrak{T}^{\{1;-1\}}(s, t, w)$ which is formed as the envelope of a family of pseudo hyperbolic hyperspheres in E_1^4 generated by timelike center curves.

Furthermore the tubular hypersurfaces $\mathfrak{T}^{\{j;0\}}(s, t, w)$ which are formed as the envelope of a family of null hypercones in E_1^4 generated by spacelike or timelike center curves can be parametrized by (2.15).

Proof. Let the center curve $\beta : I \subseteq \mathbb{R} \rightarrow E_1^4$ be a arc-length parametrized timelike or spacelike curve with non-zero curvature. Then, the parametrization of the envelope of pseudo hyperspheres

(resp. pseudo hyperbolic hyperspheres or null hypercones) defining the tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}$ in E_1^4 can be given by

$$\mathfrak{T}^{\{j;\lambda\}}(s, t, w) - \beta(s) = a_1(s, t, w)F_1(s) + a_2(s, t, w)F_2(s) + a_3(s, t, w)F_3(s) + a_4(s, t, w)F_4(s). \quad (3.2)$$

Furthermore, since $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$ lies on the pseudo hyperspheres (resp. pseudo hyperbolic hyperspheres or null hypercones), we have

$$g(\mathfrak{T}^{\{j;\lambda\}}(s, t, w) - \beta(s), \mathfrak{T}^{\{j;\lambda\}}(s, t, w) - \beta(s)) = \lambda r^2 \quad (3.3)$$

which leads to from (3.2) that

$$\varepsilon_1 a_1^2 + \varepsilon_2 a_2^2 + \varepsilon_3 a_3^2 + \varepsilon_4 a_4^2 = \lambda r^2 \quad (3.4)$$

and

$$\varepsilon_1 a_1 a_{1_s} + \varepsilon_2 a_2 a_{2_s} + \varepsilon_3 a_3 a_{3_s} + \varepsilon_4 a_4 a_{4_s} = 0, \quad (3.5)$$

where $\lambda = 1$, $\lambda = -1$ and $\lambda = 0$ if the tubular hypersurfaces are obtained by pseudo hyperspheres, pseudo hyperbolic hyperspheres and null hypercones, respectively; r is constant radius $a_i = a_i(s, t, w)$, $a_{i_s} = \frac{\partial a_i(s, t, w)}{\partial s}$, and so on.

With similar procedure in the proof of the Theorem 1, we reach that

$$a = 0. \quad (3.6)$$

Hence, using (3.6) in (3.4), we have

$$\varepsilon_2 a_2^2 + \varepsilon_3 a_3^2 + \varepsilon_4 a_4^2 = \lambda r^2. \quad (3.7)$$

From (3.2) and (3.7), we obtain the parametric expressions of the tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$ as (3.1) and (2.15). \square

Now, let us give the Gaussian and mean curvatures of tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$ given by (3.1) with similar method used for canal hypersurfaces.

Theorem 10. *The Gaussian and mean curvatures of tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$, given by (3.1) in E_1^4 , are*

$$\left. \begin{aligned} K^{\{1;1\}} &= \frac{k_1 \cos t \cos w}{r^2(1+rk_1 \cos t \cos w)}, & H^{\{1;1\}} &= \frac{2+3rk_1 \cos t \cos w}{3r(1+rk_1 \cos t \cos w)}; \\ K^{\{2;1\}} &= \frac{k_1 \cosh t \sinh w}{r^2(1+rk_1 \cosh t \sinh w)}, & H^{\{2;1\}} &= \frac{2+3rk_1 \cosh t \sinh w}{3r(1+rk_1 \cosh t \sinh w)}; \\ K^{\{2;-1\}} &= \frac{k_1 \cosh t \cosh w}{r^2(1+rk_1 \cosh t \cosh w)}, & H^{\{2;-1\}} &= \frac{2+3rk_1 \cosh t \cosh w}{3r(1+rk_1 \cosh t \cosh w)}; \\ K^{\{3;1\}} &= \frac{k_1 \sinh t \sinh w}{r^2(1-rk_1 \sinh t \sinh w)}, & H^{\{3;1\}} &= \frac{2-3rk_1 \sinh t \sinh w}{3r(-1+rk_1 \sinh t \sinh w)}; \\ K^{\{3;-1\}} &= \frac{k_1 \sinh t \cosh w}{r^2(-1+rk_1 \sinh t \cosh w)}, & H^{\{3;-1\}} &= \frac{2-3rk_1 \sinh t \cosh w}{3r(1-rk_1 \sinh t \cosh w)}; \\ K^{\{4;1\}} &= \frac{k_1 \cosh w}{r^2(1-rk_1 \cosh w)}, & H^{\{4;1\}} &= \frac{2-3rk_1 \cosh w}{3r(-1+rk_1 \cosh w)}; \\ K^{\{4;-1\}} &= \frac{k_1 \sinh w}{r^2(1-rk_1 \sinh w)}, & H^{\{4;-1\}} &= \frac{2-3rk_1 \sinh w}{3r(-1+rk_1 \sinh w)}. \end{aligned} \right\} \quad (3.8)$$

From (3.8), we can state the following theorem:

Theorem 11. *The tubular hypersurfaces $\mathfrak{T}^{\{j;\lambda\}}(s, t, w)$, given by (3.1) in E_1^4 , are $(H, K)_{tw}$ -Weingarten, $(H, K)_{ts}$ -Weingarten and $(H, K)_{sw}$ -Weingarten hypersurfaces.*

4. VISUALIZATION

In this section, we construct examples for canal hypersurfaces $\mathfrak{C}^{\{1;\lambda\}}(s, t, w)$ and $\mathfrak{C}^{\{3;\lambda\}}(s, t, w)$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 with the aid of a timelike curve and a spacelike curve with timelike binormal vector, seperately.

Firstly, let we take the unit speed timelike curve

$$\beta_1(s) = \left(2 \sinh s, 2 \cosh s, \sqrt{3} \cos s, \sqrt{3} \sin s \right) \quad (4.1)$$

in E_1^4 . The Frenet vectors and curvatures of the curve (4.1) are

$$\left. \begin{aligned} F_1^{\beta_1} &= \left(2 \cosh s, 2 \sinh s, -\sqrt{3} \sin s, \sqrt{3} \cos s \right), \\ F_2^{\beta_1} &= \left(\frac{2}{\sqrt{7}} \sinh s, \frac{2}{\sqrt{7}} \cosh s, -\sqrt{\frac{3}{7}} \cos s, -\sqrt{\frac{3}{7}} \sin s \right), \\ F_3^{\beta_1} &= \left(-\sqrt{3} \cosh s, -\sqrt{3} \sinh s, 2 \sin s, -2 \cos s \right), \\ F_4^{\beta_1} &= \left(\sqrt{\frac{3}{7}} \sinh s, \sqrt{\frac{3}{7}} \cosh s, \frac{2}{\sqrt{7}} \cos s, \frac{2}{\sqrt{7}} \sin s \right), \\ k_1^{\beta_1} &= \sqrt{7}, \quad k_2^{\beta_1} = 4\sqrt{\frac{3}{7}}, \quad k_3^{\beta_1} = \frac{1}{\sqrt{7}}. \end{aligned} \right\} \quad (4.2)$$

From (2.14), the canal hypersurfaces $\mathfrak{C}^{\{1;\lambda\}}(s, t, w)$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by the curve (4.1) are

$$\mathfrak{C}^{\{1;1\}}(s, t, w) = \left(\begin{aligned} &-2s \cosh s \left(-4 + \sqrt{15} \cos w \sin t \right) + \frac{2}{7} \sinh s \left(7 + \sqrt{35}s \left(2 \cos t \cos w + \sqrt{3} \sin w \right) \right), \\ &-2s \sinh s \left(-4 + \sqrt{15} \cos w \sin t \right) + \frac{2}{7} \cosh s \left(7 + \sqrt{35}s \left(2 \cos t \cos w + \sqrt{3} \sin w \right) \right), \\ &-4s \sin s \left(\sqrt{3} - \sqrt{5} \cos w \sin t \right) + \cos s \left(\sqrt{3} - 2\sqrt{\frac{5}{7}}s \left(\sqrt{3} \cos t \cos w - 2 \sin w \right) \right), \\ &\sin s \left(\sqrt{3} - 2\sqrt{\frac{5}{7}}s \left(\sqrt{3} \cos t \cos w - 2 \sin w \right) \right) + 4s \cos s \left(\sqrt{3} - \sqrt{5} \cos w \sin t \right) \end{aligned} \right) \quad (4.3)$$

and

$$\mathfrak{C}^{\{1;-1\}}(s, t, w) = \left(\begin{aligned} &-2s \left(4 + 3 \cos w \sin t \right) \cosh s + \frac{2}{7} \left(7 + 2\sqrt{21}s \cos t \cos w + 3\sqrt{7}s \sin w \right) \sinh s, \\ &-2s \left(4 + 3 \cos w \sin t \right) \sinh s + \frac{2}{7} \left(7 + \sqrt{21}s \left(2 \cos t \cos w + \sqrt{3} \sin w \right) \right) \cosh s, \\ &4\sqrt{3}s \left(1 + \cos w \sin t \right) \sin s + \left(\sqrt{3} - \frac{2}{\sqrt{7}}s \left(3 \cos t \cos w - 2\sqrt{3} \sin w \right) \right) \cos s, \\ &\sin s \left(\sqrt{3} - \frac{2}{\sqrt{7}}s \left(3 \cos t \cos w - 2\sqrt{3} \sin w \right) \right) - 4\sqrt{3}s \left(1 + \cos w \sin t \right) \cos s \end{aligned} \right), \quad (4.4)$$

where the radius function has been taken as $r(s) = 2s$. From (2.18), (2.19) and (2.20), the Gaussian, mean and principal curvatures of the canal hypersurfaces $\mathfrak{C}^{\{1;1\}}$ and $\mathfrak{C}^{\{1;-1\}}$ are obtained as

$$\left. \begin{aligned} K^{\{1;1\}} &= \frac{5(\sqrt{35}+14s \cos t \cos w) \cos t \cos w}{4s^2(5+2\sqrt{35}s \cos t \cos w)^2}, \\ H^{\{1;1\}} &= \frac{1}{3} \left(\frac{1}{s} + \frac{5(\sqrt{35}+14s \cos t \cos w) \cos t \cos w}{(5+2\sqrt{35}s \cos t \cos w)^2} \right), \\ \mu_1^{\{1;1\}} = \mu_2^{\{1;1\}} &= \frac{1}{2s}, \quad \mu_3^{\{1;1\}} = \frac{5(\sqrt{35}+14s \cos t \cos w) \cos t \cos w}{(5+2\sqrt{35}s \cos t \cos w)^2} \end{aligned} \right\} \quad (4.5)$$

and

$$\left. \begin{aligned} K^{\{1;-1\}} &= \frac{3(\sqrt{21}-14s \cos t \cos w) \cos t \cos w}{4s^2(3-2\sqrt{21}s \cos t \cos w)^2}, \\ H^{\{1;-1\}} &= \frac{-3+5\sqrt{21}s \cos t \cos w-42s^2 \cos^2 t \cos^2 w}{s(3-2\sqrt{21}s \cos t \cos w)^2}, \\ \mu_1^{\{1;-1\}} = \mu_2^{\{1;-1\}} &= \frac{-1}{2s}, \quad \mu_3^{\{1;-1\}} = \frac{3(\sqrt{21}-14s \cos t \cos w) \cos t \cos w}{(3-2\sqrt{21}s \cos t \cos w)^2}, \end{aligned} \right\} \quad (4.6)$$

respectively. In the following figures, one can see the projections of the canal hypersurfaces (4.3) and (4.4) for $w = 2$ and $r(s) = 2s$ into $x_2x_3x_4$ -spaces in (A) and (B), respectively.




FIGURE 1

Secondly, let us take the unit speed spacelike curve with timelike binormal vector

$$\beta_2(s) = \left(\sqrt{3} \sinh s, \sqrt{3} \cosh s, 2 \cos s, 2 \sin s \right) \quad (4.7)$$

in E_1^4 . The Frenet vectors and curvatures of the curve (4.7) are

$$\left. \begin{aligned} F_1^{\beta_2} &= \left(\sqrt{3} \cosh s, \sqrt{3} \sinh s, -2 \sin s, 2 \cos s \right), \\ F_2^{\beta_2} &= \left(\sqrt{\frac{3}{7}} \sinh s, \sqrt{\frac{3}{7}} \cosh s, -\frac{2}{\sqrt{7}} \cos s, -\frac{2}{\sqrt{7}} \sin s \right), \\ F_3^{\beta_2} &= \left(2 \cosh s, 2 \sinh s, -\sqrt{3} \sin s, \sqrt{3} \cos s \right), \\ F_4^{\beta_2} &= \left(\frac{2}{\sqrt{7}} \sinh s, \frac{2}{\sqrt{7}} \cosh s, \sqrt{\frac{3}{7}} \cos s, \sqrt{\frac{3}{7}} \sin s \right), \\ k_1^{\beta_2} &= \sqrt{7}, \quad k_2^{\beta_2} = 4\sqrt{\frac{3}{7}}, \quad k_3^{\beta_2} = \frac{1}{\sqrt{7}}. \end{aligned} \right\} \quad (4.8)$$

From (2.14), the canal hypersurfaces $\mathfrak{C}^{\{3;\lambda\}}(s, t, w)$ which are formed as the envelope of a family of pseudo hyperspheres or pseudo hyperbolic hyperspheres in E_1^4 generated by the curve (4.7) are

$$\mathfrak{C}^{\{3;1\}}(s, t, w) = \quad (4.9)$$

$$\left(\begin{array}{l} \frac{\sqrt{3}}{7} (28s(-1 + \cosh t \cosh w) \cosh s + (7 + 2\sqrt{21}s \cosh w \sinh t + 4\sqrt{7}s \sinh w)) \sinh s, \\ \frac{\sqrt{3}}{7} (28s(-1 + \cosh t \cosh w) \sinh s + (7 + 2\sqrt{21}s \cosh w \sinh t + 4\sqrt{7}s \sinh w)) \cosh s, \\ -6s \sin s \cosh t \cosh w + 2(\cos s + 4s \sin s) + \frac{2s}{\sqrt{7}} (-2\sqrt{3} \cosh w \sinh t + 3 \sinh w) \cos s, \\ 2s(-4 + 3 \cosh t \cosh w) \cos s + \frac{2}{7} (7 - 2\sqrt{21}s \cosh w \sinh t + 3\sqrt{7} \sinh w) \sin s \end{array} \right)$$

and

$$\mathfrak{C}^{\{3;-1\}}(s, t, w) = \left(\begin{array}{l} 4s (\sqrt{3} + \sqrt{5} \cosh t \cosh w) \cosh s + \left(\sqrt{3} + 2\sqrt{\frac{5}{7}}s (\sqrt{3} \cosh w \cosh t + 2 \sinh w) \right) \sinh s, \\ 4s (\sqrt{3} + \sqrt{5} \cosh t \cosh w) \sinh s + \left(\sqrt{3} + 2\sqrt{\frac{5}{7}}s (\sqrt{3} \cosh w \cosh t + 2 \sinh w) \right) \cosh s, \\ -2s (4 + \sqrt{15} \cosh t \cosh w) \sin s + \frac{2}{7} (7 + \sqrt{35}s (-2 \cosh w \sinh t + \sqrt{3} \sinh w)) \cos s, \\ 2s (4 + \sqrt{15} \cosh t \cosh w) \cos s + \frac{2}{7} (7 + \sqrt{35}s (-2 \cosh w \sinh t + \sqrt{3} \sinh w)) \sin s \end{array} \right), \quad (4.10)$$

where the radius function has been taken as $r(s) = 2s$. From (2.18), (2.19) and (2.20), the Gaussian, mean and principal curvatures of the canal hypersurfaces $\mathfrak{C}^{\{3;1\}}$ and $\mathfrak{C}^{\{3;-1\}}$ are obtained as

$$\left. \begin{array}{l} K^{\{3;1\}} = -\frac{(\sqrt{21}+14s \cosh w \sinh t) \cosh w \sinh t}{4s^2(\sqrt{3}+2\sqrt{7}s \cosh w \sinh t)^2}, \\ H^{\{3;1\}} = -\frac{3+5\sqrt{21}s \cosh w \sinh t+42s^2 \cosh^2 w \sinh^2 t}{s(3+2\sqrt{21}s \cosh w \sinh t)^2}, \\ \mu_1^{\{3;1\}} = \mu_2^{\{3;1\}} = \frac{-1}{2s}, \quad \mu_3^{\{3;1\}} = -\frac{3(\sqrt{21}+14s \cosh w \sinh t) \cosh w \sinh t}{(3+2\sqrt{21}s \cosh w \sinh t)^2} \end{array} \right\} \quad (4.11)$$

and

$$\left. \begin{array}{l} K^{\{3;-1\}} = \frac{5(-\sqrt{35}+14s \cosh w \sinh t) \cosh w \sinh t}{4s^2(5-2\sqrt{35}s \cosh w \sinh t)^2}, \\ H^{\{3;-1\}} = \frac{1}{3} \left(\frac{1}{s} + \frac{5(-\sqrt{35}+14s \cosh w \sinh t) \cosh w \sinh t}{(5-2\sqrt{35}s \cosh w \sinh t)^2} \right), \\ \mu_1^{\{3;-1\}} = \mu_2^{\{3;-1\}} = \frac{1}{2s}, \quad \mu_3^{\{3;-1\}} = \frac{5(-\sqrt{35}+14s \cosh w \sinh t) \cosh w \sinh t}{(5-2\sqrt{35}s \cosh w \sinh t)^2}, \end{array} \right\} \quad (4.12)$$

respectively. In the following figures, one can see the projections of the canal hypersurfaces (4.9) and (4.10) for $w = 2$ and $r(s) = 2s$ into $x_2x_3x_4$ -spaces in (A) and (B), respectively.




FIGURE 2

REFERENCES

- [1] M. Akyiğit, K. Eren and H.H. Kosal; *Tubular Surfaces with Modified Orthogonal Frame in Euclidean 3-Space*, Honam Math. J., 43(3), (2021), 453–463.
- [2] M. Altın, A. Kazan and H.B. Karadağ; *Monge Hypersurfaces in Euclidean 4-Space with Density*, J. Polytechnic, 23(1), (2020), 207-214.
- [3] M. Altın, A. Kazan, and D.W. Yoon; *2-Ruled hypersurfaces in Euclidean 4-space*, J. Geom. Phys., 166, (2021), 104236.
- [4] M. Altın and A. Kazan; *Rotational Hypersurfaces in Lorentz-Minkowski 4-Space*, Hacettepe J. Math and Stat., 50(5), (2021), 1409-1433.
- [5] S. Aslan and Y. Yaylı; *Canal Surfaces with Quaternions*, Adv. Appl. Clifford Algebr., 26, (2016), 31-38.
- [6] M.E. Aydın and I. Mihai; *On certain surfaces in the isotropic 4-space*, Math. Commun., 22(1), (2017), 41-51.
- [7] F. Doğan and Y. Yaylı; *Tubes with Darboux Frame*, Int. J. Contemp. Math. Sci., 7(16), (2012), 751-758.
- [8] X. Fu, S.D. Jung, J. Qian and M. Su; *Geometric Characterizations of Canal Surfaces in Minkowski 3-space I*, Bull. Korean Math. Soc., 56(4), (2019), 867-883.
- [9] R. Garcia, J. Llibre and J. Sotomayor; *Lines of Principal Curvature on Canal Surfaces in \mathbb{R}^3* , An. Acad. Brasil. Cienc., 78(3), (2006), 405-415.
- [10] E. Güler; *Helical Hypersurfaces in Minkowski Geometry E_1^4* , Symmetry, 12, (2020), 1206.
- [11] E. Hartman; *Geometry and Algorithms for Computer Aided Design*, Dept. of Math. Darmstadt Univ. of Technology, 2003.
- [12] S. Izumiya and M. Takahashi; *On caustics of submanifolds and canal hypersurfaces in Euclidean space*, Topology Appl., 159, (2012), 501-508.
- [13] M.K. Karacan, H. Es and Y. Yaylı; *Singular Points of Tubular Surfaces in Minkowski 3-Space*, Sarajevo J. Math., 2(14), (2006), 73-82.
- [14] M.K. Karacan and Y. Tuncer; *Tubular Surfaces of Weingarten Types in Galilean and Pseudo-Galilean*, Bull. Math. Anal. Appl., 5(2), (2013), 87-100.
- [15] M.K. Karacan, D.W. Yoon and Y. Tuncer; *Tubular Surfaces of Weingarten Types in Minkowski 3-Space*, Gen. Math. Notes, 22(1), (2014), 44-56.
- [16] M.K. Karacan and B. Bukcu; *An alternative moving frame for a tubular surface around a spacelike curve with a spacelike normal in Minkowski 3-space* Rendiconti del Circolo Matematico di Palermo, 57, (2008), 193-201.
- [17] A.Kazan, M. Altın and D.W. Yoon; *Geometric Characterizations of Canal Hypersurfaces in Euclidean Spaces*, arXiv:2111.04448v1 [math.DG], (2021).
- [18] S. Kızıltuğ, M. Dede and C. Ekici; *Tubular Surfaces with Darboux Frame in Galilean 3-Space*, Facta Universitatis Ser. Math. Inform., 34(2), (2019), 253-260.
- [19] Y.H. Kim, H. Liu and J. Qian; *Some Characterizations of Canal Surfaces*, Bull. Korean Math. Soc., 53(2), (2016), 461-477.
- [20] İ. Kişi, G. Öztürk and K. Arslan; *A new type of canal surface in Euclidean 4-space IE^4* , Sakarya Univ. J. Sci., 23(5), (2019), 801-809.
- [21] S.N. Krivoshapko and C.A.B. Hyeng; *Classification of Cyclic Surfaces and Geometrical Research of Canal Surfaces*, Intern. J. Research Rev. Appl. Sci., 12(3), (2012), 360-374.
- [22] W. Kuhnel; *Differential geometry: curves-surfaces-manifolds*. American Mathematical Soc., Braunschweig, Wiesbaden, 1999.
- [23] J.M. Lee; *Riemannian Manifolds-An Introduction to Curvature*, Springer-Verlag New York, Inc, 1997.
- [24] T. Maekawa, N.M. Patrikalakis, T. Sakkalis and G. Yu; *Analysis and Applications of Pipe Surfaces*, Comput. Aided Geom. Design, 15, (1998), 437-458.
- [25] M. Peternell and H. Pottmann; *Computing Rational Parametrizations of Canal Surfaces*, J. Symbolic Comput., 23, (1997), 255-266.
- [26] J.S. Ro and D.W. Yoon; *Tubes of Weingarten Type in a Euclidean 3-Space*, J. Chungcheong Math. Soc., 22(3), (2009), 359-366.
- [27] A. Uçum and K. İlarıslan; *New Types of Canal Surfaces in Minkowski 3-Space*, Adv. Appl. Clifford Algebr., 26, (2016), 449-468.
- [28] J. Walrave; *Curves and surfaces in Minkowski space*, Dissertation, K. U. Leuven, Fac. of Science, Leuven, 1995.
- [29] Z. Xu, R. Feng and J-G. Sun; *Analytic and Algebraic Properties of Canal Surfaces*, J. Comput. Appl. Math., 195, (2006), 220-228.
- [30] D.W. Yoon and Z.K. Yüzbaşı; *Tubular Surfaces with Galilean Darboux Frame in G_3* , J. Math. Phys. Anal. Geom., 15(2), (2019), 278-287.

,¹DEPARTMENT OF COMPUTER TECHNOLOGIES, DOĞANŞEHİR VAHAP KÜÇÜK VOCATIONAL SCHOOL, MALATYA TURGUT ÖZAL UNIVERSITY, MALATYA, TURKEY,,²TECHNICAL SCIENCES VOCATIONAL SCHOOL, BİNGÖL UNIVERSITY, BİNGÖL, TURKEY,,³DEPARTMENT OF MATHEMATICS EDUCATION AND RİNS, GYEONGSANG NATIONAL UNIVERSITY, JİNJU 52828, REPUBLIC OF KOREA,, *CORRESPONDING AUTHOR: AHMET.KAZAN@OZAL.EDU.TR