

Factorial-Type Recurrence Relations and p -adic Incomplete Gamma Functions

Paul Buckingham

Abstract

We introduce an automorphism \mathcal{S} of the space $C(\mathbb{Z}_p, \mathbb{C}_p)$ of continuous functions $\mathbb{Z}_p \rightarrow \mathbb{C}_p$ and show that it can be used to give an alternative construction of the p -adic incomplete Γ -functions recently introduced by O’Desky and Richman [7]. We then describe various properties of the automorphism \mathcal{S} , showing that it is self-adjoint with respect to a certain non-degenerate symmetric bilinear form defined in terms of p -adic integration, and showing that its inverse plays a role in a p -adic integral-transform space akin to the role of differentiation in the classical space of Laplace-transformed functions. We also derive an integral-transform formula for the p -adic incomplete Γ -functions.

Keywords: p -adic incomplete Gamma functions; p -adic integral transform

Mathematics Subject Classification (2020): 11S80

1 Introduction

In number theory, one is interested in finding p -adic analogues of classical functions of a real or complex variable. Important examples are p -adic L -functions and Morita’s p -adic Γ -function. These p -adic analogues are usually defined via some interpolation property. More precisely, if f is a complex function whose values at the positive integers can be viewed also as elements of the field \mathbb{C}_p , via a choice of isomorphism $\mathbb{C} \cong \mathbb{C}_p$, say, then one looks for a continuous p -adic function $F : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ that takes the same values as f on $\mathbb{Z}_{\geq 1}$. By the denseness of $\mathbb{Z}_{\geq 1}$ in \mathbb{Z}_p , the continuous function F , if it exists, is uniquely determined. In practice, the function f may need to be modified first, and the set of points at which one interpolates may be some other subset of \mathbb{Z} dense in \mathbb{Z}_p .

In a recent paper [7], O’Desky and Richman construct a p -adic analogue of the incomplete Γ -function $\Gamma(-, r)$ for each $r \in 1 + p\mathbb{Z}$. Their construction uses the combinatorial machinery of derangements.

Here, we introduce an automorphism \mathcal{S} of the space $C(\mathbb{Z}_p, \mathbb{C}_p)$ of continuous functions, restricting to an automorphism of the space $\text{la}(\mathbb{Z}_p, \mathbb{C}_p)$ of locally analytic functions $\mathbb{Z}_p \rightarrow \mathbb{C}_p$, and show that O’Desky and Richman’s p -adic incomplete Γ -function $\Gamma_p(-, r)$ can be constructed by applying \mathcal{S}^{-1} to the locally

analytic p -adic function $x \mapsto r^x$ ($r \in 1 + p\mathbb{Z}$). We thus give a construction of O’Desky and Richman’s functions that does not use derangements.

We then turn to proving properties of the automorphism \mathcal{S} . We show in Section 4 that it is self-adjoint with respect to a non-degenerate symmetric bilinear form on $C(\mathbb{Z}_p, \mathbb{C}_p)$ defined in terms of p -adic integration. We also show that, with respect to a convolution product \star on $C(\mathbb{Z}_p, \mathbb{C}_p)$, \mathcal{S} again enjoys a sort of self-adjointness; see Section 5.

Next, in Section 6, we introduce an integral transform $\mathcal{I} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$, again related to p -adic integration, describe its relationship with \mathcal{S} , derive an integral-transform formula for the p -adic incomplete Γ -functions, and give a complete description of the image of \mathcal{I} . We also prove a convolution property of \mathcal{I} .

Finally, in Section 7, we relate \mathcal{S} to solutions of the well-known differential equation $F' + F = G$. For example, if $G \in \mathbb{Z}[[t]]$, then the unique solution $F \in \mathbb{Z}_p[[t]]$ can be constructed concretely via \mathcal{S}^{-1} . We work in a more general setting than \mathbb{Z}_p . The precise result is Proposition 7.3.

Running throughout is the function $q = \mathcal{S}^{-1}(\mathbf{1})$, where $\mathbf{1} : x \mapsto 1$. Up to a certain constant multiplicative factor and a one-unit shift in the argument, q is the p -adic analogue of $\Gamma(-, 1)$. It features prominently in results on the convolution product \star and in the description of the image of \mathcal{I} . The function q has been well studied in various contexts (see Section 2.4 for a sample of works), and we hope our contribution will provide further evidence that it may have a role to play in p -adic analysis.

1.1 Notation and definitions

Throughout, $|\cdot|$ and v_p will denote, respectively, the absolute value and the valuation on \mathbb{C}_p , normalized such that $|p| = 1/p$ and $v_p(p) = 1$. We will occasionally make reference to the balls $B_{<\rho}(a, K) = \{x \in K \mid |x - a| < \rho\}$, where K is a subfield of \mathbb{C}_p , $a \in K$, and ρ is a positive real number.

Frequently used will be the functions $\beta_n : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ ($n \geq 0$) defined by $\beta_n(x) = \binom{x}{n} = \frac{1}{n!}(x)_n$, where $(x)_n = x(x-1)(x-2)\cdots(x-n+1)$. They satisfy $\nabla\beta_n = \beta_{n-1}$ for all $n \geq 1$, where ∇ is the forward-difference operator, i.e., $(\nabla\phi)(x) = \phi(x+1) - \phi(x)$ for a function ϕ on \mathbb{Z}_p . By Mahler’s Theorem, the continuous functions $\phi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ are the functions that may be expressed as $\phi = \sum_{n=0}^{\infty} b_n\beta_n$ where the elements $b_n \in \mathbb{C}_p$ converge to zero.

If $\phi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$, we will be interested in the functions

$$\begin{aligned} \widehat{\phi} : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ x &\mapsto \phi(-1-x) \end{aligned}$$

and

$$\begin{aligned} \sigma(\phi) : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ x &\mapsto \phi(x+1). \end{aligned}$$

We will also refer often to the constant function $\mathbf{1} : \mathbb{Z}_p \rightarrow \mathbb{C}_p$, $x \mapsto 1$.

2 The automorphism \mathcal{S}

Let $C(\mathbb{Z}_p, \mathbb{C}_p)$ denote the \mathbb{C}_p -vector space of continuous functions $\mathbb{Z}_p \rightarrow \mathbb{C}_p$. We define a map

$$\mathcal{S} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow C(\mathbb{Z}_p, \mathbb{C}_p)$$

as follows. If $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\mathcal{S}(\phi)$ is the function $\mathbb{Z}_p \rightarrow \mathbb{C}_p$ defined by

$$\mathcal{S}(\phi)(x) = \phi(x) - x\phi(x-1).$$

It is clear that $\mathcal{S}(\phi)$ is again continuous.

Lemma 2.1 *Suppose that $\phi_0, \phi_1, \phi_2, \dots$ is a sequence of functions in $C(\mathbb{Z}_p, \mathbb{C}_p)$ that converges uniformly to zero. Then the function $\phi = \sum_{k=0}^{\infty} \phi_k$ is in $C(\mathbb{Z}_p, \mathbb{C}_p)$, and $\mathcal{S}(\phi) = \sum_{k=0}^{\infty} \mathcal{S}(\phi_k)$.*

Proof. The continuity of ϕ follows from the assumption of uniform convergence. For the second assertion, observe that for $x \in \mathbb{Z}_p$,

$$\begin{aligned} \mathcal{S}(\phi)(x) &= \phi(x) - x\phi(x-1) \\ &= \sum_{k=0}^{\infty} \phi_k(x) - x \sum_{k=0}^{\infty} \phi_k(x-1) \\ &= \sum_{k=0}^{\infty} (\phi_k(x) - x\phi_k(x-1)) \\ &= \sum_{k=0}^{\infty} \mathcal{S}(\phi_k)(x). \end{aligned}$$

■

Proposition 2.2 *If $\phi = \sum_{n=0}^{\infty} b_n \beta_n$ is continuous, then*

$$\mathcal{S}(\phi) = \sum_{n=0}^{\infty} (b_n - nb_{n-1}) \beta_n,$$

where we have set $b_{-1} = 0$.

Proof. First note that

$$\mathcal{S}(\beta_n)(x) = \binom{x}{n} - x \binom{x-1}{n} = \binom{x}{n} - (n+1) \binom{x}{n+1},$$

so $\mathcal{S}(\beta_n) = \beta_n - (n+1)\beta_{n+1}$. Hence,

$$\mathcal{S}(\phi) = \sum_{n=0}^{\infty} b_n (\beta_n - (n+1)\beta_{n+1}) \quad \text{by Lemma 2.1}$$

Paul Buckingham

$$\begin{aligned}
&= \sum_{n=0}^{\infty} b_n \beta_n - \sum_{n=1}^{\infty} n b_{n-1} \beta_n \\
&= \sum_{n=0}^{\infty} (b_n - n b_{n-1}) \beta_n.
\end{aligned}$$

■

Lemma 2.3 *If $k \geq 0$, then*

$$\mathcal{S} \left(\sum_{n=k}^{\infty} \frac{n!}{k!} \beta_n \right) = \beta_k.$$

Proof. Let $\phi = \sum_{n=k}^{\infty} \frac{n!}{k!} \beta_n$. Note that ϕ is a well-defined function in $C(\mathbb{Z}_p, \mathbb{C}_p)$, because its sequence of Mahler coefficients converges to 0. Then by Proposition 2.2,

$$\mathcal{S}(\phi) = \beta_k + \sum_{n=k+1}^{\infty} \left(\frac{n!}{k!} - n \frac{(n-1)!}{k!} \right) \beta_n = \beta_k + \sum_{n=k+1}^{\infty} \left(\frac{n!}{k!} - \frac{n!}{k!} \right) \beta_n = \beta_k.$$

■

Theorem 2.4 *The map \mathcal{S} is an automorphism of the \mathbb{C}_p -vector space $C(\mathbb{Z}_p, \mathbb{C}_p)$. If $\psi = \sum_{k=0}^{\infty} b_k \beta_k$, then*

$$\mathcal{S}^{-1}(\psi) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{n!}{k!} b_k \right) \beta_n. \quad (2.1)$$

Proof. For injectivity, suppose that $\mathcal{S}(\phi) = 0$, i.e., $\phi(x) = x\phi(x-1)$ for all $x \in \mathbb{Z}_p$. We show by induction on n that $\phi(n) = 0$ for all $n \in \mathbb{Z}_{\geq 0}$, from which it follows that $\phi = 0$ by continuity. The base case is clear, because $\phi(0) = 0 \cdot \phi(-1) = 0$. Now let $n \geq 0$ and assume that $\phi(n) = 0$. Then $\phi(n+1) = (n+1)\phi(n) = 0$. This completes the induction and therefore the proof of injectivity.

For surjectivity, let $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ be any continuous function, and let $\psi = \sum_{k=0}^{\infty} b_k \beta_k$ be its Mahler series, so that $b_k \rightarrow 0$. We already know from Lemma 2.3 that $\mathcal{S}(\phi_k) = \beta_k$, where $\phi_k = \sum_{n=k}^{\infty} \frac{n!}{k!} \beta_n$. Because the functions ϕ_k are all maps into \mathbb{Z}_p , they have sup-norm $\|\phi_k\| \leq 1$, so the sequence of functions $b_k \phi_k$ converges uniformly to zero. Hence, by uniform convergence, we have a well-defined, continuous function

$$\phi = \sum_{k=0}^{\infty} b_k \phi_k$$

from \mathbb{Z}_p to \mathbb{C}_p . Then

$$\begin{aligned} \mathcal{S}(\phi) &= \sum_{k=0}^{\infty} \mathcal{S}(b_k \phi_k) \quad \text{by Lemma 2.1} \\ &= \sum_{k=0}^{\infty} b_k \beta_k \\ &= \psi. \end{aligned}$$

As for the formula in (2.1), if $x \in \mathbb{Z}_p$, then

$$\phi(x) = \sum_{k=0}^{\infty} b_k \sum_{n=k}^{\infty} \frac{n!}{k!} \beta_n(x) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{n!}{k!} b_k \right) \beta_n(x),$$

the interchanging of the summation signs being justified as follows. Fix $x \in \mathbb{Z}_p$ and let

$$\eta(n, k) = \begin{cases} \frac{n!}{k!} b_k \beta_n(x) & \text{if } k \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $|\eta(n, k)| \leq \frac{n!}{k!} |b_k|$. By [9, Chap. 2, Sect. 1.2, Cor.], it is enough to show that, for any $\epsilon > 0$, there are only finitely many pairs (n, k) such that $|\eta(n, k)| > \epsilon$. Let $\epsilon > 0$, then. Because $b_k \rightarrow 0$, there is $M \geq 0$ such that $|b_k| \leq \epsilon$ for all $k \geq M$. For each $k < M$, choose $N_k \geq 0$ such that $\frac{n!}{k!} |b_k| \leq \epsilon$ for all $n \geq N_k$, possible because $|n!/k!| \rightarrow 0$ as $n \rightarrow \infty$. Now let $N = \max(N_0, \dots, N_{M-1})$. If $k \geq M$, then $|\eta(n, k)| \leq |b_k| \leq \epsilon$. On the other hand, if $k < M$ and $n \geq N$, then either $k > n$, in which case $\eta(n, k) = 0$, or $k \leq n$, and then $|\eta(n, k)| \leq \frac{n!}{k!} |b_k| \leq \epsilon$ because $n \geq N \geq N_k$. Thus, $|\eta(n, k)| \leq \epsilon$ except possibly for the finitely many pairs $(n, k) \in \{0, \dots, N-1\} \times \{0, \dots, M-1\}$. ■

2.1 Locally analytic functions

We use the definition of *locally analytic* given in [3, Sect. I.4]. The following proposition is [3, Cor. I.4.8]. See also [1, Sect. 10, Cor. 1] in the case where \mathbb{Z}_p maps into \mathbb{Q}_p .

Proposition 2.5 *If $f : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ has Mahler coefficients $(a_n)_n$, then f is locally analytic on \mathbb{Z}_p if and only if $\liminf_{n \rightarrow \infty} \frac{v_p(a_n)}{n} > 0$. Equivalently, if we set $a'_n = a_n/n!$, then f is locally analytic on \mathbb{Z}_p if and only if $\liminf_{n \rightarrow \infty} \frac{v_p(a'_n)}{n} > -\frac{1}{p-1}$.*

The following equivalent form of Proposition 2.5 is an immediate consequence of the fact that, if $(x_n)_n$ is a sequence of real numbers, then the sequence $(\inf_{k \geq n} x_k)_n$ is monotone non-decreasing.

Proposition 2.6 *If $f : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ has Mahler coefficients $(a_n)_n$, then f is locally analytic on \mathbb{Z}_p if and only if there are $\beta > 0$ and $N \in \mathbb{Z}_{\geq 0}$ such that $v_p(a_n) \geq n\beta$ for all $n \geq N$. Equivalently, if we set $a'_n = a_n/n!$, then f is locally analytic on \mathbb{Z}_p if and only if there are $\beta > -\frac{1}{p-1}$ and $N \in \mathbb{Z}_{\geq 0}$ such that $v_p(a'_n) \geq n\beta$ for all $n \geq N$.*

We use Proposition 2.6 to prove the following.

Theorem 2.7 *The map \mathcal{S} restricts to an automorphism of the \mathbb{C}_p -vector space $\text{la}(\mathbb{Z}_p, \mathbb{C}_p)$.*

Proof. It is clear from the definition of \mathcal{S} that $\mathcal{S}(\phi) \in \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$ if ϕ is. It is also clear from the injectivity of \mathcal{S} as a map on $C(\mathbb{Z}_p, \mathbb{C}_p)$ that it is still injective when restricted to $\text{la}(\mathbb{Z}_p, \mathbb{C}_p)$. It remains to show that if $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is locally analytic, then the unique $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ such that $\mathcal{S}(\phi) = \psi$ is also locally analytic.

Let the Mahler coefficients of ψ be $(b_n)_{n \geq 0}$. By Theorem 2.4, the n th Mahler coefficient of ϕ is

$$a_n = \sum_{k=0}^n \frac{n!}{k!} b_k.$$

Letting $a'_n = a_n/n!$ and $b'_n = b_n/n!$, we therefore have

$$a'_n = \sum_{k=0}^n b'_k. \tag{2.2}$$

The assumption that ψ is locally analytic says, by Proposition 2.6, that there are $\beta > -\frac{1}{p-1}$ and $N \in \mathbb{Z}_{\geq 0}$ such that, for all $n \geq N$,

$$v_p(b'_n) \geq n\beta.$$

Assume now that $n \geq N$, and write

$$a'_n = s + \sum_{k=N}^n b'_k, \tag{2.3}$$

where $s = \sum_{k=0}^{N-1} b'_k$. Let us first consider the case where $\beta \geq 0$, in which case $v_p(b'_k) \geq 0$ for all $k \geq N$. In this case, $v_p(a'_n) \geq \min(v_p(s), 0)$, so $v_p(a'_n)/n \geq \frac{1}{n} \min(v_p(s), 0)$, which tends to 0 as $n \rightarrow \infty$. Therefore, if α is any negative real number, $v_p(a'_n) \geq n\alpha$ for n sufficiently large. Thus, ϕ is locally analytic by Proposition 2.6, because we may choose α such that $-1/(p-1) < \alpha < 0$.

We may thus assume for the remainder of the proof that $\beta < 0$. Then because $v_p(s)/n \rightarrow 0$, there is $K \geq 1$ such that, if $n \geq K$, then $v_p(s)/n \geq \beta$, i.e.,

$$v_p(s) \geq n\beta.$$

Further, each b'_k in the sum in (2.3) has valuation greater than or equal to $k\beta$ because $k \geq N$ in the sum. But $k\beta \geq n\beta$, because $\beta < 0$ and $k \leq n$ for k in

the sum. Thus, the summation term in (2.3) has valuation at least $n\beta$ as well. Therefore, $v_p(a'_n) \geq n\beta$ for all $n \geq \max(N, K)$, and so ϕ is locally analytic by Proposition 2.6. ■

2.2 An alternative description of $\mathcal{S}^{-1}(\psi)$

If $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is continuous, then because \mathbb{Z}_p is compact, the same is true of $\psi(\mathbb{Z}_p)$, which is therefore bounded, say $|\psi(x)| \leq M$ for all $x \in \mathbb{Z}_p$. Consequently, for each $n \geq 0$,

$$\sup_{x \in \mathbb{Z}_p} |(x)_n \psi(x - n)| = \sup_{x \in \mathbb{Z}_p} \left| n! \binom{x}{n} \psi(x - n) \right| \leq |n!|M,$$

and the real numbers $|n!|M$ converge to 0 as $n \rightarrow \infty$. Therefore, by uniform convergence, we have a well-defined, continuous function $\mathbb{Z}_p \rightarrow \mathbb{C}_p$ given by

$$x \mapsto \sum_{n=0}^{\infty} (x)_n \psi(x - n).$$

Proposition 2.8 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then*

$$\mathcal{S}^{-1}(\psi)(x) = \sum_{n=0}^{\infty} (x)_n \psi(x - n)$$

for all $x \in \mathbb{Z}_p$.

Proof. Let $\phi(x) = \sum_{n=0}^{\infty} (x)_n \psi(x - n)$. Then

$$\begin{aligned} \mathcal{S}(\phi)(x) &= \sum_{n=0}^{\infty} (x)_n \psi(x - n) - x \sum_{n=0}^{\infty} (x - 1)_n \psi(x - 1 - n) \\ &= \sum_{n=0}^{\infty} (x)_n \psi(x - n) - \sum_{n=0}^{\infty} (x)_{n+1} \psi(x - (n + 1)) \\ &= \sum_{n=0}^{\infty} (x)_n \psi(x - n) - \sum_{n=1}^{\infty} (x)_n \psi(x - n) \\ &= \psi(x). \end{aligned}$$

Thus, $\mathcal{S}(\phi) = \psi$, so $\mathcal{S}^{-1}(\psi) = \phi$. ■

2.3 Relationship to factorial-type recurrence relations

Consider a first-order linear recurrence relation

$$a_n = h(n)a_{n-1} + \psi(n) \quad (n \geq 1) \tag{2.4}$$

where $h, \psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ are continuous functions and the a_n are in \mathbb{C}_p . Let us say that a solution $(a_n)_{n \geq 0}$ to such a recurrence relation is *continuous* if there is a continuous function $\phi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ such that $\phi(n) = a_n$ for all $n \geq 0$.

By an *automatically seeded* recurrence relation, we will mean a recurrence relation as in (2.4) for which the function h has a zero $n_0 \in \mathbb{Z}$. Note that n_0 may be positive, negative, or zero. The terminology is explained via the following proposition and its proof.

Proposition 2.9 *If an automatically seeded recurrence relation admits a continuous solution, then the solution is unique.*

Proof. Suppose that $(a_n)_{n \geq 0}$ is a continuous solution to $a_n = h(n)a_{n-1} + \psi(n)$ ($n \geq 1$), so that there is a continuous function $\phi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ such that $\phi(n) = a_n$ for all $n \geq 0$. Then for $n \geq 1$,

$$\phi(n) = a_n = h(n)a_{n-1} + \psi(n) = h(n)\phi(n-1) + \psi(n),$$

so because ϕ , ψ , and h are all continuous, we in fact have $\phi(x) = h(x)\phi(x-1) + \psi(x)$ for all $x \in \mathbb{Z}_p$. In particular,

$$\phi(n_0) = h(n_0)\phi(n_0-1) + \psi(n_0) = \psi(n_0)$$

by the assumption that $h(n_0) = 0$. Hence, if $(\tilde{a}_n)_{n \geq 0}$ is another continuous solution, and if $\phi_1(n) = \tilde{a}_n$ for all $n \geq 0$, then $\phi(n_0) = \phi_1(n_0)$, so $\phi(n) = \phi_1(n)$ for all $n \geq n_0$, and so $\phi = \phi_1$ by continuity. Thus, $a_n = \tilde{a}_n$ for all $n \geq 0$. ■

In particular, the value a_{n_0} in an automatically seeded recurrence relation admitting a continuous solution is determined by the recurrence relation itself, unlike in an ordinary recurrence relation, where a_{n_0} may typically be chosen freely.

For us, the function h of interest will be $h(x) = x$, so that the recurrence relation becomes $a_n = na_{n-1} + \psi(n)$. It seems reasonable to call this a *factorial-type* recurrence relation, because one recovers the usual factorial recurrence relation by taking $\psi = 0$.

Theorem 2.4 says that every factorial-type recurrence relation $a_n = na_{n-1} + \psi(n)$, where $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, has the unique continuous solution $a_n = \mathcal{S}^{-1}(\psi)(n)$. Thus, every time we invoke the inverse map \mathcal{S}^{-1} in this paper, we are essentially considering the unique continuous solution to some factorial-type recurrence relation. Although this point of view will not be emphasized, the reader may well like to keep it in mind while reading the rest of the paper.

2.4 The function q

Central to this paper is the function $q = \mathcal{S}^{-1}(\mathbf{1})$, where $\mathbf{1} : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is the constant function $x \mapsto 1$. By Proposition 2.8, $q(x) = \sum_{n=0}^{\infty} (x)_n = \sum_{n=0}^{\infty} n! \binom{x}{n}$. By definition, it satisfies $q(x) = xq(x-1) + 1$ for all $x \in \mathbb{Z}_p$ and therefore gives the unique continuous solution to the recurrence relation $a_n = na_{n-1} + 1$. We

note as well that it is locally analytic by Theorem 2.7. (See also [8], where the function is called A .)

This function has been studied many times already, in several contexts and with various symbols to denote it. Aside from the reference [8] just mentioned, we point out just a few more: [5, Lemma 5], where a connection to the incomplete Γ -function $\Gamma(-, 1)$ is made, and [10] and [2], which consider convergents to e , the latter proving a conjecture of Sondow.

2.5 p -adic incomplete Γ -functions

In [7], O'Desky and Richman define a p -adic incomplete Γ -function $\Gamma_p(-, r)$ for each $r \in 1 + p\mathbb{Z}_p$ via the combinatorial machinery of derangements. Their function satisfies a p -adic interpolation property with respect to the usual (complex) incomplete Γ -function $\Gamma(-, r)$. We give a construction using instead the invertible map $\mathcal{S} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow C(\mathbb{Z}_p, \mathbb{C}_p)$.

For each $r \in B_{<1}(1, \mathbb{C}_p)$, define

$$\begin{aligned} g_r : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ x &\mapsto r^x, \end{aligned}$$

locally analytic by Proposition 2.5. Let $\gamma_r = \mathcal{S}^{-1}(g_r)$, also locally analytic by Theorem 2.7. Define \tilde{p} to be p if p is odd and to be 4 if $p = 2$, and define

$$\begin{aligned} \bar{\gamma}_r : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ x &\mapsto \exp(r\tilde{p})\gamma_r(x-1), \end{aligned}$$

again obviously locally analytic, because γ_r is. The function $\bar{\gamma}_r$ will give us the desired interpolation property.

By definition of \mathcal{S} and g_r ,

$$\begin{aligned} \gamma_r(x) &= x\gamma_r(x-1) + r^x \quad \text{for all } x \in \mathbb{Z}_p, \\ \text{i.e., } \exp(r\tilde{p})\gamma_r(x) &= x\exp(r\tilde{p})\gamma_r(x-1) + r^x \exp(r\tilde{p}) \quad \text{for all } x \in \mathbb{Z}_p, \\ \text{i.e., } \bar{\gamma}_r(x+1) &= x\bar{\gamma}_r(x) + r^x \exp(r\tilde{p}) \quad \text{for all } x \in \mathbb{Z}_p. \end{aligned}$$

In particular,

$$\bar{\gamma}_r(n+1) = n\bar{\gamma}_r(n) + r^n \exp(r\tilde{p}) \quad \text{for all } n \in \mathbb{Z}_{\geq 0}. \quad (2.5)$$

Note also that $\bar{\gamma}_r(1) = 0 \cdot \bar{\gamma}_r(0) + \exp(r\tilde{p}) = \exp(r\tilde{p})$.

Now suppose that $r \in B_{<1}(1, \mathbb{C}_p) \cap \mathbb{Z} = 1 + p\mathbb{Z}$. It is well known that the ordinary (complex) incomplete Γ -function $\Gamma(-, r)$ maps $\mathbb{Z}_{\geq 1}$ into $\mathbb{Q}(e)$ and satisfies

$$\Gamma(n+1, r) = n\Gamma(n, r) + r^n e^{-r} \quad \text{for all } n \in \mathbb{Z}_{\geq 1}. \quad (2.6)$$

Consider, then, the unique field map $\tau_p : \mathbb{Q}(e) \rightarrow \mathbb{Q}_p$ mapping e^{-1} to $\exp(\tilde{p})$. (The map τ_p appears already in O'Desky and Richman's paper; we do not claim τ_p as our own.) Applying τ_p to both sides of (2.6), we have

$$\tau_p(\Gamma(n+1, r)) = n\tau_p(\Gamma(n, r)) + r^n \exp(r\tilde{p}) \quad \text{for all } n \in \mathbb{Z}_{\geq 1}.$$

Thus, the values $\tau_p(\Gamma(n, r))$, $n \in \mathbb{Z}_{\geq 1}$, satisfy the same recurrence relation as the values $\bar{\gamma}_r(n)$, as in (2.5). Further,

$$\bar{\gamma}_r(1) = \exp(r\tilde{p}) = \tau_p(e^{-r}) = \tau_p(\Gamma(1, r)),$$

so in fact $\bar{\gamma}_r(n) = \tau_p(\Gamma(n, r))$ for all $n \geq 1$. Thus, the p -adic locally analytic function $\bar{\gamma}_r$ interpolates the values $\tau_p(\Gamma(n, r))$ at positive integers n . The existence of a p -adic locally analytic function with this interpolation property is what O'Desky and Richman showed, but via a different method.

Of course, since $g_1 = \mathbf{1}$, we have $\gamma_1 = q$.

Remark. We point out a small discrepancy between our function $\bar{\gamma}_r$ and O'Desky and Richman's $\Gamma_p(-, r)$ in the case where $p = 2$. They define the map $\tau_p : \mathbb{Q}(e) \rightarrow \mathbb{Q}_p$ by sending e^{-1} to $\exp(p)$, while our choice is to map e^{-1} to $\exp(\tilde{p})$. The maps are the same when p is odd, but our choice is made to ensure that the exponential series $\exp(\tilde{p})$ converges even in the case $p = 2$, where $\tilde{p} = 4$. (Note that $\exp(x)$ converges if and only if $|x| < |p|^{1/(p-1)}$.) The exceptional nature of the prime 2 often requires the introduction of an extra factor of 2, especially in Iwasawa theory. See [11, Sect. 7.2], for example. Another option might be to employ a continuous extension of \exp to \mathbb{C}_2 , so that $\exp(2)$ is defined for that extension. However, there is no canonical such extension. See [9, Chap. 5, Sect. 4.4] for a discussion.

We will provide an integral-transform formula for $\bar{\gamma}_r$ in Section 6.2.1.

2.5.1 Mahler series of $\bar{\gamma}_r(x+1)$

Proposition 2.10 *Let $r \in B_{<1}(1, \mathbb{C}_p)$. Then the Mahler series of $\bar{\gamma}_r(x+1)$ is*

$$\bar{\gamma}_r(x+1) = \sum_{n=0}^{\infty} \left(\exp(r\tilde{p}) \sum_{k=0}^n \frac{n!}{k!} (r-1)^k \right) \binom{x}{n}.$$

Proof. Because $\gamma_r = \mathcal{S}^{-1}(g_r)$ and $g_r = \sum_{n=0}^{\infty} (r-1)^n \beta_n$, Theorem 2.4 shows that

$$\gamma_r(x) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{n!}{k!} (r-1)^k \right) \binom{x}{n}.$$

Hence,

$$\bar{\gamma}_r(x+1) = \exp(r\tilde{p})\gamma_r(x) = \sum_{n=0}^{\infty} \left(\exp(r\tilde{p}) \sum_{k=0}^n \frac{n!}{k!} (r-1)^k \right) \binom{x}{n}.$$

■

2.5.2 Recovering O’Desky and Richman’s formula

Since $\bar{\gamma}_r$ and $\Gamma_p(-, r)$ interpolate the same p -adic values at the positive integers (with the above remark about the $p = 2$ case understood), they must be the same function, and indeed, the following formula for $\bar{\gamma}$ concurs with that given for $\Gamma_p(-, r)$ in [7, Theorem 1.1].

Proposition 2.11 *Let $r \in B_{<1}(1, \mathbb{C}_p)$. Then*

$$\bar{\gamma}_r(x) = \exp(r\bar{p}) \sum_{n=0}^{\infty} (x-1)_n r^{x-1-n} = \exp(r\bar{p}) \sum_{n=0}^{\infty} n! \binom{x-1}{n} r^{x-1-n}.$$

Proof. Because $\gamma_r = \mathcal{S}^{-1}(g_r)$, Proposition 2.8 shows that

$$\gamma_r(x) = \sum_{n=0}^{\infty} (x)_n r^{x-n}.$$

Therefore, since $\bar{\gamma}_r(x) = \exp(r\bar{p})\gamma_r(x-1)$, the formula follows. ■

2.6 1-Lipschitz functions

Recall that a function $\phi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is called 1-Lipschitz if $|\phi(x) - \phi(y)| \leq |x - y|$ for all $x, y \in \mathbb{Z}_p$. Such a function is obviously continuous.

Proposition 2.12 *Let $A = \{x \in \mathbb{C}_p \mid |x| \leq 1\}$. The map \mathcal{S} restricts to an automorphism of the A -module of 1-Lipschitz functions $\mathbb{Z}_p \rightarrow A$.*

Proof. First, suppose that $\phi : \mathbb{Z}_p \rightarrow A$ is 1-Lipschitz. It is clear from the definition of $\mathcal{S}(\phi)$ that its image is again in A . For the 1-Lipschitz property of $\mathcal{S}(\phi)$, we take $x, y \in \mathbb{Z}_p$. Then

$$\begin{aligned} & |\mathcal{S}(\phi)(x) - \mathcal{S}(\phi)(y)| \\ &= |\phi(x) - x\phi(x-1) - \phi(y) + y\phi(y-1)| \\ &= |(\phi(x) - \phi(y)) - x(\phi(x-1) - \phi(y-1)) + (y-x)\phi(y-1)| \\ &\leq \max(|\phi(x) - \phi(y)|, |x| |\phi(x-1) - \phi(y-1)|, |y-x| |\phi(y-1)|) \\ &\leq \max(|x-y|, |x| |x-y|, |x-y| |\phi(y-1)|) \\ &\leq |x-y| \end{aligned}$$

by the assumption that the image of ϕ is contained in A .

Conversely, suppose that $\psi : \mathbb{Z}_p \rightarrow A$ is 1-Lipschitz, and let $\phi = \mathcal{S}^{-1}(\psi)$, so that $\phi(x) = x\phi(x-1) + \psi(x)$ for all $x \in \mathbb{Z}_p$. It is clear from Proposition 2.8 that ϕ has its image in A . As for the 1-Lipschitz property of ϕ , it is equivalent to the statement that for all $a, x \in \mathbb{Z}_p$, $|\phi(x+a) - \phi(x)| \leq |a|$. To that end, let us fix $a \in \mathbb{Z}_p$ and introduce an equivalence relation \equiv_a on A defined by $\alpha \equiv_a \beta$ if $|\alpha - \beta| \leq |a|$. (Transitivity holds because of the ultrametric property.)

Paul Buckingham

We first show by induction on x that if $x \in \mathbb{Z}_{\geq 0}$, then $\phi(x+a) \equiv_a \phi(x)$. For the case $x = 0$, observe that

$$\begin{aligned} \phi(a) &= a\phi(a-1) + \psi(a) \\ &\equiv_a \psi(a) \quad \text{because } |\phi(a-1)| \leq 1 \\ &\equiv_a \psi(0) \quad \text{because } \psi \text{ is 1-Lipschitz} \\ &= \phi(0). \end{aligned}$$

Now suppose that $x \in \mathbb{Z}_{\geq 0}$ is such that $\phi(x+a) \equiv_a \phi(x)$. Then

$$\begin{aligned} \phi(x+1+a) &= (x+1+a)\phi(x+a) + \psi(x+1+a) \\ &\equiv_a (x+1)\phi(x+a) + \psi(x+1+a) \quad \text{because } |\phi(x+a)| \leq 1 \\ &\equiv_a (x+1)\phi(x) + \psi(x+1+a) \quad \text{by the inductive hyp. on } x \\ &\equiv_a (x+1)\phi(x) + \psi(x+1) \quad \text{because } \psi \text{ is 1-Lipschitz} \\ &= \phi(x+1). \end{aligned}$$

We now know that $\phi(x+a) \equiv_a \phi(x)$ for all $x \in \mathbb{Z}_{\geq 0}$, and it remains to extend this to $x \in \mathbb{Z}_p$, which we may achieve by continuity. Specifically, the function

$$\begin{aligned} \mathbb{Z}_p &\rightarrow A \\ x &\mapsto \phi(x+a) - \phi(x) \end{aligned}$$

maps $\mathbb{Z}_{\geq 0}$ into the set $V = \{\alpha \in A \mid |\alpha| \leq |a|\}$ and therefore, being continuous, maps \mathbb{Z}_p into V as well because V is closed in A . \blacksquare

We apply the above to the functions γ_r with $r \in 1 + p\mathbb{Z}_p$.

Proposition 2.13 *Suppose that $r \in \mathbb{C}_p$ satisfies $|r-1| \leq (1/3)^{1/3}$. (This condition is satisfied, in particular, if $r \in 1 + p\mathbb{Z}_p$, since $(1/3)^{1/3} > 1/p$ no matter what prime p is.) Then γ_r is 1-Lipschitz.*

Proof. Recall that $\gamma_r = \mathcal{S}^{-1}(g_r)$ where $g_r(x) = r^x$. By Proposition 2.12, the statement to be proven is equivalent to the statement that g_r is 1-Lipschitz. For that, note that if $x, y \in \mathbb{Z}_p$, and if $n \geq 0$, then

$$\left| \binom{x}{n} - \binom{y}{n} \right| \leq n|x-y|.$$

This is a weak form of the lemma in [9, Chap. 5, Sect. 1.5]. Hence,

$$\begin{aligned} |g_r(x) - g_r(y)| &= \left| \sum_{n=0}^{\infty} (r-1)^n \left(\binom{x}{n} - \binom{y}{n} \right) \right| \\ &= \left| \sum_{n=1}^{\infty} (r-1)^n \left(\binom{x}{n} - \binom{y}{n} \right) \right| \quad (\text{the } n=0 \text{ term vanishes}) \end{aligned}$$

Paul Buckingham

$$\begin{aligned} &\leq \max_{n \in \mathbb{Z}_{\geq 1}} |r - 1|^n n |x - y| \\ &= |x - y| \max_{n \in \mathbb{Z}_{\geq 1}} n |r - 1|^n. \end{aligned}$$

We claim that $\max_{n \in \mathbb{Z}_{\geq 1}} n |r - 1|^n \leq 1$. Indeed, this inequality holds if and only if $n |r - 1|^n \leq 1$ for all $n \in \mathbb{Z}_{\geq 1}$, if and only if $|r - 1| \leq (1/n)^{1/n}$ for all $n \in \mathbb{Z}_{\geq 1}$, if and only if

$$|r - 1| \leq \min_{n \in \mathbb{Z}_{\geq 1}} (1/n)^{1/n} = (1/3)^{1/3}. \quad \blacksquare$$

3 Generating functions

If $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, let

$$\begin{aligned} G_\phi(t) &= \sum_{n=0}^{\infty} \phi(n) t^n, \\ G_{*,\phi}(t) &= \sum_{n=0}^{\infty} \phi(n) \frac{t^n}{n!}, \end{aligned}$$

the ordinary and exponential generating functions, respectively, of $(\phi(n))_{n \geq 0}$. We also let

$$H_\phi(t) = G_{\widehat{\phi}}(-t) = \sum_{n=0}^{\infty} \widehat{\phi}(n) (-t)^n = \sum_{n=0}^{\infty} (-1)^n \phi(-1 - n) t^n.$$

Proposition 3.1 *Let $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ and $\psi = \mathcal{S}(\phi)$. Then*

$$\begin{aligned} G_{*,\phi}(t) &= \frac{G_{*,\psi}(t)}{1 - t} \\ \text{and } G'_{\widehat{\phi}}(t) + G_{\widehat{\phi}}(t) &= G_{\widehat{\psi}}(t). \end{aligned}$$

The second equality is equivalent to $H_\phi(t) - H'_\phi(t) = H_\psi(t)$.

Proof. Note that the equality $\mathcal{S}(\phi) = \psi$ means that $\phi(x) = x\phi(x - 1) + \psi(x)$ for all $x \in \mathbb{Z}_p$. Then

$$\begin{aligned} (1 - t)G_{*,\phi}(t) &= \sum_{n=0}^{\infty} \phi(n) \frac{t^n}{n!} - \sum_{n=0}^{\infty} \phi(n) \frac{t^{n+1}}{n!} \\ &= \sum_{n=0}^{\infty} \phi(n) \frac{t^n}{n!} - \sum_{n=1}^{\infty} \phi(n - 1) \frac{t^n}{(n - 1)!} \\ &= \sum_{n=0}^{\infty} (\phi(n) - n\phi(n - 1)) \frac{t^n}{n!} \end{aligned}$$

Paul Buckingham

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \psi(n) \frac{t^n}{n!} \\
&= G_{*,\psi}(t),
\end{aligned}$$

and

$$\begin{aligned}
G'_{\phi}(t) + G_{\phi}(t) &= \sum_{n=0}^{\infty} ((n+1)\widehat{\phi}(n+1) + \widehat{\phi}(n))t^n \\
&= \sum_{n=0}^{\infty} ((n+1)\phi(-2-n) + \phi(-1-n))t^n \\
&= \sum_{n=0}^{\infty} \psi(-1-n)t^n \\
&= G_{\widehat{\psi}}(t).
\end{aligned}$$

■

For example, let us apply the proposition to the functions $\gamma_r = \mathcal{S}^{-1}(g_r)$ appearing in Section 2.5.

Corollary 3.2 *If $r \in B_{<1}(1, \mathbb{C}_p)$, then*

$$\begin{aligned}
G_{*,\gamma_r}(t) &= \frac{\exp(rt)}{1-t} \\
\text{and } G'_{\widehat{f}_r}(t) + G_{\widehat{f}_r}(t) &= \frac{1}{r-t}.
\end{aligned}$$

Proof. The equalities follow immediately from the proposition and the straightforward observations that

$$G_{*,g_r}(t) = \exp(rt), \quad G_{\widehat{g}_r}(t) = \frac{1}{r-t}.$$

■

4 Self-adjointness of \mathcal{S}

Recall that a linear functional $\mu : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \mathbb{C}_p$ is said to be *bounded* if there is $C > 0$ such that $|\mu(\phi)| \leq C\|\phi\|$ for all $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$. In the following, we use the term *measure* to mean a bounded linear functional $\mu : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \mathbb{C}_p$. Our approach is similar to that of [6, Chap. 4, Sec. 1], except that measures there are assumed to have values in $\{x \in \mathbb{C}_p \mid |x| \leq 1\}$. We do not impose this restriction.

Paul Buckingham

If μ is a measure, then the power series $G(t) = \sum_{n=0}^{\infty} \mu(\beta_n) t^n$ has bounded coefficients. (Recall that $\beta_n(x) = \binom{x}{n}$.) Conversely, a power series $G(t) = \sum_{n=0}^{\infty} b_n t^n$ with bounded coefficients defines a measure μ satisfying

$$\mu(\phi) = \sum_{n=0}^{\infty} a_n b_n,$$

where $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ has Mahler coefficients $(a_n)_{n \geq 0}$. The measure μ associated to G in this way will be denoted $\mu(G)$.

If μ is a measure and $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, we denote $\mu(\phi)$ by $\int_{\mathbb{Z}_p} \phi d\mu$.

In the special case where $\phi : x \mapsto (1+z)^x$, where z is an element of \mathbb{C}_p with $|z| < 1$, we have

$$\int_{\mathbb{Z}_p} (1+z)^x d\mu(G)(x) = G(z). \quad (4.1)$$

This is immediate, since the function ϕ has n th Mahler coefficient $a_n = z^n$, so if $G(t) = \sum_{n=0}^{\infty} b_n t^n$, then

$$\sum_{n=0}^{\infty} a_n b_n = \sum_{n=0}^{\infty} z^n b_n = G(z).$$

We now define a \mathbb{C}_p -valued pairing on the \mathbb{C}_p -vector space $C(\mathbb{Z}_p, \mathbb{C}_p)$ by

$$\langle \phi, \psi \rangle = \int_{\mathbb{Z}_p} \phi d\mu(H_\psi).$$

The following lemma will be used several times.

Lemma 4.1 *Let $(a_k)_{k \geq 0}$ and $(b_l)_{l \geq 0}$ be sequences in \mathbb{C}_p that converge to zero. Then for any $\epsilon > 0$, there are $M, N \geq 0$ such that if either $k \geq M$ or $l \geq N$, then $|a_k b_l| \leq \epsilon$.*

Proof. Fix $\epsilon > 0$. First, choose $L \geq 0$ such that $|b_l| \leq 1$ for all $l \geq L$, possible because $b_l \rightarrow 0$ by assumption. Then, for each $l < L$, choose $M_l \geq 0$ such that $|a_k| |b_l| \leq \epsilon$ for all $k \geq M_l$, which is again possible because $a_k \rightarrow 0$. Next, choose \widetilde{M} such that $|a_k| \leq \epsilon$ for $k \geq \widetilde{M}$. Now let $M = \max\{\widetilde{M}, M_0, \dots, M_{L-1}\}$. Suppose that $k \geq M$, and let $l \geq 0$ be arbitrary. If $l < L$, then because $k \geq M \geq M_l$, we have $|a_k| |b_l| \leq \epsilon$. Otherwise, if $l \geq L$, then $|b_l| \leq 1$, while $|a_k| \leq \epsilon$ because $k \geq \widetilde{M}$, so $|a_k| |b_l| \leq \epsilon$.

In summary, M has the property that $|a_k b_l| \leq \epsilon$ whenever $k \geq M$, irrespective of the value of l . One may prove similarly the existence of an N such that $|a_k b_l| \leq \epsilon$ whenever $l \geq N$, irrespective of the value of k . ■

Proposition 4.2 *The pairing $\langle \cdot, \cdot \rangle$ is a non-degenerate symmetric bilinear form.*

Proof. Bilinearity is clear. For symmetry, we observe that if l is a non-negative integer, then $\binom{-1-x}{l} = (-1)^l \binom{x+l}{l}$. Now, if $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ have Mahler series

$$\phi(x) = \sum_{k=0}^{\infty} a_k \binom{x}{k}, \quad \psi(x) = \sum_{l=0}^{\infty} b_l \binom{x}{l},$$

then

$$H_{\psi}(t) = \sum_{k=0}^{\infty} (-1)^k \psi(-1-k) t^k = \sum_{k=0}^{\infty} (-1)^k \sum_{l=0}^{\infty} b_l \binom{-1-k}{l} t^k,$$

so

$$\langle \phi, \psi \rangle = \int_{\mathbb{Z}_p} \phi d\mu(H_{\psi}) \tag{4.2}$$

$$= \sum_{k=0}^{\infty} a_k (-1)^k \sum_{l=0}^{\infty} b_l \binom{-1-k}{l} \tag{4.3}$$

$$= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} (-1)^{k+l} a_k b_l \binom{k+l}{l} \tag{4.4}$$

$$= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} (-1)^{k+l} a_k b_l \binom{k+l}{k} \tag{4.5}$$

$$= \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} (-1)^{k+l} a_k b_l \binom{k+l}{k} \quad (\text{see below})$$

$$= \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} (-1)^{l+k} b_l a_k \binom{l+k}{k},$$

and this last expression is equal to $\langle \psi, \phi \rangle$ by a reversal of steps (4.2)–(4.4) with the roles of ϕ and ψ interchanged.

We show that the interchanging of the summation signs at (4.5) is permissible. By [9, Chap. 2, Sect. 1.2, Cor.], it is enough to show that, for any $\epsilon > 0$, there are only finitely many pairs (k, l) such that $|\eta(k, l)| > \epsilon$, where $\eta(k, l) = (-1)^{k+l} a_k b_l \binom{k+l}{k}$. Of course, $|\eta(k, l)| \leq |a_k| |b_l|$, so it is enough to show that there are $M, N \geq 0$ such that

- (i) for all $k \geq M$ and all $l \geq 0$, $|a_k| |b_l| \leq \epsilon$, and
- (ii) for all $k \geq 0$ and all $l \geq N$, $|a_k| |b_l| \leq \epsilon$.

For this, we simply invoke Lemma 4.1.

Finally, we show that $\langle \cdot, \cdot \rangle$ is non-degenerate. Fix $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, and suppose that $\langle \phi, \psi \rangle = 0$ for all $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$. If $\phi(x) = \binom{x}{n}$, then $\langle \phi, \psi \rangle$ is the coefficient of t^n in $H_{\psi}(t)$, which is $(-1)^n \psi(-1-n)$, so $n \geq 0$ being arbitrary, we see that $\psi(-1-n) = 0$ for all $n \geq 0$, and then $\psi = 0$ by continuity. \blacksquare

In fact, we have the following more symmetric version of the double sum for $\langle \phi, \psi \rangle$ appearing in the proof of Proposition 4.2. It is more symmetric because the order of summation does not privilege either direction, k or l , in the lattice of pairs (k, l) ; rather, it runs through the lattice diagonally.

Proposition 4.3 *Suppose that ϕ and ψ have Mahler coefficients $(a_k)_{k \geq 0}$ and $(b_l)_{l \geq 0}$ respectively. Then*

$$\langle \phi, \psi \rangle = \sum_{n=0}^{\infty} (-1)^n \sum_{k=0}^n \binom{n}{k} a_k b_{n-k}.$$

Proof. By (4.4), $\langle \phi, \psi \rangle = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \eta(k, l)$, where $\eta(k, l) = (-1)^{k+l} \binom{k+l}{l} a_k b_l$. Let $(\alpha_i)_{i \geq 0}$ be the sequence

$$\begin{aligned} &(\eta(0, 0), \\ &\eta(0, 1), \eta(1, 0), \\ &\eta(0, 2), \eta(1, 1), \eta(2, 0), \\ &\eta(0, 3), \eta(1, 2), \eta(2, 1), \eta(3, 0), \\ &\dots). \end{aligned}$$

We claim that this sequence converges to zero. Let $\epsilon > 0$. We again use Lemma 4.1 to obtain $M, N \geq 0$ such that if either $k \geq M$ or $l \geq N$, then $|a_k b_l| \leq \epsilon$. Hence, if $k+l \geq M+N$, then either $k \geq M$ or $l \geq N$, so $|\eta(k, l)| \leq \epsilon$. This establishes the claim that $\alpha_0, \alpha_1, \alpha_2, \dots$ converges to zero. But then [9, Chap. 2, Sect. 1.2, Prop.] implies that

$$\sum_{i=0}^{\infty} \alpha_i = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \eta(k, l),$$

i.e.,

$$\begin{aligned} \sum_{n=0}^{\infty} (-1)^n \sum_{k=0}^n \binom{n}{k} a_k b_{n-k} &= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \eta(k, l) \\ &= \langle \phi, \psi \rangle. \end{aligned}$$

■

Lemma 4.4 *If $G \in \mathbb{C}_p[[t]]$ has bounded coefficients, and if $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then*

$$\int_{\mathbb{Z}_p} \phi d\mu(G') = \int_{\mathbb{Z}_p} x\phi(x-1) d\mu(G)(x).$$

Proof. Suppose that

$$G(t) = \sum_{n=0}^{\infty} b_n t^n \quad \text{and} \quad \phi(x) = \sum_{n=0}^{\infty} a_n \binom{x}{n}.$$

Then $G'(t) = \sum_{n=0}^{\infty} (n+1)b_{n+1}t^n$, so

$$\int_{\mathbb{Z}_p} \phi d\mu(G') = \sum_{n=0}^{\infty} a_n(n+1)b_{n+1} = \sum_{n=1}^{\infty} na_{n-1}b_n. \quad (4.6)$$

But

$$\begin{aligned} \sum_{n=1}^{\infty} na_{n-1} \binom{x}{n} &= \sum_{n=1}^{\infty} a_{n-1}x \binom{x-1}{n-1} \quad (\text{a standard binomial identity}) \\ &= x \sum_{n=0}^{\infty} a_n \binom{x-1}{n} \\ &= x\phi(x-1), \end{aligned}$$

so (4.6) shows that $\int_{\mathbb{Z}_p} \phi d\mu(G') = \int_{\mathbb{Z}_p} x\phi(x-1) d\mu(G)(x)$. ■

Theorem 4.5 *The operator \mathcal{S} on $C(\mathbb{Z}_p, \mathbb{C}_p)$ is self-adjoint with respect to $\langle \cdot, \cdot \rangle$, that is,*

$$\langle \mathcal{S}(\phi), \psi \rangle = \langle \phi, \mathcal{S}(\psi) \rangle$$

for all $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$.

Proof.

$$\begin{aligned} \langle \mathcal{S}(\phi), \psi \rangle &= \int_{\mathbb{Z}_p} \mathcal{S}(\phi) d\mu(H_\psi) \\ &= \int_{\mathbb{Z}_p} (\phi(x) - x\phi(x-1)) d\mu(H_\psi)(x) \\ &= \int_{\mathbb{Z}_p} \phi(x) d\mu(H_\psi)(x) - \int_{\mathbb{Z}_p} x\phi(x-1) d\mu(H_\psi)(x) \\ &= \int_{\mathbb{Z}_p} \phi d\mu(H_\psi) - \int_{\mathbb{Z}_p} \phi d\mu(H'_\psi) \quad \text{by Lemma 4.4} \\ &= \int_{\mathbb{Z}_p} \phi d\mu(H_\psi - H'_\psi) \\ &= \int_{\mathbb{Z}_p} \phi d\mu(H_{\mathcal{S}(\psi)}) \quad \text{by Proposition 3.1} \\ &= \langle \phi, \mathcal{S}(\psi) \rangle. \end{aligned}$$

■

Corollary 4.6 *If $r, s \in B_{<1}(1, \mathbb{C}_p)$, then $\langle g_r, \gamma_s \rangle = \langle \gamma_r, g_s \rangle$, where the functions γ_r, g_r are as in Section 2.5.*

Paul Buckingham

Proof. This follows immediately from the fact that $\mathcal{S}(\gamma_r) = g_r$. ■

Corollary 4.7 *If $r, s \in B_{<1}(1, \mathbb{C}_p)$, and if $(\gamma_{r,n})_{n \geq 0}$ is the sequence of Mahler coefficients of γ_r , then*

$$\sum_{n=0}^{\infty} (1-r)^n \gamma_s(-1-n) = \sum_{n=0}^{\infty} (-1)^n \gamma_{r,n} s^{-1-n}.$$

Proof. On the one hand,

$$\begin{aligned} \langle g_r, \gamma_s \rangle &= \int_{\mathbb{Z}_p} r^x d\mu(H_{\gamma_s})(x) \\ &= H_{\gamma_s}(r-1) \quad \text{by (4.1)} \\ &= \sum_{n=0}^{\infty} (1-r)^n \gamma_s(-1-n). \end{aligned}$$

On the other hand,

$$\langle \gamma_r, g_s \rangle = \int_{\mathbb{Z}_p} \gamma_r d\mu(H_{g_s}) = \sum_{n=0}^{\infty} \gamma_{r,n} (-1)^n s^{-1-n}.$$

Now use Corollary 4.6. ■

More generally, we have the following.

Corollary 4.8 *If $r \in B_{<1}(1, \mathbb{C}_p)$ and $(\gamma_{r,n})_{n \geq 0}$ is the sequence of Mahler coefficients of γ_r , then for any $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$,*

$$\begin{aligned} \sum_{n=0}^{\infty} (-1)^n \mathcal{S}(\psi)(-1-n) \gamma_{r,n} &= H_{\psi}(r-1) \\ &= \sum_{n=0}^{\infty} \psi(-1-n) (1-r)^n. \end{aligned}$$

Proof. Taking $\phi = \gamma_r$ in the theorem gives $\langle g_r, \psi \rangle = \langle \gamma_r, \mathcal{S}(\psi) \rangle$. But

$$\langle g_r, \psi \rangle = \int_{\mathbb{Z}_p} r^x d\mu(H_{\psi})(x) = H_{\psi}(r-1)$$

by (4.1), and this is equal to $\sum_{n=0}^{\infty} \psi(-1-n) (1-r)^n$. On the other hand,

$$\langle \gamma_r, \mathcal{S}(\psi) \rangle = \int_{\mathbb{Z}_p} \gamma_r d\mu(H_{\mathcal{S}(\psi)}) = \sum_{n=0}^{\infty} (-1)^n \mathcal{S}(\psi)(-1-n) \gamma_{r,n}.$$

■

Example 4.9 If in Corollary 4.8 we take ψ to be the constant function $\mathbf{1}$, then $H_\psi(t) = 1/(1+t)$ and $\mathcal{S}(\psi)(x) = 1-x$, so we have

$$\sum_{n=0}^{\infty} (-1)^n (n+2) \gamma_{r,n} = \frac{1}{r}.$$

Note that the case $r = 1$ says that $\sum_{n=0}^{\infty} (-1)^n (n+2) \cdot n! = 1$, since $\gamma_1(x) = q(x) = \sum_{n=0}^{\infty} n! \binom{x}{n}$.

5 Convolution of Mahler coefficients

Let $\mathbb{C}_p[[t]]_0$ denote the set of power series $G(t) = \sum_{n=0}^{\infty} a_n \frac{t^n}{n!} \in \mathbb{C}_p[[t]]$ such that $a_n \rightarrow 0$.

Proposition 5.1 $\mathbb{C}_p[[t]]_0$ is a \mathbb{C}_p -subalgebra of $\mathbb{C}_p[[t]]$.

Proof. It is obvious that $\mathbb{C}_p[[t]]_0$ is a \mathbb{C}_p -subspace. Now suppose that $G(t) = \sum_{n=0}^{\infty} a_n \frac{t^n}{n!}$ and $H(t) = \sum_{n=0}^{\infty} b_n \frac{t^n}{n!}$, where $a_n \rightarrow 0$ and $b_n \rightarrow 0$. Then

$$G(t)H(t) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} a_k b_{n-k} \right) \frac{t^n}{n!}.$$

We show that the sequence of elements

$$\sum_{k=0}^n \binom{n}{k} a_k b_{n-k}$$

converges to zero as $n \rightarrow \infty$. Let $\epsilon > 0$. By Lemma 4.1 again, we may choose $M, N \geq 0$ such that if either $k \geq M$ or $l \geq N$, then $|a_k b_l| \leq \epsilon$. If $n \geq M + N$ and $0 \leq k \leq n$, then either $k \geq M$ or $n - k \geq N$, so $|a_k b_{n-k}| \leq \epsilon$, and then

$$\left| \sum_{k=0}^n \binom{n}{k} a_k b_{n-k} \right| \leq \epsilon.$$

■

Now consider the mutually inverse maps

$$\begin{aligned} C(\mathbb{Z}_p, \mathbb{C}_p) &\rightarrow \mathbb{C}_p[[t]]_0 \\ \phi &\mapsto \sum_{n=0}^{\infty} (\nabla^n \phi)(0) \frac{t^n}{n!}, \end{aligned} \tag{5.1}$$

$$\begin{aligned} \mathbb{C}_p[[t]]_0 &\rightarrow C(\mathbb{Z}_p, \mathbb{C}_p) \\ G &\mapsto \sum_{n=0}^{\infty} G^{(n)}(0) \beta_n. \end{aligned} \tag{5.2}$$

They are \mathbb{C}_p -linear maps but not algebra homomorphisms if $C(\mathbb{Z}_p, \mathbb{C}_p)$ has the usual product. We transfer the product on $\mathbb{C}_p[[t]]_0 \subseteq \mathbb{C}[[t]]$ to a new product \star on $C(\mathbb{Z}_p, \mathbb{C}_p)$ via the above maps. Concretely, if ϕ and ψ have Mahler coefficients $(a_k)_{k \geq 0}$ and $(b_l)_{l \geq 0}$ respectively, then

$$\phi \star \psi = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} a_k b_{n-k} \right) \beta_n, \quad (5.3)$$

so \star is a binomial convolution of the Mahler coefficients. Note that the identity element of $(C(\mathbb{Z}_p, \mathbb{C}_p), +, \star)$ is the constant function $\mathbf{1}$.

Remark. By [9, Chap. 4, Sect. 1.1, Comment (2)], the power series corresponding to a continuous function $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ via (5.1) and (5.2) is $\exp(-t)G_{\star, \phi}(t)$.

Note that, because $(\mathbb{C}_p[[t]]_0, +, \cdot)$ is a commutative \mathbb{C}_p -algebra with no zero divisors, the same is true of $(C(\mathbb{Z}_p, \mathbb{C}_p), +, \star)$.

We will also require the following notation. If $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ and $m \in \mathbb{Z}_{\geq 0}$, let

$$\phi^{\star m} = \underbrace{\phi \star \cdots \star \phi}_m. \quad (5.4)$$

If ϕ is a unit in $C(\mathbb{Z}_p, \mathbb{C}_p)$ with respect to the product operation \star , then for $m < 0$ we let

$$\phi^{\star m} = (\phi^{-1})^{\star(-m)}, \quad (5.5)$$

where ϕ^{-1} here denotes the inverse of ϕ with respect to \star .

Proposition 5.2 *If $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\langle \phi, \psi \rangle = (\phi \star \psi)(-1)$.*

Proof. Let the Mahler coefficients of ϕ and ψ be $(a_k)_{k \geq 0}$ and $(b_l)_{l \geq 0}$ respectively. Then by (5.3),

$$(\phi \star \psi)(-1) = \sum_{n=0}^{\infty} (-1)^n \sum_{k=0}^n \binom{n}{k} a_k b_{n-k},$$

because $\beta_n(-1) = \binom{-1}{n} = (-1)^n$. But this is equal to $\langle \phi, \psi \rangle$ by Proposition 4.3. \blacksquare

Proposition 5.3 *If $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\langle \phi, \psi \rangle = \int_{\mathbb{Z}_p} (\phi \star \psi) d\mu(H_{\mathbf{1}})$, where $\mathbf{1}$ is the function with constant value 1.*

Proof.

$$\begin{aligned} \langle \phi, \psi \rangle &= (\phi \star \psi)(-1) \quad \text{by Proposition 5.2} \\ &= ((\phi \star \psi) \star \mathbf{1})(-1) \\ &= \langle \phi \star \psi, \mathbf{1} \rangle \quad \text{by the same proposition} \end{aligned}$$

Paul Buckingham

$$= \int_{\mathbb{Z}_p} (\phi \star \psi) d\mu(H_1) \quad \text{by definition.}$$

■

Theorem 5.4 *If $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\mathcal{S}(\phi) \star \psi = \phi \star \mathcal{S}(\psi)$.*

Proof. Suppose that $\phi = \sum_{k=0}^{\infty} a_k \beta_k$ and $\psi = \sum_{l=0}^{\infty} b_l \beta_l$. On the one hand,

$$\mathcal{S}(\phi) \star \psi = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} (a_k - k a_{k-1}) b_{n-k} \right) \beta_n$$

by Proposition 2.2, while on the other hand

$$\phi \star \mathcal{S}(\psi) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} a_{n-k} (b_k - k b_{k-1}) \right) \beta_n.$$

Note that we have set $a_{-1} = b_{-1} = 0$. We have to show that

$$\sum_{k=0}^n \binom{n}{k} (a_k - k a_{k-1}) b_{n-k} = \sum_{k=0}^n \binom{n}{k} a_{n-k} (b_k - k b_{k-1}).$$

Since $\sum_{k=0}^n \binom{n}{k} a_k b_{n-k} = \sum_{k=0}^n \binom{n}{k} a_{n-k} b_k$, it remains to show that

$$\sum_{k=0}^n \binom{n}{k} k a_{k-1} b_{n-k} = \sum_{k=0}^n \binom{n}{k} k a_{n-k} b_{k-1}.$$

Now,

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k} k a_{k-1} b_{n-k} &= \sum_{k=0}^n \binom{n}{k} (n-k) a_{n-k-1} b_k \quad (k \mapsto n-k) \\ &= \sum_{k=0}^{n-1} \binom{n}{k} (n-k) a_{n-k-1} b_k \quad (\text{the } k=n \text{ term is } 0) \\ &= \sum_{k=1}^n \binom{n}{k-1} (n-k+1) a_{n-k} b_{k-1} \quad (\text{reindexing}) \\ &= \sum_{k=1}^n \binom{n}{k} k a_{n-k} b_{k-1} \quad (\text{a binomial identity}) \\ &= \sum_{k=0}^n \binom{n}{k} k a_{n-k} b_{k-1} \quad (\text{the } k=0 \text{ term is } 0), \end{aligned}$$

as desired. ■

Corollary 5.5 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then*

$$\begin{aligned} \mathcal{S}(\psi) &= (\mathbf{1} - \beta_1) \star \psi, \\ \text{and } \mathcal{S}^{-1}(\psi) &= q \star \psi. \end{aligned}$$

Therefore, $\mathcal{S}^m(\psi) = (\mathbf{1} - \beta_1)^{\star m} \star \psi$ for all $m \in \mathbb{Z}$.

Proof.

$$\begin{aligned} \mathcal{S}(\psi) &= \mathbf{1} \star \mathcal{S}(\psi) = \mathcal{S}(\mathbf{1}) \star \psi = (\mathbf{1} - \beta_1) \star \psi, \\ \text{and } \mathcal{S}^{-1}(\psi) &= \mathbf{1} \star \mathcal{S}^{-1}(\psi) = \mathcal{S}^{-1}(\mathbf{1}) \star \psi = q \star \psi. \end{aligned}$$

For the last assertion, observe that $q \star (\mathbf{1} - \beta_1) = \mathcal{S}^{-1}(\mathbf{1}) \star \mathcal{S}(\mathbf{1}) = \mathbf{1}$. ■

For example, if $r \in B_{<1}(1, \mathbb{C}_p)$, then

$$\gamma_r = \mathcal{S}^{-1}(g_r) = q \star g_r = \gamma_1 \star g_r.$$

Corollary 5.6 *Under the bijections in (5.1) and (5.2), the map $\mathcal{S} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow C(\mathbb{Z}_p, \mathbb{C}_p)$ corresponds to the map*

$$\begin{aligned} \mathbb{C}_p[[t]]_0 &\rightarrow \mathbb{C}_p[[t]]_0 \\ G(t) &\mapsto (1-t)G(t). \end{aligned}$$

Proof. The function $\mathbf{1} - \beta_1 \in C(\mathbb{Z}_p, \mathbb{C}_p)$ corresponds to the element $1 - t \in \mathbb{C}_p[[t]]_0$. Now use Corollary 5.5. ■

Corollary 5.7 *A function $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ is a unit with respect to \star if and only if $\mathcal{S}(\phi)$ is. Thus, \mathcal{S} restricts to a permutation of the unit group of the algebra $(C(\mathbb{Z}_p, \mathbb{C}_p), +, \star)$.*

Proof. If $\mathcal{S}(\phi) \star \psi = \mathbf{1}$, then $\phi \star \mathcal{S}(\psi) = \mathbf{1}$. Conversely, if $\phi \star \psi = \mathbf{1}$, then $\mathcal{S}(\phi) \star \mathcal{S}^{-1}(\psi) = \mathbf{1}$. ■

Corollary 5.8 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ and $\mathcal{S}^m(\psi) = \psi$ for some non-zero $m \in \mathbb{Z}$, then $\psi = 0$.*

Proof. By Corollary 5.5, the equality $\mathcal{S}^m(\psi) = \psi$ says that $(\mathbf{1} - \beta_1)^{\star m} \star \psi = \psi$, i.e., $((\mathbf{1} - \beta_1)^{\star m} - \mathbf{1}) \star \psi = 0$. But $\mathbf{1} - \beta_1$ corresponds to $1 - t \in \mathbb{C}_p[[t]]_0$, which has infinite multiplicative order, so because $m \neq 0$, $(\mathbf{1} - \beta_1)^{\star m} - \mathbf{1} \neq 0$. Hence, because $(C(\mathbb{Z}_p, \mathbb{C}_p), +, \star)$ has no zero divisors, $\psi = 0$. ■

Remark. Note that Theorem 5.4 and Proposition 5.2 combine to give an alternative proof of Theorem 4.5, the self-adjointness of \mathcal{S} with respect to $\langle \cdot, \cdot \rangle$.

6 An integral transform

Recall the notation $\phi^{\star m}$ defined in (5.4) and (5.5) for $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ and $m \in \mathbb{Z}$. Given $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, we may consider the function

$$\begin{aligned} \mathbb{Z} &\rightarrow \mathbb{C}_p \\ m &\mapsto \int_{\mathbb{Z}_p} (\mathbf{1} - \beta_1)^{\star m} d\mu(H_\psi). \end{aligned} \quad (6.1)$$

We show that this function, defined initially on \mathbb{Z} , can be extended to a locally analytic function $\mathbb{Z}_p \rightarrow \mathbb{C}_p$.

6.1 Iterates of \mathcal{S}

Proposition 6.1 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, $m \in \mathbb{Z}_{\geq 0}$, and $x \in \mathbb{Z}_p$, then*

$$\mathcal{S}^m(\psi)(x) = \sum_{k=0}^m (-1)^k \binom{m}{k} (x)_k \psi(x - k).$$

Proof. We proceed by induction on m , the case $m = 0$ being vacuous. Let $m \geq 0$, and assume that the statement is true for that m , i.e.,

$$\mathcal{S}^m(\psi)(y) = \sum_{k=0}^m (-1)^k \binom{m}{k} (y)_k \psi(y - k) \quad \text{for all } y \in \mathbb{Z}_p.$$

Then

$$\begin{aligned} &\mathcal{S}^{m+1}(\psi)(x) \\ &= \mathcal{S}(\mathcal{S}^m(\psi))(x) \\ &= \mathcal{S}\left(y \mapsto \sum_{k=0}^m (-1)^k \binom{m}{k} (y)_k \psi(y - k)\right)(x) \\ &= \sum_{k=0}^m (-1)^k \binom{m}{k} (x)_k \psi(x - k) - x \sum_{k=0}^m (-1)^k \binom{m}{k} (x-1)_k \psi(x-1-k) \\ &= \sum_{k=0}^m (-1)^k \binom{m}{k} (x)_k \psi(x - k) + \sum_{k=0}^m (-1)^{k+1} \binom{m}{k} (x)_{k+1} \psi(x - (k+1)) \\ &= \sum_{k=0}^m (-1)^k \binom{m}{k} (x)_k \psi(x - k) + \sum_{k=1}^{m+1} (-1)^k \binom{m}{k-1} (x)_k \psi(x - k) \\ &= \psi(x) + \sum_{k=1}^m (-1)^k \left(\binom{m}{k} + \binom{m}{k-1} \right) (x)_k \psi(x - k) \\ &\quad + (-1)^{m+1} (x)_{m+1} \psi(x - (m+1)) \\ &= \sum_{k=0}^{m+1} (-1)^k \binom{m+1}{k} (x)_k \psi(x - k), \end{aligned}$$

where in the last step we have used a standard binomial identity and reincorporated the $k = 0$ and $k = m + 1$ terms. The induction is complete. ■

Corollary 6.2 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ and $m \in \mathbb{Z}_{\geq 0}$, then*

$$\mathcal{S}^m(\psi)(-1) = \sum_{k=0}^m (m)_k \psi(-1 - k).$$

Proof. Take $x = -1$ in Proposition 6.1 and use the fact that $(-1)_k = (-1)^k k!$. ■

Example 6.3 Let $r \in B_{<1}(1, \mathbb{C}_p)$. If we take $\psi = \gamma_r$ in Corollary 6.2, then we obtain

$$\sum_{k=0}^m (m)_k \gamma_r(-1 - k) = \sum_{k=0}^{m-1} (m-1)_k r^{-1-k}$$

for all $m \in \mathbb{Z}_{\geq 1}$. Indeed, $\mathcal{S}^m(\gamma_r) = \mathcal{S}^{m-1}(g_r)$, and $g_r(x) = r^x$.

Specializing to $r = 1$ gives

$$\sum_{k=0}^m (m)_k q(-1 - k) = q(m-1) \quad (m \geq 1),$$

because $q(x) = \sum_{n=0}^{\infty} (x)_n$.

If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\psi(\mathbb{Z}_p)$ is bounded, so the sequence $(n! \psi(-1 - n))_{n \geq 0}$ converges to zero, and so we have a well-defined, continuous function

$$\begin{aligned} \mathcal{I}(\psi) : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ s &\mapsto \sum_{n=0}^{\infty} n! \psi(-1 - n) \binom{s}{n}, \end{aligned} \quad (6.2)$$

that is, $\mathcal{I}(\psi) = \sum_{n=0}^{\infty} n! \psi(-1 - n) \beta_n$.

Proposition 6.4 *Let $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ be continuous. Then $\mathcal{I}(\psi)$ is locally analytic.*

Proof. We apply Proposition 2.6 (in the a'_n notation). By the boundedness of $\psi(\mathbb{Z}_p)$, we may choose $C \in \mathbb{R}$ such that $v_p(\psi(-1 - n)) \geq C$ for all $n \geq 0$. Of course, there exist $\beta > -1/(p-1)$ and $N \geq 1$ such that $C/n \geq \beta$ for all $n \geq N$. Hence, for such n ,

$$\frac{v_p(\psi(-1 - n))}{n} \geq \frac{C}{n} \geq \beta,$$

and we are done. ■

Remark. The map $\mathcal{I} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$ is injective. Indeed, $\mathcal{I}(\psi)$ determines its own Mahler coefficients and therefore the values $\psi(-1-n)$ for all $n \geq 0$. By continuity, these values determine the function ψ .

Theorem 6.5 *Let $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$. The function*

$$\begin{aligned} \mathbb{Z} &\rightarrow \mathbb{C}_p \\ m &\mapsto \mathcal{S}^m(\psi)(-1) \end{aligned}$$

has a unique extension to a continuous function $\mathbb{Z}_p \rightarrow \mathbb{C}_p$. This continuous extension is in fact $\mathcal{I}(\psi)$ and is therefore locally analytic on \mathbb{Z}_p .

Proof. Uniqueness is clear by continuity, so we turn immediately to existence. By Corollary 6.2, if $m \in \mathbb{Z}_{\geq 0}$, then

$$\mathcal{S}^m(\psi)(-1) = \sum_{n=0}^m n! \psi(-1-n) \binom{m}{n} = \sum_{n=0}^{\infty} n! \psi(-1-n) \binom{m}{n} = \mathcal{I}(\psi)(m).$$

In summary,

$$\mathcal{S}^m(\psi)(-1) = \mathcal{I}(\psi)(m) \quad \text{for all } \psi \in C(\mathbb{Z}_p, \mathbb{C}_p) \text{ and all } m \in \mathbb{Z}_{\geq 0}. \quad (6.3)$$

It remains to show that the equality $\mathcal{S}^m(\psi)(-1) = \mathcal{I}(\psi)(m)$ holds for $m \in \mathbb{Z}_{<0}$ as well.

For the time being, we stay with $m \geq 0$, and in fact consider the situation where $m \geq 1$. In that case,

$$\begin{aligned} \mathcal{S}^m(\psi)(-1) &= \mathcal{S}^{m-1}(\mathcal{S}(\psi))(-1), \\ \text{i.e., } \mathcal{I}(\psi)(m) &= \mathcal{I}(\mathcal{S}(\psi))(m-1) \end{aligned}$$

by (6.3) applied to both (ψ, m) and $(\mathcal{S}(\psi), m-1)$. This being true for all $m \in \mathbb{Z}_{\geq 1}$, it follows by the continuity of $\mathcal{I}(\psi)$ and $\mathcal{I}(\mathcal{S}(\psi))$ that

$$\mathcal{I}(\psi)(s) = \mathcal{I}(\mathcal{S}(\psi))(s-1) \quad \text{for all } \psi \in C(\mathbb{Z}_p, \mathbb{C}_p) \text{ and } s \in \mathbb{Z}_p. \quad (6.4)$$

Applying (6.4) with ψ replaced by $\mathcal{S}^{-1}(\psi)$, we obtain

$$\mathcal{I}(\mathcal{S}^{-1}(\psi))(s) = \mathcal{I}(\psi)(s-1) \quad \text{for all } \psi \in C(\mathbb{Z}_p, \mathbb{C}_p) \text{ and } s \in \mathbb{Z}_p. \quad (6.5)$$

Iterating (6.5), i.e., invoking induction, we arrive at

$$\mathcal{I}(\mathcal{S}^{-m}(\psi))(s) = \mathcal{I}(\psi)(s-m) \quad \text{for all } \psi \in C(\mathbb{Z}_p, \mathbb{C}_p), m \in \mathbb{Z}_{\geq 0}, \text{ and } s \in \mathbb{Z}_p. \quad (6.6)$$

Finally, taking $s = 0$ in (6.6) gives $\mathcal{I}(\mathcal{S}^{-m}(\psi))(0) = \mathcal{I}(\psi)(-m)$, and since $\mathcal{I}(\mathcal{S}^{-m}(\psi))(0) = \mathcal{S}^{-m}(\psi)(-1)$ by definition, we have

$$\mathcal{S}^{-m}(\psi)(-1) = \mathcal{I}(\psi)(-m) \quad (m \geq 0),$$

which provides the desired $m < 0$ case of (6.3). ■

Corollary 6.6 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, $m \in \mathbb{Z}$, and $s \in \mathbb{Z}_p$, then*

$$\mathcal{I}(\mathcal{S}^m(\psi))(s) = \mathcal{I}(\psi)(s + m).$$

Proof. This is a corollary of the proof of the theorem. The case $m \leq 0$ is provided by (6.6). Now let $m \in \mathbb{Z}_{\geq 0}$, and apply (6.6) with ψ replaced by $\mathcal{S}^m(\psi)$. This gives

$$\mathcal{I}(\psi)(s) = \mathcal{I}(\mathcal{S}^m(\psi))(s - m) \quad \text{for all } s \in \mathbb{Z}_p.$$

Replacing s by $s + m$ gives the desired equality. ■

For a continuous function $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$, let $\Sigma\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ be its indefinite sum function, i.e., the unique continuous function with the property that

$$(\Sigma\psi)(n) = \sum_{k=0}^{n-1} \psi(k)$$

for all $n \in \mathbb{Z}_{\geq 0}$. (See [9, Chap. 4, Sect. 1.5] and [9, Chap. 4, Sect. 2.5].) Note that $(\Sigma\psi)(0) = 0$ and that $\nabla\Sigma\psi = \psi$.

Proposition 6.7 *If $\Psi \in \text{Im}(\mathcal{I})$, then $\mathcal{S}^{-1}(\Psi) \in \text{Im}(\mathcal{I})$. Specifically,*

$$\mathcal{S}^{-1}(\mathcal{I}(\psi)) = \mathcal{I}(-\Sigma\psi)$$

for all $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$.

Proof. We show that $\mathcal{S}(\mathcal{I}(\Sigma\psi)) = \mathcal{I}(-\psi)$, which is an equivalent assertion. Let $\phi = \Sigma\psi$. Then because $\phi(0) = 0$, Proposition 2.2 shows that

$$\begin{aligned} \mathcal{S}(\mathcal{I}(\Sigma\psi)) &= \sum_{n=0}^{\infty} (n! \phi(-1-n) - n! \phi(-n)) \beta_n \\ &= \sum_{n=0}^{\infty} n! (\phi(-1-n) - \phi(-n)) \beta_n \\ &= \sum_{n=0}^{\infty} n! (-\nabla\phi)(-1-n) \beta_n \\ &= \sum_{n=0}^{\infty} n! (-\psi)(-1-n) \beta_n \\ &= \mathcal{I}(-\psi). \end{aligned}$$

■

6.2 Manifesting \mathcal{I} as an integral transform

We now turn to our claim that the function $m \mapsto \int_{\mathbb{Z}_p} (\mathbf{1} - \beta_1)^{*m} d\mu(H_\psi)$ in (6.1) can be extended to a locally analytic function on \mathbb{Z}_p , showing at the same time that this extension is simply $\mathcal{I}(\psi)$. We hence derive an integral-transform expression for p -adic incomplete Γ -functions.

Define

$$\begin{aligned} T : \mathbb{Z}_p &\rightarrow C(\mathbb{Z}_p, \mathbb{C}_p) \\ s &\mapsto \sum_{n=0}^{\infty} (-1)^n \binom{s}{n} n! \beta_n, \end{aligned}$$

where, as always, $\beta_n(x) = \binom{x}{n}$. Note that, for any fixed $s \in \mathbb{Z}_p$, the function $\sum_{n=0}^{\infty} (-1)^n \binom{s}{n} n! \beta_n$ is indeed continuous, because its sequence of Mahler coefficients tends to zero.

Let $\|\cdot\|_M$ denote the norm on $C(\mathbb{Z}_p, \mathbb{C}_p)$ defined by $\|\phi\|_M = \max_{n \geq 0} |a_n|$, where $(a_n)_{n \geq 0}$ is the sequence of Mahler coefficients of ϕ .

Proposition 6.8 *The map T is uniformly continuous with respect to $\|\cdot\|_M$.*

Proof. Let $s_1, s_2 \in \mathbb{Z}_p$. Then

$$T(s_1) - T(s_2) = \sum_{n=1}^{\infty} (-1)^n \left(\binom{s_1}{n} - \binom{s_2}{n} \right) n! \beta_n,$$

and for $n \geq 1$,

$$\left| (-1)^n \left(\binom{s_1}{n} - \binom{s_2}{n} \right) n! \right| \leq |s_1 - s_2| \cdot n |n!| \quad (6.7)$$

by the lemma in [9, Chap. 5, Sect. 1.5]. (Actually, we have used only a weak form of the lemma, given in the remark that follows its proof.) An analysis of $n|n!|$ via the well-known fact that $v_p(n!) = (n - S_p(n))/(p - 1)$, where $S_p(n)$ is the p -adic digit sum of n , shows that $n|n!| \rightarrow 0$ as $n \rightarrow \infty$, so $n|n!|$ has a maximum M .

Now, let $\epsilon > 0$, and let $\delta = \epsilon/M$. If $|s_1 - s_2| < \delta$, then (6.7) shows that every Mahler coefficient of $T(s_1) - T(s_2)$ has absolute value at most $|s_1 - s_2| M < \delta M = \epsilon$, so $\|T(s_1) - T(s_2)\|_M < \epsilon$. Thus, T is uniformly continuous. \blacksquare

Proposition 6.9 *If $m \in \mathbb{Z}$, then $T(m) = (\mathbf{1} - \beta_1)^{*m}$.*

Proof. The continuous function $\mathbf{1} - \beta_1$ corresponds to the power series $1 - t$ under the bijections in (5.1) and (5.2), and if $m \geq 0$, then

$$(1 - t)^m = \sum_{n=0}^m \binom{m}{n} (-t)^n = \sum_{n=0}^{\infty} (-1)^n \binom{m}{n} n! \frac{t^n}{n!},$$

which corresponds to $T(m)$ under the same correspondence. For $m < 0$, we instead consider $T(-m)$ where $m > 0$. Observe that $(\mathbf{1} - \beta_1)^{\star(-m)}$ corresponds to

$$\begin{aligned} (1-t)^{-m} &= \sum_{n=0}^{\infty} (-1)^n \binom{-m}{n} t^n \quad (\text{see [9, Chap. 6, Sect. 3.2], for example}) \\ &= \sum_{n=0}^{\infty} (-1)^n \binom{-m}{n} n! \frac{t^n}{n!}, \end{aligned}$$

and this corresponds to $T(-m)$, as desired. ■

If $s \in \mathbb{Z}_p$, we define $(\mathbf{1} - \beta_1)^{\star s}$ to be $T(s)$. By Proposition 6.9, this definition extends $(\mathbf{1} - \beta_1)^{\star s}$ from $s \in \mathbb{Z}$ to $s \in \mathbb{Z}_p$.

Theorem 6.10 *Let $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, and let $\mathcal{I}(\psi)$ be the locally analytic function defined in (6.2). Then for all $s \in \mathbb{Z}_p$,*

$$\mathcal{I}(\psi)(s) = \int_{\mathbb{Z}_p} (\mathbf{1} - \beta_1)^{\star s} d\mu(H_\psi) \tag{6.8}$$

$$= \int_{\mathbb{Z}_p} \left((\mathbf{1} - \beta_1)^{\star s} \star \psi \right) d\mu(H_1). \tag{6.9}$$

Proof. Consider the function

$$\begin{aligned} \text{Int}(\psi) : \mathbb{Z}_p &\rightarrow \mathbb{C}_p \\ s &\mapsto \int_{\mathbb{Z}_p} (\mathbf{1} - \beta_1)^{\star s} d\mu(H_\psi). \end{aligned}$$

We first show that $\text{Int}(\psi)$ is continuous. Write $H_\psi(t) = \sum_{n=0}^{\infty} b_n t^n$, and let $N = \sup_{n \geq 0} |b_n|$. Note that the supremum is finite because ψ is bounded. If $\epsilon > 0$, then by Proposition 6.8, there is $\delta > 0$ such that $\|T(s_1) - T(s_2)\|_M < \epsilon/N$ whenever $|s_1 - s_2| < \delta$. Take such s_1, s_2 , and write $T(s_1) - T(s_2) = \sum_{n=0}^{\infty} a_n \beta_n$, where $a_n \rightarrow 0$. Then

$$\begin{aligned} \left| \int_{\mathbb{Z}_p} T(s_1) d\mu(H_\psi) - \int_{\mathbb{Z}_p} T(s_2) d\mu(H_\psi) \right| &= \left| \int_{\mathbb{Z}_p} (T(s_1) - T(s_2)) d\mu(H_\psi) \right| \\ &= \left| \sum_{n=0}^{\infty} a_n b_n \right| \\ &\leq \left(\max_{n \geq 0} |a_n| \right) \left(\sup_{n \geq 0} |b_n| \right) \\ &= \|T(s_1) - T(s_2)\|_M \cdot N \\ &< \epsilon. \end{aligned}$$

This proves continuity.

Now, for all $m \in \mathbb{Z}$,

$$\begin{aligned}
 \mathcal{I}(\psi)(m) &= \mathcal{S}^m(\psi)(-1) \quad \text{by Theorem 6.5} \\
 &= ((\mathbf{1} - \beta_1)^{\star m} \star \psi)(-1) \quad \text{by Corollary 5.5} \\
 &= \langle (\mathbf{1} - \beta_1)^{\star m}, \psi \rangle \quad \text{by Proposition 5.2} \\
 &= \int_{\mathbb{Z}_p} (\mathbf{1} - \beta_1)^{\star m} d\mu(H_\psi) \quad \text{by definition} \\
 &= \text{Int}(\psi)(m),
 \end{aligned}$$

so because both $\mathcal{I}(\psi)$ and $\text{Int}(\psi)$ are continuous, they are equal as functions on \mathbb{Z}_p . We thus have (6.8), and then (6.9) follows by Proposition 5.3. \blacksquare

Remark. We observe a similarity between the integral transform

$$\mathcal{I}(\psi)(s) = \int_{\mathbb{Z}_p} \left((\mathbf{1} - \beta_1)^{\star s} \star \psi \right) d\mu(H_1)$$

and the Laplace transform

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt = \int_0^\infty (e^{-t})^s f(t) dt.$$

The equation $\mathcal{I}(\Sigma\psi) = -\mathcal{S}^{-1}(\mathcal{I}(\psi))$ satisfied by the integral transform \mathcal{I} (see Proposition 6.7) is akin to the equation $\mathcal{L}\{tf(t)\} = -\mathcal{L}\{f\}'$ satisfied by the Laplace transform. In our situation, the derivative operator is replaced by \mathcal{S}^{-1} , and the multiplication-by- t map is replaced by the Σ operator. We will also see in Section 6.4 an analogue of the Laplace-transform formula $\mathcal{L}\{f * g\} = \mathcal{L}\{f\}\mathcal{L}\{g\}$.

6.2.1 $\Gamma_p(s, r)$ in terms of the integral transform \mathcal{I}

We show that O'Desky and Richman's p -adic incomplete Γ -functions can be expressed in terms of the integral transform \mathcal{I} . We recall the functions $\gamma_r = \mathcal{S}^{-1}(g_r)$ and $\bar{\gamma}_r$ defined in Section 2.5, where $r \in B_{<1}(1, \mathbb{C}_p)$.

Proposition 6.11 *Let $r \in B_{<1}(1, \mathbb{C}_p)$. Then $\gamma_r = \sigma(g_r)\mathcal{I}(g_r)$.*

Proof. If $s \in \mathbb{Z}_p$,

$$\begin{aligned}
 \gamma_r(s) &= \sum_{n=0}^{\infty} (s)_n r^{s-n} \quad \text{by Proposition 2.8} \\
 &= r^{s+1} \sum_{n=0}^{\infty} (s)_n r^{-1-n} \\
 &= \sigma(g_r)(s)\mathcal{I}(g_r)(s) \quad \text{by definition of } \mathcal{I}.
 \end{aligned}$$

\blacksquare

Proposition 6.12 *Let $r \in B_{<1}(1, \mathbb{C}_p)$, and define $\Gamma_p(-, r) = \bar{\gamma}_r$, where $\bar{\gamma}_r$ is as in Section 2.5. Then for all $s \in \mathbb{Z}_p$,*

$$\Gamma_p(s, r) = \exp(r\tilde{p})r^s \int_{\mathbb{Z}_p} \left((\mathbf{1} - \beta_1)^{\star(s-1)} \star g_r \right) d\mu(H_1).$$

Proof. If $s \in \mathbb{Z}_p$, then

$$\begin{aligned} \bar{\gamma}_r(s) &= \exp(r\tilde{p})\gamma_r(s-1) \\ &= \exp(r\tilde{p})r^s \mathcal{S}(g_r)(s-1) \quad \text{by Proposition 6.11} \\ &= \exp(r\tilde{p})r^s \int_{\mathbb{Z}_p} \left((\mathbf{1} - \beta_1)^{\star(s-1)} \star g_r \right) d\mu(H_1) \quad \text{by Theorem 6.10.} \end{aligned}$$

■

We compare the above formula with the definition of the complex incomplete Γ -function:

$$\Gamma(s, r) = \int_r^\infty x^{s-1} e^{-x} dx.$$

6.3 Image of \mathcal{S}

We determine which functions are in the image $\text{Im}(\mathcal{S})$ of the injective map $\mathcal{S} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$. The following proposition will help in the task.

Proposition 6.13 *If $(\psi_k)_{k \geq 0}$ is a sequence of functions in $C(\mathbb{Z}_p, \mathbb{C}_p)$ converging uniformly to some ψ , then the sequence of functions $\mathcal{S}(\psi_k)$ converges uniformly to $\mathcal{S}(\psi)$.*

Proof. We let $\|\cdot\|$ denote the sup-norm for functions $\mathbb{Z}_p \rightarrow \mathbb{C}_p$. Let $\epsilon > 0$, and choose $N \geq 0$ such that $\|\psi_k - \psi\| < \epsilon$ for all $k \geq N$. Then for the same integers k ,

$$\begin{aligned} \|\mathcal{S}(\psi_k) - \mathcal{S}(\psi)\| &= \left\| \sum_{n=0}^{\infty} n! \psi_k(-1-n)\beta_n - \sum_{n=0}^{\infty} n! \psi(-1-n)\beta_n \right\| \\ &= \left\| \sum_{n=0}^{\infty} n! (\psi_k - \psi)(-1-n)\beta_n \right\| \\ &\leq \|\psi_k - \psi\| \quad \text{because } \|\beta_n\| = 1 \\ &< \epsilon. \end{aligned}$$

■

Recall that $\beta_1(x) = x$, and that $\sigma(\psi)(x) = \psi(x+1)$.

Lemma 6.14 *If $\psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\mathcal{S}(\beta_1\sigma^{-1}(\psi)) = -\nabla(\mathcal{S}(\psi))$.*

Paul Buckingham

Proof. We begin with the definition of $\mathcal{S}(\psi)$, namely, $\mathcal{S}(\psi)(x) = \psi(x) - x\psi(x-1)$, i.e., $\mathcal{S}(\psi) = \psi - \beta_1\sigma^{-1}(\psi)$. Then

$$\begin{aligned}\mathcal{I}(\beta_1\sigma^{-1}(\psi)) &= \mathcal{I}(\psi - \mathcal{S}(\psi)) \\ &= \mathcal{I}(\psi) - \mathcal{I}(\mathcal{S}(\psi)) \\ &= \mathcal{I}(\psi) - \sigma(\mathcal{I}(\psi)) \quad \text{by Corollary 6.6} \\ &= -\nabla(\mathcal{I}(\psi)).\end{aligned}$$

■

Before giving the next proposition, we observe that the functions $\frac{1}{n!}\nabla^n q$ map \mathbb{Z}_p into itself. Indeed, $q = \sum_{k=0}^{\infty} k! \beta_k$, so

$$\frac{1}{n!}\nabla^n q = \frac{1}{n!} \sum_{k=0}^{\infty} (k+n)! \beta_k = \sum_{k=0}^{\infty} \frac{(k+n)!}{n!} \beta_k,$$

and $(n+k)!/n! \in \mathbb{Z}$.

Proposition 6.15 *If $\psi = \sum_{n=0}^{\infty} a_n \beta_n \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then*

$$\mathcal{I}(\psi) = \sum_{n=0}^{\infty} a_n \frac{(-1)^n}{n!} \nabla^n q.$$

Proof. The binomial identity $x^{\binom{x-1}{n}} = (n+1)\binom{x}{n+1}$ says that $\beta_1\sigma^{-1}(\beta_n) = (n+1)\beta_{n+1}$, so

$$\mathcal{I}(\beta_{n+1}) = -\frac{1}{n+1} \nabla(\mathcal{I}(\beta_n))$$

by Lemma 6.14. Therefore, because $\mathcal{I}(\beta_0) = q$ by definition (see (6.2)), induction on n shows that

$$\mathcal{I}(\beta_n) = \frac{(-1)^n}{n!} \nabla^n q.$$

Hence, Proposition 6.13 shows that if $a_n \rightarrow 0$, then

$$\mathcal{I}\left(\sum_{n=0}^{\infty} a_n \beta_n\right) = \sum_{n=0}^{\infty} a_n \frac{(-1)^n}{n!} \nabla^n q.$$

■

Corollary 6.16 *The image of the transform map $\mathcal{I} : C(\mathbb{Z}_p, \mathbb{C}_p) \rightarrow \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$ consists of the functions*

$$\Psi = \sum_{n=0}^{\infty} a_n \frac{(-1)^n}{n!} \nabla^n q \tag{6.10}$$

where $(a_n)_{n \geq 0}$ is a sequence in \mathbb{C}_p converging to zero. A function Ψ in the image determines the coefficients a_n in (6.10) uniquely.

Proof. The first assertion is clear by Proposition 6.15. As for the uniqueness of the coefficients for a given Ψ , if $\sum_{n=0}^{\infty} a_n \frac{(-1)^n}{n!} \nabla^n q = 0$ where $a_n \rightarrow 0$, let $\psi = \sum_{n=0}^{\infty} a_n \beta_n \in C(\mathbb{Z}_p, \mathbb{C}_p)$. Then

$$\mathcal{I}(\psi) = \sum_{n=0}^{\infty} a_n \frac{(-1)^n}{n!} \nabla^n q = 0,$$

so $\psi = 0$ by the injectivity of \mathcal{I} , and therefore the a_n are all zero. \blacksquare

Corollary 6.17 *For all $n \geq 0$, $\frac{1}{n!} \nabla^n q = \mathcal{S}^{-n}(q)$. Therefore, $\text{Im}(\mathcal{I})$ consists of the functions*

$$\Psi = \sum_{n=0}^{\infty} a_n (-1)^n \mathcal{S}^{-n}(q)$$

where $(a_n)_{n \geq 0}$ is a sequence in \mathbb{C}_p converging to zero. Again, the a_n are determined uniquely by Ψ .

Proof. We first recall Proposition 6.7, which, once iterated, says that $\mathcal{I}(\Sigma^n \psi) = (-1)^n \mathcal{S}^{-n}(\mathcal{I}(\psi))$. We apply this in the case where $\psi = \beta_0$. Then because $\beta_n = \Sigma^n \beta_0$, we have

$$\mathcal{I}(\beta_n) = \mathcal{I}(\Sigma^n \beta_0) = (-1)^n \mathcal{S}^{-n}(\mathcal{I}(\beta_0)) = (-1)^n \mathcal{S}^{-n}(q),$$

the last equality by Proposition 6.15. On the other hand, $\mathcal{I}(\beta_n) = (-1)^n \frac{1}{n!} \nabla^n q$ by Proposition 6.15 again. Therefore, $\mathcal{S}^{-n}(q) = \frac{1}{n!} \nabla^n q$, so the rest of the statement to be proven follows from Corollary 6.16. \blacksquare

We note in passing that

$$\mathcal{S}^{-1}\left(\frac{1}{n!} \nabla^n q\right) = \mathcal{S}^{-1}(\mathcal{S}^{-n}(q)) = \mathcal{S}^{-(n+1)}(q) = \frac{1}{(n+1)!} \nabla^{n+1}(q).$$

We make another observation regarding the image of \mathcal{I} .

Proposition 6.18 *If $\text{lc}(\mathbb{Z}_p, \mathbb{C}_p) \subset \text{la}(\mathbb{Z}_p, \mathbb{C}_p)$ denotes the subspace of locally constant functions, then $\text{Im}(\mathcal{I}) \cap \text{lc}(\mathbb{Z}_p, \mathbb{C}_p) = 0$.*

Proof. Let $\Psi = \mathcal{I}(\psi)$ be in the intersection. Being locally constant, Ψ is \mathbb{Z} -periodic of period $m = p^j$ for some $j \geq 0$ (see [9, Chap. 4, Sect. 3.2], for example). That is, $\Psi(s + m) = \Psi(s)$ for all $s \in \mathbb{Z}_p$. Then

$$\begin{aligned} \mathcal{I}(\mathcal{S}^m(\psi))(s) &= \mathcal{I}(\psi)(s + m) \quad \text{by Corollary 6.6} \\ &= \mathcal{I}(\psi)(s), \end{aligned}$$

so $\mathcal{I}(\mathcal{S}^m(\psi)) = \mathcal{I}(\psi)$. Hence, $\mathcal{S}^m(\psi) = \psi$ by the injectivity of \mathcal{I} , so $\psi = 0$ by Corollary 5.8. (Note that $m = p^j > 0$.) Thus, $\Psi = 0$. \blacksquare

6.4 \mathcal{I} and convolution

Recall the convolution product \star on $C(\mathbb{Z}_p, \mathbb{C}_p)$. We define a similar product on $\text{Im}(\mathcal{I})$, again via the algebra $\mathbb{C}_p[[t]]_0$ (see the beginning of Section 5). Consider the mutually inverse maps

$$\begin{aligned} \text{Im}(\mathcal{I}) &\rightarrow \mathbb{C}_p[[t]]_0 \\ \sum_{n=0}^{\infty} b_n \mathcal{S}^{-n}(q) &\mapsto \sum_{n=0}^{\infty} b_n \frac{t^n}{n!} \quad (\text{where } b_n \rightarrow 0 \text{ in } \mathbb{C}_p), \end{aligned} \quad (6.11)$$

$$\begin{aligned} \mathbb{C}_p[[t]]_0 &\rightarrow \text{Im}(\mathcal{I}) \\ \sum_{n=0}^{\infty} b_n \frac{t^n}{n!} &\mapsto \sum_{n=0}^{\infty} b_n \mathcal{S}^{-n}(q). \end{aligned} \quad (6.12)$$

These maps are well defined by Corollary 6.17. Denote by \diamond the product on $\text{Im}(\mathcal{I})$ inherited from $\mathbb{C}_p[[t]]_0$ via these maps. Concretely,

$$\left(\sum_{k=0}^{\infty} b_k \mathcal{S}^{-k}(q) \right) \diamond \left(\sum_{l=0}^{\infty} c_l \mathcal{S}^{-l}(q) \right) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} b_k c_{n-k} \right) \mathcal{S}^{-n}(q).$$

Then $(\text{Im}(\mathcal{I}), +, \diamond)$ is a commutative \mathbb{C}_p -algebra with no zero divisors.

Proposition 6.19 *If $\phi, \psi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\mathcal{I}(\phi \star \psi) = \mathcal{I}(\phi) \diamond \mathcal{I}(\psi)$.*

Proof. Let the Mahler coefficients of ϕ and ψ be $(b_k)_{k \geq 0}$ and $(c_l)_{l \geq 0}$ respectively. Then

$$\begin{aligned} \mathcal{I}(\phi \star \psi) &= \mathcal{I} \left(\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} b_k c_{n-k} \right) \beta_n \right) \\ &= \sum_{n=0}^{\infty} (-1)^n \left(\sum_{k=0}^n \binom{n}{k} b_k c_{n-k} \right) \mathcal{S}^{-n}(q) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} (-1)^k b_k (-1)^{n-k} c_{n-k} \right) \mathcal{S}^{-n}(q) \\ &= \left(\sum_{k=0}^{\infty} (-1)^k b_k \mathcal{S}^{-k}(q) \right) \diamond \left(\sum_{l=0}^{\infty} (-1)^l c_l \mathcal{S}^{-l}(q) \right) \\ &= \mathcal{I}(\phi) \diamond \mathcal{I}(\psi). \end{aligned}$$

■

Proposition 6.19 is akin to the convolution formula for the Laplace transform.

7 Connection to the equation $F' + F = G$

We describe the role of the automorphism \mathcal{S} in the solution to the well-known differential equation $F' + F = G$, where $G \in \mathbb{C}_p[[t]]$ is a given power series with bounded coefficients. Of course, our intention is not to attempt to contribute anything to the general theory of p -adic differential equations, which has a vast literature extending far beyond this equation, but simply to highlight a connection.

It is worth remarking that, trivial though the equation $F' + F = G$ appears to be, it is not at first sight obvious that if $G \in \mathbb{Z}_p[[t]]$, say, then the equation has a solution F that is also in $\mathbb{Z}_p[[t]]$ (which it does).

We have already seen that if $G(t) = \sum_{n=0}^{\infty} \psi(n)t^n \in \mathbb{C}_p[[t]]$ has coefficients given by a continuous function $\psi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$, then $F' + F = G$ where

$$F(t) = \sum_{n=0}^{\infty} \mathcal{S}^{-1}(\widehat{\psi})(-1-n)t^n,$$

$\widehat{\psi}$ being the function $x \mapsto \psi(-1-x)$. Just replace ψ by $\widehat{\psi}$ in Proposition 3.1. A similar fact holds even when the coefficients of G are not assumed to vary continuously with n and are instead assumed only to be bounded. We turn now to this generalization and show how the automorphism \mathcal{S} still appears.

Let K be a complete subfield of \mathbb{C}_p . For a positive real number M , we define $\mathcal{F}_M(\mathbb{Z}_p, K)$ to consist of the functions $\psi : \mathbb{Z}_p \rightarrow K$ having Mahler series of the form

$$\psi = \sum_{n=0}^{\infty} n! a_n \beta_n \quad \text{where } |a_n| \leq M \text{ for all } n \geq 0.$$

We also let

$$\mathcal{F}(\mathbb{Z}_p, K) = \bigcup_{M>0} \mathcal{F}_M(\mathbb{Z}_p, K).$$

Proposition 7.1 *For each $M > 0$, the \mathbb{C}_p -linear automorphisms \mathcal{S} and σ of $C(\mathbb{Z}_p, \mathbb{C}_p)$ restrict to K -linear automorphisms of $\mathcal{F}_M(\mathbb{Z}_p, K)$.*

Proof. Let $\psi = \sum_{n=0}^{\infty} n! a_n \beta_n$. Let us handle \mathcal{S} first. By Proposition 2.2,

$$\begin{aligned} \mathcal{S} \left(\sum_{n=0}^{\infty} n! a_n \beta_n \right) &= a_0 \beta_0 + \sum_{n=1}^{\infty} (n! a_n - n(n-1)! a_{n-1}) \beta_n \\ &= a_0 \beta_0 + \sum_{n=1}^{\infty} n! (a_n - a_{n-1}) \beta_n \\ &= \sum_{n=0}^{\infty} n! (a_n - a_{n-1}) \beta_n \end{aligned} \tag{7.1}$$

if we let $a_{-1} = 0$. It is now clear that if $\psi \in \mathcal{F}_M(\mathbb{Z}_p, K)$, then so is $\mathcal{S}(\psi)$. Similarly, by Theorem 2.4,

$$\begin{aligned} \mathcal{S}^{-1} \left(\sum_{n=0}^{\infty} n! a_n \beta_n \right) &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{n!}{k!} k! a_k \right) \beta_n \\ &= \sum_{n=0}^{\infty} n! \left(\sum_{k=0}^n a_k \right) \beta_n, \end{aligned}$$

so again, $\mathcal{S}^{-1}(\psi) \in \mathcal{F}_M(\mathbb{Z}_p, K)$ if ψ is.

As for σ , the easily verified equalities

$$\begin{aligned} \sigma \left(\sum_{n=0}^{\infty} n! a_n \beta_n \right) &= \sum_{n=0}^{\infty} n! ((n+1)a_{n+1} + a_n) \beta_n, \quad (7.2) \\ \sigma^{-1} \left(\sum_{n=0}^{\infty} n! a_n \beta_n \right) &= \sum_{n=0}^{\infty} n! \left(\sum_{k=n}^{\infty} (-1)^{k-n} \frac{k!}{n!} a_k \right) \beta_n \end{aligned}$$

show that if $\psi \in \mathcal{F}(\mathbb{Z}_p, K)$, then so are $\sigma(\psi)$ and $\sigma^{-1}(\psi)$. ■

Now define

$$\begin{aligned} \Lambda : \mathbb{C}_p[[t]] &\rightarrow \mathbb{C}_p[[t]] \\ F &\mapsto (1-t)(F' + F). \end{aligned}$$

Also, if $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$ with Mahler series $\phi = \sum_{n=0}^{\infty} n! a_n \beta_n$, define

$$J_\phi(t) = \sum_{n=0}^{\infty} (a_n - a_{n-1}) t^n,$$

where, for convenience, we have set $a_{-1} = 0$.

Proposition 7.2 *If $\phi \in C(\mathbb{Z}_p, \mathbb{C}_p)$, then $\Lambda(J_\phi) = J_{\sigma \circ \mathcal{S}(\phi)}$.*

Proof. Let $\phi = \sum_{n=0}^{\infty} n! a_n \beta_n$. One sees immediately from (7.1) and (7.2) that $\sigma \circ \mathcal{S}(\phi) = \sum_{n=0}^{\infty} n! b_n \beta_n$ where

$$b_n = (n+1)a_{n+1} - na_n - a_{n-1}.$$

Note that, if we define $a_{-2} = 0$, then $b_{-1} = 0$. (We have already taken a_{-1} to be 0.)

Now,

$$\begin{aligned} J'_\phi(t) + J_\phi(t) &= \sum_{n=0}^{\infty} (n+1)(a_{n+1} - a_n) t^n + \sum_{n=0}^{\infty} (a_n - a_{n-1}) t^n \\ &= \sum_{n=0}^{\infty} ((n+1)a_{n+1} - na_n - a_{n-1}) t^n \end{aligned}$$

Paul Buckingham

$$\begin{aligned}
&= \sum_{n=0}^{\infty} b_n t^n \quad \text{by the above} \\
&= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n (b_k - b_{k-1}) \right) t^n \quad (b_{-1} = 0) \\
&= \frac{1}{1-t} \sum_{n=0}^{\infty} (b_n - b_{n-1}) t^n \\
&= \frac{1}{1-t} J_{\sigma \circ \mathcal{S}(\phi)}(t).
\end{aligned}$$

■

For a real number $M > 0$, let $K[[t]]_{\leq M}$ denote the set of power series $G(t) = \sum_{n=0}^{\infty} b_n t^n$ with $|b_n| \leq M$ for all n , and let $K[[t]]_b = \bigcup_{M>0} K[[t]]_{\leq M}$, which consists of the power series with bounded coefficients.

Before stating the next proposition, we point out [4, Theorem 2.1], which considers a related differential equation in the situation where $A = \mathbb{Z}_p$, but without using the automorphism \mathcal{S} .

Proposition 7.3 *Let X be equal either to $K[[t]]_{\leq M}$ with $M > 0$, or to $K[[t]]_b$, and let $A = \{x \in K \mid |x| \leq 1\}$. Then*

$$\begin{aligned}
X &\rightarrow X \\
F &\mapsto F' + F
\end{aligned}$$

is an A -module automorphism of X . In particular, $F \mapsto F' + F$ is an automorphism of the A -module $A[[t]]$. If $G(t) = \sum_{n=0}^{\infty} b_n t^n \in X$ and $\psi = \sum_{n=0}^{\infty} n! b_n \beta_n$, then the unique $F \in X$ such that $F' + F = G$ is J_ϕ where $\phi = \mathcal{S}^{-1} \circ \sigma^{-1}(\psi)$.

Proof. Because $K[[t]]_b = \bigcup_{M>0} K[[t]]_{\leq M}$ and $A[[t]] = K[[t]]_{\leq 1}$, it is enough to prove everything in the case where $X = K[[t]]_{\leq M}$.

For injectivity, suppose that $F \in K[[t]]_{\leq M}$ satisfies $F' + F = 0$. Then $F(t) = c \exp(-t)$ for some $c \in K$, but $\exp(-t)$ does not have bounded coefficients, so $c = 0$.

For surjectivity, let $G \in K[[t]]_{\leq M}$, say $G(t) = \sum_{n=0}^{\infty} b_n t^n$. Then

$$(1-t)G(t) = \sum_{n=0}^{\infty} (b_n - b_{n-1}) t^n = J_\psi(t)$$

where $\psi = \sum_{n=0}^{\infty} n! b_n \beta_n$ (and $b_{-1} = 0$). Note that $\psi \in \mathcal{F}_M(\mathbb{Z}_p, K)$, because $|b_n| \leq M$ for all n . Now let $\phi = \mathcal{S}^{-1} \circ \sigma^{-1}(\psi)$, which is in $\mathcal{F}_M(\mathbb{Z}_p, K)$ by Proposition 7.1. Then $J_\phi \in K[[t]]_{\leq M}$, and Proposition 7.2 says that

$$\Lambda(J_\phi) = J_{\sigma \circ \mathcal{S}(\phi)} = J_\psi.$$

Hence, remembering that $J_\psi(t) = (1-t)G(t)$, we have $J'_\phi(t) + J_\phi(t) = G(t)$. ■

Acknowledgements

The author would like to thank Al Weiss and Neil Dummigan for helpful conversations.

References

- [1] Yvette Amice. Interpolation p -adique. *Bull. Soc. Math. France*, 92:117–180, 1964.
- [2] Bruce C. Berndt, Sun Kim, and Alexandru Zaharescu. Diophantine approximation of the exponential function and Sondow’s Conjecture. *Adv. Math.*, 248:1298–1331, 2013.
- [3] Pierre Colmez. Fonctions d’une variable p -adique. *Astérisque*, (330):13–59, 2010.
- [4] S. L. Gefter and A. B. Goncharuk. The Hurwitz product, p -adic topology on \mathbb{Z} , and fundamental solution to linear differential equation in the ring $\mathbb{Z}[[x]]$. *J. Math. Sci. (N.Y.)*, 228(6, Problems in mathematical analysis. No. 90 (Russian)):633–638, 2018.
- [5] Lorenz Halbeisen and Norbert Hungerbühler. Number theoretic aspects of a combinatorial function. *Notes Number Theory Discrete Math.*, 5(4):138–150, 1999.
- [6] S. Lang. *Cyclotomic fields I and II*, volume 121 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1990. With an appendix by Karl Rubin.
- [7] Andrew O’Desky and David Richman. Derangements and the p -adic incomplete gamma function. *To appear in Transactions of the AMS*.
- [8] Vicențiu Pașol and Alexandru Zaharescu. Sondow’s conjecture, convergents to e , and p -adic analytic functions. *Math. Z.*, 292(1-2):499–511, 2019.
- [9] A.M. Robert. *A Course in p -adic Analysis*. Springer, 2000.
- [10] Jonathan Sondow and Kyle Schalm. Which partial sums of the Taylor series for e are convergents to e ? (and a link to the primes 2, 5, 13, 37, 463). Part II. In *Gems in experimental mathematics*, volume 517 of *Contemp. Math.*, pages 349–363. Amer. Math. Soc., Providence, RI, 2010.
- [11] Lawrence C. Washington. *Introduction to cyclotomic fields*, volume 83 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1997.