

BICATEGORIES OF DIFFEOLOGICAL GROUPOIDS

JORDAN WATTS

ABSTRACT. We study three variants of bicategories of diffeological groupoids, and show that they are all equivalent: the bicategory with generalised morphisms, the bicategory with anafunctors, and the bicategory with bibundles. Moreover, we show that the corresponding sub-bicategories of locally subductive diffeological groupoids, as well as Lie groupoids, are (essentially) full sub-bicategories. As applications, we obtain equivalences with the corresponding 2-categories of geometric stacks, and present some Morita invariants.

CONTENTS

1.	Introduction	2
2.	Diffeological Preliminaries	6
3.	Diffeological Groupoids	9
4.	Weak Equivalences	11
5.	Pullbacks and Weak Pullbacks	17
6.	The Bicategory of Fractions	21
6.1.	Weak Equivalences Admit a Calculus of Fractions	22
6.2.	The Bicategory of Fractions $\mathbf{DGpoid}[W^{-1}]$	22
6.3.	Spanisation	27
7.	Anafunctors	29
7.1.	Subductive Weak Equivalences Induce a 2-Site	29
7.2.	The Anafunctor Bicategory $\mathbf{DGpoid}_{\text{ana}}$	31
7.3.	Anafunctorisation	36
7.4.	$\mathbf{DGpoid}[W^{-1}]$ versus $\mathbf{DGpoid}_{\text{ana}}$	38
7.5.	Optimising 2-Cells	43
8.	Bibundles	46
8.1.	The Bibundle Bicategory $\mathbf{DBiBund}$	46
8.2.	Bibundlisation	50
8.3.	$\mathbf{DGpoid}[W^{-1}]$ versus $\mathbf{DBiBund}$	52
8.4.	Optimising 1-cells	53
8.5.	Quasi-Inverses in the Bicategories	58
9.	Local Subductions & Lie Groupoids	59
9.1.	Locally Subductive Diffeological Groupoids	60
9.2.	Lie Groupoids	68
10.	Applications	74
10.1.	Of Sites and Stacks	74
10.2.	Invariants of Diffeological Groupoids	80
10.3.	Principal Bundles	83

1. INTRODUCTION

Diffeological groupoids and the application of diffeology to the study of Lie groupoids and Lie algebroids are current and important trends in geometry. They appear in work on general relativity [BFW13], in which a diffeological groupoid describes the choices of embeddings of an initial space-like hypersurface in a lorentzian spacetime, up to a prescribed equivalence. They show up in the study of singular subalgebroids of Lie algebroids [AZ20], and related, the holonomy and fundamental groupoids of a singular foliation [GV21], both of which are connected to solutions to Lie’s third problem for Lie algebroids. Diffeological groupoids have become crucial in the study of the “higher geometric” version of loop spaces: loop stacks; in [RV18a], the authors show that the stack $\text{Hom}(\mathbb{S}^1, \mathcal{X})$ is presentable by not just a diffeological groupoid, but a Fréchet-Lie groupoid, where \mathcal{X} is a differentiable stack. As Fréchet manifolds have been shown to form a full subcategory of the category of diffeological spaces [Los94, Section 3], here is an example (of many) where working in the diffeological category is very beneficial to infinite-dimensional differential geometry. In [RV18b], the authors announce that they have extended the results of [RV18a] to stacks $\text{Hom}(M, \mathcal{X})$ where M is a compact manifold. Yet another place diffeological groupoids appear is as inertia groupoids (see Example 3.9), which play an important role in the theory of Lie group actions and orbifolds.

Thus it became natural for a rigorous foundation for diffeological groupoids and their Morita equivalence to be developed. The thesis and subsequent paper [vdS20; vdS21] of van der Schaaf provide such a foundation in terms of bibundles, extending the theory of Lie groupoids and their bibundles to the diffeological realm; indeed, the bicategory of Lie groupoids, right principal bibundles, and bi-equivariant diffeomorphisms forms a sub-bicategory of the diffeological version. An important open question [vdS21, Question 7.6] is whether a Morita equivalence between two Lie groupoids *in the diffeological bicategory* is, in fact, a Lie Morita equivalence; that is, the biprincipal bibundle between the two Lie groupoids representing the Morita equivalence is actually a bibundle from the Lie bicategory (*i.e.* a smooth Hausdorff manifold). An affirmative answer Corollary 9.32 is one of the motivations for the present manuscript.

At the same time, there are other, equivalent, bicategories of Lie groupoids besides the bibundle version mentioned above. Indeed, the bicategory of fractions developed by Pronk [Pro96] is one that is utilised regularly in the literature (see also [PS21]). Another is the anafunctor bicategory, popularised by Roberts [Rob12; Rob16; Rob21]. Each of these bicategories has its pros and cons:

- The 1-cells of the bicategory of fractions, called generalised morphisms, are spans of functors between Lie groupoids $\mathcal{G} \leftarrow \mathcal{K} \rightarrow \mathcal{H}$, in which the left arrow is a “weak equivalence” (see Definition 6.3). These 1-cells have few conditions and are relatively easy to construct. Also, composition is associative, which is not always the case in a bicategory. However, the 2-cells are equivalence classes of diagrams

(see [Definition 6.5](#)), and quickly become tedious to work with, especially when the user needs to vertically or horizontally compose them.

- The 1-cells of the anafunctor bicategory, called anafunctors, have one more condition on these spans: the weak equivalence is also required to be a covering in the Grothendieck pretopological sense (see [Definition 7.8](#)); for Lie groupoids, this reduces to the weak equivalence being surjective submersive on objects. This extra condition has the wonderful property of isolating a natural representative of a 2-cell mentioned above, which in fact is just a natural transformation (see [Definition 7.10](#) and [Lemma 7.25](#)). Thus working with 2-cells in the anafunctor bicategory becomes much easier than in the bicategory of fractions. Composition also remains associative, as anafunctors are just special cases of generalised morphisms.
- The 1-cells of the bibundle bicategory are right principal bibundles (see [Definition 8.5](#)). These can be viewed as very specialised anafunctors, so specialised that the arrow space of the groupoid at the apex of the span becomes redundant, its information being absorbed into actions on the bibundle itself. The 2-cells are bi-equivariant diffeomorphisms (see [Definition 8.7](#)), thus leading to a very rigid bicategory. Consequently, a bibundle can be difficult to construct when needed: for instance, given two Lie groupoids that you suspect are Morita equivalent, how do you construct a bibundle between them? However, often this rigidity allows a bibundle to exhibit desired properties that the user may wish for the 1-cells of their bicategory that are invariant under bi-equivariant diffeomorphisms (see below). Moreover, since 2-cells are just functions, they are very easy to deal with when needing to compose vertically or horizontally. Composition of bibundles, however, is only associative up to isomorphism, although the corresponding associator is not overly complicated (see [Definition 8.6](#)).

Due to the pros and cons mentioned above, knowing how to move from one bicategory to another becomes a very useful tool. Indeed, one of the motivations for this paper is the proof of [[PS10](#), Proposition 7.1]. In this proposition, the authors begin with a generalised morphism between action groupoids of Lie group actions $G \ltimes M$ and $H \ltimes N$ representing orbifolds. They replace each functor of the span with their corresponding bibundles, compose these bibundles, and observe that the resulting bibundle induces a generalised morphism that is “equivariant” in the sense that it corresponds to a Lie group homomorphism $G \rightarrow H$. Thus the authors started with the first bicategory mentioned above, moved to the third bicategory where the composition took place, and then moved back to the first taking the desired properties exhibited by the resultant bibundle along with them. (In fact, if one looks carefully, they may notice that the authors moved from the first to the second bicategory, then from the second to the third, and then back to the first.)

“Moving from one bicategory to another” can be made more precise as follows. In [[Pro96](#)], Pronk shows that if a class W of 1-cells of a bicategory B satisfies certain properties, then B can be localised at W , meaning that a new bicategory $B[W^{-1}]$ (called the bicategory of fractions) can be constructed in which elements of W become invertible. This emulates the localisation of a ring, or the localisation procedure of [[GZ67](#)] for a

category on which the recipe of [Pro96] is based. Pronk also gives conditions under which a different bicategory C , admitting an “inclusion” of B in which elements of W become invertible in C is equivalent to $B[W^{-1}]$. This is the celebrated Comparison Theorem [Pro96, Proposition 24], the result being a pseudo-natural transformation

$$\begin{array}{ccc} B[W^{-1}] & \xrightarrow{\cong} & C \\ \uparrow & \Downarrow & \nearrow \\ B & & \end{array}$$

So, “moving” from one bicategory to another is moving along an arrow at the top of the above diagram, which is an equivalence of bicategories. There is the typical existential issue with this, however. While the existence of such an equivalence may be known, what is a 1-cell or a 2-cell of $B[W^{-1}]$ sent to when moving to C ? Or back? It turns out the pseudo-natural transformation allows for more than one correct answer, although all of the answers are isomorphic. This is fine if the user only needs to know of the existence, but what if one needs an explicit formula?

The hope is that this paper will fulfill this need of providing explicit formulas and methods for not only moving from one bicategory to another, but also formulas for operations such as vertical and horizontal composition in all of the bicategories, which can be difficult to find in the literature. One often has to look to the general theory and deduce what the operations are in their setting, which can be a laborious task. Thus this paper is partially expository, addressing this purpose. On the other hand, it generalises the known relationships between the three bicategories of Lie groupoids mentioned above to diffeological groupoids, which (except for diffeological bibundles) is new.

So, what is diffeology? It is a generalisation of the theory of smooth manifolds, developed by Souriau in the 1980s [Sou80; Sou84]. In a few more words, the category **Diffeol** of diffeological spaces and smooth maps is a complete, co-complete quasi-topos [BH11] that includes smooth manifolds as a full subcategory. This means that arbitrary subsets, quotients, fibred products, function spaces, etc., are all included in **Diffeol**, and thus one can perform differential geometry on singular spaces arising as orbit spaces of Lie group actions or non-smooth varieties, as well as infinite-dimensional spaces. Indeed, as mentioned above, Fréchet manifolds form a full subcategory of **Diffeol**. Thus it is a very convenient category to work with. When groupoids exhibiting singularities show up in differential geometry, it is therefore only natural to look to a diffeological version of groupoids to work with. Our reference for the basics of diffeology is the standard text by Iglesias-Zemmour [IZ13].

Bicategories and 2-categories are “higher” categories, which have popped up in geometry (algebraic and differential), especially as treatments of orbit spaces of group actions that capture invariants that other methods may not detect, such as quotient topologies or even diffeologies. Our main resources for bicategories and 2-categories are [B67; JY21].

This paper is broken down as follows. **Section 2** provides the reader with the basics of diffeology required for this paper, although we refer to [IZ13] for details. In **Section 3**, we define diffeological groupoids and their 2-category **DGpoid**, which contains the 2-category of Lie groupoids **LieGpoid** as a full subcategory. **Section 4** presents one of the

motivations for the localisation of the 2-category of groupoids: weak equivalences, which are “smooth” equivalences of categories between groupoids, are not typically invertible, and so we require a larger bicategory in which they are. This section develops this idea, and presents properties of weak equivalences required in the rest of the paper. [Section 5](#) develops weak and strict pullbacks of diffeological groupoids, as well as their properties, which are crucial throughout the sequel.

With all of the ingredients in place, [Section 6](#) presents the bicategory of fractions $\mathbf{DGpoid}[W^{-1}]$ à la Pronk [[Pro96](#)] for diffeological groupoids [Corollary 6.11](#), with explicit formulas for vertical and horizontal compositions, unitors, etc. It also includes an explicit formulation of how functors and natural transformations embed in $\mathbf{DGpoid}[W^{-1}]$; we call the associated pseudofunctor “spanisation”. In [Section 7](#), we repeat this presentation, but for the anafunctor category $\mathbf{DGpoid}_{\text{ana}}$ à la Roberts [[Rob21](#)], again with explicit formulas; see [Corollary 7.15](#). Spanisation is what often appears in the literature as the method of embedding a bicategory B into its anafunctor category; we present a different pseudofunctor “anafunctisation” for the embedding. We find that anafunctisation allows for easier access to explicit formulas for the various bicategorical operations. This is discussed in [Subsection 7.3](#), and $\mathbf{DGpoid}[W^{-1}]$ and $\mathbf{DGpoid}_{\text{ana}}$ are compared in [Subsection 7.4](#). In [Subsection 7.5](#), we show how passing from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DGpoid}_{\text{ana}}$ “optimises” 2-cells of $\mathbf{DGpoid}[W^{-1}]$, in the sense that there is a unique representative of each 2-cell in $\mathbf{DGpoid}[W^{-1}]$ that is a natural transformation, which in turn is a 2-cell in $\mathbf{DGpoid}_{\text{ana}}$. Next, in [Section 8](#), we present the necessary background on the bibundle bicategory $\mathbf{DBiBund}$ as presented in [[vdS21](#)]. [Subsection 8.2](#) discusses “bibundlisation”, which is the embedding of \mathbf{DGpoid} into $\mathbf{DBiBund}$. We then compare $\mathbf{DGpoid}[W^{-1}]$ and $\mathbf{DBiBund}$ in [Subsection 8.3](#), similar to the comparison made with $\mathbf{DGpoid}_{\text{ana}}$ earlier. In [Subsection 8.4](#), we indicate how passing from $\mathbf{DGpoid}[W^{-1}]$ or $\mathbf{DGpoid}_{\text{ana}}$ to $\mathbf{DBiBund}$ “optimises” 1-cells, yielding a more geometric type of generalised morphism/anafunctor in which all redundant information has been removed. These three sections are then summarised in [Theorem 8.25](#).

In [Section 9](#), we focus our attention on Lie groupoids and locally subductive diffeological groupoids; a local subduction is a smooth surjection that not only admits smooth “sections” (with a more diffeological flavour), but admits them through specified points, similar to surjective submersions in smooth manifold theory. Some geometers prefer to use locally subductive diffeological groupoids due to the nice properties that local subductions exhibit, and so in [Subsection 9.1](#) we show that such groupoids localise in the same way as standard diffeological groupoids, and the result is an essentially full sub-bicategory of $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, or $\mathbf{DBiBund}$; see [Theorem 9.19](#). We then further restrict to Lie groupoids; even though this theory is well-known, we wish to not only show that the theory is a special case of the diffeological version (and thus the formulas we presented in earlier sections are valid for Lie groupoids as well), but use the Axiom of Choice to answer the open question [[vdS21](#), Question 7.6] in [Corollary 9.32](#) confirming that a Morita equivalence in $\mathbf{DBiBund}$ between Lie groupoids is a Morita equivalence in the corresponding Lie sub-bicategory $\mathbf{LieBiBund}$. In fact, we prove this in the Lie anafunctor sub-bicategory, and “move” this to the bibundle setting. The section is summarised in [Theorem 9.30](#).

We present some applications of the theory in [Section 10](#). We would be remiss not to mention stacks in a paper such as this on groupoids, and so in [Subsection 10.1](#), we prove the expected result that the three bicategories $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, and $\mathbf{DBiBund}$ are all equivalent to geometric stacks over the site of diffeological spaces; we then repeat the result for the locally subductive case. This result is already known to experts, but pinning down an explicit proof appears to be difficult, and so we show that the recipe developed by Villatoro [[Vil18](#)] applies in our setting. In [Subsection 10.2](#), we port some of the invariants discussed in [[vdS21](#), Section 6] of diffeological groupoids in the localised bicategories (called ‘‘Morita invariants’’) to this paper, and introduce some new ones. Finally, in [Subsection 10.3](#), we show that in very special cases, we can use known classification results of principal bundles in the diffeological setting to study the category of generalised morphisms and 2-cells between them in $\mathbf{DGpoid}[W^{-1}]$ using diffeological Čech cohomology. Throughout the paper, we present examples to help illustrate the theory.

In future work, categorical properties of these localised bicategories of diffeological groupoids will be explored. This said, many more directions exist for exploration. For example, since we are working in the category of diffeological spaces, it should be possible to turn the ‘‘optimisation’’ procedures discussed in [Subsection 7.5](#) and [Subsection 8.4](#) into actual optimisation problems. This involves equipping the groupoid whose objects are 1-cells between fixed diffeological groupoids, and arrows are 2-cells, with diffeologies on objects and arrows. The first problem that arises, of course, is with size, but there are ways around this. Another direction: while [[BFW13](#)] constructs a ‘‘Lie algebroid’’ to the diffeological groupoid that they are working with, there is no formal development for this theory. Of course, the development of such a theory is not a simple task. Indeed, even in the case of diffeological groups, the relationship with Lie algebras enjoyed by finite-dimensional Lie groups does not generalise, as is well-known in the study of infinite-dimensional groups.

Acknowledgements: The author wishes to thank Dorette Pronk and Laura Scull for many conversations on subtleties of bicategorical theory and localisation, Joel Villatoro for discussions on stacks and locally subductive diffeological groupoids, and Nesta van der Schaaf for his help as this project was just getting off the ground. We are also indebted to Laura Scull for suggestions on notation, which helped make the paper much easier to read, and David Michael Roberts for his feedback on the first version of the paper.

2. DIFFEOLOGICAL PRELIMINARIES

The purpose of this section is to review the preliminary diffeological theory required for the paper. The main reference is [[IZ13](#)]. Throughout, all manifolds will be considered smooth Hausdorff second-countable manifolds with no boundary, and ‘‘smooth’’ means infinitely-differentiable.

Definition 2.1 (The Category $\mathbf{Diffeol}$). Fix a set X . A **parametrisation of X** is a map of sets $p: U \rightarrow X$ where U is an open subset of a Cartesian space \mathbb{R}^n , but n is not fixed. A **diffeology** \mathcal{D}_X on X is a family of parametrisations satisfying:

- (1) (**Covering Axiom**) Every constant parametrisation is in \mathcal{D}_X .

- (2) (**Locality Axiom**) If $p: U \rightarrow X$ is a parametrisation in which U admits an open cover $\{U_\alpha\}$ so that for each α , there exists $p_\alpha: U_\alpha \rightarrow X$ in \mathcal{D}_X satisfying $p_\alpha = p|_{U_\alpha}$, then $p \in \mathcal{D}_X$.
- (3) (**Smooth Compatibility Axiom**) If $p: U \rightarrow X$ is in \mathcal{D}_X and $f: V \rightarrow U$ is smooth, with V an open subset of a Cartesian space, then $p \circ f \in \mathcal{D}_X$.

We refer to elements of \mathcal{D}_X as **plots** of the diffeology. We often will denote the domain of a plot p by U_p . We call the pair (X, \mathcal{D}_X) a **diffeological space**, and we often will drop \mathcal{D}_X .

A function $F: X \rightarrow Y$ between diffeological spaces X and Y is a (**diffeologically smooth map**) if $F \circ p$ is a plot of Y for any plot p of X . A smooth map is a **diffeomorphism** if it has a smooth inverse. Diffeological spaces with smooth maps comprise a category, denoted **Diffeol**. \diamond

In [BH11] it is shown that the category **Diffeol** is a complete co-complete quasi-topos; in particular, it admits diffeological structures on subsets, products, disjoint unions, and quotients, as well as function spaces. We go through the definitions we need here. See [IZ13] for details.

Definition 2.2 (New Diffeological Spaces from Old). Fix a diffeological space X and a family of diffeological spaces $\{X_\beta\}$.

- (1) (**Subset Diffeology**) If $A \subseteq X$, then a plot p is in the **subset diffeology** on A if it is a plot of X with image in A . The **inclusion map** $i_A: A \rightarrow X$ is smooth with respect to this diffeology.
- (2) (**Product Diffeology**) A plot p is in the **product diffeology** on the product $\prod_\beta X_\beta$ if $\text{pr}_\beta \circ p \in \mathcal{D}_{X_\beta}$ for each β , where $\text{pr}_{\beta_0}: \prod_\beta X_\beta \rightarrow X_{\beta_0}$ is the **projection map**. The projection maps pr_β are smooth with respect to this diffeology.
- (3) (**Quotient Diffeology**) If \sim is an equivalence relation on X with **quotient map** $\pi: X \rightarrow X/\sim$, then a plot p is in the **quotient diffeology** on X/\sim if U_p admits an open cover $\{U_\alpha\}$ so that for each α , there is a plot $q_\alpha: U_\alpha \rightarrow X$ such that $\pi \circ q_\alpha = p|_{U_\alpha}$; that is, p locally lifts to a plot of X . The quotient map π is smooth with respect to this diffeology.
- (4) (**Coproduct Diffeology**) A plot p is in the **coproduct diffeology** on $\coprod_\beta X_\beta$ if $p|_{p^{-1}(X_\beta)}$ is a plot of X_β for each β . The **inclusion maps** $i_\beta: X_\beta \rightarrow \coprod_\beta X_\beta$ are smooth with respect to this diffeology.
- (5) (**Fibred Product Diffeology**) Given diffeological spaces Y and Z and smooth maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$, the **fibred product diffeology** on the fibred product of sets $X_f \times_g Y$ is the subset diffeology induced by the product diffeology on $X \times Y$. \diamond

In the case of the subset and coproduct diffeologies, the maps i_A and i_B of [Definition 2.2](#) are special cases of a more general map called an induction.

Definition 2.3 (Inductions). An injective smooth map $f: X \rightarrow Y$ is an **induction** if $f^{-1} \circ p$ is a plot of X for any plot p of Y with image in $\text{im}(f)$. \diamond

In the case of the quotient and product diffeologies, the maps pr_B and π of [Definition 2.2](#) are special cases of a more general map called a subduction that plays an important rôle throughout the paper.

Definition 2.4 (Subductions). A smooth map $f: X \rightarrow Y$ is a **subduction** if for any plot p of Y , there is an open cover $\{U_\alpha\}$ of U_p and for each α a plot $q_\alpha: U_\alpha \rightarrow X$ such that $f \circ q_\alpha = p|_{U_\alpha}$. In particular, a subduction is surjective. We refer to q_α as a **(local) lift of p to X** , and often denote a subduction with a double-headed arrow \rightarrow . It is further a **local subduction** if for any $u \in U_p$, $x \in f^{-1}(p(u))$, and each α such that $u \in U_\alpha$, we can choose q_α so that $q_\alpha(u) = x$. We refer to such a q_α as a **(local) lift of p through x** . \diamond

Checking whether a map is a subduction, and taking lifts of plots along maps that are subductions, is a major theme throughout this paper. For the sake of brevity and to help streamline proofs, we set the following notation and terminology.

Notation 2.5 (Plot and Subduction Notation and Terminology). Given a plot p of a product space such as $X \times Y \times Z$, we use subscripts to denote the plot p as in the triple $p = (x_u, y_u, z_u)$ where $u \in U_p$. Given a (local) subduction $f: X \rightarrow Y$ and a plot p , we often will “shrink” the domain U_p so that a lift of it exists to X . Typically, without loss of generality, we will assume $0 \in U_p$ and thus by “shrink” U_p , we mean to take an open neighbourhood of 0 in U_p .

This paper is about diffeological groupoids, and a special type of diffeological groupoid will involve nebulae of a diffeological space. These are constructed out of so-called generating families of the diffeology.

Definition 2.6 (Generating Families and Nebulae). Fix a diffeological space X . A **(covering) generating family** of the diffeology \mathcal{D}_X is a family of plots $\mathcal{F} \subseteq \mathcal{D}_X$ satisfying: for each $p \in \mathcal{D}_X$ there exist an open cover $\{U_\alpha\}$ of U_p and for each α , a plot $q_\alpha: V_\alpha \rightarrow X$ in \mathcal{F} and a smooth function $f_\alpha: U_\alpha \rightarrow V_\alpha$ such that $p|_{U_\alpha} = q_\alpha \circ f_\alpha$. The **nebula** $\mathcal{N}(\mathcal{F})_0$ of \mathcal{F} is the coproduct $\coprod_{q \in \mathcal{F}} U_q$, which comes equipped with the (smooth) **evaluation map** $\text{ev}: \mathcal{N}(\mathcal{F})_0 \rightarrow X$ sending $u \in U_q$ to $q(u)$ for each $q \in \mathcal{F}$. \diamond

The following is a list of properties of (local) subductions used throughout the paper.

Proposition 2.7 (Properties of (Local) Subductions).

- (1) *The composition of two (local) subductions is a subduction.*
- (2) *An injective subduction is a diffeomorphism.*

- (3) If $f: M \rightarrow N$ is a smooth map of manifolds, then f is a surjective submersion if and only if it is a local subduction.
- (4) If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are smooth with $g \circ f$ a subduction, then g is a subduction.
- (5) Given smooth functions $f: X \rightarrow Z$ and $g: Y \rightarrow Z$, if f is a (local) subduction, then so is $\text{pr}_2: X_f \times_g Y \rightarrow Y$.

Proof. **Item 1** is proved in [IZ13, Articles 1.47 and 2.17]. **Item 2** is proved in [IZ13, Article 1.49]. **Item 3** is proved in [IZ13, Article 2.16].

For **Item 4**, it is immediate that g is a surjection. Let p be a plot of Z . Since $g \circ f$ is a subduction, after shrinking U_p , there is a lift q of p to X . Then $f \circ q$ is the desired lift of p to Y .

For **Item 5**, since f is a surjection, for any $y \in Y$ there is an $x \in X$ such that $f(x) = g(y)$. Thus $(x, y) \in X_f \times_g Y$, and it follows that pr_2 is surjective. Fix a plot p of Y . Since f is a subduction, after shrinking U_p there is a lift q of $g \circ p$ to X . Then (q, p) is a lift of p to $X_f \times_g Y$. Suppose now that f is a local subduction. Fix $u \in U$ and $(x, y) \in X_f \times_g Y$ such that $y = p(u)$. We can choose q so that it is a lift of $g \circ p$ through x . Then (q, p) is a lift of p to $X_f \times_g Y$ through (x, y) . \square

3. DIFFEOLOGICAL GROUPOIDS

In this section we introduce the main mathematical object of study for this paper: the *diffeological groupoid*. The definition is a straightforward generalisation of the definition of a Lie groupoid, however, due to the nice properties of the category **Diffeol** of diffeological spaces, it is simpler to state (see also **Remark 3.2**).

Definition 3.1 (Diffeological Groupoids). A **diffeological groupoid** \mathcal{G} is a groupoid $\mathcal{G}_1 \rightrightarrows \mathcal{G}_0$ in which \mathcal{G}_1 and \mathcal{G}_0 are diffeological spaces and all structure maps are smooth. We denote by s , t , u , m , and inv the **source**, **target**, **unit**, **multiplication**, and **inversion** maps, although we will also often write $g \cdot h$ or gh instead of $m(g, h)$ for multiplication and g^{-1} instead of $\text{inv}(g)$ for inversion. A **Lie groupoid** is a diffeological groupoid in which \mathcal{G}_0 and \mathcal{G}_1 are manifolds, and s (and hence t) are local subductions (equivalently surjective submersions, by **Item 3** of **Proposition 2.7**). \diamond

Remark 3.2. The source and target maps of a diffeological groupoid are subductions, and the unit map an induction. However, it is not true that a diffeological groupoid \mathcal{G} in which \mathcal{G}_1 and \mathcal{G}_0 are manifolds is necessarily a Lie groupoid. Indeed, consider the diffeological groupoid $\mathbb{R} \amalg \mathbb{R} \rightrightarrows \mathbb{R}$, in which one copy of \mathbb{R} is the unit space, and the other copy is sent by both source and target maps to $0 \in \mathbb{R}$. Neither the source nor the target map is a submersion, although they are both subductions. \lrcorner

Definition 3.3 (Smooth Functors and Smooth Natural Transformations). A functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ between diffeological groupoids is **smooth** if the maps $\varphi_0: \mathcal{G}_0 \rightarrow \mathcal{H}_0$ and

$\varphi_1: \mathcal{G}_1 \rightarrow \mathcal{H}_1$ are smooth. A natural transformation $T: \varphi \Rightarrow \psi$ between smooth functors $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$ is **smooth** if the corresponding map $T: \mathcal{G}_0 \rightarrow \mathcal{H}_1$ is smooth. \diamond

It is straightforward to check that diffeological groupoids with smooth functors and smooth natural transformations form a strict 2-category **DGpoid**, where horizontal and vertical composition of 2-cells are the usual ones for natural transformations. Lie groupoids with smooth functors and smooth natural transformations between them form a full 2-subcategory **LieGpoid** of **DGpoid**. All functors and natural transformations henceforth will be assumed smooth unless stated otherwise. Keep in mind that all natural transformations in this 2-category are natural isomorphisms, since the objects are groupoids.

Example 3.4 (Trivial and Pair Groupoids). Let X be a diffeological space. The **trivial groupoid** is the diffeological groupoid $X \rightrightarrows X$, often just denoted by X itself, in which the source and target maps are both equal to the identity map of X . The **pair groupoid** of X is the diffeological groupoid $X^2 \rightrightarrows X$ whose source and target maps are the first and second projection maps, resp. The unit map in this case is the diagonal map. Any diffeological groupoid \mathcal{G} has a natural functor $\chi_{\mathcal{G}}$ to the pair groupoid $\mathcal{G}_0^2 \rightrightarrows \mathcal{G}_0$, called the **characteristic functor**, equal to the identity on objects and which sends arrows g to $(s(g), t(g))$. We often will simply denote $\chi_{\mathcal{G}}$ by (s, t) when the groupoid \mathcal{G} is understood. \parallel

Example 3.5 (Relation Groupoids). Let X be a diffeological space and \sim an equivalence relation on X with quotient map π . The **relation groupoid of \sim** is the diffeological groupoid $X_{\pi} \times_{\pi} X \rightrightarrows X$ with source and target the projection maps pr_1 and pr_2 to X . \parallel

Example 3.6 (Action Groupoids). Let X be a diffeological space and G a diffeological group acting on X . The **action groupoid** $G \ltimes X := (G \times X \rightrightarrows X)$ with source $s(g, x) = x$ and target $t(g, x) = g \cdot x$ is a diffeological groupoid. \parallel

We can extend the previous example to include groupoid actions, which we will need in the sequel.

Definition 3.7 (Groupoid Actions). A **(smooth) left groupoid action** $\mathcal{G} \curvearrowright^{l_X} X$ of a diffeological groupoid \mathcal{G} on a diffeological space X *along* a map $l_X: X \rightarrow \mathcal{G}_0$ is a smooth function $\text{act}: (\mathcal{G}_1)_{s \times l_X} X \rightarrow X: (g, x) \mapsto g \cdot x$, satisfying the following three conditions:

- (1) $l_X(g \cdot x) = t(g)$ for $g \in \mathcal{G}_1$ and $x \in X$ such that $s(g) = l_X(x)$.
- (2) $u_{l_X(x)} \cdot x = x$ for every $x \in X$.
- (3) $h \cdot (g \cdot x) = (hg) \cdot x$ whenever defined.

We call l_X the **(left) anchor map**; it is also called the (left) moment map in the literature.

Right actions are defined similarly: a **(smooth) right groupoid action** $X \curvearrowright^{r_X} \mathcal{G}$ of \mathcal{G} on X *along* $r_X: X \rightarrow \mathcal{G}_0$ is a smooth map $\text{act}: X_{r_X} \times_t \mathcal{G} \rightarrow X: (x, g) \mapsto x \cdot g$ satisfying the following three conditions:

- (1) $r_X(x \cdot g) = s(g)$ for $g \in \mathcal{G}_1$ and $x \in X$ such that $t(g) = r_X(x)$,
- (2) $x \cdot u_{r_X(x)} = x$ for every $x \in X$,
- (3) $(x \cdot g) \cdot h = x \cdot (gh)$ whenever defined.

We call r_X the **(right) anchor map**; it is also called the (right) moment map in the literature. A groupoid action induces an equivalence relation on X : x is equivalent to x' if they are in the same orbit of the action; that is, there is an element $g \in \mathcal{G}_1$ for which $x' = g \cdot x$ (or $x' = x \cdot g$). We call the diffeological quotient the **orbit space** of the action, denoted $\mathcal{G} \backslash^X$ in the case of a left action and X / \mathcal{G} in the case of a right action. \diamond

Example 3.8 (Groupoid Multiplication as an Action). Any diffeological groupoid $\mathcal{G} \rightrightarrows \mathcal{G}_0$ acts on its own arrow space from the left and right by composition, which gives actions $\mathcal{G} \curvearrowright^t \mathcal{G}_1$ and $\mathcal{G}_1 \curvearrowleft^s \mathcal{G}$, resp., that are both defined by $(g, h) \mapsto gh$. The corresponding orbit spaces are denoted $\mathcal{G}_1 \backslash^{\mathcal{G}_0}$ and $\mathcal{G}_0 / \mathcal{G}_1$, resp. \parallel

Example 3.9 (Inertia Groupoids). Let \mathcal{G} be a diffeological groupoid. The image of the characteristic functor $\chi_{\mathcal{G}}$ is the relation groupoid induced by the equivalence relation as in [Definition 3.7](#) with respect to either action of [Example 3.8](#). The **kernel** of $\chi_{\mathcal{G}}$ is all arrows of \mathcal{G} sent to the unit space of the relation groupoid; that is, the subgroupoid of \mathcal{G} comprising the stabilisers: arrows with equal source and target. This subgroupoid is a diffeological groupoid, denoted $\ker(\chi_{\mathcal{G}})$. However, its arrow space,

$$\ker(\chi_{\mathcal{G}})_1 = \{k \in \mathcal{G}_1 \mid s(k) = t(k)\}$$

admits a left groupoid action of \mathcal{G} . Indeed, the anchor map for the action is $t \circ i$ where i is the inclusion of $\ker(\chi_{\mathcal{G}})$ into \mathcal{G}_1 , and the action itself is given by conjugation: $\text{act}(g, k) := g \cdot k \cdot g^{-1}$. The corresponding action groupoid $\mathcal{G} \ltimes \ker(\chi_{\mathcal{G}})$ is called the **inertia groupoid** of \mathcal{G} , and is a diffeological groupoid, denoted $\mathcal{J}_{\mathcal{G}}$. \parallel

4. WEAK EQUIVALENCES

Given that diffeological groupoids are categories, one may talk about a “smooth equivalence of categories” between two diffeological groupoids. In fact, it will be more useful to introduce a so-called “smooth adjoint equivalence”.

Definition 4.1 (Adjoint Equivalence). Let \mathcal{G} and \mathcal{H} be diffeological groupoids. A functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is a **(smooth) adjoint equivalence** if there exist a functor $\psi: \mathcal{H} \rightarrow \mathcal{G}$ and natural transformations $S: \text{id}_{\mathcal{G}} \Rightarrow \psi \circ \varphi$ and $T: \varphi \circ \psi \Rightarrow \text{id}_{\mathcal{H}}$ satisfying the **Triangle Identities**:

$$(T\varphi) \circ (\varphi S) = \text{id}_{\varphi} \quad \text{and} \quad (\psi T) \circ (S\psi) = \text{id}_{\psi}. \quad (1)$$

\diamond

Example 4.2 (Principal G -Bundles I). Let G be a Lie group, and $\pi: P \rightarrow B$ a principal G -bundle over a manifold B . The action groupoid $G \ltimes P$ admits a functor $\Pi: G \ltimes P \rightarrow B: (g, p) \mapsto \pi(p)$, denoting by B the trivial groupoid $B \rightrightarrows B$. Then Π is an adjoint equivalence if and only if P is the trivial G -bundle. Indeed, given a functor $\sigma: B \rightarrow G \ltimes P$ and natural transformations $S: \sigma \circ \Pi \Rightarrow \text{id}_{G \ltimes P}$ and $T: \text{id}_B \Rightarrow \Pi \circ \sigma$, it follows that σ is a global section of π , S is induced by the division map whose value at p yields the (unique) $g \in G$ sending $\sigma \circ \pi(p)$ to p , and T is the identity. Conversely, if P is trivial, we can take σ_0 to be a global section, S to be induced by the division map, and T to be trivial. The Triangle Identities are satisfied, making π an adjoint equivalence. //

Example 4.2 shows why the definition of adjoint equivalence may be too strong for most purposes involving diffeological groupoids. In geometry, diffeological/Lie groupoids are used when more information about a quotient space is needed to be retained than topology, and even diffeology, does not detect (*e.g.* stabilisers). But differentiating between different principal G -bundles is undesirable: the resulting quotient (the base manifold) is always the same, and so we expect the corresponding groupoids to be somehow “equivalent”. A potential solution is to weaken the definition of “equivalent”. Often when working with categories, one will drop the requirement that the Triangle Identities (1) are satisfied. But examining the details in **Example 4.2** will show that the Triangle Identities are not required to obtain that σ is a global section. So even this definition of “equivalence”, which is the standard definition of an equivalence of categories, is still too strong. There is another definition of equivalence of categories (typically applied when categories are locally small): the fully faithful and essentially surjective functor. This avoids the use of natural transformations, and more importantly, a functor going in the opposite direction. This motivates the following definition.

Definition 4.3 (Weak Equivalences). Let \mathcal{G} and \mathcal{H} be diffeological groupoids. A functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is a **weak equivalence** (also called an essential equivalence in the literature) if the following two conditions are met:

(**Smooth Essential Surjectivity**) The following map is a subduction:

$$\Psi_\varphi: \mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1 \longrightarrow \mathcal{H}_0: (x, h) \longmapsto s(h).$$

(**Smooth Fully Faithfulness**) The following map is a diffeomorphism:

$$\Phi_\varphi: \mathcal{G}_1 \longrightarrow (\mathcal{G}_0^2)_{\varphi^2} \times_{(s,t)} \mathcal{H}_1: g \longmapsto (s(g), t(g), \varphi(g)).$$

We will often denote a weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ by $\varphi: \mathcal{G} \xrightarrow{\sim} \mathcal{H}$. ◇

Remark 4.4.

- (1) In the definition of smooth essential surjectivity, we could equivalently replace $\mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1$ with $\mathcal{G}_{0,\varphi} \times_{\mathfrak{s}} \mathcal{H}_1$ and $(x, h) \mapsto s(h)$ with $(x, h) \mapsto t(h)$ [**vdS20**, Proposition 5.2].

- (2) An equivalent way of stating that φ is smoothly fully faithful is that \mathcal{G}_1 is a pullback (*i.e.* diffeomorphic to the fibred product) as in the following diagram.

$$\begin{array}{ccc} \mathcal{G}_1 \cong (\mathcal{G}_0^2)_{\varphi_0^2 \times_{(s,t)}} \mathcal{H}_1 & \xrightarrow{\text{pr}_3} & \mathcal{H}_1 \\ (\text{pr}_1, \text{pr}_2) \downarrow & & \downarrow (s,t) \\ \mathcal{G}_0^2 & \xrightarrow{\varphi_0^2} & \mathcal{H}_0^2 \end{array}$$

- (3) If $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is smoothly fully faithful, then Φ_φ is invariant under left and right multiplication. In particular,

$$\Phi_\varphi^{-1}(x_2, x_3, h_2) \cdot \Phi_\varphi^{-1}(x_1, x_2, h_1) = \Phi_\varphi^{-1}(x_1, x_3, h_2 h_1). \quad (2)$$

┘

One can check that a weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is a fully faithful and essentially surjective functor between the underlying categories of \mathcal{G} and \mathcal{H} . However, to motivate that it is a “natural” definition, we provide in the following lemma another description that is much more “categorical” using natural transformations, as well as [Lemma 7.7](#). Functors satisfying the first lemma below are called “representably fully faithful” by Roberts [[Rob21](#), Definition 2.2], and simply “fully faithful” by Pronk-Scull [[PS21](#), Definition 5.1]. It resembles the left-cancellative property of a monic map of a category. Functors satisfying [Lemma 7.7](#) are called “ J -locally split” by Roberts [[Rob21](#), Definition 3.22], and indicate that essential surjectivity is the same (up to natural transformation) as being a covering in the Grothendieck (pre)topological sense.

Lemma 4.5 (Representable Fully Faithfulness). *A functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is smoothly fully faithful if and only if for any two functors $\psi, \psi': \mathcal{K} \rightarrow \mathcal{G}$ and natural transformation $S: \varphi \circ \psi \Rightarrow \varphi \circ \psi'$, there is a unique natural transformation $S': \psi \Rightarrow \psi'$ such that $S = \varphi S'$.*

Proof. Suppose $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is smoothly fully faithful, and let $\psi, \psi': \mathcal{K} \rightarrow \mathcal{G}$ be functors and $S: \varphi \circ \psi \Rightarrow \varphi \circ \psi'$ a natural transformation. Define $S': \psi \Rightarrow \psi'$ by

$$S'(z) := \Phi_\varphi^{-1}(\psi(z), \psi'(z), S(z)).$$

Since φ is smoothly fully faithful, S' is well-defined and smooth, is a natural transformation, and $S = \varphi S'$. Uniqueness also follows from smooth fully faithfulness.

Conversely, suppose that for any functors $\psi, \psi': \mathcal{K} \rightarrow \mathcal{G}$ and natural transformation $S: \varphi \circ \psi \Rightarrow \varphi \circ \psi'$, there exists a unique natural transformation $S': \psi \Rightarrow \psi'$ such that $S = \varphi S'$. Fix a point $(x_1, x_2, h) \in (\mathcal{G}_0^2)_{\varphi_0^2 \times_{(s,t)}} \mathcal{H}_1$, and set \mathcal{K} to be the trivial groupoid of a point, ψ_0 to have image x_1 , ψ'_0 to have image x_2 , and S to have image h . There is a unique $S': \psi \Rightarrow \psi'$ such that $\varphi S' = S$, from which it follows that there is a unique $g \in \mathcal{G}_1$ with source x_1 , target x_2 , and such that $\varphi(g) = h$. So Φ_φ is bijective.

Fix a plot $p = (p_1, p_2, p_3): U \rightarrow (\mathcal{G}_0^2)_{\varphi_0^2 \times_{(s,t)}} \mathcal{H}_1$, set \mathcal{K} to be the trivial groupoid of U , $\psi_0 = p_1$, $\psi'_0 = p_2$, and $S = p_3$. There is a unique $S': \psi \Rightarrow \psi'$ such that $\varphi S' = S$, from

which it follows that $\Phi_\varphi \circ S' = p$. By **Item 2** of **Proposition 2.7**, Φ_φ is a diffeomorphism, and so φ is smoothly fully faithful. \square

We now show that a weak equivalence is the type of equivalence that we desire: any action groupoid of a principal G -bundle is weakly equivalent to the trivial groupoid of the underlying base manifold.

Example 4.6 (Principal G -Bundles II). Continuing the theme of **Example 4.2**, let G be a Lie group and $\pi: P \rightarrow B$ a principal G -bundle over a manifold B . Let q be a plot of B . After shrinking U_q , there exist an open subset V containing $q(U_q)$ and a smooth section $\sigma: V \rightarrow P|_V$. The plot $r = (\sigma \circ q, q)$ is a lift of q from B to $P_\pi \times_t B$. It follows that Ψ_π is a subduction.

Since the action of G is free, Φ_π is injective. Since the division map $d: P_\pi \times_\pi P \rightarrow G$ sending (p_1, p_2) to the unique g such that $p_2 = g \cdot p_1$ is smooth, it follows that Φ_π is subductive. By **Item 2** of **Proposition 2.7**, Φ_π is a diffeomorphism, and π is smoothly fully faithful. Thus π is a weak equivalence. We will see later (see **Proposition 9.21**) that it is a weak equivalence in the Lie sense as well. \parallel

From any diffeological space, we can construct a diffeological groupoid using a generating family for the diffeology; this groupoid appears in **[KWW21]** and is similar to the structure groupoid appearing in **[IZP20]**.

Example 4.7 (Nebulaic Groupoids I). Let X be a diffeological space with generating family \mathcal{F} of the diffeology \mathcal{D}_X . Construct a diffeological groupoid $\mathcal{N}(\mathcal{F})$, called the **nebulaic groupoid of \mathcal{F}** , with object space $\mathcal{N}(\mathcal{F})_0$, and arrow space

$$\mathcal{N}(\mathcal{F})_1 := (\mathcal{N}(\mathcal{F})_0)_{\text{ev}} \times_{\text{ev}} (\mathcal{N}(\mathcal{F})_0).$$

The inclusion $i: \mathcal{N}(\mathcal{F}) \rightarrow \mathcal{N}(\mathcal{D}_X)$ is a weak equivalence. Indeed, fix a plot f of $\mathcal{N}(\mathcal{D}_X)_0$. Without loss of generality, assume U_f is connected, so that its image is contained in U_p for some $p \in \mathcal{D}_X$. Then $p \circ f \in \mathcal{D}_X$. Thus, shrinking U_f , there exist a plot $q \in \mathcal{F}$ and a smooth $g: U_f \rightarrow U_q$ so that $p \circ f = q \circ g$. Then $(g, (f, g)): U_f \rightarrow (\mathcal{N}(\mathcal{F})_0)_i \times_t \mathcal{N}(\mathcal{D})_1$ is a lift of f . It follows that Ψ_i is a subduction.

It is immediate that Φ_i is injective, so by **Item 2** of **Proposition 2.7** it suffices to show that it is a subduction. Fix a plot $q = (q_1, q_2, (q_3, q_4))$ of $(\mathcal{N}(\mathcal{F})_0)_{i^2} \times_{(s,t)} \mathcal{N}(\mathcal{D})_1$. Then $q_1 = q_3$ and $q_2 = q_4$. The map $(q_1, q_2): U_q \rightarrow \mathcal{N}(\mathcal{F})_1$ is a plot, and $\Phi_i \circ (q_1, q_2) = q$. It follows that Φ_i is subductive, and we conclude that i is a weak equivalence. This example generalises the so-called Čech groupoids of Lie groupoid theory, in which \mathcal{F} is an open cover of a manifold. \parallel

The following lemma shows how essential surjectivity on objects of a category is induced by surjectivity on objects, within the realm of diffeological groupoids.

Lemma 4.8 (Subductive Weak Equivalences). *If $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is smoothly fully faithful and φ_0 is a subduction, then φ is a weak equivalence. In this case we call φ a **subductive weak equivalence**.*

Proof. Let p be a plot of \mathcal{H}_0 . Since φ_0 is a subduction, shrinking U_p , there is a lift q of p to \mathcal{G}_0 . The plot $(q, u \circ p): U \rightarrow \mathcal{G}_{0\varphi} \times_t \mathcal{H}_1$ lifts p through Ψ_φ . \square

Remark 4.9. It follows from the definitions that a subductive weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is also a subduction on arrows $\varphi_1: \mathcal{G}_1 \rightarrow \mathcal{H}_1$. \lrcorner

Example 4.10 (Nebulaic Groupoids II). Let X be a diffeological space and \mathcal{F} a generating family of the diffeology. Let ev be the functor from $\mathcal{N}(\mathcal{F})$ to the trivial groupoid of X , sending $u \in \mathcal{N}(\mathcal{F})_0$ to $\text{ev}(u)$ and $(u_1, u_2) \in \mathcal{N}(\mathcal{F})_1$ to $\text{ev}(u_1)$. It is straightforward to check that this is a subductive weak equivalence. \parallel

The major issue with weak equivalences in **DGpoid** is that they are typically not invertible. This manifests itself in **Example 4.6** if the principal G -bundle is not trivial. However, there are three standard ways of *formally* inverting weak equivalences, and we study the three resulting bicategories in remaining sections of this paper. To this end, we collect some necessary properties of weak equivalences, including some from [vdS20].

Example 4.11 (Adjoint Equivalence are Weak Equivalences). If $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is an adjoint equivalence, then φ is a weak equivalence, as shown in [vdS20, Proposition 5.5]. Indeed, let $\psi: \mathcal{H} \rightarrow \mathcal{G}$, S , and T be as in **Definition 4.1**. A plot p of \mathcal{H}_0 has lift $(\psi \circ p, T^{-1}p)$ to $\mathcal{G}_{0\varphi} \times_t \mathcal{H}_1$. Thus Ψ_φ is a subduction. Also, the map Φ_φ has smooth inverse given by $(x_1, x_2, h) \mapsto S^{-1}(x_2)\psi(h)S(x_1)$, and so is a diffeomorphism. The result follows. Note that we did *not* use the Triangle Identities (1) in the argument.

That the composition of two weak equivalences is a weak equivalence is proved in [vdS20, Proposition 5.7]. We expand on this here, showing that if two of φ , ψ , and $\psi \circ \varphi$ are weak equivalences, then so is the third.

Proposition 4.12 (Weak Equivalences and Composition). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ be functors. If any two of φ , ψ , and $\psi \circ \varphi$ are weak equivalences, then so is the third.*

Proof. Suppose φ and ψ are smoothly essentially surjective. Fix a plot p of \mathcal{K}_0 . Since ψ is smoothly essentially surjective, after shrinking U_p , there exists a lift $q: U_p \rightarrow \mathcal{H}_{0\psi} \times_t \mathcal{K}_1$ of p . Since φ is smoothly essentially surjective, after shrinking U_p again, there exists a lift $r: U_p \rightarrow \mathcal{G}_{0\varphi} \times_t \mathcal{H}_1$ of $\text{pr}_1 \circ q$. The product $P := (\psi \circ \text{pr}_2 \circ r) \cdot (\text{pr}_2 \circ q)$ is a plot of \mathcal{K}_1 , and the plot $(\text{pr}_1 \circ r, P): U_p \rightarrow (\mathcal{G}_0)_{\psi \circ \varphi} \times_t \mathcal{K}_1$ satisfies $\Psi_{\psi \circ \varphi} \circ (\text{pr}_1 \circ r, P) = p$. It follows that $\psi \circ \varphi$ is smoothly essentially surjective.

Suppose φ and ψ are smoothly fully faithful. Smooth fully faithfulness of $\psi \circ \varphi$ follows from **Item 2** of **Remark 4.4** and the fact that the concatenation of two pullback squares is itself a pullback square:

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{\varphi} & \mathcal{H}_1 & \xrightarrow{\psi} & \mathcal{K}_1 \\ \text{(s,t)} \downarrow & \lrcorner & \text{(s,t)} \downarrow & \lrcorner & \downarrow \text{(s,t)} \\ \mathcal{G}_0 \times \mathcal{G}_0 & \xrightarrow{\varphi \times \varphi} & \mathcal{H}_0 \times \mathcal{H}_0 & \xrightarrow{\psi \times \psi} & \mathcal{K}_0 \times \mathcal{K}_0. \end{array}$$

Therefore if φ and ψ are weak equivalences, so is $\psi \circ \varphi$.

Suppose ψ is smoothly fully faithful and $\psi \circ \varphi$ is smoothly essentially surjective. Fix a plot p of \mathcal{K}_0 . After shrinking U_p , there exists a lift q of p to $\mathcal{G}_{0,\psi \circ \varphi} \times_t \mathcal{K}_1$. The plot $r := \Phi_\psi^{-1}((p, \varphi \circ \text{pr}_1 \circ q), \text{pr}_2 \circ q)$ of \mathcal{H}_1 satisfies $s \circ r = p$ and $t \circ r = \varphi \circ \text{pr}_1 \circ q$, from which it follows that $(\text{pr}_1 \circ q, r)$ is the desired lift of p to $\mathcal{G}_{0,\varphi} \times_t \mathcal{K}_1$. It follows that φ is smoothly essentially surjective.

Suppose ψ and $\psi \circ \varphi$ are smoothly fully faithful. Then smooth fully faithfulness of φ follows from the fact that if the outer and right squares in the above diagram are pullback squares, then so is the left. Therefore, if ψ and $\psi \circ \varphi$ are weak equivalences, so is φ .

Suppose φ and $\psi \circ \varphi$ are weak equivalences. Let p be a plot of \mathcal{K}_0 . Since $\psi \circ \varphi$ is smoothly essentially surjective, shrinking U_p , there is a lift (q_1, q_2) of p to $\mathcal{G}_{0,\psi \circ \varphi} \times_t \mathcal{K}_1$. Then $(\varphi \circ q_1, q_2)$ is a plot of $\mathcal{H}_{0,\psi} \times_t \mathcal{K}_1$ such that $\text{pr}_2 \circ s \circ (\varphi \circ q_1, q_2) = p$. This proves that Ψ_ψ is a subduction.

Let $h_1, h_2 \in \mathcal{H}_1$ such that $\Phi_\psi(h_1) = \Phi_\psi(h_2)$. Then $s(h_1) = s(h_2) =: y$ and $t(h_1) = t(h_2) =: y'$. Since φ is smoothly essentially surjective, there exist $x, x' \in \mathcal{G}_0$ and arrows \tilde{h} from $\varphi(x)$ to y and \tilde{h}' from $\varphi(x')$ to y' in \mathcal{H}_1 . Since φ is smoothly fully faithful, there exist unique $g_i = \Phi_\varphi^{-1}\left((x, x'), \left(\tilde{h}'\right)^{-1} h_i \tilde{h}\right)$ for $i = 1, 2$ in \mathcal{G}_1 . It follows from the smooth fully faithfulness of $\psi \circ \varphi$ and the definitions of g_i that $g_1 = g_2$, and so $h_1 = h_2$. Thus Φ_ψ is injective.

Let $p = (p_1, p_2, p_3)$ be a plot of $\mathcal{H}_{0,\psi^2}^2 \times_{(s,t)} \mathcal{K}_1$. Since φ is smoothly essentially surjective, after shrinking U_p , there are plots (q_1, q_2) and (q'_1, q'_2) of $\mathcal{G}_{0,\varphi} \times_t \mathcal{K}_1$ such that $s \circ q_2 = p_1$ and $s \circ q'_2 = p_2$. Define a smooth map $g: U_p \rightarrow \mathcal{G}_1$ by

$$g := \Phi_{\psi \circ \varphi}^{-1}(q_1, q'_1, (\psi \circ q'_2) \cdot p_3 \cdot (\text{inv} \circ \psi \circ q_2)).$$

Setting $r := (\text{inv} \circ q'_2) \cdot (\varphi \circ g) \cdot q_2$, it follows that $\Phi_\psi \circ r = p$. It follows that Φ_ψ is a subduction, and hence a diffeomorphism by [Item 2](#) of [Proposition 2.7](#). Therefore, if φ and $\psi \circ \varphi$ are weak equivalences, so is ψ . \square

The last result of this section is [[vds20](#), Proposition 5.6], which shows how weak equivalences interact with natural transformations. We include the proof for completeness.

Proposition 4.13 (Natural Transformations Preserve Weak Equivalences). *Let $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$ be smooth functors between diffeological groupoids, admitting a smooth natural transformation $S: \varphi \Rightarrow \psi$. Then φ is a weak equivalence if and only if ψ is a weak equivalence.*

Proof. Since S is a natural isomorphism, it suffices to prove one of the implications. Suppose φ is a weak equivalence. To prove ψ is smoothly essentially surjective, fix a plot $p = y_u$ of \mathcal{H}_0 . Since φ is smoothly essentially surjective, after shrinking U_p , there is a lift (x_u, h_u) of p with respect to Ψ_φ . The map $(x_u, T(x_u)h_u)$ is the desired lift of p with respect to Ψ_ψ .

To prove that Φ_ψ is a diffeomorphism, by [Item 2](#) of [Remark 4.4](#) it suffices to show that \mathcal{G} satisfies the universal property of a pullback: let \mathcal{K} be a diffeological groupoid

and let $\chi: \mathcal{K} \rightarrow \mathcal{H}$ and $(\omega_1, \omega_2): \mathcal{K} \rightarrow \mathcal{G}_0 \times \mathcal{G}_0$ be functors making the following diagram commute:

$$\begin{array}{ccc}
 \mathcal{K}_1 & \xrightarrow{\chi} & \mathcal{H} \\
 \downarrow \exists! \theta & \searrow \psi & \downarrow (s,t) \\
 \mathcal{G}_1 & \xrightarrow{\psi} & \mathcal{H} \\
 \downarrow (s,t) & & \downarrow (s,t) \\
 \mathcal{G}_0^2 & \xrightarrow{\psi^2} & \mathcal{H}_0^2
 \end{array}$$

(ω_1, ω_2) is indicated by a curved arrow from \mathcal{K}_1 to \mathcal{G}_0^2 .

The functor $\chi': \mathcal{K} \rightarrow \mathcal{H}: k \mapsto S^{-1}(\omega_2(k))\chi(k)S(\omega_1(k))$ satisfies

$$(s, t) \circ \chi' = \varphi^2 \circ (\omega_1, \omega_2),$$

and so the smooth fully faithfulness of φ implies that there exists a unique smooth map $\theta: \mathcal{K} \rightarrow \mathcal{G}$ satisfying

$$(s, t) \circ \theta = (\omega_1, \omega_2) \quad \text{and} \quad \varphi \circ \theta = \chi'.$$

It follows that $\psi \circ \theta = \chi$, and so θ in the diagram above makes it commute. Uniqueness of θ follows from its uniqueness in the corresponding diagram for φ , from which smooth fully faithfulness of ψ follows. \square

5. PULLBACKS AND WEAK PULLBACKS

Pullbacks of various kinds are an integral part of this paper; namely, they are used to compose the 1-cells of our bicategories in later sections. We begin by defining what is meant by a pullback (or fibred product) of diffeological groupoids, as well as a “base change”.

Definition 5.1 (Pullbacks of Diffeological Groupoids). Let $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ be functors. The **strict** (or **strong**) **pullback** of φ and ψ is the diffeological groupoid $\mathcal{G}_{\varphi \times_{\psi} \mathcal{H}}$ whose object (resp. arrow) space is the fibred product of the object (resp. arrow) spaces of \mathcal{G} and \mathcal{H} . \diamond

Definition 5.2 (Base Change). Let \mathcal{H} be a diffeological groupoid and $f: X \rightarrow \mathcal{H}_0$ a smooth map. The **base change of \mathcal{H} along f** , denoted $\mathcal{H}[X]$, is the groupoid whose object space is X and arrow space is $X^2_{f^2 \times_{(s,t)} \mathcal{H}_1}$; the source and target maps are the projections to the first and second copies of X , respectively. In terms of **Definition 5.1**, $\mathcal{H}[X]$ is (canonically isomorphic to) $(X^2 \rightrightarrows X)_{f^2 \times_{(s,t)} \mathcal{H}}$, where f^2 is the functor sending objects x to $f(x)$ and arrows (x_1, x_2) to $(f(x_1), f(x_2)) \in \mathcal{H}_0^2$. In the literature, $\mathcal{H}[X]$ is also called the pullback of \mathcal{H} along f , and also denoted $f^*\mathcal{H}$. \diamond

Example 5.3 (Smooth Fully Faithfulness and Base Change). As an application of base change, we reformulate **Item 2** of **Remark 4.4** in terms of pullbacks of groupoids, yielding an equivalent condition for a functor to be smoothly fully faithful. Suppose $\varphi: \mathcal{G} \rightarrow \mathcal{H}$

is a functor. Then φ is smoothly fully faithful if and only if \mathcal{G} is diffeomorphic to the pullback of the pair groupoid $\mathcal{G}_0^2 \rightrightarrows \mathcal{G}_0$ and \mathcal{H} with respect to φ^2 and (s, t) , resp.

$$\begin{array}{ccc} \mathcal{G} \cong (\mathcal{G}_0^2 \rightrightarrows \mathcal{G}_0)_{\varphi_0^2 \times_{(s,t)}} \mathcal{H} & \xrightarrow{\text{pr}_2} & \mathcal{H} \\ \text{pr}_1 \downarrow & & \downarrow (s,t) \\ \mathcal{G}_0^2 \rightrightarrows \mathcal{G}_0 & \xrightarrow{\varphi_0^2} & \mathcal{H}_0^2 \rightrightarrows \mathcal{H}_0 \end{array}$$

Thus, a weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is a smoothly essentially surjective functor in which \mathcal{G} is isomorphic to the base change $\mathcal{H}[\mathcal{G}_0]$. //

The following is a list of some observations we can make thus far.

Proposition 5.4 (Weak Equivalences and Strict Pullbacks). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ be functors.*

- (1) *If ψ (resp. φ) is smoothly fully faithful, then so is $\text{pr}_1: \mathcal{G}_{\varphi \times_{\psi}} \mathcal{H} \rightarrow \mathcal{G}$ (resp. pr_2), in which case $\mathcal{G}[\mathcal{G}_{0, \varphi \times_{\psi}} \mathcal{H}_0]$ (resp. $\mathcal{H}[\mathcal{G}_{0, \varphi \times_{\psi}} \mathcal{H}_0]$) is isomorphic as a diffeological groupoid to $\mathcal{G}_{\varphi \times_{\psi}} \mathcal{H}$.*
- (2) *If ψ (resp. φ) is a subductive weak equivalence, then so is $\text{pr}_1: \mathcal{G}_{\varphi \times_{\psi}} \mathcal{H} \rightarrow \mathcal{G}$ (resp. pr_2).*

Proof. Suppose that ψ is smoothly fully faithful (a similar argument works for φ in place of ψ). Since ψ is a weak equivalence, $h = \Phi_{\psi}^{-1}(s(h), t(h), \varphi(g))$ for any $(g, h) \in (\mathcal{G}_{\varphi \times_{\psi}} \mathcal{H})_1$. It follows that Φ_{pr_1} is injective and subductive, and hence a diffeomorphism by **Item 2** of **Proposition 2.7**. Thus ψ is smoothly fully faithful. **Item 1** follows from the definition of $\mathcal{G}[\mathcal{G}_{0, \varphi \times_{\psi}} \mathcal{H}_0]$.

Suppose that ψ is a subductive weak equivalence (a similar argument works for φ in place of ψ). By **Lemma 4.8** and **Item 5** of **Proposition 2.7**, it suffices to show that pr_1 is smoothly fully faithful. But this follows from **Item 1**. This proves **Item 2**. \square

What happens if ψ in **Item 2** of **Proposition 5.4** is a weak equivalence, but not subductive? Then pr_1 is not guaranteed to be a weak equivalence in this case, as the following example illustrates.

Example 5.5 (Strict Pullbacks do not Preserve Weak Equivalence). Let \mathcal{G} and \mathcal{H} be the trivial groupoid over a point, \mathcal{K} be the pair groupoid $\mathbb{R} \times \mathbb{R} \rightrightarrows \mathbb{R}$, let $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ be the functor such that $\text{im } \varphi_0 = \{1\}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ the functor such that $\text{im } \psi_0 = \{-1\}$. Then φ and ψ are weak equivalences, but $\mathcal{G}_{\varphi \times_{\psi}} \mathcal{H} = \emptyset$; in particular, $\text{pr}_1: \mathcal{G}_{\varphi \times_{\psi}} \mathcal{H} \rightarrow \mathcal{G}$ is not a weak equivalence. //

To get pr_1 in **Item 2** of **Proposition 5.4** to be a weak equivalence without requiring ψ to be subductive, we must weaken the notion of pullback to allow for 2-commutative squares instead of commutative squares.

Definition 5.6 (Weak Pullback). Let $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ be functors. Define the **weak pullback** of φ and ψ to be the diffeological groupoid $\mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ in which

$$(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_0 := (\mathcal{G}_0)_\varphi \times_s (\mathcal{K}_1)_t \times_\psi (\mathcal{H}_0),$$

the space of triples $(x, k, y) \in \mathcal{G}_0 \times \mathcal{K}_1 \times \mathcal{H}_0$ such that k is an arrow from $\varphi(x)$ to $\psi(y)$; and

$$(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_1 := (\mathcal{G}_1)_{\text{so}\varphi} \times_s (\mathcal{K}_1)_t \times_{\text{so}\psi} (\mathcal{H}_1),$$

the space of triples $(g, k, h) \in \mathcal{G}_1 \times \mathcal{K}_1 \times \mathcal{H}_1$ with source and target maps

$$s(g, k, h) = (s(g), k, s(h)) \quad \text{and} \quad t(g, k, h) = (t(g), \psi(h) \circ k \circ \varphi(g)^{-1}, t(h)).$$

Thus, given triples $(x_1, k_1, y_1), (x_2, k_2, y_2) \in (\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_0$, an arrow (g, k_1, h) from (x_1, k_1, y_1) to (x_2, k_2, y_2) induces a commutative diagram

$$\begin{array}{ccc} \varphi(x_1) & \xrightarrow{k_1} & \psi(y_1) \\ \varphi(g) \downarrow & & \downarrow \psi(h) \\ \varphi(x_2) & \xrightarrow{k_2} & \psi(y_2). \end{array}$$

The other structure maps are given as follows. The unit map is given by $u(x, k, y) = (u(x), k, u(y))$, multiplication is given by

$$m((g_2, k_2, h_2), (g_1, k_1, h_1)) = (g_2 g_1, k_1, h_2 h_1)$$

for composable triples (g_1, k_1, h_1) and (g_2, k_2, h_2) , and inversion is given by

$$\text{inv}(g, k, h) = (g^{-1}, h \cdot k \cdot g^{-1}, h^{-1}).$$

◇

Remark 5.7.


- (1) The diagram induced by the weak pullback is 2-commutative, with the 2-cell the natural transformation $\text{pr}_2: \varphi \Rightarrow \psi$ given by $(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_0 \rightarrow \mathcal{K}_1: (x, k, y) \mapsto k$:

$$\begin{array}{ccc} & \mathcal{G}_\varphi \times_\psi^w \mathcal{H} & \\ \text{pr}_1 \swarrow & & \searrow \text{pr}_2 \\ \mathcal{G} & & \mathcal{H} \\ \varphi \searrow & \Rightarrow_{\text{pr}_2} & \swarrow \psi \\ & \mathcal{K} & \end{array}$$

- (2) There is a subtle point in [Definition 5.6](#) that should be addressed: given the diagram of diffeological spaces below, there are three candidates for what $A_f \times_g B_h \times_k C$ should be: $(A_f \times_g B)_{h \circ \text{pr}_2} \times_k C$, $A_f \times_{g \circ \text{pr}_1} (B_h \times_k C)$, or the subset


$$\{(a, b, c) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \subseteq A \times B \times C.$$

But these are all the same space. Indeed, this follows from the fact that given subsets of diffeological spaces $S \subseteq X$ and $T \subseteq Y$, the subset diffeology induced by the product diffeology on $S \times T \subseteq X \times Y$ is equal to the product diffeology induced by the subset diffeologies on S and T .



□

Proposition 5.8 (Universal Property of the Weak Pullback). *Given two functors $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$, the weak pullback $\mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ satisfies the following universal property: if \mathcal{L} is another diffeological groupoid with functors $\alpha: \mathcal{L} \rightarrow \mathcal{G}$ and $\beta: \mathcal{L} \rightarrow \mathcal{H}$ and a natural transformation $T: \varphi \circ \alpha \Rightarrow \psi \circ \beta$, then there exists a unique functor $\theta: \mathcal{L} \rightarrow \mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ up to natural isomorphism, making the following diagram 2-commute.*



(3)

Proof. Given \mathcal{L} as in the statement of the proposition, define $\theta: \mathcal{L} \rightarrow \mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ as follows: $\theta(x) = (\alpha(x), T(x), \beta(x))$ for $x \in \mathcal{L}_0$ and $\theta(\ell) = (\alpha(\ell), T(s(\ell)), \beta(\ell))$ for $\ell \in \mathcal{L}_1$. This is a well-defined, smooth functor such that $\text{pr}_1 \circ \theta = \alpha$ and $\text{pr}_3 \circ \theta = \beta$.

Let $\theta': \mathcal{L} \rightarrow \mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ be another smooth functor with natural transformations $R: \text{pr}_1 \circ \theta' \Rightarrow \alpha$ and $S: \text{pr}_3 \circ \theta' \Rightarrow \beta$. Define $U: \mathcal{L}_0 \rightarrow (\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_1$ by

$$U(x) := (R(x), (\psi S)^{-1} \circ T \circ (\varphi R)(x), S(x)).$$

Then U is a well-defined, smooth natural transformation $U: \theta' \Rightarrow \theta$. □

Corollary 5.9. *The inclusion $\theta: \mathcal{G}_\varphi \times_\psi \mathcal{H} \rightarrow \mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H}$ sending an object (x, y) to $(x, u(\varphi(x)), y)$ and an arrow (g, h) to $(g, u(\varphi(s(g))), h)$, is the unique map (up to natural isomorphism) making the diagram (3) 2-commute.*

We now prove the associated version of **Item 2** of **Proposition 5.4** in the case that we have a weak equivalence that is not subductive. However, if it *is* subductive, then this trait is maintained as well.

Proposition 5.10 (Properties of the Weak Pullback). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ be functors.*

- (1) *If ψ is a weak equivalence, then $\text{pr}_1: \mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H} \rightarrow \mathcal{G}$ is a subductive weak equivalence.*
- (2) *If ψ_0 is a subduction, then so is $(\text{pr}_1)_0: (\mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H})_0 \rightarrow \mathcal{G}_0$.*

Similar properties hold for pr_2 if we replace ψ with φ in **Proposition 5.10**.

Proof. Suppose ψ is a weak equivalence. Fix a plot $p = x_u$ of \mathcal{G}_0 . Since ψ is a weak equivalence, after shrinking U_p , there is a lift $q = (y_u, k_u)$ of $\mathcal{H}_{0_\psi} \times_{\mathcal{K}_1} \mathcal{K}_1$ of $\varphi \circ p$. Then (x_u, k_u, y_u) is a lift of p to $(\mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H})_0$, and so $(\text{pr}_1)_0$ is a subduction.

Since Φ_ψ is a diffeomorphism, Φ_{pr_1} is injective. A plot $p = ((x_u, k_u, y_u), (x'_u, k'_u, y'_u), g_u)$ of $\left((\mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H})_0^2 \right)_{\text{pr}_1^2 \times_{(s,t)} \mathcal{G}_1}$ has a lift to $(\mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H})_1$ given by

$$(g_u, k_u, \Phi_\psi^{-1}(y_u, y'_u, k'_u \cdot \varphi(g_u) \cdot k_u^{-1})).$$

It follows that Φ_{pr_1} is a subduction, and hence a diffeomorphism. By **Lemma 4.8**, pr_1 is a subductive weak equivalence. This proves **Item 1**.

Suppose that ψ_0 is subductive. By **Item 5** of **Proposition 2.7**, $(\text{pr}_1)_0: \mathcal{G}_{0_\varphi} \times_{\psi_0} \mathcal{H}_0 \rightarrow \mathcal{G}_0$ is a subduction. By **Corollary 5.9** and **Item 4** of **Proposition 2.7**, $(\text{pr}_1)_0: (\mathcal{G}_\varphi \overset{\text{w}}{\times}_\psi \mathcal{H})_0 \rightarrow \mathcal{G}_0$ is a subduction. This proves **Item 2**. \square

6. THE BICATEGORY OF FRACTIONS

The goal of this section is to expand **DGpoid** to include “inverses” of weak equivalences. More precisely, we wish to construct a new bicategory in which to “embed” **DGpoid** in such a way that we obtain inverses to the embedded weak equivalences from **DGpoid** (up to isomorphism); we call such a procedure a **localisation** of **DGpoid** at the class W of weak equivalences. To obtain this result, we show that W admits a right calculus of fractions (**Theorem 6.2**), and then follow the recipe in Pronk-Scull [**PS21**, §3] to construct the new so-called bicategory of fractions. This recipe is based on the original recipe of Pronk [**Pro96**, §2], but streamlined with weakened conditions on what is required for a class of arrows to admit a right calculus of fractions. We end the section by discussing in **Subsection 6.3** how **DGpoid** “embeds” into the localisation, a pseudofunctor called “spanisation”.

6.1. Weak Equivalences Admit a Calculus of Fractions. We begin by defining what it means for a class of arrows of **DGpoid** to admit a right calculus of fractions, and then show that the class of all weak equivalences satisfies this.

Definition 6.1 (Admits a Right Calculus of Fractions). Let A be a class of arrows of **DGpoid**. A admits a right calculus of fractions if it satisfies the following:

- (1) A contains all identity functors.
- (2) If $\varphi, \psi \in A$ are composable, then there exists $\chi \in A$ such that $\psi \circ \varphi \circ \chi \in A$.
- (3) For every pair of functors $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ with $\psi \in A$, there is a groupoid \mathcal{L} and functors $\chi: \mathcal{L} \rightarrow \mathcal{G}$ and $\omega: \mathcal{L} \rightarrow \mathcal{H}$, with $\chi \in A$ and a natural transformation $S: \varphi \circ \chi \Rightarrow \psi \circ \omega$.
- (4) For any $\varphi \in A$ and $S: \varphi \circ \psi \Rightarrow \varphi \circ \psi'$, there exists $\chi \in A$ and $T: \psi \circ \chi \Rightarrow \psi' \circ \chi$ such that $S\chi = \varphi T$. Moreover, given two pairs (χ_i, T_i) ($i = 1, 2$) satisfying this property, there exist ω_i such that $\chi_i \circ \omega_i \in A$ for $i = 1, 2$, and there exists $U: \chi_1 \circ \omega_1 \Rightarrow \chi_2 \circ \omega_2$ such that the following diagram commutes.

$$\begin{array}{ccc} \psi_1 \circ \chi_1 \circ \omega_1 & \xrightarrow{T_1 \omega_1} & \psi_2 \circ \chi_1 \circ \omega_1 \\ \psi_1 U \downarrow & & \downarrow \psi_2 U \\ \psi_1 \circ \chi_2 \circ \omega_2 & \xrightarrow{T_2 \omega_2} & \psi_2 \circ \chi_2 \circ \omega_2. \end{array}$$

- (5) Given $\varphi \Rightarrow \psi$, if $\varphi \in A$, then so is ψ .

◇

Theorem 6.2. *The class W of weak equivalences in **DGpoid** admits a right calculus of fractions.*

Proof. Each item in **Definition 6.1** will be checked. Since identity functors are weak equivalences, **Item 1** is true. By **Proposition 4.12**, **Item 2** is true where we let χ be the identity. By **Item 1** of **Remark 5.7** and **Item 1** of **Proposition 5.10**, **Item 3** is true. By **Lemma 4.5**, **Item 4** is true, where χ is the identity, and $T_1 = T_2$ due to uniqueness (and so ω_i are each the identity and U is trivial). By **Proposition 4.13**, **Item 5** is true. □

6.2. The Bicategory of Fractions $\mathbf{DGpoid}[W^{-1}]$. The next step is to go through the construction of the bicategory of fractions, which we do following **[Pro96]** and **[PS21]**. The objects of this bicategory will be the same as those of **DGpoid**: diffeological groupoids. The arrows will be spans of smooth functors, in which one functor is a weak equivalence; these are called generalised morphisms.

Definition 6.3 (Generalised Morphisms). Let \mathcal{G} and \mathcal{H} be diffeological groupoids. A **generalised morphism** between \mathcal{G} and \mathcal{H} is a diffeological groupoid \mathcal{K} and two functors

$\varphi: \mathcal{K} \rightarrow \mathcal{G}$ and $\psi: \mathcal{K} \rightarrow \mathcal{H}$ in which φ is a weak equivalence. We denote such a generalised morphism by $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$. Given a diffeological groupoid \mathcal{G} , the **identity generalised morphism** of \mathcal{G} is the generalised morphism $\mathcal{G} \xleftarrow{=} \mathcal{G} \xrightarrow{=} \mathcal{G}$. \diamond

We use the weak pullback to compose two generalised morphisms.

Definition 6.4 (Composition of Generalised Morphisms). The **composition** of a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{L} \xrightarrow[\psi]{\simeq} \mathcal{H}$ with $\mathcal{H} \xleftarrow[\varphi']{\simeq} \mathcal{M} \xrightarrow[\psi']{\simeq} \mathcal{K}$ is the generalised morphism $\mathcal{G} \xleftarrow[\varphi \circ \text{pr}_1]{\simeq} \mathcal{L} \times_{\psi \circ \varphi'}^{\mathcal{W}} \mathcal{M} \xrightarrow[\psi' \circ \text{pr}_3]{\simeq} \mathcal{K}$. That $\varphi \circ \text{pr}_1$ is a weak equivalence follows from **Item 1** of **Proposition 5.10** and **Proposition 4.12**. \diamond

Now we describe the 2-cells of the bicategory of fractions.

Definition 6.5 (2-Cells Between Generalised Morphisms). Given generalised morphisms $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{\simeq} \mathcal{H}$, consider a diagram of the form

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \nearrow \varphi & \uparrow \alpha & \searrow \psi & \\
 \mathcal{G} & & \mathcal{L} & & \mathcal{H} \\
 & \searrow \varphi' & \downarrow \alpha' & \nearrow \psi' & \\
 & & \mathcal{K}' & &
 \end{array}$$

$\Downarrow S_1$ $\Downarrow S_2$


where $\mathcal{K} \xleftarrow[\alpha]{\simeq} \mathcal{L} \xrightarrow[\alpha']{\simeq} \mathcal{K}'$ is a generalised morphism, and $S_1: \varphi \circ \alpha \Rightarrow \varphi' \circ \alpha'$ and $S_2: \psi \circ \alpha \Rightarrow \psi' \circ \alpha'$ are natural transformations making the diagram 2-commute. Denote such a diagram by the quadruple $(\alpha, \alpha', S_1, S_2)$.

Given another such diagram $(\beta, \beta', T_1, T_2)$ with generalised morphism $\mathcal{K} \xleftarrow[\beta]{\simeq} \mathcal{M} \xrightarrow[\beta']{\simeq} \mathcal{K}'$, we say $(\alpha, \alpha', S_1, S_2)$ is equivalent to $(\beta, \beta', T_1, T_2)$ if there exist a generalised morphism $\mathcal{L} \xleftarrow[\gamma]{\simeq} \mathcal{N} \xrightarrow[\gamma']{\simeq} \mathcal{M}$ and natural transformations $U_1: \alpha \circ \gamma \Rightarrow \beta \circ \gamma'$ and $U_2: \alpha' \circ \gamma \Rightarrow \beta' \circ \gamma'$ such that

$$(T_1 \gamma') \circ (\varphi U_1) = (\varphi' U_2) \circ (S_1 \gamma) \quad \text{and} \quad (T_2 \gamma') \circ (\psi U_1) = (\psi' U_2) \circ (S_2 \gamma). \quad (4)$$

A **2-cell** from $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ to $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{\simeq} \mathcal{H}$ is defined to be an equivalence class $[\alpha, \alpha', S_1, S_2]$ of the diagrams above. That the relation between diagrams is in fact an equivalence relation follows from our conditions for W to admit a right calculus of fractions; see [**Pro96**, §§2.3] and [**PS21**, Remarks 2.1].

Given a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow{\psi} \mathcal{H}$, the **identity 2-cell** is represented by the diagram




◇

Remark 6.6. One difference between what appears in [Pro96, §§2.3] and **Definition 6.5** is that Pronk only requires $\varphi \circ \alpha$, $\varphi' \circ \alpha'$, $\varphi \circ \beta$, $\varphi' \circ \beta'$, $\varphi \circ \alpha \circ \gamma$, and $\varphi \circ \beta \circ \gamma'$ to be in W . However, this is equivalent in our setting to the definition above due to **Proposition 4.12**.


┘

For fixed diffeological groupoids \mathcal{G} and \mathcal{H} , we require their generalised morphisms with 2-cells between them to form a category. In particular, we need to define the “composition” of 2-cells.

Definition 6.7 (Vertical Composition of Generalised Morphisms). Given representatives of 2-cells as in the following diagrams



define their **vertical composition** to be the equivalence class of the diagram



where $V_1 = (T_1 \text{pr}_3) \circ (\varphi' \text{pr}_2) \circ (S_1 \text{pr}_1)$, and $V_2 = (T_2 \text{pr}_3) \circ (\psi' \text{pr}_2) \circ (S_2 \text{pr}_1)$. It is straightforward to check that this is indeed a 2-cell. We will use \circ to denote vertical composition.

◇

We also need to be able to compose 2-cells across compositions of generalised morphisms. We accomplish this by vertically composing whiskers of 2-cells. (The reader should be warned: there does not seem to be a consensus on which whiskering is left and which is right: diagrammatically, our left/right whiskering occurs on the left/right of the 2-cell; however, notationally, it is written in the opposite order, following functional composition convention.)

Definition 6.8 (Whiskering and Horizontal Composition of 2-Cells). The **left whiskering** of a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{L} \xrightarrow[\psi]{\simeq} \mathcal{H}$ with a 2-cell $[\alpha, \alpha', S_1, S_2]$ represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{M} & & \\
 & \swarrow \chi & \uparrow \alpha & \searrow \omega & \\
 \mathcal{H} & & \mathcal{N} & & \mathcal{K} \\
 & \swarrow \chi' & \downarrow \alpha' & \searrow \omega' & \\
 & & \mathcal{M}' & &
 \end{array}$$

$\Downarrow S_1$ $\Downarrow S_2$

is the 2-cell $[\beta, \beta', \text{id}_{\varphi \circ \text{pr}_1}, S_2 \text{pr}_3]$ represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{L} \overset{w}{\times}_{\psi \chi} \mathcal{M} & & \\
 & \swarrow \varphi \circ \text{pr}_1 & \uparrow \beta & \searrow \omega \circ \text{pr}_3 & \\
 \mathcal{G} & & \mathcal{L} \overset{w}{\times}_{\psi \chi \circ \alpha} \mathcal{N} & & \mathcal{K} \\
 & \swarrow \varphi \circ \text{pr}_1 & \downarrow \beta' & \searrow \omega' \circ \text{pr}_3 & \\
 & & \mathcal{L} \overset{w}{\times}_{\psi \chi'} \mathcal{M}' & &
 \end{array}$$

$\Downarrow S_2 \text{pr}_3$

where $\beta(l, h, n) = (l, h, \alpha(n))$ and $\beta'(l, h, n) = (l, S_1(s(n))h, \alpha'(n))$. That β and β' are weak equivalences follows from the fact that α and α' are. Symmetrically, the **right whiskering** of a generalised morphism $\mathcal{H} \xleftarrow[\chi]{\simeq} \mathcal{M} \xrightarrow[\omega]{\simeq} \mathcal{K}$ with a 2-cell $[\alpha, \alpha', S_1, S_2]$ represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{L} & & \\
 & \swarrow \varphi & \uparrow \alpha & \searrow \psi & \\
 \mathcal{G} & & \mathcal{N} & & \mathcal{H} \\
 & \swarrow \varphi' & \downarrow \alpha' & \searrow \psi' & \\
 & & \mathcal{L}' & &
 \end{array}$$

$\Downarrow S_1$ $\Downarrow S_2$

is the 2-cell $[\beta, \beta', S_1 \text{pr}_1, \text{id}_{\omega \text{opr}_3}]$ represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{L}_{\psi} \times_{\chi}^w \mathcal{M} & & \\
 & \nearrow \varphi \circ \text{pr}_1 & \uparrow \beta & \searrow \omega \circ \text{pr}_3 & \\
 \mathcal{G} & \Downarrow \mu \text{pr}_1 & \mathcal{N}_{\psi \circ \alpha} \times_{\chi}^w \mathcal{M} & \circlearrowleft & \mathcal{K} \\
 & \searrow \varphi' \circ \text{pr}_1 & \downarrow \beta' & \nearrow \omega \circ \text{pr}_3 & \\
 & & \mathcal{L}'_{\psi'} \times_{\chi}^w \mathcal{M} & &
 \end{array}$$

where $\beta(\ell, h, n) = (\alpha(n), h, m)$ and $\beta'(\alpha'(n), h S_2(s(n))^{-1}, m)$.

Finally, the **horizontal composition** of two 2-cells represented by the diagrams below

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \mathcal{L} \\
 & \nearrow \varphi & \uparrow \alpha & \searrow \psi & \nearrow \chi \\
 \mathcal{G} & \Downarrow S_1 & \mathcal{M} & \Downarrow S_2 & \mathcal{H} \\
 & \searrow \varphi' & \downarrow \alpha' & \nearrow \psi' & \searrow \chi' \\
 & & \mathcal{K}' & & \mathcal{L}' \\
 & & & & \nearrow \omega \\
 & & & & \mathcal{J} \\
 & & & & \searrow \omega' \\
 & & & & \mathcal{J}'
 \end{array}$$

is given by the left whiskering of the right 2-cell with the generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$, vertically composed with the right whiskering of the left 2-cell with the generalised morphism $\mathcal{H} \xleftarrow[\chi']{\simeq} \mathcal{L}' \xrightarrow[\omega']{} \mathcal{J}$. Alternatively, one could switch the order of the 2-cells in the vertical composition, using appropriate whiskerings. The end result is independent of the order; see [Pro96, Subsection 2.3] and [PS21, page 15] for details. We will use juxtaposition to denote whiskering and horizontal composition. \diamond

Remark 6.9 (Associator of Generalised Morphisms). In general, composition of 1-cells in a bicategory is not associative. However, it follows from **Item 2** of **Remark 5.7** that the composition of generalised morphisms is an associative operation; that is, the so-called ‘‘associator’’ is trivial. \lrcorner

Definition 6.10 (Unitors for Generalised Morphisms). For a pair of Lie groupoids $(\mathcal{G}, \mathcal{H})$, the **left unitor** $\lambda_{\mathcal{G}, \mathcal{H}}$ is a natural transformation assigning to each generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ the 2-cell represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K} & & \\
 & \nearrow \text{pr}_1 & \parallel & \searrow \psi \circ \text{pr}_3 & \\
 \mathcal{G} & \Downarrow \text{pr}_2 & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K} & \circlearrowleft & \mathcal{H} \\
 & \searrow \varphi & \downarrow \text{pr}_3 \simeq & \nearrow \psi & \\
 & & \mathcal{K} & &
 \end{array}$$

The **right unitor** $\rho_{\mathcal{G}, \mathcal{H}}$ of $(\mathcal{G}, \mathcal{H})$ is a natural transformation assigning to each generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow{\psi} \mathcal{H}$ the 2-cell represented by the diagram

$$\begin{array}{ccccc}
 & & \mathcal{K} \times_{\psi \times \text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H} & & \\
 & \swarrow \varphi \circ \text{pr}_1 & \parallel & \searrow \text{pr}_3 & \\
 \mathcal{G} & & \mathcal{K} \times_{\psi \times \text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H} & & \mathcal{H} \\
 & \swarrow \varphi & \downarrow \text{pr}_1 \simeq & \uparrow \text{pr}_2 & \\
 & & \mathcal{K} & &
 \end{array}$$

◇

We now have all of the ingredients to construct the bicategory of fractions.

Corollary 6.11. *The class W of weak equivalences in \mathbf{DGpoid} induces a bicategory of fractions $\mathbf{DGpoid}[W^{-1}]$, whose objects are diffeological groupoids, 1-cells are generalised morphisms, and 2-cells, vertical and horizontal compositions, and unitors are as defined above.*

Proof. Due to [Theorem 6.2](#), the result is a direct consequence of [[PS21](#), Theorem 3.6] (or, equivalently, [[Pro96](#), Proposition 20]). □

6.3. Spanisation. In the proof of [Corollary 6.11](#), we use [[PS21](#), Theorem 3.6], which describes a pseudofunctor $\mathfrak{S}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}[W^{-1}]$ that sends weak equivalences to internal adjoint equivalences. In a bicategory \mathcal{C} , a 1-cell $f: c_1 \rightarrow c_2$ is an **adjoint equivalence in \mathcal{C}** if there exist a 1-cell $g: c_2 \rightarrow c_1$ and 2-isomorphisms $\varepsilon: \text{id}_{c_1} \Rightarrow g \circ f$ and $\eta: f \circ g \Rightarrow \text{id}_{c_2}$; the 2-isomorphisms are required to satisfy the Triangle Identities (see Equation (1), noting these equations are written for \mathbf{DGpoid} , and not $\mathbf{DGpoid}[W^{-1}]$). We describe this pseudofunctor $\mathfrak{S}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}[W^{-1}]$ now, dubbing it “spanisation”.

Definition 6.12 (Spanisation). The **spanisation** of a functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is the generalised morphism given by $\mathcal{G} \xleftarrow[\text{id}_{\mathcal{G}}]{\simeq} \mathcal{G} \xrightarrow{\varphi} \mathcal{H}$. Let $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$ be functors and $T: \varphi \Rightarrow \psi$ a natural transformation. The **spanisation** of T is the 2-cell $[\text{id}_{\mathcal{G}}, \text{id}_{\mathcal{G}}, \text{id}_{\text{id}_{\mathcal{G}}}, T]$. ◇

As mentioned before, the purpose of constructing $\mathbf{DGpoid}[W^{-1}]$ is to formally invert weak equivalences. How is this done in our context? If φ is a weak equivalence, then its spanisation is $\mathcal{G} \xleftarrow[\text{id}_{\mathcal{G}}]{\simeq} \mathcal{G} \xrightarrow{\varphi} \mathcal{H}$, where both functors are weak equivalences. A **quasi-inverse** is given by $\mathcal{H} \xleftarrow[\varphi]{\simeq} \mathcal{G} \xrightarrow[\text{id}_{\mathcal{G}}]{\simeq} \mathcal{G}$; that is, there is a 2-cell from the composition of the spanisation of φ with its quasi-inverse to the spanisation of the identity functor of \mathcal{G} , as well as a 2-cell from the reversed composition with the quasi-inverse to the spanisation of the identity functor of \mathcal{H} ; these equivalences satisfy the Triangle Identities. This is a consequence of the following more general result, which is proved in [[Pro96](#), Proposition 20].

Lemma 6.13 (Morita Equivalence). *A generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ in which ψ is a weak equivalence has a quasi-inverse $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{K} \xrightarrow[\varphi]{\simeq} \mathcal{G}$.*

We say that \mathcal{G} and \mathcal{H} are **Morita equivalent** if such a generalised morphism exists [Moe02]. A diagram representing the 2-cell between the composition of $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ and $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{K} \xrightarrow[\varphi]{\simeq} \mathcal{G}$, and the spanisation of $\text{id}_{\mathcal{G}}$, is given by

$$\begin{array}{ccccc}
 & & \mathcal{K}_{\psi} \times_{\psi}^w \mathcal{K} & & \\
 & \swarrow \varphi \circ \text{pr}_1 & \parallel & \searrow \varphi \circ \text{pr}_3 & \\
 \mathcal{G} & \xrightarrow{\simeq} & \mathcal{K}_{\psi} \times_{\psi}^w \mathcal{K} & \xrightarrow{\downarrow S} & \mathcal{G} \\
 & \searrow \varphi \circ \text{pr}_1 & \downarrow \simeq & \swarrow & \\
 & & \mathcal{G} & &
 \end{array}$$

where $S(y_1, h, y_2) := \varphi \circ \Phi_{\psi}^{-1}(y_1, y_2, h)$. It follows from **Lemma 6.13**, **Item 1** of **Proposition 5.10**, and **Proposition 4.12** that Morita equivalence is an equivalence relation on diffeological groupoids. Moreover, any generalised morphism admitting a quasi-inverse is of the form as in **Lemma 6.13**; this follows from **Subsection 10.1**, although we will not need this result. And so, we have reached our goal: weak equivalences become invertible in $\mathbf{DGpoid}[W^{-1}]$. In the following section, we construct another bicategory which serves as a smaller localisation of \mathbf{DGpoid} at weak equivalences. We end this section with some examples.

Example 6.14 (Principal G -Bundles III). By **Example 4.6**, for a Lie group G , and two principal G -bundles $\pi_i: P_i \rightarrow B$ ($i = 1, 2$) over the same manifold B , the action groupoids $G \times P_1$ and $G \times P_2$ are Morita equivalent to the trivial groupoid $B \rightrightarrows B$, and hence Morita equivalent to each other. //

Example 6.15. [Nebulaic Groupoids III] By **Example 4.7**, given a diffeological space X and two generating families \mathcal{F}_1 and \mathcal{F}_2 of the diffeology, the nebulaic groupoids $\mathcal{N}(\mathcal{F}_1)$ and $\mathcal{N}(\mathcal{F}_2)$ are Morita equivalent to each other, and by **Example 4.10** to the trivial groupoid of X . //

Example 6.16 (Nebulaic Groupoids of Quotients). Let X be a diffeological space, \sim an equivalence relation, $\pi: X \rightarrow X/\sim$ the quotient map, and \mathcal{D}_X the diffeology on X . Let \mathcal{F} be the generating family of the diffeology of X/\sim made up of plots $\pi \circ p$ where $p \in \mathcal{D}_X$. Define a functor $\text{ev}_X: \mathcal{N}(\mathcal{F}) \rightarrow R$, where R is the relation groupoid of \sim (see **Example 3.5**) and $\mathcal{N}(\mathcal{F})$ is the nebulaic groupoid of \mathcal{F} (see **Example 4.7**), as follows: given $u \in U_{\pi \circ p} \subseteq \mathcal{N}(\mathcal{F})_0$, set $(\text{ev}_X)_0(u) := p(u)$, and $(\text{ev}_X)_1(u_1, u_2) := ((\text{ev}_X)_0(u_1), (\text{ev}_X)_0(u_2))$. Then $(\text{ev}_X)_0$ is a subduction and Φ_{ev_X} is a diffeomorphism. Thus ev_X is a subductive weak equivalence by **Lemma 4.8**, and so $\mathcal{N}(\mathcal{F})$ and R are Morita equivalent. By **Example 6.15**, since \mathcal{F} and $\mathcal{D}_{X/\sim}$ are both generating families of $\mathcal{D}_{X/\sim}$, the diffeological groupoids $\mathcal{N}(\mathcal{D}_{X/\sim})$ and $\mathcal{N}(\mathcal{F})$ are Morita equivalent. Finally, by

Example 4.10, $\mathcal{N}(\mathcal{D}_{X/\sim})$ is Morita equivalent to the trivial groupoid of X . We conclude that all of R , X , $\mathcal{N}(\mathcal{F})$, and $\mathcal{N}(\mathcal{D}_{X/\sim})$ are Morita equivalent. One can interpret this to mean that all of the diffeological information contained in the quotient diffeology on X/\sim is captured by the relation groupoid R . //

7. ANAFUNCTORS

The purpose of this section is to optimise the localisation $\mathbf{DGpoid}[W^{-1}]$ so that it is smaller, by inverting a smaller class of arrows. The benefit is that 2-cells in this setting are natural transformations, which are special cases of representatives of the 2-cells in $\mathbf{DGpoid}[W^{-1}]$. These natural transformations between two 1-cells form a set, whereas representatives of a 2-cell between generalised morphisms do not. To this end, following [Rob12; Rob21], we begin by defining a so-called singleton strict pretopology on \mathbf{DGpoid} , which replaces the requirement that W admit a right calculus of fractions; this is the content of [Subsection 7.1](#). We then construct the corresponding bicategory $\mathbf{DGpoid}_{\text{ana}}$ in [Subsection 7.2](#). We then construct a pseudofunctor ‘‘anafunctorisation’’ from \mathbf{DGpoid} to $\mathbf{DGpoid}_{\text{ana}}$, similar to spanisation, in [Subsection 7.3](#). We then show that $\mathbf{DGpoid}_{\text{ana}}$ is equivalent to $\mathbf{DGpoid}[W^{-1}]$ in [Subsection 7.4](#). Finally, we show that passing from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DGpoid}_{\text{ana}}$ ‘‘optimises’’ 2-cells in [Subsection 7.5](#).

7.1. Subductive Weak Equivalences Induce a 2-Site.

Definition 7.1 (Singleton Strict Pretopology). Let \mathcal{C} be a 2-category. A class of 1-cells J of \mathcal{C} forms a **singleton strict pretopology** on \mathcal{C} if J contains all identity arrows, and is closed under composition and base change; the latter more specifically means that if $c \xrightarrow{f} d$ is in J , and $b \xrightarrow{g} d$ is a 1-cell, then $b_g \times_f c$ exists and $b_g \times_f c \xrightarrow{\text{pr}_1} b$ is in J . Furthermore, we say $c \xrightarrow{f} d \in J$ is **representably fully faithful** if for any 1-cells $b \xrightarrow{g} c$, $b \xrightarrow{h} c$, and 2-cell $A: f \circ g \Rightarrow f \circ h$, there exists a unique $A': g \Rightarrow h$ so that $fA' = A$. Similarly, we say $c \xrightarrow{f} d \in J$ is **co-fully faithful** if for any 1-cells $d \xrightarrow{g} e$, $d \xrightarrow{h} e$ and 2-cell $A: g \circ f \Rightarrow h \circ f$, there exists a unique $A': g \Rightarrow h$ so that $A'f = A$. Following [Rob21], if all elements of J are representably fully faithful and co-fully faithful, we call the pair (\mathcal{C}, J) a **2-site**.

As mentioned in [Rob21], in the literature, 2-sites have a more general definition; we use the above one for convenience as it suits the purposes of this paper. We now prove that subductive weak equivalences are co-fully faithful, which will allow us to construct a 2-site structure using these.

Lemma 7.2 (Co-Fully Faithfulness). *Subductive weak equivalences are co-fully faithful.*

Proof. Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a subductive weak equivalence, $\psi, \psi': \mathcal{H} \rightarrow \mathcal{K}$ functors, and $S: \psi \circ \varphi \Rightarrow \psi' \circ \varphi$ a natural transformation. Define $S': \mathcal{H}_0 \rightarrow \mathcal{K}_1$ by $S'(y) = S(x)$ where $x \in \varphi_0^{-1}(y)$. To prove this is well-defined, suppose $\varphi(x_1) = \varphi(x_2)$. Since φ is a weak equivalence, there exists $g = \Phi_\varphi^{-1}((x_1, x_2), u_{\varphi(x_1)})$ from x_1 to x_2 , inducing the

commutative diagram

$$\begin{array}{ccc} \psi \circ \varphi(x_1) & \xrightarrow{S_{x_1}} & \psi' \circ \varphi(x_1) \\ \downarrow \psi \circ \varphi(g) & & \downarrow \psi' \circ \varphi(g) \\ \psi \circ \varphi(x_2) & \xrightarrow{S_{x_2}} & \psi' \circ \varphi(x_2). \end{array}$$

However, since $\varphi(g) = u_{\varphi(x_1)}$, we have $S_1(x_1) = S_1(x_2)$. This shows that S' is well-defined on the image of φ , which is \mathcal{H}_0 since φ is subductive. By construction, $S = S'\varphi$.

To show that S' is smooth, let p be a plot of \mathcal{H}_0 be a plot. Since φ is subductive, after shrinking U_p , there exists a lift q of p to \mathcal{G}_0 . Then $S' \circ p = S \circ q$, the latter of which is a plot of \mathcal{K}_1 .

That S' is a natural transformation follows from the fact that φ is a subductive weak equivalence, and that S is a natural transformation. Uniqueness follows from the construction. \square

At this point it is worth pausing and mentioning that while natural transformations preserve weak equivalences (by Proposition 4.13), this is no longer true of subductive weak equivalences, as the following example illustrates. However, natural transformations do preserve co-fully faithfulness, as the lemma below proves. Its corollary will be important in the definition of horizontal composition later in this section.

Example 7.3 (Natural Transformations do not Preserve Subductive Weak Equivalences). Let $\mathcal{G} = \mathcal{H} = (\mathbb{R} \times \mathbb{R} \rightrightarrows \mathbb{R})$, let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be the identity functor, and let $\psi: \mathcal{G} \rightarrow \mathcal{H}$ be the unique functor such that $\psi_0(x) = 0$ for all $x \in \mathbb{R}$ and $\psi_1(x_1, x_2) = (0, 0)$ for all $(x_1, x_2) \in \mathbb{R} \times \mathbb{R}$. There is a natural transformation $S: \varphi \Rightarrow \psi$ sending x to $(x, 0)$. While φ is a subductive weak equivalence, ψ is a weak equivalence that is not subductive. \parallel

Lemma 7.4 (Natural Transformations Preserve Co-Fully Faithfulness). *Let $\varphi, \varphi': \mathcal{G} \rightarrow \mathcal{H}$ be functors and $T: \varphi \Rightarrow \varphi'$ a natural transformation. If one of φ or φ' is co-fully faithful, then so is the other.*

Proof. Without loss of generality, suppose φ' is co-fully faithful. Fix $\psi, \psi': \mathcal{H} \rightarrow \mathcal{K}$ and a natural transformation $S: \psi \circ \varphi \Rightarrow \psi' \circ \varphi$. Since φ' is co-fully faithful, the vertical composition $(\psi'T) \circ S \circ (\psi T^{-1})$ factors uniquely as $S'\varphi'$. Thus $S = (\psi'T^{-1}) \circ (S'\varphi') \circ (\psi T)$. Since horizontal composition of natural transformation as the vertical composition of whiskerings is independent of the order of composition taken, we have $S = (S'\varphi) \circ (\psi T^{-1}) \circ (\psi T) = S'\varphi$. Uniqueness follows from the uniqueness of S' above. \square

The following corollary is [Rob21, Lemma 2.14], but with a different proof.

Corollary 7.5. *If $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ are subductive weak equivalences, and $\chi: \mathcal{G} \rightarrow \mathcal{H}$ is a functor such that $\varphi = \psi \circ \chi$, then χ is a co-fully faithful weak equivalence.*

Proof. That χ is a weak equivalence follows from [Proposition 4.12](#). The trivial natural transformation between $\psi \circ \text{pr}_2$ and $\varphi \circ \text{pr}_1 = \psi \circ \chi \circ \text{pr}_1$ from $\mathcal{G}_{\varphi \times \psi} \mathcal{H}$ to \mathcal{K} factors uniquely as ψT for a natural transformation $T: \text{pr}_2 \Rightarrow \chi \circ \text{pr}_1$, by [Lemma 4.5](#). By [Item 2](#) or [Proposition 5.4](#) and [Lemma 7.2](#), pr_2 is co-fully faithful, and thus so is $\chi \circ \text{pr}_1$ by [Lemma 7.4](#). Suppose $\omega, \omega': \mathcal{H} \rightarrow \mathcal{L}$ are functors and $S: \omega \circ \chi \Rightarrow \omega' \circ \chi$ is a natural transformation. Since $\chi \circ \text{pr}_1$ is co-fully faithful, $S \text{pr}_1$ factors uniquely as $S'(\chi \circ \text{pr}_1)$ for a natural transformation $S' \circ \omega \Rightarrow \omega'$. Since pr_1 is surjective, it follows that $S'\chi = S$. Thus χ is co-fully faithful. \square

We are now ready to equip **DGpoid** with a 2-site structure:

Theorem 7.6 (**DGpoid** is a 2-Site). **DGpoid** equipped with the class J of all subductive weak equivalences is a 2-site.

Proof. It is immediate that identity functors are contained in J . That J is closed under composition follows from [Proposition 4.12](#) and [Item 1](#) of [Proposition 2.7](#). That J is closed under base change follows from [Item 2](#) of [Proposition 5.4](#). That elements of J are representably fully faithful follows from [Lemma 4.5](#), and that they are co-fully faithful is [Lemma 7.2](#). \square

As an example of how the perspective just introduced can be used, we show how smooth essential surjectivity of a weak equivalence can be recast in terms of elements of J .

Lemma 7.7 (Locally Split). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a smoothly fully faithful functor. Then φ is a weak equivalence if and only if there is a subductive weak equivalence $\psi: \mathcal{K} \rightarrow \mathcal{H}$, a functor $\chi: \mathcal{K} \rightarrow \mathcal{G}$, and a natural transformation $S: \varphi \circ \chi \Rightarrow \psi$. (This latter condition is called *J-locally split*.)*

Proof. Suppose φ is a weak equivalence. Choose $\mathcal{K} := \mathcal{G}_{\varphi \times \text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H}$, $\psi = \text{pr}_3$, $\chi = \text{pr}_1$, and $S = \text{pr}_2$. Then ψ is a subductive weak equivalence by [Item 1](#) of [Proposition 5.10](#).

Conversely, suppose there exist a subductive weak equivalence $\psi: \mathcal{K} \rightarrow \mathcal{H}$, a functor $\chi: \mathcal{K} \rightarrow \mathcal{G}$, and a natural transformation $S: \varphi \circ \chi \Rightarrow \psi$. Let $p = y_u$ be a plot of \mathcal{H}_0 . Since ψ_0 is a subduction, after shrinking U_p , there is a lift z_u of y_u to \mathcal{K}_0 . Then $(\chi(z_u), S(z_u)^{-1})$ is the desired lift of y_u to \mathcal{G}_0 , implying that Ψ_{φ} is a subduction. \square

7.2. The Anafunctor Bicategory $\text{DGpoid}_{\text{ana}}$. Following [\[Rob21\]](#), we now construct a new bicategory of diffeological groupoids whose 1-cells are so-called ‘‘anafunctors’’. Anafunctors originally were introduced by Makkai [\[Mak96\]](#) in order to allow one to discuss 1-cells in a 2-category without using the axiom of choice. Bartels [\[Bar06\]](#) develops the theory further in terms of internal categories.

Definition 7.8 (Anafunctor). A generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \rightarrow \mathcal{H}$ is an **anafunctor** if φ is a subductive weak equivalence. The **identity anafunctor** of a diffeological groupoid \mathcal{G} is the same as the identity generalised morphism: $\mathcal{G} \xleftarrow{=} \mathcal{G} \xrightarrow{=} \mathcal{G}$. \diamond

The composition of two generalised morphisms is again a generalised morphism due to the use of the weak pullback in its definition. We now define the composition of anafunctors similarly, except that we only need to use the strict pullback.

Definition 7.9 (Composition of Anafunctors). Let $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{L} \xrightarrow[\psi]{} \mathcal{H}$ and $\mathcal{H} \xleftarrow[\chi]{\simeq} \mathcal{M} \xrightarrow[\omega]{} \mathcal{K}$ be anafunctors. Define their **composition** to be $\mathcal{G} \xleftarrow[\varphi \circ \text{pr}_1]{\simeq} \mathcal{L} \times_{\psi, \chi} \mathcal{M} \xrightarrow[\omega \circ \text{pr}_2]{} \mathcal{K}$. That the result is an anafunctor follows from **Item 2** of **Proposition 5.4**, **Proposition 4.12**, and **Item 1** of **Proposition 2.7**. \diamond

The 2-cells of the bicategory are certain natural transformations, which immediately makes the new bicategory we are constructing friendlier than $\mathbf{DGpoid}[W^{-1}]$.

Definition 7.10 (Transformation). Given anafunctors $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{} \mathcal{H}$, a **transformation** between them is a natural transformation

$$\begin{array}{ccc}
 & \mathcal{K} \times_{\varphi, \varphi'} \mathcal{K}' & \\
 \text{pr}_1 \swarrow & & \searrow \text{pr}_2 \\
 \mathcal{K} & \xrightarrow[S]{} & \mathcal{K}' \\
 \psi \searrow & & \swarrow \psi' \\
 & \mathcal{H} &
 \end{array}$$

The **identity transformation** of an anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ is given by the natural transformation


$$I_{\mathcal{G} \leftarrow \mathcal{K} \rightarrow \mathcal{H}} : (\mathcal{K} \times_{\varphi, \varphi} \mathcal{K})_0 \rightarrow \mathcal{H}_1 : (y_1, y_2) \mapsto \psi(\Phi_{\varphi}^{-1}(y_1, y_2, u_{\varphi(y_1)})).$$

\diamond

We now describe the vertical composition, horizontal composition, and unitors of our bicategory.

Definition 7.11 (Vertical Composition of Transformations). Let $\mathcal{G} \xleftarrow[\varphi_i]{\simeq} \mathcal{K}_i \xrightarrow[\psi_i]{} \mathcal{H}$ be anafunctors ($i = 1, 2, 3$), and let S be a transformation between the first and second, and T a transformation between the second and third. Define the **vertical composition** to be the unique natural transformation $T \circ S$ so that $(T \text{pr}_{23}) \circ (S \text{pr}_{12}) = (T \circ S) \text{pr}_{13}$,

where pr_{12} , pr_{13} , and pr_{23} are the projections as indicated in the diagram below:



That such a natural transformation $T \circ S$ exists is due to the co-fully faithfulness of $\text{pr}_{13} \in J$. We will use \circ to denote vertical composition. \diamond


Before continuing, note that in the definition of vertical composition above, to compute $(T \circ S)(y_1, y_3)$, one uses the formula $(T\text{pr}_{23}) \circ (S\text{pr}_{12}) = (T \circ S)\text{pr}_{13}$. But this requires introducing an auxiliary object $y_2 \in \mathcal{K}_2$ such that $(y_1, y_2, y_3) \in \mathcal{K}_{1\varphi_1} \times_{\varphi_2} \mathcal{K}_{2\varphi_2} \times_{\varphi_3} \mathcal{K}_3$ and computing

$$(T \circ S)(y_1, y_3) = T(y_2, y_3)S(y_1, y_2).$$

This is well-defined, as the result is independent of the choice of y_2 (and such a y_2 exists since pr_{13} is subductive). See [Rob21, page 191 and Proposition 3.8] for details.

Next, we again define horizontal composition using whiskering and vertical composition. Left whiskering, however, requires its definition to be broken down further.

Definition 7.12 (Whiskering and Horizontal Composition of Transformations). The **left whiskering** of an anafunctor of the form $\mathcal{L} \xleftarrow{\omega} \mathcal{L} \xrightarrow{\omega'} \mathcal{H}$ with a transformation ψ



in the diagram

$$\begin{array}{ccccc}
 & & \mathcal{M}_\omega \times_\xi \mathcal{J} & & \\
 & \swarrow \chi^{\text{opr}_1} & \uparrow \simeq & \searrow \zeta^{\text{opr}_2} & \\
 \mathcal{H} & \circlearrowleft & (\mathcal{M}_\omega \times_\chi \mathcal{J})_{\chi^{\text{opr}_1}} \times_{\chi'^{\text{opr}_1}} (\mathcal{M}'_{\omega'} \times_\xi \mathcal{J}) & \Downarrow T & \mathcal{J} \\
 & \swarrow \chi'^{\text{opr}_1} & \downarrow \simeq & \searrow \zeta^{\text{opr}_2} & \\
 & & \mathcal{M}'_{\omega'} \times_\xi \mathcal{J} & &
 \end{array}$$

The **horizontal composition** of the two transformations as in the diagrams

$$\begin{array}{ccccc}
 & & \mathcal{L} & & \mathcal{M} \\
 & \swarrow \varphi & \uparrow \simeq & \searrow \psi & \\
 \mathcal{G} & \circlearrowleft & \mathcal{L}_{\varphi \times \varphi'} \mathcal{L}' & \Downarrow S & \mathcal{H} \\
 & \swarrow \varphi' & \downarrow \simeq & \searrow \psi' & \\
 & & \mathcal{L}' & & \mathcal{M}' \\
 & & & & \uparrow \simeq \\
 & & & & \mathcal{M}_\chi \times_{\chi'} \mathcal{M}' \\
 & & & & \downarrow \simeq \\
 & & & & \mathcal{M}' \\
 & & & & \uparrow \chi \\
 & & & & \mathcal{M} \\
 & & & & \searrow \omega \\
 & & & & \mathcal{K} \\
 & & & & \uparrow \omega' \\
 & & & & \mathcal{M}' \\
 & & & & \downarrow \chi' \\
 & & & & \mathcal{H} \\
 & & & & \swarrow \omega' \\
 & & & & \mathcal{K}
 \end{array}$$

is given by the left whiskering of T with the anafunctor $\mathcal{G} \xleftarrow[\psi]{\simeq} \mathcal{L} \xrightarrow[\psi]{\simeq} \mathcal{H}$, vertically composed with the right whiskering of S with the anafunctor $\mathcal{H} \xleftarrow[\chi']{\simeq} \mathcal{M}' \xrightarrow[\omega']{\simeq} \mathcal{K}$. One could also vertically compose with the order S and T switched, using appropriate whiskerings, and obtain the same result [Rob21, Lemma 3.15]. We will use juxtaposition to denote whiskering and horizontal composition. \diamond

Remark 7.13 (Associator for Anafunctors). Similar to the case of $\mathbf{DGpoid}[W^{-1}]$, it follows from **Item 2** of **Remark 5.7** that the composition of anafunctors is associative. \lrcorner

Composing an anafunctor with an identity anafunctor does not return the original anafunctor, but there is a transformation between the two.

Definition 7.14 (Unitors for Anafunctors). The **left unitor** $\lambda_{\mathcal{G}, \mathcal{H}}^{\text{ana}}$ of a pair of diffeological groupoids $(\mathcal{G}, \mathcal{H})$ is the natural transformation assigning to each anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ the transformation

$$\lambda_{\mathcal{G}, \mathcal{H}}^{\text{ana}}(\mathcal{G} \leftarrow \mathcal{K} \rightarrow \mathcal{H}): \left((\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi} \mathcal{K})_{\text{pr}_1} \times_{\varphi} \mathcal{K} \right)_0 \rightarrow \mathcal{H}_1: ((x, y_1), y_2) \mapsto \Phi_{\varphi}^{-1}(y_1, y_2, u_x).$$

The **right unitor** $\rho_{\mathcal{G}, \mathcal{H}}^{\text{ana}}$ of a pair of diffeological groupoids $(\mathcal{G}, \mathcal{H})$ is the natural transformation assigning to each anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ the transformation

$$\rho_{\mathcal{G}, \mathcal{H}}^{\text{ana}}(\mathcal{G} \leftarrow \mathcal{H} \rightarrow \mathcal{K}): \left((\mathcal{K}_{\psi \times \text{id}_{\mathcal{H}}} \mathcal{H})_{\varphi \text{opr}_1} \times_{\varphi} \mathcal{K} \right)_0 \rightarrow \mathcal{H}_1: ((y_1, z), y_2) \mapsto \Phi_{\varphi}^{-1}(y_1, y_2, u_{\varphi(y_1)}).$$

\diamond

We are now ready for our new bicategory.

Corollary 7.15. *The class J of subductive weak equivalences in \mathbf{DGpoid} induces a bicategory $\mathbf{DGpoid}_{\text{ana}}$, whose objects are diffeological groupoids, 1-cells are anafunctors, 2-cells are transformations between anafunctors, and vertical composition, horizontal composition, and unitors are as defined above.*

Proof. By [Theorem 7.6](#), we may apply [[Rob21](#), Proposition 3.20] to our setting, which yields the result. \square

Example 7.16 (Nebulaic Groupoids IV). Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a functor and \mathcal{F} a generating family of the diffeology of \mathcal{G}_0 . Define \mathcal{U} to be the base change of \mathcal{G} along $\text{ev}: \mathcal{N}(\mathcal{F})_0 \rightarrow \mathcal{G}_0$. Then $\mathcal{G} \xleftarrow[\varepsilon]{\simeq} \mathcal{U} \xrightarrow[\varphi \circ \varepsilon]{\simeq} \mathcal{H}$ is an anafunctor, where $\varepsilon_0 = \text{ev}$ and $\varepsilon_1 = \text{pr}_3$. Indeed, ε is smoothly fully faithful, and ev a subduction; thus ε is a subductive weak equivalence by [Lemma 4.8](#).

If \mathcal{F}' is another generating family of the diffeology of \mathcal{G}_0 , with \mathcal{U}' the corresponding base change and ε' the corresponding subductive weak equivalence, then

$$\varphi \circ \varepsilon \circ \text{pr}_1, \varphi \circ \varepsilon' \circ \text{pr}_2: \mathcal{U}_{\varepsilon} \times_{\varepsilon'} \mathcal{U}' \rightarrow \mathcal{H}$$

are equal, and so we get a (trivial) transformation between $\mathcal{G} \xleftarrow[\varepsilon]{\simeq} \mathcal{U} \xrightarrow[\varphi \circ \varepsilon]{\simeq} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\varepsilon']{\simeq} \mathcal{U}' \xrightarrow[\varphi \circ \varepsilon']{\simeq} \mathcal{H}$. This fact is important later when considering Lie groupoids and (diffeological) Morita equivalence. \parallel

7.3. Anafunctisation. It turns out that spanisation already places functors and natural transformations from \mathbf{DGpoid} into the anafunctor setting.

Example 7.17 (Spanisation and Anafunctors). The spanisation of any functor between diffeological groupoids is an anafunctor. Moreover, given a natural transformation $S: \varphi \Rightarrow \psi$ in \mathbf{DGpoid} , its spanisation induces a transformation between the spanisations of φ and ψ (see [Definition 6.12](#)). \parallel

However, there is another way to associate an anafunctor to a functor and a transformation besides spanisation, as suggested by [[Li15](#)]:

Definition 7.18 (Anafunctisation). Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a functor. Define the **anafunctisation** of φ to be the anafunctor $\mathcal{G} \xleftarrow[\text{pr}_1]{\simeq} \mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H} \xrightarrow[\text{pr}_3]{\simeq} \mathcal{H}$. By [Item 1](#) of [Proposition 5.10](#), this is an anafunctor. Given another functor $\psi: \mathcal{G} \rightarrow \mathcal{H}$ with natural transformation $S: \varphi \Rightarrow \psi$, we obtain the following transformation between the anafunctisations of φ

and ψ , where $\tilde{S}((x, h_1, y_1), (x, h_2, y_2)) := h_2 S(x) h_1^{-1}$:

$$\begin{array}{ccccc}
 & & (\mathcal{G}_\varphi \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H})_{\text{pr}_1} \times_{\text{pr}_1} (\mathcal{G}_\psi \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H}) & & \\
 & \swarrow \text{pr}_1 & & \searrow \text{pr}_2 & \\
 & \simeq & & \simeq & \\
 \mathcal{G}_\varphi \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H} & & \xrightarrow{\tilde{S}} & & \mathcal{G}_\psi \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H} \\
 & \searrow \text{pr}_3 & & \swarrow \text{pr}_3 & \\
 & & \mathcal{H} & &
 \end{array}$$

◇

At this point, we should check how quasi-inverses behave in $\mathbf{DGpoid}_{\text{ana}}$. In particular, we are interested in the anafunctorisation of a subductive weak equivalence having a quasi-inverse. This result is implied by the following.

Lemma 7.19 (Anafunctors and Morita Equivalence). *An anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ in which ψ is a subductive weak equivalence has a quasi-inverse $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{K} \xrightarrow[\varphi]{\simeq} \mathcal{G}$.*

Proof. To prove this, we must show that the identity anafunctor of \mathcal{G} has a transformation to the composition of $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ with $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{K} \xrightarrow[\varphi]{\simeq} \mathcal{G}$. Such a transformation is given by

$$S: (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi \circ \text{pr}_1} (\mathcal{K}_\psi \times_\psi \mathcal{K}))_0 \rightarrow \mathcal{G}_1$$

where

$$S(x, (y_1, y_2)) := \varphi(\Phi_\psi^{-1}(y_1, y_2, \mathbf{u}_\psi(y_1))).$$

Similarly, we require a transformation from the composition of $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{K} \xrightarrow[\varphi]{\simeq} \mathcal{G}$ with $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ to the identity anafunctor of \mathcal{H} . Such a transformation is given by

$$T: ((\mathcal{K}_\varphi \times_\varphi \mathcal{K})_{\psi \circ \text{pr}_1} \times_{\text{id}_{\mathcal{H}}} \mathcal{H})_0 \rightarrow \mathcal{H}_1$$

where

$$T((y_1, y_2), z) := \psi(\Phi_\varphi^{-1}(y_1, y_2, \mathbf{u}_\varphi(y_1)))^{-1}.$$

We must verify the Triangle Identities. The first is that $I_{\mathcal{G} \leftarrow \mathcal{K} \rightarrow \mathcal{H}}$ is equal to the vertical composition of T whiskered on the left by $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ with S whiskered on the right by $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$. Unravelling definitions, we find that T whiskered on the left by

$\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ yields the transformation T' in the diagram

$$\begin{array}{ccccc}
& & \mathcal{K}_\psi \times_\psi \mathcal{K}_\varphi \times_\varphi \mathcal{K} & & \\
& \swarrow \varphi \circ \text{pr}_1 & \uparrow \simeq & \searrow \psi \circ \text{pr}_3 & \\
\mathcal{G} & \xleftarrow[\simeq]{\circlearrowleft} & (\mathcal{K}_\psi \times_\psi \mathcal{K}_\varphi \times_\varphi \mathcal{K})_{\varphi \circ \text{pr}_1} \times_{\varphi \circ \text{pr}_1} (\mathcal{K}_\psi \times_{\text{id}_{\mathcal{H}}} \mathcal{H}) & \xrightarrow[\simeq]{\Downarrow T'} & \mathcal{H} \\
& \swarrow \varphi \circ \text{pr}_1 & \downarrow \simeq & \searrow \text{pr}_2 & \\
& & \mathcal{K}_\psi \times_{\text{id}_{\mathcal{H}}} \mathcal{H} & &
\end{array}$$

where

$$T'((y_1, y_2, y_3), (y'_1, \psi(y'_1))) = \psi \left(\Phi_\varphi^{-1}(y_1, y'_1, u_{\varphi(y_1)}) \right) \psi \left(\Phi_\varphi^{-1}(y_3, y_2, u_{\varphi(y_2)}) \right).$$

On the other hand, S whiskered on the right by $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ yields the transformation S' in the diagram

$$\begin{array}{ccccc}
& & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_\varphi \mathcal{K} & & \\
& \swarrow \text{pr}_1 & \uparrow \simeq & \searrow \psi \circ \text{pr}_2 & \\
\mathcal{G} & \xleftarrow[\simeq]{\circlearrowleft} & (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_\varphi \mathcal{K})_{\text{pr}_1} \times_{\varphi \circ \text{pr}_1} (\mathcal{K}_\psi \times_\psi \mathcal{K}_\varphi \times_\varphi \mathcal{K}) & \xrightarrow[\simeq]{\Downarrow S'} & \mathcal{H} \\
& \swarrow \varphi \circ \text{pr}_1 & \downarrow \simeq & \searrow \psi \circ \text{pr}_3 & \\
& & \mathcal{K}_\psi \times_\psi \mathcal{K}_\varphi \times_\varphi \mathcal{K} & &
\end{array}$$

where

$$S'((\varphi(y'_1), y'_1), (y_1, y_2, y_3)) = \psi \left(\Phi_\varphi^{-1}(y'_1, y_3, \varphi(\Phi_\psi^{-1}(y_1, y_2, u_{\psi(y_1)}))) \right).$$

Taking the vertical composition $T' \circ S'$ yields $\psi \left(\Phi_\varphi^{-1}(y_1, y_2, u_{\varphi(y_1)}) \right)$, which is precisely $I_{\mathcal{G} \leftarrow \mathcal{K} \rightarrow \mathcal{H}}$.

A similar computation yields the second triangle inequality, completing the proof. \square

In fact, we can do better: by **Item 1** of **Proposition 5.10**, the anafunctorisation of *any* weak equivalence is an anafunctor of the form appearing in **Lemma 7.19**. We thus have:

Corollary 7.20. *The anafunctorisation of a weak equivalence has a quasi-inverse.*

In light of **Lemma 6.13**, we refer to diffeological groupoids admitting an anafunctor which has a quasi-inverse as **Morita equivalent**. This is justified by the following subsection.

7.4. $\mathbf{DGpoid}[W^{-1}]$ versus $\mathbf{DGpoid}_{\text{ana}}$. We now have need for a formal definition of a localisation.

Definition 7.21 (Localisation). A pseudofunctor $F: \mathcal{B} \rightarrow \mathcal{C}$ between bicategories \mathcal{B} and \mathcal{C} is a **localisation** of \mathcal{B} at a class of 1-cells W of \mathcal{B} if F sends elements of W to adjoint equivalences in \mathcal{C} , and is universal with respect to this property; that is, any pseudofunctor $\mathcal{B} \rightarrow \mathcal{A}$ that sends elements of W to adjoint equivalences must factor through F . In

fact, there must be an equivalence of bicategories $\text{Bicat}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Bicat}_W(\mathcal{B}, \mathcal{A})$, where the subscript W indicates the elements of W are being sent to adjoint equivalences (see [Pro96, Theorem 21]). \diamond

We have the following theorem of Roberts [Rob21, Theorem 3.24]:

Theorem 7.22. *Spanisation $\mathbf{DGpoid} \rightarrow \mathbf{DGpoid}_{\text{ana}}$, sending objects to themselves and everything else to their spanisations, is a localisation of W .*

We wish to prove the same result, but with spanisation replaced with anafunctorisation. There are two reasons for this: first, it is not immediately clear what the resulting equivalence of bicategories from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DGpoid}_{\text{ana}}$ is if one is to use spanisation: for instance, given a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$, what is the corresponding anafunctor? One may try taking the spanisation of φ and ψ , compose, and obtain $\mathcal{G} \xleftarrow[\varphi \circ \text{pr}_1]{\simeq} \mathcal{K}_{\text{id}_{\mathcal{K}}} \times_{\text{id}_{\mathcal{K}}} \mathcal{K} \xrightarrow[\psi \circ \text{pr}_2]{\simeq} \mathcal{H}$. However, this is still not an anafunctor (in fact, up to canonical isomorphism $\mathcal{K} \rightarrow \mathcal{K}_{\text{id}_{\mathcal{K}}} \times_{\text{id}_{\mathcal{K}}} \mathcal{K}$, this is the original generalised morphism). Roberts addresses this in [Rob12, Lemma 5.15] using the fact that a weak equivalence is J -locally split (see Lemma 7.7). Instead, one can take the composition of anafunctorisations of φ and ψ , and the result is an anafunctor; this process also preserves quasi-inverses as in Lemma 6.13 and Lemma 7.19 between the two bicategories (see Item 3 of Remark 7.24). What appears to be a discrepancy between the two methods is in fact captured by the equivalence of bicategories $\mathfrak{J}: \text{Bicat}_W(\mathbf{DGpoid}, \mathbf{DGpoid}_{\text{ana}}) \rightarrow \text{Bicat}(\mathbf{DGpoid}[W^{-1}], \mathbf{DGpoid}_{\text{ana}})$: this implies that there is a *pseudo-natural transformation* between the pseudofunctor spanisation to $\mathbf{DGpoid}_{\text{ana}}$, denoted $\mathfrak{S}_{\text{ana}}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}_{\text{ana}}$, and the composition of pseudofunctors $\mathfrak{S}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}[W^{-1}]$ given by spanisation and $\mathfrak{J}(\mathfrak{S}): \mathbf{DGpoid}[W^{-1}] \rightarrow \mathbf{DGpoid}_{\text{ana}}$. In particular, there is a pseudo-natural transformation between $\mathfrak{S}_{\text{ana}}$ and the (yet-to-be-proven) pseudofunctor $\mathfrak{A}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}_{\text{ana}}$ given by anafunctorisation. This is highlighted in Remark 7.24. Whatever method one chooses, the end results will always only differ by 2-cells of $\mathbf{DGpoid}_{\text{ana}}$. If this bothers the reader, then this might be motivation for them to consider bibundles Section 8, where such differences are bi-equivariant diffeomorphisms of spaces.

Theorem 7.23. *Anafunctorisation $\mathfrak{A}: \mathbf{DGpoid} \rightarrow \mathbf{DGpoid}_{\text{ana}}$, sending objects to themselves and everything else to their anafunctorisations, is a localisation of W .*

Proof. We begin by showing that anafunctorisation \mathfrak{A} is a pseudofunctor. It is defined to send diffeological groupoids to themselves, functors to their anafunctorisations, and natural transformations to their anafunctorisations (see Definition 7.18).

Fixing diffeological groupoids \mathcal{G} and \mathcal{H} , \mathfrak{A} is required to restrict to a functor from $\mathbf{DGpoid}(\mathcal{G}, \mathcal{H})$ to $\mathbf{DGpoid}_{\text{ana}}(\mathcal{G}, \mathcal{H})$. Given functors $\varphi_i: \mathcal{G} \rightarrow \mathcal{H}$ for $i = 1, 2, 3$, with natural transformations $S: \varphi_1 \Rightarrow \varphi_2$ and $T: \varphi_2 \Rightarrow \varphi_3$, the anafunctorisations of S and T

are

$$\begin{aligned}\widetilde{S}((x, h_1, y_1), (x, h_2, y_2)) &= h_2 S(x) h_1^{-1}, \text{ and} \\ \widetilde{T}((x, h_2, y_2), (x, h_3, y_3)) &= h_3 T(x) h_2^{-1}\end{aligned}$$

as defined in [Definition 7.18](#). Their vertical composition is

$$\widetilde{T} \circ \widetilde{S}((x, h_1, y_1), (x, h_3, y_3)) = h_3 T(x) S(x) h_1^{-1};$$

but this is exactly the anafunctorisation of $T \circ S$. Thus \mathfrak{A} respects vertical composition.

Given a functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$, the anafunctorisation of id_φ is $\widetilde{\text{id}}_\varphi((x, h_1, y_1), (x, h_2, y_2)) = h_2 h_1^{-1}$, whereas the identity transformation $I_{\mathcal{G} \leftarrow \mathcal{G} \times_{\mathcal{H}} \mathcal{H} \rightarrow \mathcal{H}}$ sends $((x, h_1, y_1), (x, h_2, y_2))$ to $\text{pr}_3 \left(\Phi_{\text{pr}_1}^{-1}((x, h_1, y_1), (x, h_2, y_2), u_x) \right)$. But by uniqueness this is equal to $h_2 h_1^{-1}$. Thus \mathfrak{A} respects identity 2-cells, and thus restricts to a functor as desired.

Given a diffeological groupoid \mathcal{G} , there is a transformation

$$\text{ID}_{\mathcal{G}}: \mathcal{G}_{0 \text{id}_{\mathcal{G}}} \times_{\text{pr}_1} (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\text{id}_{\mathcal{G}}}^{\text{w}} \mathcal{G})_0 \rightarrow \mathcal{G}_1$$

from the identity anafunctor $\mathcal{G} \xleftarrow{=} \mathcal{G} \xrightarrow{=} \mathcal{G}$ to the anafunctorisation of $\text{id}_{\mathcal{G}}$, equal to $\mathcal{G} \xleftarrow[\text{pr}_1]{\simeq} \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\text{id}_{\mathcal{G}}}^{\text{w}} \mathcal{G} \xrightarrow[\text{pr}_3]{\simeq} \mathcal{G}$, sending $(x, (x, g, x'))$ to g . Also, given functors $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$, there is a transformation

$$C_{\varphi, \psi}: \left((\mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H})_{\text{pr}_3} \times_{\text{pr}_1} (\mathcal{H}_{\psi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K})_{\text{pr}_1 \circ \text{pr}_1} \times_{\text{pr}_1} (\mathcal{G}_{\psi \circ \varphi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K})_0 \right) \rightarrow \mathcal{K}_1$$

from the composition of the anafunctorisations of φ and ψ to the anafunctorisation of the composition $\psi \circ \varphi$, sending $((x, h, y), (y, k_1, z_1), (x, k_2, z_2))$ to $k_2 \psi(h^{-1}) k_1^{-1}$. We need to confirm the coherence relations for a pseudofunctor; these are (4.1.3) and the pair (4.1.4) of [\[JY21, page 104\]](#), or equivalently, (M.1) and the pair (M.2) of [\[B67, page 30\]](#).

The first coherence relation (4.1.3) of [\[JY21\]](#) checks the coherence of composition, called ‘‘pseudofunctorial associativity’’. In our case, it reduces to confirming that for three functors

$$\mathcal{G} \xrightarrow[\varphi]{} \mathcal{H} \xrightarrow[\psi]{} \mathcal{K} \xrightarrow[\chi]{} \mathcal{L}$$

that we have the following equality

$$C_{\psi \circ \varphi, \chi} \circ (\mathfrak{A}(\chi) C_{\varphi, \psi}) = C_{\varphi, \chi \circ \psi} \circ (C_{\psi, \chi} \mathfrak{A}(\varphi)).$$

To check this requires very large tuples (with a lot of redundancy), but otherwise is a straightforward application of the definitions of vertical and horizontal compositions. In particular, to compute $C_{\psi \circ \varphi, \chi} \circ (\mathfrak{A}(\chi) C_{\varphi, \psi})$, begin with a point as input of the form

$$\begin{aligned}& ((x, h, y), (y, k, z), (z, \ell_1, w_1)), ((x, k', z'), (z', \ell', w')), (x, \ell_2, w_2)) \\ & \in \left((\mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H})_{\text{id}_{\mathcal{H}}} \times_{\text{id}_{\mathcal{H}}} (\mathcal{H}_{\psi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K})_{\text{id}_{\mathcal{K}}} \times_{\text{id}_{\mathcal{K}}} (\mathcal{K}_{\chi} \times_{\text{id}_{\mathcal{L}}}^{\text{w}} \mathcal{L}) \right)_{\text{pr}_1 \circ \text{pr}_1} \times_{\text{pr}_1 \circ \text{pr}_1} \\ & \quad \left((\mathcal{G}_{\psi \circ \varphi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K})_{\text{id}_{\mathcal{K}}} \times_{\text{id}_{\mathcal{K}}} (\mathcal{K}_{\chi} \times_{\text{id}_{\mathcal{L}}}^{\text{w}} \mathcal{L}) \right)_{\text{pr}_1 \circ \text{pr}_1} \times_{\text{pr}_1} (\mathcal{G}_{\chi \circ \psi \circ \varphi} \times_{\text{id}_{\mathcal{L}}}^{\text{w}} \mathcal{L});\end{aligned}$$

here, the second component with primed symbols is an auxiliary element used to connect the two natural transformations only (see [Definition 7.11](#)). The corresponding output is $\ell_2\chi((k')^{-1})(\ell')^{-1}$, contributed from $C_{\psi\circ\varphi,\chi}$, multiplied by

$$\ell'\chi(k'\psi(h^{-1})k^{-1})\ell_1^{-1},$$

contributed by $\mathfrak{A}(\chi)C_{\varphi,\psi}$; their product is equal to $\ell_2\chi(\psi(h^{-1})k^{-1})\ell_1^{-1}$. Similarly, a point as input into $C_{\varphi,\chi\circ\psi}\circ(C_{\psi,\chi}\mathfrak{A}(\varphi))$ produces an output of $\ell_2\chi(\psi((h')^{-1}))(\ell')^{-1}$, contributed by $C_{\varphi,\chi\circ\psi}$, multiplied by

$$\ell'\chi(\psi(h')\chi(\psi(h^{-1}))\chi(k^{-1})\ell_1^{-1},$$

contributed by $C_{\psi,\chi}\mathfrak{A}(\varphi)$; their product is equal to the product above, confirming the coherence relation.

The pair of coherence relations (4.1.4) of [\[JY21\]](#) check the coherence of composition with identities, using unitors, called ‘‘pseudofunctorial left and right unity’’. Given a functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$, these reduce to

$$\begin{aligned} \lambda_{\mathcal{G},\mathcal{H}}^{\text{ana}}(\mathfrak{A}(\varphi)) &= I_{\mathfrak{A}(\varphi)} \circ C_{\text{id}_{\mathcal{G}},\varphi} \circ (\mathfrak{A}(\varphi) \text{ID}_{\mathcal{G}}) \text{ and} \\ \rho_{\mathcal{G},\mathcal{H}}^{\text{ana}}(\mathfrak{A}(\varphi)) &= I_{\mathfrak{A}(\varphi)} \circ C_{\varphi,\text{id}_{\mathcal{H}}} \circ (\text{ID}_{\mathcal{H}} \mathfrak{A}(\varphi)). \end{aligned}$$

Again, unravelling definitions and using a point

$$((x, (x, h_1, y_1)), (x, h', y'), (x, h_2, y_2))$$

from the domain of the right-hand side as input (where $(x, h', y') \in \mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H}$ is auxiliary; see [Definition 7.11](#)) produces an output of $h_2(h')^{-1}$ (contributed by $I_{\mathfrak{A}(\varphi)}$) multiplied by $h'h_1^{-1}$ (contributed by $C_{\text{id}_{\mathcal{G}},\varphi} \circ (\mathfrak{A}(\varphi) \text{ID}_{\mathcal{G}})$), whose product is $h_2h_1^{-1}$. But this is exactly the output via $\lambda_{\mathcal{G},\mathcal{H}}^{\text{ana}}(\mathfrak{A}(\varphi))$, confirming the first of the coherence relations in the pair. The second one is checked similarly. Thus, all three coherence conditions hold, and we conclude that \mathfrak{A} is indeed a pseudofunctor.

We now need to show that this pseudofunctor ‘extends’ to an equivalence between bicategories $\mathbf{DGpoid}[W^{-1}] \rightarrow \mathbf{DGpoid}_{\text{ana}}$. To accomplish this, we utilise Pronk’s Comparison Theorem [\[Pro96, Proposition 24\]](#). That is, we need to show:

- (1) \mathfrak{A} sends weak equivalences to adjoint equivalences in $\mathbf{DGpoid}_{\text{ana}}$;
- (2) \mathfrak{A} is essentially surjective on objects (this is trivially true for \mathfrak{A});
- (3) for every anafunctor $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{L} \xrightarrow[\chi]{\simeq} \mathcal{K}$, there exist a weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$, a functor $\omega: \mathcal{G} \rightarrow \mathcal{K}$, and a transformation S from $\mathfrak{A}(\varphi)$ to the composition of $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{L} \xrightarrow[\chi]{\simeq} \mathcal{K}$ with $\mathfrak{A}(\varphi)$; and
- (4) \mathfrak{A} is fully faithful on 2-cells.

[Item 1](#) is precisely [Corollary 7.20](#). To prove [Item 3](#), choose φ to be ψ and ω to be χ . We claim that $\mathcal{H} \xleftarrow[\psi]{\simeq} \mathcal{L} \xrightarrow[\chi]{\simeq} \mathcal{K}$ composed with $\mathfrak{A}(\psi)$ has a transformation to the

anafunctorisation of χ . Indeed, consider the diagram

$$\begin{array}{ccccc}
 & & (\mathcal{L}_{\psi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H})_{\text{pr}_3} \times_{\psi} \mathcal{L} & & \\
 & \swarrow \text{pr}_1 \circ \text{pr}_1 & \uparrow \simeq & \searrow \psi \circ \text{pr}_2 & \\
 \mathcal{L} & \circlearrowleft & \mathcal{M} & \Downarrow S & \mathcal{K} \\
 & \swarrow \text{pr}_1 & \downarrow \simeq & \nearrow \text{pr}_3 & \\
 & & \mathcal{L}_{\chi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K} & &
 \end{array}$$

where $\mathcal{M} := \left((\mathcal{L}_{\psi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H})_{\text{pr}_3} \times_{\psi} \mathcal{L} \right)_{\text{pr}_1 \circ \text{pr}_1} \times_{\text{pr}_1} (\mathcal{L}_{\chi} \times_{\text{id}_{\mathcal{K}}}^{\text{w}} \mathcal{K})$ and

$$S(((w_1, h, \psi(w_2)), w_2), (w_1, k, y)) := k\chi(\Phi_{\psi}^{-1}(w_2, w_1, h^{-1})).$$

It follows from the fact that ψ is a weak equivalence that S is well-defined and a natural transformation; hence S induces a transformation as desired between the two anafunctors.

To prove **Item 4**, fix functors $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$. Given $S: \varphi \Rightarrow \psi$, its anafunctorisation \tilde{S} is given by (see **Definition 7.18**)

$$\tilde{S}((x, h_1, y_1), (x, h_2, y_2)) := h_2 S(x) h_1^{-1}.$$

Since $\text{pr}_1 \circ \text{pr}_1, \text{pr}_1 \circ \text{pr}_2: (\mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H})_{\text{pr}_1} \times_{\text{pr}_1} (\mathcal{G}_{\psi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H}) \rightarrow \mathcal{G}$ are subductions (hence surjective) on the level of objects, it follows that if $S_1, S_2: \varphi \Rightarrow \psi$ have the same anafunctorisations, then $S_1 = S_2$.

Suppose T is a transformation between the anafunctorisations of φ and ψ . Define $S: \mathcal{G}_0 \rightarrow \mathcal{H}_1$ by

$$S(x) := T((x, u_{\varphi(x)}, \varphi(x)), (x, u_{\psi(x)}, \psi(x))).$$

Since T is a natural transformation, so is S . We conclude that \mathfrak{A} is fully faithful on 2-cells. The theorem is proved. \square

Remark 7.24.

- (1) As a corollary to **Theorem 7.23**, the spanisation and anafunctorisation of a functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ admit a 2-cell between them in $\mathbf{DGpoid}_{\text{ana}}$:

$$\begin{array}{ccccc}
 & & \mathcal{G} & & \\
 & \swarrow & \uparrow \text{pr}_1 \simeq & \searrow \varphi & \\
 \mathcal{G} & \circlearrowleft & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\text{pr}_1} (\mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H}) & \Downarrow \text{pr}_2 \circ \text{pr}_2 & \mathcal{H} \\
 & \swarrow \text{pr}_1 & \downarrow \text{pr}_2 \simeq & \nearrow \text{pr}_3 & \\
 & & \mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H} & &
 \end{array}$$

- (2) Another corollary to **Theorem 7.23** is that the composition of two anafunctors $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{L} \xrightarrow[\psi]{\simeq} \mathcal{H}$ and $\mathcal{H} \xleftarrow[\chi]{\simeq} \mathcal{M} \xrightarrow[\omega]{\simeq} \mathcal{K}$ in $\mathbf{DGpoid}[W^{-1}]$ and in $\mathbf{DGpoid}_{\text{ana}}$ yield

generalised morphisms that admit a 2-cell between them in $\mathbf{DGpoid}[W^{-1}]$. A representative diagram of this 2-cell is given by the following, where θ is the functor given by [Corollary 5.9](#) sending arrows (ℓ, m) to $(\ell, u_{\psi \circ s}(\ell), m)$.

$$\begin{array}{ccccc}
 & & \mathcal{L}_{\psi} \times_{\chi} \mathcal{M} & & \\
 & \nearrow \varphi \circ \text{pr}_1 & \parallel & \searrow \omega \circ \text{pr}_2 & \\
 \mathcal{G} & \circlearrowleft & \mathcal{L}_{\psi} \times_{\chi} \mathcal{M} & \circlearrowright & \mathcal{K} \\
 & \searrow \varphi \circ \text{pr}_1 & \downarrow \theta \simeq & \nearrow \omega \circ \text{pr}_3 & \\
 & & \mathcal{L}_{\psi}^w \times_{\chi} \mathcal{M} & &
 \end{array}$$

- (3) As mentioned before [Theorem 7.23](#), given a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$, there is a 2-cell in $\mathbf{DGpoid}[W^{-1}]$ between this and the composition of the anafunctors of φ and ψ ; call the latter the **anafunctorisation** of $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$. The 2-cell is given by the vertical composition of the equivalence classes represented by the following diagrams.

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \nearrow \varphi & \uparrow \text{pr}_3 \simeq & \searrow \psi & \\
 \mathcal{G} & \circlearrowleft & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K} & \circlearrowright & \mathcal{H} \\
 & \nearrow \text{pr}_2 & \parallel & \searrow \psi \circ \text{pr}_3 & \\
 & \searrow \text{pr}_1 & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K} & \nearrow \psi \circ \text{pr}_3 & \\
 & & & & \\
 & & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K} & & \\
 & \nearrow \text{pr}_1 & \uparrow \text{pr}_1 \simeq & \searrow \psi \circ \text{pr}_3 & \\
 \mathcal{G} & \circlearrowleft & (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K})_{\text{pr}_3} \times_{\text{pr}_1} (\mathcal{K}_{\psi} \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H}) & \circlearrowright & \mathcal{H} \\
 & \nearrow \text{pr}_1 \circ \text{pr}_1 & \parallel & \searrow \text{pr}_2 \circ \text{pr}_2 & \\
 & \searrow \text{pr}_1 \circ \text{pr}_1 & (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^w \mathcal{K})_{\text{pr}_3} \times_{\text{pr}_1} (\mathcal{K}_{\psi} \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H}) & \nearrow \text{pr}_3 \circ \text{pr}_2 &
 \end{array}$$

7.5. Optimising 2-Cells. One of the main benefits of passing from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DGpoid}_{\text{ana}}$ is that 2-cells are replaced with transformations; in fact, a transformation can be viewed as an equivalence in $\mathbf{DGpoid}[W^{-1}]$ that is an ‘‘optimised’’ representative of the 2-cell, and is furthermore the *unique* such representative. What sets transformations apart from other representatives of a 2-cell of $\mathbf{DGpoid}[W^{-1}]$ is that the left 2-commutative ‘square’ is in fact 1-commutative.

To this end, consider the following equivalence of generalised morphisms in $\mathbf{DGpoid}[W^{-1}]$.

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \swarrow \varphi & \uparrow \alpha \simeq & \searrow \psi & \\
 \mathcal{G} & & \mathcal{L} & & \mathcal{H} \\
 & \swarrow \varphi' & \downarrow \alpha' \simeq & \searrow \psi' & \\
 & & \mathcal{K}' & &
 \end{array}
 \quad (5)$$

$\Downarrow S_1$ $\Downarrow S_2$

By [Item 3](#) of [Remark 7.24](#), we can replace the top and bottom generalised morphisms with their anafunctisations, so let us suppose they are anafunctors to begin with. We also want to replace \mathcal{L} with $\mathcal{K}_{\text{pr}_1} \times_{\text{pr}_1} \mathcal{K}'$, replace S_1 with a commuting square, and replace S_2 with a different natural transformation in order to obtain a transformation as in [Definition 7.10](#). However, it is not immediately obvious how to go about doing this, as S_2 depends on the domain of \mathcal{L} , and so it remains unclear which transformation in $\mathbf{DGpoid}_{\text{ana}}$ stands in for the equivalence above. In [\[PS21, §5\]](#), Pronk and Scull prove that for a class of arrows that admit a right calculus of fractions (see [Definition 6.1](#)), each of which is co-fully faithful (see [Definition 7.1](#)), not only is such a set of replacements possible, but the end result is unique.

Initially, this looks promising: if we follow the recipe outlined by Pronk-Scull, then we should see precisely what transformation in $\mathbf{DGpoid}_{\text{ana}}$ that the 2-cell (5) is sent to. (We already know that there is a (necessarily unique) such transformation by [Theorem 7.23](#); what we wish to see is what the image is explicitly.) Indeed, we already proved that subductive weak equivalences are co-fully faithful ([Theorem 7.6](#)). But there is a problem: the recipe of Pronk-Scull requires that a natural transformation from a subductive weak equivalence to another functor forces the second functor to be a subductive weak equivalence. But this is not true. That is, if we replace W in [Definition 6.1](#) with the class J of *subductive* weak equivalences, while the first four conditions are met, the last is not, as [Example 7.3](#) illustrates.

Thus the construction of Pronk-Scull does not directly apply. This said, the class of subductive weak equivalences has many nice properties beyond those required by [Definition 6.1](#), and it turns out these properties allow us to follow the construction used by Pronk and Scull, making necessary adjustments¹. To this end, we prove a version of [\[PS21, Lemma 5.2\]](#) for subductive weak equivalences. This yields the existence of a diagram representing the 2-cell (5), but where the left-hand ‘square’ is a strict pullback.

¹Thank you to Dorette Pronk and Laura Scull for pointing this out in a private discussion.

Lemma 7.25. *Given a 2-cell in $\mathbf{DGpoid}[W^{-1}]$ between anafunctors represented by the following diagram*

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \nearrow \varphi & \uparrow \alpha \simeq & \searrow \psi & \\
 \mathcal{G} & & \mathcal{L} & & \mathcal{H} \\
 & \searrow \varphi' & \downarrow \alpha' \simeq & \nearrow \psi' & \\
 & & \mathcal{K}' & &
 \end{array}$$

$\Downarrow S_1$ $\Downarrow S_2$

there exists a (unique) representative diagram of the form

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \nearrow \varphi & \uparrow \text{pr}_1 \simeq & \searrow \psi & \\
 \mathcal{G} & & \mathcal{K}_\varphi \times_{\varphi'} \mathcal{K}' & & \mathcal{H} \\
 & \searrow \varphi' & \downarrow \text{pr}_2 \simeq & \nearrow \psi' & \\
 & & \mathcal{K}' & &
 \end{array}$$

$\Downarrow U$

Here U is the unique natural transformation for which $U \text{pr}_1 = \tilde{U}$ where the natural transformation $\tilde{U}: \left((\mathcal{K}_\varphi \times_{\varphi'} \mathcal{K}')_{\text{pr}_1} \overset{w}{\times} \mathcal{L} \right)_0 \rightarrow \mathcal{H}_1$ is the composition given by Equation (6).

Proof. Define $\tilde{\mathcal{L}} := (\mathcal{K}_\varphi \times_{\varphi'} \mathcal{K}')_{\text{pr}_1} \overset{w}{\times} \mathcal{L}$, and consider the diagram

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \nearrow \text{pr}_1 \simeq & \uparrow \text{pr}_2 \simeq & \searrow \alpha \simeq & \\
 & & \mathcal{K}_\varphi \times_{\varphi'} \mathcal{K}' & \xleftarrow{\text{pr}_1 \simeq} & \tilde{\mathcal{L}} & \xrightarrow{\text{pr}_3 \simeq} & \mathcal{L} & \\
 & \searrow \text{pr}_2 \simeq & & & & & & \searrow \alpha' \simeq & \\
 & & \mathcal{K}' & & & & & &
 \end{array}$$

By [Lemma 4.5](#), the natural transformation

$$(S_1 \text{pr}_3) \circ (\varphi \text{pr}_2) \circ (\text{id}_{\varphi' \circ \text{pr}_2} \text{pr}_1): \varphi' \circ \text{pr}_2 \circ \text{pr}_1 \Rightarrow \varphi' \circ \alpha' \circ \text{pr}_3$$

factors as $\varphi' T$ for a unique natural transformation $T: \text{pr}_2 \circ \text{pr}_1 \Rightarrow \alpha' \circ \text{pr}_3$, making the lower 'square' in the above diagram 2-commute. It follows from the definition of T and the equality $\text{id}_{\varphi' \circ \text{pr}_2}^{-1} = \text{id}_{\varphi \circ \text{pr}_1}$ that

$$(S_1 \text{pr}_3) \circ (\varphi \text{pr}_2) = (\varphi' T) \circ (\text{id}_{\varphi \circ \text{pr}_1} \text{pr}_1).$$

By [Item 1](#) of [Proposition 5.10](#), $\text{pr}_1: \tilde{\mathcal{L}} \rightarrow \mathcal{K}_{\varphi \times \varphi} \mathcal{K}'$ and $\text{pr}_3: \tilde{\mathcal{L}} \rightarrow \mathcal{L}$ are subductive weak equivalences. In particular, pr_1 is co-fully faithful, and so the natural transformation

$$\tilde{U} := (\psi' T^{-1}) \circ (S_2 \text{pr}_3) \circ (\psi \text{pr}_2): \psi \circ \text{pr}_1 \circ \text{pr}_1 \Rightarrow \psi' \circ \text{pr}_2 \circ \text{pr}_1 \quad (6)$$

factors as $\tilde{U} = U \text{pr}_1$ for some unique $U: \psi \circ \text{pr}_1 \Rightarrow \psi' \circ \text{pr}_2$. It follows from the definition of U that

$$(S_2 \text{pr}_3) \circ (\psi \text{pr}_2) = (\psi' T) \circ (U \text{pr}_1),$$

which completes the proof. Thus, the two diagrams in the statement of the lemma are in the same equivalence class, and hence represent the same 2-cell. \square

8. BIBUNDLES

In this section we introduce a third bicategory **DBiBund** of diffeological groupoids, this time with a more geometric flavour. The presentation of the general theory and notation follow that in [\[vdS20; vdS21\]](#), which in turn is largely based on various other sources in the literature, such as [\[Blo08; dHo12\]](#). We start with the definition of diffeological groupoid bundles and bibundles, constructing the bibundle bicategory. For more details on the subtle differences between the Lie- and diffeological development of the theories, as well as proofs, we refer to [\[vdS21\]](#). In [Subsection 8.2](#), we discuss the embedding of **DGpoid** into **DBiBund**, named “bibundlisation”, and show that **DBiBund** is a localisation of **DGpoid** at W in [Subsection 8.3](#). Moving from **DGpoid** $[W^{-1}]$ to **DBiBund** “optimises” 1-cells in much the same way that moving to **DGpoid**_{ana} “optimised” 2-cells; moreover this “movement” passes through **DGpoid**_{ana}. This is the subject of [Subsection 8.4](#).

8.1. The Bibundle Bicategory DBiBund. Recall the definition of a groupoid action [Definition 3.7](#), which is essential to this section. The objects of the bicategory **DBiBund** will again be diffeological groupoids, and the 1-cells are so-called bibundles, which admit a left and right groupoid action. Our first goal is to define these.

Definition 8.1 (Principal \mathcal{G} -Bundle). Given a diffeological groupoid \mathcal{G} and two left groupoid actions $\mathcal{G} \curvearrowright^l X$ and $\mathcal{G} \curvearrowright^l Y$, a smooth map $\varphi: X \rightarrow Y$ is **\mathcal{G} -equivariant** if $l_Y = l_X \circ \varphi$ and for all $(g, x) \in \mathcal{G}_1 \times_{l_X} X$,

$$\varphi(gx) = g\varphi(x).$$

In particular, if the action on Y is trivial, then we say that φ is **\mathcal{G} -invariant**. A similar definition holds for right actions.

A **(smooth) left \mathcal{G} -bundle** is a left groupoid action $\mathcal{G} \curvearrowright^l X$ together with a \mathcal{G} -invariant smooth map $\pi: X \rightarrow B$, denoted $\mathcal{G} \curvearrowright^l X \xrightarrow{\pi} B$. **Right \mathcal{G} -bundles** are defined similarly, and denoted $B \xleftarrow{\pi} X \curvearrowright^r \mathcal{G}$.

A left \mathcal{G} -bundle $\mathcal{G} \curvearrowright^l X \xrightarrow{\pi} B$ is **subductive** if $\pi: X \rightarrow B$ is a subduction. The bundle is **pre-principal** if the **action map** $A_{\mathcal{G}}: \mathcal{G}_s \times_{l_X} X \rightarrow X_{\pi} \times_{\pi} X: (g, x) \mapsto (gx, x)$ is a diffeomorphism; in this case, there is a smooth **division map** $d_{\mathcal{G}}: X_{\pi} \times_{\pi} X \rightarrow \mathcal{G}$, sending (x, x') to the unique $g \in \mathcal{G}_1$ such that $gx = x'$. The bundle is **principal** if it is both subductive and pre-principal. A similar definition holds for right bundles. \diamond

Example 8.2 (Principal Group Bundles). Given a diffeological group G , a principal G -bundle in the standard sense [IZ13, Article 8.11] is a principal bundle in the sense above with \mathcal{G} the groupoid with one object and arrow space G . //

Example 8.3 (Quotient Structures). If a bundle $\mathcal{G} \curvearrowright^{l_X} X \xrightarrow{\pi} B$ is pre-principal, the action map induces a diffeomorphism from the quotient space induced by the equivalence relation whose equivalence classes are fibres of π , denoted X/π , to the orbit space X/\mathcal{G} . If the bundle is subductive, there is a diffeomorphism from X/π to B . Thus, if the bundle is principal, then B is diffeomorphic to X/\mathcal{G} . //

Example 8.4 (Self-Groupoid Actions). The action of any diffeological groupoid \mathcal{G} on its own arrow space (Example 3.8) forms a left principal \mathcal{G} -bundle $\mathcal{G} \curvearrowright^t \mathcal{G}_1 \xrightarrow{s} \mathcal{G}_0$. Similarly, the right action of \mathcal{G} on itself yields a right principal \mathcal{G} -bundle $\mathcal{G}_0 \xleftarrow{t} \mathcal{G}_1 \curvearrowright^s \mathcal{G}$. //

We are now ready to define the 1-cells of **DBiBund**: the right principal bibundles.

Definition 8.5 (Bibundles). Given diffeological groupoids \mathcal{G} and \mathcal{H} , a **(diffeological) $(\mathcal{G}, \mathcal{H})$ -bibundle** comprises a left action $\mathcal{G} \curvearrowright^{l_X} X$ and a right action $X \curvearrowright^{r_X} \mathcal{H}$ such that the left anchor map l_X is \mathcal{H} -invariant, the right anchor map r_X is \mathcal{G} -invariant, and the actions commute. Denote these by $\mathcal{G} \curvearrowright^{l_X} X \curvearrowright^{r_X} \mathcal{H}$. The bibundle is **left principal** if the left underlying bundle $\mathcal{G} \curvearrowright^{l_X} X \xrightarrow{r_X} \mathcal{H}_0$ is principal; it is **right principal** if the right underlying bundle $\mathcal{G}_0 \xleftarrow{l_X} X \curvearrowright^{r_X} \mathcal{H}$ is principal. It is **biprincipal** if it is both left and right principal. We often represent a bibundle diagrammatically as follows:

$$\begin{array}{ccc} \mathcal{G}_1 & X & \mathcal{H}_1 \\ \Downarrow & \swarrow l_X & \searrow r_X \\ \mathcal{G}_0 & & \mathcal{H}_0 \end{array}$$

For a fixed diffeological groupoid $\mathcal{G} \rightrightarrows \mathcal{G}_0$, the **identity bibundle** is given by $\mathcal{G} \curvearrowright^t \mathcal{G} \curvearrowright^s \mathcal{G}$, where the actions are as given in Example 3.8. \diamond

The composition of bibundles is defined using a construction known as the ‘‘balanced tensor product’’. In the literature, this is also known as the Hilsum-Skandalis tensor product, first appearing in the context of topological groupoids in a paper by Hilsum and Skandalis [HS87].

Definition 8.6 (Balanced Tensor Products). Given right principal bibundles $\mathcal{G} \curvearrowright^{l_X} X \curvearrowright^{r_X} \mathcal{H}$ and $\mathcal{H} \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} \mathcal{K}$ define the **balanced tensor product** to be the space $X \otimes_{\mathcal{H}} Y := (X_{r_X} \times_{l_Y} Y) / \mathcal{H}$, where the \mathcal{H} -action is the natural diagonal action. Equip $X \otimes_{\mathcal{H}} Y$ with a left \mathcal{G} -action

$$\mathcal{G} \curvearrowright^{L_X} X \otimes_{\mathcal{H}} Y : g \cdot (x \otimes y) := (gx) \otimes y$$

with left anchor map

$$L_X : X \otimes_{\mathcal{H}} Y \rightarrow \mathcal{G}_0 : x \otimes y \mapsto l_X(x),$$

and with a right \mathcal{K} -action

$$X \otimes_{\mathcal{H}} Y \xrightarrow{R_Y \curvearrowright \mathcal{K}} (x \otimes y)k := x \otimes (yk)$$

with right anchor map

$$R_Y: X \otimes_{\mathcal{H}} Y \rightarrow \mathcal{K}_0: x \otimes y \mapsto r_Y(y).$$

This is a well-defined right principal bibundle [vdS21, Constructions 4.6, 5.8]. \diamond

We now define the 2-cells of **DBiBund** to be bi-equivariant diffeomorphisms between bibundles.

Definition 8.7 (Bibundle Morphisms). Given diffeological groupoids \mathcal{G} and \mathcal{H} , a **\mathcal{G} -bundle morphism** from a left \mathcal{G} -bundle $\mathcal{G} \curvearrowright^{l_X} X \xrightarrow{\pi_X} B$ to a left \mathcal{G} -bundle $\mathcal{G} \curvearrowright^{l_Y} Y \xrightarrow{\pi_Y} B$ is a \mathcal{G} -equivariant smooth map $\alpha: X \rightarrow Y$ such that $\pi_X = \pi_Y \circ \alpha$. A similar definition holds for right bundles.

Given principal bibundles $\mathcal{G} \curvearrowright^{l_X} X \xrightarrow{r_X} \mathcal{H}$ and $\mathcal{G} \curvearrowright^{l_Y} Y \xrightarrow{r_Y} \mathcal{H}$, a **bibundle morphism** is a smooth map $\alpha: X \rightarrow Y$ that is a bundle morphism between both underlying bundles. In this case, α is **bi-equivariant**, that is, it is equivariant with respect to both actions and the following diagram commutes:

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & X & & \mathcal{H}_1 \\
 \Downarrow & & \downarrow & & \Downarrow \\
 \mathcal{G}_0 & \xleftarrow{l_X} & & \xrightarrow{r_X} & \mathcal{H}_0 \\
 & & \alpha & & \\
 & \xleftarrow{l_Y} & Y & \xrightarrow{r_Y} &
 \end{array}$$

The **2-cells** of **DBiBund** are precisely the bi-equivariant diffeomorphisms between the bibundles. The **identity 2-cell** of a bibundle is its identity map. \diamond

Working with bibundles allows one to have much tamer vertical and horizontal compositions than those in **DGpoid** $[W^{-1}]$ and **DGpoid** $_{\text{ana}}$.

Remark 8.8 (Vertical Composition of Bi-Equivariant Diffeomorphisms). Vertical composition in **DBiBund** is given by composition of the bi-equivariant diffeomorphisms [vdS21, Proposition 5.14]. \lrcorner

Definition 8.9 (Horizontal Composition of Bi-Equivariant Diffeomorphisms). Given four right principal bibundles with bi-equivariant diffeomorphisms

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & X & & \mathcal{H}_1 \\
 \Downarrow & & \downarrow & & \Downarrow \\
 \mathcal{G}_0 & \xleftarrow{l_X} & & \xrightarrow{r_X} & \mathcal{H}_0 \\
 & & \alpha & & \\
 & \xleftarrow{l_Y} & Y & \xrightarrow{r_Y} &
 \end{array}$$

and

$$\begin{array}{ccccc}
 \mathcal{H}_1 & & Z & & \mathcal{K}_1 \\
 \Downarrow & & \swarrow & & \Downarrow \\
 \mathcal{H}_0 & & l_Z & & r_Z \\
 & & \beta & & \\
 & & \downarrow & & \\
 & & W & & \\
 & & \swarrow & & \searrow \\
 & & l_W & & r_W \\
 & & & & \mathcal{K}_0
 \end{array}$$

define the **horizontal composition** of the two 2-cells to be

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & X \otimes_{\mathcal{H}} Z & & \mathcal{K}_1 \\
 \Downarrow & & \swarrow & & \Downarrow \\
 \mathcal{G}_0 & & L_X & & R_Z \\
 & & \alpha \otimes \beta & & \\
 & & \downarrow & & \\
 & & Y \otimes_{\mathcal{H}} W & & \\
 & & \swarrow & & \searrow \\
 & & L_Y & & R_W \\
 & & & & \mathcal{K}_0
 \end{array}$$

where $\alpha \otimes \beta: x \otimes z \mapsto \alpha(x) \otimes \beta(z)$, and the anchor maps L_X , L_Y , R_Z , and R_W are as defined in [Definition 8.6](#). See [[vdS21](#), Proposition 5.16] for details. Left and right whiskering is performed using horizontal composition with the identity 2-cell of a bibundle. \diamond

While 2-cells and their compositions are better behaved in **DBiBund**, we no longer have honest associativity of composition, but up to bi-equivariant diffeomorphism.

Definition 8.10 (Associators for Bibundles). Given right principal bibundles $\mathcal{G} \curvearrowright^{l_X} X \curvearrowright^{r_X} \mathcal{H}$, $\mathcal{H} \curvearrowright^{l_Y} Y \curvearrowright^{r_Y} \mathcal{K}$, and $\mathcal{K} \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} \mathcal{L}$, there exists a bi-equivariant diffeomorphism, called the **associator**,

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & (X \otimes_{\mathcal{H}} Y) \otimes_{\mathcal{K}} Z & & \mathcal{H}_1 \\
 \Downarrow & & \swarrow & & \Downarrow \\
 \mathcal{G}_0 & & L_{X \otimes_{\mathcal{H}} Y} & & R_Z \\
 & & A & & \\
 & & \downarrow & & \\
 & & X \otimes_{\mathcal{H}} (Y \otimes_{\mathcal{K}} Z) & & \\
 & & \swarrow & & \searrow \\
 & & L_X & & R_{Y \otimes_{\mathcal{K}} Z} \\
 & & & & \mathcal{H}_0
 \end{array}$$

where $A: (x \otimes y) \otimes z \mapsto x \otimes (y \otimes z)$ and the anchor maps are defined in [Definition 8.6](#). See [[vdS21](#), Proposition 5.13] for details. \diamond

We also have non-trivial unitors, as in the previous two bicategories.

Definition 8.11 (Unitors for Bibundles). For fixed diffeological groupoids \mathcal{G} and \mathcal{H} , the **left unitor** $\lambda_{\mathcal{G}, \mathcal{H}}^{\text{bi}}$ is a natural transformation assigning to each right principal bibundle

$\mathcal{G} \curvearrowright^{l_X} X \curvearrowleft^{r_X} \mathcal{H}$, the bi-equivariant diffeomorphism

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & \mathcal{G}_1 \otimes_{\mathcal{G}} X & & \mathcal{H}_1 \\
 \Downarrow & & \downarrow & & \Downarrow \\
 \mathcal{G}_0 & \xleftarrow{L_{\mathcal{G}}} & & \xrightarrow{R_X} & \mathcal{H}_0 \\
 & \searrow^{l_X} & \downarrow \lambda_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(X) & \nearrow^{r_X} & \\
 & & X & &
 \end{array}$$

where $\lambda_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(X)(g \otimes x) := gx$. The **right unitor** $\rho_{\mathcal{G}, \mathcal{H}}^{\text{bi}}$ is a natural transformation assigning to each right principal bibundle $\mathcal{G} \curvearrowright^{l_X} X \curvearrowleft^{r_X} \mathcal{H}$, the bi-equivariant diffeomorphism

$$\begin{array}{ccccc}
 \mathcal{G}_1 & & X \otimes_{\mathcal{H}} \mathcal{H}_1 & & \mathcal{H}_1 \\
 \Downarrow & & \downarrow & & \Downarrow \\
 \mathcal{G}_0 & \xleftarrow{L_X} & & \xrightarrow{R_{\mathcal{H}}} & \mathcal{H}_0 \\
 & \searrow^{l_X} & \downarrow \rho_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(X) & \nearrow^{r_X} & \\
 & & X & &
 \end{array}$$

where $\rho_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(X)(x \otimes h) := xh$. See [vdS21, Proposition 5.12] for details. \diamond

We are now ready for the main theorem of this subsection.

Theorem 8.12 (The Bicategory **DBiBund**). *There is a bicategory **DBiBund** consisting of diffeological groupoids as objects, right principal bibundles as 1-cells, bi-equivariant diffeomorphisms as 2-cells, with all compositions, associators, and unitors as defined as above.*

The proof follows from [vdS21, Theorem 5.17, Subsection 5.3]; the theorem focuses on 1-cells as bibundles that are not necessarily principal, and 2-cells that are not necessarily diffeomorphisms (this yields a category for diffeological groupoids, although it does not for Lie groupoids). Subsection 5.3 then shows that the restriction to the sub-bicategory as in **Theorem 8.12** satisfies the required identities. The proof of [vdS21, Theorem 5.17] is analogous to that for Lie theory; see [Blo08, Proposition 2.12].

8.2. Bibundlisation. Similar to spanisation and anafunctisation, there is a natural way to turn a functor between diffeological groupoids into a bibundle, and a natural transformation into a bi-equivariant diffeomorphism.

Definition 8.13 (Bibundlisation). Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a functor. Its **bibundlisation** (or just bundlisation, as in [Blo08]) is the bibundle

$$\mathcal{G} \curvearrowright^{\text{pr}_1} \mathcal{G}_{0, \varphi} \times_{\text{t}} \mathcal{H}_1 \xrightarrow{\Psi_{\varphi}} \mathcal{H}$$

where the \mathcal{G} -action sends $(g, (x, h))$ to $(t(g), \varphi(g)h)$ with anchor map pr_1 , and the \mathcal{H} -action sends $((x, h), h')$ to (x, hh') . If $\psi: \mathcal{G} \rightarrow \mathcal{H}$ is another functor and $S: \varphi \Rightarrow \psi$ is a natural transformation, then the **bibundlisation** of S is the bi-equivariant diffeomorphism $\alpha: \mathcal{G}_{0, \varphi} \times_{\text{t}} \mathcal{H}_1 \rightarrow \mathcal{G}_{0, \psi} \times_{\text{t}} \mathcal{H}_1$ sending (x, h) to $(x, S(x)h)$. \diamond

Remark 8.14 (Bibundlisation versus Anafunctorisation of Functors). Given a functor $\varphi: \mathcal{G} \rightarrow \mathcal{H}$, the $(\mathcal{G}\mathcal{H})$ -action groupoid corresponding to the joint \mathcal{G} - and \mathcal{H} -actions on $\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1$, denoted $\mathcal{G} \times \mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1 \rtimes \mathcal{H}$, is defined to be the action groupoid of the left groupoid action of $\mathcal{G} \times \mathcal{H}$ on $\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1$ with action map $((g', h'), (x, h)) \mapsto (t(g'), \varphi(g')h(h')^{-1})$ and anchor map $(x, h) \mapsto (x, s(h))$. It is straightforward to check that the map $\mathcal{G} \times \mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1 \rtimes \mathcal{H} \rightarrow \mathcal{G}_{\varphi} \times_{\text{id}_{\mathcal{H}}}^w \mathcal{H}$ sending $((g', h'), (x, h))$ to (g', h^{-1}, h') is an isomorphism of diffeological groupoids. In particular, the action groupoid of the bibundlisation of φ is isomorphic to the anafunctorisation of φ . \square

Lemma 8.15 (Bibundlisation of a Weak Equivalence). *The bibundlisation of a functor is biprincipal if and only if the functor is a weak equivalence.*

Proof. Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a functor. It is immediate from its definition that the bibundlisation of φ is left subductive if and only if φ is smoothly essentially surjective. It is straightforward to check that if φ is smoothly fully faithful, then the action map

$$A_{\mathcal{G}}: \mathcal{G}_s \times_{\text{pr}_1} (\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1) \rightarrow (\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1)_{\Psi_{\varphi}} \times_{\Psi_{\varphi}} (\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1)$$

is a diffeomorphism. Conversely, if $A_{\mathcal{G}}$ is a diffeomorphism, then the division map of the \mathcal{G} -bundle $\mathcal{G}_{0,\varphi} \times_t \mathcal{H}_1 \xrightarrow{\Psi_{\varphi}} \mathcal{H}_0$ is well-defined and smooth, from which it follows that Φ_{φ} is a diffeomorphism. \square

Lemma 8.15 indicates that biprincipal bibundles may be the bibundle version of the quasi-invertible generalised morphisms and anafunctors of **Lemma 6.13** and **Lemma 7.19**. The following definition and lemma confirm a more general statement.

Definition 8.16 (Opposite Bibundles). Consider a bibundle $\mathcal{G} \curvearrowright^{l_X} X \curvearrowright^{r_X} \mathcal{H}$. The **opposite bibundle** $\mathcal{H} \curvearrowright^{l_{\overline{X}}} \overline{X} \curvearrowright^{r_{\overline{X}}} \mathcal{G}$ is defined as follows. The underlying diffeological space does not change, $\overline{X} := X$, but the anchor maps switch, meaning that $l_{\overline{X}} := r_X$ and $r_{\overline{X}} := l_X$, and the actions are defined as follows:

$$\begin{aligned} \mathcal{H} \curvearrowright^{r_X} \overline{X}; & \quad h \cdot x := xh^{-1}, \\ \overline{X} \curvearrowright^{l_X} \mathcal{G}; & \quad x \cdot g := g^{-1}x. \end{aligned}$$

Here the actions on the right-hand sides are the original actions of the bibundle. It is easy to see that performing this operation twice gives the original bibundle back. Also note that for all properties defined in **Definition 8.5**, taking the opposite merely switches the words ‘left’ and ‘right’. In particular, if a bibundle is biprincipal, so is its opposite bibundle. \diamond

Lemma 8.17. *The opposite bibundle of a biprincipal bibundle is its quasi-inverse.*

Proof. This is [**vdS21**, Proposition 5.24]. The division maps induced by biprincipality are the desired bi-equivariant diffeomorphisms. \square

8.3. $\mathbf{DGpoid}[W^{-1}]$ versus $\mathbf{DBiBund}$. Similar to what we did for $\mathbf{DGpoid}_{\text{ana}}$, we show that $\mathbf{DBiBund}$ is a localisation of \mathbf{DGpoid} at W . We need the following lemma, which could be called the ‘‘anafunctorisation of a bibundle’’.

Lemma 8.18. *Let $\mathcal{G} \curvearrowright^{l_X} X \curvearrowright^{r_X} \mathcal{H}$ be a bibundle. The $(\mathcal{G} \times \mathcal{H})$ -action groupoid $\mathcal{G} \times X \times \mathcal{H}$ for the action on X admits functors \widehat{r}_X to \mathcal{H} and \widehat{l}_X to \mathcal{G} , such that $(\widehat{l}_X)_0 = l_X$, $(\widehat{r}_X)_0 = r_X$, and \widehat{l}_X is a subductive weak equivalence. That is, $\mathcal{G} \xleftarrow[\widehat{l}_X]{\cong} \mathcal{G} \times X \times \mathcal{H} \xrightarrow[\widehat{r}_X]{} \mathcal{H}$ is an anafunctor.*

Proof. The action groupoid for the $(\mathcal{G} \times \mathcal{H})$ -action on X has arrows $((g, h), x)$ with source map the second projection map, and target map the action map $((g, h), x) \mapsto gxh^{-1}$. Define $\widehat{r}_X((g, h), x) = h^{-1}$ and $\widehat{l}_X((g, h), x) = g$. Then $(\widehat{r}_X)_0 = r_X$ and $(\widehat{l}_X)_0 = l_X$. Injectivity of $\Phi_{\widehat{l}_X}$ follows from the freeness of the \mathcal{H} -action on X . If (x_u, x'_u, g_u) is a plot of $X^2 \times_{l_X^2} \times_{(s,t)} \mathcal{G}_1$, then $((g_u, h_u), x_u)$ is a lift to $(\mathcal{G} \times X \times \mathcal{H})_1$, where $h_u = d_{\mathcal{H}}(x'_u, g_u x_u)$ and $d_{\mathcal{H}}$ is the division map for the right \mathcal{H} -bundle. Thus $\Phi_{\widehat{l}_X}$ is a diffeomorphism, and since l_X is subductive, we conclude that \widehat{l}_X is a subductive weak equivalence. \square

Theorem 8.19. *Bibundlisation $\mathfrak{B}: \mathbf{DGpoid} \rightarrow \mathbf{DBiBund}$, sending objects to themselves and everything else to their bibundlisation, is a localisation of W .*

Proof. We begin by showing that bibundlisation \mathfrak{B} is a pseudofunctor. It is defined to send diffeological groupoids to themselves, functors to their bibundlisations, and natural transformations to their bibundlisations (see [Definition 8.13](#)).

Fixing diffeological groupoids \mathcal{G} and \mathcal{H} , \mathfrak{B} is required to restrict to a functor from $\mathbf{DGpoid}(\mathcal{G}, \mathcal{H})$ to $\mathbf{DBiBund}(\mathcal{G}, \mathcal{H})$. But this is immediate, since vertical composition of bi-equivariant diffeomorphisms is just their composition in the standard sense, and the bibundlisation of an identity natural transformation is equal to the identity bi-equivariant diffeomorphism.

Given a diffeological groupoid \mathcal{G} , there is a bi-equivariant diffeomorphism $\iota_{\mathcal{G}}$ from the identity bibundle of \mathcal{G} to the bibundlisation of $\text{id}_{\mathcal{G}}$, given by the inverse of the map $\text{pr}_2: \mathcal{G}_{0, \text{id}_{\mathcal{G}}} \times_t \mathcal{G}_1 \rightarrow \mathcal{G}_1$. Also, given functors $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$, there is a bi-equivariant diffeomorphism

$$\gamma_{\varphi, \psi}: (\mathcal{G}_{0, \varphi} \times_t \mathcal{H}_1) \otimes_{\mathcal{H}} (\mathcal{H}_{0, \psi} \times_t \mathcal{K}_1) \rightarrow \mathcal{G}_{0, \psi \circ \varphi} \times_t \mathcal{K}_1: (x, h) \otimes (y, k) \mapsto (x, \psi(h)k).$$

We need to confirm the coherence relations for a pseudofunctor; again, we refer to (4.1.3) and (4.1.4) of [\[JY21, page 104\]](#) or (M.1) and (M.2) of [\[B67, page 30\]](#).

For the pseudofunctorial associativity coherence relation [\[JY21, \(4.1.3\)\]](#), fix three functors

$$\mathcal{G} \xrightarrow[\varphi]{} \mathcal{H} \xrightarrow[\psi]{} \mathcal{K} \xrightarrow[\chi]{} \mathcal{L}.$$

Let A be the associator of [Definition 8.10](#). Then the coherence relation reduces to showing that

$$\gamma_{\psi \circ \varphi, \chi} \circ (\gamma_{\varphi, \psi} \otimes \text{id}_{\mathcal{K}_0 \times \mathcal{L}_1}) = \gamma_{\varphi, \chi \circ \psi} \circ (\text{id}_{\mathcal{G}_0 \times \mathcal{H}_1} \otimes \gamma_{\psi, \chi}) \circ A.$$

The left-hand side sends $((x, h) \otimes (s(h), k)) \otimes (s(k), \ell)$ to $(x, \chi(\psi(h)k)\ell)$, whereas the right-hand side sends the same point to $(x, \chi \circ \psi(h)\chi(k)\ell)$; these are equal.

For the pseudofunctorial left and right unity coherence relations [JY21, (4.1.4)], called ‘‘pseudofunctorial left and right unity’’, the first reduces to

$$\lambda_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(\mathcal{G}_{0\varphi} \times_t \mathcal{H}_1) = \mathfrak{B}(\text{id}_\varphi) \circ \gamma_{\text{id}_{\mathcal{G}}, \varphi} \circ (\iota_{\mathcal{G}} \otimes \text{id}_{\mathcal{G}_{0\varphi} \times_t \mathcal{H}_1}).$$

Both sides send $(g, (s(g), h))$ to $(t(g), \varphi(g)h)$, confirming this coherence relation. The second relation reduces to

$$\rho_{\mathcal{G}, \mathcal{H}}^{\text{bi}}(\mathcal{G}_{0\varphi} \times_t \mathcal{H}_1) = \mathfrak{B}(\text{id}_\varphi) \circ \gamma_{\varphi, \text{id}_{\mathcal{H}}} \circ (\text{id}_{\mathcal{G}_{0\varphi} \times_t \mathcal{H}_1} \otimes \iota_{\mathcal{H}}),$$

in which both sides send $((x, h), h')$ to (x, hh') , confirming the last coherence relation. We conclude that \mathfrak{B} is a pseudofunctor.

Next, we again use Pronk’s Comparison Theorem [Pro96, Proposition 24] to show that \mathfrak{B} ‘extends’ to an equivalence of bicategories $\mathbf{DGpoid}[W^{-1}]$. We need to show

- (1) \mathfrak{B} sends weak equivalences to adjoint equivalences in $\mathbf{DBiBund}$;
- (2) \mathfrak{B} is essentially surjective on objects (this is trivially true for \mathfrak{B});
- (3) for every bibundle $\mathcal{H} \curvearrowright^{l_Y} Y \curvearrowright^{r_X} \mathcal{K}$, there exist a weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$, a functor $\omega: \mathcal{G} \rightarrow \mathcal{K}$, and a bi-equivariant diffeomorphism α from $\mathfrak{B}(\omega)$ to the composition of $\mathcal{H} \curvearrowright^{l_Y} Y \curvearrowright^{r_X} \mathcal{K}$ with $\mathfrak{B}(\varphi)$; and
- (4) \mathfrak{B} is fully faithful on 2-cells.

Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a weak equivalence. By Lemma 8.15, $\mathfrak{B}(\varphi)$ is a biprincipal bibundle, which is an adjoint equivalence by Lemma 8.17. Thus Item 1 is satisfied.

For Item 3, let \mathcal{G} be the action groupoid $\mathcal{H} \times Y \rtimes \mathcal{K}$, $\varphi = \widehat{l}_X$, and $\omega = \widehat{r}_X$ (see Lemma 8.18). Define $\alpha((y, h) \otimes y') := (y, d_{\mathcal{K}}(y', h^{-1}y))$, where $d_{\mathcal{K}}$ is the division map of the underlying \mathcal{K} -bundle. This is well-defined and smooth with smooth inverse $\alpha^{-1}(y, k) = ((y, u_{r_X(y)}), yk)$. It follows from the \mathcal{H} -invariance of r_X and the \mathcal{K} -invariance of l_X that α is bi-equivariant. This proves Item 3.

Finally, for Item 4, let $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$ be functors with $S_1, S_2: \varphi \Rightarrow \psi$ natural transformations. If $\mathfrak{B}(S_1) = \mathfrak{B}(S_2)$, then by definition, $(x, S_1(x)h) = (x, S_2(x)h)$ for every $(x, h) \in \mathcal{G}_{0\varphi} \times_t \mathcal{H}_1$. In particular, $S_1(x) = S_2(x)$ for every $x \in X$, and so $S_1 = S_2$. On the other hand, for any bi-equivariant diffeomorphism $\alpha: \mathcal{G}_{0\varphi} \times_t \mathcal{H}_1 \rightarrow \mathcal{G}_{\psi} \times_t \mathcal{H}_1$, define $S(x) := h'h^{-1}$ for any $h, h' \in \mathcal{H}_1$ such that $(x, h) \in \mathcal{G}_{0\varphi} \times_t \mathcal{H}_1$ and $(x, h') = \alpha(x, h) \in \mathcal{G}_{\psi} \times_t \mathcal{H}_1$; this is well-defined by the bi-equivariance of α . Moreover, that S is a natural transformation follows from the \mathcal{G} -equivariance of α . We have shown that \mathfrak{B} acts fully faithfully on 2-cells. This completes the proof. \square

8.4. Optimising 1-cells. Similar to how passing from the bicategory of fractions $\mathbf{DGpoid}[W^{-1}]$ to the bicategory of anafunctors $\mathbf{DGpoid}_{\text{ana}}$ yields a unique representative of a 2-cell between two anafunctors (viewed as generalised morphisms in $\mathbf{DGpoid}[W^{-1}]$), passing from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DBiBund}$ yields for a fixed generalised morphism a unique bibundle (up to bi-equivariant diffeomorphism) whose action groupoid is an anafunctor

admitting a 2-cell from it to the original generalised morphism. This is already known given [Theorem 8.19](#), but the theorem hides how to obtain an explicit bibundle from a given generalised morphism. The purpose of [Proposition 8.20](#) is to give an explicit description of this bibundle, inspired by [[PS10](#), Proposition 7.1]. What is interesting is that we pass through $\mathbf{DGpoid}_{\text{ana}}$ during the construction, anafunctising the generalised morphism as in [Item 3](#) of [Remark 7.24](#) before quotienting out redundant information that the original generalised morphism contained, resulting in an optimised anafunctor made up of the action groupoid of a bibundle.

Proposition 8.20. *Given a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$, there is an anafunctor $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \xrightarrow[\omega]{\simeq} \mathcal{H}$ in which \mathcal{L} is the $(\mathcal{G}\text{-}\mathcal{H})$ -action groupoid of a bibundle $\mathcal{G} \curvearrowright^{x_0} \mathcal{L}_0 \curvearrowleft^{y_0} \mathcal{H}$, and a 2-cell from $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ to $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \xrightarrow[\omega]{\simeq} \mathcal{H}$.*

Proof. Define

$$\tilde{\mathcal{L}} := (\mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^{\text{w}} \mathcal{K})_{\text{pr}_3} \times_{\text{pr}_1} (\mathcal{K}_{\psi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H}).$$

By [Item 3](#) of [Remark 7.24](#), $\mathcal{G} \xleftarrow[\text{pr}_{11}]{\simeq} \tilde{\mathcal{L}} \xrightarrow[\text{pr}_{23}]{\simeq} \mathcal{H}$ is an anafunctor admitting a 2-cell from $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ to it. We have the following 2-commutative diagram.

$$\begin{array}{ccccc}
 & & \tilde{\mathcal{L}} & & \\
 & \swarrow \text{pr}_1 & & \searrow \text{pr}_2 & \\
 & \simeq & & \simeq & \\
 & \mathcal{G}_{\text{id}_{\mathcal{G}}} \times_{\varphi}^{\text{w}} \mathcal{K} & \circlearrowleft & \mathcal{K}_{\psi} \times_{\text{id}_{\mathcal{H}}}^{\text{w}} \mathcal{H} & \\
 \text{pr}_1 \swarrow & & \text{pr}_3 \searrow & \text{pr}_1 \swarrow & \text{pr}_3 \searrow \\
 \simeq & & \simeq & \simeq & \\
 \mathcal{G} & & \mathcal{K} & & \mathcal{H} \\
 \Rightarrow \text{pr}_2 & & \Rightarrow \text{pr}_2 & & \\
 \varphi & & \psi & & \\
 \simeq & & \simeq & & \\
 \mathcal{G} & & \mathcal{H} & & \mathcal{H}
 \end{array}$$

It is straightforward to check that $\tilde{\mathcal{L}}_1$ admits a left \mathcal{K} -action

$$k \cdot ((\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})) := ((\tilde{g}, g, k\tilde{k}), (k\tilde{k}, h, \tilde{h}))$$

with anchor map $t \circ \text{pr}_3 \circ \text{pr}_1$, and a right \mathcal{K} -action

$$((\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})) \cdot k := ((\tilde{g}, \varphi(k)^{-1}g, \tilde{k}k), (\tilde{k}k, h\psi(k), \tilde{h}))$$

with anchor map $s \circ \text{pr}_3 \circ \text{pr}_1$. The left and right \mathcal{K} -actions commute. Moreover, there is a right \mathcal{K} -action on $\tilde{\mathcal{L}}_0$

$$((x, g, y), (y, h, z)) \cdot k := (x, \varphi(k)^{-1}g, s(k)), (s(k), h\psi(k), z))$$

with anchor map $\text{pr}_3 \circ \text{pr}_1$. Define \mathcal{L}_0 to be the orbit space $\tilde{\mathcal{L}}_0/\mathcal{K}$ and \mathcal{L}_1 to be the orbit space $\mathcal{K} \backslash \tilde{\mathcal{L}}_1/\mathcal{K}$. Since $s: \tilde{\mathcal{L}}_1 \rightarrow \tilde{\mathcal{L}}_0$ is equivariant with respect to the right \mathcal{K} -actions

and \mathcal{K} -invariant with respect to the left \mathcal{K} -action on $\tilde{\mathcal{L}}_1$, it descends to a subduction $s: \mathcal{L}_1 \rightarrow \mathcal{L}_0$ given by

$$s[(\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})] := [(s(\tilde{g}), g, s(\tilde{k})), (s(\tilde{k}), h, s(\tilde{h}))].$$

Since $t: \tilde{\mathcal{L}}_1 \rightarrow \tilde{\mathcal{L}}_0$ is \mathcal{K} -invariant with respect to the right \mathcal{K} -action on $\tilde{\mathcal{L}}_1$, and

$$t(k \cdot ((\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h}))) = t((\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})) \cdot k^{-1},$$

t descends to a subduction $t: \mathcal{L}_1 \rightarrow \mathcal{L}_0$ given by

$$t[(\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})] := [(t(\tilde{g}), \varphi(\tilde{k})g\tilde{g}^{-1}, t(\tilde{k})), (t(\tilde{k}), \tilde{h}h\psi(\tilde{k})^{-1}, t(\tilde{h}))].$$

It is straightforward to check that $\mathcal{L} = (\mathcal{L}_1 \rightrightarrows \mathcal{L}_0)$ is a diffeological groupoid.

Let $\pi_i: \tilde{\mathcal{L}}_i \rightarrow \mathcal{L}_i$ ($i = 0, 1$) be the quotient maps. Then $\pi = (\pi_0, \pi_1)$ is a functor and π_0 a subduction. Since $\text{pr}_1 \circ \text{pr}_1: \tilde{\mathcal{L}} \rightarrow \mathcal{G}$ and $\text{pr}_3 \circ \text{pr}_2: \tilde{\mathcal{L}} \rightarrow \mathcal{H}$ are \mathcal{K} -invariant (with respect to both \mathcal{K} -actions), they descend to smooth functors $\chi: \mathcal{L} \rightarrow \mathcal{G}$ and $\omega: \mathcal{L} \rightarrow \mathcal{H}$, resp., so that $\chi \circ \pi = \text{pr}_1 \circ \text{pr}_1$ and $\omega \circ \pi = \text{pr}_3 \circ \text{pr}_2$. By **Item 4** of **Proposition 2.7**, χ_0 is a subduction. Moreover, the \mathcal{K} -invariance of $\text{pr}_1 \circ \text{pr}_1$ and $\text{pr}_3 \circ \text{pr}_2$, along with the smooth fully faithfulness of φ , imply that Φ_π is a diffeomorphism. By **Lemma 4.8**, π is a subductive weak equivalence. By **Proposition 4.12** and **Lemma 4.8**, χ is also a subductive weak equivalence; in particular, $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \xrightarrow[\omega]{\simeq} \mathcal{H}$ is an anafunctor. In fact, the diagram

$$\begin{array}{ccccc}
 & & \mathcal{L} & & \\
 & \swarrow \chi & \uparrow \pi & \searrow \omega & \\
 \mathcal{G} & \xrightarrow{\simeq} & \tilde{\mathcal{L}} & \xrightarrow{\simeq} & \mathcal{H} \\
 & \nwarrow \text{pr}_1 \circ \text{pr}_1 & \parallel & \nearrow \text{pr}_3 \circ \text{pr}_2 & \\
 & & \tilde{\mathcal{L}} & &
 \end{array}$$

is a representative of a 2-cell between the generalised morphisms $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \rightarrow \mathcal{H}$ and $\mathcal{G} \xleftarrow[\text{pr}_1 \circ \text{pr}_1]{\simeq} \tilde{\mathcal{L}} \rightarrow \mathcal{H}$. The inverse of this, vertically composed with the 2-cells from **Item 3** of **Remark 7.24**, yields a 2-cell from $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{K} \rightarrow \mathcal{H}$ to $\mathcal{G} \xleftarrow[\omega]{\simeq} \mathcal{L} \rightarrow \mathcal{H}$.

We now set out to prove that \mathcal{L} is isomorphic to the $(\mathcal{G}\text{-}\mathcal{H})$ -action groupoid of a right principal bibundle $\chi_0: \mathcal{L}_0 \rightarrow \mathcal{G}_0$. There is a left action of \mathcal{G} on \mathcal{L}_0 with anchor map χ_0 given by

$$\tilde{g} \cdot [(x, g, y), (y, h, z)] = [(t(\tilde{g}), g\tilde{g}^{-1}, y), (y, h, z)].$$

There is a right action of \mathcal{H} on \mathcal{L}_0 with anchor map ω_0 given by

$$[(x, g, y), (y, h, z)] \cdot \tilde{h} = [(x, g, y), (y, \tilde{h}^{-1}h, s(\tilde{h}))].$$

The two actions commute, χ_0 is \mathcal{H} -invariant, and ω_0 is \mathcal{G} -invariant, and so the groupoid $\mathcal{G} \times \mathcal{L}_0 \times \mathcal{H}$ is well-defined. Define $\Xi: \mathcal{L} \rightarrow \mathcal{G} \times \mathcal{L}_0 \times \mathcal{H}$ to be equality on objects, and on arrows:

$$\Xi_1([(\tilde{g}, g, \tilde{k}), (\tilde{k}, h, \tilde{h})]) = (\tilde{g}, [(s(\tilde{g}), g, s(\tilde{k})), (s(\tilde{k}), h, s(\tilde{h}))], \tilde{h}^{-1}).$$

This is a well-defined smooth functor, and it follows from the definitions that Ξ_1 is a diffeomorphism. Since χ_0 is a subduction, we only need to show right pre-principality to prove the claim.

Let $A_{\mathcal{H}}: \mathcal{L}_{0\omega_0} \times_t \mathcal{H}_1 \rightarrow \mathcal{L}_{0\chi_0} \times_{\chi_0} \mathcal{L}_0$ be the action map for the \mathcal{H} -action on \mathcal{L}_0 . Suppose $A_{\mathcal{H}}(w_0, \tilde{h}_0) = A_{\mathcal{H}}(w_1, \tilde{h}_1)$. Then $w_0 = w_1$ and $w_0 \cdot \tilde{h}_0 = w_1 \cdot \tilde{h}_1$. In particular, there exist $\tilde{w}_0, \tilde{w}_1 \in \tilde{\mathcal{L}}_0$ and $k, k' \in \mathcal{K}_1$ such that $\pi_0(\tilde{w}_i) = w_i$ ($i = 0, 1$), and

$$\tilde{w}_1 = \tilde{w}_0 \cdot k \quad \text{and} \quad \tilde{w}_1 \cdot \tilde{h}_1 = (\tilde{w}_0 \cdot \tilde{h}_0) \cdot k'.$$

It follows from the smooth fully faithfulness of φ that $k = k'$, from which it follows that $\tilde{h}_0 = \tilde{h}_1$. Thus $A_{\mathcal{H}}$ is injective.

Let

$$p = ([(x_u, g_u, y_u), (y_u, h_u, z_u)], [(x'_u, g'_u, y'_u), (y'_u, h'_u, z'_u)])]$$

be a plot of $\mathcal{L}_{0\chi_0} \times_{\chi_0} \mathcal{L}_0$. Define $k_u: U_p \rightarrow \mathcal{K}_1$ as $\Phi_{\varphi}^{-1}(y_u, y'_u, g'_u g_u^{-1})$ and $\hat{h}_u: U_p \rightarrow \mathcal{H}_1$ as $h_u \psi(k_u^{-1})(h'_u)^{-1}$. It follows that

$$([(x'_u, g'_u, y'_u), (y'_u, h'_u, z'_u)], [(x'_u, g'_u, y'_u), (y'_u, h'_u, z'_u)]) \cdot k_u = ([(x_u, g_u, y_u), (y_u, h_u, z_u)], [(x_u, g_u, y_u), (y_u, h_u, z_u)]) \cdot \hat{h}_u,$$

and so the plot $(\text{pr}_1 \circ p, \hat{h}_u)$ is a lift of p to $\mathcal{L}_{0\omega_0} \times_t \mathcal{H}_1$. Thus $A_{\mathcal{H}}$ is subductive, and by **Item 2** of **Proposition 2.7**, it is a diffeomorphism. We thus have a bibundle $\mathcal{G} \curvearrowright^{\chi_0} \mathcal{L}_0 \overset{\omega_0}{\curvearrowright} \mathcal{H}$. \square

We now give an explicit description of the bi-equivariant diffeomorphism in **DBiBund** one obtains from a 2-cell between generalised morphisms in **DGpoid** $[W^{-1}]$.

Proposition 8.21. *Given generalised morphisms $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{\simeq} \mathcal{H}$ admitting a 2-cell between them, there exists a bi-equivariant diffeomorphism between the associated bibundles $\mathcal{G} \curvearrowright^{\chi_0} \mathcal{L}_0 \overset{\omega_0}{\curvearrowright} \mathcal{H}$ and $\mathcal{G} \curvearrowright^{\chi'_0} \mathcal{L}'_0 \overset{\omega'_0}{\curvearrowright} \mathcal{H}$ given by **Proposition 8.20**.*

Proof. Let \mathcal{L} and \mathcal{L}' be the action groupoids of the $(\mathcal{G}\text{-}\mathcal{H})$ -actions on the bibundles $\mathcal{G} \curvearrowright^{\chi_0} \mathcal{L}_0 \overset{\omega_0}{\curvearrowright} \mathcal{H}$ and $\mathcal{G} \curvearrowright^{\chi'_0} \mathcal{L}'_0 \overset{\omega'_0}{\curvearrowright} \mathcal{H}$, and let $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \xrightarrow[\omega]{\simeq} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\chi']{\simeq} \mathcal{L}' \xrightarrow[\omega']{\simeq} \mathcal{H}$ be the corresponding anafunctors. By **Theorem 7.23** and **Lemma 7.25**, there is a unique 2-cell represented by a diagram of the form

$$\begin{array}{ccccc} & & \mathcal{L} & & \\ & \swarrow \chi & \uparrow \text{pr}_1 \simeq & \searrow \omega & \\ \mathcal{G} & \xrightarrow{\simeq} & \mathcal{L}_{\chi} \times_{\chi'} \mathcal{L}' & \xrightarrow{U \Downarrow} & \mathcal{H} \\ & \swarrow \chi' & \downarrow \text{pr}_2 \simeq & \searrow \omega' & \\ & & \mathcal{L}' & & \end{array}$$

an explicit description of U is found **Lemma 7.25**. For $(w, w') \in (\mathcal{L}_{\chi} \times_{\chi'} \mathcal{L}')_0$, define

$$\alpha: \mathcal{L}_0 \rightarrow \mathcal{L}'_0: w \mapsto w' \cdot U(w, w').$$

We claim this is well-defined. Let $w = [(x, g, y), (y, h, z)] \in \mathcal{L}_0$ and $w'_i = [(x, g'_i, y'_i), (y'_i, h'_i, z'_i)] \in \mathcal{L}'_0$ such that $(w, w'_i) \in (\mathcal{L}_{\chi} \times_{\chi'} \mathcal{L}')_0$ for $i = 1, 2$. Since φ' is smoothly fully faithful, there

exists $k' = \Phi_{\varphi'}^{-1}(y'_1, y'_2, g'_2(g'_1)^{-1})$. We have the following diagrams in \mathcal{G} (on the left) and \mathcal{H} (on the right).

$$\begin{array}{ccc}
 & \varphi'(y'_2) & \\
 g'_2 \nearrow & & \searrow \\
 x & \xrightarrow{g} & \varphi(y) & \xrightarrow{\varphi'(k')} & \varphi'(y'_1) \\
 g'_1 \searrow & & & & \\
 & \varphi'(y'_1) & & &
 \end{array}
 \quad
 \begin{array}{ccc}
 & \psi'(y'_2) & \xrightarrow{h'_2} & z'_2 \\
 \psi'(k') \nearrow & & & \uparrow U(w, w'_2) \\
 \psi(y) & \xrightarrow{h} & z & \\
 \psi'(k') \searrow & & & \downarrow U(w, w'_1) \\
 & \psi'(y'_1) & \xrightarrow{h'_1} & z'_1
 \end{array}$$

Keeping in mind that $[(x, \varphi'(k')^{-1}g'_2, y'_1), (y'_1, h'_2\psi'(k'), z'_2)] = w'_2$, the two arrows

$$\left([(u_x, g, u_y), (u_y, h, h^{-1})], [(u_x, g'_1, u_{y'_1}), (u_{y'_1}, h'_1, (h'_1)^{-1})] \right)$$

and

$$\left([(u_x, g, u_y), (u_y, h, h^{-1})], [(u_x, \varphi'(k')^{-1}g'_2, u_{y'_1}), (u_{y'_1}, h'_2\psi'(k'), (h'_2\psi'(k'))^{-1})] \right)$$

have sources (w, w'_1) and (w, w'_2) , resp., and the same target

$$v := \left([(x, g, y), (y, u_{\psi(y)}, \psi(y))], [(x, g'_1, y'_1), (y'_1, u_{\psi'(y'_1)}, \psi'(y'_1))] \right).$$

By the naturality of U

$$h'_1 U(v) = U(w, w'_1)h \quad \text{and} \quad h'_2 \psi'(k') U(v) = U(w, w'_2)h,$$

from which it follows that $U(w, w'_2)^{-1}h'_2\psi'(k') = U(w, w'_1)^{-1}h'_1$. That is, $w'_2 U(w, w'_2) = w'_1 U(w, w'_1)$, and α is well-defined.

To show that α is smooth, let p be a plot of \mathcal{L}_0 . Since $\text{pr}_1: (\mathcal{L}_\chi \times_{\chi'} \mathcal{L}')_0 \rightarrow \mathcal{L}_0$ is a subduction, after shrinking U_p , there is a lift q of p to $(\mathcal{L}_\chi \times_{\chi'} \mathcal{L}')_0$ of p , and $\alpha \circ p = (\text{pr}_2 \circ q) \cdot (U \circ q)$. Smoothness follows.

Symmetrically, there is also a well-defined smooth map $\beta: \mathcal{L}'_0 \rightarrow \mathcal{L}_0$ sending w' to $wU(w, w')^{-1}$ for some $(w, w') \in \mathcal{L}_0 \times_{\chi'} \mathcal{L}'_0$. We need to show that $\beta = \alpha^{-1}$. To this end, we make the following observations, which follow from the naturality of U and the facts that \mathcal{L} and \mathcal{L}' are isomorphic to $\mathcal{G} \times \mathcal{L}_0 \times \mathcal{H}$ and $\mathcal{G} \times \mathcal{L}'_0 \times \mathcal{H}$, resp:

$$U(w, w' \cdot \tilde{h}) = (\tilde{h})^{-1} U(w, w'), \tag{7}$$

$$U(w \cdot \tilde{h}, w') = U(w, w') \tilde{h}, \tag{8}$$

$$U(\tilde{g} \cdot w, w') = U(w, \tilde{g} \cdot w') = U(w, w'). \tag{9}$$

By Equation (7), and the definition of the fibred product $\mathcal{L}_\chi \times_{\chi'} \mathcal{L}'$, one obtains $\beta \circ \alpha(w) = w$. Symmetrically, but using Equation (8), one obtains $\alpha \circ \beta(w') = w'$, and we conclude that $\beta = \alpha^{-1}$. That is, α is a diffeomorphism from \mathcal{L}_0 to \mathcal{L}'_0 . That α is \mathcal{G} -invariant follows from Equation (9); that it is \mathcal{H} -invariant follows from Equations (7) and (8). We conclude that α induces an isomorphism of $(\mathcal{G}\text{-}\mathcal{H})$ -action groupoids; that is, α extends to an isomorphism $\mathcal{L} \rightarrow \mathcal{L}'$. \square

Through [Lemma 8.18](#) we see how to obtain an anafunctor from a bibundle. We now give an explicit way of constructing a transformation out of a bi-equivariant diffeomorphism between bibundles.

Proposition 8.22. *Given bibundles $\mathcal{G} \curvearrowright^l X \curvearrowright^r \mathcal{H}$ and $\mathcal{G} \curvearrowright^{l_Y} Y \curvearrowright^r \mathcal{H}$ and a bi-equivariant diffeomorphism $\alpha: X \rightarrow Y$, there is a transformation S between the associated anafunctors as given in [Lemma 8.18](#).*

Proof. Since α is a bi-equivariant diffeomorphism, it extends to an isomorphism between the action groupoids $\mathcal{K} := \mathcal{G} \times X \times \mathcal{H}$ and $\mathcal{L} := \mathcal{G} \times Y \times \mathcal{H}$, commuting with functors \widehat{l}_X , \widehat{r}_X , \widehat{l}_Y , and \widehat{r}_Y as described in [Lemma 8.18](#). Define $S: \left(\mathcal{K}_{\widehat{l}_X \times \widehat{l}_Y} \times \widehat{\mathcal{L}}\right)_0 \rightarrow \mathcal{H}_1$ by $S(y, w) := d_{\mathcal{H}}(\alpha(y), w)$ where $d_{\mathcal{H}}$ is the division map for the \mathcal{H} -bundle $Y \xrightarrow{r_Y} \mathcal{G}_0$. It follows from the right principality of this bundle that S is well-defined, smooth, and is a natural transformation $\widehat{r}_X \circ \text{pr}_1 \Rightarrow \widehat{r}_Y \circ \text{pr}_2$. That is, we have a transformation between anafunctors:

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \widehat{l}_X & \uparrow & \widehat{r}_X & \\
 \mathcal{G} & \xrightarrow{\simeq} & \mathcal{K} & \xrightarrow{\simeq} & \mathcal{H} \\
 & \circlearrowleft & \mathcal{K}_{\widehat{l}_X \times \widehat{l}_Y} \times \widehat{\mathcal{L}} & \xrightarrow{S \downarrow} & \\
 & \widehat{l}_Y & \downarrow & \widehat{r}_Y & \\
 & & \mathcal{L} & &
 \end{array}$$

□

8.5. Quasi-Inverses in the Bicatagories. We showed in [Lemma 6.13](#) (resp. [Lemma 7.19](#)) that a generalised morphism (resp. anafunctor) in which both functors are weak equivalences admits a quasi-inverse; however, we never showed the converse: a generalised morphism (resp. anafunctor) admitting a quasi-inverse is a span of two weak equivalences. Moreover, in the case of an anafunctor admitting a quasi-inverse, is it a span of two *subductive* weak equivalences? The answer is in one sense no: [Lemma 6.13](#) shows that an anafunctor *as a generalised morphism* has a quasi-inverse if the right functor is only a weak equivalence. However, there will be a transformation from such an anafunctor to an anafunctor in which both weak equivalences are subductive. All of these claims are proven below using bibundles, and the following result of van der Schaaf [[vdS21](#), Theorem 5.31] (see also [[Vil18](#), Lemma 1.3.22] for the same result but in a general context).

Proposition 8.23 (Quasi-Inverses in **DBiBund**). *A right principal bibundle admits a quasi-inverse if and only if it is biprincipal.*

We obtain the equivalent result in $\mathbf{DGpoid}[W^{-1}]$ and $\mathbf{DGpoid}_{\text{ana}}$ in the following proposition.

Proposition 8.24 (Quasi-Inverses in $\mathbf{DGpoid}[W^{-1}]$ and $\mathbf{DGpoid}_{\text{ana}}$). *A generalised morphism admits a quasi-inverse if and only if both of its functors are weak equivalences.*

Moreover, if the generalised morphism is an anafunctor, it admits a transformation to an anafunctor in which both functors are subductive weak equivalences.

Proof. A generalised morphism in which both of its functors are weak equivalences yields a quasi-inverse by [Lemma 6.13](#). Conversely, suppose $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ admits a quasi-inverse. By [Proposition 8.20](#), there is a bibundle $\mathcal{G} \curvearrowright^{l_x} X \curvearrowright^{r_x} \mathcal{H}$ and a 2-cell between its action groupoid and $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$. Since $\mathbf{DGpoid}[W^{-1}]$ and $\mathbf{DBiBund}$ are equivalent bicategories by [Theorem 8.19](#), $\mathcal{G} \curvearrowright^{l_x} X \curvearrowright^{r_x} \mathcal{H}$ also admits a quasi-inverse, and so is biprincipal by [Proposition 8.24](#). A straightforward check similar to the proof of [Lemma 8.15](#) yields that $\widehat{r_X}$ is a (subductive) weak equivalence. That ψ is a weak equivalence now follows from [Proposition 4.13](#) and [Proposition 4.12](#).

If $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ is an anafunctor in which ψ is a subductive weak equivalence, then it admits a quasi-inverse in $\mathbf{DGpoid}_{\text{ana}}$ by [Lemma 7.19](#). Conversely, if $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ is an anafunctor admitting a quasi-inverse, then the proof above shows that it admits a 2-cell to an anafunctor in which both functors are subductive weak equivalences; by [Lemma 7.25](#), we can assume this 2-cell is a transformation. \square

We now end this section with a summary of what we have accomplished so far in this paper.

Theorem 8.25. *The three bicategories of diffeological groupoids $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, and $\mathbf{DBiBund}$ are all equivalent, and the equivalence from $\mathbf{DGpoid}[W^{-1}]$ to $\mathbf{DBiBund}$ can be chosen to factor through $\mathbf{DGpoid}_{\text{ana}}$.*

Proof. This theorem is the combination of [Theorem 7.23](#), [Theorem 8.19](#), [Lemma 8.18](#), and the proofs of [Proposition 8.20](#), [Proposition 8.21](#), and [Proposition 8.22](#). \square

9. LOCAL SUBDUCTIONS & LIE GROUPOIDS

For many purposes, it is useful to restrict one's attention to diffeological groupoids in which the source and target are local subductions; for instance, Lie groupoids satisfy this property: recall that in the category of smooth manifolds, local subductions are exactly surjective submersions [Item 3](#) of [Proposition 2.7](#). In this case, it makes sense to go further and modify the definition of a weak equivalence between such diffeological groupoids [Definition 4.3](#) by requiring Ψ_φ to be a local subduction, and not just a subduction. But it turns out that this modification is unnecessary, as it is automatically satisfied by weak equivalences between these particular diffeological groupoids ([Proposition 9.6](#)). Similarly, when turning to anafunctors, one may wish to replace subductive weak equivalences between these diffeological groupoids with ones that are locally subductive on objects; again, this will be shown to be unnecessary, as they automatically will be locally subductive on objects ([Proposition 9.7](#)).

We give examples indicating that the property of the source and target maps being local subductions is not a Morita invariant; that is, not preserved by Morita equivalences. [Example 9.4](#) provides an example in which there is a weak equivalence from a diffeological

groupoid with locally subductive source and target to a diffeological groupoid whose source and target are not locally subductive. Also, [Example 9.5](#) provides an example of a diffeological groupoid that is not Morita equivalent to any locally subductive diffeological groupoid. Thus it makes sense to restrict to the sub-2-category of **DGpoid** made up of diffeological groupoids whose source and target are local subductions. As suggested by the previous paragraph, the localisation of this sub-2-category is exactly the bicategory in which subductions are required to be local subductions in our definition of diffeological groupoid, weak equivalence, anafunctor, etc., which in turn is an essentially full sub-bicategory of the bicategories studied in the previous sections (see [Theorem 9.19](#)).

This now leads us to Lie groupoids ([Subsection 9.2](#)): we replace locally subductive diffeological groupoids with Lie groupoids and obtain well-known results, but via the diffeological theory established already. This indicates that the study of Lie groupoids is indeed just a special case of the study of diffeological groupoids in all aspects. This is summarised in [Theorem 9.30](#). In this subsection, we also answer the open question [[vdS21](#), Question 7.6] of van der Schaaf in [Corollary 9.32](#).

9.1. Locally Subductive Diffeological Groupoids. Throughout this section, we implement the notation, terminology, and assumption of [Notation 2.5](#); namely, when proving a map is a local subduction by showing that a plot p lifts, *we will always assume that* $0 \in U_p$. Moreover, when we shrink U_p in order to lift the plot, we will always assume that we are replacing the domain U_p with an open neighbourhood of 0, and recycling the symbol U_p . Thus given a smooth map $f: X \rightarrow Y$, to show that it is a local subduction, fix a plot $p = y_u$ and $x_0 \in X$ such that $f(x_0) = y_0$. We wish to find a lift x_u of p to X (and thus implicitly, a lift through x_0). And so on.

We now introduce some terminology inspired by [[vdS20](#); [vdS21](#)].

Definition 9.1 (Locally Subductive Groupoids). A diffeological groupoid is **locally subductive** if its source (and hence target) map is a local subduction. A weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is **pure** if Ψ_φ is a local subduction. A weak equivalence is **locally subductive** if it is locally subductive on objects. Denote by **DGpoid^{ls}** the full sub-2-category of **DGpoid** whose objects are locally subductive diffeological groupoids, by W_{ls} the class of pure weak equivalences in **DGpoid^{ls}**, and by J_{ls} the class of locally subductive weak equivalences in **DGpoid^{ls}**. \diamond

Example 9.2 (Lie Groupoids are Locally Subductive). Since the source map of a Lie groupoid is a surjective submersion, it is immediate that Lie groupoids are locally subductive. //

Example 9.3 (Kernel Groupoid may not be Locally Subductive). Let $\mathbb{Z}/2$ act on \mathbb{R} by reflection. The corresponding action groupoid \mathcal{G} admits a kernel $\ker(\chi_{\mathcal{G}})$ to its characteristic functor $\chi_{\mathcal{G}} = (s, t): \mathbb{Z}/2 \ltimes \mathbb{R} \rightarrow (\mathbb{R}^2 \rightrightarrows \mathbb{R})$ whose source fibres are made up of units except for the source fibre of 0, which is isomorphic to $\mathbb{Z}/2$. It follows that the source map of $\ker(\chi_{\mathcal{G}})$ is not locally subductive, as one cannot lift a non-trivial path through 0 of the object space \mathbb{R} to a path through the non-trivial arrow in the stabiliser at 0. //

The first question one might ask regarding the local subductivity is whether a diffeological groupoid being locally subductive (or perhaps *not* being locally subductive) is a Morita invariant. A second question is whether a diffeological groupoid that is not locally subductive is Morita equivalent to one that is. The following two examples show the answer to these questions is negative in all cases.

Example 9.4 (Weak Equivalences Versus Local Subductivity I). Let $X = \mathbb{R} \amalg \{\infty\}$ be the real line and an extra isolated point denoted ∞ . Let \mathcal{H} be the diffeological groupoid with object space X and arrow space $\{u_x \mid x \in X\} \amalg \{h, h^{-1}\}$ where h is an arrow from $0 \in \mathbb{R}$ to ∞ . Let \mathcal{G} be the trivial groupoid of \mathbb{R} , and let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be the unique functor that on objects is the inclusion of \mathbb{R} to itself in X . Then φ is a weak equivalence (not subductive), \mathcal{H} is not locally subductive (the path $t \mapsto t \in \mathbb{R} \subset X$ does not lift to a path through h with respect to the source map), but \mathcal{G} is locally subductive. It follows that neither local subductivity of diffeological groupoids, nor the lack of this property, are Morita invariants. //

Example 9.5 (Weak Equivalence Versus Local Subductivity II). Let $\mathcal{H} = \ker(\chi_{\mathcal{G}})$ be the kernel groupoid of the action groupoid $\mathcal{G} = \mathbb{Z}/2 \ltimes \mathbb{R}$ as in [Example 9.3](#). Then \mathcal{H} has \mathbb{R} as object space and $\mathbb{R} \amalg \{\tau\}$ as arrow space, where $\{u_0, \tau\} \cong \mathbb{Z}/2$ is the source fibre (and target fibre and stabiliser) of 0. Suppose $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is a weak equivalence. Since φ is smoothly essentially surjective, some connected component of \mathcal{G}_0 must contain a smooth curve C that is sent by φ_0 inductively to $(-\varepsilon, \varepsilon) \subset \mathcal{H}_0$ for some $\varepsilon > 0$ (see [Definition 2.3](#)). Since φ is smoothly fully faithful, the union of the source fibres of \mathcal{G} along C is diffeomorphic to $C \amalg \{\sigma\}$ where $\varphi(\sigma) = \tau$. In particular, there is a smooth path $p: (-\varepsilon, \varepsilon) \rightarrow \mathcal{G}_0$ with image in C that has no lift through σ with respect to the source map. Therefore, \mathcal{G} is not locally subductive.

Suppose further that $\psi: \mathcal{G} \rightarrow \mathcal{K}$ is a weak equivalence. Let $x_1, x_2 \in C$ such that $\psi(x_1) = \psi(x_2)$. Setting $g = \Phi_{\psi}^{-1}(x_1, x_2, u_{\psi(x_1)})$, we have $g = \Phi_{\varphi}^{-1}(x_1, x_2, \varphi(g))$. But $\varphi(g)$ is in some stabiliser of \mathcal{H} , and so $x_1 = x_2$ since $\varphi_0|_C$ is injective. Thus $\psi_0|_C$ is injective. If $q = z_u$ is a plot of $\psi_0(C)$, then smooth essential surjectivity of ψ implies that after shrinking U_q , there is a lift (x'_u, k_u) of z_u . Define $x_u := \psi_0|_C^{-1}(z_u)$ and $g_u := \Phi_{\psi}^{-1}(x_u, x'_u, k_u)$; these *a priori* may not be smooth. But $\varphi(g_u)$ has equal source and target $\varphi(x_u) = \varphi(x'_u)$, and x'_u is smooth. Thus $\varphi(x_u)$ is a plot of $\varphi_0(C)$, and since $\varphi_0|_C$ is an induction, we conclude that x_u is smooth, from which it follows that $\psi_0|_C$ is an induction.

If there is a lift k_u of $\psi \circ p$ to \mathcal{K}_1 through $\psi(\sigma)$, then $g'_u := \Phi_{\psi}^{-1}(\psi_0|_C^{-1}(s(k_u)), \psi_0|_C^{-1}(t(k_u)), k_u)$ is a lift of p through σ , a contradiction. Thus there is no such lift of $\psi \circ p$ through $\psi(\sigma)$, and we conclude that \mathcal{K} is not locally subductive. We conclude that there is no locally subductive diffeological groupoid that is Morita equivalent to \mathcal{H} . //

Due to the above examples, it makes sense to restrict our attention to **DGpoid**^{ls}. But before we apply this restriction, we develop some properties of weak equivalences, pull-backs, and natural transformations detailing their interactions with local subductivity.

Proposition 9.6 (Weak Equivalences and Purenness). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a weak equivalence. Then \mathcal{G} is locally subductive if and only if φ is pure. Consequently, $W_{\text{ls}} = W \cap \mathbf{DGpoid}_1^{\text{ls}}$.*

Proof. Suppose \mathcal{G} is locally subductive. Let $p = y_u$ be a plot of \mathcal{H}_0 , and fix $(x_0, h_0) \in \mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1$ such that $\Psi_\varphi(x_0, h_0) = y_0$. Since Ψ_φ is subductive, after shrinking U_p , there exists a lift (x'_u, h'_u) of p to $\mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1$. Let $g_0 := \Phi_\varphi^{-1}(x_0, x'_0, h'_0 h_0^{-1})$. Since the target map of \mathcal{G} is locally subductive, after shrinking U_p again, there is a lift g_u of x'_u with respect to \mathfrak{t} through g_0 . Then $r: U_p \rightarrow \mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1$ defined by

$$r := (s(g_u), \varphi(g_u)^{-1} h'_u)$$

is a lift of p through (x_0, h_0) . Thus Ψ_φ is a local subduction.

Conversely, suppose Ψ_φ is locally subductive. By **Item 2** of **Remark 4.4**, it suffices to show that given a plot $p = x_u$ of \mathcal{H}_0 and $(x_0, x'_0, h'_0) \in \mathcal{G}_{0,\varphi^2}^2 \times_{(s,\mathfrak{t})} \mathcal{H}_1$, that there is a lift (x_u, x'_u, h_u) of p to $\mathcal{G}_{0,\varphi^2}^2 \times_{(s,\mathfrak{t})} \mathcal{H}_1$. Since Ψ_φ is locally subductive, after shrinking U_p , there is a lift (x'_u, h'_u) of p to $\mathcal{G}_{0,\varphi} \times_{\mathfrak{t}} \mathcal{H}_1$ through (x'_0, h'_0) . Thus (x_u, x'_u, h'_u) is the desired lift. \square

Proposition 9.7 (Subductive Weak Equivalences and Local Subductivity). *Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a subductive weak equivalence. Then \mathcal{G} is locally subductive if and only if \mathcal{H} and φ are. Consequently, $J_{\text{ls}} = J \cap \mathbf{DGpoid}_1^{\text{ls}} \subset W_{\text{ls}}$.*

Proof. Suppose \mathcal{G} is locally subductive. Let $p = y_u$ be a plot of \mathcal{H}_0 and let $h_0 \in \mathcal{H}_1$ such that $s(h_0) = y_0$. Since φ is subductive, after shrinking U_p , there is a lift x_u of y_u to \mathcal{G}_0 ; there is also some $x'_0 \in \mathcal{G}_0$ such that $\varphi(x'_0) = \mathfrak{t}(h_0)$. Let $g_0 = \Phi_\varphi^{-1}(x_0, x'_0, h_0)$. Since \mathcal{G} is locally subductive, after shrinking U_p , there is a lift g_u of x_u with respect to the source map through g_0 . Then $\varphi(g_u)$ is the desired lift of y_u . Thus \mathcal{H} is locally subductive.

Continuing with the same plot $p = y_u$, let $x_0 \in \mathcal{G}_0$ such that $\varphi(x_0) = y_0$. After shrinking U_p , there is a lift x'_u of p to \mathcal{G}_0 , since φ is subductive. Let $g_0 = \Phi_\varphi^{-1}(x_0, x'_0, u_{y_0})$. Since \mathcal{G} is locally subductive, after shrinking U_p , there is a lift g_u of x'_u with respect to the target map of \mathcal{G} through g_0 . Then $s(g_u)$ is the desired lift of p , from which it follows that φ is locally subductive.

Conversely, suppose \mathcal{H} and φ are locally subductive. Let $p = x_u$ be a plot of \mathcal{G} and $g_0 \in \mathcal{G}_1$ with source x_0 . Since \mathcal{H} is locally subductive, after shrinking U_p , there exists a lift h_u of $\varphi(x_u)$ with respect to the source map of \mathcal{H} through $\varphi(g_0)$. Since φ is locally subductive, after shrinking U_p , there exists a lift x'_u of $\mathfrak{t}(h_u)$ through $\mathfrak{t}(g_0)$. Then $g_u = \Phi_\varphi^{-1}(x_u, x'_u, h_u)$ is the desired lift, from which it follows that \mathcal{G} is locally subductive.

The last statement follows from **Proposition 9.6**. \square

Due to **Proposition 9.6** and **Proposition 9.7**, if one wishes to work with pure and/or locally subductive weak equivalence between locally subductive diffeological groupoids, then one only need to restrict to $\mathbf{DGpoid}^{\text{ls}}$ and use the theory from the previous sections. Purenness and/or local subductivity of weak equivalences is obtained for free. We now show pullbacks are well behaved in this setting.

Proposition 9.8 (Weak Pullbacks and Local Subductivity). *Let \mathcal{G} be locally subductive, $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ be a weak equivalence, and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ a functor. Then $\text{pr}_3: \mathcal{G}_{\varphi} \times_{\psi}^{\text{w}} \mathcal{H} \rightarrow \mathcal{H}$ is a locally subductive weak equivalence. If furthermore \mathcal{H} is locally subductive, then so is $\mathcal{G}_{\varphi} \times_{\psi}^{\text{w}} \mathcal{H}$.*

Proof. By **Item 1** of **Proposition 5.10**, $\text{pr}_3: \mathcal{G}_{\varphi} \times_{\psi}^{\text{w}} \mathcal{H} \rightarrow \mathcal{H}$ is a subductive weak equivalence. Fix a plot $p = y_u$ of \mathcal{H}_0 and $(x_0, k_0, y_0) \in (\mathcal{G}_{\varphi} \times_{\psi}^{\text{w}} \mathcal{H})_0$. By **Proposition 9.6**, Ψ_{φ} is a local subduction, and so after shrinking U_p , there is a lift (x_u, k_u^{-1}) of $\psi \circ p$ through (x_0, k_0^{-1}) . Then (x_u, k_u, y_u) is the desired lift of p , from which it follows that pr_3 is locally subductive. The second statement follows from **Proposition 9.7**. \square

Proposition 9.9 (Strict Pullbacks and Local Subductivity). *Let \mathcal{G} be locally subductive, $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ a subductive weak equivalence, and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ a functor. Then $\text{pr}_2: \mathcal{G}_{\varphi} \times_{\psi} \mathcal{H} \rightarrow \mathcal{H}$ is a locally subductive weak equivalence. If furthermore \mathcal{H} is locally subductive, then so is $\mathcal{G}_{\varphi} \times_{\psi} \mathcal{H}$.*

Proof. That $\text{pr}_2: \mathcal{G}_{\varphi} \times_{\psi} \mathcal{H}$ is a locally subductive weak equivalence follows from **Item 5** of **Proposition 2.7** and **Item 2** of **Proposition 5.4**. Suppose further that \mathcal{H} is locally subductive. Let $p = (x_u, y_u)$ be a plot of $(\mathcal{G}_{\varphi} \times_{\psi} \mathcal{H})_0$ and $(g_0, h_0) \in (\mathcal{G}_{\varphi} \times_{\psi} \mathcal{H})_1$ such that $s(g_0, h_0) = (x_0, y_0)$. After shrinking U_p , there is a lift h_u of y_u with respect to the source map of \mathcal{H} through h_0 . By **Proposition 9.7**, φ is locally subductive. Thus after shrinking U_p again, there is a lift x'_u of $\psi(t(h_u))$ through $t(g_0)$. Let $g_u := \Phi_{\varphi}^{-1}(x_u, x'_u, \psi(h_u))$. Then (g_u, h_u) is a desired lift of (x_u, y_u) through (g_0, h_0) . The result follows. \square

Example 7.3 provides an example of how natural transformations do not necessarily preserve locally subductive weak equivalences. However, they do preserve pureness of weak equivalences.

Proposition 9.10 (Natural Transformations and Pureness). *Let $\varphi, \psi: \mathcal{G} \rightarrow \mathcal{H}$ be two functors and let $S: \varphi \Rightarrow \psi$ be a natural transformation. If one of φ or ψ is a pure weak equivalence, so is the other.*

Proof. Suppose that φ is a pure weak equivalence. That ψ is a weak equivalence is **Proposition 4.13**. By **Proposition 9.6**, \mathcal{G} is locally subductive, and consequently, ψ is pure. \square

In the remainder of this subsection, an important result will be that any generalised morphism between locally subductive diffeological groupoids admits a 2-cell to a generalised morphism made up of locally subductive diffeological groupoids. To accomplish this, we will apply the principle mentioned in the introduction: the bibundle between two locally subductive diffeological groupoids exhibits the properties we require. In **[vdS21, Subsection 4.4.3]**, van der Schaaf develops how the theory of bibundles interacts with local subductivity. We generalise some of those results in the following proposition.

Proposition 9.11 (Bibundles and Local Subductivity). *Let $\mathcal{G} \overset{l}{\curvearrowright} Z \overset{r}{\curvearrowright} \mathcal{H}$ be a right principal bibundle.*

- (1) If \mathcal{H} is locally subductive, then l_Z is a local subduction.
- (2) If \mathcal{G} and \mathcal{H} are locally subductive, then l_Z and the action groupoid $\mathcal{G} \times Z \rtimes \mathcal{H}$ are locally subductive.

Proof. Suppose \mathcal{H} is locally subductive. Let $p = x_u$ be a plot of \mathcal{G}_0 , and $z_0 \in Z$ such that $l_Z(z_0) = x_0$. Since l_Z is a subduction, after shrinking U_p , there is a lift z'_u of x_u to Z . Since $l_Z(z_0) = l_Z(z'_0)$, there is a unique $h_0 \in \mathcal{H}_1$ so that $z_0 = z'_0 h_0$. Since \mathcal{H} is locally subductive, after shrinking U_p again, there is a lift h_u of $r_Z(z'_u)$ through h_0 with respect to the target map of \mathcal{H} . Letting $z_u := z'_u h_u$, we obtain

$$l_Z(z_u) = l_Z(z'_u h_u) = l_Z(z'_u) = x_u$$

using the \mathcal{H} -invariance of l_Z . Thus z_u is the desired lift. This proves [Item 1](#). [Item 2](#) follows from [Item 1](#), [Lemma 8.18](#), and [Proposition 9.7](#). \square

We are now in a position to develop the main results of this subsection. Denote by \mathfrak{S}^{ls} the restrict of the spanisation pseudofunctor \mathfrak{S} to $\mathbf{DGpoid}^{\text{ls}}$.

Theorem 9.12. *The class W_{ls} admits a right calculus of fractions, the bicategory of fractions $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is an essentially full sub-bicategory of $\mathbf{DGpoid}[W^{-1}]$, and the following diagram of bicategories commutes.*

$$\begin{array}{ccc} \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{S}^{\text{ls}}} & \mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}] \\ \downarrow & & \downarrow \\ \mathbf{DGpoid} & \xrightarrow{\mathfrak{S}} & \mathbf{DGpoid}[W^{-1}]. \end{array}$$

Proof. We begin by following [Definition 6.1](#), checking each item. All identity functors of locally subductive diffeological groupoids are contained in W_{ls} ; this is [Item 1](#) of [Definition 6.1](#). That W_{ls} is closed under composition is inherited from W ; this is [Item 2](#) of [Definition 6.1](#). That [Item 3](#) of [Definition 6.1](#) is satisfied is [Proposition 9.8](#). [Item 4](#) and [Item 5](#) of [Definition 6.1](#) are inherited from W . Thus W_{ls} admits a right calculus of fractions.

It is immediate that $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is a sub-bicategory of $\mathbf{DGpoid}[W^{-1}]$. To show that it is an essentially full sub-bicategory, we need to show that the inclusion pseudofunctor is essentially full on generalised morphisms and fully faithful on 2-cells.

Fix a generalised morphism $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ in which \mathcal{G} and \mathcal{H} are locally subductive diffeological groupoids. By [Proposition 8.20](#), there is an anafunctor $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{\simeq} \mathcal{H}$ in which \mathcal{K}' is the action groupoid $\mathcal{G} \times Z \rtimes \mathcal{H}$ of a bibundle $\mathcal{G} \curvearrowright^Z Z \curvearrowright^r \mathcal{H}$, $\varphi' = \widehat{l}_Z$ and $\psi' = \widehat{r}_Z$, using the notation of [Lemma 8.18](#); moreover there is a 2-cell from $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$ to $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{\simeq} \mathcal{H}$. By [Item 1](#) of [Proposition 9.11](#), φ' is local subductive. By [Proposition 9.7](#), \mathcal{K}' is locally subductive. Thus the inclusion of bicategories is essentially full on generalised morphisms.

Fix two generalised morphisms $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow{\psi} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow{\psi'} \mathcal{H}$ in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, and a 2-cell between them in $\mathbf{DGpoid}[W^{-1}]$ represented by the following diagram:

$$\begin{array}{ccccc}
 & & \mathcal{K} & & \\
 & \swarrow \varphi & \uparrow \alpha \simeq & \searrow \psi & \\
 \mathcal{G} & & \mathcal{L} & & \mathcal{H} \\
 & \swarrow \varphi' & \downarrow \alpha' \simeq & \searrow \psi & \\
 & & \mathcal{K}' & &
 \end{array}$$

$\Downarrow S_1$ $\Downarrow S_2$

By [Item 3](#) of [Remark 7.24](#), there are 2-cells between the two generalised morphisms and their anafunctisations $\mathcal{G} \xleftarrow[\tilde{\varphi}]{\simeq} \tilde{\mathcal{K}} \xrightarrow{\tilde{\psi}} \mathcal{H}$ and $\mathcal{G} \xleftarrow[\tilde{\varphi}']{\simeq} \tilde{\mathcal{K}}' \xrightarrow{\tilde{\psi}'} \mathcal{H}$. By [Proposition 9.8](#) and [Proposition 9.9](#), $\tilde{\mathcal{K}}$ and $\tilde{\mathcal{K}}'$ are locally subductive. A vertical composition of these three 2-cells (with the last one inverted) yields a 2-cell between the two anafunctisations. By [Lemma 7.25](#), this diagram is equivalent to a transformation $U: (\tilde{\mathcal{K}}_{\tilde{\varphi}} \times_{\tilde{\varphi}'} \tilde{\mathcal{K}}')_0 \rightarrow \mathcal{H}_1$; by [Proposition 9.9](#), $\tilde{\mathcal{K}}_{\tilde{\varphi}} \times_{\tilde{\varphi}'} \tilde{\mathcal{K}}'$ is locally subductive. Now compose this transformation U with the inverses of the 2-cells we vertically composed with originally, undoing the anafunctisations of the generalised morphisms. This uses the weak pullback on the level of diagrams (see [Definition 6.7](#)), but [Proposition 9.8](#) maintains the local subductivity of the groupoids involved. We now have a diagram representing the original 2-cell between the original generalised morphisms, contained entirely in the category $\mathbf{DGpoid}^{\text{ls}}$. We conclude that $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is an essentially full sub-bicategory of $\mathbf{DGpoid}[W^{-1}]$.

Since the spanisation of a functor between two locally subductive diffeological groupoids is a generalised morphism in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, that the diagram commutes is immediate, and the proof is complete. \square

We now strive to see how local subductivity interacts with anafunctors. Let \mathfrak{A}^{ls} be the restriction of the anafunctor pseudofunctor \mathfrak{A} to $\mathbf{DGpoid}^{\text{ls}}$.

Theorem 9.13. *$(\mathbf{DGpoid}^{\text{ls}}, J_{\text{ls}})$ is a 2-site, induces a bicategory $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ of anafunctors which is an essentially full sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}$ and a localisation of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} , and the following diagram commutes.*

$$\begin{array}{ccc}
 \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{A}^{\text{ls}}} & \mathbf{DGpoid}_{\text{ana}}^{\text{ls}} \\
 \downarrow & & \downarrow \\
 \mathbf{DGpoid} & \xrightarrow{\mathfrak{A}} & \mathbf{DGpoid}_{\text{ana}}
 \end{array}$$

Proof. To prove that $(\mathbf{DGpoid}^{\text{ls}}, J_{\text{ls}})$ is a 2-site, we check each condition in [Definition 7.1](#). All identity functors of locally subductive diffeological groupoids are contained in J_{ls} . That J_{ls} is closed under composition and base change is inherited from J and [Proposition 9.9](#). That elements of J_{ls} are representably fully faithful and co-fully faithful is also inherited from J .

That the corresponding anafunctor bicategory $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ is a sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}$ is immediate. To show that the inclusion pseudofunctor is essentially full on anafunctors, one follows the corresponding proof for the inclusion of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ into $\mathbf{DGpoid}[W^{-1}]$ of [Theorem 9.12](#), but starting with an anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ instead of an arbitrary generalised morphism. The 2-cell can be represented by a transformation by [Lemma 7.25](#). The inclusion pseudofunctor is full on transformations due to [Proposition 9.9](#), completing the proof that $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ is an essentially full sub-bicategory. Therefore, the application of Pronk's Comparison Theorem in the proof of [Theorem 7.23](#) still applies when our focus is restricted to $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, proving that $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ is a localisation of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} .

By [Proposition 9.8](#), the anafunctorisation of a functor between locally subductive diffeological groupoids is an anafunctor in $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, and so the diagram commutes, and the proof is complete. \square

We point out as an aside that in Roberts' terminology, a so-called J_{ls} -locally split functor is a J -locally split functor between locally subductive diffeological groupoids that fits appropriately in the locally subductive setting. We give details in the following example.

Example 9.14 (J_{ls} -Locally Split). Suppose \mathcal{G} and \mathcal{H} are locally subductive and $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ is smoothly fully faithful. Then φ is a pure weak equivalence if and only if it is J_{ls} -locally split; that is, using the notation of [Lemma 7.7](#), φ is J -locally split, \mathcal{K} is locally subductive, and ψ is a pure locally subductive weak equivalence. Indeed, if φ is a pure weak equivalence, then we follow the proof of [Lemma 7.7](#). [Proposition 9.8](#) implies that \mathcal{K} is locally subductive, and [Proposition 9.6](#) and [Proposition 9.7](#) imply that ψ is pure and locally subductive. The converse is immediate from [Lemma 7.7](#) and [Proposition 9.6](#). \parallel

We now move onto bibundles, borrowing terminology from [[vds20](#); [vds21](#)].

Definition 9.15 (Purely Right Principal Bibundles). A right principal bibundle is **purely** right principal if the right anchor map is a local subduction. If the bibundle is biprincipal and the right anchor map is also a local subduction, then call the bibundle a **pure Morita equivalence**. Let $\mathbf{DBiBund}^{\text{ls}}$ be the sub-bicategory of $\mathbf{DBiBund}$ of locally subductive diffeological groupoids with purely right principal bibundles as 1-cells. \diamond

Let \mathfrak{B}^{ls} be the restriction of the bibundlisation pseudofunctor \mathfrak{B} to $\mathbf{DGpoid}^{\text{ls}}$.

Theorem 9.16. $\mathbf{DBiBund}^{\text{ls}}$ is a full sub-bicategory of $\mathbf{DBiBund}$ and a localisation of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} , and the following diagram commutes.

$$\begin{array}{ccc} \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{B}^{\text{ls}}} & \mathbf{DBiBund}^{\text{ls}} \\ \downarrow & & \downarrow \\ \mathbf{DGpoid} & \xrightarrow{\mathfrak{B}} & \mathbf{DBiBund} \end{array}$$

Proof. To show that the inclusion pseudofunctor of $\mathbf{DBiBund}^{\text{ls}}$ into $\mathbf{DBiBund}$ is full, we must show that every right principal bibundle between locally subductive diffeological groupoids is purely right principal. But this is immediate from [Item 1](#) of [Proposition 9.11](#). Therefore, the application of Pronk’s Comparison Theorem in the proof of [Theorem 8.19](#) still applies when our focus is restricted to $\mathbf{DBiBund}^{\text{ls}}$, proving that $\mathbf{DBiBund}^{\text{ls}}$ is a localisation of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} . By [Item 1](#) of [Proposition 9.11](#), bibundlisation of a functor between locally subductive diffeological groupoids yields a purely right principal bibundle, and so the diagram commutes. \square

Remark 9.17. As a consequence of [Theorems 9.12](#), [9.13](#) and [9.16](#), the pseudonatural transformations making the triangles below 2-commute

$$\begin{array}{ccccc} \mathbf{DGpoid}[W^{-1}] & \xrightarrow{\simeq} & \mathbf{DGpoid}_{\text{ana}} & \xrightarrow{\simeq} & \mathbf{DBiBund} \\ \uparrow \mathfrak{G} & \searrow \mathfrak{A} & \searrow \mathfrak{B} & \nearrow & \\ \mathbf{DGpoid} & & & & \end{array}$$

restrict to the locally subductive sub-bicategories, yielding the following 2-commutative triangles

$$\begin{array}{ccccc} \mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}] & \xrightarrow{\simeq} & \mathbf{DGpoid}_{\text{ana}}^{\text{ls}} & \xrightarrow{\simeq} & \mathbf{DBiBund}^{\text{ls}} \\ \uparrow \mathfrak{G}^{\text{ls}} & \searrow \mathfrak{A}^{\text{ls}} & \searrow \mathfrak{B}^{\text{ls}} & \nearrow & \\ \mathbf{DGpoid}^{\text{ls}} & & & & \end{array}$$

┘

Remark 9.18. In [Proposition 8.23](#), we saw that a right principal bibundle admits a quasi-inverse if and only if it is biprincipal. In $\mathbf{DBiBund}^{\text{ls}}$, one could ask for a similar result, but where “biprincipal” is replaced with “a pure Morita equivalence”. It turns out a biprincipal bundle between locally subductive diffeological groupoids is automatically a pure Morita equivalence, as proven in [[vdS21](#), Proposition 7.5]. We give a proof of this in [Subsection 10.1](#) (see [Lemma 10.4](#)).

By [Proposition 8.24](#), a generalised morphism in $\mathbf{DGpoid}[W^{-1}]$ admits a quasi-inverse if and only if both of its functors are weak equivalences. In $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, these are both pure weak equivalences by [Proposition 9.6](#). The same proposition in conjunction with [Proposition 9.7](#) shows if the generalised morphism is an anafunctor, then there is a transformation from it to an anafunctor in which both functors are locally subductive weak equivalences. \square

We conclude this subsection with the following summary of the above.

Theorem 9.19. $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, and $\mathbf{DBiBund}^{\text{ls}}$ are all equivalent bicategories, are all localisations of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} , and are essentially full sub-bicategories of $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, and $\mathbf{DBiBund}$, resp.

9.2. Lie Groupoids. We now restrict our attention to Lie groupoids, which are examples of locally subductive diffeological groupoids. We will constantly throughout this subsection use the fact that the local subductions between manifolds are exactly the surjective submersions (Item 3 of Proposition 2.7). Many of the results of this section are already well-known; however, we show that these results come directly from the theory developed thus far for diffeological groupoids.

Definition 9.20 (Lie and Surjective Submersive Weak Equivalences). A weak equivalence $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ between Lie groupoids is **Lie** if Φ_φ and Ψ_φ are smooth maps between manifolds and φ is pure (that is, Ψ_φ is a surjective submersion). A weak equivalence between Lie groupoids is **surjective submersive** if it is locally subductive; that is, φ_0 is surjective submersive. Denote by $\mathbf{LieGpoid}$ the full sub-2-category of \mathbf{DGpoid} whose objects are Lie groupoids, by W_{Lie} the class of Lie weak equivalences in $\mathbf{LieGpoid}$, and by J_{Lie} the class of surjective submersive weak equivalences in $\mathbf{LieGpoid}$. \diamond

Similar to the property of local subductivity of diffeological groupoids, a diffeological groupoid being Lie (or *not* being Lie) is not a Morita invariant. For instance, the trivial groupoid of a point, which is Lie, is Morita equivalent to the trivial groupoid of any diffeological space X . And not surprisingly, some diffeological groupoids that are not Lie do not admit a Morita equivalence to a Lie groupoid: Example 9.5 gives an example of a diffeological groupoid that is not locally subductive, and any diffeological groupoid that is Morita equivalent to it cannot be locally subductive either, and hence not Lie. We now move onto how weak equivalences and pullbacks interact with the Lie and surjective submersive properties.

Proposition 9.21 (Weak Equivalence between Lie Groupoids). *Let \mathcal{G} and \mathcal{H} be Lie groupoids and let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a weak equivalence. Then φ is a Lie weak equivalence. Consequently, $W_{\text{Lie}} = W_{\text{ls}} \cap \mathbf{LieGpoid}_1 = W \cap \mathbf{LieGpoid}_1$. Furthermore, if φ also subductive, then it is surjective submersive. Consequently, $J_{\text{Lie}} = J_{\text{ls}} \cap \mathbf{LieGpoid}_1 = J \cap \mathbf{LieGpoid}_1$.*

Proof. Since \mathcal{G}_1 is a manifold and Φ_φ is a diffeomorphism, its codomain is also a manifold. Since the target map of \mathcal{H} is a surjective submersion, $\mathcal{G}_{0_\varphi \times_t \mathcal{H}_1}$ is a manifold. Since \mathcal{G} is Lie, it is locally subductive, and so Ψ_φ is a local subduction by Proposition 9.6, and hence a surjective submersion. Finally, if φ is subductive, then it is locally subductive by Proposition 9.7, and hence a surjective submersion. \square

Pullbacks work in the Lie category as well.

Proposition 9.22 (Weak Pullbacks of Lie Groupoids). *Let \mathcal{G} , \mathcal{H} , and \mathcal{K} be Lie groupoids and $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ functors with φ a weak equivalence. The weak pullback $\mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ is a Lie groupoid and pr_3 is a surjective submersive weak equivalence.*

Proof. By [Item 2](#) of [Remark 5.7](#),

$$(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_0 = ((\mathcal{G}_0)_\varphi \times_s (\mathcal{K}_1))_{\text{topr}_2} \times_\psi \mathcal{H}_0.$$

Since s is a surjective submersion, $(\mathcal{G}_0)_\varphi \times_s (\mathcal{K}_1)$ is a manifold. The map $(\text{id}_{\mathcal{G}_0}, \text{inv}): \mathcal{G}_0 \times \mathcal{K}_1 \rightarrow \mathcal{G}_0 \times \mathcal{K}_1$ restricts to a diffeomorphism making the following diagram commute.

$$\begin{array}{ccc} (\mathcal{G}_0)_\varphi \times_s (\mathcal{K}_1) & \xrightarrow{(\text{id}_{\mathcal{G}_0}, \text{inv})} & (\mathcal{G}_0)_\varphi \times_t (\mathcal{K}_1) \\ & \searrow \text{topr}_2 & \swarrow \Psi_\varphi \\ & & \mathcal{K}_0 \end{array}$$

The map Ψ_φ is a surjective submersion by [Proposition 9.21](#), and so it follows that topr_2 in the diagram is a surjective submersion. This implies that $(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_0$ is a manifold.

Similarly, to show that $(\mathcal{G}_\varphi \times_\psi^w \mathcal{H})_1$ is a manifold, it suffices to show that

$$t \circ \text{pr}_2: (\mathcal{G}_1)_{s \circ \varphi} \times_s (\mathcal{K}_1) \rightarrow \mathcal{K}_0$$

is a surjective submersion. But this map factors as

$$(\mathcal{G}_1)_{s \circ \varphi} \times_s (\mathcal{K}_1) \xrightarrow{s \times \text{id}_{\mathcal{K}_1}} (\mathcal{G}_0)_\varphi \times_s (\mathcal{K}_1) \xrightarrow{\text{topr}_2} \mathcal{K}_0,$$

where the second map is the map topr_2 from the previous paragraph, which is a surjective submersion. Thus it suffices to show that $s \times \text{id}_{\mathcal{K}_1}$ is a surjective submersion, which is immediate since the source map of \mathcal{G} is surjective submersive.

That $\mathcal{G}_\varphi \times_\psi^w \mathcal{H}$ is a locally subductive diffeological groupoid and pr_3 is a surjective submersion follows from [Proposition 9.8](#). \square

Proposition 9.23. *Let \mathcal{G} , \mathcal{H} , and \mathcal{K} be Lie groupoids and $\varphi: \mathcal{G} \rightarrow \mathcal{K}$ and $\psi: \mathcal{H} \rightarrow \mathcal{K}$ functors with φ a subductive weak equivalence. Then $\mathcal{G}_\varphi \times_\psi \mathcal{H}$ is a Lie groupoid and $\text{pr}_2: \mathcal{G}_\varphi \times_\psi \mathcal{H}$ is a surjective submersive weak equivalence.*

Proof. Since φ is a subductive weak equivalence and \mathcal{G} is locally subductive, φ is locally subductive by [Proposition 9.7](#), and hence surjective submersive. Thus $(\mathcal{G}_\varphi \times_\psi \mathcal{H})_0$ is a manifold. To show that $(\mathcal{G}_\varphi \times_\psi \mathcal{H})_1$ is a manifold, it suffices to show that $\varphi_1: \mathcal{G}_1 \rightarrow \mathcal{K}_1$ is a surjective submersion. That φ_1 is surjective follows from φ being a subductive weak equivalence. To show that it is locally subductive, fix a plot $p = k_u$ of \mathcal{K}_1 and $g_0 \in \mathcal{G}_1$ such that $\varphi(g_0) = k_0$. Let $z_u = s(k_u)$ and $z'_u = t(k_u)$. Since φ_0 is locally subductive, after shrinking U_p , there exist lifts x_u and x'_u of z_u and z'_u through $s(g_0)$ and $t(g_0)$, resp. Define $g_u := \Phi_\varphi^{-1}(x_u, x'_u, k_u)$; this is the desired lift.

By [Proposition 9.9](#), $\mathcal{G}_\varphi \times_\psi \mathcal{H}$ is locally subductive, and hence a Lie groupoid. Also by [Proposition 9.9](#), pr_2 is a surjective submersive weak equivalence. \square

Similar to the previous subsection, we show that any generalised morphism between Lie groupoids admits a 2-cell to a generalised morphism made up of Lie groupoids. The proof works in $\mathbf{DGpoid}_{\text{ana}}$, and uses the idea behind [Example 7.16](#) as well as the Axiom of Choice.

Proposition 9.24. *Let \mathcal{G} be a Lie groupoid and $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ a generalised morphism in $\mathbf{DGpoid}[W^{-1}]$. There is a Lie groupoid \mathcal{K}' , an anafunctor $\mathcal{G} \xleftarrow[\varphi']{\simeq} \mathcal{K}' \xrightarrow[\psi']{} \mathcal{H}$, and a 2-cell between the two generalised morphisms.*

Proof. By [Item 3](#) of [Remark 7.24](#), without loss of generality, we may assume that $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{} \mathcal{H}$ is an anafunctor. Let $\mathcal{U} = \{i_\mu: U_\mu \rightarrow \mathcal{G}_0\}_{\mu \in A}$ be a countable open cover of \mathcal{G}_0 . Without loss of generality, assume that each U_μ is an open subset of \mathbb{R}^n where $n = \dim \mathcal{G}_0$. Then \mathcal{U} is a generating family of the diffeology on \mathcal{G}_0 . Since φ_0 is a subduction, for each μ there is a countable open cover $\mathcal{V}_\mu = \{V_{\mu,\nu}\}_{\nu \in B_\mu}$ of U_μ such that $i_\mu|_{V_{\mu,\nu}}$ admits a lift $j_{\mu,\nu}$ to \mathcal{K}_0 . Let $\mathcal{V} := \{\varphi \circ j_{\mu,\nu}\}_{\mu \in A, \nu \in B_\mu}$. Then \mathcal{V} is a generating family of the diffeology on \mathcal{G}_0 , $\mathcal{N}(\mathcal{V})_0$ is a manifold, and $\text{ev}_\mathcal{V}: \mathcal{N}(\mathcal{V})_0 \rightarrow \mathcal{G}_0$ is a surjective submersion. Hence the base change $\mathcal{G}[\mathcal{N}(\mathcal{V})_0]$ is a Lie groupoid and $\varphi' := (\text{ev}_\mathcal{V}, \text{pr}_3)$ sending objects $v \in V_{\mu,\nu}$ to $\varphi(j_{\mu,\nu}(v))$ and arrows $(v_1, v_2, g) \in (\mathcal{N}(\mathcal{V})_{0, \text{ev}_\mathcal{V}}^2 \times_{(s,t)} \mathcal{G}_1)$ to g is a subductive weak equivalence.

Due to its construction, there is a local diffeomorphism I from $\mathcal{N}(\mathcal{V})_0$ to $\mathcal{N}(\mathcal{D}_{\mathcal{X}_0})_0$, the nebula of the diffeology $\mathcal{D}_{\mathcal{X}_0}$ with evaluation map $\text{ev}_\mathcal{X}$, defined as follows. Since $\mathcal{N}(\mathcal{V})_0 := \coprod_{q \in \mathcal{V}} U_q$, and for each $q \in \mathcal{V}$, there is a plot p_q of \mathcal{K}_0 so that $q = \varphi \circ p_q$, in which case $U_q = U_{p_q} \subseteq \mathcal{N}(\mathcal{D}_{\mathcal{X}_0})_0$. Define $I|_{U_q} = \text{id}_{U_{p_q}}$ (this is where the Axiom of Choice is utilised). Let $\psi': \mathcal{G}[\mathcal{N}(\mathcal{V})_0] \rightarrow \mathcal{H}$ be the functor given by $\psi'_0 := \psi \circ \text{ev}_\mathcal{X} \circ I$ and

$$\psi'_1(v_1, v_2, g) := \psi(\Phi_\varphi^{-1}(\text{ev}_\mathcal{X}(v_1), \text{ev}_\mathcal{X}(v_2), g)).$$

Set $\mathcal{K}' = \mathcal{G}[\mathcal{N}(\mathcal{V})_0]$ and let $\mathcal{L} := \mathcal{K}_\varphi \times_{\varphi'} \mathcal{K}'$ and $S: \psi \circ \text{pr}_1 \Rightarrow \psi' \circ \text{pr}_2$ be given by the smooth map

$$S: \mathcal{L}_0 \rightarrow \mathcal{H}_1: (y, v) \mapsto \psi(\Phi_\varphi^{-1}(y, \text{ev}_\mathcal{X}(v), u_{\varphi(y)})).$$

By [Item 2](#) of [Proposition 5.4](#), it suffices to show that S is a natural transformation.

Fix an arrow $(k, (v_1, v_2, g))$ from (y_1, v_1) to (y_2, v_2) in \mathcal{L}_1 . We need to show that

$$\psi(\Phi_\varphi^{-1}(\text{ev}_\mathcal{X}(v_1), \text{ev}_\mathcal{X}(v_2), g) \cdot \Phi_\varphi^{-1}(y_1, \text{ev}_\mathcal{X}(v_1), u_{\varphi(y_1)})) = \psi(\Phi_\varphi^{-1}(y_2, \text{ev}_\mathcal{X}(v_2), u_{\varphi(y_2)}) \cdot k).$$

But this follows from the fact that $\varphi(k) = g$ and [Equation \(2\)](#). \square

We now emulate [Subsection 9.1](#), and consider localisations of $\mathbf{LieGpoid}$ as sub-bicategories of the localisations of \mathbf{DGpoid} . Let $\mathfrak{S}^{\text{Lie}}$ be the restriction of the spanisation pseudofunctor \mathfrak{S} to $\mathbf{LieGpoid}$.

Theorem 9.25. *The class W_{Lie} admits a right calculus of fractions; the bicategory $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ is an essentially full sub-bicategory of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ and hence of*

$\mathbf{DGpoid}[W^{-1}]$, and is a localisation of $\mathbf{LieGpoid}$ at W_{Lie} ; and the following diagram of bicategories commutes.

$$\begin{array}{ccc} \mathbf{LieGpoid} & \xrightarrow{\mathfrak{S}^{\text{Lie}}} & \mathbf{LieGpoid}[W_{\text{Lie}}^{-1}] \\ \downarrow & & \downarrow \\ \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{S}^{\text{ls}}} & \mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}] \end{array}$$

Proof. To prove W_{Lie} admits a right calculus of fractions, we follow [Definition 6.1](#), checking each item. All identity functors of Lie groupoids are contained in W_{Lie} ; this is [Item 1](#) of [Definition 6.1](#). That W_{Lie} is closed under composition is inherited from W ; this is [Item 2](#) of [Definition 6.1](#). That [Item 3](#) of [Definition 6.1](#) is satisfied is [Proposition 9.22](#). W_{Lie} inherits [Item 4](#) of [Definition 6.1](#) from W , and [Item 5](#) of [Definition 6.1](#) is [Proposition 9.10](#) and the fact that any weak equivalence between Lie groupoids is Lie. Thus W_{Lie} admits a right calculus of fractions.

It is immediate that $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ is a sub-bicategory of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$. To show that it is essentially full on 1-cells, we need to prove that for any generalised morphism in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ between Lie groupoids, there is a generalised morphism in $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ and a 2-cell in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ between them. The existence of the generalised morphism in $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ is given by [Proposition 9.24](#), which gives a 2-cell in $\mathbf{DGpoid}[W^{-1}]$ between the two generalised morphisms. That this 2-cell is in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is implied by [Theorem 9.12](#).

We also need to show that the inclusion pseudofunctor $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}] \rightarrow \mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is full on 2-cells; that is, any 2-cell of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ between Lie groupoids has a representative diagram in $\mathbf{LieGpoid}$. To prove this, one follows the relevant part of the proof of [Theorem 9.12](#) and notices that one remains in the Lie category throughout the construction due to [Proposition 9.22](#) and [Proposition 9.23](#). Thus $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ is an essentially full sub-bicategory of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, and by [Theorem 9.12](#) of $\mathbf{DGpoid}[W^{-1}]$ as well. Similar to the proof that $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ is a localisation of $\mathbf{DGpoid}^{\text{ls}}$ at W_{ls} , it follows that $\mathbf{LieGpoid}[W_{\text{Lie}}^{-1}]$ is a localisation of $\mathbf{LieGpoid}$ at W_{Lie} .

For the commutative diagram, note that $\mathfrak{S}^{\text{Lie}}$ takes values in $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, and so the diagram is well-defined; that it commutes is immediate, and the proof is complete. \square

Moving onto surjective submersive weak equivalences, we establish an anafunctorial version of the above. Let $\mathfrak{A}^{\text{Lie}}$ be the restriction of the anafunctorisation pseudofunctor \mathfrak{A} to $\mathbf{LieGpoid}$.

Theorem 9.26. *($\mathbf{LieGpoid}, J_{\text{Lie}}$) is a 2-site; induces a bicategory $\mathbf{LieGpoid}_{\text{ana}}$ of anafunctors which is an essentially full sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ and hence of $\mathbf{DGpoid}_{\text{ana}}$,*

and is a localisation of $\mathbf{LieGpoid}$ at $W_{\mathbf{Lie}}$; and the following diagram commutes.

$$\begin{array}{ccc} \mathbf{LieGpoid} & \xrightarrow{\mathfrak{J}_{\mathbf{Lie}}} & \mathbf{LieGpoid}_{\text{ana}} \\ \downarrow & & \downarrow \\ \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{J}_{\text{ls}}} & \mathbf{DGpoid}_{\text{ana}}^{\text{ls}} \end{array}$$

Proof. To prove that $(\mathbf{LieGpoid}, J_{\mathbf{Lie}})$ is a 2-site, we check each condition in [Definition 7.1](#). All identity functors of Lie groupoids are surjective submersive weak equivalences contained in $J_{\mathbf{Lie}}$. That $J_{\mathbf{Lie}}$ is closed under composition and base change is inherited from J and [Proposition 9.23](#). That elements of $J_{\mathbf{Lie}}$ are representably fully faithful and co-fully faithful is also inherited from J .

That the corresponding anafunctor bicategory $\mathbf{LieGpoid}_{\text{ana}}$ is a sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$ is immediate. To show that the inclusion pseudofunctor is essentially full on anafunctors, one follows the corresponding proof for the inclusion of $\mathbf{LieGpoid}[W_{\mathbf{Lie}}^{-1}]$ into $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$ of [Theorem 9.25](#), but starting with an anafunctor instead of an arbitrary generalised morphism, which is already working in the anafunctor bicategory. The inclusion pseudofunctor is full on 2-cells due to [Proposition 9.23](#), completing the proof that $\mathbf{LieGpoid}_{\text{ana}}$ is an essentially full sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, from which it follows that $\mathbf{LieGpoid}_{\text{ana}}$ is a localisation of $\mathbf{LieGpoid}$ at $W_{\mathbf{Lie}}$, following the same logic as in [Theorem 9.13](#).

By [Proposition 9.22](#), the anafunctorisation of a functor between Lie groupoids is an anafunctor in $\mathbf{LieGpoid}_{\text{ana}}$, which is a sub-bicategory of $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, and so the diagram is well-defined; that it commutes is straightforward to check. \square

We now consider Lie groupoids with bibundles between them, again borrowing terminology from [\[vdS20; vdS21\]](#), and show that such bibundles are already in the Lie category.

Definition 9.27 (Lie Right Principal Bibundles). A right principal bibundle between Lie groupoids $\mathcal{G} \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} \mathcal{H}$ is **Lie** if Z is a manifold and l_Z a surjective submersion. If the bibundle is biprincipal and r_Z is a surjective submersion, then the bibundle is a **Lie Morita equivalence**. Let $\mathbf{LieBiBund}$ be the sub-bicategory of $\mathbf{DBiBund}$ of Lie groupoids with Lie right principal bibundles as 1-cells. \diamond

Proposition 9.28. *Let $\mathcal{G} \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} \mathcal{H}$ be a right principal bibundle between Lie groupoids. Then l_Z is a surjective submersion and the action groupoid $\mathcal{G} \times Z \times \mathcal{H}$ a Lie groupoid.*

Proof. By [Item 1](#) of [Proposition 9.11](#), we only need to show that the action groupoid is Lie; in particular, that its arrow and objects spaces are manifolds. Suppose Z is a manifold. Then $(\mathcal{G}_1)_s \times_{l_Z} Z$ is a manifold since the source map of \mathcal{G} is a surjective submersion, and similarly $((\mathcal{G}_1)_s \times_{l_Z} Z)_{r_Z \circ \text{opr}_2} \times_t \mathcal{H}_1$ is a manifold since the target map of \mathcal{H} is a surjective submersion. This space is diffeomorphic to $(\mathcal{G} \times Z \times \mathcal{H})_1$ by definition, and so it now suffices to prove that Z is a manifold.

By [Proposition 9.24](#), there is a Lie groupoid \mathcal{K} , an anafunctor $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$, and a transformation S from $\mathcal{G} \xleftarrow[\widehat{\iota}_Z]{\simeq} \mathcal{G} \times Z \times \mathcal{H} \xrightarrow[\widehat{r}_Z]{\simeq} \mathcal{H}$ to $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \xrightarrow[\psi]{\simeq} \mathcal{H}$. Let $\widetilde{\mathcal{L}}$ be as in the proof of [Proposition 8.20](#). It follows from [Proposition 9.22](#) and [Proposition 9.23](#) that $\widetilde{\mathcal{L}}$ is Lie. It follows from the smooth fully faithfulness of φ that the right \mathcal{K} -action on $\widetilde{\mathcal{L}}_0$ is free.

Fix a sequence $((x_i, g_i, y_i), (y_i, h_i, z_i), k_i) \in \widetilde{\mathcal{L}}_{0\text{pr}_3\text{opr}_1} \times_t \mathcal{K}_1$ such that

$$(((x_i, g_i, y_i), (y_i, k_i, z_i)), ((x_i, \varphi(k_i)^{-1}g_i, s(k_i)), (s(k_i), h_i\psi(k_i), z_i)))$$

converges to $((x, g, y), (y, g, z)), ((x, g', y'), (y', h', z))$ in $\widetilde{\mathcal{L}}_0^2$. Then $g' = g''g$ for some g'' to which $\varphi(k_i^{-1})$ converges. Define $k = \Phi_{\varphi}^{-1}(y', y, (g'')^{-1})$. Then $k_i \rightarrow k$ by the continuity of φ , and since the arrow spaces of Lie groupoids are Hausdorff. Thus the sequence $((x_i, g_i, y_i), (y_i, h_i, z_i), k_i)$ converges to $((x, g, y), (y, h, z), k)$, from which it follows that the map $(s, t): \widetilde{\mathcal{L}}_{0\text{pr}_3\text{opr}_1} \times_t \mathcal{K}_1 \rightarrow \widetilde{\mathcal{L}}_0^2$ is proper. Thus the \mathcal{K} -action on $\widetilde{\mathcal{L}}_0$ is free and proper. Thus $\mathcal{L}_0 := \widetilde{\mathcal{L}}_0/\mathcal{K}$ is a manifold (see [\[dHo13, Proposition 3.6.2\]](#)).

By [Proposition 8.20](#), there is a 2-cell from $\mathcal{G} \xleftarrow[\varphi]{\simeq} \mathcal{K} \rightarrow \mathcal{H}$ to $\mathcal{G} \xleftarrow[\omega]{\simeq} \mathcal{L} \rightarrow \mathcal{H}$ where \mathcal{L} is the $(\mathcal{G}\text{-}\mathcal{H})$ -action groupoid of the bibundle $\mathcal{G} \curvearrowright^{x_0} \mathcal{L}_0 \curvearrowright^{y_0} \mathcal{H}$. Vertically composing this with S (viewed as a 2-cell between generalised morphisms) yields a 2-cell from $\mathcal{G} \xleftarrow[\widehat{\iota}_Z]{\simeq} \mathcal{G} \times Z \times \mathcal{H} \xrightarrow[\widehat{r}_Z]{\simeq} \mathcal{H}$ to $\mathcal{G} \xleftarrow[\chi]{\simeq} \mathcal{L} \xrightarrow[\omega]{\simeq} \mathcal{H}$. By [Theorem 8.25](#), this 2-cell induces a bi-equivariant diffeomorphism $Z \rightarrow \mathcal{L}_0$. Since \mathcal{L}_0 is a manifold, we conclude that Z is as well. \square

Let $\mathfrak{B}^{\text{Lie}}$ be the restriction of the bibundlisation pseudofunctor \mathfrak{B} to **LieGpoid**.

Theorem 9.29. *LieBiBund is a full sub-bicategory of DBiBund^{ls}, and hence of DBiBund, and is a localisation of LieGpoid at W_{Lie} . Also, the following diagram commutes.*

$$\begin{array}{ccc} \mathbf{LieGpoid} & \xrightarrow{\mathfrak{B}^{\text{Lie}}} & \mathbf{LieBiBund} \\ \downarrow & & \downarrow \\ \mathbf{DGpoid}^{\text{ls}} & \xrightarrow{\mathfrak{B}^{\text{ls}}} & \mathbf{DBiBund} \end{array}$$

Proof. To show that the inclusion pseudofunctor of **LieBiBund** into **DBiBund^{ls}** is full, we must show that every purely right principal bibundle between Lie groupoids is Lie. But this is immediate from [Proposition 9.28](#). Therefore the application of Pronk's Comparison Theorem in the proof of [Theorem 8.19](#) still applied when our focus is restricted to **LieBiBund**, proving that **LieBiBund** is a localisation of **LieGpoid** at W_{Lie} . By [Proposition 9.28](#), the bibundlisation of a functor between Lie groupoids yields a Lie right principal bibundle, and so the diagram commutes. \square

We conclude this subsection with the following summary of the above.

Theorem 9.30. $\mathbf{LieGpoid}[W_{\mathbf{Lie}}^{-1}]$, $\mathbf{LieGpoid}_{\text{ana}}$, and $\mathbf{LieBiBund}$ are all equivalent categories, are all localisations of $\mathbf{LieGpoid}$ at $W_{\mathbf{Lie}}$, and are essentially full sub-bicategories of $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, and $\mathbf{DBiBund}^{\text{ls}}$, resp., and hence of $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, and $\mathbf{DBiBund}$, resp.

Remark 9.31. As a consequence of [Theorem 9.25](#), [Theorem 9.26](#), and [Theorem 9.29](#), the pseudonatural transformations making the triangles in [Remark 9.17](#) 2-commute restrict to the Lie sub-bicategories, yielding the following 2-commutative triangles

$$\begin{array}{ccccc}
 \mathbf{LieGpoid}[W_{\mathbf{Lie}}^{-1}] & \xrightarrow{\simeq} & \mathbf{LieGpoid}_{\text{ana}} & \xrightarrow{\simeq} & \mathbf{LieBiBund} \\
 \uparrow \mathfrak{S}^{\mathbf{Lie}} & \searrow \Downarrow & & \searrow \Downarrow & \\
 \mathbf{LieGpoid} & \xrightarrow{\mathfrak{Q}^{\mathbf{Lie}}} & & \xrightarrow{\mathfrak{B}^{\mathbf{Lie}}} & \\
 & \nearrow \mathfrak{A}^{\mathbf{Lie}} & & \nearrow &
 \end{array}$$

⌋

Before concluding this section, we provide an affirmative answer to the open question [[vdS21](#), Question 7.6], which is a corollary of [Proposition 9.28](#).

Corollary 9.32. *A diffeological Morita equivalence between Lie groupoids is a Lie Morita equivalence.*

Proof. Let $\mathcal{G} \overset{l_Z}{\curvearrowright} Z \overset{r_Z}{\curvearrowleft} \mathcal{H}$ be a right principal bibundle between Lie groupoids. By [Proposition 9.28](#), we only need to show that r_Z is a surjective submersion. But since the bibundle is biperincipal and Lie groupoids are locally subductive diffeological groupoids, the bibundle is a pure Morita equivalence by [[vdS21](#), Proposition 7.5] (see also [Remark 9.18](#)). In particular, r_Z is a local subduction; that is, r_Z is a surjective submersion. \square

Remark 9.33. [Remark 9.18](#) now holds in the Lie setting as well. In particular, by [Proposition 8.23](#) and [Corollary 9.32](#), a right principal bibundle between Lie groupoids admits a quasi-inverse if and only if it is a Lie Morita equivalence. Also, a generalised morphism in $\mathbf{LieGpoid}[W_{\mathbf{Lie}}^{-1}]$ admits a quasi-inverse if and only if both of its functors are weak equivalences, and by [Proposition 9.21](#), both of these are automatically Lie. Similarly, if the generalised morphism is an anafunctor, then there is a transformation from it to an anafunctor in which both functors are surjective submersive weak equivalences. \square

10. APPLICATIONS

In this section, we present a few applications of the above theory.

10.1. Of Sites and Stacks. Grothendieck (pre)topologies on the category of diffeological spaces is becoming an important tool in the study of loop groups and loop stacks [[RV18a](#); [RV18b](#); [Wal12b](#); [Wal12a](#)]. Besides having used these (somewhat implicitly) to construct $\mathbf{DGpoid}_{\text{ana}}$ and $\mathbf{DGpoid}_{\text{ana}}^{\text{ls}}$, we use these here in our first application to show that all three bicategories $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, and $\mathbf{DBiBund}$ are equivalent to geometric stacks over the site of diffeological spaces. This is already known to experts (e.g. [[RV18a](#), Theorem 2.10]) but a proof does not exist in the literature

as far as the author is aware. We give a proof using the set up of [Vil18], and refer the reader to this thesis for all definitions of terminology. To begin, we point out that since subductions, as well as local subductions, include identity maps, are closed under composition, and are closed under pullback (see Item 1 and Item 5 of Proposition 2.7), these two classes of maps between diffeological spaces induce Grothendieck pretopologies on **Diffeol**.

We want to show that these are both so-called “good” sites (see [Vil18, Definition 1.3.2]). We start with the site of diffeological spaces with subductions as covering families.

Proposition 10.1. *The site of diffeological spaces with covering families equal to subductions is a good site.*

Proof. We go through each condition required of a good site.

- (1) **Diffeol** contains an initial object: the empty set.
- (2) Every covering sieve is generated by some set of cartesian morphisms. In fact, every smooth map is cartesian (*i.e.* admits a pullback in the category **Diffeol**), and our covering sieves are generated by a subduction by definition.
- (3) We must show that for every pair of diffeological spaces X and Y together with a covering sieve $S = \{s_i: U_i \rightarrow X\}_{i \in I}$ of X and a family of maps $\{f_i: U_i \rightarrow Y\}_{i \in I}$ making the following diagrams commute

$$\begin{array}{ccc} U_{i_{s_i} \times_{s_j} U_j} & \longrightarrow & U_i \\ \downarrow & & \downarrow f_i \\ U_j & \xrightarrow{f_j} & Y, \end{array}$$

there exists a unique smooth $f: X \rightarrow Y$ such that $f \circ s_i = f_i$ for all i . Construct such an f as follows. Suppose S is generated by a subduction $F: Z \rightarrow X$. Fix $x \in X$. There exists $t_x: \mathbb{R}^0 = \{\text{pt}\} \rightarrow F^{-1}(x)$ since F is surjective. Define $s_x := F \circ t_x$, which does not depend on the choice of t_x . Since $s_x \in S$, there exists $i_x \in I$ so that $s_x = s_{i_x}$. Define $f(x) := f_{i_x}(\text{pt})$. Then f is a well-defined function.

Fix $i \in I$ and $u \in U_i$. Then $f_i(u) = f_j(\text{pt})$ where $j = i_{s_i(u)}$. But by definition, $f \circ s_i(u) = f_j(\text{pt})$, and so $f \circ s_i(u) = f_i(u)$. Since u is arbitrary, we conclude that $f \circ s_i = f_i$, and this is true for all i . Fix a plot p of X . Since F is a subduction, after shrinking U_p , there is a lift q of p to Z with respect to F . Since $p = F \circ q$, then $p = s_i$ for some i . Thus $f \circ p = f \circ s_i = f_i$, and so $f \circ p$ is smooth. It follows from the locality axiom of diffeology that $f \circ p$ is a plot of Y for general plots of p , without shrinking the domain. Thus, f is smooth.

(4) We need to show that subductions form a stack over diffeological spaces (we know already that they form a category fibred in groupoids [Vil18, Section 1.3]). In particular, we need to show two conditions (see [Vil18, Definition 1.2.8])²:

(a) Let X be a diffeological space, $S = \{s_i: U_i \rightarrow X\}_{i \in I}$ a covering sieve generated by a subduction $F: Y \rightarrow X$, $f: P \rightarrow X$ and $g: Q \rightarrow X$ subductions, and $\{\phi_i: f|_{U_i} \rightarrow g|_{U_i}\}_{i \in I}$ a family of maps covering id_X satisfying $\phi_i|_{U_{ij}} = \phi_j|_{U_{ij}}$ for every $i, j \in I$. Here, we are using the shorthand for pullbacks $f|_{U_i} := \text{pr}_2: P_f \times_{s_i} U_i \rightarrow U_i$, $U_{ij} := U_{i s_i} \times_{s_j} U_j$, and $\phi_i|_{U_{ij}}: f|_{U_{ij}} \rightarrow g|_{U_{ij}}$ equal to the pullback square

$$\begin{array}{ccc} P|_{U_{ij}} & \longrightarrow & P|_{U_i} \\ \downarrow & & \downarrow \\ U_{ij} & \longrightarrow & U_i, \end{array}$$

where the top map is $(p, u_i, u_j) \mapsto (\text{pr}_1(\phi_i(p, u_i)), u_i, u_j)$; note that by **Item 2** of **Remark 5.7**, we may identify

$$(P|_{U_i})|_{U_j} = (P|_{U_j})|_{U_i} = P|_{U_{ij}}.$$

The goal is to construct $\phi: P \rightarrow Q$ such that $\phi|_{P|_{U_i}} = \phi_i$ for each i .

We construct ϕ the same way we constructed f above: define $\phi(p) := \phi_{i_x}(p)$ where $s_{i_x} = F \circ t_x$ for some $t_x: \mathbb{R}^0 \rightarrow F^{-1}(x)$. Then ϕ is a well-defined function. An argument similar to the above shows that $\text{pr}_1 \circ \phi|_{U_i} = \phi_i$ for all i .

Let r be a plot of P . Then $f \circ r$ is a plot of X , and since F is a subduction, after shrinking U_r , there is a lift \tilde{r} of r to Y . Then $F \circ \tilde{r} \in S$, and so is equal to s_i for some i . Define $\hat{r}: U_r = U_i \rightarrow P|_{U_i}$ sending u to $(u, r(u))$. Then \hat{r} is smooth, and

$$\phi \circ r = \phi \circ \text{pr}_2 \circ \hat{r} = \text{pr}_2 \circ \phi|_{U_i} \circ \hat{r}.$$

Since $\phi|_{U_i} = \phi_i$, it follows that $\phi \circ r$ is smooth, hence a plot of Q . We conclude that ϕ is smooth. (This shows that subductions form a prestack over diffeological spaces.)

(b) Let X be a diffeological space, $\{s_i: U_i \rightarrow X\}_{i \in I}$ a covering sieve generated by a subduction $F: Y \rightarrow X$, $\{f_i: P_i \rightarrow U_i\}_{i \in I}$ a family of subductions, and $\{\phi_{ij}: P_i|_{U_{ij}} \rightarrow P_j|_{U_{ij}}\}_{i, j \in I}$ a family of diffeomorphisms satisfying the cocycle condition $\phi_{ij}|_{U_{ijk}} \circ \phi_{jk}|_{U_{ijk}} = \phi_{ik}|_{U_{ijk}}$ for all triples $(i, j, k) \in I^3$. The goal is to construct a subduction $g: Q \rightarrow X$ and a family of maps $\{\phi_i: Q|_{U_i} \rightarrow P_i\}_{i \in I}$ such that $\phi_{ij} \circ \phi_j|_{U_{ij}} = \phi_i|_{U_{ij}}$ for all i, j .

To this end, let $Q := (\coprod_i P_i) / \sim$, where the equivalence relation \sim is defined as follows: $p_i \in P_i$ is equivalent to $p_j \in P_j$ if $\phi_{ij}(p_i, f_i(p_i), f_j(p_j)) =$

²In a private conversation, Villatoro suggested that (S2) of [Vil18, Definition 1.2.8] should in fact have ‘‘covering family’’ replaced with ‘‘covering sieve’’; both are true in our setting, but we use ‘‘covering sieve’’.

$(p_j, f_i(p_i), f_j(p_j))$. The cocycle condition on $\{\phi_{ij}\}$ guarantees that this is, in fact, an equivalence relation. Let $g: Q \rightarrow X$ be the map $g[p_i] = s_i(f_i(p_i))$ for $p_i \in P_i$, for all i . This is well-defined and smooth.

Fix a plot r of X . Since $\{s_i\}$ is a sieve, there exists some $i_Y \in I$ for which $Y = U_{i_Y}$ and $F = s_{i_Y}$. Shrinking U_r , there is a lift \tilde{r} of r to P_{i_Y} with respect to $F \circ f_{i_Y}$. Then $g([\tilde{r}]) = s_{i_Y} \circ f_{i_Y} \circ \tilde{r} = r$, and we conclude that g is subductive.

For each $i \in I$, define $\psi_i: P_i \rightarrow Q_{g \times_{s_i} U_i}$. It follows from the cocycle condition that ϕ_{ii} is the identity, from which it follows that ψ_i is injective. Let $v \mapsto (q_v, u_v)$ be a plot of $Q_{g \times_{s_i} U_i}$; without loss of generality, assume the domain of the plot V is connected. By definition of the quotient diffeology, after shrinking V , there exists $j \in I$ and a lift p_v of q_v to P_j . Define $p'_v := \text{pr}_1 \circ \phi_{ji}(p_v, f_j(p_v), u_v)$. This is well-defined since $s_i(u_v) = g(q_v) = s_j(f_j(p_j))$, smooth, and has image in P_i . Moreover, $\psi_i(p'_v) = (q_v, u_v)$, and so ψ_i is subductive, and hence a diffeomorphism. Defining $\phi_i := \psi_i^{-1}$, it follows that $\{\phi_i\}$ are the desired morphisms. Indeed, for any $i, j, k \in I$,

$$\begin{aligned} \phi_j \circ \phi_i|_{U_{ij}}([p], u_i, u_j) &= \phi_{ij}(\text{pr}_1 \circ \phi_{ki}(p, f_k(p), u_i), u_i, u_j) \\ &= (\text{pr}_1 \circ \phi_{kj}(p, f_k(p), u_j), u_i, u_j) \\ &= \phi_j([p], u_i, u_j) \end{aligned}$$

where the second last line follows from the cocycle condition, and $p \in P_k$ is a representative of $[p]$.

- (5) The so-called ‘‘submersions’’ of **Diffeol** are exactly the subductions, and so it is required that if $g \circ f$ is a subduction, then so is g . But this is **Item 4** of **Proposition 2.7**.

This completes the proof. □

Let **B** be the pseudofunctor sending a diffeological groupoid \mathcal{G} to its classifying stack **B** \mathcal{G} (whose objects over the diffeological space X are right principal \mathcal{G} -bundles $P \rightarrow X$ and arrows are \mathcal{G} -equivariant bundle maps [Vil18, Subsection 1.3.4]), and defined on bibundles and bi-equivariant diffeomorphisms as defined in [Vil18, Subsection 1.3.5]. Recall that a stack \mathcal{S} over diffeological spaces is **geometric** if it admits a **representable submersion** $\overline{X} \rightarrow \mathcal{S}$, where X is a diffeological space [Vil18, Subsection 1.2.3]. Then we apply [Vil18, Theorem 1.3.27] and obtain:

Corollary 10.2. ***B** is an equivalence of bicategories between **DBiBund** and **DiffeolStacks**, the 2-category of geometric stacks over the site of diffeological spaces with subductions as covering families.*

This result generalises the well-known equivalence between **LieBiBund** and geometric stacks over the site of smooth manifolds (whose covering families are open covers, or equivalently surjective submersions).

Next, we want a similar result for the site of diffeological spaces in which the covering families are local subductions. However, [Vil18, (GS5) of Definition 1.3.2] requires that if $g \circ f$ is a local subduction, then so is g . (Note that this is true, however, if f and g are local subductions, *i.e.* surjective submersions, between manifolds.) This is true for subductions (see [Item 4](#) of [Proposition 2.7](#)), but false for local subductions: consider $s \circ u$ of the kernel groupoid in [Example 9.5](#). This said, Villatoro uses this to show that a bibundle admits a quasi-inverse if and only if it is biprincipal. We now replace this condition with the following lemma, and use it to reproduce Villatoro's result for the locally subductive case (which is the same as [[vdS21](#), Proposition 7.5]).

Lemma 10.3. *Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be smooth so that f is a surjective and $g \circ f$ is a local subduction. Then g is a local subduction.*

Proof. Let $p = z_u$ be a plot of Z and $y_0 \in g^{-1}(z_0)$. Since f is surjective, there exists $x_0 \in f^{-1}(y_0)$. Since $g \circ f$ is a local subduction, after shrinking U_p , there is a lift x_u of p through x_0 . Then $y_u := f(x_u)$ is the desired lift. \square

To illustrate how this result is used, we reproduce [[vdS21](#), Proposition 7.5] below.

Lemma 10.4. *A right principal bibundle between locally subductive diffeological groupoids admits a quasi-inverse if and only if it is a pure Morita equivalence.*

Proof. It follows from [[vdS21](#), Theorem 5.31] (see also [Vil18, Lemma 1.3.22]) that a bibundle in **DBiBund** admits a quasi-inverse if and only if it is biprincipal, and biprincipal bibundles between locally subductive diffeological groupoids are pure Morita equivalences by [[vdS21](#), Proposition 7.5]. Conversely, let \mathcal{G} and \mathcal{H} be locally subductive diffeological groupoids and $\mathcal{G} \overset{l_Z}{\curvearrowright} Z \overset{r_Z}{\curvearrowright} \mathcal{H}$ a bibundle admitting a quasi-inverse $\mathcal{H} \overset{l_W}{\curvearrowright} W \overset{r_W}{\curvearrowright} \mathcal{G}$. To complete the proof, we need to show that r_Z is locally subductive. The following commutative diagram will be useful:

$$\begin{array}{ccccc}
 & & W_{r_W} \times_{l_Z} Z & & \\
 & \swarrow \text{pr}_1 & \downarrow \pi_{\mathcal{G}} & \searrow \text{pr}_2 & \\
 & W & W \otimes_{\mathcal{G}} Z & Z & \\
 \mathcal{H}_0 & \swarrow l_W & \downarrow L_W & \searrow R_Z & \mathcal{H}_0 \\
 & & \cong \beta & & \\
 & \swarrow t & \downarrow \beta & \searrow s & \\
 & & \mathcal{H}_1 & &
 \end{array}$$

Here, $\pi_{\mathcal{G}}$ is the quotient map by the diagonal \mathcal{G} -action on $W_{r_W} \times_{l_Z} Z$, L_W and R_Z are the left and right anchor maps, resp., of the composed bibundles (see [Definition 8.6](#)), and β is the bi-equivariant diffeomorphism $W \otimes_{\mathcal{G}} Z \rightarrow \mathcal{H}_1$, which exists by definition of a quasi-inverse. Since \mathcal{H} is locally subductive, $s: \mathcal{H}_1 \rightarrow \mathcal{H}_0$ is locally subductive, and since β is a diffeomorphism, R_Z is locally subductive, too. As mentioned earlier, $\mathcal{G} \overset{l_Z}{\curvearrowright} Z \overset{r_Z}{\curvearrowright} \mathcal{H}$ and its quasi-inverse are biprincipal, and thus r_W is a subduction. By [Item 5](#) of [Proposition 2.7](#), $\text{pr}_2: W_{r_W} \times_{l_Z} Z \rightarrow Z$ is a subduction and hence surjective.

Thus, by [Lemma 10.3](#), if $r_Z \circ \text{pr}_2$ is locally subductive, so is r_Z . Since $r_Z \circ \text{pr}_2 = R_Z \circ \pi_{\mathcal{G}}$, and R_Z is locally subductive, the proof reduces to showing that $\pi_{\mathcal{G}}$ is locally subductive.

Fix a plot $p = w_u \otimes z_u \in W \otimes_{\mathcal{G}} Z$; we choose the domain small enough so that there is a lift (w_u, z_u) of it to $W_{r_W} \times_{l_Z} Z$. Let $(w'_0, z'_0) \in W_{r_W} \times_{l_Z} Z$ so that $w'_0 \otimes z'_0 = w_0 \otimes z_0$. Then there is some $g_0 \in \mathcal{G}_1$ so that $(w'_0, z'_0) = (w_0 g_0^{-1}, g_0 z_0)$. Since \mathcal{G} is locally subductive, after shrinking U_p , there is a lift g_u of $l_Z(z_u) = r_W(w_u)$ through g_0 with respect to the source map of \mathcal{G} . Then $(w'_u, z'_u) = (w_u g_u^{-1}, g_u z_u)$ is a lift of $w_u \otimes z_u$ through (w'_0, z'_0) . Thus $\pi_{\mathcal{G}}$ is a local subduction, completing the proof. \square

Before continuing with our goal, another subtlety is that Villatoro’s “submersions”, when covering families are equal to local subductions, form a sub-class of subductions that include local subductions, along with other subductions as well. We wish to focus solely on the local subductions. As per the remark on [[Vil18](#), pages 13–14], provided that the local subductions include isomorphisms (they do), form a stack (they do, as per the proof below), and satisfy [Lemma 10.3](#), it is fine to restrict to them: the correspondence between geometric stacks over the site of diffeological spaces with local subductions as covering families and locally subductive diffeological groupoids goes through.

Proposition 10.5. *The site of diffeological spaces with covering families and submersions equal to local subductions satisfy the first four conditions of a good site, with the fifth condition replaced by [Lemma 10.3](#).*

Proof. This proof follows the proof of [Proposition 10.1](#), since local subductions are subductions. The only addition that is needed is to (4)(b), where we require $g: Q \rightarrow X$ to be a *local* subduction. Using the same notation, but with all f_i and F local subductions, fix the plot $r = x_v$ of X and $[p_0] \in Q$ such that $g[p_0] = x_0$. Let $i \in I$ so that $p_0 \in P_i$. Then since s_i factors as $F \circ t_i$ for some smooth $t_i: U_i \rightarrow Y$, setting $y_0 := t_i(f_i(p_0))$. Then $F(y_0) = x_0$, and since F is a local subduction, after shrinking U_r , there is a lift y_v of x_v through y_0 . Since F is in the covering sieve, there is some $i_Y \in I$ such that $U_{i_Y} = Y$ and $s_{i_Y} = F$. Since $f_{i_Y}: P_{i_Y} \rightarrow Y$ is a local subduction, after shrinking U_r , there is a lift p'_v of y_v to P_{i_Y} through $\phi_{i_Y}(p_0, f_i(p_0), y_0)$, noting that $s_i(p_0) = F(y_0)$. Then $[p'_0] = [p_0]$, and $g(p'_v) = r$. \square

Corollary 10.6. *\mathbf{B} restricts to an equivalence of bicategories between $\mathbf{DBiBund}^{\text{ls}}$ and $\mathbf{DiffeolStacks}^{\text{ls}}$, the 2-category of geometric stacks over the site of diffeological spaces with local subductions as covering families and submersions.*

Of course, one needs to take care with what is meant by “restricting B ” in [Corollary 10.6](#), as the covering families are different. We conclude this subsection with the following theorem.

Theorem 10.7. *The bicategories $\mathbf{DGpoid}[W^{-1}]$, $\mathbf{DGpoid}_{\text{ana}}$, $\mathbf{DBiBund}$, and $\mathbf{DiffeolStacks}$ are all equivalent, as are $\mathbf{DGpoid}^{\text{ls}}[W_{\text{ls}}^{-1}]$, $\mathbf{DGpoid}^{\text{ls}}_{\text{ana}}$, $\mathbf{DBiBund}^{\text{ls}}$, and $\mathbf{DiffeolStacks}^{\text{ls}}$.*

Of course, the same statement but for Lie groupoids is well-known; see, for instance, [[Blo08](#); [Ler10](#)].

10.2. Invariants of Diffeological Groupoids. In [vdS21, Section 6], three types of Morita invariants (*i.e.* properties of diffeological groupoids maintained by Morita equivalences) are discussed in the context of diffeological bibundles. Namely, their orbit spaces, whether the groupoid is “fibrating”, and their action categories of left \mathcal{G} -spaces and \mathcal{G} -equivariant maps. **Theorem 8.25** now allows us to extend these invariants beyond **DBiBund**, in the sense that we can take “Morita equivalent” as defined in any of the bicategories **DGpoid** $[W^{-1}]$, **DGpoid** $_{\text{ana}}$, or **DBiBund**.

We restate the extended results stated in [vdS21, Section 6] with appropriate definitions.

Definition 10.8 (Fibrating Groupoid). A diffeological groupoid \mathcal{G} is **fibrating** if its characteristic map $\chi_{\mathcal{G}} = (s, t): \mathcal{G}_1 \rightarrow \mathcal{G}_0^2$ is a subduction on arrows. \diamond

Definition 10.9 (Action Category). The **action category** of a diffeological groupoid \mathcal{G} is the category whose objects are diffeological spaces admitting a left \mathcal{G} -action, and whose morphisms are the \mathcal{G} -equivariant maps between these spaces. \diamond

Theorem 10.10 (Morita Invariants). *Suppose that \mathcal{G} and \mathcal{H} are diffeological groupoids admitting a Morita equivalence in **DGpoid** $[W^{-1}]$, **DGpoid** $_{\text{ana}}$, or **DBiBund**.*

- (1) *The orbit spaces $\mathcal{G}_0/\mathcal{G}_1$ and $\mathcal{H}_0/\mathcal{H}_1$ are diffeomorphic.*
- (2) *\mathcal{G} is fibrating if and only if \mathcal{H} is fibrating.*
- (3) *The action category of \mathcal{G} is equivalent to the action category of \mathcal{H} .*

Proof. These are proved in **DBiBund** in [vdS21, Theorems 6.1, 6.3, 6.4]. Thus the result follows from **Theorem 8.25**. \square

We now restrict **Item 3** of **Theorem 10.10** to the case of linear representations of diffeological groupoids. For this, we will need some definitions (see, for instance, [CW16, Definition 4.5]).

Definition 10.11 (Diffeological Vector Pseudo-Bundle). A **diffeological vector pseudo-bundle** $\rho: E \rightarrow X$ is a smooth map between diffeological spaces satisfying for each $x \in X$ that the fibre $\rho^{-1}(x)$ admits a vector space structure such that addition $E_{\rho} \times_{\rho} E \rightarrow E$, scalar multiplication $\mathbb{R} \times E \rightarrow E$, and the zero section $\zeta: X \rightarrow E: x \mapsto 0_x$ are smooth. Given a diffeological groupoid \mathcal{G} , a **linear representation** of \mathcal{G} is a left \mathcal{G} -action on a diffeological vector pseudo-bundle $\rho: E \rightarrow \mathcal{G}_0$, in which ρ is the anchor map, and the action is linear; thus for any $x \in \mathcal{G}_0$, $g \in s^{-1}(x)$, $e, e' \in \rho^{-1}(x)$, and $r \in \mathbb{R}$, we have

$$\begin{aligned} g(e + e') &= ge + ge', \\ g(re) &= r(ge), \quad \text{and} \\ g0_x &= 0_x. \end{aligned}$$

Denote by $\text{Rep}(\mathcal{G})$ the category of left linear \mathcal{G} -representations and \mathcal{G} -equivariant maps between them. \diamond

Proposition 10.12 (Diffeological Groupoids Representations). *Given Morita equivalent diffeological groupoids \mathcal{G} and \mathcal{H} , the categories $\text{Rep}(\mathcal{G})$ and $\text{Rep}(\mathcal{H})$ are equivalent.*

Proof. By [Proposition 8.23](#), there exists a biprincipal bibundle $\mathcal{G} \curvearrowright^{L_X} X \curvearrowright^{r_X} \mathcal{H}$, with opposite bibundle $\mathcal{H} \curvearrowright^{\overline{L_X}} \overline{X} \curvearrowright^{\overline{r_X}} \mathcal{G}$. By [Item 3 of Theorem 10.10](#), there is an equivalence of categories between the action categories of \mathcal{G} and \mathcal{H} . The equivalence is given by the functors

$$\begin{aligned} X \otimes_{\mathcal{H}} \cdot &: \mathcal{H} \curvearrowright^a Y \mapsto \mathcal{G} \curvearrowright^{L_X} X \otimes_{\mathcal{H}} Y \\ &(\varphi: Y_1 \rightarrow Y_2) \mapsto (\text{id}_X \otimes_{\mathcal{H}} \varphi: X \otimes_{\mathcal{H}} Y_1 \rightarrow X \otimes_{\mathcal{H}} Y_2), \quad \text{and} \\ \overline{X} \otimes_{\mathcal{G}} \cdot &: \mathcal{G} \curvearrowright^{a'} Y' \mapsto \mathcal{H} \curvearrowright^{\overline{L_X}} \overline{X} \otimes_{\mathcal{G}} Y' \\ &(\psi: Y'_1 \rightarrow Y'_2) \mapsto (\text{id}_{\overline{X}} \otimes_{\mathcal{G}} \psi: \overline{X} \otimes_{\mathcal{G}} Y'_1 \rightarrow \overline{X} \otimes_{\mathcal{G}} Y'_2). \end{aligned}$$

The associated natural transformations are given by compositions of associators, unitors, etc.

To complete the proof, it suffices to show that the first functor restricted to the full subcategory $\text{Rep}(\mathcal{G})$ of the \mathcal{G} -action category maps to $\text{Rep}(\mathcal{H})$. To this end, fix a left linear \mathcal{H} -representation $\rho: E \rightarrow \mathcal{H}_0$. We need to show that $L_X: X \otimes_{\mathcal{H}} E \rightarrow \mathcal{G}_0$ is a vector pseudo-bundle, and that the \mathcal{G} -action on $X \otimes_{\mathcal{H}} E$ is a linear representation.

Let $[x, e], [x', e'] \in X \otimes_{\mathcal{H}} E$ such that $L_X[x, e] = L_X[x', e']$. Then $l_X(x_1) = l_X(x_2)$, and by \mathcal{H} -principality, there exists a unique $h = d_{\mathcal{H}}(x, x') \in \mathcal{H}_1$ such that $x' = xh$, where $d_{\mathcal{H}}$ is the division maps of the underlying right principal \mathcal{H} -bundle. Thus

$$[x', e'] = [xh, e'] = [x, he'],$$

and by uniqueness, (x, he') is the unique representative of $[x', e']$ for which $L_X[x', e'] = x$. Define fibre-wise addition of $L_X: X \otimes_{\mathcal{H}} E \rightarrow \mathcal{G}_0$ by

$$[x, e] + [x', e'] := [x, e + he'].$$

This is well-defined and smooth. For $r \in \mathbb{R}$ define scalar multiplication of $L_X: X \otimes_{\mathcal{H}} E \rightarrow \mathcal{G}_0$ by

$$r[x, e] := [x, re].$$

This is also well-defined and smooth. Finally, if $\zeta_{\rho}: \mathcal{H}_0 \rightarrow E$ is the zero section of ρ , define the zero section $\zeta_{L_X}: \mathcal{G}_0 \rightarrow X \otimes_{\mathcal{H}} E$ by

$$\zeta_{L_X}(z) := [x, 0]$$

where $x \in l_X^{-1}(z)$. Given another point $x' \in l_X^{-1}(z)$, there exists a unique $h \in \mathcal{H}_1$ such that $x' = xh$. Thus

$$[x', 0] = [xh, 0] = [x, h0] = [x, 0].$$

This zero section is well-defined and smooth. Thus L_X is a vector pseudo-bundle. It now follows from the fact that the \mathcal{G} - and \mathcal{H} -actions on X commute that the \mathcal{G} -action on $X \otimes_{\mathcal{H}} E$ is a linear representation. \square

We next show that the inertia groupoid of a diffeological groupoid \mathcal{G} is a Morita invariant, made more precise below, using the notation established in [Example 3.9](#).

Proposition 10.13. *Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a weak equivalence.*

- (1) The kernels of the characteristic maps $\ker(\chi_{\mathcal{G}})$ and $\ker(\chi_{\mathcal{H}})$ are Morita equivalent.
- (2) The inertia groupoids $\mathcal{J}_{\mathcal{G}}$ and $\mathcal{J}_{\mathcal{H}}$ are Morita equivalent.

Proof. Since φ is a weak equivalence, its anafunctorisation $\mathcal{G} \xleftarrow{\simeq} \mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}} \mathcal{H} \xrightarrow{\simeq} \mathcal{H}$ has subductive weak equivalences for both left and right functors. It follows that $\text{pr}_1: \ker(\chi_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}}) \rightarrow \ker(\chi_{\mathcal{G}})$ and $\text{pr}_3: \ker(\chi_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}}) \rightarrow \ker(\chi_{\mathcal{H}})$ are both subductive on objects. It is straightforward to show that pr_1 is also smoothly fully faithful, and hence a subductive weak equivalence. A similar computation using the smooth fully faithfulness of φ yields the smooth fully faithfulness of pr_3 , and we conclude that it is a subductive weak equivalence. This proves [Item 1](#).

The objects of $\mathcal{J}_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}}$ are exactly the arrows of $\ker(\chi_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}})$; these triples (κ, h', λ) satisfy

$$\kappa \in \ker(\chi_{\mathcal{G}}), \quad \lambda \in \ker(\chi_{\mathcal{H}}), \quad \text{and} \quad \lambda h' \varphi(\kappa)^{-1} = h'. \quad (10)$$

The arrows of $\mathcal{J}_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}}$ are pairs of triples

$$((g, h', h), (\kappa, h', \lambda)) \in ((\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}} \mathcal{H})_1)_{\text{s} \times \text{s}} \left(\ker(\chi_{\mathcal{G}})_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}} \ker(\chi_{\mathcal{H}}) \right)_1.$$

Define functors $\psi: \mathcal{J}_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}} \rightarrow \mathcal{J}_{\mathcal{G}}$ and $\omega: \mathcal{J}_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}} \rightarrow \mathcal{J}_{\mathcal{H}}$ by

$$\begin{aligned} \psi((g, h', h), (\kappa, h', \lambda)) &= (g, \kappa) \quad \text{and} \\ \omega((g, h', h), (\kappa, h', \lambda)) &= (h, \lambda). \end{aligned}$$

ψ_0 and ω_0 are equal to the maps between arrow spaces of the functors pr_1 and pr_3 , resp., from $\ker(\chi_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}})$; thus by [Remark 4.9](#), ψ_0 and ω_0 are subductive.

It is straightforward to check that Φ_{ψ} is injective. Fix a plot $p = ((\kappa_u, h'_u, \lambda_u), (\kappa'_u, h''_u, \lambda'_u), (g_u, \kappa_u))$ of the codomain of Φ_{ψ} . Then $\kappa'_u = g_u \kappa_u g_u^{-1}$, and if we define $h_u := h''_u \varphi(g_u) (h'_u)^{-1}$, it follows from (10) (applied twice) that $\lambda'_u = h_u \lambda_u h_u^{-1}$. Thus $((g_u, h'_u, h_u), (\kappa_u, h'_u, \lambda_u))$ is a lift of p to the domain of Φ_{ψ} . It follows that Φ_{ψ} is a diffeomorphism, and so ψ is a subductive weak equivalence.

For arrow $((g, h', h), (\kappa, h', \lambda))$, since φ is smoothly fully faithful, $g = \Phi_{\varphi}^{-1}(\text{s}(\kappa), \text{t}(g\kappa g^{-1}), \varphi(g))$. It follows that Φ_{ω} is injective. Fix a plot $p = ((\kappa_u, h'_u, \lambda_u), (\kappa'_u, h''_u, \lambda'_u), (h_u, \lambda_u))$. Then $\lambda'_u = h_u \lambda_u h_u^{-1}$. Setting $g_u = \Phi_{\varphi}^{-1}(\text{s}(\kappa_u), \text{t}(\kappa'_u), (h''_u)^{-1} h_u h'_u)$, it follows from the smooth fully faithfulness of φ and (10) (applied twice) that $\kappa'_u = g_u \kappa_u g_u^{-1}$. Thus $((g_u, h'_u, h_u), (\kappa_u, h'_u, \lambda_u))$ is a lift of p to the domain of Φ_{ω} . It follows that Φ_{ω} is a diffeomorphism, and so ω is a subductive weak equivalence. We now have a Morita equivalence $\mathcal{J}_{\mathcal{G}} \xleftarrow[\psi]{\simeq} \mathcal{J}_{\mathcal{G}_{\varphi \times_{\text{id}_{\mathcal{H}}}}^{\text{w}}} \xrightarrow[\omega]{\simeq} \mathcal{J}_{\mathcal{H}}$. \square

Corollary 10.14. *Morita equivalent diffeological groupoids have Morita equivalent kernels of their characteristic functors and Morita equivalent inertia groupoids.*

Proof. If \mathcal{G} and \mathcal{H} are Morita equivalent, there exists a diffeological groupoid \mathcal{K} and weak equivalences $\varphi: \mathcal{K} \rightarrow \mathcal{G}$ and $\psi: \mathcal{K} \rightarrow \mathcal{H}$. Since Morita equivalences compose to form Morita equivalences, the result follows from [Proposition 10.13](#). \square

10.3. Principal Bundles. Let X be a diffeological space and \mathcal{G} a diffeological groupoid. Consider the category of all generalised morphisms from the trivial groupoid of X to \mathcal{G} . By [Theorem 8.25](#), this category is equivalent to the category of all anafunctors between the two diffeological groupoids, as well as the category of all right principal bibundles between them. But these bibundles take on a very specific form.

Lemma 10.15. *Let X be a diffeological space and \mathcal{G} a diffeological groupoid. The groupoid of bibundles from X to \mathcal{G} with bi-equivariant diffeomorphisms as arrows is isomorphic to the groupoid of right principal G -bundles of X with bundle isomorphisms as arrows.*

Proof. Let $X \curvearrowright^{l_Z} Z \curvearrowright^{r_Z} \mathcal{G}$ be a right principal bibundle. That $l_Z: Z \rightarrow X$ is a right principal \mathcal{G} -bundle follows immediately from the definition. Conversely, if $\rho: P \rightarrow X$ is a right principal \mathcal{G} -bundle with anchor map a , then there is a trivial action of the trivial groupoid of X on P , and we have a bibundle $X \curvearrowright^{\rho} P \curvearrowright^a \mathcal{G}$.

It also follows from the definition of bi-equivariant diffeomorphisms that these are exactly the bundle isomorphisms between the right principal \mathcal{G} -bundles. The result follows. \square

If we specialise to the case of a diffeological group $\mathcal{G} = (G \rightrightarrows \mathbb{R}^0)$, then these right principal \mathcal{G} -bundles are exactly the right principal G -bundles (see [\[KWW21, Definition 5.1\]](#)). If we specialise further to an abelian diffeological group G , then a result of [\[KWW21\]](#) is that principal G -bundles over X are classified by the first Čech cohomology group $\check{H}^1(X; G)$. In fact, if \mathcal{F} is a generating family of the diffeology of X , then [\[KWW21, Lemma 5.11, Remark 5.12, Corollary 5.16\]](#) imply that the category of right principal bibundles from $\mathcal{N}(\mathcal{F})$ to \mathcal{G} is equivalent to the category $\check{\mathcal{H}}^1(\mathcal{F}, G)$ whose objects are Čech 1-cocycles in $\check{C}^1(\mathcal{F}, G)$, and arrows from 1-cocycle f_1 to 1-cocycle f_2 are the 0-cochains α in $\check{C}^0(\mathcal{F}, G)$, such that $\partial\alpha = f_2 - f_1$. It now follows from [Example 4.10](#) that:

Theorem 10.16. *The category of generalised morphisms between X and the abelian group G (viewed as groupoids) is equivalent to category $\check{\mathcal{H}}^1(X, G)$ whose objects are 1-cocycles in $\check{C}^1(X, G)$ and arrows are 0-cochains as described above.*

It would be interesting to see generalisations of this: if we replace the trivial groupoid of X with a diffeological groupoid \mathcal{H} , do we get a classification of equivariant bundles? If we generalise G to a non-abelian group, does the Čech result still go through? What if we go further and let \mathcal{G} be a general diffeological groupoid? There is much to explore in this direction.

REFERENCES

- [AZ20] Iakovos Androulidakis and Marco Zambon. “Integration of singular subalgebroids”. 2020. ([preprint](#)).
- [BH11] John C. Baez and Alexander E. Hoffnung. “Convenient categories of smooth spaces”. In: *Trans. Amer. Math. Soc.* 363 (2011), pp. 5789–5825.
- [Bar06] Toby Bartels. “Higher gauge theory I: 2-bundles”. PhD Dissertation. PhD thesis. University of California Riverside, 2006.

- [B67] Jean Bénabou. *Introduction to Bicategories*. Lectures Notes in Mathematics, 47. Springer, 1967.
- [Blo08] Christian Blohmann. “Stacky Lie groups”. In: *Int. Math. Res. Not.* (2008), 51 pages.
- [BFW13] Christian Blohmann, Marco Cezar Barbosa Fernandes and Alan Weinstein. “Groupoid symmetry and constraints in general relativity”. In: *Commun. Contemp. Math.* 1.1250061 (2013), 25 pages.
- [CW16] J. Daniel Christensen and Enxin Wu. “Tangent spaces and tangent bundles for diffeological spaces”. In: *Cah. Topol. Géom. Différ. Catég.* 57 (2016), pp. 3–50.
- [dHo12] Matias L. del Hoyo. “Lie groupoids and differentiable stacks”. 2012. ([preprint](#)).
- [dHo13] Matias L. del Hoyo. “Lie groupoids and their orbispaces”. In: *Portugal. Math.* 70 (2013), pp. 161–209.
- [GZ67] Peter Gabriel and Michel Zisman. *Calculus of Fractions and Homotopy Theory*. Springer-Verlag, 1967.
- [GV21] Alfonso Garmendia and Joel Villatoro. “Integration of singular foliations via paths”. 2021. ([preprint](#)).
- [HS87] Michel Hilsun and Georges Skandalis. “Morphismes K -orientés d’espaces de feuilles et fonctorialité en théorie de Kasparov (d’après une conjecture d’A. Connes)”. In: *Annales scientifiques de l’École normale supérieure*. Vol. 20. 3. 1987, pp. 325–390.
- [IZ13] Patrick Iglesias-Zemmour. *Diffeology*. Vol. 185. Mathematical Surveys and Monographs. American Mathematical Soc., 2013.
- [IZP20] Patrick Iglesias-Zemmour and Elisa Prato. “Quasifolds, diffeology and noncommutative geometry”. In: *J. Noncommut. Geom.* (2020). (to appear).
- [JY21] Niles Johnson and Donald Yau. *2-Dimensional Categories*. Oxford University Press, 2021.
- [KWW21] Derek Krespki, Jordan Watts and Seth Wolbert. “Sheaves, principal bundles, and Čech cohomology for diffeological spaces”. 2021. ([preprint](#)).
- [Ler10] Eugene Lerman. “Orbifolds as stacks?”. In: *Enseign. Math. (2)* 56 (2010), pp. 315–363.
- [Li15] Du Li. “Higher Groupoid Actions, Bibundles, and Differentiation”. PhD thesis. Georg-August-Universität Göttingen, 2015.
- [Los94] Mark Losik. “Categorical differential geometry”. In: *Cah. Topol. Géom. Différ. Catég.* 35 (1994), pp. 274–290.
- [Mak96] Michael Makkai. “Avoiding the axiom of choice in general category theory”. In: *J. Pure Appl. Algebra* 108 (1996), pp. 109–173.
- [Moe02] Ieke Moerdijk. “Orbifolds as groupoids: an introduction”. In: *Orbifolds in mathematics and physics (Madison, WI, 2001)*, *Contemp. Math.* 310 (2002), pp. 205–222.
- [Pro96] Dorette Pronk. “Etendues and stacks as bicategories of fractions”. In: *Compositio Mathematica* 102.3 (1996), pp. 243–303.
- [PS10] Dorette Pronk and Laura Scull. “Translation groupoids and orbifold cohomology”. In: *Canad. J. Math.* 62.3 (2010), pp. 615–645.
- [PS21] Dorette Pronk and Laura Scull. “Bicategories of fractions revisited: towards small Homs and canonical 2-cells”. 2021. ([preprint](#)).
- [Rob12] David Michael Roberts. “Internal categories, anafunctors and localisations”. In: *Theory and Applications of Categories* 26 (2012), pp. 788–829.
- [Rob16] David Michael Roberts. “On certain 2-categories admitting localisation by bicategories of fractions”. In: *Appl. Categ. Structures* 24 (2016), pp. 373–384.
- [Rob21] David Michael Roberts. “The elementary construction of formal anafunctors”. In: *Categ. Gen. Algebr. Struct. Appl.* 15 (2021), pp. 183–229.
- [RV18a] David Michael Roberts and Raymond F. Vozzo. “Smooth loop stacks of differentiable stacks and gerbes”. In: *Cah. Topol. Géom. Différ. Catég.* 59 (2018), pp. 95–141.
- [RV18b] David Michael Roberts and Raymond F. Vozzo. “The smooth Hom-stack of an orbifold”. In: *2016 MATRIX Annals*. MATRIX Book Ser., 1 (2018), pp. 43–47.
- [Sou80] Jean-Marie Souriau. “Groupes différentiels”. In: *Differential geometrical methods in mathematical physics* (1980), pp. 91–128.
- [Sou84] Jean-Marie Souriau. “Groupes différentiels et physique mathématique”. In: *Group Theoretical Methods in Physics* (1984), pp. 511–513.

- [vdS20] Nesta van der Schaaf. “Diffeology, Groupoids & Morita Equivalence”. MA thesis. Radboud University, 2020.
- [vdS21] Nesta van der Schaaf. “Diffeological Morita Equivalence”. In: *Cahiers de Topologie et Géométrie Différentielle Catégoriques* LXII.2 (2021), pp. 177–238.
- [Vil18] Joel Villatoro. “Stacks in Poisson Geometry”. PhD thesis. University of Illinois, 2018.
- [Wal12a] Konrad Waldorf. “A construction of string 2-group models using a transgression-regression technique”. In: *Analysis, geometry and quantum field theory*. Contemp. Math., 584 (2012), pp. 99–115.
- [Wal12b] Konrad Waldorf. “Transgression to loop spaces and its inverse I: diffeological bundles and fusion maps”. In: *Cah. Topol. Géom. Différ. Catég.* 53 (2012), pp. 162–210.

DEPARTMENT OF MATHEMATICS, CENTRAL MICHIGAN UNIVERSITY, PEARCE HALL 214, MOUNT PLEASANT, MI 48859, USA

Email address: `jordan.watts@cmich.edu`