

ON THE DISTRIBUTION OF POLYNOMIALS HAVING A GIVEN NUMBER OF IRREDUCIBLE FACTORS OVER FINITE FIELDS

ARGHYA DATTA

ABSTRACT. Let $q \geq 2$ be a fixed prime power. We prove an asymptotic formula for counting the number of monic polynomials that are of degree n and have exactly k irreducible factors over the finite field \mathbb{F}_q . We also compare our results with the analogous existing ones in the integer case, where one studies all the natural numbers up to x with exactly k prime factors. In particular, we show that the number of monic polynomials grows at a surprisingly higher rate when k is a little larger than $\log n$ than what one would speculate from looking at the integer case.

1. INTRODUCTION

How many integers are there up to 1000 billion (or any large number) with exactly 2022 (or any given integer) prime factors? This exciting question fascinated mathematicians for ages. Gauss initially started the investigation and was later pursued by several eminent mathematicians, including Landau, Hardy and Ramanujan, Sathe and Selberg. Thanks to their compound effort, we entirely understand these answers in the integer setting. The main goal of this work is to study the analogous question for polynomials over finite fields. We begin our discussion with a brief historical interlude and then state our main results.

Let $m \leq x$ be positive integers and $\Omega(m)$ denotes the number of prime factors counted with multiplicity that divide m . In 1900, Landau [2] showed using the prime number theorem that for $k \in \mathbb{N}$ fixed we have

$$\Pi(x, k) \sim \frac{x}{\log x} \frac{(\log \log x)^{k-1}}{(k-1)!},$$

where

$$\Pi(x, k) := \#\left\{m \leq x : \Omega(m) = k\right\}.$$

The above result was predicted by Gauss [7] long back. However, understanding the asymptotic behaviour of $\Pi(x, k)$ for growing k uniformly with x remained an open problem for a long time. In 1953, Sathe [4] ingeniously showed that if $\epsilon \in (0, 2)$ is given, then uniformly for $1 \leq k \leq (2 - \epsilon) \log \log x$ we have

$$(1.1) \quad \Pi(x, k) \sim F\left(\frac{k}{\log \log x}\right) \frac{(\log \log x)^{k-1}}{(k-1)!} \frac{x}{\log x},$$

where

$$F(z) = \frac{1}{\Gamma(z+1)} \prod_p \left(1 - \frac{1}{p}\right)^z \left(1 - \frac{z}{p}\right)^{-1} \quad (|z| < 2).$$

This beautiful result was further extended by Selberg [5] when k is a little larger and varies

2020 *Mathematics Subject Classification.* 11T06, 11N37, 11T55, 11N56.

Key words and phrases. polynomials over finite field, fixed number of irreducible factors, Selberg-Delange method, multiplicative functions, saddle point approximation.

uniformly in a slightly wider range $(2 + \epsilon) \log \log x \leq k \leq B^* \log \log x$, where $B^* > 2$ is a real constant. In his paper Selberg only discussed the main ideas behind the proof. He observed a significant change in the asymptotic behaviour of $\Pi(x, k)$. He showed for this range we have

$$(1.2) \quad \Pi(x, k) \sim \frac{Cx \log(x/2^k)}{2^k},$$

where C is an absolute constant. Finally Nicolas [1] settled this question for the entire range $(2 + \epsilon) \log \log x \leq k < \log x / \log 2$ by proving the same asymptotic result holds as (1.2) in this range. Unlike the previous attempts his main idea behind the proof was less analytic and more of combinatorial nature. As mentioned earlier, we will be working in the polynomial ring $\mathbb{F}_q[t]$. Henceforth q is assumed to be fixed and all the implied constants are allowed to depend on q . For an easy reference later on, we employ some standard notation that will be used throughout the paper.

1.1. Notation. We set

$$\begin{aligned} \mathcal{M} &:= \left\{ \text{set of all monics in } \mathbb{F}_q[t] \right\}, \\ \widetilde{\mathcal{M}} &:= \left\{ \text{set of all monics in } \mathbb{F}_q[t] \text{ having no root in } \mathbb{F}_q \right\}, \\ \mathcal{P} &:= \left\{ \text{set of all monic irreducibles in } \mathbb{F}_q[t] \right\}. \end{aligned}$$

\mathcal{M}_n consists of only those monics that have degree = n and similarly we define $\widetilde{\mathcal{M}}_n, \mathcal{P}_n$. Let $\mathcal{P}_{\geq j}$ denotes the set of all monic irreducibles whose degree is at least j .

Given $f \in \mathbb{F}_q[t]$, let $\Omega(f)$ denotes the number of irreducible polynomials counted with multiplicity that divide f . We also set

$$\begin{aligned} N(n, k) &:= \# \left\{ f \in \mathcal{M}_n : \Omega(f) = k \right\}, \\ N'(n, k) &:= \# \left\{ f \in \widetilde{\mathcal{M}}_n : \Omega(f) = k \right\}. \end{aligned}$$

As discussed initially, we are interested in understanding the object $N(n, k)$ for different ranges of k . Surprisingly one of our main results reveals that the situation in $\mathbb{F}_q[t]$ is quite different than what one could naively guess from the knowledge in the integer case. This is remarkable since these two worlds are significantly different in only a few instances.

Our first result is fairly what one could speculate in light of (1.1).

Theorem 1.1. *Given $\epsilon \in (0, q)$, then uniformly for $1 \leq k \leq (q - \epsilon) \log n$ we have*

$$N(n, k) \sim \frac{q^n}{n} H \left(\frac{k}{\log n} \right) \frac{(\log n)^{k-1}}{(k-1)!},$$

where

$$H(z) = \frac{1}{\Gamma(z+1)} \prod_{p \in \mathcal{P}} \left(1 - \frac{1}{q^{\deg(p)}} \right)^z \left(1 - \frac{z}{q^{\deg(p)}} \right)^{-1}$$

for $|z| < q$.

We may notice that the expression for the function H is quite similar to F in the integer case. The appearance of q^n/n is also no surprise as letting $q^n \approx x$ yields $q^n/n \approx x/\log x$ up to constant factors. While the above result is more or less an application of the Selberg-Delange method developed in [6], the following surprising outcome of our work is where a substantial amount of new techniques goes in.

Theorem 1.2. *Let $B \geq 2$ be a real constant and $q > 2$ be a prime power. Let $\xi : \mathbb{N} \rightarrow \mathbb{R}$ be a function such that $\xi(n) \rightarrow \infty$ however slowly as $n \rightarrow \infty$. Then uniformly for $\xi(n) \log n \leq k \leq n/B$ we have*

$$N(n, k) \sim C(q) \frac{q^n k^{q-1} (n-k)^{q-1}}{q^k},$$

where

$$C(q) = \frac{1}{((q-1)!)^2} \left(1 - \frac{1}{q}\right)^{q^2} \prod_{p \in \mathcal{P}_{\geq 2}} \left(1 - \frac{1}{q^{\deg(p)-1}}\right)^{-1} \left(1 - \frac{1}{q^{\deg(p)}}\right)^q,$$

an absolute constant that depends only on q .

Here “however slowly” means that $\xi(n)$ could be any function that grows to infinity with an additional constraint that ξ is sufficiently small so that the interval $[\xi(n) \log n, n/B]$ contains at least one integer for n large enough. We remind the reader that $x/2^k$ is replaced by q^n/q^k as was expected. The appearance of $(n-k)$ is not also surprising in light of the main theorem proved in [1] and (1.2) where a factor of $\log(x/2^k)$ is present in the statement. However, the extra factor k^{q-1} and also the presence of higher powers of $(n-k)$ are what seem surprising at the moment, which shows $N(n, k)$ grows at a much larger rate than what we would predict from (1.2). The proofs of the above theorems are discussed in detail in the following sections. The main ingredients of our proofs are the following two key propositions.

Proposition 1.3. *For all $\epsilon \in (0, q)$ and uniformly for $|z| \leq q - \epsilon$ we have*

$$M_n(z) := \sum_{f \in \mathcal{M}_n} z^{\Omega(f)} = q^n n^{z-1} \left\{ zH(z) + O_\epsilon\left(\frac{1}{n}\right) \right\},$$

with $H(z)$ as described in Theorem 1.1.

Proposition 1.4. *For all $\delta \in (0, 1)$ and uniformly for $|z| \leq q^2 - \delta$ we have*


$$\widetilde{M}_n(z) := \sum_{f \in \widetilde{\mathcal{M}}_n} z^{\Omega(f)} = q^n n^{z-1} \left\{ zh(z) + O_\delta\left(\frac{1}{n}\right) \right\},$$

where $h(z) = \frac{1}{\Gamma(z+1)} \left(1 - \frac{1}{q}\right)^{qz} \prod_{p \in \mathcal{P}_{\geq 2}} \left(1 - \frac{z}{q^{\deg(p)}}\right)^{-1} \left(1 - \frac{1}{q^{\deg(p)}}\right)^z$, an analytic function for $|z| < q^2$.

The connection between Theorem 1.1 and Proposition 1.3 becomes clear once we notice $N(n, k)$ is precisely given by the coefficient of z^k in $M_n(z)$. In other words, if we could extract the information about $N(n, k)$ from $M_n(z)$, we would have our result. Furthermore, we note that $M_n(z)$ is polynomial of degree at most n and therefore, we could hope to study the coefficients

using Cauchy's integral formula. This is going to be our main strategy to attack Theorem 1.1. The connection between $\widetilde{M}_n(z)$ and Theorem 1.2 is not as immediate as the previous case since the coefficient of z^k in $\widetilde{M}_n(z)$ is $N'(n, k)$ and not $N(n, k)$. As we shall see, in section 4 the two objects $N(n, k)$ and $N'(n, k)$ are intimately entangled, and information about one could be transferred into another. We also note the relation $C(q) = qh(q)/(q-1)!$ holds.

1.2. Organization of the paper. The proofs of the above results rest on a technique developed by Sathe and Selberg along with a combination of an observation made by Nicolas [1] in his work and an analogue of the Selberg-Delange style argument for function fields developed in Porrit [6, section 2]. We will also use ideas from Tenenbaum [9, Chapter II.6], where the author outlines an application of the Selberg-Delange method to integers with a given number of prime factors. In the following section, we give a detailed proof of Theorem 1.1 and Proposition 1.3 as a direct application of the Selberg-Delange method discussed in [6]. The flow of the argument and preliminary lemmas leading to Theorem 1.2 are fairly delicate. Proof of Proposition 1.4 will be fundamentally the same as that of Proposition 1.3 apart from some technical modifications. We will begin section 4 by establishing Lemma 4.1, an odd-even decomposition for polynomials. We will then discuss the role of the auxiliary results Lemma 4.2, Lemma 4.5, Lemma 4.6, Lemma 4.4 behind the main proof of the Theorem 1.2. Next, we establish two important results, Proposition 4.3 and Proposition 4.7 which will finally lead us to Theorem 1.2. Detailed proofs of the technical lemmas and Proposition 1.4 will be presented in the appendix. The graphical dependency between the results in section 4 is described below:



Acknowledgements. The author would like to thank his supervisor, Prof. Andrew Granville, for suggesting that he work on this problem and also his co-supervisor, Prof. Dimitris Koukoulopoulos, for many helpful conversations and continuous support during this project and for their valuable suggestions and comments on the previous versions of this manuscript. The author was financially supported by his supervisors and the fellowships provided by Faculté des études supérieures et postdoctorales (FESP), Bourse d'exemption of Université de Montréal, Centre interuniversitaire en calcul mathématique algébrique (CICMA) while carrying out this work.

2. SET-UP AND THE PROOF OF PROPOSITION 1.3

2.1. Preparatory results for Proposition 1.3. In this subsection we aim to develop the framework for a Selberg-Delange type argument that will be used to establish Proposition 1.3. As

already discussed we are interested in the quantity

$$M_z(n) := \sum_{f \in \mathcal{M}_n} z^{\Omega(f)} = \sum_{k \geq 0} z^k N(n, k).$$

We also want to introduce an auxiliary quantity

$$(2.1) \quad G_z(u) = \sum_{f \in \mathcal{M}} z^{\Omega(f)} u^{\deg(f)} = \prod_{p \in \mathcal{P}} (1 - zu^{\deg(p)})^{-1}.$$

The function $G_z(u)$ defines an analytic function since the Euler product converges absolutely for $|u| < \min\{|z|^{-1}, q^{-1}\}$ which we argue now. We have that

$$|1 - zu^{\deg(p)}| \geq 1 - |zu^{\deg(p)}| \geq 1 - |zu| > 0,$$

showing there is no pole in the concerned region.

We now recall a well known result from complex analysis that says for a sequence of complex numbers $\{a_n\} \subset \mathbb{C}$ with $a_n \neq -1$ if $\sum_{n=1}^{\infty} |a_n| < \infty$, then

$$\prod_{n=1}^{\infty} (1 + a_n) \text{ converges.}$$

Using this result we see the product in (2.1) converges whenever $0 < |u| < q^{-1}$. Indeed we have

$$\begin{aligned} \sum_{p \in \mathcal{P}} |zu^{\deg(p)}| &\leq \sum_{p \in \mathcal{P}} |u|^{\deg(p)-1} \quad (|uz| < 1) \\ &\leq q + \sum_{p \in \mathcal{P}_{\geq 2}} |u|^{\deg(p)-1} \\ &\leq q + \sum_{n=2}^{\infty} q^n |u|^{n-1} \quad (\text{since } \#\mathcal{P}_n \leq q^n) \\ &\leq q + \frac{1}{|u|} \sum_{n=2}^{\infty} |qu|^n < \infty. \quad (|qu| < 1) \end{aligned}$$

Our job is to understand $M_n(z)$, which is precisely given by the coefficient of u^n in the power series of $G_z(u)$. Hence information about $M_n(z)$ can be obtained from $G_z(u)$ by applying Cauchy's integral formula. We took this leap from $M_n(z)$ to G_z because it seems to have a nicer arithmetic structure (i.e. an Euler product expansion), making it more amenable from the analytic point of view. Therefore it is natural to seek to meromorphically extend $G_z(u)$ beyond the region of absolute convergence discussed above to collect the contribution coming from the singularity of G_z in the extended domain. We do this job by introducing another function $F_z(u)$ via the Euler product

$$(2.2) \quad F_z(u) = \prod_{p \in \mathcal{P}} (1 - zu^{\deg(p)})^{-1} (1 - u^{\deg(p)})^z.$$

We intend to analyze the above Euler product in the following open region and show it is holomorphic.

$$\mathcal{R} := \left\{ (u, z) \in \mathbb{C}^2 : |z| < q, |u| < |z|^{-1}, |u| < q^{-1/2} \right\} \subset \mathbb{C}^2.$$

The following lemma shows $F_z(u)$ does converge absolutely in \mathcal{R} which is what we wanted to establish.

Lemma 2.1. *We have that $F_z(u)$ defined in (2.2) is holomorphic in \mathcal{R} .*

Proof. We take the standard branch of complex logarithm defined over $\mathbb{C} \setminus (-\infty, 0]$. Since $|u|^n, |zu^n| < 1$ we could make use of their Taylor series and write

$$\begin{aligned} z \log(1 - u^n) - \log(1 - zu^n) &= -zu^n + O(|zu^{2n}|) + zu^n + O(|z^2u^{2n}|) \\ &= O(|q^2u^{2n}|) \quad (|z| < q) \end{aligned} \tag{2.3}$$

and hence using $\exp(O(x)) = 1 + O(x)$ for $x = O(1)$ we get

$$(1 - u^n)^z (1 - zu^n)^{-1} = 1 + O(|q^2u^{2n}|).$$

We now observe the product in (2.2) converges absolutely in \mathcal{R} . Since $\#\mathcal{P}_n \leq q^n$, we have that

$$\sum_{p \in \mathcal{P}} |u|^{2n} \leq \sum_{n \geq 1} |qu^2|^n < \infty.$$

□

We recall the Riemann zeta function in the context of polynomials over finite field, defined for $|u| < q^{-1}$

$$\zeta(u) = \frac{1}{1 - qu} = \prod_{p \in \mathcal{P}} (1 - u^{\deg(p)})^{-1}.$$

As we can see $\zeta(u)$ can be continued meromorphically on \mathbb{C} with a simple pole at $u = 1/q$. Furthermore considering the Euler products in (2.1) and (2.2) we can connect $G_z(u)$ and $F_z(u)$ in the following way whenever $|u| < \min\{|z|^{-1}, q^{-\frac{1}{2}}\}$


$$G_z(u) = \zeta(u)^z F_z(u). \tag{2.4}$$

We are now in a position to prove our first Proposition. However, first, we want to record the following estimate [6, Lemma 1], which is going to be helpful in the further calculation of the contribution that comes from the singularity of $G_z(u)$ at $u = 1/q$.

Lemma 2.2. *Let $A, \delta > 0$. Let \mathcal{H} be the Hankel contour of radius 1 around 0 going in clockwise direction along the negative real axis to $-n\delta$. Then uniformly for $|z| \leq A$ we have that*

$$\frac{1}{2\pi i} \int_{\mathcal{H}} w^z \frac{dw}{\left(1 - \frac{w}{n}\right)^{n+1}} = -\frac{1}{\Gamma(-z)} + O_{\delta, A} \left(\frac{1}{n} \right).$$

FIGURE 1. THE CONTOUR



2.2. Deduction of Proposition 1.3.

Proof of Proposition 1.3. Let $|u| = r < q^{-1}$ (small green circle in Fig-1 above). Using Cauchy's integral formula on (2.1) we get for each fixed z satisfying the hypothesis $|z| \leq q - \epsilon$,

$$(2.5) \quad M_z(n) = \frac{1}{2\pi i} \int_{|u|=r} G_z(u) \frac{du}{u^{n+1}} = \frac{1}{2\pi i} \int_{|u|=r} \zeta(u)^z F_z(u) \frac{du}{u^{n+1}}.$$

We wish to get past the point $u = 1/q$ so that we could collect the contribution coming from the singularity of $\zeta(u)$ there. We shift the smaller (green) circle $|u| = r$ to a bigger circle (red) $|u| = q^{-1} + \eta$ and a portion of Hankel contour around the point q^{-1} (Fig-1). Let \mathcal{H}' be the contour that consists of a circle of radius $(qn)^{-1}$ traversed clockwise around q^{-1} and the two line segments on the ray 0 to q^{-1} joining this small circle to the bigger circle $|u| = q^{-1} + \eta$ where $\eta > 0$ is a fixed number. (to be specified shortly) However we must have $q^{-1} + \eta < q^{-\frac{1}{2}}$ as we cannot go beyond the region \mathcal{R} in (2.4) where $G_z(u)$ makes sense. We recall that $|z| \leq (q - \epsilon)$ inside \mathcal{R} while performing the above integral. Keeping this in mind we get another constraint on η , $|uz| \leq (q^{-1} + \eta)(q - \epsilon) \leq 1$. This shows taking any $0 < \eta < \min\{1/(q - \epsilon) - 1/q, 1/\sqrt{q} - 1/q\}$ suffices. We then plan to use Lemma 2.2 to evaluate the integral in (2.5) with $\eta = \min\{1/(q - \epsilon) - 1/q, 1/\sqrt{q} - 1/q\}/2$ for $|z| \leq q - \epsilon$. We observe that for z fixed, $F_z(u)$ and $\zeta(u)$ are both analytic in an open neighbourhood containing the circle $|u| = q^{-1} + \eta$ which is compact. This implies $F_z(u)$, $\zeta(u)$ and hence $G_z(u)$ are

uniformly bounded (bound depends on q, η) on this circle. Thus we obtain

$$M_z(n) = \frac{1}{2\pi i} \int_{\mathcal{H}'} G_z(u) \frac{du}{u^{n+1}} + O \left(\int_{|u|=\frac{1}{q}+\eta} \left| G_z(u) \frac{du}{u^{n+1}} \right| \right) := I + O_{q,\epsilon} \left(\frac{q^n}{(1+q\eta)^n} \right),$$

where at the last step we used the fact that, $\int_{|u|=R} |du/u^{n+1}| = \frac{1}{R^{n+1}} \int_{|u|=R} |du| = 2\pi/R^n$ and the big-O-uniform bound here depends only on q, η and η according our choice depends only on q and ϵ . Now we focus on the integral I .

Analyticity of F_z near q^{-1} allows us to consider the Taylor series around q^{-1} and write

$$F_z(u) = F_z(1/q) + a_1(u - 1/q) + a_2(u - 1/q)^2 + \dots$$

Hence we have that

$$I = F_z \left(\frac{1}{q} \right) \frac{1}{2\pi i} \int_{\mathcal{H}'} \zeta(u)^z \frac{du}{u^{n+1}} + O \left(\int_{\mathcal{H}'} \left| \left(u - \frac{1}{q} \right) \zeta(u)^z \frac{du}{u^{n+1}} \right| \right).$$

We use a change of variable so that our contour \mathcal{H}' transforms to \mathcal{H} and we could then use our Lemma 2.2. The change of variable is given by, $w = n(1 - uq) \implies u = 1/q(1 - w/n)$. We can quickly check the transformation is indeed a valid one. If u is parameterized by $u = 1/q + (1/qn)e^{-it}$, $t \in [\epsilon', 2\pi - \epsilon']$ for some small enough $\epsilon' > 0$ near the tiny circle around $1/q$ then $w = -e^{-it}$ is a circle revolving around 0 of radius 1 in the clockwise direction and similarly the horizontal ray part also gets reflected and \mathcal{H}' indeed transforms into \mathcal{H} and as $|u|$ goes up to $1/q + \eta$, we have w goes up to $-n\delta$ with $\delta = \eta q$. We note for this change of variable $du = -dw/nq$. First we show the alleged error term is small. The O -term is

$$\begin{aligned} \ll_q \int_{\mathcal{H}'} \left| \frac{(1-qu)du}{u^{n+1}(1-qu)^z} \right| &= \int_{\mathcal{H}} \left| \frac{1}{\left(\frac{w}{n}\right)^{z-1} \frac{1}{q^{n+1}} \left(1 - \frac{w}{n}\right)^{n+1}} \frac{-dw}{nq} \right| \\ &= \frac{q^n}{n} \int_{\mathcal{H}} \left| \frac{1}{\left(\frac{w}{n}\right)^{z-1} \left(1 - \frac{w}{n}\right)^{n+1}} dw \right| \quad (\text{using Lemma 2.2}) \\ &\ll q^n n^{\Re(z)-2}. \end{aligned}$$

Now, we evaluate the main term using Lemma 2.2

$$\begin{aligned} F_z \left(\frac{1}{q} \right) \frac{1}{2\pi i} \int_{\mathcal{H}'} \zeta(u)^z \frac{du}{u^{n+1}} &= F_z \left(\frac{1}{q} \right) \frac{1}{2\pi i} \int_{\mathcal{H}'} \frac{1}{(1-qu)^z} \frac{du}{u^{n+1}} \\ &= F_z \left(\frac{1}{q} \right) \frac{q^n}{n} \frac{1}{2\pi i} \int_{\mathcal{H}} \frac{1}{\left(\frac{w}{n}\right)^z} \frac{-dw}{\left(1 - \frac{w}{n}\right)^{n+1}} \\ &= F_z \left(\frac{1}{q} \right) q^n n^{z-1} \left(\frac{1}{\Gamma(z)} + O_\epsilon \left(\frac{1}{n} \right) \right). \end{aligned}$$

Hence we have that

$$(2.6) \quad M_z(n) = F_z \left(\frac{1}{q} \right) \frac{q^n n^{z-1}}{\Gamma(z)} + O(q^n n^{\Re(z)-2}) = q^n n^{z-1} \left\{ zH(z) + O_\epsilon \left(\frac{1}{n} \right) \right\},$$

where $H(z) = \frac{1}{\Gamma(z+1)}F_z\left(\frac{1}{q}\right)$ is an analytic function for $|z| < q$. □

3. DEDUCTION OF THEOREM 1.1

With Proposition 1.3 at our disposal, we are now ready to prove Theorem 1.1. We also note three other straightforward results which will help us do so. We will verify the following claims in the appendix A.

Lemma 3.1. *For $t \in [-\pi, \pi]$ we have $\cos t - 1 \leq -\frac{t^2}{5}$ and $|1 - e^{it}|^2 \leq t^2$.*

Lemma 3.2. *We have the following upper bound*

$$\int_{|z|=r} |n^{z-r}| |dz| \ll \frac{r}{\sqrt{j}},$$

where $j = r \log n$.

Lemma 3.3. *We have the following estimate*

$$\int_{-\infty}^{\infty} \theta^2 \exp\left(-\frac{k\theta^2}{5}\right) d\theta \ll k^{-\frac{3}{2}}.$$

Deduction of Theorem 1.1. We plan to recover $N(n, k)$ with another application of Cauchy's integral formula,

$$(3.1) \quad N(n, k) = \frac{1}{2\pi i} \int_{|z|=\frac{k}{\log n}} M_z(n) \frac{dz}{z^{k+1}}.$$

Given $\epsilon \in (0, q)$ by the hypothesis of Theorem 1.1, from (2.6) we see the main term of $M_z(n)$ which is $F_z\left(\frac{1}{q}\right) \frac{q^n n^{z-1}}{\Gamma(z)}$ is analytic in the disk $|z| = r = k/\log n \leq (q - \epsilon)$ since $F_z\left(\frac{1}{q}\right)$ is analytic for $|z| < q$ and $n^z, 1/\Gamma$ are entire functions. Now writing $M_z(n)/q^n = n^{z-1}(f(z) + O(1/n))$ where $f(z) = zH(z) = F_z(1/q) \frac{1}{\Gamma(z)}$ and considering the Taylor expansion of $f(z)$ near $z = r$, $f(z) = f(r) + f'(r)(z - r) + O(|z - r|^2)$ we get

$$(3.2) \quad \frac{1}{q^n} N(n, k) = \frac{1}{2\pi i} \int_{|z|=r} n^{z-1} f(z) \frac{dz}{z^{k+1}} + O(E),$$

where E is given by $\frac{1}{n} \int_{|z|=r} \left| \frac{n^{z-1}}{z^{k+1}} \right| |dz|$. We see that

$$\frac{1}{2\pi i} \int_{|z|=r} (z - r) n^{z-1} \frac{dz}{z^{k+1}} = \frac{1}{n} \left\{ \frac{(\log n)^{k-1}}{(k-1)!} - r \frac{(\log n)^k}{k!} \right\} = 0.$$

Hence

$$\begin{aligned} \frac{1}{q^n} N(n, k) &= f(r) \frac{1}{2\pi i} \int_{|z|=r} n^{z-1} \frac{dz}{z^{k+1}} + O\left(r^{-k-1} \int_{|z|=r} |n^{z-1}(z-r)^2| |dz| + E\right) \\ &= f(r) \frac{(\log n)^k}{k!n} + \text{Error term.} \end{aligned}$$

By using Lemma 3.1 and Lemma 3.3, we see the integral in the Error term is bounded by

$$\frac{r^3}{n} \int_{-\pi}^{\pi} |1 - e^{i\theta}|^2 e^{k \cos \theta} d\theta \ll \frac{r^3}{n} \int_{-\infty}^{\infty} \theta^2 e^{k(1 - \frac{\theta^2}{5})} d\theta \ll \frac{r^3}{n} e^k k^{-\frac{3}{2}}.$$

Since there is a factor of r^{-k-1} outside, we have the above expression (final step by Stirling's formula)

$$\ll \frac{1}{n} r^{2-k} e^k k^{-\frac{3}{2}} \ll (\log n)^{k-2} \frac{e^k \sqrt{k}}{nk^k} \ll \frac{1}{n} \frac{(\log n)^{k-2}}{(k-1)!}.$$

Noting $e^k = n^r \iff k = r \log n$ and using Lemma 3.2, we also see the term E is bounded by

$$\ll \frac{r^{-k-1} e^k}{n^2} \int_{|z|=r} |n^{z-r}| |dz| \ll \frac{r^{-k-1} e^k}{n^2} \left(\frac{r}{\sqrt{k}}\right) \ll \frac{1}{n^2} \frac{(e \log n)^k}{k^{k+\frac{1}{2}}} \ll \frac{1}{n^2} \frac{(\log n)^k}{k!},$$

where in the last step we have used Stirling. Thus we finally obtain

$$N(n, k) \sim q^n f(r) \frac{(\log n)^k}{k!n} = \frac{q^n}{n} F_r \left(\frac{1}{q}\right) \frac{1}{\Gamma(r+1)} \frac{(\log n)^{k-1}}{(k-1)!}$$

and the proof follows once we recall $r = k/\log n$. The derived expression is identical to the integer case for $k \leq (2 - \epsilon) \log \log x$ as mentioned in [1] on the first page, a result due to Sathe(1953) [4]. □

4. PREPARATORY RESULTS AND AN ODD-EVEN DECOMPOSITION

4.1. An odd-even decomposition for polynomials. We begin this subsection by introducing a decomposition for $\Pi(x, k)$ into k smaller parts used by Nicolas [1] to prove (1.2). We first note that any integer $n \leq x$ can be uniquely written as $n = 2^j \ell$ with ℓ odd. We note the following identity

$$(4.1) \quad \Pi(x, k) = \sum_{0 \leq j \leq k} \Pi'(x/2^{k-j}, j),$$

where

$$\Pi'(x, k) := \#\left\{m \leq x : m \text{ odd and } \Omega(m) = k\right\}$$

holds. The key idea discussed in [1] says the major contribution in the former sum comes if we restrict $0 \leq j \leq \alpha \log \log x$ for some explicit constant $\alpha > 2$ whenever k ranges from $2 \log \log x$ to $\log x / \log 2$.

Inspired from the idea above we wish to write $f = hg$ for any $f \in \mathcal{M}_n$, where the polynomials g and h are expected to play the role of ‘‘odd’’ and ‘‘even’’ respectively in some appropriate sense. Let us define the set

$$\mathcal{S}(j) := \left\{h \in \mathcal{M}_j : h \text{ has only degree 1 irreducible factors}\right\}.$$

Given $f \in \mathcal{M}_n$ with $\Omega(f) = k$, we can uniquely write $f = hg$ where $h \in \mathcal{S}(j)$ for some $0 \leq j \leq k$ and g has no degree 1 irreducible factor. Elements of the set $\mathcal{S}(j)$ mimic the even part while the odd part is mimicked by the polynomial g . Using this heuristic we could prove the following decomposition result for $N(n, k)$.

Lemma 4.1. *We have*

$$N(n, k) = \sum_{1 \leq j \leq k} \binom{k-j+q-1}{q-1} N'(n+j-k, j)$$

for $1 \leq k \leq n/2$.

Proof. We observe that for any $h \in \mathcal{S}(j)$, one can find non negative integers a_1, a_2, \dots, a_q such that $h(t) = (t - \alpha_1)^{a_1} (t - \alpha_2)^{a_2} \dots (t - \alpha_q)^{a_q}$ with $\Omega(h) = a_1 + a_2 + \dots + a_q = j$. Here $\{\alpha_i\}_{1 \leq i \leq q}$ are the q -elements in \mathbb{F}_q . Counting the number of non negative integral solutions we readily see that, $\#\mathcal{S}(j) = \binom{j+q-1}{q-1}$. If we consider the unique decomposition $f = hg$ for some $f \in \mathcal{M}_n$ with $\Omega(f) = k$ then we note that $j \neq k$. For if $j = k$ then $f = hg$ will imply $f = h$ and hence they both have the same degree, a contradiction since $k \leq n/2$ by the hypothesis. Therefore we obtain

$$N(n, k) = \sum_{0 \leq j \leq k-1} \sum_{h \in \mathcal{S}(j)} N'(n-j, k-j) = \sum_{0 \leq j \leq k-1} \binom{j+q-1}{q-1} N'(n-j, k-j).$$

The statement of the lemma follows after we make the change of variable $j \mapsto k-j$. \square

The rough sketch of our strategy to prove the Theorem 1.2 is: (a) In light of the approach discussed in (4.1) in the integer setting, we start by splitting the whole sum in the above decomposition up to $j \ll \log(n-k)$ and hence we end up with two parts so that we can write, $T_1 = \sum_{1 \leq j \ll \log(n-k)} \binom{k-j+q-1}{q-1} N'(n+j-k, j)$ and $T_2 = \sum_{\log(n-k) \ll j \leq k} \binom{k-j+q-1}{q-1} N'(n+j-k, j)$.

(b) We then show the main contribution comes from T_1 and is of order $q^{n-k} k^{q-1} (n-k)^{q-1}$ while the term T_2 is significantly smaller and is of order $q^{n-k} (n-k)^{q-2}$. Unlike the ideas discussed in the integer case, which is very combinatorial, our proof will be more analytical. The central idea to use a Selberg-Delange type argument followed by an application of Cauchy's integral formula shall remain the same. The situation over $\mathbb{F}_q[t]$ is quite different due to the presence of many degree 1 irreducibles which is why an extra binomial term and other technical difficulties arise.

We begin our preparation with the easier task, a non-trivial upper bound on T_2 . To do so, we would like an upper bound on each summand of T_2 . The following result accomplishes this job.

Lemma 4.2. *Let $0 < \delta < 1, 0 < \eta \leq q^2 - q - \delta$ be given. We have the following upper bound*

$$N'(n, j) \ll_{\delta} q^n n^{q+\eta-1} \left(\frac{1}{q+\eta} \right)^j.$$

Throughout our discussion, the implied constant solely depends on q, δ , which will be a fixed quantity. (the value of δ is soon to be decided in the following result) It is possible to show the above result as a direct application Proposition 1.4. The proof of Proposition 1.4 will roughly be in the same spirit as that of Proposition 1.3. We will present this proof followed by an argument

leading to Lemma 4.2 in the appendix B.1, B.2. We now shift our attention to check that the term T_2 is small as claimed.

Proposition 4.3. *Let $q > 2$ be an integer and $Y = \log(n - k)$. We have the following upper bound*

$$T_2 = \sum_{eqY < j \leq k} \binom{q-1+k-j}{q-1} N'(n+j-k, j) \ll_B q^{n-k} (n-k)^{q-2}.$$

Proof. We plan to use $\binom{\alpha}{\beta} \leq \alpha^\beta / \beta!$. We apply Lemma 4.2 with $\eta = (e-1)q$ and $\delta = 0.6$ so that $(q+\eta) = eq$. First we check this choice satisfies the hypothesis of Lemma 4.2. We see that $(e-1)q < q^2 - q - 0.6$ is true whenever $q \geq 3$. We now have that

$$\begin{aligned} T_2 &\ll \sum_{j > (q+\eta)Y} \frac{(q-1+k-j)^{q-1}}{(q-1)!} q^{n-k+j} (n+j-k)^{q+\eta-1} \left(\frac{1}{q+\eta}\right)^j \\ &\ll q^{n-k} \sum_{j > (q+\eta)Y} (q-1+k-j)^{q-1} (n+j-k)^{q+\eta-1} \left(\frac{q}{q+\eta}\right)^j \\ &= q^{n-k} \sum_{j > eqY} (n-k)^{q-1} \left(\frac{q-1}{n-k} + \frac{k-j}{n-k}\right)^{q-1} (n-k)^{q+\eta-1} \left(1 + \frac{j}{n-k}\right)^{q+\eta-1} e^{-j}, \end{aligned}$$

where we used $q+\eta = eq$. Since $k \leq n/B$ we have $\left(\frac{q-1}{n-k} + \frac{k-j}{n-k}\right)^{q-1} \leq \left(q-1 + \frac{k}{n-k}\right)^{q-1} \leq \left(q-1 + \frac{1}{B-1}\right)^{q-1} \ll_B 1$. Also $k+j \leq 2k \leq n$ gives $\left(1 + \frac{j}{n-k}\right)^{eq-1} \ll 1$. Recalling $Y = \log(n-k)$ and continuing from the previous step we get

$$T_2 \ll_B q^{n-k} (n-k)^{eq+q-2} \sum_{j > eqY} e^{-j} \ll_B q^{n-k} (n-k)^{eq+q-2} e^{-eqY} = q^{n-k} (n-k)^{q-2}.$$

□

We are now only left to estimate T_1 and show it dominates in the asymptotic of $N(n, k)$. For the ease of exploration, the calculation of T_1 is broken down into three smaller sub-parts. To have a good estimate for $T_1 = \sum_{1 \leq j \leq \log(n-k)} \binom{k-j+q-1}{q-1} N'(n+j-k, j)$, first it is natural to want to understand the terms $N'(n+j-k, j)$ when the range for j is restricted to $\ll \log(n-k)$. With these observations in mind we could show the following result.

Lemma 4.4. *Let $Y = \log(n-k)$. We have uniformly for $j \leq eqY$,*

$$N'(n+j-k, j) = \frac{q^{n+j-k}}{n-k} \left\{ Q_j(Y) + O\left(\frac{(\log(n-k))^{j+1}}{j!(n-k)}\right) \right\}$$

where

$$Q_j(X) := \sum_{m+\ell=j-1} \frac{1}{m!\ell!} h^{(m)}(0) X^\ell.$$

If we are given the above result in our toolbox, it is then evident that we are just a few steps away from proving an estimate for T_1 once we take care of what happens with the sum when the terms $N'(n + j - k, j)$ are twisted by appropriate binomial coefficients for different j 's. However, before we proceed to do that, we would like to mention a few words about what goes into the proof of Lemma 4.4. The proof of the above result rests on another two fairly technical auxiliary lemmas, which we shall state next. Details of these three claims are postponed till the appendix B.3, B.4, B.5 and can be skipped at the moment as they do not offer many insights into our actual goal.

Lemma 4.5. *We have uniformly for $j \leq eq \log n$,*

$$N'(n, j) = \frac{q^n}{n} \left\{ Q_j(\log n) + O\left(\frac{(\log n)^j}{j!n}\right) \right\}.$$

Lemma 4.6. *Let $m \geq 1$ be an integer. We have the following upper bounds*

$$h^{(m)}(0)/m! \ll \left(\frac{1}{2.9q}\right)^m,$$

$$Q_j(X) \ll \frac{X^{j-1}}{(j-1)!} \text{ and } Q'_j(X) \ll \frac{X^{j-2}}{(j-2)!},$$

for $1 \leq j \leq eqX$ uniformly, where X is a parameter that tends to ∞ with n .

We can now set up the stage for proving Theorem 1.2. As mentioned earlier we need to have a precise estimate for the twisted binomial sum appearing in T_1 . Therefore the following Proposition plays a key role in establishing Theorem 1.2.

Proposition 4.7. *Let $Y = \log(n - k)$. We have the following estimate*

$$\sum_{j \leq eqY} \binom{q-1+k-j}{q-1} q^j Q_j(Y) = \frac{k^{q-1} q e^{qY}}{(q-1)!} h(q) \{1 + o(1)\}.$$

Proof. We start with an observation first. By the hypothesis of Theorem 1.2 we have $k > \xi(n) \log n$ yielding $k - j \geq \xi(n) \log n - eq \log(n - k) \geq (\xi(n) - eq) \log n \rightarrow \infty$. Therefore using the standard fact $\binom{\alpha}{\beta} = (1 + o(1)) \frac{\alpha^\beta}{\beta!}$ where β is assumed to be fixed and $\alpha \rightarrow \infty$ we obtain

$$\begin{aligned} \sum_{j \leq eqY} \binom{q-1+k-j}{q-1} q^j Q_j(Y) &= \frac{1}{(q-1)!} \sum_{j \leq eqY} \left\{ q^j (q-1+k-j)^{q-1} Q_j(Y) \right\} (1 + o(1)) \\ &= \frac{k^{q-1}}{(q-1)!} \sum_{j \leq eqY} \left\{ q^j Q_j(Y) \right\} (1 + o(1)), \end{aligned}$$

where we used the Taylor expansion of $(k + q - j - 1)^{q-1} = k^{q-1}(1 + o(1))$ since $j = o(k)$.

We now evaluate the sum

$$\begin{aligned}
\sum_{j \leq eqY} q^j Q_j(Y) &= \left(\sum_{j \geq 1} q^j \sum_{\substack{m+\ell=j-1 \\ m, \ell \geq 0}} \frac{h^m(0)}{m! \ell!} Y^\ell \right) - R \\
&= \left(q \sum_{\ell \geq 0} \frac{(qY)^\ell}{\ell!} \sum_{j-m=\ell+1} \frac{h^m(0)}{m!} q^{j-\ell-1} \right) - R \quad (\text{after interchanging the sum}) \\
&= \left(q \sum_{\ell \geq 0} \frac{(qY)^\ell}{\ell!} \sum_{m \geq 0} \frac{h^m(0)}{m!} q^m \right) - R \quad (\text{since } m = j - \ell - 1 \text{ and } j \geq \ell + 1) \\
&= qe^{qY} h(q) - R, \quad (h(z) \text{ is analytic for } |z| < q^2)
\end{aligned}$$

where

$$\begin{aligned}
R &= \sum_{j > [eqY]} q^j Q_j(Y) \quad (\text{where } [x] \text{ denotes the greatest integer } \leq x) \\
&= \sum_{j \geq [eqY]+1} q^j \sum_{\substack{m+\ell=j-1 \\ m, \ell \geq 0}} \frac{h^m(0)}{m! \ell!} Y^\ell \\
&= q \left(\sum_{0 \leq \ell \leq [eqY]} \frac{(qY)^\ell}{\ell!} \sum_{m \geq [eqY]-\ell} \frac{h^m(0)}{m!} q^m \right) + q \left(\sum_{\ell \geq [eqY]} \frac{(qY)^\ell}{\ell!} \sum_{m \geq 0} \frac{h^m(0)}{m!} q^m \right) \\
&\ll_q S_1 + S_2. \quad (\text{say})
\end{aligned}$$

We have the following estimates,

$$\begin{aligned}
S_1 &\leq \sum_{0 \leq \ell \leq [eqY]} \frac{(qY)^\ell}{\ell!} \sum_{m \geq [eqY]-\ell} \left(\frac{1}{2} \right)^m \quad (\text{using } h^{(m)}(0)/m! < (1/2q)^m \text{ from Lemma 4.6}) \\
&= \sum_{0 \leq \ell \leq [eqY]} \frac{(qY)^\ell}{\ell!} \left(\frac{1}{2} \right)^{[eqY]-\ell} \\
&= \left(\frac{1}{2} \right)^{[eqY]} \sum_{0 \leq \ell \leq [eqY]} \frac{(2qY)^\ell}{\ell!} \\
&\leq e^{0.12qY}. \quad \left(\text{since } \left(\frac{1}{2} \right)^{[eqY]} = e^{-1.88qY} \text{ and the rest is bounded by the complete sum } e^{2qY} \right)
\end{aligned}$$

Similarly

$$\begin{aligned}
 S_2 &= h(q) \sum_{\ell \geq \lfloor eqY \rfloor} \frac{(qY)^\ell}{\ell!} \\
 &\ll (qY)^{\lfloor eqY \rfloor} \sum_{\ell \geq 0} \frac{(qY)^\ell}{(\ell + \lfloor eqY \rfloor)!} \quad (\text{dropping } h(q) \text{ and using a change of variable } \ell \mapsto \ell - \lfloor eqY \rfloor) \\
 &\leq \frac{(qY)^{\lfloor eqY \rfloor}}{\lfloor eqY \rfloor!} \sum_{\ell \geq 0} \frac{(qY)^\ell}{\ell!} \left(\text{since } \binom{\ell + \lfloor eqY \rfloor}{\ell} \geq 1 \text{ so that } \frac{1}{(\ell + \lfloor eqY \rfloor)!} \leq \frac{1}{\ell! \lfloor eqY \rfloor!} \right) \\
 &\ll \frac{1}{\sqrt{\lfloor eqY \rfloor}} e^{qY},
 \end{aligned}$$

where at the last step we used Stirling. Indeed letting $p = eqY$, we have

$$\frac{(qY)^{\lfloor eqY \rfloor}}{\lfloor eqY \rfloor!} \ll \frac{\left(\frac{p}{e}\right)^{\lfloor p \rfloor}}{\left(\frac{\lfloor p \rfloor}{e}\right)^{\lfloor p \rfloor} \sqrt{\lfloor p \rfloor}} = \left(\frac{p}{\lfloor p \rfloor}\right)^{\lfloor p \rfloor} \frac{1}{\sqrt{\lfloor p \rfloor}} \leq \left(1 + \frac{1}{\lfloor p \rfloor}\right)^{\lfloor p \rfloor} \frac{1}{\sqrt{\lfloor p \rfloor}} \leq \frac{e}{\sqrt{\lfloor p \rfloor}}.$$

□

Proof of Theorem 1.2. We have that

$$\begin{aligned}
 N(n, k) &= \sum_{1 \leq j \leq eqY} \binom{k-j+q-1}{q-1} N'(n+j-k, j) + \sum_{j > eqY} \binom{k-j+q-1}{q-1} N'(n+j-k, j) \\
 &= T_1 + T_2.
 \end{aligned}$$

Proposition 4.3 shows the tail T_2 is small and is $\ll q^{n-k}(n-k)^{q-2}$.

Lemma 4.4 and Proposition 4.7 implies,

$$\begin{aligned}
 T_1 &= \sum_{j \leq eqY} \binom{k-j+q-1}{q-1} \frac{q^{n+j-k}}{n-k} Q_j(Y) + O\left(\sum_{j \leq eqY} \binom{k-j+q-1}{q-1} \frac{(\log(n-k))^{j+1}}{j!(n-k)}\right) \\
 &= \frac{q^{n-k}}{n-k} \sum_{j \leq eqY} \binom{k-j+q-1}{q-1} q^j Q_j(Y) + O\left(\sum_{j \leq eqY} \binom{k-j+q-1}{q-1} \frac{(\log(n-k))^{j+1}}{j!(n-k)}\right) \\
 &= \frac{q^{n-k+1} k^{q-1} e^{qY}}{(q-1)!(n-k)} h(q) \{1 + o(1)\} + O\left((1 + o(1)) \frac{k^{q-1}}{(q-1)!} \sum_{j=1}^{\infty} \frac{(\log(n-k))^{j+1}}{j!(n-k)}\right) \\
 &= \frac{q^{n-k+1} k^{q-1} e^{qY}}{(q-1)!(n-k)} h(q) \{1 + o(1)\} + O\left(k^{q-1} \frac{\log(n-k) e^{\log(n-k)}}{n-k}\right) \\
 &= \frac{q^{n-k+1} k^{q-1} e^{qY}}{(q-1)!(n-k)} h(q) \{1 + o(1)\} + O(k^{q-1} \log(n-k)).
 \end{aligned}$$

Thus we obtain

$$T_1 \sim \frac{q^{n-k} k^{q-1} e^{qY}}{(n-k)} \frac{qh(q)}{(q-1)!} \sim C(q) \frac{q^n k^{q-1} (n-k)^{q-1}}{q^k},$$

where we used $e^{qY}/(n-k) = (n-k)^{q-1}$ when $Y = \log(n-k)$ and

$$C(q) = \frac{qh(q)}{(q-1)!} = \frac{1}{((q-1)!)^2} \left(1 - \frac{1}{q}\right)^{q^2} \prod_{p \in \mathcal{P}_{\geq 2}} \left(1 - \frac{1}{q^{\deg(p)-1}}\right)^{-1} \left(1 - \frac{1}{q^{\deg(p)}}\right)^q,$$

an absolute constant that only depends on q . \square

APPENDIX A. AUXILIARY RESULTS TOWARDS THEOREM 1.1

We establish the technical Lemmas 3.1, 3.2 and 3.3, used in the proofs of Theorem 1.1 and will again be used in showing Proposition 1.4.

Proof of Lemma 3.1. We define the function $f(t) := 1 - \frac{t^2}{5} - \cos t$ for $t \in [0, \pi]$ and observe that $f'(t) = 0 \implies \sin t = \frac{2t}{5}$. A quick graph plot or plugging this equation into a computer reveals that it has only two solutions in the interval $[0, \pi]$, namely $t = 0, 2.125$. Checking that $f''(0) = -\frac{2}{5} + \cos 0 = \frac{3}{5} > 0$ and hence $t = 0$ is the global minima in the interval $[0, \pi]$ since $0 = f(0) < f(2.125)$ which can also be verified on a computer. Finally since $\cos t$ and t^2 are both even functions we have for all $t \in [-\pi, \pi]$ the following inequality,

$$f(t) \geq 0 \implies 1 - \frac{t^2}{5} - \cos t \geq 0 \implies \cos t - 1 \leq -\frac{t^2}{5}.$$

For $0 \leq x, t \leq \pi$ it is well known that,

$$\sin x \leq x \implies \int_0^t \sin x dx \leq \int_0^t x dx \implies 1 - \cos t \leq \frac{t^2}{2}.$$

Again the above inequality is true for $-\pi \leq t \leq \pi$ since \cos and t^2 are both even functions. Finally we observe $|1 - e^{it}|^2 = (1 - \cos t)^2 + \sin^2 t = 2(1 - \cos t) \leq t^2$ and the claim follows. \square

Proof of Lemma 3.2. Let $I_0 = \int_{|z|=r} |n^{z-r}| |dz|$. After substituting $z = re^{2\pi it}$ where $t \in \left[-\frac{1}{2}, \frac{1}{2}\right)$ we get

$$\begin{aligned} I_0 &\ll \int_{-\frac{1}{2}}^{\frac{1}{2}} |n^{re^{2\pi it}-r}| r dt \\ &= r \int_{-\frac{1}{2}}^{\frac{1}{2}} |n^{r(\cos 2\pi t-1)}| dt \quad (|n^{ir \sin 2\pi t}| = 1) \\ &\ll r \int_{-\frac{1}{2}}^{\frac{1}{2}} n^{-\frac{4\pi^2 r t^2}{5}} dt \quad (\text{by Lemma 3.1}) \\ &\ll r \int_{-\infty}^{\infty} e^{-t^2} \frac{dt}{\sqrt{r \log n}} \quad \left(t \mapsto 2\pi t \sqrt{\frac{r \log n}{5}}\right) \\ &\ll r \frac{1}{\sqrt{r \log n}} = \frac{r}{\sqrt{j}}. \quad (j = r \log n) \end{aligned}$$

\square

Proof of Lemma 3.3.

$$\int_{-\infty}^{\infty} \theta^2 \exp\left(-\frac{k\theta^2}{5}\right) d\theta = \frac{5\sqrt{5}}{k^{\frac{3}{2}}} \int_0^{\infty} \sqrt{z} e^{-z} dz \quad \left(z = \frac{k\theta^2}{5}\right)$$

$$\ll k^{-\frac{3}{2}}. \quad \left(\text{since } \int_0^{\infty} \sqrt{z} e^{-z} dz \text{ is convergent}\right)$$

□

APPENDIX B. TOWARDS PROPOSITION 1.4 AND AUXILIARY RESULTS FOR THEOREM 1.2

B.1. A Selberg-Delange style argument for $\widetilde{M}_n(n)$.

Proof of Proposition 1.4. We recall some definitions for convenience,

$$\widetilde{\mathcal{M}} := \left\{ \text{set of all monics in } \mathbb{F}_q[t] \text{ having no root in } \mathbb{F}_q \right\},$$

$$\widetilde{\mathcal{M}}_n := \left\{ \text{set of all of degree } n \text{ monics in } \mathbb{F}_q[t] \text{ having no root in } \mathbb{F}_q \right\},$$

$$N'(n, k) := \#\left\{ f \in \widetilde{\mathcal{M}}_n : \Omega(f) = k \right\}.$$

We proceed as the proof of Theorem 1.1 by first defining the following quantities,

$$\widetilde{M}_n(z) = \sum_{f \in \widetilde{\mathcal{M}}_n} z^{\Omega(f)} = \sum_{k \geq 0} z^k N'(n, k) \text{ and } \widetilde{G}_z(u) = \sum_{f \in \widetilde{\mathcal{M}}} z^{\Omega(f)} u^{\deg(f)}.$$

Our Euler product decomposition for $\widetilde{G}_z(u)$ now becomes

$$\widetilde{G}_z(u) = \prod_{p \in \mathcal{P}_{\geq 2}} (1 - zu^{\deg(p)})^{-1}.$$

Following the same spirit we now have

$$\widetilde{G}_z(u) = \zeta(u)^z \widetilde{F}_z(u),$$

where \widetilde{F}_z is given by the Euler product below

$$\widetilde{F}_z(u) = (1 - u)^{qz} \prod_{p \in \mathcal{P}_{\geq 2}} (1 - zu^{\deg(p)})^{-1} (1 - u^{\deg(p)})^z.$$

We first show the above Euler product converges absolutely in the region defined by,

$$\widetilde{\mathcal{R}} := \left\{ (u, z) \in \mathbb{C}^2 : |z| < q^2, |u|^2 < |z|^{-1}, |u| < q^{-1/2} \right\} \subset \mathbb{C}^2.$$

Since $|1 - zu^{\deg(p)}| \geq 1 - |zu^{\deg(p)}| \geq 1 - |zu^2| > 0$, we have none of the $(1 - zu^{\deg(p)})^{-1}$ factors contribute to a pole in $\widetilde{\mathcal{R}}$.

We take the standard branch of complex logarithm defined over $\mathbb{C} \setminus (-\infty, 0]$ and get for each integer $n \geq 2$, (since $|u|^n < 1, |zu^n| \leq |zu^2| < 1$)

$$\begin{aligned} z \log(1 - u^n) - \log(1 - zu^n) &= -zu^n + O(|zu^{2n}|) + zu^n + O(|z^2u^{2n}|) \\ &= O(|q^4u^{2n}|) \quad (|z| < q^2). \end{aligned}$$

(B.1)

and hence using $\exp(O(x)) = 1 + O(x)$ for $x = O(1)$ we get,

$$(1 - u^n)^z (1 - zu^n)^{-1} = 1 + O(|q^4u^{2n}|).$$

We now observe the Euler product for \widetilde{F}_z converges absolutely in $\widetilde{\mathcal{R}}$ as

$$\sum_{p \in \mathcal{P}_{\geq 2}} |u|^{2n} \leq \sum_{n \geq 2} |qu^2|^n < \infty. \quad (\text{where we used, } \#\mathcal{P}_n \leq q^n)$$

Also $(1 - u)^{qz}$ is holomorphic in $\widetilde{\mathcal{R}}$ and this proves $\widetilde{F}_z(u)$ is holomorphic inside $\widetilde{\mathcal{R}}$. We determine the coefficient of u^n in $\widetilde{G}_z(u)$ using Cauchy's integral formula and recover $\widetilde{M}_z(n)$ exactly as in the proof of Theorem 1.1. We have that, if $|u| = r < q^{-1}$ (refer to the small green circle in the same Fig-1), then using Cauchy's residue formula on (2.1) gives

$$(B.2) \quad \widetilde{M}_z(n) = \frac{1}{2\pi i} \int_{|u|=r} \widetilde{G}_z(u) \frac{du}{u^{n+1}} = \frac{1}{2\pi i} \int_{|u|=r} \zeta(u)^z \widetilde{F}_z(u) \frac{du}{u^{n+1}}.$$

Now we again want to collect the contribution coming from the singularity of $\zeta(u)^z$ at $u = 1/q$ to evaluate the above integral. We shift the contour $|u| = r$ to a bigger circle $|u| = 1/q + \widetilde{\eta}$ with a different choice of $\widetilde{\eta}$ here (not the same choice for η in Proposition 1.3) and with $|z| \leq q^2 - \delta$. Since the region of absolute convergence for $\widetilde{F}_z(u)$ here is $\widetilde{\mathcal{R}}$, we have a different set of constraints on $\widetilde{\eta}$ than before. Here we get for $\widetilde{\eta}$, $|zu^2| \leq (q^{-1} + \widetilde{\eta})^2 (q^2 - \delta) \leq 1$ and $|u| = 1/q + \widetilde{\eta} < q^{-\frac{1}{2}}$. This shows taking any $0 < \widetilde{\eta} < \min\{1/\sqrt{q^2 - \delta} - 1/q, 1/\sqrt{q} - 1/q\}$ suffices. We then plan to use Lemma 2.2 to evaluate the integral in (B.2) with, $\widetilde{\eta} = \min\{1/\sqrt{q^2 - \delta} - 1/q, 1/\sqrt{q} - 1/q\}/2$ for $|z| \leq q^2 - \delta$ just as we did in the proof of Proposition 1.3.

Remark B.1. $F_z(u)$ was absolutely convergent in \mathcal{R} which gave us the appropriate choice for η . All the calculations are identical here with \widetilde{F}_z instead of F_z . The main difference is, here we have $|z| \leq q^2 - \delta$ instead of $|z| \leq q - \epsilon$. Earlier $F_z\left(\frac{1}{q}\right)$ had a pole at $z = q, q^2, \dots$ but we got rid of the $z = q$ pole here. This is what essentially allows us to take $|z| \leq q^2 - \delta$.

We therefore get

$$\begin{aligned} \widetilde{M}_z(n) &= \frac{1}{2\pi i} \int_{\{|u|=\frac{1}{q}+\widetilde{\eta}\} \cup \mathcal{H}'} \zeta(u)^z \widetilde{F}_z(u) \frac{du}{u^{n+1}} \\ &= \widetilde{F}_z\left(\frac{1}{q}\right) q^n n^{z-1} \left(\frac{1}{\Gamma(z)} + O_\delta\left(\frac{1}{n}\right) \right) \\ &= q^n n^{z-1} \left(zh(z) + O_\delta\left(\frac{1}{n}\right) \right), \end{aligned}$$

where

$$h(z) = \frac{1}{z} \tilde{F}_z \left(\frac{1}{q} \right) \frac{1}{\Gamma(z)} = \frac{1}{\Gamma(z+1)} \left(1 - \frac{1}{q} \right)^{qz} \prod_{p \in \mathcal{P}_{\geq 2}} \left(1 - \frac{z}{q^{\deg(p)}} \right)^{-1} \left(1 - \frac{1}{q^{\deg(p)}} \right)^z.$$

Also we have $h(z) = \frac{1}{\Gamma(z+1)} \tilde{F}_z \left(\frac{1}{q} \right)$. Since, $\tilde{F}_z \left(\frac{1}{q} \right)$ is analytic in $|z| < q^2$, so is h . \square

B.2. An useful upper bound for $N'(n, j)$.

Proof of Lemma 4.2. We use Proposition 1.4 with $z = q + \eta$ and if $a_j(f)$ denotes the indicator function of polynomials in \tilde{M}_n with $\Omega(f) = j$ then, we see that every time $a_j = 1$ occurs, it contributes an amount of $(q + \eta)^j$ on the left hand side of the sum of Proposition 1.4 and all the other terms are positive. We have by the previous lemma $h(z)$ is analytic in an open set containing the disc $|z| \leq q^2 - \delta$. Since the disk is compact and h is continuous there, we have $h(z) \ll_{q, \delta} 1$ for every $|z| \leq q^2 - \delta$. Consequently for every $|z| \leq q^2 - \delta$, $|zh(z)| \leq |(q + \eta)h(q + \eta)| \ll_{q, \delta} 1$. This observation leads to an upper bound $N'(n, j) \ll_{q, \delta} q^n n^{q+\eta-1} \left(\frac{1}{q+\eta} \right)^j$. \square

B.3. An uniform estimate for $N'(n, j)$ when $j \leq eq \log n$.

Proof of Lemma 4.5. We use Proposition 1.4. The coefficient of z^j in $\tilde{M}_z(n)$ is what we want to evaluate. We have for $r < q^2$,

$$N'(n, j) = \frac{1}{2\pi i} \int_{|z|=r} \tilde{M}_z(n) \frac{dz}{z^{j+1}}.$$

We recall from Proposition 1.4 $\tilde{M}_z(n) = \frac{q^n}{n} \left\{ zh(z)n^z + O(n^{z-1}) \right\}$. Plugging this back in the integral we get that

$$\frac{n}{q^n} N'(n, j) = \frac{1}{2\pi i} \int_{|z|=r} zh(z)n^z \frac{dz}{z^{j+1}} + O(E_0),$$

where E_0 is given by, $\int_{|z|=r} \left| \frac{n^{z-1}}{z^{j+1}} \right| |dz|$. At this point we choose $r = \frac{j}{\log n} \leq eq < q^2$ since by hypothesis $j \leq eq \log n$ and $q \geq 3$.

Noting $e^j = n^r \iff j = r \log n$ and using Lemma 3.2, we get the term E_0 is bounded by

$$\ll \frac{r^{-j-1} e^j}{n} \int_{|z|=r} |n^{z-r}| |dz| \ll \frac{r^{-j-1} e^j}{n} \frac{r}{j^{\frac{1}{2}}} \ll \frac{1}{n} \frac{(e \log n)^j}{j^{j+\frac{1}{2}}} \ll \frac{1}{n} \frac{(\log n)^j}{j!},$$

where the final step of bounding follows by Stirling. This is the same trick that we used while bounding above E on page 10 in the proof of Theorem 1.1. We now return to the main term

$$M_0 = \frac{1}{2\pi i} \int_{|z|=r} zh(z)n^z \frac{dz}{z^{j+1}} = \frac{1}{2\pi i} \int_{|z|=r} h(z)n^z \frac{dz}{z^j},$$

where $h(z)$ is an analytic function in the region $|z| < q^2$, so the integrand has a pole at $z = 0$ of order j . We evaluate this integral using residue theorem and obtain

$$M_0 = \lim_{z \rightarrow 0} \frac{1}{(j-1)!} \frac{d^{j-1}}{dz^{j-1}} h(z)n^z.$$

Using binomial theorem for higher order differentiation of products of two complex functions

$$\begin{aligned} \frac{1}{(j-1)!} \frac{d^{j-1}}{dz^{j-1}} h(z) n^z \Big|_{z=0} &= \frac{1}{(j-1)!} \sum_{m+\ell=j-1} \frac{(j-1)!}{m!\ell!} h^{(m)}(z) (n^z)^\ell \Big|_{z=0} \\ &= \sum_{m+\ell=j-1} \frac{1}{m!\ell!} h^{(m)}(0) (\log n)^\ell. \end{aligned}$$

where we have taken the limits inside the argument of the higher order derivatives which is allowed because all the functions here are analytic and hence infinitely differentiable. Plugging everything back we obtain the desired result. \square

B.4. A technical upper bound for $h(z)$, $Q_j(X)$ and their derivatives.

Proof of Lemma 4.6. From Proposition 1.4 we have that $h(z)$ is holomorphic for $|z| < q^2$ which in particular implies $h(z) \ll_q 1$ for $|z| \leq 2.9q$. Therefore we have by Cauchy's integral formula,

$$h^{(m)}(0)/m! = \left| \int_{|z|=2.9q} h(z) \frac{dz}{z^{m+1}} \right| \ll \left(\frac{1}{2.9q} \right)^m \quad (2.9q < q^2 \text{ whenever } q \geq 3).$$

We have from Lemma 4.4

$$Q_j(X) = \sum_{m+\ell=j-1} \frac{1}{m!\ell!} h^{(m)}(0) X^\ell \ll \sum_{m+\ell=j-1} \left(\frac{1}{2.9q} \right)^m \frac{X^\ell}{\ell!}$$

We now take out the top term $\frac{X^{j-1}}{(j-1)!}$ to bound the sum from above and observe for each $m \geq 1$, $\frac{X^\ell}{\ell!} = \frac{X^{j-m-1}}{(j-m-1)!} \leq (eq)^m \frac{X^{j-1}}{(j-1)!}$ which is true after cross multiplying whenever $j \leq eqX$ since $(j-1)(j-2)\cdots(j-m) \leq j^m \leq (eqX)^m$. Hence we arrive at the following upper bound,

$$Q_j(X) \ll \frac{X^{j-1}}{(j-1)!} \sum_{0 \leq m \leq j-1} \left(\frac{eq}{2.9q} \right)^m \ll \frac{X^{j-1}}{(j-1)!}.$$

Likewise when $j \leq eqX$,

$$\begin{aligned} Q'_j(X) &= \sum_{m+\ell=j-1} \frac{1}{m!(\ell-1)!} h^{(m)}(0) X^{\ell-1} \\ &\ll \sum_{m+\ell=j-1} \left(\frac{1}{2.9q} \right)^m \frac{X^{\ell-1}}{(\ell-1)!} \\ &\ll \frac{X^{j-2}}{(j-2)!} \sum_{0 \leq m \leq j-1} \left(\frac{eq}{2.9q} \right)^m \\ &\ll \frac{X^{j-2}}{(j-2)!}. \end{aligned}$$

\square

B.5. One of the main ingredients of Proposition 4.7 and Theorem 1.2.

Proof of Lemma 4.4. By Lemma 4.5 we have that

$$N'(n+j-k, j) = \frac{q^{n+j-k}}{n+j-k} \left\{ Q_j(\log(n+j-k)) + O\left(\frac{(\log(n+j-k))^j}{j!(n+j-k)}\right) \right\}.$$

We observe $n-k \geq n(1 - \frac{1}{B}) \rightarrow \infty$ as $n \rightarrow \infty$. We also note if $E_1 = \frac{(\log(n-k))^{j+1}}{j!(n-k)^2}$ and $E_2 = \frac{(\log(n+j-k))^j}{j!(n+j-k)^2}$ then as $n \rightarrow \infty$

$$\begin{aligned} \frac{E_2}{E_1} &= \frac{1}{(\log(n-k))} \left(\frac{\log(n+j-k)}{\log(n-k)} \right)^j \frac{(n-k)^2}{(n+j-k)^2} \\ &\leq \left(1 + \frac{\log\left(1 + \frac{j}{n-k}\right)}{\log(n-k)} \right)^j \frac{1}{(\log(n-k))} \rightarrow 0 \\ &\implies E_2 = o(E_1), \end{aligned}$$

where we have at the second step the term inside the bracket is bounded, because $\frac{j}{n-k} \leq \frac{eq \log(n-k)}{(n-k)} \rightarrow 0$ and hence for n large enough it is bounded by $(1 + \log 2 / \log(n-k))^{eq \log(n-k)} \leq e^{eq \log 2} = 2^{eq}$.

We show at a small cost it is possible to replace $\frac{Q_j(\log(n+j-k))}{n+j-k}$ by $\frac{Q_j(\log(n-k))}{n-k}$. We use mean value theorem to study the error.

Let $F(x) := \frac{Q_j(\log x)}{x}$ which is a regular function if $x > 0$. The difference caused by this replacement

$$D = \frac{Q_j(\log(n+j-k))}{n+j-k} - \frac{Q_j(\log(n-k))}{n-k} = F(n+j-k) - F(n-k).$$

By mean value theorem one can pick $y \in (n-k, n+j-k)$ such that

$$D = jF'(y) = j \frac{Q'_j(\log y) - Q_j(\log y)}{y^2} = \frac{j}{y^2} Q'_j(\log y) - \frac{j}{y^2} Q_j(\log y) = F_1 + F_2.$$

We apply Lemma 4.6 with the choice $X = \log y$. We note from the hypothesis that $j \leq eqY \leq eq \log y$ and $y > n-k$, so that we must have $X = \log y \rightarrow \infty$ with n . Using this observation, we upper bound the first term of F_1 which gives $F_1 = \frac{j}{y^2} Q'_j(\log y) \ll \frac{j}{y^2} \frac{(\log(n+j-k))^{j-2}}{(j-2)!}$. Thus we obtain

$$\begin{aligned} \frac{F_1}{E_1} &= \frac{(n-k)^2 j!}{(\log(n-k))^{j+1}} \frac{j}{y^2} \frac{(\log(n+j-k))^{j-2}}{(j-2)!} \\ &\leq \left(\frac{\log(n+j-k)}{\log(n-k)} \right)^{j-2} \frac{j^2(j-1)}{(\log(n-k))^3} \\ &\ll 1, \end{aligned}$$

since $j \leq eq \log(n-k)$ and $y > n-k$. Hence $F_1 = O(E_1)$ and likewise from Lemma 4.6 one can show $|F_2| = \frac{j}{y^2} Q_j(\log y) \ll \frac{j(\log(n+j-k))^{j-1}}{y^2(j-1)!}$.

Thus we infer

$$\begin{aligned} \frac{|F_2|}{E_1} &= \frac{j!(n-k)^2}{(\log(n-k))^{j+1}} \frac{j(\log(n+j-k))^{j-1}}{y^2(j-1)!} \\ &\leq \left(\frac{\log(n+j-k)}{\log(n-k)} \right)^{j-1} \frac{j^2}{(\log(n-k))^2} \\ &\ll 1, \end{aligned}$$

which completes the proof of Lemma 4.4. □

REFERENCES

- [1] J. L. Nicolas, *Sur la distribution des nombres entiers ayant une quantité fixée de facteurs premiers*, Acta Arithmetica, 44, 191–200 (1984).
- [2] E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, Chealsea Publishing Company (1953).
- [3] S. Ramanujan, *Collected Papers*, Chealsea Publishing Company (1953).
- [4] L.G. Sathe, *On a problem of Hardy on the distribution of integers having a given number of prime factors*, J. Indian Math Soc. 17 (1953), p. 63–141; 18 (1954); p. 27–81.
- [5] A Selberg, *Note on a paper by L.G. Sathe*, *ibid.* 18, J. Indian Math Soc., p. 83–87 (1954).
- [6] S. Porrit, *Character sums over products of prime polynomials*, [arXiv:2003.12002](https://arxiv.org/abs/2003.12002). (2020).
- [7] P.D.T.A. Elliot, *Probabilistic Number Theory I et II*, Springer Verlag , Grundlehren der mathematischen Wissenschaften (1980).
- [8] M. Rosen, *Number Theory in Function Fields*, Springer-Verlag (2002).
- [9] G. Tenenbaum, *An Introduction to Analytic and Probabilistic Number Theory*, Cambridge University Press (2015).

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6192 SUCC. CENTRE-VILLE, MONTRÉAL, QC H3C 3J7, CANADA

Email address: arghya.datta@umontreal.ca