

## NOVIKOV DIALGEBRAS AND PERM ALGEBRAS

F. MASHUROV, B. K. SARTAYEV

ABSTRACT. In this paper, we consider Perm algebra with the derivation  $d$ . The algebra itself is equipped with the new operation  $a \succ b = d(a)b$ . We construct a linear basis of the free Novikov dialgebra in terms of new operations. Also, we prove that the class of algebras under the new operation form a variety. Finally, we find the defining identities of the variety.

## 1. INTRODUCTION

An associative algebra with the left-commutative identity is called perm algebra [3]. In [11] subalgebras of the perm algebra under the commutator and the anti-commutator are considered. It turned out that every metabelian Lie algebra can be embedded into perm algebra under the commutator. In [8] was considered a subalgebra of perm algebra with derivation under the operation  $x_1 \prec x_2 = x_1 d(x_2)$ . We denote the perm algebra with derivation and its subalgebra under the operation  $\prec$  by  $\text{DerPerm}\langle X \rangle$  and  $\text{DerPerm}_\prec\langle X \rangle$ , respectively. It turn out that  $\text{DerPerm}_\prec\langle X \rangle$  is a pre-Lie algebra with additional identities of the following form:

$$((x_1 \prec x_2) \prec x_3) \prec x_4 = ((x_1 \prec x_3) \prec x_2) \prec x_4, \quad (1.1)$$

$$(x_1, x_2 \prec x_3, x_4) = (x_2, x_1 \prec x_3, x_4). \quad (1.2)$$

where  $(x_1, x_2, x_3)$  stands for the associator  $(x_1 \prec x_2) \prec x_3 - x_1 \prec (x_2 \prec x_3)$ . Moreover, there is proved that every Novikov dialgebra can be embedded into perm algebra with derivation as follows:

$$x_1 \vdash x_2 = x_1 d(x_2), \quad x_1 \dashv x_2 = d(x_2)x_1.$$

In this paper, we consider a new operation  $\succ$  on  $\text{DerPerm}\langle X \rangle$  algebra which is defined as follows:

$$x_1 \succ x_2 = d(x_1)x_2.$$

We denote that algebra by  $\text{DerPerm}_\succ\langle X \rangle$ . In our case the operations  $\vdash$  and  $\dashv$  correspond to the operations  $\prec$  and  $\succ$ , respectively. The main aim of the paper is to describe the subalgebra of  $\text{DerPerm}\langle X \rangle$  under the operation  $\succ$ .

---

*Key words and phrases.* Free algebras, Perm algebras, polynomial identities, Novikov dialgebras.

This work was supported by the Program of MES RK.

Let us call that an algebra  $A$  is special if it can be embedded into some algebra from fixed class of algebras, otherwise, it is called exceptional. It was proved that there exists pre-Lie algebra with additional identities (1.1) and (1.2) that is exceptional i.e. can not be embedded into the class of DerPerm algebras. For the case of the class of DerPerm $\succ$  algebras, we have the opposite result. In addition, it turns out that the class of DerPerm $\succ$  algebras forms a variety. Moreover, we describe the defining identities of that variety and construct a monomial basis of free DerPerm $\succ$  algebra. The analogical result can be traced in the class of subalgebras of commutative and Poisson algebras with derivation [2], [9].

The main motivation of this work comes from the theory of operads. It is well-known that for the fixed binary quadratic operad denoted by Var, we can define the Manin white product with operad Nov governed by the variety of Novikov algebras and obtain the operad DerVar [7]. That operad is obtained from operad Var under the operations  $\succ$  and  $\prec$  which are defined as follows:

$$x_1 \succ x_2 = d(x_1)x_2, \quad x_1 \prec x_2 = x_1d(x_2).$$

In the other words, we have

$$\text{Var} \circ \text{Nov} = \text{DerVar}.$$

On the other hand [10], [13],

$$\text{Perm} \circ \text{Var} = \text{di-Var}.$$

In this paper and [8], a special case is considered. That is

$$\text{Perm} \circ \text{Nov} = \text{di-Nov} = \text{DerPerm}.$$

Let us now pay close attention to Novikov dialgebra. Every dialgebra satisfies to the *0-identities*:

$$\begin{aligned} (x_1 \dashv x_2) \vdash x_3 &= (x_1 \vdash x_2) \dashv x_3, \\ x_1 \dashv (x_2 \vdash x_3) &= x_1 \dashv (x_2 \dashv x_3). \end{aligned} \tag{1.3}$$

**Definition 1.** The Novikov dialgebra is defined by (1.3) together with

$$\begin{aligned} (x_1 \vdash x_2) \vdash x_3 - x_1 \vdash (x_2 \vdash x_3) &= (x_2 \vdash x_1) \vdash x_3 - x_2 \vdash (x_1 \vdash x_3), \\ (x_1 \dashv x_2) \dashv x_3 &= (x_1 \dashv x_3) \dashv x_2, \end{aligned} \tag{1.4}$$

$$\begin{aligned} (x_1 \dashv x_2) \dashv x_3 - x_1 \dashv (x_2 \dashv x_3) &= (x_2 \vdash x_1) \dashv x_3 - x_2 \vdash (x_1 \dashv x_3), \\ (x_1 \vdash x_2) \dashv x_3 &= (x_1 \vdash x_3) \vdash x_2. \end{aligned} \tag{1.5}$$

We see that the Novikov dialgebra is left-symmetric and right-commutative under the operations  $\vdash$  and  $\dashv$ , respectively. Since a free basis for these algebras has been constructed (for example, see [12]), it is not difficult to construct a basis of the free algebra with identities (1.4). At times it is difficult to solve the same problem for algebras with complementary identities that contains both

multiplications as (1.3) and (1.5). For the free Novikov dialgebras, we solve this problem in terms of  $\succ$  and  $\prec$ .

## 2. MAIN RESULTS

Firstly, let us recall a free basis of  $\text{Perm}\langle X \rangle$  algebra. That is monomials of the following form:

$$x_{i_1} \cdots x_{i_{n-1}} x_{i_n}, \quad (2.1)$$

where  $i_1 \leq \dots \leq i_{n-1}$ . Since the generators of  $\text{DerPerm}\langle X \rangle$  are

$$X^{(\omega)} = X \cup X' \cup X'' \cdots \cup X^{(n)} \cup \dots,$$

every monomial  $a \in \text{DerPerm}\langle X \rangle$  can be written as follows:

$$a = x_{i_1}^{(k_1)} \cdots x_{i_{n-1}}^{(k_{n-1})} x_{i_n}^{(k_n)}. \quad (2.2)$$

To state the main results let us recall the crucial definition that used in [5], [7]:

**Definition 2.** The weight function is defined on monomials of  $\text{DerPerm}\langle X \rangle$  as follows:

$$\begin{aligned} \text{wt} : \text{DerPerm}\langle X \rangle &\rightarrow \mathbb{Z}, \\ \text{wt}(x^{(j)}) &= j - 1, \quad x \in X, \quad j \geq 0, \\ \text{wt}(uv) &= \text{wt}(u) + \text{wt}(v) \end{aligned}$$

for all monomials  $u, v \in \text{DerPerm}\langle X \rangle$ .

**Theorem 1.** *Let  $a$  be a monomial of  $\text{DerPerm}\langle X \rangle$  of the form (2.2). A monomial  $a \in \text{DerPerm}_{\succ}\langle X \rangle$  if and only if  $\text{wt}(a) = -1$  and  $k_n = 0$ .*

Using Theorem 1, we obtain the dimension of operad  $\text{DerPerm}_{\succ}$ :

**Corollary 1.**  $\dim(\text{DerPerm}_{\succ}(n)) = n \binom{2n-3}{n-1}$ .

**Theorem 2.** *Every element of the free Novikov dialgebra can be written in a unique way as a sum of elements  $\text{DerPerm}_{\succ}\langle X \rangle$  and  $\text{DerPerm}_{\prec}\langle X \rangle$ , that is,*

$$\text{di-Nov}\langle X \rangle = \text{DerPerm}_{\succ}\langle X \rangle \oplus \text{DerPerm}_{\prec}\langle X \rangle.$$

as a vector space.

**Theorem 3.** *Every homomorphic image of  $\text{DerPerm}_{\succ}\langle X \rangle$  is special.*

Using Theorem 3, we obtain the following result:

**Corollary 2.** *The class of  $\text{DerPerm}_{\succ}\langle X \rangle$  algebras form a variety.*

Since the class of  $\text{DerPerm}_{\succ}\langle X \rangle$  algebras is a variety and thus defined by some set of identities, the interesting problem is to determine a list of defining identities of  $\text{DerPerm}_{\succ}\langle X \rangle$ . The next result answers that question:

**Theorem 4.** *The defining identities of the variety  $\text{DerPerm}_\succ$  are*

$$x_1 \succ (x_2 \succ x_3) = x_2 \succ (x_1 \succ x_3), \quad (2.3)$$

$$\langle x_1, x_2 \succ x_3, x_4 \rangle - \langle x_1, x_3, x_2 \succ x_4 \rangle = 0, \quad (2.4)$$

$$\langle x_1, x_2, x_3 \rangle \succ x_4 - \langle x_1, x_3, x_2 \rangle \succ x_4 = 0, \quad (2.5)$$

where  $\langle a, b, c \rangle = (a \succ b) \succ c - a \succ (b \succ c)$ .

### 3. THE PROOF OF THEOREM 1 AND THEOREM 2

*Proof.* It is clear that every monomial of  $\text{DerPerm}_\succ \langle X \rangle$  can be written as a sum of monomials of the form (2.2), where  $\text{wt}(a) = -1$  and  $k_n = 0$ .

It remains to show that the monomial of the form (2.2) with conditions  $\text{wt}(a) = -1$  and  $k_n = 0$  can be written in terms of  $\succ$ . Let us prove it by induction on the length of monomial (2.2). The base of induction is  $n = 2$  and we have  $x'_1 x_2 = x_1 \succ x_2$ .

For the monomial of the length  $n$ , we consider two cases depending on the appearance of the derivation degree 1 in  $(k_1), \dots, (k_{n-1})$ . If there is  $r$  such that  $k_r = 1$ , then we have

$$x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} x'_{i_r} x_{i_{r+1}}^{(k_{r+1})} \dots x_{i_{n-1}}^{(k_{n-1})} x_{i_n}.$$

By using left-commutative identity and definition of product  $\succ$  we have

$$x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} x'_{i_r} x_{i_{r+1}}^{(k_{r+1})} \dots x_{i_{n-1}}^{(k_{n-1})} x_{i_n} = x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} x_{i_{r+1}}^{(k_{r+1})} \dots x_{i_{n-1}}^{(k_{n-1})} (x_{i_r} \succ x_{i_n}).$$

Since,  $(x_{i_r} \succ x_{i_n}) \in \text{DerPerm}_\succ \langle X \rangle$  the remaining part can be rewritten by inductive hypothesis.

Now, consider the second case such that there is no generator with derivation degree 1. That is,

$$x_{i_1}^{(k_1)} \dots x_{i_{n-1}}^{(k_{n-1})} x_{i_n},$$

where  $k_1, \dots, k_{n-1} \neq 1$  and  $k_1 + \dots + k_{n-1} = n - 1$ . Then we may assume that

$$x_{i_1}^{(k_1)} \dots x_{i_{n-1}}^{(k_{n-1})} x_{i_n} = x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} x_{i_r}^{(k_r)} x_{i_{r+1}} \dots x_{i_{n-1}} x_{i_n},$$

where  $k_1, \dots, k_{r-1} \neq 1$  and  $k_1 + \dots + k_{r-1} = n - 1$ . We choose any generator  $x_{i_j}$  with minimal derivation degree greater than 1. Without loss of generality, we assume that the generator  $x_{i_r}$  has minimal derivation degree, that is  $k_r = \min(\{k_1, k_2, \dots, k_r\})$ . Then we rewrite it as follows:

$$\begin{aligned} & x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} x_{i_r}^{(k_r)} x_{i_{r+1}} \dots x_{i_{n-1}} x_{i_n} = \\ & x_{i_1}^{(k_1)} \dots x_{i_{r-1}}^{(k_{r-1})} ((\dots (x_{i_r} \succ x_{i_{r+1}}) \succ \dots) \succ x_{j_{r+k_r}}) x_{j_{r+k_r+1}} \dots x_{i_n} \\ & - \sum_{t, t_1, \dots, t_{k_r} < k_r} x_{i_1}^{(k_1)} \dots x_{i_{n-1}}^{(k_{n-1})} x_{i_r}^{(t)} x_{j_{r+1}}^{(t_1)} \dots x_{j_{r+k_r}}^{(t_{k_r})} x_{j_{r+k_r+1}} \dots x_{i_n}. \end{aligned}$$

The monomial

$$x_{i_1}^{(k_1)} \cdots x_{i_{r-1}}^{(k_{r-1})} ((\cdots (x_{i_r} \succ x_{i_{r+1}}) \succ \cdots) \succ x_{j_{r+k_r}}) x_{j_{r+k_r+1}} \cdots x_{i_n}$$

can be written in terms of  $\succ$  by inductive hypothesis. For the monomials of the form

$$x_{i_1}^{(k_1)} \cdots x_{i_{n-1}}^{(k_{n-1})} x_{i_r}^{(t)} x_{j_{r+1}}^{(t_1)} \cdots x_{j_{r+k_r}}^{(t_{k_r})} x_{j_{r+k_r+1}} \cdots x_{i_n},$$

where  $t, t_1, \dots, t_{k_r} < k_r$ , we have if  $t, t_1, \dots, t_{k_r} > 1$ , we use the same way which is given above several times and finally obtain the generator with derivation degree 1 which correspond to the first case.  $\square$

Let us recall some facts about Novikov dialgebra(see [8]):

- Every Novikov dialgebra can be embedded into perm algebra with derivation relative to the operations

$$x_1 \vdash x_2 = x_1 d(x_2), \quad x_1 \dashv x_2 = d(x_2) x_1.$$

- If  $a \in \text{DerPerm}\langle X \rangle$  of the form (2.2) then  $a$  can be written in terms of  $\vdash$  and  $\dashv$  if and only if  $\text{wt}(a) = -1$ .
- If  $a \in \text{DerPerm}\langle X \rangle$  of the form (2.2) then  $a$  can be written in terms of  $\prec$  if and only if  $\text{wt}(a) = -1$  and  $k_n \neq 0$ .

Now, we are ready to prove the Theorem 2.

*Proof.* Firstly, we equalize the operations of Novikov dialgebra as follows:

$$x_1 \succ x_2 = x_2 \dashv x_1, \quad x_1 \prec x_2 = x_1 \vdash x_2.$$

We denote by  $P_{(-1)}$  the subspace of monomials  $\text{DerPerm}\langle X \rangle$  of the weight  $-1$ . Also, we denote by  $P_{(-1, k_n=0)}$  and  $P_{(-1, k_n \neq 0)}$  the subspaces of monomials  $P_{(-1)}$  of the form (2.2) with  $k_n = 0$  and  $k_n \neq 0$ , respectively.

By the facts that the free di-Nov $\langle X \rangle$  isomorphic to  $\text{DerPerm}\langle X \rangle$  and a linear basis of di-Nov $\langle X \rangle$  in  $\text{DerPerm}\langle X \rangle$  are monomials of the weight  $-1$ , we can represent a linear basis of di-Nov $\langle X \rangle$  as follows:

$$P_{(-1)} = P_{(-1, k_n=0)} \oplus P_{(-1, k_n \neq 0)}.$$

By Theorem 1 and the third fact, all monomials of subspaces  $P_{(-1, k_n=0)}$  and  $P_{(-1, k_n \neq 0)}$  can be written in terms of  $\succ$  and  $\prec$  respectively, i.e. in terms of  $\dashv$  and  $\vdash$ .  $\square$

#### 4. PROOF OF THEOREM 3

*Proof.* Let us prove it by Cohn's criterion on homomorphic images of special algebras [4]. Suppose that  $\alpha$  is an ideal of  $\text{DerPerm}_\succ\langle X \rangle$  and  $\{\alpha\}$  is an ideal of  $\text{DerPerm}\langle X \rangle$  generated by the set  $\alpha$ . It is enough to prove that  $\text{DerPerm}_\succ\langle X \rangle \cap \{\alpha\} \subseteq \alpha$ .

Assume that  $g_i$  ( $i \in I$ ) are generators of the ideal  $\alpha$ . Let  $w$  be a non-zero element of  $\text{DerPerm}_{\succ}\langle X \rangle \cap \{\alpha\}$ . Then

$$w = \sum_k \mu_k w_k, \quad (\mu_k \in \mathbb{K})$$

is a linear combination of monomials  $w_k$  in  $X^{(w)}$  and  $g_i$  ( $i \in I$ ) such that each monomial is linear by at least one generator of  $\alpha$ .

Let  $w_k = a_1 \cdots a_n$  be a term of  $w$  in the linear combination. Assume that there is  $r$  such that  $a_r = g_i^{(k_r)}$ , where  $g_i$  is a generator of the ideal  $\alpha$ . Then express  $a_1, \dots, a_{r-1}, a_{r+1}, \dots, a_n$  in terms of  $x_{i_j}^{(i_j)} \in X^{(w)}$ , therefore we can assume that  $a_1, \dots, a_{r-1}, a_{r+1}, \dots, a_n \in X^{(w)}$ .

Since  $w \in \text{DerPerm}_{\succ}\langle X \rangle$ , by Theorem 1 we have  $\text{wt}(w_k) = -1$  and  $a_n \in X$ . That is

$$w = \sum_k \mu_k w_k, \quad (\mu_k \in \mathbb{K})$$

where

$$w_k = x_{i_1}^{(k_1)} \cdots g_i^{(k_r)} \cdots x_{i_{n-1}}^{(k_{n-1})} x_{i_n},$$

where  $g_i$  is a generator of the ideal  $\alpha$ . But by Theorem 1 we can express each  $w_k$  as linear combination of monomials in terms of product  $\succ$  with generators  $x_{i_1}, \dots, x_{i_n}$  and  $g_i$  ( $i \in I$ ), where  $g_i$  appears in each monomial at least once. Therefore, every  $w_k \in \alpha$ , and we have  $w \in \alpha$ . □

## 5. A LINEAR BASIS OF THE FREE $\text{DerPerm}_{\succ}$ ALGEBRA

A monomial basis of the free  $\text{DerPerm}_{\prec}$  algebra was constructed in [8]. By Theorem 2, to construct a free basis of Novikov dialgebra it is enough to solve the same problem for  $\text{DerPerm}_{\succ}$  algebra. In this section, we construct a monomial basis of the free  $\text{DerPerm}_{\succ}$  algebra. Let us define an order on generators of  $\text{DerPerm}\langle X \rangle$  as follows:

$$x_i^{(r_m)} > x_j^{(r_n)}$$

if  $r_m > r_n$ , or  $r_m = r_n$  and  $i > j$ .

Let us define a normal form for monomials of the form (2.2) of weight  $-1$  and  $k_n = 0$  as follows:

$$x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}, \quad (5.1)$$

where

$$\begin{aligned} r_1, \dots, r_n &\neq 1, & x_{i_1} &\geq \dots \geq x_{i_2}, \\ x_{j_n}^{(r_n)} &\geq \dots \geq x_{j_1}^{(r_1)}, & k_m &\geq \dots \geq k_1. \end{aligned}$$

Denote by  $P(X)$  the set of all monomials of normal forms (2.2) and denote by  $\text{Magma}\langle X \rangle$  the free magma algebra with binary operation  $\succ$  generated by  $X$ .

We define a mapping  $\phi$  inductively on monomials of the form (5.1) as follows:

$$\phi : \text{DerPerm}\langle X \rangle \rightarrow \text{Magma}\langle X \rangle,$$

$$\phi(x_{i_n}^{(n-1)} x_{i_{n-1}} \cdots x_{i_2} x_{i_1}) = ((\cdots (x_{i_n} \succ x_{i_{n-1}}) \succ \cdots) \succ x_{i_2}) \succ x_{i_1},$$

where  $x_{i_{n-1}} \geq \cdots \geq x_{i_2}$ .

$$\begin{aligned} \phi(x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}) = \\ \phi(x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_{n-1}}^{(r_{n-1})} P_1 x_{i_{1-r_n}} \cdots x_{i_2} x_{i_1}), \end{aligned}$$

where  $P_1 = \phi(x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_{1+r_n}})$ . We may assume that  $P_1$  is a new generator i.e.  $X_1 = X \cup \{P_1\}$  such that  $\text{wt}(P_1) = -1$  and  $P_1 \geq x_{i_{1-r_n}} \geq \cdots \geq x_1$ . Calculating by the given rule, finally, we get

$$\begin{aligned} \phi(x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}) = \cdots \\ = x_{k_1} \succ (\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots). \end{aligned} \quad (5.2)$$

We denote by  $[a]$  the image  $\phi(a)$ . Define a map  $\tau : \text{Magma}\langle X \rangle \rightarrow \text{DerPerm}\langle X \rangle$  by the formula  $\tau(x) = x$ ,  $x \in X$ , and  $\tau(a \succ b) = a'b$ . For example, if  $x, y, z \in X$ , then  $\tau((x \succ y) \succ z) = (x'y)'z = x''yz + x'y'z$ .

Let us define a sequence on a monomial of the set  $P(X)$  of the form (5.1) as follows:

$$S(a) = (\underbrace{1, \dots, 1}_m, r_1, \dots, r_{n-1}, r_n),$$

i.e. that is a sequence of positive numbers in a non-decreasing order. Define an order on that sequences lexicographically. Define an order on monomials of the set  $P(X)$  as follows:

$$a > b \quad \text{if} \quad S(a) > S(b)$$

and call that order by *der-lex*.

**Lemma 1.** *Let  $a = x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}$  be from  $P(X)$ . Then  $\tau([a]) = a + \sum_j b_j$ , where  $a > b_j$  for all  $j$ .*

*Proof.* By definition of the mapping  $\tau$ , it is enough to prove it for

$$a = x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}.$$

By Leibniz rule for derivation, we have

$$\begin{aligned} \tau([x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_l} \cdots x_{i_2} x_{i_1}]) &= \tau((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1}) = \\ &= x_{j_1}^{(r_1)} \tau(P_{n-1}) x_{i_l} \cdots x_{i_2} x_{i_1} + \sum_{p < r_1} x_{j_1}^{(p)} \tau(P_{n-1}) \cdots x_{i_1}. \end{aligned}$$

By der-lex order,  $a$  is a bigger than all monomials of  $\sum_{p < r_1} x_{j_1}^{(p)} \tau(P_{n-1}) \cdots x_{i_1}$ . In the same way, applying Leibniz rule to  $\tau(P_{n-1})$  we obtain the same result as above. Finally, we obtain only one monomial which is not less than  $a$ . That is  $a$  itself.  $\square$

Under the map  $\phi$  for each monomial of  $P(X)$  correspond a unique monomial of magma algebra. Let us denote by  $[P(X)]$  the set of all such monomials obtained under the mapping  $\phi$ . Also, we define a linear mapping  $\tau$  as above for the following algebras:

$$\tau : \text{DerPerm}_{\succ} \rightarrow \text{DerPerm}.$$

**Theorem 5.** *The set  $[P(X)]$  is a linear basis of  $\text{DerPerm}_{\succ}\langle X \rangle$ .*

*Proof.* It is enough to show that for every weight-homogeneous differential polynomial  $a \in \text{DerPerm}\langle X \rangle$  of the form (5.1) there exists  $[p_1], \dots, [p_r] \in [P(X)]$  such that

$$a = \sum_{i=1}^r \lambda_i \tau([p_i]), \quad \lambda_i \in \mathbb{k}. \quad (5.3)$$

Indeed, assume (5.3) holds and  $w \in \text{DerPerm}_{\succ}\langle X \rangle$  is a homogeneous element of degree  $n$ . Then  $a = \tau(w)$  is a weight-homogeneous differential polynomial which can be presented by (5.3). Since  $\tau$  is injective, we obtain

$$w = \sum_{i=1}^r \lambda_i [v_i], \quad [v_i] \in [P_n(X)].$$

Hence, the set  $[P_n(X)]$  is linearly complete in the space  $F_n$  of degree  $n$  elements in  $\text{DerPerm}_{\succ}\langle X \rangle$ . Thus this is a basis since  $|P_n(X)| = |[P_n(X)]| = \dim F_n$ .

The statement (5.3) is enough to prove for monomials only. Let us prove it by induction on the degree of  $a$ . The base of induction is  $n = 2$ . Suppose that the statement holds for the degree up to  $n - 1$ . Let us prove that statement for monomial of degree  $n$ .

Case 1: if

$$x'_{k_1} \cdots x'_{k_m} x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_l} \cdots x_{i_2} x_{i_1},,$$

such that  $m \neq 0$ , then by inductive hypothesis for the subword of  $a$ , we have

$$x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_l} \cdots x_{i_2} x_{i_1} = \sum_i \lambda_i \tau([p_i])$$



and

$$a = \sum_i \lambda_i \tau(x_{k_1} \succ (\dots \succ (x_{k_m} \succ ([p_i]))) \dots),$$

where  $x_{k_1} \succ (\dots \succ (x_{k_m} \succ ([p_i]))) \dots \in [P(X)]$ .

Case 2: It is enough to consider only monomials of the form

$$a = x_{j_1}^{(r_1)} \cdots x_{j_n}^{(r_n)} x_{i_1} \cdots x_{i_2} x_{i_1}, \quad (5.4)$$

such that  $r_t \neq 1$  for all  $t$ .

For  $a$ , we have the following equality:

$$a = \tau([a]) - \sum_i b_i \quad (5.5)$$

and by Lemma 1,  $a > b_i$  for all  $i$ . Applying the equality (5.5) to  $b_i$  several times, finally, by der-lex order we obtain a generator with derivation degree 1 which correspond to the first case.  $\square$

## 6. THE DEFINING IDENTITIES OF VARIETY $\text{DerPerm}_\succ$

**Definition 3.** An algebra with identities (2.3), (2.4) and (2.5) is called SLC-algebra(special left-commutative).

Let the set  $N(X)$  be a linear basis of the free Novikov algebra. Every non-associative word  $u$  in  $X$  can be written as

$$u = L(u_1, \dots, u_k, x) := u_1 \succ (u_2 \succ (\dots \succ (u_k \succ x) \dots)), \quad (6.1)$$

where  $u_j$  are non-associative words,  $x \in X$ . Let  $k$  be the *length* of  $u$ . For a linear combination of (6.1), the length is the maximal length of its summands.

**Lemma 2.** *Every element of  $\text{SLC}\langle X \rangle$  may be written as a linear combination of words of the form (6.1) with  $u_j \in N(X)$ ,  $x \in X$ .*

*Proof.* It follows from identities (2.3) and (2.5).  $\square$

Let us recall a free basis of Novikov algebra [6]: that is the monomials of the form (5.2) with additional condition  $i_2 \geq i_1$ . By (5.2),  $u_i$  has a form

$$x_{k_1} \succ (\dots (x_{k_m} \succ ((\dots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots).$$

Now, we are ready to prove Theorem 4.

*Proof.* To prove that the defining identities of the variety  $\text{DerPerm}_\succ$  are (2.3), (2.4) and (2.5) we have to show that any monomial of the free SLC-algebra can be written as a sum of monomials from the set  $[P(X)]$ . In the other words, it means that  $\text{SLC}\langle X \rangle \cong \text{DerPerm}_\succ \langle X \rangle$ . Let us prove it by induction on the degree of monomial

$$u = u_1 \succ (u_2 \succ (\dots \succ (u_k \succ x) \cdots)),$$

where  $u_i \in N(X)$ . The base of the induction is the case when the degree of  $u$  is 4. For monomial of degree  $n$ , we consider several cases:

Case 1:  $u_i = x_j \in X$ . For that case, by (2.3) we have

$$u = u_1 \succ (u_2 \succ (\cdots \succ (u_k \succ x) \cdots)) = x_j \succ (u_1 \succ (\cdots \succ (u_k \succ x) \cdots))$$

and the monomial

$$u_1 \succ (\cdots \succ (u_{i-1} \succ (u_{i+1} \succ \cdots \succ (u_k \succ x) \cdots)) \cdots)$$

can be rewritten by inductive hypothesis.

Case 2:  $u_i \neq x_j$ . By (2.3), without lose of generality we can assume that  $u_1 = x_{k_1} \succ (\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots)$  and we have

$$u = (x_{k_1} \succ (\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots)) \succ R,$$

where  $m \geq 2$  and  $R = u_2 \succ (\cdots \succ (u_k \succ x) \cdots)$ . By (2.4),

$$\begin{aligned} u &= x_{k_1} \succ ((\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots) \succ R) \\ &+ x_{k_2} \succ ((x_{k_1} \succ (\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots)) \succ R) \\ &x_{k_1} \succ (x_{k_2} \succ ((\cdots (x_{k_m} \succ ((\cdots (x_{j_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1})) \cdots) \succ R)) \end{aligned}$$

which correspond to the first case.

Case 3.  $u_i \neq x_j$ , and

$$u_i = (\cdots ((x_{j_1} \succ P_{n-1}) \succ x_{i_p}) \cdots \succ x_{i_2}) \succ x_{i_1} \quad (6.2)$$

or

$$u_i = x_{k_1} \succ ((\cdots ((x_{j_1} \succ P_{n-1}) \succ x_{i_p}) \cdots \succ x_{i_2}) \succ x_{i_1}). \quad (6.3)$$

The identity (2.4) can be represented as follows:

$$\begin{aligned} (x_1 \succ x_2) \succ ((x_3 \succ x_4) \succ x_5) &= ((x_1 \succ x_2) \succ (x_3 \succ x_4)) \succ x_5 \\ &- ((x_1 \succ x_2) \succ x_4) \succ (x_3 \succ x_5) + x_4 \succ ((x_1 \succ x_2) \succ (x_3 \succ x_5)). \end{aligned} \quad (6.4)$$

For

$$u = L(u_1, \dots, u_k, x) = u_1 \succ (u_2 \succ (R)),$$

suppose that  $u_1 = (\cdots (x_{p_1} \succ P_{n-1}) \cdots \succ x_{i_2}) \succ x_{i_1}$ ,  $u_2 = (\cdots (x_{t_1} \succ P_{m-1}) \cdots \succ x_{j_2}) \succ x_{j_1}$  and without lose of generality we assume that the degree of  $u_1$  bigger or equal than degree of  $u_2$ . We use (6.4) on  $u$  and obtain

$$\begin{aligned} u_1 \succ (u_2 \succ R) &= (u_1 \succ u_2) \succ R \\ &- (u_1 \succ x_{j_1}) \succ (\overline{u_2} \succ R) + x_{j_1} \succ (u_1 \succ (\overline{u_2} \succ R)), \end{aligned}$$

where  $\overline{u_2} = (\cdots (x_{t_1} \succ P_{m-1}) \cdots) \succ x_{j_2}$ . The monomial  $x_{j_1} \succ (u_1 \succ (\overline{u_2} \succ R))$  correspond to the first case. For the monomial  $(u_1 \succ x_{j_1}) \succ (\overline{u_2} \succ R)$  we use (6.4) in the same way several times until  $\overline{u_2}$  comes to the form  $x_{t_1} \succ P_{m-1}$ . Finally, we obtain

$$\overline{u_1} \succ ((x_{t_1} \succ P_{m-1}) \succ R)$$

and by (6.4) we have

$$\begin{aligned} \overline{u_1} \succ ((x_{t_1} \succ P_{m-1}) \succ R) &= (\overline{u_1} \succ (x_{t_1} \succ P_{m-1})) \succ R \\ -(\overline{u_1} \succ P_{m-1}) \succ (x_{t_1} \succ R) + x_{t_1} \succ (P_{m-1} \succ (\overline{u_1} \succ R)) &= (\overline{u_1} \succ (x_{t_1} \succ P_{m-1})) \succ R \\ &\quad - x_{t_1} \succ ((\overline{u_1} \succ P_{m-1}) \succ R) + x_{t_1} \succ (P_{m-1} \succ (\overline{u_1} \succ R)). \end{aligned}$$

The monomials  $x_{t_1} \succ ((\overline{u_1} \succ B) \succ R)$  and  $x_{t_1} \succ (P_{m-1} \succ (\overline{u_1} \succ R))$  correspond to the first case. Notice that for the remaining monomials  $(u_1 \succ u_2) \succ R$  and  $(\overline{u_1} \succ (x_{t_1} \succ P_{m-1})) \succ R$  the number of left adjoint operators decreased to one i.e. each monomial has the form

$$L(v_1, u_3, \dots, u_k, x) \quad \text{and} \quad L(v_2, u_3, \dots, u_k, x),$$

respectively, where  $v_1 = u_1 \succ u_2$  and  $v_2 = \overline{u_1} \succ (x_{t_1} \succ P_{m-1})$ .

In the same way we consider the case that

$$u_1 = x_{k_1} \succ ((\dots(x_{p_1} \succ P_{n-1}) \dots \succ x_{i_2}) \succ x_{i_1})$$

and  $u_2$  has arbitrary form. By (6.4), we have

$$\begin{aligned} u_1 \succ (u_2 \succ R) &= u_2 \succ (u_1 \succ R) = (u_2 \succ u_1) \succ R \\ &\quad - (u_2 \succ \overline{u_1}) \succ (x_{k_1} \succ R) + \overline{u_1} \succ (u_2 \succ (x_{k_1} \succ R)) = \\ &\quad (u_2 \succ u_1) \succ R - x_{k_1} \succ ((u_2 \succ \overline{u_1}) \succ R) + x_{k_1} \succ (u_2 \succ (\overline{u_1} \succ R)), \end{aligned}$$

where the second and the third monomials correspond to the first case and the number of left adjoint operator of monomial  $(u_2 \succ u_1) \succ R$  decreased to one.

Applying the above method several times, we end up with monomials of the form

$$L(x_{s_1}, \dots, x_{s_r}, N, x_j) = x_{s_1} \succ (\dots(x_{s_r} \succ (N \succ x_j)) \dots),$$

where  $N$  is a Novikov word of the form (6.2) and (6.3). We obtain the following result:

**Lemma 3.** *Every monomial of  $\text{SLC}\langle X \rangle$  algebra can be written as a sum of monomials of the following form:*

$$x_{s_1} \succ (\dots(x_{s_r} \succ (N \succ x_j)) \dots),$$

where  $N$  is a Novikov word of the form (6.2) and (6.3).

Now, we consider the monomials of  $\text{SLC}\langle X \rangle$  of the form

$$u = u_1 \succ x.$$

Let us prove that

$$u = \sum_i \alpha_i p_i, \quad p_i \in [P(X)].$$

Firstly, we consider the case that  $u_1$  has the form (6.3). By (5.1) and (5.2),  $u \notin P(X)$  if and only if  $p = 1$  and  $k_1 > j_1$ :

$$u = (x_{k_1} \succ ((x_{j_1} \succ P_{n-1}) \succ x_{i_1})) \succ x \notin [P(X)].$$

Notice that the identity

$$\begin{aligned} (x_{k_1} \succ ((x_{j_1} \succ P_{n-1}) \succ x_{i_1})) \succ x &= (x_{j_1} \succ ((x_{k_1} \succ P_{n-1}) \succ x_{i_1})) \succ x - \\ & x_{j_1} \succ (((x_{k_1} \succ P_{n-1}) \succ x_{i_1}) \succ x) - x_{k_1} \succ ((x_{j_1} \succ (P_{n-1} \succ x_{i_1})) \succ x) \\ & + x_{k_1} \succ (x_{j_1} \succ ((P_{n-1} \succ x_{i_1}) \succ x)) + x_{k_1} \succ (((x_{j_1} \succ P_{n-1}) \succ x_{i_1}) \succ x) \\ & + x_{j_1} \succ ((x_{k_1} \succ (P_{n-1} \succ x_{i_1})) \succ x) - x_{j_1} \succ (x_{k_1} \succ ((P_{n-1} \succ x_{i_1}) \succ x)) \end{aligned}$$

holds in  $\text{SLC}\langle X \rangle$  algebra. It can be proved using computer algebra as software programs Albert [1]. The first monomial of the right side belongs to  $[P(X)]$  and the other monomials correspond to the first case. So,  $u$  can be written as a sum of monomials of  $[P(X)]$ .

It remain to consider the case that  $u_1$  has the form (6.2). By (5.1) and (5.2), we may assume that

$$u_1 = (\cdots((x_j \succ P_{n-1}) \succ x_{i_r}) \cdots \succ x_{i_2}) \succ x_{i_1}$$

and

$$P_{n-1} = (\cdots((x_t \succ P_{n-2}) \succ x_{l_k}) \cdots \succ x_{l_2}) \succ x_{l_1},$$

where  $k > r$ , or  $k = r$  and  $t > j$ . If  $k > r + 2$ , or  $k = r + 1$  and  $t > j$  then

$$u_1 \succ x \in [P(X)].$$

To save space we omit the multiplication  $\succ$ . We have only two cases for which

$$u_1 \succ x \notin [P(X)].$$

There are  $k = r$ , or  $k = r + 1$  and  $t < j$ . Let us start the second induction on  $r$  for the monomial  $u_1$ . The base of induction is  $r = 0$  and this case is considered above i.e.

$$u = u_1 \succ x = (x_j \succ ((\cdots((x_t \succ P_{n-2}) \succ x_{l_k}) \cdots \succ x_{l_2}) \succ x_{l_1})) \succ x.$$

The “ $\equiv$ ” symbol means that we are excluding all monomials for which  $r$  is smaller than given.

Now, we are ready to consider the general case of  $u$ . By (2.4) we rewrite the monomial  $u$  as follows:

$$\begin{aligned} u_1 x &= ((\cdots((x_j P_{n-1}) x_{i_r}) \cdots x_{i_2}) x_{i_1}) x \equiv \\ & (((\cdots(x_j x_{i_r}) \cdots x_{i_2}) x_{i_1}) P_{n-1}) x = \\ & ((\cdots(x_j x_{i_r}) \cdots x_{i_2}) x_{i_1}) (((\cdots((x_t P_{n-2}) x_{l_k}) \cdots x_{l_2}) x_{l_1}) x) + \\ & (\cdots((x_t P_{n-2}) x_{l_k}) \cdots x_{l_2}) (((\cdots(x_j x_{i_r}) \cdots x_{i_2}) x_{i_1}) x) + \\ & x_{l_1} ((\cdots((x_t P_{n-2}) x_{l_k}) \cdots x_{l_2}) (((\cdots(x_j x_{i_r}) \cdots x_{i_2}) x_{i_1}) x)). \end{aligned}$$

The last term correspond to the first case. For the first and second terms we use (2.3) and (6.4), and obtain

$$\begin{aligned}
& (((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1})x) + \\
& \quad (((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1})x_{l_1})((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x) = \\
& \quad (((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1}))x - \\
& \quad (\cdots(x_j x_{i_r}) \cdots x_{i_2})(((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})x_{i_1})x) \\
& + x_{i_1}((\cdots(x_j x_{i_r}) \cdots x_{i_2})(((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})x)) + \\
& \quad (((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1})x_{l_1})(\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2}))x - \\
& (\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_3})(((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1})x_{l_1})x_{l_2})x) + \\
& \quad x_{l_2}(((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1})x_{l_1})((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_3})x)).
\end{aligned}$$

The third and sixth terms correspond to the first case. The first term we rewrite as follows:

$$\begin{aligned}
& (((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})((\cdots(x_j x_{i_r}) \cdots x_{i_2})x_{i_1}))x \equiv \\
& \quad ((\cdots(x_j x_{i_r}) \cdots x_{i_2})(((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})x_{i_1}))x \equiv \\
& \quad ((\cdots(x_j((\cdots((x_t P_{n-2})x_{l_k}) \cdots x_{l_2})x_{l_1})x_{i_1}))x_{i_r}) \cdots x_{i_2}))x \in [P(X)].
\end{aligned}$$

For the second, forth and fifth terms we use the same rewriting method as given above and finally obtain representation in terms of  $[P(X)]$ .  $\square$

## REFERENCES

- [1] Albert version 4.0M6; <https://web.osu.cz/Zusmanovich/soft/albert/>.
- [2] L. A. Bokut, Y. Chen, Z. Zhang, Gröbner-Shirshov bases method for Gelfand-Dorfman-Novikov algebras, *Journal of Algebra and Its Applications*, 16(1) (2017) p. 1-22. .
- [3] F. Chapoton, Un endofoncteur de la catégorie des opérades, In: *Dialgebras and Related Operads. Lecture Notes in Mathematics 1763*, Springer, Berlin, Heidelberg, 2001.
- [4] P. M. Cohn, On homomorphic images of special Jordan algebras, *Canadian Journal of Mathematics* 6 (1954), 253–264.
- [5] A. S. Dzhumadil'daev, C. Löfwall, Trees, free right-symmetric algebras, free Novikov algebras and identities, *Homology, Homotopy Appl.*, 4 (2), 165–190 (2002).
- [6] V. Gubarev, B. K. Sartayev, Free special Gelfand–Dorfman algebras, to appear.
- [7] P. S. Kolesnikov, B. Sartayev, A. Orazgaliev, Gelfand–Dorfman algebras, derived identities, and the Manin product of operads, *Journal of Algebra* 539, 260–284 (2019).
- [8] P. S. Kolesnikov, B. Sartayev, On the embedding of left-symmetric algebras into differential Perm-algebras, *Communications in Algebra*, 50 (2022), 3246–3260
- [9] P. S. Kolesnikov, B. Sartayev, On the special identities of Gelfand–Dorfman algebras, *Experimental Mathematics*, DOI: 10.1080/10586458.2022.2041134
- [10] J.-L. Loday, Dialgebras, In: Loday J.-L. , Frabetti A., Chapoton F., Goichot F. (Eds), *Dialgebras and related operads*, Springer-Verl., Berlin, 2001, pp. 105–110. (Lectures Notes in Math., vol. 1763).
- [11] F. Mashurov, B. K. Sartayev, Metabelian Lie and Perm algebras, <https://arxiv.org/abs/2110.06032>, (2022).

- [12] Rabigul Tuniyaz, L. A. Bokut, Marhaba Xiryazidin, Abdukadir Obul, Gröbner-Shirshov bases for free Gelfand-Dorfman-Novikov algebras and for right ideals of free right Leibniz algebras, *Communications in Algebra* 2018, VOL. 46, NO. 10, 4392–4402.
- [13] B. Vallette, Manin products, Koszul duality, Loday algebras and Deligne conjecture, *J. Reine Angew. Math.*, 620, 105–164 (2008).

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, RUSSIA

SULEYMAN DEMIREL UNIVERSITY, KASKELEN, KAZAKHSTAN