

Strict Log-Subadditivity for Overpartition Rank

Helen W.J. Zhang^{1,2} and Ying Zhong¹

¹School of Mathematics
Hunan University
Changsha 410082, P. R. China

²Hunan Provincial Key Laboratory of
Intelligent Information Processing and Applied Mathematics
Changsha 410082, P. R. China

Emails: helenzhang@hnu.edu.cn, YingZhong@hnu.edu.cn

Abstract

Bessenrodt and Ono initially found the strict log-subadditivity of partition function $p(n)$, that is, $p(a+b) < p(a)p(b)$ for $a, b > 1$ and $a+b > 9$. Many other important statistics of partitions are proved to enjoy similar properties. Lovejoy introduced the overpartition rank as an analog of Dyson's rank for partitions from the q -series perspective. Let $\overline{N}(a, c, n)$ denote the number of overpartitions with rank congruent to a modulo c . Ciolan computed the asymptotic formula of $\overline{N}(a, c, n)$ and showed that $\overline{N}(a, c, n) > \overline{N}(b, c, n)$ for $c \geq 7$ and n large enough. In this paper, we derive an upper bound and a lower bound of $\overline{N}(a, c, n)$ for each $c \geq 3$ by using the asymptotics of Ciolan. Consequently, we establish the strict log-subadditivity of $\overline{N}(a, c, n)$ analogous to the partition function $p(n)$.

Mathematics Subject Classification: 05A20, 11P82, 11P83

Keywords: Rank, Overpartition, Strict Log-Subadditivity

1 Introduction

Strict log-subadditivity phenomena of partition statistics oriented from the work of Bessenrodt and Ono [4]. Recall that a partition of a positive integer n is a sequence of non-increasing positive integers whose sum equals n . Let $p(n)$ denote the number of partitions of n . Bessenrodt and Ono showed that $p(n)$ satisfies the strict log-subadditivity result, that is, for $a, b > 1$ and $a+b > 9$

$$p(a+b) < p(a)p(b).$$

Since then different partition functions are proven to enjoy the strict log-subadditivity as $p(n)$, for instance, the overpartition function [18], the k -regular partition function [3] and the spt-function [10, 12].

Strict log-subadditivity of other statistics concerned with partitions comes into sight. Dyson [13] introduced the rank of a partition to interpret Ramanujan's congruences for $p(n)$ combinatorially. This is defined as the largest part of the partition minus the number of parts. Let $N(a, c, n)$ denote the number of partitions of n with rank congruent to a modulo c . Hou and Jagadeesan [17] showed that

$$N(r, 3, a + b) < N(r, 3, a)N(r, 3, b)$$

when $r = 0$ (resp. $r = 1, 2$) and $a + b \geq 12$ (resp. 11, 12). They also conjectured the general strict log-subadditivity: for $0 \leq r < t$ and $t \geq 2$,

$$N(r, t, a + b) < N(r, t, a)N(r, t, b) \tag{1.1}$$

holds for sufficiently large a and b . This conjecture is confirmed by Males [20]. Gomez and Zhu [15] proved that (1.1) holds for $t = 2$ and $a, b \geq 12$. The strict log-subadditivity for other statistics are referred to [2, 7, 16].

In this paper, we show the strict log-subadditivity rank of overpartitions. Recall that an overpartition of a nonnegative integer n is a partition of n where the first occurrence of each distinct part may be overlined. Let $\ell(\lambda)$ denote the largest part of λ , $\#(\lambda)$ denote the number of parts of λ . Lovejoy [19] defined the rank of an overpartiton λ as

$$rank(\lambda) = \ell(\lambda) - \#(\lambda).$$

Let $\overline{N}(a, c, n)$ be the number of overpartitions of n with rank congruent to a modulo c . Our main result as stated as follows.

Theorem 1.1. *Given any residue $a \pmod{c}$, we have*

$$\overline{N}(a, c, n_1 + n_2) < \overline{N}(a, c, n_1)\overline{N}(a, c, n_2) \tag{1.2}$$

if $3 \leq c \leq 5$ and $n_1, n_2 \geq 9$ or if $c \geq 6$ and $n_1, n_2 \geq M_c$, where

$$M_c = 1.691 \times 10^{13} c^{20} \exp\left(\frac{16c^2(2c^2 + \pi)}{\pi^2}\right).$$

This paper is organized as follows. In Section 2, we state the asymptotic behavior of $\overline{N}(a, c, n)$ due to Ciolan. In Section 3, we estimate the upper bounds for both main terms and error terms in the asymptotic formula. Based on the estimates, we finally prove Theorem 1.1 in Section 4.

2 Asymptotic formula of Ciolan

Our approach to the strict log-subadditivity of overpartition rank relies on the asymptotic behavior of generating function of $\overline{N}(a, c, n)$. In this section, we recall the asymptotic formula due to Ciolan [8, 9].

We begin with some notations that we need. Let

$$\omega_{h,k} := \exp(\pi i s(h, k)),$$

where the Dedekind sum $s(h, k)$ is defined by

$$s(h, k) := \sum_{u \pmod{k}} \left(\left(\frac{u}{k} \right) \right) \left(\left(\frac{hu}{k} \right) \right).$$

Here $((\cdot))$ is the sawtooth function defined by

$$((x)) := \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \in \mathbb{R} \setminus \mathbb{Z}, \\ 0, & \text{if } x \in \mathbb{Z}. \end{cases}$$

In what follows, $0 \leq h < k$ are coprime integers, and $h' \in \mathbb{Z}$ is defined by the congruence $hh' \equiv 1 \pmod{k}$. Further, let $0 < a < c$ are coprime positive integers with $c > 2$, $c_1 := \frac{c}{(c,k)}$ and $k_1 := \frac{k}{(c,k)}$. Let the integer $0 \leq l < c_1$ be the solution to $l \equiv ak_1 \pmod{c_1}$.

We then introduce several Kloosterman sums. Here and throughout we write \sum'_h to denote summation over the integers $0 \leq h < k$ that are coprime to k .

If $c|k$, let

$$A_{a,c,k}(n, m) := (-1)^{k_1+1} \tan\left(\frac{\pi a}{c}\right) \sum'_h \frac{\omega_{h,k}^2}{\omega_{h,k/2}} \cdot \cot\left(\frac{\pi ah'}{c}\right) \cdot e^{-\frac{2\pi i h' a^2 k_1}{c}} \cdot e^{\frac{2\pi i}{k}(nh+mh')},$$

and

$$B_{a,c,k}(n, m) := -\frac{1}{\sqrt{2}} \tan\left(\frac{\pi a}{c}\right) \sum'_h \frac{\omega_{h,k}^2}{\omega_{2h,k}} \cdot \frac{1}{\sin\left(\frac{\pi ah'}{c}\right)} \cdot e^{-\frac{2\pi i h' a^2 k_1}{c}} \cdot e^{\frac{2\pi i}{k}(nh+mh')}.$$

If $c \nmid k$ and $0 < \frac{l}{c_1} \leq \frac{1}{4}$, let

$$D_{a,c,k}(n, m) := \frac{1}{\sqrt{2}} \tan\left(\frac{\pi a}{c}\right) \sum_h' \frac{\omega_{h,k}^2}{\omega_{2h,k}} \cdot e^{\frac{2\pi i}{k}(nh+mh')},$$

and if $c \nmid k$ and $\frac{3}{4} < \frac{l}{c_1} < 1$, let

$$D_{a,c,k}(n, m) := -\frac{1}{\sqrt{2}} \tan\left(\frac{\pi a}{c}\right) \sum_h' \frac{\omega_{h,k}^2}{\omega_{2h,k}} \cdot e^{\frac{2\pi i}{k}(nh+mh')}.$$

Finally, if $c \nmid k$, set

$$\delta_{c,k,r} := \begin{cases} \frac{1}{16} - \frac{l}{2c_1} + \frac{l^2}{c_1^2} - r\frac{l}{c_1}, & \text{if } 0 < \frac{l}{c_1} \leq \frac{1}{4}, \\ 0, & \text{if } \frac{1}{4} < \frac{l}{c_1} \leq \frac{3}{4}, \\ \frac{1}{16} - \frac{3l}{2c_1} + \frac{l^2}{c_1^2} + \frac{1}{2} - r\left(1 - \frac{l}{c_1}\right), & \text{if } \frac{3}{4} < \frac{l}{c_1} < 1, \end{cases}$$

and

$$m_{a,c,k,r} := \begin{cases} -\frac{1}{2c_1^2}(2(ak_1 - l)^2 + c_1(ak_1 - l) + 2rc_1(ak_1 - l)), & \text{if } 0 < \frac{l}{c_1} \leq \frac{1}{4}, \\ 0, & \text{if } \frac{1}{4} < \frac{l}{c_1} \leq \frac{3}{4}, \\ -\frac{1}{2c_1^2}(2(ak_1 - l)^2 + 3c_1(ak_1 - l) - 2rc_1(ak_1 - l) - c_1^2(2r - 1)), & \text{if } \frac{3}{4} < \frac{l}{c_1} < 1. \end{cases}$$

Let $\overline{N}(m, n)$ denote the number of overpartitions of n with rank m . We adopt the notation $\mathcal{O}(u; q)$ in [19] to denote the generating function of $\overline{N}(m, n)$, that is,

$$\mathcal{O}(u; q) := \sum_{n \geq 0} \sum_m \overline{N}(m, n) u^m q^n. \quad (2.1)$$

If $0 < a < c$ are coprime positive integers, and if by $\zeta_n = e^{\frac{2\pi i}{n}}$ we denote the primitive n -th root of unity, we set

$$\mathcal{O}\left(\frac{a}{c}; q\right) := \mathcal{O}(\zeta_c^a; q) = 1 + \sum_{n \geq 1} A\left(\frac{a}{c}; n\right) q^n. \quad (2.2)$$

Ciolan obtained the following asymptotic formula for $A\left(\frac{a}{c}; n\right)$.

Theorem 2.1. [8, Theorem 1] *If $0 < a < c$ are coprime positive integers with $c > 2$, and $\varepsilon > 0$ is arbitrary, then*

$$A\left(\frac{a}{c}; n\right) = i\sqrt{\frac{2}{n}} \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k, 2 \nmid k}} \frac{B_{a,c,k}(-n, 0)}{\sqrt{k}} \cdot \sinh\left(\frac{\pi\sqrt{n}}{k}\right)$$

$$+ 2\sqrt{\frac{2}{n}} \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k, 2 \nmid k, c_1 \neq 4 \\ r \geq 0, \delta_{c,k,r} > 0}} \frac{D_{a,c,k}(-n, m_{a,c,k,r})}{\sqrt{k}} \cdot \sinh\left(\frac{4\pi\sqrt{\delta_{c,k,r}n}}{k}\right) + O(n^\varepsilon). \quad (2.3)$$

To obtain the inequality of $\overline{N}(a, c, n)$, Ciolan [9] bounded the main term and the error term in above theorem. Here we list these bounds respectively.

- Bounds in the main terms: Ciolan [9, § 4.2] gave upper bounds for Kloosterman sums and hyperbolic sine function in (2.3).

$$|B_{j,c,k}(-n, 0)| \leq \cot\left(\frac{\pi}{2c}\right) \frac{2k(1 + \log(\frac{c-1}{2}))}{\pi(1 - \frac{\pi^2}{24})}, \quad (2.4)$$

$$|D_{j,c,k}(-n, m_{j,c,k,r})| \leq \frac{k}{\sqrt{2}} \cot\left(\frac{\pi}{2c}\right), \quad (2.5)$$

$$\sinh\left(\frac{4\pi\sqrt{\delta_{c,k,r}n}}{k}\right) \leq \sinh\left(\pi\sqrt{n}\left(1 - \frac{4}{c}\right)\right). \quad (2.6)$$

- Bounds in the error terms: the error term $O(n^\varepsilon)$ is constituted of S_1, S_2, \dots, S_8 , the error terms of $\sum_1, \sum_2, \dots, \sum_8$ respectively. Here we strictly follows the notations of Ciolan in [9, § 4.3]. Let $\overline{p}(n)$ denote the number of overpartitions of n . Then

$$\begin{aligned} S_1 &< 4C_3 e^{2\pi} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, & S_2 &< 4C_1 e^{2\pi} \sqrt{2} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, \\ S_3 &< 2C_4 e^{2\pi} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, & S_4 &< C_5 e^{2\pi} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, \\ S_5 &< C_2 e^{2\pi} \sqrt{2} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, & S_6 &< C_2 e^{2\pi} \frac{1}{\sqrt{2}} \cot\left(\frac{\pi}{2c}\right) \sum_k k^{-\frac{1}{2}}, \\ S_7 &< \frac{n^{\frac{3}{4}} \log\left(\frac{n}{4}\right)}{2\pi\left(1 - \frac{\pi^2}{24}\right) \sin\left(\frac{\pi}{c}\right)}, & S_8 &< \frac{n^{\frac{3}{4}} \log\left(\frac{n}{4}\right)}{2\pi\left(1 - \frac{\pi^2}{24}\right) \sin\left(\frac{\pi}{c}\right)}, \end{aligned} \quad (2.7)$$

where

$$C_1 = \sum_{r \geq 1} \overline{p}(r) \left(e^{-\frac{(16r-1)\pi}{16}} + e^{-\frac{(16r+7)\pi}{16}} \right), \quad C_2 = 2 \sum_{r \geq 1} \overline{p}(r) e^{-\frac{(c^2-8)\pi r}{16c^2}}, \quad (2.8)$$

$$C_3 = \sum_{r \geq 1} \overline{p}(r) e^{-\pi r}, \quad C_4 = \sum_{r \geq 1} \overline{p}(r) e^{-\frac{\pi r}{2c^2}}, \quad C_5 = \sum_{r \geq 1} \overline{p}(r) e^{-\frac{\pi(2r+1)}{8}}. \quad (2.9)$$

To our need, we adopt the following notations. The errors arising from integrating over the remaining parts of the interval are denoted by S_{2err} , S_{5err} , S_{6err} , and errors introduced by integrating along a smaller arc are denoted by I_{2err} , I_{5err} , I_{6err} . Ciolan [9, § 4.3] provided the following bounds on each of those pieces:

$$\begin{aligned}
S_{2err} &< \sqrt{2}e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) n^{-\frac{1}{2}} \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} \sum_k k^{-\frac{1}{2}}, \\
S_{5err} &< 2\sqrt{2}e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) n^{-\frac{1}{2}} \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} \sum_k k^{-\frac{1}{2}}, \\
S_{6err} &< \sqrt{2}e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) n^{-\frac{1}{2}} \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} \sum_k k^{-\frac{1}{2}}
\end{aligned} \tag{2.10}$$

and

$$\begin{aligned}
I_{2err} &< 2\sqrt{2} \left(\frac{4}{3} + 2^{\frac{5}{4}}\right) e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} n^{\frac{1}{4}}, \\
I_{5err} &< 4\sqrt{2} \left(\frac{4}{3} + 2^{\frac{5}{4}}\right) e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} n^{\frac{1}{4}}, \\
I_{6err} &< 2\sqrt{2} \left(\frac{4}{3} + 2^{\frac{5}{4}}\right) e^{2\pi+\frac{\pi}{8}} \cot\left(\frac{\pi}{2c}\right) \frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} n^{\frac{1}{4}}.
\end{aligned} \tag{2.11}$$

3 Bounds for $\overline{N}(a, c, n)$

In this section, we show that $\overline{N}(a, c, n)$ can be bounded by a constant multiple of $\overline{p}(n)$ for each $c \geq 3$.

Theorem 3.1. *We have*

$$0.0019 \overline{p}(n) < \overline{N}(a, 3, n) < 0.6648 \overline{p}(n), \quad n \geq 2089, \tag{3.1}$$

$$0.0091 \overline{p}(n) < \overline{N}(a, 4, n) < 0.4909 \overline{p}(n), \quad n \geq 272, \tag{3.2}$$

$$0.0103 \overline{p}(n) < \overline{N}(a, 5, n) < 0.3897 \overline{p}(n), \quad n \geq 449. \tag{3.3}$$

For $c \geq 6$ we define the constant

$$M_c := 1.691 \times 10^{13} c^{20} \exp\left(\frac{16c^2(2c^2 + \pi)}{\pi^2}\right), \tag{3.4}$$

we have

$$\left(\frac{1}{2c}\right)\bar{p}(n) < \bar{N}(a, c, n) < \left(\frac{3}{2c}\right)\bar{p}(n), \quad n \geq M_c. \quad (3.5)$$

Our main mission of this section is to prove the above theorem. We sketch our strategy here. Setting $u = \zeta_c^a$ in (2.1), we have

$$\mathcal{O}\left(\frac{a}{c}; q\right) = \sum_{n \geq 0} \sum_m \bar{N}(m, n) \zeta_c^{am} q^n. \quad (3.6)$$

Comparing the coefficient of q^n in (2.2), (3.6) and with the orthogonality of the roots of unity, we obtain

$$\bar{N}(a, c, n) = \frac{1}{c}\bar{p}(n) + \frac{1}{c} \sum_{1 \leq j \leq c-1} \zeta_c^{-aj} A\left(\frac{j}{c}; n\right). \quad (3.7)$$

The second summation in right-hand side of (3.7) is denoted by $\bar{R}(a, c, n)$, that is,

$$\bar{N}(a, c, n) = \frac{1}{c}\bar{p}(n) + \bar{R}(a, c, n). \quad (3.8)$$

Due to (3.8), we need to estimate $\bar{R}(a, c, n)$ and $\bar{p}(n)$ separately. For $\bar{R}(a, c, n)$, we need to give a bound to the main terms as well as the error terms in view of Theorem 2.1. These are completed in Theorem 3.3 and Theorem 3.5 respectively. After then, we give an upper bound and lower bound for $\bar{p}(n)$ in Theorem 3.6 based on the asymptotic formula of Engel [14].

From the view of Theorem 2.1, we have the following estimates to bound the main terms of $\bar{R}(a, c, n)$.

Lemma 3.2. *For integer $c > 2$, then we have*

$$|B_{j,c,k}(-n, 0)| \leq 0.3444kc^2, \quad (3.9)$$

$$|D_{j,c,k}(-n, m_{j,c,k,r})| \leq 0.4503kc, \quad (3.10)$$

$$\sinh\left(\frac{4\pi\sqrt{\delta_{c,k,r}n}}{k}\right) \leq \frac{1}{2}e^{\pi\sqrt{n}(1-\frac{4}{c})}. \quad (3.11)$$

Proof. We shall estimate (3.9). Using the fact in [6, P. 159],

$$\log x \leq a(x^{1/a} - 1), \quad a, x > 0, \quad (3.12)$$

we have

$$\frac{1 + \log\left(\frac{c-1}{2}\right)}{\pi\left(1 - \frac{\pi^2}{24}\right)} < 0.2704c. \quad (3.13)$$

Substituting (3.13) and

$$\cot\left(\frac{\pi}{2c}\right) \leq \frac{2c}{\pi} < 0.6367c \quad (3.14)$$

into (2.4), we get the first inequality (3.9). By means of the estimates in (2.5) and (3.14), we arrive at the second inequality (3.10). The third inequality (3.11) immediately follows from the definition of hyperbolic sine function. This completes the proof. \blacksquare

We can give a bound for the main term of $\bar{R}(a, c, n)$.

Theorem 3.3. *A bound for the main term of $\bar{R}(a, c, n)$ is*

$$0.1624e^{\frac{\pi\sqrt{n}}{c}}n^{\frac{1}{4}}c + (0.0266c + 0.2123)e^{\pi\sqrt{n}\left(1-\frac{4}{c}\right)}n^{\frac{1}{4}}c.$$

Proof. Immediately, the main term of (2.3) gives the main term of $\bar{R}(a, c, n)$

$$\begin{aligned} & \frac{1}{c} \sum_{1 \leq j \leq c-1} \zeta_c^{-aj} i \sqrt{\frac{2}{n}} \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k, 2 \nmid k}} \frac{B_{j,c,k}(-n, 0)}{\sqrt{k}} \cdot \sinh\left(\frac{\pi\sqrt{n}}{k}\right) \\ & + \frac{1}{c} \sum_{1 \leq j \leq c-1} \zeta_c^{-aj} 2 \sqrt{\frac{2}{n}} \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k, 2 \nmid k, c_1 \neq 4 \\ r \geq 0, \delta_{c,k,r} > 0}} \frac{D_{j,c,k}(-n, m_{j,c,k,r})}{\sqrt{k}} \cdot \sinh\left(\frac{4\pi\sqrt{\delta_{c,k,r}n}}{k}\right). \end{aligned} \quad (3.15)$$

Let G_1 and G_2 denote the summations of (3.15), respectively. By the bounds in (3.9), we have

$$|G_1| \leq \sqrt{\frac{2}{n}} \sinh\left(\frac{\pi\sqrt{n}}{c}\right) \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k}} 0.3444k^{\frac{1}{2}}c^2 \leq 0.1624e^{\frac{\pi\sqrt{n}}{c}}n^{\frac{1}{4}}c, \quad (3.16)$$

where the last inequality is due to

$$\sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k}} k^{\frac{1}{2}} \leq c^{\frac{1}{2}} \sum_{1 \leq m \leq \lfloor \frac{\sqrt{n}}{c} \rfloor} m^{\frac{1}{2}} \leq c^{\frac{1}{2}} \int_1^{\lfloor \frac{\sqrt{n}}{c} \rfloor} x^{\frac{1}{2}} dx \leq \frac{2}{3c} n^{\frac{3}{4}}.$$

Now we consider the bound of $|G_2|$. For fixed k , it is easy to see that

$$\sum_{\substack{r \geq 0 \\ c|k, 2 \nmid k, c_1 \neq 4 \\ \delta_{c,k,r} > 0}} 1 \leq \frac{c+8}{16}. \quad (3.17)$$

Then by (3.10), (3.11) and (3.17), we get

$$|G_2| \leq 0.4503c \sqrt{\frac{2}{n}} e^{\pi\sqrt{n}(1-\frac{4}{c})} \frac{c+8}{16} \sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k}} k^{\frac{1}{2}}.$$

Since

$$\sum_{\substack{1 \leq k \leq \sqrt{n} \\ c|k}} k^{\frac{1}{2}} \leq \int_1^{\sqrt{n}} x^{\frac{1}{2}} dx \leq \frac{2}{3} n^{\frac{3}{4}},$$

we have

$$\begin{aligned} |G_2| &\leq 0.4503c \sqrt{\frac{2}{n}} e^{\pi\sqrt{n}(1-\frac{4}{c})} \cdot \frac{c+8}{16} \cdot \frac{2}{3} n^{\frac{3}{4}} \\ &\leq (0.0266c + 0.2123) e^{\pi\sqrt{n}(1-\frac{4}{c})} n^{\frac{1}{4}} c. \end{aligned} \quad (3.18)$$

Plugging (3.16) and (3.18) into (3.15), we complete the proof. \blacksquare

Next we shall give a bound for the error terms of $\bar{R}(a, c, n)$. To this end, we need to analyse the terms in (2.7), (2.10) and (2.11) more detailedly.

Lemma 3.4. *For integer $c > 2$, then we have*

$$\begin{aligned} S_1 &\leq 1496.9n^{\frac{1}{4}}c, & S_2 &\leq 3111.36n^{\frac{1}{4}}c, & S_3 &\leq 1363.79\bar{C}_4n^{\frac{1}{4}}c, \\ S_4 &\leq 82469.8n^{\frac{1}{4}}c, & S_5 &\leq 964.35\bar{C}_2n^{\frac{1}{4}}c, & S_6 &\leq 482.18\bar{C}_2n^{\frac{1}{4}}c, \\ S_7 &\leq 0.9093n^{\frac{7}{8}}c, & S_8 &\leq 0.9093n^{\frac{7}{8}}c \end{aligned}$$

and

$$\begin{aligned} S_{2err} &\leq 386.18n^{-\frac{1}{4}}c^2, & S_{5err} &\leq 772.36n^{-\frac{1}{4}}c^2, & S_{6err} &\leq 386.18n^{-\frac{1}{4}}c^2, \\ I_{2err} &\leq 1433.39n^{\frac{1}{4}}c^2, & I_{5err} &\leq 2866.78n^{\frac{1}{4}}c^2, & I_{6err} &\leq 1433.39n^{\frac{1}{4}}c^2, \end{aligned}$$

where

$$\bar{C}_2 = 2 \exp\left(\frac{32c^2(16c^2 + (c^2 - 8)\pi)}{\pi^2(c^2 - 8)^2}\right), \quad \bar{C}_4 = \exp\left(\frac{4c^2(2c^2 + \pi)}{\pi^2}\right).$$

Proof. First, we give the estimations of C_1, C_2, C_3, C_4, C_5 in (2.8) and (2.9). By the fact

$$\bar{p}(n) < e^{\pi\sqrt{n}}, \quad (3.19)$$

it is easy to check that

$$C_1 \leq 0.8066, \quad C_3 \leq 0.5488, \quad C_5 \leq 120.942. \quad (3.20)$$

To estimate C_2 , we claim that

$$\sum_{n \geq 0} \bar{p}(n) e^{-2\pi n y} \leq \exp\left(\frac{2e^{-2\pi y}}{(1 - e^{-2\pi y})^2}\right), \quad (3.21)$$

where y is a positive real number. Recall that the generating function of $\bar{p}(n)$ is given by Corteel and Lovejoy [11]

$$F(q) := \sum_{n \geq 0} \bar{p}(n) q^n = \prod_{n \geq 1} \frac{(1 + q^n)}{(1 - q^n)}, \quad (3.22)$$

where $q = e^{2\pi iz}$ with $z = x + iy$. Taking logarithm of both sides of (3.22), we have

$$\begin{aligned} \log F(q) &= \sum_{n \geq 1} (\log(1 + q^n) - \log(1 - q^n)) \\ &= \sum_{n \geq 1} \sum_{m \geq 1} \left(\frac{q^{nm}}{m} - \frac{(-1)^m q^{nm}}{m} \right) \\ &< \frac{2q}{(1 - q)^2}. \end{aligned}$$

Hence

$$F(|q|) \leq \exp\left(\frac{2|q|}{(1 - |q|)^2}\right),$$

together with $|q| = e^{-2\pi y}$, which gives (3.21) as claimed. Since integer $c > 2$, $c^2 - 8 > 0$. Take $y = (c^2 - 8)/32c^2$ in (3.21) and we have

$$C_2 \leq 2 \exp\left(\frac{2e^{-\frac{(c^2-8)\pi}{16c^2}}}{\left(1 - e^{-\frac{(c^2-8)\pi}{16c^2}}\right)^2}\right).$$

For $x > -1$, it can be easily derived from

$$\frac{x}{1+x} < 1 - e^{-x} < x$$

that

$$\frac{e^{-x}}{(1 - e^{-x})^2} < \frac{1 + x}{x^2}. \quad (3.23)$$

Therefore

$$C_2 \leq 2 \exp\left(\frac{32c^2(16c^2 + (c^2 - 8)\pi)}{\pi^2(c^2 - 8)^2}\right) := \overline{C}_2. \quad (3.24)$$

By the same argument as above, one can derive

$$C_4 \leq \exp\left(\frac{4c^2(2c^2 + \pi)}{\pi^2}\right) := \overline{C}_4 \quad (3.25)$$

from (3.21) and (3.23).

Since

$$\sum_k k^{-\frac{1}{2}} = \int_1^{\sqrt{n}} x^{-\frac{1}{2}} dx \leq 2n^{\frac{1}{4}}. \quad (3.26)$$

Substituting (3.14), (3.26) and the bounds for C_1, \dots, C_5 as given in (3.20), (3.24) and (3.25) into (2.7), we finally derive estimates for S_1, \dots, S_6 in Lemma 3.4.

As for S_7 and S_8 , first we note that when $x \in (0, \frac{\pi}{2})$,

$$\sin x \geq \frac{2x}{\pi}.$$

Then we have

$$\sin\left(\frac{\pi}{c}\right) \geq \frac{2}{c}. \quad (3.27)$$

Meanwhile, setting $x = \frac{n}{4}$ and $a = 8$ in (3.12), we have

$$\log\left(\frac{n}{4}\right) \leq 8\left(\frac{n}{4}\right)^{\frac{1}{8}}. \quad (3.28)$$

Thus S_7 and S_8 is bounded by $0.9093n^{\frac{7}{8}}c$ as we substitute (3.27) and (3.28) into (2.7). The asymptotic estimates $S_{2err}, S_{5err}, S_{6err}, I_{2err}, I_{5err}$ and I_{6err} easily follow from (3.13), (3.14) and (3.26). This completes the proof. \blacksquare

A bound for the error terms of $\overline{R}(a, c, n)$ immediately follows from the above lemma.

Theorem 3.5. *The bound for the error term of $\overline{R}(a, c, n)$ is*

$$1544.72n^{-\frac{1}{4}}c^2 + 87078.1n^{\frac{1}{4}}c + 5733.56n^{\frac{1}{4}}c^2 + 1.8186n^{\frac{7}{8}}c \\ + 1363.79\overline{C}_4n^{\frac{1}{4}}c + 1446.53\overline{C}_2n^{\frac{1}{4}}c. \quad (3.29)$$

Proof. Adding the estimates in Lemma 3.4 together and applying to (2.3), we get the bound for the error term of $\overline{R}(a, c, n)$ as desired. ■

The following bounds for $\overline{p}(n)$ serves as a connection between $\overline{p}(n)$ and $\overline{R}(a, c, n)$.

Theorem 3.6. *We have*

$$\frac{1}{8n} \left(1 - \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}} \leq \overline{p}(n) \leq \frac{1}{8n} \left(1 + \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}}. \quad (3.30)$$

Proof. Recall that Engel [14] provided an error term for the overpartition function

$$\overline{p}(n) = \frac{1}{2\pi} \sum_{\substack{1 \leq k \leq N \\ 2|k}} \sqrt{k} \sum_{\substack{0 \leq h \leq k \\ (h,k)=1}} \frac{\omega(h, k)^2}{\omega(2h, k)} e^{-\frac{2\pi i n h}{k}} \frac{d}{dn} \left(\frac{\sinh\left(\frac{\pi\sqrt{n}}{k}\right)}{\sqrt{n}} \right) + R_2(n, N),$$

where

$$|R_2(n, N)| \leq \frac{N^{\frac{5}{2}}}{\pi n^{\frac{3}{2}}} \sinh\left(\frac{\pi\sqrt{n}}{N}\right).$$

In particular, when $N = 3$, we have

$$\overline{p}(n) = \frac{1}{8n} \left[\left(1 + \frac{1}{\pi\sqrt{n}}\right) e^{-\pi\sqrt{n}} + \left(1 - \frac{1}{\pi\sqrt{n}}\right) e^{\pi\sqrt{n}} \right] + R_2(n, 3), \quad (3.31)$$

where

$$|R_2(n, 3)| \leq \frac{3^{\frac{5}{2}}}{\pi n^{\frac{3}{2}}} \sinh\left(\frac{\pi\sqrt{n}}{3}\right) \leq \frac{3^{\frac{5}{2}} e^{\frac{\pi\sqrt{n}}{3}}}{2\pi n^{\frac{3}{2}}} \leq \frac{\pi e^{\pi\sqrt{n}}}{16\pi n^{\frac{3}{2}}}. \quad (3.32)$$

By (3.31), we get the following upper bound for $\overline{p}(n)$

$$\overline{p}(n) \leq \frac{1}{8n} \left(1 + \frac{1}{\pi\sqrt{n}}\right) e^{\pi\sqrt{n}} + \frac{\pi e^{\pi\sqrt{n}}}{16\pi n^{\frac{3}{2}}}.$$

The right hand side of above inequality can be rewritten as

$$\frac{1}{8n} \left(1 + \frac{1}{\pi\sqrt{n}}\right) e^{\pi\sqrt{n}} + \frac{\pi e^{\pi\sqrt{n}}}{16\pi n^{\frac{3}{2}}} = \frac{1}{8n} \left(1 + \frac{2 + \pi}{2\pi\sqrt{n}}\right) e^{\pi\sqrt{n}},$$

which arrives at the right-hand side of (3.30). The left hand-side of (3.30) can be verified easily.

This completes the proof. ■

We are now ready to complete the proof of Theorem 3.1.

Proof of Theorem 3.1. Combining Theorems 3.3 and 3.5, we have

$$\begin{aligned} |\overline{R}(a, c, n)| \leq & \left| 0.1624e^{\frac{\pi\sqrt{n}}{c}} n^{\frac{1}{4}}c + (0.0266c + 0.2123)e^{\pi\sqrt{n}(1-\frac{4}{c})} n^{\frac{1}{4}}c \right. \\ & + 1544.72n^{-\frac{1}{4}}c^2 + 87078.1n^{\frac{1}{4}}c + 5733.56n^{\frac{1}{4}}c^2 + 1.8186n^{\frac{7}{8}}c \\ & \left. + 1363.79\overline{C}_4n^{\frac{1}{4}}c + 1446.53\overline{C}_2n^{\frac{1}{4}}c \right|. \end{aligned} \quad (3.33)$$

We note that for $n \geq 2$,

$$\frac{\sqrt{n}}{\sqrt{n}-1} \leq \frac{\sqrt{2}}{\sqrt{2}-1} \leq 3.415.$$

By Theorem 3.6, the lower bound of $\overline{p}(n)^{-1}$ can be estimated as

$$\overline{p}(n)^{-1} \leq 8n \frac{\sqrt{n}}{\sqrt{n}-1} e^{-\pi\sqrt{n}} \leq 27.32ne^{-\pi\sqrt{n}}.$$

Combining above inequality, we have for $c > 2$

$$\begin{aligned} \left| \frac{\overline{R}(a, c, n)}{\overline{p}(n)} \right| \leq & 4.44ce^{(\frac{1}{c}-1)\pi\sqrt{n}} n^{\frac{5}{4}} + (0.73c + 5.81)ce^{-\frac{4}{c}\pi\sqrt{n}} n^{\frac{5}{4}} + e^{-\pi\sqrt{n}} \left[42201.8c^2n^{\frac{3}{4}} \right. \\ & + 156641c^2n^{\frac{5}{4}} + 2.379 \times 10^6 cn^{\frac{5}{4}} + 49.69cn^{\frac{15}{8}} + 37258.7c\overline{C}_4n^{\frac{5}{4}} \\ & \left. + 39519.2c\overline{C}_2n^{\frac{5}{4}} \right]. \end{aligned} \quad (3.34)$$

For $c = 3$, recalling the definition of C_2 in (2.8), then using (3.19), we compute that $C_2 < 4.5303 \times 10^{52}$. Similarly, using (2.9), we have $C_4 < 1.0535 \times 10^8$. So replacing \overline{C}_2 and \overline{C}_4 in (3.34) with 4.5303×10^{52} and 1.0535×10^8 , we obtain that

$$\begin{aligned} \left| \frac{\overline{R}(a, 3, n)}{\overline{p}(n)} \right| \leq & 13.32e^{-\frac{2\pi\sqrt{n}}{3}} n^{\frac{5}{4}} + 24e^{-\frac{4}{3}\pi\sqrt{n}} n^{\frac{5}{4}} + e^{-\pi\sqrt{n}} \left(379816.2n^{\frac{3}{4}} \right. \\ & \left. + 5.3711 \times 10^{57} n^{\frac{5}{4}} + 149.07n^{\frac{15}{8}} \right) := R_3. \end{aligned} \quad (3.35)$$

Plugging (3.35) into (3.8), we have

$$\overline{p}(n) \left(\frac{1}{3} - R_3 \right) < \overline{N}(a, 3, n) < \overline{p}(n) \left(\frac{1}{3} + R_3 \right). \quad (3.36)$$

Note that R_3 is decreasing for $n > 1$, then we have

$$R_3 < 0.33142, \quad n \geq 2089. \quad (3.37)$$

Combining (3.36) and (3.37), we are led to (3.1).

Similar to $c = 3$, we first confirm the estimations of C_2 and C_4 . When $c = 4$, $C_2 < 2.977 \times 10^{13}$ and $C_4 < 1.4885 \times 10^{13}$, for $c = 5$, we have $C_2 < 2.4221 \times 10^{10}$ and $C_4 < 4.0102 \times 10^{19}$. Then we obtain

$$\left| \frac{\overline{R}(a, 4, n)}{\overline{p}(n)} \right| \leq 17.76e^{-\frac{3\pi\sqrt{n}}{4}} n^{\frac{5}{4}} + e^{-\pi\sqrt{n}} \left(675228.9n^{\frac{3}{4}} + 6.9244 \times 10^{18} n^{\frac{5}{4}} + 198.76n^{\frac{15}{8}} \right) := R_4, \quad (3.38)$$

and

$$\left| \frac{\overline{R}(a, 5, n)}{\overline{p}(n)} \right| \leq 69.5e^{-\frac{4\pi\sqrt{n}}{5}} n^{\frac{5}{4}} + e^{-\pi\sqrt{n}} \left(1.0551 \times 10^6 n^{\frac{3}{4}} + 7.4708 \times 10^{24} n^{\frac{5}{4}} + 248.45n^{\frac{15}{8}} \right) := R_5. \quad (3.39)$$

Substituting (3.38) and (3.39) into (3.8), respectively, we have

$$\overline{p}(n) \left(\frac{1}{4} - R_4 \right) < \overline{N}(a, 4, n) < \overline{p}(n) \left(\frac{1}{4} + R_4 \right), \quad (3.40)$$

$$\overline{p}(n) \left(\frac{1}{5} - R_5 \right) < \overline{N}(a, 5, n) < \overline{p}(n) \left(\frac{1}{5} + R_5 \right). \quad (3.41)$$

Since

$$R_4 < 0.24084, \quad n \geq 272, \quad (3.42)$$

$$R_5 < 0.1897, \quad n \geq 449. \quad (3.43)$$

Applying the estimates (3.42) into (3.40) and (3.43) into (3.41), we reach (3.2) and (3.3).

Next, we deal with the case $c \geq 6$. Here we combine all the terms in (3.34) except the term of $49.69ce^{-\pi\sqrt{n}} n^{\frac{15}{8}}$ into an upper bound. We take the sum of their coefficients, the highest order exponential, and the highest power of n from the all terms and put them together in one term. For $c \geq 6$, $4/c < 1 - 1/c$, so we have

$$\left| \frac{\overline{R}(a, c, n)}{\overline{p}(n)} \right| \leq 37259c \exp\left(\frac{4c^2(2c^2 + \pi)}{\pi^2}\right) e^{-\frac{4}{c}\pi\sqrt{n}} n^{\frac{5}{4}} + 49.69ce^{-\pi\sqrt{n}} n^{\frac{15}{8}} := R_c. \quad (3.44)$$

In view of (3.8) and (3.44), we get the following bounds of $\overline{N}(a, c, n)$

$$\overline{p}(n) \left(\frac{1}{c} - R_c \right) < \overline{N}(a, c, n) < \overline{p}(n) \left(\frac{1}{c} + R_c \right). \quad (3.45)$$

We claim that

$$R_c < \frac{1}{2c}, \quad n \geq M_c, \quad (3.46)$$

where M_c is defined in (3.4).

To verify (3.46), we only need to show that

$$37259c \exp\left(\frac{4c^2(2c^2 + \pi)}{\pi^2}\right) e^{-\frac{4}{c}\pi\sqrt{n}n^{\frac{5}{4}}} < \frac{1}{4c} \quad (3.47)$$

and

$$49.69ce^{-\pi\sqrt{n}n^{\frac{15}{8}}} < \frac{1}{4c}, \quad (3.48)$$

which are equivalent to prove that

$$\frac{e^{\frac{4}{c}\pi\sqrt{n}}}{n^{\frac{5}{4}}} > 4 \times 37259c^2 \exp\left(\frac{4c^2(2c^2 + \pi)}{\pi^2}\right) \quad (3.49)$$

and

$$\frac{e^{\pi\sqrt{n}}}{n^{\frac{15}{8}}} > 4 \times 49.69c^2. \quad (3.50)$$

In addition, we recall the following inequality [21, P. 76]

$$e^x > \left(1 + \frac{x}{y}\right)^y, \quad x, y > 0.$$

Hence, letting $x = \frac{4}{c}\pi\sqrt{n}$, $y = 3$ and $x = \pi\sqrt{n}$, $y = 4$ respectively, we obtain

$$\frac{e^{\frac{4}{c}\pi\sqrt{n}}}{n^{\frac{5}{4}}} > \frac{1}{n^{\frac{5}{4}}} \left(1 + \frac{4\pi\sqrt{n}}{3c}\right)^3 > \frac{4^3\pi^3}{3^3c^3} n^{\frac{1}{4}} \quad (3.51)$$

and

$$\frac{e^{\pi\sqrt{n}}}{n^{\frac{15}{8}}} > \frac{1}{n^{\frac{15}{8}}} \left(1 + \frac{\pi\sqrt{n}}{4}\right)^4 > \frac{\pi^4}{4^4} n^{\frac{1}{8}}. \quad (3.52)$$

By (3.51) and (3.52), we know that the inequalities (3.49) and (3.50) hold when $n \geq M_c$ and $n \geq M_c'$, respectively, where

$$M_c' := 5.544 \times 10^{21} c^{16}.$$

Furthermore, we see that (3.47) and (3.48) hold when $n \geq M_c$ and $n \geq M_c'$, respectively. Therefore, (3.46) is verified for $n \geq \max\{M_c, M_c'\} = M_c$. Plugging (3.46) into (3.45), we complete the proof. \blacksquare

4 Strict log-subadditivity of $\overline{N}(a, c, n)$

In this section, we present a proof of Theorem 1.1 based on the intermediate inequalities in the previous sections.

Proof of Theorem 1.1. We begin with the case of $c = 3$. Substituting (3.30) into (3.1), we obtain that for $n \geq 2089$

$$(0.0019) \frac{1}{8n} \left(1 - \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}} < \overline{N}(a, 3, n) < (0.6648) \frac{1}{8n} \left(1 + \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}}. \quad (4.1)$$

Letting $n_2 = Cn_1$ for some $C \geq 1$, then by (4.1), we have

$$\overline{N}(a, 3; n_1) \overline{N}(a, 3; n_2) > (0.0019)^2 \frac{1}{64Cn_1^2} \left(1 - \frac{1}{\sqrt{n_1}}\right) \left(1 - \frac{1}{\sqrt{Cn_1}}\right) e^{\pi(\sqrt{n_1} + \sqrt{Cn_1})},$$

and

$$\overline{N}(a, 3; n_1 + n_2) < (0.6648) \frac{1}{8(n_1 + Cn_1)} \left(1 + \frac{1}{\sqrt{n_1 + Cn_1}}\right) e^{\pi\sqrt{n_1 + Cn_1}}.$$

It is sufficient to show that

$$T_{n_1}(C) > \log(V_{n_1}(C)) + \log(S_{n_1}(C)),$$

where

$$T_{n_1}(C) := \pi(\sqrt{n_1} + \sqrt{Cn_1}) - \pi\sqrt{n_1 + Cn_1},$$

$$V_{n_1}(C) := \frac{0.6648 \times 8Cn_1}{(0.0019)^2(C + 1)},$$

$$S_{n_1}(C) := \frac{1 + \frac{1}{\sqrt{n_1 + Cn_1}}}{\left(1 - \frac{1}{\sqrt{n_1}}\right) \left(1 - \frac{1}{\sqrt{Cn_1}}\right)}.$$

As function of C , it can be shown that $T_{n_1}(C)$ is increasing and $S_{n_1}(C)$ is decreasing for $C \geq 1$, and combined with

$$V_{n_1}(C) < \frac{0.6648 \times 8n_1}{(0.0019)^2},$$

it suffices to show that

$$T_{n_1}(1) = 2\pi\sqrt{n_1} - \pi\sqrt{2n_1}$$

$$> \log \left(\frac{0.6648 \times 8n_1}{(0.0019)^2} \right) + \log \left(\frac{1 + \frac{1}{\sqrt{2n_1}}}{\left(1 - \frac{1}{\sqrt{n_1}}\right)^2} \right). \quad (4.2)$$

By a short computation, we find that (4.2) holds for all $n_1 \geq 109$. Hence, if $n_1, n_2 \geq 2089$, we have

$$\bar{N}(a, 3, n_1 + n_2) < \bar{N}(a, 3, n_1)\bar{N}(a, 3, n_2).$$

It is routine to check that (1.2) is true for $c = 3$ and $9 \leq n \leq 2088$.

The proofs of $c = 4, 5$ are similar to that of $c = 3$, and hence, they are omitted.

We proceed to prove the case for $c \geq 6$. Plugging (3.30) into (3.5), we obtain that

$$\frac{1}{16nc} \left(1 - \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}} < \bar{N}(a, c, n) < \frac{3}{16nc} \left(1 + \frac{1}{\sqrt{n}}\right) e^{\pi\sqrt{n}}, \quad n \geq M_c, \quad (4.3)$$

where M_c is defined in (3.4).

Setting $n_2 = Cn_1$ for some $C \geq 1$, one can easily find that for $n_1 \geq M_c$

$$\bar{N}(a, c; n_1)\bar{N}(a, c; n_2) > \frac{1}{256Cn_1^2c^2} \left(1 - \frac{1}{\sqrt{n_1}}\right) \left(1 - \frac{1}{\sqrt{n_2}}\right) e^{\pi(\sqrt{n_1} + \sqrt{Cn_1})},$$

and

$$\bar{N}(a, c; n_1 + n_2) < \frac{3}{16(n_1 + Cn_1)c} \left(1 + \frac{1}{\sqrt{n_1 + Cn_1}}\right) e^{\pi\sqrt{n_1 + Cn_1}}.$$

It remains to show that for $n_1 \geq M_c$

$$T_{n_1}(C) > \log(W_{n_1}(C)) + \log(S_{n_1}(C)),$$

where

$$W_{n_1}(C) := (48c) \frac{Cn_1}{C+1}.$$

Obviously,

$$W_{n_1}(C) < 48cn_1.$$

Therefore, we only need show

$$T_{n_1}(1) = 2\pi\sqrt{n_1} - \pi\sqrt{2n_1}$$

$$> \log(48cn_1) + \log\left(\frac{1 + \frac{1}{\sqrt{2n_1}}}{\left(1 - \frac{1}{\sqrt{n_1}}\right)^2}\right). \quad (4.4)$$

If $n_1 \geq 2$, we obtain

$$\log(48cn_1) + \log\left(\frac{1 + \frac{1}{\sqrt{2n_1}}}{\left(1 - \frac{1}{\sqrt{n_1}}\right)^2}\right) < \log(48cn_1) + \log\left(\frac{1 + \frac{1}{\sqrt{4}}}{\left(1 - \frac{1}{\sqrt{2}}\right)^2}\right) < \log(840cn_1). \quad (4.5)$$

Moreover, if $n_1 \geq (840c)^2 \geq (840 \times 6)^2$, then

$$T_{n_1}(1) = 2\pi\sqrt{n_1} - \pi\sqrt{2n_1} > 2\log(n_1) \geq \log(n_1) + 2\log(840c) > \log(840cn_1). \quad (4.6)$$

Combined with (4.4), (4.5) and (4.6), choosing $n_1, n_2 \geq (840c)^2$, we complete the proof. ■

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant Nos. 12001182 and 12171487), the Fundamental Research Funds for the Central Universities (Grant No. 531118010411) and Hunan Provincial Natural Science Foundation of China (Grant No. 2021JJ40037).

References

- [1] G.E. Andrews, *The Theory of Partitions*, Addison-Wesley, 1976.
- [2] A. Baker and J. Males, Asymptotics, Turán inequalities, and the distribution of the BG-rank and 2-quotient rank of partitions, [arXiv:2203.00747](https://arxiv.org/abs/2203.00747).
- [3] O. Beckwith, C. Bessenrodt, Multiplicative properties of the number of k -regular partitions, *Ann. Comb.* 20 (2) (2016) 231–250.
- [4] C. Bessenrodt and K. Ono, Maximal multiplicative properties of partitions, *Ann. Comb.* 20(1) (2016) 59–64.
- [5] K. Bringmann and J. Lovejoy, Dyson’s rank, overpartitions, and weak Maass forms, *Int. Math. Res. Not.* 2007 (19) (2007) rnm063.
- [6] P.S. Bullen, *A dictionary of inequalities*. Pitman Monographs and Surveys in Pure and Applied Mathematics, 97. Longman, Harlow, 1998.
- [7] G. Cesana, W. Craig and J. Males, Asymptotic equidistribution for partition statistics and topological invariants, [arXiv:2111.13766](https://arxiv.org/abs/2111.13766).

- [8] A. Ciolan, Ranks of overpartitions: asymptotics and inequalities, *J. Math. Anal. Appl.* 480 (2) (2019) 123444, 28 pp.
- [9] A. Ciolan, Inequalities between overpartition ranks for all moduli, *Ramanujan J.* 1 58 (2022) 463–489.
- [10] W.Y.C. Chen, The spt-function of Andrews, *Surveys in Combinatorics 2017*, 141–203, London Math. Soc. Lecture Note Ser., 440, Cambridge Univ. Press, Cambridge, 2017.
- [11] S. Corteel and J. Lovejoy, Overpartitions, *Trans. Amer. Math. Soc.* 356 (2004) 1623–1635.
- [12] M.L. Dawsey and R. Masri, Effective bounds for the Andrews spt-function, *Forum Math.* 31 (2019) 743–767.
- [13] F.J. Dyson, Some guesses in the theory of partitions, *Eureka (Cambridge)* 8 (1944) 10–15.
- [14] B. Engel, Log-concavity of the overpartition function, *Ramanujan J.* 43 (2) (2017) 229–241.
- [15] K. Gomez and E. Zhu, Bounds for coefficients of the $f(q)$ mock theta function and applications to partition ranks, *J. Number Theory* 226 (2021) 1–23.
- [16] A. Hamakiotes, A. Kriegman and W.L. Tsai, Asymptotic distribution of the partition crank, *Ramanujan J.* 56 (3) (2021) 803–820.
- [17] E. Hou and M. Jagadeesan, Dyson’s partition ranks and their multiplicative extensions, *Ramanujan J.* 45 (3) (2018) 817–839.
- [18] E.Y.S. Liu and H.W.J. Zhang, Inequalities for the overpartition function, *Ramanujan J.* 54 (3) (2021) 485–509.
- [19] J. Lovejoy, Rank and conjugation for the Frobenius representation of an overpartition, *Ann. Comb.* 9 (3) (2005) 321–334.
- [20] J. Males, Asymptotic equidistribution and convexity for partition ranks, *Ramanujan J.* 54 (2) (2021) 397–413.
- [21] D.S. Mitrinović, Elementary Inequalities. In cooperation with E.S. Barnes, D.C.B. Marsh, J.R.M. Radok. Tutorial Text No. 1 P. Noordhoff Ltd., Groningen 1964 159 pp.