

BACKSTRÖM ALGEBRAS

YURIY DROZD

ABSTRACT. We introduce *Backström pairs* and *Backström rings*, study their derived categories and construct for them a sort of *categorical resolutions*. For the latter we define the global dimension, construct a sort of semi-orthogonal decompositions of the derived category and deduce that the derived dimension of a Backström ring is at most 2. Using this semi-orthogonal decomposition, we define a description of the module category as the category of elements of a special bimodule. We also construct a partial tilting for a Backström pair to a ring of triangular matrices and define the global dimension of the latter.

CONTENTS

Introduction	1
1. Partial tilting	3
2. Backström pairs	5
3. The structure of derived categories	10
4. Semi-orthogonal decomposition	12
5. Relation to bimodule categories	14
6. Partial tilting for Backström pairs	16
References	21

INTRODUCTION

Backström orders were introduced in [27], where it was shown that their representations are in correspondence with those of quivers or species. A special class of Backström orders are *nodal orders*, which appeared (without this name) in [13] as such pure noetherian algebras that the classification of their finitely generated modules is tame. In

2020 *Mathematics Subject Classification.* 16E35, 16D90, 18G80, 16E30, 16S38.

Key words and phrases. Backström rings, Backström pairs, Backström curves, derived categories, partial tilting.

This work was accomplished during the visits of the author to the Max-Planck-Institut of Mathematics and the University of Paderborn, and he is grateful to these institutions for their hospitality and financial support.

[6] tameness was also proved for the derived categories of nodal orders. Global analogues of nodal algebras, called *nodal curves*, were considered in [8, 15, 29]. Namely, in [8] a sort of tilting theory for such curves was developed, which related them to some quasihereditary finite dimensional algebras. In [15] a criterion was found for a nodal curve to be tame with respect to the classification of vector bundles, and in [29] it was proved that the same class of curves has tame derived categories. It was clear that the tilting theory of [8] can be extended to a general situation, namely, to *Backström curves*, i.e. non-commutative curves having Backström orders as their localizations. Nodal orders and related gentle algebras appear in studying mirror symmetry, see, for instance, [23]. Finite dimensional analogues of nodal orders, called *nodal algebras*, were introduced in [16, 30]. In the latter paper their structure was completely described. In [31] it was shown that such important classes as gentle and skewed-gentle algebras are nodal. In [9] a tilting theory was developed for nodal algebras, which was applied to the study of derived categories of gentle and skewed-gentle algebras.

This paper is devoted to a tilting theory for *Backström rings*, which are a straightforward generalization of Backström orders and algebras. In Section 1 we propose a variant of partial tilting, which generalizes the technique of minors from [10]. In Section 2 we introduce *Backström pairs* as the pairs of semiperfect rings $\mathbf{H} \supseteq \mathbf{A}$ with common radical and *Backström rings* (*piecewise Backström rings*) as the rings \mathbf{A} that occur in Backström pairs with hereditary (respectively, piecewise hereditary) \mathbf{H} . We construct the *Auslander envelope* $\tilde{\mathbf{A}}$ of a Backström pair and calculate its global dimension. Actually, this global dimension only depends on the global dimension of \mathbf{H} . In particular, Auslander envelopes for Backström rings are of global dimension at most 2. In Section 3 we apply the tilting technique to show that the derived category of the algebra \mathbf{A} is connected by recollement with the derived category of its Auslander envelope. It implies that the derived dimension of \mathbf{A} in the sense of Rouquier [28] is not greater than that of the Auslander envelope. In Section 4 we consider a recollement between the derived categories of the algebra \mathbf{H} and of the Auslander envelope. It is used to calculate the derived dimension of the Auslander envelope, thus obtaining an upper bound for the derived dimension of the algebra \mathbf{A} . In particular, we prove that the derived dimension of a Backström or piecewise Backström algebra is at most 2. Moreover, if \mathbf{A} is a Backström or piecewise Backström algebra of Dynkin type which is not piecewise hereditary of Dynkin type, then $\text{der.dim } \mathbf{A} = 1$. In Section 5 we establish an equivalence between the category $\mathcal{D}(\tilde{\mathbf{A}})$ and a

bimodule category. Such equivalence gives a useful instrument for calculations in this derived category (see, for instance, [4, 5, 6, 7, 9, 29]). In Section 6 we consider another partial tilting for the algebra $\tilde{\mathbf{A}}$ from a Backström pair, relating its derived category by a recollement to the derived category of an algebra \mathbf{B} of triangular matrices which looks simpler than the Auslander algebra. In this case we calculate explicitly the global dimension of \mathbf{B} and the kernel of the partial tilting functor $F : \mathcal{D}(\mathbf{B}) \rightarrow \mathcal{D}(\mathbf{A})$.

1. PARTIAL TILTING

Let \mathcal{T} be a triangulated category, $\mathfrak{X} \subseteq \text{Ob } \mathcal{T}$. We denote by $\text{Tri}(\mathfrak{X})$ the smallest strictly full triangulated subcategory containing \mathfrak{X} and closed under coproducts (it means that if a coproduct of objects from $\text{Tri}(\mathfrak{X})$ exists in \mathcal{T} , it belongs to $\text{Tri}(\mathfrak{X})$). For a DG-category \mathcal{R} we denote by $\mathcal{D}(\mathcal{R})$ its derived category [20]. The following result is a generalization of [24, Prop. 2.6].

Theorem 1.1. *Let \mathfrak{X} be a subset of compact objects of $\text{Ob } \mathcal{D}(\mathcal{A})$, where \mathcal{A} is a Grothendieck category. We consider the DG-category \mathcal{R} with the set of objects \mathfrak{X} and the sets of morphisms $\mathcal{R}(T, R) = \text{RHom}(T, R)$. Define the functor $F : \mathcal{D}(\mathcal{A}) \rightarrow \mathcal{D}(\mathcal{R}^{\text{op}})$ mapping a complex C to the DG-module $FC = \text{RHom}_{\mathcal{D}(\mathcal{A})}(-, C)$ restricted onto \mathfrak{X} .*

- (1) *The restriction of F onto $\text{Tri}(\mathfrak{X})$ is an equivalence $\text{Tri}(\mathfrak{X}) \rightarrow \mathcal{D}(\mathcal{R}^{\text{op}})$.*
- (2) *The functor F has a left adjoint F^* and a right adjoint $F^!$ such that both adjunction morphisms $\eta : \text{Id}_{\mathcal{D}(\mathcal{R}^{\text{op}})} \rightarrow FF^*$ and $\zeta : FF^! \rightarrow \text{Id}_{\mathcal{D}(\mathcal{A})}$ are isomorphisms.*
- (3) *There is a recollement diagram in the sense of [3]*

$$(1.1) \quad \text{Ker } F \begin{array}{c} \xleftarrow{l^*} \\ \xrightarrow{l} \\ \xleftarrow{l^!} \end{array} \mathcal{D}(\mathcal{A}) \begin{array}{c} \xleftarrow{F^*} \\ \xrightarrow{F} \\ \xleftarrow{F^!} \end{array} \mathcal{D}(\mathcal{R}^{\text{op}}),$$

where l is the embedding, l^* and $l^!$ are, respectively, its left and right adjoints.

If \mathfrak{X} generates $\mathcal{D}(\mathcal{A})$, we obtain an equivalence $\mathcal{D}(\mathcal{A}) \simeq \mathcal{D}(\mathcal{R}^{\text{op}})$, as in [24]. If \mathfrak{X} consists of one object R , we obtain an equivalence $\text{Tri}(R) \simeq \mathcal{D}(\mathbf{R}^{\text{op}})$, where $\mathbf{R} = \text{RHom}(R, R)$.

Proof. (1): We identify $\mathcal{D}(\mathcal{A})$ with the homotopy category $\mathcal{I}(\mathcal{A})$ of K -injective complexes, i.e. such complexes I that $\text{Hom}(C, I)$ is acyclic for every acyclic complex C , and suppose that $\mathfrak{X} \subseteq \mathcal{I}(c\mathcal{A})$.

Then RHom coincides with Hom within the category $\mathcal{I}(\mathcal{A})$ so, for $C \in \mathcal{I}(\mathcal{A})$, $\mathrm{FC} = \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(-, C)$ restricted onto \mathfrak{X} . The full subcategory of $\mathcal{I}(\mathcal{A})$ consisting of such complexes C that the natural map $\mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(R, C) \rightarrow \mathrm{Hom}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}(\mathrm{FR}, \mathrm{FC})$ is bijective for all $R \in \mathfrak{X}$ contains \mathfrak{X} , is strictly full, triangulated and closed under coproducts, since all objects from \mathcal{R} are compact. Therefore, it contains $\mathrm{Tri}(\mathfrak{X})$. Quite analogously, the full subcategory of such complexes C that the natural map $\mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(C, C') \rightarrow \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(\mathrm{FC}, \mathrm{FC}')$ is bijective for every $C' \in \mathrm{Tri}(\mathfrak{X})$ also contains $\mathrm{Tri}(\mathfrak{X})$. Hence the restriction of F onto $\mathrm{Tri}(\mathfrak{X})$ is fully faithful. Moreover, as the functors $\mathrm{Hom}_{\mathcal{R}}(-, R)$, where R runs through \mathfrak{X} , generate $\mathcal{D}(\mathcal{R}^{\mathrm{op}})$, the functor F is essentially surjective. Therefore, restricted to $\mathrm{Tri}(\mathfrak{X})$, it gives an equivalence $\mathrm{Tri}(\mathfrak{X}) \rightarrow \mathcal{D}(\mathcal{R})$.

(2) and (3): Note that $\mathcal{D}(\mathcal{R}^{\mathrm{op}})$ is cocomplete and compactly generated, hence satisfies the Brown representability theorem [25, Thm. 8.3.3]. Therefore, it is true for $\mathrm{Tri}(\mathfrak{X})$ too. Then [25, Prop. 9.1.19] implies that a Bousfield localization functor exists for $\mathrm{Tri}(\mathfrak{X}) \subseteq \mathcal{D}(\mathcal{A})$ and [25, Prop. 9.1.18] implies that the embedding $\mathrm{E} : \mathrm{Tri}(\mathfrak{X}) \rightarrow \mathcal{D}(\mathcal{A})$ has a right adjoint $\Theta : \mathcal{D}(\mathcal{A}) \rightarrow \mathrm{Tri}(\mathfrak{X})$. Let $\mathrm{F}' : \mathcal{D}(\mathcal{R}^{\mathrm{op}}) \rightarrow \mathrm{Tri}(\mathfrak{X})$ be a quasi-inverse to the restriction of F onto $\mathrm{Tri}(\mathfrak{X})$. In particular, F' is a left adjoint to this restriction and the adjunction $\mathrm{FF}' \rightarrow \mathrm{Id}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}$ is an isomorphism. Then $\mathrm{FC} = \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(-, C)|_{\mathfrak{X}} \simeq \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(-, \Theta C)|_{\mathfrak{X}} = \mathrm{F}\Theta C$. Set $\mathrm{F}^* = \mathrm{E}\mathrm{F}'$. Since $\mathrm{F}'M \in \mathrm{Tri}(\mathfrak{X})$ for every $M \in \mathcal{D}(\mathcal{R}^{\mathrm{op}})$,

$$\begin{aligned} \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(\mathrm{F}^*M, C) &\simeq \mathrm{Hom}_{\mathrm{Tri}(\mathfrak{X})}(\mathrm{F}'M, \Theta C) \simeq \\ &\simeq \mathrm{Hom}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}(M, \mathrm{F}\Theta C) \simeq \mathrm{Hom}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}(M, \mathrm{FC}), \end{aligned}$$

for any $M \in \mathcal{D}(\mathcal{R}^{\mathrm{op}})$ and $C \in \mathcal{I}(\mathcal{A})$. Hence F^* is a left adjoint to F . If, moreover, $C \in \mathrm{Tri}(\mathfrak{X})$, we obtain

$$\mathrm{Hom}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}(\mathrm{FF}^*M, \mathrm{FC}) \simeq \mathrm{Hom}_{\mathcal{I}(\mathcal{A})}(\mathrm{F}^*M, C) \simeq \mathrm{Hom}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}(M, \mathrm{FC}).$$

As F is essentially surjective, it implies that $\eta : \mathrm{FF}^* \rightarrow \mathrm{Id}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}$ is an isomorphism. As all objects from \mathfrak{X} are compact, F respects coproducts, hence has a right adjoint $\mathrm{F}^!$ [25, Thm. 8.4.4]. Now it follows from [10, Cor. 2.3] that $\zeta : \mathrm{FF}^! \rightarrow \mathrm{Id}_{\mathcal{D}(\mathcal{R}^{\mathrm{op}})}$ is an isomorphism and there is a recollement diagram (1.1). □

Note that $\mathrm{Im} \mathrm{F}^* = \mathrm{Tri}(\mathfrak{X})$ by construction, but usually $\mathrm{Im} \mathrm{F}^! \neq \mathrm{Tri}(\mathfrak{X})$, though it is equivalent to $\mathrm{Tri}(\mathfrak{X})$.

Corollary 1.2. *Under conditions and notations of the preceding theorem, suppose that $\mathrm{Hom}_{\mathcal{D}(\mathcal{A})}(R, T[m]) = 0$ for $R, T \in \mathfrak{X}$ and $m \neq 0$.*

Then the functor F induces an equivalence $\text{Tri}(R) \xrightarrow{\sim} \mathcal{D}(\mathcal{R}^{\text{op}})$, where \mathcal{R} is the category with the set of objects \mathfrak{R} and the sets of morphisms $\mathcal{R}(A, B) = \text{Hom}_{\mathcal{D}(\mathcal{A})}(A, B)$.

In this situation we call the functor F a *partial tilting functor*.

2. BACKSTRÖM PAIRS

Recall [2, 22] that a *semiperfect ring* is a ring \mathbf{A} such that $\mathbf{A}/\text{rad } \mathbf{A}$ is a semisimple artinian ring and idempotents can be lifted modulo $\text{rad } \mathbf{A}$. Equivalently, as a left (or as a right) \mathbf{A} -module, \mathbf{A} decomposes into a direct sum of modules with local endomorphism rings.

Definition 2.1. (1) A *Backström pair* is a pair of semiperfect rings $\mathbf{H} \supseteq \mathbf{A}$ such that $\text{rad } \mathbf{A} = \text{rad } \mathbf{H}$. We denote by $\mathbf{C}(\mathbf{H}, \mathbf{A})$ the *conductor* of \mathbf{H} in \mathbf{A} :

$$\mathbf{C}(\mathbf{H}, \mathbf{A}) = \{\alpha \in \mathbf{A} \mid \mathbf{H}\alpha \subseteq \mathbf{A}\} = \text{ann}(\mathbf{H}/\mathbf{A})_{\mathbf{A}}$$

(the right subscript \mathbf{A} means that we consider \mathbf{H}/\mathbf{A} as a right \mathbf{A} -module). Obviously, $\mathbf{C}(\mathbf{H}, \mathbf{A}) \supseteq \text{rad } \mathbf{A}$.

- (2) We call a ring \mathbf{A} a (left) *Backström ring* (*piecewise Backström ring*) if there is a Backström pair $\mathbf{H} \supseteq \mathbf{A}$, where the ring \mathbf{H} is left hereditary (respectively, \mathbf{H} is *left piecewise hereditary* [18], i.e. derived equivalent to a left hereditary ring). If, moreover, both \mathbf{A} and \mathbf{H} are finite dimensional algebras over a field \mathbb{k} , we call \mathbf{A} a *Backström algebra* (respectively, *piecewise Backström algebra*).

Remark 2.2. If e is an idempotent in \mathbf{A} , then $\text{rad}(e\mathbf{A}e) = e(\text{rad } \mathbf{A})e$, hence, if $\mathbf{H} \supseteq \mathbf{A}$ is a Backström pair, so is also $e\mathbf{H}e \supseteq e\mathbf{A}e$. It implies that if P is a finitely generated projective \mathbf{A} -module, $\mathbf{A}' = \text{End}_{\mathbf{A}} P$ and $\mathbf{H}' = \text{End}_{\mathbf{H}}(\mathbf{H} \otimes_{\mathbf{A}} P)$, then $\mathbf{H}' \supseteq \mathbf{A}'$ is also a Backström pair. Note that if \mathbf{H} is left hereditary (or piecewise hereditary), so is \mathbf{H}' , hence \mathbf{A}' is a Backström ring (piecewise Backström ring) if so is \mathbf{A} . In particular, the notion of Backström (or piecewise Backström) ring is Morita invariant. Note also that if \mathbf{H} is left hereditary and noetherian, it is also right hereditary, so \mathbf{A}^{op} is also a Backström ring (piecewise Backström ring).

Example 2.3. (1) An important example of Backström algebras are *nodal algebras* introduced in [16, 30]. By definition, they are finite dimensional algebras such that there is a Backström pair $\mathbf{H} \supseteq \mathbf{A}$, where \mathbf{H} is a hereditary algebra and $\text{length}_{\mathbf{A}}(\mathbf{H} \otimes_{\mathbf{A}} U) \leq 2$ for every simple \mathbf{A} -module U . Their structure was completely described in [30].

- (2) Recall that a \mathbb{k} -algebra \mathbf{A} is called *gentle* [1] if $\mathbf{A} \simeq \mathbb{k}\Gamma/J$, where Γ is a finite quiver (oriented graph) and J is an ideal in the path algebra $\mathbb{k}\Gamma$ such that $(J_+)^2 \supseteq J \supseteq (J_+)^k$ for some k , where J_+ is the ideal generated by all arrows, and the following conditions hold:
- (a) For every vertex $i \in \text{Ver } \Gamma$ there are at most 2 arrows starting at i and at most 2 arrows ending at i .
 - (b) If an arrow a starts at i (ends at i) and arrows b_1, b_2 end at i (respectively, start at i), then either $ab_1 = 0$ or $ab_2 = 0$ (respectively, either $b_1a = 0$ or $b_2a = 0$), but not both.
 - (c) The ideal J is generated by products of arrows of the sort ab .

It is proved in [31] that such algebras are nodal, hence Backström algebras. The same is true for skewed-gentle algebras [17] obtained from gentle algebras by blowing up some vertices.

- (3) *Backström orders* are orders \mathbf{A} over a discrete valuation ring such that there is a Backström pair $\mathbf{H} \supseteq \mathbf{A}$, where \mathbf{H} is a hereditary order. They were considered in [27].
- (4) $\Lambda_n = \mathbb{k}[x_1, x_2, \dots, x_n]/(x_1, x_2, \dots, x_n)^2$ embeds into $\mathbf{H} = \prod_{i=1}^n \mathbb{k}\Gamma_i$, where $\Gamma_i = \cdot \xrightarrow{a_i} \cdot$ (x_i maps to a_i). Obviously, under this embedding $\text{rad } \Lambda_n = \text{rad } \mathbf{H}$, so Λ_n is a Backström algebra.

We consider a fixed Backström pair $\mathbf{H} \supseteq \mathbf{A}$, set $\mathfrak{r} = \text{rad } \mathbf{A} = \text{rad } \mathbf{H}$ and denote by \mathbf{C} the conductor $\mathbf{C}(\mathbf{H}, \mathbf{A})$. Obviously, \mathbf{C} is a two-sided \mathbf{A} -ideal and the biggest left \mathbf{H} -ideal contained in \mathbf{A} . Actually, it even turns out to be a two-sided \mathbf{H} -ideal and its definition is left-right symmetric.

Lemma 2.4. *Let $R \subseteq S$ be semisimple rings, $I = \{\alpha \in R \mid S\alpha \subseteq R\}$. Then I is a two-sided S -ideal.*

Proof. Obviously, I is a left S -ideal and a two-sided R -ideal. As R is semisimple, $I = Re$ for some central idempotent $e \in R$. Then $Se \subseteq Re$, so $Se = Re$ and $(1-e)Se = 0$. Hence $eS(1-e)$ is a left ideal in S and $(eS(1-e))^2 = 0$, so $eS(1-e) = 0$ and $I = Se = eS$ is also a right S -ideal. \square

Proposition 2.5. *\mathbf{C} is a two-sided \mathbf{H} -ideal. It is the biggest \mathbf{H} -ideal contained in \mathbf{A} . Therefore, it coincides with the set $\{\alpha \in \mathbf{A} \mid \alpha\mathbf{H} \subseteq \mathbf{A}\}$ or with $\text{ann}_{\mathbf{A}}(\mathbf{H}/\mathbf{A})$ considered as left \mathbf{A} -module.*

Proof. It follows from the preceding lemma applied to the rings $\mathbf{A}/\text{rad } \mathbf{A}$ and $\mathbf{H}/\text{rad } \mathbf{H}$. \square

In what follows we assume that $\mathbf{A} \neq \mathbf{H}$, so $\mathbf{C} \neq \mathbf{A}$. To calculate \mathbf{C} , we consider a decomposition $\mathbf{A} = \bigoplus_{i=1}^m A_i$, where A_i are indecomposable projective left \mathbf{A} -modules. Arrange them so that $\mathbf{H}A_i \neq A_i$ for $1 \leq i \leq r$ and $\mathbf{H}A_i = A_i$ for $r < i \leq m$, and set $A^0 = \bigoplus_{i=1}^r A_i$, $H^0 = \mathbf{H}A^0$ and $A^1 = \bigoplus_{i=r+1}^m A_i = \mathbf{H}A^1$. Then $\mathbf{A} = A^0 \oplus A^1$ and $\mathbf{H} = H^0 \oplus A^1$ (possibly, $r = m$, so $A^0 = \mathbf{A}$ and $H^0 = \mathbf{H}$). Let $A^0 = \mathbf{A}e_0$ and $A^1 = \mathbf{A}e_1$, where e_0 and e_1 are orthogonal idempotents and $e_0 + e_1 = 1$. Set $A_b^a = e_b \mathbf{A} e_a$ and $H_b^a = e_b \mathbf{H} e_a$, where $a, b \in \{0, 1\}$. Note that $A_b^1 = H_b^1$ and $A_1^0 = H_1^0$. As A^0 and A^1 have no isomorphic direct summands, $A_b^a \subseteq \text{rad } \mathbf{A}$ if $a \neq b$. Hence, if we set $\mathfrak{r}^a = \text{rad } A^a$ ($a = 0, 1$) and consider the Pierce decomposition of the ring \mathbf{A} :

$$\mathbf{A} = \begin{pmatrix} A_0^0 & A_0^1 \\ A_0^1 & A_1^1 \end{pmatrix},$$

the Pierce decomposition of the ideal \mathfrak{r} becomes

$$\mathfrak{r} = \begin{pmatrix} \mathfrak{r}_0^0 & A_0^1 \\ A_0^1 & \mathfrak{r}_1^1 \end{pmatrix},$$

where $\mathfrak{r}_a^a = \text{rad } A_a^a$ ($a = 0, 1$). It implies, in particular, that H^0 and H^1 have no isomorphic direct summands, the Pierce decomposition of \mathbf{H} is

$$\mathbf{H} = \begin{pmatrix} H_0^0 & A_0^1 \\ A_0^1 & A_1^1 \end{pmatrix}$$

and $\mathfrak{r}_0^0 = \text{rad } H_0^0$. Now one easily sees that an element $a = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ belongs to \mathbf{C} if and only if $H^0 a \subseteq A^0$. We claim that then $H^0 a \subseteq \text{rad } A^0$. Otherwise $H^0 a$ contains an idempotent, hence a direct summand of A^0 , which is isomorphic to some A_i with $1 \leq i \leq r$. It is impossible, since $\mathbf{H}A_i \neq A_i$. Therefore, $\alpha \in \mathfrak{r}_0^0$ and we obtain the following result.

Proposition 2.6. *The Pierce decomposition of the ideal \mathbf{C} is*

$$\mathbf{C} = \begin{pmatrix} \mathfrak{r}_0^0 & A_0^1 \\ A_0^1 & A_1^1 \end{pmatrix}.$$

Definition 2.7. We define the *Auslander envelope* of the Backström pair $\mathbf{H} \supseteq \mathbf{A}$ as the ring $\tilde{\mathbf{A}}$ of 2×2 matrices of the form

$$\tilde{\mathbf{A}} = \begin{pmatrix} \mathbf{A} & \mathbf{H} \\ \mathbf{C} & \mathbf{H} \end{pmatrix}$$

Using Pierce decompositions of \mathbf{A} , \mathbf{H} and \mathbf{C} , we also present $\tilde{\mathbf{A}}$ as the ring of 4×4 matrices

$$(2.1) \quad \tilde{\mathbf{A}} = \begin{pmatrix} A_0^0 & A_0^1 & H_0^0 & A_0^1 \\ A_1^0 & A_1^1 & A_1^0 & A_1^1 \\ \mathfrak{r}_0^0 & A_0^1 & H_0^0 & A_0^1 \\ A_1^0 & A_1^1 & A_1^0 & A_1^1 \end{pmatrix}$$

We also define $\tilde{\mathbf{H}}$ as the ring of 4×4 matrices of the form

$$\tilde{\mathbf{H}} = \begin{pmatrix} \mathbf{H} & \mathbf{H} \\ \mathbf{C} & \mathbf{H} \end{pmatrix}$$

or

$$\tilde{\mathbf{H}} = \begin{pmatrix} H_0^0 & A_0^1 & H_0^0 & A_0^1 \\ A_1^0 & A_1^1 & A_1^0 & A_1^1 \\ \mathfrak{r}_0^0 & A_0^1 & H_0^0 & A_0^1 \\ A_1^0 & A_1^1 & A_1^0 & A_1^1 \end{pmatrix}$$

Obviously, $\text{rad } \tilde{\mathbf{H}} = \text{rad } \tilde{\mathbf{A}}$, so $\tilde{\mathbf{H}} \supseteq \tilde{\mathbf{A}}$ is also a Backström pair. $\tilde{\mathbf{A}}$ is left noetherian if and only if \mathbf{A} is left noetherian and \mathbf{H} is finitely generated as left \mathbf{A} -module.

In the noetherian case one can calculate the global dimensions of $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{H}}$. It turns out that it only depends on \mathbf{H} .

Theorem 2.8. *Suppose that either \mathbf{A} (hence also \mathbf{H}) is left perfect or \mathbf{A} is left noetherian and \mathbf{H} is finitely generated as left \mathbf{A} -module (hence also left noetherian). Then*

$$\begin{aligned} \text{l.gl.dim } \tilde{\mathbf{A}} &= 1 + \max(1 + \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0, \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^1) = \\ &= \begin{cases} 1 + \text{l.gl.dim } \mathbf{H} & \text{if } \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0 \geq \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^1, \\ \text{l.gl.dim } \mathbf{H} & \text{if } \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0 < \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^1 \end{cases} \end{aligned}$$

and

$$\text{l.gl.dim } \tilde{\mathbf{H}} = \text{l.gl.dim } \mathbf{H}.$$

In particular, if \mathbf{A} is a Backström ring, so is also $\tilde{\mathbf{A}}$, and if \mathbf{A} is not left hereditary, then $\text{l.gl.dim } \tilde{\mathbf{A}} = 2$.

For instance, it is the case for nodal (in particular, gentle or skewed-gentle) algebras (Example 2.3).

Proof. Under these conditions $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{H}}$ are either left perfect or left noetherian. We recall that if a ring Λ is left perfect or left noetherian and semiperfect, then $\text{l.gl.dim } \Lambda = \text{pr.dim}_\Lambda(\Lambda/\text{rad } \Lambda) = 1 + \text{pr.dim}_\Lambda \text{rad } \Lambda$. The 4×4 matrix presentation (2.1) of $\tilde{\mathbf{A}}$ implies that the corresponding presentation of $\text{rad } \tilde{\mathbf{A}}$ is

$$(2.2) \quad \text{rad } \tilde{\mathbf{A}} = \begin{pmatrix} \mathfrak{r}_0^0 & A_0^1 & H_0^0 & A_0^1 \\ A_1^0 & \mathfrak{r}_1^1 & A_1^0 & \mathfrak{r}_1^1 \\ \mathfrak{r}_0^0 & A_0^1 & \mathfrak{r}_0^0 & A_0^1 \\ A_1^0 & \mathfrak{r}_1^1 & A_1^0 & \mathfrak{r}_1^1 \end{pmatrix}$$

An $\tilde{\mathbf{A}}$ -module M is given by a quadruple (M', M'', ϕ, ψ) , where M' is an \mathbf{A} -module, M'' is an \mathbf{H} -module, $\psi : M'' \rightarrow M'$ is a homomorphism of \mathbf{A} -modules and $\phi : \mathbf{C} \otimes_{\mathbf{A}} M' \rightarrow M''$ is a homomorphism of \mathbf{H} -modules. We usually write $M = \begin{pmatrix} M' \\ M'' \end{pmatrix}$ not mentioning ϕ and ψ . For an \mathbf{H} -module N we define the $\tilde{\mathbf{A}}$ -module $N^+ = \begin{pmatrix} N \\ N \end{pmatrix}$. Then $N \mapsto N^+$ is an exact functor mapping projective modules to projective ones.

We denote by L^i and by R^i , respectively, the i -th column of the presentations (2.1) and (2.2). Then $R^1 = (\mathfrak{r}^0)^+$ and $R^2 = R^4 = (\mathfrak{r}^1)^+$, where $\mathfrak{r}^a = \mathfrak{r}e_a$. If

$$\cdots \rightarrow F_k \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow N \rightarrow 0$$

is a minimal projective resolution of an \mathbf{H} -module N ,

$$\cdots \rightarrow F_k^+ \rightarrow \cdots \rightarrow F_1^+ \rightarrow F_0^+ \rightarrow N^+ \rightarrow 0$$

is a minimal projective resolution of N^+ , so $\text{pr.dim}_{\tilde{\mathbf{A}}} N^+ = \text{pr.dim}_{\mathbf{H}} N$. In particular, $\text{pr.dim}_{\tilde{\mathbf{A}}} R^1 = \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0$ and $\text{pr.dim}_{\tilde{\mathbf{A}}} R^2 = \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^1$. For the module R^3 we have an exact sequence

$$(2.3) \quad 0 \rightarrow (\mathfrak{r}^0)^+ \rightarrow R^3 \rightarrow \begin{pmatrix} H^0/\mathfrak{r}^0 \\ 0 \end{pmatrix} \rightarrow 0.$$

Note that H^0/\mathfrak{r}^0 is a semisimple \mathbf{A} -module and $e_1(H^0/\mathfrak{r}^0) = 0$, hence it contains the same simple direct summands as $\mathbf{A}^0/\mathfrak{r}^0$. The same is true for $\begin{pmatrix} H^0/\mathfrak{r}^0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} A^0/\mathfrak{r}^0 \\ 0 \end{pmatrix} = L^1/R^1$. Hence

$$\text{pr.dim}_{\tilde{\mathbf{A}}} \begin{pmatrix} H^0/\mathfrak{r}^0 \\ 0 \end{pmatrix} = 1 + \text{pr.dim}_{\tilde{\mathbf{A}}} R^1 = 1 + \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0.$$

Therefore, the exact sequence (2.3) shows that $\text{pr.dim}_{\tilde{\mathbf{A}}} R^3 = 1 + \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0$ and

$$\text{pr.dim}_{\tilde{\mathbf{A}}} \text{rad } \tilde{\mathbf{A}} = \max(1 + \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^0, \text{pr.dim}_{\mathbf{H}} \mathfrak{r}^1),$$

which gives the necessary result for $\tilde{\mathbf{A}}$. On the other hand, R^3 is a projective $\tilde{\mathbf{H}}$ -module, whence $\text{l.gl.dim } \tilde{\mathbf{H}} = \text{l.gl.dim } \mathbf{H}$. \square

3. THE STRUCTURE OF DERIVED CATEGORIES

In what follows we denote by $\mathcal{D}(\mathbf{A})$ the derived category $\mathcal{D}(\mathbf{A}\text{-Mod})$. We denote by $\mathcal{D}_f(\mathbf{A})$ the full subcategory of $\mathcal{D}(\mathbf{A})$ consisting of complexes quasi-isomorphic to complexes of finitely generated projective modules. If \mathbf{A} is left noetherian, it coincides with the derived category of the category $\mathbf{A}\text{-mod}$ of finitely generated \mathbf{A} -modules. We also use the usual superscripts $+$, $-$, b . By $\text{Perf}(\mathbf{A})$ we denote the full subcategory of perfect complexes from $\mathcal{D}(\mathbf{A})$, i.e. complexes quasi-isomorphic to finite complexes of finitely generated projective modules. It coincides with the full subcategory of compact objects in $\mathcal{D}(\mathbf{A})$ [28]. If \mathbf{A} is left noetherian, an \mathbf{A} -module M belongs to $\text{Perf}(\mathbf{A})$ if and only if it is finitely generated and of finite projective dimension.

There are close relations between the categories $\mathcal{D}(\mathbf{A})$, $\mathcal{D}(\mathbf{H})$ and $\mathcal{D}(\tilde{\mathbf{A}})$ based on the following construction [10].

Let $\mathbf{P} = \begin{pmatrix} \mathbf{A} \\ \mathbf{C} \end{pmatrix}$. It is a projective $\tilde{\mathbf{A}}$ -module and $\text{End } \mathbf{P} \simeq \mathbf{A}^{\text{op}}$, so it can be considered as a right \mathbf{A} -module. Consider the functors

$$\begin{aligned} \mathbf{F} &= \text{Hom}_{\tilde{\mathbf{A}}}(\mathbf{P}, -) \simeq \mathbf{P}^\vee \otimes_{\tilde{\mathbf{A}}} - : \tilde{\mathbf{A}}\text{-Mod} \rightarrow \mathbf{A}\text{-Mod}, \\ \mathbf{F}^* &= \mathbf{P} \otimes_{\mathbf{A}} - : \mathbf{A}\text{-Mod} \rightarrow \tilde{\mathbf{A}}\text{-Mod}, \\ \mathbf{F}^\dagger &= \text{Hom}_{\mathbf{A}}(\mathbf{P}^\vee, -) : \mathbf{A}\text{-Mod} \rightarrow \tilde{\mathbf{A}}\text{-Mod}, \end{aligned}$$

where $\mathbf{P}^\vee = \text{Hom}_{\tilde{\mathbf{A}}}(\mathbf{P}, \tilde{\mathbf{A}}) \simeq (\mathbf{A} \ \mathbf{H})$ is the dual right projective $\tilde{\mathbf{A}}$ -module. The functor \mathbf{F} is exact, \mathbf{F}^* is its left adjoint and \mathbf{F}^\dagger is its right adjoint. Moreover, the adjunction morphisms $\mathbf{F}\mathbf{F}^* \rightarrow \text{Id}_{\tilde{\mathbf{A}}\text{-Mod}}$ and $\text{Id}_{\mathbf{A}\text{-Mod}} \rightarrow \mathbf{F}^\dagger\mathbf{F}$ are isomorphisms [10, Thm. 4.3]. The functors \mathbf{F}^* and \mathbf{F}^\dagger are fully faithful and \mathbf{F} is essentially surjective, i.e. every \mathbf{A} -module is isomorphic to $\mathbf{F}M$ for some $\tilde{\mathbf{A}}$ -module M . $\text{Ker } \mathbf{F}$ is a Serre subcategory of $\tilde{\mathbf{A}}\text{-Mod}$ equivalent to $\tilde{\mathbf{H}}\text{-Mod}$, where $\tilde{\mathbf{H}} = \mathbf{H}/\mathbf{C} \simeq \tilde{\mathbf{A}} / \begin{pmatrix} \mathbf{A} & \mathbf{H} \\ \mathbf{C} & \mathbf{C} \end{pmatrix}$. The embedding functor $\mathbf{l} : \text{Ker } \mathbf{F} \rightarrow \tilde{\mathbf{A}}\text{-Mod}$ has a left adjoint \mathbf{l}^* and a right adjoint \mathbf{l}^\dagger and we obtain a recollement diagram

$$\begin{array}{ccccc} & \xleftarrow{\mathbf{l}^*} & & \xleftarrow{\mathbf{F}^*} & \\ \text{Ker } \mathbf{F} & \xrightarrow{\mathbf{l}} & \tilde{\mathbf{A}}\text{-Mod} & \xrightarrow{\mathbf{F}} & \mathbf{A}\text{-Mod} \\ & \xleftarrow{\mathbf{l}^\dagger} & & \xleftarrow{\mathbf{F}^\dagger} & \end{array}$$

As the functor \mathbf{F} is exact, it extends to the functor between the derived categories $\mathbf{DF} : \mathcal{D}(\tilde{\mathbf{A}}) \rightarrow \mathcal{D}(\mathbf{A})$ acting on complexes componentwise. The derived functors \mathbf{LF}^* and \mathbf{RF}^\dagger are, respectively, its left and right

adjoints, the adjunction morphisms $\text{Id}_{\mathcal{D}(\mathbf{A})} \rightarrow \text{DF} \cdot \text{LF}^*$ and $\text{DF} \cdot \text{LF}^* \rightarrow \text{Id}_{\mathcal{D}(\mathbf{A})}$ are again isomorphisms and we have a recollement diagram

$$\begin{array}{ccccc} & \xleftarrow{\text{LI}^*} & & \xleftarrow{\text{LF}^*} & \\ \text{Ker DF} & \xrightarrow{\text{DI}} & \mathcal{D}(\tilde{\mathbf{A}}) & \xrightarrow{\text{DF}} & \mathcal{D}(\mathbf{A}) \\ & \xleftarrow{\text{RI}^*} & & \xleftarrow{\text{RF}^*} & \end{array}$$

(It also follows from Corollary 1.2.) Here $\text{Ker DF} = \mathcal{D}_{\bar{\mathbf{H}}}(\tilde{\mathbf{A}})$, the full subcategory of complexes whose cohomologies are $\bar{\mathbf{H}}$ -modules, i.e. are annihilated by the ideal $\begin{pmatrix} \mathbf{A} & \mathbf{H} \\ \mathbf{C} & \mathbf{C} \end{pmatrix}$. Note that, as a rule, it is not equivalent to $\mathcal{D}(\bar{\mathbf{H}})$. From the definition of F it follows that

$$\text{Ker DF} = \mathbf{P}^\perp = \{C \in \mathcal{D}(\tilde{\mathbf{A}}) \mid \text{Hom}_{\mathcal{D}(\tilde{\mathbf{A}})}(\mathbf{P}, C[k])\} = 0 \text{ for all } k.$$

Obviously, DF maps $\mathcal{D}^\sigma(\tilde{\mathbf{A}})$ to $D^\sigma(\mathbf{A})$ for $\sigma \in \{+, -, b\}$, LF^* maps $\mathcal{D}^-(\mathbf{A})$ to $\mathcal{D}^-(\tilde{\mathbf{A}})$ and RF^1 maps $\mathcal{D}^+(\mathbf{A})$ to $\mathcal{D}^+(\tilde{\mathbf{A}})$. If $\tilde{\mathbf{A}}$ is left noetherian, DF maps $\mathcal{D}_f(\tilde{\mathbf{A}})$ to $\mathcal{D}_f(\mathbf{A})$ and LF^* maps $\mathcal{D}_f(\mathbf{A})$ to $\mathcal{D}_f(\tilde{\mathbf{A}})$. Finally, both DF and LF^* map compact objects (i.e. perfect complexes) to compact ones, since they have right adjoints. On the contrary, usually LF^* does not map $\mathcal{D}^b(\mathbf{A})$ to $\mathcal{D}^b(\tilde{\mathbf{A}})$. For instance, it is definitely so if $\text{l.gl.dim } \tilde{\mathbf{A}} < \infty$ while $\text{l.gl.dim } \mathbf{A} = \infty$ as in Example 2.3(4). If $\text{l.gl.dim } \mathbf{H}$ is finite, so is $\text{l.gl.dim } \tilde{\mathbf{A}}$, thus this recollement can be considered as a sort of categorical resolution of the category $\mathcal{D}(\mathbf{A})$. In any case, it is useful for studying the categories $\mathbf{A}\text{-Mod}$ and $\mathcal{D}(\mathbf{A})$ if we know the structure of the categories $\tilde{\mathbf{A}}\text{-Mod}$ and $\mathcal{D}(\tilde{\mathbf{A}})$. For instance, it is so if we are interested in the *derived dimension*, i.e. the dimension of the category $\mathcal{D}_f^b(\mathbf{A})$ in the sence of Rouquier [28]. We recall the definition.

Definition 3.1. Let \mathcal{T} be a triangular category and \mathfrak{M} be a set of objects from \mathcal{T} .

- (1) We denote by $\langle \mathfrak{M} \rangle$ the smallest full subcategory of \mathcal{T} containing \mathfrak{M} and closed under direct sums, direct summands and shifts (not closed under cones, so not a triangular subcategory).
- (2) If \mathfrak{N} is another subset of \mathcal{T} , we denote by $\mathfrak{M} \dagger \mathfrak{N}$ the set of objects C from \mathcal{T} such that there is an exact triangle $A \rightarrow B \rightarrow C \xrightarrow{+}$, where $A \in \mathfrak{M}$, $B \in \mathfrak{N}$.
- (3) We define $\langle \mathfrak{M} \rangle_k$ recursively, setting $\langle \mathfrak{M} \rangle_1 = \langle \mathfrak{M} \rangle$ and $\langle \mathfrak{M} \rangle_{k+1} = \langle \langle \mathfrak{M} \rangle \dagger \langle \mathfrak{M} \rangle_k \rangle$.
- (4) The *dimension* $\dim \mathcal{T}$ of \mathcal{T} is the smallest k such that there is a finite set of objects \mathfrak{M} such that $\langle \mathfrak{M} \rangle_{k+1} = \mathcal{T}$ (if it exists). We call the dimension $\dim \mathcal{D}_f^b(\mathbf{A})$ the *derived dimension* of the ring \mathbf{A} and denote it by $\text{der.dim } \mathbf{A}$.

As the functor F is exact and essentially surjective, the next result is evident.

Proposition 3.2. $\text{der.dim } \mathbf{A} \leq \text{der.dim } \tilde{\mathbf{A}}$. *Namely, if $\mathcal{D}_f^b(\tilde{\mathbf{A}}) = \langle \mathfrak{M} \rangle_{k+1}$, then $\mathcal{D}_f^b(\mathbf{A}) = \langle \text{DF}(\mathfrak{M}) \rangle_{k+1}$.*

4. SEMI-ORTHOGONAL DECOMPOSITION

There is another recollement diagram for $\mathcal{D}(\tilde{\mathbf{A}})$ related to the projective module $\mathbf{Q} = \begin{pmatrix} \mathbf{H} \\ \mathbf{H} \end{pmatrix}$ with $\text{End } \mathbf{Q} \simeq \mathbf{H}^{\text{op}}$. Namely, we set

$$\begin{aligned} \mathbf{G} &= \text{Hom}_{\tilde{\mathbf{A}}}(\mathbf{Q}, -) \simeq \mathbf{Q}^\vee \otimes_{\tilde{\mathbf{A}}} - : \tilde{\mathbf{A}}\text{-Mod} \rightarrow \mathbf{H}\text{-Mod}, \\ \mathbf{G}^* &= \mathbf{Q} \otimes_{\mathbf{H}} - : \mathbf{H}\text{-Mod} \rightarrow \tilde{\mathbf{A}}\text{-Mod}, \\ \mathbf{G}^\dagger &= \text{Hom}_{\mathbf{H}}(\mathbf{Q}^\vee, -) : \mathbf{H}\text{-Mod} \rightarrow \tilde{\mathbf{A}}\text{-Mod}, \end{aligned}$$

where $\mathbf{Q}^\vee = \text{Hom}_{\tilde{\mathbf{A}}}(\mathbf{Q}, \tilde{\mathbf{A}}) \simeq (\mathbf{C} \ \mathbf{H})$,

$$\begin{aligned} \text{DG} &: \mathcal{D}(\tilde{\mathbf{A}}) \rightarrow \mathcal{D}(\mathbf{H}) \text{ is } \mathbf{G} \text{ applied componentwise,} \\ \text{LG}^* &: \mathcal{D}(\mathbf{A}) \rightarrow \mathcal{D}(\tilde{\mathbf{A}}) \text{ is the left adjoint of } \text{DG}, \\ \text{RG}^\dagger &: \mathcal{D}(\mathbf{A}) \rightarrow \mathcal{D}(\tilde{\mathbf{A}}) \text{ is the right adjoint of } \text{DG}. \end{aligned}$$

We also set $\bar{\mathbf{A}} = \mathbf{A}/\mathbf{C} \simeq \tilde{\mathbf{A}}/(\begin{smallmatrix} \mathbf{C} \\ \mathbf{C} \end{smallmatrix} \ \mathbf{H})$. Then we have recollement diagrams

$$\begin{array}{ccccc} & \begin{array}{c} \leftarrow \text{J}^* \rightarrow \\ \leftarrow \text{J} \rightarrow \\ \leftarrow \text{J}^\dagger \rightarrow \end{array} & \tilde{\mathbf{A}}\text{-Mod} & \begin{array}{c} \leftarrow \text{G}^* \rightarrow \\ \leftarrow \text{G} \rightarrow \\ \leftarrow \text{G}^\dagger \rightarrow \end{array} & \mathbf{H}\text{-Mod} \\ \text{Ker } \mathbf{G} & \xrightarrow{\quad} & & \xrightarrow{\quad} & \end{array}$$

and

$$\begin{array}{ccccc} & \begin{array}{c} \leftarrow \text{LJ}^* \rightarrow \\ \leftarrow \text{DJ} \rightarrow \\ \leftarrow \text{RJ}^\dagger \rightarrow \end{array} & \mathcal{D}(\tilde{\mathbf{A}}) & \begin{array}{c} \leftarrow \text{LG}^* \rightarrow \\ \leftarrow \text{DG} \rightarrow \\ \leftarrow \text{RG}^\dagger \rightarrow \end{array} & \mathcal{D}(\mathbf{H}), \\ \text{Ker } \text{DG} & \xrightarrow{\quad} & & \xrightarrow{\quad} & \end{array}$$

where $\text{Ker } \mathbf{G} \simeq \bar{\mathbf{A}}\text{-Mod}$. Since the $\tilde{\mathbf{A}}$ -ideal $(\mathbf{C} \ \mathbf{H})$ is projective as right $\tilde{\mathbf{A}}$ -module, [10, Thm. 4.6] implies that $\text{Ker } \text{DG} \simeq \mathcal{D}(\bar{\mathbf{A}})$.

As usually, this recollement diagram gives semi-orthogonal decompositions [10, Cor. 2.6]

$$(4.1) \quad \mathcal{D}(\tilde{\mathbf{A}}) = (\text{Ker } \text{DG}, \text{Im } \text{LG}^*) = (\text{Im } \text{RG}^\dagger, \text{Ker } \text{DG})$$

with $\text{Ker } \text{DG} \simeq \mathcal{D}(\bar{\mathbf{A}})$ and $\text{Im } \text{LG}^* \simeq \text{Im } \text{RG}^\dagger \simeq \mathcal{D}(\mathbf{H})$ (though usually $\text{Im } \text{LG}^* \neq \text{Im } \text{RG}^\dagger$).

Recall that a semi-orthogonal decomposition $\mathcal{S} = (\mathcal{S}_1, \mathcal{S}_2)$ means that $\text{Hom}_{\mathcal{S}}(T_2, T_1) = 0$ if $T_i \in \mathcal{T}_i$ and for every object $T \in \mathcal{S}$ there is

an exact triangle $T_1 \rightarrow T_2 \rightarrow T \xrightarrow{+}$, where $T_i \in \mathcal{T}_i$. Then the following fact holds.

Lemma 4.1. *If $\mathcal{T} = (\mathcal{T}_1, \mathcal{T}_2)$ is a semi-orthogonal decomposition of a triangulated category \mathcal{T} , then*

$$\dim \mathcal{T} \leq \dim \mathcal{T}_1 + \dim \mathcal{T}_2 + 1.$$

Proof. First we show that for any subsets $\mathfrak{M}, \mathfrak{N}$ of objects of the category \mathcal{T}

$$(4.2) \quad \begin{aligned} \langle \mathfrak{M} \rangle_{k+1} \dagger \mathfrak{N} &\subseteq \langle \mathfrak{M} \rangle \dagger \langle \langle \mathfrak{M} \rangle_k \dagger \mathfrak{N} \rangle \subseteq \\ &\subseteq \underbrace{\langle \mathfrak{M} \rangle \dagger \langle \langle \mathfrak{M} \rangle \dagger \langle \langle \mathfrak{M} \rangle \dagger \dots \langle \langle \mathfrak{M} \rangle \dagger \mathfrak{N} \rangle \dots \rangle}_{k+1} \end{aligned}$$

Indeed, let $C \in \langle \mathfrak{M} \rangle_{k+1} \dagger \mathfrak{N}$, i.e there is an exact triangle $A \rightarrow B \rightarrow C \xrightarrow{+}$, where $A \in \langle \mathfrak{M} \rangle_{k+1}, B \in \mathfrak{N}$. There is also an exact triangle $A_1 \rightarrow A \rightarrow A_2 \xrightarrow{+}$, where $A_1 \in \langle \mathfrak{M} \rangle_k, A_2 \in \langle \mathfrak{M} \rangle$. The octahedron axiom implies that there are exact triangles $A_1 \rightarrow B \rightarrow B' \xrightarrow{+}$ and $A_2 \rightarrow B' \rightarrow C \xrightarrow{+}$. Therefore, $B' \in \langle \mathfrak{M} \rangle_k \dagger \mathfrak{N}$ and $C \in \langle \mathfrak{M} \rangle \dagger \langle \langle \mathfrak{M} \rangle_k \dagger \mathfrak{N} \rangle$.

Now, let $\langle \mathfrak{M} \rangle_{k+1} = \mathcal{T}_1$ and $\langle \mathfrak{N} \rangle_{l+1} = \mathcal{T}_2$. Then, for every $T \in \mathcal{T}$ there is an exact triangle $T_1 \rightarrow T_2 \rightarrow T \xrightarrow{+}$, where $T_1 \in \langle \mathfrak{M} \rangle_{k+1}, T_2 \in \langle \mathfrak{N} \rangle_{l+1}$. But, according to (4.2), $\langle \mathfrak{M} \rangle_{k+1} \dagger \langle \mathfrak{N} \rangle_{l+1} \subseteq \langle \mathfrak{M} \cup \mathfrak{N} \rangle_{k+l+2}$, so $\mathcal{T} = \langle \mathfrak{M} \cup \mathfrak{N} \rangle_{k+l+2}$ and $\dim \mathcal{T} \leq k + l + 1$. \square

As $\bar{\mathbf{A}}$ is semisimple, any indecomposable object from $\mathcal{D}(\bar{\mathbf{A}})$ is just a shifted simple module, so $\mathcal{D}_f^b(\bar{\mathbf{A}}) = \langle \bar{\mathbf{A}} \rangle$ and $\dim \mathcal{D}^b(\bar{\mathbf{A}}) = 0$. If \mathbf{H} is hereditary, every indecomposable object from $\mathcal{D}_f^b(\mathbf{H})$ is a shift of a module. For every module M there is an exact sequence $0 \rightarrow P' \rightarrow P \rightarrow M \rightarrow 0$ with projective modules P, P' and, since \mathbf{H} is semiperfect, every indecomposable projective \mathbf{H} -module is a direct summand of \mathbf{H} . Hence $\mathcal{D}_f^b(\mathbf{H}) = \langle \mathbf{H} \rangle_2$ and $\text{der.dim } \mathbf{H} \leq 1$.

Corollary 4.2. *$\text{der.dim } \mathbf{A} \leq \text{der.dim } \mathbf{H} + 1$. In particular, if \mathbf{A} is a Backström (or piecewise Backström) ring, $\text{der.dim } \mathbf{A} \leq 2$.*

A finite dimensional hereditary algebra is said to be of *Dynkin type* if it has finitely many isomorphism classes of indecomposable modules. Such algebras, up to Morita equivalence, correspond to Dynkin diagrams [12]. If the derived category of an algebra \mathbf{H} is equivalent to the derived category of a hereditary algebra of Dynkin type, we say that \mathbf{H} is *piecewise hereditary of Dynkin type*.¹ We say that a Backström (or piecewise Backström) algebra \mathbf{A} is of *Dynkin type* if

¹It is proved in [18] that piecewise hereditary algebras of Dynkin type are just iterated tilted algebras of Dynkin type.

there is a Backström pair $\mathbf{H} \supseteq \mathbf{A}$, where \mathbf{H} is a hereditary (piecewise hereditary) algebra of Dynkin type. For instance, it is so if \mathbf{A} is a gentle or skewed-gentle algebra [31], or if it is the algebra Λ_n from Example 2.3(4). In this case $\mathcal{D}_f^b(\mathbf{H}) = \langle M_1, M_2, \dots, M_m \rangle_1$, where M_1, M_2, \dots, M_m are all pairwise non-isomorphic indecomposable \mathbf{H} -modules, so $\text{der.dim } \mathbf{H} = 0$.

In [11] it was proved that $\text{der.dim } \mathbf{A} = 0$ for a finite dimensional algebra \mathbf{A} if and only if \mathbf{A} is a piecewise hereditary algebra of Dynkin type.

Corollary 4.3. *If \mathbf{A} is a Backström (or piecewise Backström) algebra of Dynkin type (for instance, gentle or skewed-gentle), but is not piecewise hereditary of Dynkin type, then $\text{der.dim } \mathbf{A} = 1$.*

Example 4.4. The path algebra of the commutative quiver

$$\begin{array}{ccccc}
 & & 2 & & 4 \\
 & \nearrow^{\alpha_0} & & \searrow^{\alpha_1} & \\
 1 & & & & 3 \\
 & \searrow_{\beta_0} & & \nearrow_{\beta_1} & \\
 & & 2' & & 4' \\
 & & & & \searrow_{\gamma'} \\
 & & & & 4
 \end{array}
 \qquad \alpha_1 \alpha_0 = \beta_1 \beta_0$$

is a tilted (hence piecewise hereditary) algebra of type \tilde{D}_5 . At the same time it is Backström algebra of type A_4 . Namely, it is a skewed-gentle algebra obtained from the path algebra of the quiver $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ by blowing up the vertices 2 and 4.

5. RELATION TO BIMODULE CATEGORIES

Let \mathcal{A} and \mathcal{B} be additive categories, \mathcal{U} be an \mathcal{A} - \mathcal{B} -bimodule, i.e. a biadditive functor $\mathcal{A}^{\text{op}} \times \mathcal{B} \rightarrow \mathbf{Ab}$. Recall [12] that the *bimodule category* or the *category of elements* of the bimodule \mathcal{U} is the category $\text{El}(\mathcal{U})$ with the set of objects $\bigcup_{A \in \mathcal{A}, B \in \mathcal{B}} \mathcal{U}(A, B)$ and the set of morphisms $u \rightarrow v$, where $u \in \mathcal{U}(A, B), v \in \mathcal{U}(A', B')$ being the set of pairs $\{(\alpha, \beta) \mid \alpha : A' \rightarrow A, \beta : B \rightarrow B', u\alpha = \beta v\}$. Here we write, as usually, $u\alpha$ and βv , respectively, instead of $\mathcal{U}(\alpha, 1_B)u$ and $\mathcal{U}(1_{A'}, \beta)v$. Bimodule categories appear when there is a semi-orthogonal decomposition of a triangulated category.

Theorem 5.1. *Let $(\mathcal{A}, \mathcal{B})$ be a semiorthogonal decomposition of a triangulated category \mathcal{C} . Consider the \mathcal{A} - \mathcal{B} -bimodule \mathcal{U} such that $\mathcal{U}(A, B) = \text{Hom}_{\mathcal{C}}(A, B)$ ($A \in \mathcal{A}, B \in \mathcal{B}$). For every $f : A \rightarrow B$ fix a cone Cf , that is an exact triangle $A \xrightarrow{f} B \xrightarrow{f_1} Cf \xrightarrow{f_2} A[1]$. The map $f \mapsto Cf$ induces an equivalence of categories $\mathbf{C} : \text{El}(\mathcal{U}) \xrightarrow{\sim} \mathcal{C}/\mathcal{J}$, where \mathcal{J} is the ideal of \mathcal{C} consisting of morphisms η such that there are factorizations $\eta = \eta'\xi = \zeta\eta''$, where the source of η' is in \mathcal{A} and*

the target of η'' is in \mathcal{B} . Moreover, $\mathcal{J}^2 = 0$, so \mathbf{C} induces a bijection between isomorphism classes of objects from $\text{El}(\mathcal{U})$ and from \mathcal{C} .²

Proof. As $(\mathcal{A}, \mathcal{B})$ is a semi-orthogonal decomposition of \mathcal{C} , every object from \mathcal{C} occurs in an exact triangle $A \rightarrow B \rightarrow C \xrightarrow{+}$, where $A \in \mathcal{A}, B \in \mathcal{B}$, so f is an object from $\text{El}(\mathcal{U})$ and $C \simeq Cf$. Let $f' : A' \rightarrow B'$ be another object of $\text{El}(\mathcal{U})$ and $(\alpha, \beta) : f \rightarrow f'$ be a morphism from $\text{El}(\mathcal{U})$. Fix a commutative diagram

$$(5.1) \quad \begin{array}{ccccccc} A & \xrightarrow{f} & B & \xrightarrow{f_1} & Cf & \xrightarrow{f_2} & A[1] \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \alpha[1] \downarrow \\ A' & \xrightarrow{f'} & B' & \xrightarrow{f'_1} & Cf' & \xrightarrow{f'_2} & A'[1] \end{array}$$

It exists, though is not unique. Let γ' be another morphism making the diagram (5.1) commutative and set $\eta = \gamma - \gamma'$. Then $\eta f_1 = 0$, hence η factors through f_2 , and $f'_2 \eta = 0$, hence η factors through f'_1 . Thus $\eta \in \mathcal{J}$. On the other hand, if $\eta : Cf \rightarrow Cf'$ is in \mathcal{J} , the decomposition $\eta = \eta' \xi$ implies that $\eta f_1 = \eta' \xi f_1 = 0$ and the decomposition $\eta = \zeta \eta''$ implies that $\alpha[1] \eta = \alpha[1] \zeta \eta'' = 0$, hence the morphism $\gamma' = \gamma + \eta$ makes the diagram (5.1) commutative. Therefore, the class $\mathbf{C}(\alpha, \beta)$ of γ modulo \mathcal{J} is uniquely defined, so the maps $f \mapsto Cf$ and $(\alpha, \beta) \mapsto \mathbf{C}(\alpha, \beta)$ define a functor $\mathbf{C} : \text{El}(\mathcal{U}) \rightarrow \mathcal{C}/\mathcal{J}$.

Let now $\gamma : Cf \rightarrow Cf'$ be any morphism. Then $f'_2 \gamma f_1 = 0$, so $\gamma f_1 = f'_1 \beta$ for some $\beta : B \rightarrow B'$. Hence there is a morphism $\alpha : A \rightarrow A'$ making the diagram (5.1) commutative, i.e. defining a morphism $(\alpha, \beta) : f \rightarrow f'$ such that $\gamma \equiv \mathbf{C}(\alpha, \beta) \pmod{\mathcal{J}}$. If (α', β') is another such morphism, $f'_1(\beta - \beta') = 0$, so $\beta - \beta' = f' \xi$ for some $\xi : B \rightarrow A$. But $\xi = 0$, so $\beta = \beta'$. In the same way $\alpha = \alpha'$. Hence the functor \mathbf{C} is fully faithful. As we have already noticed, it is essentially surjective, therefore defines an equivalence $\text{El}(\mathcal{U}) \xrightarrow{\sim} \mathcal{C}/\mathcal{J}$. The equality $\mathcal{J}^2 = 0$ follows immediately from the definition and the conditions of the theorem. \square

We apply Theorem 5.1 to Backström pairs $\mathbf{H} \subseteq \mathbf{A}$ such that \mathbf{A} is left noetherian and \mathbf{H} is left hereditary and finitely generated as left \mathbf{A} -module. For instance, it is so in the case of Backström algebras or Backström orders. Then the ring $\tilde{\mathbf{A}}$ is also noetherian and \mathbf{C} is projective as left \mathbf{H} -module. According to (4.1), $(\text{Ker DG}, \text{Im LG}^*)$ is a semi-orthogonal decomposition of $\mathcal{D}(\tilde{\mathbf{A}})$. Moreover, both \mathbf{G} and \mathbf{G}^* map finitely generated modules to finitely generated, so the same is

²This theorem is actually a partial case of [14, Thm. 1.1].

valid if we consider their restrictions onto $\mathcal{D}_f(\tilde{\mathbf{A}})$ and $\mathcal{D}_f(\mathbf{H})$. Note also that \mathbf{G}^* is exact, so \mathbf{G}^* can be applied to complexes componentwise. The $\tilde{\mathbf{A}}$ -module \mathbf{G}^*M can be identified with the module of columns $M^+ = \begin{pmatrix} M \\ M \end{pmatrix}$ with the action of $\tilde{\mathbf{A}}$ given by the matrix multiplication. It gives an equivalence of $\mathcal{D}(\mathbf{H})$ with $\text{Im } \mathbf{L}\mathbf{G}^*$. As \mathbf{H} is left hereditary, every complex from $\mathcal{D}(\mathbf{H})$ is equivalent to a direct sum of shifted modules (see [21, Sec. 2.5]). On the other hand, $\text{Ker } \mathbf{D}\mathbf{G} \simeq \mathcal{D}(\bar{\mathbf{A}})$ and $\bar{\mathbf{A}}$ is semisimple, so every complex from $\mathcal{D}(\bar{\mathbf{A}})$ is isomorphic to a direct sum of shifted simple $\bar{\mathbf{A}}$ -modules. So, to calculate the bimodule \mathcal{U} , we only have to calculate $\text{Ext}_{\bar{\mathbf{A}}}^i(\bar{\mathbf{A}}, M^+)$, where M is an \mathbf{H} -module. Note also that \mathbf{C}^+ is a projective $\tilde{\mathbf{A}}$ -module, since \mathbf{C} is a projective \mathbf{H} -module. Therefore, a projective resolution of $\bar{\mathbf{A}}$ is $0 \rightarrow \mathbf{C}^+ \xrightarrow{\varepsilon} \mathbf{P} \rightarrow \bar{\mathbf{A}} \rightarrow 0$ and $\text{pr.dim}_{\bar{\mathbf{A}}} \bar{\mathbf{A}} = 1$. Hence we only have to calculate $\text{Hom}_{\bar{\mathbf{A}}}(\bar{\mathbf{A}}, M^+)$ and $\text{Ext}_{\bar{\mathbf{A}}}^1(\bar{\mathbf{A}}, M^+)$.

Theorem 5.2. (1) $\text{Hom}_{\bar{\mathbf{A}}}(\bar{\mathbf{A}}, M^+) \simeq \text{ann}_M \mathbf{C} = \{u \in M \mid \mathbf{C}u = 0\}$.
(2) $\text{Ext}_{\bar{\mathbf{A}}}^1(\bar{\mathbf{A}}, M^+) \simeq \text{Hom}_{\mathbf{H}}(\mathbf{C}, M)/(\text{Hom}_{\mathbf{H}}(\mathbf{C}, M)/\text{ann}_M \mathbf{C})$, where the quotient $\text{Hom}_{\mathbf{H}}(\mathbf{C}, M)/\text{ann}_M \mathbf{C}$ embeds in $\text{Hom}_{\mathbf{H}}(\mathbf{C}, M)$ if we map an element $u \in M$ to the homomorphism $\mu_u : c \mapsto cu$.

Proof. (1) $\text{Hom}_{\bar{\mathbf{A}}}(\bar{\mathbf{A}}, M^+)$ is identified with the set of homomorphisms $\phi : \mathbf{P} \rightarrow M^+$ such that $\phi\varepsilon = 0$. A homomorphism $\phi : \mathbf{P} \rightarrow M^+$ is uniquely defined by an element $u \in M$ such that $\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} u \\ 0 \end{pmatrix}$. Namely, $\phi \begin{pmatrix} a \\ c \end{pmatrix} = \begin{pmatrix} au \\ cu \end{pmatrix}$. Obviously, $\phi\varepsilon = 0$ if and only if $\mathbf{C}u = 0$, i.e. $u \in \text{ann}_M \mathbf{C}$.

(2) $\text{Ext}_{\bar{\mathbf{A}}}^1(\bar{\mathbf{A}}, M^+) \simeq \text{Hom}_{\bar{\mathbf{A}}}(\mathbf{C}^+, M^+)/\text{Hom}_{\bar{\mathbf{A}}}(\mathbf{P}, M^+)\varepsilon$. As the functor \mathbf{G}^* is fully faithful, $\text{Hom}_{\bar{\mathbf{A}}}(\mathbf{C}^+, M^+) \simeq \text{Hom}_{\mathbf{H}}(\mathbf{C}, M)$. Namely, $\psi : \mathbf{C} \rightarrow M$ induces $\psi^+ : \mathbf{C}^+ \rightarrow M^+$ mapping $\begin{pmatrix} a \\ b \end{pmatrix}$ to $\begin{pmatrix} \psi(a) \\ \psi(b) \end{pmatrix}$. Let $\phi : \mathbf{P} \rightarrow M^+$ corresponds, as above, to an element $u \in M$. Then $\phi\varepsilon \begin{pmatrix} a \\ c \end{pmatrix} = \begin{pmatrix} au \\ cu \end{pmatrix}$, so it equals μ_u , and $\text{Hom}_{\bar{\mathbf{A}}}(\mathbf{P}, M^+)\varepsilon$ is identified with $\text{Hom}_{\mathbf{H}}(\mathbf{C}, M)/\text{ann}_M \mathbf{C}$ embedded in $\text{Hom}_{\mathbf{H}}(\mathbf{C}, M)$ as above. \square

6. PARTIAL TILTING FOR BACKSTRÖM PAIRS

Let $\mathbf{H} \subseteq \mathbf{A}$ be a Backström pair. Consider the ring \mathbf{B} of triangular matrices of the form

$$\mathbf{B} = \begin{pmatrix} \bar{\mathbf{A}} & \bar{\mathbf{H}} \\ 0 & \mathbf{H} \end{pmatrix}.$$

Let $e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $B^1 = \mathbf{B}e_1$ and $B^2 = \mathbf{B}e_2$ be projective \mathbf{B} -modules given by the first and the second column of \mathbf{B} , i.e.

$$B^1 = \begin{pmatrix} \bar{\mathbf{A}} \\ 0 \end{pmatrix}, \quad B^2 = \begin{pmatrix} \bar{\mathbf{H}} \\ \mathbf{H} \end{pmatrix}.$$

A \mathbf{B} -module M is defined by a triple $\begin{pmatrix} M_1 \\ M_2 \\ \chi_M \end{pmatrix}$, where $M_1 = e_1M$ is an $\bar{\mathbf{A}}$ -module, $M_2 = e_2M$ is an \mathbf{H} -module and $\chi_M : M_2 \rightarrow M_1$ is an \mathbf{A} -homomorphism such that $\text{Ker } \chi_M \supseteq \mathbf{C}M_2$. We write an element $u \in M$ as a column $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$, where $u_1 = e_1u, u_2 = e_2u$. Then

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} au_1 + \chi_M(bu_2) \\ cu_2 \end{pmatrix}.$$

A homomorphism $\alpha : M \rightarrow N$ is defined by two homomorphisms $\alpha_1 : M_1 \rightarrow N_1$ and $\alpha_2 : M_2 \rightarrow N_2$ such that $\alpha_1\chi_M = \chi_N\alpha_2$. We write $\alpha = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}$.

Proposition 6.1. $\text{l.gl.dim } \mathbf{B} = \max\{\text{l.gl.dim } \mathbf{H}, \text{w.dim } \bar{\mathbf{H}}_{\mathbf{H}} + 1\}$.
In particular, if \mathbf{H} is left hereditary and $\bar{\mathbf{H}}$ is not flat as right \mathbf{H} module, then $\text{l.gl.dim } \mathbf{B} = 2$.

Proof. [26, Thm. 5] shows that $\text{l.gl.dim } \mathbf{B} \leq n$ if and only if

$$\text{l.gl.dim } \mathbf{H} \leq n \quad \text{and} \quad \text{R}^n \text{Hom}_{\bar{\mathbf{A}}}(\bar{\mathbf{H}} \otimes_{\mathbf{H}} -, -) = 0.$$

As the ring $\bar{\mathbf{A}}$ is semisimple,

$$\text{R}^n \text{Hom}_{\bar{\mathbf{A}}}(\bar{\mathbf{H}} \otimes_{\mathbf{H}} -, -) = \text{Hom}_{\bar{\mathbf{A}}}(\text{Tor}_n^{\mathbf{H}}(\bar{\mathbf{H}}, -), -).$$

It implies the first assertion. The second is obvious, since $\text{Tor}_1^{\mathbf{H}}(\bar{\mathbf{H}}, -) = 0$ if and only if $\bar{\mathbf{H}}$ is flat as right \mathbf{H} -module. \square

We denote by R the \mathbf{B} -module given by the triple $\begin{pmatrix} \mathbf{H}/\mathbf{A} \\ \mathbf{H} \\ \pi \end{pmatrix}$, where $\pi : \mathbf{H} \rightarrow \mathbf{H}/\mathbf{A}$ is the natural surjection.

Proposition 6.2. (1) $\text{End}_{\mathbf{B}} R \simeq \mathbf{A}^{\text{op}}$.
(2) $\text{pr.dim}_{\mathbf{B}} R = 1$.
(3) $\text{Ext}_{\mathbf{B}}^1(R, R) = 0$.

Recall that the conditions (2) and (3) mean that R is a *partial tilting* \mathbf{B} -module.

Proof. The minimal projective resolution of R is

$$0 \rightarrow B^1 \xrightarrow{\varepsilon} B^2 \rightarrow R \rightarrow 0,$$

where ε is the embedding, which gives (2). Any endomorphism γ of R induces a commutative diagram

$$\begin{array}{ccc} B^1 & \xrightarrow{\varepsilon} & B^2 \\ \gamma_1 \downarrow & & \downarrow \gamma_2 \\ B^1 & \xrightarrow{\varepsilon} & B_2 \end{array}$$

As $\text{End}_{\mathbf{B}} B^2 \simeq \mathbf{H}^{\text{op}}$, γ_2 is given by multiplication with an element $h \in \mathbf{H}$ on the right. If there is a commutative diagram as above, necessarily $h \in \mathbf{A}$, which proves (1). Finally, a homomorphism $\alpha : B^1 \rightarrow R$ maps the generator $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ of B^1 to an element $\begin{pmatrix} \bar{h} \\ 0 \end{pmatrix} \in R$. If h is a preimage of \bar{h} in \mathbf{H} , then α extends to the homomorphism $B^2 \rightarrow R$ that maps the generator $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ of B^2 to $\begin{pmatrix} 0 \\ h \end{pmatrix} \in R$. It implies (3). \square

Together with Theorem 1.1 it gives the following result.

- Theorem 6.3.** (1) *The functor $\mathbf{F} = \text{RHom}(R, -)$ induces an equivalence $\text{Tri}(R) \xrightarrow{\sim} \mathcal{D}(\mathbf{A})$.*
(2) *$\text{Ker } \mathbf{F}$ consists of complexes C such that the map $\chi_{H^k(C)}$ is bijective for all k .*
(3) *There is a recollement diagram*

$$\text{Ker } \mathbf{F} \begin{array}{c} \xleftarrow{\mathbf{l}^*} \\ \xrightarrow{\mathbf{l}} \\ \xleftarrow{\mathbf{l}'} \end{array} \mathcal{D}(\mathbf{B}) \begin{array}{c} \xleftarrow{\mathbf{F}^*} \\ \xrightarrow{\mathbf{F}} \\ \xleftarrow{\mathbf{F}'} \end{array} \mathcal{D}(\mathbf{A})$$

where \mathbf{F}^* and \mathbf{F}' are, respectively, left and right adjoints to \mathbf{F} , \mathbf{l} is the embedding, \mathbf{l}^* and \mathbf{l}' are, respectively, left and right adjoints to \mathbf{l} , and both adjunction morphisms $\eta :: \mathbf{F}\mathbf{F}^* \rightarrow \text{Id}_{\mathcal{D}(\mathbf{A})}$ and $\zeta : \text{Id}_{\mathcal{D}(\mathbf{A})} \rightarrow \mathbf{F}\mathbf{F}'$ are isomorphisms.

Actually, the claim (2) means that a complex C is in $\text{Ker } \mathbf{F}$ if and only if its cohomologies are direct sums of \mathbf{B} -modules of the form $\begin{pmatrix} U \\ U \end{pmatrix}$, where U is a simple $\bar{\mathbf{H}}$ -module.

Proof. (1) and (3) follow from Theorem 1.1, since the complex $P : 0 \rightarrow B^1 \xrightarrow{\varepsilon} B^2 \rightarrow 0$ is perfect, hence compact, and isomorphic to R in $\mathcal{D}(\mathcal{B})$. To find $\text{Ker } \mathbf{F}$, consider a complex

$$C : \dots \rightarrow C^{k-1} \xrightarrow{d^{k-1}} C^k \xrightarrow{d^k} C^{k+1} \rightarrow \dots$$

where C^k is defined by a triple $\begin{pmatrix} C_1^k \\ C_2^k \\ \chi_k \end{pmatrix}$ and $d^k = \begin{pmatrix} d_1^k \\ d_2^k \end{pmatrix}$, where $d_1^k \chi_k = \chi_{k+1} d_2^k$ for all k . Note that $C_i = (C_i^k, d_i^k)$ ($i = 1, 2$) are complexes, (χ_k) is a homomorphism of complexes and $H^k(C) = \begin{pmatrix} H^k(C_1) \\ H^k(C_2) \\ \bar{\chi}_k \end{pmatrix}$, where $\bar{\chi}_k = \chi_{H^k(C)}$ is induced by χ_k . A homomorphism $P \rightarrow C[k]$ is a pair of homomorphisms $\alpha : B^2 \rightarrow C^k$, $\beta : B^1 \rightarrow C^{k-1}$ such that

$\alpha_1\pi = \chi_k\alpha_2$, $\beta_2 = 0$, $d_i^k\alpha_i = 0$ ($i = 1, 2$) and $d^{k-1}\beta_1 = \alpha_1|_{\bar{\mathbf{A}}}$. Let $\alpha_2(1) = x \in C_2^k$ and $\beta_1(1) = y \in C_1^{k-1}$. These values completely define α and β . The conditions for α and β mean that $d_2^kx = 0$ and $d^{k-1}y = \chi_kx$.

This morphism is homotopic to zero if and only if there are maps $\sigma : B^2 \rightarrow C^{k-1}$ and $\tau : B^1\mathfrak{B}C^{k-2}$ such that $\alpha = d^{k-1}\sigma$ and $\beta = \sigma\varepsilon + d^{k-2}\tau$. Again σ is defined by the element $z = \sigma_2(1) \in C_2^{k-1}$ and τ is defined by the element $t = \tau_1(1) \in C_1^{k-2}$. Then the conditions for α and β mean that $x = d_2^{k-1}z$ and $y = \chi_{k-1}z + d_1^{k-2}t$.

Suppose that any homomorphism $P \rightarrow C[k]$ is homotopic to zero. Let $\bar{x} \in H^k(C^2)$ be such that $\bar{\chi}_k(\bar{x}) = 0$ and $x \in \text{Ker } d_2^k$ be a representative of \bar{x} . Then $\chi_k(x) = d_1^{k-1}y$ for some $y \in C^{k-1}$, so the pair (x, y) defines a homomorphism $P \rightarrow C[k]$. Therefore, there must be $z \in C_2^{k-1}$ such that $x = d^{k-1}z$, thus $\bar{x} = 0$ and $\bar{\chi}_k$ is injective. Let now $\bar{y} \in H^{k-1}(C_2)$ and $y \in C_2^{k-1}$ be its representative. Then the pair $(0, y)$ defines a homomorphism $P \rightarrow C[k]$, so there must be elements $z \in C_2^{k-1}$ and $t \in C_1^{k-2}$ such that $d_1^{k-1}z = 0$ and $y = \chi_{k-1}z + d_1^{k-2}t$. Hence $\bar{y} = \bar{\chi}_{k-1}(\bar{z})$, so $\bar{\chi}_{k-1}$ is surjective. As it holds for all k , we have that all maps $\bar{\chi}_k$ are bijective.

On the contrary, suppose that all $\bar{\chi}_k$ are bijective. If a pair (x, y) defines a homomorphism $P \rightarrow C[k]$, then $\chi_k(x) = d_1^{k-1}y$, so $\bar{\chi}_k(x) = 0$. Therefore, $\bar{x} = 0$, i.e. $x = d_2^{k-1}z$ for some $z \in C_2^{k-1}$ and $\chi_kx = d_1^{k-1}\chi_{k-1}z$. Then $d_1^{k-1}(y - \chi_{k-1}z) = 0$, hence there is an element $z' \in C_2^{k-1}$ such that $d_2^{k-1}z' = 0$ and the cohomology class of $y - \chi_{k-1}z$ equals $\bar{\chi}_{k-1}\bar{z}'$, i.e. $y - \chi_{k-1}z = \chi_{k-1}z' + d_1^{k-2}t$ for some t . Then $x = d_2^{k-1}(z + z')$ and $y = \chi_{k-1}(z + z') + d_1^{k-2}t$, so this homomorphism is homotopic to zero. \square

As usually, we identify the category $\mathbf{A}\text{-Mod}$ with the full subcategory of $\mathcal{D}(\mathbf{A})$ consisting of the complexes C concentrated in degree 0. The following result shows how the partial tilting functor \mathbf{F} behaves with respect to modules.

Corollary 6.4. *Let a \mathbf{B} -module M be given by the triple $(\begin{smallmatrix} M_1 \\ M_2 \end{smallmatrix} \chi_M)$.*

- (1) $\mathbf{F}M \in \mathbf{A}\text{-Mod}$ if and only if χ_M is surjective. Namely, then $\mathbf{F}M \simeq \text{Ker } \chi_M$.
- (2) $\mathbf{F}M \in \mathbf{A}\text{-Mod}[1]$ if and only if χ_M is injective. Namely, then $\mathbf{F}M \simeq \text{Cok } \chi_M[1]$.

Proof. Note that $\text{Hom}_{\mathbf{B}}(B^1, M) \simeq M_1$, $\text{Hom}_{\mathbf{B}}(B_2, M) \simeq M_2$ and if $\phi : B_2 \rightarrow M$ maps $(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix})$ to $(\begin{smallmatrix} 0 \\ x \end{smallmatrix})$, then $\phi\varepsilon$ maps $(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix})$ to $(\begin{smallmatrix} \chi_M(x) \\ 0 \end{smallmatrix})$. Therefore,

$\mathrm{RHom}_{\mathbf{B}}(R, M)$ is the complex

$$0 \rightarrow M_2 \xrightarrow{\chi_M} M_1 \rightarrow 0,$$

which proves the claim. \square

Remark 6.5. There are several derived equivalences related to $\tilde{\mathbf{A}}$.

- (1) If \mathbf{A} is a Backström order, it is known (see [10]) that the complex $T = B^1[1] \oplus \mathbf{H}^+$, where $B_1 = \begin{pmatrix} \tilde{\mathbf{A}} \\ 0 \end{pmatrix}$, is a tilting complex for $\tilde{\mathbf{A}}$ and $(\mathrm{End}_{\mathcal{D}(\tilde{\mathbf{A}})} T)^{\mathrm{op}} \simeq \mathbf{B}$, hence $\tilde{\mathbf{A}}$ is derived equivalent to \mathbf{B} . Nevertheless, in general situation of Backström rings (even of Backström algebras) it is not so. First of all, $\mathrm{Hom}_{\tilde{\mathbf{A}}}(B^1, \mathbf{H}^+) \simeq \mathrm{ann}_{\mathbf{H}} \mathbf{C}$, so it can happen that $\mathrm{Hom}_{\mathcal{D}(\tilde{\mathbf{A}})}(T, T[1]) \neq 0$. Moreover, even if $\mathrm{ann}_{\mathbf{H}} \mathbf{C} = 0$, one can see that $\tilde{\mathbf{H}}' = \mathrm{Ext}_{\tilde{\mathbf{A}}}^1(B^1, \mathbf{H}^+) \simeq \mathbf{C}^{-1}/{}_{\mathbf{C}}\mathbf{H}$, where $\mathbf{C}^{-1} = \mathrm{Hom}_{\mathbf{H}}(\mathbf{C}, \mathbf{H})$ and ${}_{\mathbf{C}}\mathbf{H} = \mathbf{H}/\mathrm{ann}_{\mathbf{H}} \mathbf{C}$ is naturally embedded into \mathbf{C}^{-1} . Therefore, in this case

$$(\mathrm{End}_{\mathcal{D}(\tilde{\mathbf{A}})} T)^{\mathrm{op}} \simeq \mathbf{B}' = \begin{pmatrix} \tilde{\mathbf{A}} & \tilde{\mathbf{H}}' \\ 0 & \mathbf{H} \end{pmatrix},$$

which need not coincide with \mathbf{B} . If \mathbf{H} is a hereditary order, then $\mathrm{ann}_{\mathbf{H}} \mathbf{C} = 0$ and $\tilde{\mathbf{H}}' \simeq \tilde{\mathbf{H}}$, hence $\mathbf{B}' \simeq \mathbf{B}$, in accordance with [10].

- (2) On the other hand, set $T' = \begin{pmatrix} \mathbf{A} & \mathbf{H}/\mathbf{A} \\ \mathbf{C} & \tilde{\mathbf{H}} \end{pmatrix}$ considered as left $\tilde{\mathbf{A}}$ -module. One can check that it is a tilting module for $\tilde{\mathbf{A}}$ and

$$(\mathrm{End}_{\mathcal{D}(\tilde{\mathbf{A}})} T')^{\mathrm{op}} \simeq \begin{pmatrix} \mathbf{A} & \mathbf{H}/\mathbf{A} \\ 0 & \tilde{\mathbf{A}} \end{pmatrix} = \mathbf{B}',$$

hence $\tilde{\mathbf{A}}$ is derived equivalent to \mathbf{B}' .

- (3) One more observation. Consider the right $\tilde{\mathbf{A}}$ -modules $(\tilde{\mathbf{A}} \ 0)$ and $(\mathbf{C} \ \mathbf{H})$. One can check that $T'' = (\tilde{\mathbf{A}} \ 0)[1] \oplus (\mathbf{C} \ \mathbf{H})$ is a tilting complex for $\mathcal{D}(\tilde{\mathbf{A}}^{\mathrm{op}})$ and

$$\mathrm{End}_{\mathcal{D}(\tilde{\mathbf{A}}^{\mathrm{op}})} T'' \simeq \mathbf{B}'' = \begin{pmatrix} \tilde{\mathbf{A}} & 0 \\ \tilde{\mathbf{H}} & \mathbf{H} \end{pmatrix},$$

hence $\tilde{\mathbf{A}}^{\mathrm{op}}$ is derived equivalent to $(\mathbf{B}'')^{\mathrm{op}}$.

Note that the functor $P \mapsto \mathrm{Hom}_{\mathbf{R}}(P, \mathbf{R})$ induces an exact duality $\mathrm{Perf}(\tilde{\mathbf{A}}) \rightarrow \mathrm{Perf}(\mathbf{B}'')$ for any ring \mathbf{R} . Hence $1 \ \mathrm{Perf}(\tilde{\mathbf{A}}) \simeq \mathrm{Perf}(\mathbf{B}'')$.

REFERENCES

- [1] Assem, I., Skowroński, A.: Iterated tilted algebras of type \tilde{A}_n . *Math. Z.* **195**, 269–290 (1987). DOI 10.1007/BF01166463
- [2] Bass, H.: Finitistic dimension and a homological generalization of semiprimary rings. *Trans. Am. Math. Soc.* **95**, 466–488 (1960). DOI 10.2307/1993568
- [3] Beilinson, A.A., Bernstein, J., Deligne, P.: Perverse sheaves. *Astérisque* **100** (1982)
- [4] Bekkert, V., Marcos, E.N., Merklen, H.A.: Indecomposables in derived categories of skewed-gentle algebras. *Commun. Algebra* **31**(6), 2615–2654 (2003). DOI 10.1081/AGB-120021885
- [5] Bekkert, V., Merklen, H.A.: Indecomposables in derived categories of gentle algebras. *Algebr. Represent. Theory* **6**(3), 285–302 (2003). DOI 10.1023/A:1025142023594
- [6] Burban, I., Drozd, Y.: Derived categories of nodal algebras. *J. Algebra* **272**(1), 46–94 (2004). DOI 10.1016/j.jalgebra.2003.07.025
- [7] Burban, I., Drozd, Y.: Derived categories for nodal rings and projective configurations. In: *Noncommutative algebra and geometry. Selected papers of a conference in honor of Z. Borevich, St. Petersburg, Russia.*, pp. 23–46. Boca Raton, FL: CRC Press (2005)
- [8] Burban, I., Drozd, Y.: Tilting on non-commutative rational projective curves. *Math. Ann.* **351**(3), 665–709 (2011). DOI 10.1007/s00208-010-0585-4
- [9] Burban, I., Drozd, Y.: On the derived categories of gentle and skew-gentle algebras: homological algebra and matrix problems. arXiv:1706.08358 [math.RT] (2017). DOI 10.48550/ARXIV.1706.08358
- [10] Burban, I., Drozd, Y., Gavran, V.: Minors and resolutions of non-commutative schemes. *Eur. J. Math.* **3**(2), 311–341 (2017). DOI:10.1007/s40879-017-0128-6
- [11] Chen, X.W., Ye, Y., Zhang, P.: Algebras of derived dimension zero. *Commun. Algebra* **36**(1), 1–10 (2008). DOI 10.1080/00927870701649184
- [12] Dlab, V., Ringel, C.M.: Indecomposable representations of graphs and algebras. *Mem. Am. Math. Soc.* **173**, 1–57 (1976)
- [13] Drozd, Y.A.: Finite modules over pure Noetherian algebras. *Proc. Steklov Inst. Math.* **183**, 97–108 (1990)
- [14] Drozd, Y.A.: Matrix problems, triangulated categories and stable homotopy types. *São Paulo J. Math. Sci.* **4**(2), 209–249 (2010). DOI 10.11606/issn.2316-9028.v4i2p209-249
- [15] Drozd, Y.A., Voloshyn, D.E.: Vector bundles over noncommutative nodal curves. *Ukr. Math. J.* **64**(2), 208–224 (2012). DOI 10.1007/s11253-012-0639-8
- [16] Drozd, Y.A., Zembyk, V.V.: Representations of nodal algebras of type A . *Algebra Discrete Math.* **15**(2), 179–200 (2013)
- [17] Geiß, C., de la Peña, J.A.: Auslander-Reiten components for clans. *Bol. Soc. Mat. Mex., III. Ser.* **5**(2), 307–326 (1999)
- [18] Happel, D.: Triangulated categories in the representation theory of finite dimensional algebras, *Lond. Math. Soc. Lect. Note Ser.*, vol. 119. Cambridge (UK) etc.: Cambridge University Press (1988)
- [19] Happel, D., Zacharia, D.: A homological characterization of piecewise hereditary algebras. *Math. Z.* **260**(1), 177–185 (2008). DOI 10.1007/s00209-007-0268-3

- [20] Keller, B.: Deriving DG categories. *Ann. Sci. Éc. Norm. Supér.* (4) **27**(1), 63–102 (1994). DOI 10.24033/asens.1689
- [21] Keller, B.: Derived categories and tilting. In: *Handbook of Tilting Theory, Lond. Math. Soc. Lect. Note Ser.*, vol. 332, pp. 49–104. Cambridge: Cambridge University Press (2007)
- [22] Lambek, J.: *Lectures on rings and modules*. New York, NY: Chelsea Publishing Company (1976)
- [23] Lekili, Y., Polishchuk, A.: Auslander orders over nodal stacky curves and partially wrapped Fukaya categories. *J. Topol.* **11**(3), 615–644 (2018). DOI 10.1112/topo.12064
- [24] Lunts, V.A.: Categorical resolution of singularities. *J. Algebra* **323**(10), 2977–3003 (2010). DOI 10.1016/j.jalgebra.2009.12.023
- [25] Neeman, A.: *Triangulated categories*, *Ann. Math. Stud.*, vol. 148. Princeton, NJ: Princeton University Press (2001). DOI 10.1515/9781400837212
- [26] Palmér, I., Roos, J.E.: Explicit formulae for the global homological dimensions of trivial extensions of rings. *J. Algebra* **60**, 11–42 (1979). DOI 10.1016/0021-8693(79)90106-6
- [27] Ringel, C.M., Roggenkamp, K.W.: Diagrammatic methods in the representation theory of orders. *J. Algebra* **60**, 11–42 (1979). DOI 10.1016/0021-8693(79)90106-6
- [28] Rouquier, R.: Dimensions of triangulated categories. *J. K-Theory* **1**(2), 193–256 (2008)
- [29] Voloshyn, D.E., Drozd, Y.A.: Derived categories of nodal curves. *Ukr. Math. J.* **64**(8), 1177–1184 (2013). DOI 10.1007/s11253-013-0708-7
- [30] Zembyk, V.V.: Structure of finite-dimensional nodal algebras. *Ukr. Math. J.* **66**(3), 467–472 (2014). DOI 10.1007/s11253-014-0944-5
- [31] Zembyk, V.V.: Skewed-gentle algebras are nodal. *Ukr. Math. J.* **67**(4), 648–651 (2015). DOI 10.1007/s11253-015-1129-6

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE,
 TERESCHENKIVSKA STR. 3, 01024 KYIV, UKRAINE
Email address: y.a.drozd@gmail.com, drozd@imath.kiev.ua
URL: www.imath.kiev.ua/~drozd