

A C^0 FINITE ELEMENT APPROXIMATION OF PLANAR OBLIQUE DERIVATIVE PROBLEMS IN NON-DIVERGENCE FORM *

GUANGWEI GAO SHUONAN WU[†]

Abstract. This paper proposes a C^0 (non-Lagrange) primal finite element approximation of the linear elliptic equations in non-divergence form with oblique boundary conditions in planar, curved domains. As an extension of [Calcolo, 58 (2022), No. 9], the Miranda-Talenti estimate for oblique boundary conditions at a discrete level is established by enhancing the regularity on the vertices. Consequently, the coercivity constant for the proposed scheme is exactly the same as that from PDE theory. The quasi-optimal order error estimates are established by carefully studying the approximation property of the finite element spaces. Numerical experiments are provided to verify the convergence theory and to demonstrate the accuracy and efficiency of the proposed methods.

Key words. Elliptic PDEs in non-divergence form, oblique derivative problems, Cordes condition, C^0 (non-Lagrange) finite element methods.

AMS subject classifications. 65N12, 65N15, 65N30, 35J15, 35D35

1. Introduction. We consider the elliptic equations in non-divergence form in a planar domain subject to the oblique boundary conditions. The model problem is to find $u : \Omega \rightarrow \mathbb{R}$ with $\int_{\Omega} u dx = 0$ such that

$$(1.1) \quad \begin{aligned} A : D^2 u &= f && \text{a.e. in } \Omega, \\ \ell \cdot \nabla u &= c && \text{on } \partial\Omega. \end{aligned}$$

The domain $\Omega \subset \mathbb{R}^2$ is assumed to have a C^2 boundary. Here, the coefficient matrix $A \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$ satisfies the uniform ellipticity condition (2.1), ℓ is a given unit vector field defined on $\partial\Omega$ (hereby called the “oblique vector field”), $A : B := \sum_{i,j=1}^2 A_{i,j} B_{i,j}$ denotes the Frobenius inner product of two matrices. The constant c appearing in the boundary conditions is intended to absorb the compatibility condition, i.e., c is a priori unknown. The problem (1.1) considered in this paper corresponds to the homogeneous oblique boundary conditions.

The oblique derivative problems arise from the linearization of transport boundary conditions for the Monge-Ampère equation [25]. Further applications of oblique derivative problems include the problem of determining the gravitational fields of celestial bodies [21], and the study of systems of certain conservation laws [15, 29].

Due to the non-divergence structure of (1.1), the concept of weak solutions based on the integration by part is no longer applicable. Instead, solution concepts such as classical solutions, viscosity solutions, and strong solutions are applicable. For these different solution concepts, numerical methods for the linear elliptic equations in non-divergence form have recently experienced rapid developments; See [23, 24, 24, 11, 4, 13, 14, 8, 10, 22, 28] for approximating the H^2 strong solutions, and [6, 7, 20, 16, 17, 26, 10] for others.

This article considers the H^2 strong solutions of non-divergence form with oblique boundary conditions (1.1). Here, the oblique vector field ℓ satisfies some technical assumptions (c.f. [18]), which are specified in Assumption 2.6. The coefficient matrix A is allowable to be discontinuous. As compensation, it is required to satisfy the Cordes

*The work of Shuonan Wu is supported in part by the National Natural Science Foundation of China grant No. 11901016.

[†]Corresponding author. School of Mathematical Sciences, Peking University, Beijing 100871, China .

condition (2.2), which is equivalence to the uniform ellipticity (2.1) in the planar case. In addition, the well-posedness of (1.1) hinges on a variant of the Miranda-Talenti estimate. We refer the reader to [18] for the analysis of PDEs with discontinuous coefficients under the Cordes condition.

For the numerical approximations of (1.1), a discontinuous Galerkin (DG) method was proposed in [12], which is applicable when choosing suitably large penalization parameters. A mixed method was proposed in [9], where the function approximating ∇u is confined to the piecewise linear space. More examples on the numerical approximations of oblique derivative problems are discussed in [1, 27, 5, 19], where [1, 5] apply to a particular geodetic and free boundary problem.

This paper proposed a C^0 primal finite element approximation of (1.1) without introducing any penalization term. Following the principle in [28], we established a discrete version of the Miranda-Talenti estimate for the oblique boundary condition by adopting the C^0 finite element with the enhanced regularity on the vertices. A typical family of finite elements that meets this requirement is the family of \mathcal{P}_k -Hermite finite elements ($k \geq 3$). Since the problem (1.1) is solved in a planar C^2 domain, we apply the techniques in [30, 2] and use the curved Hermite element; See Section 3.2 for details. The jump and boundary terms in the discrete Miranda-Talenti-type identity naturally induce the stabilization term in the proposed numerical scheme.

Thanks to the Miranda-Talenti estimate at a discrete level, we show the well-posedness of the proposed numerical scheme, which mimics the analysis of H^2 solutions to a great extent. A striking feature of the proposed scheme is that the coercivity constant at the discrete level is exactly the same as that from PDE theory. Another interesting feature is that the proposed method also gives an approximation of the unknown constant that arises in the compatibility condition. The proposed scheme is proved to be consistent, coercive, and bounded. Moreover, we constructed a quasi-interpolation operator that preserves the oblique boundary conditions at a discrete level with optimal order in the energy norm, which naturally leads to the energy norm error estimates.

The remaining parts of this paper are organized as follows. Section 2 reviews the H^2 strong solution theory of (1.1). In Section 3, we state the finite element spaces and present the proof of the discrete Miranda-Talenti-type identity. The numerical scheme for approximating (1.1) is proposed and analyzed in Section 4. Numerical experiments are presented in Section 5. Some technical proofs can be found in Appendixes A and B.

We use D to denote a generic subdomain of Ω , and ∂D denotes its boundary. $W_p^s(D)$ denotes the standard Sobolev space for $s \geq 0$ and $1 \leq p \leq \infty$, $W_p^0(D) = L^p(D)$ and $W_2^s(D) = H^s(D)$. $(\cdot, \cdot)_D$ denotes the standard inner product on $L^2(D)$. For convenience, we use C to denote a generic positive constant independent of mesh size h . The notation $X \lesssim Y$ means $X \leq CY$. $X \approx Y$ means $X \lesssim Y$ and $Y \lesssim X$. We also denote $|\cdot|$ as the Euclidean norm for vectors and the Frobenius norm for matrices.

2. Review of the H^2 strong solutions. This section reviews the H^2 strong solutions to the linear elliptic equations in non-divergence form with the oblique boundary conditions.

The coefficient $A \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$ is assumed to satisfy the uniform ellipticity, i.e., there exist $\underline{\nu}, \bar{\nu} > 0$ such that

$$(2.1) \quad \underline{\nu}|\boldsymbol{\xi}|^2 \leq \boldsymbol{\xi}^t A(x)\boldsymbol{\xi} \leq \bar{\nu}|\boldsymbol{\xi}|^2 \quad \forall \boldsymbol{\xi} \in \mathbb{R}^2, \text{ a.e. in } \Omega.$$

It is well known that in two dimensions, the uniform ellipticity (2.1) implies the

following Cordes condition (2.2) with $\varepsilon = 2\nu\bar{\nu}/(\nu^2 + \bar{\nu}^2)$, see [24].

DEFINITION 2.1 (Cordes condition). *The coefficient satisfies that there is an $\varepsilon \in (0, 1]$ such that*

$$(2.2) \quad \frac{|A|^2}{(\operatorname{tr} A)^2} \leq \frac{1}{1 + \varepsilon} \quad \text{a.e. in } \Omega.$$

Define the strictly positive function $\gamma \in L^\infty(\Omega)$ by $\gamma := \frac{\operatorname{tr} A}{|A|^2}$. The H^2 well-posedness of the problem (1.1) hinges on the following lemma; see [23, Lemma 1] for the proof.

LEMMA 2.2 (property of Cordes condition). *Under the Cordes condition (2.2), for any $v \in H^2(\Omega)$ and open set $U \subset \Omega$, the following inequality holds a.e. in U*

$$(2.3) \quad |\gamma A : D^2 v - \Delta v| \leq \sqrt{1 - \varepsilon} |D^2 v|.$$

The bounded domain $\Omega \subset \mathbb{R}^2$ is assumed to have a C^2 boundary. Moreover, $\partial\Omega$ is parametrized by the arc length φ , i.e., $\mathbf{x} : [0, L] \rightarrow \mathbb{R}^2$

$$(2.4) \quad \mathbf{x}(\varphi) = \begin{pmatrix} x_1(\varphi) \\ x_2(\varphi) \end{pmatrix} \quad \text{with } \mathbf{x}(0) = \mathbf{x}(L).$$

For a function $v : [0, L] \rightarrow \mathbb{R}$, let \dot{v} be its derivative with respect to the arc length parameter φ . Similarly, \ddot{v} denotes the second order derivative. Note that $\mathbf{x} \in C^2([0, L]; \mathbb{R}^2)$, we denote $\chi(\varphi) := \ddot{x}_1(\varphi)\dot{x}_2(\varphi) - \ddot{x}_2(\varphi)\dot{x}_1(\varphi)$ the curvature of $\partial\Omega$ at $\mathbf{x}(\varphi)$. Let n be the unit outward normal vector of $\partial\Omega$, and $t = \dot{\mathbf{x}}$ be the unit tangent vector.

The oblique vector field $\ell : [0, L] \rightarrow \mathbb{R}^2$, which satisfies $\ell(0) = \ell(L)$, is assumed to have C^1 regularity. We denote θ the oriented angle (anticlockwise) from n to ℓ , then $\theta : [0, L] \rightarrow \mathbb{R}$ is of class C^1 . We further assume an additional condition on the winding number of ℓ , namely

$$(2.5) \quad \frac{\theta(L) - \theta(0)}{2\pi} = 0.$$

This means that ℓ does not make a full turn around the normal n . Note that ℓ is defined on the interval $[0, L]$, and it can be identified with a function on $\partial\Omega$. Notation like $\ell(\mathbf{x}(\varphi))$ instead of $\ell(\varphi)$ will sometimes be used for convenience.

We define the following subspace of $H^2(\Omega)$:

$$(2.6) \quad \begin{aligned} H_\ell^2(\Omega) &:= \{v \in H^2(\Omega) : \ell \cdot \nabla v \text{ is constant on } \partial\Omega\}, \\ H_{\ell,0}^2(\Omega) &:= \{v \in H_\ell^2(\Omega) : \int_\Omega v dx = 0\}. \end{aligned}$$

The analysis of the H^2 well-posedness of the problem (1.1) hinges on several lemmas introduced below, which extend the important Miranda-Talenti estimate to the case of oblique boundary conditions. The proofs of these lemmas are given in [18], and we sketch the proof here for completeness.

LEMMA 2.3. *For any $v \in H_\ell^2(\Omega)$, it holds that*

$$(2.7) \quad \int_\Omega (\Delta v)^2 - |D^2 v|^2 dx = \int_0^L |\nabla v|^2 (\dot{\theta} - \chi) d\varphi.$$

Proof. First, a direct calculation gives

$$(\Delta v)^2 - |D^2 v|^2 = \frac{\partial}{\partial x_1} (\partial_1 v \partial_{22} v - \partial_2 v \partial_{12} v) - \frac{\partial}{\partial x_2} (\partial_1 v \partial_{12} v - \partial_2 v \partial_{11} v),$$

for sufficiently smooth function v . By the divergence Theorem, we obtain

$$\int_{\Omega} (\Delta v)^2 - |D^2 v|^2 dx = \int_0^L (\partial_1 v \partial_{22} v - \partial_2 v \partial_{12} v) n_1 - (\partial_1 v \partial_{12} v - \partial_2 v \partial_{11} v) n_2 d\varphi.$$

Next, we express the first and second order derivative of v as the directional derivative along ℓ and its perpendicular direction $\ell^\perp := (-\ell_2, \ell_1)$. Using the condition that $\ell \cdot \nabla v$ is constant on the boundary, we can obtain (2.7). We refer to [18, Page. 51] for more details of the proof. \square

We are now ready to give the Miranda-Talenti estimate in the case of oblique boundary conditions.

LEMMA 2.4 (Miranda-Talenti estimate). *Assume the domain Ω and the oblique vector field ℓ satisfy*

$$(2.8) \quad \chi_0 := \min_{\partial\Omega} (\dot{\theta} - \chi) \geq 0.$$

Then for any $v \in H_\ell^2(\Omega)$, it holds that

$$(2.9) \quad \int_{\Omega} |D^2 v|^2 dx \leq \int_{\Omega} (\Delta v)^2 dx.$$

Under a stronger assumption on Ω and ℓ , we obtain the following gradient estimate (see [18, Page. 53]).

LEMMA 2.5 (gradient estimate). *Assume the domain Ω and the oblique vector field ℓ satisfy $\chi_0 = \min_{\partial\Omega} (\dot{\theta} - \chi) > 0$. Then for any $v \in H_\ell^2(\Omega)$, it holds that*

$$(2.10) \quad \int_{\Omega} |\nabla v|^2 dx \leq C \int_{\Omega} (\Delta v)^2 dx,$$

where the constant C only depends on Ω and χ_0 .

Proof. Applying the divergence Theorem to the field $(x_1(\partial_1 v)^2, x_2(\partial_2 v)^2)$ and employing the Young inequality, we get the existence of two positive constants C_1 and C_2 (depend on Ω), such that

$$(2.11) \quad \int_{\Omega} |\nabla v|^2 dx \leq C_1 \int_{\Omega} |D^2 v|^2 dx + C_2 \int_{\partial\Omega} |\nabla v|^2 ds.$$

For any $A \geq 1$, using Lemma 2.3, we have

$$(2.12) \quad \int_{\Omega} |D^2 v|^2 dx \leq A \int_{\Omega} |D^2 v|^2 dx = A \int_{\Omega} (\Delta v)^2 dx - A \int_{\partial\Omega} |\nabla v|^2 (\dot{\theta} - \chi) ds.$$

Substituting (2.12) into (2.11) and combining it with the condition $\chi_0 = \min_{\partial\Omega} (\dot{\theta} - \chi) > 0$, we have

$$\int_{\Omega} |\nabla v|^2 dx \leq C_1 A \int_{\Omega} (\Delta v)^2 dx + (C_2 - C_1 A \chi_0) \int_{\partial\Omega} |\nabla v|^2 ds.$$

Then (2.10) is obtained by choosing $A = \max\{\frac{C_2}{C_1 \chi_0}, 1\}$. \square

We find that the semi-norm $|\cdot|_{H^2(\Omega)}$ is indeed a norm on $H_{\ell,0}^2(\Omega)$ by combining Lemma 2.5 (gradient estimate) and the Poincaré inequality, i.e., for any $v \in H_{\ell,0}^2(\Omega)$

$$\begin{aligned} \|v\|_{H^2(\Omega)}^2 &= \|v\|_{L^2(\Omega)}^2 + |v|_{H^1(\Omega)}^2 + |v|_{H^2(\Omega)}^2 \\ &\lesssim |v|_{H^1(\Omega)}^2 + |v|_{H^2(\Omega)}^2 \quad (\text{Poincaré inequality}) \\ &\lesssim \|\Delta v\|_{L^2(\Omega)}^2 + |v|_{H^2(\Omega)}^2 \lesssim |v|_{H^2(\Omega)}^2. \quad (\text{Lemma 2.5}) \end{aligned}$$

Now in the Hilbert space $(H_{\ell,0}^2(\Omega), |\cdot|_{H^2(\Omega)})$, we are ready to define the bilinear form $b : H_{\ell,0}^2(\Omega) \times H_{\ell,0}^2(\Omega) \rightarrow \mathbb{R}$ as

$$(2.13) \quad b(w, v) := \int_{\Omega} \gamma A : D^2 w \Delta v dx.$$

Lemma 2.2 (property of Cordes condition) and Lemma 2.4 (Miranda-Talenti estimate) imply the coercivity, i.e., for any $v \in H_{\ell,0}^2(\Omega)$

$$(2.14) \quad \begin{aligned} b(v, v) &= \int_{\Omega} (\Delta v)^2 dx + \int_{\Omega} (\gamma A - I) : D^2 v \Delta v dx \\ &\geq \|\Delta v\|_{L^2(\Omega)}^2 - \sqrt{1-\varepsilon} \|D^2 v\|_{L^2(\Omega)} \|\Delta v\|_{L^2(\Omega)} \geq (1 - \sqrt{1-\varepsilon}) |v|_{H^2(\Omega)}^2. \end{aligned}$$

The variational form of problem (1.1) reads: Find $u \in H_{\ell,0}^2(\Omega)$ such that

$$(2.15) \quad b(u, v) = l(v) \quad \forall v \in H_{\ell,0}^2(\Omega),$$

where $l(v) = \int_{\Omega} \gamma f \Delta v dx$ is a linear functional on $H_{\ell,0}^2(\Omega)$. Before claiming the well-posedness result for (1.1), we summarize the assumptions on the data as follows.

Assumption 2.6. Let $A \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$, satisfy the uniform ellipticity (2.1). Ω is assumed to be a bounded domain with C^2 boundary. The oblique vector field ℓ is of class C^1 with unit length, and (2.5) is satisfied. Furthermore, it is assumed that $\chi_0 = \min_{\partial\Omega}(\theta - \chi) > 0$.

THEOREM 2.7 (well-posedness). *Under Assumption 2.6, for any given $f \in L^2(\Omega)$, there exists a unique solution $u \in H_{\ell,0}^2(\Omega)$ to the variational problem (2.15). Furthermore, u is the H^2 strong solution to the oblique derivative problem (1.1).*

Proof. It is easy to verify that the linear form $l(\cdot)$ and the bilinear form $b(\cdot, \cdot)$ are bounded on the Hilbert space $(H_{\ell,0}^2(\Omega), |\cdot|_{H^2(\Omega)})$. Then the coercivity of $b(\cdot, \cdot)$ and the Lax-Milgram Theorem imply the existence of a unique solution $u \in H_{\ell,0}^2(\Omega)$ to the variational problem (2.15).

Note that for any given $z \in L^2(\Omega)$, the condition (2.5) implies that there exists a unique $v_z \in H_{\ell,0}^2(\Omega)$ such that $\Delta v_z = z$, a.e. in Ω (see [18, Page. 49-50] for details). Therefore,

$$\int_{\Omega} (\gamma A : D^2 u - \gamma f) z dx = \int_{\Omega} (\gamma A : D^2 u - \gamma f) \Delta v_z dx = b(u, v_z) - l(v_z) = 0.$$

Then the fundamental Lemma of the calculus of variations yields $\gamma A : D^2 u - \gamma f = 0$, which leads to $A : D^2 u = f$, a.e. in Ω since γ is uniformly positive. \square

3. Finite element space and discrete Miranda-Talenti-type identity.

In this section, we shall construct the finite element space for approximating (1.1). More precisely, we adopt the curved Hermite elements on exact triangulations of the domain Ω . The discrete Miranda-Talenti-type identity for oblique boundary conditions is therefore established through the C^1 -continuity on the vertices.

3.1. Exact triangulation of the curved domain. Since ℓ is only defined on $\partial\Omega$, the triangulation needs to strictly fit the curved boundary of the domain Ω . For this purpose, the domain Ω is exactly triangulated by \mathcal{T}_h , i.e., \mathcal{T}_h consists of curved triangles near the boundary with at most one truly curved edge (fits the curved boundary). In the interior of Ω , \mathcal{T}_h consists of straight triangles. A formal definition of the curved triangle is stated as follows [2].

DEFINITION 3.1 (curved triangle). *A closed set $K \subset \mathbb{R}^2$ is a curved triangle if there exists a C^1 mapping F_K that maps a straight reference triangle \hat{K} onto K and that is of the form*

$$(3.1) \quad F_K = \tilde{F}_K + \Phi_K,$$

where $\tilde{F}_K : \hat{x} \mapsto B_K \hat{x} + b_K$ is an invertible affine mapping and Φ_K is a C^1 mapping satisfying

$$(3.2) \quad c_K := \sup_{\hat{x} \in \hat{K}} \|D\Phi_K B_K^{-1}\| < 1.$$

Let $\tilde{K} := \tilde{F}_K(\hat{K})$ be a straight triangle that ‘‘approximates’’ the curved triangle K . Note that if the mapping Φ_K is equal to zero, $K = \tilde{K}$ is a straight triangle. The mesh size of K is defined as $h_K := \text{diam}(\tilde{K})$. A simple calculation shows that (c.f. [2, Lemma 3.1])

$$(3.3) \quad (1 - c_K)h_K \leq \text{diam}(K) \leq (1 + c_K)h_K.$$

Denote $h := \max_{K \in \mathcal{T}_h} h_K$. Let \mathcal{N}_h be the set of vertices in \mathcal{T}_h , $\mathcal{N}_h^\partial := \mathcal{N}_h \cap \partial\Omega$ and $\mathcal{N}_h^i := \mathcal{N}_h \setminus \mathcal{N}_h^\partial$. Denote \mathcal{F}_h the set of edges in \mathcal{T}_h . Let $\mathcal{F}_h^\partial := \mathcal{F}_h \cap \partial\Omega$ and $\mathcal{F}_h^i := \mathcal{F}_h \setminus \mathcal{F}_h^\partial$. For each edge $F \in \mathcal{F}_h^i$, we define $h_F := \min(h_K, h_{K'})$ where K and K' are such that $F = \partial K \cap \partial K'$. If $F \in \mathcal{F}_h^\partial$, we define $h_F := h_K$ where K is such that $F = \partial K \cap \partial\Omega$.

A family of conforming triangulations $\{\mathcal{T}_h\}_{h>0}$ is said to be shape-regular if there exist two constants σ and c_1 independent of h such that,

$$\sup_h \sup_{K \in \mathcal{T}_h} \frac{h_K}{\rho_K} \leq \sigma \quad \text{and} \quad \sup_h \sup_{K \in \mathcal{T}_h} c_K \leq c_1 < 1.$$

Here, ρ_K is the diameter of the sphere inscribed in \tilde{K} . In the standard finite element theory, the Sobolev space $W_p^m(K)$ is usually related to $W_p^m(\hat{K})$ by the push-forward mapping, i.e., $\hat{v} = F_K^* v = v \circ F_K$. To reproduce this in the curved triangle case, we need to assume some smoothness on F_K .

DEFINITION 3.2 (class m triangulation). *The family $\{\mathcal{T}_h\}_{h>0}$ is said to be regular of order m if it is shape-regular and if, for each h , any $K \in \mathcal{T}_h$ the mapping F_K is of class C^{m+1} , with*

$$\sup_h \sup_{K \in \mathcal{T}_h} c_i(K) \leq c_i < \infty, \quad 2 \leq i \leq m + 1,$$

where $c_i(K) := \|F_K|_{W_\infty^i(\hat{K}; \mathbb{R}^2)}\| \|B_K\|^{-i}$.

Remark 3.3 (construction of F_K). The construction of F_K is specified in [2, Section 6]. Here, we emphasize that the construction of F_K guarantees the following fact: if F is a straight edge of the curved triangle K , then for any edge \hat{F} of \hat{K} ,

$$F_K|_{\hat{F}} : \hat{F} \rightarrow F \text{ is an affine mapping.}$$

Further, to construct $\{\mathcal{T}_h\}_{h>0}$ with order m , the mapping F_K needs to be C^{m+1} , which requires the boundary $\partial\Omega$ to be piecewise C^{m+1} .

3.2. Finite element space. Following the idea in [28], the key to designing the finite element space is the C^1 -continuity on the vertices. Since the discretization is based on the exact triangulation of the domain Ω , we adopt the technique in [2] to consider the curved Hermite element. To this end, we start with the standard Hermite element on straight triangles.

Hermite element on the straight reference triangle. For the straight reference triangle \hat{K} , the shape function space is given as $\mathcal{P}_k(\hat{K})$ ($k \geq 3$), where $\mathcal{P}_k(\hat{K})$ denotes the set of polynomials with total degree not exceeding k on \hat{K} . The set of degrees of freedom $\hat{\Sigma}$ is defined as follows:

- Function value $\hat{v}(\hat{\mathbf{a}})$ and first order derivatives $\hat{\partial}_i \hat{v}(\hat{\mathbf{a}})$, $i = 1, 2$ at each vertex;
- Function values at $(k - 3)$ nodes on the interior of each edge \hat{e} ;
- Function values at $\frac{(k-1)(k-2)}{2}$ nodes in the interior of \hat{K} .

It is simple to check that the degrees of freedom given above form a unisolvent set [3].

Curved Hermite element. The shape function space for a curved triangle K is given as

$$P_K := \{v = \hat{v} \circ F_K^{-1}, \hat{v} \in \mathcal{P}_k(\hat{K})\}, \quad k \geq 3.$$

The set of degrees of freedom Σ_K is defined as follows:

- Function value $v(\mathbf{a})$ and first order derivatives $\partial_i v(\mathbf{a})$, $i = 1, 2$ at each vertex;
- Function values at $(k - 3)$ nodes on the interior of each edge e ;
- Function values at $\frac{(k-1)(k-2)}{2}$ nodes in the interior of K .

In Σ_K , the choice of the nodes on the edge or in the triangle are induced by the nodes in $\hat{\Sigma}$ through F_K .

We now verify the unisolvent property. First, the dimension of the shape function space is $\dim P_K = \dim \mathcal{P}_k(\hat{K}) = \frac{(k+1)(k+2)}{2}$, which coincides with the number of the degrees of freedom. Therefore, it suffices to show that $v \in P_K$ vanishes if it vanishes at all the degrees of freedom in Σ_K . By the definition of P_K , we know that $v = \hat{v} \circ F_K^{-1}$ for some $\hat{v} \in \mathcal{P}_k(\hat{K})$. Note that v and its first derivatives vanish at the three vertices of K , by the chain rule, we immediately know that \hat{v} and its first derivatives also vanish at the three vertices of \hat{K} . Next, the vanishing of the remaining degrees of freedom in Σ_K ensures that \hat{v} has $\frac{(k-1)(k-2)}{2}$ zero points inside \hat{K} and $(k - 3)$ zero points inside each edge \hat{e} . Following the unisolvent property of $(\hat{K}, \mathcal{P}_k(\hat{K}), \hat{\Sigma})$, \hat{v} vanishes. Therefore, $v = \hat{v} \circ F_K^{-1} \equiv 0$.

The degrees of freedom ensure that the global finite element space has the C^0 -continuity in Ω and C^1 -continuity on the vertices for every triangulation \mathcal{T}_h , i.e.,

$$V_h = \{v \in H^1(\Omega) : v|_K \in P_K, \forall K \in \mathcal{T}_h, v \text{ is } C^1 \text{ at all vertices}\}.$$

Enforcing the boundary conditions, we define

$$V_{h,0}^\ell := \{v_h \in V_h, \int_{\Omega} v_h dx = 0, \nabla v_h(x_i) \cdot \ell \text{ is constant } \forall x_i \in \mathcal{N}_h^\partial\}.$$

Remark 3.4 (regularity of \mathcal{T}_h). To get the optimal approximation property, we need to assume that $\{\mathcal{T}_h\}_{h>0}$ is regular of order k , which is consistent with the degree of the polynomial involved in V_h .

The following lemmas deal with the scaling argument and the trace estimate. We refer to [2] for the proof.

LEMMA 3.5 (scaling on curved triangles, see Lemma 2.3 in [2]). *Let $\{\mathcal{T}_h\}_{h>0}$ be regular of order k . Let s be an integer, $0 \leq s \leq k+1$, and $p \in [1, \infty]$. For any $K \in \mathcal{T}_h$, a function v belongs to $W_p^s(K)$ if and only if the function $\hat{v} = v \circ F_K$ belongs to $W_p^s(\hat{K})$, and we have*

$$(3.4a) \quad |v|_{W_p^s(K)} \leq C_S |\det B_K|^{1/p} \|B_K^{-1}\|^s \left(\sum_{i=\min(s,1)}^s \|B_K\|^{(s-i)} |\hat{v}|_{W_p^i(\hat{K})} \right),$$

$$(3.4b) \quad |\hat{v}|_{W_p^s(\hat{K})} \leq C_S |\det B_K|^{-1/p} \|B_K\|^s \left(\sum_{i=\min(s,1)}^s |v|_{W_p^i(K)} \right),$$

where C_S depends continuously on c_1, c_2, \dots, c_{k+1} given in Definition 3.2.

By the shape regularity of $\{\mathcal{T}_h\}_{h>0}$, we know $\|B_K\| \approx h_K$ and $|\det B_K| \approx h_K^2$. Thus we obtain

$$|v|_{W_p^s(K)} \lesssim h_K^{2/p-s} \sum_{i=\min(s,1)}^s h_K^{(s-i)} |\hat{v}|_{W_p^i(\hat{K})} \quad \text{and} \quad |\hat{v}|_{W_p^s(\hat{K})} \lesssim h_K^{s-2/p} \sum_{i=\min(s,1)}^s |v|_{W_p^i(K)},$$

with hidden constants depend on the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

LEMMA 3.6 (trace estimate, see Lemma 2.4 in [2]). *Let $\{\mathcal{T}_h\}_{h>0}$ be shape-regular. For any $K \in \mathcal{T}_h$ we have*

$$(3.5) \quad \|v\|_{L^2(\partial K)}^2 \leq C_{\text{Tr}} (h_K^{-1} \|v\|_{L^2(K)}^2 + h_K |v|_{H^1(K)}^2) \quad \forall v \in H^1(K),$$

where C_{Tr} depends on c_1 and the shape-regular constant σ .

LEMMA 3.7 (inverse estimate). *Let $\{\mathcal{T}_h\}_{h>0}$ is regular of order k . For any $K \in \mathcal{T}_h$, the following inverse estimate holds with $0 \leq m < s \leq k+1$*

$$(3.6) \quad |v|_{H^s(K)} \leq C_{\text{Inv}} h_K^{m-s} \sum_{i=\min(1,m)}^m |v|_{H^i(K)}, \quad \forall v \in P_K,$$

where C_{Inv} depends on the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

Proof. If $m \geq 1$, by Lemma 3.5 (scaling on curved triangles) and the norm equivalence on finite dimensional space, we have

$$(3.7) \quad |v|_{H^s(K)} \lesssim h_K^{1-s} \sum_{i=1}^s |\hat{v}|_{H^i(\hat{K})} h_K^{(s-i)} \lesssim h_K^{1-s} \sum_{i=1}^m |\hat{v}|_{H^i(\hat{K})} h_K^{(m-i)}.$$

Next for each $1 \leq i \leq m$, by Lemma 3.4 (scaling on curved triangles) again, we have $|\hat{v}|_{H^i(\hat{K})} \lesssim h_K^{i-1} \sum_{j=1}^i |v|_{H^j(K)}$. Substitute this into (3.7), we obtained (3.6). For the case of $m = 0$, similar arguments lead to

$$|v|_{H^s(K)} \lesssim h_K^{1-s} \sum_{i=1}^s |\hat{v}|_{H^i(\hat{K})} h_K^{(s-i)} \lesssim h_K^{1-s} \|\hat{v}\|_{L^2(\hat{K})} \lesssim h_K^{-s} \|v\|_{L^2(K)}.$$

This completes the proof. \square

As one of the main focuses of our work, the following theorem gives the approximation property with oblique boundary conditions. The proof is a nontrivial, and is therefore postponed to Appendix A for the reader's convenience.

THEOREM 3.8 (approximation property of $V_{h,0}^\ell$). *Assume that $\{\mathcal{T}_h\}_{h>0}$ is regular of order k , $u \in H^s(\Omega) \cap H_{\ell,0}^2(\Omega)$, and the oblique vector field ℓ is piecewise C^{s-1} with $2 \leq s \leq k+1$. Then there exists a quasi-interpolation $\Pi_h^\ell : H_{\ell,0}^2(\Omega) \rightarrow V_{h,0}^\ell$ satisfying*

$$(3.8a) \quad \|u - \Pi_h^\ell u\|_{H^m(K)} \lesssim h_K^{s-m} \|u\|_{H^s(\omega_K)}, \quad m = 1, 2,$$

$$(3.8b) \quad \|\nabla(u - \Pi_h^\ell u)\|_{L^2(\partial K)} \lesssim h_K^{s-1-1/2} \|u\|_{H^s(\omega_K)}.$$

Here, ω_K is the local neighborhood of $K \in \mathcal{T}_h$ and the hidden constants depend on $\|\ell\|_{W_\infty^{s-1}(\partial\omega_K \cap \partial\Omega)}$, the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

3.3. The discrete Miranda-Talenti-type identity for oblique boundary conditions. In this subsection, we establish the Miranda-Talenti estimate for oblique boundary conditions in the discrete level by using the C^1 -continuity on the vertices of the finite element space.

Define the jump of the normal derivative on an interior edge $F = \partial K^+ \cap \partial K^-$ as

$$\left[\frac{\partial v}{\partial n}\right] := \frac{\partial v^+}{\partial n^+} + \frac{\partial v^-}{\partial n^-} \quad \text{where} \quad v^\pm := v|_{K^\pm},$$

where n^\pm is the unit outward normal vector of K^\pm . For any interior edge $F \in \mathcal{F}_h^i$, we specify a tangent direction denoted as t_F . Let $\mathbf{x}_F^{(1)}$ and $\mathbf{x}_F^{(2)}$ be the two vertices of F (along the direction t_F). Recall $\ell = (\ell_1, \ell_2)$ is the oblique vector field defined on $\partial\Omega$, and we denote

$$\frac{\partial v}{\partial \ell} := \ell_1 \partial_1 v + \ell_2 \partial_2 v \quad \text{and} \quad \frac{\partial v}{\partial \ell^\perp} := -\ell_2 \partial_1 v + \ell_1 \partial_2 v$$

as the directional derivatives along ℓ and its perpendicular direction $\ell^\perp = (-\ell_2, \ell_1)$. Moreover, we denote $\left(\frac{\partial v}{\partial \ell}\right) := \frac{d}{d\varphi} \left(\frac{\partial v}{\partial \ell}(\mathbf{x}(\varphi))\right)$.

THEOREM 3.9 (discrete Miranda-Talenti-type identity for oblique boundary conditions). *Let Ω be a bounded domain with C^2 boundary. The oblique vector field ℓ is of class C^1 . Let \mathcal{T}_h be an exact conforming triangulation of Ω . For any $v_h \in V_h$, it holds that*

$$(3.9) \quad \begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K (\Delta v_h)^2 dx &= \sum_{K \in \mathcal{T}_h} \int_K |D^2 v_h|^2 dx + 2 \sum_{F \in \mathcal{F}_h^i} \int_F \left[\frac{\partial v_h}{\partial n}\right] \frac{\partial^2 v_h}{\partial t_F^2} ds \\ &+ \sum_{F \in \mathcal{F}_h^\partial} \int_F |\nabla v_h|^2 (\theta - \chi) - 2 \frac{\partial v_h}{\partial \ell^\perp} \left(\frac{\partial v_h}{\partial \ell}\right) ds. \end{aligned}$$

Proof. Step 1: For a straight triangle $K \in \mathcal{T}_h$, the normal vector of ∂K is piecewise constant. Using integration by part, we obtain (see [23, Equ. (3.7)])

$$(3.10) \quad \begin{aligned} \int_K (\Delta v_h)^2 dx &= \int_K |D^2 v_h|^2 dx + \int_{\partial K} \frac{\partial v_h}{\partial n} \Delta v_h - \nabla \frac{\partial v_h}{\partial n} \cdot \nabla v_h ds \\ &= \int_K |D^2 v_h|^2 dx + \int_{\partial K} \frac{\partial v_h}{\partial n} \frac{\partial^2 v_h}{\partial t^2} - \frac{\partial^2 v_h}{\partial t \partial n} \frac{\partial v_h}{\partial t} ds. \end{aligned}$$

Here, the common term $\int_{\partial K} \frac{\partial^2 v_h}{\partial n^2} \frac{\partial v_h}{\partial n} ds$ is cancelled in the last step. Using integration by part on the edge $F \subset \partial K$, we have

$$\begin{aligned} \int_{\partial K} \frac{\partial^2 v_h}{\partial t \partial n} \frac{\partial v_h}{\partial t} ds &= \sum_{F \subset \partial K} \int_F \frac{\partial^2 v_h}{\partial t_F \partial n} \frac{\partial v_h}{\partial t_F} ds \\ &= \sum_{F \subset \partial K} \left(- \int_F \frac{\partial v_h}{\partial n} \frac{\partial^2 v_h}{\partial t_F^2} ds + \frac{\partial v_h}{\partial n} \frac{\partial v_h}{\partial t_F} \Big|_{\mathbf{x}_F^{(1)}}^{\mathbf{x}_F^{(2)}} \right). \end{aligned}$$

Hence, (3.10) can be reformulated as

$$(3.11) \quad \int_K (\Delta v_h)^2 dx = \int_K |D^2 v_h|^2 dx + \sum_{F \subset \partial K} \left(2 \int_F \frac{\partial v_h}{\partial n} \frac{\partial^2 v_h}{\partial t_F^2} ds - \frac{\partial v_h}{\partial n} \frac{\partial v_h}{\partial t_F} \Big|_{\mathbf{x}_F^{(1)}}^{\mathbf{x}_F^{(2)}} \right).$$

Step 2: A direct calculation shows that

$$(\Delta v_h)^2 - |D^2 v_h|^2 = \frac{\partial}{\partial x_1} (\partial_1 v_h \partial_{22} v_h - \partial_2 v_h \partial_{12} v_h) - \frac{\partial}{\partial x_2} (\partial_1 v_h \partial_{12} v_h - \partial_2 v_h \partial_{11} v_h).$$

For any curved triangle $K \in \mathcal{T}_h$, apply the divergence Theorem

$$\begin{aligned} &\int_K (\Delta v_h)^2 - |D^2 v_h|^2 dx \\ &= \int_{\partial K} n_1 (\partial_1 v_h \partial_{22} v_h - \partial_2 v_h \partial_{12} v_h) - n_2 (\partial_1 v_h \partial_{12} v_h - \partial_2 v_h \partial_{11} v_h) ds \\ &= \sum_{i=1, F_i \subset \partial K}^3 \underbrace{\int_{F_i} n_1 (\partial_1 v_h \partial_{22} v_h - \partial_2 v_h \partial_{12} v_h) - n_2 (\partial_1 v_h \partial_{12} v_h - \partial_2 v_h \partial_{11} v_h) ds}_{I_i}. \end{aligned}$$

In the last step, we denote $\partial K = \cup_{i=1}^3 F_i$ where F_1 and F_2 are the two straight edges and $F_3 \subset \partial \Omega$ is the curved edge. On the straight edge (i.e. $i = 1, 2$), the same calculation as in *Step 1* leads to

$$(3.12) \quad I_i = 2 \int_{F_i} \frac{\partial v_h}{\partial n} \frac{\partial^2 v_h}{\partial t_{F_i}^2} ds - \left(\frac{\partial v_h}{\partial n} \frac{\partial v_h}{\partial t_{F_i}} \right) \Big|_{\mathbf{x}_{F_i}^{(1)}}^{\mathbf{x}_{F_i}^{(2)}}, \quad i = 1, 2.$$

On the curved edge F_3 , we first use the directional derivatives $\frac{\partial v_h}{\partial \ell}$ and $\frac{\partial v_h}{\partial \ell^\perp}$ to represent $\partial_1 v_h$ and $\partial_2 v_h$, i.e.,

$$\partial_1 v_h = \frac{\partial v_h}{\partial \ell} \ell_1 - \frac{\partial v_h}{\partial \ell^\perp} \ell_2 \quad \text{and} \quad \partial_2 v_h = \frac{\partial v_h}{\partial \ell} \ell_2 + \frac{\partial v_h}{\partial \ell^\perp} \ell_1.$$

Then, substituting above formula into I_3 and parametrizing $F_3 \subset \partial \Omega$ by the arc length φ through the C^2 mapping, $\mathbf{x} : [L_1, L_2] \rightarrow \mathbb{R}^2$, we obtain

$$\begin{aligned} (3.13) \quad I_3 &= \int_{F_3} n_1 (\partial_1 v_h \partial_{22} v_h - \partial_2 v_h \partial_{12} v_h) - n_2 (\partial_1 v_h \partial_{12} v_h - \partial_2 v_h \partial_{11} v_h) ds \\ &= \int_{L_1}^{L_2} \frac{\partial v_h}{\partial \ell} (\ell_2 \partial_{11} v_h n_2 - \ell_2 \partial_{12} v_h n_1 - \ell_1 \partial_{21} v_h n_2 + \ell_1 \partial_{22} v_h n_1) \\ &\quad + \frac{\partial v_h}{\partial \ell^\perp} (\ell_1 \partial_{11} v_h n_2 - \ell_1 \partial_{12} v_h n_1 + \ell_2 \partial_{21} v_h n_2 - \ell_2 \partial_{22} v_h n_1) d\varphi. \end{aligned}$$

Taking the derivative of $\frac{\partial v_h}{\partial \ell}$ and $\frac{\partial v_h}{\partial \ell^\perp}$ with respect to the arc length φ , and noting that $(n_1, n_2) = (\dot{\mathbf{x}}_2, -\dot{\mathbf{x}}_1)$, we obtain

$$(3.14) \quad \begin{aligned} \left(\frac{\partial \dot{v}_h}{\partial \ell} \right) &= (\ell_1 \partial_{11} v_h \dot{\mathbf{x}}_1 + \ell_1 \partial_{12} v_h \dot{\mathbf{x}}_2 + \ell_2 \partial_{21} v_h \dot{\mathbf{x}}_1 + \ell_2 \partial_{22} v_h \dot{\mathbf{x}}_2) \\ &\quad + \dot{\ell}_1 \partial_1 v_h + \dot{\ell}_2 \partial_2 v_h \\ &= (-\ell_1 \partial_{11} v_h n_2 + \ell_1 \partial_{12} v_h n_1 - \ell_2 \partial_{21} v_h n_2 + \ell_2 \partial_{22} v_h n_1) \\ &\quad + \dot{\ell}_1 \partial_1 v_h + \dot{\ell}_2 \partial_2 v_h, \end{aligned}$$

and similarly

$$(3.15) \quad \begin{aligned} \left(\frac{\partial \dot{v}_h}{\partial \ell^\perp} \right) &= (\ell_2 \partial_{11} v_h n_2 - \ell_2 \partial_{12} v_h n_1 - \ell_1 \partial_{21} v_h n_2 + \ell_1 \partial_{22} v_h n_1) \\ &\quad - \dot{\ell}_2 \partial_1 v_h + \dot{\ell}_1 \partial_2 v_h. \end{aligned}$$

Substituting (3.14) and (3.15) into (3.13), we arrive at

$$I_3 = \int_{L_1}^{L_2} \left(\left(\frac{\partial v_h}{\partial \ell} \right)^2 + \left(\frac{\partial v_h}{\partial \ell^\perp} \right)^2 \right) (\ell_1 \dot{\ell}_2 - \ell_2 \dot{\ell}_1) + \frac{\partial v_h}{\partial \ell} \left(\frac{\partial \dot{v}_h}{\partial \ell^\perp} \right) - \frac{\partial v_h}{\partial \ell^\perp} \left(\frac{\partial \dot{v}_h}{\partial \ell} \right) d\varphi.$$

Using integration by part, $|\nabla v_h|^2 = \left(\frac{\partial v_h}{\partial \ell} \right)^2 + \left(\frac{\partial v_h}{\partial \ell^\perp} \right)^2$ and the identity $(\ell_1 \dot{\ell}_2 - \ell_2 \dot{\ell}_1) = \dot{\theta} - \chi$ (see [18, Page. 48]), we have

$$(3.16) \quad I_3 = \int_{L_1}^{L_2} |\nabla v_h|^2 (\dot{\theta} - \chi) - 2 \frac{\partial v_h}{\partial \ell^\perp} \left(\frac{\partial \dot{v}_h}{\partial \ell} \right) d\varphi + \left(\frac{\partial v_h}{\partial \ell} \frac{\partial v_h}{\partial \ell^\perp} \right) \Bigg|_{\mathbf{x}_{F_3}^{(1)}}^{\mathbf{x}_{F_3}^{(2)}}.$$

Finally, combing (3.12) and (3.16), we have, on the boundary curved element K ,

$$(3.17) \quad \begin{aligned} \int_K (\Delta v_h)^2 dx &= \int_K |D^2 v_h|^2 dx + \sum_{i=1}^2 \left(2 \int_{F_i} \frac{\partial v_h}{\partial n} \frac{\partial^2 v_h}{\partial t_{F_i}^2} ds - \left(\frac{\partial v_h}{\partial n} \frac{\partial v_h}{\partial t_{F_i}} \right) \Bigg|_{\mathbf{x}_{F_i}^{(1)}}^{\mathbf{x}_{F_i}^{(2)}} \right) \\ &\quad + \int_{F_3} |\nabla v_h|^2 (\dot{\theta} - \chi) - 2 \frac{\partial v_h}{\partial \ell^\perp} \left(\frac{\partial \dot{v}_h}{\partial \ell} \right) ds + \left(\frac{\partial v_h}{\partial \ell} \frac{\partial v_h}{\partial \ell^\perp} \right) \Bigg|_{\mathbf{x}_{F_3}^{(1)}}^{\mathbf{x}_{F_3}^{(2)}}. \end{aligned}$$

Step 3: Using (3.11) and (3.17), we now sum over all the triangles $K \in \mathcal{T}_h$ to obtain

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K (\Delta v_h)^2 dx &= \sum_{K \in \mathcal{T}_h} \int_K |D^2 v_h|^2 dx \\ &\quad + 2 \sum_{F \in \mathcal{F}_h^i} \int_F \left[\frac{\partial v_h}{\partial n} \right] \frac{\partial^2 v_h}{\partial t_F^2} ds - \underbrace{\sum_{F \in \mathcal{F}_h^i} \left(\frac{\partial v_h^+}{\partial n^+} \frac{\partial v_h^+}{\partial t_F} + \frac{\partial v_h^-}{\partial n^-} \frac{\partial v_h^-}{\partial t_F} \right)}_{J_1} \Bigg|_{\mathbf{x}_F^{(1)}}^{\mathbf{x}_F^{(2)}} \\ &\quad + \sum_{F \in \mathcal{F}_h^\partial} \int_F |\nabla v_h|^2 (\dot{\theta} - \chi) - 2 \frac{\partial v_h}{\partial \ell^\perp} \left(\frac{\partial \dot{v}_h}{\partial \ell} \right) ds + \underbrace{\sum_{F \in \mathcal{F}_h^\partial} \left(\frac{\partial v_h}{\partial \ell} \frac{\partial v_h}{\partial \ell^\perp} \right)}_{J_2} \Bigg|_{\mathbf{x}_F^{(1)}}^{\mathbf{x}_F^{(2)}}. \end{aligned}$$

For any interior edge $F = K^+ \cap K^-$, the C^1 -continuity on the vertices implies that

$$\frac{\partial v_h^+}{\partial t_F}(\mathbf{x}) = \frac{\partial v_h^-}{\partial t_F}(\mathbf{x}), \quad \frac{\partial v_h^+}{\partial n^+}(\mathbf{x}) = -\frac{\partial v_h^-}{\partial n^-}(\mathbf{x}),$$

here $\mathbf{x} \in \mathcal{N}_h$ is the vertex of F . This shows that $J_1 = 0$. Let $\mathbf{x} = F^+ \cap F^-$ be a boundary vertex with $F^+, F^- \in \mathcal{F}_h^\partial$, we have

$$J_2 = \sum_{\mathbf{x} \in \mathcal{N}_h^\partial} \left(\left(\frac{\partial v_h}{\partial \ell} \frac{\partial v_h}{\partial \ell^\perp} \right) \Big|_{F^+} - \left(\frac{\partial v_h}{\partial \ell} \frac{\partial v_h}{\partial \ell^\perp} \right) \Big|_{F^-} \right) (\mathbf{x}).$$

Similarly, the C^1 -continuity on the vertices leads to $J_2 = 0$, which concludes the proof. \square

4. Numerical scheme and analysis. In this section, we will give the numerical scheme for approximating (1.1). We first define a broken norm as

$$\|v\|_h^2 := \sum_{K \in \mathcal{T}_h} |v|_{H^2(K)}^2 + \chi_0 \sum_{F \in \mathcal{F}_h^\partial} \int_F |\nabla v|^2 ds,$$

where we recall $\chi_0 = \min_{\partial\Omega}(\theta - \chi)$. For any $v \in V_{h,0}^\ell$, $\|v\|_h = 0$ implies that $D^2v|_K \equiv 0$ on each triangle $K \in \mathcal{T}_h$. Due to C^1 -continuity on the vertices, we immediately know that v is a linear polynomial on Ω . Since $\sum_{F \in \mathcal{F}_h^\partial} \int_F |\nabla v|^2 ds = 0$ and $\int_\Omega v dx = 0$, we conclude that $v \equiv 0$. Hence, $\|\cdot\|_h$ is a norm on $V_{h,0}^\ell$.

In light of (2.13), we define the bilinear form $b_h : V_h \times V_h \rightarrow \mathbb{R}$ as

$$(4.1) \quad b_h(w_h, v_h) := \sum_{K \in \mathcal{T}_h} (\gamma A : D^2 w_h, \Delta v_h)_K + \frac{2 - \sqrt{1 - \varepsilon}}{2} s_h(w_h, v_h),$$

where the stabilized term $s_h : V_h \times V_h \rightarrow \mathbb{R}$ is defined by

$$s_h(w_h, v_h) := -2 \sum_{F \in \mathcal{F}_h^i} \int_F \left[\frac{\partial w_h}{\partial n} \right] \frac{\partial^2 v_h}{\partial t_F^2} ds + 2 \sum_{F \in \mathcal{F}_h^\partial} \int_F \left(\frac{\partial w_h}{\partial \ell} \right) \frac{\partial v_h}{\partial \ell^\perp} ds,$$

which is naturally induced by the jump and boundary terms in the discrete Miranda-Talenti-type identity (3.9). We are now ready to define the numerical scheme as follows: Find $u_h \in V_{h,0}^\ell$ such that

$$(4.2) \quad b_h(u_h, v_h) = l_h(v_h) \quad \forall v_h \in V_{h,0}^\ell,$$

where the linear form $l_h : V_h \rightarrow \mathbb{R}$ is defined by $l_h(v_h) := \sum_{K \in \mathcal{T}_h} (\gamma f, \Delta v_h)_K$.

LEMMA 4.1 (consistence). *Let $u \in H_{\ell,0}^2(\Omega)$ be the H^2 strong solution of (1.1), then u satisfies the numerical scheme (4.2), i.e., $b_h(u, v_h) = l_h(v_h)$ for any $v_h \in V_{h,0}^\ell$.*

Proof. $u \in H_{\ell,0}^2(\Omega)$ leads to

$$\left[\frac{\partial u}{\partial n} \right]_F \equiv 0, \quad \forall F \in \mathcal{F}_h^i \quad \text{and} \quad \left(\frac{\partial u}{\partial \ell} \right) \Big|_F \equiv 0, \quad \forall F \in \mathcal{F}_h^\partial.$$

Thus we know $s_h(u, v_h) = 0$. Since u is the H^2 strong solution of (1.1), we have

$$b_h(u, v_h) = \sum_{K \in \mathcal{T}_h} (\gamma A : D^2 u, \Delta v_h)_K = \sum_{K \in \mathcal{T}_h} (\gamma f, \Delta v_h)_K = l_h(v_h).$$

That conclude the proof. \square

The following coercivity result follows directly from Theorem 3.9 (discrete Miranda-Talenti-type identity for oblique boundary conditions).

LEMMA 4.2 (coercivity). *Under the Assumption 2.6, let \mathcal{T}_h be a exact conforming triangulation of Ω , then*

$$(4.3) \quad b_h(v_h, v_h) \geq (1 - \sqrt{1 - \varepsilon}) \|v_h\|_h^2 \quad v_h \in V_h.$$

Here, recall ε is the parameter in the Cordes condition (2.1).

Proof. Combining Lemma 2.2 (property of Cordes condition) and the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} b_h(v_h, v_h) &= \sum_{K \in \mathcal{T}_h} \|\Delta v_h\|_{L^2(K)}^2 + \sum_{K \in \mathcal{T}_h} ((\gamma A - I) : D^2 v_h, \Delta v_h)_K \\ &\quad + \frac{2 - \sqrt{1 - \varepsilon}}{2} s_h(v_h, v_h) \\ &\geq \sum_{K \in \mathcal{T}_h} \|\Delta v_h\|_{L^2(K)}^2 - \sqrt{1 - \varepsilon} \sum_{K \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(K)} \|\Delta v_h\|_{L^2(K)} \\ &\quad + \frac{2 - \sqrt{1 - \varepsilon}}{2} s_h(v_h, v_h) \quad (\text{by Lemma 2.2}) \\ &\geq \frac{2 - \sqrt{1 - \varepsilon}}{2} \left(\sum_{K \in \mathcal{T}_h} \|\Delta v_h\|_{L^2(K)}^2 + s_h(v_h, v_h) \right) \\ &\quad - \frac{\sqrt{1 - \varepsilon}}{2} \sum_{K \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(K)}^2. \quad (\text{by Cauchy-Schwarz}) \end{aligned}$$

Applying (3.9) in Theorem 3.9 (discrete Miranda-Talenti-type identity for oblique boundary conditions), we have

$$\begin{aligned} b_h(v_h, v_h) &\geq (1 - \sqrt{1 - \varepsilon}) \sum_{K \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(K)}^2 \\ &\quad + \frac{2 - \sqrt{1 - \varepsilon}}{2} \chi_0 \sum_{F \in \mathcal{F}_h^\partial} \int_F |\nabla v|^2 ds \geq (1 - \sqrt{1 - \varepsilon}) \|v_h\|_h^2. \end{aligned}$$

This concludes the proof. \square

We note that the coercivity constant (namely, $1 - \sqrt{1 - \varepsilon}$) under the broken norm $\|\cdot\|_h$ is exactly the same as that for the PDE theory.

Remark 4.3 (approximation of ε in Cordes condition). We note that the bilinear form (4.1) explicitly uses the constant ε in the Cordes condition (2.2). As discussed in [28], if the optimal value of ε is not easy to compute, a simple modification of (4.1) reads

$$\tilde{b}_h(w_h, v_h) := \sum_{K \in \mathcal{T}_h} (\gamma A : D^2 w_h, \Delta v_h)_K + \frac{2 - \sqrt{1 - \tilde{\varepsilon}}}{2} s_h(w_h, v_h),$$

where $\tilde{\varepsilon}$ is an approximation of ε that satisfies $\sqrt{1 - \tilde{\varepsilon}} + \frac{1 - \varepsilon}{\sqrt{1 - \tilde{\varepsilon}}} < 2$. As a result, the coercivity constant becomes $1 - \frac{\sqrt{1 - \tilde{\varepsilon}}}{2} - \frac{1 - \varepsilon}{2\sqrt{1 - \tilde{\varepsilon}}}$. Even if there is no a priori estimate of ε , one may simply take $\tilde{\varepsilon} = 0$, which leads to the coercivity constant $\frac{\varepsilon}{2}$. We refer to [28, Section 4.4] for more details.

Next, we introduce a Poincaré-type estimate which is helpful in the analysis. The proof is given in Appendix B for the reader's convenience.

LEMMA 4.4 (Poincaré-type estimate). *Let $K \in \mathcal{T}_h$ be a curved triangle with a curved edge F . Let $u \in H^1(K)$, we have*

$$\|u\|_{L^2(K)}^2 \leq 2(1 + c_K)^2 \left(h_K \|u\|_{L^2(F)}^2 + h_K^2 |u|_{H^1(K)}^2 \right),$$

where c_K is defined in (3.2).

We are now ready to give the boundedness result by combing Lemma 3.7 (inverse estimate), Lemma 3.6 (trace estimate), and Lemma 4.4 (Poincaré-type estimate).

LEMMA 4.5 (boundedness). *Under the Assumption 2.6, let $\{\mathcal{T}_h\}_{h>0}$ be regular of order 2. Then, it holds that*

$$(4.4) \quad b_h(w_h, v_h) \leq C_b \|w_h\|_h \|v_h\|_h \quad w_h, v_h \in V_{h,0}^\ell.$$

Here the constant C_b depends on the the polynomial degree k , the shape-regular constant σ and c_1, c_2, c_3 .

Proof. Step 1: For the first term of $b_h(w_h, v_h)$, by Lemma 2.2 (property of Cordes condition) we have

$$I_1 := \left| \sum_{K \in \mathcal{T}_h} ((\gamma A - I) : D^2 w_h, \Delta v_h)_K + \sum_{K \in \mathcal{T}_h} (\Delta w_h, \Delta v_h)_K \right| \lesssim \|w_h\|_h \|v_h\|_h.$$

Step 2: We estimate the first term of $s_h(w_h, v_h)$. Note that $[\frac{\partial w_h}{\partial n}]$ vanishes on the two vertices of $F \in \mathcal{F}_h^i$, a Poincaré inequality on F leads to

$$(4.5) \quad \begin{aligned} I_2 &:= \left| \sum_{F \in \mathcal{F}_h^i} \int_F [\frac{\partial w_h}{\partial n}] \frac{\partial^2 v_h}{\partial t_F^2} ds \right| \\ &\leq \left(\sum_{F \in \mathcal{F}_h^i} h_F^{-1} \| [\frac{\partial w_h}{\partial n}] \|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^i} h_F \| \frac{\partial^2 v_h}{\partial t_F^2} \|_{L^2(F)}^2 \right)^{1/2} \\ &\lesssim \left(\sum_{F \in \mathcal{F}_h^i} h_F \| \frac{\partial}{\partial t_F} [\frac{\partial w_h}{\partial n}] \|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^i} h_F \| \frac{\partial^2 v_h}{\partial t_F^2} \|_{L^2(F)}^2 \right)^{1/2}. \end{aligned}$$

We define an indicator function $\delta : \mathcal{T}_h \rightarrow \{0, 1\}$ as $\delta(K) := \begin{cases} 1, & |\partial K \cap \partial \Omega| \neq 0 \\ 0, & |\partial K \cap \partial \Omega| = 0 \end{cases}$. For an interior edge $F = K^+ \cap K^-$, Lemma 3.6 (trace estimate) leads to

$$\begin{aligned} h_F \| \frac{\partial}{\partial t_F} [\frac{\partial w_h}{\partial n}] \|_{L^2(F)}^2 &\lesssim \sum_{K \in \{K^+, K^-\}} |w_h|_{H^2(K)}^2 + h_K^2 |w_h|_{H^3(K)}^2 \\ &\lesssim \sum_{K \in \{K^+, K^-\}} |w_h|_{H^2(K)}^2 + \delta(K) |w_h|_{H^1(K)}^2. \end{aligned}$$

In the last step, we use the standard inverse estimate [3] for the term $|w_h|_{H^3(K)}^2$ if K is a straight triangle. If K is a curved triangle, we apply Lemma 3.7 (inverse

estimate), which requires $\{T_h\}_{h>0}$ to be regular of order 2. Then, applying Lemma 4.4 (Poincaré-type estimate), we arrive at

$$(4.6) \quad \begin{aligned} \sum_{F \in \mathcal{F}_h^i} h_F \left\| \frac{\partial}{\partial t_F} \left[\frac{\partial w_h}{\partial n} \right] \right\|_{L^2(F)}^2 &\lesssim \sum_{K \in \mathcal{T}_h} |w_h|_{H^2(K)}^2 + \delta(K) |w_h|_{H^1(K)}^2 \\ &\lesssim \sum_{K \in \mathcal{T}_h} |w_h|_{H^2(K)}^2 + \sum_{F \in \mathcal{F}_h^\partial} \|\nabla w_h\|_{L^2(F)}^2 \lesssim \|w_h\|_h^2. \end{aligned}$$

Similar arguments as (4.6) lead to $\sum_{F \in \mathcal{F}_h^i} h_F \left\| \frac{\partial^2 v_h}{\partial t_F^2} \right\|_{L^2(F)}^2 \lesssim \|v_h\|_h^2$. This, combined with (4.6), leads to $I_2 \lesssim \|w_h\|_h \|v_h\|_h$.

Step 3: Now we estimate the second term of $s_h(w_h, v_h)$. By the boundary condition of $V_{h,0}^\ell$, we know $\int_F \left(\frac{\partial w_h}{\partial \ell} \right) ds = \frac{\partial w_h}{\partial \ell}(\mathbf{x}_F^{(2)}) - \frac{\partial w_h}{\partial \ell}(\mathbf{x}_F^{(1)}) = 0$ for any boundary edge F . Denote $c_F := \int_F \frac{\partial v_h}{\partial \ell^\perp} ds$, a Poincaré inequality on F leads to

$$(4.7) \quad \begin{aligned} I_3 &:= \left| \sum_{F \in \mathcal{F}_h^\partial} \int_F \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \frac{\partial v_h}{\partial \ell^\perp} ds \right| = \left| \sum_{F \in \mathcal{F}_h^\partial} \int_F \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \left(\frac{\partial v_h}{\partial \ell^\perp} - c_F \right) ds \right| \\ &\leq \left(\sum_{F \in \mathcal{F}_h^\partial} h_F \left\| \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \right\|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^\partial} h_F^{-1} \left\| \frac{\partial v_h}{\partial \ell^\perp} - c_F \right\|_{L^2(F)}^2 \right)^{1/2} \\ &\lesssim \left(\sum_{F \in \mathcal{F}_h^\partial} h_F \left\| \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \right\|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^\partial} h_F \left\| \left(\frac{\partial v_h}{\partial \ell^\perp} \right) \right\|_{L^2(F)}^2 \right)^{1/2}. \end{aligned}$$

For a boundary edge $F = \partial K \cap \partial \Omega$, combining Lemma 3.6 (trace estimate), Lemma 3.7 (inverse estimate) and (3.14), we have

$$\begin{aligned} h_F \left\| \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \right\|_{L^2(F)}^2 &\lesssim h_F \left(\|\nabla w_h\|_{L^2(F)}^2 + \|D^2 w_h\|_{L^2(F)}^2 \right) && \text{(by (3.14))} \\ &\lesssim |w_h|_{H^1(K)}^2 + (h_K^2 + 1) |w_h|_{H^2(K)}^2 + h_K^2 |w_h|_{H^3(K)}^2 && \text{(Lemma 3.6)} \\ &\lesssim |w_h|_{H^1(K)}^2 + |w_h|_{H^2(K)}^2. && \text{(Lemma 3.7)} \end{aligned}$$

By Lemma 4.4 (Poincaré-type estimate), we obtain

$$(4.8) \quad h_F \left\| \left(\frac{\partial \dot{w}_h}{\partial \ell} \right) \right\|_{L^2(F)}^2 \lesssim |w_h|_{H^2(K)}^2 + \|\nabla w_h\|_{L^2(F)}^2.$$

Similarly $h_F \left\| \left(\frac{\partial v_h}{\partial \ell^\perp} \right) \right\|_{L^2(F)}^2 \lesssim |v_h|_{H^2(K)}^2 + \|\nabla v_h\|_{L^2(F)}^2$. This, combined with (4.8), gives $I_3 \lesssim \|w_h\|_h \|v_h\|_h$. Then, the proof is concluded by combing the estimate for I_i , $i = 1, 2, 3$. \square

THEOREM 4.6. *Under the Assumption 2.6, let $u \in H_{\ell,0}^2(\Omega)$ be the H^2 strong solution of (1.1). Let $\{\mathcal{T}_h\}_{h>0}$ be regular of order 2. Then, there exists a unique*

solution $u_h \in V_{h,0}^\ell$ to (4.2) satisfying

$$\begin{aligned} & \left(\sum_{K \in \mathcal{T}_h} |u - u_h|_{H^2(K)}^2 \right)^{1/2} \\ & \leq \frac{C_{\text{err}}}{1 - \sqrt{1 - \varepsilon}} \inf_{w_h \in V_{h,0}^\ell} \left(\sum_{K \in \mathcal{T}_h} |u - w_h|_{H^2(K)}^2 + h_K^{-1} \|\nabla(u - w_h)\|_{L^2(\partial K)}^2 \right)^{1/2}. \end{aligned}$$

Here, the C_{err} depends on the polynomial degree k , the shape-regular constant σ and c_1, c_2, c_3 .

Proof. The existence and uniqueness of the numerical solution $u_h \in V_{h,0}^\ell$ follow from Lemma 4.2 (coercivity), Lemma 4.5 (boundedness), and the Lax-Milgram Theorem. For any $w_h \in V_{h,0}^\ell$, denote $\psi_h := u_h - w_h \in V_{h,0}^\ell$. Lemma 4.2 (coercivity) and Lemma 4.1 (consistence) lead to

$$\|\psi_h\|_h^2 \leq \frac{1}{1 - \sqrt{1 - \varepsilon}} b_h(u_h - w_h, \psi_h) = \frac{1}{1 - \sqrt{1 - \varepsilon}} b_h(u - w_h, \psi_h).$$

Here, we denote

$$\begin{aligned} b_h(u - w_h, \psi_h) & \lesssim \underbrace{\left| \sum_{K \in \mathcal{T}_h} (\gamma A : D^2(u - w_h), \Delta \psi_h)_K \right|}_{I_1} + \underbrace{\left| \sum_{F \in \mathcal{F}_h^i} \int_F \left[\frac{\partial(u - w_h)}{\partial n} \right] \frac{\partial^2 \psi_h}{\partial t_F^2} ds \right|}_{I_2} \\ & \quad + \underbrace{\left| \sum_{F \in \mathcal{F}_h^\partial} \int_F \left(\frac{\partial(u - w_h)}{\partial \ell} \right) \frac{\partial \psi_h}{\partial \ell^\perp} ds \right|}_{I_3}. \end{aligned}$$

For I_1 , similar arguments as *Step 1* in Lemma 4.5 (boundedness) lead to

$$I_1 \lesssim \left(\sum_{K \in \mathcal{T}_h} |u - w_h|_{H^2(K)}^2 \right)^{1/2} \|\psi_h\|_h.$$

For I_2 , we have

$$\begin{aligned} I_2 & \leq \left(\sum_{F \in \mathcal{F}_h^i} h_F^{-1} \left\| \left[\frac{\partial(u - w_h)}{\partial n} \right] \right\|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^i} h_F \left\| \frac{\partial^2 \psi_h}{\partial t_F^2} \right\|_{L^2(F)}^2 \right)^{1/2} \\ & \lesssim \left(\sum_{F \in \mathcal{F}_h^i} h_F^{-1} \left\| \left[\frac{\partial(u - w_h)}{\partial n} \right] \right\|_{L^2(F)}^2 \right)^{1/2} \|\psi_h\|_h. \end{aligned}$$

In the last step, we use the same argument as *Step 2* (4.6) in Lemma 4.5 (boundedness).

For I_3 , we first apply the integration by part on edge $F \in \mathcal{F}_h^\partial$. Due to the C^1 -continuity on vertices of the finite element space and the boundary condition of

$u \in H_{\ell,0}^2(\Omega)$, we obtain

$$\begin{aligned} I_3 &= \left| \sum_{F \in \mathcal{F}_h^\partial} \int_F \left(\frac{\partial(u - w_h)}{\partial \ell} \right) \frac{\partial \psi_h}{\partial \ell^\perp} ds \right| = \left| \sum_{F \in \mathcal{F}_h^\partial} \int_F \frac{\partial(u - w_h)}{\partial \ell} \left(\frac{\partial \dot{\psi}_h}{\partial \ell^\perp} \right) ds \right| \\ &\leq \left(\sum_{F \in \mathcal{F}_h^\partial} h_F^{-1} \left\| \frac{\partial(u - w_h)}{\partial \ell} \right\|_{L^2(F)}^2 \right)^{1/2} \left(\sum_{F \in \mathcal{F}_h^\partial} h_F \left\| \left(\frac{\partial \dot{\psi}_h}{\partial \ell^\perp} \right) \right\|_{L^2(F)}^2 \right)^{1/2} \\ &\lesssim \left(\sum_{F \in \mathcal{F}_h^\partial} h_F^{-1} \|\nabla(u - w_h)\|_{L^2(F)}^2 \right)^{1/2} \|\psi_h\|_h. \end{aligned}$$

In the last step, we use the same argument as *Step 3* (4.8) in Lemma 4.5 (boundedness). By combing the estimate of I_i , $i = 1, 2, 3$, we arrive

$$\|\psi_h\|_h \lesssim \left(\sum_{K \in \mathcal{T}_h} |u - w_h|_{H^2(K)}^2 + h_K^{-1} \|\nabla(u - w_h)\|_{L^2(\partial K)}^2 \right)^{1/2}.$$

This, combined with the triangular inequality, concludes the proof. \square

Taking $w_h = \Pi_h^\ell u$ in Theorem 4.6, and combining with (3.8b) and (3.8a) in Theorem 3.8 (approximation property of $V_{h,0}^\ell$), we have the following error estimate result.

COROLLARY 4.7. *Under the Assumption 2.6, assume that $\{\mathcal{T}_h\}_{h>0}$ is regular of order k , $u \in H^s(\Omega) \cap H_{\ell,0}^2(\Omega)$ is the H^2 strong solution of (1.1), and the oblique vector field ℓ is piecewise C^{s-1} with $2 \leq s \leq k+1$. It holds that*

$$(4.9) \quad \left(\sum_{K \in \mathcal{T}_h} |u - u_h|_{H^2(K)}^2 \right)^{1/2} \lesssim \frac{1}{1 - \sqrt{1 - \varepsilon}} \left(\sum_{K \in \mathcal{T}_h} h_K^{2s-4} \|u\|_{H^s(\omega_K)}^2 \right)^{1/2},$$

where the hidden constant depends on ℓ , the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

5. Numerical experiments. This section presents numerical experiments of the proposed methods for (1.1). We apply the curved Hermite element discussed in Section 3 with polynomial degree $k = 3$. The convergence history plots are logarithmically scaled in all the convergence order experiments.

5.1. Experiment 1. In the first experiment, we consider the problem (1.1) in the unit disk $\Omega := \{|x|^2 < 1\}$. The coefficient matrix is set to be $A := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. Take the oblique vector field ℓ to be the rotated normal vector, i.e.,

$$\ell := \sqrt{1/2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} n,$$

so that the rotation angle is $\theta = \pi/4$. To test the convergence rate, we consider the smooth solution

$$u = \sin(\pi(x_1^2 + x_2^2)) e^{(x_1^2 + x_2^2)} - \frac{\pi(e+1)}{\pi^2 + 1}.$$

The function $f := A : D^2u$ is directly calculated from the coefficient matrix and solution. A straightforward calculation shows that the compatibility constant $c = -\sqrt{2}\pi e$ and $\chi_0 = 1$.

We apply the numerical scheme (4.2) to the problem on a sequence of uniform triangulations $\{\mathcal{T}_h\}_{h>0}$. The discretization errors in the L^2 , H^1 and H^2 norm are reported in Table 1. We also display the approximations of the compatibility constant. The expected optimal convergence rate $\|D^2(u - u_h)\|_{L^2(\mathcal{T}_h)} = \mathcal{O}(h^{k-1})$ is observed, agreeing with Corollary 4.7. Further, the experiments indicate that the scheme converges with (sub-optimal) second-order convergence in both H^1 and L^2 norm. Convergence to the compatibility constant $c = -\sqrt{2}\pi e$ is also observed for the approximation c_h .

h	$\ u - u_h\ _{L^2}$	Order	$ u - u_h _{H^1}$	Order	$ u - u_h _{H^2}$	Order	c_h
2^{-2}	2.80e-02	0.00	2.88e-01	0.00	4.95	0.00	-12.128
2^{-3}	7.67e-03	1.87	6.84e-02	2.07	2.81	0.82	-12.086
2^{-4}	1.43e-03	2.43	1.20e-02	2.51	8.48e-01	1.73	-12.077
2^{-5}	2.97e-04	2.26	2.36e-03	2.35	2.21e-01	1.94	-12.077
2^{-6}	7.15e-05	2.05	5.58e-04	2.08	5.55e-02	1.99	-12.077

Table 1: Errors and observed convergence orders for Experiment 1.

5.2. Experiment 2. In the second experiment, let the domain $\Omega := \{|x|^2 < 1\}$. The coefficient matrix is set to be

$$A := \begin{pmatrix} 2 & \frac{x_1 x_2}{|x_1 x_2|} \\ \frac{x_1 x_2}{|x_1 x_2|} & 2 \end{pmatrix}.$$

A straightforward calculation shows the Cordes condition (2.2) is satisfied with $\varepsilon = 3/5$. We note that the coefficient matrix is discontinuous across the set $\{(x_1, x_2) \in \Omega : x_1 = 0 \text{ or } x_2 = 0\}$. Take the oblique vector field ℓ to be the anti-clockwise rotation of the normal vector by the angle $\theta = s + \pi/4$, i.e.,

$$\ell := (\cos(2s + \pi/4), \sin(2s + \pi/4))^T,$$

where $s := \arctan(x_2/x_1)$. The function f is chosen so that the solution is given by

$$u = 1/6(x_1^2 + x_2^2)^3 - 1/2(x_1^2 + x_2^2) + 5/24.$$

Directly calculation shows that the compatibility constant $c = 0$ and $\chi_0 = 2$.

The numerical result is displayed in Figure 1, which is in agreement with Corollary 4.7. The convergence orders are similar to Experiment 5.1. This experiment demonstrates the robustness of this proposed scheme with respect to the choice of oblique vector field and the discontinuous coefficient matrix.

5.3. Experiment 3. In the third experiment, let the domain $\Omega := \{|x|^2 < 1\}$. The coefficient matrix A is the same as in Experiment 5.2. The oblique vector field is given as

$$\ell := \sqrt{1/2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} n,$$




Fig. 1: Convergence orders for Experiment 2.

so that the rotation angle is $\theta = \pi/4$. Take f so that the solution is

$$u = (x_1^2 + x_2^2)e^{(x_1^2 + x_2^2)} - 1.$$

Here the compatibility constant $c = 2\sqrt{2}e$ and $\chi_0 = 1$. The numerical results on uniformly refined meshes are shown in Table 2. Similar convergence orders to Experiments 5.1 and 5.2 are observed. The convergence to the compatibility constant $c = 2\sqrt{2}e$ is also observed.

h	$\ u - u_h\ _{L^2}$	Order	$ u - u_h _{H^1}$	Order	$ u - u_h _{H^2}$	Order	c_h
2^{-2}	9.29e-03	0.00	1.11e-01	0.00	2.52	0.00	7.6997
2^{-3}	1.48e-03	2.65	1.80e-02	2.62	6.95e-01	1.86	7.6882
2^{-4}	5.51e-04	1.43	3.40e-03	2.41	1.70e-01	2.03	7.6880
2^{-5}	1.73e-04	1.67	8.25e-04	2.04	4.04e-02	2.08	7.6883
2^{-6}	4.81e-05	1.85	2.15e-04	1.94	9.64e-03	2.07	7.6884

Table 2: Errors and observed convergence orders for Experiment 3.

5.4. Experiment 4. In the fourth experiment, let $\Omega := \{x_1^2/4 + x_2^2 < 1\}$ be an ellipse domain. The coefficient matrix is the same as in Experiment 5.2. The oblique vector field ℓ is taken to be the tangential vector. We take the function f so that the solution is given by $u = 1/4 \sin(\pi(x_1^2/4 + x_2^2)) - 1/(2\pi)$. In this case, the compatibility constant $c = 0$ and $\chi_0 \geq 1/4$. We apply the numerical scheme (4.2) to the problem and refine the mesh uniformly. Similar convergence orders to the above experiments are observed. Numerical results and the mesh are shown in Figure 2.

Appendix A. Approximation property of finite element space. This appendix considers the construction of the quasi-interpolation operator $\Pi_h^\ell : H_{\ell,0}^2(\Omega) \rightarrow V_{h,0}^\ell$ and its optimal approximation property.

A.1. Construction of Π_h^ℓ . We consistently use the following correspondence in the sequel: $\mathbf{x} = F_K(\hat{\mathbf{x}})$, $v = \hat{v} \circ F_K^{-1}$. Let $N_h := \dim V_h$ and $\mu_i(\cdot)$, $1 \leq i \leq N_h$, be the global degrees of freedom of V_h . Denote \mathbf{a}_i , the node where $\mu_i(\cdot)$ takes its value.




Fig. 2: Convergence orders and mesh for Experiment 4.

Let s_i equal to 0 or 1, indicating whether $\mu_i(\cdot)$ takes a function or derivative value. Denote $\phi_i \in V_h$, the basis function of $\mu_i(\cdot)$. $\omega_i := \bigcup\{K : \hat{K} \cap \text{supp}(\phi_i) \neq \emptyset, K \in \mathcal{T}_h\}$, and $\#\omega_i := \#\{K : \hat{K} \cap \text{supp}(\phi_i) \neq \emptyset, K \in \mathcal{T}_h\}$. Note that $\#\omega_i$ is uniformly bounded if $\{\mathcal{T}_h\}_{h>0}$ is shape-regular. For any $K \in \mathcal{T}_h$, ω_K is the union of all ω_i containing K .

LEMMA A.1 (scale of ϕ_i). *Assume $\{\mathcal{T}_h\}_{h>0}$ is regular of order k . Then for $1 \leq i \leq N_h$,*

$$(A.1) \quad \|\phi_i\|_{H^m(K)} \lesssim h_K^{1-m+s_i} \quad 0 \leq m \leq k+1,$$

where $K \in \mathcal{T}_h$ is contained in ω_i and the hidden constant is depends on the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

Proof. Case 1: $s_i = 0$. In this case, $\mu_i(v) = v(\mathbf{a}_i)$. By Lemma 3.5 (scaling on curved triangles), we have $\|\phi_i\|_{H^m(K)} \lesssim h_K^{1-m} \|\hat{\phi}_i\|_{H^m(\hat{K})}$. It is easy to check that $\hat{\phi}_i = \phi_i \circ F_K$ is a basis function of $(\hat{K}, \mathcal{P}_k(\hat{K}), \hat{\Sigma})$, hence $\|\hat{\phi}_i\|_{H^m(\hat{K})} = \mathcal{O}(1)$.

Case 2: $s_i = 1$. In this case, $\mu_i(v) = \partial_j v(\mathbf{a}_i)$, where $j = 1$ or 2 and $\mathbf{a}_i \in \mathcal{N}_h$ is a vertex. The chain rule shows that $\hat{\phi}_i = \phi_i \circ F_K$ vanishes on all the degrees of freedom in $\hat{\Sigma}$ except

$$\hat{\partial}_j \hat{\phi}_i(\hat{\mathbf{a}}_i) = \partial_1 \phi_i(\mathbf{a}_i)(DF_K(\hat{\mathbf{a}}_i))_{1,j} + \partial_2 \phi_i(\mathbf{a}_i)(DF_K(\hat{\mathbf{a}}_i))_{2,j} \quad j = 1, 2.$$

Then, the norm equivalence on $\mathcal{P}_k(\hat{K})$ leads to

$$\begin{aligned} \|\hat{\phi}_i\|_{H^m(\hat{K})}^2 &\approx \sum_{\hat{\mu} \in \hat{\Sigma}} \hat{\mu}(\hat{\phi}_i)^2 = (\hat{\partial}_1 \hat{\phi}_i(\hat{\mathbf{a}}_i))^2 + (\hat{\partial}_2 \hat{\phi}_i(\hat{\mathbf{a}}_i))^2 \\ &\leq \sup_{\hat{x} \in \hat{K}} \|DF_K(\hat{x})\|^2 ((\partial_1 \phi_i(\mathbf{a}_i))^2 + (\partial_2 \phi_i(\mathbf{a}_i))^2) \lesssim h_K^2. \end{aligned}$$

This, combined with the scaling argument, yields $\|\phi_i\|_{H^m(K)} \lesssim h_K^{2-m}$. \square

Let $u \in L^2(K)$ and $\hat{u} := u \circ F_K$. Denote $Q_{\hat{K}} \hat{u}$ the L^2 projection of \hat{u} on $\mathcal{P}_k(\hat{K})$. Then, we define the operator $Q_K : L^2(K) \rightarrow P_K$ by

$$Q_K u := Q_{\hat{K}} \hat{u} \circ F_K^{-1}.$$

By the Lemma 3.5 (scaling on curved triangles) and the the Bramble-Hilbert Lemma [3], we have the following approximation property of Q_K .

LEMMA A.2 (approximation property of Q_K). *Let $\{\mathcal{T}_h\}_{h>0}$ is of regular of k , $u \in H^s(\Omega)$ with $2 \leq s \leq k+1$. Then we have*

$$(A.2) \quad \|u - Q_K u\|_{H^m(K)} \lesssim h_K^{s-m} \|u\|_{H^s(K)}, \quad m = 0, 1, 2.$$

Here the hidden constant depends on the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

We are now ready to define the quasi-interpolation operator $\Pi_h : L^2(\Omega) \rightarrow V_h$ by averaging the local L^2 projection, i.e.,

$$(A.3) \quad \mu_i(\Pi_h u) := \frac{1}{\#\omega_i} \sum_{j=1}^{\#\omega_i} \mu_i(Q_{K_j} u), \quad 1 \leq i \leq N_h.$$

Here, we write $\omega_i = \bigcup_{j=1}^{\#\omega_i} K_j$ with K_j and K_{j+1} sharing a common edge.

THEOREM A.3 (approximation property of Π_h). *Assume $\{\mathcal{T}_h\}_{h>0}$ is of regular of order k . Let $u \in H^s(\Omega)$ with $2 \leq s \leq k+1$. Then we have*

$$(A.4) \quad \|u - \Pi_h u\|_{H^m(K)} \lesssim h_K^{s-m} \|u\|_{H^s(\omega_K)}, \quad m = 0, 1, 2.$$

Here, the hidden constant depends on the polynomial degree k , the shape-regular constant σ and c_1, c_2, \dots, c_{k+1} .

Proof. For now, we number $\omega_1, \dots, \omega_n$ the sets ω_i such that $K \subset \omega_i$. Then, we have

$$\|Q_K u - \Pi_h u\|_{H^m(K)} \leq \sum_{i=1}^n |\mu_i(Q_K u) - \mu_i(\Pi_h u)| \|\phi_i\|_{H^m(K)}.$$

In the case of $\#\omega_i = 1$, we know that $\omega_i = K$ and $\mu_i(Q_K u) - \mu_i(\Pi_h u) = 0$. Below, we assume $\#\omega_i \geq 2$. By (A.3) and triangle inequality, we have

$$(A.5) \quad \begin{aligned} |\mu_i(Q_K u) - \mu_i(\Pi_h u)| &= \left| \frac{1}{\#\omega_i} \sum_{j=1}^{\#\omega_i} (\mu_i(Q_K u) - \mu_i(Q_{K_j} u)) \right| \\ &\lesssim \sum_{j=1}^{\#\omega_i-1} |\mu_i(Q_{K_j} u) - \mu_i(Q_{K_{j+1}} u)| \lesssim \sum_{\substack{F \in \mathcal{F}_h^i \\ \text{with } \mathbf{a}_i \in \bar{F}}} |Q_{K^+} u - Q_{K^-} u|_{W_\infty^{s_i}(F)}, \end{aligned}$$

where we denote $F = K^+ \cap K^-$, and recall \mathbf{a}_i is the node where μ_i takes its value, $s_i = 0$ or 1 . Note $(Q_{K^\pm} u)|_F = Q_{\hat{K}} \hat{u} \circ (F_{K^\pm}^{-1}|_F)$, where $F_{K^\pm}^{-1}|_F$ is an affine mapping (see Remark 3.3). Therefore, it is clear that $(Q_{K^\pm} u)|_F$ is indeed a polynomial. Hence, we have

$$(A.6) \quad \begin{aligned} |Q_{K^+} u - Q_{K^-} u|_{W_\infty^{s_i}(F)} &\lesssim h_F^{-\frac{1}{2}-s_i} \|Q_{K^+} u - Q_{K^-} u\|_{L^2(F)} \\ &\leq h_F^{-\frac{1}{2}-s_i} (\|Q_{K^+} u - u\|_{L^2(F)} + \|Q_{K^-} u - u\|_{L^2(F)}). \end{aligned}$$

Lemma 3.6 (trace estimate) and Lemma A.2 (approximation property of Q_K) lead to

$$(A.7) \quad \begin{aligned} \|Q_{K^\pm} u - u\|_{L^2(F)} &\lesssim \left(h_{K^\pm}^{-\frac{1}{2}} \|Q_{K^\pm} u - u\|_{L^2(K^\pm)} + h_{K^\pm}^{\frac{1}{2}} |Q_{K^\pm} u - u|_{H^1(K^\pm)} \right) \\ &\lesssim h_K^{s-\frac{1}{2}} \|u\|_{H^s(\omega_K)}. \end{aligned}$$

Combing the above estimates, we obtain $|\mu_i(Q_K u) - \mu_i(\Pi_h u)| \lesssim h_K^{s-1-s_i} \|u\|_{H^s(\omega_K)}$. This, combined with Lemma A.1 (scale of ϕ_i), leads to $\|Q_K u - \Pi_h u\|_{H^m(K)} \lesssim h_K^{s-m} \|u\|_{H^s(\omega_K)}$. Then the proof is concluded by triangle inequality and Lemma A.2 (approximation property of Q_K). \square

Next, we reorder the degrees of freedom such that for $i = N_0 + 1, \dots, N_h$, $\mu_i(v) = (\nabla v \cdot \mathbf{e}_i)(\mathbf{a}_i)$, where $\mathbf{a}_i \in \mathcal{N}_h^\partial$ is a boundary vertex and $\mathbf{e}_i = (1, 0)^T$ or $(0, 1)^T$. Then for any $u \in H_{\ell,0}^2(\Omega)$, denote $C_u := (\nabla u \cdot \ell)|_{\partial\Omega}$. We first define $w_h \in V_h$ as follows.

- For $i = 1, \dots, N_0$,

$$(A.8a) \quad \mu_i(w_h) := \mu_i(\Pi_h u) = \frac{1}{\#\omega_i} \sum_{j=1}^{\#\omega_i} \mu_i(Q_{K_j} u).$$

- For $i = N_0 + 1, \dots, N_h$,

$$(A.8b) \quad \mu_i(w_h) := \left((\mathbf{e}_i \cdot \ell) C_u + (\mathbf{e}_i \cdot \ell^\perp) \frac{1}{\#\omega_i} \sum_{j=1}^{\#\omega_i} \frac{\partial Q_{K_j} u}{\partial \ell^\perp} \right) (\mathbf{a}_i).$$

It can be easily verified that $(\nabla w_h \cdot \ell)(\mathbf{x}) = C_u$ for any $\mathbf{x} \in \mathcal{N}_h^\partial$. Finally, the quasi-interpolation operator $\Pi_h^\ell : H_{\ell,0}^2(\Omega) \rightarrow V_{h,0}^\ell$ is defined by

$$(A.8c) \quad \Pi_h^\ell u := w_h - \frac{1}{|\Omega|} \int_\Omega w_h dx.$$

A.2. Proof of Theorem 3.8. By (A.8c), we know $|\Pi_h u - \Pi_h^\ell u|_{H^m(K)} = |\Pi_h u - w_h|_{H^m(K)}$, for $m = 1, 2$. Note that if K does not contain boundary vertex, then $(\Pi_h u - w_h)|_K \equiv 0$ by (A.8a). Below, we assume K contains a boundary vertex. We number $\omega_1, \dots, \omega_n$ the sets ω_i satisfying $K \subset \omega_i$ and $N_0 + 1 \leq i \leq N_h$. Then,

$$|\Pi_h u - w_h|_{H^m(K)} \leq \sum_{i=1}^n |\mu_i(\Pi_h u) - \mu_i(w_h)| |\phi_i|_{H^m(K)}.$$

For any $1 \leq i \leq n$, there exists $K^{(i)} \subset \omega_i$ with edge $F^{(i)}$ satisfying $\mathbf{a}_i \in F^{(i)} \subset \partial\Omega$. Then, (A.8b) and triangle inequality leads to

$$\begin{aligned} |\mu_i(\Pi_h u) - \mu_i(w_h)| &\leq \left| \frac{1}{\#\omega_i} \sum_{j=1}^{\#\omega_i} \frac{\partial Q_{K_j} u}{\partial \ell}(\mathbf{a}_i) - C_u \right| \\ &\lesssim \underbrace{\sum_{j=1}^{\#\omega_i-1} \left| \frac{\partial Q_{K_j} u}{\partial \ell}(\mathbf{a}_i) - \frac{\partial Q_{K_{j+1}} u}{\partial \ell}(\mathbf{a}_i) \right|}_{I_1} + \underbrace{\left| \frac{\partial Q_{K^{(i)}} u}{\partial \ell}(\mathbf{a}_i) - C_u \right|}_{I_2}. \end{aligned}$$

Similar arguments as (A.6) and (A.7) in Theorem A.3 (approximation property of Π_h) yield

$$I_1 \lesssim \sum_{\substack{F \in \mathcal{F}_h^i \\ \text{with } \mathbf{a}_i \in \bar{F}}} |Q_{K^+} u - Q_{K^-} u|_{W_\infty^1(F)} \lesssim h_K^{s-2} \|u\|_{H^s(\omega_K)}.$$

For the estimate of I_2 , let $\hat{F} := F_{K^{(i)}}^{-1}(F^{(i)})$ with vertex $\hat{\mathbf{a}} := F_{K^{(i)}}^{-1}(\mathbf{a}_i)$. Define $\hat{\ell} : \hat{F} \rightarrow \mathbb{R}^2$ and $\hat{\eta} : \hat{F} \rightarrow \mathbb{R}^2$ by

$$(A.9) \quad \hat{\ell}(\hat{x}) := (\ell \circ F_{K^{(i)}})(\hat{x}) \quad \text{and} \quad \hat{\eta}(\hat{x}) := (DF_{K^{(i)}}(\hat{x}))^{-1} \hat{\ell}(\hat{x}) \quad \hat{x} \in \hat{F}.$$

The regularity of $\{\mathcal{T}_h\}_{h>0}$ and ℓ implies that $\hat{\eta} \in C^{s-1}(\hat{F}, \mathbb{R}^2)$. Now, pulling back I_2 to \hat{F} , and applying the chain rule yields

$$(A.10a) \quad I_2 = |(\nabla Q_{K^{(i)}} u \cdot \ell)(\mathbf{a}_i) - C_u| = |(\hat{\nabla} Q_{\hat{K}} \hat{u} \cdot \hat{\eta})(\hat{\mathbf{a}}) - C_u|,$$

and

$$(A.10b) \quad C_u = (\nabla u \cdot \ell)|_{F^{(i)}} = (\hat{\nabla} \hat{u} \cdot \hat{\eta})|_{\hat{F}}.$$

Let $\hat{\mathcal{I}}\hat{\eta} \in \mathcal{P}_k(\hat{F}; \mathbb{R}^2)$ be the Lagrange interpolation of $\hat{\eta}$ satisfying $\hat{\mathcal{I}}\hat{\eta}(\hat{\mathbf{a}}) = \hat{\eta}(\hat{\mathbf{a}})$. By (A.10) and the norm equivalence on the polynomial space, we arrive at

$$\begin{aligned} I_2 &= |(\hat{\nabla} Q_{\hat{K}} \hat{u} \cdot \hat{\eta})(\hat{\mathbf{a}}) - C_u| = |(\hat{\nabla} Q_{\hat{K}} \hat{u} \cdot \hat{\mathcal{I}}\hat{\eta})(\hat{\mathbf{a}}) - C_u| \\ &\lesssim \|\hat{\nabla} Q_{\hat{K}} \hat{u} \cdot \hat{\mathcal{I}}\hat{\eta} - C_u\|_{L^2(\hat{F})} = \|\hat{\nabla} Q_{\hat{K}} \hat{u} \cdot \hat{\mathcal{I}}\hat{\eta} - \hat{\nabla} \hat{u} \cdot \hat{\eta}\|_{L^2(\hat{F})} \\ &\leq \underbrace{\|\hat{\nabla} Q_{\hat{K}} \hat{u} - \hat{\nabla} \hat{u}\|_{L^2(\hat{F})} \|\hat{\mathcal{I}}\hat{\eta}\|_{L^\infty(\hat{F})}}_{J_1} + \underbrace{\|\hat{\nabla} \hat{u}\|_{L^2(\hat{F})} \|\hat{\mathcal{I}}\hat{\eta} - \hat{\eta}\|_{L^\infty(\hat{F})}}_{J_2}. \end{aligned}$$

For J_1 , we have $\|\hat{\mathcal{I}}\hat{\eta}\|_{L^\infty(\hat{F})} \leq \|\hat{\eta}\|_{L^\infty(\hat{F})} = \|(DF_{K^{(i)}})^{-1} \hat{\ell}\|_{L^\infty(\hat{F})} \lesssim h_{K^{(i)}}^{-1} \|\ell\|_{L^\infty(F^{(i)})}$. Lemma 3.6 (trace estimate), Lemma 3.5 (scaling on curved triangles), and Lemma A.2 (approximation property of Q_K) give

$$\begin{aligned} \|\hat{\nabla} Q_{\hat{K}} \hat{u} - \hat{\nabla} \hat{u}\|_{L^2(\hat{F})} &\lesssim |Q_{\hat{K}} \hat{u} - \hat{u}|_{H^1(\hat{K})} + |Q_{\hat{K}} \hat{u} - \hat{u}|_{H^2(\hat{K})} \\ &\lesssim \|Q_{K^{(i)}} u - u\|_{H^1(K^{(i)})} + h_{K^{(i)}} \|Q_{K^{(i)}} u - u\|_{H^2(K^{(i)})} \\ &\lesssim h_{K^{(i)}}^{s-1} \|u\|_{H^s(\omega_K)}, \end{aligned}$$

which leads to $J_1 \lesssim h_{K^{(i)}}^{s-2} \|u\|_{H^s(\omega_K)}$.

Similarly, for J_2 , we have

$$\|\hat{\nabla} \hat{u}\|_{L^2(\hat{F})} \lesssim |\hat{u}|_{H^1(\hat{K})} + |\hat{u}|_{H^2(\hat{K})} \lesssim \|u\|_{H^2(K^{(i)})} \leq \|u\|_{H^s(\omega_K)}.$$

Recall the definition of $\hat{\eta}$ in (A.9), the Leibniz rule leads to

$$|\hat{\eta}|_{W_\infty^{s-1}(\hat{F})} \lesssim \sum_{j=0}^{s-1} |(DF_{K^{(i)}}(\hat{x}))^{-1}|_{W_\infty^j(\hat{F})} |\hat{\ell}(\hat{x})|_{W_\infty^{s-1-j}(\hat{F})}.$$


By the fact that ℓ is piecewise C^{s-1} and $\{\mathcal{T}_h\}_{h>0}$ is regular of order k , the standard scaling argument on the curved element (cf. [2, Lemma 2.2, Lemma 2.3]) gives

$$|(DF_{K^{(i)}}(\hat{x}))^{-1}|_{W_\infty^j(\hat{F})} \lesssim h_{F^{(i)}}^{j-1} \quad \text{and} \quad |\hat{\ell}(\hat{x})|_{W_\infty^{s-1-j}(\hat{F})} \lesssim h_{F^{(i)}}^{s-1-j} \|\ell\|_{W_\infty^{s-1-j}(F^{(i)})}.$$

Therefore, we have $\|\hat{\eta} - \hat{\mathcal{I}}\hat{\eta}\|_{L^\infty(\hat{F})} \lesssim |\hat{\eta}|_{W_\infty^{s-1}(\hat{F})} \lesssim h_{F^{(i)}}^{s-2}$. Combining the two above estimates, we obtain $J_2 \lesssim h_{K^{(i)}}^{s-2} \|u\|_{H^s(\omega_K)}$. Thus, we conclude $I_2 \lesssim J_1 + J_2 \lesssim h_{K^{(i)}}^{s-2} \|u\|_{H^s(\omega_K)}$.

Combining the estimate of I_1 , I_2 , and (A.1) in Lemma A.1 (scale of ϕ_i), we obtain $|\Pi_h u - w_h|_{H^m(K)} \lesssim h_K^{s-m} \|u\|_{H^s(\omega_K)}$. Then, (3.8a) is obtained by triangle inequality and Theorem A.3 (approximation property of Π_h). (3.8b) follows from Lemma 3.6 (trace estimate) and (3.8a). The proof is thus complete.

Appendix B. Proof of Lemma 4.4. Let K be a curved triangle with vertices P_j , $j = 1, 2, 3$ (see Figure 3), $F = \widehat{P_2 P_3}$ be the curved edge.

Fig. 3: curved triangle K

Let (z_1, z_2) be the coordinates of point P_1 . We have

$$\operatorname{div}\left(\begin{pmatrix} x_1 - z_1 \\ x_2 - z_2 \end{pmatrix} u^2\right) = 2u^2 + 2u((x_1 - z_1)\partial_1 u + (x_2 - z_2)\partial_2 u).$$

By integrating over K , we have

$$\begin{aligned} & \int_K u^2 dx \\ &= \frac{1}{2} \int_K \operatorname{div}\left(\begin{pmatrix} x_1 - z_1 \\ x_2 - z_2 \end{pmatrix} u^2\right) dx - \int_K u((x_1 - z_1)\partial_1 u + (x_2 - z_2)\partial_2 u) dx \\ &\leq \frac{1}{2} \int_{\partial K} \mathbf{n} \cdot \begin{pmatrix} x_1 - z_1 \\ x_2 - z_2 \end{pmatrix} u^2 ds + \operatorname{diam}(K) \int_K |u|(|\partial_1 u| + |\partial_2 u|) dx. \end{aligned}$$

By Cauchy-Schwarz inequality, we get

$$\operatorname{diam}(K) \int_K |u|(|\partial_1 u| + |\partial_2 u|) dx \leq \frac{1}{2} \int_K u^2 dx + \operatorname{diam}(K)^2 \int_K |\nabla u|^2 dx.$$

Note that $\mathbf{n} \cdot (x_1 - z_1, x_2 - z_2)^T \equiv 0$ on $\overline{P_1 P_2}$ and $\overline{P_3 P_1}$, thus

$$\int_{\partial K} \mathbf{n} \cdot \begin{pmatrix} x_1 - z_1 \\ x_2 - z_2 \end{pmatrix} u^2 ds = \int_F \mathbf{n} \cdot \begin{pmatrix} x_1 - z_1 \\ x_2 - z_2 \end{pmatrix} u^2 ds \leq \operatorname{diam}(K) \int_F u^2 ds.$$

In summary, we have

$$\int_K u^2 dx \leq \frac{1}{2} \operatorname{diam}(K) \int_F u^2 ds + \frac{1}{2} \int_K u^2 dx + \operatorname{diam}(K)^2 \int_K |\nabla u|^2 dx,$$

or

$$\int_K u^2 dx \leq \operatorname{diam}(K) \int_F u^2 ds + 2 \operatorname{diam}(K)^2 \int_K |\nabla u|^2 dx.$$

We then conclude the proof by applying (3.3).

Acknowledgments. The authors would like to express their gratitude to Prof. Jun Hu in Peking University for his helpful discussions.

REFERENCES

- [1] J. W. BARRETT AND C. M. ELLIOTT, *Fixed mesh finite element approximations to a free boundary problem for an elliptic equation with an oblique derivative boundary condition*, Computers & mathematics with applications, 11 (1985), pp. 335–345.

- [2] C. BERNARDI, *Optimal finite-element interpolation on curved domains*, SIAM Journal on Numerical Analysis, 26 (1989), pp. 1212–1240.
- [3] S. BRENNER AND R. SCOTT, *The Mathematical Theory of Finite Element Methods*, vol. 15, Springer Science & Business Media, Switzerland, 2007.
- [4] S. C. BRENNER AND E. L. KAWECKI, *Adaptive c^0 interior penalty methods for hamilton–jacobi–bellman equations with cordes coefficients*, Journal of Computational and Applied Mathematics, (2020), p. 113241.
- [5] Z. FAŠKOVÁ, R. ČUNDERLÍK, AND K. MIKULA, *Finite element method for solving geodetic boundary value problems*, Journal of Geodesy, 84 (2010), pp. 135–144.
- [6] X. FENG, L. HENNINGS, AND M. NEILAN, *Finite element methods for second order linear elliptic partial differential equations in non-divergence form*, Mathematics of Computation, 86 (2017), pp. 2025–2051.
- [7] X. FENG, M. NEILAN, AND S. SCHNAKE, *Interior penalty discontinuous galerkin methods for second order linear non-divergence form elliptic pdes*, Journal of Scientific Computing, 74 (2018), pp. 1651–1676.
- [8] D. GALLISTL, *Variational formulation and numerical analysis of linear elliptic equations in nondivergence form with Cordes coefficients*, SIAM Journal on Numerical Analysis, 55 (2017), pp. 737–757.
- [9] D. GALLISTL, *Numerical approximation of planar oblique derivative problems in nondivergence form*, Mathematics of computation, 88 (2019), pp. 1091–1119.
- [10] D. GALLISTL AND E. SÜLI, *Mixed finite element approximation of the Hamilton–Jacobi–Bellman equation with Cordes coefficients*, SIAM Journal on Numerical Analysis, 57 (2019), pp. 592–614.
- [11] E. L. KAWECKI, *A dgfem for nondivergence form elliptic equations with cordes coefficients on curved domains*, Numerical Methods for Partial Differential Equations, 35 (2019), pp. 1717–1744.
- [12] E. L. KAWECKI, *A discontinuous galerkin finite element method for uniformly elliptic two dimensional oblique boundary-value problems*, SIAM Journal on Numerical Analysis, 57 (2019), pp. 751–778.
- [13] E. L. KAWECKI AND I. SMEARS, *Convergence of adaptive discontinuous galerkin and c^0 -interior penalty finite element methods for hamilton–jacobi–bellman and isaacs equations*, Foundations of Computational Mathematics, (2021).
- [14] E. L. KAWECKI AND I. SMEARS, *Unified analysis of discontinuous galerkin and c^0 -interior penalty finite element methods for hamilton–jacobi–bellman and isaacs equations*, ESAIM: Mathematical Modelling and Numerical Analysis, 55 (2021), pp. 449–478.
- [15] B. L. KEYFITZ AND G. M. LIEBERMAN, *A proof of existence of perturbed steady transonic shocks via a free boundary problem*, Communications on Pure and Applied Mathematics, 53 (2000), pp. 0001–0028.
- [16] O. LAKKIS AND T. PRYER, *A finite element method for second order nonvariational elliptic problems*, SIAM Journal on Scientific Computing, 33 (2011), pp. 786–801.
- [17] O. LAKKIS AND T. PRYER, *A finite element method for nonlinear elliptic problems*, SIAM Journal on Scientific Computing, 35 (2013), pp. A2025–A2045.
- [18] A. MAUGERI, D. K. PALAGACHEV, AND L. G. SOFTOVA, *Elliptic and Parabolic Equations with Discontinuous Coefficients*, vol. 109, WILEY-VCH Verlag GmbH & Co., Berlin, 2000.
- [19] M. MEDL’A, K. MIKULA, R. ČUNDERLÍK, AND M. MACÁK, *Numerical solution to the oblique derivative boundary value problem on non-uniform grids above the earth topography*, Journal of Geodesy, 92 (2018), pp. 1–19.
- [20] R. H. NOCHETTO AND W. ZHANG, *Discrete abp estimate and convergence rates for linear elliptic equations in non-divergence form*, Foundations of Computational Mathematics, 18 (2018), pp. 537–593.
- [21] D. K. PALAGACHEV, *The poincaré problem in l p -sobolev spaces ii: full dimension degeneracy*, Communications in Partial Differential Equations, 33 (2008), pp. 209–234.
- [22] W. QIU AND S. ZHANG, *Adaptive first-order system least-squares finite element methods for second-order elliptic equations in nondivergence form*, SIAM Journal on Numerical Analysis, 58 (2020), pp. 3286–3308.
- [23] I. SMEARS AND E. SÜLI, *Discontinuous Galerkin finite element approximation of nondivergence form elliptic equations with Cordes coefficients*, SIAM Journal on Numerical Analysis, 51 (2013), pp. 2088–2106.
- [24] I. SMEARS AND E. SÜLI, *Discontinuous Galerkin finite element approximation of Hamilton–Jacobi–Bellman equations with Cordes coefficients*, SIAM Journal on Numerical Analysis, 52 (2014), pp. 993–1016.
- [25] J. URBAS, *On the second boundary value problem for equations of Monge–Ampère type*, J. Reine

- Angew. Math., 487 (1997), pp. 115–124.
- [26] C. WANG AND J. WANG, *A primal-dual weak galerkin finite element method for second order elliptic equations in non-divergence form*, Mathematics of Computation, 87 (2018), pp. 515–545.
 - [27] G. WEN AND C. YANG, *Finite element solutions of the oblique derivative problem for elliptic complex equations of second order*, Sichuan Shifan Daxue Xuebao Ziran Kexue Ban, 17 (1994), pp. 20–28.
 - [28] S. WU, *C^0 finite element approximations of linear elliptic equations in non-divergence form and Hamilton–Jacobi–Bellman equations with Cordes coefficients*, Calcolo, 58 (2021).
 - [29] Y. ZHENG, *A global solution to a two-dimensional riemann problem involving shocks as free boundaries*, Acta Mathematicae Applicatae Sinica, 19 (2003), pp. 559–572.
 - [30] M. ZLÁMAL, *Curved elements in the finite element method. i*, SIAM Journal on Numerical Analysis, 10 (1973), pp. 229–240.