

Results for the Mobius Function and Liouville Function over the Shifted Primes

N. A. Carella

Abstract: This article provides new asymptotic results for the summatory Mobius function $\sum_{p \leq x} \mu(p+a) = O(x(\log x)^{-c})$ and the summatory Liouville function $\sum_{p \leq x} \lambda(p+a) = O(x(\log x)^{-c})$ over the shifted primes, where $a \neq 0$ is a fixed parameter, and $c > 1$ is an arbitrary constant. These results improve the current estimates $\sum_{p \leq x} \mu(p+a) = (1-\delta)\pi(x)$, and $\sum_{p \leq x} \lambda(p+a) = (1-\delta)\pi(x)$ for $\delta > 0$, respectively. Furthermore, a conditional proof for the autocorrelation function $\sum_{p \leq x} \mu(p+a)\mu(p+b) = O(x(\log x)^{-c})$, and an unconditional proof for the autocorrelation function $\sum_{p \leq x} \lambda(p+a)\lambda(p+b) = O(x(\log x)^{-c})$ over the shifted primes, where $a \neq b$, are also included.

Contents

1	Introduction	3
2	Results for the Mobius Function over the Shifted Primes	5
2.1	Average Orders of Mobius Functions	5
2.2	Mean Values and Hypotheses over the Shifted Primes	5
2.3	Squarefree Shifted Primes	6
2.4	Nonlinear Autocorrelation Functions Results	7
2.5	Single-Sign Pattern Mobius Characteristic Functions	9
2.6	Single-Sign Pattern Mobius Counting Functions	10
2.7	Result for the Summatory Mobius Function	11
2.8	Double-Sign Patterns Mobius Characteristic Functions	12
2.9	Double-Sign Patterns Mobius Counting Functions	12
2.10	Result for the Mobius Autocorrelation Function	14
3	Higher Order Mobius Autocorrelation Functions	15
3.1	Triple-Sign Patterns Mobius Characteristic Functions	15
3.2	Triple-Sign Patterns Mobius Counting Functions	16
3.3	Result for the Mobius Autocorrelation Function	18

July 6, 2022

AMS MSC: Primary 11N37, Secondary 11L20.

Keywords: Shifted prime; Arithmetic function; Mobius function; Liouville function; vonMangoldt function; Chowla conjecture; Sarnak conjecture.

4	Results for the Liouville Function over the Shifted Primes	18
4.1	Average Orders of Liouville Functions	18
4.2	Single-Sign Patterns Liouville Characteristic Functions	19
4.3	Single-Sign Pattern Liouville Counting Functions	20
4.4	Result for Liouville Summatory Function	21
4.5	Double-Sign Patterns Liouville Characteristic Functions	22
4.6	Double-Sign Patterns in Liouville Counting Functions	22
4.7	Result for the Liouville Autocorrelation Function	24
5	Higher Order Liouville Autocorrelation Functions	24
5.1	Triple Patterns Liouville Characteristic Functions	24
5.2	Triple-Sign Patterns Liouville Counting Functions	25
5.3	Result for the Liouville Autocorrelation Function	27
6	Hardy-Littlewood-Chowla Conjecture	27
6.1	Twisted vonMangoldt Function over the Squarefree Shifted Primes . .	28
6.2	Signed vonMangoldt Function over the Shifted Primes	29
6.3	Twisted vonMangoldt Function over the Shifted Primes	30
7	Results for the Mobius Autocorrelation Function over the Integers	30
7.1	Nonlinear Autocorrelation Functions Results	31
7.2	Double-Sign Patterns Characteristic Functions	33
7.3	Double-Sign Patterns Counting Functions	33
7.4	Result for the Mobius Autocorrelation Function over the Integers . .	35
8	Results for the Liouville Autocorrelation Function over the Integers	36
8.1	Double-Sign Patterns Characteristic Functions	37
8.2	Double-Sign Patterns Counting Functions	37
8.3	Result for the Liouville Autocorrelation Function over the Integers . .	39
9	Open Problems	40
9.1	Mobius autocorrelation function problems	40
9.2	Liouville autocorrelation function problems	42
9.3	Intercorrelation functions problems	43
9.4	Hardy-Littlewood-Chowla conjecture problems	43

1 Introduction

The Mobius function $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ is defined by

$$\mu(n) = \begin{cases} (-1)^v & n = p_1 p_2 \cdots p_v \\ 0 & n \neq p_1 p_2 \cdots p_v, \end{cases} \quad (1)$$

where the $p_i \geq 2$ are primes. Similarly, the Liouville function $\lambda : \mathbb{N} \rightarrow \{-1, 1\}$ is defined by

$$\lambda(n) = (-1)^{p_1^{v_1} p_2^{v_2} \cdots p_w^{v_w}}, \quad (2)$$

where the $p_i \geq 2$ are primes, and $n = p_1^{v_1} p_2^{v_2} \cdots p_w^{v_w}$. The Mobius autocorrelation function

$$\sum_{n \leq x} \mu(n) \mu(n+a) \quad (3)$$

and the Liouville autocorrelation function

$$\sum_{n \leq x} \lambda(n) \lambda(n+a) \quad (4)$$

are topics of current research in several area of Mathematics, [32], [30], [21], et alii. Restricting the autocorrelation functions of multiplicative functions (3) over the integers to the shifted primes reduce these functions to standard arithmetic averages over the shifted primes. For example, (3) reduces to

$$\sum_{p \leq x} \mu(p) \mu(p+a) = - \sum_{p \leq x} \mu(p+a). \quad (5)$$

Similarly, an autocorrelation function of degree 3,

$$\sum_{n \leq x} \mu(n) \mu(n+a) \mu(n+b), \quad (6)$$

where $a, b \neq 0$ such that $a \neq b$ are small fixed integers, over the integers reduces to an autocorrelation function of degree 2,

$$- \sum_{p \leq x} \mu(p+a) \mu(p+b) \quad (7)$$

over the shifted primes. Accordingly, these two open problems are equivalent.

Currently, the best asymptotic result for this summatory function is

$$\sum_{p \leq x} \mu(p+a) = (1-\delta)\pi(x), \quad (8)$$

where $\delta > 0$ is a constant, see [8, Theorem 1], and the restricted average

$$\sum_{a \leq z} \sum_{p \leq x} \mu(p+a) = o(z\pi(x)), \quad (9)$$

see [16] for extensive details on recent developments on this topic. It is expected that $\sum_{p \leq x} \mu(p+a) = o(\pi(x))$. This note proposes the first nontrivial upper bound.

Theorem 1.1. *Let $c > 1$ be an arbitrary constant, and let $x > 1$ be a large number. If $a \neq 0$ is a small fixed integer, then*

$$\sum_{p \leq x} \mu(p + a) = O\left(\frac{x}{(\log x)^c}\right).$$

There is a similar result for the Liouville function, but it has a sharper error term.

Theorem 1.2. *Let $x > 1$ be a large number. If $a \neq 0$ is a small fixed integer, then*

$$\sum_{p \leq x} \lambda(p + a) = O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 0$ is an absolute constant.

Furthermore, a conditional result for the Mobius autocorrelation function over the shifted primes achieves the followings asymptotic formula.

Theorem 1.3. *Assume Hypothesis 2.1. Let $x > 1$ be a large number, and let $a, b \in \mathbb{Z}$ be small fixed integers such that $a \neq b$. Then,*

$$\sum_{p \leq x} \mu(p + a)\mu(p + b) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 1$ is an arbitrary constant.

Unexpectedly, the matching result for the Liouville autocorrelation function is unconditional and has a sharper error term.

Theorem 1.4. *Let $x > 1$ be a large number, and let $a, b \in \mathbb{Z}$ be small fixed integers such that $a \neq b$. Then,*

$$\sum_{p \leq x} \lambda(p + a)\lambda(p + b) = O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 0$ is an absolute constant.

The essential foundational materials are covered in Section 2.1 to Section 2.6. The proof of Theorem 1.1 for the summatory Mobius function over the shifted primes is assembled in Section 2.7, and the proof of Theorem 1.2 for the summatory Liouville function over the shifted primes is assembled in Section 4.4. The verification of Theorem 1.3 for the Mobius autocorrelation function over the shifted primes is assembled in Section 2.10. The proof of Theorem 1.4 for the Liouville autocorrelation function over the shifted primes appears in Section 4.7

For completeness some results on the Mobius autocorrelation function and the Liouville autocorrelation function over the integers, based on the sign patterns technique, are included in Section 7 and Section 8 respectively.

2 Results for the Mobius Function over the Shifted Primes

2.1 Average Orders of Mobius Functions

Theorem 2.1. *If $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ is the Mobius function, then, for any large number $x > 1$, the following statements are true.*

- (i) $\sum_{n \leq x} \mu(n) = O\left(xe^{-c\sqrt{\log x}}\right)$, where $c > 0$ is an absolute constant, unconditionally,
- (ii) $\sum_{n \leq x} \mu(n) = O\left(x^{1/2+\varepsilon}\right)$, where $\varepsilon > 0$ is an arbitrarily small number, conditional on the RH.

Proof. See [3, p. 6], [24, p. 182], [10, p. 347], et alii. ■

There are many sharp bounds of the summatory function of the Mobius function, say, $O(xe^{-c(\log x)^\delta})$, and the conditional estimate $O(x^{1/2+\varepsilon})$ presupposes that the nontrivial zeros of the zeta function $\zeta(\rho) = 0$ in the critical strip $\{0 < \Re(s) < 1\}$ are of the form $\rho = 1/2 + it, t \in \mathbb{R}$. However, the simpler notation will be used whenever it is convenient.

2.2 Mean Values and Hypotheses over the Shifted Primes

There are several mean values and equidistribution results for arithmetic functions over arithmetic progressions of level of distribution $\theta < 1/2$. The best known case is the Bombieri-Vinogradov theorem, see [3, Theorem 15.4], the case for the Mobius function is proved in [31, Theorem 1] and [28] states the following.

Corollary 2.1. ([28, Corollary 1]) *Let $a \geq 1$ be a fixed parameter, and let $x \geq 1$ be a large number. If $C > 0$ is a constant, then*

$$\sum_{q \leq x^{1/2}/\log^B x} \max_{a \bmod q} \max_{z \leq x} \left| \sum_{\substack{n \leq z \\ n \equiv a \pmod q}} \mu(n) \right| \ll \frac{x}{(\log x)^C}, \quad (10)$$

where the constant $B > 0$ depends on C .

However, there is no literature on the mean values and equidistribution for arithmetic functions over arithmetic progressions of shifted prime. A comparable result is expected to hold.

Hypothesis 2.1. *Let $a \geq 1$ be a fixed parameter, and let $x \geq 1$ be a large number. If $C > 0$ is a constant, then*

$$\sum_{q \leq x^{1/2}/\log^B x} \max_{a \bmod q} \max_{z \leq x} \left| \sum_{\substack{p \leq z \\ p \equiv d \pmod q}} \mu(p+a) \right| \ll \frac{x}{(\log x)^C}, \quad (11)$$

where the constant $B > 0$ depends on C .

A similar mean value for the autocorrelation over arithmetic progression is of interest in the theory of arithmetic correlation functions.

Hypothesis 2.2. *Let $a, b \in \mathbb{Z}$ be a fixed pair of small integers such that $a \neq b$, and let $x \geq 1$ be a large number. If $C > 0$ is a constant, then*

$$\sum_{q \leq x^{1/2}/\log^B x} \max_{d \bmod q} \max_{z \leq x} \left| \sum_{\substack{p \leq z \\ p \equiv d \bmod q}} \mu(p+a)\mu(p+b) \right| \ll \frac{x}{(\log x)^C}, \quad (12)$$

where the constant $B > 0$ depends on C .

2.3 Squarefree Shifted Primes

Let $p \geq 2$ be prime, and let $a \neq 0$ be a fixed integer. The number of squarefree integers of the form $p+a$, or totient integers, have the following asymptotic formula expressed in terms of the logarithm integral $\text{li}(x) = \int_2^x (\log t)^{-1} dt$.

Theorem 2.2. *Let $x \geq 1$ be a large number, and let $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ be the Mobius function. If $a \neq 0$ is a fixed integer, then,*

$$\sum_{p \leq x} \mu(p+a)^2 = s_0 \text{li}(x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$ is a constant, and $c > 1$ is an arbitrary constant.

Proof. Substituting the identity $\mu(n)^2 = \sum_{d^2|n} \mu(d)$, and switching the order of summation yield

$$\begin{aligned} \sum_{p \leq x} \mu(p+a)^2 &= \sum_{p \leq x} \sum_{d^2|p+a} \mu(d) & (13) \\ &= \sum_{d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2|p+a}} 1 \\ &= \sum_{d^2 \leq x_0} \mu(d) \sum_{\substack{p \leq x \\ d^2|p+a}} 1 + \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2|p+a}} 1, \end{aligned}$$

where $x_0 = (\log x)^{2c}$, and $c > 1$ is a constant. Applying the Siegel-Walfisz theorem, see [6, p. 405], [3, Theorem 15.3], et cetera, to the first subsum in the partition yields

$$\begin{aligned} \sum_{d^2 \leq x_0} \mu(d) \sum_{\substack{p \leq x \\ d^2|p+a}} 1 &= \sum_{d^2 \leq x_0} \mu(d) \left(\frac{\text{li}(x)}{\varphi(d^2)} + O\left(xe^{-c_0\sqrt{\log x}}\right) \right) & (14) \\ &= \text{li}(x) \sum_{n \geq 1} \frac{\mu(n)}{\varphi(n^2)} + O\left(\frac{x}{(\log x)^c}\right), \end{aligned}$$

where $c_0 > 0$ is an absolute constant. An estimate of the second subsum in the partition yields

$$\begin{aligned}
 \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} 1 &\leq \sum_{x_0 < d^2 \leq x} \sum_{\substack{p \leq x \\ d^2 | p+a}} 1 & (15) \\
 &\ll x \sum_{x_0 < d^2 \leq x} \frac{1}{d^2} \\
 &\ll \frac{x}{(\log x)^c}.
 \end{aligned}$$

Summing (14) and (15) completes the verification. ■

The well known constant has the numerical approximation

$$s_0 = \sum_{n \geq 1} \frac{\mu(n)}{\varphi(n^2)} = \prod_{p \geq 2} \left(1 - \frac{1}{p(p-1)} \right) = 0.373955838964330040631201 \dots \quad (16)$$

2.4 Nonlinear Autocorrelation Functions Results

The asymptotic formula for squarefree autocorrelation function over the shifted primes is evaluated now.

Theorem 2.3. *Let $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ be the Mobius function, and let $a, b \neq 0$ be integers such that $a \neq b$. Then, for any sufficiently large number $x \geq 1$,*

$$\sum_{p \leq x} \mu^2(p+a) \mu^2(p+b) = s_0^2 \operatorname{li}(x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955 \dots$, and $c > 1$ is an arbitrary constant.

Proof. Substitute the identity $\mu(n)^2 = \sum_{d^2 | n} \mu(d)$, and switching the order of summation yield

$$\begin{aligned}
 \sum_{p \leq x} \mu^2(p+a) \mu^2(p+b) &= \sum_{p \leq x} \sum_{d^2 | p+a} \mu(d) \sum_{e^2 | p+b} \mu(e) & (17) \\
 &= \sum_{d^2 \leq x} \mu(d) \sum_{e^2 \leq x} \mu(e) \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1 \\
 &= \sum_{d^2 \leq x_0} \mu(d) \sum_{e^2 \leq x_0} \mu(e) \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1 \\
 &\quad + \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{x_0 < e^2 \leq x} \mu(e) \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1,
 \end{aligned}$$

where $x_0 = (\log x)^c$, and $c > 1$ is an arbitrary constant. Let $q = d^2 e^2$, $\gcd(d, e) = 1$, and $p \equiv f \pmod q$. Applying the Siegel-Walfisz theorem, see [6, p. 405], [3, Theorem

15.3], et cetera, to the first subsum in the partition yields

$$\begin{aligned}
 T(x) &= \sum_{d^2 \leq x_0} \mu(d) \sum_{e^2 \leq x_0} \mu(e) \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1 & (18) \\
 &= \sum_{d^2 \leq x_0} \mu(d) \sum_{e^2 \leq x_0} \mu(e) \left(\frac{\text{li}(x)}{\varphi(d^2 e^2)} + O\left(xe^{-c\sqrt{\log x}}\right) \right) \\
 &= \sum_{d^2 \leq x_0} \frac{\mu(d)}{\varphi(d^2)} \sum_{e^2 \leq x_0} \frac{\mu(e)}{\varphi(e^2)} \left(\text{li}(x) + O\left(xe^{-c_0\sqrt{\log x}}\right) \right) \\
 &= \text{li}(x) \left(\sum_{n \geq 1} \frac{\mu(n)}{\varphi(n^2)} \right)^2 + O\left(\frac{x}{(\log x)^c}\right),
 \end{aligned}$$

where the constant $s_0 = \sum_{n \geq 1} \mu(n)\varphi(n^2)^{-1}$ is computed in (16), and $c_0 > 0$ is an absolute constant. An estimate of the second subsum in the partition yields

$$\begin{aligned}
 \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{x_0 < e^2 \leq x} \mu(e) \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1 &\leq \sum_{x_0 < d^2 \leq x} \sum_{x_0 < e^2 \leq x} \sum_{\substack{p \leq x \\ d^2 | p+a, e^2 | p+b}} 1 \\
 &\ll x \sum_{x_0 < d^2 \leq x} \frac{1}{d^2} \sum_{x_0 < e^2 \leq x} \frac{1}{e^2} & (19) \\
 &\ll \frac{x}{(\log x)^c}.
 \end{aligned}$$

Summing (18) and (19) completes the verification. ■

Theorem 2.4. *Assume Hypothesis 2.1. Let $x \geq 1$ be a large number, and let $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ be the Mobius function. If $a, b \neq 0$ are small fixed integers such that $a \neq b$, then, for any sufficiently large number $x \geq 1$,*

$$\sum_{p \leq x} \mu(p+a)^2 \mu(p+b) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant.

Proof. Substitute the identity $\mu(n)^2 = \sum_{d^2 | n} \mu(d)$, and switching the order of summation yield

$$\begin{aligned}
 \sum_{p \leq x} \mu(p+a)^2 \mu(p+b) &= \sum_{p \leq x} \mu(p+b) \sum_{d^2 | p+a} \mu(d) & (20) \\
 &= \sum_{d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} \mu(p+b) \\
 &= \sum_{d^2 \leq x^{2\varepsilon}} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} \mu(p+b) + \sum_{x^{2\varepsilon} < d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} \mu(p+b),
 \end{aligned}$$

where $\varepsilon \in (0, 1/4)$. Applying Hypothesis 2.1 to the first subsum in the partition yields

$$\begin{aligned} \sum_{d^2 \leq x^{2\varepsilon}} \mu(d) \sum_{\substack{n \leq x \\ d^2 | n}} \mu(n+t) &\leq \sum_{q \leq x^\varepsilon} \left| \sum_{\substack{p \leq x \\ p \equiv -a \pmod{q}}} \mu(p+b) \right| \\ &= O\left(\frac{x}{(\log x)^c}\right), \end{aligned} \quad (21)$$

where $q = d^2$. An estimate of the second subsum in the partition yields

$$\begin{aligned} \sum_{x^{2\varepsilon} < d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} \mu(p+b) &\leq \sum_{x^{2\varepsilon} < d^2 \leq x} \sum_{\substack{p \leq x \\ d^2 | p+a}} 1 \\ &\ll x \sum_{x^{2\varepsilon} < d^2 \leq x} \frac{1}{d^2} \\ &\ll x^{1-\varepsilon}. \end{aligned} \quad (22)$$

Summing (21) and (22) completes the verification. \blacksquare

Theorem 2.5. *Assume Hypothesis 2.2. Let $x \geq 1$ be a large number, and let $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ be the Mobius function. If $a_0, a_1, a_2 \neq 0$ are small fixed integers such that $a_0 < a_1 < a_2$, then, for any sufficiently large number $x \geq 1$,*

$$\sum_{p \leq x} \mu(p+a_0)^2 \mu(p+a_1) \mu(p+a_2) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant.

Proof. Except for minor changes, the verification is similar to the previous proof, for example, repeat (20) to (22), but uses Hypothesis 2.2. \blacksquare

2.5 Single-Sign Pattern Mobius Characteristic Functions

The analysis of single-Sign pattern characteristic function

$$\mu^\pm(n) = \mu^2(n) \left(\frac{1 \pm \mu(n)}{2} \right) = \begin{cases} 1 & \text{if } \mu(n) = \pm 1, \\ 0 & \text{if } \mu(n) \neq \pm 1, \end{cases} \quad (23)$$

of the subset of integers

$$\mathcal{A}_0^\pm = \{n \geq 1 : \mu(n) = \pm 1\} \quad (24)$$

is well known. Here, the same idea is extended to the shifted primes.

Lemma 2.1. *Let $a \neq 0$ be an integer, and let $\mu(n) \in \{-1, 0, 1\}$ be the Mobius function. Then,*

$$\begin{aligned} \mu^\pm(a, p) &= \mu^2(p+a) \left(\frac{1 \pm \mu(p+a)}{2} \right) \\ &= \begin{cases} 1 & \text{if } \mu(p+a) = \pm 1, \\ 0 & \text{if } \mu(p+a) \neq \pm 1, \end{cases} \end{aligned} \quad (25)$$

are the characteristic functions of the subset of primes

$$\mathcal{A}^\pm(a) = \{p \geq 2 : \mu(p+a) = \pm 1\}. \quad (26)$$

2.6 Single-Sign Pattern Mobius Counting Functions

The single-sign patterns $\mu(n) = 1$ and $\mu(n) = -1$ counting functions over the integers have the forms

$$R^+(x) = \sum_{\substack{n \leq x \\ \mu(n)=1}} 1 = \sum_{n \leq x} \left(\frac{1 + \mu(n)}{2} \right) \mu^2(n) = \frac{1}{2} \frac{x}{\zeta(2)} + O\left(xe^{-c\sqrt{\log x}}\right), \quad (27)$$

and

$$R^-(x) = \sum_{\substack{n \leq x \\ \mu(n)=-1}} 1 = \sum_{n \leq x} \left(\frac{1 - \mu(n)}{2} \right) \mu^2(n) = \frac{1}{2} \frac{x}{\zeta(2)} + O\left(xe^{-c\sqrt{\log x}}\right), \quad (28)$$

respectively. In terms of these functions, the summatory Mobius function has the asymptotic formula

$$R(x) = \sum_{n \leq x} \mu(n) = R^+(x) - R^-(x) = O\left(xe^{-c\sqrt{\log x}}\right). \quad (29)$$

Basically, it is a different form of the Prime Number Theorem

$$\pi(x) = \text{li}(x) + O\left(xe^{-c\sqrt{\log x}}\right), \quad (30)$$

where $\text{li}(x) = \int_2^x (\log t)^{-1} dt$ is the logarithm integral, and $c > 0$ is an absolute constant, see [4, Eq. 27.12.5], [5, Theorem 3.10], et alii.

The same idea extends to the single-sign patterns $\mu(p+a) = 1$ and $\mu(p+a) = -1$ of the shifted primes. Let $R(a, x) = \sum_{p \leq x} \mu(p+a)$. The single-sign pattern counting functions are defined by

$$R^+(a, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=1}} 1 = \sum_{n \in \mathcal{A}^+(a)} 1, \quad (31)$$

$$R^-(a, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=-1}} 1 = \sum_{p \in \mathcal{A}^-(a)} 1. \quad (32)$$

The single-sign pattern counting functions (31) to (32) are precisely the cardinalities of the subsets of integers

1. $\mathcal{A}^+(a) \subset \mathbb{P}$,
2. $\mathcal{A}^-(a) \subset \mathbb{P}$,

defined in (26). The symbol $\mathbb{P} = \{2, 3, 5, \dots\}$ denotes the set of prime numbers.

The next result is required to complete the analysis of the asymptotic formula for $R(a, x) = \sum_{p \leq x} \mu(p+a)$, which is completed in the next section.

Lemma 2.2. *Let $x \geq 1$ be a large number, and let $a \neq 0$ be a fixed integer. Then,*

$$R^\pm(a, x) = \frac{1}{2}s_0 \operatorname{li}(x) + \frac{1}{2}R(a, x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$, and $c > 0$ are constants.

Proof. Consider the pattern $\mu(p+a) = +1$. Now, use Lemma 2.1 to express the single-sign pattern counting function as

$$\begin{aligned} 2R^+(a, x) &= \sum_{p \leq x} \mu^+(a, p) \\ &= \sum_{p \leq x} \mu^2(p+a) (1 + \mu(p+a)) \\ &= \sum_{p \leq x} \mu^2(p+a) + \sum_{p \leq x} \mu^3(p+a). \end{aligned} \tag{33}$$

The two finite sums have the following evaluations or estimates.

$$1. \sum_{p \leq x} \mu^2(p+a) = s_0 \operatorname{li}(x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$ is a constant, see Theorem 2.2.

$$2. \sum_{n \leq x} \mu^3(p+a) = \sum_{p \leq x} \mu(p+a),$$

since $\mu^{2k+1}(n) = \mu(n)$ for any integer k .

Summing these evaluations or estimates verifies the claim for $R^+(a, x) \geq 0$. The verification for the other single-sign pattern counting function $R^-(a, x) \geq 0$, is similar. ■

2.7 Result for the Summatory Mobius Function

The analysis of the average order over the shifted primes

$$R(a, x) = \sum_{p \leq x} \mu(p+a) \tag{34}$$

is currently viewed as an intractable problem, the restricted double average order

$$\sum_{a \leq z} \sum_{p \leq x} \mu(p+a) = o(z\pi(x)) \tag{35}$$

is the only result available in the literature, see [16, Theorem 1.1] for the exact details. However, the introduction of the partial summatory function $R^+(a, x)$ and $R^-(a, x)$ transforms the problem into an elementary problem.

Proof. (Theorem 1.1) Rewriting the the summatory function in terms of the single-sign counting functions, and applying Lemma 2.2 lead to the asymptotic formula

$$\begin{aligned}
 R(a, x) &= \sum_{p \leq x} \mu(p+a) & (36) \\
 &= R^+(a, x) - R^-(a, x) \\
 &= \frac{1}{2}s_0 \operatorname{li}(x) + \frac{1}{2}R(a, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &\quad - \frac{1}{2}s_0 \operatorname{li}(x) - \frac{1}{2}R(a, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &= O\left(\frac{x}{(\log x)^c}\right),
 \end{aligned}$$

where $s_0 > 0$ is a constant, and $c > 1$ is an arbitrary constant. ■

2.8 Double-Sign Patterns Mobius Characteristic Functions

The analysis of single-sign pattern characteristic functions is extended here to the double-sign patterns

$$\mu(p+a) = \pm 1 \quad \text{and} \quad \mu(p+b) = \pm 1, \quad (37)$$

where $a, b \neq 0$ such that $a \neq b$, and $p \geq 2$ is prime. The current research has emphasized the sign patterns over the integers, confer [11], [29, Corollary 1.7], [14], and there is some literature on the patterns over the shifted primes, [25], and similar literature, for details.

Lemma 2.3. *Let $a, b \neq 0$ such that $a \neq b$ be small fixed integers, and let $\mu(n) \in \{-1, 0, 1\}$ be the Mobius function. Then,*

$$\begin{aligned}
 \mu^{\pm\pm}(a, b, p) &= \mu^2(p+a)\mu^2(p+b) \left(\frac{1 \pm \mu(p+a)}{2}\right) \left(\frac{1 \pm \mu(p+b)}{2}\right) & (38) \\
 &= \begin{cases} 1 & \text{if } \mu(p+a) = \pm 1, \mu(p+b) = \pm 1, \\ 0 & \text{if } \mu(p+a) \neq \pm 1, \mu(p+b) \neq \pm 1, \end{cases}
 \end{aligned}$$

are the characteristic functions of the subset of integers

$$\mathcal{A}^{\pm\pm}(a, b) = \{p \geq 2 : \mu(p+a) = \pm 1, \mu(p+b) = \pm 1\}. \quad (39)$$

2.9 Double-Sign Patterns Mobius Counting Functions

The single-sign counting functions for the single patterns $\mu(p+a) = 1$ and $\mu(p+a) = -1$ are extended to the double-sign counting functions for the double signs patterns

$$(\mu(p+a), \mu(p+b)) = (\pm 1, \pm 1). \quad (40)$$

The double-sign counting functions are defined by

$$R^{++}(a, b, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=1, \mu(p+b)=1}} 1 = \sum_{p \in \mathcal{A}^{++}(a, b)} 1, \quad (41)$$

$$R^{+-}(a, b, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=1, \mu(p+b)=-1}} 1 = \sum_{p \in \mathcal{A}^{+-}(a, b)} 1, \quad (42)$$

$$R^{-+}(a, b, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=-1, \mu(p+b)=1}} 1 = \sum_{p \in \mathcal{A}^{-+}(a, b)} 1, \quad (43)$$

$$R^{--}(a, b, x) = \sum_{\substack{p \leq x \\ \mu(p+a)=-1, \mu(p+b)=-1}} 1 = \sum_{p \in \mathcal{A}^{--}(a, b)} 1. \quad (44)$$

The double-sign counting functions (41) to (44) are precisely the cardinalities of the subsets of integers

1. $\mathcal{A}^{++}(a, b) \subset \mathbb{P}$,
2. $\mathcal{A}^{+-}(a, b) \subset \mathbb{P}$,
3. $\mathcal{A}^{-+}(a, b) \subset \mathbb{P}$,
4. $\mathcal{A}^{--}(a, b) \subset \mathbb{P}$,

defined in (39). In terms of these functions, the double autocorrelation function has form

$$\begin{aligned} R(a, b, x) &= \sum_{p \leq x} \mu(p+a)\mu(p+b) \\ &= R^{++}(a, b, x) - R^{+-}(a, b, x) + R^{-+}(a, b, x) - R^{--}(a, b, x). \end{aligned} \quad (45)$$

The next result is required to complete the analysis of the asymptotic formula for $R(a, b, x)$, which is completed in the next section.

Lemma 2.4. *Assume Hypothesis 2.1. Let $x \geq 1$ be a large number, and let $a, b \in \mathbb{Z}$ such that $a \neq b$, be fixed integers. Then,*

$$R^{\pm\pm}(a, b, x) = \frac{1}{4}s_0^2 \operatorname{li}(x) + \frac{1}{4}R(a, b, x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$, and $c > 1$ is an arbitrary constants.

Proof. Without loss in generality, consider the pattern $(\mu(p+a), \mu(p+b)) = (+1, +1)$. Now, use Lemma 2.3 to express the double-sign pattern counting function

as

$$\begin{aligned}
 4R^{++}(a, b, x) &= \sum_{p \leq x} \mu^{++}(a, b, p) & (46) \\
 &= \sum_{p \leq x} \mu^2(p+a)\mu^2(p+b) (1 + \mu(p+a)) (1 + \mu(p+b)) \\
 &= \sum_{p \leq x} \mu^2(p+a)\mu^2(p+b) (1 + \mu(p+a) + \mu(p+b) + \mu(p+a)\mu(p+b)) \\
 &= \sum_{p \leq x} \mu^2(p+a)\mu^2(p+b) + \sum_{p \leq x} \mu^3(p+a)\mu^2(p+b) \\
 &\quad + \sum_{p \leq x} \mu(p+a)^2\mu^3(p+b) + \sum_{p \leq x} \mu^3(p+a)\mu^3(p+b) \\
 &\geq 0.
 \end{aligned}$$

The last four finite sums have the following evaluations or estimates.

$$1. \sum_{p \leq x} \mu^2(p+a)\mu^2(p+b) = s_0^2 \operatorname{li}(x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 > 0$ is a constant, see Theorem 2.3.

$$2. \sum_{p \leq x} \mu^3(p+a)\mu^2(p+b) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant, see Theorem 2.4.

$$3. \sum_{p \leq x} \mu^2(p+a)\mu^3(p+b) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant, see Theorem 2.4.

$$4. \sum_{p \leq x} \mu^3(p+a)\mu^3(p+b) = \sum_{p \leq x} \mu(p+a)\mu(p+b),$$

since $\mu^{2k+1}(n) = \mu(n)$ for any integer k .

Summing these evaluations or estimates verifies the claim for $R^{++}(a, b, x) \geq 0$. The verifications for the next three double-sign pattern counting functions $R^{+-}(a, b, x) \geq 0$, $R^{-+}(a, b, x) \geq 0$, and $R^{--}(a, b, x) \geq 0$ are similar. \blacksquare

2.10 Result for the Mobius Autocorrelation Function

Proof. (Theorem 1.3) Assume Hypothesis 2.1. Rewriting the the autocorrelation function in terms of the double-sign counting functions, and applying Lemma 2.4

lead to the asymptotic formula

$$\begin{aligned}
 R(a, b, x) &= \sum_{p \leq x} \mu(p+a)\mu(p+b) & (47) \\
 &= R^{++}(a, b, x) - R^{+-}(a, b, x) + R^{--}(a, b, x) - R^{-+}(a, b, x) \\
 &= \frac{1}{4}s_0^2 \operatorname{li}(x) + \frac{1}{4}R(a, b, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &\quad - \frac{1}{4}s_0^2 \operatorname{li}(x) - \frac{1}{4}R(a, b, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &\quad + \frac{1}{4}s_0^2 \operatorname{li}(x) + \frac{1}{4}R(a, b, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &\quad - \frac{1}{4}s_0^2 \operatorname{li}(x) - \frac{1}{4}R(a, b, x) + O\left(\frac{x}{(\log x)^c}\right) \\
 &= O\left(\frac{x}{(\log x)^c}\right),
 \end{aligned}$$

where $s_0 = 0.373955\dots$, and $c > 1$ is an arbitrary constant. ■

3 Higher Order Mobius Autocorrelation Functions

Let $a_0 < a_1 < \dots < a_{k-1}$ be a subset of small integers. The higher degree Mobius autocorrelation functions of order $k \geq 3$ over the shifted primes, is defined by

$$R(a_0, \dots, a_{k-1}, x) = \sum_{p \leq x} \mu(p+a_0)\mu(p+a_1)\cdots\mu(p+a_{k-1}). \quad (48)$$

The Mobius autocorrelation functions can be computed or estimated, but conditional on some hypotheses on the distribution of primes in arithmetic progressions, see Hypothesis 2.1, and Hypothesis 2.2. Moreover, the maximal degree $k \leq k_{max}$, required to have a meaningful definition in (48), seems to be unknown. The last three subsections are dedicated to demonstrate this technique for the smallest case of $k = 3$. The verification for any other degree $k \leq k_{max}$ is similar.

3.1 Triple-Sign Patterns Mobius Characteristic Functions

The results for single-sign patterns and double-sign patterns characteristic functions has a seamless extension to the triple-sign patterns characteristic functions.

$$\mu(p+a_0) = \pm 1 \quad \mu(p+a_1) = \pm 1, \quad \text{and} \quad \mu(p+a_2) = \pm 1, \quad (49)$$

where $a_0, a_1, a_2 \in \mathbb{Z}$ are small integers such that $a_0 < a_1 < a_2$, and $p \geq 2$ is prime. The current research has emphasized the sign patterns over the integers, confer [11], [29, Corollary 1.7], [14], and there is some literature on the patterns over the shifted primes, [25], and similar literature, for details.

Lemma 3.1. *Let $a_0, a_1, a_2 \in \mathbb{Z}$ be small integers such that $a_0 < a_1 < a_2$, and let $\mu(n) \in \{-1, 0, 1\}$ be the Mobius function. Then,*

$$\begin{aligned} \mu^{\pm\pm\pm}(a_0, a_1, a_2, p) &= \prod_{0 \leq i \leq 2} \mu^2(p + a_i) \times \prod_{0 \leq i \leq 2} \left(\frac{1 \pm \mu(p + a_i)}{2} \right) \\ &= \begin{cases} 1 & \text{if } \mu(p + a_0) = \pm 1, \mu(p + a_1) = \pm 1, \mu(p + a_2) = \pm 1, \\ 0 & \text{if } \mu(p + a_0) \neq \pm 1, \mu(p + a_1) \neq \pm 1, \mu(p + a_2) \neq \pm 1, \end{cases} \end{aligned} \quad (50)$$

are the characteristic functions of the subset of integers

$$\mathcal{A}^{\pm\pm\pm}(a_0, a_1, a_2) = \{p \geq 2 : \mu(p + a_0) = \pm 1, \mu(p + a_1) = \pm 1, \mu(p + a_2) = \pm 1\}. \quad (51)$$

3.2 Triple-Sign Patterns Mobius Counting Functions

The counting functions for the single-sign patterns, and double-sign patterns are extended to the counting functions for the triple-sign patterns

$$\mu(p + a_0) = \pm 1, \quad \mu(p + a_1) = \pm 1, \quad \text{and} \quad \mu(p + a_2) = \pm 1. \quad (52)$$

The pattern counting functions are defined by

$$R^{+++}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \mu(p+a_0)=1, \mu(p+a_1)=1, \mu(p+a_2)=1}} 1 = \sum_{p \in \mathcal{A}^{+++}(a_0, a_1, a_2)} 1, \quad (53)$$

$$R^{++-}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \mu(p+a_0)=1, \mu(p+a_1)=1, \mu(p+a_2)=-1}} 1 = \sum_{p \in \mathcal{A}^{++-}(a_0, a_1, a_2)} 1, \quad (54)$$

$$\vdots \qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots$$

$$R^{---}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \mu(p+a_0)=-1, \mu(p+a_1)=-1, \mu(p+a_2)=-1}} 1 = \sum_{p \in \mathcal{A}^{---}(a_0, a_1, a_2)} 1 \quad (55)$$

The triple-sign pattern counting functions (53) to (55) are precisely the cardinalities of the subsets of integers

1. $\mathcal{A}^{+++}(a_0, a_1, a_2) \subset \mathbb{P}$,
2. $\mathcal{A}^{++-}(a_0, a_1, a_2) \subset \mathbb{P}$,
3. $\vdots \quad \vdots \quad \vdots$
4. $\mathcal{A}^{---}(a_0, a_1, a_2) \subset \mathbb{P}$,

defined in (51).

Lemma 3.2. *Assume Hypothesis 2.1. Let $x \geq 1$ be a large number, and let $a_0, a_1, a_2 \in \mathbb{Z}$ be small integers such that $a_0 < a_1 < a_2$. Then,*

$$R^{\pm\pm\pm}(a_0, a_1, a_2, x) = \frac{1}{2^3} s_0^3 \text{li}(x) + \frac{1}{2^3} R(a_0, a_1, a_2, x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$, and $c > 1$ is an arbitrary constants.

Proof. Without loss in generality, consider the pattern

$$(\mu(p + a_0), \mu(p + a_1), \mu(p + a_2)) = (+1, +1, +1). \quad (56)$$

Now, use Lemma 3.1 to express the triple pattern counting function as

$$\begin{aligned} 2^3 R^{+++}(a_0, a_1, a_2, x) &= \sum_{p \leq x} \mu^{+++}(a_0, a_1, a_2, p) \\ &= \sum_{p \leq x} \prod_{0 \leq i \leq 2} \mu^2(p + a_i) \times \prod_{0 \leq i \leq 2} (1 + \mu(p + a_i)) \\ &= \sum_{p \leq x} \mu^2(p + a_0) \mu^2(p + a_1) \mu^2(p + a_2) \\ &\quad + \sum_{p \leq x} \prod_{\substack{0 \leq i, j \leq 2 \\ i \neq j}} \mu^2(p + a_i) \mu(p + a_j) \\ &\quad + \sum_{p \leq x} \mu^3(p + a_0) \mu^3(p + a_1) \mu^3(p + a_2) \\ &\geq 0. \end{aligned} \quad (57)$$

The last eight finite sums have the following evaluations or estimates.

$$1. \sum_{p \leq x} \mu^2(p + a_0) \mu^2(p + a_1) \mu^2(p + a_2) = c_0^3 \operatorname{li}(x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 > 0$ is a constant, see Theorem 2.3.

$$2. \sum_{p \leq x} \prod_{\substack{0 \leq i, j \leq 2 \\ i \neq j}} \mu^2(p + a_i) \mu(p + a_j) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant, see Theorem 2.4.

$$3. \sum_{p \leq x} \prod_{\substack{0 \leq i, j, k \leq 2 \\ i \neq j \neq k}} \mu^2(p + a_i) \mu(p + a_j) \mu(p + a_k) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant, see Theorem 2.5.

$$4. \sum_{p \leq x} \mu^3(p + a_0) \mu^3(p + a_1) \mu^3(p + a_2) = \sum_{p \leq x} \mu(p + a_0) \mu(p + a_1) \mu(p + a_2),$$

since $\mu^{2k+1}(n) = \mu(n)$ for any integer k .

Summing these evaluations or estimates verifies the claim for $R^{+++}(a_0, a_1, a_2, x) \geq 0$. The verifications for the remaining seven triple-sign pattern counting functions $R^{++-}(a_0, a_1, a_2, x) \geq 0$, $R^{+-+}(a_0, a_1, a_2, x) \geq 0$, \dots , $R^{---}(a_0, a_1, a_2, x) \geq 0$ are similar. \blacksquare

- (i) $\sum_{\substack{n \leq x \\ \text{ally}}} \lambda(n) = O\left(xe^{-c\sqrt{\log x}}\right)$, where $c > 0$ is an absolute constant, unconditionally,
- (ii) $\sum_{n \leq x} \lambda(n) = O\left(x^{1/2+\varepsilon}\right)$, where $\varepsilon > 0$ is an arbitrarily small number, conditional on the RH.

Proof. (i) Substitute the identity $\lambda(n) = \sum_{d^2|n} \mu(n/d^2)$ to convert the summatory function to

$$\begin{aligned} \sum_{n \leq x} \lambda(n) &= \sum_{n \leq x} \sum_{d^2|n} \mu(n/d^2) \\ &= \sum_{d^2 \leq x} M(x/d^2), \end{aligned} \quad (59)$$

where $M(t) = \sum_{n \leq t} \mu(n)$. Applying Theorem 2.1 yields

$$\begin{aligned} \sum_{d^2 \leq x} M(x/d^2) &= O\left(xe^{-c\sqrt{\log x}} \sum_{d \leq \sqrt{x}} \frac{1}{d^2}\right) \\ &= O\left(xe^{-c\sqrt{\log x}}\right), \end{aligned} \quad (60)$$

where $c > 0$ is an absolute constant. ■

4.2 Single-Sign Patterns Liouville Characteristic Functions

The analysis of single-sign pattern characteristic function

$$\lambda^\pm(n) = \left(\frac{1 \pm \lambda(n)}{2}\right) = \begin{cases} 1 & \text{if } \lambda(n) = \pm 1, \\ 0 & \text{if } \lambda(n) \neq \pm 1, \end{cases} \quad (61)$$

of the subset of integers

$$\mathcal{B}_0^\pm = \{n \geq 1 : \mu(n) = \pm 1\} \quad (62)$$

is well known. Here, the same idea is extended to the shifted primes.

Lemma 4.1. *Let $a \neq 0$ be an integer, and let $\lambda(n) \in \{-1, 1\}$ be the Liouville function. Then,*

$$\begin{aligned} \lambda^\pm(a, p) &= \left(\frac{1 \pm \lambda(p+a)}{2}\right) \\ &= \begin{cases} 1 & \text{if } \lambda(p+a) = \pm 1, \\ 0 & \text{if } \lambda(p+a) \neq \pm 1, \end{cases} \end{aligned} \quad (63)$$

are the characteristic functions of the subset of primes

$$\mathcal{B}^\pm(a) = \{p \geq 2 : \lambda(p+a) = \pm 1\}. \quad (64)$$

4.3 Single-Sign Pattern Liouville Counting Functions

The single-sign patterns $\lambda(n) = 1$ and $\lambda(n) = -1$ single-sign pattern counting functions over the integers have the forms

$$Q^+(x) = \sum_{\substack{n \leq x \\ \lambda(n)=1}} 1 = \sum_{n \leq x} \left(\frac{1 + \lambda(n)}{2} \right) = \frac{1}{2}[x] + O\left(xe^{-c\sqrt{\log x}}\right), \quad (65)$$

and

$$Q^-(x) = \sum_{\substack{n \leq x \\ \lambda(n)=-1}} 1 = \sum_{n \leq x} \left(\frac{1 - \lambda(n)}{2} \right) = \frac{1}{2}[x] + O\left(xe^{-c\sqrt{\log x}}\right), \quad (66)$$

where $[x]$ is the largest integer function, respectively. In terms of the single-sign pattern counting functions, the summatory function has the asymptotic formula

$$Q(x) = \sum_{n \leq x} \lambda(n) = Q^+(x) - Q^-(x) = O\left(xe^{-c\sqrt{\log x}}\right). \quad (67)$$

Basically, it is a form of the prime number theorem, stated in (30).

The same idea extends to the single-sign patterns $\lambda(p+a) = 1$ and $\lambda(p+a) = -1$ of the shifted primes. Let $Q(a, x) = \sum_{p \leq x} \lambda(p+a)$. The single-sign patterns counting functions are defined by

$$Q^+(a, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=1}} 1 = \sum_{\substack{p \leq x \\ n \in \mathcal{B}^+(a)}} 1, \quad (68)$$

$$Q^-(a, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=-1}} 1 = \sum_{\substack{p \leq x \\ p \in \mathcal{B}^-(a)}} 1. \quad (69)$$

The single-sign patterns counting functions (31) to (32) are precisely the cardinalities of the subsets of integers

1. $\mathcal{B}^+(a) \subset \mathbb{P}$,
2. $\mathcal{B}^-(a) \subset \mathbb{P}$,

defined in (26). The symbol $\mathbb{P} = \{2, 3, 5, \dots\}$ denotes the set of prime numbers.

The next result is required to complete the analysis of the asymptotic formula for $Q(a, x) = \sum_{p \leq x} \lambda(p+a)$, which is completed in the next section.

Lemma 4.2. *Let $x \geq 1$ be a large number, and let $a \neq 0$ be a fixed integer. Then,*

$$Q^\pm(a, x) = \frac{1}{2} \text{li}(x) + \frac{1}{2} Q(a, x) + O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 0$ is an absolute constant.

Proof. Consider the pattern $\lambda(p+a) = +1$. Now, use Lemma 4.1 to express the single-sign pattern counting function as

$$\begin{aligned} 2Q^+(a, x) &= \sum_{p \leq x} \lambda^+(a, p) \\ &= \sum_{p \leq x} (1 + \lambda(p+a)) \\ &= \sum_{p \leq x} 1 + \sum_{p \leq x} \lambda(p+a). \end{aligned} \tag{70}$$

The first finite sum is $\sum_{p \leq x} 1 = \pi(x)$, see (30) for more details, and the second sum is $Q(a, x)/2$. Summing these evaluations or estimates verifies the claim for $Q^+(a, x) \geq 0$. The verification for the next single-sign pattern counting function $Q^-(a, x) \geq 0$, is similar. ■

4.4 Result for Liouville Summatory Function

The analysis of the average order over the shifted primes

$$Q(a, x) = \sum_{p \leq x} \lambda(p+a) \tag{71}$$

is currently viewed as an intractable problem, the restricted double average order

$$\sum_{a \leq z} \sum_{p \leq x} \lambda(p+a) = o(z\pi(x)) \tag{72}$$

is the only result available in the literature, see [25], and [16, Theorem 1.1] for the exact details. However, the introduction of the single-sign counting function $Q^+(a, x)$ and $Q^-(a, x)$ transforms the problem into an elementary problem.

Proof. (Theorem 1.2) Rewriting the summatory Liouville function in terms of the single-sign pattern counting functions, and applying Lemma 4.2 lead to the asymptotic formula

$$\begin{aligned} Q(a, x) &= \sum_{p \leq x} \lambda(p+a) \\ &= Q^+(a, x) - Q^-(a, x) \\ &= \frac{1}{2} \text{li}(x) + \frac{1}{2} Q(a, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad - \frac{1}{2} \text{li}(x) - \frac{1}{2} Q(a, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &= O\left(xe^{-c\sqrt{\log x}}\right), \end{aligned} \tag{73}$$

where $c > 0$ is an absolute constant. ■

4.5 Double-Sign Patterns Liouville Characteristic Functions

The analysis of single-sign pattern characteristic functions is extended here to the double-sign patterns

$$\lambda(p+a) = \pm 1 \quad \text{and} \quad \lambda(p+b) = \pm 1, \quad (74)$$

where $a, b \neq 0$ such that $a \neq b$, and $p \geq 2$ is prime. The current research has emphasized the sign patterns over the integers, confer [11], [29, Corollary 1.7], [14], [27], and there is some literature on the sign patterns over the shifted primes, [25], and similar literature, for details.

Lemma 4.3. *Let $a, b \in \mathbb{Z}$ such that $a \neq b$ be small fixed integers, and let $\lambda(n) \in \{-1, 1\}$ be the Liouville function. Then,*

$$\begin{aligned} \lambda^{\pm\pm}(a, b, p) &= \left(\frac{1 \pm \lambda(p+a)}{2} \right) \left(\frac{1 \pm \lambda(p+b)}{2} \right) \\ &= \begin{cases} 1 & \text{if } \lambda(p+a) = \pm 1, \mu(p+b) = \pm 1, \\ 0 & \text{if } \lambda(p+a) \neq \pm 1, \mu(p+b) \neq \pm 1, \end{cases} \end{aligned} \quad (75)$$

are the characteristic functions of the subset of integers

$$\mathcal{B}^{\pm\pm}(a, b) = \{p \geq 2 : \lambda(p+a) = \pm 1, \lambda(p+b) = \pm 1\}. \quad (76)$$

4.6 Double-Sign Patterns in Liouville Counting Functions

The counting functions for the single-sign patterns $\lambda(p+a) = 1$ and $\lambda(p+a) = -1$ are extended to the counting functions for the double-sign patterns

$$(\lambda(p+a), \lambda(p+b)) = (\pm 1, \pm 1). \quad (77)$$

The double-sign patterns counting functions are defined by

$$Q^{++}(a, b, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=1, \lambda(p+b)=1}} 1 = \sum_{p \in \mathcal{B}^{++}(a, b)} 1, \quad (78)$$

$$Q^{+-}(a, b, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=1, \lambda(p+b)=-1}} 1 = \sum_{p \in \mathcal{B}^{+-}(a, b)} 1, \quad (79)$$

$$Q^{-+}(a, b, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=-1, \lambda(p+b)=1}} 1 = \sum_{p \in \mathcal{B}^{-+}(a, b)} 1, \quad (80)$$

$$Q^{--}(a, b, x) = \sum_{\substack{p \leq x \\ \lambda(p+a)=-1, \lambda(p+b)=-1}} 1 = \sum_{p \in \mathcal{B}^{--}(a, b)} 1. \quad (81)$$

The double-sign patterns counting functions (41) to (44) are precisely the cardinalities of the subsets of integers

1. $\mathcal{B}^{++}(a, b) \subset \mathbb{P}$,
2. $\mathcal{B}^{+-}(a, b) \subset \mathbb{P}$,
3. $\mathcal{B}^{-+}(a, b) \subset \mathbb{P}$,
4. $\mathcal{B}^{--}(a, b) \subset \mathbb{P}$,

defined in (76). The symbol $\mathbb{P} = \{2, 3, 5, \dots\}$ denotes the set of prime numbers. In terms of the double-sign patterns counting functions, the autocorrelation function has form

$$\begin{aligned} Q(a, b, x) &= \sum_{p \leq x} \lambda(p+a)\lambda(p+b) \\ &= Q^{++}(a, b, x) - Q^{+-}(a, b, x) + Q^{-+}(a, b, x) - Q^{--}(a, b, x). \end{aligned} \quad (82)$$

The next result is required to complete the analysis of the asymptotic formula for $Q(a, b, x)$, which is completed in the next section.

Lemma 4.4. *Let $x \geq 1$ be a large number, and let $a, b \in \mathbb{Z}$ such that $a \neq b$, be fixed integers. If $\lambda : \mathbb{Z} \rightarrow \{-1, 1\}$ is the Liouville function, then,*

$$Q^{\pm\pm}(a, b, x) = \frac{1}{4} \text{li}(x) + \frac{1}{4} Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 0$ is an absolute constant.

Proof. Without loss in generality, consider the pattern $(\lambda(p+a), \lambda(p+b)) = (+1, +1)$. Now, use Lemma 4.3 to express the double-sign pattern counting function as

$$\begin{aligned} 4Q^{++}(a, b, x) &= \sum_{p \leq x} \lambda^{++}(a, b, p) \\ &= \sum_{p \leq x} (1 + \lambda(p+a))(1 + \lambda(p+b)) \\ &= \sum_{p \leq x} (1 + \lambda(p+a) + \lambda(p+b) + \lambda(p+a)\lambda(p+b)) \\ &= \sum_{p \leq x} 1 + \sum_{p \leq x} \lambda(p+a) + \sum_{p \leq x} \lambda(p+b) + \sum_{p \leq x} \lambda(p+a)\lambda(p+b) \\ &\geq 0. \end{aligned} \quad (83)$$

The first three finite sums on the last line have the following evaluations or estimates.

1. $\sum_{p \leq x} 1 = \text{li}(x) + O\left(xe^{-c\sqrt{\log x}}\right)$, see (30).
2. $\sum_{p \leq x} \lambda(p+a) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 4.1.
3. $\sum_{p \leq x} \lambda(p+b) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 4.1.

In all cases $c > 0$ is an absolute constant, unconditionally. Summing these evaluations or estimates verifies the claim for $Q^{++}(a, b, x) \geq 0$. The verifications for the next three double-sign pattern counting functions $Q^{+-}(a, b, x) \geq 0$, $Q^{-+}(a, b, x) \geq 0$, and $Q^{--}(a, b, x) \geq 0$ are similar. ■

4.7 Result for the Liouville Autocorrelation Function

Unlike the conditional result for the Mobius autocorrelation function in Theorem 1.3, the matching result for the Liouville autocorrelation function is unconditional, and has a sharper error term.

Proof. (Theorem 1.4) Rewriting the autocorrelation function in terms of the double-sign patterns counting functions, and applying Lemma 4.4 lead to the asymptotic formula

$$\begin{aligned}
 Q(a, b, x) &= \sum_{p \leq x} \lambda(p+a)\lambda(p+b) & (84) \\
 &= Q^{++}(a, b, x) - Q^{+-}(a, b, x) + Q^{--}(a, b, x) - Q^{-+}(a, b, x) \\
 &= \frac{1}{4} \operatorname{li}(x) + \frac{1}{4} Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\
 &\quad - \frac{1}{4} \operatorname{li}(x) - \frac{1}{4} Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\
 &\quad + \frac{1}{4} \operatorname{li}(x) + \frac{1}{4} Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\
 &\quad - \frac{1}{4} \operatorname{li}(x) - \frac{1}{4} Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\
 &= O\left(xe^{-c\sqrt{\log x}}\right),
 \end{aligned}$$

where $a < b$ are small integers, and $c > 0$ is an absolute constant. ■

5 Higher Order Liouville Autocorrelation Functions

Let $a_0 < a_1 < \dots < a_{k-1}$ be a subset of small integers. The higher degree Liouville autocorrelation functions of order $k \geq 3$, is defined by

$$Q(a_0, \dots, a_{k-1}, x) = \sum_{p \leq x} \lambda(p+a_0)\lambda(p+a_1) \cdots \lambda(p+a_{k-1}). \quad (85)$$

The higher degree Liouville autocorrelation functions can be computed recursively and unconditionally. However, the maximal degree $k \leq k_{max}$, for a meaningful definition in (85), seems to be unknown. The last three subsections are dedicated to demonstrate this technique for the smallest case of $k = 3$. The verification for any other degree $k \leq k_{max}$ is similar.

5.1 Triple Patterns Liouville Characteristic Functions

The analysis of single-sign patterns and double-sign patterns characteristic functions are extended here to the triple-sign patterns

$$\lambda(p+a_0) = \pm 1 \quad \lambda(p+a_1) = \pm 1, \quad \text{and} \quad \lambda(p+a_2) = \pm 1, \quad (86)$$

where $a_0, a_1, a_2 \in \mathbb{Z}$ are small integers such that $a_0 < a_1 < a_2$, and $p \geq 2$ is prime. The current research has emphasized the sign patterns over the integers, confer [11], [29, Corollary 1.7], [14], and there is some literature on the sign patterns over the shifted primes, [25], and similar literature, for details.

Lemma 5.1. *Let $a_0, a_1, a_2 \in \mathbb{Z}$ be small integers such that $a_0 < a_1 < a_2$, and let $\lambda(n) \in \{-1, 1\}$ be the Liouville function. Then,*

$$\begin{aligned} \lambda^{\pm\pm\pm}(a_0, a_1, a_2, p) &= \prod_{0 \leq i \leq 2} \left(\frac{1 \pm \lambda(p + a_i)}{2} \right) & (87) \\ &= \begin{cases} 1 & \text{if } \lambda(p + a_0) = \pm 1, \lambda(p + a_1) = \pm 1, \lambda(p + a_2) = \pm 1, \\ 0 & \text{if } \lambda(p + a_0) \neq \pm 1, \lambda(p + a_1) \neq \pm 1, \lambda(p + a_2) \neq \pm 1, \end{cases} \end{aligned}$$

are the characteristic functions of the subset of integers

$$\mathcal{B}^{\pm\pm\pm}(a_0, a_1, a_2) = \{p \geq 2 : \lambda(p + a_0) = \pm 1, \lambda(p + a_1) = \pm 1, \lambda(p + a_2) = \pm 1\}. \quad (88)$$

5.2 Triple-Sign Patterns Liouville Counting Functions

The counting functions for the single-sign patterns, and double-sign patterns are extended to the counting functions for the triple-sign patterns

$$\lambda(p + a_0) = \pm 1, \quad \lambda(p + a_1) = \pm 1, \quad \text{and} \quad \lambda(p + a_2) = \pm 1. \quad (89)$$

The sign pattern counting functions are defined by

$$Q^{+++}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \lambda(p+a_0)=1, \lambda(p+a_1)=1, \lambda(p+a_2)=1}} 1 = \sum_{p \in \mathcal{B}^{+++}(a_0, a_1, a_2)} 1, \quad (90)$$

$$Q^{++-}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \lambda(p+a_0)=1, \lambda(p+a_1)=1, \lambda(p+a_2)=-1}} 1 = \sum_{p \in \mathcal{B}^{++-}(a_0, a_1, a_2)} 1, \quad (91)$$

\vdots \vdots \vdots

$$Q^{---}(a_0, a_1, a_2, x) = \sum_{\substack{p \leq x \\ \lambda(p+a_0)=-1, \lambda(p+a_1)=-1, \lambda(p+a_2)=-1}} 1 = \sum_{p \in \mathcal{B}^{---}(a_0, a_1, a_2)} 1 \quad (92)$$

The triple-sign pattern counting functions (53) to (92) are precisely the cardinalities of the subsets of integers

1. $\mathcal{B}^{+++}(a_0, a_1, a_2) \subset \mathbb{P}$, 3. \vdots \vdots \vdots
2. $\mathcal{B}^{++-}(a_0, a_1, a_2) \subset \mathbb{P}$, 4. $\mathcal{B}^{---}(a_0, a_1, a_2) \subset \mathbb{P}$,

defined in (88).

Lemma 5.2. *Let $x \geq 1$ be a large number, and let $a_0, a_1, a_2 \in \mathbb{Z}$ be small integers such that $a_0 < a_1 < a_2$. Then,*

$$Q^{\pm\pm\pm}(a_0, a_1, a_2, x) = \frac{1}{2^3} s_0^3 \operatorname{li}(x) + \frac{1}{2^3} R(a_0, a_1, a_2, x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$, and $c > 1$ is an arbitrary constants.

Proof. Without loss in generality, consider the sign pattern

$$(\lambda(p + a_0), \lambda(p + a_1), \lambda(p + a_2)) = (+1, +1, +1). \quad (93)$$

Now, use Lemma 5.1 to express the triple pattern counting function as

$$\begin{aligned} 2^3 Q^{+++}(a_0, a_1, a_2, x) &= \sum_{p \leq x} \lambda^{+++}(a_0, a_1, a_2, p) \\ &= \sum_{p \leq x} \prod_{0 \leq i \leq 2} (1 + \lambda(p + a_i)) \\ &= \sum_{p \leq x} 1 + \sum_{0 \leq i \leq 2, p \leq x} \lambda(p + a_i) \\ &\quad + \sum_{\substack{0 \leq i, j \leq 2, p \leq x \\ i \neq j}} \lambda(p + a_i) \lambda(p + a_j) \\ &\quad + \sum_{p \leq x} \lambda^3(p + a_0) \lambda^3(p + a_1) \lambda^3(p + a_2) \\ &\geq 0. \end{aligned} \quad (94)$$

The last eight finite sums have the following evaluations or estimates.

1. $\sum_{p \leq x} 1 = \operatorname{li}(x) + O\left(xe^{-c\sqrt{\log x}}\right)$, see (30).
2. $\sum_{0 \leq i \leq 2, p \leq x} \lambda(p + a_i) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 4.1.
3. $\sum_{\substack{0 \leq i, j \leq 2, p \leq x \\ i \neq j}} \lambda(p + a_i) \lambda(p + a_j) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 1.4, this is the recursive step: a lower degree autocorrelation is used to compute a higher degree autocorrelation.
4. $\sum_{p \leq x} \lambda^3(p + a_0) \lambda^3(p + a_1) \lambda^3(p + a_2) = \sum_{p \leq x} \lambda(p + a_0) \lambda(p + a_1) \lambda(p + a_2)$, since $\mu^{2k+1}(n) = \mu(n)$ for any integer k .

In all cases $c > 0$ is an absolute constant.

Summing these evaluations or estimates verifies the claim for $Q^{+++}(a_0, a_1, a_2, x) \geq 0$. The verifications for the remaining seven triple-sign pattern counting functions $Q^{++-}(a_0, a_1, a_2, x) \geq 0$, $Q^{+-+}(a_0, a_1, a_2, x) \geq 0$, \dots , $Q^{---}(a_0, a_1, a_2, x) \geq 0$ are similar. ■

5.3 Result for the Liouville Autocorrelation Function

Theorem 5.1. *Let $x > 1$ be a large number, and let $a_0, a_1, a_2 \in \mathbb{Z}$ be small fixed integers such that $a_0 < a_1 < a_2$. Then,*

$$\sum_{p \leq x} \lambda(p + a_0)\lambda(p + a_1)\lambda(p + a_2) = O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 1$ is an absolute constant.

Proof. Rewriting the the autocorrelation function in terms of the triple-sign pattern counting functions, and applying Lemma 5.2 lead to the asymptotic formula

$$\begin{aligned} Q(a_0, a_1, a_2, x) &= \sum_{p \leq x} \lambda(p + a_0)\lambda(p + a_1)\lambda(p + a_2) & (95) \\ &= Q^{+++}(a_0, a_1, a_2, x) - Q^{++-}(a_0, a_1, a_2, x) \\ &\quad + Q^{+--}(a_0, a_1, a_2, x) - Q^{+-+}(a_0, a_1, a_2, x) \\ &\quad + \cdots + Q^{-++}(a_0, a_1, a_2, x) - Q^{-+-}(a_0, a_1, a_2, x) \\ &= \frac{1}{8} \operatorname{li}(x) + \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad - \frac{1}{8} \operatorname{li}(x) - \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad + \frac{1}{8} \operatorname{li}(x) + \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(\frac{x}{(\log x)^c}\right) \\ &\quad - \frac{1}{8} \operatorname{li}(x) - \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \\ &\quad + \frac{1}{8} \operatorname{li}(x) + \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad - \frac{1}{8} \operatorname{li}(x) - \frac{1}{8} Q(a_0, a_1, a_2, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &= O\left(xe^{-c\sqrt{\log x}}\right), \end{aligned}$$

where $c > 0$ is an absolute constant. ■

6 Hardy-Littlewood-Chowla Conjecture

The Hardy-Littlewood-Chowla Conjecture is a hybrid of the Hardy-Littlewood conjecture, see [2], [12, Conjecture B], and the Chowla conjecture, see [1], [26]. A version of the merged conjectures has the structure described below.

Conjecture 6.1. *Let $x \geq 1$ be a large number, let a_0, a_1, \dots, a_{k-1} be a fixed admissible k -tuple, and let $b_0 < b_1 < \cdots < b_{l-1}$ be a fixed subset of small integers. Then,*

$$\sum_{n \leq x} \Lambda(n + a_0)\Lambda(n + a_1) \cdots \Lambda(n + a_{k-1})\mu(n + b_0)\mu(n + b_1) \cdots \mu(n + b_{l-1}) = o(x).$$

The special case $k = 0$ reduces to the Chowla conjecture, and the special case $l = 0$ reduces to the Hardy-Littlewood conjecture. The other cases for $(k, l) \neq (0, 0)$ has been proved on average for almost all combinations of the admissible k -tuple a_0, a_1, \dots, a_{k-1} , and the small subset of integers $b_0 < b_1 < \dots < b_{l-1}$, the details of the proof are provided in [17].

The first proof for the case $k = 1$ and $l = 1$ for any fixed pair of small integers $a_0 = 0$ and $b_0 \neq 0$ is proved in Theorem 6.1, unconditionally.

6.1 Twisted vonMangoldt Function over the Squarefree Shifted Primes

Lemma 6.1. *Let $x \geq 1$ be a large number, and let $a \neq 0$ be a fixed integer. Then,*

$$\sum_{n \leq x} \Lambda(n) \mu^2(n+a) = s_0 x + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 = 0.373955\dots$, and $c > 1$ is a constant.

Proof. Substituting the identity $\mu(n)^2 = \sum_{d^2|n} \mu(d)$, and switching the order of summation yield

$$\begin{aligned} \sum_{n \leq x} \Lambda(n) \mu^2(n+a) &= \sum_{n \leq x} \Lambda(n) \sum_{d^2|n+a} \mu(d) & (96) \\ &= \sum_{d^2 \leq x} \mu(d) \sum_{\substack{n \leq x \\ d^2|n+a}} \Lambda(n) \\ &= \sum_{d^2 \leq x_0} \mu(d) \sum_{\substack{n \leq x \\ d^2|n+a}} \Lambda(n) + \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{\substack{n \leq x \\ d^2|n+a}} \Lambda(n), \end{aligned}$$

where $x_0 = (\log x)^{2c_1}$, with $c_1 > 1$. Applying the Siegel-Walfisz theorem, see [6, p. 405], [3, Theorem 15.3], et cetera, to the first subsum in the above partition yields

$$\begin{aligned} \sum_{d^2 \leq x_0} \mu(d) \sum_{\substack{n \leq x \\ d^2|n+a}} \Lambda(n) &= \sum_{d^2 \leq x_0} \mu(d) \left(\frac{x}{\varphi(d^2)} + O\left(xe^{-c_0\sqrt{\log x}}\right) \right) & (97) \\ &= s_0 x + O\left(xe^{-c_1\sqrt{\log x}}\right), \end{aligned}$$

where $c_0, c_2 > 0$ are absolute constants, and

$$s_0 = \sum_{n \geq 1} \frac{\mu(n)}{\varphi(n^2)} = \prod_{p \geq 2} \left(1 - \frac{1}{p(p-1)}\right), \quad (98)$$

see (16). An estimate of the second subsum in the partition yields

$$\begin{aligned} \sum_{x_0 < d^2 \leq x} \mu(d) \sum_{\substack{p \leq x \\ d^2 | p+a}} \Lambda(n) &\leq (\log x) \sum_{x_0 < d^2 \leq x} \sum_{\substack{p \leq x \\ d^2 | p+a}} 1 \\ &\ll x(\log x) \sum_{x_0 < d^2 \leq x} \frac{1}{d^2} \\ &\ll \frac{x}{(\log x)^{c_1-1}}. \end{aligned} \quad (99)$$

Summing (96) and (97), and setting $c = c_1 - 1 > 0$ complete the verification. \blacksquare

6.2 Signed vonMangoldt Function over the Shifted Primes

Define the pattern indicator function

$$\mu^+(n) = \mu^2(n) \left(\frac{1 + \mu(n)}{2} \right) = \begin{cases} 1 & \text{if } \mu(n) = 1, \\ 0 & \text{if } \mu(n) \neq 1, \end{cases} \quad (100)$$

and

$$\mu^-(n) = \mu^2(n) \left(\frac{1 - \mu(n)}{2} \right) = \begin{cases} 1 & \text{if } \mu(n) = -1, \\ 0 & \text{if } \mu(n) \neq -1. \end{cases} \quad (101)$$

These are the same as the characteristic functions stated in (23).

Lemma 6.2. *Let $x \geq 1$ be a large number, and let $a \neq 0$ be a fixed integer. Then,*

$$\sum_{n \leq x} \Lambda(n) \mu^\pm(n+a) = \frac{1}{2} s_0 x + \frac{1}{2} \sum_{n \leq x} \Lambda(n) \mu(n+a) + O\left(\frac{x}{(\log x)^c}\right),$$

where $s_0 > 0$ is a constant, and $c > 1$ is an absolute constant.

Proof. Without loss in generality, consider the pattern $\mu^+(n+a) = +1$. Now, use Lemma 2.3 to express the double pattern counting function as

$$\begin{aligned} \sum_{n \leq x} \Lambda(n) \mu^+(n+a) &= \sum_{n \leq x} \Lambda(n) \mu^2(n+a) (1 + \mu(n+a)) \\ &= \sum_{n \leq x} \Lambda(n) (\mu^2(n+a) + \mu^3(n+a)) \\ &= \sum_{n \leq x} \Lambda(n) \mu^2(n+a) + \sum_{n \leq x} \Lambda(n) \mu(n+a) \\ &\geq 0. \end{aligned} \quad (102)$$

By Lemma 6.1, the first finite sums has asymptotic evaluation

$$\sum_{n \leq x} \Lambda(n) \mu^2(n+a) = s_0 x + O\left(x e^{-c_0 \sqrt{\log x}}\right) \quad (103)$$

where $s_0 > 0$ is a constant, and $c_0 > 0$ is an absolute constant. Summing these evaluations or estimates verifies the claim for $\sum_{n \leq x} \Lambda(n) \mu^+(n+a) \geq 0$. The verifications for the other single-sign pattern counting function $\sum_{n \leq x} \Lambda(n) \mu^-(n+a) \geq 0$ is similar. \blacksquare

6.3 Twisted vonMangoldt Function over the Shifted Primes

The first case of the hybrid finite, confer Conjecture (6.1), involving the vonMangoldt function and the Mobius function is verified here.

Theorem 6.1. *Let $x \geq 1$ be a large number, and let $a \neq 0$ be a fixed integer. Then,*

$$\sum_{n \leq x} \Lambda(n) \mu(n+a) = O\left(\frac{x}{(\log x)^c}\right), \quad (104)$$

where $c > 1$ is an arbitrary constant.

Proof. Rewrite it in terms of the single-sign pattern counting function computed in Lemma 6.2 as

$$\begin{aligned} \sum_{n \leq x} \Lambda(n) \mu(n+a) &= \sum_{n \leq x} \Lambda(n) \mu^+(n+a) - \sum_{n \leq x} \Lambda(n) \mu^-(n+a) \quad (105) \\ &= \frac{1}{2} s_0 x + \frac{1}{2} \sum_{n \leq x} \Lambda(n) \mu(n+a) + O\left(\frac{x}{(\log x)^c}\right) \\ &\quad - \frac{1}{2} s_0 x - \frac{1}{2} \sum_{n \leq x} \Lambda(n) \mu(n+a) + O\left(\frac{x}{(\log x)^c}\right) \\ &= O\left(\frac{x}{(\log x)^c}\right), \end{aligned}$$

where $c > 1$ is an arbitrary constant. ■

For an interesting application of Theorem 6.1, set $a = 2k$. Then, a result in [23, Theorem 1.1] proves that (104) and

$$\sum_{n \leq x} \Lambda(n) \Lambda(n+2k) = \mathfrak{G}(2k)x + o(x) \quad (106)$$

are equivalent.

7 Results for the Mobius Autocorrelation Function over the Integers

The current estimates of the logarithmic average order of the autocorrelation function of the Mobius function μ has the asymptotic formula

$$\frac{1}{\log x} \sum_{n \leq x} \frac{\mu(n) \mu(n+t)}{n} = O\left(\frac{1}{(\log \log \log x)^c}\right), \quad (107)$$

where $t \neq 0$ is a fixed parameter, and $c > 0$ is a constant, as $x \rightarrow \infty$, confer [22], [29], [30, Chapter 5], et alii. Another attempt to prove the arithmetic average order of (108) was made in [18], and an improved error term $O((\sqrt{\log \log x})^{-2})$ was just

proposed in [9, Corollary 2]. Observe that a stronger error term, approximately $O((\log x)^{-2})$, is required to compute the arithmetic average orders (108) directly from (107), see [7, Exercise 2.12] for explanation. Here, the following result is considered.

Theorem 7.1. *Let $\mu : \mathbb{N} \rightarrow \{-1, 0, 1\}$ be the Mobius function. Then, for any sufficiently large number $x > 1$, and a fixed integer $t \neq 0$,*

$$\sum_{n \leq x} \mu(n)\mu(n+t) = O\left(\frac{x}{(\log x)^c}\right), \quad (108)$$

where $c > 0$ is a constant.

Section 7.1 to Section 7.3 cover the basic standard materials in analytic number theory, and new materials. The proof of Theorem 7.1 is assembled in Section 7.4.

7.1 Nonlinear Autocorrelation Functions Results

The number of squarefree integers have the following asymptotic formulas.

Lemma 7.1. *Let $\mu : \mathbb{Z} \rightarrow \{-1, 0, 1\}$ be the Mobius function. Then, for any sufficiently large number $x \geq 1$,*

$$\sum_{n \leq x} \mu^2(n) = \frac{6}{\pi^2}x + O(x^{1/2}).$$

Proof. Use identity $\mu(n)^2 = \sum_{d^2|n} \mu(d)$, and other elementary routines, or confer to the literature. ■

The constant coincides with the density of squarefree integers. Its approximate numerical value is

$$\frac{6}{\pi^2} = \prod_{p \geq 2} \left(1 - \frac{1}{p^2}\right) = 0.607988295164627617135754\dots, \quad (109)$$

where $p \geq 2$ ranges over the primes. The remainder term

$$E(x) = \sum_{n \leq x} \mu^2(n) - \frac{6}{\pi^2}x \quad (110)$$

is a topic of current research, its optimum value is expected to satisfies the upper bound $E(x) = O(x^{1/4+\varepsilon})$ for any small number $\varepsilon > 0$. Currently, $E(x) = O(x^{1/2}e^{-\sqrt{\log x}})$ is the best unconditional remainder term.

The earliest result on the autocorrelation of the squarefree indicator function $\mu^2(n)$ appears to be

$$\sum_{n \leq x} \mu^2(n)\mu^2(n+t) = cx + O(x^{2/3}), \quad (111)$$

where $c > 0$, is studied in [20]. Except for minor adjustments, the generalization to the k -tuple autocorrelation function has nearly the same structure.

Theorem 7.2. *Let $q \neq 0$, a_0, a_1, \dots, a_{k-1} be small integers, such that $0 \leq a_0 < a_1 < \dots < a_{k-1}$. Let $x \geq 1$ be a large number, and let $\mu : \mathbb{Z} \rightarrow \{-1, 0, 1\}$ be the Mobius function. Then,*

$$\sum_{n \leq x} \mu^2(n + a_0) \mu^2(n + a_1) \cdots \mu^2(n + a_{k-1}) = \prod_{p \geq 2} \left(1 - \frac{\varpi(p)}{p^2}\right) x + O(x^{2/3+\varepsilon}),$$

where

$$\varpi(p) = \#\{m \leq p^2 : qm + a_i \equiv 0 \pmod{p^2} \text{ for } i = 0, 1, 2, \dots, k-1\}. \quad (112)$$

The small number $\varepsilon > 0$ and the implied constant depends on $q \neq 0$.

Proof. Consult [20], [19, Theorem 1.2], and the literature. ■

Lemma 7.2. *Let $x \geq 1$ be a large number, and let $\mu : \mathbb{Z} \rightarrow \{-1, 0, 1\}$ be the Mobius function. If $t \neq 0$ is a fixed integer, then,*

$$\sum_{n \leq x} \mu(n)^2 \mu(n + t) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is an arbitrary constant.

Proof. Substitute the identity $\mu(n)^2 = \sum_{d^2|n} \mu(d)$, and switching the order of summation yield

$$\begin{aligned} \sum_{n \leq x} \mu(n)^2 \mu(n + t) &= \sum_{n \leq x} \mu(n + t) \sum_{d^2|n} \mu(d) & (113) \\ &= \sum_{d^2 \leq x} \mu(d) \sum_{\substack{n \leq x \\ d^2|n}} \mu(n + t) \\ &= \sum_{d^2 \leq x^{2\varepsilon}} \mu(d) \sum_{\substack{n \leq x \\ d^2|n}} \mu(n + t) + \sum_{x^{2\varepsilon} < d^2 \leq x} \mu(d) \sum_{\substack{n \leq x \\ d^2|n}} \mu(n + t), \end{aligned}$$

where $\varepsilon \in (0, 1/4)$. Applying Corollary 2.1 to the first subsum in the partition yields

$$\begin{aligned} \sum_{d^2 \leq x^{2\varepsilon}} \mu(d) \sum_{\substack{n \leq x \\ d^2|n}} \mu(n + t) &\leq \sum_{q \leq x^\varepsilon} \left| \mu(d) \sum_{\substack{n \leq x \\ m \equiv b \pmod{q}}} \mu(m) \right| & (114) \\ &= O\left(\frac{x}{(\log x)^c}\right), \end{aligned}$$

where $q = d^2$. An estimate of the second subsum in the partition yields

$$\begin{aligned} \sum_{x^{2\varepsilon} < d^2 \leq x} \mu(d) \sum_{\substack{n \leq x \\ d^2|n}} \mu(n + t) &\leq \sum_{x^{2\varepsilon} < d^2 \leq x} \sum_{\substack{n \leq x \\ d^2|n}} 1 & (115) \\ &\ll x \sum_{d^2 \leq x} \frac{1}{d^2} \\ &\ll x^{1-\varepsilon}. \end{aligned}$$

Summing (21) and (115) completes the verification. ■

7.2 Double-Sign Patterns Characteristic Functions

The analysis of single-sign pattern characteristic function

$$\mu^+(n) = \mu^2(n) \left(\frac{1 + \mu(n)}{2} \right) = \begin{cases} 1 & \text{if } \mu(n) = 1, \\ 0 & \text{if } \mu(n) \neq 1, \end{cases} \quad (116)$$

of the subset of integers

$$\mathcal{A}_0^+ = \{n \geq 1 : \mu(n) = 1\} \quad (117)$$

is well known. This idea is extended to the double-sign patterns $(\mu(n+a), \mu(n+b)) = (\pm 1, \pm 1)$, where $a, b \in \mathbb{Z}$ is a pair of small integers such that $a \neq b$. Other sign patterns are topics of current research, confer [11], [29, Corollary 1.7], [14], and similar literature, for details. A new and different approach to the analysis of double-sign patterns, triple-sign patterns, et cetera, based on elementary methods, is provided here.

Lemma 7.3. *Let $t \neq 0$ be an integer, and let $\mu(n) \in \{-1, 0, 1\}$ be the Mobius function. Then,*

$$\begin{aligned} \mu^{\pm\pm}(n, t) &= \mu^2(n)\mu^2(n+t) \left(\frac{1 \pm \mu(n)}{2} \right) \left(\frac{1 \pm \mu(n+t)}{2} \right) \\ &= \begin{cases} 1 & \text{if } \mu(n) = \pm 1, \mu(n+t) = \pm 1, \\ 0 & \text{if } \mu(n) \neq \pm 1, \mu(n+t) \neq \pm 1, \end{cases} \end{aligned} \quad (118)$$

are the characteristic functions of the subset of integers

$$\mathcal{A}^{\pm\pm}(t) = \{n \geq 1 : \mu(n) = \pm 1, \mu(n+t) = \pm 1\}. \quad (119)$$

7.3 Double-Sign Patterns Counting Functions

The single-sign patterns $\mu(n) = 1$ and $\mu(n) = -1$ have single-sign pattern counting functions of the forms

$$R^+(x) = \sum_{\substack{n \leq x \\ \mu(n)=1}} 1 = \sum_{n \leq x} \left(\frac{1 + \mu(n)}{2} \right) \mu^2(n) = \frac{1}{2} \frac{x}{\zeta(2)} + O\left(xe^{-c\sqrt{\log x}}\right), \quad (120)$$

and

$$R^-(x) = \sum_{\substack{n \leq x \\ \mu(n)=-1}} 1 = \sum_{n \leq x} \left(\frac{1 - \mu(n)}{2} \right) \mu^2(n) = \frac{1}{2} \frac{x}{\zeta(2)} + O\left(xe^{-c\sqrt{\log x}}\right), \quad (121)$$

these follow from Lemma 7.1 and Theorem 2.1. In terms of these functions, the summatory Mobius function has the asymptotic formula

$$\begin{aligned} R(x) &= \sum_{n \leq x} \mu(n) \\ &= R^+(x) - R^-(x) \\ &= \left(\left(\frac{1}{2} \frac{x}{\zeta(2)} \right) + O\left(xe^{-c\sqrt{\log x}}\right) \right) - \left(\left(\frac{1}{2} \frac{x}{\zeta(2)} \right) + O\left(xe^{-c\sqrt{\log x}}\right) \right) \\ &= O\left(xe^{-c\sqrt{\log x}}\right). \end{aligned} \quad (122)$$

Basically, it is a form of the prime number theorem.

The same principle is applied to the double-sign patterns $(\mu(n), \mu(n+t)) = (\pm 1, \pm 1)$ to derive the extended results provided here.

The double-sign pattern counting functions are defined by

$$R^{++}(t, x) = \sum_{\substack{n \leq x \\ \mu(n)=1, \mu(n+t)=1}} 1 = \sum_{n \in \mathcal{A}^{++}(t)} 1, \quad (123)$$

$$R^{+-}(t, x) = \sum_{\substack{n \leq x \\ \mu(n)=1, \mu(n+t)=-1}} 1 = \sum_{n \in \mathcal{A}^{+-}(t)} 1, \quad (124)$$

$$R^{-+}(t, x) = \sum_{\substack{n \leq x \\ \mu(n)=-1, \mu(n+t)=1}} 1 = \sum_{n \in \mathcal{A}^{-+}(t)} 1, \quad (125)$$

$$R^{--}(t, x) = \sum_{\substack{n \leq x \\ \mu(n)=-1, \mu(n+t)=-1}} 1 = \sum_{n \in \mathcal{A}^{--}(t)} 1. \quad (126)$$

The double-sign pattern counting functions (123) to (126) are precisely the cardinalities of the subsets of integers

1. $\mathcal{A}^{++}(t) \subset \mathbb{N}$,
2. $\mathcal{A}^{+-}(t) \subset \mathbb{N}$,
3. $\mathcal{A}^{-+}(t) \subset \mathbb{N}$,
4. $\mathcal{A}^{--}(t) \subset \mathbb{N}$,

defined in (119). In terms of the double-sign pattern counting functions, the Mobius autocorrelation function has form

$$\begin{aligned} R(t, x) &= \sum_{n \leq x} \mu(n)\mu(n+t) \\ &= R^{++}(t, x) - R^{+-}(t, x) + R^{-+}(t, x) - R^{--}(t, x). \end{aligned} \quad (127)$$

The next result is required to complete the analysis of the asymptotic formula for $R(t, x)$, which is completed in the next section.

Lemma 7.4. *Let $x \geq 1$ be a large number, and let $t \neq 0$ be a fixed integer. Then,*

$$R^{\pm\pm}(t, x) = \frac{1}{4}a_0(t)x + \frac{1}{4}R(t, x) + O\left(\frac{x}{(\log x)^c}\right),$$

where $a_0(t) > 0$, and $c > 0$ are constants.

Proof. Without loss in generality, consider the double-sign pattern $(\mu(n), \mu(n+t)) = (+1, +1)$. Now, use Lemma 7.3 to express the double-sign pattern counting function

as

$$\begin{aligned}
 4R^{++}(t, x) &= \sum_{n \leq x} \mu^{++}(t, n) & (128) \\
 &= \sum_{n \leq x} \mu^2(n) \mu^2(n+t) (1 + \mu(n)) (1 + \mu(n+t)) \\
 &= \sum_{n \leq x} \mu^2(n) \mu^2(n+t) (1 + \mu(n) + \mu(n+t) + \mu(n)\mu(n+t)) \\
 &= \sum_{n \leq x} \mu^2(n) \mu^2(n+t) + \sum_{n \leq x} \mu^3(n) \mu^2(n+t) \\
 &\quad + \sum_{n \leq x} \mu(n)^2 \mu^3(n+t) + \sum_{n \leq x} \mu^3(n) \mu^3(n+t) \\
 &\geq 0.
 \end{aligned}$$

The last four finite sums have the following evaluations or estimates.

$$1. \sum_{n \leq x} \mu^2(n) \mu^2(n+t) = a_0(t)x + O(x^{2/3}),$$

where $a_0(t) > 0$ is a constant, see Theorem 7.2.

$$2. \sum_{n \leq x} \mu^3(n) \mu^2(n+t) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is a constant, see Lemma 7.2.

$$3. \sum_{n \leq x} \mu^2(n) \mu^3(n+t) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$ is a constant, see Lemma 7.2.

$$4. \sum_{n \leq x} \mu^3(n) \mu^3(n+t) = \sum_{n \leq x} \mu(n) \mu(n+t),$$

since $\mu^{2k+1}(n) = \mu(n)$ for any integer k .

Summing these evaluations or estimates verifies the claim for $R^{++}(t, x) \geq 0$. The verifications for the next three double-sign pattern counting functions $R^{+-}(t, x) \geq 0$, $R^{-+}(t, x) \geq 0$, and $R^{--}(t, x) \geq 0$ are similar. \blacksquare

7.4 Result for the Mobius Autocorrelation Function over the Integers

The elementary results presented in the previous sections are utilized here to prove an effective form of the autocorrelation function $\sum_{n \leq x} \mu(n) \mu(n+t)$, which is a

significant improvement over the current results in the literature, see (107). The average result claims that

$$\sum_{t \leq T} \sum_{n \leq x} \mu(n)\mu(n+t) = o(Tx), \quad (129)$$

see [22, Theorem 1.1] for the precise details, and the conjecture due to Chowla, see [1], and [26], claims that

$$\sum_{n \leq x} \mu(n)\mu(n+t) = o(x), \quad (130)$$

for any small fixed integer $t \neq 0$.

Proof: (Theorem 7.1). Replace the asymptotic estimates for the double-sign pattern counting functions $R^{\pm\pm}(t, x) \geq 0$ computed in Lemma 7.4 into the Mobius autocorrelation function, see (127), to obtain

$$\begin{aligned} R(t, x) &= \sum_{n \leq x} \mu(n)\mu(n+t) & (131) \\ &= R^{++}(t, x) - R^{+-}(t, x) + R^{--}(t, x) - R^{-+}(t, x) \\ &= \frac{1}{4}a_0(t)x + \frac{1}{4}R(t, x) + O\left(\frac{x}{(\log x)^c}\right) \\ &\quad - \frac{1}{4}a_0(t)x - \frac{1}{4}R(t, x) + O\left(\frac{x}{(\log x)^c}\right) \\ &\quad + \frac{1}{4}a_0(t)x + \frac{1}{4}R(t, x) + O\left(\frac{x}{(\log x)^c}\right) \\ &\quad - \frac{1}{4}a_0(t)x - \frac{1}{4}R(t, x) + O\left(\frac{x}{(\log x)^c}\right) \\ &= O\left(\frac{x}{(\log x)^c}\right), \end{aligned}$$

where $a_0(t) > 0$, and $c > 0$ are constants. Quod erat demonstrandum. ■

8 Results for the Liouville Autocorrelation Function over the Integers

The current estimates of the logarithmic average orders of the autocorrelations of the Liouville function has the asymptotic formulae

$$\frac{1}{\log x} \sum_{n \leq x} \frac{\lambda(n)\lambda(n+t)}{n} = O\left(\frac{1}{(\log \log \log x)^c}\right), \quad (132)$$

respectively, where $t \neq 0$ is a fixed parameter, and $c > 0$ is a constant, as $x \rightarrow \infty$, confer [22], [29], [30, Chapter 5], et alii. Another attempt to prove the arithmetic average order of (133) was made in [18], and an improved error term $O((\sqrt{\log \log x})^{-2})$ was just proposed in [9, Corollary 2]. Observe that a stronger error term, approximately $O((\log x)^{-2})$, is required to compute the arithmetic average orders (133)

directly from (132), see [7, Exercise 2.12] for explanation. Here, the following result is considered.

Theorem 8.1. *Let $\lambda : \mathbb{N} \rightarrow \{-1, 1\}$ be the Liouville function, and let $a, b \in \mathbb{Z}$ be a pair of fixed integers such that $a \neq b$. Then, for any sufficiently large number $x > 1$,*

$$\sum_{n \leq x} \lambda(n+a)\lambda(n+b) = O\left(xe^{-c\sqrt{\log x}}\right), \quad (133)$$

where $c > 0$ is an absolute constant.

Section 8.1 to Section 8.2 cover basic materials in analytic number theory, and new materials. The proof of Theorem 8.1 is spliced together in Section 8.3.

8.1 Double-Sign Patterns Characteristic Functions

The analysis of single-sign pattern characteristic function

$$\lambda^+(n) = \left(\frac{1 + \lambda(n)}{2}\right) = \begin{cases} 1 & \text{if } \lambda(n) = 1, \\ 0 & \text{if } \lambda(n) \neq 1, \end{cases} \quad (134)$$

of the subset of integers

$$\mathcal{B}_0^+ = \{n \geq 1 : \lambda(n) = 1\} \quad (135)$$

is well known. A simple extension to the double-sign patterns $(\lambda(n+a), \lambda(n+b)) = (\pm 1, \pm 1)$ is considered here. Other sign patterns are topics of current research, confer [11], [29, Corollary 1.7], [14], [27] and similar literature, for details.

Lemma 8.1. *Let $a, b \in \mathbb{Z}$ be small integers, such that $a \neq b$, and let $\lambda(n) \in \{-1, 1\}$ be the Liouville function. Then,*

$$\begin{aligned} \lambda^{\pm\pm}(a, b, n) &= \left(\frac{1 \pm \lambda(n+a)}{2}\right) \left(\frac{1 \pm \lambda(n+b)}{2}\right) \\ &= \begin{cases} 1 & \text{if } \lambda(n+a) = \pm 1, \lambda(n+b) = \pm 1, \\ 0 & \text{if } \lambda(n+a) \neq \pm 1, \lambda(n+b) \neq \pm 1, \end{cases} \end{aligned} \quad (136)$$

are the characteristic functions of the subset of integers

$$\mathcal{B}^{\pm\pm}(a, b) = \{n \geq 1 : \lambda(n+a) = \pm 1, \lambda(n+b) = \pm 1\}. \quad (137)$$

8.2 Double-Sign Patterns Counting Functions

The single-sign patterns $\lambda(n) = 1$ and $\lambda(n) = -1$ have single-sign pattern counting functions of the forms

$$Q^+(x) = \sum_{\substack{n \leq x \\ \lambda(n)=1}} 1 = \sum_{n \leq x} \left(\frac{1 + \lambda(n)}{2}\right) = \frac{1}{2}[x] + O\left(xe^{-c\sqrt{\log x}}\right), \quad (138)$$

and

$$Q^-(x) = \sum_{\substack{n \leq x \\ \lambda(n) = -1}} 1 = \sum_{n \leq x} \left(\frac{1 - \lambda(n)}{2} \right) = \frac{1}{2}[x] + O\left(xe^{-c\sqrt{\log x}}\right), \quad (139)$$

where $[x]$ is the largest integer function, these follow from Theorem 4.1. The same principle is applied to the double-sign patterns $(\lambda(n+a), \lambda(n+b)) = (\pm 1, \pm 1)$ to derive the extended results provided here.

The double-sign pattern counting functions are defined by

$$Q^{++}(a, b, x) = \sum_{\substack{n \leq x \\ \lambda(n+a)=1, \lambda(n+b)=1}} 1 = \sum_{n \in \mathcal{B}^{++}(a,b)} 1, \quad (140)$$

$$Q^{+-}(a, b, x) = \sum_{\substack{n \leq x \\ \lambda(n)=1, \lambda(n+t)=-1}} 1 = \sum_{n \in \mathcal{B}^{+-}(a,b)} 1, \quad (141)$$

$$Q^{-+}(a, b, x) = \sum_{\substack{n \leq x \\ \lambda(n+a)=-1, \lambda(n+b)=1}} 1 = \sum_{n \in \mathcal{B}^{-+}(a,b)} 1, \quad (142)$$

$$Q^{--}(a, b, x) = \sum_{\substack{n \leq x \\ \lambda(n+a)=-1, \lambda(n+b)=-1}} 1 = \sum_{n \in \mathcal{B}^{--}(a,b)} 1. \quad (143)$$

The double-sign pattern counting functions (140) to (143) are precisely the cardinalities of the subsets of integers

1. $\mathcal{B}^{++}(a, b) \subset \mathbb{N}$,
2. $\mathcal{B}^{+-}(a, b) \subset \mathbb{N}$,
3. $\mathcal{B}^{-+}(a, b) \subset \mathbb{N}$,
4. $\mathcal{B}^{--}(a, b) \subset \mathbb{N}$,

defined in (137). In terms of the double-sign pattern counting functions, the Liouville autocorrelation function has form

$$\begin{aligned} Q(a, b, x) &= \sum_{n \leq x} \lambda(n+a)\lambda(n+b) \\ &= Q^{++}(a, b, x) - Q^{+-}(a, b, x) + Q^{-+}(a, b, x) - Q^{--}(a, b, x). \end{aligned} \quad (144)$$

The next result is required to complete the analysis of the asymptotic formula for $Q(a, b, x)$, which is completed in the next section.

Lemma 8.2. *Let $x \geq 1$ be a large number, and let $a, b \in \mathbb{Z}$ be small fixed integer, such that $a \neq b$. Then,*

$$Q^{\pm\pm}(a, b, x) = \frac{1}{4}[x] + \frac{1}{4}Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right),$$

where $c > 0$ is an absolute constant.

Proof. Without loss in generality, consider the double-sign pattern $(\lambda(n+a), \lambda(n+b)) = (+1, +1)$. Now, use Lemma 8.1 to express the double-sign pattern counting function as

$$\begin{aligned}
 4Q^{++}(a, b, x) &= \sum_{n \leq x} \lambda^{++}(a, b, n) & (145) \\
 &= \sum_{n \leq x} (1 + \lambda(n+a))(1 + \lambda(n+b)) \\
 &= \sum_{n \leq x} (1 + \lambda(n+a) + \lambda(n+b) + \lambda(n+a)\lambda(n+b)) \\
 &= \sum_{n \leq x} 1 + \sum_{n \leq x} \lambda(n+a) \\
 &\quad + \sum_{n \leq x} \lambda(n+b) + \sum_{n \leq x} \lambda(n+a)\lambda(n+b) \\
 &\geq 0.
 \end{aligned}$$

The last four finite sums have the following evaluations or estimates.

1. $\sum_{n \leq x} 1 = [x]$, where $[x]$ is the largest integer function.
2. $\sum_{n \leq x} \lambda(n+a) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 4.1.
3. $\sum_{n \leq x} \lambda(n+b) = O\left(xe^{-c\sqrt{\log x}}\right)$, see Theorem 4.1.
4. $\sum_{n \leq x} \lambda(n+a)\lambda(n+b) = Q(a, b, x)$.

In all cases $c > 0$ is an absolute constant. Summing these evaluations or estimates verifies the claim for $Q^{++}(a, b, x) \geq 0$. The verifications for the next three double-sign pattern counting functions $Q^{+-}(a, b, x) \geq 0$, $Q^{-+}(a, b, x) \geq 0$, and $Q^{--}(a, b, x) \geq 0$ are similar. \blacksquare

8.3 Result for the Liouville Autocorrelation Function over the Integers

The elementary results presented in the previous sections are utilized here to prove an effective form of the autocorrelation function $\sum_{n \leq x} \lambda(n+a)\lambda(n+b)$, which is a significant improvement over the current results in the literature, see (132). The average result claims that

$$\sum_{t \leq T} \sum_{n \leq x} \lambda(n+a)\lambda(n+b) = o(Tx), \quad (146)$$

see [22, Theorem 1.1] for the precise details, and the conjecture due to Chowla, see [1], and [26], claims that

$$\sum_{n \leq x} \lambda(n+a)\lambda(n+b) = o(x), \quad (147)$$

for any pair of small fixed integers $a, b \in \mathbb{Z}$, such that $a \neq b$.

Proof: (Theorem 8.1). Replace the asymptotic estimates for the double-sign pattern counting functions $Q^{\pm\pm}(a, b, x) \geq 0$ computed in Lemma 8.2 into the Liouville autocorrelation function, see (144), to obtain

$$\begin{aligned} Q(a, b, x) &= \sum_{n \leq x} \lambda(n+a)\lambda(n+b) & (148) \\ &= Q^{++}(a, b, x) - Q^{+-}(a, b, x) + Q^{--}(a, b, x) - Q^{-+}(a, b, x) \\ &= \frac{1}{4}[x] + \frac{1}{4}Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad - \frac{1}{4}[x] - \frac{1}{4}Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad + \frac{1}{4}[x] + \frac{1}{4}Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &\quad - \frac{1}{4}[x] - \frac{1}{4}Q(a, b, x) + O\left(xe^{-c\sqrt{\log x}}\right) \\ &= O\left(xe^{-c\sqrt{\log x}}\right), \end{aligned}$$

where $[x]$ is the largest integer function, and $c > 0$ is an absolute constant. Quod erat demonstrandum. \blacksquare

9 Open Problems

9.1 Mobius autocorrelation function problems

Problem 9.1. Let $x \geq x_0$ be a large number, and let $a < b$ be small integers. Determine the oscillation results for the autocorrelation functions

$$\sum_{n \leq x} \mu(n+a)\mu(n+b) = \Omega(x^\beta)$$

over the integers, and

$$\sum_{p \leq x} \mu(p+a)\mu(p+b) = \Omega(x^\beta)$$

over the shifted primes, where $\beta \in (1/2, 1)$. In [15], there is some evidence and material for random autocorrelation functions.

Problem 9.2. Let $x \geq x_0$ be a large number, and let $a < b$ be small integers. Assume the RH. Determine the conditional results for the autocorrelation functions

$$\sum_{p \leq x} \mu(p+a)\mu(p+b) = O(x^\beta)$$

over the integers, and

$$\sum_{p \leq x} \mu(p+a)\mu(p+b) = O(x^\beta)$$

over the shifted primes, where $\beta \in (1/2, 1)$.

Problem 9.3. Let $x \geq x_0$ be a large number, let $y \geq x^{1/2}$, and let $a < b$ be small integers. Determine a nontrivial upper bound for the autocorrelation function

$$\sum_{x \leq n \leq x+y} \mu(n+a)\mu(n+b)$$

over the integers, and

$$\sum_{x \leq p \leq x+y} \mu(p+a)\mu(p+b)$$

over the shifted primes in the short interval $[x, x+y]$.

Problem 9.4. Let $x \geq x_0$ be a large number, and let $a_0 < a_1 < a_2$ be small integers. Either conditionally or unconditionally, use sign patterns techniques to verify the asymptotic for the autocorrelation function

$$\sum_{n \leq x} \mu(n+a_0)\mu(n+a_1)\mu(n+a_2) = O\left(\frac{x}{(\log x)^c}\right)$$

over the integers.

Problem 9.5. Let $a_0 < a_1 < \dots < a_{k-1}$ be a subset of small integers. Determine the maximal value of $k \geq 2$ for which the autocorrelations

$$\sum_{n \leq x} \mu(n+a_0)\mu(n+a_1) \cdots \mu(n+a_{k-1})$$

over the integers, and

$$\sum_{p \leq x} \mu(p+a_0)\mu(p+a_1) \cdots \mu(p+a_{k-1})$$

over the shifted primes, have nontrivial upper bounds.

Problem 9.6. Let $x \geq x_0$ be a large number, let $1 \leq a < q \ll (\log x)^c$, with $c \geq 0$, and let $a_0 < a_1$ be small integers. Determine a nontrivial upper bound for the autocorrelation function

$$\sum_{\substack{x \leq n \\ n \equiv a \pmod q}} \mu(n+a_0)\mu(n+a_1)$$

over the integers in arithmetic progression, and

$$\sum_{\substack{x \leq n \\ n \equiv a \pmod q}} \mu(p+a_0)\mu(p+a_1)$$

over the shifted primes in arithmetic progression.

9.2 Liouville autocorrelation function problems

Problem 9.7. Let $x \geq x_0$ be a large number, and let $a < b$ be small integers. Determine the oscillation results for the autocorrelation functions

$$\sum_{n \leq x} \lambda(n+a)\lambda(n+b) = \Omega(x^\beta)$$

over the integers, and

$$\sum_{p \leq x} \lambda(p+a)\lambda(p+b) = \Omega(x^\beta)$$

over the shifted primes, where $\beta \in (1/2, 1)$. In [15], there is some evidence and material for random autocorrelation functions.

Problem 9.8. Let $x \geq x_0$ be a large number, and let $a < b$ be small integers. Assume the RH. Determine the conditional results for the autocorrelation functions

$$\sum_{n \leq x} \lambda(n+a)\lambda(n+b) = O(x^\beta)$$

over the integers, and

$$\sum_{p \leq x} \lambda(p+a)\lambda(p+b) = O(x^\beta)$$

over the shifted primes, where $\beta \in (1/2, 1)$.

Problem 9.9. Let $x \geq x_0$ be a large number, let $y \geq x^{1/2}$, and let $a < b$ be small integers. Determine a nontrivial upper bounds for the autocorrelation function

$$\sum_{x \leq n \leq x+y} \lambda(n+a)\lambda(n+b)$$

over the integers, and

$$\sum_{x \leq p \leq x+y} \lambda(p+a)\lambda(p+b)$$

over the shifted primes in the short interval $[x, x+y]$.

Problem 9.10. Let $x \geq x_0$ be a large number, and let $a_0 < a_1 < a_2$ be small integers. Use sign patterns techniques to verify the asymptotic for the autocorrelation function

$$\sum_{n \leq x} \lambda(n+a_0)\lambda(n+a_1)\lambda(n+a_2) = O\left(xe^{-c\sqrt{\log x}}\right)$$

over the integers

Problem 9.11. Let $a_0 < a_1 < \dots < a_{k-1}$ be a subset of small integers. Determine the maximal value of $k \geq 2$ for which the autocorrelations

$$\sum_{n \leq x} \lambda(n+a_0)\lambda(n+a_1)\cdots\lambda(n+a_{k-1})$$

over the integers, and

$$\sum_{p \leq x} \lambda(p+a_0)\lambda(p+a_1)\cdots\lambda(p+a_{k-1})$$

over the shifted primes, have nontrivial upper bounds.

Problem 9.12. Let $x \geq x_0$ be a large number, let $1 \leq a < q \ll (\log x)^c$, with $c \geq 0$, and let $a_0 < a_1$ be small integers. Determine a nontrivial upper bound for the autocorrelation function

$$\sum_{\substack{x \leq n \\ n \equiv a \pmod{q}}} \lambda(n + a_0)\lambda(n + a_1)$$

over the integers in arithmetic progression, and

$$\sum_{\substack{x \leq n \\ n \equiv a \pmod{q}}} \lambda(p + a_0)\lambda(p + a_1)$$

over the shifted primes in arithmetic progression.

9.3 Intercorrelation functions problems

Problem 9.13. Let $a < b$ be a pair of small integers. Use the identities $\mu(n) = \lambda(n)\mu^2(n)$ and $\lambda(n) = \sum_{d^2|n} \mu(n/d^2)$, etc., to find an easy way to prove that

$$\sum_{n \leq x} \mu(n + a)\mu(n + b) = O\left(\frac{x}{(\log x)^c}\right)$$

implies that

$$\sum_{n \leq x} \lambda(n + a)\lambda(n + b) = O\left(\frac{x}{(\log x)^c}\right),$$

where $c > 0$. Is the converse true?

9.4 Hardy-Littlewood-Chowla conjecture problems

Problem 9.14. Let $a < b$ be a pair of small integers. Use sign patterns techniques to estimate the twisted finite sum

$$\sum_{n \leq x} \Lambda(n)\mu(n + a)\mu(n + b).$$

Problem 9.15. Let $a < b$ be a pair of small integers. Use sign patterns techniques to estimate the twisted finite sum

$$\sum_{n \leq x} \Lambda(n)\lambda(n + a)\lambda(n + b).$$

References

- [1] Chowla, S. *The Riemann hypothesis and Hilbert's tenth problem*. Mathematics and Its Applications, Vol. 4. Gordon and Breach Science Publishers, New York-London-Paris, 1965. [27](#), [36](#), [40](#)
 - [2] dePolignac, A. *Six propositions arithmologiques d'eduites du crible d'Eratosthene*. Nouv. Ann. Math. 8 (1849), 423-429. [27](#)
 - [3] De Koninck, Jean-Marie; Luca, Florian. *Analytic number theory. Exploring the anatomy of integers*. Graduate Studies in Mathematics, 134. American Mathematical Society, Providence, RI, 2012. [5](#), [6](#), [8](#), [28](#)
 - [4] NIST *Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, 2019. F. W. J. Olver, ..., and M. A. McClain, eds. [10](#)
 - [5] Ellison, William; Ellison, Fern. *Prime numbers*. A Wiley-Interscience Publication. John Wiley and Sons, Inc., New York; Hermann, Paris, 1985. [10](#)
 - [6] Friedlander, John; Iwaniec, Henryk. *Opera de cribro*. American Mathematical Society Colloquium Publications, 57. American Mathematical Society, Providence, RI, 2010. [6](#), [7](#), [28](#)
 - [7] Hildebrand, Adolph. *Introduction to Analytic Number Theory Math 531 Lecture Notes, Fall 2005*. <http://www.math.uiuc.edu/hildebr/ant>. [31](#), [37](#)
 - [8] Hildebrand, Adolph. *Additive and Multiplicative Functions on Shifted Primes*. Volumes 3-59, Issue 2, September 1989, Pages 209-232. [3](#)
 - [9] Helfgott, Harald Andres. *Expansion, divisibility and parity: an explanation*. <http://arxiv.org/abs/2201.00799>. [31](#), [36](#)
 - [10] Hardy, G. H.; Wright, E. M. *An introduction to the theory of numbers*. Sixth edition. Oxford University Press, Oxford, 2008. [5](#)
 - [11] Hildebrand, Adolph. *On consecutive values of the Liouville function*. Enseign. Math. (2) 32 (1986), no. 3-4, 219-226. [12](#), [15](#), [22](#), [25](#), [33](#), [37](#)
 - [12] Hardy, G. H.; Littlewood, J. E. *Some problems of Partitio Numerorum, III: On the expression of a number as a sum of primes*. Acta Math. 44 (1923), 1-70. [27](#)
 - [13] Iwaniec, Henryk; Kowalski, Emmanuel. *Analytic number theory*. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.
 - [14] Kaczorowski, J.; Pintz, J. *Oscillatory properties of arithmetic functions I*. Acta Math. Acad. Sci. Hung. 48 (1986), 173-185. doi: 10.1007/BF01949062) [12](#), [15](#), [22](#), [25](#), [33](#), [37](#)
 - [15] Klurman, O.; Shkredov, I. D.; Xu, M. *On the random Chowla conjecture*. <http://arxiv.org/abs/2202.08767>. [40](#), [42](#)
-

- [16] Lichtman, Jared. *Averages of the Mobius function on shifted primes*. <http://arxiv.org/abs/2009.08969>. 3, 11, 21
 - [17] Lichtman, Jared; Teräväinen, Joni. *On the Hardy-Littlewood-Chowla conjecture on average*. <http://arxiv.org/abs/2111.08912>. 28
 - [18] Mangerel, A. *On the Bivariate Erdos-Kac Theorem and Correlations of the Mobius Function*. <http://arxiv.org/abs/1612.09544>. 30, 36
 - [19] Mennema, Ingela. *The distribution of consecutive square-free numbers*. Master Thesis, Leiden University, 2017. 32
 - [20] Mirsky, L. *Note on an asymptotic formula connected with r -free integers*. Quart. J. Math., Oxford Ser. 18, (1947). 178-182. 31, 32
 - [21] McNamara, Redmond. *Dynamical Methods for the Sarnak and Chowla Conjectures*. University Of California, Los Angeles, Dissertation 2021. 3
 - [22] Matomaki, Kaisa, Radziwiłł, M., Tao, T. *An averaged form of Chowla's conjecture*. <http://arxiv.org/abs/1503.05121>. 30, 36, 40
 - [23] Murty, M. R.; Vatwani, Akshaa. *Twin primes and the parity problem*. Journal of Number Theory Volume 180, November 2017, Pages 643-659. 30
 - [24] Montgomery, Hugh L.; Vaughan, Robert C. *Multiplicative number theory. I. Classical theory*. Cambridge University Press, Cambridge, 2007. 5
 - [25] Pintz, J. *An approximation to the twin prime conjecture and the parity phenomenon*. Indagationes Mathematicae, Volume 26, Issue 5, December 2015, Pages 883-896. 12, 15, 21, 22, 25
 - [26] Ramare, Olivier. *Chowla's Conjecture: From the Liouville Function to the Moebius Function*. Part of the Lecture Notes in Mathematics book series (LNM, volume 2213), 16 June 2018. 27, 36, 40
 - [27] Srinivasan, Anitha. *Infinitely many sign changes of the Liouville function*. Proc. Amer. Math. Soc. 150 (2022), 3799-3809. 22, 37
 - [28] Siebert, Hartmut; Wolke, D. *über einige Analoga zum Bombierischen Primzahlsatz*. Math. Z. 122 (1971), no. 4, 327-341. 5
 - [29] Tao, T. *The logarithmically averaged Chowla and Elliott conjectures for two-point correlations*. <http://arxiv.org/abs/1509.05422>. 12, 15, 22, 25, 30, 33, 36, 37
 - [30] Teravainen, T. *Topics in Multiplicative Number Theory*. University of Turku Dissertation 2018. 3, 30, 36
 - [31] Wolke, Dieter. *über die mittlere Verteilung der Werte zahlentheoretischer Funktionen auf Restklassen. I*. Math. Ann. 202 (1973), 1-25. 5
-

- [32] Wabnitz, Paul. *Sarnak's Conjecture about Mobius Function Randomness in Deterministic Dynamical Systems*. Universitat Leipzig, Dissertation 2016. [3](#)

ResultMobiusFunctionShiftedPrimes-SignPatterns-07-04-22-65-arxiv.tex.