

# RÉNYI'S ENTROPY ON LORENTZIAN SPACES. TIMELIKE CURVATURE-DIMENSION CONDITIONS

MATHIAS BRAUN

*Dedicated to Professor Karl-Theodor Sturm on the occasion of his sixtieth birthday*

ABSTRACT. For a measured Lorentzian space  $(M, \mathbf{d}, \mathbf{m}, \ll, \leq, \tau)$  in the sense of Kunzinger, Sämann, Cavalletti, and Mondino, we introduce and study synthetic notions of timelike lower Ricci curvature bounds by  $K \in \mathbf{R}$  and upper dimension bounds by  $N \in [1, \infty)$ , namely the timelike curvature-dimension conditions  $\text{TCD}_p(K, N)$  and  $\text{TCD}_p^*(K, N)$  in weak and strong forms, and the timelike measure-contraction properties  $\text{TMCP}_p(K, N)$  and  $\text{TMCP}_p^*(K, N)$ ,  $p \in (0, 1)$ . These are formulated by convexity properties of the Rényi entropy with respect to  $\mathbf{m}$  along  $\ell_p$ -geodesics of probability measures. This complements recent information-theoretic approaches to spacetime curvature.

We show many features of these notions, including their compatibility with the smooth setting, sharp geometric inequalities, stability, equivalence of the named weak and strong versions, local-to-global properties, and uniqueness of chronological  $\ell_p$ -optimal couplings and  $\ell_p$ -geodesics. Moreover, we prove the equivalence of  $\text{TCD}_p^*(K, N)$  and  $\text{TMCP}_p^*(K, N)$  to their respective entropic counterparts in the sense of Cavalletti and Mondino.

Some of these results are obtained under the assumption of timelike  $p$ -essential nonbranching, a newly introduced Lorentzian concept conjectured to be strictly weaker than timelike nonbranching.

## CONTENTS

1. Introduction	2
2. Preliminaries	9
2.1. Volume distortion coefficients	9
2.2. Nonsmooth Lorentzian spaces	10
2.3. Optimal transport on Lorentzian spaces	12
2.4. Timelike essential nonbranching	16
3. Timelike curvature-dimension condition	17
3.1. Definition and basic properties	17
3.2. Geometric inequalities	19
3.3. Stability	22
3.4. Good geodesics	33
3.5. Equivalence with the entropic TCD condition	35
3.6. From local to global	40
3.7. Universal cover	44
4. Timelike measure-contraction property	46

*Date:* August 8, 2022.

2010 *Mathematics Subject Classification.* 49J52, 53C50, 58E10, 58Z05, 83C99.

*Key words and phrases.* Timelike geodesics; Timelike curvature-dimension condition; Rényi entropy; Strong energy condition; Lorentzian pre-length spaces.

The author is sincerely grateful to Robert McCann for help- and fruitful discussions. This research is supported in part by Canada Research Chairs Program funds and a Natural Sciences and Engineering Research Council of Canada Grant (2020-04162) held by Robert McCann.

4.1. Definition and basic properties	46
4.2. Stability	48
4.3. Good geodesics	52
4.4. Uniqueness of $\ell_p$ -optimal couplings and $\ell_p$ -geodesics	53
4.5. Equivalence with the entropic TMCP condition	58
Appendix A. Smooth Lorentzian spacetimes	59
A.1. Smooth timelike curvature-dimension condition	59
A.2. Smooth timelike measure-contraction property	63
Appendix B. Geodesics of probability measures	63
B.1. Causal and timelike geodesics minimizing a Lagrangian action	63
B.2. Proper-time parametrizations	65
References	67

## 1. INTRODUCTION

Throughout this paper, let  $(M, \mathbf{d})$  be a nonempty metric space. All topological and measure-theoretic properties will be understood with respect to the topology induced by  $\mathbf{d}$  and its induced Borel  $\sigma$ -algebra. The completeness of  $\mathbf{d}$  is *not* assumed; every subsequent result which requires this regularity is phrased relative to compact (hence complete and separable) subsets of  $M$ . Moreover, let  $\mathbf{m}$  be a given Radon measure on  $M$  with full support, symbolically  $\text{spt } \mathbf{m} = M$ .

**Background.** Mathematical general relativity has recently developed a promising novel research direction. McCann [66] pioneered lower boundedness of the Ricci tensor  $\text{Ric}$  of a smooth Lorentzian spacetime  $M$  in timelike directions to be equivalent to convexity properties of the Boltzmann entropy  $\text{Ent}_{\mathbf{m}}$ , for  $\mathbf{m}$  a possibly weighted Lorentzian volume measure, along “chronological geodesics” on the space of probability measures  $\mathcal{P}(M)$  in the entropic sense [32]. This includes the *strong energy condition* of Hawking and Penrose [49, 51, 73] which, for a vanishing cosmological constant, reduces to timelike nonnegativity of  $\text{Ric}$ . Mondino and Suhr [69] provided a similar description of the Einstein equations, i.e. timelike lower and upper Ricci curvature bounds, through synthetic optimal transport means. In a different direction, Kunzinger and Sämman [58], see also [82] for a related approach, set up the theory of Lorentzian (pre-length, length, and geodesic) spaces. These are singular analogues of smooth Lorentzian spacetimes, comparable to metric spaces generalizing Riemannian manifolds. These discoveries let Cavalletti and Mondino [24] introduce the *entropic timelike curvature-dimension condition*  $\text{TCD}_p^e(K, N)$  and the *entropic timelike measure-contraction property*  $\text{TMCP}_p^e(K, N)$ ,  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Both are ways to concretize the meaning of a Lorentzian space measured by  $\mathbf{m}$  to admit timelike Ricci curvature bounded from below by  $K$  and dimension bounded from above by  $N$ . As indicated above, the inherent formulations are phrased by convexity properties of  $\text{Ent}_{\mathbf{m}}$  along  $\ell_p$ -geodesics, thus make sense a priori in far more general frameworks than smooth Lorentzian spacetimes. Here,  $\ell_p$  is the so-called *p-Lorentz-Wasserstein distance* on  $\mathcal{P}(M)$ , see (1.1), first introduced in [31] and studied further in [24, 53, 66, 69, 87]. Recently [16], we have furthermore introduced an “infinite-dimensional” Lorentzian analogue of [85].

This line of thought to axiomatize synthetic Ricci curvature bounds is not new. In metric geometry, results for Riemannian manifolds [28, 72, 93] analogous to [66, 69] lead Sturm [85, 86], independently Lott and Villani [61], to paraphrase lower Ricci curvature and upper dimension bounds for metric measure spaces by convexity of certain entropy functionals along geodesics in  $\mathcal{P}(M)$  with respect to

the 2-Wasserstein distance  $W_2$ . This was the starting point of an intense activity on so-called CD or RCD spaces, vivid to date, which produced remarkable outcomes and singular generalizations of known smooth results. Among others, we mention the splitting theorem [42], structure theory [18, 68], a first and second order Sobolev calculus [5, 6, 7, 43, 44, 79], rigidity [33, 54, 55], and stability [8, 45, 88].

Synthetic timelike Ricci curvature bounds for Lorentzian spaces through optimal transport as initiated by the mentioned works [24, 66, 69] are expected to lead to an equally flourishing theory. There are many smooth results, both classical and recent, that might (or, very occasionally, already do) admit generalizations to such spaces, notably singularity theorems à la Hawking and Penrose [2, 24, 46, 50, 57, 62, 94] and à la Gannon and Lee [38, 39, 59, 80], as well as splitting theorems [11, 34, 37, 63]. See the recent survey article [83]. Already from the conceptual viewpoint, [24] sheds a novel light to recent efforts to understand Lorentzian spacetimes with low regularity metrics, where curvature bounds are phrased distributionally thus far [41, 46, 84]; the setting of [24, 58] allows not only the potential metric tensor, but also the base space to be irregular. This extension is desirable since general relativity generally *predicts* the presence of singularities under extreme physical conditions, e.g. in black holes, causal Fermion systems [35, 36], or causal sets [15] which, even worse, might not be approximable by smooth structures (even if so, they can well be irregular). See also [58, 65] and the introduction of [24]. Moreover, by what we learned from [69] such a theory is a further step to understand the Einstein equations [56, 78] in more singular, yet physically relevant settings [20, 47, 60, 64, 74, 90, 91], which also relates to the *censorship conjectures* [26, 29]. For recently introduced synthetic notions of timelike *upper* Ricci curvature bounds, see [25].

Information-theoretic approaches to spacetime curvature, towards which we take a new direction, are based on vast physical evidences and connections of gravity and entropies already discovered up to half a decade ago [10, 12, 52, 89].

**Objective.** The goal of this paper is to introduce and investigate new timelike curvature-dimension conditions and timelike measure-contraction properties for measured Lorentzian spaces by optimal transport, different from [24]: instead of the Boltzmann entropy, we use the *Rényi entropy*, cf. (1.4) below. Our treatise is the Lorentzian counterpart of the groundbreaking works [9, 86], as is [24] relative to [32]. It sets up an especially geometrically valuable viewpoint to synthetic timelike lower Ricci curvature and upper dimension bounds. In fact, even in the smooth case, we are the first to study the Rényi entropy within general relativity.

**Setting.** Before making all above indicated objects, notions, and results precise, we outline our framework, see Section 2.2 and Section 2.3 for details.

Let  $(M, d, \ll, \leq, \tau)$  be a Lorentzian pre-length space in the sense of [58]. Here,  $\ll$  and  $\leq$  are generalized notions of *chronology* and *causality* between points in  $M$ . Moreover,  $\tau: M^2 \rightarrow [0, \infty]$  is the so-called *time separation function*. Roughly speaking,  $\tau(x, y)$  measures the maximal proper-time a point  $x \in M$  in spacetime needs to travel to  $y \in M$ . Compared to [9, 86],  $\tau$  takes over the role of the metric in the formulation of our timelike curvature-dimension conditions and timelike measure-contraction properties, yet this analogy should be handled carefully; for instance,  $\tau$  satisfies the *reverse* triangle inequality (2.4), it is symmetric only in pathological cases, and  $\tau(x, y) = 0$  as soon as  $y$  is not in the chronological future of  $x$ . With few exceptions, we naturally assume  $(M, d, \ll, \leq, \tau)$  to be causally closed,  $\mathcal{K}$ -globally hyperbolic, and geodesic in addition, cf. Assumption 3.1. Inter alia, this hypothesis ensures that  $\tau$  is finite and continuous. When endowed with a reference measure  $\mathfrak{m}$  as initially fixed, for terminological cleanness, in this introduction we call  $(M, d, \mathfrak{m}, \ll, \leq, \tau)$  *measured Lorentzian space*.

For  $p \in (0, 1]$ , define the total transport cost  $\ell_p: \mathcal{P}(M)^2 \rightarrow [0, \infty] \cup \{-\infty\}$  by

$$\ell_p(\mu, \nu) := \sup \|\tau\|_{L^p(M^2, \pi)}. \quad (1.1)$$

The supremum is taken over all *causal* couplings  $\pi \in \Pi(\mu, \nu)$ , i.e.  $x \leq y$  for  $\pi$ -a.e.  $(x, y) \in M^2$ , subject to the usual convention  $\sup \emptyset := -\infty$ . (If such a  $\pi$  satisfies  $x \ll y$  for  $\pi$ -a.e.  $(x, y) \in M^2$ , it is called *chronological*.) A causal coupling attaining this supremum is called  $\ell_p$ -*optimal*. Similarly as for  $\tau$ , we interpret the quantity  $\ell_p$  as measuring the maximal proper-time a spacetime mass distribution  $\mu$  needs to be moved to a mass distribution  $\nu$  in a future-directed, causal manner. Moreover,  $\ell_p$  satisfies the reverse triangle inequality (2.8).

The convexity properties making up our main definitions are understood along *timelike proper-time parametrized  $\ell_p$ -geodesics* in  $\mathcal{P}(M)$ , axiomatized as now described in Subsection 2.2.4, Subsection 2.3.6, and Appendix B. A timelike geodesic on  $M$  is a curve  $\gamma \in C([0, 1]; M)$  which is  $d$ -Lipschitz continuous [sic], obeys  $\gamma_s \ll \gamma_t$  for every  $s, t \in [0, 1]$  with  $s < t$ , and maximizes the  $\tau$ -length (2.5), symbolically  $\gamma \in \text{TGeo}(M)$ . As for maximizing geodesics in metric geometry [19], every timelike geodesic has a proper-time reparametrization  $\eta \in C([0, 1]; M)$ , i.e.

$$\tau(\eta_s, \eta_t) = (t - s) \tau(\eta_0, \eta_1) > 0 \quad (1.2)$$

for every  $s, t \in [0, 1]$  with  $s < t$  [58]. The inherent reparametrization procedure induces a map  $r: \text{TGeo}(M) \rightarrow C([0, 1]; M)$  which is continuous, cf. Lemma B.4. Given any  $\mu_0, \mu_1 \in \mathcal{P}(M)$ , let  $\text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$  be the set of all  $\pi \in \mathcal{P}(\text{TGeo}(M))$  such that  $\pi := (\mathbf{e}_0, \mathbf{e}_1)_{\#} \pi$  — where  $\mathbf{e}_t$  is the usual evaluation map at  $t \in [0, 1]$  — is a chronological  $\ell_p$ -optimal coupling of  $\mu_0$  and  $\mu_1$ . (Our geodesy hypothesis from Assumption 3.1 below ensures that every  $x, y \in M$  with  $x \ll y$  can be connected by an element of  $\text{TGeo}(M)$ .) Set  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1) := r_{\#} \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$ , and observe that all measures belonging to this set are concentrated on  $\text{TGeo}^{\tau}(M) := r(\text{TGeo}(M))$  whose elements, in turn, obey (1.2) for every  $s, t \in [0, 1]$  with  $s < t$ .

**Definition 1.1.** *A weakly continuous curve  $(\mu_t)_{t \in [0, 1]}$  in  $\mathcal{P}(M)$  is called*

- a. *timelike  $\ell_p$ -geodesic if there exists a measure  $\pi \in \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$  such that  $\mu_t = (\mathbf{e}_t)_{\#} \pi$  for every  $t \in [0, 1]$ , and*
- b. *timelike proper-time parametrized  $\ell_p$ -geodesic provided there is a measure  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that  $\mu_t = (\mathbf{e}_t)_{\#} \pi$  for every  $t \in [0, 1]$ .*

We point out that our definition of a bare  $\ell_p$ -geodesic differs from [66]. Indeed, every timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  satisfies

$$\ell_p(\mu_s, \mu_t) = (t - s) \ell_p(\mu_0, \mu_1) > 0 \quad (1.3)$$

for every  $s, t \in [0, 1]$  with  $s < t$  and is thus an  $\ell_p$ -geodesic as in [66]. On the other hand, timelike  $\ell_p$ -geodesics in our sense are implicit maximizers of some length functional, cf. Proposition B.3. Thus, Definition 1.1 reflects how geodesics are set up at the base space terminologically and conceptually [58]. In general, it is more restrictive, as [66] does not even assume weak continuity of the considered paths in  $\mathcal{P}(M)$ . (It is remarkable that this regularity is implied solely by (1.3) in all relevant smooth cases, cf. Remark B.8.) Our approach seems right for  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  to admit good compactness properties, cf. Proposition B.9. Indeed, proper-time reparametrizations of timelike geodesics in  $M$  are *not* Lipschitz continuous [58], which makes it difficult to find compact subsets of  $C([0, 1]; M)$  satisfying (1.2). On the other hand, by Assumption 3.1 — more precisely, a property termed *non-total imprisonment* — all elements of  $\text{TGeo}(M)$  passing through a fixed causal diamond have uniformly bounded Lipschitz constants. This entails excellent compactness properties of  $\text{TGeo}(M)$  and  $\text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$  which transfer to  $\text{TGeo}^{\tau}(M)$  and  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  by continuity of  $r$  and  $r_{\#}$ , respectively.

Lastly, the basic object which will be hypothesized to have some convexity along the named curves in  $\mathcal{P}(M)$  is the  $N$ -Rényi entropy

$$\mathfrak{S}_N(\mu) := - \int_M \rho^{-1/N} d\mu = - \int_M \rho^{1-1/N} d\mathbf{m}, \quad (1.4)$$

where  $\mu = \rho \mathbf{m} + \mu_\perp$  is the Lebesgue decomposition of  $\mu \in \mathcal{P}(M)$  with respect to  $\mathbf{m}$ . Its properties are summarized in Subsection 2.3.7. It only relates to  $\text{Ent}_{\mathbf{m}}$  in the limit case  $N \rightarrow \infty$ , see (2.10).

**Contributions.** Now we come to the communication of our novelties. The respective advantages of our different conditions are outlined along the way.

► Given  $p \in (0, 1)$ , in Definition 3.2 and Definition 3.3 we introduce the subsequent synthetic notions of the timelike Ricci curvature of  $(M, \mathbf{d}, \mathbf{m}, \ll, \leq, \tau)$  being bounded from below by  $K \in \mathbf{R}$ , and its dimension being bounded from above by  $N \in [1, \infty)$ , respectively:

- the reduced timelike curvature-dimension condition  $\text{TCD}_p^*(K, N)$ ,
- the weak reduced timelike curvature-dimension condition  $\text{wTCD}_p^*(K, N)$ ,
- the timelike curvature-dimension condition  $\text{TCD}_p(K, N)$ , and
- the weak timelike curvature-dimension condition  $\text{wTCD}_p(K, N)$ .

These are Lorentzian versions of [9, 86] and are all formulated via convexity properties of the Rényi entropy along timelike proper-time parametrized  $\ell_p$ -geodesics  $(\mu_t)_{t \in [0, 1]}$  between pairs  $(\mu_0, \mu_1) \in \mathcal{P}(M)^2$  of compactly supported,  $\mathbf{m}$ -absolutely continuous mass distributions, symbolically  $\mu_0, \mu_1 \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$ . The inherent inequalities defining  $\text{TCD}_p^*(K, N)$  and  $\text{wTCD}_p^*(K, N)$  use the distortion coefficients  $\sigma_{K, N}^{(r)}$ , while  $\text{TCD}_p(K, N)$  and  $\text{wTCD}_p(K, N)$  employ the quantities  $\tau_{K, N}^{(r)}$ ; we refer to Definition 2.1 below. The strong versions require the relevant estimate for Rényi's entropy to hold for all *timelike  $p$ -dualizable* endpoint pairs  $(\mu_0, \mu_1)$ , meaning that there exists an appropriate chronological  $\ell_p$ -optimal coupling  $\pi \in \Pi(\mu_0, \mu_1)$ . The weak versions require  $(\mu_0, \mu_1)$  to be *strongly timelike  $p$ -dualizable*, which essentially asks *all*  $\ell_p$ -optimal couplings of  $\mu_0$  and  $\mu_1$  to be chronological. The latter notions have been introduced in [24], see Definition 2.11 for their precise definition.

In Proposition 3.6, we establish the general relations

$$\begin{aligned} \text{TCD}_p(K, N) &\implies \text{TCD}_p^*(K, N), \\ \text{wTCD}_p(K, N) &\implies \text{wTCD}_p^*(K, N), \\ \text{TCD}_p^*(K, N) &\implies \text{TCD}_p(K^*, N), \\ \text{wTCD}_p^*(K, N) &\implies \text{wTCD}_p(K^*, N) \end{aligned} \quad (1.5)$$

between these conditions, where  $K^* := K(N - 1)/N$ , and assuming  $K > 0$  in the latter two cases.

► Our first major result is the compatibility of the above notions with timelike lower Ricci curvature bounds in the smooth framework, cf. Theorem A.1. In fact, we also cover the weighted case which has recently lead to singularity and splitting theorems for so-called Bakry–Émery spacetimes [94].

**Theorem 1.2.** *Let  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [n, \infty)$ , where  $n \in \mathbf{N}_{\geq 2}$ . For the Lorentzian space induced by a smooth,  $n$ -dimensional Lorentzian spacetime  $M$  and measured by the Lorentzian volume measure, any of  $\text{TCD}_p^*(K, N)$ ,  $\text{wTCD}_p^*(K, N)$ ,  $\text{TCD}_p(K, N)$ , and  $\text{wTCD}_p(K, N)$  holds if and only if the Ricci curvature of  $M$  is bounded from below by  $K$  in all timelike directions of the tangent bundle  $TM$ .*

In particular, for  $K = 0$  this gives a new characterization of the strong energy condition à la Hawking and Penrose, paralleling [66], and a new description of one inequality in the Einstein equations [69].

► The conditions  $\text{wTCD}_p(K, N)$  and  $\text{wTCD}_p^*(K, N)$  are proven to imply important timelike geometric estimates à la

- Brunn–Minkowski, see Proposition 3.11 and Proposition 3.20,
- Bonnet–Myers, see Corollary 3.14 and Corollary 3.21, as well as
- Bishop–Gromov, see Theorem 3.16 and Theorem 3.22.

Here the key advantage of  $\text{wTCD}_p(K, N)$  is that it yields these estimates in *sharp* form. Indeed, for a continuous, globally hyperbolic,  $n$ -dimensional spacetime  $M$ ,  $n \in \mathbf{N}_{\geq 2}$ , which is additionally causally plain [27] — our running example of a Lorentzian space of low regularity satisfying Assumption 3.1 below, cf. Example 2.7 —, the timelike Bishop–Gromov inequality obtained from  $\text{wTCD}_p(K, N)$  yields  $n = \dim^\tau M \leq N$  for the so-called geometric dimension  $\dim^\tau M$  recently introduced in [67], see Corollary 3.18. As inferred in Remark 3.19, the same reasoning starting from  $\text{wTCD}_p^*(K, N)$  only gives  $n = \dim^\tau M \leq N + 1$ ; compare with (1.5). Corresponding geometric inequalities have been obtained from the weak entropic timelike curvature-dimension condition  $\text{wTCD}_p^e(K, N)$  in [24] as well, but do not yield sharp versions either (unless timelike nonbranching is assumed, which we avoid).

► Another important general feature of our timelike curvature-dimension conditions is their weak stability according to a natural Lorentzian notion of convergence introduced in [24] following [45], concretized in Theorem 3.29.

**Theorem 1.3.** *Let  $(M_k, \mathbf{d}_k, \mathbf{m}_k, \ll_k, \leq_k, \tau_k)_{k \in \mathbf{N}}$  be a sequence of measured Lorentzian spaces obeying  $\text{TCD}_p^*(K, N)$  for fixed  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$  which converges to a measured Lorentzian space  $(M_\infty, \mathbf{d}_\infty, \mathbf{m}_\infty, \ll_\infty, \leq_\infty, \tau_\infty)$  in the sense of Definition 3.24. Then the latter satisfies  $\text{wTCD}_p^*(K, N)$ .*

*The same conclusion holds true when replacing  $\text{TCD}_p^*(K, N)$  and  $\text{wTCD}_p^*(K, N)$  by  $\text{TCD}_p(K, N)$  and  $\text{wTCD}_p(K, N)$ , respectively.*

Hence, limits according to Definition 3.24 of sequences of smooth Lorentzian spacetimes satisfying the hypothesis of Theorem 1.2 are  $\text{wTCD}_p(K, N)$ .

Theorem 1.3 is where the a priori distinction between weak and strong versions of our timelike curvature-dimension conditions, as in its entropic counterpart from [24], comes into play. There are essentially two technical reasons which arise from the proof of Theorem 3.29. First, given a strongly timelike  $p$ -dualizable pair in  $\mathcal{P}_c^{\text{ac}}(M_\infty, \mathbf{m}_\infty)^2$ , the members  $\mu_{k,0}, \mu_{k,1} \in \mathcal{P}_c^{\text{ac}}(M_k, \mathbf{m}_k)$ ,  $k \in \mathbf{N}$ , of corresponding recovery sequences between which  $\text{TCD}_p^*(K, N)$  applies can only be constructed to be timelike  $p$ -dualizable in general. Second, in the passage as  $k \rightarrow \infty$ , our timelike curvature-dimension condition requires the involved limit objects (namely, timelike proper-time parametrized  $\ell_p$ -geodesics and  $\ell_p$ -optimal couplings), obtained by a compactness argument, to be chronological. However, unlike causality, chronology of either objects is not stable under weak limits in general, unless we know that every  $\ell_p$ -optimal coupling of the limit marginals is chronological.

► In [24], the most powerful consequences of  $\text{TCD}_p^e(K, N)$  — e.g. uniqueness of  $\ell_p$ -optimal couplings and  $\ell_p$ -geodesics, localization, and singularity theorems — were obtained under the assumption of *timelike nonbranching*. Roughly speaking, the latter means that elements of  $\text{TGeo}^\tau(M)$  do not admit forward or backward branching at intermediate times. A drawback of this definition is that it is expected to fail e.g. for spacetimes with regularity less than  $C^{1,1}$  [24]. Hence, it may well be lost in limit cases, e.g. in the framework of Theorem 1.3.

Thus, we newly introduce the following notion called *timelike  $p$ -essential nonbranching*. See Definition 2.16 for the precise definition.

**Definition 1.4.** *Given  $p \in (0, 1]$ , we term the space  $(M, \mathbf{d}, \ll, \leq, \tau)$  timelike  $p$ -essentially nonbranching if every element of  $\text{OptTGeo}_{\ell_p}^{\tau}(\mathcal{P}(M)^2)$  is concentrated on a timelike nonbranching subset of  $\text{TGeo}^{\tau}(M)$ .*

Every timelike nonbranching Lorentzian pre-length space is timelike  $p$ -essentially nonbranching for every  $p \in (0, 1]$ , cf. Remark 2.17.

Definition 1.4 is based on the notion of *essentially nonbranching* metric measure spaces [77]. In [77], every RCD space was proven to be essentially nonbranching. Since the RCD condition is stable with respect to measured Gromov–Hausdorff (mGH) convergence [45, 85], this implicitly gives a stability result for the essential nonbranching condition. For this reason and the suspected issues for low regularity spacetimes, we believe Definition 1.4 to be relevant for the setting of nonsmooth general relativity as well, given that a suitable Lorentzian version of infinitesimal Hilbertianity [5, 6, 43] is found.

Here, we obtain a local-to-global property for the  $\text{TCD}_p^*(K, N)$  condition under timelike  $p$ -essential nonbranching, cf. Theorem 3.45. This constitutes a Lorentzian variant of [9] and is a feature of the reduced timelike curvature-dimension condition; indeed, as detailed in Remark 3.50, such a property seems subtle for  $\text{TCD}_p(K, N)$ . Furthermore, after [32, Thm. 3.14] we expect such a local-to-global property for  $\text{TCD}_p^e(K, N)$  to hold *a priori* only in a strong form.

► Another cornerstone result of our paper is the equivalence proof of *all* our reduced timelike curvature-dimension conditions with *all* entropic ones from [24] under timelike  $p$ -essential nonbranching as follows, see Theorem 3.35. Remarkably, none of these conditions is *a priori* weaker than the other.

**Theorem 1.5.** *Assume  $(M, \mathbf{d}, \mathbf{m}, \ll, \leq, \tau)$  to be timelike  $p$ -essentially nonbranching for some  $p \in (0, 1)$ . Then the following are equivalent.*

- (i) *The condition  $\text{TCD}_p^*(K, N)$  holds.*
- (ii) *The condition  $\text{wTCD}_p^*(K, N)$  holds.*
- (iii) *The condition  $\text{TCD}_p^e(K, N)$  holds.*
- (iv) *The condition  $\text{wTCD}_p^e(K, N)$  holds.*
- (v) *For every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})^2$ , there exists some  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , the measure  $(\mathbf{e}_t)_\# \pi$  is  $\mathbf{m}$ -absolutely continuous, and its density  $\rho_t$  with respect to  $\mathbf{m}$  obeys, for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}$$

for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^{\tau}(M)$ .

Even for the timelike nonbranching case and apart from the smooth case [66, 69], the equivalence of  $\text{TCD}_p^e(K, N)$  and  $\text{wTCD}_p^e(K, N)$  has not been proven in [24]. In fact, we establish both inherent equivalences of  $\text{TCD}_p^*(K, N)$  and  $\text{wTCD}_p^*(K, N)$  as well as  $\text{TCD}_p^e(K, N)$  and  $\text{wTCD}_p^e(K, N)$  only through their respective one-to-one correspondence with (v). The latter point constitutes a *pathwise* form of our reduced timelike curvature-dimension conditions. Contrary to the entropic case, such an interpretation being *a priori* meaningful — even when timelike branching occurs — thus makes our notions especially geometrically valuable. Moreover, point (v) provides density bounds for interpolating timelike proper-time parametrized  $\ell_p$ -geodesics which, under  $\text{wTCD}_p^e(K, N)$ , were obtained in [16] using variational methods. (In [16], we have not assumed timelike essential nonbranching, though.) Compare with Section 3.4 and Section 4.3 below, where we use similar methods to construct “good geodesics”.

An analogous pathwise form of the condition  $\text{TCD}_p(K, N)$  and  $\text{wTCD}_p(K, N)$  — however, with no direct link to [24] — is established in Theorem 3.41.

- Lastly, we introduce the synthetic notions of
- the reduced timelike measure-contraction property  $\text{TMCP}_p^*(K, N)$ , and
  - the timelike measure-contraction property  $\text{TMCP}_p(K, N)$ .

These are versions of the respective timelike curvature-dimension conditions where convexity of the Rényi entropy is not required along the chronological  $\ell_p$ -optimal transport between two  $\mathfrak{m}$ -absolutely continuous measures, but rather from an  $\mathfrak{m}$ -absolutely continuous distribution to a Dirac mass. For metric measure spaces, the corresponding notion was introduced in [70, 86]; the entropic Lorentzian version called  $\text{TMCP}_p^e(K, N)$  is due to [24].

In Proposition 4.4 and Proposition 4.8 we prove the following correspondences, where  $K^* := K(N - 1)/N$  is defined as for (1.5):

$$\begin{aligned} \text{TMCP}_p(K, N) &\implies \text{TMCP}_p^*(K, N), \\ \text{TMCP}_p^*(K, N) &\implies \text{TMCP}_p(K^*, N), \\ \text{wTCD}_p^*(K, N) &\implies \text{TMCP}_p^*(K, N), \\ \text{wTCD}_p(K, N) &\implies \text{TMCP}_p(K, N), \end{aligned}$$

again assuming  $K > 0$  in the latter two cases. Furthermore, we extend the above mentioned geometric inequalities, cf. Remark 4.9, we prove the full stability of both properties under the notion of convergence from Definition 3.24, cf. Theorem 4.10, and we show the equivalence of  $\text{TMCP}_p^*(K, N)$  with  $\text{TMCP}_p^e(K, N)$  under timelike  $p$ -essential nonbranching, cf. Theorem 4.20. We also provide pathwise characterizations of  $\text{TMCP}_p^*(K, N)$  and  $\text{TMCP}_p(K, N)$ , see Theorem 4.20 and Theorem 4.21.

Lastly, in Theorem 4.16 and Theorem 4.17 we establish the uniqueness of chronological  $\ell_p$ -optimal couplings as well as timelike proper-time parametrized  $\ell_p$ -optimal geodesics for large classes of initial measures, again under timelike essential nonbranching. (In fact, this was our main motivation to introduce the named weaker timelike measure-contraction properties.) This extends the smooth outcomes from [66]. Our reasoning follows [24]; in fact, we extend the corresponding uniqueness results obtained therein under  $\text{TMCP}_p^e(K, N)$  from the timelike nonbranching to the timelike essentially nonbranching case, cf. Remark 4.18. However, let us stress that even in the timelike nonbranching case, our uniqueness results do *not* follow by combining Theorem 1.5 and [24]. Rather, Theorem 4.16 and Theorem 4.17 are *prerequisites* for the proof of Theorem 1.5, cf. Remark 3.37.

**Outlook.** Finally, we outline various possible directions the notions and results of our work could lead to in the future.

From the viewpoint of general relativity, it is natural to ask whether singularity theorems à la Hawking and Penrose, such as those from [24] under  $\text{TMCP}_p^e(K, N)$  and the assumption of timelike nonbranching, hold in our setting as well, possibly under the weaker hypothesis of timelike essential nonbranching. It is also worth seeking for nonsmooth analogues of the above mentioned singularity theorems à la Gannon and Lee. Note that the universal cover of a spacetime appears in regular formulations of these, cf. [80, Thm. 2.6] for the most recent result in this direction for  $C^1$ -metrics. To this aim, in Theorem 3.55 we make a first step by showing that  $\text{TCD}_p^*(K, N)$  lifts to the universal cover of a measured Lorentzian space, in case of its existence. Coupled with an extension of the Lorentzian localization technique, initiated in [24], to the timelike essentially nonbranching case, this will be the goal of our forthcoming work [17].

Second, our work is a further contribution to our long-term goal already outlined in [16], namely to obtain a splitting theorem for measured Lorentzian spaces. In the corresponding result for metric measure spaces [42], Rényi's entropy plays a crucial role, and we expect a similar importance of it in our relativistic framework.



As mentioned, what is still missing to date in order to get started is a Lorentzian notion of infinitesimal Hilbertianity.

Analogously to [18, 68], such a splitting theorem could then provide insights into the structure of TCD spaces, e.g. giving that they look like Minkowski space  $\mathbf{m}$ -a.e.

We learned from (1.5) that  $\text{TCD}_p^*(K, N)$  is a priori weaker than the condition  $\text{TCD}_p(K, N)$ . Recently, for essentially nonbranching metric measure spaces, Cavalletti and Milman [22] proved the *equivalence* of  $\text{CD}^*(K, N)$  [9] and  $\text{CD}(K, N)$  [86] by using sophisticated localization tools. The analogous step from  $\text{TCD}_p^*(K, N)$  to  $\text{TCD}_p(K, N)$ , possibly under timelike  $p$ -essential nonbranching, together with Theorem 1.5 would close the circle in (1.5) and show the equivalence of all (finite-dimensional) TCD conditions known thus far.

A further possible direction is to compare different notions of convergence for measured Lorentzian spaces, different from Definition 3.24, and to establish their equivalence in analogy to [45] for metric measure spaces. Our proof of Theorem 1.3 suggests a Lorentzian adaptation of Sturm's  $\mathbf{D}$ -convergence [86] to be potentially interesting, see Remark 3.31 (and also Remark 3.30). In general, finding a good definition of convergence of Lorentzian spaces is a central question in relativity theory; for instance, what it should provide is a Lorentzian precompactness theorem à la Gromov [48], which is missing thus far. In particular, it remains open whether any of the classes of spaces obeying  $\text{TCD}_p^e(K, N)$ ,  $\text{TCD}_p^*(K, N)$ , or  $\text{TCD}_p(K, N)$  (or suitable subsets thereof) are compact in any reasonable topology.

Lastly, based on a similar question asked in [25] and the results from [1], it is natural to expect that  $\text{TCD}_p(K, N)$  and  $\text{TCD}_q(K, N)$  are equivalent if the space  $(M, \mathbf{d}, \ll, \leq, \tau)$  is timelike  $p$ - and  $q$ -essentially nonbranching,  $p, q \in (0, 1)$ . Conceptually, this would be important in Lorentzian geometry where, unlike the metric setting [61, 85, 86], there is a priori no exponent  $p \in (0, 1)$  which seems most natural to ground the theory of synthetic timelike Ricci curvature bounds on.

**Organization.** In Chapter 2, we collect all necessary background material about measured Lorentzian spaces and optimal transport theory on these. Chapter 3 is devoted to a detailed investigation of our timelike curvature-dimension conditions. Chapter 4 studies the timelike measure-contraction property. Appendix A contains the smooth proof of Theorem 1.2, and Appendix B collects some technical results evolving around Definition 1.1.

## 2. PRELIMINARIES

**2.1. Volume distortion coefficients.** We start by introducing the *volume distortion coefficients* that make up our main definitions, and by listing their basic properties. We refer to [9, 32, 86] for details.

Given  $\kappa \in \mathbf{R}$ ,  $t \in [0, 1]$ , and  $\vartheta \geq 0$ , we define the quantities

$$\mathfrak{s}_\kappa(\vartheta) := \begin{cases} \frac{\sin(\sqrt{\kappa} \vartheta)}{\sqrt{\kappa}} & \text{if } \kappa > 0, \\ \vartheta & \text{if } \kappa = 0, \\ \frac{\sinh(\sqrt{-\kappa} \vartheta)}{\sqrt{-\kappa}} & \text{otherwise,} \end{cases} \quad (2.1)$$

$$\sigma_\kappa^{(t)}(\vartheta) := \begin{cases} \infty & \text{if } \kappa \vartheta^2 \geq \pi^2, \\ t & \text{if } \kappa \vartheta^2 = 0, \\ \frac{\mathfrak{s}_\kappa(t \vartheta)}{\mathfrak{s}_\kappa(\vartheta)} & \text{otherwise,} \end{cases}$$

**Definition 2.1.** For  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ , slightly abusing notation we set

$$\begin{aligned}\sigma_{K,N}^{(t)}(\vartheta) &:= \sigma_{K/N}^{(t)}(\vartheta), \\ \tau_{K,N}^{(t)}(\vartheta) &:= t^{1/N} \sigma_{K,N-1}^{(t)}(\vartheta)^{1-1/N}.\end{aligned}$$

Note that for every  $t \in (0, 1)$  and every  $\vartheta > 0$ ,  $\sigma_{K,N}^{(t)}(\vartheta)$  is continuous in  $(K, N) \in \mathbf{R} \times [1, \infty)$ , nondecreasing in  $K$ , and nonincreasing in  $N$  [9, Rem. 2.2]. Analogous claims apply to the quantity  $\tau_{K,N}^{(t)}(\vartheta)$  [86, p. 138]. Furthermore, for every  $t \in [0, 1]$ , every  $\vartheta \geq 0$ , and every  $\kappa \in (-\infty, \pi^2/\vartheta^2)$ ,

$$\sigma_{\kappa}^{(t)}(\vartheta) = \sigma_{\kappa\vartheta^2}^{(t)}(1). \quad (2.2)$$

*Remark 2.2.* We recall the following elementary inequality from [23, Rem. 2.3]: for every  $K \in \mathbf{R}$ , every  $N \in [1, \infty)$ , every  $t \in [0, 1]$ , and every  $\vartheta \geq 0$ ,

$$\sigma_{K,N}^{(t)}(\vartheta) \geq t e^{-(1-t)\vartheta\sqrt{K^-/N}}.$$

Lastly, in view of Theorem 3.35 and Theorem 4.20, given any  $t \in [0, 1]$  let us define  $G_t: \mathbf{R}^2 \times (-\infty, \pi^2) \rightarrow (-\infty, \infty]$  and  $H_t: \mathbf{R} \times (-\infty, \pi^2) \rightarrow (-\infty, \infty]$  by

$$\begin{aligned}G_t(x, y, \kappa) &:= \log[\sigma_{\kappa}^{(1-t)}(1) e^x + \sigma_{\kappa}^{(t)}(1) e^y] \\ H_t(x, \kappa) &:= \log[\sigma_{\kappa}^{(1-t)}(1) e^x] = \log \sigma_{\kappa}^{(1-t)}(1) + x.\end{aligned} \quad (2.3)$$

Then the functions  $G_t$  and  $H_t$  are jointly convex [32, Lem. 2.11].

**2.2. Nonsmooth Lorentzian spaces.** We continue with a concise digression on the theory of nonsmooth Lorentzian (pre-length, length, and geodesic) spaces. We refer to [24, 58] for details, proofs, and examples about the corresponding notions.

**2.2.1. Chronology and causality.** Let us fix a preorder  $\leq$  and a transitive relation  $\ll$ , contained in  $\leq$ , on  $M$ . The triple  $(M, \ll, \leq)$  is called *causal space* [58, Def. 2.1]. We say that  $x, y \in M$  are *timelike* or *causally* related if  $x \ll y$  or  $x \leq y$ , respectively. We write  $x < y$  whenever  $x \leq y$  yet  $x \neq y$ . Define the sets  $M_{\ll}^2$  and  $M_{\leq}^2$  to consist of all pairs  $(x, y) \in M^2$  with  $x \ll y$  and  $x \leq y$ , respectively.

**Definition 2.3.** We term the triple  $(M, \ll, \leq)$

- causally closed if the relation  $\leq$  is closed, i.e.  $M_{\leq}^2$  is closed in  $M^2$ , and
- locally causally closed if every point in  $M$  has some open neighborhood  $U \subset M$  such that  $\bar{U}^2 \cap M_{\leq}^2$  is relatively closed in  $\bar{U}^2$ .

Given any  $A \subset M$ , we define the *chronological future*  $I^+(A) \subset M$  and the *causal future*  $J^+(A) \subset M$  of  $A$  [58, Def. 2.3] through

$$\begin{aligned}I^+(A) &:= \{y \in M : x \ll y \text{ for some } x \in A\}, \\ J^+(A) &:= \{y \in M : x \leq y \text{ for some } x \in A\}.\end{aligned}$$

Analogously, we define the *chronological past*  $I^-(A)$  and the *causal past*  $J^-(A)$  of  $A$ . Employing a slight abuse of notation, given  $x \in M$  we write  $I^{\pm}(x) := I^{\pm}(\{x\})$  and  $J^{\pm}(x) := J^{\pm}(\{x\})$ . For a Borel measure  $\mu$  on  $M$ , we shall write  $I^{\pm}(\mu) := I^{\pm}(\text{spt } \mu)$  and  $J^{\pm}(\mu) := J^{\pm}(\text{spt } \mu)$ . For all these objects, set  $I(A, B) := I^+(A) \cap I^-(B)$ , and define  $I(x, y)$ ,  $I(\mu, \nu)$ ,  $J(A, B)$ ,  $J(x, y)$ , and  $J(\mu, \nu)$  in an analogous way.

**2.2.2. Lorentzian pre-length spaces.** A function  $\tau: M^2 \rightarrow [0, \infty]$  is called *time separation function* [58, Def. 2.8] if it is lower semicontinuous, and for every  $x, y, z \in M$  it satisfies the properties

- $\tau(x, y) = 0$  if  $x \not\leq y$ ,
- $\tau(x, y) > 0$  if and only if  $x \ll y$ , and

c. if  $x \leq y \leq z$  we have the *reverse triangle inequality*

$$\tau(x, z) \geq \tau(x, y) + \tau(y, z). \quad (2.4)$$

The existence of such a  $\tau$  implies that  $\ll$  is an *open relation* [58, Prop. 2.13]; in particular, the set  $I^\pm(A)$  is open for every  $A \subset M$  [58, Lem. 2.12].

**Definition 2.4.** A Lorentzian pre-length space is a quintuple  $(M, \mathbf{d}, \ll, \leq, \tau)$  which consists of a causal space  $(M, \ll, \leq)$  endowed with a metric  $\mathbf{d}$  and a time separation function  $\tau$  as introduced above.

**2.2.3. Length of curves.** Let  $(M, \mathbf{d}, \ll, \leq, \tau)$  be a given Lorentzian pre-length space. A *path* designates a map  $\gamma: [a, b] \rightarrow M$ , where  $a, b \in \mathbf{R}$  with  $a < b$ ; if  $\gamma$  is continuous, we call  $\gamma$  a *curve*. Such a curve  $\gamma$  is (*future-directed*) *timelike* or (*future-directed*) *causal* if it is  $\mathbf{d}$ -Lipschitz continuous [sic], and  $\gamma_s \ll \gamma_t$  or  $\gamma_s \leq \gamma_t$  for every  $s, t \in [a, b]$  with  $s < t$ , respectively. It is *null* if it is causal and  $\gamma_a \not\ll \gamma_b$ . Similar notions evidently make sense for *past-directed* curves and their *causal character* (that is, their property of being chronological, causal, or null). Unless explicitly stated otherwise, any curve of a specified causal character is assumed future-directed.

The *length* with respect to  $\tau$  of a path  $\gamma: [a, b] \rightarrow M$  is

$$\text{Len}_\tau(\gamma) := \inf\{\tau(\gamma_{t_0}, \gamma_{t_1}) + \cdots + \tau(\gamma_{t_{n-1}}, \gamma_{t_n})\}, \quad (2.5)$$

where the infimum is taken over all integers  $n \in \mathbf{N}$  and all times  $t_0, \dots, t_n \in [a, b]$  with  $t_0 = a$ ,  $t_n = b$ , and  $t_i < t_{i+1}$  for every  $i \in \{0, \dots, n-1\}$  [58, Def. 2.24]. It is additive [58, Lem. 2.25], and reparametrizations do neither change causal characters [58, Lem. 2.27] nor the  $\tau$ -length itself [58, Lem. 2.28].

**2.2.4. Geodesics.** A causal curve  $\gamma: [0, 1] \rightarrow M$  is called *geodesic* (or *maximal*) provided  $\text{Len}_\tau(\gamma) = \tau(\gamma_0, \gamma_1)$  [58, Def. 2.33]. The space of all such curves is denoted  $\text{Geo}(M)$ , and its subset of timelike curves is written  $\text{TGeo}(M)$ .

**Definition 2.5.** We call  $(M, \mathbf{d}, \ll, \leq, \tau)$  *geodesic* if for every  $x, y \in M$  with  $x < y$ , there exists a geodesic  $\gamma \in \text{Geo}(M)$  starting at  $x$  and ending at  $y$ .

Every *timelike* geodesic has a weak parametrization [58, Def. 3.31] by  $\tau$ -arclength provided  $\tau$  is continuous as well as  $\tau(x, x) = 0$  for every  $x \in M$  [58, Cor. 3.35]. More precisely, given  $\gamma \in \text{TGeo}(M)$  define  $\psi_\gamma \in C([0, 1]; [0, 1])$  by

$$\psi_\gamma(t) := \frac{\text{Len}_\tau(\gamma|_{[0, t]})}{\tau(\gamma_0, \gamma_1)} = \frac{\tau(\gamma_0, \gamma_t)}{\tau(\gamma_0, \gamma_1)}.$$

Consider the *reparametrization map*  $r: \text{TGeo}(M) \rightarrow C([0, 1]; M)$  given by

$$r(\gamma)_t := \gamma_{\psi_\gamma^{-1}(t)};$$

see also Appendix B. Then for every  $\gamma \in \text{TGeo}(M)$ , the curve  $\eta := r(\gamma)$  obeys

$$\tau(\eta_s, \eta_t) = (t - s) \tau(\eta_0, \eta_1) > 0 \quad (2.6)$$

for every  $s, t \in [0, 1]$  with  $s < t$ . Note that elements of  $\text{TGeo}^\tau(M) := r(\text{TGeo}(M))$  might not be  $\mathbf{d}$ -Lipschitz continuous any more [58, p. 424]. On the other hand, as shown in Lemma B.4,  $r$  is continuous with respect to the uniform topology, which will be important for our axiomatization of  $\ell_p$ -geodesics and the “transference” of compactness properties in  $\text{TGeo}(M)$  to  $\text{TGeo}^\tau(M)$ , see Subsection 2.3.6 below. In order to not interrupt the exposition, we have deferred corresponding definitions and results to Appendix B.

2.2.5. *Global hyperbolicity.* Following [24, Sec. 1.1], we term  $(M, d, \ll, \leq, \tau)$  *non-totally imprisoning* if for every compact  $C \subset M$  there exists a constant  $c > 0$  such that the  $d$ -arclength of any causal curve contained in  $C$  is no larger than  $c$ .

**Definition 2.6.** *The space  $(M, d, \ll, \leq, \tau)$  is*

- a. *globally hyperbolic if it is non-totally imprisoning and the causal diamond  $J(x, y)$  is compact for every  $x, y \in M$ , and*
- b.  *$\mathcal{K}$ -globally hyperbolic if it is non-totally imprisoning and the causal diamond  $J(C_0, C_1)$  is compact for all compact  $C_0, C_1 \subset M$ .*

We list some properties of ( $\mathcal{K}$ -)globally hyperbolic Lorentzian pre-length spaces which will lead to our standing Assumption 3.1 below as well as useful consequences thereof. If  $(M, d, \ll, \leq, \tau)$  is locally causally closed, globally hyperbolic, and  $I^\pm(x) \neq \emptyset$  for every  $x \in M$ , then  $\mathcal{K}$ -global hyperbolicity holds [24, Lem. 1.5]. On the other hand, every locally causally closed,  $\mathcal{K}$ -globally hyperbolic Lorentzian geodesic space is in fact causally closed [24, Lem. 1.6]. By [58, Def. 3.25, Thm. 3.26], global hyperbolicity implies the nonsmooth analogue of the *strong causality condition* for smooth Lorentzian spacetimes, cf. e.g. [71, Def. 14.11]. On every globally hyperbolic Lorentzian length space (see [58, Def. 3.22] for the corresponding definition, we will not really need it), the time separation function  $\tau$  is finite and continuous [58, Thm. 3.28]; in particular, we have  $\tau(x, x) = 0$  for every  $x \in M$ . Furthermore, every such space is geodesic by the nonsmooth analogue of Avez–Seifert’s theorem [58, Thm. 3.30]. In [21], a singular analogue of Geroch’s characterization [40] of global hyperbolicity via Cauchy time functions is proven.

*Example 2.7* (Spacetimes of low regularity). Let  $M$  be a smooth spacetime endowed with a continuous, globally hyperbolic, causally plain [27, Def. 1.16] metric  $\langle \cdot, \cdot \rangle$ . Causal plainness is implied if  $\langle \cdot, \cdot \rangle$  is of class  $C^{0,1}$  [27, Def. 1.16], which is the borderline regularity to pathological causal phenomena such as *bubbling* [27, Thm. 1.20]. In this setting,  $(M, \langle \cdot, \cdot \rangle)$  canonically induces a causally closed,  $\mathcal{K}$ -globally hyperbolic Lorentzian geodesic space, cf. [24, Rem. 1.12] and [58, Prop. 3.5, Thm. 3.30, Thm. 5.12], which will in particular satisfy our standing Assumption 3.1 below.

**2.3. Optimal transport on Lorentzian spaces.** Next, let us briefly review the theory of optimal transport on the class of spaces introduced above [24]. We refer to [31, 53, 66, 69, 87] for prior developments in the smooth case.

2.3.1. *Basic probabilistic notation.* Let  $\mathcal{P}(M)$  denote the set of all Borel probability measures on  $M$ . Let  $\mathcal{P}_c(M)$  and  $\mathcal{P}^{\text{ac}}(M, \mathfrak{m})$  be its subsets consisting of all compactly supported and  $\mathfrak{m}$ -absolutely continuous elements, respectively; set  $\mathcal{P}_c^{\text{ac}}(M, \mathfrak{m}) := \mathcal{P}_c(M) \cap \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$ . Given any  $\mu \in \mathcal{P}(M)$ , by  $\mu_\perp$  we mean the  $\mathfrak{m}$ -singular part in the corresponding Lebesgue decomposition of  $\mu \in \mathcal{P}(M)$ .

For a Borel map  $F: M \rightarrow M'$  into a metric space  $(M', d')$ , given any  $\mu \in \mathcal{P}(M)$  the measure  $F_\# \mu \in \mathcal{P}(M')$  designates the usual *push-forward* of  $\mu$  under  $F$ , defined by the formula  $F_\# \mu[B] := \mu[F^{-1}(B)]$  for every Borel set  $B \subset M'$ .

Given  $\mu, \nu \in \mathcal{P}(M)$ , let  $\Pi(\mu, \nu)$  denote the set of all *couplings* of  $\mu$  and  $\nu$ , i.e. all  $\pi \in \mathcal{P}(M^2)$  with  $\pi[\cdot \times M] = \mu$  and  $\pi[M \times \cdot] = \nu$ .

With  $C([0, 1]; M)$  denoting the set of all curves  $\gamma: [0, 1] \rightarrow M$ , endowed with the uniform topology, for  $t \in [0, 1]$  the so-called *evaluation map*  $e_t: C([0, 1]; M) \rightarrow M$  is defined through  $e_t(\gamma) := \gamma_t$ .

2.3.2. *Weak convergence.* For convenience, let us briefly recall basic notions and results about weak convergence of probability measures now.

A sequence  $(\mu_n)_{n \in \mathbf{N}}$  in  $\mathcal{P}(M)$  converges *weakly* to  $\mu \in \mathcal{P}(M)$  [13, p. 7] if for every bounded continuous function  $\varphi: M \rightarrow \mathbf{R}$ , symbolically  $\varphi \in C_b(M)$ ,

$$\lim_{n \rightarrow \infty} \int_M \varphi d\mu_n = \int_M \varphi d\mu.$$

Recall that weak convergence is induced by a metric [4, Rem. 5.1.1].

The following two classical results will be used frequently, e.g. in the proofs of Theorem 3.29 and Theorem 4.10. Theorem 2.8, often called *Portmanteau's theorem* [13, Thm. 2.1], relates weak convergence to “setwise convergence” of sequences of probability measures, whereas Theorem 2.9, the *Prokhorov theorem* [13, Thm. 5.1, Thm. 5.2], arms us with a very useful compactness criterion.

**Theorem 2.8.** *For every sequence  $(\mu_n)_{n \in \mathbf{N}}$  in  $\mathcal{P}(M)$  and every  $\mu \in \mathcal{P}(M)$ , the following are equivalent.*

- (i) **Weak convergence.** *The sequence  $(\mu_n)_{n \in \mathbf{N}}$  converges weakly to  $\mu$ .*
- (ii) **Upper semicontinuity on closed sets.** *For every closed  $C \subset M$ ,*

$$\limsup_{n \rightarrow \infty} \mu_n[C] \leq \mu[C].$$

- (iii) **Lower semicontinuity on open sets.** *For every open  $U \subset M$ ,*

$$\mu[U] \leq \liminf_{n \rightarrow \infty} \mu_n[U].$$

- (iv) **Continuity sets.** *For every Borel set  $E \subset M$  with  $\mu[\partial E] = 0$ ,*

$$\lim_{n \rightarrow \infty} \mu_n[E] = \mu[E].$$

**Theorem 2.9.** *Assume  $\mathbf{d}$  to be complete and separable. Then a set  $\mathcal{C} \subset \mathcal{P}(M)$  is relatively compact with respect to the weak topology if and only if it is tight, i.e. for every  $\varepsilon > 0$  there exists a compact set  $C \subset M$  with*

$$\sup\{\mu[C^c] : \mu \in \mathcal{C}\} \leq \varepsilon.$$

**2.3.3. Chronological and causal couplings.** Let  $(M, \mathbf{d}, \ll, \leq, \tau)$  be a Lorentzian pre-length space. We define the (possibly empty) set  $\Pi_{\ll}(\mu, \nu)$  of all *chronological couplings* of two probability measures  $\mu, \nu \in \mathcal{P}(M)$  to consist of all  $\pi \in \Pi(\mu, \nu)$  such that  $\pi[M_{\ll}^2] = 1$ . The set  $\Pi_{\leq}(\mu, \nu)$  of all *causal couplings* of  $\mu$  and  $\nu$  is defined analogously. Under causal closedness, clearly  $\pi \in \Pi_{\leq}(\mu, \nu)$  if and only if  $\pi \in \Pi(\mu, \nu)$  as well as  $\text{spt } \pi \subset M_{\leq}^2$ ; an analogous statement holds for the locally causally closed case if  $\mu, \nu \in \mathcal{P}_c(M)$ .

A chronological or causal coupling of  $\mu, \nu \in \mathcal{P}(M)$  intuitively describes a way of transporting an infinitesimal mass portion  $d\mu(x)$  to an infinitesimal mass portion  $d\nu(y)$  as to guarantee  $x \ll y$  or  $x \leq y$ , respectively.

We call  $\mu, \nu \in \mathcal{P}(M)$  *chronologically related* if  $\Pi_{\ll}(\mu, \nu) \neq \emptyset$ .

**2.3.4. The  $\ell_p$ -optimal transport problem.** Given an exponent  $p \in (0, 1]$  and following the expositions in [24, 66] we will adopt the conventions

$$\begin{aligned} \sup \emptyset &:= -\infty, \\ (-\infty)^p &:= (-\infty)^{1/p} := -\infty, \\ \infty - \infty &:= -\infty. \end{aligned}$$

Let the total transport cost function  $\ell_p: \mathcal{P}(M)^2 \rightarrow [0, \infty] \cup \{-\infty\}$  be given by

$$\begin{aligned} \ell_p(\mu, \nu) &:= \sup\{\|\tau\|_{L^p(M^2, \pi)} : \pi \in \Pi_{\leq}(\mu, \nu)\} \\ &= \sup\{\|\ell\|_{L^p(M^2, \pi)} : \pi \in \Pi(\mu, \nu)\}, \end{aligned} \tag{2.7}$$

where  $l: M^2 \rightarrow [0, \infty] \cup \{-\infty\}$  is defined through

$$l^p(x, y) := \begin{cases} \tau^p(x, y) & \text{if } x \leq y, \\ -\infty & \text{otherwise.} \end{cases}$$

*Remark 2.10.* The sets of maximizers for both suprema defining  $\ell_p(\mu, \nu)$  coincide, including the case  $\Pi_{\leq}(\mu, \nu) = \emptyset$ . One advantage of the formulation via  $l$  is that under (local) causal closedness and global hyperbolicity,  $l^p$  is upper semicontinuous. In this case, customary optimal transport techniques [3, 92] are applicable to study the second problem, which in turn yields corresponding results for the first [24, Rem. 2.2]. Note that the preimages  $l^{-1}([0, \infty))$  and  $l^{-1}((0, \infty))$  encode causality and chronology of points in  $M^2$ , respectively.

A coupling  $\pi \in \Pi(\mu, \nu)$  of  $\mu, \nu \in \mathcal{P}(M)$  is called  $\ell_p$ -optimal if  $\pi \in \Pi_{\leq}(\mu, \nu)$  and

$$\ell_p(\mu, \nu) = \|\tau\|_{L^p(M^2, \pi)} = \|l\|_{L^p(M^2, \pi)}.$$

For existence of such couplings, for our work it will suffice to keep the following in mind. If  $(M, \mathbf{d}, \ll, \leq, \tau)$  is locally causally closed and globally hyperbolic, and if  $\mu, \nu \in \mathcal{P}_c(M)$  with  $\Pi_{\leq}(\mu, \nu) \neq \emptyset$ , existence of an  $\ell_p$ -optimal coupling  $\pi$  of  $\mu$  and  $\nu$  holds, and  $\ell_p(\mu, \nu) < \infty$  [24, Prop. 2.3].

A key property of  $\ell_p$  is the *reverse triangle inequality* [24, Prop. 2.5] somewhat reminiscent of (2.4) and of  $l$  satisfying the reverse triangle inequality for *every*, i.e. not necessarily causally related,  $x, y, z \in M$ : for *every*  $\mu, \nu, \sigma \in \mathcal{P}(M)$ ,

$$\ell_p(\mu, \sigma) \geq \ell_p(\mu, \nu) + \ell_p(\nu, \sigma). \quad (2.8)$$

**2.3.5. Timelike  $p$ -dualizability.** The concept of so-called (*strong*) *timelike  $p$ -dualizability*, where  $p \in (0, 1]$ , of pairs  $(\mu, \nu) \in \mathcal{P}(M)^2$  originates in [24] and generalizes the notion of  *$p$ -separation* introduced in [66, Def. 4.1]. Pairs satisfying this condition admit good duality properties [24, Prop. 2.19, Prop. 2.21, Thm. 2.26], which leads to a characterization of  $\ell_p$ -geodesics (see Subsection 2.3.6 below) in the smooth case [66, Thm. 4.3, Thm. 5.8], cf. Theorem A.1.

In view of the following Definition 2.11, cf. [24, Def. 2.18, Def. 2.27], we refer to [24, Def. 2.6] for the definition of *cyclical monotonicity* of a set in  $M_{\leq}^2$  with respect to  $l^p$ , see also [92, Def. 5.1]. It will not be relevant in our work.

Given  $a, b: M \rightarrow \mathbf{R}$  we define  $a \oplus b: M^2 \rightarrow \mathbf{R}$  by  $(a \oplus b)(x, y) := a(x) + b(y)$ .

**Definition 2.11.** *Given any  $p \in (0, 1]$ , a pair  $(\mu, \nu) \in \mathcal{P}(M)$  is called*

- a. *timelike  $p$ -dualizable by  $\pi \in \Pi_{\ll}(\mu, \nu)$  if  $\pi$  is  $\ell_p$ -optimal, and there exist Borel functions  $a, b: M \rightarrow \mathbf{R}$  such that  $a \oplus b \in L^1(M^2, \mu \otimes \nu)$  as well as  $l^p \leq a \oplus b$  on  $\text{spt } \mu \times \text{spt } \nu$ ,*
- b. *strongly timelike  $p$ -dualizable by  $\pi \in \Pi_{\ll}(\mu, \nu)$  if the pair  $(\mu, \nu)$  is timelike  $p$ -dualizable by  $\pi$ , and there is an  $l^p$ -cyclically monotone Borel set  $\Gamma \subset M_{\leq}^2 \cap (\text{spt } \mu_0 \times \text{spt } \mu_1)$  such that every given  $\sigma \in \Pi_{\leq}(\mu_0, \mu_1)$  is  $\ell_p$ -optimal if and only if  $\sigma[\Gamma] = 1$  (in particular,  $\sigma$  is chronological), and*
- c. *timelike  $p$ -dualizable if  $(\mu, \nu)$  is timelike  $p$ -dualizable by some coupling  $\pi \in \Pi_{\ll}(\mu, \nu)$ ; analogously for strong timelike  $p$ -dualizability.*

*Moreover, any  $\pi$  as in the above items is called timelike  $p$ -dualizing.*

*Remark 2.12.* If  $\mu, \nu \in \mathcal{P}_c(M)$  then the pair  $(\mu, \nu)$  is timelike  $p$ -dualizable if and only if there is an  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\leq}(\mu, \nu)$  which is concentrated on  $M_{\leq}^2$ .

If  $\mu, \nu \in \mathcal{P}_c(M)$  on a causally closed, globally hyperbolic Lorentzian geodesic space  $(M, \mathbf{d}, \ll, \leq, \tau)$  with  $\text{spt } \mu \times \text{spt } \nu \subset M_{\leq}^2$ , then  $(\mu, \nu)$  is automatically strongly timelike  $p$ -dualizable,  $p \in (0, 1]$  [24, Cor. 2.29], see also [66, Lem. 4.4, Thm. 7.1].

**2.3.6. Geodesics revisited.** The definitions we consider need the notion of an  $\ell_p$ -geodesic,  $p \in (0, 1]$ , to be made precise. We propose Definition 2.13 as Lorentzian version of Wasserstein geodesics in metric spaces. Precise technical results will be deferred to Appendix B.

Solely in this subsection, for simplicity we assume compactness of  $M$ ; the results derived here and in Appendix B will mostly be used by replacing  $M$  by a causal diamond of two compact sets, which is compact under  $\mathcal{K}$ -global hyperbolicity.

Recall the continuous reparametrization map  $r: \text{TGeo}(M) \rightarrow \text{TGeo}^\tau(M)$  and the definition  $\text{TGeo}^\tau(M) := r(\text{TGeo}(M))$  from Subsection 2.2.4. If  $(M, d, \ll, \leq, \tau)$  is causally closed, the set  $\text{Geo}(M)$  is Polish [58, Prop. 3.17]. Since  $\text{TGeo}(M)$  is open,  $\text{TGeo}^\tau(M)$  is a Suslin set, hence universally measurable.

Given any  $\mu_0, \mu_1 \in \mathcal{P}(M)$ , we define

$$\text{OptGeo}_{\ell_p}(\mu_0, \mu_1) := \{\pi \in \mathcal{P}(\text{Geo}(M)) : (\mathbf{e}_0, \mathbf{e}_1)_\# \pi \in \Pi_{\leq}(\mu_0, \mu_1) \\ \text{is } \ell_p\text{-optimal}\},$$

$$\text{OptTGeo}_{\ell_p}(\mu_0, \mu_1) := \{\pi \in \text{OptGeo}_{\ell_p}(\mu_0, \mu_1) : (\mathbf{e}_0, \mathbf{e}_1)_\# \pi[M_{\ll}^2] = 1\},$$

$$\text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1) := r_\# \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1).$$

We say that a measure  $\pi \in \mathcal{P}(C([0, 1]; M))$  represents a curve  $(\mu_t)_{t \in [0, 1]}$  in  $\mathcal{P}(M)$  provided  $\mu_t = (\mathbf{e}_t)_\# \pi$  for every  $t \in [0, 1]$ .

**Definition 2.13.** We term a weakly continuous path  $(\mu_t)_{t \in [0, 1]}$  in  $\mathcal{P}(M)$

- causal  $\ell_p$ -geodesic if it is represented by some  $\pi \in \text{OptGeo}_{\ell_p}(\mu_0, \mu_1)$ ,
- timelike  $\ell_p$ -geodesic if it is represented by some  $\pi \in \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$ , and
- timelike proper-time parametrized  $\ell_p$ -geodesic if it is represented by some  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ .

A measure  $\pi$  as in the last item will be called  $\ell_p$ -optimal geodesic plan.

*Remark 2.14.* If  $(M, d, \ll, \leq, \tau)$  is  $\mathcal{K}$ -globally hyperbolic, every causal or timelike  $\ell_p$ -geodesic connecting compactly supported  $\mu_0$  and  $\mu_1$  is Lipschitz continuous with respect to the  $q$ -Wasserstein metric  $W_q$  induced by  $(M, d)$  for every  $q \in [1, \infty]$ .

By construction, we have  $\pi[\text{TGeo}(M)] = 1$  for every  $\pi \in \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$ . Consequently,  $\pi[\text{TGeo}^\tau(M)] = 1$  for every  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ , and therefore  $\pi$ -a.e.  $\gamma \in C([0, 1]; M)$  obeys  $\tau(\gamma_s, \gamma_t) = (t - s)\tau(\gamma_0, \gamma_1) > 0$  for every  $s, t \in [0, 1]$  with  $s < t$  by (2.6). Since  $r(\gamma)_r = \gamma_r$  for every  $\gamma \in \text{TGeo}(M)$  and every  $r \in \{0, 1\}$  by definition of  $r$ ,  $(\mathbf{e}_0, \mathbf{e}_1)_\# \pi \in \Pi_{\ll}(\mu_0, \mu_1)$  is  $\ell_p$ -optimal for  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ . A standard argument using the reverse triangle inequality (2.4) thus yields

$$\ell_p(\mu_s, \mu_t) = (t - s)\ell_p(\mu_0, \mu_1) > 0$$

for every timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  and every  $s, t \in [0, 1]$  with  $s < t$ . Hence every such curve  $(\mu_t)_{t \in [0, 1]}$  is an  $\ell_p$ -geodesic in the sense of [24, Def. 2.13] and [66, Def. 1.1], but the converse is unclear to us.

**2.3.7. Rényi entropy.** From now on, we draw our attention to *measured* Lorentzian pre-length spaces, defined as follows.

**Definition 2.15.** We call a sextuple  $(M, d, \mathbf{m}, \ll, \leq, \tau)$  a measured Lorentzian pre-length space if  $(M, d, \ll, \leq, \tau)$  is a Lorentzian pre-length space according to Definition 2.4, and  $\mathbf{m}$  is a fully supported Radon measure on  $M$  as in Chapter 1. Accordingly, measured Lorentzian length spaces and measured Lorentzian geodesic spaces are defined.

All over the sequel, we use the convenient abbreviation

$$\mathcal{X} := (M, \mathbf{d}, \mathbf{m}, \ll, \leq, \tau). \quad (2.9)$$

All Lorentzian notions from Section 2.2 phrased for  $\mathcal{X}$  will be evidently understood with respect to the Lorentzian structure  $(M, \mathbf{d}, \ll, \leq, \tau)$ .

Given any  $N \in [1, \infty)$ , define the  $N$ -Rényi entropy  $\mathcal{S}_N: \mathcal{P}(M) \rightarrow [-\infty, 0]$  with respect to  $\mathbf{m}$  through

$$\mathcal{S}_N(\mu) := - \int_M \rho^{-1/N} \mathrm{d}\mu = - \int_M \rho^{1-1/N} \mathrm{d}\mathbf{m}$$

subject to the decomposition  $\mu = \rho \mathbf{m} + \mu_\perp$ . This functional will play a central role in our work. Observe that  $\mathcal{S}_1(\mu) = -\mathbf{m}[\mathrm{spt}(\rho \mathbf{m})]$ . More generally, for every  $N \in [1, \infty)$ , by Jensen's inequality we have  $\mathcal{S}_N(\mu) \geq -\mathbf{m}[\mathrm{spt} \mu]^{1/N}$  for every  $\mu \in \mathcal{P}_c(M)$ . Moreover,  $\mathcal{S}_N$  is jointly weakly lower semicontinuous in the following sense [61, Thm. B.33] provided (for notational convenience, cf. Subsection 2.3.6)  $M$  is compact. Assume two sequences  $(\mu_n)_{n \in \mathbf{N}}$  and  $(\mathbf{m}_n)_{n \in \mathbf{N}}$  in  $\mathcal{P}(M)$  to converge weakly to  $\mu_\infty \in \mathcal{P}(M)$  and  $\mathbf{m}_\infty \in \mathcal{P}(M)$ , respectively. Then for the Rényi entropies  $\mathcal{S}_N^n$  with respect to  $\mathbf{m}_n$ ,  $n \in \mathbf{N}_\infty$ , we have

$$\mathcal{S}_N^\infty(\mu_\infty) \leq \liminf_{n \rightarrow \infty} \mathcal{S}_N^n(\mu_n).$$

**2.3.8. Boltzmann entropy.** Another functional on  $\mathcal{P}(M)$  we occasionally consider, cf. Lemma 3.8, Section 3.5, and Section 4.5, and whose relativistic consequences have been studied deeply in [24, 66, 69], is the *Boltzmann entropy*  $\mathrm{Ent}_\mathbf{m}: \mathcal{P}(M) \rightarrow [-\infty, \infty]$  with respect to  $\mathbf{m}$ . It is defined by

$$\mathrm{Ent}_\mathbf{m}(\mu) := \begin{cases} \int_M \rho \log \rho \mathrm{d}\mathbf{m} & \text{if } \mu = \rho \mathbf{m} \in \mathcal{P}^{\mathrm{ac}}(M, \mathbf{m}), (\rho \log \rho)^+ \in L^1(M, \mathbf{m}), \\ \infty & \text{otherwise.} \end{cases}$$

Let  $\mathcal{D}(\mathrm{Ent}_\mathbf{m})$  denote the usual finiteness domain of  $\mathrm{Ent}_\mathbf{m}$ .

For every  $\mu \in \mathcal{P}_c(M)$ , we have  $\mathrm{Ent}_\mathbf{m}(\mu) \geq -\log \mathbf{m}[\mathrm{spt} \mu]$ . Moreover, as for the Rényi entropy  $\mathrm{Ent}_\mathbf{m}$  is weakly lower semicontinuous on  $\mathcal{P}(C)$  for every closed  $C \subset M$  with  $\mathbf{m}[C] < \infty$  [85, Lem. 4.1]; in fact, for every  $\mu \in \mathcal{P}_c(M)$ ,

$$\mathrm{Ent}_\mathbf{m}(\mu) = \lim_{N \rightarrow \infty} [N - N \mathcal{S}_N(\mu)] = \sup\{N - N \mathcal{S}_N(\mu) : N \in [1, \infty)\}. \quad (2.10)$$

**2.4. Timelike essential nonbranching.** Finally, we introduce the following Lorentzian version of essential nonbranching for metric measure spaces [77], termed *timelike essential nonbranching*. The ‘‘curvewise nonbranching’’ part is adapted from [24, Def. 1.10].

Given  $s, t \in [0, 1]$  with  $s < t$ , let  $\mathrm{restr}_s^t: C([0, 1]; M) \rightarrow C([0, 1]; M)$  be the map restricting a curve  $\gamma: [0, 1] \rightarrow M$  to  $[s, t]$  and then stretching it to  $[0, 1]$ , i.e.

$$\mathrm{restr}_s^t(\gamma)_r := \gamma_{(1-r)s+rt}. \quad (2.11)$$

**Definition 2.16.** A subset  $G \subset \mathrm{TGeo}^\tau(M)$  is *timelike nonbranching* if for every  $r \in (0, 1)$ ,  $\mathrm{restr}_s^t$  is injective for every  $(s, t) \in \{(0, r), (r, 1)\}$ .

We call  $(M, \mathbf{d}, \ll, \leq, \tau)$  *timelike  $p$ -essentially nonbranching*,  $p \in (0, 1]$ , if every plan  $\pi$  representing some timelike proper-time parametrized  $\ell_p$ -geodesic is concentrated on a timelike nonbranching subset of  $\mathrm{TGeo}^\tau(M)$ .

*Remark 2.17.* The set  $G := \mathrm{TGeo}^\tau(M)$  is timelike nonbranching if the Lorentzian space  $(M, \mathbf{d}, \ll, \leq, \tau)$  is timelike nonbranching according to [24, Def. 1.10]. In particular, timelike nonbranching Lorentzian pre-length spaces are timelike  $p$ -essentially nonbranching for every  $p \in (0, 1]$ .



Important results derived later which assume  $p$ -timelike essential nonbranching are the local-to-global property, Theorem 3.45, uniqueness of  $\ell_p$ -optimal couplings and timelike proper-time parametrized  $\ell_p$ -geodesics, Theorem 4.16 and Theorem 4.17, and the equivalence of some of our curvature-dimension conditions and measure-contraction properties to their entropic versions set up by Cavalletti and Mondino, Theorem 3.35 and Theorem 4.20.

The following result will be useful at many occasions. It follows directly from the definition of  $p$ -timelike essential nonbranching. See [9, Lem. 2.6], [32, Lem. 3.11], and [85, Lem. 2.11] for related results for metric measure spaces.

**Lemma 2.18.** *Assume timelike  $p$ -essential nonbranching for some  $p \in (0, 1]$ . Let  $\pi$  be an  $\ell_p$ -optimal geodesic plan which can be decomposed as  $\pi = \lambda_1 \pi^1 + \dots + \lambda_n \pi^n$  for some  $n \in \mathbf{N}$ ,  $\lambda_1, \dots, \lambda_n \in (0, 1)$ , and  $\ell_p$ -optimal geodesic plans  $\pi^1, \dots, \pi^n$ . If the measures  $(e_0)_\# \pi^1, \dots, (e_0)_\# \pi^n$  or  $(e_1)_\# \pi^1, \dots, (e_1)_\# \pi^n$  are mutually singular, so are  $(e_t)_\# \pi^1, \dots, (e_t)_\# \pi^n$  for every  $t \in (0, 1)$ .*

### 3. TIMELIKE CURVATURE-DIMENSION CONDITION

**Assumption 3.1.** *From now on, in the notation of (2.9), unless stated otherwise we assume  $\mathcal{X}$  to be a causally closed,  $\mathcal{K}$ -globally hyperbolic, measured Lorentzian geodesic space.*

Of course, our main notions from Definition 3.2, Definition 3.3, and Definition 4.1 a priori make sense for general Lorentzian pre-length spaces. We always mention explicitly if other results hold in greater generality than under Assumption 3.1.

**3.1. Definition and basic properties.** Definition 3.2 is the Lorentzian counterpart of the *reduced* curvature-dimension condition [9, Def. 2.3]. On the other hand, Definition 3.3 is the natural Lorentzian adaptation of the  $\text{CD}(K, N)$  condition for metric measure spaces from [86, Def. 1.3]. Both conditions deserve their names from their compatibility with smooth Lorentzian case, cf. Theorem A.1.

**Definition 3.2.** *Let  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ .*

- a. *We say that  $\mathcal{X}$  satisfies the reduced timelike curvature-dimension condition  $\text{TCD}_p^*(K, N)$  if for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$ , there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  connecting  $\mu_0$  to  $\mu_1$  as well as a timelike  $p$ -dualizing coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,*

$$\begin{aligned} \mathcal{S}_{N'}(\mu_t) \leq & - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1) \\ & - \int_{M^2} \sigma_{K, N'}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N'} d\pi(x^0, x^1). \end{aligned}$$

- b. *If the previous statement holds only for every strongly timelike  $p$ -dualizable  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$ , we say that  $\mathcal{X}$  obeys the weak reduced timelike curvature-dimension condition  $\text{wTCD}_p^*(K, N)$ .*

**Definition 3.3.** *Let  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ .*

- a. *We say that  $\mathcal{X}$  obeys the timelike curvature-dimension condition  $\text{TCD}_p(K, N)$  if for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$ , there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  connecting  $\mu_0$  to  $\mu_1$  as well as a timelike  $p$ -dualizing coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,*

$$\mathcal{S}_{N'}(\mu_t) \leq - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1)$$

$$- \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N'} d\pi(x^0, x^1).$$

- b. If the previous statement holds only for every strongly timelike  $p$ -dualizable  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$ , we say that  $\mathcal{X}$  obeys the weak timelike curvature-dimension condition  $\text{wTCD}_p(K, N)$ .

*Remark 3.4.* The conditions from Definition 3.2 and Definition 3.3 are a priori formulated for compactly supported endpoints  $\mu_0$  and  $\mu_1$  only. In the timelike  $p$ -essentially nonbranching case, cf. Definition 2.16, this does in fact imply the respective defining statements for more general  $\mu_0, \mu_1 \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  according to Theorem 3.41 and Theorem 3.35.

*Remark 3.5.* Note that  $\text{TCD}_p^*(0, N)$  and  $\text{wTCD}_p^*(0, N)$  coincide with  $\text{TCD}_p(0, N)$  and  $\text{wTCD}_p(0, N)$ , respectively, for every  $p \in (0, 1)$  and every  $N \in [1, \infty)$ , since

$$\sigma_{0,N'}^{(r)}(\vartheta) = \tau_{0,N'}^{(r)}(\vartheta) = r$$

for every  $r \in [0, 1]$ , every  $N' \in [1, \infty)$ , and every  $\vartheta \geq 0$ .

In the remainder of this section, we examine elementary properties of the conditions introduced above, and their respective relations. Proposition 3.6 as well as Proposition 3.7 hold on any measured Lorentzian pre-length space.

**Proposition 3.6.** *The following hold for every  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ ; set  $K^* := K(N - 1)/N$ .*

- (i) *The condition  $\text{TCD}_p(K, N)$  implies  $\text{TCD}_p^*(K, N)$ .*
- (ii) *The condition  $\text{wTCD}_p(K, N)$  implies  $\text{wTCD}_p^*(K, N)$ .*
- (iii) *If  $K > 0$ , the condition  $\text{TCD}_p^*(K, N)$  implies  $\text{TCD}_p(K^*, N)$ .*
- (iv) *If  $K > 0$ , the condition  $\text{wTCD}_p^*(K, N)$  implies  $\text{wTCD}_p(K^*, N)$ .*

*Proof.* The first two statements are a direct consequence of the inequality

$$\tau_{K,N'}^{(r)}(\vartheta) \geq \sigma_{K,N'}^{(r)}(\vartheta)$$

for every  $r \in [0, 1]$ , every  $N' \geq N$ , and every  $\vartheta \geq 0$ , cf. the proof of [9, Prop. 2.5]. The last two statements follow similarly by noting that

$$\tau_{K^*,N'}^{(t)}(\vartheta) \leq \sigma_{K^*,N'}^{(t)}(\vartheta). \quad \square$$

**Proposition 3.7.** *Assume  $\text{TCD}_p^*(K, N)$  for  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Then the following statements hold.*

- (i) **Consistency.** *The condition  $\text{TCD}_p^*(K', N')$  holds for every  $K' \leq K$  and every  $N' \geq N$ .*
- (ii) **Scaling.** *Given any  $a, b, \theta > 0$ , the rescaled measured Lorentzian pre-length space  $(M, \text{ad}, \text{bm}, \ll, \leq, \theta\tau)$  satisfies  $\text{TCD}_p^*(K/\theta^2, N)$ .*

*Analogous statements are obeyed by the conditions  $\text{wTCD}_p^*(K, N)$ ,  $\text{TCD}_p(K, N)$ , and  $\text{wTCD}_p(K, N)$ .*

*Proof.* Concerning (i), for all four conditions consistency in  $K$  follows by nondecreasingness of  $\sigma_{K,N}^{(r)}(\vartheta)$  and  $\tau_{K,N}^{(r)}(\vartheta)$  in  $K \in \mathbf{R}$  for fixed  $r \in [0, 1]$ ,  $N \in [1, \infty)$ , and  $\vartheta \geq 0$ . Consistency in  $N$  is clear since the defining inequality, in either case, is asked to hold for every  $N'' \geq N$ , so is particularly satisfied for every  $N'' \geq N'$ .

Item (ii) follows from the scaling properties

$$\begin{aligned} \sigma_{K/\theta^2, N'}^{(r)}(\theta \vartheta) &= \sigma_{K, N'}^{(r)}(\vartheta), \\ \tau_{K/\theta^2, N'}^{(r)}(\theta \vartheta) &= \tau_{K, N'}^{(r)}(\vartheta). \end{aligned} \quad \square$$

Lastly, recall that the  $N$ -Rényi entropy  $\mathcal{S}_N(\mu)$ ,  $N \in [1, \infty)$ , at  $\mu \in \mathcal{P}(M)$  only depends on the  $\mathfrak{m}$ -absolutely continuous part of  $\mu$ , which might be trivial. Along those timelike proper-time parametrized  $\ell_p$ -geodesics in Definition 3.2 or Definition 3.3, this does not happen; in fact, under mild additional hypotheses, these always consist of  $\mathfrak{m}$ -absolutely continuous measures. This is addressed in Lemma 3.8 and will be useful at various occasions. (We also use variants of it without explicit notice occasionally, e.g. in Section 3.6.) Its proof uses (2.10) and is analogous to the one of [86, Prop. 1.6], hence omitted.

**Lemma 3.8.** *Fix  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Let  $(\mu_0, \mu_1) = (\rho_0 \mathfrak{m}, \mu_1 \mathfrak{m}) \in (\mathcal{P}_c(M) \cap \mathcal{D}(\text{Ent}_{\mathfrak{m}}))^2$ , let  $\mu \in \mathcal{P}_c(M)$ , and let  $\pi \in \Pi(\mu_0, \mu_1)$  be a coupling of  $\mu_0$  and  $\mu_1$  such that for every  $N' \geq N$ ,*

$$\begin{aligned} \mathcal{S}_{N'}(\mu) \leq & - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1) \\ & - \int_{M^2} \sigma_{K, N'}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N'} d\pi(x^0, x^1). \end{aligned}$$

Then  $\mu = \rho \mathfrak{m} \in \mathcal{D}(\text{Ent}_{\mathfrak{m}})$  with

$$\text{Ent}_{\mathfrak{m}}(\mu) \leq (1-t) \text{Ent}_{\mathfrak{m}}(\mu_0) + t \text{Ent}_{\mathfrak{m}}(\mu_1) - \frac{K}{2} t(1-t) \|\tau\|_{L^2(M^2, \pi)}^2.$$

*Remark 3.9.* Using Proposition B.9, the assumptions in Lemma 3.8 are satisfied for  $\mu := \mu_t$ ,  $t \in [0, 1]$ , and  $\pi$  if these objects are coming from a timelike proper-time parametrized  $\ell_p$ -geodesic as well as a timelike  $p$ -dualizing coupling, respectively, witnessing the inequality defining  $\text{TCD}_p^*(K, N)$ .

**Corollary 3.10.** *For every  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ , the  $\text{wTCD}_p^*(K, N)$  condition implies  $\text{wTCD}_p(K, \infty)$  in the sense of [16, Def. 4.1].*

**3.2. Geometric inequalities.** Now we derive fundamental geometric inequalities from the conditions introduced in Chapter 3. We start with the more restrictive  $\text{wTCD}_p(K, N)$ -case which implies sharp versions of these facts, cf. Corollary 3.18; Subsection 3.2.4 provides nonsharp extensions for  $\text{wTCD}_p^*(K, N)$ -spaces. All proofs are variations of standard arguments [86], see also [9, 24].

Some of the geometric inequalities to follow hold in fact under weaker timelike measure-contraction properties, cf. Proposition 4.8 and Remark 4.9.

Moreover, all results in this section also hold if  $\mathcal{X}$  is a locally causally closed and globally hyperbolic measured Lorentzian geodesic space.

**3.2.1. Sharp timelike Brunn–Minkowski inequality.** To introduce the *timelike Brunn–Minkowski inequality* in the next Proposition 3.11, given  $A_0, A_1 \subset M$  and  $t \in [0, 1]$ , define the set of timelike  $t$ -intermediate points between  $A_0$  and  $A_1$  as

$$A_t := \{\gamma_t : \gamma \in \text{TGeo}^\tau(M), \gamma_0 \in A_0, \gamma_1 \in A_1\}.$$

Furthermore, define

$$\Theta := \begin{cases} \sup \tau(A_0 \times A_1) & \text{if } K < 0, \\ \inf \tau(A_0 \times A_1) & \text{otherwise.} \end{cases} \quad (3.1)$$

**Proposition 3.11.** *Assume  $\text{wTCD}_p(K, N)$  for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Let  $A_0, A_1 \subset M$  be relatively compact Borel sets with  $\mathfrak{m}[A_0] \mathfrak{m}[A_1] > 0$ , and assume strong timelike  $p$ -dualizability of*

$$(\mu_0, \mu_1) := (\mathfrak{m}[A_0]^{-1} \mathfrak{m} \llcorner A_0, \mathfrak{m}[A_1]^{-1} \mathfrak{m} \llcorner A_1).$$

Then for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\mathfrak{m}[A_t]^{1/N'} \geq \tau_{K, N'}^{(1-t)}(\Theta) \mathfrak{m}[A_0]^{1/N'} + \tau_{K, N'}^{(t)}(\Theta) \mathfrak{m}[A_1]^{1/N'}. \quad (3.2)$$

Assuming  $\text{TCD}_p(K, N)$  in place of  $\text{wTCD}_p(K, N)$ , the same conclusion holds if  $(\mu_0, \mu_1)$  is merely timelike  $p$ -dualizable.

*Proof.* Without loss of generality, we assume  $N' > 1$  and  $\mathfrak{m}[A_t] < \infty$ . Let a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  connecting  $\mu_0$  to  $\mu_1$  and a timelike  $p$ -dualizing  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  witness the semiconvexity inequality defining  $\text{wTCD}_p(K, N)$ . Note that  $\text{spt } \mu_t \subset A_t$  and, by  $\text{wTCD}_p(K, N)$ , that  $\mathfrak{m}[A_t] > 0$ . Let  $\rho_t$  denote the density of the absolutely continuous part of  $\mu_t$  with respect to  $\mathfrak{m}$ . Employing Jensen's inequality, and  $\text{wTCD}_p(K, N)$  again we obtain

$$\begin{aligned} & \tau_{K,N}^{(1-t)}(\Theta) \mathfrak{m}[A_0]^{1/N'} + \tau_{K,N}^{(t)}(\Theta) \mathfrak{m}[A_1]^{1/N'} \\ & \leq \int_{A_t} \rho_t^{1-1/N'} \, d\mathfrak{m} \leq \mathfrak{m}[A_t] \left[ \int_{A_t} \rho_t \, d\mathfrak{m} \right]^{1-1/N'} \leq \mathfrak{m}[A_t]^{-1/N'}, \end{aligned}$$

which is the desired claim.

The proof of the last statement is completely analogous.  $\square$

*Remark 3.12.* Note that in the case  $K \geq 0$ , (3.2) implies

$$\mathfrak{m}[A_t]^{1/N'} \geq (1-t) \mathfrak{m}[A_0]^{1/N'} + t \mathfrak{m}[A_1]^{1/N'}.$$

*Remark 3.13.* Recall from Remark 2.12 that the strong timelike  $p$ -dualizability hypothesis for  $(\mu_0, \mu_1)$  in Proposition 3.11 is satisfied if  $A_0 \times A_1 \subset M_{\ll}^2$  provided the space  $\mathcal{X}$  is locally causally closed, globally hyperbolic, and geodesic.

3.2.2. *Sharp timelike Bonnet–Myers inequality.* Next, an immediate consequence of Proposition 3.11 is the subsequent *timelike Bonnet–Myers inequality*.

**Corollary 3.14.** *Assume the  $\text{wTCD}_p(K, N)$  condition for some  $p \in (0, 1)$ ,  $K > 0$ , and  $N \in [1, \infty)$ . Then*

$$\sup \tau(M^2) \leq \pi \sqrt{\frac{N-1}{K}}.$$

*Proof.* Suppose to the contrapositive that for some  $\varepsilon > 0$ , we have

$$\tau(z_0, z_1) \geq \pi \sqrt{\frac{N-1}{K}} + 4\varepsilon$$

for two given points  $z_0, z_1 \in M$ . By continuity of  $\tau$ , we fix  $\delta > 0$  and  $x_0, x_1 \in M$  such that  $A_0 := \mathbb{B}^d(x_0, \delta) \subset I^+(z_0)$ ,  $A_1 := \mathbb{B}^d(x_1, \delta) \subset I^-(z_1)$ , and

$$\inf \tau(\mathbb{B}^d(x_0, \delta) \times \mathbb{B}^d(x_1, \delta)) \geq \pi \sqrt{\frac{N-1}{K}} + \varepsilon.$$

This implies strong timelike  $p$ -dualizability of  $(\mathfrak{m}[A_0]^{-1} \mathfrak{m} \llcorner A_0, \mathfrak{m}[A_1]^{-1} \mathfrak{m} \llcorner A_1)$  by Remark 2.12. Hence Proposition 3.11 is applicable, and the inherent inequality  $\Theta \geq \pi \sqrt{(N-1)/K} + \varepsilon$  together with the definition of  $\tau_{K,N}^{(r)}$  gives  $\mathfrak{m}[A_{1/2}] = \infty$ . On the other hand, the set  $A_{1/2}$  is relatively compact since  $A_{1/2} \subset J(z_0, z_1)$  by global hyperbolicity, which makes the situation  $\mathfrak{m}[A_{1/2}] = \infty$  impossible.  $\square$

*Remark 3.15.* Corollary 3.14 implies that, under its assumptions, the  $\tau$ -length of every causal curve in  $M$  is no larger than  $\pi \sqrt{(N-1)/K}$ . Moreover, if  $K > 0$  and  $N = 1$  then  $M$  does not contain any chronologically related pair of points.

3.2.3. *Sharp timelike Bishop–Gromov inequality.* To prove volume growth estimates à la *Bishop–Gromov*, cf. Theorem 3.16 below, we recall the notion of  $\tau$ -star-shaped sets from [24, Sec. 3.1]; this is used to localize the volume of  $\tau$ -balls

$$\mathbb{B}^\tau(x, r) := \{y \in I^+(x) : \tau(x, y) < r\} \cup \{x\}, \quad (3.3)$$

with  $x \in M$  and  $r > 0$ , which typically have infinite volume.

A set  $E \subset I^+(x) \cup \{x\}$  is termed  $\tau$ -star-shaped with respect to  $x \in M$  if for every  $\gamma \in \text{TGeo}^\tau(M)$  with  $\gamma_0 = x$  and  $\gamma_1 \in E$ , we have  $\gamma_t \in E$  for every  $t \in (0, 1)$ . For such  $E$  and  $x$ , and given  $r > 0$ , we define the quantities

$$\begin{aligned} v_r &:= \mathfrak{m}[\bar{\mathbf{B}}^\tau(x, r) \cap E], \\ s_r &:= \limsup_{\delta \rightarrow 0} \delta^{-1} \mathfrak{m}[(\bar{\mathbf{B}}^\tau(x, r + \delta) \setminus \mathbf{B}^\tau(x, r)) \cap E]. \end{aligned}$$

Whenever confusion is excluded, we avoid making notationally explicit the dependency of  $v_r$  and  $s_r$  on  $E$  and  $x$ . Recalling (2.1), we set  $\mathfrak{s}_{K,N} := \mathfrak{s}_{K/N}$  and

$$\mathfrak{v}_{K,N}(r) := \left[ \int_0^r \mathfrak{s}_{K,N-1}(r)^{N-1} dr \right]^{1/N}. \quad (3.4)$$

In the next result, we employ the convention  $1/0 := \infty$ .

**Theorem 3.16.** *Assume  $\text{wTCD}_p(K, N)$  for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in (1, \infty)$ . Let  $E \subset I^+(x) \cup \{x\}$  be a compact set which is  $\tau$ -star-shaped with respect to  $x \in M$ . Then for every  $r, R > 0$  with  $r < R \leq \pi\sqrt{(N-1)/\max\{K, 0\}}$ ,*

$$\begin{aligned} \frac{s_r}{s_R} &\geq \left[ \frac{\mathfrak{s}_{K,N-1}(r)}{\mathfrak{s}_{K,N-1}(R)} \right]^{N-1}, \\ \frac{v_r}{v_R} &\geq \left[ \frac{\mathfrak{v}_{K,N}(r)}{\mathfrak{v}_{K,N}(R)} \right]^N. \end{aligned}$$

*Proof.* We assume  $K > 0$ , the rest is argued similarly. Define  $t := r/R$ , let  $\delta > 0$ , and let  $\varepsilon > 0$  be small enough such that  $A_0 \times A_1 \subset M_{\ll}^2$ , where  $A_0 := \mathbf{B}^\tau(x, \varepsilon) \cap E$  and  $A_1 := (\bar{\mathbf{B}}^\tau(x, R + \delta R) \setminus \mathbf{B}^\tau(x, R)) \cap E$ . Thus  $(\mathfrak{m}[A_0]^{-1} \mathfrak{m} \llcorner A_0, \mathfrak{m}[A_1]^{-1} \mathfrak{m} \llcorner A_1)$  is strongly timelike  $p$ -dualizable according to Remark 2.12. Now we observe that  $A_t \subset (\mathbf{B}^\tau(x, r + \delta r) \setminus \mathbf{B}^\tau(x, r)) \cap E$ . By the reverse triangle inequality for  $\tau$ , we get  $\Theta \leq R + \delta R$ . Employing Proposition 3.11, we therefore obtain

$$\begin{aligned} &\mathfrak{m}[(\bar{\mathbf{B}}^\tau(x, r + \delta r) \setminus \mathbf{B}^\tau(x, r)) \cap E]^{1/N} \\ &\geq \tau_{K,N}^{(1-r/R)}(R + \delta R) \mathfrak{m}[\mathbf{B}^\tau(x, \varepsilon) \cap E]^{1/N} \\ &\quad + \tau_{K,N}^{(r/R)}(R + \delta R) \mathfrak{m}[(\bar{\mathbf{B}}^\tau(x, R + \delta R) \setminus \mathbf{B}^\tau(x, R)) \cap E]^{1/N} \\ &\geq \tau_{K,N}^{(r/R)}(R + \delta R) \mathfrak{m}[(\bar{\mathbf{B}}^\tau(x, R + \delta R) \setminus \mathbf{B}^\tau(x, R)) \cap E]^{1/N}. \end{aligned}$$

Writing out the expression  $\tau_{K,N}^{(r/R)}(R + \delta R)^N$  and finally sending  $\delta \rightarrow 0$  implies the claimed lower bound for  $s_r/s_R$ .

As in the proof of [86, Thm. 2.3], we argue that  $v$  is locally Lipschitz continuous on  $(0, \infty)$ . Hence, it is differentiable  $\mathcal{L}^1$ -a.e. on  $(0, \infty)$ , and  $v_r = \int_0^r \dot{v}_s ds$  for every  $r > 0$ . Therefore, the claimed lower bound on  $v_r/v_R$  follows from the previous part in combination with [92, Lem. 18.9].  $\square$

*Remark 3.17.* If  $K = 0$ , the estimates from Theorem 3.16 become

$$\begin{aligned} \frac{s_r}{s_R} &\geq \left[ \frac{r}{R} \right]^{N-1}, \\ \frac{v_r}{v_R} &\geq \left[ \frac{r}{R} \right]^N. \end{aligned}$$

In fact, these estimates hold for  $N = 1$  as well.

Incidentally, from Theorem 3.16 we deduce the following natural and sharp upper bound on the  $\tau$ -Hausdorff dimension as introduced and studied in [67]. Our results improve [67, Thm. 5.2, Cor. 5.3]. We refer to [67] for details about the notions presented in the subsequent Corollary 3.18.

**Corollary 3.18.** *Let  $(M, \langle \cdot, \cdot \rangle)$  be a continuous, globally hyperbolic, causally plain spacetime of dimension  $n \in \mathbf{N}$ , cf. Example 2.7. Assume its induced Lorentzian geodesic space, with  $\mathfrak{m}$  being the Lorentzian volume measure induced by  $\langle \cdot, \cdot \rangle$ , to obey  $\text{wTCD}_p(K, N)$  for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Then the geometric dimension  $\dim^\tau M$  in the sense of [67, Def. 3.1] satisfies*

$$n = \dim^\tau M \leq N.$$

*Remark 3.19.* As in [67, Thm. 5.2], which is proven under  $\text{wTCD}_p^e(K, N)$ , assuming  $\text{wTCD}_p^*(K, N)$  instead of  $\text{wTCD}_p(K, N)$  in Corollary 3.18 would only lead to the conclusion  $n = \dim^\tau M \leq N + 1$  by Theorem 3.22.

**3.2.4. Versions in the reduced case.** With evident modifications, the following non-sharp versions of Proposition 3.11, Corollary 3.14, Theorem 3.16, and Corollary 3.18 are readily verified under the more general  $\text{wTCD}_p^*(K, N)$  condition. The deduced inequalities are analogous to their counterparts drawn from the  $\text{wTCD}_p^e(K, N)$  condition in [24, Prop. 3.4, Prop. 3.5, Prop. 3.6].

In view of Theorem 3.22 below, recall the definition of  $\mathfrak{v}_{K, N}$  from (3.4).

**Proposition 3.20.** *Assume  $\text{wTCD}_p^*(K, N)$  for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Let  $A_0, A_1 \subset M$  be relatively compact Borel sets which satisfy  $\mathfrak{m}[A_0] \mathfrak{m}[A_1] > 0$ , and assume strong timelike  $p$ -dualizability of*

$$(\mu_0, \mu_1) := (\mathfrak{m}[A_0]^{-1} \mathfrak{m} \llcorner A_0, \mathfrak{m}[A_1]^{-1} \mathfrak{m} \llcorner A_1).$$

Let  $\Theta$  be as in (3.1). Then for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\mathfrak{m}[A_t]^{1/N'} \geq \sigma_{K, N'}^{(1-t)}(\Theta) \mathfrak{m}[A_0]^{1/N'} + \sigma_{K, N'}^{(t)}(\Theta) \mathfrak{m}[A_1]^{1/N'}.$$

Assuming  $\text{TCD}_p^*(K, N)$  in place of  $\text{wTCD}_p^*(K, N)$ , the same conclusion holds if  $(\mu_0, \mu_1)$  is merely timelike  $p$ -dualizable.

**Corollary 3.21.** *Assume the  $\text{wTCD}_p^*(K, N)$  condition for some  $p \in (0, 1)$ ,  $K > 0$ , and  $N \in [1, \infty)$ . Then*

$$\sup \tau(M^2) \leq \pi \sqrt{\frac{N}{K}}.$$

**Theorem 3.22.** *Assume  $\text{wTCD}_p^*(K, N)$  for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in (1, \infty)$ . Let  $E \subset I^+(x) \cup \{x\}$  be a compact set which is  $\tau$ -star-shaped with respect to  $x \in M$ . Then for every  $r, R > 0$  with  $r < R \leq \pi \sqrt{N / \max\{K, 0\}}$ ,*

$$\begin{aligned} \frac{s_r}{s_R} &\geq \left[ \frac{\mathfrak{s}_{K, N}(r)}{\mathfrak{s}_{K, N}(R)} \right]^N, \\ \frac{\mathfrak{v}_r}{\mathfrak{v}_R} &\geq \left[ \frac{\mathfrak{v}_{K, N+1}(r)}{\mathfrak{v}_{K, N+1}(R)} \right]^{N+1}. \end{aligned}$$

**3.3. Stability.** In this section, we prove a key property of our timelike curvature-dimension conditions, namely the weak stability of the notions from Definition 3.2 and Definition 3.3, cf. Theorem 3.29. The relevant notion of convergence of measured Lorentzian pre-length spaces, see Definition 3.24, is due to [24, Sec. 3.3].

**Definition 3.23.** *For Lorentzian pre-length spaces  $(M^i, \mathfrak{d}^i, \ll^i, \leq^i, \tau^i)$ ,  $i \in \{0, 1\}$ , we term a map  $\iota: M^0 \rightarrow M^1$  a Lorentzian isometric embedding if  $\iota$  is a topological embedding such that for every  $x, y \in M^0$ ,*

- a.  $x \leq^0 y$  if and only if  $\iota(x) \leq^1 \iota(y)$ , and
- b.  $\tau^1(\iota(x), \iota(y)) = \tau^0(x, y)$ .

Given any  $k \in \mathbf{N}_\infty := \mathbf{N} \cup \{\infty\}$  let  $(M_k, \mathfrak{d}_k, \mathfrak{m}_k, \ll_k, \leq_k, \tau_k)$  be a fixed measured Lorentzian pre-length space which, as in (2.9), we will abbreviate by  $\mathcal{X}_k$ .

**Definition 3.24.** We define  $(\mathcal{X}_k)_{k \in \mathbf{N}}$  to converge to  $\mathcal{X}_\infty$  if the following holds.

- a. There is a proper, causally closed,  $\mathcal{K}$ -globally hyperbolic measured Lorentzian geodesic space  $(M, \mathbf{d}, \ll, \leq, \tau)$  such that for every  $k \in \mathbf{N}_\infty$  there exists a Lorentzian isometric embedding  $\iota_k: M_k \rightarrow M$ .
- b. In terms of the maps  $\iota_k$  from a., the sequence  $((\iota_k)_\# \mathbf{m}_k)_{k \in \mathbf{N}}$  converges to  $(\iota_\infty)_\# \mathbf{m}_\infty$  in duality with  $C_c(M)$ , i.e. for every  $\varphi \in C_c(M)$  we have

$$\lim_{k \rightarrow \infty} \int_M \varphi d(\iota_k)_\# \mathbf{m}_k = \int_M \varphi d(\iota_\infty)_\# \mathbf{m}_\infty.$$

*Remark 3.25.* If  $(\mathcal{X}_k)_{k \in \mathbf{N}}$  converges to  $\mathcal{X}_\infty$  according to Definition 3.24,  $\mathcal{X}_k$  is automatically causally closed,  $\mathcal{K}$ -globally hyperbolic, and geodesic for every  $k \in \mathbf{N}_\infty$ .

The following Lemma 3.26 is an implicit consequence from the proof of [24, Lem. 3.15] and will be useful for the proof of Theorem 3.29.

**Lemma 3.26.** Assume that  $\mathcal{X}$  is causally closed, globally hyperbolic, and geodesic. Let the pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$  admit some  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$ ,  $p \in (0, 1]$ . Then there exist sequences  $(\pi^n)_{n \in \mathbf{N}}$  in  $\mathcal{P}(M^2)$  and  $(a_n)_{n \in \mathbf{N}}$  in  $[1, \infty)$  with the properties

- (i)  $(a_n)_{n \in \mathbf{N}}$  converges to 1,
- (ii)  $\pi^n[M_{\ll}^2] = 1$  for every  $n \in \mathbf{N}$ ,
- (iii)  $\pi^n = \rho^n \mathbf{m}^{\otimes 2} \in \mathcal{P}^{\text{ac}}(M^2, \mathbf{m}^{\otimes 2})$  with  $\rho^n \in L^\infty(M^2, \mathbf{m}^{\otimes 2})$  for every  $n \in \mathbf{N}$ ,
- (iv)  $(\pi^n)_{n \in \mathbf{N}}$  converges weakly to  $\pi$ ,
- (v) the density  $\rho_0^n$  and  $\rho_1^n$  of the first and second marginal of  $\pi^n$  with respect to  $\mathbf{m}$  is no larger than  $a_n \rho_0$  and  $a_n \rho_1$ , respectively, for every  $n \in \mathbf{N}$ , and
- (vi)  $\rho_0^n \rightarrow \rho_0$  and  $\rho_1^n \rightarrow \rho_1$  in  $L^1(M, \mathbf{m})$  as  $n \rightarrow \infty$ .

Furthermore, in the notation of Lemma 3.26, given any  $K \in \mathbf{R}$ ,  $N \in [1, \infty)$ , and  $t \in [0, 1]$ , as well as a measure  $\pi \in \Pi(\mu_0, \mu_1)$  with marginals  $\mu_0 = \rho_0 \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  and  $\mu_1 = \rho_1 \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ , for brevity we define

$$\begin{aligned} \mathcal{J}_{K,N}^{(t)}(\pi) := & - \int_{M^2} \tau_{K,N}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N} d\pi(x^0, x^1) \\ & - \int_{M^2} \tau_{K,N}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N} d\pi(x^0, x^1). \end{aligned} \quad (3.5)$$

We note the following variant of [86, Lem. 3.2]. The latter proof easily carries over to the setting of Lemma 3.27, where the marginal densities, instead of being assumed to be the same throughout the sequence as in [86], are allowed to be perturbed by constants vanishing in the limit. Compare with Lemma 4.7.

**Lemma 3.27.** Suppose  $\mathbf{m} \in \mathcal{P}(M)$ . Let  $(a_k)_{k \in \mathbf{N}}$  be a sequence in  $[1, \infty)$  converging to 1, and let  $(b_k)_{k \in \mathbf{N}}$  be a sequence of nonnegative real numbers converging to 0. Define  $\nu_{k,0}, \nu_{k,1} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ ,  $k \in \mathbf{N}$ , by

$$\begin{aligned} \nu_{k,0} &:= (a_k + b_k)^{-1} (a_k \mu_0 + b_k \mathbf{m}) = (a_k + b_k)^{-1} (a_k \rho_0 + b_k) \mathbf{m}, \\ \nu_{k,1} &:= (a_k + b_k)^{-1} (a_k \mu_1 + b_k \mathbf{m}) = (a_k + b_k)^{-1} (a_k \rho_1 + b_k) \mathbf{m}. \end{aligned}$$

Then for every sequence  $(\pi_k)_{k \in \mathbf{N}}$  of measures  $\pi_k \in \Pi(\nu_{k,0}, \nu_{k,1})$  which converges weakly to  $\pi \in \Pi(\mu_0, \mu_1)$ , every  $K \in \mathbf{R}$ , every  $N \in [1, \infty)$ , and every  $t \in [0, 1]$ ,

$$\limsup_{k \rightarrow \infty} \mathcal{J}_{K,N}^{(t)}(\pi_k) \leq \mathcal{J}_{K,N}^{(t)}(\pi).$$

*Remark 3.28.* Of course, the assertion from Lemma 3.27 remains valid if  $\tau_{K,N}^{(1-t)}$  and  $\tau_{K,N}^{(t)}$  are replaced by  $\sigma_{K,N}^{(1-t)}$  and  $\sigma_{K,N}^{(t)}$  in (3.5), respectively.

The subsequent proof of Theorem 3.29 below follows [86, Thm. 3.1]. However, by the nature of our timelike curvature-dimension conditions, we additionally have to ensure chronological relations of many measures under consideration. Hence, the proof becomes much longer and quite technical. It may be skipped at first reading; alternatively, the reader may consult the proof of the counterpart Theorem 4.10 of Theorem 3.29 below for a somewhat easier argument in which one does not have to trace chronological  $\ell_p$ -optimal couplings.

Let  $S_N^n$  denote the  $N$ -Rényi entropy with respect to  $\mathbf{m}_n$ ,  $n \in \mathbf{N}_\infty$ .

**Theorem 3.29.** *Assume the convergence of  $(\mathcal{X}_k)_{k \in \mathbf{N}}$  to  $\mathcal{X}_\infty$  as in Definition 3.24. Moreover, let  $(K_k, N_k)_{k \in \mathbf{N}}$  be a sequence in  $\mathbf{R} \times [1, \infty)$  converging to  $(K_\infty, N_\infty) \in \mathbf{R} \times [1, \infty)$ . Suppose the existence of  $p \in (0, 1)$  such that  $\mathcal{X}_k$  obeys  $\text{TCD}_p(K_k, N_k)$  for every  $k \in \mathbf{N}$ . Then  $\mathcal{X}_\infty$  satisfies  $\text{wTCD}_p(K_\infty, N_\infty)$ .*

*The analogous statement in which  $\text{TCD}_p(K_k, N_k)$  and  $\text{wTCD}_p(K_\infty, N_\infty)$  are replaced by  $\text{TCD}_p^*(K_k, N_k)$  and  $\text{wTCD}_p^*(K_\infty, N_\infty)$ ,  $k \in \mathbf{N}$ , respectively, holds too.*

*Proof.* We only prove the first implication, the second is similar. To relax notation, without restriction and further notice we identify  $M_k$  with its image  $\iota_k(M_k)$  in  $M$ , and the measure  $\mathbf{m}_k$  with its push-forward  $(\iota_k)_\# \mathbf{m}_k$  for every  $k \in \mathbf{N}_\infty$ .

**Step 1.** *Reduction to compact  $M$ .* Fix any strongly timelike  $p$ -dualizable pair  $(\mu_{\infty,0}, \mu_{\infty,1}) = (\rho_{\infty,0} \mathbf{m}_\infty, \rho_{\infty,1} \mathbf{m}_\infty) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m}_\infty)$ . By compactness of  $\text{spt } \mu_{\infty,0}$  and  $\text{spt } \mu_{\infty,1}$ , and by  $\mathcal{K}$ -global hyperbolicity of  $M$ , we restrict all arguments below to a compact subset  $C \subset M$  with  $\mathbf{m}_k[C]^{-1} \mathbf{m}_k \llcorner C \rightarrow \mathbf{m}_\infty[C]^{-1} \mathbf{m}_\infty \llcorner C$  weakly as  $k \rightarrow \infty$ . To relax notation, we will thus assume with no loss of generality that  $M$  itself is compact, and that  $\mathbf{m}_k \in \mathcal{P}(M)$  for every  $k \in \mathbf{N}_\infty$ . In particular, without notationally expressing this property all the time, all measures will henceforth be compactly supported.

**Step 2.** *Restriction of the assumptions on  $\mu_{\infty,0}$  and  $\mu_{\infty,1}$ .* We will first assume that  $\rho_{\infty,0}, \rho_{\infty,1} \in L^\infty(M, \mathbf{m}_\infty)$ . The general case is discussed in Step 8 below; we note for now that this conclusion will not conflict with our reductions from Step 1.

**Step 3.** *Construction of a chronological recovery sequence.* The goal of this step is the construction of a sequence  $(\mu_{k,0}, \mu_{k,1})_{k \in \mathbf{N}}$  of timelike  $p$ -dualizable pairs  $(\mu_{k,0}, \mu_{k,1}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)^2$ ,  $k \in \mathbf{N}$ , such that  $\mu_{k,0} \rightarrow \mu_{\infty,0}$  and  $\mu_{k,1} \rightarrow \mu_{\infty,1}$  weakly as  $k \rightarrow \infty$ , up to a subsequence. The highly nontrivial part we address here is that such a pair can in fact be constructed to be chronological.

**Step 3.1.** Let  $W_2$  denote the 2-Wasserstein distance on  $(M, d)$ . Since  $(\mathbf{m}_k)_{k \in \mathbf{N}}$  converges weakly to  $\mathbf{m}_\infty$  and since  $M$  is compact, we have  $W_2(\mathbf{m}_k, \mathbf{m}_\infty) \rightarrow 0$  as  $k \rightarrow \infty$ . Let  $\mathbf{q}_k \in \mathcal{P}(M^2)$  be a  $W_2$ -optimal coupling of  $\mathbf{m}_k$  and  $\mathbf{m}_\infty$ ,  $k \in \mathbf{N}$ . For later use, we disintegrate  $\mathbf{q}_k$  with respect to  $\text{proj}_1$  and  $\text{proj}_2$ , respectively, writing

$$d\mathbf{q}_k(x, y) = d\mathbf{p}_x^k(y) d\mathbf{m}_k(x) = d\mathbf{p}_y^\infty(x) d\mathbf{m}_\infty(y)$$

for certain Borel maps  $\mathbf{p}^k: M \rightarrow \mathcal{P}(M)$  and  $\mathbf{p}^\infty: M \rightarrow \mathcal{P}(M)$ . (Although  $\mathbf{p}^\infty$  depends on  $k$  as well, we do not make this explicit to not overload our notation.) These disintegrations induce nonrelabeled Borel maps  $\mathbf{p}^k: \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty) \rightarrow \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  and  $\mathbf{p}^\infty: \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k) \rightarrow \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty)$  defined by

$$\begin{aligned} d\mathbf{p}^k(f \mathbf{m}_\infty)(x) &:= \left[ \int_M f(y) d\mathbf{p}_x^k(y) \right] d\mathbf{m}_k(x), \\ d\mathbf{p}^\infty(g \mathbf{m}_k)(y) &:= \left[ \int_M g(x) d\mathbf{p}_y^\infty(x) \right] d\mathbf{m}_\infty(y). \end{aligned}$$

**Step 3.2.** Let  $\pi_\infty \in \Pi_{\ll}(\mu_{\infty,0}, \mu_{\infty,1})$  be timelike  $p$ -dualizing, and let  $(\pi_\infty^n)_{n \in \mathbf{N}}$  be an associated sequence in  $\mathcal{P}^{\text{ac}}(M^2, \mathbf{m}_\infty^{\otimes 2})$  as in Lemma 3.26, where we write

$$\pi_\infty^n = \rho_\infty^n \mathbf{m}_\infty^{\otimes 2},$$



$$\begin{aligned}\mu_{\infty,0}^n &:= (\text{proj}_1)_\# \pi_\infty^n = \rho_{\infty,0}^n \mathbf{m}_\infty, \\ \mu_{\infty,1}^n &:= (\text{proj}_2)_\# \pi_\infty^n = \rho_{\infty,1}^n \mathbf{m}_\infty.\end{aligned}$$

For  $k \in \mathbf{N}$ , define  $\mu_{k,0}^n, \mu_{k,1}^n \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  by

$$\begin{aligned}\mu_{k,0}^n &:= \mathfrak{p}^k(\mu_{\infty,0}^n) = \rho_{k,0}^n \mathbf{m}_k, \\ \mu_{k,1}^n &:= \mathfrak{p}^k(\mu_{\infty,1}^n) = \rho_{k,1}^n \mathbf{m}_k\end{aligned}$$

as well as  $\pi_k^n \in \mathcal{P}^{\text{ac}}(M^2, \mathbf{m}_k^{\otimes 2})$  by

$$d\pi_k^n(x^0, x^1) := \left[ \int_{M^2} \rho_\infty^n(y^0, y^1) d(\mathfrak{p}_{x^0}^k \otimes \mathfrak{p}_{x^1}^k)(y^0, y^1) \right] d\mathbf{m}_k^{\otimes 2}(x^0, x^1),$$

or in other words,

$$\pi_k^n = (\text{proj}_1, \text{proj}_3)_\# [(\rho_\infty^n \circ (\text{proj}_2, \text{proj}_4)) \mathbf{q}_k \otimes \mathbf{q}_k].$$

A straightforward computation entails  $\pi_k^n \in \Pi(\mu_{k,0}^n, \mu_{k,1}^n)$ . Also, for given  $n \in \mathbf{N}$ , weak convergence of  $(\mathbf{q}_k)_{k \in \mathbf{N}}$  to the unique  $W_2$ -optimal (diagonal) coupling of  $\mathbf{m}_\infty$  and  $\mathbf{m}_\infty$  by tightness [92, Lem. 4.3, Lem. 4.4], up to a nonreabeled subsequence, imply weak convergence of  $(\pi_k^n)_{k \in \mathbf{N}}$  to  $\pi_\infty^n$  up to a nonreabeled subsequence.

**Step 3.3.** By Lemma 3.26, a tightness and a diagonal argument, passage to a subsequence, and relabeling of indices, we find  $n: \mathbf{N} \rightarrow \mathbf{N}$  such that, setting

$$\tilde{\pi}_k := \pi_k^{n_k},$$

the sequence  $(\tilde{\pi}_k)_{k \in \mathbf{N}}$  converges weakly to  $\pi_\infty$ . By Portmanteau's theorem,

$$1 = \pi_\infty[M_{\ll}^2] \leq \liminf_{k \rightarrow \infty} \tilde{\pi}_k[M_{\ll}^2]. \quad (3.6)$$

Hence, up passing to a subsequence, we henceforth assume that  $\tilde{\pi}_k[M_{\ll}^2] > 0$  for every  $k \in \mathbf{N}$ . We write the marginals  $\tilde{\mu}_{k,0}, \tilde{\mu}_{k,1} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  of  $\tilde{\pi}_k$  as

$$\begin{aligned}\tilde{\mu}_{k,0} &= \tilde{\rho}_{k,0} \mathbf{m}_k = \rho_{k,0}^{n_k} \mathbf{m}_k, \\ \tilde{\mu}_{k,1} &= \tilde{\rho}_{k,1} \mathbf{m}_k = \rho_{k,1}^{n_k} \mathbf{m}_k.\end{aligned}$$

**Step 3.4.** Now we define  $\hat{\pi}_k \in \mathcal{P}^{\text{ac}}(M^2)$  by

$$\hat{\pi}_k := \tilde{\pi}_k[M_{\ll}^2]^{-1} \tilde{\pi}_k \llcorner M_{\ll}^2.$$

Note that  $(\hat{\pi}_k)_{k \in \mathbf{N}}$  converges weakly to  $\pi_\infty$ , whence

$$\lim_{k \rightarrow \infty} \int_{M^2} \tau^p d\hat{\pi}_k = \int_{M^2} \tau^p d\pi_\infty. \quad (3.7)$$

Let  $\hat{\mu}_{k,0}, \hat{\mu}_{k,1} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  denote the marginals of  $\hat{\pi}_k$ , with

$$\begin{aligned}\hat{\mu}_{k,0} &= \hat{\rho}_{k,0} \mathbf{m}_k, \\ \hat{\mu}_{k,1} &= \hat{\rho}_{k,1} \mathbf{m}_k.\end{aligned}$$

These measures are chronologically related, but we do not know whether they have a chronological  $\ell_p$ -optimal coupling, i.e. whether they are timelike  $p$ -dualizable. To achieve this, note that  $\ell_p(\hat{\mu}_{k,0}, \hat{\mu}_{k,1}) \in (0, \infty)$  by chronology of  $\hat{\pi}_k$  as well as compactness of  $M^2$ . Let  $\check{\pi}_k \in \Pi_{\leq}(\hat{\mu}_{k,0}, \hat{\mu}_{k,1})$  be an  $\ell_p$ -optimal coupling. By weak convergence of  $(\hat{\pi}_k)_{k \in \mathbf{N}}$ , its marginal sequences  $(\hat{\mu}_{k,0})_{k \in \mathbf{N}}$  and  $(\hat{\mu}_{k,1})_{k \in \mathbf{N}}$  are tight, and so is  $(\check{\pi}_k)_{k \in \mathbf{N}}$  [92, Lem. 4.4]. By Prokhorov's theorem, the latter converges weakly to some  $\check{\pi}_\infty \in \Pi(\mu_{\infty,0}, \mu_{\infty,1})$ ; causal closedness of  $M$  and Portmanteau's theorem imply  $\check{\pi}_\infty[M_{\leq}^2] = 1$ . Moreover, (3.7) gives

$$\begin{aligned}\ell_p(\mu_{\infty,0}, \mu_{\infty,1})^p &= \int_{M^2} \tau^p d\pi_\infty = \lim_{k \rightarrow \infty} \int_{M^2} \tau^p d\hat{\pi}_k \\ &\leq \lim_{k \rightarrow \infty} \int_{M^2} \tau^p d\check{\pi}_k = \int_{M^2} \tau^p d\check{\pi}_\infty.\end{aligned} \quad (3.8)$$

Here the inequality follows from  $\ell_p$ -optimality of  $\check{\pi}_k$ . Thus,  $\check{\pi}_\infty$  is an  $\ell_p$ -optimal coupling of  $\mu_{\infty,0}$  and  $\mu_{\infty,1}$ . Since the latter pair is strongly timelike  $p$ -dualizable, every such coupling is chronological, whence  $\check{\pi}_\infty[M_{\ll}^2] = 1$ ; thus,

$$1 = \check{\pi}_\infty[M_{\ll}^2] \leq \liminf_{k \rightarrow \infty} \check{\pi}_k[M_{\ll}^2].$$

As usual, we may and will assume  $\check{\pi}_k[M_{\ll}^2] > 0$  for every  $k \in \mathbf{N}$ .

**Step 3.5.** Lastly, we define  $\bar{\pi}_k \in \mathcal{P}(M^2)$  by

$$\bar{\pi}_k := \check{\pi}_k[M_{\ll}^2]^{-1} \check{\pi}_k \llcorner M_{\ll}^2.$$

Since the restriction of an  $\ell_p$ -optimal coupling is  $\ell_p$ -optimal [24, Lem. 2.10],  $\bar{\pi}_k$  is a chronological  $\ell_p$ -optimal coupling of its marginals  $\mu_{k,0}, \mu_{k,1} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m}_k)$  with

$$\begin{aligned} \mu_{k,0} &= \rho_{k,0} \mathfrak{m}_k, \\ \mu_{k,1} &= \rho_{k,1} \mathfrak{m}_k. \end{aligned}$$

The pair  $(\mu_{k,0}, \mu_{k,1})$  is timelike  $p$ -dualizable by  $\bar{\pi}_k$  for every  $k \in \mathbf{N}$ , as desired.

**Step 4.** *Invoking the TCD condition.* Fix  $K \in \mathbf{R}$  and  $N \in (1, \infty)$  with  $K < K_\infty$  and  $N > N_\infty$ . Then  $K < K_k$  and  $N > N_k$  for large enough  $k \in \mathbf{N}$ . Hence, again up to a subsequence, we may and will assume that the previous strict inequalities hold for every  $k \in \mathbf{N}$ , respectively.

By Proposition 3.7, for every  $k \in \mathbf{N}$  there are a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{k,t})_{t \in [0,1]}$  connecting  $\mu_{k,0}$  to  $\mu_{k,1}$  as well as a timelike  $p$ -dualizing coupling  $\pi_k \in \Pi_{\ll}(\mu_{k,0}, \mu_{k,1})$  such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\mathfrak{S}_{N'}^k(\mu_{k,t}) \leq \mathcal{J}_{K,N'}^{(t)}(\pi_k). \quad (3.9)$$

**Step 5.** *Estimating  $-\mathcal{J}_{K,N'}^{(t)}(\pi_k)$  from below.* Tracing back the construction from Step 2 and Step 3, we infer the inequalities

$$\begin{aligned} \rho_{k,0} &\leq \check{\pi}_k[M_{\ll}^2]^{-1} \bar{\pi}_k[M_{\ll}^2]^{-1} \tilde{\rho}_{k,0} \quad \mathfrak{m}\text{-a.e.}, \\ \rho_{k,1} &\leq \check{\pi}_k[M_{\ll}^2]^{-1} \bar{\pi}_k[M_{\ll}^2]^{-1} \tilde{\rho}_{k,1} \quad \mathfrak{m}\text{-a.e.} \end{aligned} \quad (3.10)$$

However, when using these estimates to bound  $-\mathcal{J}_{K,N'}^{(t)}(\pi_k)$  from below, we change the involved densities in the integrands, but not the coupling with respect to which it is integrated; we thus have to make up for this in the coupling as well. In turn, this requires us to take into account that both  $\bar{\pi}_k$  and  $\check{\pi}_k$  are not chronological, but only close to being so for large  $k \in \mathbf{N}$ .

**Step 5.1.** We modify  $\pi_k$  into a measure with marginals

$$\begin{aligned} \nu_{k,0} &:= (1 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (\tilde{\rho}_{k,0} + \delta_k + \varepsilon_k + \zeta_k) \mathfrak{m}_k = \varrho_{k,0} \mathfrak{m}_k, \\ \nu_{k,1} &:= (1 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (\tilde{\rho}_{k,1} + \delta_k + \varepsilon_k + \zeta_k) \mathfrak{m}_k = \varrho_{k,1} \mathfrak{m}_k, \end{aligned}$$

where  $\delta_k, \varepsilon_k \in [0, 1]$  and  $\zeta_k \geq 0$  are defined by

$$\begin{aligned} \delta_k &:= \bar{\pi}_k[\{\tau = 0\}], \\ \varepsilon_k &:= \check{\pi}_k[\{\tau = 0\}], \\ \zeta_k &:= W_2(\mathfrak{m}_k, \mathfrak{m}_\infty). \end{aligned} \quad (3.11)$$

Note that  $(\delta_k)_{k \in \mathbf{N}}$ ,  $(\varepsilon_k)_{k \in \mathbf{N}}$ , and  $(\zeta_k)_{k \in \mathbf{N}}$  converge to 0. Define  $\varpi_k \in \mathcal{P}(M^2)$  by

$$\begin{aligned} \varpi_k &:= (1 + \delta_k + \varepsilon_k + \zeta_k)^{-1} [\bar{\pi}_k[M_{\ll}^2] \check{\pi}_k[M_{\ll}^2] \pi_k + \bar{\pi}_k \llcorner \{\tau = 0\}] \\ &\quad + (\delta_k + \varepsilon_k + \zeta_k) \mathfrak{m}_k^{\otimes 2} + \bar{\pi}_k[M_{\ll}^2] \check{\pi}_k \llcorner \{\tau = 0\}. \end{aligned}$$

By tracing back the definitions of  $\bar{\pi}_k$  and  $\check{\pi}_k$  in Step 2, one verifies that  $\varpi_k$  is a coupling, not necessarily  $\ell_p$ -optimal, of  $\nu_{k,0}$  and  $\nu_{k,1}$ .

**Step 5.2.** By (3.10), we have

$$\rho_{k,0} \leq \check{\pi}_k[M_{\ll}^2]^{-1} \bar{\pi}_k[M_{\ll}^2]^{-1} (1 + \delta_k + \varepsilon_k + \zeta_k) \varrho_{k,0} \quad \mathfrak{m}\text{-a.e.},$$

$$\rho_{k,1} \leq \check{\pi}_k[M_{\ll}^2]^{-1} \tilde{\pi}_k[M_{\ll}^2]^{-1} (1 + \delta_k + \varepsilon_k + \zeta_k) \varrho_{k,1} \quad \mathbf{m}\text{-a.e.}$$

In the sequel, to clear up notation, given  $a, b \in \mathbf{R}$  we write  $a \geq_k b$  if there exists a sequence  $(c_k)_{k \in \mathbf{N}}$  of positive real numbers converging to 1 with  $a \geq c_k b$  for every  $k \in \mathbf{N}$ . By definition of the inherent distortion coefficients, we obtain

$$\begin{aligned} -\mathcal{J}_{K,N'}^{(t)}(\pi_k) &= \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \rho_{k,0}(x^0)^{-1/N'} d\pi_k(x^0, x^1) \\ &\quad + \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \rho_{k,1}(x^1)^{-1/N'} d\pi_k(x^0, x^1) \\ &\geq_k \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\pi_k(x^0, x^1) \\ &\quad + \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \varrho_{k,1}(x^1)^{-1/N'} d\pi_k(x^0, x^1) \\ &\geq_k \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &\quad + \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &\quad - (1-t) \int_{\{\tau=0\}} \varrho_{k,0}(x^0)^{-1/N'} d\tilde{\pi}_k(x^0, x^1) \\ &\quad - t \int_{\{\tau=0\}} \varrho_{k,1}(x^1)^{-1/N'} d\tilde{\pi}_k(x^0, x^1) \\ &\quad - (1-t) \int_{\{\tau=0\}} \varrho_{k,0}(x^0)^{-1/N'} d\check{\pi}_k(x^0, x^1) \\ &\quad - t \int_{\{\tau=0\}} \varrho_{k,1}(x^1)^{-1/N'} d\check{\pi}_k(x^0, x^1) \\ &\quad - c(\delta_k + \varepsilon_k + \zeta_k) \int_M \varrho_{k,0}(x^0)^{-1/N'} d\mathbf{m}_k(x^0) \\ &\quad - c(\delta_k + \varepsilon_k + \zeta_k) \int_M \varrho_{k,1}(x^1)^{-1/N'} d\mathbf{m}_k(x^1), \end{aligned}$$

where, recalling our choice  $K < K_k$  for large enough  $k \in \mathbf{N}$  and Corollary 3.14,

$$c := \max\{\sup \tau_{K,N'}^{(1-t)} \circ \tau(M^2), \sup \tau_{K,N'}^{(t)} \circ \tau(M^2)\}. \quad (3.12)$$

Since  $\varrho_{k,0} \geq_k \delta_k + \varepsilon_k + \zeta_k$   $\mathbf{m}_k$ -a.e. and  $\varrho_{k,1} \geq_k \delta_k + \varepsilon_k + \zeta_k$   $\mathbf{m}_k$ -a.e., by definition of  $\delta_k$  and  $\varepsilon_k$  we obtain the estimates

$$\begin{aligned} (\delta_k + \varepsilon_k + \zeta_k)^{1-1/N'} &\geq_k (1-t) \int_{\{\tau=0\}} \varrho_{k,1}(x^0)^{-1/N'} d\tilde{\pi}_k(x^0, x^1) \\ &\quad + t \int_{\{\tau=0\}} \varrho_{k,1}(x^0)^{-1/N'} d\tilde{\pi}_k(x^0, x^1), \\ (\delta_k + \varepsilon_k + \zeta_k)^{1-1/N'} &\geq_k (1-t) \int_{\{\tau=0\}} \varrho_{k,1}(x^0)^{-1/N'} d\check{\pi}_k(x^0, x^1) \\ &\quad + t \int_{\{\tau=0\}} \varrho_{k,1}(x^0)^{-1/N'} d\check{\pi}_k(x^0, x^1), \\ 2(\delta_k + \varepsilon_k + \zeta_k)^{1-1/N'} &\geq_k (\delta_k + \varepsilon_k + \zeta_k) \int_M \varrho_{k,0}(x^0)^{-1/N'} d\mathbf{m}_k(x^0) \\ &\quad + (\delta_k + \varepsilon_k + \zeta_k) \int_M \varrho_{k,1}(x^1)^{-1/N'} d\mathbf{m}_k(x^1), \end{aligned}$$

and therefore

$$\begin{aligned} -\mathcal{J}_{K,N'}^{(t)}(\pi_k) &\geq_k \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &\quad + \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &\quad - 2(c+1)(\delta_k + \varepsilon_k + \zeta_k)^{1-1/N'}. \end{aligned}$$

**Step 5.3.** Let  $\mathfrak{r}^0, \mathfrak{r}^1: M \rightarrow \mathcal{P}(M)$  denote the ( $k$ -dependent) disintegrations of  $\varpi_k$  with respect to  $\text{proj}_1$  and  $\text{proj}_2$ , respectively, i.e.

$$d\varpi_k(x^0, x^1) = d\mathfrak{r}_{x^0}^0(x^1) d\nu_{k,0}(x^0) = d\mathfrak{r}_{x^1}^1(x^0) d\nu_{k,1}(x^1),$$

and define

$$\begin{aligned} v_0(x^0) &:= \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) d\mathfrak{r}_{x^0}^0(x^1), \\ v_1(x^1) &:= \int_M \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) d\mathfrak{r}_{x^1}^1(x^0). \end{aligned} \tag{3.13}$$

Moreover, define  $\nu_{\infty,0}^k, \nu_{\infty,1}^k \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m}_\infty)$  by

$$\begin{aligned} \nu_{\infty,0}^k &:= (1 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (\rho_{\infty,0}^{n_k} + \delta_k + \varepsilon_k + \zeta_k) \mathfrak{m}_\infty = \varrho_{\infty,0}^k \mathfrak{m}_\infty, \\ \nu_{\infty,1}^k &:= (1 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (\rho_{\infty,1}^{n_k} + \delta_k + \varepsilon_k + \zeta_k) \mathfrak{m}_\infty = \varrho_{\infty,1}^k \mathfrak{m}_\infty, \end{aligned}$$

and observe that

$$\begin{aligned} \mathfrak{p}^k(\nu_{\infty,0}^k) &= \nu_{k,0} = \varrho_{k,0} \mathfrak{m}_k, \\ \mathfrak{p}^k(\nu_{\infty,1}^k) &= \nu_{k,1} = \varrho_{k,1} \mathfrak{m}_k. \end{aligned} \tag{3.14}$$

Then by Jensen's inequality,

$$\begin{aligned} &\int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &\quad + \int_{M^2} \tau_{K,N'}^{(t)}(\tau(x^0, x^1)) \varrho_{k,0}(x^0)^{-1/N'} d\varpi_k(x^0, x^1) \\ &= \sum_{i=0}^1 \int_M \varrho_{k,i}(x^i)^{1-1/N'} v_i(x^i) d\mathfrak{m}_k(x^i) \\ &= \sum_{i=0}^1 \int_M \left[ \int_M \varrho_{\infty,i}^k(y^i) d\mathfrak{p}_{x^i}^k(y^i) \right]^{1-1/N'} v_i(x^i) d\mathfrak{m}_k(x^i) \\ &\geq \sum_{i=0}^1 \int_{M^2} \varrho_{\infty,i}^k(y^i)^{1-1/N'} v_i(x^i) d\mathfrak{p}_{x^i}^k(y^i) d\mathfrak{m}_k(x^i) \\ &= \sum_{i=0}^1 \int_{M^2} \varrho_{\infty,i}^k(y^i)^{1-1/N'} v_i(x^i) d\mathfrak{p}_{y^i}^\infty(x^i) d\mathfrak{m}_\infty(y^i). \end{aligned}$$

**Step 5.4.** Next, we estimate the latter sum from below. Using (3.13) and (3.14),

$$\begin{aligned} &\int_{M^2} \varrho_{\infty,0}^k(y^0)^{1-1/N'} v_0(x^0) d\mathfrak{p}_{y^0}^\infty(x^0) d\mathfrak{m}_\infty(y^0) \\ &= \int_{M^4} d\mathfrak{p}_{x^1}^k(y^1) d\mathfrak{r}_{x^0}^0(x^1) d\mathfrak{p}_{y^0}^\infty(x^0) d\mathfrak{m}_\infty(y^0) \\ &\quad \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right]. \end{aligned}$$

Let  $\varepsilon > 0$ . Since  $\tau_{K,N'}^{(t)} \circ \tau$  is uniformly continuous by compactness of  $M^2$ , our choice of  $K$ , and possibly invoking Corollary 3.14, we fix  $\delta > 0$  such that

$$\begin{aligned} |\tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) - \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1))| &\leq \varepsilon, \\ |\tau_{K,N'}^{(t)}(\tau(x^0, x^1)) - \tau_{K,N'}^{(t)}(\tau(y^0, y^1))| &\leq \varepsilon \end{aligned}$$

for every  $(x^0, y^0, x^1, y^1) \in A_\delta$ , where

$$A_\delta := \{z \in M^4 : d(z_1, z_2) + d(z_3, z_4) \leq \delta\}. \quad (3.15)$$

Lastly, somewhat suggestively, for  $k \in \mathbf{N}$  we define a coupling  $\eta_k \in \Pi(\nu_{\infty,0}^k, \nu_{\infty,1}^k)$ , which is notably independent of  $\varepsilon$ , by

$$\begin{aligned} d\eta_k(y^0, y^1) &:= \int_{M^2} \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,0}(x^0) \varrho_{k,1}(x^1)} d\mathbf{p}_{x^0}^k(y^0) d\mathbf{p}_{x^1}^k(y^1) d\varpi_k(x^0, x^1) \\ &= \int_{M^2} \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} d\mathbf{p}_{x^1}^k(y^1) d\mathbf{r}_{x^0}^0(x^1) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \\ &= \int_{M^2} \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,0}(x^0)} d\mathbf{p}_{x^0}^k(y^0) d\mathbf{r}_{x^1}^1(x^0) d\mathbf{p}_{y^1}^\infty(x^1) d\mathbf{m}_\infty(y^1). \end{aligned}$$

The previous computations then entail

$$\begin{aligned} &\int_{M^2} \varrho_{\infty,0}^k(y^0)^{1-1/N'} v_0(x^0) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \\ &\geq -\varepsilon + \int_{A_\delta} d\mathbf{p}_{x^1}^k(y^1) d\mathbf{r}_{x^0}^0(x^1) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \\ &\quad \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right] \\ &\quad + \int_{A_\delta^c} d\mathbf{p}_{x^1}^k(y^1) d\mathbf{r}_{x^0}^0(x^1) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \\ &\quad \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right] \\ &= -\varepsilon + \int_{M^2} \varrho_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) d\eta_k(y^0, y^1) \\ &\quad + \int_{A_\delta^c} d\mathbf{p}_{x^1}^k(y^1) d\mathbf{r}_{x^0}^0(x^1) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \\ &\quad \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} [\tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) - \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1))] \right. \\ &\quad \left. \times \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right]. \end{aligned}$$

**Step 5.5.** Let us estimate the latter error term. We first recall the definition  $\zeta_k := W_2(\mathbf{m}_k, \mathbf{m}_\infty)$  from (3.11). By definition of  $A_\delta$  and the value  $c$  from (3.12), the  $\mathbf{m}$ -a.e. valid inequality for the density  $\rho_{\infty,1}^k$  involved in the definition of  $\varrho_{\infty,1}^k$  from Lemma 3.26, as well as Hölder's inequality,

$$\begin{aligned} &\left| \int_{A_\delta^c} d\mathbf{p}_{x^1}^k(y^1) d\mathbf{r}_{x^0}^0(x^1) d\mathbf{p}_{y^0}^\infty(x^0) d\mathbf{m}_\infty(y^0) \right. \\ &\quad \left. \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} [\tau_{K,N'}^{(1-t)}(\tau(x^0, x^1)) - \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1))] \right. \right. \\ &\quad \left. \left. \times \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right] \right| \end{aligned}$$

$$\begin{aligned}
&\leq 2c\delta^{-1} \int_{M^4} \mathrm{d}\mathbf{p}_{x^1}^k(y^1) \mathrm{d}\tau_{x^0}^0(x^1) \mathrm{d}\mathbf{p}_{y^0}^\infty(x^0) \mathrm{d}\mathbf{m}_\infty(y^0) \\
&\quad \left[ \varrho_{\infty,0}^k(y^0)^{-1/N'} [\mathbf{d}(x^0, y^0) + \mathbf{d}(x^1, y^1)] \frac{\varrho_{\infty,0}^k(y^0) \varrho_{\infty,1}^k(y^1)}{\varrho_{k,1}(x^1)} \right] \\
&\leq_k 2c\delta^{-1} \int_{M^2} \varrho_{\infty,0}^k(y^0)^{1-1/N'} \mathbf{d}(x^0, y^0) \mathrm{d}\mathbf{q}_k(x^0, y^0) \\
&\quad + 2c\delta^{-1} \zeta_k^{-1/N'} \int_M \rho_{\infty,1}(y^1) \mathbf{d}(x^1, y^1) \mathrm{d}\mathbf{q}_k(x^1, y^1) \\
&\leq_k 2c\delta^{-1} \|\rho_{\infty,0}\|_{L^\infty(M, \mathbf{m}_\infty)}^{1-1/N'} \zeta_k + 2c\delta^{-1} \|\rho_{\infty,1}\|_{L^\infty(M, \mathbf{m}_\infty)} \zeta_k^{1-1/N'}.
\end{aligned}$$

**Step 5.6.** Taking the estimates from Step 5.4 and Step 5.5 together, we get

$$\begin{aligned}
&\int_{M^2} \varrho_{\infty,0}^k(y^0)^{1-1/N'} v_0(x^0) \mathrm{d}\mathbf{p}_{y^0}^\infty(x^0) \mathrm{d}\mathbf{m}_\infty(y^0) \\
&\geq_k -\varepsilon + \int_{M^2} \varrho_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) \mathrm{d}\eta_k(y^0, y^1) \\
&\quad - 2c\delta^{-1} \|\rho_{\infty,0}\|_{L^\infty(M, \mathbf{m}_\infty)}^{1-1/N'} \zeta_k \\
&\quad - 2c\delta^{-1/N'} \|\rho_{\infty,1}\|_{L^\infty(M, \mathbf{m}_\infty)} \zeta_k^{1-1/N'}.
\end{aligned}$$

Analogously, for the second summand in the last part of Step 5.3 we obtain

$$\begin{aligned}
&\int_{M^2} \varrho_{\infty,1}^k(y^1)^{1-1/N'} v_1(x^1) \mathrm{d}\mathbf{p}_{y^1}^\infty(x^1) \mathrm{d}\mathbf{m}_\infty(y^1) \\
&\geq_k -\varepsilon + \int_{M^2} \varrho_{\infty,1}^k(y^1)^{-1/N'} \tau_{K,N'}^{(t)}(\tau(y^0, y^1)) \mathrm{d}\eta_k(y^0, y^1) \\
&\quad - 2c\delta^{-1} \|\rho_{\infty,1}\|_{L^\infty(M, \mathbf{m}_\infty)}^{1-1/N'} \zeta_k \\
&\quad - 2c\delta^{-1} \|\rho_{\infty,0}\|_{L^\infty(M, \mathbf{m}_\infty)} \zeta_k^{1-1/N'}.
\end{aligned}$$

**Step 6. Passage to the limit.** Let  $(a_k)_{k \in \mathbf{N}}$  be a given sequence of normalization constants in  $[1, \infty)$  as provided by Lemma 3.26, i.e.

$$\begin{aligned}
\rho_{\infty,0}^{n_k} &\leq a_k \rho_{\infty,0} \quad \mathbf{m}\text{-a.e.}, \\
\rho_{\infty,1}^{n_k} &\leq a_k \rho_{\infty,1} \quad \mathbf{m}\text{-a.e.}
\end{aligned}$$

**Step 6.1.** For  $k \in \mathbf{N}$ , we define  $\tilde{\nu}_{\infty,0}^k, \tilde{\nu}_{\infty,1}^k \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty)$  by

$$\begin{aligned}
\tilde{\nu}_{\infty,0}^k &:= (a_k^2 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (a_k^2 \rho_{\infty,0} + \delta_k + \varepsilon_k + \zeta_k) \mathbf{m}_\infty = \tilde{\varrho}_{\infty,0}^k \mathbf{m}_\infty, \\
\tilde{\nu}_{\infty,1}^k &:= (a_k^2 + \delta_k + \varepsilon_k + \zeta_k)^{-1} (a_k^2 \rho_{\infty,1} + \delta_k + \varepsilon_k + \zeta_k) \mathbf{m}_\infty = \tilde{\varrho}_{\infty,1}^k \mathbf{m}_\infty.
\end{aligned}$$

We turn  $\eta_k$  into a coupling  $\alpha_k \in \Pi(\nu_{\infty,0}^k, \nu_{\infty,1}^k)$  by setting

$$\begin{aligned}
\alpha_k &:= (a_k^2 + \delta_k + \varepsilon_k + \zeta_k)^{-1} [(1 + \delta_k + \varepsilon_k + \zeta_k) \eta_k \\
&\quad + a_k^2 \mu_{\infty,0} \otimes \mu_{\infty,1} - \tilde{\mu}_{k,0} \otimes \tilde{\mu}_{k,1}].
\end{aligned}$$

This construction yields

$$\begin{aligned}
&\int_{M^2} \varrho_{k,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) \mathrm{d}\eta_k(y^0, y^1) \\
&\quad + \int_{M^2} \varrho_{k,1}^k(y^1)^{-1/N'} \tau_{K,N'}^{(t)}(\tau(y^0, y^1)) \mathrm{d}\eta_k(y^0, y^1) \\
&\geq_k \int_{M^2} \tilde{\varrho}_{\infty,0}^k(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) \mathrm{d}\alpha_k(y^0, y^1)
\end{aligned}$$

$$\begin{aligned}
& -c \int_M \varrho_{\infty,0}^k(y^0)^{-1/N'} |\rho_{\infty,0}(y^0) - \tilde{\rho}_{k,0}(y^0)| \, d\mathbf{m}_{\infty}(y^0) \\
& + \int_{M^2} \tilde{\varrho}_{\infty,1}^k(y^1)^{-1/N'} \tau_{K,N'}^{(t)}(\tau(y^0, y^1)) \, d\alpha_k(y^0, y^1) \\
& -c \int_M \varrho_{\infty,1}^k(y^1)^{-1/N'} |\rho_{\infty,1}(y^1) - \tilde{\rho}_{k,1}(y^1)| \, d\mathbf{m}_{\infty}(y^1).
\end{aligned}$$

**Step 6.2.** Taking together the above estimates with those obtained in Step 5 and invoking the  $L^1$ -convergence asserted in Lemma 3.26, we obtain

$$-\limsup_{k \rightarrow \infty} \mathcal{J}_{K,N'}^{(t)}(\pi_k) \geq -\limsup_{k \rightarrow \infty} \mathcal{J}_{K,N'}^{(t)}(\alpha_k) - 2\varepsilon.$$

Now, by Prokhorov's theorem, the sequence  $(\alpha_k)_{k \in \mathbf{N}}$  converges weakly to some  $\alpha \in \Pi(\mu_{\infty,0}, \mu_{\infty,1})$  up to a nonrelabeled subsequence. Since  $\varepsilon$  was arbitrary and did not influence the construction of  $\alpha_k$ , by Lemma 3.27 we thus get

$$-\limsup_{k \rightarrow \infty} \mathcal{J}_{K,N'}^{(t)}(\pi_k) \geq -\mathcal{J}_{K,N'}^{(t)}(\alpha). \quad (3.16)$$

**Step 6.3.** Now we send  $k \rightarrow \infty$  in (3.9). For every  $k \in \mathbf{N}$ , there exists some plan  $\pi_k \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_{k,0}, \mu_{k,1})$  representing the curve  $(\mu_{k,t})_{t \in [0,1]}$ . Proposition B.9 allows us to extract a nonrelabeled subsequence of  $(\pi_k)_{k \in \mathbf{N}}$  converging weakly to some  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_{\infty,0}, \mu_{\infty,1})$ . The assignment  $\mu_{\infty,t} := (\mathbf{e}_t)_{\#} \pi_{\infty}$  produces a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{\infty,t})_{t \in [0,1]}$  connecting  $\mu_{\infty,0}$  to  $\mu_{\infty,1}$ . Joint lower semicontinuity of Rényi's entropy on  $\mathcal{P}(M)$  by compactness of  $M$  and (3.16) thus yield, for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\mathcal{S}_{N'}^{\infty}(\mu_{\infty,t}) \leq \limsup_{k \rightarrow \infty} \mathcal{S}_{N'}^k(\mu_{k,t}) \leq \limsup_{k \rightarrow \infty} \mathcal{J}_{K,N'}^{(t)}(\pi_k) \leq \mathcal{J}_{K,N'}^{(t)}(\alpha). \quad (3.17)$$

**Step 7.** *Proof of the  $\ell_p$ -optimality of  $\alpha$ .* To conclude the desired property leading towards  $\text{TCD}_p(K, N)$  for  $\mathcal{X}_{\infty}$ , at least under the restrictions of Step 2, we have to prove the causality and the  $\ell_p$ -optimality of  $\alpha$ . Both is argued similarly to Step 5.4; we concentrate on the proof of the estimate entailing  $\ell_p$ -optimality later, and then outline how to modify this argument to prove  $\alpha[M_{\ll}^2] = 1$ .

Given any  $\varepsilon > 0$ , fix  $\delta > 0$  with the property

$$|\tau(x^0, x^1) - \tau(y^0, y^1)| \leq \varepsilon$$

for every  $(x^0, y^0, x^1, y^1) \in M^4$  belonging to the set  $A_{\delta}$  from (3.15). As in Step 5.5 and tracing back the definitions of all considered couplings, we obtain

$$\begin{aligned}
\int_{M^2} \tau^p(y^0, y^1) \, d\alpha(y^0, y^1) &= \lim_{k \rightarrow \infty} \int_{M^2} \tau^p(y^0, y^1) \, d\alpha_k(y^0, y^1) \\
&\geq \liminf_{k \rightarrow \infty} \int_{M^2} \tau^p(y^0, y^1) \, d\eta_k(y^0, y^1) \\
&\geq \liminf_{k \rightarrow \infty} \int_{M^2} \tau^p(x^0, x^1) \, d\varpi_k(x^0, x^1) - \varepsilon \\
&\quad - 2c\delta^{-1} \|\rho_{\infty,0}\|_{L^{\infty}(M, \mathbf{m}_{\infty})} \limsup_{k \rightarrow \infty} W_2(\mathbf{m}_k, \mathbf{m}_{\infty}) \\
&\quad - 2c\delta^{-1} \|\rho_{\infty,1}\|_{L^{\infty}(M, \mathbf{m}_{\infty})} \limsup_{k \rightarrow \infty} W_2(\mathbf{m}_k, \mathbf{m}_{\infty}) \\
&\geq \liminf_{k \rightarrow \infty} \int_{M^2} \tau^p(x^0, x^1) \, d\pi_k(x^0, x^1) - \varepsilon \\
&= \liminf_{k \rightarrow \infty} \int_{M^2} \tau^p(x^0, x^1) \, d\bar{\pi}_k(x^0, x^1) - \varepsilon
\end{aligned}$$

$$\geq \liminf_{k \rightarrow \infty} \int_{M^2} \tau^p(x^0, x^1) d\tilde{\pi}_k(x^0, x^1) - \varepsilon \geq \ell_p(\mu_{\infty,0}, \mu_{\infty,1})^p - \varepsilon.$$

In the equality in the third last step, we have used that both  $\pi_k, \bar{\pi}_k \in \Pi_{\ll}(\mu_{k,0}, \mu_{k,1})$  are  $\ell_p$ -optimal; the last inequality is due to (3.8).

The relation  $\alpha[M_{\ll}^2] = 1$  is argued similarly by replacing, given  $\varepsilon > 0$ ,  $\tau$  by a nonnegative function  $\phi \in C(M^2)$  obeying  $\phi(M_{\ll}^2) = \{1\}$ ,  $\sup \phi(M) \leq 1$ , and

$$\alpha[M_{\ll}^2] \geq \int_{M^2} \phi d\alpha - \varepsilon.$$

Both results together and the arbitrariness of  $\varepsilon$  yield the claim.

**Step 8.** *Relaxation of the assumptions on  $\mu_{\infty,0}$  and  $\mu_{\infty,1}$ .* Now we outline how to get rid of the assumption  $\rho_{\infty,0}, \rho_{\infty,1} \in L^\infty(M, \mathfrak{m}_\infty)$  from Step 2.

If  $\rho_{\infty,0}$  and  $\rho_{\infty,1}$  are not  $\mathfrak{m}$ -essentially bounded, given any  $i \in \mathbf{N}$  we set

$$E_i := \{\rho_{\infty,0} \leq i, \rho_{\infty,1} \leq i\}.$$

Let  $\pi \in \Pi_{\ll}(\mu_{\infty,0}, \mu_{\infty,1})$  be an  $\ell_p$ -optimal coupling, and set

$$\pi_i := \pi[E_i]^{-1} \pi \llcorner E_i$$

provided  $\pi[E_i] > 0$ . By restriction [24, Lem. 2.10],  $\pi_i$  is an  $\ell_p$ -optimal coupling of its marginals  $\lambda_{\infty,0}^i, \lambda_{\infty,1}^i \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m}_\infty)$ ; moreover, the pair  $(\lambda_{\infty,0}^i, \lambda_{\infty,1}^i)$  is strongly timelike  $p$ -dualizable. Let  $\tilde{\pi}_i \in \Pi_{\ll}(\lambda_{\infty,0}^i, \lambda_{\infty,1}^i)$  be an  $\ell_p$ -optimal coupling as constructed in the previous steps. Define  $\beta_i \in \Pi_{\ll}(\mu_{\infty,0}, \mu_{\infty,1})$  by

$$\beta_i := \pi[E_i] \tilde{\pi}_i + \pi \llcorner E_i^c,$$

and note that  $\beta_i$  is an  $\ell_p$ -optimal coupling of its marginals. Moreover, we observe that, for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{J}_{K,N'}^{(t)}(\tilde{\pi}_i) &\leq i - \int_{M^2} \rho_{\infty,0}(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) d\tilde{\pi}_i(y^0, y^1) \\ &\quad - \int_{M^2} \rho_{\infty,1}(y^1)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) d\tilde{\pi}_i(y^0, y^1) \\ &\leq i - \int_{M^2} \rho_{\infty,0}(y^0)^{-1/N'} \tau_{K,N'}^{(1-t)}(\tau(y^0, y^1)) d\beta_i(y^0, y^1) \\ &\quad + c \int_{E_i^c} \rho_{\infty,0}(y^0)^{-1/N'} d\pi(y^0, y^1) \\ &\quad - \int_{M^2} \rho_{\infty,1}(y^1)^{-1/N'} \tau_{K,N'}^{(t)}(\tau(y^0, y^1)) d\beta_i(y^0, y^1) \\ &\quad + c \int_{E_i^c} \rho_{\infty,1}(y^1)^{-1/N'} d\pi(y^0, y^1). \end{aligned}$$

By Prokhorov's theorem,  $(\beta_i)_{i \in \mathbf{N}}$  converges weakly to some  $\beta \in \Pi(\mu_{\infty,0}, \mu_{\infty,1})$  which, thanks to Portmanteau's theorem, is a causal coupling. By stability [24, Thm. 2.14], we thus infer the  $\ell_p$ -optimality of  $\beta$ . Together with Lemma 3.27 and Hölder's inequality, this addresses the right-hand side of the desired inequality; the left-hand side is treated with the compactness argument from Step 6.3.

**Step 9.** *Passing from  $K$  and  $N$  to  $K_\infty$  and  $N_\infty$ .* Lastly, the restrictions from Step 4 the previous arguments imply  $\text{TCD}_p(K, N)$  for  $\mathcal{X}_\infty$  for every  $K < K_\infty$  and every  $N > N_\infty$ . Given  $(\mu_{\infty,0}, \mu_{\infty,1}) \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m}_\infty)$  strongly timelike  $p$ -dualizable, letting  $K$  and  $N$  approach  $K_\infty$  and  $N_\infty$ , respectively, and using compactness arguments as in Step 6.2 and Step 6.3 gives the existence of a proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{\infty,t})_{t \in [0,1]}$  from  $\mu_{\infty,0}$  to  $\mu_{\infty,1}$  and a timelike  $p$ -dualizing coupling



$\pi \in \Pi_{\ll}(\mu_{\infty,0}, \mu_{\infty,1})$  witnessing the desired inequality for the Rényi entropy for every  $K < K_{\infty}$  and every  $N' > N_{\infty}$ . The semicontinuity properties of both sides in the respective parameters yields  $\text{TCD}_p(K_{\infty}, N_{\infty})$ . The proof is terminated.  $\square$

*Remark 3.30.* Very roughly speaking, the relevant convergences in the above proof are justified by *uniform continuity*, recall e.g. Step 5.4. This is considerably weaker than *Lipschitz continuity* of the quantity inside the distortion coefficients, which has been used in the proof of [86, Thm. 3.1] in the metric case at a similar point, yet is useless here since  $\tau$  is no distance. Our proof thus has the potential to admit straightforward adaptations to settings with continuous potentials other than  $\tau$  which do not come from a distance either.

*Remark 3.31.* A remarkable byproduct of the above proof is that both  $\text{TCD}_p^*(K, N)$  and  $\text{TCD}_p(K, N)$  are weakly stable under convergence of *normalized* Lorentzian pre-length spaces with respect to the following Lorentzian modification of Sturm's transport distance  $\mathbf{D}$  [85, Def. 3.2]: it is still defined in terms of  $\mathbf{d}$ , but the inherent metric measure isometric embeddings should also respect the given Lorentzian structures, and are required to map into spaces obeying the regularity assumptions from item **a.** of Definition 3.24.

**3.4. Good geodesics.** In this section, following [16], see also [23, 75, 76], we show the existence of timelike proper-time parametrized  $\ell_p$ -geodesics with  $\mathbf{m}$ -densities uniformly in  $L^{\infty}$  in time under mild assumptions on their endpoints. This treatise does not require any nonbranching condition on the underlying space. The proofs of the results to follow are mostly analogous to those of [16] and hence only outlined.

We regard the corresponding result from Theorem 3.33 as a key to develop the notion of *timelike weak gradients* on Lorentzian geodesic spaces by using so-called *test plans* — many of which should exist by Theorem 3.33 — following [5], and to prove a Lorentzian analogue of the *Sobolev-to-Lipschitz property* [42, Subsec. 4.1.3].

The crucial result providing the critical threshold for (3.18) is the following.

**Lemma 3.32.** *Assume  $\text{wTCD}_p^*(K, N)$  for  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Let  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  be strongly timelike  $p$ -dualizable with  $\rho_0, \rho_1 \in L^{\infty}(M, \mathbf{m})$ . Let  $D$  be any real number no smaller than  $\sup \tau(\text{spt } \mu_0 \times \text{spt } \mu_1)$ . Then there exists a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1$  such that  $\mu_t = \rho_t \mathbf{m} \in \mathcal{D}(\text{Ent}_{\mathbf{m}})$  for every  $t \in [0, 1]$ , and*

$$\mathbf{m}[\{\rho_{1/2} > 0\}] \geq e^{-D\sqrt{K-N}/2} \max\{\|\rho_0\|_{L^{\infty}(M, \mathbf{m})}, \|\rho_1\|_{L^{\infty}(M, \mathbf{m})}\}^{-1}.$$

*Proof.* Recall that  $\text{wTCD}_p^*(K, N)$  implies  $\text{wTCD}_p^*(K^-, N)$  by Proposition 3.7. Let  $(\mu_t)_{t \in [0,1]}$  be a timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0$  to  $\mu_1$  and  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  be some  $\ell_p$ -optimal coupling obeying the inequality defining the condition  $\text{wTCD}_p^*(K^-, N)$  as in Definition 3.2. Thanks to Lemma 3.8, we obtain  $\mu_t \in \mathcal{D}(\text{Ent}_{\mathbf{m}})$  for every  $t \in [0, 1]$ ; in particular, we have  $\mu_t = \rho_t \mathbf{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$ . Moreover, by Remark 2.2,

$$\begin{aligned} \mathcal{S}_N(\mu_{1/2}) &\leq - \int_{M^2} \sigma_{K^-, N}^{(1/2)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N} d\pi(x^0, x^1) \\ &\quad - \int_{M^2} \sigma_{K^-, N}^{(1/2)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N} d\pi(x^0, x^1) \\ &\leq -e^{-D\sqrt{K^-/N}/2} \max\{\|\rho_0\|_{L^{\infty}(M, \mathbf{m})}, \|\rho_1\|_{L^{\infty}(M, \mathbf{m})}\}^{-1/N}. \end{aligned}$$

On the other hand, we have  $\mathcal{S}_N(\mu_{1/2}) \geq -\mathbf{m}[\{\rho_{1/2} > 0\}]^{1/N}$  by Jensen's inequality, and rearranging terms provides the claim.  $\square$

**Theorem 3.33.** *Under the hypotheses of Lemma 3.32, there exists a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1$  such that for every  $t \in [0, 1]$ ,  $\mu_t = \rho_t \mathbf{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  and*

$$\|\rho_t\|_{L^\infty(M, \mathbf{m})} \leq e^{D\sqrt{K-N}/2} \max\{\|\rho_0\|_{L^\infty(M, \mathbf{m})}, \|\rho_1\|_{L^\infty(M, \mathbf{m})}\}, \quad (3.18)$$

where

$$D := \sup \tau(\text{spt } \mu_0 \times \text{spt } \mu_1).$$

*Proof.* We construct the required geodesic at dyadic times in  $[0, 1]$  by a bisection argument and then define the rest of the geodesic by completion. By induction, given  $n \in \mathbf{N}_0$  we assume that measures  $\alpha^0, \dots, \alpha^n \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  have already been constructed with the following properties. For every  $k \in \{0, \dots, 2^n\}$ , the pair  $((\mathbf{e}_{(k-1)2^{-n}})_\# \alpha^n, (\mathbf{e}_{k2^{-n}})_\# \alpha^n) \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})^2$  is strongly timelike  $p$ -dualizable, and

$$\sup \tau(\text{spt } (\mathbf{e}_{(k-1)2^{-n}})_\# \alpha^n \times \text{spt } (\mathbf{e}_{k2^{-n}})_\# \alpha^n) \leq 2^{-n} D.$$

**Step 1.** *Minimization of an appropriate functional.* Set

$$c_{n+1} := e^{2^{-n-2}D\sqrt{K-N}} \max\{\|\rho_0\|_{L^\infty(M, \mathbf{m})}, \|\rho_1\|_{L^\infty(M, \mathbf{m})}\}$$

and define the functional  $\mathcal{F}_{c_{n+1}}: \mathcal{P}(M) \rightarrow [0, 1]$  by

$$\mathcal{F}_{c_{n+1}}(\mu) := \|(\rho - c_{n+1})^+\|_{L^1(M, \mathbf{m})} + \mu_\perp[M]. \quad (3.19)$$

subject to the decomposition  $\mu = \rho \mathbf{m} + \mu_\perp$ . It measures how far  $\rho$  deviates from being bounded from above by  $c$ , and how much the input  $\mu$  fails to be  $\mathbf{m}$ -absolutely continuous. Let  $k \in \{1, \dots, 2^{n+1} - 1\}$  be odd. As for [16, Lem. 3.13], we prove that  $\mathcal{F}_{c_{n+1}} \circ \mathbf{e}_{1/2}$  admits a minimizer  $\pi_k^{n+1} \in \text{OptTGeo}_{\ell_p}^\tau(\mu_{(k-1)2^{-n-1}}, \mu_{(k+1)2^{-n-1}})$ . By gluing, we find a measure  $\alpha^{n+1} \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  such that

$$(\text{restr}_{(k-1)2^{-n-1}}^{(k+1)2^{-n-1}})_\# \alpha^{n+1} = \pi_k^{n+1}$$

for every  $k$  as above. Here, given  $s, t \in [0, 1]$  with  $s < t$ ,  $\text{restr}_s^t$  is defined in (2.11).

For every odd  $k \in \{1, \dots, 2^{n+1} - 1\}$ , the pairs

$$\begin{aligned} &((\mathbf{e}_{(k-1)2^{-n-1}})_\# \alpha^{n+1}, (\mathbf{e}_{k2^{-n-1}})_\# \alpha^{n+1}), \\ &((\mathbf{e}_{k2^{-n-1}})_\# \alpha^{n+1}, (\mathbf{e}_{(k+1)2^{-n-1}})_\# \alpha^{n+1}) \end{aligned}$$

are both strongly timelike  $p$ -dualizable [16, Lem. 3.1]. Furthermore, since  $\pi_k^{n+1}$  is concentrated on curves in  $\text{TGeo}^\tau(M)$  with  $\tau$ -length at most  $2^{-n} D$ , we obtain

$$\begin{aligned} \sup \tau(\text{spt } (\mathbf{e}_{(k-1)2^{-n-1}})_\# \alpha^{n+1} \times \text{spt } (\mathbf{e}_{k2^{-n-1}})_\# \alpha^{n+1}) &\leq 2^{-n-1} D, \\ \sup \tau(\text{spt } (\mathbf{e}_{k2^{-n-1}})_\# \alpha^{n+1} \times \text{spt } (\mathbf{e}_{(k+1)2^{-n-1}})_\# \alpha^{n+1}) &\leq 2^{-n-1} D. \end{aligned}$$

After Step 2, this will allow us to iteratively produce a sequence  $(\alpha^n)_{n \in \mathbf{N}}$  in  $\text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  according to the preceding scheme.

**Step 2.** *Uniform density bound.* Given any  $n \in \mathbf{N}_0$ , any  $c' > c_{n+1}$ , and any odd  $k \in \{1, \dots, 2^{n+1} - 1\}$ , arguing as in [16, Prop. 3.14] and using Lemma 3.32 we get

$$\begin{aligned} \inf\{\mathcal{F}_{c'}((\mathbf{e}_{1/2})_\# \pi) : \pi \in \text{OptTGeo}_{\ell_p}^\tau((\mathbf{e}_{(k-1)2^{-n-1}})_\# \alpha^{n+1}, \\ (\mathbf{e}_{(k+1)2^{-n-1}})_\# \alpha^{n+1})\} = 0. \end{aligned} \quad (3.20)$$

The support of all considered measures belongs to the compact set  $J(\mu_0, \mu_1)$ ; thus, using [16, Cor. 3.15] we get (3.20) for  $c'$  replaced by  $c_{n+1}$ . Thus,  $(\mathbf{e}_{k2^{-n-1}})_\# \alpha^{n+1} = \rho_{k2^{-n-1}} \mathbf{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  and, computing the geometric sum in the exponent,

$$\|\rho_{k2^{-n-1}}\|_{L^\infty(M, \mathbf{m})} \leq e^{D\sqrt{K-N}/2} \max\{\|\rho_0\|_{L^\infty(M, \mathbf{m})}, \|\rho_1\|_{L^\infty(M, \mathbf{m})}\}. \quad (3.21)$$

**Step 3.** *Completion.* Let  $(\alpha^n)_{n \in \mathbf{N}}$  be the sequence in  $\text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  iteratively constructed according to Step 1 and Step 2. Using Proposition B.9, this sequence converges weakly, up to a subsequence, to some  $\ell_p$ -optimal geodesic plan

$\alpha \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ . In particular, the assignment  $\mu_t := (\mathbf{e}_t)_{\#}\alpha$  defines a time-like proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1$ . Since the functional  $\mathcal{F}_c$  given in terms of the threshold

$$c := e^{D\sqrt{K-N}/2} \max\{\|\rho_0\|_{L^\infty(M, \mathfrak{m})}, \|\rho_1\|_{L^\infty(M, \mathfrak{m})}\}$$

analogously to (3.19) is weakly lower semicontinuous in  $\mathcal{P}(J(\mu_0, \mu_1))$  [16, Lem. 3.13], (3.21) yields  $\mathcal{F}_c(\mu_t) = 0$  for every  $t \in [0, 1]$ , which is the claim.  $\square$

**3.5. Equivalence with the entropic TCD condition.** In this section, we prove the equivalence of  $\text{TCD}_p^*(K, N)$  with its entropic counterpart  $\text{TCD}_p^e(K, N)$  introduced in [24], provided the underlying space is timelike  $p$ -essentially nonbranching according to Definition 2.16. We also obtain equivalence of the respective weak with their strong versions; see Theorem 3.35. Yet another characterization will be obtained in Proposition 3.43 below.

In this section, in addition to our standing assumptions on  $\mathcal{X}$  we suppose the latter to be timelike  $p$ -essentially nonbranching for a fixed  $p \in (0, 1)$ .

For the convenience of the reader, let us recall the following notions [24, Def. 3.2, Prop. 3.3], for which we define  $\mathcal{U}_N: \mathcal{P}(M) \rightarrow [0, \infty]$ ,  $N \in (0, \infty)$ , by

$$\mathcal{U}_N(\mu) := e^{-\text{Ent}_{\mathfrak{m}}(\mu)/N}. \quad (3.22)$$

**Definition 3.34.** *Let  $K \in \mathbf{R}$  and  $N \in (0, \infty)$ .*

- a. *We say that  $\mathcal{X}$  satisfies the entropic timelike curvature-dimension condition  $\text{TCD}_p^e(K, N)$  if for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) \in \mathcal{D}(\text{Ent}_{\mathfrak{m}})^2$ , there exist a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  connecting  $\mu_0$  and  $\mu_1$  as well as a timelike  $p$ -dualizing coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ ,*

$$\mathcal{U}_N(\mu_t) \geq \sigma_{K,N}^{(1-t)}[\|\tau\|_{L^2(M^2, \pi)}] \mathcal{U}_N(\mu_0) + \sigma_{K,N}^{(t)}[\|\tau\|_{L^2(M^2, \pi)}] \mathcal{U}_N(\mu_1).$$

- b. *If the previous claim holds for every strongly timelike  $p$ -dualizable  $(\mu_0, \mu_1) \in (\mathcal{P}_c(M) \cap \mathcal{D}(\text{Ent}_{\mathfrak{m}}))^2$ , we say that  $\mathcal{X}$  satisfies the weak entropic timelike curvature-dimension condition  $\text{wTCD}_p^e(K, N)$ .*

**Theorem 3.35.** *The following statements are equivalent for every given  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ .*

- (i) *The condition  $\text{TCD}_p^*(K, N)$  holds.*
- (ii) *The condition  $\text{wTCD}_p^*(K, N)$  holds.*
- (iii) *For every timelike  $p$ -dualizable  $(\mu_0, \mu_1) = (\rho_0 \mathfrak{m}, \rho_1 \mathfrak{m}) \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})^2$  there exists an  $\ell_p$ -optimal geodesic plan  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_{\#}\pi = \rho_t \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \sigma_{K,N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \sigma_{K,N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}$$

*holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^{\tau}(M)$ .*

- (iv) *The condition  $\text{TCD}_p^e(K, N)$  holds.*
- (v) *The condition  $\text{wTCD}_p^e(K, N)$  holds.*

*Remark 3.36.* The exceptional set in (iii) may depend on  $t$  and  $N'$ .

By Proposition 3.39 and the proofs of Proposition 3.38 and Proposition 3.40, any of the claims in Theorem 3.35 is equivalent to the following weaker version of (iii): for every pair  $(\mu_0, \mu_1) = (\rho_0 \mathfrak{m}, \rho_1 \mathfrak{m}) \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})^2$  such that  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$  and  $\rho_0, \rho_1 \in L^\infty(M, \mathfrak{m})$ , there exists  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_{\#}\pi \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$  and

$$\rho_t(\gamma_t)^{-1/N} \geq \sigma_{K,N}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N} + \sigma_{K,N}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N}$$

holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ , where  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m}$ .

An analogous note applies to Theorem 3.41, Theorem 4.20, and Theorem 4.21.

It is clear that (i) implies (ii), and that (iv) yields (v). Moreover, (iii) implies (i) by integration. The proofs of (ii) implying (iii), (iii) implying (iv), and (v) implying (iii) are, for the sake of clarity, outsourced to Proposition 3.38, Proposition 3.39, and Proposition 3.40, respectively.

*Remark 3.37.* Below, we occasionally use two results which are only established later in Chapter 4, namely the following.

- First, the condition  $\text{wTCD}_p^*(K, N)$  implies the timelike measure-contraction property  $\text{TMCP}_p^*(K, N)$  according to Definition 4.1, cf. Proposition 4.8.
- Second, on timelike  $p$ -essentially nonbranching spaces satisfying the latter condition — or  $\text{TMCP}_p^e(K, N)$  according to Definition 4.19 —, chronological  $\ell_p$ -optimal couplings and  $\ell_p$ -optimal geodesic plans with sufficiently well-behaved endpoints are unique, cf. Theorem 4.16, Theorem 4.17, and Remark 4.18.

As no result of this section is used in Chapter 4, no circular reasoning occurs.

The key point of these results here is that the timelike proper-time parametrized  $\ell_p$ -geodesics and the  $\ell_p$ -optimal couplings from Definition 3.2, Definition 3.3, as well as Definition 4.1, which are a priori unrelated, are in fact induced by the same  $\ell_p$ -optimal geodesic plan.

This means that we must have these uniqueness results at our disposal *a priori* to establish Theorem 3.35. In particular, these do *not* follow from Theorem 3.35, Remark 4.18, and [24, Thm. 3.19, Thm. 3.20].

**Proposition 3.38.** *Assume  $\text{wTCD}_p^*(K, N)$  for some  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Then for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})^2$  there exists some  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}$$

holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ , where  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m}$ .

*Proof. Step 1. Reinforcement of the assumptions on  $\mu_0$  and  $\mu_1$ .* We first assume that  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$ , that  $\mu_0, \mu_1 \in \mathcal{P}_c(M)$ , and that  $\rho_0, \rho_1 \in L^\infty(M, \mathbf{m})$ . As mentioned above, timelike  $p$ -dualizable couplings and  $\ell_p$ -optimal geodesic plans will then be unique all over Step 1, Step 2, and Step 3 without further notice.

**Step 2. Restriction.** Observe that the pair  $(\mu_0, \mu_1)$  is in fact strongly timelike  $p$ -dualizable thanks to Remark 2.12. Fix a  $\cap$ -stable generator  $\{M_n : n \in \mathbf{N}\}$  of the Borel  $\sigma$ -field of  $M$ ; to be precise, the  $M_n$ 's should be chosen to generate the relative Borel  $\sigma$ -field of the compact set  $\text{spt } \mu_0 \cup \text{spt } \mu_1$ , but we ignore this minor technicality by assuming compactness of  $M$  instead until Step 4. Given  $n \in \mathbf{N}$  we cover  $M$  by the mutually disjoint Borel sets

$$\begin{aligned} L_1 &:= M_1 \cap \cdots \cap M_n, \\ L_2 &:= M_1 \cap \cdots \cap M_{n-1} \cap M_n^c, \dots \\ L_{2^n} &:= M_1^c \cap \cdots \cap M_n^c. \end{aligned}$$

Given the unique timelike  $p$ -dualizing coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  of  $\mu_0$  and  $\mu_1$ , let us define  $\mu_0^{ij}, \mu_1^{ij} \in \mathcal{P}_c(M) \cap \mathcal{D}(\text{Ent}_{\mathbf{m}})$  by

$$\begin{aligned} \mu_0^{ij}[A] &:= \lambda_{ij}^{-1} \pi[(A \cap L_i) \times L_j] = \varrho_0^{ij} \mathbf{m}, \\ \mu_1^{ij}[A] &:= \lambda_{ij}^{-1} \pi[L_i \times (A \cap L_j)] = \varrho_1^{ij} \mathbf{m} \end{aligned}$$

for  $i, j \in \{1, \dots, 2^n\}$  provided  $\lambda_{ij} := \pi[L_i \times L_j] > 0$ . Then

$$\begin{aligned} \mu_0^{ij} \perp \mu_0^{i'j'}, \\ \mu_1^{ij} \perp \mu_1^{i'j'} \end{aligned} \quad (3.23)$$

for every  $i, i', j, j' \in \{1, \dots, 2^n\}$  with  $i \neq i'$  or  $j \neq j'$ . Also,  $(\mu_0^{ij}, \mu_1^{ij})$  is strongly timelike  $p$ -dualizable by Remark 2.12. By the consistency part in Proposition 3.7 and the above mentioned uniqueness results, and  $\text{wTCD}_p^*(K, N)$ , there exists some  $\pi^{ij} \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0^{ij}, \mu_1^{ij})$  with

$$\begin{aligned} \int_{\text{TGeo}^\tau(M)} \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \varrho_0^{ij}(\gamma_0)^{-1/N'} d\pi^{ij}(\gamma) \\ + \int_{\text{TGeo}^\tau(M)} \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \varrho_1^{ij}(\gamma_1)^{-1/N'} d\pi^{ij}(\gamma) \\ \leq \int_{\text{TGeo}^\tau(M)} \varrho_t^{ij}(\gamma_t)^{-1/N'} d\pi^{ij}(\gamma) \end{aligned} \quad (3.24)$$

for every  $t \in [0, 1]$  and every  $N' \geq N$ ; note that since  $\mu_0^{ij}, \mu_1^{ij} \in \mathcal{P}_c(M) \cap \mathcal{D}(\text{Ent}_{\mathfrak{m}})$  as a consequence of the assumption  $\rho_0, \rho_1 \in L^\infty(M, \mathfrak{m})$ , we have  $(\mathbf{e}_t)_\# \pi^{ij} = \rho_t^{ij} \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$  thanks to Lemma 3.8.

**Step 3.** *Pasting plans together and conclusion.* By uniqueness, we get

$$\pi = \lambda_{11} \pi^{11} + \lambda_{12} \pi^{12} + \dots + \lambda_{2^n 2^n} \pi^{2^n 2^n}$$

for every  $n \in \mathbf{N}$ , since the right-hand side belongs to  $\text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ . In particular, setting  $\mu_t := (\mathbf{e}_t)_\# \pi = \rho_t \mathfrak{m}$ ,  $t \in [0, 1]$  — the  $\mathfrak{m}$ -absolute continuity stemming from Lemma 3.8 and again uniqueness —, we have

$$\rho_t = \lambda_{11} \varrho_t^{11} + \lambda_{12} \varrho_t^{12} + \dots + \lambda_{2^n 2^n} \varrho_t^{2^n 2^n} \quad \mathfrak{m}\text{-a.e.}$$

Moreover, (3.23) and Lemma 2.18 imply

$$(\mathbf{e}_t)_\# \pi^{ij} \perp (\mathbf{e}_t)_\# \pi^{i'j'}$$

for every  $t \in (0, 1)$  and every  $i, i', j, j' \in \{1, \dots, 2^n\}$  with  $i \neq i'$  or  $j \neq j'$ , i.e.

$$\mathfrak{m}[\{\varrho_t^{ij} > 0\} \cap \{\varrho_t^{i'j'} > 0\}] = 0.$$

Setting  $G_{ij} := (\mathbf{e}_0, \mathbf{e}_1)^{-1}(L_i \times L_j)$ , by construction we obtain from (3.24) that

$$\begin{aligned} \int_{G_{ij}} \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} d\pi(\gamma) \\ + \int_{G_{ij}} \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'} d\pi(\gamma) \\ \leq \int_{G_{ij}} \rho_t(\gamma_t)^{-1/N'} d\pi(\gamma). \end{aligned}$$

Since  $i, j \in \{1, \dots, 2^n\}$  and  $n \in \mathbf{N}$  were arbitrary, the claim follows.

**Step 4.** *Removing the restrictions on  $\mu_0$  and  $\mu_1$ .* We initially address the first two restrictions in Step 1, then the last one.

**Step 4.1.** If the pair  $(\mu_0, \mu_1) \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})^2$  with  $\rho_0, \rho_1 \in L^\infty(M, \mathfrak{m})$  is merely timelike  $p$ -dualizable, let  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  be a corresponding timelike  $p$ -dualizing coupling. Since  $M_{\ll}^2$  is open and  $\mathfrak{d}$  is separable, we cover  $\text{spt } \pi \cap M_{\ll}^2$  by countably many (not necessarily disjoint) relatively compact rectangles  $A_i \times B_i$  with  $\bar{A}_i \times \bar{B}_i \subset M_{\ll}^2$ , where  $A_i, B_i \subset M$  are open,  $i \in \mathbf{N}$ . Set  $\pi^0 := 0$ , and define the measure  $\pi^i$  on  $M$  inductively by

$$\pi^i := (\pi - \pi^{i-1}) \mathbf{L}(A_i \times B_i).$$

Then  $\pi = \pi^1 + \pi^2 + \dots$  by construction. Define  $\hat{\pi}^i \in \mathcal{P}_c(M^2)$  by

$$\hat{\pi}^i := \pi^i[M^2]^{-1} \pi^i$$

provided  $\pi^i[M^2] > 0$ , and consider its marginals

$$\begin{aligned} \mu_0^i &:= (\text{proj}_1)_\# \hat{\pi}^i = \varrho_0^i \mathbf{m}, \\ \mu_1^i &:= (\text{proj}_2)_\# \hat{\pi}^i = \varrho_1^i \mathbf{m}. \end{aligned}$$

By Remark 2.12, the pair  $(\mu_0^i, \mu_1^i)$  is strongly timelike  $p$ -dualizable for every  $i \in \mathbf{N}$ . Thus, the previous steps imply the existence of  $\boldsymbol{\pi}^i \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0^i, \mu_1^i)$  such that for every  $t \in (0, 1)$ , we have  $(\mathbf{e}_t)_\# \boldsymbol{\pi}^i = \varrho_t^i \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ , and for every  $N' \geq N$ ,

$$\varrho_t^i(\gamma_t)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \varrho_0^i(\gamma_0)^{-1/N'} + \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \varrho_1^i(\gamma_1)^{-1/N'} \quad (3.25)$$

holds for  $\boldsymbol{\pi}^i$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ . By construction,

$$\begin{aligned} \mu_0^i &\perp \mu_0^j, \\ \mu_1^i &\perp \mu_1^j \end{aligned}$$

for every  $i, j \in \mathbf{N}$  with  $i \neq j$ , and from Lemma 2.18 we derive

$$(\mathbf{e}_t)_\# \boldsymbol{\pi}^i \perp (\mathbf{e}_t)_\# \boldsymbol{\pi}^j.$$

Note that the measure

$$\boldsymbol{\pi} := \pi^1[M^2] \boldsymbol{\pi}^1 + \pi^2[M^2] \boldsymbol{\pi}^2 + \dots$$

is an  $\ell_p$ -optimal geodesic plan interpolating  $\mu_0$  and  $\mu_1$ . It satisfies  $(\mathbf{e}_t)_\# \boldsymbol{\pi} = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  for every  $t \in [0, 1]$ , and

$$\rho_t = \pi^1[M^2] \varrho_t^1 + \pi^2[M^2] \varrho_t^2 + \dots \quad \mathbf{m}\text{-a.e.}$$

Together with (3.25) this discussion implies, for  $\boldsymbol{\pi}$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ ,

$$\rho_t(\gamma)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}.$$

**Step 4.2.** Finally, given any timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})^2$ , for every  $i, j \in \mathbf{N}$  we set

$$\begin{aligned} E_i &:= \{i - 1 \leq \rho_0^i < i\}, \\ F_j &:= \{j - 1 \leq \rho_1^j < j\}. \end{aligned}$$

By restriction [24, Lem. 2.10],  $(\mu_0^i, \mu_1^j)$  is timelike  $p$ -dualizable, where

$$\begin{aligned} \mu_0^i &:= \mu_0[E_i]^{-1} \mu_0 \llcorner E_i = \rho_0^i \mathbf{m}, \\ \mu_1^j &:= \mu_1[F_j]^{-1} \mu_1 \llcorner F_j = \rho_1^j \mathbf{m} \end{aligned}$$

provided  $\mu_0[E_i] > 0$  as well as  $\mu_1[F_j] > 0$ . Clearly  $\rho_0^i, \rho_1^j \in L^\infty(M, \mathbf{m})$ , and a similar argument as above leads to the conclusion. We omit the details.  $\square$

**Proposition 3.39.** *Let  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Assume that for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{D}(\text{Ent}_{\mathbf{m}})^2$ , there is  $\boldsymbol{\pi} \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_\# \boldsymbol{\pi} = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  and*

$$\rho_t(\gamma_t)^{-1/N} \geq \sigma_{K, N}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N} + \sigma_{K, N}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N} \quad (3.26)$$

*holds for  $\boldsymbol{\pi}$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ . Then  $\mathcal{X}$  obeys  $\text{TCD}_p^e(K, N)$ .*

*Proof.* The assignment  $\mu_t := (\mathbf{e}_t)_\# \boldsymbol{\pi}$  creates a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  from  $\mu_0$  to  $\mu_1$ . Moreover,  $\pi := (\mathbf{e}_0, \mathbf{e}_1)_\# \boldsymbol{\pi} \in \Pi_{\ll}(\mu_0, \mu_1)$  forms a timelike  $p$ -dualizing coupling of  $\mu_0$  and  $\mu_1$ . Taking logarithms on both sides of (3.26) and recalling (2.2), we get

$$-\frac{1}{N} \log \rho_t(\gamma_t) \geq \text{G}_t \left[ -\frac{1}{N} \log \rho_0(\gamma_0), -\frac{1}{N} \log \rho_1(\gamma_1), \frac{K}{N} \tau^2(\gamma_0, \gamma_1) \right]$$

for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ ; recall that the function  $G_t$  is defined in (2.3). Integrating this inequality with respect to  $\pi$  and using Jensen's inequality together with joint convexity of  $G_t$  yields

$$-\frac{1}{N} \text{Ent}_m(\mu_t) \geq G_t \left[ -\frac{1}{N} \text{Ent}_m(\mu_0), -\frac{1}{N} \text{Ent}_m(\mu_1), \frac{K}{N} \|\tau\|_{L^2(M^2, \pi)}^2 \right].$$

Here, in the case  $K > 0$  and  $\|\tau\|_{L^2(M^2, \pi)} = \infty$  the right-hand side is consistently interpreted as  $\infty$ , which implies  $\text{Ent}_m(\mu_t) = -\infty$ ; for  $K < 0$  and  $\|\tau\|_{L^2(M^2, \pi)} = \infty$  the right-hand side has an evident interpretation. Exponentiating both sides now yields the desired  $\text{TCD}_p^e(K, N)$  condition.  $\square$

**Proposition 3.40.** *Assume  $\text{wTCD}_p^e(K, N)$  for some  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Then for every timelike  $p$ -dualizable pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})^2$  there exists  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m}$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \sigma_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}$$

holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ , where  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m}$ .

*Proof.* The proof is similar to the one of Proposition 3.38 — whose notation we retain here —, whence we only outline the main differences. Again, it suffices to consider the case  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$ ,  $\mu_0, \mu_1 \in \mathcal{P}_c(M)$ , and  $\rho_0, \rho_1 \in L^\infty(M, \mathbf{m})$ . In particular, these restrictions imply  $\mu_0, \mu_1 \in \mathcal{D}(\text{Ent}_m)$ . The parallel uniqueness results outlined in Remark 3.37 under  $\text{TCD}_p^e(K, N)$  which are implicitly used below are due to [24, Thm. 3.19, Thm. 3.20], see also Remark 4.18.

Let  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  be an  $\ell_p$ -optimal geodesic plan interpolating  $\mu_0$  and  $\mu_1$ . Given any  $i, j \in \{1, \dots, 2^n\}$ , we define

$$\begin{aligned} \mu_0^{ij} &:= \lambda_{ij}^{-1} (\text{proj}_1)_\# [(\mathbf{e}_0, \mathbf{e}_1)_\# \pi \llcorner (L_i \times L_j)] = \varrho_0^{ij} \mathbf{m} \\ \mu_1^{ij} &:= \lambda_{ij}^{-1} (\text{proj}_2)_\# [(\mathbf{e}_0, \mathbf{e}_1)_\# \pi \llcorner (L_i \times L_j)] = \varrho_1^{ij} \mathbf{m} \end{aligned}$$

provided  $\lambda_{ij} := \pi[G_{ij}] > 0$ . Recall that  $\text{wTCD}_p^e(K, N)$  implies  $\text{wTCD}_p^e(K, N')$  by [24, Lem. 3.10]. Arguing as for Proposition 3.38, invoking  $\text{wTCD}_p^e(K, N')$  for the strongly timelike  $p$ -dualizable pair  $(\mu_0^{ij}, \mu_1^{ij}) \in (\mathcal{P}_c(M) \cap \mathcal{D}(\text{Ent}_m))^2$ , employing the uniqueness both of  $\pi$  as well as the involved timelike  $p$ -dualizing coupling  $\pi^{ij} \in \Pi_{\ll}(\mu_0^{ij}, \mu_1^{ij})$  by Remark 4.18, and taking logarithms on both sides of the resulting “pasted inequality” we infer that

$$\begin{aligned} & -\frac{\lambda_{ij}^{-1}}{N'} \int_{G_{ij}} \log \rho_t(\gamma_t) d\pi(\gamma) \\ & \geq G_t \left[ -\frac{\lambda_{ij}^{-1}}{N'} \int_{G_{ij}} \log \rho_0(\gamma_0) d\pi(\gamma), -\frac{\lambda_{ij}^{-1}}{N'} \int_{G_{ij}} \log \rho_1(\gamma_1) d\pi(\gamma), \right. \\ & \quad \left. \lambda_{ij}^{-1} \frac{K}{N'} \int_{G_{ij}} \tau^2(\gamma_0, \gamma_1) d\pi(\gamma) \right] \\ & \geq \lambda_{ij}^{-1} \int_{G_{ij}} G_t \left[ -\frac{1}{N'} \log \rho_0(\gamma_0), -\frac{1}{N'} \log \rho_1(\gamma_1), \frac{K}{N'} \tau^2(\gamma_0, \gamma_1) \right] d\pi(\gamma). \end{aligned}$$

Here the deduction that  $(\mathbf{e}_t)_\# \pi = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  comes directly from the definition of the functional  $\mathcal{U}_N$ , and the last inequality follows from joint convexity of  $G_t$  and Jensen's inequality. Since  $i, j \in \{1, \dots, 2^n\}$  and  $n \in \mathbf{N}$  were arbitrary, the claim follows by taking exponentials of both integrands, respectively.  $\square$

By evident adaptations of the above arguments, we also obtain the following result for the conditions from Definition 3.3.

**Theorem 3.41.** *The following statements are equivalent for every given  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ .*

- (i) *The condition  $\text{TCD}_p(K, N)$  holds.*
- (ii) *The condition  $\text{wTCD}_p(K, N)$  holds.*
- (iii) *For every timelike  $p$ -dualizable  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})^2$  there exists an  $\ell_p$ -optimal geodesic plan  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_{\#} \pi = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \tau_{K, N'}^{(1-t)}(\tau(\gamma_0, \gamma_1)) \rho_0(\gamma_0)^{-1/N'} + \tau_{K, N'}^{(t)}(\tau(\gamma_0, \gamma_1)) \rho_1(\gamma_1)^{-1/N'}$$

*holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^{\tau}(M)$ .*

**3.6. From local to global.** The goal of this section is to establish the equivalence of  $\text{TCD}_p^*(K, N)$  to its corresponding local version in Definition 3.42 below, i.e. to establish a Lorentzian analogue of the local-to-global property from [9, Ch. 5].

To this aim, in addition to our standing assumptions on the base space, suppose again  $\mathcal{X}$  to be timelike  $p$ -essentially nonbranching for  $p \in (0, 1)$ .

**Definition 3.42.** *Let  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ .*

- a. *We say that  $\mathcal{X}$  satisfies  $\text{TCD}_p^*(K, N)$  locally, briefly  $\text{TCD}_{p, \text{loc}}^*(K, N)$ , if there exists an open cover  $(M_i)_{i \in I}$  of  $M$  with the following property. For every  $i \in I$  and every pair  $(\mu_0, \mu_1) = (\rho_0 \mathbf{m}, \rho_1 \mathbf{m}) \in \mathcal{P}_c^{\text{ac}}(M_i, \mathbf{m})^2$  such that  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset (M_i)_{\ll}^2$ , there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  in  $\mathcal{P}(M)$  from  $\mu_0$  to  $\mu_1$  and an  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,*

$$\begin{aligned} \mathcal{S}_{N'}(\mu_t) \leq & - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1) \\ & - \int_{M^2} \sigma_{K, N'}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N'} d\pi(x^0, x^1). \end{aligned} \quad (3.27)$$

- b. *If the previous statement holds for  $\sigma_{K, N'}^{(1-t)}$  and  $\sigma_{K, N'}^{(t)}$  replaced by  $\tau_{K, N'}^{(1-t)}$  and  $\tau_{K, N'}^{(t)}$ , respectively,  $\mathcal{X}$  is termed to satisfy  $\text{TCD}_p(K, N)$  locally, briefly  $\text{TCD}_{p, \text{loc}}(K, N)$ .*

**Proposition 3.43.** *Given any  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ , the property  $\text{TCD}_p^*(K, N)$  holds if and only if the following does. For every  $\mu_0, \mu_1 \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  with  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$  there exists  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $N' \geq N$ ,*

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2})_{\#} \pi) \leq \sigma_{K, N'}^{(1/2)}(\theta) \mathcal{S}_{N'}(\mu_0) + \sigma_{K, N'}^{(1/2)}(\theta) \mathcal{S}_{N'}(\mu_1),$$

where

$$\theta := \begin{cases} \sup \tau(\text{spt } \mu_0 \times \text{spt } \mu_1) & \text{if } K < 0, \\ \inf \tau(\text{spt } \mu_0 \times \text{spt } \mu_1) & \text{otherwise.} \end{cases}$$

*Proof.* We only outline the main differences of the proof to its counterpart from [9, Prop. 2.10]. The forward implication is clear. Conversely, similarly to the proof of Theorem 3.33, by successively gluing plans corresponding to ‘‘midpoints’’ for which the Rényi entropy obeys the given inequality as in the proof of [9, Prop. 2.10], we inductively construct a sequence  $(\alpha^n)_{n \in \mathbf{N}}$  in  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that for every  $n \in \mathbf{N}$  and every odd  $k \in \{1, \dots, 2^n - 1\}$ ,

$$\begin{aligned} \mathcal{S}_{N'}((\mathbf{e}_{k2^{-n}})_{\#} \alpha^n) & \leq \sigma_{K, N'}^{(1/2)}(2^{-n+1} \theta) \mathcal{S}_{N'}((\mathbf{e}_{(k-1)2^{-n}})_{\#} \alpha^n) \\ & \quad + \sigma_{K, N'}^{(1/2)}(2^{-n+1} \theta) \mathcal{S}_{N'}((\mathbf{e}_{(k+1)2^{-n}})_{\#} \alpha^n) \\ & \leq \sigma_{K, N'}^{(1/2)}(2^{-n+1} \theta) \sigma_{K, N'}^{(1-(k-1)2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_0) \end{aligned}$$



$$\begin{aligned}
& + \sigma_{K,N'}^{(1/2)}(2^{-n+1}\theta) \sigma_{K,N'}^{((k-1)2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_1) \\
& + \sigma_{K,N'}^{(1/2)}(2^{-n+1}\theta) \sigma_{K,N'}^{(1-(k+1)2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_0) \\
& + \sigma_{K,N'}^{(1/2)}(2^{-n+1}\theta) \sigma_{K,N'}^{((k+1)2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_1) \\
& = \sigma_{K,N'}^{(1-k2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_0) + \sigma_{K,N'}^{(k2^{-n})}(\theta) \mathcal{S}_{N'}(\mu_1).
\end{aligned}$$

holds for every  $N' \geq N$ . The latter equality is a crucial property of the distortion coefficients  $\sigma_{K,N'}^{(r)}$  [76, Lem. 3.1] which does not hold for  $\tau_{K,N'}^{(r)}$ . Moreover, note that to proceed by induction in this strategy, it is important to know that the respective “midpoints” come from  $\ell_p$ -optimal geodesic plans, which ensures that all chronology relations are preserved through this process.

As in the proof of Theorem 3.33,  $(\alpha^n)_{n \in \mathbf{N}}$  admits an accumulation point  $\alpha \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ , giving rise to a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1$  via  $\mu_t := (\mathbf{e}_t)_{\#} \alpha$ . Thanks to the weak lower semicontinuity of the Rényi entropy for probability measures with uniformly compact support, the latter satisfies, for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\mathcal{S}_{N'}(\mu_t) \leq \sigma_{K,N'}^{(1-t)}(\theta) \mathcal{S}_{N'}(\mu_0) + \sigma_{K,N'}^{(t)}(\theta) \mathcal{S}_{N'}(\mu_1).$$

As for [9, Prop. 2.8], where the replacement of [9, Rem. 2.9] is Lemma 3.8, and Proposition 3.38, and using Theorem 3.35 we obtain  $\text{TCD}_p^*(K, N)$ .  $\square$

**Definition 3.44.** We term  $\mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  chronologically  $\ell_p$ -geodesic if every  $\mu_0, \mu_1 \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  with  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$  are joined by a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_r)_{r \in [0,1]}$  which consists of  $\mathbf{m}$ -absolutely continuous measures.

**Theorem 3.45.** Let  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Then  $\text{TCD}_p^*(K, N)$  holds if and only if  $\text{TCD}_{p,\text{loc}}^*(K, N)$  is satisfied and  $\mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  is chronologically  $\ell_p$ -geodesic.

We will prove Theorem 3.45 only in the case  $K \geq 0$ , the other situation follows by analogous computations. To this aim, we formulate the following property in Definition 3.46 for which, given any  $m \in \mathbf{N}_0$ , we define

$$I_m := \{k 2^{-m} : k \in \{0, \dots, 2^m\}\}.$$

Moreover, given a collection  $\mu := (\mu_t)_{t \in [0,1]}$  in  $\mathcal{P}(M)$ , let  $G_{\mu}^m$  denote the set of all  $\gamma \in \text{TGeo}^{\tau}(M)$  with  $\gamma_t \in \text{spt } \mu_t$  for every  $t \in I_m$ .

**Definition 3.46.** Let  $C \subset M$  be a compact set. Given  $m \in \mathbf{N}_0$ , we say that the property  $\text{P}_m(C)$  is satisfied if for every timelike proper-time parametrized  $\ell_p$ -geodesic  $\mu := (\mu_r)_{r \in [0,1]}$  consisting of  $\mathbf{m}$ -absolutely continuous measures with  $\mu_0, \mu_1 \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  with  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2 \cap C^2$ , and every  $s, t \in I_n$  with  $t - s = 2^{-m}$  there is an  $\ell_p$ -optimal geodesic plan  $\pi_{s,t} \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_s, \mu_t)$  with

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2})_{\#} \pi_{s,t}) \leq \sigma_{K,N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\mu_s) + \sigma_{K,N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\mu_t),$$

where

$$\theta_{s,t}^m := \inf \{ \tau(\gamma_s, \gamma_t) : \gamma \in G_{\mu}^m \}. \quad (3.28)$$

**Lemma 3.47.** Retaining the notation of Definition 3.46,  $\text{P}_m(C)$  implies  $\text{P}_{m-1}(C)$  for every  $m \in \mathbf{N}$ .

*Proof.* Let a timelike proper-time parametrized  $\ell_p$ -geodesic  $\mu := (\mu_r)_{r \in [0,1]}$  according to  $\text{P}_n(C)$  be given. Let  $s, t \in I_{m-1}$  such that  $t - s = 2^{1-m}$ , and let  $\theta := \theta_{s,t}^{m-1}$  be defined with respect to  $\mu$ . Inductively, we build a sequence  $(\mu^i)_{i \in \mathbf{N}_0}$  of timelike proper-time parametrized  $\ell_p$ -geodesics  $\mu^i := (\mu_r^i)_{r \in [0,1]}$  obeying  $\mu_r^i = \mu_r$  for every  $i \in \mathbf{N}_0$  and every  $r \in [0, s] \cup [t, 1]$  as follows. Initially, set  $\mu^0 := \mu$ .

**Step 1.** *Construction for odd  $i \in \mathbf{N}_0$ .* If the element  $\mu^{2i}$  has been constructed for a given  $i \in \mathbf{N}_0$ , as in the proof of [3, Thm. 2.10] and using that the respective “midpoints” result from an  $\ell_p$ -optimal geodesic plan interpolating its endpoints, we construct a timelike proper-time parametrized  $\ell_p$ -geodesic  $\mu^{2i+1} := (\mu_r^{2i+1})_{r \in [0,1]}$  with the subsequent properties. For every  $r \in [0, s] \cup [t, 1]$ , we have  $\mu_r^{2i+1} = \mu_r$ , and moreover, for every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}(\mu_{s+2^{-m-1}}^{2i+1}) &\leq \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_s) + \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i}), \\ \mathcal{S}_{N'}(\mu_{s+3 \times 2^{-m-1}}^{2i+1}) &\leq \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i}) + \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_t). \end{aligned}$$

Here we have used the property  $P_m(C)$ , the inequalities

$$\begin{aligned} 2\theta_{s,s+2^{-m}}^m &\geq \theta, \\ 2\theta_{s+2^{-m},t}^m &\geq \theta, \end{aligned}$$

where both quantities on the left-hand sides are defined with respect to  $\mu^{2i}$ , and nonincreasingness of  $\sigma_{K,N'}^{(1/2)}(\vartheta)$  in  $\vartheta \geq 0$  by our assumption  $K \geq 0$ .

**Step 2.** *Construction for even  $i \in \mathbf{N}_0$ .* Given  $\mu^{2i+1}$  exhibited in Step 1 above, similarly to this step we construct a timelike proper-time parametrized  $\ell_p$ -geodesic  $\mu^{2i+2} := (\mu_r^{2i+2})_{r \in [0,1]}$  with the property that  $\mu_r^{2i+2} = \mu_r^{2i+1}$  for every  $r \in [0, s + 2^{-m-1}] \cup [s + 3 \times 2^{-m-1}, 1]$  such that for every  $N' \geq N$ ,

$$\mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i+2}) \leq \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_{s+2^{-m-1}}^{2i+1}) + \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right] \mathcal{S}_{N'}(\mu_{s+3 \times 2^{-m-1}}^{2i+1}).$$

**Step 3.** *Conclusion.* Pasting together the inequalities from Step 1 and Step 2 yields, for every  $i \in \mathbf{N}_0$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i+2}) &\leq 2\sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right]^2 \mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i}) \\ &\quad + \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right]^2 \mathcal{S}_{N'}(\mu_s) + \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right]^2 \mathcal{S}_{N'}(\mu_t). \end{aligned}$$

Iterating this inequality gives

$$\begin{aligned} \mathcal{S}_{N'}(\mu_{s+2^{-m}}^{2i}) &\leq 2^i \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right]^2 \mathcal{S}_{N'}(\mu_{s+2^{-m}}) \\ &\quad + \frac{1}{2} \sum_{k=1}^i 2^k \sigma_{K,N'}^{(1/2)} \left[ \frac{\theta}{2} \right]^{2k} [\mathcal{S}_{N'}(\mu_s) + \mathcal{S}_{N'}(\mu_t)]. \end{aligned}$$

As  $\mu_0^{2i} = \mu_0$  and  $\mu_1^{2i} = \mu_1$  for every  $i \in \mathbf{N}_0$  by construction, using Proposition B.9 we get the existence of a sequence  $(\pi^{2i})_{i \in \mathbf{N}_0}$  in  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  with  $\mu_r^{2i} = (\mathbf{e}_r)_{\#} \pi^{2i}$  for every  $r \in [0, 1]$  which converges weakly, up to a nonrelabeled subsequence, to an  $\ell_p$ -optimal geodesic plan  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ . By  $\mathcal{K}$ -global hyperbolicity, lower semicontinuity of  $\mathcal{S}_{N'}$  on  $\mathcal{P}(J(\mu_0, \mu_1))$ , and the same computations as for [9, Clm. 5.2] for the distortion coefficients on the right-hand side of the previous inequality, setting  $\pi_{s,t} := (\text{restr}_s^t)_{\#} \pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_s, \mu_t)$  we get

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2})_{\#} \pi_{s,t}) \leq \sigma_{K,N'}^{(1/2)}(\theta) \mathcal{S}_{N'}(\mu_s) + \sigma_{K,N'}^{(1/2)}(\theta) \mathcal{S}_{N'}(\mu_t).$$

Here  $\text{restr}_s^t$  is defined in (2.11). This accomplishes  $P_{n-1}(C)$ .  $\square$

*Proof of Theorem 3.45.* The forward implication is a consequence of Lemma 3.8.

Concerning the backward implication, by the partition procedure in the proof of Proposition 3.38 it suffices to prove the desired inequality defining the condition  $\text{TCD}_p^*(K, N)$  for every  $\mu_0, \mu_1 \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  with  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$ .

Thanks to  $\mathcal{K}$ -global hyperbolicity, every member of any timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0$  to  $\mu_1$  has support in  $J(\mu_0, \mu_1)$ . By  $\text{TCD}_{p,\text{loc}}^*(K, N)$  and by compactness of  $J(\mu_0, \mu_1)$ , there exist  $\delta > 0$ , a disjoint cover  $L_1, \dots, L_n \subset M$

of  $J(\mu_0, \mu_1)$ , where  $n \in \mathbf{N}$ , and closed sets  $C_k \subset M$  containing  $\mathbf{B}^d(L_k, \delta)$ ,  $k \in \{1, \dots, n\}$ , such that every two measures in  $\mathcal{P}_c^{\text{ac}}(C_k, \mathfrak{m})$  are joined by a timelike proper-time parametrized  $\ell_p$ -geodesic obeying (3.27). Let  $m \in \mathbf{N}$  with

$$2^{-m} \leq \delta. \quad (3.29)$$

We claim  $\mathbf{P}_m(J(\mu_0, \mu_1))$ . Let  $(\mu_r)_{r \in [0,1]}$  be a timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0$  to  $\mu_1$  consisting of  $\mathfrak{m}$ -absolutely continuous measures. Let  $\boldsymbol{\pi} \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  represent  $(\mu_r)_{r \in [0,1]}$ , and define  $\varpi \in \mathcal{P}(M^{2^m+1})$  by

$$\varpi := (\mathbf{e}_0, \mathbf{e}_{2^{-m}}, \mathbf{e}_{2^{-m+1}}, \dots, \mathbf{e}_{(2^m-1)2^{-m}}, \mathbf{e}_1) \# \boldsymbol{\pi}. \quad (3.30)$$

Let  $s, t \in I_m$  with  $t - s = 2^{-m}$ . For  $k \in \{1, \dots, n\}$ , define  $\nu_s^k, \nu_t^k \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  by

$$\begin{aligned} \nu_s^k &:= \alpha_k^{-1} (\text{proj}_{2^m s}) \# \varpi \llcorner L_k, \\ \nu_t^k &:= \alpha_k^{-1} (\text{proj}_{2^m t}) \# [(\text{proj}_{2^m s}, \text{proj}_{2^m t}) \# \boldsymbol{\pi} \llcorner (L_k \times M)], \end{aligned}$$

provided  $\alpha_k := \mu_s[L_k] > 0$ . Note that  $\text{spt } \nu_s^k \subset \bar{L}_k$  and  $\text{spt } \nu_s^k \times \text{spt } \nu_t^k \subset M_{\ll}^2$ . Since  $r := \inf \tau(\text{spt } \mu_0 \times \text{spt } \mu_1) > 0$ , every curve in  $\text{TGeo}^{\tau}(M)$  starting in  $\text{spt } \mu_0$  and ending in  $\text{spt } \mu_1$  belongs to the set  $G_r$  defined in Corollary B.5, which is uniformly equicontinuous. Hence, using (3.29) and (3.30) we get  $\text{spt } \nu_s^k, \text{spt } \nu_t^k \subset \bar{\mathbf{B}}^d(L_k, \delta) \subset C_k$ . By  $\text{TCD}_{p, \text{loc}}^*(K, N)$  in the form described above, there is some  $\ell_p$ -optimal geodesic plan  $\boldsymbol{\pi}_{s,t}^k \in \text{OptTGeo}_{\ell_p}^{\tau}(\nu_s^k, \nu_t^k)$  such that

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2}) \# \boldsymbol{\pi}_{s,t}^k) \leq \sigma_{K, N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\nu_s^k) + \sigma_{K, N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\nu_t^k) \quad (3.31)$$

for every  $N' \geq N$ , where  $\theta_{s,t}^m$  is defined as in (3.28) with respect to  $\mu := (\mu_r)_{r \in [0,1]}$ .

Define  $\boldsymbol{\pi}_{s,t} \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_s, \mu_t)$  by

$$\boldsymbol{\pi}_{s,t} := \alpha_1 \boldsymbol{\pi}_{s,t}^1 + \dots + \alpha_n \boldsymbol{\pi}_{s,t}^n.$$

By construction and Lemma 2.18, respectively,  $(\mathbf{e}_r) \# \boldsymbol{\pi}_{s,t}^1, \dots, (\mathbf{e}_r) \# \boldsymbol{\pi}_{s,t}^n$  are mutually singular for every fixed  $r \in \{0, 1/2\}$ , whence

$$\mathcal{S}_{N'}(\mu_s) = \alpha_1^{1-1/N'} \mathcal{S}_{N'}(\nu_s^1) + \dots + \alpha_n^{1-1/N'} \mathcal{S}_{N'}(\nu_s^n),$$

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2}) \# \boldsymbol{\pi}_{s,t}) = \alpha_1^{1-1/N'} \mathcal{S}_{N'}((\mathbf{e}_{1/2}) \# \boldsymbol{\pi}_{s,t}^1) + \dots + \alpha_n^{1-1/N'} \mathcal{S}_{N'}((\mathbf{e}_{1/2}) \# \boldsymbol{\pi}_{s,t}^n).$$

On the other hand, since mutual singularity of  $\nu_t^1, \dots, \nu_t^n$  may fail,

$$\mathcal{S}_{N'}(\mu_t) \geq \alpha_1^{1-1/N'} \mathcal{S}_{N'}(\nu_t^1) + \dots + \alpha_n^{1-1/N'} \mathcal{S}_{N'}(\nu_t^n).$$

Merging these inequalities with (3.31) yields

$$\mathcal{S}_{N'}((\mathbf{e}_{1/2}) \# \boldsymbol{\pi}_{s,t}) \leq \sigma_{K, N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\mu_s) + \sigma_{K, N'}^{(1/2)}(\theta_{s,t}^m) \mathcal{S}_{N'}(\mu_t),$$

and the desired property  $\mathbf{P}_m(J(\mu_0, \mu_1))$  follows.

To close the proof of Theorem 3.45, let  $(\mu_r)_{r \in [0,1]}$  be a timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0$  to  $\mu_1$  which consists of  $\mathfrak{m}$ -absolutely continuous measures, as hypothesized. Iteratively, the property  $\mathbf{P}_m(J(\mu_0, \mu_1))$  proven above implies  $\mathbf{P}_0(J(\mu_0, \mu_1))$  by Lemma 3.47. By Proposition 3.20, this already implies the condition  $\text{TCD}_p^*(K, N)$ .  $\square$

The following Corollary 3.48 easily follows from Theorem 3.45 and the stability result from Theorem 3.29. Proposition 3.49 is proven along the same lines as [9, Prop. 5.5] by using uniform continuity of  $\tau$  on compact subsets of  $M^2$ ; the point is that for small  $\vartheta \geq 0$ , the quantities  $\sigma_{K, N}^{(r)}(\vartheta)$  and  $\tau_{K, N}^{(r)}(\vartheta)$  “almost coincide”. See also the computations in [30].

**Corollary 3.48.** *Let  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Then chronological  $\ell_p$ -geodesy of  $\mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  and the condition  $\text{TCD}_{p, \text{loc}}^*(K', N)$  hold for every  $K' < K$  if and only if  $\text{TCD}_p^*(K, N)$  is satisfied.*

**Proposition 3.49.** *Assume  $\mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  to be chronologically  $\ell_p$ -geodesic. Given any  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ ,  $\text{TCD}_{p,\text{loc}}^*(K', N)$  holds for every  $K' < K$  if and only if  $\text{TCD}_{p,\text{loc}}(K', N)$  is satisfied for every  $K' < K$ .*

*Remark 3.50.* We do not know whether Theorem 3.45 holds for the stronger conditions  $\text{TCD}_{p,\text{loc}}(K, N)$  and  $\text{TCD}_p(K, N)$ , respectively; compare with [9, Rem. 5.6]. It would follow from Theorem 3.45 if we succeeded in proving that  $\text{TCD}_p^*(K, N)$  and  $\text{TCD}_p(K, N)$  are *equivalent* on timelike  $p$ -essentially nonbranching spaces, similarly to the metric measure result [22, Thm. 1.1]; recall Proposition 3.6.

**3.7. Universal cover.** Now we discuss an application of Theorem 3.45 to the universal cover of a Lorentzian pre-length space, assuming that the latter exists. By construction of the Lorentzian structure on the universal cover, the latter inherits the curvature properties of its base space locally, and hence globally if  $\mathcal{X}$  is timelike  $p$ -essentially nonbranching; see Theorem 3.55 below for details. The universal cover plays a key role in the smooth *Gannon–Lee singularity theorems* [38, 39, 59]. (See also the recent work [80] for spacetimes with  $C^1$ -metrics, and the references therein.) More elementary, our discussion also creates further examples of spaces satisfying the  $\text{TCD}_p^*(K, N)$  condition.

**3.7.1. Covering spaces.** We start by recalling the following basic definition due to [81, p. 62, p. 80] (given for general topological spaces therein).

**Definition 3.51.** *Let  $(M, d)$  be a metric space.*

- a. *Given a topological space  $\hat{M}$ , a map  $\mathfrak{p}: \hat{M} \rightarrow M$  is called covering map if every point in  $M$  has an open neighborhood  $U \subset M$  such that  $\mathfrak{p}^{-1}(U)$  is the disjoint union of open subsets of  $\hat{M}$  on each of which  $\mathfrak{p}$  restricts to a homeomorphism.*
- b. *A topological space  $\hat{M}$  is called covering space for  $M$  if there exists a covering map  $\mathfrak{p}: \hat{M} \rightarrow M$ .*
- c. *A topological space  $\hat{M}$  is called universal cover for  $M$  if it is a covering space for  $M$ , and for every covering space  $\check{M}$  for  $M$  (with covering map  $\mathfrak{q}$ ) there exists a continuous map  $F: \hat{M} \rightarrow \check{M}$  such that*

$$\mathfrak{p} = F \circ \mathfrak{q}.$$

*Remark 3.52.* A universal cover might fail to exist in general [81, Ex. 2.5.17], but if it exists, it is unique up to homeomorphism. If  $M$  is connected, locally path-connected [81, p. 65], and semi-locally simply connected [81, p. 78], then it admits a simply connected universal cover [81, Cor. 2.5.14]. If one drops semi-local simple connectedness, a universal cover might still exist, but fail to be simply connected [81, Ex. 2.5.18].

Every covering projection  $\mathfrak{p}: \hat{M} \rightarrow M$  is a local homeomorphism [81, Lem. 2.1.8]. Hence, defining  $\hat{d}: \hat{M}^2 \rightarrow [0, \infty)$  by

$$\hat{d}(\hat{x}, \hat{y}) := d(\mathfrak{p}(\hat{x}), \mathfrak{p}(\hat{y})), \quad (3.32)$$

the covering space  $\hat{M}$  of a metric space is automatically a metric space with respect to  $\hat{d}$ , and  $\hat{d}$  induces the topology of  $\hat{M}$ .

*Remark 3.53.* By definition of a covering map and (3.32), for every compact  $C \subset M$  there exists an open subset  $U \subset M$  containing  $C$  such that  $\mathfrak{p}|_{\mathfrak{p}^{-1}(U)}$  is an isometric isomorphism, in particular bi-Lipschitz continuous. This implies that properness of  $d$  lifts to  $\hat{d}$ . Moreover, a curve  $\hat{\gamma}: [a, b] \rightarrow \hat{M}$ ,  $a, b \in \mathbf{R}$  with  $a < b$ , is  $\hat{d}$ -rectifiable if and only if  $\gamma := \mathfrak{p} \circ \hat{\gamma}$  is  $d$ -rectifiable, and in either case  $\text{Len}_{\hat{d}}(\hat{\gamma}) = \text{Len}_d(\gamma)$ .

**3.7.2. Lift of Lorentzian structures.** In this subsection, we fix a Lorentzian pre-length space  $(M, \mathbf{d}, \ll, \leq, \tau)$ . In the remainder of this section, we assume the existence of a universal cover  $\hat{M}$  for  $M$  with covering map  $\mathbf{p}$ .

We introduce a natural lifted Lorentzian pre-length structure on  $\hat{M}$ ; various Lorentzian properties will lift to  $\hat{M}$  as well. We confine ourselves to those introduced in Section 2.2 and needed for our work, although we expect more of the vast list of properties from [24, 58] to lift to  $\hat{M}$ .

The following Proposition 3.54 essentially follows from Remark 3.53 and the way  $\ll, \leq$ , and  $\hat{\tau}$  below are defined. (In the language of Definition 3.23,  $\mathbf{p}$  is a Lorentzian isometric embedding.) In particular, a curve  $\hat{\gamma}: [a, b] \rightarrow \hat{M}$ ,  $a, b \in \mathbf{R}$  with  $a < b$ , is causal or timelike if and only if  $\gamma := \mathbf{p} \circ \hat{\gamma}$  is causal or timelike, respectively, and in either case  $\text{Len}_{\hat{\tau}}(\hat{\gamma}) = \text{Len}_{\tau}(\gamma)$ . Moreover, the former is a geodesic if and only if the latter is. We omit the straightforward proof.

**Proposition 3.54.** *Define two relations  $\hat{\leq}$  and  $\hat{\ll}$  on  $\hat{M}$  by  $\hat{x} \hat{\leq} \hat{y}$  provided  $\mathbf{p}(\hat{x}) \leq \mathbf{p}(\hat{y})$ , and  $\hat{x} \hat{\ll} \hat{y}$  provided  $\mathbf{p}(\hat{x}) \ll \mathbf{p}(\hat{y})$ . Moreover, define  $\hat{\tau}: \hat{M}^2 \rightarrow [0, \infty]$  by*

$$\hat{\tau}(\hat{x}, \hat{y}) := \tau(\mathbf{p}(\hat{x}), \mathbf{p}(\hat{y})).$$

*Then  $(\hat{M}, \hat{\mathbf{d}}, \hat{\ll}, \hat{\leq}, \hat{\tau})$  is a Lorentzian pre-length space. If  $(M, \mathbf{d}, \ll, \leq, \tau)$  is*

- (i) *causally closed,*
- (ii) *locally causally closed,*
- (iii) *causally path-connected,*
- (iv) *non-totally imprisoning,*
- (v) *localizable,*
- (vi) *globally hyperbolic,*
- (vii)  *$\mathcal{K}$ -globally hyperbolic,*
- (viii) *length, or*
- (ix) *geodesic,*

*then the respective property holds for  $(\hat{M}, \hat{\mathbf{d}}, \hat{\ll}, \hat{\leq}, \hat{\tau})$  as well.*

**3.7.3. Lift of curvature properties.** Given a proper measured Lorentzian pre-length space  $\mathcal{X}$ , using the covering map  $\mathbf{p}$  the Lorentzian structure on the universal cover from Proposition 3.54 can be naturally endowed with a reference measure  $\hat{\mathbf{m}}$  as follows. Let  $\hat{\Sigma}$  be the family of all sets  $\hat{B} \subset \hat{M}$  such that  $\mathbf{p}|_{\hat{B}}$  is a local isometry with Borel image  $\mathbf{p}(\hat{B}) \subset M$ . This family is closed under any number of intersections, and since  $\mathbf{d}$  is proper by Remark 3.53, the  $\sigma$ -algebra  $\sigma(\hat{\Sigma})$  coincides with the Borel  $\sigma$ -algebra on  $\hat{M}$ . The definition  $\hat{\mathbf{m}}[\hat{B}] := \mathbf{m}[\mathbf{p}(\hat{B})]$ ,  $B \in \hat{\Sigma}$ , thus evidently gives rise to the desired (lifted) Borel measure  $\hat{\mathbf{m}}$  on  $\hat{M}$ .

For the subsequent theorem, we add the hypothesis of timelike  $p$ -essential nonbranching,  $p \in (0, 1)$ , to our standing assumptions on  $\mathcal{X}$ .

**Theorem 3.55.** *Let  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Assume  $\text{TCD}_p^*(K, N)$  for  $\mathcal{X}$ . Then the causally closed,  $\mathcal{K}$ -globally hyperbolic measured Lorentzian geodesic space  $\hat{\mathcal{X}} := (\hat{M}, \hat{\mathbf{d}}, \hat{\mathbf{m}}, \hat{\ll}, \hat{\leq}, \hat{\tau})$  according to Proposition 3.54 is timelike  $p$ -essentially nonbranching, and it obeys  $\text{TCD}_p^*(K, N)$ .*

*Proof.* Composition with the covering map  $\mathbf{p}$  yields a canonical nonrelabeled and continuous map  $\mathbf{p}: \text{TGeo}^{\tau}(\hat{M}) \rightarrow \text{TGeo}^{\tau}(M)$ . In an evident notation, given any  $\hat{\ell}_p$ -optimal geodesic plan  $\hat{\pi} \in \mathcal{P}(\text{TGeo}^{\tau}(\hat{M}))$ , the plan  $\pi := \mathbf{p}_{\#} \hat{\pi}$  lies in  $\mathcal{P}(\text{TGeo}^{\tau}(M))$ . Given a timelike nonbranching set  $G \subset \text{TGeo}^{\tau}(M)$  with  $\pi[G] = 1$ , the set  $\hat{G} := \mathbf{p}^{-1}(G) \subset \text{TGeo}^{\tau}(\hat{M})$  is timelike nonbranching as well, and  $\hat{\pi}[\hat{G}] = 1$ . Hence, timelike  $p$ -essential nonbranching lifts to the Lorentzian structure from Proposition 3.54.

Thanks to the construction given in Proposition 3.54 as well as Remark 3.53, the spaces  $\hat{\mathcal{X}}$  and  $\mathcal{X}$  are locally isometric as metric and Lorentzian pre-length spaces; recall Section 3.3. Therefore, the condition  $\text{TCD}_p^*(K, N)$  lifts and makes  $\hat{\mathcal{X}}$  obey

$\text{TCD}_p^*(K, N)$  locally. By lifting of timelike proper-time parametrized  $\ell_p$ -geodesics witnessing the inequality defining  $\text{TCD}_p^*(K, N)$  for  $\mathcal{X}$ ,  $\mathcal{P}_c^{\text{ac}}(\hat{M}, \hat{\mathfrak{m}})$  is chronologically  $\ell_p$ -geodesic according to Definition 3.44. Using Proposition 3.54 again with Theorem 3.45,  $\text{TCD}_p^*(K, N)$  thus holds globally on the universal cover.  $\square$

#### 4. TIMELIKE MEASURE-CONTRACTION PROPERTY

In this chapter, we introduce and study Lorentzian analogues of the measure-contraction property for metric measure spaces [70, 86]. An entropic version has recently been introduced in [24], cf. Definition 4.19 below. Again, for smooth spacetimes the respective conditions are tightly linked to timelike lower Ricci curvature bounds, cf. Theorem A.5.

##### 4.1. Definition and basic properties.

**Definition 4.1.** *Let  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ .*

- a. *We say that  $\mathcal{X}$  obeys the reduced timelike measure-contraction property  $\text{TMCP}_p^*(K, N)$  if for every  $\mu_0 \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  and every  $x_1 \in I^+(\mu_0)$  there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  from  $\mu_0$  to  $\mu_1 := \delta_{x_1}$  such that for every  $t \in [0, 1)$  and every  $N' \geq N$ ,*

$$\mathcal{S}_{N'}(\mu_t) \leq - \int_M \sigma_{K, N'}^{(1-t)}(\tau(x^0, x_1)) \rho_0(x^0)^{-1/N'} d\mu_0(x^0).$$

- b. *If the previous statement holds for  $\tau_{K, N'}^{(1-t)}$  instead of  $\sigma_{K, N'}^{(1-t)}$ ,  $\mathcal{X}$  is said to satisfy the timelike measure-contraction property  $\text{TMCP}_p(K, N)$ .*

*Remark 4.2.* As in Remark 3.5, the conditions  $\text{TMCP}_p^*(0, N)$  and  $\text{TMCP}_p(0, N)$  coincide for every  $p \in (0, 1)$  and every  $N \in [1, \infty)$ .

*Remark 4.3.* Under our standing assumptions on  $\mathcal{X}$ ,  $\text{TMCP}_p^*(K, N)$  holds if and only if the statement from Definition 4.1 is merely satisfied for  $\mathfrak{m}$ -a.e.  $x_1 \in M$  thanks to Proposition B.9 and Fatou's lemma. The analogous statement holds for  $\text{TMCP}_p(K, N)$  as well.

As for Proposition 3.6 and Proposition 3.7, the statements from Proposition 4.4 and Proposition 4.5 below are readily established (in all generality, the proofs work for general measured Lorentzian pre-length spaces).

**Proposition 4.4.** *The following hold for every  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ ; as in Proposition 3.6, set  $K^* := K(N - 1)/N$ .*

- (i) *The condition  $\text{TMCP}_p(K, N)$  implies  $\text{TMCP}_p^*(K, N)$ .*
- (ii) *If  $K > 0$ , the condition  $\text{TMCP}_p^*(K, N)$  implies  $\text{TMCP}_p(K^*, N)$ .*

**Proposition 4.5.** *Assume  $\text{TMCP}_p^*(K, N)$  for  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Then the following statements hold.*

- (i) **Consistency.** *The condition  $\text{TMCP}_p^*(K', N')$  holds for every  $K' \leq K$  and every  $N' \geq N$ .*
- (ii) **Scaling.** *Given any  $a, b, \theta > 0$ , the rescaled measured Lorentzian pre-length space  $(M, \text{ad}, \text{bm}, \ll, \leq, \theta\tau)$  satisfies  $\text{TMCP}_p^*(K/\theta^2, N)$ .*

*Analogous statements hold for the  $\text{TMCP}_p(K, N)$  condition.*

*Remark 4.6.* An analogue of Lemma 3.8 is unlikely to hold under  $\text{TMCP}_p^*(K, N)$  or  $\text{TMCP}_p(K, N)$ . In particular, we do not know a priori whether the measures  $\mu_t$ ,  $t \in [0, 1)$ , of a timelike proper-time parametrized  $\ell_p$ -geodesic as in Definition 4.1 are always — or can be chosen to be —  $\mathfrak{m}$ -absolutely continuous. Instead, we establish the existence of timelike proper-time parametrized  $\ell_p$ -geodesics consisting of  $\mathfrak{m}$ -absolutely continuous measures whose densities are uniformly  $L^\infty$  in time

under  $\text{TMCP}_p^*(K, N)$ , at least if  $\mu_0 \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  has  $\mathfrak{m}$ -essentially bounded density, by a variational method following [16], see also [23, 75, 76] in Section 4.3; cf. Theorem 4.12. By uniqueness of timelike proper-time parametrized  $\ell_p$ -geodesics under timelike essential nonbranching conditions, see Theorem 4.17, this suffices to show an equivalence result analogous to Theorem 3.35 for our timelike measure-contraction property, see Theorem 4.20 and Theorem 4.21.

The link of  $\text{TMCP}_p^*(K, N)$  and  $\text{TMCP}_p(K, N)$  to their respective TCD counterparts is more subtle. The proof of the corresponding Proposition 4.8 follows [24, Prop. 3.11]. We need the following result whose proof is a straightforward adaptation of the argument for [86, Lem. 3.3], recall also Lemma 3.27. The difference to [86] is that we only require the marginal whose density is contained in the respective integrand to be constant.

**Lemma 4.7.** *Let  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Let  $\rho: M \rightarrow [0, \infty)$  be a Borel function with  $\mu := \rho \mathfrak{m} \in \mathcal{P}(M)$ . Moreover, let  $(\pi_n)_{n \in \mathbf{N}}$  be a sequence in  $\mathcal{P}(M^2)$  converging weakly to  $\pi \in \mathcal{P}(M^2)$  such that for some  $i \in \{1, 2\}$ , we have*

$$(\text{proj}_i)_\# \pi_n = \mu$$

for every  $n \in \mathbf{N}$ . Then for every every  $r \in [0, 1]$ ,

$$\begin{aligned} & \int_{M^2} \tau_{K, N}^{(r)}(\tau(x^0, x^1)) (\rho \circ \text{proj}_i)(x^0, x^1)^{-1/N} d\pi(x^0, x^1) \\ & \leq \liminf_{n \rightarrow \infty} \int_{M^2} \tau_{K, N}^{(r)}(\tau(x^0, x^1)) (\rho \circ \text{proj}_i)(x^0, x^1)^{-1/N} d\pi_n(x^0, x^1). \end{aligned}$$

An analogous assertion holds with  $\tau_{K, N}^{(r)}$  replaced by  $\sigma_{K, N}^{(r)}$ .

**Proposition 4.8.** *The following hold for every  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ .*

- (i) *The condition  $\text{wTCD}_p(K, N)$  implies  $\text{TMCP}_p(K, N)$ .*
- (ii) *The condition  $\text{wTCD}_p^*(K, N)$  implies  $\text{TMCP}_p^*(K, N)$ .*

*Proof.* We only prove (i), the proof of (ii) is similar.

**Step 1.** *Approximation of  $\mu_0$  and  $\mu_1$ .* Given any  $\varepsilon \in (0, 1)$ , let  $C_\varepsilon \subset M$  be a compact set with  $\mu_0[C_\varepsilon] \geq 1 - \varepsilon$  and  $C_\varepsilon \times \{x_1\} \subset M_{\ll}^2$ . Define  $\mu_0^\varepsilon \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  by

$$\mu_0^\varepsilon := \mu_0[C_\varepsilon]^{-1} \mu_0 \mathbf{L} C_\varepsilon = \rho_0^\varepsilon \mathfrak{m}$$

Moreover, since  $M_{\ll}^2$  is open while  $C_\varepsilon \times \{x_1\}$  is compact, there exists  $\eta > 0$  such that  $C_\varepsilon \times \mathbf{B}^d(x_1, \eta) \subset M_{\ll}^2$ . For  $\delta \in (0, \eta)$ , we set

$$\mu_1^\delta := \mathfrak{m}[\mathbf{B}^d(x_1, \delta)]^{-1} \mathfrak{m} \mathbf{L} \mathbf{B}^d(x_1, \delta) = \rho_1^\delta \mathfrak{m}.$$

By Remark 2.12, the pair  $(\mu_0^\varepsilon, \mu_1^\delta)$  is strongly timelike  $p$ -dualizable, and using  $\text{wTCD}_p(K, N)$  we find a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t^{\varepsilon, \delta})_{t \in [0, 1]}$  connecting  $\mu_0^\varepsilon$  to  $\mu_1^\delta$  as well as a timelike  $p$ -dualizing  $\pi^{\varepsilon, \delta} \in \Pi_{\ll}(\mu_0^\varepsilon, \mu_1^\delta)$ , with support in  $C_\varepsilon \times \mathbf{B}^d(x_1, \eta)$ , such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}(\mu_t^{\varepsilon, \delta}) & \leq - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0^\varepsilon(x^0)^{-1/N'} d\pi^{\varepsilon, \delta}(x^0, x^1) \\ & \quad - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_1^\delta(x^1)^{-1/N'} d\pi^{\varepsilon, \delta}(x^0, x^1) \\ & \leq - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0^\varepsilon(x^0)^{-1/N'} d\pi^{\varepsilon, \delta}(x^0, x^1). \end{aligned}$$

**Step 2.** *Sending  $\delta \rightarrow 0$ .* Given any  $\varepsilon > 0$  and a fixed sequence  $(\delta_n)_{n \in \mathbf{N}}$  decreasing to 0, from  $(\mu_t^{\varepsilon, \delta_n})_{t \in [0, 1]}$ ,  $n \in \mathbf{N}$ , we construct a timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0^\varepsilon$  to  $\mu_1$  as follows. Let  $\pi^{\varepsilon, \delta_n} \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0^\varepsilon, \mu_1^{\delta_n})$  represent  $(\mu_t^{\varepsilon, \delta_n})_{t \in [0, 1]}$ . As  $(\mu_1^{\delta_n})_{n \in \mathbf{N}}$  converges weakly to  $\mu_1$ , this sequence is tight,

and so is  $(\pi^{\varepsilon, \delta_n})_{n \in \mathbf{N}}$  by Proposition B.9. Let  $\pi^\varepsilon \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0^\varepsilon, \mu_1)$  be a weak limit of a nonrelabeled subsequence. Then the assignment  $\mu_t^\varepsilon := (\mathbf{e}_t)_\# \pi^\varepsilon$ ,  $t \in [0, 1]$ , gives rise to a timelike proper-time parametrized  $\ell_p$ -geodesic connecting  $\mu_0^\varepsilon$  to  $\mu_1$ .

Every weak limit point of  $(\pi^{\varepsilon, \delta_n})_{n \in \mathbf{N}}$  equals  $\mu_0^\varepsilon \otimes \mu_1 = \mu_0^\varepsilon \otimes \delta_{x_1}$ . Therefore,  $\mathcal{K}$ -global hyperbolicity, weak lower semicontinuity of  $\mathcal{S}_{N'}$  on measures with uniformly bounded support, and Lemma 4.7 yield, for every such  $t$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}(\mu_t^\varepsilon) &\leq \limsup_{n \rightarrow \infty} \mathcal{S}_{N'}(\mu_t^{\varepsilon, \delta_n}) \\ &\leq - \liminf_{n \rightarrow \infty} \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x_1)) \rho_0^\varepsilon(x^0)^{-1/N'} d\pi^{\varepsilon, \delta_n}(x^0, x_1) \\ &\leq - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x_1)) \rho_0^\varepsilon(x^0)^{-1/N'} d\mu_0^\varepsilon(x^0) \end{aligned}$$

**Step 3.** *Sending  $\varepsilon \rightarrow 0$ .* Given a sequence  $(\varepsilon_n)_{n \in \mathbf{N}}$  decreasing to 0, note that  $(\mu_0^{\varepsilon_n})_{n \in \mathbf{N}}$  converges weakly to  $\mu_0$ . Similarly to Step 2, from  $(\mu_t^{\varepsilon_n})_{n \in \mathbf{N}}$  we construct a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  connecting  $\mu_0$  to  $\mu_1$ . As in Step 2 and using Fatou's lemma we get, for every  $t \in [0, 1]$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}(\mu_t) &\leq \limsup_{n \rightarrow \infty} \mathcal{S}_{N'}(\mu_t^{\varepsilon_n}) \\ &\leq - \liminf_{n \rightarrow \infty} \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x_1)) \rho_0^{\varepsilon_n}(x^0)^{-1/N'} d\mu_0^{\varepsilon_n}(x^0) \\ &\leq - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x_1)) \rho_0(x^0)^{-1/N'} d\mu_0(x^0). \quad \square \end{aligned}$$

*Remark 4.9.* With essentially identical proofs, the respective versions of the timelike Bonnet–Myers inequality, Corollary 3.14 and Corollary 3.21, and the timelike Bishop–Gromov inequality, Theorem 3.16 and Theorem 3.22, hold as well under  $\text{TMCP}_p(K, N)$  and  $\text{TMCP}_p^*(K, N)$ . We only need to assume  $\mathcal{X}$  to be a causally closed and globally hyperbolic Lorentzian geodesic space.

**4.2. Stability.** Next, we discuss the stability of the notions from Definition 4.1. In contrast to the weak stability from Theorem 3.29, both timelike measure-contraction properties are stable under the convergence introduced in Definition 3.24.

**Theorem 4.10.** *Assume the convergence of  $(\mathcal{X}_k)_{k \in \mathbf{N}}$  to  $\mathcal{X}_\infty$  as in Definition 3.24. Moreover, let  $(K_k, N_k)_{k \in \mathbf{N}}$  be a sequence in  $\mathbf{R} \times [1, \infty)$  converging to  $(K_\infty, N_\infty) \in \mathbf{R} \times [1, \infty)$ . Suppose the existence of  $p \in (0, 1)$  such that  $\mathcal{X}_k$  obeys  $\text{TMCP}_p(K_k, N_k)$  for every  $k \in \mathbf{N}$ . Then  $\mathcal{X}_\infty$  satisfies  $\text{TMCP}_p(K_\infty, N_\infty)$ .*

*The analogous statement in which  $\text{TMCP}_p(K_k, N_k)$  is respectively replaced by  $\text{TMCP}_p^*(K_k, N_k)$ ,  $k \in \mathbf{N}_\infty$ , holds as well.*

*Proof.* It suffices to prove the first statement, the second is argued analogously.

In this proof, we often adopt notations from the proof of Theorem 3.29 without explicit notice. In particular, we again identify  $M_k$  with its image  $\iota_k(M_k)$  in  $M$  and  $\mathbf{m}_k$  with its push-forward  $(\iota_k)_\# \mathbf{m}_k$  for every  $k \in \mathbf{N}_\infty$ .

**Step 1.** *Reduction to compact  $M$ .* Owing to Remark 4.3, given any  $\mu_{\infty, 0} = \rho_{\infty, 0} \mathbf{m}_\infty \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m}_\infty)$  and  $x_{\infty, 1} \in I^+(\mu_{\infty, 0}) \cap \text{spt } \mathbf{m}_\infty$ , as for Theorem 3.29 we use the compactness of  $\text{spt } \mu_{\infty, 0}$  and  $\text{spt } \mu_{\infty, 1}$ , where  $\mu_{\infty, 1} := \delta_{x_{\infty, 1}}$ , to assume without restriction that  $M$  is compact, and that  $\mathbf{m}_k \in \mathcal{P}(M)$  for every  $k \in \mathbf{N}_\infty$ . All measures considered below will thus be compactly supported. Moreover, we may and will suppose that  $W_2(\mathbf{m}_k, \mathbf{m}_\infty) \rightarrow 0$  as  $k \rightarrow \infty$ .

**Step 2.** *Restriction of the assumptions on  $\mu_{\infty, 0}$  and  $\mu_{\infty, 1}$ .* We will first assume that  $\tau(\cdot, x_{\infty, 1})$  is bounded away from zero on  $\text{spt } \mu_{\infty, 0}$ , that  $\rho_{\infty, 0} \in L^\infty(M, \mathbf{m}_\infty)$ , and that  $x_{\infty, 1}$  can be approximated with respect to  $d$  by a sequence  $(x_{k, 1})_{k \in \mathbf{N}}$  of points  $x_{k, 1} \in \text{spt } \mathbf{m}_k$  such that  $x_{\infty, 1} \in I^-(x_{k, 1})$  for every  $k \in \mathbf{N}$ . The general case



is discussed in Step 7 below; we note for now that this conclusion will not conflict with our reductions from Step 1.

**Step 3.** *Construction of a chronological recovery sequence.* In this step, given the sequence  $(x_{k,1})_{k \in \mathbf{N}}$  from Step 2 we construct a sequence  $(\mu_{k,0})_{k \in \mathbf{N}}$  of measures  $\mu_{k,0} = \rho_{k,0} \mathbf{m}_k \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$ ,  $k \in \mathbf{N}$ , such that  $\mu_{k,0} \rightarrow \mu_{\infty,0}$  weakly as  $k \rightarrow \infty$  possibly up to extracting a subsequence, and  $x_{k,1} \in I^+(\mu_{k,0})$  for every  $k \in \mathbf{N}$ . The constructed sequence will allow for the correct behavior of all functionals under consideration, cf. Step 5 below.

Given any  $k \in \mathbf{N}$ , let  $\mathbf{q}_k \in \mathcal{P}(M^2)$  be a  $W_2$ -optimal coupling of  $\mathbf{m}_k$  and  $\mathbf{m}_\infty$ . We disintegrate  $\mathbf{q}_k$  with respect to  $\text{proj}_1$ , writing

$$d\mathbf{q}_k(x, y) = d\mathbf{p}_x^k(y) d\mathbf{m}_k(x).$$

Let  $\mathbf{p}^k : \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty) \rightarrow \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  denote the canonically induced map.

Given any  $k \in \mathbf{N}$ , define  $\tilde{\mu}_{k,0} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  by

$$\tilde{\mu}_{k,0} := \mathbf{p}^k(\mu_{\infty,0}) = \tilde{\rho}_{k,0} \mathbf{m}_k.$$

Note that the measure  $\mathbf{r}_k := (\rho_{\infty,0} \circ \text{proj}_2) \mathbf{q}_k \in \mathcal{P}(M^2)$  constitutes a coupling of  $\tilde{\mu}_{k,0}$  and  $\mu_{\infty,0}$ . With the  $W_2$ -optimality of  $\mathbf{q}_k$ , this implies

$$W_2(\tilde{\mu}_{k,0}, \mu_{\infty,0}) \leq \|\rho_{\infty,0}\|_{L^\infty(M, \mathbf{m}_\infty)}^{1/2} W_2(\mathbf{m}_k, \mathbf{m}_\infty),$$

and consequently  $\tilde{\mu}_{k,0} \rightarrow \mu_{\infty,0}$  weakly as  $k \rightarrow \infty$ . Hence, by our assumption on the sequence  $(x_{k,1})_{k \in \mathbf{N}}$  from Step 2 and Portmanteau's theorem,

$$\liminf_{k \rightarrow \infty} \tilde{\mu}_{k,0}[I^-(x_{k,1})] \geq \liminf_{k \rightarrow \infty} \tilde{\mu}_{k,0}[I^-(x_{\infty,1})] \geq \mu_{\infty,0}[I^-(x_{\infty,1})] = 1.$$

Up to passing to a subsequence we may and will thus assume  $\tilde{\mu}_{k,0}[I^-(x_{k,1})] > 0$  for every  $k \in \mathbf{N}$ . We then define  $\mu_{k,0} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_k)$  by

$$\mu_{k,0} := \tilde{\mu}_{k,0}[I^-(x_{k,1})]^{-1} \tilde{\mu}_{k,0} \llcorner I^-(x_{k,1}).$$

By construction, the sequence  $(\mu_{k,0})_{k \in \mathbf{N}}$  converges to  $\mu_{\infty,0}$  weakly as  $k \rightarrow \infty$ , and we have  $x_{k,1} \in I^+(\mu_{k,0})$  for every  $k \in \mathbf{N}$ , as desired.

**Step 4.** *Invoking the TMCP condition.* Fix  $K \in \mathbf{R}$  and  $N \in (1, \infty)$  such that  $K < K_\infty$  and  $N > N_\infty$ . Up to passing to a subsequence if necessary, we may and will thus assume that  $K < K_k$  and  $N > N_k$  for every  $k \in \mathbf{N}$ .

By Proposition 4.5, for every  $k \in \mathbf{N}$  there exists a timelike proper-time parametrized  $l_p$ -geodesic  $(\mu_{k,t})_{t \in [0,1]}$  connecting  $\mu_{k,0}$  and  $\mu_{k,1} = \delta_{x_{k,1}}$  such that for every  $t \in [0, 1)$  and every  $N' \geq N$ ,

$$\mathcal{S}_{N'}^k(\mu_{k,t}) \leq - \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{k,1})) \rho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0). \quad (4.1)$$

**Step 5.** *Estimating the previous right-hand side.* We estimate the negative of the right-hand side of (4.1) from below, up to errors which become arbitrarily small. By construction of  $\mu_{k,0}$ , we find a sequence  $(a_k)_{k \in \mathbf{N}}$  of normalization constants converging to 1 such that  $\rho_{k,0} \leq a_k \tilde{\rho}_{k,0}$   $\mathbf{m}_k$ -a.e. for every  $k \in \mathbf{N}$ .

**Step 5.1.** First, we argue that  $x_{k,1}$  can be replaced  $x_{\infty,1}$  up to a small error. By our choice of  $K$  and  $N$  and the timelike Bonnet–Myers inequality from Remark 4.9,

$$c := \sup \tau_{K,N'}^{(1-t)} \circ \tau(M^2)$$

is finite. Furthermore, the involved function is jointly uniformly continuous. Thus, given any  $\varepsilon > 0$ , by uniform continuity we obtain, for sufficiently large  $k \in \mathbf{N}$ ,

$$\begin{aligned} & \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{k,1})) \rho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0) \\ & \geq \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \rho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0) - \varepsilon. \end{aligned}$$

**Step 5.2.** We modify  $\tilde{\mu}_{k,0}$  into the measure

$$\nu_{k,0} := (1 + \delta_k)^{-1} (\tilde{\rho}_{k,0} + \delta_k) \mathbf{m}_k = \varrho_{k,0} \mathbf{m}_k,$$

where  $\delta_k \in [0, 1]$  is defined by

$$\delta_k = \tilde{\mu}_{k,0}[I^-(x_{k,1})^c].$$

Note that  $(\delta_k)_{k \in \mathbf{N}}$  converges to 0. Employing the inequality  $\rho_{k,0} \leq a_k \varrho_{k,0}$   $\mathbf{m}_k$ -a.e., the definition of  $\mu_{k,0}$ , and a similar notation as in the proof of Theorem 3.29,

$$\begin{aligned} & \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \rho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0) \\ & \geq \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0) \\ & \geq \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\nu_{k,0}(x^0) \quad (4.2) \\ & \quad - \int_{I^-(x_{k,1})^c} \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\tilde{\mu}_{k,0}(x^0) \\ & \quad - \delta_k \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\mathbf{m}_k(x^0). \end{aligned}$$

**Step 5.3.** Possibly invoking the timelike Bishop–Gromov inequality outlined in Remark 4.9, we obtain the estimates

$$\begin{aligned} c \delta_k^{1-1/N'} & \geq \int_{I^-(x_{k,1})^c} \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\tilde{\mu}_{k,0}(x^0) \\ c \delta_k^{1-1/N'} & \geq \delta_k \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\mathbf{m}_k(x^0). \end{aligned}$$

Hence, our main task is to estimate the integral in (4.2) from below. To this aim, by definition of  $\nu_{k,0}$  and Jensen's inequality, we obtain

$$\begin{aligned} & \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\nu_{k,0}(x^0) \\ & = \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \left[ \int_M (\rho_{\infty,0}(y^0) + \delta_k) d\mathbf{p}_{x^0}^k(y^0) \right]^{1-1/N'} d\mathbf{m}_k(x^0) \\ & \geq \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \rho_{\infty,0}(y^0)^{1-1/N'} d\mathbf{q}_k(x^0, y^0). \end{aligned}$$

Let  $(\phi_i)_{i \in \mathbf{N}}$  be a sequence in  $C_b(M)$  such that

$$\|\phi_i - \rho_{\infty,0}^{1-1/N'}\|_{L^1(M, \mathbf{m}_{\infty})} \leq 2^{-i}$$

as well as  $\sup \phi_i(M) \leq \|\rho_{\infty,0}\|_{L^\infty(M, \mathbf{m}_{\infty})}$  for every  $i \in \mathbf{N}$ . Then we get

$$\begin{aligned} & \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\nu_{k,0}(x^0) \\ & \geq \int_{M^2} \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \phi_i(y^0) d\mathbf{q}_k(x^0, y^0) - 2^{-i} c. \end{aligned}$$

**Step 5.4.** By tightness and stability of  $W_2$ -optimal couplings [92, Lem. 4.3, Lem. 4.4],  $(\mathbf{q}_k)_{k \in \mathbf{N}}$  converges weakly to the diagonal coupling of  $\mathbf{m}_{\infty}$  and  $\mathbf{m}_{\infty}$  along a nonrelabeled subsequence. In particular, by Lebesgue's theorem,

$$\begin{aligned} & \liminf_{k \rightarrow \infty} \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \varrho_{k,0}(x^0)^{-1/N'} d\nu_{k,0}(x^0) \\ & \geq \liminf_{i \rightarrow \infty} \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \phi_i(x^0) d\mathbf{m}_{\infty}(x^0) - c \limsup_{i \rightarrow \infty} 2^{-i} \end{aligned}$$

$$= \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \rho_{\infty,0}(x^0)^{-1/N'} d\mu_{\infty,0}(x^0).$$

**Step 6. Conclusion.** Let  $(\pi_k)_{k \in \mathbf{N}}$  be a sequence of  $\pi_k \in \text{OptTGeo}_{\ell_p}^\tau(\mu_{k,0}, \mu_{k,1})$  representing the  $\ell_p$ -geodesic  $(\mu_{k,t})_{t \in [0,1]}$  in Step 4,  $k \in \mathbf{N}$ . By compactness of  $M$  and Proposition B.9, this sequence converges to an  $\ell_p$ -optimal geodesic plan  $\pi_\infty \in \text{OptTGeo}_{\ell_p}^\tau(\mu_{\infty,0}, \mu_{\infty,1})$  along a nonrelabeled subsequence. In particular, the assignment  $\mu_{\infty,t} := (\mathbf{e}_t)_\# \pi_\infty$  gives rise to a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{\infty,t})_{t \in [0,1]}$  from  $\mu_{\infty,0}$  to  $\mu_{\infty,1}$ ; note that every optimal coupling of  $\mu_{\infty,0}$  and  $\mu_{\infty,1}$  is concentrated on  $M_{\leq}^2$ . By joint weak lower semicontinuity of the Rényi entropy and (4.1), given any  $\varepsilon > 0$  this yields, for every  $t \in [0, 1)$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathcal{S}_{N'}^\infty(\mu_{\infty,t}) &\leq \limsup_{k \rightarrow \infty} \mathcal{S}_{N'}^k(\mu_{k,t}) \\ &\leq - \liminf_{k \rightarrow \infty} \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{k,1})) \rho_{k,0}(x^0)^{-1/N'} d\mu_{k,0}(x^0) \\ &\leq \varepsilon - \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1})) \rho_{\infty,0}(x^0)^{-1/N'} d\mu_{\infty,0}(x^0). \end{aligned}$$

**Step 7. Relaxation of the assumptions on  $\mu_{\infty,0}$  and  $\mu_{\infty,1}$ .** First, we argue how to construct the hypothesized sequence  $(x_{k,1})_{k \in \mathbf{N}}$  from Step 2. The idea is to approximate points which lie “in between”  $\text{spt } \mu_{\infty,0}$  and  $x_{\infty,1}$  from the future. For these points, the above discussion applies, and we will be able to conclude the desired TMCP property by a tightness argument.

**Step 7.1.** As  $\inf \tau(\text{spt } \mu_{\infty,0}, x_{\infty,1}) > 0$ , given any  $i \in \mathbf{N}$  we may and will fix a sequence  $(x_{\infty,1}^i)_{i \in \mathbf{N}}$  of points  $x_{\infty,1}^i \in I(\mu_{\infty,0}, \mu_{\infty,1}) \cap \text{spt } \mathbf{m}_\infty$  converging to  $x_{\infty,1}$ . Since  $I(x_{\infty,1}^i, x_{\infty,1})$  is open and nonempty, for every  $i \in \mathbf{N}$  we construct a sequence  $(x_{\infty,1}^{i,j})_{j \in \mathbf{N}}$  such that  $x_{\infty,1}^{i,j} \in I(x_{\infty,1}^i, x_{\infty,1}) \cap \text{spt } \mathbf{m}_\infty$  and

$$x_{\infty,1}^{i,j} \ll x_{\infty,1}^{i,j-1}$$

for every  $j \in \mathbf{N}$ . Since  $x_{\infty,1}^{i,j} \in \text{spt } \mathbf{m}_\infty$ , the weak convergence of  $(\mathbf{m}_k)_{k \in \mathbf{N}}$  implies the existence of a sequence  $(x_{k,1}^{i,j})_{k \in \mathbf{N}}$  of points  $x_{k,1}^{i,j} \in \text{spt } \mathbf{m}_k$  which converges to  $x_{\infty,1}^{i,j}$ . In particular, for a sufficiently large integer  $k_j \in \mathbf{N}$ , we have

$$x_{\infty,1}^i \ll x_{\infty,1}^{i,j} \ll x_{k_j,1}^{i,j}$$

for every  $i, j \in \mathbf{N}$ , as well as  $d(x_{k_j,1}^{i,j}, x_{\infty,1}^i) \rightarrow 0$  as  $j \rightarrow \infty$  for every  $i \in \mathbf{N}$ .

The above arguments applied for a fixed  $i \in \mathbf{N}$  (along a suitable subsequence in  $k$  which depends on  $i$ ) yield the existence of a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{\infty,t}^i)_{t \in [0,1]}$  from  $\mu_{\infty,0}$  to  $\mu_{\infty,1}^i := \delta_{x_{\infty,1}^i}$  such that for every  $t \in [0, 1)$  and every  $N' \geq N_\infty$ ,

$$\mathcal{S}_{N'}^\infty(\mu_{\infty,t}^i) \leq - \int_M \tau_{K,N'}^{(1-t)}(\tau(x^0, x_{\infty,1}^i)) \rho_{\infty,0}(x^0)^{-1/N'} d\mu_{\infty,0}(x^0).$$

Similarly to Step 6 and using lower semicontinuity of Rényi's entropy and Fatou's lemma, we find a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_{\infty,t})_{t \in [0,1]}$  connecting  $\mu_{\infty,0}$  to  $\mu_{\infty,1}$  such that the previous inequality holds for  $\mu_{\infty,t}^i$  and  $x_{\infty,1}^i$  replaced by  $\mu_{\infty,t}$  and  $x_{\infty,1}$ , respectively, for every  $t \in [0, 1)$  and every  $N' \geq N_\infty$ .

**Step 7.2.** Finally, we remove the two assumptions on  $\mu_{\infty,0}$  from Step 2. Let  $\mu_{\infty,0} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty)$  and  $x_{\infty,1} \in I^+(\mu_{\infty,0}) \cap \text{spt } \mathbf{m}_\infty$ . Given sufficiently large  $n, m \in \mathbf{N}$ , define  $\mu_{\infty,0}^n, \mu_{\infty,0}^{n,m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m}_\infty)$  by

$$\begin{aligned} \mu_{\infty,0}^n &:= \mu_{\infty,0} [\{\tau(\cdot, x_{\infty,1}) \geq 2^{-n}\}]^{-1} \mu_{\infty,0} \mathbf{L} \{\tau(\cdot, x_{\infty,1}) \geq 2^{-n}\} = \rho_{\infty,0}^n \mathbf{m}_\infty, \\ \mu_{\infty,0}^{n,m} &:= \|\min\{\rho_{\infty,0}^n, m\}\|_{L^1(M, \mathbf{m}_\infty)}^{-1} \min\{\rho_{\infty,0}^n, m\} \mathbf{m}_\infty. \end{aligned}$$

By continuity of  $\tau$ ,  $\mu_{\infty,0}^{n,m}$  obeys the hypotheses from Step 2 for every  $n, m \in \mathbf{N}$ . A diagonal procedure gives rise to a map  $m: \mathbf{N} \rightarrow \mathbf{N}$  such that  $(\tilde{\mu}_{\infty,0}^n)_{n \in \mathbf{N}}$ , where

$$\tilde{\mu}_{\infty,0}^n := \mu_{\infty,0}^{n,m_n},$$

converges weakly to  $\mu_{\infty,0}$  as  $n \rightarrow \infty$ .

The above discussion thus yields the desired inequality along some timelike proper-time parametrized  $\ell_p$ -geodesic connecting  $\tilde{\mu}_{\infty,0}^n$  to  $\mu_{\infty,1} := \delta_{x_{\infty,1}}$ . The usual tightness and lower semicontinuity argument directly gives the claim; note that the convergence of the corresponding right-hand sides is granted by Levi's theorem.

**Step 8. Passage from  $K$  and  $N$  to  $K_\infty$  and  $N_\infty$ .** According to Step 4, all in all we thus infer  $\text{TMCP}_p(K, N)$  for  $\mathcal{X}_\infty$  for every  $K < K_\infty$  and every  $N > N_\infty$ , and we deduce  $\text{TMCP}_p(K_\infty, N_\infty)$  as in Step 9 in the proof of Theorem 3.29.  $\square$

**4.3. Good geodesics.** Now we prove an analogue of Theorem 3.33, assuming the  $\text{TMCP}_p^*(K, N)$  condition. As indicated in Remark 4.6, this result will be used in the next Section 4.5 to prove the equivalence of the latter condition with the entropic timelike measure-contraction property from [24].

As in Section 3.4, we only outline the proof and refer to [16, Sec. 4.3], see also [23], for a similar discussion in the entropic case.

The following result is proven similarly to Lemma 3.32.

**Lemma 4.11.** *Assume the  $\text{wTCD}_p^*(K, N)$  condition for some  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [1, \infty)$ . Let  $\mu_0 = \rho_0 \mathbf{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$ , and let  $x_1 \in I^+(\mu_0)$ . Finally, let  $D$  be any real number no smaller than  $\sup \tau(\text{spt } \mu_0 \times \{x_1\})$ . Then there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1 := \delta_{x_1}$  such that for every  $t \in [0, 1)$ , denoting by  $\rho_t$  the  $\mathbf{m}$ -absolutely continuous part of  $\mu_t$ ,*

$$\mathbf{m}[\{\rho_t > 0\}] \geq (1-t)^N e^{-tD\sqrt{K-N}} \|\rho_0\|_{L^\infty(M, \mathbf{m})}^{-1}.$$

For the proof of Theorem 4.12, we introduce some further terminology. Following [24, Sec. 1.1], the *causally reversed* Lorentzian structure  $(M, \mathbf{d}, \ll^\leftarrow, \leq^\leftarrow, \tau^\leftarrow)$  corresponding to  $(M, \mathbf{d}, \ll, \leq, \tau)$  is given by

- $x \ll^\leftarrow y$  if and only if  $y \ll x$ ,
- $x \leq^\leftarrow y$  if and only if  $y \leq x$ , and
- $\tau^\leftarrow(x, y) := \tau(y, x)$ .

Let  $\ell_p^\leftarrow$  be the total transport cost function (2.7) induced by  $\tau^\leftarrow$ ,  $p \in (0, 1]$ .

**Theorem 4.12.** *Under the hypotheses of Lemma 4.11, there exists a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1 := \delta_{x_1}$  such that for every  $t \in [0, 1]$ ,  $\mu_t = \rho_t \mathbf{m} \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  and*

$$\|\rho_t\|_{L^\infty(M, \mathbf{m})} \leq \frac{1}{(1-t)^N} e^{Dt\sqrt{K-N}} \|\rho_0\|_{L^\infty(M, \mathbf{m})},$$

where

$$D := \sup \tau(\text{spt } \mu_0 \times \{x_1\}).$$

*Proof.* As for [16, Thm. 4.18], we first observe that the bisection argument from the proof of Theorem 3.33 does not apply here since  $\mu_1 \notin \mathcal{P}^{\text{ac}}(M, \mathbf{m})$ , while we aim to establish that  $\mu_t \in \mathcal{P}^{\text{ac}}(M, \mathbf{m})$  for every  $t \in [0, 1)$ .

Given  $n \in \mathbf{N}$  and  $k \in \mathbf{N}_0$ , we set  $s_n^k := (1 - 2^{-n})^k$ . Fix  $n \in \mathbf{N}$ , and assume that a measure  $\beta_n^k \in \text{OptTGeo}_{\ell_p^\leftarrow}^{\tau^\leftarrow}(\mu_1, \mu_0)$  has already been defined in such a way that for every  $i \in \{1, \dots, k\}$ , we have  $(e_{s_n^i})_\# \beta_n^k \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  and

$$\sup \tau^\leftarrow(\{x_1\} \times \text{spt}(e_{s_n^i})_\# \beta_n^k) \leq 2^{-n} s_n^{i-1} D.$$

**Step 1.** *Minimization of an appropriate functional.* Set

$$c_n^{k+1} := \frac{1}{(1-2^{-n})^N} e^{2^{-n} s_n^k D \sqrt{K-N}} \|\rho_0\|_{L^\infty(M, \mathfrak{m})}$$

and let the functional  $\mathcal{F}_{c_n^{k+1}}$  be defined as in (3.19) above. As in [16, Lem. 3.13], the functional  $\mathcal{F}_{c_n^{k+1}} \circ \mathbf{e}_{2^{-n}}$  admits some minimizer  $\pi_n^{k+1} \in \text{OptTGeo}_{\ell_p}^\tau((\mathbf{e}_{s_n^k})_\# \beta_n^k, \mu_1)$ . Let  $\sigma_n^{k+1} \in \text{OptTGeo}_{\ell_p}^{\tau \leftarrow}(\mu_1, (\mathbf{e}_{s_n^k})_\# \beta_n^k)$  be the  $\ell_p^{\leftarrow}$ -optimal geodesic plan obtained by “time-reversal” of  $\pi_n^{k+1}$ . By gluing, we build  $\beta_n^{k+1} \in \text{OptTGeo}_{\ell_p}^{\tau \leftarrow}(\mu_1, \mu_0)$  with

$$\begin{aligned} (\text{restr}_0^{s_n^k})_\# \beta_n^{k+1} &= \pi_n^{k+1}, \\ (\text{restr}_{s_n^k}^1)_\# \beta_n^{k+1} &= (\text{restr}_{s_n^k}^1)_\# \beta_n^k. \end{aligned}$$

Here, given any  $s, t \in [0, 1]$  with  $s < t$ , the map  $\text{restr}_s^t$  is defined in (2.11). Moreover,

$$\sup \tau^{\leftarrow}(\{x_1\} \times \text{spt}(\mathbf{e}_{s_n^{k+1}})_\# \beta_n^{k+1}) \leq 2^{-n} s_n^k D.$$

**Step 2.** *Uniform density bound.* Given any  $n \in \mathbf{N}$  and any  $k \in \mathbf{N}_0$ , arguing as in [16, Prop. 3.14, Cor. 3.15] and using Lemma 4.11,

$$\inf\{\mathcal{F}_{c_n^{k+1}}((\mathbf{e}_{2^{-n}})_\# \pi) : \pi \in \text{OptTGeo}_{\ell_p}^\tau((\mathbf{e}_{s_n^k})_\# \beta_n^k, \mu_1)\} = 0. \quad (4.3)$$

Note that Lemma 4.11 does *not* assert absolute  $\mathfrak{m}$ -continuity of the  $t$ -slice of the timelike proper-time parametrized  $\ell_p$ -geodesic under consideration. However, this does not affect the proof of [16, Prop. 3.14] as a possible further  $\mathfrak{m}$ -singular part gives an upper bound for the difference  $\mathcal{E}_{c'}(\pi) - \mathcal{E}_{c'}(\alpha)$  in Step 3 therein. In particular,  $(\mathbf{e}_{s_n^{k+1}})_\# \beta_n^{k+1} = \rho_{s_n^{k+1}} \mathfrak{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  and, by geometric summation,

$$\|\rho_{s_n^{k+1}}\|_{L^\infty(M, \mathfrak{m})} \leq \frac{1}{(s_n^{k+1})^N} e^{(1-s_n^{k+1})D\sqrt{K-N}} \|\rho_0\|_{L^\infty(M, \mathfrak{m})}. \quad (4.4)$$

**Step 3.** *Completion.* Iterating the procedure in Step 1 and Step 2, we construct a family  $\{\beta_n^k : n \in \mathbf{N}, k \in \mathbf{N}_0\} \subset \text{OptTGeo}_{\ell_p}^{\tau \leftarrow}(\mu_1, \mu_0)$ . Let  $(\beta^n)_{n \in \mathbf{N}}$  be an enumeration of the elements of the former set. By Proposition B.9 — observe that  $(M, \mathfrak{d}, \ll^{\leftarrow}, \leq^{\leftarrow}, \tau^{\leftarrow})$  inherits the regularity properties of its causally reversed structure — the latter admits an accumulation point  $\beta \in \text{OptTGeo}_{\ell_p}^{\tau \leftarrow}(\mu_1, \mu_0)$ . Let  $\alpha \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$  be the “time-reversal” of  $\beta$ , which induces a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  from  $\mu_0$  to  $\mu_1$  by definition. By weak lower semicontinuity of  $\mathcal{F}_c$  in  $\mathcal{P}(J(\mu_0, \mu_1))$  for appropriate values  $c > 0$ , recall (4.3) and (4.4), we get  $\mu_t = \rho_t \mathfrak{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$ , and  $\|\rho_t\|_{L^\infty(M, \mathfrak{m})}$  obeys the desired upper bound for every  $t \in [0, 1]$ .  $\square$

**4.4. Uniqueness of  $\ell_p$ -optimal couplings and  $\ell_p$ -geodesics.** In this section, we prove uniqueness of *chronological*  $\ell_p$ -optimal couplings (if existent), Theorem 4.16, as well as of  $\ell_p$ -optimal geodesic plans, Theorem 4.17. We follow [24, Sec. 3.4], see also [23]. A byproduct of our discussion is an extension of the results from [24] from the timelike nonbranching to the timelike *essential* nonbranching case, cf. Remark 4.18.

In this section, in addition to our standing assumptions, let  $\mathcal{X}$  be timelike  $p$ -essentially nonbranching for some fixed  $p \in (0, 1)$ .

**Lemma 4.13.** *Assume  $\text{TMCP}_p^*(K, N)$  for some  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Let  $\mu_0 = \rho_0 \mathfrak{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$ , and define  $\mu_1 := \lambda_1 \delta_{x_1} + \dots + \lambda_n \delta_{x_n}$  for  $\lambda_1, \dots, \lambda_n \in (0, 1]$  with  $\lambda_1 + \dots + \lambda_n = 1$  and pairwise distinct  $x_1, \dots, x_n \in M$ . Let  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  be an  $\ell_p$ -optimal coupling. Then there is a  $\mu_0$ -measurable map  $T : \text{spt} \mu_0 \rightarrow M$  with*

$$\pi = (\text{Id}, T)_\# \mu_0,$$

and consequently,

$$\ell_p(\mu_0, \mu_1)^p = \int_M \tau(x, T(x))^p d\mu_0(x).$$

In particular,  $\pi$  is the unique chronological  $\ell_p$ -optimal coupling of  $\mu_0$  and  $\mu_1$ .

*Proof.* By a standard argument, cf. the proof of [24, Lem. 3.17], it suffices to prove the existence of a  $\mu_0$ -measurable map  $T: \text{spt } \mu_0 \rightarrow M$  such that  $\pi = (\text{Id}, T)_\# \mu_0$ . To this aim, define the compact set  $E \subset \text{spt } \mu_0$  by

$$E := \{x \in M : \#[(\{x\} \times M) \cap \text{spt } \pi] \geq 2\}.$$

We claim that  $\mu_0[E] = 0$ , which directly gives the desired  $T$ .

Assume to the contrapositive that  $\mu_0[E] > 0$ . We first reduce the discussion to the uniform distribution on some subset of  $M$  as follows. Up to shrinking  $E$ , we assume without restriction that  $\varepsilon \leq \rho_0 \leq 1/\varepsilon$   $\mathfrak{m}$ -a.e. on  $E$  for some  $\varepsilon > 0$ . A further possible shrinking of  $E$  entails the existence of distinct points  $z_1, z_2 \in \{x_1, \dots, x_n\}$  and well-defined maps  $T_1, T_2: E \rightarrow M$  with  $T_1(x) = z_1$  and  $T_2(x) = z_2$  for every  $x \in E$ . We may and will additionally assume that  $E \times \{z_1\}, E \times \{z_2\} \subset M_{\ll}^2$ . By restriction [24, Lem. 2.10], the couplings

$$\begin{aligned} \pi_1 &:= \mathfrak{m}[E]^{-1} 1_{E \times \{z_1\}} (\rho_0 \circ \text{proj}_1)^{-1} \pi, \\ \pi_2 &:= \mathfrak{m}[E]^{-1} 1_{E \times \{z_2\}} (\rho_0 \circ \text{proj}_1)^{-1} \pi \end{aligned}$$

are  $\ell_p$ -optimal with  $\pi_1 \in \Pi_{\ll}(\nu_0, \delta_{z_1})$  and  $\pi_2 \in \Pi_{\ll}(\nu_0, \delta_{z_2})$ , respectively, where

$$\nu_0 := \mathfrak{m}[E]^{-1} \mathfrak{m} \llcorner E = \varrho_0 \mathfrak{m}.$$

From now on, we will work with  $\nu_0$  instead of  $\mu_0$ . Let  $\pi_1 \in \text{OptTGeo}_{\ell_p}^{\tau}(\nu_0, \delta_{z_1})$  and  $\pi_2 \in \text{OptTGeo}_{\ell_p}^{\tau}(\nu_0, \delta_{z_2})$  be  $\ell_p$ -optimal geodesic plans which represent time-like proper-time parametrized  $\ell_p$ -geodesics witnessing the defining inequality of  $\text{TMCP}_p^*(K, N)$ . Note that  $(\mathbf{e}_0)_\# \pi_1$  and  $(\mathbf{e}_0)_\# \pi_2$  are mutually singular since

$$\pi_i[\{\gamma \in \text{TGeo}^{\tau}(M) : \gamma_1 = z_i\}] = 1$$

for every  $i \in \{1, 2\}$ , and  $z_1 \neq z_2$ . By Lemma 2.18,

$$(\mathbf{e}_t)_\# \pi_1 \perp (\mathbf{e}_t)_\# \pi_2 \tag{4.5}$$

for every  $t \in (0, 1]$ . Given  $i \in \{1, 2\}$ , let  $\varrho_t^i$  denote the density of the  $\mathfrak{m}$ -absolutely continuous part of  $(\mathbf{e}_t)_\# \pi_1$ , which is nontrivial by  $\text{TMCP}_p^*(K, N)$ . By Remark 2.2,

$$\begin{aligned} \int_M (\varrho_t^i)^{1-1/N'} d\mathfrak{m} &\geq (1-t) e^{-tD\sqrt{K^-/N'}} \int_M \varrho_0^{1-1/N'} d\mathfrak{m} \\ &= (1-t) e^{-tD\sqrt{K^-/N'}} \mathfrak{m}[E]^{1/N'} \end{aligned}$$

for some  $N' > 1$ . On the other hand, by Jensen's inequality,

$$\begin{aligned} \int_M (\varrho_t^i)^{1-1/N'} d\mathfrak{m} &\leq \mathfrak{m}[\{\varrho_t^i > 0\}] \mathfrak{m}[\{\varrho_t^i > 0\}]^{-1} \int_{\{\varrho_t^i > 0\}} (\varrho_t^i)^{1-1/N'} d\mathfrak{m} \\ &\leq \mathfrak{m}[\{\varrho_t^i > 0\}] \left[ \mathfrak{m}[\{\varrho_t^i > 0\}]^{-1} \int_{\{\varrho_t^i > 0\}} \varrho_t^i d\mathfrak{m} \right]^{1-1/N'} \\ &\leq \mathfrak{m}[\{\varrho_t^i > 0\}]^{1/N'}. \end{aligned}$$

These inequalities imply

$$\liminf_{t \rightarrow 0} \mathfrak{m}[\{\varrho_t^i > 0\}] \geq \mathfrak{m}[E] = \mathfrak{m}[\{\varrho_0^i > 0\}]. \tag{4.6}$$

By  $\mathcal{K}$ -global hyperbolicity, the sets

$$F := \{x \in M : x \in \text{spt } (\mathbf{e}_t)_\# \pi_i \text{ for some } t \in [0, 1], i \in \{1, 2\}\},$$

$$G_\delta := \{x \in F : \tau(y, x) \leq \delta \text{ for some } y \in E\}$$

are compact for every  $\delta > 0$ . Since  $\mathbf{m}[G_\delta] \rightarrow \mathbf{m}[E]$  as  $\delta \rightarrow 0$  by Lebesgue's theorem, there exists some  $\delta_0 \in (0, 1)$  such that

$$\mathbf{m}[G_{\delta_0}] \leq \frac{3}{2} \mathbf{m}[E]. \quad (4.7)$$

By construction, for every  $i \in \{1, 2\}$ , every  $t \in (0, 1)$ , and  $(\mathbf{e}_t)_\# \pi_i$ -a.e.  $x \in M$  there exists  $\gamma \in \text{TGeo}^\tau(M)$  such that  $x = \gamma_t$ ,  $\gamma_0 \in E$ , and  $\gamma_1 = z_i$ . By definition of  $G_{\delta_0}$ , we thus obtain  $(\mathbf{e}_t)_\# \pi_i[G_{\delta_0}] = 1$  for every  $t \in [0, \delta_0]$ . Hence, (4.6) and (4.6) imply

$$\mathbf{m}[\{\rho_s^1 > 0\} \cap \{\rho_s^2 > 0\}] > 0$$

for some  $s \in (0, \delta_0)$ . However, this contradicts (4.5).  $\square$

**Lemma 4.14.** *Let  $\text{TMCP}_p^*(K, N)$  hold for some  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Let  $\mu_0 = \rho_0 \mathbf{m} \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$  and  $\mu_1 \in \mathcal{P}_c(M)$  with  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$ . Then there exist a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  connecting  $\mu_0$  to  $\mu_1$  as well as an  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1)$  and every  $N' \geq N$ ,*

$$\mathcal{S}_{N'}(\mu_t) \leq - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1). \quad (4.8)$$

*Proof. Step 1. Approximation of  $\mu_1$ .* Let  $B_1, \dots, B_n \subset M$ ,  $n \in \mathbf{N}$ , be a given Borel partition of  $\text{spt } \mu_1$  with  $\mu_1[B_i] > 0$  for every  $i \in \{1, \dots, n\}$ . Given such an  $i$ , fix  $x_1^i \in \text{spt } \mu_1$  and set  $\lambda_i := \mu_1[B_i]$  as well as

$$\mu_1^n := \lambda_1 \delta_{x_1^1} + \dots + \lambda_n \delta_{x_1^n}.$$

Since every coupling of  $\mu_0$  and  $\mu_1^n$  is chronological, an  $\ell_p$ -optimal coupling of these exists uniquely by Lemma 4.13. Let  $\pi_n \in \Pi_{\ll}(\mu_0, \mu_1^n)$  be this coupling, given by a  $\mu_0$ -measurable map  $T_n : \text{spt } \mu_0 \rightarrow M$ . For  $i \in \{1, \dots, n\}$ , we define

$$A_i := T_n^{-1}(\{x_1^i\}) \times \{x_1^i\},$$

and observe that  $A_1, \dots, A_n \subset M^2$  constitutes a Borel partition of  $\text{spt } \pi_n$ .

Define  $\pi_n^i \in \mathcal{P}(M^2)$ ,  $\nu_0^i \in \mathcal{P}_c^{\text{ac}}(M, \mathbf{m})$ , and  $\nu_1^i \in \mathcal{P}_c(M)$  by

$$\pi_n^i := \lambda_i^{-1} \pi_n \llcorner A_i,$$

$$\nu_0^i := (\text{proj}_1)_\# \pi_n^i = \varrho_0^i \mathbf{m},$$

$$\nu_1^i := (\text{proj}_2)_\# \pi_n^i = \delta_{x_1^i}.$$

By construction, we have  $\mu_0 = \lambda_1 \nu_0^1 + \dots + \lambda_n \nu_0^n$  and thus

$$\rho_0 = \lambda_1 \varrho_0^1 + \dots + \lambda_n \varrho_0^n \quad \mathbf{m}\text{-a.e.}$$

Moreover  $\text{spt } \nu_0^i \cap \text{spt } \nu_0^j = \emptyset$  for every  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ , whence

$$\nu_0^i \perp \nu_0^j. \quad (4.9)$$

**Step 2. Invoking the TMCP condition.** As  $x_1^i \in I^+(\nu_0^i)$  for every  $i \in \{1, \dots, n\}$ , using the  $\text{TMCP}_p^*(K, N)$  condition there exists a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\nu_t^i)_{t \in [0, 1]}$  from  $\nu_0^i$  to  $\nu_1^i$  such that for every  $t \in [0, 1)$  and every  $N' \geq N$ ,

$$\mathcal{S}_{N'}(\nu_t^i) \leq - \int_M \sigma_{K, N'}^{(1-t)}(\tau(x^0, x_1^i)) \rho_0^i(x^0)^{1-1/N'} d\mathbf{m}(x^0). \quad (4.10)$$

With this information, we now build a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t^n)_{t \in [0, 1]}$  from  $\mu_0$  to  $\mu_1^n$  for which (4.8) holds with  $\pi$  replaced by  $\pi_n$ . By definition,  $(\nu_t^i)_{t \in [0, 1]}$  is represented by some  $\alpha^i \in \text{OptTGeo}_{\ell_p}^\tau(\nu_0^i, \nu_1^i)$ . Thanks to (4.9) and Lemma 2.18, we obtain

$$(\mathbf{e}_t)_\# \alpha^i \perp (\mathbf{e}_t)_\# \alpha^j \quad (4.11)$$

for every  $t \in (0, 1]$  and every  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ . Then

$$\boldsymbol{\pi}_n := \lambda_1 \boldsymbol{\alpha}^1 + \dots + \lambda_n \boldsymbol{\alpha}^n$$

is an  $\ell_p$ -optimal geodesic plan from  $\mu_0$  to  $\mu_1^n$ , and the assignment  $\mu_t^n := (\mathbf{e}_t)_\# \boldsymbol{\pi}_n$  gives rise to a timelike proper-time parametrized  $\ell_p$ -geodesic connecting these probability measures. Let  $\varrho_t^i$  denote the nontrivial density of the  $\mathbf{m}$ -absolutely continuous part of  $(\mathbf{e}_t)_\# \boldsymbol{\alpha}^i$ . Then we obtain that

$$\rho_t^n \mathbf{m} := [\lambda_1 \varrho_t^1 + \dots + \lambda_n \varrho_t^n] \mathbf{m}$$

is the  $\mathbf{m}$ -absolutely continuous part of  $\mu_t^n$ ,  $t \in [0, 1]$ . Now (4.11) ensures

$$\mathbf{m}[\{\varrho_t^i > 0\} \cap \{\varrho_t^j > 0\}] = 0.$$

for every  $t \in [0, 1]$  and every  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ . Hence, using (4.10) and (4.9) we get, for every  $t \in [0, 1)$  and every  $N' \geq N$ ,

$$\begin{aligned} \mathfrak{S}_{N'}(\mu_t^n) &= \sum_{i=1}^n \lambda_i^{1-1/N'} \mathfrak{S}_{N'}(\mu_t^i) \\ &\leq - \sum_{i=1}^n \lambda_i^{1-1/N'} \int_M \sigma_{K, N'}^{(1-t)}(\tau(x^0, x_1^i)) \varrho_0^i(x^0)^{-1/N'}(x^0) \, d\nu_0^i(x^0) \\ &= - \sum_{i, i'=1}^n \int_M \sigma_{K, N'}^{(1-t)}(\tau(x^0, x_1^i)) [\lambda_{i'} \varrho_0^{i'}(x^0)]^{-1/N'} \, d[\lambda_i \nu_0^i](x^0) \\ &= - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} \, d\pi_n(x^0, x^1). \end{aligned}$$

**Step 3. Conclusion.** If  $\text{spt } \mu_1$  consists of finitely many points, the claim simply follows by choosing  $n \in \mathbf{N}$  such that  $\mu_1 = \mu_1^n$ .

Therefore, only the case  $\#\text{spt } \mu_1 = \infty$  remains to be studied. By [24, Thm. 2.16], there exists a sequence  $(\bar{\mu}_1^n)_{n \in \mathbf{N}}$  in  $\mathcal{P}_c(M)$  which converges weakly to  $\mu_1$  such that  $\bar{\mu}_1^n \in \mathcal{P}^{\text{ac}}(M, \mu_1^n)$  for every  $n \in \mathbf{N}$ , and for every sequence  $(\pi_n)_{n \in \mathbf{N}}$  of  $\ell_p$ -optimal couplings  $\pi_n \in \Pi_{\leq}(\mu_0, \bar{\mu}_1^n)$ , every weak limit of any subsequence of the latter is  $\ell_p$ -optimal for  $\mu_0$  and  $\mu_1$ . In particular, the previous discussion applies to  $\bar{\mu}_1^n$  in place of  $\mu_1^n$ ,  $n \in \mathbf{N}$ , and yields sequences  $(\boldsymbol{\pi}_n)_{n \in \mathbf{N}}$  and  $(\pi_n)_{n \in \mathbf{N}}$  of  $\ell_p$ -optimal geodesic plans  $\boldsymbol{\pi}_n \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \bar{\mu}_1^n)$  and  $\ell_p$ -optimal couplings  $\pi_n \in \Pi_{\ll}(\mu_0, \mu_1^n)$  as in Step 2, respectively. By  $\mathcal{K}$ -global hyperbolicity, Prokhorov's theorem, and Proposition B.9 these sequences converge weakly to some  $\boldsymbol{\pi} \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  and an  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$ , respectively, up to a nonrelabeled subsequence. Define a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  from  $\mu_0$  to  $\mu_1$  by  $\mu_t := (\mathbf{e}_t)_\# \boldsymbol{\pi}$ . Employing weak lower semicontinuity of  $\mathfrak{S}_{N'}$  on measures with uniformly bounded support and Lemma 4.7, we then obtain, for every  $t \in [0, 1)$ , every  $N' \geq N$ , and with  $\mu_t^n := (\mathbf{e}_t)_\# \boldsymbol{\pi}_n$ ,

$$\begin{aligned} \mathfrak{S}_{N'}(\mu_t) &\leq \limsup_{n \rightarrow \infty} \mathfrak{S}_{N'}(\mu_t^n) \\ &\leq - \liminf_{n \rightarrow \infty} \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} \, d\pi_n(x^0, x^1) \\ &\leq - \int_{M^2} \sigma_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} \, d\pi(x^0, x^1). \quad \square \end{aligned}$$

*Remark 4.15.* Arguing as in Step 4 in the proof of [24, Prop. 3.18] as well as Step 2 above, the conclusion of Lemma 4.14 holds under the following weaker assumption: the measures  $\mu_0$  and  $\mu_1$  admit an  $\ell_p$ -optimal coupling  $\varpi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that  $\text{spt } \pi \subset M_{\ll}^2$ . We will not need this extension in the sequel.



The proof of the following Theorem 4.16 follows the lines of Lemma 4.13 modulo some modifications we briefly discuss.

**Theorem 4.16.** *Assume  $\text{TMCP}_p^*(K, N)$  for  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Suppose the pair  $(\mu_0, \mu_1) \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m}) \times \mathcal{P}_c(M)$  to be timelike  $p$ -dualizable by  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$ . Then there exists a  $\mu_0$ -measurable map  $T: M \rightarrow M$  such that*

$$\pi = (\text{Id}, T)_{\#} \mu_0,$$

and consequently,

$$\ell_p(\mu_0, \mu_1)^p = \int_M \tau(x, T(x))^p d\mu_0(x).$$

In particular,  $\pi$  is the unique chronological  $\ell_p$ -optimal coupling of  $\mu_0$  and  $\mu_1$ .

*Proof.* As in the proof of Lemma 4.13, it suffices to prove the existence of  $T$ . Let  $\Gamma \subset M_{\ll}^2$  be an  $\ell_p$ -cyclically monotone set with  $\pi[\Gamma] = 1$  [24, Prop. 2.8], and set

$$E := \{x \in M : \text{proj}_2[\Gamma \cap (\{x\} \times M)] \geq 2\},$$

which is a Suslin set. We claim that  $\mu_0[S] = 0$ , which directly gives the desired  $T$ .

Assume to the contrapositive that  $\mu_0[S] > 0$ . By the von Neumann selection theorem [14, Thm. 9.1.3], there exist  $\mu_0$ -measurable maps  $T_1, T_2: E \rightarrow M$  whose graphs are both contained in  $\Gamma$  and such that  $T_1(x) \neq T_2(x)$  for every  $x \in E$ . By Lusin's theorem, there exists a compact set  $C \subset E$  with  $\mu_0[C] > 0$  such that the restrictions  $T_1|_C$  and  $T_2|_C$  are continuous. This yields

$$\min\{d(T_1(x), T_2(x)) : x \in C\} > 0.$$

In particular, there exist  $z_1, z_2 \in M$  and  $r > 0$  with  $d(z_1, z_2) > r$ , and a compact set  $C' \subset C$  with  $\mu_0[C'] > 0$  as well as  $T_1(C') \subset \mathbf{B}^d(z_1, r/2)$  and  $T_2(C') \subset \mathbf{B}^d(z_2, r/2)$ . Up to possibly shrinking the radius  $r$ , we may and will assume without restriction that  $C' \times [\bar{\mathbf{B}}^d(z_1, r/2) \cup \bar{\mathbf{B}}^d(z_2, r/2)] \subset M_{\ll}^2$ .

As in the proof of Lemma 4.13, we may further shrink  $C'$ , hence assume that  $\rho_0$  is bounded and bounded away from zero on  $C'$ . Consider  $\nu_0 \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  with

$$\nu_0 := \mathfrak{m}[C']^{-1} \mathfrak{m} \llcorner C'.$$

Furthermore, define  $\mu_1^1, \mu_1^2 \in \mathcal{P}_c(M)$  by

$$\begin{aligned} \mu_1^1 &:= (T_1)_{\#} \mu_0, \\ \mu_1^2 &:= (T_2)_{\#} \mu_0. \end{aligned}$$

By construction  $\text{spt } \mu_1^1 \cap \text{spt } \mu_1^2 = \emptyset$ , and the pairs  $(\nu_0, \mu_1^1)$  and  $(\nu_0, \mu_1^2)$  are both strongly  $p$ -timelike dualizable by Remark 2.12. Lemma 4.14 applies to both pairs; following the proof of Lemma 4.13 from now on gives the claim.  $\square$

**Theorem 4.17.** *Assume  $\text{TMCP}_p^*(K, N)$  for some  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ . Suppose the pair  $(\mu_0, \mu_1) \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m}) \times \mathcal{P}_c(M)$  to be timelike  $p$ -dualizable. Then for every  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  there is a  $\mu_0$ -measurable map  $\mathfrak{T}: \text{spt } \mu_0 \rightarrow \text{TGeo}^{\tau}(M)$  with*

$$\pi = \mathfrak{T}_{\#} \mu_0.$$

In particular, the set  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  is a singleton.

*Proof.* By the usual argument outlined in the proof of Lemma 4.13, it suffices to prove that every  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  is induced by a map  $\mathfrak{T}$  as above.

Assume to the contrapositive that one such  $\pi$ , henceforth fixed, is not given by a map. We disintegrate  $\pi$  with respect to  $\mathfrak{e}_0$ , writing, with some abuse of notation,

$$d\pi(\gamma) = d\pi_x(\gamma) d\mu_0(x)$$

for a  $\mu_0$ -measurable map  $\pi: \text{spt } \mu_0 \rightarrow \mathcal{P}(\text{TGeo}^{\tau}(M))$ . By assumption on  $\pi$ , there is a compact set  $C \subset \text{spt } \mu_0$  such that  $\mu_0[C] > 0$  and  $\#\text{spt } \pi_x \geq 2$  for every  $x \in C$ .

Hence, for every  $x \in C$  there exists  $t_x \in (0, 1)$  such that  $\#\text{spt}(\mathbf{e}_t)_\# \pi_x \geq 2$ ; since  $\text{TGeo}^\tau(M) \subset C([0, 1]; M)$  and  $\text{spt} \pi_x \subset \text{TGeo}^\tau(M)$ , there exists an open interval  $I_x \subset (0, 1)$  with  $t_x \in I_x$  such that  $\#\text{spt}(\mathbf{e}_t)_\# \pi_x \geq 2$  for every  $t \in I_x$ .

Given any  $t \in \mathbf{Q} \cap (0, 1)$ , we define

$$C_t := \{x \in C : \#\text{spt}(\mathbf{e}_t)_\# \pi_x \geq 2\}.$$

Since  $C_t$  unites to  $C$  when ranging over  $t \in \mathbf{Q} \cup (0, 1)$ , we find  $s \in \mathbf{Q} \cap (0, 1)$  with  $\mu_0[C_s] > 0$ . Finally, we define  $\alpha \in \mathcal{P}(\text{TGeo}^\tau(M))$  by

$$d\alpha(\gamma) = \mu_0[C_s]^{-1} 1_{C_s}(x) d\pi_x(\gamma) d\mu_0(x).$$

By restriction [24, Lem. 2.10],  $(\mathbf{e}_0, \mathbf{e}_1)_\# \alpha$  is an  $\ell_p$ -optimal coupling of its marginals, the first one of which is  $\mu_0[C_s]^{-1} \mu_0 \llcorner C_s \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$ . Since  $\alpha$  is concentrated on  $\text{TGeo}^\tau(M)$ , a standard argument using the reverse triangle inequality (2.4) for  $\tau$  implies that  $(\mathbf{e}_0, \mathbf{e}_s)_\# \alpha$  is a chronological  $\ell_p$ -optimal coupling of its marginals. By definition of  $C_s$ , it is not given by a map, which contradicts Theorem 4.16.  $\square$

*Remark 4.18.* The above arguments — which, as mentioned, are based on [24] — show that the uniqueness results of [24, Thm. 3.19, Thm. 3.20] remain true if the timelike nonbranching assumption therein is replaced by the hypothesis of timelike  $p$ -essential nonbranching for the considered exponent  $p \in (0, 1)$ .

**4.5. Equivalence with the entropic TMCP condition.** In this section, additionally to our standing assumptions we suppose  $\mathcal{X}$  to be timelike  $p$ -essentially nonbranching for some fixed  $p \in (0, 1)$ . Building upon the results in Section 4.3 and Section 4.4, we establish the equivalence of  $\text{TMCP}_p^*(K, N)$  with the entropic timelike measure-contraction property from [24, Def. 3.7], restated in Definition 4.19. A byproduct of our argumentation is a pathwise characterization of  $\text{TMCP}_p^*(K, N)$  and  $\text{TMCP}_p^e(K, N)$ , cf. Theorem 4.20, which is also available for  $\text{TMCP}_p(K, N)$  according to Theorem 4.21.

Recall the definition (3.22) of the exponentiated Boltzmann entropy  $\mathcal{U}_N$ .

**Definition 4.19.** *Let  $K \in \mathbf{R}$  and  $N \in (0, \infty)$ . We say that  $\mathcal{X}$  satisfies the entropic timelike measure-contraction property  $\text{TMCP}_p^e(K, N)$  if for every  $\mu_0 \in \mathcal{P}_c^{\text{ac}}(M, \mathfrak{m})$  and every  $x_1 \in I^+(\mu_0)$ , there exists an  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0, 1]}$  connecting  $\mu_0$  and  $\mu_1 := \delta_{x_1}$  such that for every  $t \in [0, 1]$ ,*

$$\mathcal{U}_N(\mu_t) \geq \sigma_{K, N}^{(1-t)} [\|\tau\|_{L^2(M^2, \mu_0 \otimes \mu_1)}] \mathcal{U}_N(\mu_0).$$

**Theorem 4.20.** *The following statements are equivalent for every given  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ .*

- (i) *The condition  $\text{TMCP}_p^*(K, N)$  holds.*
- (ii) *For every  $\mu_0 = \rho_0 \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$  and every  $x_1 \in I^+(\mu_0)$  there is an  $\ell_p$ -optimal geodesic plan  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ , where  $\mu_1 := \delta_{x_1}$ , such that for every  $t \in [0, 1]$ , we have  $(\mathbf{e}_t)_\# \pi = \rho_t \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \sigma_{K, N'}^{(1-t)}(\tau(\gamma_0, x_1)) \rho_0(\gamma_0)^{-1/N'}$$

*holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ .*

- (iii) *The condition  $\text{TMCP}_p^e(K, N)$  holds.*

*Proof.* We outline the necessary adaptations of the arguments in Section 3.5 and leave the details to the reader.

By integration, (ii) implies (i). The implication of (i) to (ii) is argued analogously to Proposition 3.38. Note that to obtain the asserted absolute continuity of  $(\mathbf{e}_t)_\# \pi$

with respect to  $\mathfrak{m}$  for every  $t \in [0, 1)$ , we have combined Theorem 4.12 with the uniqueness Theorem 4.17. The step from (ii) to (iii) follows since

$$-\frac{1}{N} \log \rho_t(\gamma_t) \geq H_t \left[ -\frac{1}{N} \log \rho_0(\gamma_0), \frac{K}{N} \tau^2(\gamma_0, x_1) \right]$$

for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ , using that the function  $H_t$  defined in (2.3) has similar convexity properties as the function  $G_t$  used to derive Proposition 3.39, and then arguing as in the proof of the latter result. Similarly, the implication from (iii) to (ii) is shown analogously to the proof of Proposition 3.40.  $\square$

Evident adaptations of the previous arguments give the following.

**Theorem 4.21.** *The following statements are equivalent for every given  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ .*

- (i) *The condition  $\text{TMCP}_p(K, N)$  holds.*
- (ii) *For every  $\mu_0 = \rho_0 \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$  and every  $x_1 \in I^+(\mu_0)$  there is an  $\ell_p$ -optimal geodesic plan  $\pi \in \text{OptTGeo}_{\ell_p}^\tau(\mu_0, \mu_1)$ , where  $\mu_1 := \delta_{x_1}$ , such that for every  $t \in [0, 1)$ , we have  $(e_t)_\# \pi = \rho_t \mathfrak{m} \in \mathcal{P}^{\text{ac}}(M, \mathfrak{m})$ , and for every  $N' \geq N$ , the inequality*

$$\rho_t(\gamma_t)^{-1/N'} \geq \tau_{K, N'}^{(1-t)}(\tau(\gamma_0, x_1)) \rho_0(\gamma_0)^{-1/N'}$$

*holds for  $\pi$ -a.e.  $\gamma \in \text{TGeo}^\tau(M)$ .*

#### APPENDIX A. SMOOTH LORENTZIAN SPACETIMES

In this appendix, we study the properties from Definition 3.2, Definition 3.3, and Definition 4.1 on smooth Lorentzian manifolds. In Theorem A.1 and Remark A.2, we prove the equivalence of all notions from the former two to upper dimension and timelike lower Ricci curvature bounds, thereby justifying the terminology ‘‘curvature-dimension condition’’ for the inherent notions. A similar equivalence for the timelike measure-contraction property is due to Theorem A.5.

We follow the conventions of [66]. See [71] for an exhaustive account on semi-Riemannian geometry, and [31, 53, 66, 69, 87] for details about optimal transport on smooth Lorentzian spacetimes.

In this chapter, let  $(M, \langle \cdot, \cdot \rangle)$  be a *smooth Lorentzian spacetime* of dimension  $n \in \mathbf{N}_{\geq 2}$ , that is,  $M$  is a smooth, connected, time-oriented manifold with smooth metric tensor  $\langle \cdot, \cdot \rangle$  of signature  $+, -, \dots, -$ . We always assume  $M$  to be globally hyperbolic [71, p. 411]. The space  $M$  canonically induces a Lorentzian pre-length structure [58, Ex. 2.11] which is locally causally closed by strong causality [58, Prop. 3.5], geodesic [71, Prop. 14.19], and  $\mathcal{K}$ -globally hyperbolic [24, Lem. 1.5]. Hence, all our previously obtained results apply to the setting in this chapter.

**A.1. Smooth timelike curvature-dimension condition.** For  $N > n$  and arbitrary  $V \in C^2(M)$  we define the  *$N$ -Bakry–Émery Ricci tensor*

$$\text{Ric}^{N, V} := \text{Ric} + \text{Hess } V - \frac{1}{N-n} DV \otimes DV. \quad (\text{A.1})$$

In the borderline case  $N = n$ , we conventionally assume  $V$  to be constant, in which case  $\text{Ric}^{N, V} := \text{Ric}$ . Given any  $K \in \mathbf{R}$ , we say that ‘‘ $\text{Ric}^{N, V} \geq K$  in every timelike direction’’ if  $\text{Ric}^{N, V}(\xi, \xi) \geq K |\xi|^2$  for every  $x \in M$  and every  $\xi \in T_x M$  with  $|\xi| > 0$ .

Set  $\text{vol}^V = e^V \text{vol}$ , and note that  $\mathcal{P}^{\text{ac}}(M, \text{vol}^V) = \mathcal{P}^{\text{ac}}(M, \text{vol})$ .

In the following main theorem of this section, the conditions in (iv) to (iii) are understood in terms of the  $N$ -Rényi entropy  $S_N^V$  with respect to  $\text{vol}^V$ ; the same applies to Theorem A.5 below.

**Theorem A.1.** *For every given  $p \in (0, 1)$ ,  $K \in \mathbf{R}$ , and  $N \in [n, \infty)$ , the subsequent statements are equivalent.*

- (i)  $\text{Ric}^{N,V} \geq K$  in every timelike direction.
- (ii) The condition  $\text{TCD}_p^*(K, N)$  holds.
- (iii) The condition  $\text{wTCD}_p^*(K, N)$ .
- (iv) The condition  $\text{TCD}_p(K, N)$  holds.
- (v) The condition  $\text{wTCD}_p(K, N)$  holds.

In particular, if any of the conditions in (ii) to (v) holds for some  $p \in (0, 1)$ , then these statements are satisfied for every  $p \in (0, 1)$ .

*Remark A.2.* Since  $M$  is causally plain (recall Example 2.7), any of (iv) and (v) does in fact imply  $N \geq n$  by Corollary 3.18. Hence, (iv) and (v) hold if and only if  $\text{Ric}^{N,V} \geq K$  in every timelike direction and  $N \geq n$ .

The only implication that still has to be established is (i) implying (v). Indeed, the equivalence of (iv) and (v) as well as (ii) and (iii) holds by Theorem 3.41, Theorem 3.35, and Remark 2.17, respectively. The latter also gives the equivalence of (ii) and (iii) to  $\text{TCD}_p^e(K, N)$ , which in turn implies (i) by [66, Thm. 8.5]. Moreover, (v) yields (iii), and (iv) yields (ii) by Proposition 3.6.

Let us now prove the remaining implication in Proposition A.3. Basically, the argument is a Lorentzian adaptation of the Riemannian proof of [86, Thm. 1.7]. The conclusion of Proposition A.3 is indeed sufficient to derive  $\text{wTCD}_p(K, N)$  by an argument as for Proposition 3.38, see also [66, Thm. 7.5].

**Proposition A.3.** *Let  $K \in \mathbf{R}$  and  $N \in [n, \infty)$ . Assume the inequality  $\text{Ric}^{N,V} \geq K$  in every timelike direction. Then for every  $p \in (0, 1)$ , every  $\mu_0, \mu_1 \in \mathcal{P}_c^{\text{ac}}(M, \text{vol}^V)$  such that  $\text{spt } \mu_0 \times \text{spt } \mu_1 \subset M_{\ll}^2$  and  $\rho_0, \rho_1 \in L^\infty(M, \text{vol}^V)$ , there is a timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$  from  $\mu_0$  to  $\mu_1$  and a timelike  $p$ -dualizing coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  such that for every  $t \in [0, 1]$  and every  $N' \geq N$ ,*

$$\begin{aligned} \mathcal{S}_{N'}^V(\mu_t) \leq & - \int_{M^2} \tau_{K, N'}^{(1-t)}(\tau(x^0, x^1)) \rho_0(x^0)^{-1/N'} d\pi(x^0, x^1) \\ & - \int_{M^2} \tau_{K, N'}^{(t)}(\tau(x^0, x^1)) \rho_1(x^1)^{-1/N'} d\pi(x^0, x^1). \end{aligned}$$

*Proof. Step 1. Setup.* We start with a short digression about Lorentzian optimal transport from [66]. Since  $\mu_0$  and  $\mu_1$  have compact support and  $\text{spt } \mu_0 \times \text{spt } \mu_1$  is chronological, the pair  $(\mu_0, \mu_1)$  is  $p$ -separated according to [66, Def. 4.1] by [66, Lem. 4.4]. Since also  $\mu_0 \ll \text{vol}$ ,  $\mu_0$  and  $\mu_1$  are both joined by a unique timelike proper-time parametrized  $\ell_p$ -geodesic  $(\mu_t)_{t \in [0,1]}$ , cf. [66, Cor. 5.9] and Remark B.8, and coupled by a unique  $\ell_p$ -optimal coupling  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$  [66, Thm. 5.8]. Additionally, these results yield the existence of some sufficiently regular vector field  $X$  on  $M$  such that, defining  $T: [0, 1] \times M \rightarrow M$  by

$$T_t(x) := \exp_x tX(x),$$

for every  $t \in [0, 1]$  we have

$$\begin{aligned} \mu_t &= (T_t)_\# \mu_0, \\ \pi &= (\text{Id}, T_1)_\# \mu_0. \end{aligned} \tag{A.2}$$

Thanks to [66, Prop. 6.1], the approximate derivative  $A_t := \tilde{D}T_t: TM \rightarrow (T_t)_*TM$  as introduced e.g. in [66, Def. 3.8] exists, is invertible, and depends smoothly on  $t \in [0, 1]$  vol-a.e. Given any  $x \in M$  and  $t \in [0, 1]$ , define

$$\begin{aligned} j_t(x) &:= |\det A_t(x)| e^{V(T_t(x))}, \\ \varphi_t(x) &:= \log j_t(x) = \log |\det A_t(x)| + V(T_t(x)), \\ B_t(x) &:= \dot{A}_t(x) A_t^{-1}(x). \end{aligned}$$

The function  $j_t$  should be thought of as a Jacobian, cf. Step 4 below.

**Step 2.** *Invoking the timelike Ricci curvature bound.* We henceforth suppose that  $N' \geq N > n$ ; in the case  $N' \geq N = n$  and recalling our convention of  $V$  being constant in this situation, the arguments to follow, e.g. the inequalities in (A.4), are somewhat easier. To relax notation, the dependency of the point  $x \in M$  at which the following computations may and will be performed is not made explicit.

By [66, Prop. 6.1], we know the formulas

$$\begin{aligned}\dot{\varphi}_t &= \operatorname{tr} B_t + \langle \nabla V, \dot{T}_t \rangle \circ T_t, \\ \ddot{\varphi}_t &= -\operatorname{Ric}(\dot{T}_t, \dot{T}_t) - \operatorname{Hess} V(\dot{T}_t, \dot{T}_t) - \operatorname{tr} B_t^2.\end{aligned}\tag{A.3}$$

Since  $(\operatorname{tr} B_t)^2 \leq n \operatorname{tr} B_t^2$  by the Cauchy–Schwarz inequality,

$$\begin{aligned}\frac{1}{N} \dot{\varphi}_t^2 &\leq \frac{1+\varepsilon}{N} (\operatorname{tr} B_t)^2 + \frac{1+\varepsilon^{-1}}{N} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &\leq \frac{n(1+\varepsilon)}{N} \operatorname{tr} B_t^2 + \frac{1+\varepsilon^{-1}}{N} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &= \operatorname{tr} B_t^2 + \frac{1}{N-n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t\end{aligned}$$

by choosing  $\varepsilon := (N-n)/n > 0$ . This entails

$$\ddot{\varphi}_t + \frac{1}{N'} \dot{\varphi}_t^2 \leq \ddot{\varphi}_t + \frac{1}{N} \dot{\varphi}_t^2 \leq -\operatorname{Ric}^{N,V}(\dot{T}_t, \dot{T}_t) \leq -K \vartheta^2,\tag{A.4}$$

where  $\vartheta := |\dot{T}_t| = \tau(\cdot, T_1) > 0$  by geodesy [66, Thm. 6.4]. This easily implies

$$\frac{\partial^2}{\partial t^2} J_t^{1/N'} \leq -\frac{K \vartheta^2}{N'} J_t^{1/N'}$$

which in turn entails

$$J_t^{1/N'} \geq \sigma_{K,N'}^{(1-t)}(\vartheta) J_0^{1/N'} + \sigma_{K,N'}^{(t)}(\vartheta) J_1^{1/N'}.$$

**Step 3.** *Replacement of  $\sigma_{K,N'}^{(r)}$  by  $\tau_{K,N'}^{(r)}$ .* Since  $(A_t)_{t \in [0,1]}$  is a Jacobi field along  $(T_t)_{t \in [0,1]}$  [66, Lem. 5.7], we obtain the Riccati equation

$$\dot{B}_t + B_t^2 + \operatorname{R}(\cdot, \dot{T}_t) \dot{T}_t = 0;\tag{A.5}$$

here  $\operatorname{R}$  is the Riemann curvature tensor of  $M$  with respect to  $\langle \cdot, \cdot \rangle$ .

Given any  $t \in [0, 1]$ , let  $\{e_1, \dots, e_n\}$  be a basis of  $T_{T_t} M$  with

$$e_1 := \dot{T}_t,\tag{A.6}$$

and let  $\{e^1, \dots, e^n\}$  denote the associated dual basis of  $T_{T_t}^* M$ . Let  $B_t^*: T_{T_t}^* M \rightarrow T_{T_t}^* M$  be the dual map of  $B_t$ , and let  $\sharp: T_{T_t}^* M \rightarrow T_{T_t} M$  be the musical isomorphism induced by  $\langle \cdot, \cdot \rangle$ . To simplify notation, we will suppress the dependency of most of these quantities on  $t$ . For  $i, i' \in \{1, \dots, n\}$ , set

$$\begin{aligned}u_t^{ii'} &:= e^i(B_t e_{i'}), \\ a_t &:= e^{1+\int_0^t u_s^{11} ds}, \\ \alpha_t &:= \log a_t.\end{aligned}$$

Employing (A.6), geodesy of  $(T_t)_{t \in [0,1]}$ , (A.5), and symmetry of  $B_t$ ,

$$\begin{aligned}-\dot{u}_t^{11} &= -\frac{\partial}{\partial t} \langle (e^1)^\sharp, B_t e_1 \rangle = -\langle (e^1)^\sharp, \dot{B}_t e_1 \rangle = \langle (e^1)^\sharp, B_t^2 e_1 \rangle \\ &= B_t^* e^1(B_t e_1) = \sum_{i'=1}^n (u_t^{1i'})^2 \geq (u_t^{11})^2.\end{aligned}\tag{A.7}$$

This easily implies  $-\dot{\alpha}_t \geq \alpha_t^2$  and in turn  $\ddot{a}_t \leq 0$ , whence

$$a_t \geq (1-t) a_0 + t a_1.\tag{A.8}$$

Let  $W_t$  be the matrix field with entries  $u_t^{ii'}$ ,  $i, i' \in \{2, \dots, n\}$ , and set

$$\begin{aligned}\delta_t &:= \varphi_t - \alpha_t, \\ d_t &:= e^{\delta_t} = a_t^{-1} j_t.\end{aligned}$$

Noting that  $\dot{\delta}_t = \text{tr } W_t + \langle \nabla V, \dot{T}_t \rangle \circ T_t$  thanks to (A.3), choosing  $\varepsilon := (N - n)/(n - 1) > 0$ , and employing the Cauchy–Schwarz inequality, the last equality of (A.7), and again (A.3), we arrive at

$$\begin{aligned}\frac{1}{N' - 1} \dot{\delta}_t^2 &\leq \frac{1}{N - 1} \dot{\delta}_t^2 \leq \frac{1 + \varepsilon}{N - 1} (\text{tr } W_t)^2 + \frac{1 + \varepsilon^{-1}}{N - 1} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &= \frac{1}{n - 1} (\text{tr } W_t)^2 + \frac{1}{N - n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &\leq \text{tr } W_t^2 + \frac{1}{N - n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &= \sum_{i, i'=2}^n (u_t^{ii'})^2 + \frac{1}{N - n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &\leq \sum_{i, i'=1}^n (u_t^{ii'})^2 - \sum_{i'=1}^n (u_t^{1i'})^2 + \frac{1}{N - n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &= \text{tr } B_t^2 + \ddot{\alpha}_t + \frac{1}{N - n} \langle \nabla V, \dot{T}_t \rangle^2 \circ T_t \\ &= -\ddot{\varphi}_t - \text{Ric}^{N, V}(\dot{T}_t, \dot{T}_t) + \ddot{\alpha}_t \leq -\ddot{\delta}_t - K \vartheta^2.\end{aligned}$$

This easily implies

$$\frac{\partial^2}{\partial t^2} d_t^{1/(N'-1)} \leq -\frac{K \vartheta^2}{N' - 1} d_t^{1/(N'-1)},$$

which in turn entails

$$d_t^{1/(N'-1)} \geq \sigma_{K, N'-1}^{(1-t)}(\vartheta) d_0^{1/(N'-1)} + \sigma_{K, N'-1}^{(t)}(\vartheta) d_1^{1/(N'-1)}. \quad (\text{A.9})$$

Finally, using  $j_t = a_t d_t$  and pasting (A.8) and (A.9) together yields

$$\begin{aligned}j_t^{1/N'} &\geq [(1 - t) a_0 + t a_1]^{1/N'} \\ &\quad \times [\sigma_{K, N'-1}^{(1-t)}(\vartheta) d_0^{1/(N'-1)} + \sigma_{K, N'-1}^{(t)}(\vartheta) d_1^{1/(N'-1)}]^{(N'-1)/N'} \\ &\geq (1 - t)^{1/N'} \sigma_{K, N'-1}^{(1-t)}(\vartheta) a_0^{1/N'} d_0^{1/N'} + t^{1/N'} \sigma_{K, N'-1}^{(t)}(\vartheta) a_1^{1/N'} d_1^{1/N'} \\ &= \tau_{K, N'}^{(1-t)}(\vartheta) j_0^{1/N'} + \tau_{K, N'}^{(t)}(\vartheta) j_1^{1/N'}.\end{aligned}$$

**Step 4. Conclusion.** Writing  $\mu_t = \rho_t \text{vol}^V = \rho_t e^V \text{vol}$ ,  $t \in [0, 1]$ , the change of variables formula with respect to  $\text{vol}$  gives  $(\rho_t \circ T_t) j_t = \rho_0 e^V \text{vol}$ -a.e. Combining this with the previous inequality and the identity  $\vartheta = \tau(\cdot, T_1)$  implies

$$\begin{aligned}\mathcal{S}_{N'}^V(\mu_t) &= - \int_M \rho_t^{1-1/N'} \text{dvol}^V = - \int_M (\rho_t \circ T_t)^{1-1/N'} j_t \text{dvol} \\ &= - \int_M \rho_0^{1-1/N'} e^{-V/N'} j_t^{1/N'} \text{dvol}^V \\ &\leq - \int_M \tau_{K, N'}^{(1-t)}(\tau \circ (\text{Id}, T_1)) j_0^{1/N'} \rho_0^{1-1/N'} e^{-V/N'} \text{dvol}^V \\ &\quad - \int_M \tau_{K, N'}^{(t)}(\tau \circ (\text{Id}, T_1)) j_1^{1/N'} \rho_0^{1-1/N'} e^{-V/N'} \text{dvol}^V\end{aligned}$$

$$\begin{aligned}
 &= - \int_M \tau_{K,N'}^{(1-t)}(\tau \circ (\text{Id}, T_1)) \rho_0^{-1/N'} d\mu_0 \\
 &\quad - \int_M \tau_{K,N'}^{(t)}(\tau \circ (\text{Id}, T_1)) (\rho_1 \circ T_1)^{-1/N'} d\mu_0
 \end{aligned}$$

for every  $t \in [0, 1]$ ; employing (A.2) gives the desired result.  $\square$

*Remark A.4.* We made the preceding assumption  $\rho_0, \rho_1 \in L^\infty(M, \text{vol}^V)$  only for cosmetic reasons in view of Proposition 3.38 and possible adaptations of Proposition A.3 to other settings. It is not used in the above proof.

**A.2. Smooth timelike measure-contraction property.** With all results shown thus far, it is now straightforward to relate the timelike measure-contraction properties from Definition 4.1 to timelike lower Ricci curvature bounds.

Recall our convention of  $V$  being constant when  $N = n$  in (A.1).

**Theorem A.5.** *For every given  $p \in (0, 1)$  and  $K \in \mathbf{R}$ , the subsequent statements are equivalent.*

- (i)  $\text{Ric} \geq K$  in every timelike direction.
- (ii) The condition  $\text{TMCP}_p^*(K, n)$  holds.
- (iii) The condition  $\text{TMCP}_p(K, n)$  holds.

*Proof.* Assuming (i), we know from Theorem A.1 that  $\text{TCD}_p(K, n)$  holds, which implies (iii) thanks to Proposition 4.8. The implication from (iii) to (ii) follows from Proposition 4.4. Lastly, (ii) implies  $\text{TMCP}_p^e(K, n)$  according to Theorem 4.20, whence [24, Thm. A.1] entails (i).  $\square$

*Remark A.6.* For general  $K \in \mathbf{R}$  and  $N \in [1, \infty)$ , even for smooth spacetimes the condition  $\text{TMCP}_p^*(K, N)$  is strictly weaker than  $\text{TCD}_p^*(K, N)$ , cf. [24, Rem. A.3], Theorem 3.35, and Theorem 4.20.

## APPENDIX B. GEODESICS OF PROBABILITY MEASURES

Here, we develop the theory of  $\ell_p$ -geodesics in  $\mathcal{P}(M)$ ,  $p \in (0, 1]$ . As indicated in Subsection 2.2.4 and Subsection 2.3.6, it is tailored to allow for a good compactness theory, which might be out of reach for the notions of  $\ell_p$ -geodesics from [24, 66] where not even weak continuity in time is a priori assumed or given.

In this chapter, in addition to Assumption 3.1, as in Subsection 2.3.6 it will not be restrictive to assume the compactness of  $M$  for notational convenience.

**B.1. Causal and timelike geodesics minimizing a Lagrangian action.** We first recall parts of Definition 2.13, defining

$$\begin{aligned}
 \text{OptGeo}_{\ell_p}(\mu_0, \mu_1) &:= \{ \boldsymbol{\pi} \in \mathcal{P}(\text{Geo}(M)) : (\mathbf{e}_0, \mathbf{e}_1) \# \boldsymbol{\pi} \in \Pi_{\leq}(\mu_0, \mu_1) \\
 &\quad \text{is } \ell_p\text{-optimal} \}.
 \end{aligned}$$

**Definition B.1.** *We term a weakly continuous path  $(\mu_t)_{t \in [0, 1]}$  in  $\mathcal{P}(M)$  a causal  $\ell_p$ -geodesic if it is represented by some  $\boldsymbol{\pi} \in \text{OptGeo}_{\ell_p}(\mu_0, \mu_1)$ .*

In Proposition B.3, we examine causal  $\ell_p$ -geodesics to precisely minimize some *Lagrangian action* according to the general theory developed in [92, Ch. 7]. (The *minimization* theory from [92] can easily be linked to our present Lorentzian setting which *maximizes* costs by flipping signs.) Analogously to the metric Wasserstein situation, cf. [3, Thm. 2.10, Rem. 2.14] and [92, Thm. 6.18], this is an instance of informally saying that  $\mathcal{P}(M)$  is a Lorentzian geodesic space with respect to  $\ell_p$  if and only if  $M$  is so with respect to  $\tau$ .

We consider a family  $\mathcal{A}$  of functionals  $\mathcal{A}^{s,t}: C([s,t]; M) \rightarrow (-\infty, \infty]$ ,  $s, t \in [0, 1]$  with  $s < t$ , defined by

$$\mathcal{A}^{s,t}(\gamma) := \begin{cases} -\text{Len}_\tau(\gamma)^p & \text{if } \gamma \text{ is causal in the sense of Subsection 2.2.3,} \\ \infty & \text{otherwise.} \end{cases}$$

In particular, by non-total imprisonment there exists  $c > 0$  such that  $\mathcal{A}^{s,t}(\gamma) < \infty$  only on curves  $\gamma: [s,t] \rightarrow M$  with  $\text{Lip } \gamma \leq c$ . Analogously, we define a family  $\mathbf{c}$  of cost functions  $c^{s,t}: M^2 \rightarrow (-\infty, \infty]$ ,  $s, t \in [0, 1]$  with  $s < t$ , by

$$c^{s,t}(x, y) := \begin{cases} -\tau(x, y)^p & \text{if } x \leq y, \\ \infty & \text{otherwise.} \end{cases}$$

**Lemma B.2.** *The family  $\mathcal{A}$  together with  $\mathbf{c}$  is a coercive Lagrangian action according to [92, Def. 7.11, Def. 7.13].*

*Proof.* We only outline the argument. For every  $s, t \in [0, 1]$  with  $s < t$ ,  $\mathcal{A}^{s,t}$  and  $c^{s,t}$  are lower semicontinuous (with respect to the uniform topology and the natural one on  $M^2$ , respectively) by causal closedness, [58, Prop. 3.17, Thm. 3.26], and continuity of  $\tau$ . Property (i) in [92, Def. 7.11] follows from additivity of  $\text{Len}_\tau$  [58, Lem. 2.25] and from  $\gamma$  failing to be causal if and only if this property fails on some subinterval of its domain of definition. The properties (ii) and (iii) in [92, Def. 7.11] follow from  $\mathcal{X}$  being a Lorentzian geodesic space. Boundedness of  $\tau$  on  $M$  ensures uniform lower boundedness of  $\mathcal{A}$  according to (i) in [92, Def. 7.13]. The compactness property (ii) from [92, Def. 7.13] follows from non-total imprisonment, compactness of  $M$ , and Arzelà–Ascoli’s theorem, and by existence of causal geodesics implicit in the definition of a Lorentzian geodesic space, cf. Definition 2.5.  $\square$

The following follows directly from Lemma B.2, cf. [92, Thm. 7.21]; for further consequences we do not need, we refer to [92, Prop. 7.16, Thm. 7.30].

Given  $s, t \in [0, 1]$  with  $s < t$  as well as  $\mu, \nu \in \mathcal{P}(M)$ , set

$$C^{s,t}(\mu, \nu) := \inf \left\{ \int_{M^2} c^{s,t}(x, y) d\pi(x, y) : \pi \in \Pi(\mu, \nu) \right\}.$$

**Proposition B.3.** *For a weakly continuous path  $(\mu_t)_{t \in [0,1]}$  in  $\mathcal{P}(M)$ , the following properties are equivalent.*

- (i) *The curve  $(\mu_t)_{t \in [0,1]}$  is a causal  $\ell_p$ -geodesic according to Definition B.1.*
- (ii) *For every  $r, s, t \in [0, 1]$  with  $r < s < t$ ,*

$$C^{r,s}(\mu_r, \mu_s) + C^{s,t}(\mu_s, \mu_t) = C^{r,t}(\mu_r, \mu_t).$$

- (iii) *For every  $s, t \in [0, 1]$  with  $s < t$ , the curve  $(\mu_r)_{r \in [s,t]}$  is a minimizer of the following coercive action  $\mathbf{A}^{s,t}$  defined by*

$$\mathbf{A}^{s,t}(\nu) := \sup \{ C^{t_0, t_1}(\nu_{t_0}, \nu_{t_1}) + C^{t_1, t_2}(\nu_{t_1}, \nu_{t_2}) + \cdots + C^{t_{n-1}, t_n}(\nu_{t_{n-1}}, \nu_{t_n}) \}$$

*for curves  $\nu := (\nu_r)_{r \in [s,t]}$  in  $\mathcal{P}(M)$ , where the supremum is taken over all  $n \in \mathbf{N}$  and all  $t_0, \dots, t_n \in [s, t]$  with  $t_i < t_{i+1}$  for every  $i \in \{0, \dots, n-1\}$ .*

*Given any  $\mu_0, \mu_1 \in \mathcal{P}(M)$  with  $\Pi_{\leq}(\mu_0, \mu_1) \neq \emptyset$ , there exists at least one curve  $(\mu_t)_{t \in [0,1]}$  satisfying either of the above properties.*

*Finally, if  $C_0, C_1 \subset \mathcal{P}(M)$  are weakly compact subsets with  $C^{0,1}(C_0 \times C_1) \subset \mathbf{R}$ , then the set  $\text{OptGeo}_{\ell_p}(C_0 \times C_1)$  is weakly compact.*



**B.2. Proper-time parametrizations.** Now we use Proposition B.3 and [92, Thm. 7.30] to establish properties of timelike (proper-time parametrized)  $\ell_p$ -geodesics. To set up these notions, let  $\mathbf{d}_\infty$  denote the uniform metric on  $C([0, 1]; M)$  induced by  $\mathbf{d}$ . As in Subsection 2.2.4, for  $\gamma \in \text{TGeo}(M)$  define  $\psi_\gamma \in C([0, 1]; [0, 1])$  by

$$\psi_\gamma(t) := \frac{\text{Len}_\tau(\gamma|_{[0,t]})}{\tau(\gamma_0, \gamma_1)} = \frac{\tau(\gamma_0, \gamma_t)}{\tau(\gamma_0, \gamma_1)}.$$

This function is strictly increasing [58, Prop. 2.34] and hence admits an inverse  $\psi_\gamma^{-1} \in C([0, 1]; [0, 1])$ . Define the  $\mathbf{r}: \text{TGeo}(M) \rightarrow C([0, 1]; M)$  by

$$\mathbf{r}(\gamma)_t := \gamma_{\psi_\gamma^{-1}(t)}.$$

Recall from [58, Cor. 3.35] that the map  $\mathbf{r}$  parametrizes  $\gamma \in \text{TGeo}(M)$  proportional to  $\tau$ -arclength, i.e. the identity (2.6) holds.

**Lemma B.4.** *The map  $\mathbf{r}$  is continuous with respect to  $\mathbf{d}_\infty$ .*

*Proof.* Let  $\varepsilon > 0$  and  $\gamma \in \text{TGeo}(M)$ , and note that for every  $\eta \in \text{TGeo}(M)$ ,

$$\begin{aligned} \mathbf{d}_\infty(\mathbf{r}(\gamma), \mathbf{r}(\eta)) &= \sup_{t \in [0,1]} \mathbf{d}(\gamma_{\psi_\gamma^{-1}(t)}, \eta_{\psi_\eta^{-1}(t)}) = \sup_{s \in [0,1]} \mathbf{d}(\gamma_{\psi_\gamma^{-1}(\psi_\eta(s))}, \eta_s) \\ &\leq \sup_{s \in [0,1]} \mathbf{d}(\gamma_{\psi_\gamma^{-1}(\psi_\eta(s))}, \gamma_s) + \mathbf{d}_\infty(\gamma, \eta) \\ &\leq \text{Lip } \gamma \sup_{s \in [0,1]} |\psi_\gamma^{-1}(\psi_\eta(s)) - s| + \mathbf{d}_\infty(\gamma, \eta). \end{aligned}$$

Uniform continuity of  $\psi_\gamma^{-1}$  implies the existence of  $\zeta > 0$  such that

$$|\psi_\gamma^{-1}(r) - \psi_\gamma^{-1}(r')| \leq (\text{Lip } \gamma)^{-1} \frac{\varepsilon}{2}$$

provided  $r, r' \in [0, 1]$  satisfy  $|r - r'| \leq \zeta$ . In turn, using that the set

$$\{\alpha \in \text{TGeo}(M) : 2\tau(\alpha_0, \alpha_1) \geq \tau(\gamma_0, \gamma_1), \alpha_0 \in \bar{\mathbf{B}}(\gamma_0, 1), \alpha_1 \in \bar{\mathbf{B}}(\gamma_1, 1)\}$$

is compact with respect to  $\mathbf{d}_\infty$ , we get the existence of  $\lambda > 0$  with

$$\sup_{s \in [0,1]} |\psi_\eta(s) - \psi_\gamma(s)| \leq \zeta$$

provided  $\eta \in \text{TGeo}(M)$  satisfies  $\mathbf{d}_\infty(\gamma, \eta) \leq \lambda$ . Hence, setting  $\delta := \min\{1, \lambda, \varepsilon/2\} > 0$ , by the above considerations we obtain that if  $\eta \in \text{TGeo}(M)$  obeys  $\mathbf{d}_\infty(\gamma, \eta) \leq \delta$ , then  $\mathbf{d}_\infty(\mathbf{r}(\gamma), \mathbf{r}(\eta)) \leq \varepsilon$ , which is the desired claim.  $\square$

**Corollary B.5.** *For every  $r > 0$ , the set*

$$G_r := \{\gamma \in \text{TGeo}^\tau(M) : \tau(\gamma_0, \gamma_1) \geq r\}$$

*is compact and uniformly equicontinuous.*

*Proof.* The compactness of the preimage of  $G_r$  under  $\mathbf{r}$  follows similarly as in the above proof and is a consequence of the compactness of  $M$ , non-total imprisonment, Arzelà–Ascoli's theorem, causal closedness, continuity of  $\tau$ , and [58, Prop. 3.17]. In particular,  $G_r$  is compact by continuity of  $\mathbf{r}$ .

To show uniform equicontinuity, let  $\varepsilon > 0$ , and let  $c > 0$  with  $\text{Lip } \gamma \leq c$  for every  $\gamma \in \text{Geo}(M)$ . Write a given  $\gamma \in G_r$  as  $\gamma := \mathbf{r}(\eta)$ , where  $\eta \in \text{TGeo}(M)$ . Since  $\psi_\eta^{-1}$  is uniformly continuous on  $[0, 1]$ , there exists  $\delta > 0$  such that

$$|\psi_\eta^{-1}(t) - \psi_\eta^{-1}(s)| \leq c^{-1} \varepsilon$$

provided  $s, t \in [0, 1]$  satisfy  $|t - s| \leq \delta$ . Therefore,

$$\mathbf{d}(\gamma_t, \gamma_s) = \mathbf{d}(\eta_{\psi_\eta^{-1}(t)}, \eta_{\psi_\eta^{-1}(s)}) \leq c |\psi_\eta^{-1}(t) - \psi_\eta^{-1}(s)| \leq \varepsilon.$$

By compactness of  $G_r$ ,  $\delta$  can be chosen independently of  $\gamma$ .  $\square$

**Corollary B.6.** *The map  $r_{\sharp} : \mathcal{P}(\text{TGeo}(M)) \rightarrow \mathcal{P}(\text{TGeo}^{\tau}(M))$  is weakly continuous.*

Here, recall the definition  $\text{TGeo}^{\tau}(M) := r(\text{TGeo}(M))$  from Subsection 2.3.6. For the convenience of the reader, we also record parts of Definition 2.13, defining

$$\begin{aligned} \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1) &:= \{\pi \in \text{OptGeo}_{\ell_p}(\mu_0, \mu_1) : \pi[M_{\ll}^2] = 1\}, \\ \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1) &:= r_{\sharp} \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1). \end{aligned}$$

**Definition B.7.** *We term a weakly continuous path  $(\mu_t)_{t \in [0,1]}$  in  $\mathcal{P}(M)$*

- a. *timelike  $\ell_p$ -geodesic if it is represented by some  $\pi \in \text{OptTGeo}_{\ell_p}(\mu_0, \mu_1)$ , and*
- b. *timelike proper-time parametrized  $\ell_p$ -geodesic if it is represented by some  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ .*

*Remark B.8.* In the smooth case, our notion of timelike proper-time parametrized  $\ell_p$ -geodesics is consistent with the notion of an  $\ell_p$ -geodesic from [66, Def. 1.1] as follows. Every timelike proper-time parametrized  $\ell_p$ -geodesic is an  $\ell_p$ -geodesic by Subsection 2.3.6, and if  $(\mu_0, \mu_1) \in \mathcal{P}_c(M)^2$  with  $\mu_0 \in \mathcal{P}^{\text{ac}}(M, \text{vol})$  is  $p$ -separated [66, Def. 4.1], these notions agree by uniqueness of  $\ell_p$ -geodesics with chronologically coupled endpoints [66, Cor. 5.9].

Recall the definition of the restriction map from (2.11).

**Proposition B.9.** *Let  $p \in (0, 1]$  and suppose that  $(M, \mathbf{d}, \ll, \leq, \tau)$  is a compact, causally closed,  $\mathcal{K}$ -globally hyperbolic Lorentzian geodesic space. Suppose that the pair  $(\mu_0, \mu_1) \in \mathcal{P}(M)^2$  is timelike  $p$ -dualizable. Then the following hold.*

- (i) *For every  $\ell_p$ -optimal  $\pi \in \Pi_{\ll}(\mu_0, \mu_1)$ , there exists  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  such that  $\pi = (e_0, e_1)_{\sharp} \pi$ .*
- (ii) *There exists at least one timelike proper-time parametrized  $\ell_p$ -geodesic from  $\mu_0$  to  $\mu_1$ .*
- (iii) *For every  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ , we have*

$$(\text{restr}_s^t)_{\sharp} \pi \in \text{OptTGeo}_{\ell_p}^{\tau}((e_s)_{\sharp} \pi, (e_t)_{\sharp} \pi)$$

*for every  $s, t \in [0, 1]$  with  $s < t$ .*

- (iv) *If  $\pi \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  and if  $\sigma$  is any nontrivial measure on  $C([0, 1]; M)$  with  $\sigma \leq \pi$ , then  $\sigma[C([0, 1]; M)]^{-1} \sigma$  is an element of  $\text{OptTGeo}_{\ell_p}^{\tau}(\sigma_0, \sigma_1)$ , where  $\sigma_i := \sigma[C([0, 1]; M)]^{-1} (e_i)_{\sharp} \sigma \in \mathcal{P}(M)$ ,  $i \in \{0, 1\}$ .*

*Furthermore, let  $(\mu_0^n)_{n \in \mathbb{N}}$  and  $(\mu_1^n)_{n \in \mathbb{N}}$  be sequences in  $\mathcal{P}(M)$  which converge weakly to  $\mu_0 \in \mathcal{P}(M)$  and  $\mu_1 \in \mathcal{P}(M)$ , respectively. If all  $\ell_p$ -optimal couplings of  $\mu_0$  and  $\mu_1$  are chronological (in particular, if  $(\mu_0, \mu_1)$  is strongly timelike  $p$ -dualizable), every sequence  $(\pi^n)_{n \in \mathbb{N}}$  with  $\pi^n \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0^n, \mu_1^n)$  has an accumulation point, and any such point belongs to  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ .*

*Proof.* All statements follow by pushing forward the corresponding statements from Proposition B.3 as well as [92, Prop. 7.16, Thm. 7.30] from  $\text{OptTGeo}_{\ell_p}(\mu_0, \mu_1) \subset \text{OptGeo}_{\ell_p}(\mu_0, \mu_1)$  to  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  by  $r_{\sharp}$ .

The only result requiring an additional comment is the last. Consider a sequence  $(\alpha^n)_{n \in \mathbb{N}}$  with  $\alpha^n \in \text{OptTGeo}_{\ell_p}(\mu_0^n, \mu_1^n)$  and  $\pi^n = r_{\sharp} \alpha^n$ . This sequence admits an accumulation point  $\alpha \in \text{OptGeo}_{\ell_p}(\mu_0, \mu_1)$  by Proposition B.3. In fact, we have  $\alpha[\text{TGeo}(M)] = 1$  by assumption on  $\mu_0$  and  $\mu_1$ . Thus  $\pi := r_{\sharp} \alpha \in \text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$  is an accumulation point of  $(\pi^n)_{n \in \mathbb{N}}$  by Corollary B.6. An analogous argument shows that all accumulation points of  $(\pi^n)_{n \in \mathbb{N}}$  belong to  $\text{OptTGeo}_{\ell_p}^{\tau}(\mu_0, \mu_1)$ .  $\square$

## REFERENCES

- [1] A. AKDEMIR, A. COLINET, R. MCCANN, F. CAVALLETTI, F. SANTARCANGELO. *Independence of synthetic curvature dimension conditions on transport distance exponent*. Trans. Amer. Math. Soc. **374** (2021), no. 8, 5877–5923.
- [2] S. B. ALEXANDER, M. GRAF, M. KUNZINGER, C. SÄMANN. *Generalized cones as Lorentzian length spaces: causality, curvature, and singularity theorems*. Preprint, [arXiv:1909.09575](https://arxiv.org/abs/1909.09575), 2019.
- [3] L. AMBROSIO, N. GIGLI. *A user's guide to optimal transport*. Modelling and optimisation of flows on networks, 1–155, Lecture Notes in Math., 2062, Fond. CIME/CIME Found. Subser., Springer, Heidelberg, 2013.
- [4] L. AMBROSIO, N. GIGLI, G. SAVARÉ. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2005. viii+333 pp.
- [5] ———. *Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below*. Invent. Math. **195** (2014), no. 2, 289–391.
- [6] ———. *Metric measure spaces with Riemannian Ricci curvature bounded from below*. Duke Math. J. **163** (2014), no. 7, 1405–1490.
- [7] ———. *Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds*. Ann. Probab. **43** (2015), no. 1, 339–404.
- [8] L. AMBROSIO, S. HONDA. *New stability results for sequences of metric measure spaces with uniform Ricci bounds from below*. Measure theory in non-smooth spaces, 1–51, Partial Differ. Equ. Meas. Theory, De Gruyter Open, Warsaw, 2017.
- [9] K. BACHER, K.-T. STURM. *Localization and tensorization properties of the curvature-dimension condition for metric measure spaces*. J. Funct. Anal. **259** (2010), no. 1, 28–56.
- [10] J. M. BARDEEN, B. CARTER, S. W. HAWKING. *The four laws of black hole mechanics*. Comm. Math. Phys. **31** (1973), 161–170.
- [11] J. K. BEEM, P. E. EHRLICH, S. MARKVORSEN, G. J. GALLOWAY. *Decomposition theorems for Lorentzian manifolds with nonpositive curvature*. J. Differential Geom. **22** (1985), no. 1, 29–42.
- [12] J. D. BEKENSTEIN. *Black holes and entropy*. Phys. Rev. D (3) **7** (1973), 2333–2346.
- [13] P. BILLINGSLEY. *Convergence of probability measures*. Second edition. Wiley Series in Probability and Statistics: Probability and Statistics. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1999. x+277 pp.
- [14] V. I. BOGACHEV. *Measure theory. II*. Springer-Verlag, Berlin, 2007, xiv+575 pp.
- [15] L. BOMBELLI, J. LEE, D. MEYER, R. D. SORKIN. *Space-time as a causal set*. Phys. Rev. Lett. **59** (1987), no. 5, 521–524.
- [16] M. BRAUN. *Good geodesics satisfying the timelike curvature-dimension condition*. Preprint, [arXiv:2205.06950](https://arxiv.org/abs/2205.06950), 2022.
- [17] ———. *Rényi's entropy on Lorentzian spaces. Singularity theorems*. In preparation.
- [18] E. BRUÉ, D. SEMOLA. *Constancy of the dimension for  $\text{RCD}(K, N)$  spaces via regularity of Lagrangian flows*. Comm. Pure Appl. Math. **73** (2020), no. 6, 1141–1204.
- [19] D. BURAGO, YU. BURAGO, S. IVANOV. *A course in metric geometry*. Graduate Studies in Mathematics, 33. American Mathematical Society, Providence, RI, 2001. xiv+415 pp.
- [20] A. BURTSCHER, P. G. LEFLOCH. *The formation of trapped surfaces in spherically-symmetric Einstein-Euler spacetimes with bounded variation*. J. Math. Pures Appl. (9) **102** (2014), no. 6, 1164–1217.
- [21] A. BURTSCHER, L. GARCÍA-HEVELING. *Time functions on Lorentzian length spaces*. Preprint, [arXiv:2108.02693](https://arxiv.org/abs/2108.02693), 2021.
- [22] F. CAVALLETTI, E. MILMAN. *The globalization theorem for the curvature-dimension condition*. Invent. Math. **226** (2021), no. 1, 1–137.
- [23] F. CAVALLETTI, A. MONDINO. *Optimal maps in essentially non-branching spaces*. Commun. Contemp. Math. **19** (2017), no. 6, 1750007, 27 pp.
- [24] ———. *Optimal transport in Lorentzian synthetic spaces, synthetic timelike Ricci curvature lower bounds and applications*. Preprint, [arXiv:2004.08934](https://arxiv.org/abs/2004.08934), 2020.
- [25] ———. *A review of Lorentzian synthetic theory of timelike Ricci curvature bounds*. Preprint, [arXiv:2204.13330](https://arxiv.org/abs/2204.13330), 2022.
- [26] D. CHRISTODOULOU. *The formation of black holes in general relativity*. EMS Monographs in Mathematics. European Mathematical Society (EMS), Zürich, 2009. x+589 pp.
- [27] P. T. CHRUSCIEL, J. D. E. GRANT. *On Lorentzian causality with continuous metrics*. Classical Quantum Gravity **29** (2012), no. 14, 145001, 32 pp.

- [28] D. CORDERO-ERAUSQUIN, R. J. MCCANN, M. SCHMUCKENSCHLÄGER. *A Riemannian interpolation inequality à la Borell, Brascamp and Lieb*. Invent. Math. **146** (2001), no. 2, 219–257.
- [29] M. DAFERMOS. *The mathematical analysis of black holes in general relativity*. Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. III, 747–772, Kyung Moon Sa, Seoul, 2014.
- [30] Q. DENG, K.-T. STURM. *Localization and tensorization properties of the curvature-dimension condition for metric measure spaces, II*. J. Funct. Anal. **260** (2011), no. 12, 3718–3725.
- [31] M. ECKSTEIN, T. MILLER. *Causality for nonlocal phenomena*. Ann. Henri Poincaré **18** (2017), no. 9, 3049–3096.
- [32] M. ERBAR, K. KUWADA, K.-T. STURM. *On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces*. Invent. Math. **201** (2015), no. 3, 993–1071.
- [33] M. ERBAR, K.-T. STURM. *Rigidity of cones with bounded Ricci curvature*. J. Eur. Math. Soc. (JEMS) **23** (2021), no. 1, 219–235.
- [34] J.-H. ESCHENBURG. *The splitting theorem for space-times with strong energy condition*. J. Differential Geom. **27** (1988), no. 3, 477–491.
- [35] F. FINSTER. *The continuum limit of causal fermion systems*. From Planck scale structures to macroscopic physics. Fundamental Theories of Physics, 186. Springer, Cham, 2016. xi+548 pp.
- [36] ———. *Causal Fermion systems: a primer for Lorentzian geometers*. J. Phys. Conf. Ser. **968** (2018), 012004, 14 pp.
- [37] G. J. GALLOWAY. *The Lorentzian splitting theorem without the completeness assumption*. J. Differential Geom. **29** (1989), no. 2, 373–387.
- [38] D. GANNON. *Singularities in nonsimply connected space-times*. J. Mathematical Phys. **16** (1975), no. 12, 2364–2367.
- [39] ———. *On the topology of spacelike hypersurfaces, singularities, and black holes*. Gen. Relativity Gravitation **7** (1976), no. 2, 219–232.
- [40] R. GEROCH. *Domain of dependence*. J. Mathematical Phys. **11** (1970), 437–449.
- [41] R. GEROCH, J. TRASCHEN. *Strings and other distributional sources in general relativity*. Phys. Rev. D (3) **36** (1987), no. 4, 1017–1031.
- [42] N. GIGLI. *The splitting theorem in non-smooth context*. Preprint, [arXiv:1302.5555](https://arxiv.org/abs/1302.5555), 2013.
- [43] ———. *On the differential structure of metric measure spaces and applications*. Mem. Amer. Math. Soc. **236** (2015), no. 1113, vi+91 pp.
- [44] ———. *Nonsmooth differential geometry — an approach tailored for spaces with Ricci curvature bounded from below*. Mem. Amer. Math. Soc. **251** (2018), no. 1196, v+161 pp.
- [45] N. GIGLI, A. MONDINO, G. SAVARÉ. *Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows*. Proc. Lond. Math. Soc. (3) **111** (2015), no. 5, 1071–1129.
- [46] M. GRAF. *Singularity theorems for  $C^1$ -Lorentzian metrics*. Comm. Math. Phys. **378** (2020), no. 2, 1417–1450.
- [47] J. B. GRIFFITHS, J. PODOLSKÝ. *Exact space-times in Einstein’s general relativity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2009. xviii+525 pp.
- [48] M. GROMOV. *Metric structures for Riemannian and non-Riemannian spaces*. With appendices by M. KATZ, P. PANSU and S. SEMMES. Translated from the French by S. M. BATES. Progress in Mathematics, 152. Birkhäuser Boston, Inc., Boston, MA, 1999. xx+585 pp.
- [49] S. W. HAWKING. *The occurrence of singularities in cosmology. I*. Proc. Roy. Soc. London Ser. A **294** (1966), 511–521.
- [50] S. W. HAWKING, G. F. R. ELLIS. *The large scale structure of space-time*. Cambridge Monographs on Mathematical Physics, No. 1. Cambridge University Press, London-New York, 1973. xi+391 pp.
- [51] S. W. HAWKING, R. PENROSE. *The singularities of gravitational collapse and cosmology*. Proc. Roy. Soc. London Ser. A **314** (1970), 529–548.
- [52] T. JACOBSON. *Thermodynamics of spacetime: the Einstein equation of state*. Phys. Rev. Lett. **75** (1995), no. 7, 1260–1263.
- [53] M. KELL, S. SUHR. *On the existence of dual solutions for Lorentzian cost functions*. Ann. Inst. H. Poincaré C Anal. Non Linéaire **37** (2020), no. 2, 343–372.
- [54] C. KETTERER. *Cones over metric measure spaces and the maximal diameter theorem*. J. Math. Pures Appl. (9) **103** (2015), no. 5, 1228–1275.
- [55] ———. *Obata’s rigidity theorem for metric measure spaces*. Anal. Geom. Metr. Spaces **3** (2015), no. 1, 278–295.

- [56] S. KLAINERMAN, I. RODNIANSKI, J. SZEFTTEL. *The bounded  $L^2$  curvature conjecture*. Invent. Math. **202** (2015), no. 1, 91–216.
- [57] M. KUNZINGER, R. STEINBAUER, M. STOJKOVIĆ, J. A. VICKERS. *Hawking's singularity theorem for  $C^{1,1}$ -metrics*. Classical Quantum Gravity **32** (2015), no. 7, 075012, 19 pp.
- [58] M. KUNZINGER, C. SÄMANN. *Lorentzian length spaces*. Ann. Global Anal. Geom. **54** (2018), no. 3, 399–447.
- [59] C. W. LEE. *A restriction on the topology of Cauchy surfaces in general relativity*. Comm. Math. Phys. **51** (1976), no. 2, 157–162.
- [60] A. LICHNEROWICZ. *Théories relativistes de la gravitation et de l'électromagnétisme. Relativité générale et théories unitaires*. Masson et Cie, Paris, 1955. xii+298 pp.
- [61] J. LOTT, C. VILLANI. *Ricci curvature for metric-measure spaces via optimal transport*. Ann. of Math. (2) **169** (2009), no. 3, 903–991.
- [62] Y. LU, E. MINGUZZI, S.-I. OHTA. *Geometry of weighted Lorentz-Finsler manifolds I: singularity theorems*. J. Lond. Math. Soc. (2) **104** (2021), no. 1, 362–393.
- [63] ———. *Geometry of weighted Lorentz-Finsler manifolds II: a splitting theorem*. Preprint, arXiv:2110.10308, 2021.
- [64] M. MARS, J. M. M. SENOVILLA. *Geometry of general hypersurfaces in spacetime: junction conditions*. Classical Quantum Gravity **10** (1993), no. 9, 1865–1897.
- [65] K. MARTIN, P. PANANGADEN. *A domain of spacetime intervals in general relativity*. Comm. Math. Phys. **267** (2006), no. 3, 563–586.
- [66] R. J. MCCANN. *Displacement convexity of Boltzmann's entropy characterizes the strong energy condition from general relativity*. Camb. J. Math. **8** (2020), no. 3, 609–681.
- [67] R. J. MCCANN, C. SÄMANN. *A Lorentzian analog for Hausdorff dimension and measure*. Preprint, arXiv:2110.04386, 2021.
- [68] A. MONDINO, A. NABER. *Structure theory of metric measure spaces with lower Ricci curvature bounds*. J. Eur. Math. Soc. (JEMS) **21** (2019), no. 6, 1809–1854.
- [69] A. MONDINO, S. SUHR. *An optimal transport formulation of the Einstein equations of general relativity*. Preprint, arXiv:1810.13309, 2018.
- [70] S.-I. OHTA. *On the measure contraction property of metric measure spaces*. Comment. Math. Helv. **82** (2007), no. 4, 805–828.
- [71] B. O'NEILL. *Semi-Riemannian geometry. With applications to relativity*. Pure and Applied Mathematics, 103. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983. xiii+468 pp.
- [72] F. OTTO, C. VILLANI. *Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality*. J. Funct. Anal. **173** (2000), no. 2, 361–400.
- [73] R. PENROSE. *Gravitational collapse and space-time singularities*. Phys. Rev. Lett. **14** (1965), 57–59.
- [74] ———. *The geometry of impulsive gravitational waves*. General relativity (papers in honour of J. L. Synge), pp. 101–115. Clarendon Press, Oxford, 1972.
- [75] T. RAJALA. *Interpolated measures with bounded density in metric spaces satisfying the curvature-dimension conditions of Sturm*. J. Funct. Anal. **263** (2012), no. 4, 896–924.
- [76] ———. *Improved geodesics for the reduced curvature-dimension condition in branching metric spaces*. Discrete Contin. Dyn. Syst. **33** (2013), no. 7, 3043–3056.
- [77] T. RAJALA, K.-T. STURM. *Non-branching geodesics and optimal maps in strong  $CD(K, \infty)$ -spaces*. Calc. Var. Partial Differential Equations **50** (2014), no. 3-4, 831–846.
- [78] A. D. RENDALL. *Theorems on existence and global dynamics for the Einstein equations*. Living Rev. Relativ. **5** (2002), 2002–6, 62 pp.
- [79] G. SAVARÉ. *Self-improvement of the Bakry-Émery condition and Wasserstein contraction of the heat flow in  $RCD(K, \infty)$  metric measure spaces*. Discrete Contin. Dyn. Syst. **34** (2014), no. 4, 1641–1661.
- [80] B. SCHINNERL, R. STEINBAUER. *A note on the Gannon-Lee theorem*. Lett. Math. Phys. **111** (2021), no. 6, Paper No. 142, 17 pp.
- [81] E. H. SPANIER. *Algebraic topology*. McGraw-Hill Book Co., New York-Toronto, Ont.-London 1966 xiv+528 pp.
- [82] C. SORMANI, C. VEGA. *Null distance on a spacetime*. Classical Quantum Gravity **33** (2016), no. 8, 085001, 29 pp.
- [83] R. STEINBAUER. *The singularity theorems of general relativity and their low regularity extensions*. Preprint, arXiv:2206.05939, 2022.
- [84] R. STEINBAUER, J. A. VICKERS. *On the Geroch-Traschen class of metrics*. Classical Quantum Gravity **26** (2009), no. 6, 065001, 19 pp.
- [85] K.-T. STURM. *On the geometry of metric measure spaces. I*. Acta Math. **196** (2006), no. 1, 65–131.

- [86] ———. *On the geometry of metric measure spaces. II.* Acta Math. **196** (2006), no. 1, 133–177.
- [87] S. SUHR. *Theory of optimal transport for Lorentzian cost functions.* Münster J. Math. **11** (2018), no. 1, 13–47.
- [88] K. SUZUKI. *Convergence of Brownian motions on metric measure spaces under Riemannian curvature-dimension conditions.* Electron. J. Probab. **24** (2019), Paper No. 102, 36 pp.
- [89] E. VERLINDE. *On the origin of gravity and the laws of Newton.* J. High Energy Phys. **2011** (2011), no. 4, 029, 27 pp.
- [90] J. A. VICKERS. *Quasi-regular singularities and cosmic strings.* Classical Quantum Gravity **7** (1990), no. 5, 731–741.
- [91] J. A. VICKERS, J. P. WILSON. *Generalized hyperbolicity in conical spacetimes.* Classical Quantum Gravity **17** (2000), no. 6, 1333–1360.
- [92] C. VILLANI. *Optimal transport. Old and new.* Grundlehren der mathematischen Wissenschaften, 338. Springer-Verlag, Berlin, 2009. xxii+973 pp.
- [93] M.-K. VON RENESSE, K.-T. STURM. *Transport inequalities, gradient estimates, entropy, and Ricci curvature.* Comm. Pure Appl. Math. **58** (2005), no. 7, 923–940.
- [94] E. WOOLGAR, W. WYLIE. *Cosmological singularity theorems and splitting theorems for  $N$ -Bakry-Émery spacetimes.* J. Math. Phys. **57** (2016), no. 2, 022504, 12 pp.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO BAHEN CENTRE, 40 ST. GEORGE STREET ROOM 6290, TORONTO, ONTARIO M5S 2E4, CANADA

*Email address:* braun@math.toronto.edu