

The Frobenius number for sequences of arithmetic progressions associated with number of solutions

Takao Komatsu

Department of Mathematical Sciences, School of Science
Zhejiang Sci-Tech University
Hangzhou 310018 China
komatsu@zstu.edu.cn

Haotian Ying

Department of Mathematical Sciences, School of Science
Zhejiang Sci-Tech University
Hangzhou 310018 China
tomyinght@gmail.com

MR Subject Classifications: Primary 11D07; Secondary 05A15, 05A17, 05A19,
11B68, 11D04, 11P81, 20M14

Abstract

For given positive integers a_1, a_2, \dots, a_k with $\gcd(a_1, a_2, \dots, a_k) = 1$, consider the number of nonnegative solutions (x_1, x_2, \dots, x_k) of the linear equation $a_1x_1 + a_2x_2 + \dots + a_kx_k = n$ for a positive integer n . For a given nonnegative integer p , there is a maximum n such that the number of nonnegative integer solutions is at most p , and it is very attractive to find the explicit formula of the maximum n . In fact, when $p = 0$, such a problem of finding the maximum integer n is called the linear Diophantine problem of Frobenius, and this maximum n is called the Frobenius number. The explicit formula for two variables is known not only for $p = 0$ but also for $p > 0$, but when there are three or more variables, it is difficult even in the special case of $p = 0$. For $p > 0$, it is not only more difficult, but no explicit formula has been found. In this paper, an explicit formula of such a generalized Frobenius number is given for the sequence of arithmetic progressions, in particular, for three variables. Further, explicit formulas are given not only for the maximum value of n such that the number of solutions is at most p , but also for the total number, the sum, and the weighted sum of such solutions.

Keywords: Frobenius problem, Frobenius numbers, number of representations, arithmetic progressions

1 Introduction

For integer $k \geq 2$, consider a set of positive integers $A = \{a_1, \dots, a_k\}$ with $\gcd(A) = \gcd\{a_1, \dots, a_k\} = 1$. Let $\text{NR}(A) = \text{NR}(a_1, a_2, \dots, a_k)$ denote the set of positive integers nonrepresentable in terms of a_1, a_2, \dots, a_k . Then, the well-known linear Diophantine problem, posed by Sylvester [24], but known as the Frobenius problem, is the problem to determine the Frobenius number $g(A)$ and the Sylvester number (genus) $n(A)$, defined by

$$g(A) = \max \text{NR} \quad \text{and} \quad n(A) = |\text{NR}(A)|,$$

respectively. The Frobenius Problem has been also known as the *Coin Exchange Problem* (or Postage Stamp Problem / Chicken McNugget Problem), which has a long history and is one of the problems that has attracted many people as well as experts. For two variables $A = \{a, b\}$, it is shown that

$$g(a, b) = (a - 1)(b - 1) - 1 \quad \text{and} \quad n(a, b) = \frac{(a - 1)(b - 1)}{2}. \quad (1)$$

([23, 24]). The related problem of determining the function

$$s(A) = \sum_{n \in \text{NR}(A)} n,$$

which is called Sylvester sum nowadays, is introduced by Brown and Shiue [9] and for two variables $A = \{a, b\}$, it is shown that

$$s(a, b) = \frac{1}{12}(a - 1)(b - 1)(2ab - a - b - 1). \quad (2)$$

However, for three or more variables, it very complicated to find a general explicit formula for Frobenius number, Sylvester number and Sylvester sum. Only for some special cases, explicit formulas have been found, including arithmetic, geometric-like, Fibonacci, Mersenne, and triangular (see, e.g., [20] and references therein). Some inexplicit formulas for the Frobenius number in three variables can be seen in [27].

Concerning arithmetic numbers, let a, d and k be positive integers with $\gcd(a, d) = 1$ and $2 \leq k \leq a$. Roberts [19] found the Frobenius number for arithmetic sequences:

$$g(a, a + d, \dots, a + (k - 1)d) = \left\lfloor \frac{a - 2}{k - 1} \right\rfloor a + (a - 1)d.$$

The case $d = 1$ had been found by Brauer [6]. Selmer [21] generalized Roberts' result by giving a formula for almost arithmetic sequences. For a positive integer h ,

$$g(a, ha + d, \dots, ha + (k - 1)d) = \left(h \left\lfloor \frac{a - 2}{k - 1} \right\rfloor + h - 1 \right) a + (a - 1)d.$$

Selmer also gave an explicit formula for the Sylvester number $n(a, ha + d, \dots, ha + (k - 1)d)$. Some formulas for the Sylvester sum $s(a, ha + d, \dots, ha + (k - 1)d)$ and its variations are given in [13].

On the other hand, to find the number of representations $d(n; A) = d(n; a_1, a_2, \dots, a_k)$ to $a_1x_1 + a_2x_2 + \dots + a_kx_k = n$ for a given positive integer n is also one of the most important and interesting topics. This number is equal to the coefficient of x^n in $1/(1 - x^{a_1})(1 - x^{a_2}) \dots (1 - x^{a_k})$. Sylvester [22] and Cayley [10] showed that $d(n; a_1, a_2, \dots, a_k)$ can be expressed as the sum of a polynomial in n of degree $k - 1$ and a periodic function of period $a_1a_2 \dots a_k$. For two variables, a formula for $d(n; a_1, a_2)$ is obtained in [25]. For three variables, in [12], the periodic function part is expressed in terms of trigonometric functions, and its results have been improved in [5] by using floor functions so that three variables case can be easily worked with in his formula. In [3], the explicit formula for the polynomial part is derived by using Bernoulli numbers.

In this paper, we are interested in one of the most general and most natural types of Frobenius numbers, which focuses on the number of solutions. For a given nonnegative integer p , consider the largest integer $g_p(A) = g_p(a_1, a_2, \dots, a_k)$ such that the number of expressions that can be represented by a_1, a_2, \dots, a_k is at most p . That is, all integers larger than $g_p(A)$ have at least the number of representations of $p + 1$ or more. This generalized Frobenius number $g_p(A)$ is called the p -Frobenius number [14, 15], which is also called the k -Frobenius number [8] or the s -Frobenius number [11]. When $p = 0$, $g(A) = g_0(A)$ is the original Frobenius number. One can consider the largest integer $g_p^*(a_1, a_2, \dots, a_k)$ that has exactly p distinct representations (see, e.g., [8, 11]). However, in this case, the ordering $g_0^* \leq g_1^* \leq \dots$ may not hold. For example, $g_{17}^*(2, 5, 7) = 43 > g_{18}^*(2, 5, 7) = 42$. In addition, for some j , g_j^* may not exist. For example, $g_{22}^*(2, 5, 7)$ does not exist because there is no positive integer whose number of representations is exactly 22. Therefore, in this paper we treat with $g_p(A)$. In the same stream, we define the generalized Sylvester number (p -genus) $n_p(A) = n_p(a_1, a_2, \dots, a_k)$ and generalized Sylvester sum (p -Sylvester sum) $s_p(A) = s_p(a_1, a_2, \dots, a_k)$ by

$$n_p(A) = \sum_{d(n; A) \leq p} 1 \quad \text{and} \quad s_p(A) = \sum_{d(n; A) \leq p} n,$$

respectively.

For two variables, it is easy to find explicit formulas of $g_p(a, b)$, $n_p(a, b)$ and $s_p(a, b)$ (see, e.g., [2, 4]). However, when there are three or more variables, it is sufficiently difficult to obtain the explicit formula of g_p or n_p even in a special case even when $p = 0$, and it is even more difficult when $p > 0$, and in the case of a special triplet even one example had not been found. However, very recently, in [14] and [15], for $p \geq 0$ we finally succeeded in obtaining explicit formulas in the sequences of triangular numbers and repunits, respectively.

In this paper, we give explicit formulas of $g_p(a, a + d, \dots, a + (k - 1)d)$ with $\gcd(a, d) = 1$, in particular, for three variables. When $p = 0$, the Frobenius number of the sequence of arithmetic progressions is give in [21]. Let a and d be positive integers with $\gcd(a, d) = 1$ and $a \geq 3$.

2 Preliminaries

For a positive integer p and a set of positive integers $A = \{a_1, a_2, \dots, a_k\}$ with $\gcd(A) = 1$, denote by $R_p(A)$ the set of all nonnegative integers whose representations in terms of a_2, \dots, a_k with nonnegative integral coefficients have at least p ways. Note that when $p = 0$, $R_1 \cup \text{NR}(A) = \mathbb{N} \cup \{0\}$ (the set of nonnegative integers). We introduce the Apéry set (see [1]) below in order to obtain the formulas for $g_p(A)$, $n_p(A)$ and $s_p(A)$. Without loss of generality, we assume that $a_1 = \min(A)$.

Definition 1. *Let p be a nonnegative integer. For a set of positive integers $A = \{a_1, a_2, \dots, a_k\}$ with $\gcd(A) = 1$ and $a_1 = \min(A)$ we denote by*

$$\text{Ap}_p(A) = \text{Ap}_p(a_1, a_2, \dots, a_k) = \{m_0^{(p)}, m_1^{(p)}, \dots, m_{a_1-1}^{(p)}\},$$

the p -Apéry set of A , where $m_i^{(p)}$ is the least positive integer of $R_{p+1}(A)$ satisfying $m_i^{(p)} \equiv i \pmod{a_1}$ ($1 \leq i \leq a_1 - 1$). Note that $m_0^{(0)}$ is defined to be 0.

It follows that for each p ,

$$\text{Ap}_p(A) \equiv \{0, 1, \dots, a_1 - 1\} \pmod{a_1}.$$

When $k \geq 3$, it is hard to find any explicit form of $g_p(A)$ as well as $n_p(A)$ and $s_p(A)$. Nevertheless, the following convenient formulas are known (For a more general case, see [16]). Though finding $m_j^{(p)}$ is enough hard in general, we can obtain it for some special sequences (a_1, a_2, \dots, a_k) .

Lemma 1. *Let k, p and μ be integers with $k \geq 2, p \geq 0$ and $\mu \geq 1$. Assume that $\gcd(a_1, a_2, \dots, a_k) = 1$. We have*

$$g_p(a_1, a_2, \dots, a_k) = \left(\max_{0 \leq j \leq a_1-1} m_j^{(p)} \right) - a_1, \quad (3)$$

$$n_p(a_1, a_2, \dots, a_k) = \frac{1}{a_1} \sum_{j=0}^{a_1-1} m_j^{(p)} - \frac{a_1-1}{2}, \quad (4)$$

$$s_p(a_1, a_2, \dots, a_k) = \frac{1}{2a_1} \sum_{j=0}^{a_1-1} (m_j^{(p)})^2 - \frac{1}{2} \sum_{j=0}^{a_1-1} m_j^{(p)} + \frac{a_1^2-1}{12}. \quad (5)$$

Remark. When $p = 0$, the formulas (3), (4) and (5) reduce to the formulas by Brauer and Shockley [7], Selmer [21], and Tripathi [26], respectively:

$$\begin{aligned} g(a_1, a_2, \dots, a_k) &= \left(\max_{1 \leq j \leq a_1-1} m_j \right) - a_1, \\ n(a_1, a_2, \dots, a_k) &= \frac{1}{a_1} \sum_{j=0}^{a_1-1} m_j - \frac{a_1-1}{2}, \\ s(a_1, a_2, \dots, a_k) &= \frac{1}{2a_1} \sum_{j=0}^{a_1-1} (m_j)^2 - \frac{1}{2} \sum_{j=0}^{a_1-1} m_j + \frac{a_1^2-1}{12}, \end{aligned}$$

where $m_j = m_j^{(0)}$ ($1 \leq j \leq a_1 - 1$) with $m_0 = m_0^{(0)} = 0$.

3 Three variables

Theorem 1. *Let a and d be integers with $a \geq 3, d > 0$ and $\gcd(a, d) = 1$. Then for $0 \leq p \leq \lfloor a/2 \rfloor$,*

$$\begin{aligned} g_p(a, a+d, a+2d) &= (a+2d)p + \left\lfloor \frac{a-2}{2} \right\rfloor a + (a-1)d, \\ n_p(a, a+d, a+2d) &= \begin{cases} (2a+2d-1-p)p + \frac{(a-1)(a+2d-1)}{4} & \text{if } a \text{ is odd;} \\ (2a+2d-1-p)p + \frac{(a-1)(a+2d-1)+1}{4} & \text{if } a \text{ is even.} \end{cases} \end{aligned}$$

Remark. When $p = 0$, the formulas reduce to

$$g_0(a, a + d, a + 2d) = \left\lfloor \frac{a - 2}{2} \right\rfloor a + (a - 1)d$$

and

$$n_0(a, a + d, a + 2d) = \begin{cases} \frac{(a-1)(a+2d-1)}{4} & \text{if } a \text{ is odd;} \\ \frac{(a-1)(a+2d-1)+1}{4} & \text{if } a \text{ is even.} \end{cases}$$

respectively, which are [21, (3.9),(3.10)] when $k = 3$.

Let $r_{x_2, x_3} = (a + d)x_2 + (a + 2d)x_3$ for nonnegative integers x_2 and x_3 . In the following tables, denote it by (x_2, x_3) for simplicity.

When a is odd, as seen in [21, (3.6)], $\text{Ap}_0(A_3)$ ($A_3 = \{a, a + d, a + 2d\}$) as the complete residue system (minimal system) modulo a is given by Table 1.

$(0, 0)$	$(1, 0)$
$(0, 1)$	$(1, 1)$
\vdots	\vdots
$(0, \frac{a-3}{2})$	$(1, \frac{a-3}{2})$
$(0, \frac{a-1}{2})$	

Table 1: Complete residue system $\text{Ap}_0(A_3)$ for odd a

Concerning the complete residue system $\text{Ap}_1(A_3)$, each congruent value modulo a moves up one line to the upper right block. However, only the two values in the top row move to fill the gap below the first block. Namely, for $x_2 \geq 1$

$$r_{0, x_2} \equiv r_{2, x_2 - 1}, \quad r_{1, x_2} \equiv r_{3, x_2 - 1} \pmod{a}$$

and

$$r_{0, 0} \equiv r_{1, \frac{a-1}{2}}, \quad r_{1, 0} \equiv r_{0, \frac{a+1}{2}} \pmod{a}.$$

Since

$$\begin{aligned} 3(a + d) + (a + 2d)x_3 &= a + (a + d) + (a + 2d)(x_3 + 1), \\ 2(a + d) + (a + 2d)x_3 &= a + (a + 2d)(x_3 + 1) \quad (x_3 \geq 0), \\ (a + d) + \frac{a - 1}{2}(a + 2d) &= \left(\frac{a + 1}{2} + d \right) a, \\ \frac{a + 1}{2}(a + 2d) &= \left(\frac{a - 1}{2} + d \right) a + (a + d), \end{aligned}$$

we can know that each element in $\text{Ap}_1(A_3)$ has exactly two expressions in terms of $(a, a + d, a + 2d)$. Note that each element minus a has only one expression which is yielded from the right-hand side, that is,

$$(a + d) + (a + 2d)(x_3 + 1), \quad (a + 2d)(x_3 + 1) \quad (x_3 \geq 0),$$

$$\left(\frac{a-1}{2} + d\right)a, \quad \left(\frac{a-3}{2} + d\right)a + (a + d).$$

Considering the maximal value, it is clear that

$$r_{0, \frac{a+1}{2}} > r_{1, \frac{a-1}{2}} > r_{2, \frac{a-3}{2}} > r_{3, \frac{a-5}{2}}.$$

$(0, 0)$	$(1, 0)$	$(2, 0)$	$(3, 0)$
$(0, 1)$	$(1, 1)$	$(2, 1)$	$(3, 1)$
\vdots	\vdots	\vdots	\vdots
$(0, \frac{a-5}{2})$	$(1, \frac{a-5}{2})$	$(2, \frac{a-5}{2})$	$(3, \frac{a-5}{2})$
$(0, \frac{a-3}{2})$	$(1, \frac{a-3}{2})$	$(2, \frac{a-3}{2})$	
$(0, \frac{a-1}{2})$	$(1, \frac{a-1}{2})$		
$(0, \frac{a+1}{2})$			

Table 2: Complete residue system $\text{Ap}_1(A_3)$ from $\text{Ap}_0(A_3)$ for odd a

Concerning the complete residue system $\text{Ap}_2(A_3)$, each congruent value modulo a moves up one line to the upper right block (the third block). However, only the two values in the top row in the second block move to fill the gap below the first block. Namely, for $x_2 \geq 1$

$$r_{2, x_2} \equiv r_{4, x_2-1}, \quad r_{3, x_2} \equiv r_{5, x_2-1} \pmod{a}$$

and

$$r_{2, 0} \equiv r_{1, \frac{a+1}{2}}, \quad r_{3, 0} \equiv r_{0, \frac{a+3}{2}} \pmod{a}.$$

Since

$$\begin{aligned} 5(a + d) + (a + 2d)x_3 &= a + 3(a + d) + (a + 2d)(x_3 + 1) \\ &= 2a + (a + d) + (a + 2d)(x_3 + 2), \\ 4(a + d) + (a + 2d)x_3 &= a + 2(a + d) + (a + 2d)(x_3 + 1) \\ &= 2a + (a + 2d)(x_3 + 2) \quad (x_3 \geq 0), \\ 3(a + d) + \frac{a-3}{2}(a + 2d) &= a + (a + d) + \frac{a-1}{2}(a + 2d) \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{a+3}{2} + d \right) a, \\
2(a+d) + \frac{a-1}{2}(a+2d) &= a + \frac{a+1}{2}(a+2d) \\
&= \left(\frac{a+1}{2} + d \right) a + (a+d), \\
(a+d) + \frac{a+1}{2}(a+2d) &= \left(\frac{a-1}{2} + d \right) a + 2(a+d) \\
&= \left(\frac{a+1}{2} + d \right) a + (a+2d), \\
\frac{a+3}{2}(a+2d) &= \left(\frac{a-3}{2} + d \right) a + 3(a+d) \\
&= \left(\frac{a-1}{2} + d \right) a + (a+d) + (a+2d),
\end{aligned}$$

we can know that each element in $\text{Ap}_1(A_3)$ has exactly three expressions in terms of $(a, a+d, a+2d)$. Note that each element minus a has two expressions which are yielded from the right-hand side, that is,

$$\begin{aligned}
3(a+d) + (a+2d)(x_3+1) &= a + (a+d) + (a+2d)(x_3+2), \\
2(a+d) + (a+2d)(x_3+1) &= a + (a+2d)(x_3+2) \quad (x_3 \geq 0), \\
(a+d) + \frac{a-1}{2}(a+2d) &= \left(\frac{a+1}{2} + d \right) a, \\
\frac{a+1}{2}(a+2d) &= \left(\frac{a-1}{2} + d \right) a + (a+d), \\
\left(\frac{a-3}{2} + d \right) a + 2(a+d) &= \left(\frac{a-1}{2} + d \right) a + (a+2d), \\
\left(\frac{a-3}{2} + d - 1 \right) a + 3(a+d) &= \left(\frac{a-3}{2} + d \right) a + (a+d) + (a+2d).
\end{aligned}$$

Considering the maximal value, it is clear that

$$r_{0, \frac{a+3}{2}} > r_{1, \frac{a+1}{2}} > r_{2, \frac{a-1}{2}} > r_{3, \frac{a-3}{2}} > r_{4, \frac{a-5}{2}} > r_{5, \frac{a-7}{2}}.$$

If this process is continued for $p = 3, 4, \dots$, when $p = (a-1)/2$, the state shown in Table 4 is reached. Here, the shaded cell parts show the elements of $\text{Ap}_{\frac{a-1}{2}}(A_3)$. Elements with the same residues modulo a move in the following positions according to $p = 0, 1, 2, \dots, (a-3)/2, (a-1)/2$.

$$(0, 0) \rightarrow \left(1, \frac{a-1}{2}\right) \rightarrow \left(3, \frac{a-3}{2}\right) \rightarrow \dots \rightarrow (a-4, 2) \rightarrow (a-2, 1)$$

		(2, 0)	(3, 0)	(4, 0)	(5, 0)
		(2, 1)	(3, 1)	(4, 1)	(5, 1)
		⋮	⋮	⋮	⋮
		⋮	⋮	(4, $\frac{a-7}{2}$)	(5, $\frac{a-7}{2}$)
		(2, $\frac{a-5}{2}$)	(3, $\frac{a-5}{2}$)	(4, $\frac{a-5}{2}$)	
		(2, $\frac{a-3}{2}$)	(3, $\frac{a-3}{2}$)		
	(1, $\frac{a-1}{2}$)	(2, $\frac{a-1}{2}$)			
(0, $\frac{a+1}{2}$)	(1, $\frac{a+1}{2}$)				
(0, $\frac{a+3}{2}$)					

Table 3: Complete residue system $\text{Ap}_2(A_3)$ from $\text{Ap}_1(A_3)$ for odd a

$$\begin{aligned}
(1, 0) &\rightarrow (0, \frac{a+1}{2}) \rightarrow (2, \frac{a-1}{2}) \rightarrow \cdots \rightarrow (a-5, 3) \rightarrow (a-3, 2) \\
(0, 1) &\rightarrow (2, 0) \rightarrow (1, \frac{a+1}{2}) \rightarrow \cdots \rightarrow (a-6, 4) \rightarrow (a-4, 3) \\
(1, 1) &\rightarrow (3, 0) \rightarrow (0, \frac{a+3}{2}) \rightarrow \cdots \rightarrow (a-7, 5) \rightarrow (a-5, 4) \\
&\dots \\
(0, \frac{a-3}{2}) &\rightarrow (2, \frac{a-5}{2}) \rightarrow (4, \frac{a-7}{2}) \rightarrow \cdots \rightarrow (a-3, 0) \rightarrow (1, a-2) \\
(1, \frac{a-3}{2}) &\rightarrow (3, \frac{a-5}{2}) \rightarrow (5, \frac{a-7}{2}) \rightarrow \cdots \rightarrow (a-2, 0) \rightarrow (0, a-1) \\
(0, \frac{a-1}{2}) &\rightarrow (2, \frac{a-3}{2}) \rightarrow (4, \frac{a-5}{2}) \rightarrow \cdots \rightarrow (a-3, 1) \rightarrow (a-1, 0)
\end{aligned}$$

(0, 0)	(1, 0)	(2, 0)	(3, 0)	⋯	⋯	(a-3, 0)	(a-2, 0)	(a-1, 0)
(0, 1)	(1, 1)	(2, 1)	(3, 1)			(a-3, 1)	(a-2, 1)	
⋮	⋮	⋮	⋮			(a-3, 2)		
⋮	⋮	⋮	⋮					
(0, $\frac{a-3}{2}$)	(1, $\frac{a-3}{2}$)	(2, $\frac{a-3}{2}$)	(3, $\frac{a-3}{2}$)					
(0, $\frac{a-1}{2}$)	(1, $\frac{a-1}{2}$)	(2, $\frac{a-1}{2}$)						
(0, $\frac{a+1}{2}$)	(1, $\frac{a+1}{2}$)							
(0, $\frac{a+3}{2}$)						(3, a-4)		
⋮						(2, a-3)		
⋮						(1, a-2)		
(0, a-1)								

Table 4: Complete residue system $\text{Ap}_p(A_3)$ for odd a

Indeed, each element $m_j^{(\frac{a-1}{2})}$ in $\text{Ap}_{\frac{a-1}{2}}(A_3)$ has exactly $(a+1)/2$ expres-

sions because

$$\begin{aligned}
(a-1)(a+2d) &= (d+i)a + (a-2i)(a+d) + (i-1)(a+2d) \\
&\quad \left(1 \leq i \leq \frac{a-1}{2}\right), \\
(a+d) + (a-2)(a+2d) &= (d+i+1)a + (a-2i-1)(a+d) \\
&\quad + (i-1)(a+2d) \quad \left(1 \leq i \leq \frac{a-1}{2}\right), \\
2(a+d) + (a-3)(a+2d) &= a + (a-2)(a+2d) \\
&= (d+i+2)a + (a-2i-2)(a+d) \\
&\quad + (i-1)(a+2d) \quad \left(1 \leq i \leq \frac{a-3}{2}\right), \\
3(a+d) + (a-4)(a+2d) &= a + (a+d) + (a-3)(a+2d) \\
&= (d+i+3)a + (a-2i-3)(a+d) \\
&\quad + (i-1)(a+2d) \quad \left(1 \leq i \leq \frac{a-3}{2}\right), \\
&\quad \dots \\
(a-3)(a+d) + 2(a+2d) &= ia + (a-2i-3)(a+d) + (i+2)(a+2d) \\
&\quad \left(1 \leq i \leq \frac{a-3}{2}\right) \\
&= (a+d-2)a + (a+d), \\
(a-2)(a+d) + (a+2d) &= ia + (a-2i-2)(a+d) + (i+1)(a+2d) \\
&\quad \left(1 \leq i \leq \frac{a-3}{2}\right) \\
&= (a+d-1)a, \\
(a-1)(a+d) &= ia + (a-2i-1)(a+d) + i(a+2d) \\
&\quad \left(1 \leq i \leq \frac{a-1}{2}\right).
\end{aligned}$$

But any element $m_j^{\binom{a-1}{2}} - a$ has $(a-1)/2$ expressions.

Considering the maximal value, it is clear that

$$r_{0,a-1} > r_{1,a-2} > r_{2,a-3} > r_{3,a-4} > \dots > r_{a-3,2} > r_{a-2,1} > r_{a-1,0}.$$

However, such a process until $0 \leq p \leq (a-1)/2$ cannot be continued further. When $p = (a+1)/2$, the same residue of $r_{a-1,0}$ modulo a comes to the position $(1, a-1)$ and no element comes to the bottom left position $(0, a)$. Thus, after that, the pattern shifts and is complicated, so it becomes difficult to determine where the maximum element of $\text{Ap}_p(A_3)$ is.

Therefore, when a is odd, for $0 \leq p \leq (a-1)/2$,

$$\begin{aligned} g_p(a, a+d, a+2d) &= \binom{\frac{a-1}{2} + p}{2} (a+2d) - a \\ &= (a+2d)p + \frac{a(a-3) + 2(a-1)d}{2}. \end{aligned}$$

Concerning the number of the representations, we need the summation of the elements in $\text{Ap}_p(A_3)$. The elements in the staircase are

$$\begin{aligned} \left(0, \frac{a-1}{2} + p\right), \left(1, \frac{a-1}{2} + p - 1\right), \dots, \\ \left(2p-2, \frac{a-1}{2} - p + 2\right), \left(2p-1, \frac{a-1}{2} - p + 1\right), \end{aligned}$$

and the elements in the large block are

$$(2p, 0), \dots, \left(2p, \frac{a-1}{2} - p\right), (2p+1, 0), \dots, \left(2p+1, \frac{a-1}{2} - p - 1\right).$$

Hence,

$$\begin{aligned} &\sum_{j=0}^{a-1} m_j^{(p)} \\ &= (0+1+\dots+(2p-1))(a+d) \\ &\quad + \left(\left(\frac{a-1}{2} - p + 1\right) + \left(\frac{a-1}{2} - p + 2\right) + \dots + \left(\frac{a-1}{2} + p\right)\right)(a+2d) \\ &\quad + 2p\left(\frac{a+1}{2} - p\right)(a+d) + \left(0+1+\dots+\left(\frac{a-1}{2} - p\right)\right)(a+2d) \\ &\quad + (2p+1)\left(\frac{a+1}{2} - p - 1\right)(a+d) \\ &\quad + \left(0+1+\dots+\left(\frac{a-1}{2} - p - 1\right)\right)(a+2d) \\ &= \frac{(2p-1)(2p)}{2}(a+d) \\ &\quad + \left(\frac{(a-1+2p)(a+1+2p)}{8} - \frac{(a-1-2p)(a+1-2p)}{8}\right)(a+2d) \\ &\quad + p(a+1-2p)(a+d) + \frac{(a-1-2p)(a+1-2p)}{8}(a+2d) \end{aligned}$$

$$\begin{aligned}
& + \frac{2p+1}{2}(a-2p-1)(a+d) + \frac{(a-3-2p)(a-1-2p)}{8}(a+2d) \\
& = \frac{a}{4}((a+d)^2 - (d+1)^2 - 4p^2 + 4(2a+2d-1)p). \tag{6}
\end{aligned}$$

Hence, by Lemma 1 (4)

$$\begin{aligned}
& n_p(a, a+d, a+2d) \\
& = \frac{1}{4}((a+d)^2 - (d+1)^2 - 4p^2 + 4(2a+2d-1)p) - \frac{a-1}{2} \\
& = \frac{(a-1)(a+2d-1)}{4} - p^2 + (2a+2d-1)p.
\end{aligned}$$

When a is even, by [21], $\text{Ap}_0(A_3)$ ($A_3 = \{a, a+d, a+2d\}$) as the complete residue system modulo a is given by the following figure.

$(0, 0)$	$(1, 0)$
$(0, 1)$	$(1, 1)$
\vdots	\vdots
$(0, \frac{a}{2} - 1)$	$(1, \frac{a}{2} - 1)$

Table 5: Complete residue system $\text{Ap}_0(A_3)$ for even a

Each congruent value modulo a moves up one line to the upper right block. However, only the two values in the top row move to fill the gap below the first block. Namely, for $x_2 \geq 1$

$$r_{0,x_2} \equiv r_{2,x_2-1}, \quad r_{1,x_2} \equiv r_{3,x_2-1} \pmod{a}$$

and

$$r_{0,0} \equiv r_{0,\frac{a}{2}}, \quad r_{1,0} \equiv r_{1,\frac{a}{2}} \pmod{a}.$$

When the process continued for $p = 1, 2, \dots, a/2$, the state shown in Table 6 is reached. Here, the shaded cell parts show the elements of $\text{Ap}_{\frac{a}{2}}(A_3)$. Elements with the same residues modulo a move in the following positions according to $p = 0, 1, 2, \dots, a/2 - 1, a/2$.

$$\begin{aligned}
(0, 0) & \rightarrow (1, \frac{a}{2}) \rightarrow (2, \frac{a}{2} - 1) \rightarrow \dots \rightarrow (a-4, 2) \rightarrow (a-2, 1) \\
(1, 0) & \rightarrow (1, \frac{a}{2}) \rightarrow (3, \frac{a}{2} - 1) \rightarrow \dots \rightarrow (a-3, 2) \rightarrow (a-1, 1) \\
(0, 1) & \rightarrow (2, 0) \rightarrow (0, \frac{a}{2} + 1) \rightarrow \dots \rightarrow (a-6, 4) \rightarrow (a-4, 3) \\
(1, 1) & \rightarrow (3, 0) \rightarrow (1, \frac{a}{2} + 1) \rightarrow \dots \rightarrow (a-5, 4) \rightarrow (a-3, 3)
\end{aligned}$$

$$\dots$$

$$(0, \frac{a}{2} - 1) \rightarrow (2, \frac{a}{2} - 2) \rightarrow (4, \frac{a}{2} - 3) \rightarrow \dots \rightarrow (a - 2, 0) \rightarrow (0, a - 1)$$

$$(1, \frac{a}{2} - 1) \rightarrow (3, \frac{a}{2} - 2) \rightarrow (5, \frac{a}{2} - 3) \rightarrow \dots \rightarrow (a - 1, 0) \rightarrow (1, a - 1)$$

(0, 0)	(1, 0)	(2, 0)	(3, 0)	(4, 0)	(5, 0)	(a - 4, 0)	(a - 3, 0)	(a - 2, 0)	(a - 1, 0)
(0, 1)	(1, 1)	(2, 1)	(3, 1)	(4, 1)	(5, 1)			(a - 4, 1)	(a - 3, 1)	(a - 2, 1)	(a - 1, 1)
⋮		⋮		⋮				(a - 4, 2)	(a - 3, 2)		
⋮		(2, $\frac{a}{2} - 3$)	(3, $\frac{a}{2} - 3$)	(4, $\frac{a}{2} - 3$)	(5, $\frac{a}{2} - 3$)			(a - 4, 3)	(a - 3, 3)		
(0, $\frac{a}{2} - 2$)	(1, $\frac{a}{2} - 2$)	(2, $\frac{a}{2} - 2$)	(3, $\frac{a}{2} - 2$)								
(0, $\frac{a}{2} - 1$)	(1, $\frac{a}{2} - 1$)	(2, $\frac{a}{2} - 1$)	(3, $\frac{a}{2} - 1$)								
(0, $\frac{a}{2}$)	(1, $\frac{a}{2}$)										
(0, $\frac{a}{2} + 1$)	(1, $\frac{a}{2} + 1$)										
⋮											
⋮		(2, a - 5)	(3, a - 5)	(4, a - 5)	(5, a - 5)						
⋮		(2, a - 4)	(3, a - 4)								
(0, a - 3)	(1, a - 3)	(2, a - 3)	(3, a - 3)								
(0, a - 2)	(1, a - 2)										
(0, a - 1)	(1, a - 1)										

Table 6: Complete residue system $\text{Ap}_{\frac{a}{2}}(A_3)$ for even a

Each element $m_j^{(\frac{a}{2})}$ in $\text{Ap}_{\frac{a}{2}}(A_3)$ has exactly $a/2 + 1$ expressions because

$$(a - 1)(a + 2d) = (d + i)a + (a - 2i)(a + d) + (i - 1)(a + 2d)$$

$$(1 \leq i \leq \frac{a}{2}),$$

$$(a + d) + (a - 1)(a + 2d) = (d + i)a + (a - 2i + 1)(a + d)$$

$$+ (i - 1)(a + 2d) \quad (1 \leq i \leq \frac{a}{2}),$$

$$2(a + d) + (a - 3)(a + 2d) = a + (a - 2)(a + 2d)$$

$$= (d + i + 2)a + (a - 2i - 2)(a + d)$$

$$+ (i - 1)(a + 2d) \quad (1 \leq i \leq \frac{a}{2} - 1),$$

$$3(a + d) + (a - 3)(a + 2d) = a + (a + d) + (a - 2)(a + 2d)$$

$$= (d + i + 2)a + (a - 2i - 1)(a + d)$$

$$+ (i - 1)(a + 2d) \quad (1 \leq i \leq \frac{a}{2} - 1),$$

$$4(a + d) + (a - 5)(a + 2d) = a + 2(a + d) + (a - 4)(a + 2d)$$

$$= 2a + (a - 3)(a + 2d)$$

$$= (d + i + 4)a + (a - 2i - 4)(a + d)$$

$$+ (i - 1)(a + 2d) \quad (1 \leq i \leq \frac{a}{2} - 2),$$

$$5(a + d) + (a - 5)(a + 2d) = a + 3(a + d) + (a - 4)(a + 2d)$$

$$\begin{aligned}
&= 2a + (a + d) + (a - 3)(a + 2d) \\
&= (d + i + 4)a + (a - 2i - 3)(a + d) \\
&\quad + (i - 1)(a + 2d) \quad (1 \leq i \leq \frac{a}{2} - 2), \\
&\quad \dots \\
(a - 2)(a + d) + (a + 2d) &= ia + (a - 2i - 2)(a + d) + (i + 1)(a + 2d) \\
&\quad (1 \leq i \leq \frac{a}{2} - 1) \\
&= (a + d - 1)a, \\
(a - 1)(a + d) + (a + 2d) &= ia + (a - 2i - 1)(a + d) + (i + 1)(a + 2d) \\
&\quad (1 \leq i \leq \frac{a - 1}{2}) \\
&= (a + d - 1)a + (a + d).
\end{aligned}$$

But any element $m_j^{(\frac{a}{2})} - a$ has $a/2$ expressions.

Since the maximal value in $\text{Ap}_p(A_3)$ is at the position $(1, a/2 - 1 + p)$, when a is even, for $0 \leq p \leq a/2$,

$$\begin{aligned}
g_p(a, a + d, a + 2d) &= (a + d) + \left(\frac{a}{2} - 1 + p\right)(a + 2d) - a \\
&= (a + 2d)p + \frac{a(a - 2) + 2(a - 1)d}{2}.
\end{aligned}$$

However, when $p = a/2 + 1$, no element comes to the position $(0, a)$ or $(1, a)$ because there is no element of $\text{Ap}_{\frac{a}{2}}(A_3)$ in the top row. Hence, after $p = a/2 + 1$, the pattern is shifted and the situation becomes irregular and complicated.

Concerning the number of the representations, the elements in the staircase are

$$\begin{aligned}
&\left(0, \frac{a}{2} + p - 1\right), \left(1, \frac{a}{2} + p - 1\right), \\
&\left(2, \frac{a}{2} + p - 3\right), \left(3, \frac{a}{2} + p - 3\right), \\
&\quad \dots \\
&\left(2p - 4, \frac{a}{2} - p + 3\right), \left(2p - 3, \frac{a}{2} - p + 3\right), \\
&\left(2p - 2, \frac{a}{2} - p + 1\right), \left(2p - 1, \frac{a}{2} - p + 1\right),
\end{aligned}$$

and the elements in the large block are

$$(2p, 0), (2p, 1), \dots, \left(2p, \frac{a}{2} - p - 1\right),$$

$$(2p+1, 0), (2p+1, 1), \dots, \left(2p+1, \frac{a}{2} - p - 1\right).$$

Hence,

$$\begin{aligned}
& \sum_{j=0}^{a-1} m_j^{(p)} \\
&= (0+2+\dots+(2p-2))(a+d) + (1+3+\dots+(2p-1))(a+d) \\
&\quad + 2\left(\left(\frac{a}{2}-p+1\right) + \left(\frac{a}{2}-p+3\right) + \dots + \left(\frac{a}{2}+p-1\right)\right)(a+2d) \\
&\quad + \left(\frac{a}{2}-p\right)(2p)(a+d) + \left(\frac{a}{2}-p\right)(2p+1)(a+d) \\
&\quad + 2\left(0+1+\dots+\left(\frac{a}{2}-p-1\right)\right)(a+2d) \\
&= (2p-1)p(a+d) + ap(a+2d) \\
&\quad + \left(\frac{a}{2}-p\right)(4p+1)(a+d) + \left(\frac{a}{2}-p-1\right)\left(\frac{a}{2}-p\right)(a+2d) \\
&= \frac{a}{4}\left((a+d)^2 - (d+1)^2 + 1 - 4p^2 + 4(2a+2d-1)p\right). \tag{7}
\end{aligned}$$

Hence, by Lemma 1 (4)

$$\begin{aligned}
& n_p(a, a+d, a+2d) \\
&= \frac{1}{4}\left((a+d)^2 - (d+1)^2 + 1 - 4p^2 + 4(2a+2d-1)p\right) - \frac{a-1}{2} \\
&= \frac{(a-1)(a+2d-1) + 1}{4} - p^2 + (2a+2d-1)p.
\end{aligned}$$

In [21, (3,9)],

$$g(a, a+d, \dots, a+(k-1)d) = \left\lfloor \frac{a-2}{k-1} \right\rfloor a + (a-1)d,$$

In [21, (3,10)],

$$n(a, a+d, \dots, a+(k-1)d) = \frac{(a-1)(q+d) + r(q+1)}{2},$$

where integers q and r are determined as

$$a-1 = q(k-1) + r, \quad 0 \leq r < k-1.$$

3.1 Generalized Sylvester sum

Concerning the sum of the representations, we need the summation of square of the elements in $\text{Ap}_p(A_3)$. When a is odd, we have

$$\begin{aligned}
& \sum_{j=0}^{(p)} (m_j^{(p)})^2 \\
&= (1^2 + 2^2 + \cdots + (2p-1)^2)(a+d)^2 \\
&\quad + 2 \left(1 \cdot \left(\frac{a-1}{2} + p - 1 \right) + 2 \left(\frac{a-1}{2} + p - 2 \right) \right. \\
&\quad \quad \left. + \cdots + (2p-1) \left(\frac{a-1}{2} - p + 1 \right) \right) (a+d)(a+2d) \\
&\quad + \left(\left(\frac{a-1}{2} + p \right)^2 + \left(\frac{a-1}{2} + p - 1 \right)^2 \right. \\
&\quad \quad \left. + \left(\frac{a-1}{2} + p - 2 \right)^2 + \cdots + \left(\frac{a-1}{2} - p + 1 \right)^2 \right) (a+2d)^2 \\
&\quad + (2p)^2 \left(\frac{a+1}{2} - p \right) (a+d)^2 \\
&\quad + 2 \cdot 2p \left(1 + 2 + \cdots + \left(\frac{a-1}{2} - p \right) \right) (a+d)(a+2d) \\
&\quad + \left(1^2 + 2^2 + \cdots + \left(\frac{a-1}{2} - p \right)^2 \right) (a+2d)^2 \\
&\quad + (2p+1)^2 \left(\frac{a-1}{2} - p \right) (a+d)^2 \\
&\quad + 2(2p+1) \left(1 + 2 + \cdots + \left(\frac{a-1}{2} - p - 1 \right) \right) (a+d)(a+2d) \\
&\quad + \left(1^2 + 2^2 + \cdots + \left(\frac{a-1}{2} - p - 1 \right)^2 \right) (a+2d)^2 \\
&= \frac{p(2p-1)(4p-1)}{3} (a+d)^2 + \frac{p(2p-1)(3a-2p-1)}{3} (a+d)(a+2d) \\
&\quad + \frac{4p^3 + 3a^2p - p}{6} (a+2d)^2 + \frac{(4p^2)(a-2p+1)}{2} (a+d)^2 \\
&\quad + \frac{p(a-2p-1)(a-2p+1)}{2} (a+d)(a+2d) \\
&\quad + \frac{(a-2p-1)(a-2p)(a-2p+1)}{24} (a+2d)^2
\end{aligned}$$

$$\begin{aligned}
& + \frac{(4p^2 + 4p + 1)(a - 2p - 1)}{2}(a + d)^2 \\
& + \frac{(2p + 1)(a - 2p - 3)(a - 2p - 1)}{4}(a + d)(a + 2d) \\
& + \frac{(a - 2p - 3)(a - 2p - 2)(a - 2p - 1)}{24}(a + 2d)^2 \\
= & \frac{a}{12}((a - 1)(a^3 + a^2(4d + 1) + ad(4d + 1) - d(2d + 3)) \\
& + 4(3a^3 + 3a^2(3d - 1) + a(6d^2 - 12d - 1) - 2(3d - 1)d)p \\
& + 12(3a^2 + 2(3d - 1)a + d(4d - 1))p^2 - 32(a + d)p^3).
\end{aligned}$$

Hence, by Lemma 1 (5) and the relation (6), we have we have the formula for the generalized Sylvester sum.

When a is even, we have

$$\begin{aligned}
& \sum_{j=0}^{(p)} (m_j^{(p)})^2 \\
= & (1^2 + 2^2 + \cdots + (2p - 1)^2)(a + d)^2 \\
& + 2 \left(1 \cdot \left(\frac{a}{2} + p - 1 \right) + (2 + 3) \left(\frac{a}{2} + p - 3 \right) \right. \\
& \quad \left. + \cdots + ((2p - 2) + (2p - 1)) \left(\frac{a}{2} - p + 1 \right) \right) (a + d)(a + 2d) \\
& + 2 \left(\left(\frac{a}{2} + p - 1 \right)^2 + \left(\frac{a - 1}{2} + p - 3 \right)^2 \right. \\
& \quad \left. + \cdots + \left(\frac{a}{2} - p + 1 \right)^2 \right) (a + 2d)^2 \\
& + ((2p)^2 + (2p + 1)^2) \left(\frac{a}{2} - p \right) (a + d)^2 \\
& + 2((2p) + (2p + 1)) \left(1 + 2 + \cdots + \left(\frac{a}{2} - p - 1 \right) \right) (a + d)(a + 2d) \\
& + 2 \left(1^2 + 2^2 + \cdots + \left(\frac{a}{2} - p - 1 \right)^2 \right) (a + 2d)^2 \\
= & \frac{p(2p - 1)(4p - 1)}{3}(a + d)^2 + \frac{p(4 - 3a + 6ap - 4p^2)}{3}(a + d)(a + 2d) \\
& + \frac{p(3a^2 + 4p^2 - 4)}{6}(a + 2d)^2 + \frac{(8p^2 + 4p + 1)(a - 2p)}{2}(a + d)^2 \\
& + \frac{(4p + 1)(a - 2p - 2)(a - 2p)}{4}(a + d)(a + 2d)
\end{aligned}$$

$$\begin{aligned}
& + \frac{(a-2p-2)(a-2p-1)(a-2p)}{12}(a+2d)^2 \\
= & \frac{a}{12}(a^4 + 4a^3d + a^2(4d^2 - 3d + 2) - 2ad(3d-1) + 2d^2 \\
& + 4(3a^3 + 3a^2(3d-1) + 2a(3d^2 - 6d + 1) - 6d^2 + 5d)p \\
& + 12(3a^2 + 2a(3d-1) + 4d^2 - d)p^2 - 32(a+d)p^3).
\end{aligned}$$

Hence, by Lemma 1 (5) and the relation (7), we have the formula for the generalized Sylvester sum.

Theorem 2. *Let a and d be integers with $a \geq 3$, $d > 0$ and $\gcd(a, d) = 1$. Then for $0 \leq p \leq \lfloor a/2 \rfloor$, when a is odd, we have*

$$\begin{aligned}
s_p(A_3) &= \frac{1}{24}((a-1)(a^3 + a^2(4d+1) + ad(4d+1) - d(2d+3)) \\
& + 4(3a^3 + 3a^2(3d-1) + a(6d^2 - 12d - 1) - 2(3d-1)d)p \\
& + 12(3a^2 + 2(3d-1)a + d(4d-1))p^2 - 32(a+d)p^3) \\
& - \frac{a}{8}((a+d)^2 - (d+1)^2 - 4p^2 + 4(2a-2d-1)p) \\
& + \frac{a^2-1}{12} \\
= & \frac{(a-1)(a+2d-1)(a^2 + 2ad - a - d - 2)}{24} \\
& + \frac{3a^3 + 9a^2(d-1) + 2a(3d^2 - 9d + 1) - 6d^2 + 2d}{6}p \\
& + \frac{3a^2 + a(6d-1) + 4d^2 - d}{2}p^2 - \frac{3(a+d)}{2}p^3.
\end{aligned}$$

When a is even, we have

$$\begin{aligned}
s_p(A_3) &= \frac{1}{24}(a^4 + 4a^3d + a^2(4d^2 - 3d + 2) - 2ad(3d-1) + 2d^2 \\
& + 4(3a^3 + 3a^2(3d-1) + 2a(3d^2 - 6d + 1) - 6d^2 + 5d)p \\
& + 12(3a^2 + 2a(3d-1) + 4d^2 - d)p^2 - 32(a+d)p^3) \\
& - \frac{a}{8}((a+d)^2 - (d+1)^2 + 1 - 4p^2 + 4(2a-2d-1)p) \\
& + \frac{a^2-1}{12} \\
= & \frac{(a-1)(a+2d-1)(a^2 + 2ad - a - d - 2) + 3(a^2 + 2ad - d)}{24} \\
& + \frac{3a^3 + 9a^2(d-1) + a(6d^2 - 18d + 5) - 6d^2 + 5d}{6}p
\end{aligned}$$

$$+ \frac{3a^2 + a(6d - 1) + 4d^2 - d}{2} p^2 - \frac{3(a + d)}{2} p^3.$$

4 Power sums

More generally, we can consider the generalized Sylvester power sum

$$s_p^{(\mu)}(a_1, a_2, \dots, a_k) = \sum_{d(n) \leq p} n^\mu \quad (\mu \geq 1),$$

so that $s_p(a_1, a_2, \dots, a_k) = s_p^{(1)}(a_1, a_2, \dots, a_k)$ and $n_p(a_1, a_2, \dots, a_k) = s_p^{(0)}(a_1, a_2, \dots, a_k)$.

Then the generalized Sylvester power sum is given as follows [16].

Lemma 2. *Let k, p and μ be integers with $k \geq 2, p \geq 0$ and $\mu \geq 1$. Assume that $\gcd(a_1, a_2, \dots, a_k) = 1$. We have*

$$\begin{aligned} s_p^{(\mu)}(a_1, a_2, \dots, a_k; p) &:= \sum_{d(n) \leq p} n^\mu \\ &= \frac{1}{\mu + 1} \sum_{\kappa=0}^{\mu} \binom{\mu + 1}{\kappa} B_\kappa a_1^{\kappa-1} \sum_{i=0}^{a_1-1} (m_i^{(p)})^{\mu+1-\kappa} + \frac{B_{\mu+1}}{\mu + 1} (a_1^{\mu+1} - 1), \end{aligned}$$

where B_κ are Bernoulli numbers, defined by the generating function

$$\frac{x}{e^x - 1} = \sum_{\kappa=0}^{\infty} B_\kappa \frac{x^\kappa}{\kappa!}.$$

In order to obtain the formula for the generalized Sylvester power sum, we need to calculate $(m_i^{(p)})^\nu$ for $\nu \geq 0$. From the previous section, when n is odd,

$$\begin{aligned} (m_i^{(p)})^\nu &= \sum_{j=0}^{\nu} \binom{\nu}{j} \sum_{k=0}^{2p-1} k^{\nu-j} \left(\frac{a-1}{2} + p - k \right)^j (a+d)^{\nu-j} (a+2d)^j \\ &\quad + \sum_{j=0}^{\nu} \binom{\nu}{j} (2p)^{\nu-j} \sum_{k=0}^{\frac{a-1}{2}-p} k^j (a+d)^{\nu-j} (a+2d)^j \end{aligned}$$

$$\begin{aligned}
& + \sum_{j=0}^{\nu} \binom{\nu}{j} (2p+1)^{\nu-j} \sum_{k=0}^{\frac{a-1}{2}-p-1} k^j (a+d)^{\nu-j} (a+2d)^j \\
& = \sum_{j=0}^{\nu} \binom{\nu}{j} \left(\sum_{k=0}^{2p-1} k^{\nu-j} \left(\frac{a-1}{2} + p - k \right)^j \right. \\
& \quad \left. + (2p)^{\nu-j} \sum_{k=0}^{\frac{a-1}{2}-p} k^j + (2p+1)^{\nu-j} \sum_{k=0}^{\frac{a-1}{2}-p-1} k^j \right) (a+d)^{\nu-j} (a+2d)^j.
\end{aligned}$$

When n is even

$$\begin{aligned}
(m_i^{(p)})^{\nu} & = \sum_{j=0}^{\nu} \binom{\nu}{j} \sum_{k=1}^p ((2k-2)^{\nu-j} + (2k-1)^{\nu-j}) \left(\frac{a}{2} + p - 2k + 1 \right)^j \\
& \quad \times (a+d)^{\nu-j} (a+2d)^j \\
& + \sum_{j=0}^{\nu} \binom{\nu}{j} (2p)^{\nu-j} \sum_{k=0}^{\frac{a}{2}-p-1} k^j (a+d)^{\nu-j} (a+2d)^j \\
& + \sum_{j=0}^{\nu} \binom{\nu}{j} (2p+1)^{\nu-j} \sum_{k=0}^{\frac{a}{2}-p-1} k^j (a+d)^{\nu-j} (a+2d)^j \\
& = \sum_{j=0}^{\nu} \binom{\nu}{j} \left(\sum_{k=1}^p ((2k-2)^{\nu-j} + (2k-1)^{\nu-j}) \left(\frac{a}{2} + p - 2k + 1 \right)^j \right. \\
& \quad \left. + ((2p)^{\nu-j} + (2p+1)^{\nu-j}) \sum_{k=0}^{\frac{a}{2}-p-1} k^j \right) (a+d)^{\nu-j} (a+2d)^j.
\end{aligned}$$

Hence, by Lemma 2, we obtain the generalized Sylvester power sum for $(a, a+d, a+2d)$.

Theorem 3. *Let a, d, p and μ be integers with $a \geq 3, d > 0, p \geq 0, \mu \geq 1$ and $\gcd(a, d) = 1$. Then, when a is odd, we have*

$$\begin{aligned}
& s_p^{(\mu)}(a, a+d, a+2d) \\
& = \frac{1}{\mu+1} \sum_{\kappa=0}^{\mu} \binom{\mu+1}{\kappa} B_{\kappa} a^{\kappa-1} \sum_{i=0}^{a-1} \sum_{j=0}^{\mu+1-\kappa} \binom{\mu+1-\kappa}{j} \\
& \quad \times \left(\sum_{k=0}^{2p-1} k^{\mu+1-\kappa-j} \left(\frac{a-1}{2} + p - k \right)^j \right)
\end{aligned}$$

$$\begin{aligned}
& + (2p)^{\mu+1-\kappa-j} \sum_{k=0}^{\frac{a-1}{2}-p} k^j + (2p+1)^{\mu+1-\kappa-j} \sum_{k=0}^{\frac{a-1}{2}-p-1} k^j \Big) \\
& \times (a+d)^{\mu+1-\kappa-j} (a+2d)^j \\
& + \frac{B_{\mu+1}}{\mu+1} (a^{\mu+1} - 1).
\end{aligned}$$

When a is even, we have

$$\begin{aligned}
& s_p^{(\mu)}(a, a+d, a+2d) \\
& = \frac{1}{\mu+1} \sum_{\kappa=0}^{\mu} \binom{\mu+1}{\kappa} B_{\kappa} a^{\kappa-1} \sum_{i=0}^{a-1} \sum_{j=0}^{\mu+1-\kappa} \binom{\mu+1-\kappa}{j} \\
& \quad \times \left(\sum_{k=1}^p ((2k-2)^{\mu+1-\kappa-j} + (2k-1)^{\mu+1-\kappa-j}) \left(\frac{a}{2} + p - 2k + 1 \right)^j \right. \\
& \quad \left. + ((2p)^{\mu+1-\kappa-j} + (2p+1)^{\mu+1-\kappa-j}) \sum_{k=0}^{\frac{a}{2}-p-1} k^j \right) \\
& \quad \times (a+d)^{\mu+1-\kappa-j} (a+2d)^j \\
& + \frac{B_{\mu+1}}{\mu+1} (a^{\mu+1} - 1).
\end{aligned}$$

5 Weighted sums

In this section, we consider the weighted sums whose numbers of representations are less than or equal to p :

$$s_{\lambda,p}^{(\mu)}(a_1, \dots, a_k) := \sum_{d(p) \leq p} \lambda^n n^{\mu},$$

where $\lambda \neq 1$ and μ is a positive integer.

Here, Eulerian numbers $\langle \frac{n}{m} \rangle$ appear in the generating function

$$\sum_{k=0}^{\infty} k^n x^k = \frac{1}{(1-x)^{n+1}} \sum_{m=0}^{n-1} \langle \frac{n}{m} \rangle x^{m+1} \quad (n \geq 1) \quad (8)$$

with $0^0 = 1$ and $\langle \frac{0}{0} \rangle = 1$, and have an explicit formula

$$\langle \frac{n}{m} \rangle = \sum_{k=0}^m (-1)^k \binom{n+1}{k} (m-k+1)^n.$$

Lemma 3. Assume that $\lambda \neq 1$ and $\lambda^{a_1} \neq 1$. Then for a positive integer μ ,

$$\begin{aligned} & s_{\lambda,p}^{(\mu)}(a_1, \dots, a_k) \\ &= \sum_{n=0}^{\mu} \frac{(-a_1)^n}{(\lambda^{a_1} - 1)^{n+1}} \binom{\mu}{n} \sum_{j=0}^n \left\langle \begin{matrix} n \\ n-j \end{matrix} \right\rangle \lambda^{ja_1} \sum_{i=0}^{a_1-1} (m_i^{(p)})^{\mu-n} \lambda^{m_i^{(p)}} \\ & \quad + \frac{(-1)^{\mu+1}}{(\lambda - 1)^{\mu+1}} \sum_{j=0}^{\mu} \left\langle \begin{matrix} \mu \\ \mu-j \end{matrix} \right\rangle \lambda^j. \end{aligned}$$

In order to obtain the formula for the generalized Sylvester weighted power sum, we need to calculate $(m_i^{(p)})^{\nu} \lambda^{m_i^{(p)}}$ for $\nu \geq 0$, $\lambda^a \neq 1$ and $\lambda^d \neq 1$. We use the formula

$$a^{\nu} x^a = \sum_{i=0}^{\nu} \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} x^i \frac{d^i}{dx^i} x^a,$$

where $\left\{ \begin{matrix} n \\ m \end{matrix} \right\}$ denote the Stirling numbers of the second kind, calculated as

$$\left\{ \begin{matrix} n \\ m \end{matrix} \right\} = \frac{1}{m!} \sum_{i=0}^m (-1)^i \binom{m}{i} (m-i)^n.$$

From the previous section, when n is odd,

$$\begin{aligned} & (m_i^{(p)})^{\nu} \lambda^{m_i^{(p)}} \\ &= \sum_{i=0}^{\nu} \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^i \left[\frac{d^i}{dx^i} \left(\sum_{l=0}^{2p-1} x^{l(a+d) + (\frac{a-1}{2} + p - l)(a+2d)} \right. \right. \\ & \quad \left. \left. + \sum_{l=0}^{\frac{a-1}{2}-p} x^{2p(a+d) + l(a+2d)} + \sum_{l=0}^{\frac{a-1}{2}-p-1} x^{(2p+1)(a+d) + l(a+2d)} \right) \right]_{x=\lambda} \\ &= \sum_{i=0}^{\nu} \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^i \left[\frac{d^i}{dx^i} \left(\frac{x^{a(\frac{a-1}{2} + d + p)} (x^{2dp} - 1)}{x^d - 1} \right. \right. \\ & \quad \left. \left. + \frac{x^{2p(a+d)} (x^{(a+2d)(\frac{a+1}{2}-p)} - 1)}{x^{a+2d} - 1} + \frac{x^{(2p+1)(a+d)} (x^{(a+2d)(\frac{a-1}{2}-p)} - 1)}{x^{a+2d} - 1} \right) \right]_{x=\lambda}. \end{aligned}$$

When n is even,

$$(m_i^{(p)})^{\nu} \lambda^{m_i^{(p)}}$$

$$\begin{aligned}
&= \sum_{i=0}^{\nu} \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^i \left[\frac{d^i}{dx^i} \left(\sum_{l=0}^{p-1} x^{2l(a+d) + (\frac{a}{2} + p - 2l - 1)(a+2d)} \right. \right. \\
&\quad + \sum_{l=0}^{p-1} x^{(2l+1)(a+d) + (\frac{a}{2} + p - 2l - 1)(a+2d)} \\
&\quad \left. \left. + \sum_{l=0}^{\frac{a}{2} - p - 1} x^{2p(a+d) + l(a+2d)} + \sum_{l=0}^{\frac{a}{2} - p - 1} x^{(2p+1)(a+d) + l(a+2d)} \right) \right]_{x=\lambda} \\
&= \sum_{i=0}^{\nu} \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^i \left[\frac{d^i}{dx^i} \left(\frac{x^{\frac{a}{2}(a+2d-2+2p)}(x^{a+d} + 1)(x^{2dp} - 1)}{x^{2d} - 1} \right. \right. \\
&\quad \left. \left. + \frac{x^{2(a+d)p}(x^{a+d} + 1)(x^{(a+2d)(\frac{a}{2}-p)} - 1)}{x^{a+2d} - 1} \right) \right]_{x=\lambda}.
\end{aligned}$$

Hence, by Lemma 3, we obtain the generalized Sylvester weighted power sum for $(a, a + d, a + 2d)$.

Theorem 4. *Let a, d, p, λ and μ be integers with $a \geq 3, d > 0, p \geq 0, \lambda \neq 1, \lambda^d \neq 1, \mu \geq 1$ and $\gcd(a, d) = 1$. Then, when a is odd, we have*

$$\begin{aligned}
&s_{\lambda, p}^{(\mu)}(a, a + d, a + 2d) \\
&= \sum_{n=0}^{\mu} \sum_{j=0}^n \sum_{i=0}^{\nu} \frac{(-a)^n}{(\lambda^a - 1)^{n+1}} \binom{\mu}{n} \left\langle \begin{matrix} n \\ n - j \end{matrix} \right\rangle \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^{ja+i} \\
&\quad \times \left[\frac{d^i}{dx^i} \left(\frac{x^{a(\frac{a-1}{2} + d + p)}(x^{2dp} - 1)}{x^d - 1} \right. \right. \\
&\quad \left. \left. + \frac{x^{2p(a+d)}(x^{(a+2d)(\frac{a+1}{2} - p)} - 1)}{x^{a+2d} - 1} + \frac{x^{(2p+1)(a+d)}(x^{(a+2d)(\frac{a-1}{2} - p)} - 1)}{x^{a+2d} - 1} \right) \right]_{x=\lambda} \\
&\quad + \frac{(-1)^{\mu+1}}{(\lambda - 1)^{\mu+1}} \sum_{j=0}^{\mu} \left\langle \begin{matrix} \mu \\ \mu - j \end{matrix} \right\rangle \lambda^j.
\end{aligned}$$

When a is even, we have

$$\begin{aligned}
&s_{\lambda, p}^{(\mu)}(a, a + d, a + 2d) \\
&= \sum_{n=0}^{\mu} \sum_{j=0}^n \sum_{i=0}^{\nu} \frac{(-a)^n}{(\lambda^a - 1)^{n+1}} \binom{\mu}{n} \left\langle \begin{matrix} n \\ n - j \end{matrix} \right\rangle \left\{ \begin{matrix} \nu \\ i \end{matrix} \right\} \lambda^{ja+i}
\end{aligned}$$

$$\begin{aligned} & \times \left[\frac{d^i}{dx^i} \left(\frac{x^{\frac{a}{2}(a+2d-2+2p)}(x^{a+d}+1)(x^{2dp}-1)}{x^{2d}-1} \right. \right. \\ & \quad \left. \left. + \frac{x^{2(a+d)p}(x^{a+d}+1)(x^{(a+2d)(\frac{a}{2}-p)}-1)}{x^{a+2d}-1} \right) \right]_{x=\lambda} \\ & + \frac{(-1)^{\mu+1}}{(\lambda-1)^{\mu+1}} \sum_{j=0}^{\mu} \left\langle \begin{matrix} \mu \\ \mu-j \end{matrix} \right\rangle \lambda^j. \end{aligned}$$

Remark. The case $\lambda = 1$ is not included in Theorem 4, but in Theorem 3.

6 Examples

For (11, 15, 19), that is, $a = 11$ and $d = 4$, when $q = 5$, by Theorems 1, 2, 3 and 4, we have

$$\begin{aligned} g_5(11, 15, 19) &= 179, \\ n_5(11, 15, 19) &= 165, \\ s_5(11, 15, 19) &= 13605, \\ s_5^{(3)}(11, 15, 19) &= 189158535 \quad (\mu = 3), \\ s_{2,5}^{(3)}(11, 15, 19) &= 46691295420476497563538523364517263 \\ & \quad 55433630648909109181546522 \quad (\lambda = 2). \end{aligned}$$

For (6, 11, 16), that is, $a = 6$ and $d = 5$, when $q = 3$, we have

$$\begin{aligned} g_3(6, 11, 16) &= 85, \\ n_3(6, 11, 16) &= 73, \\ s_3(6, 11, 16) &= 2675, \\ s_3^{(3)}(6, 11, 16) &= 7652009 \quad (\mu = 3), \\ s_{2,3}^{(3)}(6, 11, 16) &= 24083450837052351738334815453210 \quad (\lambda = 2). \end{aligned}$$

In fact, the integers whose representations in terms of (6, 11, 16) are less than or equal to 3 are

$$0, 1, 2, \dots, \underbrace{59}_{59}, 61, 62, 63, 64, 65, 67, 68, 69, 73, 74, 75, 79, 85.$$

Hence, for example,

$$\begin{aligned}
s_{2,3}^{(3)}(6, 11, 16) &= 2^0 \cdot 0^3 + 2^1 \cdot 1^3 + 2^2 \cdot 2^3 + \dots + 2^{59} \cdot 59^3 + 2^{61} \cdot 61^3 \\
&\quad + 2^{62} \cdot 62^3 + 2^{63} \cdot 63^3 + 2^{64} \cdot 64^3 + 2^{65} \cdot 65^3 \\
&\quad + 2^{67} \cdot 67^3 + 2^{68} \cdot 68^3 + 2^{69} \cdot 69^3 + 2^{73} \cdot 73^3 \\
&\quad + 2^{74} \cdot 74^3 + 2^{75} \cdot 75^3 + 2^{79} \cdot 79^3 + 2^{85} \cdot 85^3 \\
&= 24083450837052351738334815453210.
\end{aligned}$$

7 Final comments

It is hoped to have explicit formulas for $g_p(a, a+d, a+2d, a+3d)$, and more generally, $g_p(a, a+d, \dots, a+(k-1)d)$ because explicit formulas for general k has been known when $p=0$. Indeed, in [15], for $p \geq 0$ explicit formulas about four and five repunits are obtained, though the structures are even more complicated than that of three variables. In the case of three variables, for any j ($j=0, 1, \dots, a-1$), $m_j^{(0)} < m_j^{(1)} < \dots$. However, in the case of four and more variables, for some j 's, $m_j^{(p)} = m_j^{(p+1)}$. It means that some elements in $\text{Ap}_p(A)$ and in $\text{Ap}_{p+1}(A)$ are overlapped.

Another problem is whether we can find any convenient formula when $p > \lfloor a/2 \rfloor$.

Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- [1] R. Apéry, *Sur les branches superlinéaires des courbes algébriques*, C. R. Acad. Sci. Paris **222** (1946), 1198–1200.
- [2] L. Bardomero and M. Beck, *Frobenius coin-exchange generating functions*, Amer. Math. Monthly **127** (2020), no. 4, 308–315.
- [3] M. Beck, I. M. Gessel and T. Komatsu, *The polynomial part of a restricted partition function related to the Frobenius problem*, Electron. J. Combin. **8** (No.1) (2001), #N7.

- [4] M. Beck and C. Kifer, *An extreme family of generalized Frobenius numbers*, *Integers* **11** (2011), A24, 639–645.
- [5] D. S. Binner, *The number of solutions to $ax + by + cz = n$ and its relation to quadratic residues*, *J. Integer Seq.* **23**, No. 6, (2020), Article 20.6.5, 19 p.
- [6] A. Brauer, *On a problem of partitions*, *Amer. J. Math.* **64** (1942), 299–312.
- [7] A. Brauer and B. M. Shockley, *On a problem of Frobenius*, *J. Reine. Angew. Math.* **211** (1962), 215–220.
- [8] A. Brown, E. Dannenberg, J. Fox, J. Hanna, K. Keck, A. Moore, Z. Robbins, B. Samples and J. Stankewicz, *On a generalization of the Frobenius number*, arXiv:1001.0207 (2010).
- [9] T. C. Brown and P. J. Shiue, *A remark related to the Frobenius problem*, *Fibonacci Quart.* **31** (1993), 32–36.
- [10] A. Cayley, *On a problem of double partitions*, *Philos. Mag.* **XX** (1860), 337–341.
- [11] L. Fukshansky and A. Schurmann, *Bounds on generalized Frobenius numbers*, *Eur. J. Comb.* **32** (2011), No. 3, 361–368.
- [12] T. Komatsu, *On the number of solutions of the Diophantine equation of Frobenius—General case*, *Math. Commun.* **8** (2003), 195–206.
- [13] T. Komatsu, *Sylvester sums on the Frobenius set in arithmetic progression*, In: F. Yilmaz et al. (eds.), *Mathematical Methods for Engineering Applications*, Springer Proceedings in Mathematics & Statistics, vol. 384. Springer, Cham., 2022. pp.1–23.
- [14] T. Komatsu, *The Frobenius number for sequences of triangular numbers associated with number of solutions*, *Ann. Comb.* (to appear).
- [15] T. Komatsu, *The Frobenius number associated with the number of representations for sequences of repunits*, *C. R., Math., Acad. Sci. Paris* (to appear).
- [16] T. Komatsu, *Frobenius numbers, Sylvester numbers and sums associated with number of solutions*, arXiv:2111.11021 (2021).

- [17] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set*, Irish Math. Soc. Bull. **87** (2021), 21–29.
- [18] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set in more variables*, Kyushu J. Math. **76** (2022), 163–175.
- [19] J. B. Roberts, *Notes on linear forms*, Proc. Amer. Math. Soc. **7** (1956), 465–469.
- [20] A. M. Robles-Pérez and J. C. Rosales, *The Frobenius number for sequences of triangular and tetrahedral numbers*, J. Number Theory **186** (2018), 473–492.
- [21] E. S. Selmer, *On the linear diophantine problem of Frobenius*, J. Reine Angew. Math. **293/294** (1977), 1–17.
- [22] J. J. Sylvester, *On the partition of numbers*, Quart. J. Pure Appl. Math. **1** (1857), 141–152.
- [23] J. J. Sylvester, *On subinvariants, i.e. semi-invariants to binary quantics of an unlimited order*, Am. J. Math. **5** (1882), 119–136.
- [24] J. J. Sylvester, *Mathematical questions with their solutions*, Educational Times **41** (1884), 21.
- [25] A. Tripathi, *The number of solutions to $ax + by = n$* , Fibonacci Quart. **38** (2000), 290–293.
- [26] A. Tripathi, *On sums of positive integers that are not of the form $ax+by$* , Amer. Math. Monthly **115** (2008), 363–364.
- [27] A. Tripathi, *Formulae for the Frobenius number in three variables*, J. Number Theory **170** (2017), 368–389.