

# The Frobenius number for sequences of triangular numbers associated with number of solutions

Takao Komatsu

Department of Mathematical Sciences, School of Science  
Zhejiang Sci-Tech University  
Hangzhou 310018 China  
komatsu@zstu.edu.cn

MR Subject Classifications: Primary 11D07; Secondary 05A15, 05A17, 05A19, 11B68, 11D04, 11P81

## Abstract

The famous linear diophantine problem of Frobenius is the problem to determine the largest integer (Frobenius number) whose number of representations in terms of  $a_1, \dots, a_k$  is at most zero, that is not representable. In other words, all the integers greater than this number can be represented for at least one way. One of the natural generalizations of this problem is to find the largest integer (generalized Frobenius number) whose number of representations is at most a given nonnegative integer  $p$ . It is easy to find the explicit form of this number in the case of two variables. However, no explicit form has been known even in any special case of three variables. In this paper we are successful to show explicit forms of the generalized Frobenius numbers of the triples of triangular numbers. When  $p = 0$ , their Frobenius number is give by Robles-Pérez and Rosales in 2018.

**Keywords:** Frobenius problem, Frobenius numbers, number of representations, triangular numbers

## 1 Introduction

For integer  $k \geq 2$ , consider a set of positive integers  $A = \{a_1, \dots, a_k\}$  with  $\gcd(A) = \gcd\{a_1, \dots, a_k\} = 1$ . Let  $\text{NR}(A) = \text{NR}(a_1, a_2, \dots, a_k)$  denote the set of positive integers nonrepresentable as nonnegative integer combinations of  $a_1, a_2, \dots, a_k$ . Then, the well-known linear Diophantine problem, posed by Sylvester [19], but known as the Frobenius problem, is the problem

to determine the Frobenius number  $g(A)$  and the Sylvester number  $n(A)$ , defined by

$$g(A) = \max \text{NR}(A) \quad \text{and} \quad n(A) = |\text{NR}(A)|,$$

respectively. The Frobenius Problem has been also known as the *Coin Exchange Problem* (or Postage Stamp Problem / Chicken McNugget Problem), which has a long history and is one of the problems that has attracted many people as well as experts. For two variables  $A = \{a, b\}$ , it is shown that

$$g(a, b) = (a - 1)(b - 1) - 1 \quad \text{and} \quad n(a, b) = \frac{(a - 1)(b - 1)}{2} \quad (1)$$

([18, 19]). However, for three or more variables,  $g(A)$  cannot be given by any set of closed formulas which can be reduced to a finite set of certain polynomials ([8]). Only some special cases, explicit closed formulas have been found, including arithmetic, geometric-like, Fibonacci, Mersenne, and triangular (see [14] and references therein). The related problem of determining the function

$$s(A) = \sum_{n \in \text{NR}(A)} n,$$

which is called Sylvester sum nowadays, is introduced by Brown and Shiue [6] and for two variables  $A = \{a, b\}$ , it is shown that

$$s(a, b) = \frac{1}{12}(a - 1)(b - 1)(2ab - a - b - 1). \quad (2)$$

On the other hand, to find the number of representations  $d(n; A) = d(n; a_1, a_2, \dots, a_k)$  to  $a_1x_1 + a_2x_2 + \dots + a_kx_k = n$  for a given positive integer  $n$  is also one of the most important and interesting topics. This number is equal to the coefficient of  $x^n$  in  $1/(1 - x^{a_1})(1 - x^{a_2}) \dots (1 - x^{a_k})$  ([18]). Sylvester [17] and Cayley [7] showed that  $d(n; a_1, a_2, \dots, a_k)$  can be expressed as the sum of a polynomial in  $n$  of degree  $k - 1$  and a periodic function of period  $a_1a_2 \dots a_k$ . In [3], the explicit formula for the polynomial part is derived by using Bernoulli numbers.

In this paper, we are interested in one of the general types of Frobenius numbers, which focuses on the number of solutions. For a given nonnegative integer  $p$ , consider the largest integer  $g(A; p) = g(a_1, a_2, \dots, a_k; p)$  such that the number of expressions that can be represented by  $a_1, a_2, \dots, a_k$  is at most  $p$ . That is, all integers larger than  $g(A; p)$  have at least  $p + 1$  representations or more. When  $p = 0$ ,  $g(A) = g(A; 0)$  is the original Frobenius number. One can consider a slightly modified number, that is, the largest integer  $g'(A; p)$  such that the number of expressions is exactly  $p$  ([4]). However,

for some cases any positive integer does not have exactly  $p$  representations, in particular, when  $p$  becomes larger. In addition,  $p_1 < p_2$  does not necessarily imply  $g'(A; p_1) < g'(A; p_2)$ . See Remark 4 below. Therefore, it would be better to treat with  $g(A; p)$ . Of course, after knowing  $g(A; p)$  and  $g(A; p - 1)$ , we can also get  $g'(A; p)$ . In the same stream, the generalized Sylvester number  $n(A; p) = n(a_1, a_2, \dots, a_k; p)$  and generalized Sylvester sum  $s(A; p) = s(a_1, a_2, \dots, a_k; p)$  are defined by

$$n(A; p) = \sum_{d(n; A) \leq p} 1 \quad \text{and} \quad s(A; p) = \sum_{d(n; A) \leq p} n,$$

respectively.

For two variables, it is very easy to see that

$$g(a, b; p) = (p + 1)ab - a - b, \quad (3)$$

$$n(a, b; p) = \frac{1}{2}((2p + 1)ab - a - b + 1), \quad (4)$$

$$s(a, b; p) = \frac{1}{2}abp(ab(p + 1) - a - b) + \frac{1}{12}(a - 1)(b - 1)(2ab - a - b - 1). \quad (5)$$

(see [2, 4]). When  $p = 0$ , the classical results of  $g(a, b) = g(a, b; 0)$ ,  $n(a, b) = n(a, b; 0)$  in (1) and  $s(a, b) = s(a, b; 0)$  in (2) are recovered. However, for three variables, any formula even for special triples has not been known. Recently in [10], some closed formulas of  $g(A, p)$  and  $n(A, p)$  are given when  $A$  is a triple of repunits.

In this paper, we give explicit formulas of  $g(t_n, t_{n+1}, t_{n+2}; p)$ , where  $t_n = \binom{n+1}{2}$  are triangular numbers, which count the number of dots composing an equilateral triangle. When  $p = 0$ , the Frobenius number of three consecutive triangular numbers is give in [14, Proposition 6]:

$$g(t_n, t_{n+1}, t_{n+2}) = \left\lfloor \frac{n}{2} \right\rfloor (t_n + t_{n+1} + t_{n+2} - 1) - 1 \quad (6)$$

or

$$g(t_n, t_{n+1}, t_{n+2}) = \begin{cases} \frac{(3n-3)(n+1)(n+2)}{4} - 1 & \text{if } n \text{ is odd;} \\ \frac{(3n)(n+1)(n+2)}{4} - 1 & \text{if } n \text{ is even.} \end{cases} \quad (7)$$

## 2 Preliminaries

In order to prove the results, we need the following lemmas.

**Lemma 1** ([14, Lemma 3]).

$$\gcd(t_{n+1}, t_{n+2}) = \begin{cases} \frac{n+2}{2} & \text{if } n \text{ is even;} \\ n+2 & \text{if } n \text{ is odd.} \end{cases}$$

**Lemma 2** ([14, Lemma 4]).

$$\gcd(t_n, t_{n+1}, t_{n+2}) = 1.$$

Without loss of generality, we assume that  $a_1 < a_2 < \dots < a_k$ . For each  $0 \leq i \leq a_1 - 1$ , we introduce the positive integer  $m_i^{(p)}$  congruent to  $i$  modulo  $a_1$  such that the number of representations of  $m_i^{(p)}$  is bigger than or equal to  $p + 1$  and that of  $m_i^{(p)} - a_1$  is less than or equal to  $p$ . Note that  $m_0^{(0)}$  is defined to be 0. The set

$$\text{Ap}(A; p) = \text{Ap}(a_1, a_2, \dots, a_k; p) = \{m_0^{(p)}, m_1^{(p)}, \dots, m_{a_1-1}^{(p)}\},$$

is called the  $p$ -Apéry set of  $A = \{a_1, a_2, \dots, a_k\}$  for a nonnegative integer  $p$  (see, [1]), which is congruent to the set

$$\{0, 1, \dots, a_1 - 1\} \pmod{a_1}.$$

**Lemma 3** ([11]). For  $k \geq 2$ , let  $A = \{a_1, a_2, \dots, a_k\}$  with  $\gcd(a_1, a_2, \dots, a_k) = 1$ . For integers  $p$  and  $\mu$  with  $p \geq 0$  and  $\mu \geq 1$ , we have

$$\begin{aligned} s_\mu(A; p) &:= \sum_{d(n; A) \leq p} n^\mu \\ &= \frac{1}{\mu + 1} \sum_{\kappa=0}^{\mu} \binom{\mu + 1}{\kappa} B_\kappa a_1^{\kappa-1} \sum_{i=0}^{a_1-1} (m_i^{(p)})^{\mu+1-\kappa} + \frac{B_{\mu+1}}{\mu + 1} (a_1^{\mu+1} - 1), \end{aligned}$$

where  $B_n$  are Bernoulli numbers defined by

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

When  $\mu = 0, 1$  in Lemma 3, together with  $g(A; p)$  we have known formulae for the generalized Frobenius number, the generalized Sylvester number and the generalized Sylvester sum (see [4]).

**Lemma 4.** *Let  $k$  and  $p$  be integers with  $k \geq 2$  and  $p \geq 0$ . Assume that  $\gcd(A) = 1$ . We have*

$$g(A; p) = \max_{0 \leq i \leq a_1 - 1} m_i^{(p)} - a_1, \quad (8)$$

$$n(A; p) = \frac{1}{a_1} \sum_{i=0}^{a_1 - 1} m_i^{(p)} - \frac{a_1 - 1}{2}, \quad (9)$$

$$s(A; p) = \frac{1}{2a_1} \sum_{i=0}^{a_1 - 1} (m_i^{(p)})^2 - \frac{1}{2} \sum_{i=0}^{a_1 - 1} m_i^{(p)} + \frac{a_1^2 - 1}{12}. \quad (10)$$

**Remark 1.** *When  $p = 0$ , the identities (8), (9) and (10) are reduced to those in [5], [15] and [22], respectively.*

### 3 Triangular numbers

When the number of variables  $k \geq 3$ , it is not easy to find an explicit form of Frobenius number, Sylvester number or Sylvester sum. Nevertheless, for  $p = 0$ , explicit forms have been discovered in some particular cases of sequences, including arithmetic, geometric-like, Fibonacci, Mersenne, and triangular (see [14] and references therein) and so on. However, for  $p \geq 1$ , no explicit form has been found even in such particular cases because it is not easy to find  $m_i^{(p)}$  in order to use Lemma 3 or Lemma 4.

The triangular numbers  $t_n = n(n + 1)/2$  ([16, A000217]), which are elementary class of polygonal numbers described by the early Pythagoreans, represent the numbers of points used to portray equilateral triangular patterns [20]. From Lemma 2, three consecutive triangular numbers satisfy the condition  $g(A) = 1$ . However, because of Lemma 1, they are not relatively prime and the situation is different for even and odd numbers. Hence, more detailed discussion is needed.

#### 3.1 $(t_n, t_{n+1}, t_{n+2})$ for general $p$

We can find explicit formulas for general  $n$  and smaller  $p$ . First, we discuss some general arguments. Later, we show several explicit formulas for smaller  $p$ .

When  $p = 0$ , from the formula of  $g_p(n) := g(t_n, t_{n+1}, t_{n+2}; p)$  given in (6) or (7), we have

$$g_0(2) = 17, g_0(3) = 29, g_0(4) = 89, g_0(5) = 125, g_0(6) = 251, g_0(7) = 323, \dots$$

When  $p = 1$ , from Theorem 2, we can have

$$g_1(2) = 23, g_1(3) = 59, g_1(4) = 119, g_1(5) = 209, g_1(6) = 335, g_1(7) = 503, \dots$$

When  $p = 2$ , from Theorem 3, we can have

$$g_2(2) = 29, g_2(4) = 149, g_2(5) = 230, g_2(6) = 419, g_2(7) = 575, g_2(8) = 899, \dots$$

Before mentioning the formulas for small  $p$ , we show the main result for general  $p \geq 0$ .

**Theorem 1.** *Let  $p$  be a nonnegative integer. For even  $n$ ,*

$$\begin{aligned} g \left( t_n, t_{n+1}, t_{n+2}; p(p+1) + \sum_{j=1}^p \left\lceil \frac{6j}{n} \right\rceil \right) \\ = \frac{(n+1)(n+2)(2(n+3)p+3n)}{4} - 1. \end{aligned}$$

For odd  $n \geq 3$ ,

$$\begin{aligned} g \left( t_n, t_{n+1}, t_{n+2}; \sum_{j=1}^p \left\lceil \frac{j}{2} \left( 1 + \frac{3}{n} \right) \right\rceil \right) \\ = \frac{(n+1)(n+2)((n+3)p+3(n-1))}{4} - 1. \end{aligned}$$

**Remark 2.** *This theorem implies that any positive integer  $N$  is expressed more than*

$$P' := \begin{cases} p(p+1) + \sum_{j=1}^p \left\lceil \frac{6j}{n} \right\rceil & (n:\text{even}) \\ \sum_{j=1}^p \left\lceil \frac{j}{2} \left( 1 + \frac{3}{n} \right) \right\rceil & (n:\text{odd}) \end{cases}$$

ways in terms of three consecutive triangular numbers  $t_n, t_{n+1}, t_{n+2}$  if

$$N > N' := \begin{cases} \frac{(n+1)(n+2)(2(n+3)p+3n)}{4} - 1 & (n:\text{even}) \\ \frac{(n+1)(n+2)((n+3)p+3(n-1))}{4} - 1 & (n:\text{odd}). \end{cases}$$

And  $N'$  itself is expressed exactly  $P'$  ways.

*Proof of Theorem 1.* First, let  $n$  be even. Then by Lemma 1,

$$\left\langle \frac{t_{n+1}}{\frac{n+2}{2}}, \frac{t_{n+2}}{\frac{n+2}{2}} \right\rangle = \langle n+1, n+3 \rangle,$$

where  $\langle x_1, x_2 \rangle = \{a_1x_1 + a_2x_2 | a_1, a_2 \geq 0\}$  denotes the all integers generated by the linear combination of  $x_1$  and  $x_2$  with nonnegative integral coefficients. By (3), for  $p \geq 0$ ,

$$g(n+1, n+3; p) = (n+1)(n+3)p + (n^2 + 2n - 1). \quad (11)$$

Thus, the length of the period of the numbers of the representation in terms of  $n+1$  and  $n+3$  is  $(n+1)(n+3)$ , and

$$\begin{aligned} p &= d(g(n+1, n+3; p); n+1, n+3) \\ &= d\left(\frac{n+2}{2}g(n+1, n+3; p); t_{n+1}, t_{n+2}\right) \end{aligned}$$

with

$$d(\nu; t_{n+1}, t_{n+2}) = 0 \quad \left( \text{if } \nu \not\equiv 0 \pmod{\frac{n+2}{2}} \right).$$

Therefore, for a positive integer  $m$ ,

$$\begin{aligned} &d\left(\frac{n+2}{2}m; t_n, t_{n+1}, t_{n+2}\right) \\ &= d\left(\frac{n+2}{2}m; t_{n+1}, t_{n+2}\right) + d\left(\frac{n+2}{2}(m - t_n); t_{n+1}, t_{n+2}\right) \\ &\quad + d\left(\frac{n+2}{2}(m - 2t_n); t_{n+1}, t_{n+2}\right) + \cdots \\ &\quad + d\left(\frac{n+2}{2}\left(m - \left\lfloor \frac{m}{t_n} \right\rfloor t_n\right); t_{n+1}, t_{n+2}\right) \\ &= d(m; n+1, n+3) + d(m - t_n; n+1, n+3) \\ &\quad + d(m - 2t_n; n+1, n+3) + \cdots + d\left(m - t_n \left\lfloor \frac{m}{t_n} \right\rfloor; n+1, n+3\right). \quad (12) \end{aligned}$$

Hence, when  $m = g(n+1, n+3; p) = (n+1)(n+3)p + (n^2 + 2n - 1)$ , we have

$$d\left(\frac{n+2}{2}g(n+1, n+3; p); t_n, t_{n+1}, t_{n+2}\right)$$

$$= \sum_{j=0}^{\lfloor g(n+1, n+3; p)/t_n \rfloor} d(g(n+1, n+3; p) - jt_n; n+1, n+3).$$

Here, each value of  $d(g(n+1, n+3; p) - jt_n; n+1, n+3)$  must be equal to any of  $0, 1, \dots, p$ . For convenience, put

$$g'_j = g'_{j,p} := g(n+1, n+3; p) - jt_n.$$

Remember that the largest integer whose number of representations in terms of  $n+1$  and  $n+3$  is equal to  $p$  is given in (11). For example, assume that we want to know the values  $j$  such that  $d((n+1)(n+3)p + (n^2 + 2n - 1) - jt_n; n+1, n+3) = p - 2$ . This has been achieved by solving  $g_{p-3} < g'_j \leq g_{p-2}$ . By

$$-3(n+1)(n+3) < -jt_n \leq -2(n+1)(n+3),$$

we get

$$6 + \frac{18}{n} > j \geq 4 + \frac{12}{n}.$$

Hence,

$$j = 4 + \left\lceil \frac{12}{n} \right\rceil, 5 + \left\lceil \frac{12}{n} \right\rceil, \dots, 5 + \left\lceil \frac{18}{n} \right\rceil.$$

It implies that the number of value  $(p-2)$  is given by

$$2 + \left\lceil \frac{18}{n} \right\rceil - \left\lceil \frac{12}{n} \right\rceil$$

among the sequence  $d(g(n+1, n+3; p) - jt_n; n+1, n+3)$  for  $0 \leq j \leq g(n+1, n+3; p)/t_n$ . Similarly, the numbers of values  $p, (p-1), \dots, 1$  are given by

$$\begin{aligned} & 2 + \left\lceil \frac{6}{n} \right\rceil, \\ & 2 + \left\lceil \frac{12}{n} \right\rceil - \left\lceil \frac{6}{n} \right\rceil, \\ & \dots, \\ & 2 + \left\lceil \frac{6p}{n} \right\rceil - \left\lceil \frac{6(p-1)}{n} \right\rceil, \end{aligned}$$

respectively. Therefore,

$$d\left(\frac{n+2}{2}g(n+1, n+3; p) + \left(\frac{n+2}{2} - 1\right)t_n; t_n, t_{n+1}, t_{n+2}\right)$$



$$\begin{aligned}
&= d\left(\frac{(n+1)(n+2)(2(n+3)p+3n)}{4} - 1; t_n, t_{n+1}, t_{n+2}\right) \\
&= p\left(2 + \left\lfloor \frac{6}{n} \right\rfloor\right) + (p-1)\left(2 + \left\lfloor \frac{12}{n} \right\rfloor - \left\lfloor \frac{6}{n} \right\rfloor\right) \\
&\quad + (p-2)\left(2 + \left\lfloor \frac{18}{n} \right\rfloor - \left\lfloor \frac{12}{n} \right\rfloor\right) + \cdots + 1 \cdot \left(2 + \left\lfloor \frac{6p}{n} \right\rfloor - \left\lfloor \frac{6(p-1)}{n} \right\rfloor\right) \\
&= p(p+1) + \sum_{j=1}^p \left\lfloor \frac{6j}{n} \right\rfloor.
\end{aligned}$$

Next, let  $n$  be odd. Then by Lemma 1,

$$\left\langle \frac{t_{n+1}}{n+2}, \frac{t_{n+2}}{n+2} \right\rangle = \left\langle \frac{n+1}{2}, \frac{n+3}{2} \right\rangle.$$

By (3), for  $p \geq 0$ ,

$$g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right) = \frac{(n+1)(n+3)}{4}p + \frac{n^2-5}{4}. \quad (13)$$

Thus, the length of the period of the numbers of the representation in terms of  $n+1$  and  $n+3$  is  $(n+1)(n+3)/4$ , and

$$\begin{aligned}
p &= d\left(g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right); \frac{n+1}{2}, \frac{n+3}{2}\right) \\
&= d\left((n+2)g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right); t_{n+1}, t_{n+2}\right)
\end{aligned}$$

with

$$d(\nu; t_{n+1}, t_{n+2}) = 0 \quad (\text{if } \nu \not\equiv 0 \pmod{n+2}).$$

Therefore, for a positive integer  $m$ ,

$$\begin{aligned}
&d((n+2)m; t_n, t_{n+1}, t_{n+2}) \\
&= d((n+2)m; t_{n+1}, t_{n+2}) + d((n+2)(m-t_n); t_{n+1}, t_{n+2}) \\
&\quad + d((n+2)(m-2t_n); t_{n+1}, t_{n+2}) + \cdots \\
&\quad + d\left((n+2)\left(m - \left\lfloor \frac{m}{t_n} \right\rfloor t_n\right); t_{n+1}, t_{n+2}\right) \\
&= d\left(m; \frac{n+1}{2}, \frac{n+3}{2}\right) + d\left(m-t_n; \frac{n+1}{2}, \frac{n+3}{2}\right)
\end{aligned}$$

$$+ d\left(m - 2t_n; \frac{n+1}{2}, \frac{n+3}{2}\right) + \cdots + d\left(m - t_n \left\lfloor \frac{m}{t_n} \right\rfloor; \frac{n+1}{2}, \frac{n+3}{2}\right). \quad (14)$$

Hence, when  $m = g((n+1)/2, (n+3)/2; p) = (n+1)(n+3)p/4 + (n^2 - 5)/4$ , we have

$$\begin{aligned} & d\left((n+2)g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right); t_n, t_{n+1}, t_{n+2}\right) \\ &= \sum_{j=0}^{\lfloor g(\frac{n+1}{2}, \frac{n+3}{2}; p)/t_n \rfloor} d\left(g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right) - jt_n; \frac{n+1}{2}, \frac{n+3}{2}\right). \end{aligned}$$

Here, each value of  $d(g((n+1)/2, (n+3)/2; p) - jt_n; (n+1)/2, (n+3)/2)$  must be equal to any of  $0, 1, \dots, p$ . For convenience, put

$$g'_j = g'_{j,p} := g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right) - jt_n.$$

Remember that the largest integer whose number of representations in terms of  $(n+1)/2$  and  $(n+3)/2$  is equal to  $p$  is given in (11). For example, assume that we want to know the values  $j$  such that  $d((n+1)(n+3)p + (n^2 + 2n - 1) - jt_n; (n+1)/2, (n+3)/2) = p - 2$ . This has been achieved by solving  $g_{p-3} < g'_j \leq g_{p-2}$ . By

$$-\frac{3(n+1)(n+3)}{4} < -jt_n \leq -\frac{2(n+1)(n+3)}{4},$$

we get

$$\frac{3}{2} \left(1 + \frac{3}{n}\right) > j \geq \frac{2}{2} \left(1 + \frac{3}{n}\right).$$

Hence, the number of value  $(p-2)$  is given by

$$\left\lceil \frac{3}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{2}{2} \left(1 + \frac{3}{n}\right) \right\rceil$$

among the sequence  $d(g((n+1)/2, (n+3)/2; p) - jt_n; (n+1)/2, (n+3)/2)$  for  $0 \leq j \leq g((n+1)/2, (n+3)/2; p)/t_n$ . Similarly, the numbers of values  $p, (p-1), \dots, 1$  are given by

$$\left\lceil \frac{1}{2} \left(1 + \frac{3}{n}\right) \right\rceil,$$

$$\left\lceil \frac{2}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{1}{2} \left(1 + \frac{3}{n}\right) \right\rceil,$$

$$\dots,$$

$$\left\lceil \frac{p}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{p-1}{2} \left(1 + \frac{3}{n}\right) \right\rceil,$$

respectively. Therefore,

$$\begin{aligned} & d\left((n+2)g\left(\frac{n+1}{2}, \frac{n+3}{2}; p\right) + (n+1)t_n; t_n, t_{n+1}, t_{n+2}\right) \\ &= d\left(\frac{(n+1)(n+2)((n+3)p + 3(n-1))}{4} - 1; t_n, t_{n+1}, t_{n+2}\right) \\ &= \left\lceil \frac{1}{2} \left(1 + \frac{3}{n}\right) \right\rceil + (p-1) \left( \left\lceil \frac{2}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{1}{2} \left(1 + \frac{3}{n}\right) \right\rceil \right) \\ &\quad + (p-2) \left( \left\lceil \frac{3}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{2}{2} \left(1 + \frac{3}{n}\right) \right\rceil \right) + \dots \\ &\quad + 1 \cdot \left( \left\lceil \frac{p}{2} \left(1 + \frac{3}{n}\right) \right\rceil - \left\lceil \frac{p-1}{2} \left(1 + \frac{3}{n}\right) \right\rceil \right) \\ &= \sum_{j=1}^p \left\lceil \frac{j}{2} \left(1 + \frac{3}{n}\right) \right\rceil. \end{aligned}$$

□

### 3.2 $(t_n, t_{n+1}, t_{n+2})$ for $p = 1$

We consider the general  $n$  case with the number of representations is at most 1 to give an explicit formula.

**Theorem 2.**

$$g(t_n, t_{n+1}, t_{n+2}; 1) = n(n+1)(n+2) - 1 \quad (n \geq 2).$$

*Proof.* When  $n$  is even, we can choose

$$m = \frac{(3n-1)(n+2)}{2} = g(n+1, n+3; 0) + t_n.$$

in (12). Since the only nonnegative integral solution of

$$m = a(n+1) + b(n+3) \quad (n:\text{even})$$

is given by  $a = (n - 2)/2$  and  $b = n$ , together with the definition of  $g(n + 1, n + 3; 0)$ , we see that

$$d(m; n + 1, n + 3) = 1 \quad \text{and} \quad d(m - jt_n; n + 1, n + 3) = 0 \quad (j \geq 1).$$

Since  $g(n + 1, n + 3; 1) - t_n = 2n^2 + 6n + 2 - n(n + 1)/2 > g(n + 1, n + 3; 0)$ , the number of representations of  $g(n + 1, n + 3; 1) - t_n$  cannot be 0. So,  $m = g(n + 1, n + 3; 1)$  is impossible. Hence, by (12), we have

$$\begin{aligned} g(t_n, t_{n+1}, t_{n+2}; 1) &= \frac{n+2}{2}m + \left(\frac{n-2}{2} + 1\right)t_n \\ &= n^3 + 3n^2 + 2n - 1. \end{aligned}$$

When  $n$  is odd with  $n \geq 3$ , we can choose

$$m = \frac{n^2 + 2n - 1}{2} = g\left(\frac{n+1}{2}, \frac{n+3}{2}; 1\right).$$

in (14). Now,

$$m - t_n = \frac{n-1}{2}$$

cannot be represented in terms of  $(n + 1)/2$  and  $(n + 3)/2$ . Hence, by (14), we have

$$\begin{aligned} g(t_n, t_{n+1}, t_{n+2}; 1) &= (n + 2)m + (n + 2 - 1)t_n \\ &= n^3 + 3n^2 + 2n - 1. \end{aligned}$$

□

### 3.3 $(t_n, t_{n+1}, t_{n+2})$ for $p = 2$

When  $p = 2$ , we have the following explicit form.

**Theorem 3.**

$$g(t_n, t_{n+1}, t_{n+2}; 2) = \begin{cases} \frac{(5n-3)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 5 \text{ is odd;} \\ \frac{(5n)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 2 \text{ is even.} \end{cases}$$

**Remark 3.** When  $n = 3$ , no positive integer can be represented in exactly two ways in terms of  $(t_3, t_4, t_5)$ . In this case, we can include the case where  $n = 3$  as  $g(t_3, t_4, t_5; 2) = g(t_3, t_4, t_5; 1) = 59$ .

*Proof of Theorem 3.* Let  $n$  be even. The number  $p = 2$  may be achieved in (12) as  $d(m; n+1, n+3) = 2$  and  $d(m - jt_n; n+1, n+3) = 0$  ( $j \geq 1$ ). Then the number  $p = 2$  may be achieved in (12) as  $d(m; n+1, n+3) = d(m - t_n; n+1, n+3) = 1$  and  $d(m - jt_n; n+1, n+3) = 0$  ( $j \geq 2$ ). There are two possibilities. But if  $m = g(n+1, n+3; 1) = 2n^2 + 6n + 2$ , by  $g(n+1, n+3; 1) - 2t_n = n^2 + 5n + 2 > g(n+1, n+3; 0) = n^2 + 2n - 1$ , this case is invalid. If  $m = 2n^2 + 3n - 1 = g(n+1, n+3; 0) + 2t_n$ ,

$$m = (n-1)(n+1) + n(n+3) \quad \text{and} \quad m - t_n = \frac{n-2}{2}(n+1) + n(n+3)$$

are the only 1 expressions, respectively. Therefore, we have

$$\begin{aligned} g(t_n, t_{n+1}, t_{n+2}; 2) &= \frac{n+2}{2}m + \frac{n}{2}t_n \\ &= \frac{5n^3 + 15n^2 + 10n - 4}{4}. \end{aligned}$$

Let  $n$  be odd. We can only choose

$$m = \frac{(3n+5)(n-1)}{4} = g\left(\frac{n+1}{2}, \frac{n+3}{2}; 0\right) + t_n.$$

in (14). In this case, the number  $p = 2$  is achieved in (12) as  $d(m; (n+1)/2, (n+3)/2) = 2$  and  $d(m - jt_n; n+1, n+3) = 0$  ( $j \geq 1$ ). In fact,  $m$  can be expressed in two ways as

$$m = (n-1)\frac{n+1}{2} + \frac{n+1}{2}\frac{n+3}{2} = \frac{n-5}{2}\frac{n+1}{2} + n\frac{n+3}{2} \quad (n \geq 5)$$

and  $g(\frac{n+1}{2}, \frac{n+3}{2}; 0) = m - t_n = (n^2 - 5)/4$  for  $j = 1$ . Therefore, we have

$$\begin{aligned} g(t_n, t_{n+1}, t_{n+2}; 1) &= (n+2)m + (n+2-1)t_n \\ &= \frac{5n^3 + 12n^2 + n - 10}{4}. \end{aligned}$$

□

### 3.4 $(t_n, t_{n+1}, t_{n+2})$ for $p \geq 3$

When  $3 \leq p \leq 10$ , we have the following explicit forms. We omit the details. When  $p$  is larger, the arguments become more complicated. If  $n$  is small in the following formulas, they can be calculated individually, so the details are also omitted.

**Theorem 4.**

$$g(t_n, t_{n+1}, t_{n+2}; 3) = \begin{cases} \frac{(5n+3)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 3 \text{ is odd;} \\ \frac{(5n+6)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 6 \text{ is even.} \end{cases}$$

For  $n = 5$  and  $n \geq 7$ , we have

$$g(t_n, t_{n+1}, t_{n+2}; 4) = \frac{3n(n+1)(n+2)}{2} - 1.$$

For  $n = 6$  and  $n \geq 8$ , we have

$$g(t_n, t_{n+1}, t_{n+2}; 5) = \frac{3(n+1)^2(n+2)}{2} - 1.$$

We have

$$g(t_n, t_{n+1}, t_{n+2}; 6) = \begin{cases} \frac{(7n-3)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 11 \text{ is odd;} \\ \frac{7n(n+1)(n+2)}{4} - 1 & \text{if } n \geq 8 \text{ is even,} \end{cases}$$

$$g(t_n, t_{n+1}, t_{n+2}; 7) = \begin{cases} \frac{(7n+3)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 5 \text{ is odd;} \\ \frac{(7n+6)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 6 \text{ is even,} \end{cases}$$

$$g(t_n, t_{n+1}, t_{n+2}; 8) = \begin{cases} \frac{(7n+9)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 9 \text{ is odd;} \\ \frac{(7n+12)(n+1)(n+2)}{4} - 1 & \text{if } n \geq 12 \text{ is even.} \end{cases}$$

For  $n = 11$  and  $n \geq 13$ , we have

$$g(t_n, t_{n+1}, t_{n+2}; 9) = 2n(n+1)(n+2) - 1.$$

For  $n = 3, 4$  and  $n \geq 8$ , we have

$$g(t_n, t_{n+1}, t_{n+2}; 10) = \frac{(4n+3)(n+1)(n+2)}{2} - 1.$$

**Conjecture.** For some nonnegative integer  $p$ , there exists an odd integer  $q$  and integers  $n_j$  ( $j = 1, 2, 3, 4$ ) such that

$$g(t_n, t_{n+1}, t_{n+2}; p) = \begin{cases} \frac{(qn-3)(n+1)(n+2)}{4} & \text{if } n \geq n_1 \text{ is odd;} \\ \frac{(qn)(n+1)(n+2)}{4} & \text{if } n \geq n_2 \text{ is even} \end{cases}$$

and

$$g(t_n, t_{n+1}, t_{n+2}; p+1) = \begin{cases} \frac{(qn+3)(n+1)(n+2)}{4} & \text{if } n \geq n_3 \text{ is odd;} \\ \frac{(qn+6)(n+1)(n+2)}{4} & \text{if } n \geq n_4 \text{ is even} \end{cases}$$

and so on. For some nonnegative integer  $p'$ , there exists an even integer  $q'$  and integers  $n_5$  and  $n_6$  such that

$$g(t_n, t_{n+1}, t_{n+2}; p') = \frac{(q'n)(n+1)(n+2)}{4} \quad (n \geq n_5)$$

and

$$g(t_n, t_{n+1}, t_{n+2}; p' + 1) = \frac{(q'n + 6)(n+1)(n+2)}{4} \quad (n \geq n_6)$$

and so on.

## 4 Triangular triples for smaller $n$

On the contrary, if  $n$  is smaller, we can find some explicit formulas for larger  $p$ . In this section, we consider the cases for  $n = 2, 3, 4$ .

### 4.1 $(t_3, t_4, t_5)$

When  $p$  is also a triangular number, we can obtain explicit formulas for general positive integer  $p$ .

**Proposition 1.** *For  $p \geq 1$ ,*

$$\begin{aligned} g(t_3, t_4, t_5; t_{p-1}) &= 30p - 1, \\ n(t_3, t_4, t_5; t_{p-1}) &= 30p - 16, \\ s(t_3, t_4, t_5; t_{p-1}) &= 15(30p^2 - 31p + 12). \end{aligned}$$

In fact, there does not exist a positive integer  $n$  such that for some nonnegative integer  $m$   $d(n; 6, 10, 15) \neq t_m$ .

*Sketch of the proof of Proposition 1.* Since  $(t_3, t_4, t_5) = (6, 10, 15)$ , we consider the representation  $10x_2 + 15x_3 = 5(2x_2 + 3x_3)$  ( $x_2, x_3 \geq 0$ ). As the generating function

$$\begin{aligned} & \frac{1}{(1-z^2)(1-z^3)} \\ &= \frac{1}{2(1-z^2)} + \frac{1}{3(1-z^3)} - \frac{z}{3(1-z^3)} + \frac{1}{6(1-z)^2} \\ &= \frac{1}{2} \sum s_1(n)z^n + \frac{1}{3} \sum s_2(n)z^n - \frac{1}{3} \sum s_3(n)z^n + \frac{1}{6} \sum (n+1)z^n, \end{aligned}$$

where

$$s_1(n) = \begin{cases} 1 & n \equiv 0 \pmod{2}; \\ 0 & \text{otherwise,} \end{cases} \quad s_2(n) = \begin{cases} 1 & n \equiv 0 \pmod{3}; \\ 0 & \text{otherwise,} \end{cases}$$

$$s_3(n) = \begin{cases} 1 & n \equiv 1 \pmod{3}; \\ 0 & \text{otherwise,} \end{cases}$$

the sequence is given by

$$\begin{aligned} & 1, 0, 1, 1, 1, 1, 2, 1, 2, 2, 2, 2, 3, 2, 3, 3, 3, 3, 4, 3, 4, 4, 4, 4, 5, 4, 5, 5, 5, 5, 6, 5, \dots \\ & = 1, 0, \overline{m, m, m, m, m+1, m_{m=1}^\infty} \\ & = \left\{ \left\lfloor \frac{n+2}{2} \right\rfloor - \left\lfloor \frac{n+2}{3} \right\rfloor \right\}_{n \geq 0} \end{aligned} \quad (15)$$

(Cf. [16, A008615]). Hence, for  $n \geq 0$

$$d(n; 2, 3) = \left\lfloor \frac{n+2}{2} \right\rfloor - \left\lfloor \frac{n+2}{3} \right\rfloor = d(5n; 10, 15),$$

and for  $p \geq 0$ ,  $g(2, 3; p) = 6p + 1$ . Thus,

$$m_5^{(p')} \pmod{6} = \max_{0 \leq i \leq 5} m_i^{(p')} = 5(6p + 1).$$

Here, because the number of representation of  $30p + 5$  in terms of  $(6, 10, 15)$  is just  $p$  more than that of  $30p - 1$ , by applying Corollary 4 (8), we have

$$g(t_3, t_4, t_5; p') = 30p - 1,$$

where  $p' = 1 + 2 + \dots + (p - 1) = t_{p-1}$ . Similarly, from the sequence (15), we see that for  $p' = t_{p-1}$

$$\begin{aligned} m_1^{(p')} &= 5(6p - 1), & m_2^{(p')} &= 5(6p - 2), & m_3^{(p')} &= 5(6p - 3), \\ m_4^{(p')} &= 5(6p - 4), & m_0^{(p')} &= 5(6p - 6). \end{aligned}$$

By applying Corollary 4 (9) and (10), we have the second and the third identities.  $\square$

For example, since

$$\{(x_2, x_3) \mid 2x_2 + 3x_3 = 19, x_2, x_3 \geq 0\} = \{(2, 5), (5, 3), (8, 1)\},$$



we see that

$$\begin{aligned} & \{(x_1, x_2, x_3) \mid 6x_1 + 5(2x_2 + 3x_3) = 89, x_1, x_2, x_3 \geq 0\} \\ & = \{(4, 2, 3), (4, 5, 1), (9, 2, 1)\}, \\ & \{(x_1, x_2, x_3) \mid 6x_1 + 5(2x_2 + 3x_3) = 95, x_1, x_2, x_3 \geq 0\} \\ & = \{(5, 2, 3), (5, 5, 1), (10, 2, 1), (0, 2, 5), (0, 5, 3), (0, 8, 1)\}. \end{aligned}$$

## 4.2 $(t_2, t_3, t_4)$

Next, we consider the case  $n = 2$ .

**Proposition 2.** For  $n \geq 0$  and  $j = 1, 2, 3, 4, 5$ ,

$$\begin{aligned} g(t_2, t_3, t_4; (n+1)(5n+2j)/2) &= 30n + 6j + 17, \\ n(t_2, t_3, t_4; (n+1)(5n+2j)/2) &= 30n + 6j + 8, \\ s(t_2, t_3, t_4; (n+1)(5n+2j)/2) &= 3(150n^2 + 5(12j + 17)n + 6j^2 + 17j + 23). \end{aligned}$$

*Proof.* By applying Lemma 4 for Lemma 5, we get the results.  $\square$

**Lemma 5.** For  $n \geq 0$  and  $j = 1, 2, 3, 4, 5$ ,

$$\begin{aligned} m_0^{((n+1)(5n+2j)/2)} &= 30n + 6j, \\ m_1^{((n+1)(5n+2j)/2)} &= 30n + 6j + 10, \\ m_2^{((n+1)(5n+2j)/2)} &= 30n + 6j + 20. \end{aligned}$$

*Proof.* By the generating function

$$\frac{1}{(1-z^3)(1-z^5)} = \sum_{n=0}^{\infty} d(n)z^n,$$

for  $n \geq 0$ , the sequence of number of representations  $d(n)$  of  $n = 3x_2 + 5x_3$  ( $x_2, x_3 \geq 0$ ) can be given as

$$(1, \overline{m-1, m-1, m, m-1, m, m, m-1, \underbrace{m, m, m, m, m, m, m}_7, m+1})_{m=1}^{\infty}.$$

Hence, for two variables  $(3, 5)$ , we see that

$$g(3, 5; p) = 15p + 7 \quad (p \geq 0).$$

For two variables  $(6, 10)$ , the same representation appears at  $2(15p + 7)$ . In this case, odd numbers cannot be represented in terms of 6 and 10. Hence,

$$\begin{aligned} d(2n; 3, 6, 10) &= d(2n + 3; 3, 6, 10) \\ &= d(2n; 6, 10) + d(2n - 6; 6, 10) + d(2n - 12; 6, 10) + \cdots + \begin{cases} d(4; 6, 10) \\ d(2; 6, 10) \\ d(0; 6, 10) \end{cases} \\ &= d(n; 3, 5) + d(n - 3; 3, 5) + d(n - 6; 3, 5) + \cdots + \begin{cases} d(2; 3, 5) \\ d(1; 3, 5) \\ d(0; 3, 5) \end{cases}. \end{aligned}$$

Thus, for  $p \geq 0$ , the number of representations of  $30p + 17$  (or of  $30p + 14$ ) in terms of  $(3, 6, 10)$  is given by

$$\begin{aligned} &d(15p + 7; 3, 5) + d(15p + 4; 3, 5) + d(15p + 1; 3, 5) + \\ &\quad \cdots + d(4; 3, 5) + d(1; 3, 5) \\ &= \underbrace{p + \cdots + p}_5 + \underbrace{(p - 1) + \cdots + (p - 1)}_5 + \cdots + \underbrace{1 + \cdots + 1}_5 + 0 + 0 + 0 \\ &= \frac{5p(p + 1)}{2}. \end{aligned}$$

Similarly, for  $p \geq 1$ , the numbers of representations of  $30p + 11$  (or of  $30p + 8$ ),  $30p + 5$  (or of  $30p + 2$ ),  $30p - 1$  (or of  $30p - 4$ ) and  $30p - 7$  (or of  $30p - 10$ ) in terms of  $(3, 6, 10)$  are given by

$$\begin{aligned} \frac{5p(p + 1)}{2} - p &= \frac{p(5p + 3)}{2}, & \frac{5p(p + 1)}{2} - 2p &= \frac{p(5p + 1)}{2}, \\ \frac{5p(p + 1)}{2} - 3p &= \frac{p(5p - 1)}{2}, & \frac{5p(p + 1)}{2} - 4p &= \frac{p(5p - 3)}{2}, \end{aligned} \quad (16)$$

respectively. These are all the cases for represented integers equivalent to 2 (mod 3). Next, consider the cases for represented integers equivalent to 0 (mod 3). For  $p \geq 1$ , the number of representations of  $30p - 3$  (or of  $30p - 6$ ) in terms of  $(3, 6, 10)$  is given by

$$d(15p - 3; 3, 5) + d(15p - 6; 3, 5) + d(15p - 9; 3, 5)$$

$$\begin{aligned}
& + \cdots + d(6; 3, 5) + d(3; 3, 5) + d(0; 3, 5) \\
& = \underbrace{p + \cdots + p}_5 + \underbrace{(p-1) + \cdots + (p-1)}_5 + \cdots + \underbrace{2 + \cdots + 2}_5 + 1 + 1 + 1 + 1 + 1 \\
& = \frac{5p(p+1)}{2}.
\end{aligned}$$

Similarly, for  $p \geq 1$ , the numbers of representations of  $30p-9$  (or of  $30p-12$ ),  $30p-15$  (or of  $30p-18$ ),  $30p-21$  (or of  $30p-24$ ) and  $30p-27$  (or of  $30p-30$ ) in terms of  $(3, 6, 10)$  are given as in (16), respectively. Finally, consider the cases for represented integers equivalent to  $1 \pmod{3}$ . For  $p \geq 1$ , the number of representations of  $30p+7$  (or of  $30p+4$ ) in terms of  $(3, 6, 10)$  is given by

$$\begin{aligned}
& d(15p+2; 3, 5) + d(15p-1; 3, 5) + d(15p-4; 3, 5) \\
& + \cdots + d(5; 3, 5) + d(2; 3, 5) \\
& = \underbrace{p + \cdots + p}_5 + \underbrace{(p-1) + \cdots + (p-1)}_5 + \cdots + \underbrace{1 + \cdots + 1}_5 + 0 \\
& = \frac{5p(p+1)}{2}.
\end{aligned}$$

Similarly, for  $p \geq 1$ , the numbers of representations of  $30p+1$  (or of  $30p-2$ ),  $30p-5$  (or of  $30p-8$ ),  $30p-11$  (or of  $30p-14$ ) and  $30p-17$  (or of  $30p-20$ ) in terms of  $(3, 6, 10)$  are given as in (16), respectively. Therefore, for  $p \geq 1$  and  $i = 0, 1, 2, 3, 4$ , we have  $m_2^{(5p(p+1)/2-ip)} = 30p + 20 - 6i$ ,  $m_1^{(5p(p+1)/2-ip)} = 30p + 10 - 6i$ ,  $m_0^{(5p(p+1)/2-ip)} = 30p - 6i$  and

$$g(3, 6, 10; 5p(p+1)/2 - ip) = 30p + 17 - 6i.$$

□

### 4.3 $(t_4, t_5, t_6)$

Consider the case  $(10, 15, 21)$  when  $n = 4$ . For simplicity, for nonnegative integers  $n$  and  $j$ , put

$$\begin{aligned}
r_e &= r_e(n, j) = \frac{n(7n+8-2j)}{4}, \\
r_o &= r_o(n, j) = \frac{7n^2 + (8-4j)n + 1}{4}, \\
r_{oo} &= r_{oo}(n, j) = \frac{7n^2 + (2-4j)n - 1}{4}.
\end{aligned}$$

**Proposition 3.** *For even  $n$ , when  $n \geq 0$  and  $j = 0$ , or when  $n \geq 2$  and  $j = 2, 3, 4, 5, 6, 8$ , we have*

$$\begin{aligned} g(t_4, t_5, t_6; r_e) &= 105n + 89 - 15j, \\ n(t_4, t_5, t_6; r_e) &= 105n + 44 - 15j, \\ s(t_4, t_5, t_6; r_e) &= \frac{15}{2}(735n^2 - 7(60j - 89)n + 60j^2 - 178j + 194). \end{aligned}$$

*For odd  $n$ , when  $n \geq 1$  and  $j = 0$ , or when  $n \geq 3$  and  $j = 1, 2, 3, 4$ , we have*

$$\begin{aligned} g(t_4, t_5, t_6; r_o) &= 105n + 89 - 30j, \\ n(t_4, t_5, t_6; r_o) &= 105n + 44 - 30j, \\ s(t_4, t_5, t_6; r_o) &= \frac{15}{2}(735n^2 - 7(30j - 89)n + 60j^2 - 89j + 194). \end{aligned}$$

*For odd  $n \geq 1$  and  $j = 0, 1$ , we have*

$$\begin{aligned} g(t_4, t_5, t_6; r_{oo}) &= 105n + 44 - 30j, \\ n(t_4, t_5, t_6; r_{oo}) &= 105n - 1 - 30j, \\ s(t_4, t_5, t_6; r_{oo}) &= \frac{15}{2}(735n^2 - 7(60j + 1)n + 60j^2 + 2j + 62). \end{aligned}$$

The results in Proposition 3 immediately follows from the next Lemma.

**Lemma 6.** *For even  $n$ , when  $n \geq 0$  and  $j = 0$ , or when  $n \geq 2$  and  $j = 2, 4, 6, 8$ , we have*

$$\begin{aligned} m_9^{(r_e)} &= 105n + 99 - 15j, & m_4^{(r_e)} &= 105n + 84 - 15j, \\ m_8^{(r_e)} &= 105n + 78 - 15j, & m_3^{(r_e)} &= 105n + 63 - 15j, \\ m_7^{(r_e)} &= 105n + 57 - 15j, & m_2^{(r_e)} &= 105n + 42 - 15j, \\ m_6^{(r_e)} &= 105n + 36 - 15j, & m_1^{(r_e)} &= 105n + 21 - 15j, \\ m_5^{(r_e)} &= 105n + 15 - 15j, & m_0^{(r_e)} &= 105n - 15j. \end{aligned}$$

*For even  $n$ , when  $n \geq 2$  and  $j = 3, 5$ , we have*

$$\begin{aligned} m_4^{(r_e)} &= 105n + 99 - 15j, & m_9^{(r_e)} &= 105n + 84 - 15j, \\ m_3^{(r_e)} &= 105n + 78 - 15j, & m_8^{(r_e)} &= 105n + 63 - 15j, \\ m_2^{(r_e)} &= 105n + 57 - 15j, & m_7^{(r_e)} &= 105n + 42 - 15j, \end{aligned}$$

$$\begin{aligned}
m_1^{(re)} &= 105n + 36 - 15j, & m_6^{(re)} &= 105n + 21 - 15j, \\
m_0^{(re)} &= 105n + 15 - 15j, & m_5^{(re)} &= 105n - 15j.
\end{aligned}$$

For odd  $n$ , when  $n \geq 1$  and  $j = 0$ , or when  $n \geq 3$  and  $j = 1, 2, 3, 4$ , we have

$$\begin{aligned}
m_4^{(ro)} &= 105n + 99 - 30j, & m_9^{(ro)} &= 105n + 84 - 30j, \\
m_3^{(ro)} &= 105n + 78 - 30j, & m_8^{(ro)} &= 105n + 63 - 30j, \\
m_2^{(ro)} &= 105n + 57 - 30j, & m_7^{(ro)} &= 105n + 42 - 30j, \\
m_1^{(ro)} &= 105n + 36 - 30j, & m_6^{(ro)} &= 105n + 21 - 30j, \\
m_0^{(ro)} &= 105n + 15 - 30j, & m_5^{(ro)} &= 105n - 30j.
\end{aligned}$$

For odd  $n \geq 1$  and  $j = 0, 1$ , we have

$$\begin{aligned}
m_9^{(r_{oo})} &= 105n + 54 - 30j, & m_4^{(r_{oo})} &= 105n + 39 - 30j, \\
m_8^{(r_{oo})} &= 105n + 33 - 30j, & m_3^{(r_{oo})} &= 105n + 18 - 30j, \\
m_7^{(r_{oo})} &= 105n + 12 - 30j, & m_2^{(r_{oo})} &= 105n - 3 - 30j, \\
m_6^{(r_{oo})} &= 105n - 9 - 30j, & m_1^{(r_{oo})} &= 105n - 24 - 30j, \\
m_5^{(r_{oo})} &= 105n - 30 - 30j, & m_0^{(r_{oo})} &= 105n - 45 - 30j.
\end{aligned}$$

**Remark 4.** For simplicity, for example, write  $g_n = g(t_4, t_5, t_6; n)$ . We have  $g_0 = 89, g_1 = 119, g_2 = 149, g_3 = 179, g_4 = 194, g_5 = 209, g_6 = 224, g_7 = 239, g_8 = 254, g_9 = 269, g_{10} = 284, g_{11} = 299, g_{13} = 314, \dots, g_{49} = 584, g_{51} = 599, g_{54} = 614, \dots, g_{201} = 1154, g_{206} = 1169, g_{212} = 1184$ . Note that the number of representations of  $n$  in terms of  $t_4, t_5, t_6$  cannot be  $12, \dots, 50, 52, 53, \dots, 202, 203, 204, 205, 207, 208, 209, 210, 211, \dots$ . Hence, for example,  $g_{206} = g_{207} = g_{208} = g_{209} = g_{210} = g_{211}$ .

We have  $n_0 = 45, n_1 = 74, n_2 = 104, n_3 = 134, n_4 = 149, n_5 = 164, n_6 = 179, n_7 = 194, n_8 = 209, n_9 = 224, n_{10} = 239, n_{11} = 254, n_{13} = 269, \dots, n_{49} = 539, n_{51} = 554, n_{54} = 569, \dots, n_{201} = 1109, n_{206} = 1124, n_{212} = 1139$ .

We have  $s_0 = 1455, s_1 = 3240, s_2 = 5925, s_3 = 9510, s_4 = 11640, s_5 = 13995, s_6 = 16575, s_7 = 19380, s_8 = 22410, s_9 = 25665, s_{10} = 29145, s_{11} = 32850, s_{13} = 36780, \dots, s_{49} = 145995, s_{51} = 154200, s_{54} = 162630, \dots, s_{201} = 615960, s_{206} = 632715, s_{212} = 649695$ .

*Proof of Lemma 6.* The sequence of the generating function

$$\frac{1}{(1-z^5)(1-z^7)} = \sum_{n=0}^{\infty} d_n z^n$$

has the period 35, and  $g(5, 7; p) = 35p + 23$  ( $p \geq 0$ ). Since  $d(35p + 23; 5, 7) = p$ , we have  $d(3(35p + 23); 15, 21) = p$ . As  $d(n; 15, 21) = 0$  for  $n \not\equiv 0 \pmod{3}$ , we see that

$$\begin{aligned} d(3n; 10, 15, 21) &= d(3n + 10; 10, 15, 21) = d(3n + 20; 10, 15, 21) \\ &= d(3n; 15, 21) + d(3n - 30; 15, 21) + d(3n - 60; 15, 21) \\ &\quad + \cdots + d(3(n - 10 \lfloor n/10 \rfloor); 15, 21) \\ &= d(n; 5, 7) + d(n - 10; 5, 7) + d(n - 20; 5, 7) + \cdots + d(n - 10 \lfloor n/10 \rfloor; 5, 7). \end{aligned}$$

Hence, for even  $p = 2p'$ , we have

$$\begin{aligned} d(105p + 69; 10, 15, 21) &= d(105p + 79; 10, 15, 21) = d(105p + 89; 10, 15, 21) \\ &= d(35p + 23; 5, 7) + d(35p + 13; 5, 7) + d(35p + 3; 5, 7) + \cdots \\ &\quad + d(23; 5, 7) + d(13; 5, 7) + d(3; 5, 7) \\ &= \underbrace{p + \cdots + p}_4 + \underbrace{(p-1) + \cdots + (p-1)}_3 + \underbrace{(p-2) + \cdots + (p-2)}_4 \\ &\quad + \underbrace{(p-3) + \cdots + (p-3)}_3 + \cdots + 0 + 0 + 0 \\ &= 4((2p') + (2p' - 2) + \cdots + 2) + 3((2p' - 1) + (2p' - 3) + \cdots + 1) \\ &= p'(7p' + 4) = \frac{p(7p + 8)}{4}. \end{aligned}$$

For odd  $p = 2p' + 1$ , we have

$$\begin{aligned} d(105p + 69; 10, 15, 21) &= d(105p + 79; 10, 15, 21) = d(105p + 89; 10, 15, 21) \\ &= d(35p + 23; 5, 7) + d(35p + 13; 5, 7) + d(35p + 3; 5, 7) + \cdots \\ &\quad + d(18; 5, 7) + d(8; 5, 7) \\ &= 4((2p' + 1) + (2p' - 1) + \cdots + 1) + 3((2p') + (2p' - 2) + \cdots + 2) + 0 + 0 \\ &= (p' + 1)(7p' + 4) = \frac{(p+1)(7p+1)}{4}. \end{aligned}$$

Hence, for even  $p$ ,

$$g(10, 15, 21; p(7p + 8)/4) = 105p + 89 \quad \text{and} \quad m_9^{(p(7p+8)/4)} = 105p + 99.$$

For odd  $p$ ,

$$g(10, 15, 21; (p+1)(7p+1)/4) = 105p+89 \quad \text{and} \quad m_4^{((p+1)(7p+1)/4)} = 105p+99.$$

Similarly, for even  $p$ ,  $d(105p + 39; 10, 15, 21) = d(105p + 49; 10, 15, 21) = d(105p + 59; 10, 15, 21), d(105p + 9; 10, 15, 21) = d(105p + 19; 10, 15, 21) = d(105p + 29; 10, 15, 21), d(105p - 21; 10, 15, 21) = d(105p - 11; 10, 15, 21) = d(105p - 1; 10, 15, 21)$  and  $d(105p - 51; 10, 15, 21) = d(105p - 41; 10, 15, 21) = d(105p - 31; 10, 15, 21)$  are given by

$$\begin{aligned} \frac{p(7p+8)}{4} - p &= \frac{p(7p+4)}{4}, & \frac{p(7p+8)}{4} - 2p &= \frac{7p^2}{4}, \\ \frac{p(7p+8)}{4} - 3p &= \frac{p(7p-4)}{4}, & \frac{p(7p+8)}{4} - 4p &= \frac{p(7p-8)}{4}, \end{aligned}$$

respectively. For odd  $p$ ,  $d(105p + 39; 10, 15, 21) = d(105p + 49; 10, 15, 21) = d(105p + 59; 10, 15, 21), d(105p + 9; 10, 15, 21) = d(105p + 19; 10, 15, 21) = d(105p + 29; 10, 15, 21), d(105p - 21; 10, 15, 21) = d(105p - 11; 10, 15, 21) = d(105p - 1; 10, 15, 21)$  and  $d(105p - 51; 10, 15, 21) = d(105p - 41; 10, 15, 21) = d(105p - 31; 10, 15, 21)$  are given by

$$\begin{aligned} \frac{(p+1)(7p+1)}{4} - p &= \frac{7p^2 + 4p + 1}{4}, \\ \frac{(p+1)(7p+1)}{4} - 2p &= \frac{7p^2 + 1}{4}, \\ \frac{(p+1)(7p+1)}{4} - 3p &= \frac{7p^2 - 4p + 1}{4}, \\ \frac{(p+1)(7p+1)}{4} - 4p &= \frac{(p-1)(7p-1)}{4}, \end{aligned}$$

respectively.

Next, for even  $p$ , we have

$$\begin{aligned} d(105p + 24; 10, 15, 21) &= d(105p + 34; 10, 15, 21) = d(105p + 44; 10, 15, 21) \\ &= d(35p + 8; 5, 7) + d(35p - 2; 5, 7) + d(35p - 12; 5, 7) + \cdots \\ &\quad + d(24; 5, 7) + d(14; 5, 7) + d(4; 5, 7) \\ &= p + p + \underbrace{(p-1) + \cdots + (p-1)}_4 + \underbrace{(p-2) + \cdots + (p-2)}_3 \\ &\quad + \cdots + \underbrace{2 + \cdots + 2}_3 + \underbrace{1 + \cdots + 1}_4 \\ &= \frac{p(7p+2)}{4}. \end{aligned}$$

For odd  $p$ , we have

$$\begin{aligned}
d(105p + 24; 10, 15, 21) &= d(105p + 34; 10, 15, 21) = d(105p + 44; 10, 15, 21) \\
&= p + p + \underbrace{(p-1) + \cdots + (p-1)}_4 + \underbrace{(p-2) + \cdots + (p-2)}_3 \\
&\quad + \cdots + \underbrace{2 + \cdots + 2}_4 + \underbrace{1 + \cdots + 1}_3 \\
&= \frac{7p^2 + 2p - 1}{4}.
\end{aligned}$$

Hence, for even  $p$ ,

$$g(10, 15, 21; p(7p + 2)/4) = 105p + 44 \quad \text{and} \quad m_4^{(p(7p+2)/4)} = 105p + 54.$$

For odd  $p$ ,

$$g(10, 15, 21; (7p^2 + 2p - 1)/4) = 105p + 44 \quad \text{and} \quad m_9^{((7p^2+2p-1)/4)} = 105p + 54.$$

Similarly, for even  $p$ ,  $d(105p + 14; 10, 15, 21) = d(105p + 4; 10, 15, 21) = d(105p - 6; 10, 15, 21)$  is given by

$$\frac{p(7p + 2)}{4} - p = \frac{p(7p - 2)}{4}.$$

For odd  $p$ ,  $d(105p + 14; 10, 15, 21) = d(105p + 4; 10, 15, 21) = d(105p - 6; 10, 15, 21)$  is given by

$$\frac{7p^2 + 2p - 1}{4} - p = \frac{7p^2 - 2p - 1}{4}.$$

□

#### 4.4 Larger $n$

We can continue to obtain explicit formulas of the functions  $g$ ,  $n$  and  $s$  for  $n = 5, 6, \dots$ , but the situation becomes more complicated. For example, concerning  $g_p := g(t_5, t_6, t_7; p)$ , we see that

$$\begin{aligned}
g_0 &= 125, \quad g_1 = 209, \quad g_2 = 230, \quad g_3 = 293, \quad g_4 = 314, \quad g_5 = 335, \\
g_6 &= 377, \quad g_7 = 398, \quad g_8 = 419, \quad g_9 = 440, \quad g_{10} = 461, \quad g_{11} = 482, \\
g_{12} &= 503, \quad g_{13} = 524, \quad g_{14} = 545, \quad g_{16} = 566, \quad \dots
\end{aligned}$$

Note that  $g_{15} = g_{14}$  because the (largest) integer whose number of representation is exactly 15 does not exist.



## 5 Conclusion

In this paper, we are successful to obtain certain kinds of natural generalizations of Frobenius numbers together with Sylvester numbers and sums for three variables  $(a_1, a_2, a_3)$ , when they are consecutive triangular numbers. In other words, up to the present, any explicit formula for the maximum number, total number, and sum of numbers that can be expressed at most in  $p$  ways have not been found for three variables or more. In this paper, explicit formulas are given when the triangular number is small, or when  $p$  is small. By continuing the method presented here, we will be able to obtain explicit formulas for larger triplets and larger  $p$ 's, but the discussion must become more complicated.

## Conflict of interest

The author declares to have no conflict of interests.

## Acknowledgement

The author thanks the anonymous reviewers for their careful reading of the manuscript and their insightful comments and suggestions.

## References

- [1] R. Apéry, *Sur les branches superlinéaires des courbes algébriques*, C. R. Acad. Sci. Paris **222** (1946), 1198–1200.
- [2] L. Bardomero and M. Beck, *Frobenius coin-exchange generating functions*, Amer. Math. Monthly **127** (2020), no. 4, 308–315.
- [3] M. Beck, I. M. Gessel and T. Komatsu, *The polynomial part of a restricted partition function related to the Frobenius problem*, Electron. J. Combin. **8** (No.1) (2001), #N7.
- [4] M. Beck and C. Kifer, *An extreme family of generalized Frobenius numbers*, Integers **11** (2011), A24, 639–645.
- [5] A. Brauer and B. M. Shockley, *On a problem of Frobenius*, J. Reine. Angew. Math. **211** (1962), 215–220.

- [6] T. C. Brown and P. J. Shiue, *A remark related to the Frobenius problem*, Fibonacci Quart. **31** (1993), 32–36.
- [7] A. Cayley, *On a problem of double partitions*, Philos. Mag. **XX** (1860), 337–341.
- [8] F. Curtis, *On formulas for the Frobenius number of a numerical semi-group*, Math. Scand. **67** (1990), 190–192.
- [9] T. Komatsu, *On the number of solutions of the Diophantine equation of Frobenius—General case*, Math. Commun. **8** (2003), 195–206.
- [10] T. Komatsu, *The Frobenius number associated with the number of representations for sequences of repunits*, C. R. Math. Acad. Sci. Paris (accepted for publication).
- [11] T. Komatsu, *On  $p$ -Frobenius and related numbers due to  $p$ -Apéry set*, arXiv:2111.11021.
- [12] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set*, Irish Math. Soc. Bull. **87** (2021), 21–29.
- [13] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set in more variables*, Kyushu J. Math. **76** (2022), 163–175.
- [14] A. M. Robles-Pérez and J. C. Rosales, *The Frobenius number for sequences of triangular and tetrahedral numbers*, J. Number Theory **186** (2018), 473–492.
- [15] E. S. Selmer, *On the linear diophantine problem of Frobenius*, J. Reine Angew. Math. **293/294** (1977), 1–17.
- [16] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, available at [oeis.org](http://oeis.org). (2021).
- [17] J. J. Sylvester, *On the partition of numbers*, Quart. J. Pure Appl. Math. **1** (1857), 141–152.
- [18] J. J. Sylvester, *On subinvariants, i.e. semi-invariants to binary quantics of an unlimited order*, Am. J. Math. **5** (1882), 119–136.
- [19] J. J. Sylvester, *Mathematical questions with their solutions*, Educational Times **41** (1884), 21.

- [20] J. J. Tattersall, *Elementary number theory in nine chapters*, Second ed. Cambridge Univ. Press, Cambridge, 2005. xii+430 pp. ISBN: 978-0-521-61524-2; 0-521-61524-0
- [21] A. Tripathi, *The number of solutions to  $ax + by = n$* , Fibonacci Quart. **38** (2000), 290–293.
- [22] A. Tripathi, *On sums of positive integers that are not of the form  $ax+by$* , Amer. Math. Monthly **115** (2008), 363–364.
- [23] A. Tripathi, *Formulae for the Frobenius number in three variables*, J. Number Theory **170** (2017), 368–389.
- [24] A. Tripathi, *On a special case of the Frobenius problem*, J. Integer Seq. **20** (2017), Article 17.7.2, 12 pp.