

ON THE RELATIVE MORRISON-KAWAMATA CONE CONJECTURE

ZHAN LI AND HANG ZHAO

ABSTRACT. We relate the Morrison-Kawamata cone conjecture for Calabi-Yau fiber spaces to the existence of Shokurov polytopes. For K3 fibrations, the existence of (weak) fundamental domains for movable cones is established. The relationship between the relative cone conjecture and the cone conjecture for its geometric or generic fibers is studied.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Geometry of convex cones	15
4. Generic properties of fibrations and structures of cones	18
5. A variant of the cone conjecture	26
6. Generic and Geometric cone conjectures	31
References	37

1. INTRODUCTION

The purpose of this paper is to study the following (relative) Morrison-Kawamata cone conjecture [Mor93, Mor96, Kaw97, Tot09].

Conjecture 1.1. *Let $(X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Let Γ_B, Γ_A be the images of the pseudo-automorphism group $\text{PsAut}(X/S, \Delta)$ and the automorphism group $\text{Aut}(X/S, \Delta)$ under the natural group homomorphism $\text{PsAut}(X/S, \Delta) \rightarrow \text{GL}(N^1(X/S)_{\mathbb{R}})$ respectively.*

- (1) *The cone $\overline{\text{Mov}}^e(X/S)$ has a (weak) rational polyhedral fundamental domain under the action of Γ_B .*
- (2) *The cone $\overline{\text{Amp}}^e(X/S)$ has a (weak) rational polyhedral fundamental domain under the action of Γ_A .*

Relevant notions in Conjecture 1.1 are explained in Section 2 and Section 3. There are different choices of cones in the cone conjecture, see Remark 5.6 for the reason of the above choice.

At the expense of some ambiguity, for simplicity, we call Conjecture 1.1 (1) and (2) the (weak) cone conjecture for movable cones and the (weak) cone conjecture for ample cones respectively. Although our primary interest is in complex varieties, we need to work with non-algebraically closed fields. When X is a smooth Calabi-Yau variety over a field K , the analogous cone conjecture still makes sense, and we also call it the cone conjecture.

The cone conjecture is beyond merely predicting the shape of cones of Calabi-Yau varieties. In fact, for an arbitrary klt pair, if (X, Δ) is its minimal model and $(X, \Delta) \rightarrow S$ is the morphism to its canonical model, then the cone conjecture for movable cones predicts finiteness of minimal models (see Proposition 5.3 for the precise statement). Moreover, compared with the weaker predictions of finitely many $\text{PsAut}(X/S, \Delta)$ - or $\text{Aut}(X/S, \Delta)$ -equivalence classes, it seems that the existence of (weak) fundamental domains is crucial in the proof of the cone conjecture (see the proof of Proposition 6.4).

When $X \rightarrow S$ is a birational morphism, [BCHM10] established the finiteness of $\text{PsAut}(X/S)$ -equivalence classes. Finiteness of $\text{PsAut}(X/S)$ -equivalence classes is also known when $\dim X \leq 3$, $\dim S > 0$ ([Kaw97]) and elliptic fibrations ([FHS21]). When S is a point, Conjecture 1.1 is known for surfaces ([Tot09]), abelian varieties ([PS12]) and large classes of Calabi-Yau manifolds with Picard number 2 ([Ogu14, LP13]). Analogous cone conjecture of $\text{Mov}(X/\mathbb{C})_+$ (see Definition 3.1) is also known for a projective hyperkähler manifold X ([Mar11]). Over arbitrary fields of characteristic $\neq 2$, the cone conjecture is known for K3 surfaces [BLvL20]. Analogous cone conjecture of $\text{Mov}(X/K)_+$ is also known for a hyperkähler variety X over a field K with characteristic 0 ([Tak21], cf. Remark 6.3). On the other hand, it is known that Conjecture 1.1 no longer holds true for lc pairs (see [Tot09]). We recommend [LOP18] for a survey of relevant results.

The new ingredient of the current paper is to study the cone conjecture from the perspective of Shokurov polytopes. We propose the following conjecture which seems to be more tractable.

Conjecture 1.2. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space.*

(1) *There exists a polyhedral cone $P_M \subset \text{Eff}(X/S)$ such that*

$$\bigcup_{g \in \text{PsAut}(X/S, \Delta)} g \cdot P_M \supset \text{Mov}(X/S).$$

(2) *There exists a polyhedral cone $P_A \subset \text{Eff}(X/S)$ such that*

$$\bigcup_{g \in \text{Aut}(X/S, \Delta)} g \cdot P_A \supset \text{Amp}(X/S).$$

Using results of [Loo14] and assuming standard conjectures of log minimal model program (LMMP), we show that Conjecture 1.2 is nearly equivalent to the cone conjecture (when S is a point, they are indeed equivalent).

Theorem 1.3. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assume that good minimal models exist for effective klt pairs in dimension $\dim(X/S)$.*

- (1) *If $R^1 f_* \mathcal{O}_X$ is a torsion sheaf, then the weak cone conjecture for $\overline{\text{Mov}}^e(X/S)$ is equivalent to the Conjecture 1.2 (1).*
- (2) *If $\overline{\text{Mov}}(X/S)$ is non-degenerate, then the cone conjecture for $\overline{\text{Mov}}^e(X/S)$ is equivalent to the Conjecture 1.2 (1);*
- (3) *The cone conjecture for $\overline{\text{Amp}}^e(X/S)$ is equivalent to the Conjecture 1.2 (2).*

It seems that Conjecture 1.2 is more fundamental as it incorporates in both the finiteness of models or contractions, and the existence of fundamental domains.

Using this circle of ideas, we study a Calabi-Yau fiber space $X \rightarrow S$ fibered by K3 surfaces. This means that for a general closed point $t \in S$, its fiber X_t is a smooth K3 surface. We establish the (weak) cone conjecture of $\overline{\text{Mov}}^e(X/S)$ for K3 fibrations.

Theorem 1.4. *Let $f : X \rightarrow S$ be a Calabi-Yau fiber space such that X has terminal singularities.*

If f is fibered by K3 surfaces, then the weak cone conjecture of $\overline{\text{Mov}}^e(X/S)$ holds true.

Moreover, if $\overline{\text{Mov}}(X/S)$ is non-degenerate, then the cone conjecture holds true for $\overline{\text{Mov}}^e(X/S)$. In particular, if S is \mathbb{Q} -factorial, then the cone conjecture holds true for $\overline{\text{Mov}}^e(X/S)$.

We discuss the contents of the paper. Section 2 gives necessary background materials and fixes notation. Section 3 develops the geometry of convex cones following [Loo14]. Section 4 establishes properties of generic and geometric fibers which will be used in Section 6. Section 5 studies the relationship between the cone conjecture and Conjecture 1.2. In particular, Theorem 1.3 is proven. Section 6 studies the cone

conjecture by assuming that it holds true for geometric or generic fibers. Theorem 1.4 is shown in Section 6.1.

Acknowledgements. We benefit from discussions with Lie Fu, Yong Hu, Zhiyuan Li, Yannan Qiu, Hao Sun, and Jinsong Xu. Hang Zhao is partially supported by the grant No. ST20210105. Both authors are partially supported by a grant from SUSTech.

2. PRELIMINARIES

Let $f : X \rightarrow S$ be a projective morphism between normal quasi-projective varieties over \mathbb{C} . Then f is called a fibration if f has connected fibers. We write X/S to mean that X is over S .

By divisors, we mean Weil divisors. For $\mathbb{K} = \mathbb{Z}, \mathbb{Q}, \mathbb{R}$ and two \mathbb{K} -divisors A, B on X , $A \sim_{\mathbb{K}} B/S$ means that A and B are \mathbb{K} -linearly equivalent over S . If A, B are \mathbb{R} -Cartier divisors, then $A \equiv B/S$ means that A and B are numerically equivalent over S .

We use $\text{Supp } E$ to denote the support of the divisor E . A divisor E on X is called a vertical divisor (over S) if $f(\text{Supp } E) \neq S$. A vertical divisor E is called a very exceptional divisor if for any prime divisor P on S , over the generic point of P , we have $\text{Supp } f^*P \not\subset \text{Supp } E$ (see [Bir12, Definition 3.1]). If f is a birational morphism, then the notion of very exceptional divisor coincides with that of exceptional divisor.

Let X be a normal complex variety and Δ be an \mathbb{R} -divisor on X , then (X, Δ) is called a log pair. We assume that $K_X + \Delta$ is \mathbb{R} -Cartier for a log pair (X, Δ) . Then $f : (X, \Delta) \rightarrow S$ is called a Calabi-Yau fibration/fiber space if $X \rightarrow S$ is a fibration, X is \mathbb{Q} -factorial and $K_X + \Delta \sim_{\mathbb{R}} 0/S$. When (X, Δ) has lc singularities (see Section 2.2), then $K_X + \Delta \sim_{\mathbb{R}} 0/S$ is equivalent to the weaker condition $K_X + \Delta \equiv 0/S$ by [HX16, Corollary 1.4].

2.1. Movable cones and ample cones. Let V be a finite-dimensional real vector space. A set $C \subset V$ is called a cone if for any $x \in C$ and $\lambda \in \mathbb{R}_{>0}$, we have $\lambda \cdot x \in C$. We use $\text{Int}(C)$ to denote the relative interior of C and call $\text{Int}(C)$ the open cone. By convention, the origin is an open cone. A cone is called a polyhedral cone (resp. rational polyhedral cone) if it is a closed convex cone generated by finite vectors (resp. rational vectors). If $S \subset V$ is a subset, then $\text{Conv}(S)$ denotes the convex hull of S , and $\text{Cone}(S)$ denotes the closed convex cone generated by S . As we are only concerned about convex cones in this paper, we also call them cones.

Let $\text{Pic}(X/S)$ be the relative Picard group. Let

$$N^1(X/S) := \text{Pic}(X/S)/\cong$$

be the lattice. Set $\text{Pic}(X/S)_{\mathbb{K}} := \text{Pic}(X/S) \otimes_{\mathbb{Z}} \mathbb{K}$ and $N^1(X/S)_{\mathbb{K}} := N^1(X/S) \otimes_{\mathbb{Z}} \mathbb{K}$ for $\mathbb{K} = \mathbb{Q}$ or \mathbb{R} . If D is an \mathbb{R} -Cartier divisor, then $[D] \in N^1(X/S)_{\mathbb{R}}$ denotes the corresponding divisor class. To abuse the terminology, we also call $[D]$ an \mathbb{R} -Cartier divisor.

Recall that a Cartier divisor D is called movable if the base locus of the linear system $|D|$ has codimension > 1 . We list relevant cones of $N^1(X/S)_{\mathbb{R}}$ appeared in the paper:

- (1) $\text{Eff}(X/S)$: the cone generated by effective \mathbb{Q} -Cartier divisors (also generated by effective \mathbb{R} -Cartier divisors);
- (2) $\overline{\text{Eff}}(X/S)$: the closure of $\text{Eff}(X/S)$;
- (3) $\text{Mov}(X/S)$: the cone generated by movable divisors;
- (4) $\overline{\text{Mov}}(X/S)$: the closure of $\text{Mov}(X/S)$;
- (5) $\overline{\text{Mov}}^e(X/S) := \overline{\text{Mov}}(X/S) \cap \text{Eff}(X/S)$;
- (6) $\text{Mov}(X/S)_+ := \text{Conv}(\overline{\text{Mov}}(X/S) \cap N^1(X/U)_{\mathbb{Q}})$ (see Definition 3.1);
- (7) $\text{Amp}(X/S)$: the cone generated by ample divisors;
- (8) $\overline{\text{Amp}}(X/S)$: the closure of $\text{Amp}(X/S)$;
- (9) $\overline{\text{Amp}}^e(X/S) := \overline{\text{Amp}}(X/S) \cap \text{Eff}(X/S)$;
- (10) $\text{Amp}(X/S)_+ := \text{Conv}(\overline{\text{Amp}}(X/S) \cap N^1(X/U)_{\mathbb{Q}})$.

If K is a field of characteristic zero and X is a variety over K , then the above cones still make sense for X . We use $\text{Mov}(X/K)$, $\text{Amp}(X/K)$, etc. to denote the corresponding cones.

Recall that for a birational map $g : X \dashrightarrow Y/S$, if D is an \mathbb{R} -Cartier divisor on X , then the pushforward of D , g_*D , is defined as follows. Let $p : W \rightarrow X$, $q : W \rightarrow Y$ be birational morphisms such that $g \circ p = q$, then $g_*D := q_*(p^*D)$. This is independent of the choice of p and q .

Let Δ be a divisor on a \mathbb{Q} -factorial variety X . We use $\text{Bir}(X/S, \Delta)$ to denote the birational automorphism group of (X, Δ) over S . To be precise, $\text{Bir}(X/S, \Delta)$ consists of birational maps $g : X \dashrightarrow X/S$ such that $g_* \text{Supp } \Delta = \text{Supp } \Delta$. A birational map is called a pseudo-automorphism if it is isomorphic in codimension 1. Let $\text{PsAut}(X/S, \Delta)$ be the subgroup of $\text{Bir}(X/S, \Delta)$ consisting of pseudo-automorphisms. Let $\text{Aut}(X/S, \Delta)$ be the subgroup of $\text{Bir}(X/S, \Delta)$ consisting of automorphisms of X/S . For a field K , if X is a variety over K and Δ is a divisor on X , then we still use $\text{Bir}(X/K, \Delta)$, $\text{PsAut}(X/K, \Delta)$ and $\text{Aut}(X/K, \Delta)$ to denote the birational automorphism group, the

pseudo-automorphism group and the automorphism group of X/K respectively.

Let $g \in \text{Bir}(X/S, \Delta)$ and D be an \mathbb{R} -Cartier divisor on a \mathbb{Q} -factorial variety X . Because the pushforward map g_* preserves numerical equivalence classes, there is a linear map

$$g_* : N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X/S)_{\mathbb{R}}, \quad [D] \mapsto [g_*D].$$

However, if $g \in \text{Bir}(X/S)$ is not isomorphic in codimension 1, then for a $[D] \in \text{Mov}(X/S)$, $[g_*D]$ may not be in $\text{Mov}(X/S)$. Moreover, $(g, [D]) \mapsto [g_*D]$ is not a group action of $\text{Bir}(X/S, \Delta)$ on $N^1(X/S)_{\mathbb{R}}$. For one thing, if D is a divisor contracted by g , then $g_*^{-1}(g_*[D]) = 0 \neq (g^{-1} \circ g)_*[D]$.

On the other hand, it is straightforward to check that

$$\begin{aligned} \text{PsAut}(X/S, \Delta) \times N^1(X/S)_{\mathbb{R}} &\rightarrow N^1(X/S)_{\mathbb{R}} \\ (g, [D]) &\mapsto [g_*D], \end{aligned}$$

is a group action. We use $g \cdot D, g \cdot [D]$ to denote $g_*D, [g_*D]$ respectively. Let Γ_B and Γ_A be the images of $\text{PsAut}(X/S, \Delta)$ and $\text{Aut}(X/S, \Delta)$ under the natural group homomorphism

$$\iota : \text{PsAut}(X/S, \Delta) \rightarrow \text{GL}(N^1(X/S)_{\mathbb{R}}).$$

Because $\Gamma_B, \Gamma_A \subset \text{GL}(N^1(X/S))$, Γ_B and Γ_A are discrete subgroups. By abusing the notation, we also write g for $\iota(g) \in \Gamma_B$, and denote $\iota(g)([D])$ by $g \cdot [D]$. Then $\text{Mov}(X/S), \overline{\text{Mov}}(X/S), \overline{\text{Mov}}^e(X/S)$ and $\text{Mov}(X/S)_+$ are all invariant under the action of $\text{PsAut}(X/S, \Delta)$. Similarly, $\text{Amp}(X/S), \text{Amp}^e(X/S)$ and $\text{Amp}(X/S)_+$ are all invariant under the action of $\text{Aut}(X/S, \Delta)$.

When X is not a \mathbb{Q} -factorial variety, if X admits a small \mathbb{Q} -factorization $\tilde{X} \rightarrow X$, then the (weak) cone conjecture of $\overline{\text{Mov}}^e(X/S)$ is referred to the (weak) cone conjecture of $\overline{\text{Mov}}^e(\tilde{X}/S)$. The validity of the conjecture is independent of the choice of \tilde{X} .

The following example gives a birational map which is not a pseudo-automorphism.

Example 2.1. *Let $f(x, y, z)$ be a general homogenous cubic polynomial. Let $D := \{f(x, y, z) = 0\} \subset \mathbb{P}^2$ and $B := \{f(-x, y, z) = 0\} \subset \mathbb{P}^2$. Then $(\mathbb{P}^2, \frac{1}{2}D + \frac{1}{2}B)$ is a klt Calabi-Yau pair. Let*

$$p_1 = [a : b : c], p_2 = [-a : b : c] \in D \cap B$$

be two distinct points. Let $\pi_i : X_i \rightarrow \mathbb{P}^2, i = 1, 2$ be the blowing up of p_i such that $E_i, i = 1, 2$ are corresponding exceptional divisors. If D_i, B_i

are the strict transforms of D, B on X_i , then

$$K_{X_i} + \frac{1}{2}D_i + \frac{1}{2}B_i = \pi_i^*(K_{\mathbb{P}^2} + \frac{1}{2}D + \frac{1}{2}B).$$

Therefore, each $(X_i, \frac{1}{2}D_i + \frac{1}{2}B_i)$ is a klt Calabi-Yau pair. Moreover, $(X_1, \frac{1}{2}D_1 + \frac{1}{2}B_1)$ is isomorphic to $(X_2, \frac{1}{2}D_2 + \frac{1}{2}B_2)$ through $\pi_2^{-1} \circ \tau \circ \pi_1$, where $\tau : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is given by $[x : y : z] \mapsto [-x : y : z]$. However, the birational map

$$\pi_2^{-1} \circ \pi_1 : (X_1, \frac{1}{2}D_1 + \frac{1}{2}B_1) \dashrightarrow (X_2, \frac{1}{2}D_2 + \frac{1}{2}B_2)$$

is not isomorphic in codimension 1. In fact, this map contracts E_1 and extracts E_2 .

2.2. Minimal models of varieties. Let (X, Δ) be a log pair. For a divisor D over X , if $f : Y \rightarrow X$ is a birational morphism from a smooth variety Y such that D is a divisor on Y , then the log discrepancy of D with respect to (X, Δ) is defined to be

$$a(D; X, \Delta) := \text{mult}_D(K_Y - f^*(K_X + \Delta)) + 1.$$

This definition is independent of the choice of Y . A log pair (X, Δ) (or its singularity) is called sub-klt (resp. sub-lc) if the log discrepancy of any divisor over X is > 0 (resp. ≥ 0). If $\Delta \geq 0$, then a sub-klt (resp. sub-lc) pair (X, Δ) is called klt (resp. lc). If $\Delta = 0$ and the log discrepancy of any divisor over X is > 1 , then X is said to have terminal singularities. A fibration/fiber space $(X, \Delta) \rightarrow S$ is called a klt (resp. terminal) fibration/fiber space if (X, Δ) is klt (resp. terminal). In the sequel, we will use a well-known fact that if X has terminal singularities with K_X nef/ S , then $\text{Bir}(X/S) = \text{PsAut}(X/S)$.

Let $X \rightarrow S$ be a projective morphism of normal quasi-projective varieties. Suppose that (X, Δ) is klt. Let $\phi : X \dashrightarrow Y/S$ be a birational contraction (i.e. ϕ does not extract divisors) of normal quasi-projective varieties over S , where Y is projective over S . We write $\Delta_Y := \phi_*\Delta$ for the strict transform of Δ . Then $(Y/S, \Delta_Y)$ is a weak log canonical model of $(X/S, \Delta)$ if $K_Y + \Delta_Y$ is nef/ S and $a(D; X, \Delta) \geq a(D; Y, \Delta_Y)$ for any divisor D over X .

Lemma 2.2. *Let $(X/S, \Delta)$ be a klt pair with $[K_X + \Delta] \in \overline{\text{Mov}}(X/S)$. Suppose that $(Y/S, \Delta_Y)$ is a weak log canonical model of $(X/S, \Delta)$. Then $(X/S, \Delta)$ admits a weak log canonical model $(Y'/S, \Delta_{Y'})$ such that Y' is \mathbb{Q} -factorial and X, Y' are isomorphic in codimension 1.*

Proof. Let $\tau : X \dashrightarrow Y$ and $p : W \rightarrow X, q : W \rightarrow Y$ be birational morphisms such that $\tau = q \circ p^{-1}$. Then $p^*(K_X + \Delta) = q^*(K_Y + \Delta_Y) + E$,

where $E \geq 0$ is q -exceptional. As $[K_X + \Delta] \in \overline{\text{Mov}}(X/S)$, we have $E = 0$. Otherwise, there is a family of curves $\{C_t\}$ which covers a component of E such that $q_*C_t = 0$ and $E \cdot C_t < 0$. But this contradicts with $(K_X + \Delta) \cdot C_t \geq 0$. If F is a prime divisor contracted by τ , then $a(F; Y, \Delta_Y) = a(F; X, \Delta) \leq 1$. By [BCHM10, Corollary 1.4.3], there exist a \mathbb{Q} -factorial variety Y' and a birational morphism $\nu : Y' \rightarrow Y$ which extracts all the divisors contracted by τ . Hence $K_{Y'} + \Delta_{Y'} = \nu^*(K_Y + \Delta_Y)$ where $\Delta_{Y'}$ is the strict transform of Δ on Y' . Therefore,

$$\nu^{-1} \circ \tau : (X/S, \Delta) \dashrightarrow (Y'/S, \Delta_{Y'})$$

is also a weak log canonical model of $(X/S, \Delta)$. In particular, X, Y' are isomorphic in codimension 1. \square

A weak log canonical model $(Y/S, \Delta_Y)$ of $(X/S, \Delta)$ is called a good minimal model of $(X/S, \Delta)$ if $K_Y + \Delta_Y$ is semi-ample/ S . It is well-known that the existence of a good minimal model of $(X/S, \Delta)$ implies that any weak log canonical model of $(X/S, \Delta)$ is a good minimal model (for example, see [Bir12, Remark 2.7]).

By saying that “good minimal models of effective klt pairs exist in dimension n ”, we mean that for any projective variety X of dimension n over \mathbb{C} , if (X, Δ) is klt with Δ a \mathbb{Q} -divisor such that the Kodaira dimension $\kappa(K_X + \Delta) \geq 0$, then (X, Δ) has a good minimal model.

Theorem 2.3 ([HX13, Theorem 2.12]). *Let $f : X \rightarrow S$ be a surjective projective morphism and (X, Δ) a klt pair such that for a very general closed point $s \in S$, the fiber $(X_s, \Delta_s = \Delta|_{X_s})$ has a good minimal model. Then (X, Δ) has a good minimal model over S .*

Notice that the assumption (2) in [HX13, Theorem 2.12] is automatically satisfied when (X, Δ) is klt by [BCHM10, Theorem 1.2].

2.3. Shokurov polytopes. Let V a finite-dimensional \mathbb{R} -vector space. A polytope (resp. rational polytope) $P \subset V$ is the convex hull of finite points (resp. rational points) in V . In particular, a polytope is always closed and bounded. We use $\text{Int}(P)$ to denote the relative interior of P and call $\text{Int}(P)$ the open polytope. By convention, a single point is an open polytope. Therefore, $\mathbb{R}_{>0} \cdot P$ is an open polyhedral cone iff P is an open polytope.

Theorem 2.4 ([SC11, Theorem 3.4]). *Let X be a \mathbb{Q} -factorial variety and $f : X \rightarrow S$ be a fibration. Assume that good minimal models exist for effective klt pairs in dimension $\dim(X/S)$. Let $D_i, i = 1, \dots, k$ be effective \mathbb{Q} -divisors on X . Suppose that $P \subset \bigoplus_{i=1}^k [0, 1)D_i$ is a rational*

polytope such that for any $\Delta \in P$, (X, Δ) is klt and $\kappa(K_F + \Delta|_F) \geq 0$, where F is a general fiber of f .

Then P can be decomposed into a disjoint union of finitely many open rational polytopes $P = \sqcup_{i=1}^m Q_i^\circ$ such that for any $B, D \in Q_i^\circ$, if $(Y/S, B_Y)$ is a weak log canonical model of $(X/S, B)$, then $(Y/S, D_Y)$ is also a weak log canonical model of $(X/S, D)$.

For the convenience of the reader, we give the proof of Theorem 2.4. The argument essentially follows from [BCHM10, Lemma 7.1]. However, we need to take care of the weaker assumption on the existence of weak log canonical models, as opposed to log terminal models.

Proof of Theorem 2.4. We proceed by induction on the dimension of P . Note that by Theorem 2.3, $(X/S, \Delta)$ has a good minimal model/ S for any $\Delta \in P$.


Step 1. If there exists a $\Delta_0 \in \text{Int}(P)$ such that $K_X + \Delta_0 \equiv 0/S$, then we show the claim. In fact, let P' be a face of P . By the induction hypothesis, $P' = \sqcup_j \tilde{Q}_j^\circ$ such that each \tilde{Q}_j° is an open rational polytope, and for $B, D \in \tilde{Q}_j^\circ$, if $(Y/S, B_Y)$ is a weak log canonical model of $(X/S, B)$, then $(Y/S, D_Y)$ is also a weak log canonical model of $(X/S, D)$. For a $t \in [0, 1]$,

$$\begin{aligned} K_X + tB + (1-t)\Delta_0 &= t(K_X + B) + (1-t)(K_X + \Delta_0) \equiv t(K_X + B)/S, \\ K_X + tD + (1-t)\Delta_0 &= t(K_X + D) + (1-t)(K_X + \Delta_0) \equiv t(K_X + D)/S. \end{aligned}$$

Hence, $(Y/S, tB_Y + (1-t)\Delta_{0,Y})$ is a weak log canonical model of $(X/S, tB + (1-t)\Delta_0)$ iff $(Y/S, B_Y)$ is a weak log canonical model of $(X/S, B)$ iff $(Y/S, D_Y)$ is a weak log canonical model of $(X/S, D)$ iff $(Y/S, tD_Y + (1-t)\Delta_{0,Y})$ is a weak log canonical model of $(X/S, tD + (1-t)\Delta_0)$. Therefore,

$$P = \left(\sqcup_j \text{Int}(\text{Conv}(\tilde{Q}_j^\circ, \Delta_0)) \right) \sqcup \{\Delta_0\}$$

satisfies the claim.



Step 2. Next, we show the general case. It suffices to show the result locally around any point $\Delta_0 \in P$. We can assume that Δ_0 is a rational point and P is a small rational polytope containing Δ_0 . During the argument, by saying that shrinking P , we mean that replacing P by a sufficiently small rational polytope $P' \subset P$ such that $P' \supset P \cap \mathbb{B}(\Delta_0, \epsilon)$, where $\mathbb{B}(\Delta_0, \epsilon) \subset \bigoplus_{i=1}^k \mathbb{R} \cdot D_i$ is the ball centered at Δ_0 with radius $\epsilon \in \mathbb{R}_{>0}$.

Let $(X'/S, \Delta'_0)$ be a weak log canonical model of $(X/S, \Delta_0)$. By Theorem 2.3, there exist a contraction $\pi : X' \rightarrow Z'/S$ and an ample/ S \mathbb{Q} -Cartier divisor A on Z' such that $K_{X'} + \Delta'_0 \sim_{\mathbb{Q}} \pi^* A/S$.

Let $p : W \rightarrow X, q : W \rightarrow X'$ be birational morphisms such that $q \circ p^{-1}$ is the natural map $X \dashrightarrow X'$. Moreover, we assume that p is a log resolution of $(X, \sum_{i=1}^k D_i)$. Let $\tilde{D}_i, i = 1, \dots, k$ be the strict transforms of $D_i, i = 1, \dots, k$ on W , and $E_j, j = 1, \dots, l$ be prime q -exceptional divisors. Shrinking P , there is a linear bijective map defined over \mathbb{Q} ,

$$L : P \rightarrow P_W, \quad \Delta \mapsto L(\Delta),$$

such that $P_W \subset (\bigoplus_{i=1}^k \tilde{D}_i) \oplus (\bigoplus_{j=1}^l E_j)$ is a rational polytope, and

$$(2.3.1) \quad K_W + L(\Delta) = p^*(K_X + \Delta) + E(\Delta),$$

such that $E(\Delta) \geq 0$ is p -exceptional and $(W, L(\Delta))$ is still klt for each $\Delta \in P$. Let $\Delta_{W,0} := L(\Delta_0)$. Run a $(K_W + \Delta_{W,0})$ -LMMP with scaling of an ample divisor over X' , then it terminates with $(W'/X', \Delta_{W',0})$ by [BCHM10, Corollary 1.4.2]. As $(X'/S, \Delta'_0)$ is a weak log canonical model of $(X/S, \Delta_0)$, there exists a q -exceptional divisor $E_0 \geq 0$ such that

$$p^*(K_X + \Delta_0) = q^*(K_{X'} + \Delta'_0) + E_0.$$

Hence

$$K_W + \Delta_{W,0} = q^*(K_{X'} + \Delta'_0) + E(\Delta_0) + E_0.$$

Because $E(\Delta_0) + E_0 \geq 0$ is q -exceptional, we have

$$K_{W'} + \Delta_{W',0} = q'^*(K_{X'} + \Delta'_0),$$

where $q' : W' \rightarrow X'$ is the natural morphism. In particular,

$$K_{W'} + \Delta_{W',0} \equiv 0/Z'.$$

Let $\theta : W \dashrightarrow W'/X$ be the natural map. Shrinking P , we can assume that θ is $(K_W + L(\Delta))$ -negative (see [BCHM10, Definition 3.6.1]) for each $\Delta \in P$.

Step 3. By Step 1, $P_W = \sqcup Q_i^\circ$ can be decomposed into a disjoint union of finitely many open rational polytopes such that for any $B', D' \in Q_i^\circ$, if $(W_i''/Z', B'')$ is a weak log canonical model of $(W'/Z', B')$ then $(W_i''/Z', D'')$ is a weak log canonical model of $(W'/Z', D')$, where B'', D'' are the strict transforms of B', D' respectively. In the sequel, we fix a W_i'' for each Q_i° .

We claim that after shrinking P_W , for any $\Delta_i \in Q_i^\circ$, $K_{W_i''} + \Delta_i''$ is nef over S , where Δ_i'' is the strict transform of Δ_i . Let Δ_i be a vertex of Q_i° . By Theorem 2.3, $(W_i''/Z', \Delta_i)$ is semi-ample/ Z' . Let $\tau : W_i'' \rightarrow T_i/Z'$ be the morphism such that $K_{W_i''} + \Delta_i'' \sim_{\mathbb{Q}} \tau^* H_i/Z'$, where H_i is an ample/ Z' \mathbb{Q} -Cartier divisor on T_i . Hence, there is a \mathbb{Q} -Cartier divisor Θ on Z' such that $K_{W_i''} + \Delta_i'' = \tau^* H_i + (\mu \circ \tau)^* \Theta$, where $\mu : T_i \rightarrow Z'$. Then $t(H_i + \mu^* \Theta) + (1-t)\mu^* A$ is nef over S when $t \in [0, t_0]$ for some rational number $0 < t_0 \ll 1$. Note that

$$t(K_{W_i''} + \Delta_i'') + (1-t)(K_{W_i''} + \Delta_{W,0}'') \sim_{\mathbb{R}} \tau^*(t(H_i + \mu^* \Theta) + (1-t)\mu^* A) / S,$$

where $\Delta_{W,0}''$ is the strict transform of $\Delta_{W,0}$ on W_i'' . Replacing Δ_i by $t_0 \Delta_i + (1-t_0) \Delta_{W,0}$ and repeating this process for each vertex of Q_i° , we get a polytope satisfying the desired claim.

Step 4. Let $P := L^{-1}(P_W)$ be the polytope corresponding to P_W under the map L . Let $Q_i^\circ := L^{-1}(Q_i^\circ)$ be the corresponding open rational polytope. To complete the proof, it suffices to show that for any $B, D \in Q_i^\circ$, if $(Y/S, B_Y)$ is a weak log canonical model of $(X/S, B)$ then $(Y/S, D_Y)$ is a weak log canonical model of $(X/S, D)$.

Let $r : V \rightarrow W, s : V \rightarrow W', \tilde{p} : V \rightarrow Y, \tilde{q} : V \rightarrow W_i''$ be birational morphisms which commute with the existing maps. Moreover, $r, s, \tilde{p}, \tilde{q}$ can be assumed to be log resolutions. By (2.3.1), for a $\Delta \in Q_i^\circ$,

$$(2.3.2) \quad r^*(K_W + L(\Delta)) = r^* p^*(K_X + \Delta) + r^*(E(\Delta)).$$

As $\theta : W \dashrightarrow W'$ is $(K_W + L(\Delta))$ -negative, $(W_i''/S, L(\Delta)'')$ is also a weak log canonical model of $(W/S, L(\Delta))$, where $L(\Delta)''$ is the strict

transform of $L(\Delta)$. Then there is a \tilde{q} -exceptional divisor $F(\Delta) \geq 0$ such that

$$r^*(K_W + L(\Delta)) = \tilde{q}^*(K_{W_i''} + L(\Delta)'') + F(\Delta).$$

Combining with (2.3.2), we have

$$\tilde{q}^*(K_{W_i''} + L(\Delta)'') + F(\Delta) = r^*p^*(K_X + \Delta) + r^*(E(\Delta)).$$

Hence $-F(\Delta) + r^*(E(\Delta))$ is nef over X . As

$$(p \circ r)_*(-F(\Delta) + r^*(E(\Delta))) = (p \circ r)_*(-F(\Delta)) \leq 0,$$

we have

$$-F(\Delta) + r^*(E(\Delta)) \leq 0$$

by the negativity lemma (see [KM98, Lemma 3.39]). Let

$$(2.3.3) \quad r^*p^*(K_X + \Delta) = \tilde{p}^*(K_Y + \Delta_Y) + \Theta(\Delta),$$

where $\Theta(\Delta)$ is \tilde{p} -exceptional. Hence

$$\tilde{q}^*(K_{W_i''} + L(\Delta)'') + F(\Delta) = \tilde{p}^*(K_Y + \Delta_Y) + \Theta(\Delta) + r^*(E(\Delta)).$$

As $-F(\Delta) + \Theta(\Delta) + r^*(E(\Delta))$ is nef over Y and

$$\tilde{p}_*(-F(\Delta) + \Theta(\Delta) + r^*(E(\Delta))) = \tilde{p}_*(-F(\Delta)) \leq 0,$$

we have $-F(\Delta) + \Theta(\Delta) + r^*(E(\Delta)) \leq 0$ by the negativity lemma.

Now we use that $(Y/S, B_Y)$ is a weak log canonical model of $(X/S, B)$. As $K_Y + B_Y$ is nef/ S , $F(B) - \Theta(B) - r^*(E(B))$ is nef over W_i'' . As $F(B)$ is \tilde{q} -exceptional, we have

$$\tilde{q}_*(F(B) - \Theta(B) - r^*(E(B))) \leq 0.$$

By the negativity lemma again, we have $F(B) - \Theta(B) - r^*(E(B)) \leq 0$. Therefore, $F(B) - \Theta(B) - r^*(E(B)) = 0$. Note that $F(\Delta) - \Theta(\Delta) - r^*(E(\Delta))$ is linear for $\Delta \in Q_i^\circ$. Because Q_i° is open and $F(\Delta) - \Theta(\Delta) - r^*(E(\Delta)) \geq 0$ for each $\Delta \in Q_i^\circ$, we have $F(\Delta) - \Theta(\Delta) - r^*(E(\Delta)) = 0$ for each $\Delta \in Q_i^\circ$. Thus $\Theta(\Delta) = F(\Delta) - r^*(E(\Delta)) \geq 0$ and

$$\tilde{q}^*(K_{W_i''} + L(\Delta)'') = \tilde{p}^*(K_Y + \Delta_Y)$$

is nef/ S for any $\Delta \in Q_i^\circ$. As $X \dashrightarrow Y$ does not extract divisors, (Y, D_Y) is also a weak log canonical model of (X, D) by (2.3.3). \square

Remark 2.5. *In the proof of Theorem 2.4, we use the existence of good minimal models in Step 1 and Step 3. In Step 1, this is needed to ensure that the statement of Theorem 2.4 holds true for lower dimensional polytopes. In Step 3, let $L(\Delta)'$ be the strict transform of $L(\Delta)$ on W' (see (2.3.1)), then we need that $(F', \Delta'|_{F'})$ has a good minimal model, where F' is a general fiber of $\pi \circ q' : W' \rightarrow Z'$. Therefore, the assumption that “good minimal models exist for effective klt pairs*

in dimension $\dim(X/S)$ ” can be replaced by the following more precise form:

Suppose that $h : X \dashrightarrow X'/S$ is a birational contraction and $p : W \rightarrow X/S, q : W \rightarrow X'/S$ are birational morphisms such that $h = q \circ p^{-1}$. Suppose that $X' \rightarrow S$ factors through a projective morphism $\pi : X' \rightarrow Z'/S$ between normal varieties. Let $\theta : W \dashrightarrow W'$ be a birational contraction and $q' : W' \rightarrow X'$ be a morphism such that $q' = \theta \circ q^{-1}$. For each $\Delta \in P$, let

$$K_W + \tilde{\Delta} = p^*(K_X + \Delta) + E$$

such that $(W, \tilde{\Delta})$ is klt and $E \geq 0$ is p -exceptional. Let $\tilde{\Delta}'$ be the strict transform of $\tilde{\Delta}$ on W' and F' be a general fiber of $\pi \circ q' : W' \rightarrow Z'$. Then $(F', \tilde{\Delta}'|_{F'})$ has a good minimal model.

This remark will be needed in the proof of Proposition 4.4 (3) and (4).

Theorem 2.6. *Let $(X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assume that good minimal models of effective klt pairs exist in dimension $\dim(X/S)$. Let $P \subset \text{Eff}(X/S)$ be a rational polyhedral cone. Then P is a finite union of open rational polyhedral cones $P = \cup_{i=0}^m P_i^\circ$ such that whenever*

- (1) B, D are effective divisors with $[B], [D] \in P_i^\circ$, and
- (2) $(X, \Delta + \epsilon B), (X, \Delta + \epsilon D)$ are klt for some $\epsilon \in \mathbb{R}_{>0}$,

then if $(Y/S, \Delta_Y + \epsilon B_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon B)$, then $(Y/S, \Delta_Y + \epsilon D_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon D)$.

Proof. Let $\Delta = \sum_{i=1}^m c_i \Delta_i$ be the decomposition into irreducible components. Then

$$\{\Theta \in \oplus_{i=1}^m \mathbb{R} \cdot \Delta_i \mid K_X + \Theta \equiv 0/S\}$$

is a subspace of $\oplus_{i=1}^m \mathbb{R} \cdot \Delta_i$ defined over \mathbb{Q} . Hence, there exists a \mathbb{Q} -Cartier divisor $\tilde{\Delta}$ such that $(X, \tilde{\Delta}) \rightarrow S$ is a klt Calabi-Yau fiber space. Let $\tilde{\epsilon} \in \mathbb{R}_{>0}$ such that $(X, \tilde{\Delta} + \tilde{\epsilon} B)$ is klt. By

$$K_X + \Delta + \epsilon B \equiv \frac{\epsilon}{\tilde{\epsilon}}(K_X + \tilde{\Delta} + \tilde{\epsilon} B)/S,$$

$(Y/S, \Delta_Y + \epsilon B_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon B)$ iff $(Y/S, \tilde{\Delta}_Y + \tilde{\epsilon} B_Y)$ is a weak log canonical model of $(X/S, \tilde{\Delta} + \tilde{\epsilon} B)$. Therefore, replacing Δ by $\tilde{\Delta}$, we can assume that Δ is a \mathbb{Q} -Cartier divisor.

Let $\tilde{P} \subset \text{Eff}(X/S)$ be a rational polytope such that $\mathbb{R}_{\geq 0} \cdot \tilde{P} = P$. We can choose \tilde{P} such that $0 \notin \tilde{P}$. Let $\Delta_j \geq 0, j = 1 \dots k$ be effective

\mathbb{Q} -Cartier divisors such that $\tilde{P} = \text{Conv}(\Delta_j \mid j = 1 \dots k)$. Replacing Δ_j by $\epsilon \Delta_j$ for some $\epsilon \in \mathbb{R}_{>0}$, we can assume that $(X, \Delta + \Delta_j)$ is klt for each j . Let

$$\tilde{P} + \Delta = \sqcup_{i=1}^m (Q_i^\circ + \Delta)$$

be the decomposition as in Theorem 2.4. For each open rational polytope $Q_i^\circ + \Delta$, and $\Theta_1 + \Delta, \Theta_2 + \Delta \in Q_i^\circ + \Delta$, if $(Y/S, \Theta_{1,Y} + \Delta_Y)$ is a weak log canonical model of $(X, S, \Theta_1 + \Delta)$, then $(Y/S, \Theta_{2,Y} + \Delta_Y)$ is a weak log canonical model of $(X, S, \Theta_2 + \Delta)$. Let P_i° be the image of the open rational polyhedral cone $\mathbb{R}_{>0} \cdot Q_i^\circ$ in $N^1(X/S)_\mathbb{R}$. Set $P_0^\circ = \{0\}$, then $P = \cup_{i=0}^m P_i^\circ$. Note that this union may not be disjoint.

The claim certainly holds true for P_0° . For effective divisors B, D with $[B], [D] \in P_i^\circ, i > 0$, there exist $\Delta_B, \Delta_D \in Q_i^\circ$ such that

$$[B] = t[\Delta_B], \quad [D] = s[\Delta_D] \quad \text{for some } t, s \in \mathbb{R}_{>0}.$$

By $K_X + \Delta \equiv 0/S$,

$$(2.3.4) \quad \begin{aligned} K_X + \Delta + \Delta_B &\equiv \frac{1}{\epsilon t} (K_X + \Delta + \epsilon B)/S \\ K_X + \Delta + \Delta_D &\equiv \frac{1}{\epsilon s} (K_X + \Delta + \epsilon D)/S. \end{aligned}$$

Therefore, $(Y/S, \Delta_Y + \epsilon B_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon B)$ iff $(Y/S, \Delta_Y + \Delta_{B,Y})$ is a weak log canonical model of $(X/S, \Delta + \Delta_B)$. By Theorem 2.4, this implies that $(Y/S, \Delta_Y + \Delta_{D,Y})$ is a weak log canonical model of $(X/S, \Delta + \Delta_D)$. Hence $(Y/S, \Delta_Y + \epsilon D_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon D)$ by (2.3.4) again. \square

Theorem 2.7 ([Sho96, §6.2. First main theorem]). *Let $(X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Let $P \subset \text{Eff}(X/S)$ be a rational polyhedral cone. Then*

$$P_N := P \cap \overline{\text{Amp}}(X/S)$$

is a rational polyhedral cone.

Proof. Let $D_j, 1 \leq j \leq k$ be effective \mathbb{Q} -Cartier divisors on X such that $P = \text{Cone}([D_j] \mid 1 \leq j \leq k)$. Replacing D_j by ϵD_j for some $\epsilon \in \mathbb{Q}_{>0}$, we can assume that $(X, \Delta + D_j)$ is klt for each j . Then

$$\mathcal{N} = \{D \in \oplus_{j=1}^k [0, 1]D_j \mid D \text{ is nef over } S\}$$

is a rational polytope by [Sho96, §6.2. First main theorem] (also see [Bir11, Proposition 3.2 (3)]). The image $[\mathcal{N}]$ of \mathcal{N} in $N^1(X/S)_\mathbb{R}$ is still a rational polytope. By the construction,

$$P_N = \text{Cone}([\mathcal{N}]).$$

Thus P_N is a rational polyhedral cone. \square

3. GEOMETRY OF CONVEX CONES

Let $V(\mathbb{Z})$ be a lattice and $V(\mathbb{Q}) := V(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q}$, $V := V(\mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R}$. A cone $C \subset V$ is non-degenerate if it does not contain an affine line. This is equivalent to say that its closure \bar{C} does not contain a non-trivial vector space.

In the following, we assume that Γ is a group and $\rho : \Gamma \rightarrow \mathrm{GL}(V)$ is a group homomorphism. The group Γ acts on V through ρ . For a $\gamma \in \Gamma$ and an $x \in V$, we write $\gamma \cdot x$ or γx for the action. For a set $S \subset V$, set $\Gamma \cdot S := \{\gamma \cdot x \mid \gamma \in \Gamma, x \in S\}$. Suppose that this action leaves a convex cone C and some lattice in $V(\mathbb{Q})$ invariant. We assume that $\dim C = \dim V$. The following definition slightly generalizes [Loo14, Proposition-Definition 4.1].

Definition 3.1. *Under the above notation and assumptions.*

- (1) *Suppose that $C \subset V$ is an open convex cone (may be degenerate). Let*

$$C_+ := \mathrm{Conv}(\bar{C} \cap V(\mathbb{Q}))$$

be the convex hull of rational points in \bar{C} .

- (2) *We say that (C_+, Γ) is of polyhedral type if there is a polyhedral cone $\Pi \subset C_+$ such that $\Gamma \cdot \Pi \supset C$.*

Remark 3.2. *Recall that a polyhedral cone is closed by definition (see Section 2.1).*

Proposition 3.3 ([Loo14, Proposition-Definition 4.1]). *Under the above notation and assumptions. If C is non-degenerate, then the following conditions are equivalent:*

- (1) *there exists a polyhedral cone $\Pi \subset C_+$ with $\Gamma \cdot \Pi = C_+$;*
 (2) *there exists a polyhedral cone $\Pi \subset C_+$ with $\Gamma \cdot \Pi \supset C$.*

Moreover, in case (2), we necessarily have $\Gamma \cdot \Pi = C_+$.

Definition 3.4. *Let $\rho : \Gamma \hookrightarrow \mathrm{GL}(V)$ be an injective group homomorphism and $C \subset V$ be a cone (may not necessarily be open). Let $\Pi \subset C$ be a (rational) polyhedral cone. Suppose that Γ acts on C . Then Π is called a weak (rational) polyhedral fundamental domain for C under the action Γ if*

- (1) *$\Gamma \cdot \Pi = C$, and*
 (2) *for each $\gamma \in \Gamma$, either $\gamma\Pi = \Pi$ or $\gamma\Pi \cap \mathrm{Int}(\Pi) = \emptyset$.*

Moreover, for $\Gamma_{\Pi} := \{\gamma \in \Gamma \mid \gamma\Pi = \Pi\}$, if $\Gamma_{\Pi} = \{\mathrm{id}\}$, then Π is called a (rational) polyhedral fundamental domain.

Lemma 3.5 ([Loo14, Theorem 3.8 & Application 4.14]). *Under the notation and assumptions of Definition 3.1. Suppose that $\rho : \Gamma \hookrightarrow \mathrm{GL}(V)$ is injective. Let (C_+, Γ) be of polyhedral type with C non-degenerate. Then under the action of Γ , C_+ admits a rational polyhedral fundamental domain.*

Proof. Let V^* be the dual vector space of V with pairing

$$\langle -, - \rangle : V \times V^* \rightarrow \mathbb{R}.$$

Let

$$C^* := \{y \in V^* \mid \langle x, y \rangle \geq 0 \text{ for all } x \in C\}$$

be the dual cone of C , and $(C^*)^\circ$ be the relative interior of C^* . By C non-degenerate and $\dim C = \dim V$, we still have $\dim(C^*)^\circ = \dim V$.

The group Γ naturally acts on V^* . In fact, for a $\gamma \in \Gamma$ and a $y \in V^*$, $\gamma \cdot y$ is defined by the relation $\langle x, \gamma \cdot y \rangle = \langle \gamma \cdot x, y \rangle$ for all $x \in V$. It is straightforward to check that this action gives an injective group homomorphism $\Gamma \hookrightarrow \mathrm{GL}(V^*)$ which leaves C^* and a lattice in $V^*(\mathbb{Q})$ invariant. Therefore, by [Loo14, Theorem 3.8], Γ acts properly discontinuously on $(C^*)^\circ$.

By [Loo14, Application 4.14], for each $\xi \in (C^*)^\circ \cap V(\mathbb{Q})^*$, there is a rational polyhedral cone σ associated with ξ , such that σ is a rational polyhedral fundamental domain for the action of Γ on C_+ whenever the stabilizer subgroup $\Gamma_\xi = \{1\}$. It suffices to find such ξ to complete the proof. As Γ acts properly discontinuously on $(C^*)^\circ$, for any polyhedral cone $P \subset (C^*)^\circ$ such that $\dim P = \dim(C^*)^\circ = \dim V$, the set

$$\{\gamma \in \Gamma \mid \gamma P^\circ \cap P^\circ \neq \emptyset\}$$

is a finite set. Then a general $\xi \in P^\circ \cap V^*(\mathbb{Q})$ satisfies $\Gamma_\xi = \{1\}$. \square

The following consequence of having a polyhedral fundamental domain is well-known (see [Loo14, Corollary 4.15] or [Mor15, (4.7.7) Proposition])

Theorem 3.6. *Let $\rho : \Gamma \hookrightarrow \mathrm{GL}(V)$ be an injective group homomorphism and $C \subset V$ be a cone. Suppose that C is Γ -invariant. If C admits a polyhedral fundamental domain under the action of Γ , then Γ is finitely presented.*

For a possibly degenerate open convex cone C , let $W \subset \bar{C}$ be the maximal \mathbb{R} -linear vector space. We say that W is defined over \mathbb{Q} if $W = W(\mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R}$ where $W(\mathbb{Q}) = W \cap V(\mathbb{Q})$. In this case, $V/W = (V(\mathbb{Q})/W(\mathbb{Q})) \otimes_{\mathbb{Q}} \mathbb{R}$ has a natural lattice structure, and we denote everything in V/W by $\widetilde{(-)}$. For example, $\widetilde{(C_+)}$ is the image

of C_+ under the projection $p : V \rightarrow V/W$. By the maximality, W is Γ -invariant, and thus $V/W, \tilde{C}$ admit natural Γ -actions.

Lemma 3.7. *Under the above notation and assumptions,*

- (1) $\tilde{\tilde{C}} = \tilde{C}$,
- (2) $(\tilde{C})_+ = \widetilde{(\tilde{C}_+)}$, which is denoted by \tilde{C}_+ , and
- (3) if (C_+, Γ) is of polyhedral type, then (\tilde{C}_+, Γ) is still of polyhedral type. More precisely, if $\Pi \subset C_+$ is a polyhedral cone with $\Gamma \cdot \Pi \supset C$, then $\tilde{\Pi} \subset \tilde{C}_+$ and $\Gamma \cdot \tilde{\Pi} \supset \tilde{C}$.

Proof. For (1), $\tilde{\tilde{C}} \supset \tilde{C}$ trivially holds. The converse does not hold for an arbitrary linear projection (see [Loo14, Remark 2.5]). In our case, let $j : V/W \rightarrow V$ be a spitting of $p : V \rightarrow V/W$. For an $x \in \tilde{\tilde{C}}$, let $\tilde{a}_n \rightarrow x$ with $a_n \in C$. Then $a_n - j(\tilde{a}_n) \in W$ as $p(a_n - j(\tilde{a}_n)) = 0$. By $W \subset \tilde{C}$, $j(\tilde{a}_n) \in \tilde{C}$. Moreover, as $\{\tilde{a}_n\}_{n \in \mathbb{N}}$ converges and j is continuous, $\{j(\tilde{a}_n)\}_{n \in \mathbb{N}}$ converges to $\alpha \in \tilde{C}$. Thus $\tilde{a}_n = p(j(\tilde{a}_n)) \rightarrow p(\alpha)$. Hence $x = p(\alpha) \in \tilde{C}$.

For (2), we first show

$$\tilde{\tilde{C}} \cap (V/W)(\mathbb{Q}) = (\tilde{C} \cap \widetilde{V(\mathbb{Q})}).$$

The “ \supset ” follows from definition. For the converse, let $\tilde{a} \in \tilde{\tilde{C}}$ be a rational point in V/W . Then by $W \subset \tilde{C}$, we can assume that $a \in \tilde{C}$ is a rational point in V . This gives “ \subset ”. Next, we show

$$\text{Conv} \left(\tilde{C} \cap \widetilde{V(\mathbb{Q})} \right) = \widetilde{\text{Conv}(\tilde{C} \cap V(\mathbb{Q}))}.$$

As the image of a convex set is still convex, we have “ \subset ”.

For a set $S \subset V$, we have

$$\text{Conv}(S) = \left\{ \sum_{i \in I} \lambda_i s_i \mid \sum_{i \in I} \lambda_i = 1, \lambda_i > 0, |I| < \infty, s_i \in S \right\}.$$

For an $a \in \text{Conv}(\tilde{C} \cap V(\mathbb{Q}))$, take finitely many $s_i \in \tilde{C} \cap V(\mathbb{Q})$, and $\lambda_i > 0, \sum_{i \in I} \lambda_i = 1$, so that $a = \sum \lambda_i s_i$. Thus $\tilde{a} = \sum \lambda_i \tilde{s}_i \in \text{Conv} \left(\tilde{C} \cap \widetilde{V(\mathbb{Q})} \right)$. This shows the converse inclusion.

Finally, by (1),

$$(\tilde{C})_+ = \text{Conv}(\tilde{\tilde{C}} \cap (V/W)(\mathbb{Q})) = \text{Conv}(\tilde{C} \cap (V/W)(\mathbb{Q})).$$

Then (2) follows from

$$\text{Conv}(\tilde{\tilde{C}} \cap (V/W)(\mathbb{Q})) = \text{Conv} \left(\tilde{C} \cap \widetilde{V(\mathbb{Q})} \right) = \widetilde{\text{Conv}(\tilde{C} \cap V(\mathbb{Q}))} = \widetilde{(\tilde{C}_+)}$$

For (3), let $\Pi \subset C_+$ be a polyhedral cone such that $\Gamma \cdot \Pi \supset C$. By (2), $\tilde{\Pi} \subset \tilde{C}_+$. Moreover, $\Gamma \cdot \tilde{\Pi} \supset \tilde{C}$. \square

Proposition 3.8. *Let (C_+, Γ) be of polyhedral type. Let $W \subset \tilde{C}$ be the maximal vector space. Suppose that W is defined over \mathbb{Q} . Then there is a rational polyhedral cone $\Pi \subset C_+$ such that $\Gamma \cdot \Pi = C_+$, and for each $\gamma \in \Gamma$, either $\gamma\Pi \cap \text{Int}(\Pi) = \emptyset$ or $\gamma\Pi = \Pi$. Moreover,*

$$\{\gamma \in \Gamma \mid \gamma\Pi = \Pi\} = \{\gamma \in \Gamma \mid \gamma \text{ acts trivially on } V/W\}.$$

Proof. By Lemma 3.7 (3), (\tilde{C}_+, Γ) is still of polyhedral type. By Lemma 3.5, there is a rational polyhedral cone $\tilde{\Pi}$ as a fundamental domain of \tilde{C}_+ under the action of $\tilde{\rho}(\Gamma)$, where $\tilde{\rho} : \Gamma \rightarrow \text{GL}(V/W)$ is the natural group homomorphism. By Lemma 3.7 (2), let $\Pi' \subset C_+$ be a rational polyhedral cone such that $p(\Pi') = \tilde{\Pi}$, where $p : V \rightarrow V/W$. Let $\Pi := \Pi' + W$ which is a rational polyhedral cone. As $\gamma(\Pi' + W) = (\gamma\Pi') + W$, by Lemma 3.7 (2), we have $\Gamma \cdot \Pi = C_+$.

If $\gamma\tilde{\Pi} \cap \text{Int}(\tilde{\Pi}) = \emptyset$, then $\gamma\Pi \cap \text{Int}(\Pi) = \emptyset$ as $\text{Int}(\Pi)$ maps to $\text{Int}(\tilde{\Pi})$. If $\gamma\tilde{\Pi} = \tilde{\Pi}$, then we claim that $\gamma\Pi = \Pi$. In fact, for some $a \in \Pi'$, we have $(\gamma \cdot a) = \gamma \cdot \tilde{a} \in \tilde{\Pi}$ and thus $\gamma \cdot a = b + w$ for some $b \in \Pi', w \in W$. Thus $\gamma\Pi \subset \Pi$. Similarly, $\gamma^{-1}\Pi \subset \Pi$. This shows the claim. Moreover, $\gamma\Pi = \Pi$ iff γ acts trivially on $\tilde{\Pi}$ iff γ acts trivially on V/W because $\tilde{\Pi}$ is a fundamental domain under the action of $\tilde{\rho}(\Gamma)$. \square

4. GENERIC PROPERTIES OF FIBRATIONS AND STRUCTURES OF CONES

Let $(X, \Delta) \rightarrow S$ be a fiber space. Let $K := K(S)$ be the field of rational functions on S and \bar{K} be the algebraic closure of K . Then $X_{\bar{K}} := X \times_S \text{Spec } \bar{K}$ is the geometric fiber of f . Set $\Delta_{\bar{K}} := \Delta \times_S \text{Spec } \bar{K}$.

Proposition 4.1. *Let $f : X \rightarrow S$ be a fibration.*

- (1) *If (X, Δ) has klt singularities, then $(X_{\bar{K}}, \Delta_{\bar{K}})$ still has klt singularities. Moreover, if $f : (X, \Delta) \rightarrow S$ is a klt Calabi-Yau fiber space, then $(X_{\bar{K}}, \Delta_{\bar{K}})$ is a klt Calabi-Yau pair over $\text{Spec } \bar{K}$.*
- (2) *For a finite base change $h : T \rightarrow S$ between varieties, let $U \subset S$ be a non-empty open set and $V = h^{-1}(U)$. Then we can shrink U such that $X_V := X \times_U V$ satisfies the following properties.*

If (X, Δ) has klt singularities, then (X_V, Δ_V) still has klt singularities, where $\Delta_V := \Delta \times_T V$. Moreover, if $f : (X, \Delta) \rightarrow S$ is a klt Calabi-Yau fiber space, then (X_V, Δ_V) has klt singularities and $K_{X_V} + \Delta_V \sim_{\mathbb{R}} 0/V$.

Proof. For (1), we first show that $X_{\bar{K}}$ is normal. This is a local statement for both source and target, hence we can assume that $f : \text{Spec } A \rightarrow \text{Spec } B$. The collection of affine open sets $\{\text{Spec } B_i \subset \text{Spec } B \mid i\}$ forms a direct system such that $\varinjlim B_i = K$. Then $A \otimes_B \varinjlim B_i = \varinjlim (A \otimes_B B_i)$. As $A \otimes_B B_i$ is normal, by [Sta22, Lemma 037D], $A \otimes_B K$ is also normal. Then $X_{\bar{K}} = X_K \otimes_K \bar{K}$ is normal by [Sta22, Lemma 0C3M]. Let $v : \text{Spec } \bar{K} \rightarrow S$, $u : X_{\bar{K}} \rightarrow X$ and $\bar{f} : X_{\bar{K}} \rightarrow \text{Spec } \bar{K}$ be natural morphisms. Then $\bar{f}_*(u^*\mathcal{O}_X) = v^*(f_*\mathcal{O}_X)$. By $u^*\mathcal{O}_X = \mathcal{O}_{X_{\bar{K}}}$ and $f_*\mathcal{O}_X = \mathcal{O}_S$, we see that $X_{\bar{K}}$ is connected. Hence $X_{\bar{K}}$ is an irreducible normal variety over \bar{K} .

Next, we show that $X_{\bar{K}}$ has klt singularities. Let X_{reg} be the smooth part of X . Shrinking S , we can assume that S is smooth and $X_{\text{reg}} \rightarrow S$ is a smooth morphism. Then the sequence

$$0 \rightarrow f^*\Omega_S \rightarrow \Omega_{X_{\text{reg}}} \rightarrow \Omega_{X_{\text{reg}}/S} \rightarrow 0$$

is exact. Let $r = \dim(X/S)$. By

$$(\Omega_{X_{\text{reg}}/S})_{\bar{K}} = \Omega_{(X_{\text{reg}})_{\bar{K}}} \text{ and } \mathcal{O}_X(K_{X_{\text{reg}}/S}) = \wedge^r \Omega_{X_{\text{reg}}/S},$$

we have

$$(K_{X_{\text{reg}}/S})_{\bar{K}} \sim K_{(X_{\text{reg}})_{\bar{K}}}.$$

By $\text{codim}(X \setminus X_{\text{reg}}) \geq 2$, we have

$$(K_{X/S})_{\bar{K}} - K_{X_{\bar{K}}} \sim 0.$$

Take a log resolution $g : \tilde{X} \rightarrow X$, then

$$K_{\tilde{X}/S} + \tilde{\Delta} = g^*(K_{X/S} + \Delta)$$

with coefficients of $\tilde{\Delta} < -1$. The natural morphism $\bar{g} : \tilde{X}_{\bar{K}} \rightarrow X_{\bar{K}}$ is also a log resolution and the above argument implies that

$$K_{\tilde{X}_{\bar{K}}} + \tilde{\Delta}_{\bar{K}} = \bar{g}^*(K_{X_{\bar{K}}} + \Delta_{\bar{K}}).$$

As coefficients of $\tilde{\Delta}_{\bar{K}} < -1$, $(X_{\bar{K}}, \Delta_{\bar{K}})$ has klt singularities.

When $f : (X, \Delta) \rightarrow S$ is a klt Calabi-Yau fiber space. We only need to note that $K_X + \Delta \sim_{\mathbb{R}} 0/S$ implies that $K_{X_{\bar{K}}} + \Delta_{\bar{K}} \sim_{\mathbb{R}} 0$.

For (2), shrinking U , we can assume that $V \rightarrow S$ is étale. We first show that X_V is normal. Note that $\phi : X_V \rightarrow X$ is also étale. Let $x \in X_V$ be a point (not necessarily a closed point) and $y = \phi(x)$. Set $\mathcal{O}_x := \mathcal{O}_{X_V, x}$ (resp. $\mathcal{O}_y := \mathcal{O}_{X_U, y}$). Let $\hat{\mathcal{O}}_x$ (resp. $\hat{\mathcal{O}}_y$) be the completion with respect to the maximal ideal. By [Har77, III, Exercise 10.4],

$$\hat{\mathcal{O}}_y \otimes_{k(y)} k(x) \simeq \hat{\mathcal{O}}_x,$$

where $k(y) \subset \hat{\mathcal{O}}_y$ and $k(x) \subset \hat{\mathcal{O}}_x$ are fields of representatives. Note that $k(y)$ and $k(x)$ are of characteristic zero. We claim that $\hat{\mathcal{O}}_x$ is normal. In fact, as X has klt singularities, X is Cohen-Macaulay. Hence $\hat{\mathcal{O}}_y$ is Cohen-Macaulay by [Sta22, Lemma 07NX]. In particular, it satisfies Serre's condition S_2 . As $\hat{\mathcal{O}}_y$ is certainly regular in codimension 1, it is normal. Then [Sta22, Lemma 0C3M] shows that $\hat{\mathcal{O}}_y \otimes_{k(y)} k(x)$ is normal. Thus \mathcal{O}_x is normal by [Sta22, Lemma 0FIZ]. This shows that X_V is normal.

Let $f_V : X_V \rightarrow V$ be the natural map. By $V \rightarrow U$ flat and $f_*\mathcal{O}_X = \mathcal{O}_S$, $(f_V)_*\mathcal{O}_{X_V} = \mathcal{O}_V$, and thus $H^0(X_V, \mathcal{O}_{X_V}) = H^0(V, \mathcal{O}_V)$ is an integral domain. This shows that X_V is an irreducible normal variety.

Let $\pi : W \rightarrow X$ be a log resolution of (X, Δ) with natural morphisms $\pi_V : W_V \rightarrow X_V$ and $\tilde{\phi} : W_V \rightarrow W$. Set $K_W + \Delta_W := \pi^*(K_X + \Delta)$ and $K_{W_V} + \Delta_{W_V} := \pi_V^*(K_{X_V} + \Delta_V)$. As ϕ is étale, we have

$$K_{X_V} + \Delta_V = \phi^*(K_X + \Delta).$$

Therefore, $K_{W_V} + \Delta_{W_V} = \tilde{\phi}^*(K_W + \Delta_W)$. By (X, Δ) klt, coefficients of Δ_W are < -1 . As $\tilde{\phi}$ is étale, $\Delta_{W_V} = \tilde{\phi}^*\Delta_W$, and the coefficients of Δ_{W_V} are < -1 . As $W_V \rightarrow X_V$ is a log resolution of (X_V, Δ_V) , (X_V, Δ_V) is still klt.

When $f : (X, \Delta) \rightarrow S$ is a klt Calabi-Yau fiber space, then $K_X + \Delta \sim_{\mathbb{R}} 0/S$ implies that $K_{X_V} + \Delta_V = \phi^*(K_X + \Delta) \sim_{\mathbb{R}} 0/V$. \square

Remark 4.2. *Even if X is \mathbb{Q} -factorial, $X_{\bar{K}}$ and X_V may not be \mathbb{Q} -factorial.*

Proposition 4.3. *Let $f : X \rightarrow S$ be a fibration. For any sufficiently small open set $U \subset S$, there exists a natural inclusion*

$$N^1(X_U/U)_{\mathbb{R}} \hookrightarrow N^1(X_{\bar{K}})_{\mathbb{R}}, \quad [D] \mapsto [D_{\bar{K}}].$$

Proof. First, we show that there is a natural map

$$N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X_{\bar{K}})_{\mathbb{R}}, \quad [D] \mapsto [D_{\bar{K}}].$$

Because $\{D \in \text{Pic}(X/S)_{\mathbb{R}} \mid D \equiv 0/S\}$ is defined over \mathbb{Q} , we only need to show that if a Cartier divisor $D \equiv 0/S$, then $D_{\bar{K}} \equiv 0$. Replacing X by a resolution $\tilde{X} \rightarrow X$ and $D, D_{\bar{K}}$ by their pullbacks on $\tilde{X}, \tilde{X}_{\bar{K}}$ respectively, we can assume that X is smooth.

Let $C_{\bar{K}} \rightarrow X_{\bar{K}}$ be a smooth curve, we will show $D_{\bar{K}} \cdot C_{\bar{K}} = 0$. By definition, this is to show that the coefficient of m in the polynomial $\chi(C_{\bar{K}}, mD_{\bar{K}})$ is 0. Let C be a spreading out of $C_{\bar{K}}$ over a variety T such that $h : T \rightarrow S$ is a finite morphism (see [Poo17, Chapter 3.2]). We

can assume that C is smooth over T . Shrinking S , we can assume that S is smooth and h is étale. By Proposition 4.1 (2), we can assume that X_T is normal. Shrinking T further, we may assume that $T = \text{Spec } A$ is affine. Moreover, as $\text{Spec } \bar{K} \rightarrow T$ is flat, [Har77, III Prop 9.3] implies that

$$H^i(C, mD_T) \otimes_A \bar{K} \simeq H^i(C_{\bar{K}}, mD_{\bar{K}}).$$

Thus

$$\chi(C_{\bar{K}}, mD_{\bar{K}}) = \sum (-1)^k \dim_{\bar{K}} H^i(C, mD_T) \otimes_A \bar{K}.$$

Shrinking T , by [Har77, III Prop 12.9], we have

$$H^i(C, mD_T) \otimes_A k(t) \simeq H^i(C_t, mD_t),$$

where $t \in T$ is a closed point. [Har77, III Prop 12.9] also implies that $H^i(C, mD_T)$ is a free A -module. Thus

$$\begin{aligned} \dim H^i(C, mD_T) \otimes_A \bar{K} &= \dim H^i(C, mD_T) \otimes_A k(t) \\ &= \dim H^i(C_t, mD_t). \end{aligned}$$

Let $\phi : X_T \rightarrow X$ be the natural morphism. Then $D_T \cdot C_t = \phi^* D \cdot C_t = D \cdot \phi_* C_t = 0$. Therefore, the coefficient of m in

$$\chi(C_{\bar{K}}, mD_{\bar{K}}) = \chi(C_t, mD_t)$$

is 0. This shows that $D_{\bar{K}} \equiv 0$.

Next, to get the desired inclusion for any sufficiently small open set, it suffices to find one such open set.

Suppose that $D_{\bar{K}} \equiv 0$, we want to find U such that $D \equiv 0/U$ (this U may depend on D). Let $\tilde{X} \rightarrow X$ be a resolution, and \tilde{D} be the pullback of D . We have $\tilde{D}_{\bar{K}} \equiv 0$ on $\tilde{X}_{\bar{K}}$. If $\tilde{D} \equiv 0/U$, then $D \equiv 0/U$. Therefore, we can assume that X is smooth.

By [Kle05, Theorem 9.6.3 (a) and (b)], there exists an $m \in \mathbb{Z}_{>0}$ such that $mD_{\bar{K}}$ is algebraically equivalent to $\mathcal{O}_{X_{\bar{K}}}$. That is, there exist connected \bar{K} -schemes of finite type $\bar{B}_i, 1 \leq i \leq n$, invertible sheaves \bar{M}_i on $X_{\bar{B}_i}$ and closed points s_i, t_i of \bar{B}_i such that

$$\begin{aligned} \mathcal{O}(mD_{\bar{K}})_{s_1} &\simeq \bar{M}_{1,s_1}, \bar{M}_{1,t_1} \simeq \bar{M}_{2,s_2}, \dots \\ &\dots, \bar{M}_{n-1,t_{n-1}} \simeq \bar{M}_{n,s_n}, \bar{M}_{n,t_n} \simeq \mathcal{O}_{X_{\bar{K}},t_n} \end{aligned}$$

(see [Kle05, Definition 9.5.9]). Moreover, connecting s_i, t_i by the image of a smooth curve, we can further assume that \bar{B}_i is a smooth curve. All the above schemes, sheaves, and isomorphisms are defined in a finite extension of K .

By generic smoothness, shrinking S to U , we can assume that $X_U \rightarrow U$ is smooth. There is a finite morphism $T \rightarrow U$ such that every

objects and relations mentioned above over \bar{K} is defined on $X_T \rightarrow T$. In particular, there are dominant morphisms of finite type $B_i \rightarrow T$, $1 \leq i \leq n$ with sections $\tilde{s}_i, \tilde{t}_i : T \rightarrow B_i$, and invertible sheaves M_i on X_{B_i} such that

$$\begin{aligned} \mathcal{O}(mD)_{\tilde{s}_1(T)} &\simeq M_{1,\tilde{s}_1(T)}, M_{1,\tilde{t}_1(T)} \simeq M_{2,\tilde{s}_2(T)}, \cdots \\ &\cdots, M_{n-1,\tilde{t}_{n-1}(T)} \simeq M_{n,\tilde{s}_n(T)}, M_{n,\tilde{t}_n(T)} \simeq \mathcal{O}_{\tilde{t}_n(T)}, \end{aligned}$$

where $\mathcal{O}(mD)_{\tilde{s}_1(T)} = \tau_1^* \mathcal{O}(mD)|_{\theta_1^{-1}(\tilde{s}_1(T))}$ with $\tau_1 : X_{B_1} \rightarrow X_T \rightarrow X_U$, $\theta_1 : X_{B_1} \rightarrow B_1$ (other sheaves are defined similarly). Note that $X_T \rightarrow T$ is isomorphic to both $\theta_i^{-1}(\tilde{s}_i(T)) \rightarrow \tilde{s}_i(T)$ and $\theta_i^{-1}(\tilde{t}_i(T)) \rightarrow \tilde{t}_i(T)$, where $\theta_i : X_{B_i} \rightarrow B_i$. Shrinking U, T further, we can assume that each B_i is also smooth.

We will show that $mg^*D \equiv 0/T$, where $g : X_T \rightarrow X_U$. Because the intersection is taken in the singular cohomology groups, this can be checked in the analytic topology. First, as B_i is smooth, shrinking T (hence also U), we can assume that $B_i \rightarrow T$ is smooth. As $X_T \rightarrow T$ is a smooth morphism, $X_{B_i} \rightarrow B_i$ is also a smooth morphism between smooth varieties. Thus $X_{B_i} \rightarrow B_i$ is locally trivial in the analytic topology by Ehresmann's theorem. Let $\ell \subset \theta_1^{-1}(\tilde{s}_1(T))$ be a curve which maps to a point on $\tilde{s}_1(T)$. Let $\ell' \subset \theta_1^{-1}(\tilde{t}_1(T))$ be a manifold which is a deformation of ℓ in the analytic topology (we do not need ℓ' to be an algebraic curve). By induction on i , it is enough to show

$$M_{1,\tilde{s}_1(T)} \cdot \ell = M_{1,\tilde{t}_1(T)} \cdot \ell'.$$

As $M_{1,\tilde{s}_1(T)} \cdot \ell = M_1 \cdot \ell$ and $M_{1,\tilde{t}_1(T)} \cdot \ell' = M_1 \cdot \ell'$, the desired result follows. Hence $D \equiv 0/U$.

To obtain an open set U which is independent of divisors, we can use one of the following two approaches:

(A) By $\dim N^1(X/S)_{\mathbb{R}} < \infty$, we have

$$\text{Ker}(N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X_{\bar{K}})_{\mathbb{R}}) < \infty.$$

Let $[D_1], \dots, [D_e]$ be a basis of $\text{Ker}(N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X_{\bar{K}})_{\mathbb{R}})$. By the above construction, there exists an open set U_i such that $D_i \equiv 0/U_i$. Then $U := \bigcap_{i=1}^e U_i$ satisfies the desired property.

(B) Replacing X by a resolution, it is enough to show the claim for smooth X . Shrinking U , we can assume that $X_U \rightarrow U$ is smooth. We show that U satisfies the desired property. Let D be any divisor such that $D_{\bar{K}} \equiv 0$. By the above construction, there exists an open set $V \subset U$ such that $D_V \equiv 0/V$. We claim that $D \equiv 0/U$. It is enough to show that for any curve ℓ such that ℓ maps to a point in $U - V$, we have $D \cdot \ell = 0$. By Ehresmann's theorem, ℓ can be deformed to

a complex manifold ℓ' in the analytic topology such that ℓ' maps to a point in V under f . Thus $D \cdot \ell' = D \cdot \ell$. By the dual form of the Lefschetz theorem on $(1, 1)$ -classes, there exists an algebraic curve ℓ'' such that ℓ'' maps to a point in V under f and $D \cdot \ell' = D \cdot \ell''$. Therefore, $D \cdot \ell = D \cdot \ell'' = 0$. \square

Proposition 4.4. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assume that $R^1 f_* \mathcal{O}_X$ is a torsion sheaf. Then the following results hold true.*

(1) *There is a natural surjective linear map*

$$r : N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X_U/U)_{\mathbb{R}} \quad [D] \mapsto [D|_U].$$

When U is sufficiently small, we have

$$(4.0.1) \quad \text{Ker}(r) = \text{Span}_{\mathbb{R}}\{[D] \mid \text{Supp } D \subset \text{Supp}(X - X_U)\}.$$

- (2) *The maximal vector space $W \subset \overline{\text{Eff}}(X/S)$ is generated by divisors in $\text{Ker}(r)$. In particular, W is defined over \mathbb{Q} .*
 (3) *The maximal vector space of $\overline{\text{Mov}}(X/S)$ is defined over \mathbb{Q} .*
 (4) *If S is \mathbb{Q} -factorial, then $\overline{\text{Mov}}(X/S)$ is non-degenerate.*

Proof. (1) Note that $r([D]) = [D|_U]$ is well-defined. If D_U is a divisor on X_U such that $D_U = \sum c_i B_i$ is the decomposition into irreducible components, then $\overline{D}_U := \sum c_i \overline{B}_i$ is a divisor on X such that $(\overline{D}_U)|_U = D_U$. Hence r is surjective.

Let $\bar{f} : X_{\bar{K}} \rightarrow \text{Spec } \bar{K}$. As $\text{Spec } \bar{K} \rightarrow S$ is flat, $R^1 \bar{f}_* \mathcal{O}_{X_{\bar{K}}} = (R^1 f_* \mathcal{O}_X)_{\bar{K}} = 0$. Thus $N^1(X_{\bar{K}})_{\mathbb{Q}} \simeq \text{Pic}(X_{\bar{K}})_{\mathbb{Q}}$. As r is defined over \mathbb{Q} , $\text{Ker}(r)$ is also defined over \mathbb{Q} . It is enough to show (4.0.1) for Cartier divisors. Take D to be a Cartier divisor such that $[D] \in \text{Ker}(r)$. Shrinking S to U as in Proposition 4.3, then by Proposition 4.3, we have $D_{\bar{K}} \equiv 0$. Possibly replacing D by a multiple, we can assume $D_{\bar{K}} \sim 0$. Thus $D_{\bar{K}} = \text{div}(\bar{\alpha})$ for some $\bar{\alpha} \in K(X_{\bar{K}})$. Shrinking U further, we can assume that there is a finite Galois morphism $T \rightarrow U$ such that the above relation is defined on X_T/T . In particular, $D_T := D|_T = \text{div}(\alpha)$ for some $\alpha \in K(X_T)$. As D_T is $\text{Gal}(X_T/X_U)$ -invariant, we have

$$mD_T = \text{div}(\tau) \text{ with } \tau := \prod_{\theta \in \text{Gal}(X_T/X_U)} \theta(\alpha),$$

where $m = |\text{Gal}(X_T/X_U)|$. As τ is $\text{Gal}(X_T/X_U)$ -invariant, there exists a $\beta \in K(X)$ whose pullback is τ under the morphism $X_T \rightarrow X_U$. Thus $mD_U = \text{div}(\beta)$ on X_U . Therefore,

$$\text{Supp}(mD - \text{div}(\beta)) \subset X - X_U.$$

This shows “ \subset ” in (4.0.1). The converse inclusion is trivial.

(2) For any $[D] \in \overline{\text{Eff}}(X/S)$, we know $r([D]) \in \overline{\text{Eff}}(X_U/U)$. We claim that if $r([D]) \neq 0$, then $[D] \notin W$. Otherwise, $r([D]) \neq 0$ implies that $[D_{\bar{K}}] \neq 0$ by Proposition 4.3. If $[D] \in W$, then $[D_{\bar{K}}], -[D_{\bar{K}}] \in \overline{\text{Eff}}(X_{\bar{K}})$. Hence $\overline{\text{Eff}}(X_{\bar{K}})$ is degenerate. This is a contradiction as $X_{\bar{K}}$ is projective. Therefore, $[D] \in W$ implies that $[D] \in \text{Ker}(r)$.

Conversely, let D be an \mathbb{R} -Cartier divisor such that

$$\text{Supp } D \subset \text{Supp}(X - X_U).$$

Then $f(\text{Supp } D) \neq S$. There is an ample divisor $H > 0$ on S such that $f(\text{Supp } D) \subset \text{Supp } H$. Thus $D + kf^*H > 0$ for a $k \gg 1$ and $[D + kf^*H] = [D] \in \text{Eff}(X/S)$.

(3) Let $W' \subset \overline{\text{Mov}}(X/S)$ be the maximal vector space. By (2), $W' \subset W$. In particular, if $[D] \in W'$, then D can be assumed to be an effective divisor. To show that W' is defined over \mathbb{Q} , it is enough to show that for any $[D] \in W'$, there are finitely many rational divisors $[D_j] \in \overline{\text{Mov}}(X/S), j \in I$ such that $[D] \in \text{Conv}([D_j] \mid j \in I)$. In fact, let $\{L_k = 0\} \subset N^1(X/S)_{\mathbb{R}}, k \in T$ be supporting hyperplanes of $\overline{\text{Mov}}(X/S)$ and assume that $\overline{\text{Mov}}(X/S) = \bigcap_{k \in T} \{L_k \geq 0\}$. Then $W' = \bigcap_{k \in T} \{L_k = 0\}$. If $[D] \in \text{Conv}([D_j] \mid j \in I)$, then $L_k([D_j]) = 0$ for all $k \in T$. Thus $[D_j] \in W'$ and $W' = (W' \cap V(\mathbb{Q})) \otimes \mathbb{R}$.

By the same argument as in the first paragraph of Theorem 2.6, there is a \mathbb{Q} -Cartier divisor $\tilde{\Delta}$ such that $(X, \tilde{\Delta}) \rightarrow S$ is a klt Calabi-Yau fiber space. Replacing Δ by $\tilde{\Delta}$, we can assume that Δ is \mathbb{Q} -Cartier.

Now let $\Delta + D = \sum a_i P_i$ with $a_i \in \mathbb{Q}_{>0}$. Suppose that $(X, \Delta + \sum \delta P_i)$ is klt for some $\delta \in \mathbb{Q}_{>0}$. Let $\tilde{P} = \bigoplus_i [0, \delta] P_i$ be the cube generated by P_j and

$$P := \tilde{P} + \Delta$$

be the transform of \tilde{P} by Δ . Then P is a rational polytope. Replacing D by ϵD for some $\epsilon \in \mathbb{Q}_{>0}$, we can assume that $\Delta + D \in P$. We claim that each $B = \Xi + \Delta \in P$ satisfies the weaker assumption in Remark 2.5. Note that Ξ is vertical over S by the construction of \tilde{P} .

Under the notation of Remark 2.5 and let $q'(F') = \tilde{F}$ which is also a general fiber of $\pi : X' \rightarrow Z'$. Let $q'|_{F'} : F' \rightarrow \tilde{F}$ be the restriction of q' to F' . Then

$$K_{F'} + \tilde{B}'|_{F'} = (K_{W'} + \tilde{B}')|_{F'} = (q'^*(K_{X'} + \Delta' + \Xi') + E')|_{F'},$$

where $E' \geq 0$ is q' -exceptional and Δ', Ξ' are strict transforms of Δ, Ξ on W' respectively. As Ξ' is also vertical over Z' and $K_{X'} + \Delta' \sim_{\mathbb{R}} 0/S$, we have

$$K_{F'} + \tilde{B}'|_{F'} \sim_{\mathbb{R}} E'|_{F'}.$$

As E' is exceptional over X' , $E'|_{F'}$ is exceptional over \tilde{F} . In particular, the negative part of the Nakayama-Zariski decomposition $N_\sigma(K_{F'} + \tilde{B}'|_{F'}) = E'|_{F'}$. Run a $(K_{F'} + \tilde{B}'|_{F'})$ -LMMP with scaling of an ample divisor. It is well-known that $N_\sigma(K_{F'} + \tilde{B}'|_{F'})$ will be contracted in finite steps. Thus, this LMMP terminates with F'' such that $K_{F''} + \tilde{B}'|_{F''} \sim_{\mathbb{R}} 0$. Therefore, $(F', \tilde{B}'|_{F'})$ has a good minimal model. This shows the claim.

By Theorem 2.4 and Remark 2.5, there is an open rational polytope $Q^\circ \subset P$ such that $\Delta + D \in Q^\circ$, and if $(Y/S, \Delta_Y + D_Y)$ is a weak log canonical model of $(X/S, \Delta + D)$, then for any $\Delta + B \in Q^\circ$, $(Y/S, \Delta_Y + B_Y)$ is also a weak log canonical model of $(X/S, \Delta + B)$. Note that the existence of such weak log canonical models is guaranteed by Theorem 2.3. As $D \in \overline{\text{Mov}}(X/S)$, by Lemma 2.2, we can assume that X, Y are isomorphic in codimension 1. As B is the strict transform of B_Y and $[B_Y] \in \overline{\text{Mov}}(Y/S)$, we still have $[B] \in \overline{\text{Mov}}(X/S)$. This shows $[\Theta] \in \overline{\text{Mov}}(X/S)$ for any $\Theta \in \bar{Q}^\circ$.

(4) Assume that S is \mathbb{Q} -factorial. We claim that if $0 \neq [D] \in W'$, then there exists a family of curves $\{C_t \mid t \in R\}$ which covers a divisor such that $[C_t] \in N_1(X/S)$ and $D \cdot C_t \neq 0$. By (2), we can assume that D is vertical over S . Replacing D by $D - f^*B$ for some \mathbb{R} -Cartier divisor B on S and shrinking S , we can assume that D is a very exceptional divisor. Write $D = D^+ - D^-$ such that $D^+, D^- \geq 0$ do not have common components. If $D^+ \neq 0$ (resp. $D^- \neq 0$), then by the standard reduce-to-surface argument (for example, see [Bir12, Lemma 3.3]), there exists a family of curves $\{C_t \mid t \in R\}$ covering an irreducible component of $\text{Supp } D^+$ (resp. $\text{Supp } D^-$) such that $[C_t] \in N_1(X/S)$ and $D^+ \cdot C_t < 0$ (resp. $D^- \cdot C_t < 0$). Thus $D \cdot C_t < 0$ (resp. $D \cdot C_t > 0$). This shows the claim.

Possibly replacing D by $-D \in W'$, we can assume that $D \cdot C_t < 0$. This contradicts with $D \in \overline{\text{Mov}}(X/S)$. \square

Question 4.5. *Do the claims in Proposition 4.4 still hold true for an arbitrary fibration $f : X \rightarrow S$?*

Recall that Γ_B is the image of $\text{PsAut}(X/S, \Delta)$ under the natural group homomorphism $\iota : \text{PsAut}(X/S, \Delta) \rightarrow \text{GL}(N^1(X/S)_{\mathbb{R}})$.

Lemma 4.6. *Let $f : X \rightarrow S$ be a Calabi-Yau fiber space such that X has terminal singularities. Assume that $R^1 f_* \mathcal{O}_X$ is a torsion sheaf. Let $W \subset \overline{\text{Mov}}(X/S)$ be the maximal vector space. Then*

$$\Gamma_W := \{\gamma \in \Gamma_B \mid \gamma \text{ acts trivially on } N^1(X/S)_{\mathbb{R}}/W\}$$

is a finite group.

Proof. As $R^1 f_* \mathcal{O}_X$ is a torsion sheaf, we have $H^1(X_{\bar{K}}, \mathcal{O}_{X_{\bar{K}}}) = 0$ and thus $\text{Pic}(\bar{X})_{\mathbb{Q}} \simeq N^1(X_{\bar{K}})_{\mathbb{Q}}$. Let $G := \{g \in \text{PsAut}(X/S) \mid \iota(g) \in \Gamma_W\}$. It suffices to show that G is a finite set. By Proposition 4.4 (1) and (2), there exists an open set $U \subset S$ such that $N^1(X/S)_{\mathbb{R}}/W \rightarrow N^1(X_U/U)_{\mathbb{R}}$ is surjective. Let H be an ample/ S divisor on X_U . Then $g \cdot H \equiv H/U$ for any $g \in G$. Thus $g_{\bar{K}} \cdot H_{\bar{K}} \equiv H_{\bar{K}}$ in $N^1(X_{\bar{K}})$, where $g_{\bar{K}}$ and $H_{\bar{K}}$ correspond to g and H respectively after the base change. Replacing H be a multiple, we can assume that $g_{\bar{K}} \cdot H_{\bar{K}} \sim H_{\bar{K}}$. The same argument as [Ogu14, Proposition 2.4] shows that $\{g_{\bar{K}} \mid g \in G\}$ is a finite set. Note that the smoothness assumption on $X_{\bar{K}}$ in [Ogu14, Proposition 2.4] can be relaxed to the assumption that $X_{\bar{K}}$ has terminal singularities by [Han87, (3.14) Theorem]. For $g, h \in G$, if $g_{\bar{K}} = h_{\bar{K}}$, then $g = h$. Thus G is also a finite set. \square

Remark 4.7. *The group Γ_W may not be trivial. There exists a sequence of flops which is also a sequence of birational automorphisms (see [Kaw97, Example 3.8 (4)]). Then for each $X \rightarrow S$, we have $N^1(X/S)_{\mathbb{R}} = \text{Mov}(X/S) = W = \mathbb{R}$. $\Gamma_W = \{\pm 1\}$ which acts trivially on $N^1(X/S)_{\mathbb{R}}/W$.*

Conjecture 4.8. *For a klt Calabi-Yau fibration $f : (X, \Delta) \rightarrow S$ such that $R^1 f_* \mathcal{O}_X$ is a torsion sheaf, Γ_W is always a finite group.*

5. A VARIANT OF THE CONE CONJECTURE

In this section, we study the relationship between the cone conjecture and Conjecture 1.2. Note that in Conjecture 1.2, by enlarging P_M and P_A , we can always assume that P_M and P_A are rational polyhedral cones. Recall that a polyhedral cone is closed by definition and Γ_B (resp. Γ_A) is the image of $\text{PsAut}(X/S, \Delta)$ (resp. $\text{Aut}(X/S, \Delta)$) under the group homomorphism $\text{PsAut}(X/S, \Delta) \rightarrow \text{GL}(N^1(X/S)_{\mathbb{R}})$. By Definition 3.1, we set

$$\begin{aligned} \text{Mov}(X/S)_+ &:= \text{Conv}(\overline{\text{Mov}}(X/S) \cap N^1(X/S)_{\mathbb{Q}}), \\ \text{Amp}(X/S)_+ &:= \text{Conv}(\overline{\text{Amp}}(X/S) \cap N^1(X/S)_{\mathbb{Q}}). \end{aligned}$$

Lemma 5.1. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space.*

- (1) *We have $\overline{\text{Amp}}^e(X/S) \subset \text{Amp}(X/S)_+$.*
- (2) *Assume the existence of good minimal models for effective klt pairs in $\dim(X/S)$, then $\overline{\text{Mov}}^e(X/S) \subset \text{Mov}(X/S)_+$*

Proof. For a $[D] \in \text{Eff}(X/S)$, replacing D by a divisor which is numerically equivalent to D , we can assume that the irreducible decomposition

of $D = \sum_{i=1}^k a_i D_i$ with $a_i > 0$. Let $P := \text{Cone}([D_i] \mid i = 1, \dots, k) \subset \text{Eff}(X/S)$ be a rational polyhedral cone.

For (1), assume that $[D] \in \overline{\text{Amp}}^e(X/S)$. By Theorem 2.7, $P_N = P \cap \overline{\text{Amp}}(X/S)$ is a rational polyhedral cone. Thus

$$[D] \in P_N \subset \text{Amp}(X/S)_+.$$

For (2), assume that $[D] \in \overline{\text{Mov}}^e(X/S)$. Let $P = \cup_{i=1}^m P_i^\circ$ be the union of finitely many open rational polyhedral cones satisfying the claim of Theorem 2.6. Suppose that $[D] \in P_i^\circ$. Let $[B] \in P_i^\circ$ such that $B \geq 0$. Choose $\epsilon \in \mathbb{Q}_{>0}$ such that both $(X, \Delta + \epsilon D)$ and $(X, \Delta + \epsilon B)$ are klt. By Theorem 2.3, let $(Y/S, \Delta_Y + \epsilon D_Y)$ be a weak log canonical model of $(X/S, \Delta + \epsilon D)$. As $[D] \in \overline{\text{Mov}}(X/S)$, we can assume that Y is \mathbb{Q} -factorial and X, Y are isomorphic in codimension 1 by Lemma 2.2. By Theorem 2.6, $(Y/S, \Delta_Y + \epsilon B_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon B)$. In particular, $[B_Y] \in \overline{\text{Mov}}(Y/S)$. Thus $[B] \in \overline{\text{Mov}}(X/S)$. As $[B] \in P_i^\circ$ is an arbitrary point, the above shows $P_i^\circ \subset \overline{\text{Mov}}(X/S)$. Because P_i° is a rational polyhedral cone, we have $[D] \in P_i^\circ \subset \text{Mov}(X/S)_+$. \square

Lemma 5.2. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assume the existence of good minimal models for effective klt pairs in $\dim(X/S)$. If there exists a rational polyhedral cone $P_M \subset \text{Eff}(X/S)$ satisfying Conjecture 1.2 (1), then there is a rational polyhedral cone $Q_M \subset \text{Mov}(X/S) \cap P_M$ such that*

$$(5.0.1) \quad \bigcup_{g \in \text{PsAut}(X/S, \Delta)} g \cdot Q_M = \text{Mov}(X/S).$$

Similarly, if there exists a rational polyhedral cone $P_A \subset \text{Eff}(X/S)$ satisfying Conjecture 1.2 (2), then there is a rational polyhedral cone $Q_A \subset \overline{\text{Amp}}^e(X/S) \cap P_A$ such that

$$(5.0.2) \quad \bigcup_{g \in \text{Aut}(X/S, \Delta)} g \cdot Q_A = \overline{\text{Amp}}^e(X/S).$$

Proof. For (5.0.1), by Theorem 2.6, $P_M = \cup_{i=1}^m P_i^\circ$ is a union of finitely many open rational polyhedral cones. Let $P_1^\circ, \dots, P_k^\circ$ be the polyhedral cones such that $P_j^\circ \cap \overline{\text{Mov}}(X/S) \neq \emptyset$.

We claim that $P_j := \overline{P_j^\circ} \subset \text{Mov}(X/S)$. Let $D \geq 0$ such that $[D] \in P_j^\circ \cap \overline{\text{Mov}}(X/S)$. Assume that $(Y/S, \Delta_Y + \epsilon D_Y)$ is a weak log canonical model of $(X/S, \Delta + \epsilon D)$ for some $\epsilon \in \mathbb{Q}_{>0}$. By Lemma 2.2, we can assume that X, Y are isomorphism in codimension 1. Take $[B'] \in P_j$, by $P_M \subset \text{Eff}(X/S)$, there exists a sequence $\{[B_l]\}_{l \in \mathbb{N}} \subset P_j^\circ$ with $B_l \geq 0$

such that $\lim[B_l] = [B']$ and $\lim B_l = B$ as the limit of Weil divisors. Thus $[B'] = [B]$. By Theorem 2.6, there exists a $\delta \in \mathbb{Q}_{>0}$ such that $(Y/S, \Delta_Y + \delta B_{l,Y})$ is a weak log canonical model of $(X/S, \Delta + \delta B_l)$ for each l . Thus $(Y/S, \Delta_Y + \delta B_Y)$ is also a weak log canonical model of $(X/S, \Delta + \delta B)$. By Theorem 2.3, B_Y is semi-ample/ S . Hence B is movable, and thus $[B] \in \text{Mov}(X/S)$.

Let $Q_M := \text{Cone}(P_1, \dots, P_k)$ be the cone generated by $P_j, 1 \leq j \leq k$. Then $Q_M \subset \text{Mov}(X/S)$. For an $[M] \in \text{Mov}(X/S)$, there exists a $g \in \text{PsAut}(X/S, \Delta)$ such that $g \cdot [M] \in P_M$. Thus $g \cdot [M] \in P_j^\circ$ for some j and hence $g \cdot [M] \in Q_M$. This shows (5.0.1).

For (5.0.2), Theorem 2.7 shows that $Q_A := P_A \cap \overline{\text{Amp}}(X/S)$ is a rational polyhedral cone. By $P_A \subset \text{Eff}(X/S)$, we have $Q_A \subset \overline{\text{Amp}}^e(X/S)$. For any $[H] \in \text{Amp}(X/S)$, there exist an $[H'] \in P_A$ and a $g \in \text{Aut}(X/S, \Delta)$ such that $g \cdot [H'] = [H]$. Hence $[H'] \in Q_A$. Thus $\Gamma_A \cdot Q_A \supset \text{Amp}(X/S)$. As $\overline{\text{Amp}}(X/S)$ is non-degenerate and $Q_A \subset \text{Amp}(X/S)_+$ by Lemma 5.1, Proposition 3.3 implies that $\Gamma_A \cdot Q_A = \text{Amp}(X/S)_+ \supset \overline{\text{Amp}}^e(X/S)$. The “ \subset ” of (5.0.2) follows from definition. \square

Proposition 5.3. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Let $W \subset \overline{\text{Mov}}(X/S)$ be the maximal vector space. Assume that good minimal models exist for effective klt pairs in dimension $\dim(X/S)$. Suppose that there is a polyhedral cone $P \subset \text{Mov}(X/S)$ such that*

$$\text{PsAut}(X/S, \Delta) \cdot P = \text{Mov}(X/S).$$

We have the following results.

- (1) *If either $R^1 f_* \mathcal{O}_X$ is a torsion sheaf or $W = 0$, then we have*

$$\text{Mov}(X/S) = \overline{\text{Mov}}^e(X/S) = \text{Mov}(X/S)_+.$$

- (2) *There are finitely many varieties $Y_j/S, j \in J$ such that if $X \dashrightarrow Y/S$ is isomorphic in codimension 1 with Y a \mathbb{Q} -factorial variety, then $Y \simeq Y_j/S$ for some $j \in J$.*
- (3) *If $\overline{\text{Mov}}(X/S)$ is non-degenerate, then $\overline{\text{Mov}}^e(X/S)$ has a rational polyhedral fundamental domain under the action of Γ_B .*
- (4) *If $R^1 f_* \mathcal{O}_X$ is a torsion sheaf, then $\overline{\text{Mov}}^e(X/S)$ has a weak rational polyhedral fundamental domain (may be degenerate) under the action of Γ_B .*

Proof. Possibly enlarging P , we can assume that P is a rational polyhedral cone.

For (1), we have “ \subset ” for the above three cones by Lemma 5.1. By Proposition 4.4 (2), W is defined over \mathbb{Q} . By definition, $\text{Mov}(X/S) \supset \text{Int}(\overline{\text{Mov}}(X/S))$. Then $\Gamma_B \cdot P \supset \text{Int}(\overline{\text{Mov}}(X/S))$. Thus $(\text{Mov}(X/S)_+, \Gamma_B)$

is of polyhedral type. We follow the notation of Lemma 3.7. By Lemma 3.7 (3) and Proposition 3.3, we have

$$(5.0.3) \quad \Gamma_B \cdot \tilde{P} = \widetilde{(\text{Mov}(X/S))_+} = (\text{Mov}(X/S)_+)^{\sim},$$

where the last equality follows from Lemma 3.7 (2).

We claim that $W \subset \text{Mov}(X/S)$. By Proposition 4.4 (1) and (2), $W \subset \text{Eff}(X/S)$. Let $[D] \in W$ be a rational point such that $D \geq 0$. Then for a sufficiently small $\epsilon \in \mathbb{Q}_{>0}$, $(X/S, \Delta + \epsilon D)$ has a weak log canonical model $(Y/S, \Delta + \epsilon D_Y)$. Because $[D] \in \overline{\text{Mov}}(X/S)$, by Lemma 2.2, we can assume that X, Y are isomorphic in codimension 1. Note that D_Y is semi-ample/ S by Theorem 2.3. Thus $[D] \in \text{Mov}(X/S)$. As W is Γ -invariant and $\Gamma_B \cdot (P + W) = \text{Mov}(X/S)_+$ by (5.0.3), we have $\text{Mov}(X/S) = \text{Mov}(X/S)_+$.

For (2), there exists a decomposition $P = \cup_{i=1}^k P_i^\circ$ as in Theorem 2.6. For each j , by Lemma 2.2 and Theorem 2.3, we can choose a $f_j : X \dashrightarrow Y_j/S$ which is isomorphic in codimension 1 such that if $[D] \in P_j^\circ$ with $D \geq 0$, then $(Y_j/S, \Delta_{Y_j} + \epsilon D_{Y_j})$ is a \mathbb{Q} -factorial weak log canonical model of $(X/S, \Delta + \epsilon D)$ for some $\epsilon \in \mathbb{Q}_{>0}$. We claim that if $g : Y \dashrightarrow X/S$ is isomorphic in codimension 1, then $Y \simeq Y_j/S$ for some j . In fact, let $A \geq 0$ be an ample/ S divisor on Y . Then $g_*A \in \text{Mov}(X/S)$. Let $\sigma \in \text{PsAut}(X/S, \Delta)$ such that $\sigma \cdot g_*A \in P$. Then $\sigma \cdot g_*A \in P_j^\circ$ for some j . Note that Y, Y_j are \mathbb{Q} -factorial varieties. Because $(\sigma \cdot g_*A)_{Y_j} = (f_j \circ \sigma \circ g)_*A$ is nef/ S and

$$f_j \circ \sigma \circ g : Y \dashrightarrow X \dashrightarrow X \dashrightarrow Y_j/S$$

is isomorphic in codimension 1, we have $Y \simeq Y_j/S$.

For (3) and (4), as $(\text{Mov}(X/S)_+, \Gamma_B)$ is of polyhedral type, by Proposition 3.8, there is a rational polyhedral cone Π such that $\Gamma_B \cdot \Pi = \text{Mov}(X/S)_+$, and for each $\gamma \in \Gamma_B$, either $\gamma\Pi \cap \text{Int}(\Pi) = \emptyset$ or $\gamma\Pi = \Pi$. Moreover,

$$\{\gamma \in \Gamma_B \mid \gamma\Pi = \Pi\} = \{\gamma \in \Gamma_B \mid \gamma \text{ acts trivially on } N^1(X/S)_{\mathbb{R}}/W\}.$$

Hence Π is a weak rational polyhedral fundamental domain. In particular, if $W = 0$, then Π is a rational polyhedral fundamental domain. \square

Remark 5.4. *The assumption in Proposition 5.3 (1) is necessary. [Kaw97, Example 3.8 (2)] gives an elliptic fibration (hence $R^1 f_* \mathcal{O}_X$ is not a torsion sheaf) with $W \neq 0$ such that $\text{Mov}(X/S) = \overline{\text{Mov}}^e(X/S) \neq \text{Mov}(X/S)_+$. In this example, W is defined over \mathbb{Q} but $W \not\subset \text{Mov}(X/S)$.*

Proposition 5.5. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Suppose that there is a polyhedral cone $P \subset \overline{\text{Amp}}^e(X/S)$ such that $\text{Aut}(X/S, \Delta) \cdot P = \overline{\text{Amp}}^e(X/S)$. We have the following results.*

- (1) *There are finitely many varieties $Y_j/S, j \in J$ such that if $X \rightarrow Z/S$ is a surjective fibration to a normal variety Z , then $Y_j \simeq Z/S$ for some $j \in J$.*
- (2) *The cone $\overline{\text{Amp}}^e(X/S)$ has a rational polyhedral fundamental domain.*

This result can be shown analogously as Proposition 5.3 and thus we only sketch the proof.

Sketch of the Proof. For (1), let A be an ample/ S divisor on Z . Then for a morphism $g : X \rightarrow Z/S$, g^*A lies in $\overline{\text{Amp}}^e(X/S)$. There exists $\theta \in \text{Aut}(X/S, \Delta)$ such that $[\theta \cdot g^*A]$ lies in the interior of a face $F \subset P$. The morphism $g \circ \theta^{-1} : X \rightarrow Z$ corresponds to the contraction of F . As P is a polyhedral cone, there are only finitely many faces.

(2) follows from Lemma 3.5 as $\text{Amp}(X/S)$ is non-degenerate. \square

We have the following remark regarding the cones chosen in the statement of the cone conjecture (cf. [LOP18, Section 3]):

Remark 5.6. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assuming that good minimal models of effective klt pairs exist in dimension $\dim(X/S)$ and either $R^1 f_* \mathcal{O}_X$ is a torsion sheaf or $\overline{\text{Mov}}(X/S)$ is non-degenerate, Lemma 5.2 and Proposition 5.3 (1) imply that $\text{Mov}(X/S)$ has a (weak) rational polyhedral fundamental domain iff $\overline{\text{Mov}}^e(X/S)$ has a (weak) rational polyhedral fundamental domain.*

Therefore, at least when S is a point, modulo the standard conjectures in the minimal model program, there is no difference to state the cone conjecture of movable cones for either $\text{Mov}(X/S)$ or $\overline{\text{Mov}}^e(X/S)$.

If $\text{Amp}(X/S)$ has a (weak) rational polyhedral fundamental domain, then $(\text{Amp}(X/S)_+, \Gamma_A)$ is of polyhedral type. Proposition 3.3 and Lemma 5.1 imply that

$$\text{Amp}(X/S) = \overline{\text{Amp}}^e(X/S) = \text{Amp}(X/S)_+.$$

Therefore, $\overline{\text{Amp}}^e(X/S)$ has a rational polyhedral fundamental domain by Lemma 3.5. In summary, the cone conjecture for $\text{Amp}(X/S)$ implies that for $\overline{\text{Amp}}^e(X/S)$.

However, a priori, $\text{Mov}(X/S)_+$ (resp. $\text{Amp}(X/S)_+$) has a rational polyhedral fundamental domain Π may not imply that $\overline{\text{Mov}}^e(X/S)$ (resp. $\overline{\text{Amp}}^e(X/S)$) has a rational polyhedral fundamental domain. More importantly, under this assumption, we only know $\Pi \subset \overline{\text{Eff}}(X/S)$, hence Theorem 2.6 and Theorem 2.7 do not apply in this setting. Therefore, the argument of finiteness of birational models which are isomorphic in codimension 1 (resp. finiteness of contraction morphisms)

breaks. It is for this reason that we do not state the cone conjectures for $\text{Mov}(X/S)_+$ and $\text{Amp}(X/S)_+$.

The above discussions lead to the proof of Theorem 1.3.

Proof of Theorem 1.3. The (1) and (2) follow from Lemma 5.2 and Proposition 5.3 (4) and (3). The (3) follows from Lemma 5.2 and Proposition 5.5 (2). \square

6. GENERIC AND GEOMETRIC CONE CONJECTURES

6.1. Generic cone conjecture. For a Calabi-Yau fiber space, we study the relationship between the relative cone conjecture and the cone conjecture of its generic fiber. Conjecture 1.2 is especially convenient to study movable cones in the relative setting. Hence we only focus on the cone conjecture for movable cones in this section.

Let $f : X \rightarrow S$ be a Calabi-Yau fiber space. Recall that $K := K(S)$ is the field of rational functions of S , and $X_K := X \times_S \text{Spec } K$.

Theorem 6.1. *Let $f : X \rightarrow S$ be a Calabi-Yau fiber space such that X has terminal singularities. Suppose that good minimal models of effective klt pairs exist in dimension $\dim(X/S)$. Assume that $R^1 f_* \mathcal{O}_X$ is a torsion sheaf.*

If the weak cone conjecture holds true for $\overline{\text{Mov}}^e(X_K/K)$, then the weak cone conjecture holds true for $\overline{\text{Mov}}^e(X/S)$.

Moreover, if $\text{Mov}(X/S)$ is non-degenerate, then the cone conjecture holds true for $\overline{\text{Mov}}^e(X/S)$. In particular, if S is \mathbb{Q} -factorial, then the cone conjecture holds true for $\overline{\text{Mov}}^e(X/S)$.

Proof. Let $\Pi_K \subset \overline{\text{Mov}}^e(X_K/K)$ be a polyhedral cone such that

$$\text{PsAut}(X_K/K) \cdot \Pi_K = \text{Mov}(X_K/K).$$

Let $\Pi \subset \text{Eff}(X/S)$ be a polyhedral cone which is a lift of Π_K . In other words, Π maps to Π_K under $N^1(X/S)_{\mathbb{R}} \rightarrow N^1(X_K/K)_{\mathbb{R}}$.

If $g_K \in \text{PsAut}(X_K/K)$, then g_K can be viewed as a birational morphism g of X over S . Then $g \in \text{Bir}(X/S) = \text{PsAut}(X/S)$ as K_X is nef/ S and X has terminal singularities.

Let $W \subset \overline{\text{Mov}}(X/S)$ be the maximal vector space. We claim that for $P := \text{Cone}(\Pi \cup W)$,

$$(6.1.1) \quad P \subset \text{Eff}(X/S) \text{ and } \text{PsAut}(X/S) \cdot P \supset \text{Mov}(X/S).$$

By Proposition 4.4 (2), W is generated by vertical divisors and thus $W \subset \text{Eff}(X/S)$. This shows $P \subset \text{Eff}(X/S)$. Next, for any $[M] \in$

$\text{Mov}(X/S)$ such that M is an \mathbb{R} -Cartier divisor. There exist an \mathbb{R} -Cartier divisor D on X and a $g \in \text{PsAut}(X/S)$ such that $[D] \in \Pi$ and $g_K \cdot [D_K] = [M_K]$. As $R^1 f_* \mathcal{O}_X$ is a torsion sheaf, $g_K \cdot D_K \sim_{\mathbb{R}} M_K$. Therefore, there exists a vertical divisor B on X such that $g \cdot D + B \sim_{\mathbb{R}} M/S$. Thus $D + g^{-1} \cdot B \in P$ and $g \cdot [D + g^{-1} \cdot B] = [M]$. This shows $\text{PsAut}(X/S) \cdot P \supset \text{Mov}(X/S)$.

The (6.1.1) shows that Conjecture 1.2 (1) is satisfied. Then Theorem 1.3 (1) and (2) imply the desired claim. Note that by Proposition 4.4 (4), if S is \mathbb{Q} -factorial, then $\overline{\text{Mov}}(X/S)$ is non-degenerate. \square

Remark 6.2. *The above argument does not work for a log pair $(X/S, \Delta)$ because $g \in \text{PsAut}(X_K/K, \Delta_K)$ may not lift to $\text{PsAut}(X/S, \Delta)$.*

Now Theorem 1.4 follows from Theorem 6.1 and the cone conjecture of K3 surfaces over arbitrary fields with characteristic $\neq 2$ ([BLvL20]).

Proof of Theorem 1.4. We have $R^1 f_* \mathcal{O}_X \otimes k(t) \simeq H^1(X_t, \mathcal{O}_{X_t}) = 0$, where $t \in S$ is a general closed point. Hence $R^1 f_* \mathcal{O}_X$ is a torsion sheaf.

We claim that X_K is a smooth K3 surface. Let $U \subset S$ be a smooth open set such that $X_U \rightarrow U$ is flat and for any closed point $t \in U$, X_t is a K3 surface. By [Sta22, Lemma 01V8], $f_U : X_U \rightarrow U$ is a smooth morphism. Thus X_K/K is smooth. Note that

$$\text{Spec } K(S) \rightarrow U, \quad \text{Spec } k(t) \rightarrow U$$

are flat morphisms, where $t \in U$ is a closed point. Then [Har77, III Prop 9.3] implies that for a quasi-coherent sheaf \mathcal{F} on X_U and $i \geq 0$,

$$(6.1.2) \quad \begin{aligned} H^i(X_U, \mathcal{F}) \otimes_U K &\simeq H^i(X_K, \mathcal{F}_{X_K}), \\ H^i(X_U, \mathcal{F}) \otimes_U k(t) &\simeq H^i(X_t, \mathcal{F}_{X_t}). \end{aligned}$$

First, applying (6.1.2) to $\omega_{X_U/U}, \omega_{X_U/U}^{-1}$ and $i = 0$, we have $\mathcal{O}_{X_K}(K_{X_K}) \simeq \mathcal{O}_{X_K}$. Next, applying (6.1.2) to \mathcal{O}_{X_U} and $i = 1$, we have $H^1(X_K, \mathcal{O}_{X_K}) = 0$. This shows that X_K/K is a K3 surface.

We claim that $\text{Amp}(X_K/K)_+ = \overline{\text{Amp}}^e(X_K/K)$. It suffices to show $\text{Amp}(X_K/K)_+ \subset \overline{\text{Eff}}(X_K/K)$. Let D_K be a Cartier divisor on X_K such that $[D_K] \in \overline{\text{Amp}}(X_K/K) \cap N^1(X_K/K)_{\mathbb{Q}}$. A similar argument as above shows that $X_{\bar{K}}$ is a K3 surface over \bar{K} . An application of Riemann-Roch shows that there exists an effective divisor $E_{\bar{K}}$ such that $D_{\bar{K}} \sim E_{\bar{K}}$. To see $[D_K] \in \overline{\text{Eff}}(X_K/K)$, we can argue similarly as Proposition 4.4 (1). Hence, we only sketch the argument. Shrinking S , there is a finite Galois morphism $T \rightarrow S$ such that everything above can be lifted to X_T . Let $\mu : X_T \rightarrow X_S$. In particular, if D is a divisor

on X which is D_K after base change, then there is a divisor $E > 0$ on X_T which is $E_{\bar{K}}$ after base change, and $D_T \sim E/T$. Then

$$m\mu^*D = mD_T \sim \sum_{\theta \in \text{Gal}(X_T/X)} \theta \cdot E / S,$$

where $m = |\text{Gal}(X_T/X)|$. Hence $[D] \in \text{Eff}(X/S)$ and thus $[D_K] \in \text{Eff}(X_K/K)$.

By [BLvL20, Corollary 3.15], there is a rational polyhedral cone $\Pi \subset \text{Amp}(X_K/K)_+$ which is a fundamental domain of $\text{Amp}(X_K/K)_+$ under the action of $\text{Aut}(X_K/K)$. By $\text{Amp}(X_K/K)_+ = \overline{\text{Amp}}^e(X_K/K) = \overline{\text{Mov}}^e(X_K/K)$ and $\text{Aut}(X_K/K) = \text{PsAut}(X_K/K)$, Theorem 6.1 implies the desired result. \square

Remark 6.3. *For a projective hyperkähler manifold X over a characteristic zero field k , [Tak21, Theorem 1.0.5] showed that $\text{Mov}(X/k)_+$ has a rational polyhedral fundamental domain Π under the action of $\text{Bir}(X/k)$. However, this is not sufficient to deduce the cone conjecture for movable cones of hyperkähler fibrations. Indeed, we do not know $\Pi \subset \text{Eff}(X/k)$. On the other hand, when $k = \mathbb{C}$, it is conjectured that $\text{Mov}(X/k)_+ \subset \text{Eff}(X/k)$ (see [BM14, Conjecture 1.4]).*

6.2. Geometric cone conjecture. For a klt Calabi-Yau fiber space, we study the relationship between the relative cone conjecture and the cone conjecture of its geometric fiber. We show that the cone conjecture for the movable cone of the geometric fiber implies the relative cone conjecture for the movable cone of a finite Galois base change.

Recall that if X is a non- \mathbb{Q} -factorial variety over S and $\tilde{X} \rightarrow X$ is a small \mathbb{Q} -factorization, then the (weak) cone conjecture for $\overline{\text{Mov}}^e(X/S)$ is referred to that for $\overline{\text{Mov}}^e(\tilde{X}/S)$. By [BCHM10, Corollary 1.4.3], such small \mathbb{Q} -factorization exists if there is a divisor Δ such that (X, Δ) is klt.

For a klt Calabi-Yau fiber space $f : (X, \Delta) \rightarrow S$. Let $K = K(S)$ and \bar{K} be the algebraic closure of K . For a $g \in \text{Bir}(X/S)$, let $g_{\bar{K}} \in \text{Bir}(X_{\bar{K}}/\bar{K})$ be the extension of g under the base change $\text{Spec } \bar{K} \rightarrow S$. Let $X_{\bar{K}} := X \times_S \text{Spec } \bar{K}$ be the geometric fiber of f . Set $\Delta_{\bar{K}} := \Delta \times_S \text{Spec } \bar{K}$. By Proposition 4.1 (1), $(X_{\bar{K}}, \Delta_{\bar{K}})$ is still klt. Let $\tilde{\pi} : \widetilde{X}_{\bar{K}} \rightarrow X_{\bar{K}}$ be a small \mathbb{Q} -factorization. Set $\widetilde{\Delta}_{\bar{K}}$ be the strict transform of $\Delta_{\bar{K}}$. Let $\bar{\Gamma}_B$ be the image of $\text{PsAut}(\widetilde{X}_{\bar{K}}/\bar{K}, \widetilde{\Delta}_{\bar{K}})$ ($= \text{PsAut}(X_{\bar{K}}/\bar{K}, \Delta_{\bar{K}})$) under the group homomorphism

$$(6.2.1) \quad \iota_{\bar{K}} : \text{PsAut}(\widetilde{X}_{\bar{K}}/\bar{K}, \widetilde{\Delta}_{\bar{K}}) \rightarrow \text{GL}(N^1(\widetilde{X}_{\bar{K}}/\bar{K})_{\mathbb{R}}).$$

Proposition 6.4. *Under the above notation and assumptions.*

- (1) If the weak cone conjecture of $\overline{\text{Mov}}^e(X_{\bar{K}}/\bar{K})$ holds true, then there is a dominant étale Galois morphism $T \rightarrow S$ such that for any $\bar{g} \in \text{PsAut}(X_{\bar{K}}/\bar{K}, \Delta_{\bar{K}})$, there exists a $g \in \text{PsAut}(\widetilde{X}_T/T, \widetilde{\Delta}_T)$ such that $g_{\bar{K}}$ and \bar{g} induce the same action on $N^1(\widetilde{X}_{\bar{K}}/\bar{K})_{\mathbb{R}}$. Here \widetilde{X}_T is a small \mathbb{Q} -factorization of X_T and $\widetilde{\Delta}_T$ is the strict transform of Δ_T .
- (2) If the weak cone conjecture of $\overline{\text{Amp}}^e(X_{\bar{K}}/\bar{K})$ holds true, then there is a dominant étale Galois morphism $T \rightarrow S$ such that for any $\bar{g} \in \text{Aut}(X_{\bar{K}}/\bar{K}, \Delta_{\bar{K}})$, there exists a $g \in \text{Aut}(X_T/T, \Delta_T)$ such that $g_{\bar{K}}$ and \bar{g} induce the same action on $N^1(X_{\bar{K}}/\bar{K})_{\mathbb{R}}$.

Proof. We only show (1) as (2) can be shown analogously.

By assumption, Conjecture 1.2 (1) is satisfied for $\text{Mov}(\widetilde{X}_{\bar{K}}/\bar{K})$. As $\overline{\text{Mov}}(\widetilde{X}_{\bar{K}}/\bar{K})$ is non-degenerate, Theorem 1.3 (2) shows that the cone conjecture holds true for $\overline{\text{Mov}}^e(\widetilde{X}_{\bar{K}}/\bar{K})$. By Theorem 3.6, $\bar{\Gamma}_B$ is finitely presented (the following argument only needs it to be finitely generated). Choose

$$\bar{g}_1, \dots, \bar{g}_m \in \text{PsAut}(\widetilde{X}_{\bar{K}}/\bar{K}, \Delta_{\bar{K}})$$

such that $\iota_{\bar{K}}(\bar{g}_1), \dots, \iota_{\bar{K}}(\bar{g}_m)$ (see (6.2.1)) are generators of $\bar{\Gamma}_B$. As $N^1(\widetilde{X}_{\bar{K}}/\bar{K})_{\mathbb{R}}$ is of finite dimension, let $\widetilde{D}_1, \dots, \widetilde{D}_\rho$ be divisors such that $[\widetilde{D}_1], \dots, [\widetilde{D}_\rho]$ is a basis. There is a finite base change $T \rightarrow S$ such that \bar{g}_j and $\widetilde{D}_i := \tilde{\pi}_* \widetilde{D}_i$ can be defined on $X_T \rightarrow T$. In other words, there exist a $g_j \in \text{Bir}(X_T/T, \Delta_T)$ and a D_i on X_T , such that $(g_j)_{\bar{K}} = \bar{g}_j$ and $(D_i)_{\bar{K}} = \widetilde{D}_i$. Shrinking T , (X_T, Δ_T) has klt singularities by Proposition 4.1 (2). Let $\mu : \widetilde{X}_T \rightarrow X_T$ be a small \mathbb{Q} -factorization. Let $\widetilde{D}_i := \mu_*^{-1} D_i$. Shrinking T further, there exists a natural inclusion $N^1(\widetilde{X}_T/T)_{\mathbb{R}} \hookrightarrow N^1((\widetilde{X}_T)_{\bar{K}}/\bar{K})_{\mathbb{R}}$ by Proposition 4.3. Because $((\widetilde{X}_T)_{\bar{K}}, (\widetilde{\Delta}_T)_{\bar{K}})$ has klt singularities by Proposition 4.1 (1) and $(\widetilde{X}_T)_{\bar{K}} \rightarrow (X_T)_{\bar{K}} = X_{\bar{K}}$ is a small morphism, a small \mathbb{Q} -factorization $Y_{\bar{K}} \rightarrow (\widetilde{X}_T)_{\bar{K}}$ is still a small \mathbb{Q} -factorization of $X_{\bar{K}}$. Thus

$$N^1(\widetilde{X}_T/T)_{\mathbb{R}} \hookrightarrow N^1((\widetilde{X}_T)_{\bar{K}}/\bar{K})_{\mathbb{R}} \hookrightarrow N^1(Y_{\bar{K}}/\bar{K})_{\mathbb{R}} \simeq N^1(\widetilde{X}_{\bar{K}}/\bar{K})_{\mathbb{R}}.$$

By the choice of T , this is also a surjective map. Hence

$$N^1((\widetilde{X}_T)_{\bar{K}}/\bar{K})_{\mathbb{R}} \simeq N^1(Y_{\bar{K}}/\bar{K})_{\mathbb{R}}$$

and thus $(\widetilde{X_T})_{\bar{K}}$ is \mathbb{Q} -factorial. As $(\widetilde{X_T})_{\bar{K}} \rightarrow X_{\bar{K}}$ is a small \mathbb{Q} -factorization, it suffices to show the claim for $N^1((\widetilde{X_T})_{\bar{K}}/\bar{K})_{\mathbb{R}}$. Possibly taking a Galois cover of T and shrinking T , we can assume that $T \rightarrow S$ is a dominant étale Galois morphism.

We claim that after shrinking T , we have $g_j \in \text{PsAut}(X_T/T, \Delta_T)$ for each j . If $g_j \in \text{Bir}(X_T/T, \Delta_T) \setminus \text{PsAut}(X_T/T, \Delta_T)$, then there are finitely many divisors $B_l, l \in J$ which are contracted by g_j or g_j^{-1} . As $(g_j)_{\bar{K}}$ and $(g_j^{-1})_{\bar{K}}$ do not contract $(B_l)_{\bar{K}}$, B_l is vertical over T . Therefore, shrinking T , we can assume that g_j and g_j^{-1} do not contract divisors for each j . This shows the claim.

Finally, let $\bar{D}_i := (\widetilde{D}_i)_{\bar{K}}$, and for $g, h \in \{g_j \mid 1 \leq j \leq m\}$, let $\bar{g} := g_{\bar{K}}, \bar{h} := h_{\bar{K}}$. Then for each i ,

$$\bar{g}_*(\bar{h}_*(\bar{D}_i)) = \overline{(g \circ h)}_*(\bar{D}_i).$$

This implies that

$$\iota_{\bar{K}}(\bar{g})(\iota_{\bar{K}}(\bar{h}) \cdot [\bar{D}_i]) = \iota_{\bar{K}}(\overline{g \cdot h}) \cdot [\bar{D}_i].$$

As $[\bar{D}_i], i = 1, \dots, \rho$ is a basis of $N^1((\widetilde{X_T})_{\bar{K}}/\bar{K})_{\mathbb{R}}$, we have

$$\iota_{\bar{K}}(\bar{g})\iota_{\bar{K}}(\bar{h}) = \iota_{\bar{K}}(\overline{g \cdot h}).$$

Now the desired result follows as $\iota_{\bar{K}}(\bar{g}_j), 1 \leq j \leq m$ generate $\bar{\Gamma}_B$. \square

Remark 6.5. *Let $T' \rightarrow S$ be a dominant étale Galois morphism which factors through $T \rightarrow S$. By the proof of Proposition 6.4, after shrinking T' , the claims in Proposition 6.4 still hold true for $T' \rightarrow S$.*

Theorem 6.6. *Let $f : (X, \Delta) \rightarrow S$ be a klt Calabi-Yau fiber space. Assume that good minimal models of effective klt pairs exist in dimension $\dim(X/S)$.*

- (1) *Assume that the weak cone conjecture holds true for $\overline{\text{Mov}}^e(X_{\bar{K}}/\bar{K})$. Then there is a dominant étale Galois morphism $T \rightarrow S$ such that the cone conjecture holds true for $\overline{\text{Mov}}^e(X_T/T)$.*
- (2) *Assume that the weak cone conjecture holds true for $\overline{\text{Amp}}^e(X_{\bar{K}}/\bar{K})$. Then there is a dominant étale Galois morphism $T \rightarrow S$ such that the cone conjecture holds true for $\overline{\text{Amp}}^e(X_T/T)$.*

Proof. We only show (1) as (2) can be shown analogously.

By the proof of Proposition 6.4, there exist a dominant étale Galois morphism $T \rightarrow S$ and a small \mathbb{Q} -factorization $\widetilde{X_T} \rightarrow X_T$ such that $(\widetilde{X_T}, \widetilde{\Delta_T}) \rightarrow T$ is a klt Calabi-Yau fiber space and $(\widetilde{X_T})_{\bar{K}} \rightarrow X_{\bar{K}}$ is a

small \mathbb{Q} -factorization. Replacing $(X, \Delta) \rightarrow S$ by $(\widetilde{X}_T, \widetilde{\Delta}_T) \rightarrow T$, we can assume that $X_{\bar{K}}$ is \mathbb{Q} -factorial.

Let $\Pi_{\bar{K}} \subset \overline{\text{Mov}}^e(X_{\bar{K}}/\bar{K})$ be a rational polyhedral cone such that

$$(6.2.2) \quad \text{PsAut}(X_{\bar{K}}/\bar{K}, \Delta_{\bar{K}}) \cdot \Pi_{\bar{K}} = \overline{\text{Mov}}^e(X_{\bar{K}}/\bar{K}).$$

There exist a finite morphism $T \rightarrow S$ and finitely many effective divisors $D_j, j \in J$ on X_T such that $\text{Cone}([(D_j)_{\bar{K}}] \mid j \in J) = \Pi_{\bar{K}}$. Replacing T by a higher finite morphism and shrinking T , we can further assume that $T \rightarrow S$ is a dominant étale Galois morphism which satisfies Proposition 6.4 (1) (see Remark 6.5). We can further assume that (X_T, Δ_T) has klt singularities with $K_{X_T} + \Delta_T \sim_{\mathbb{R}} 0/T$ by Proposition 4.1 (2).

Let $\mu : \widetilde{X}_T \rightarrow X_T$ be a small \mathbb{Q} -factorization and $\widetilde{D}_j := \mu_*^{-1} D_j, j \in J$. Shrinking T , there is a natural inclusion $N^1(\widetilde{X}_T/T)_{\mathbb{R}} \hookrightarrow N^1((\widetilde{X}_T)_{\bar{K}}/\bar{K})_{\mathbb{R}}$. Because $(\widetilde{X}_T)_{\bar{K}} \rightarrow (X_T)_{\bar{K}} = X_{\bar{K}}$ is a small morphism and $((\widetilde{X}_T)_{\bar{K}}, (\widetilde{\Delta}_T)_{\bar{K}})$ is klt by Proposition 4.1 (1), there exists a small \mathbb{Q} -factorization $Y_{\bar{K}} \rightarrow (\widetilde{X}_T)_{\bar{K}}$ which is also a small \mathbb{Q} -factorization of $X_{\bar{K}}$. By $X_{\bar{K}}$ \mathbb{Q} -factorial, we have $Y_{\bar{K}} = X_{\bar{K}}$ and thus $(\widetilde{X}_T)_{\bar{K}} = X_{\bar{K}}$. Hence

$$(6.2.3) \quad \text{Mov}(\widetilde{X}_T/T) \hookrightarrow \text{Mov}(X_{\bar{K}}/\bar{K}).$$

Let $\Pi := \text{Cone}([\widetilde{D}_j] \mid j \in J) \subset \text{Eff}(\widetilde{X}_T/T)$. We claim that

$$\text{PsAut}(\widetilde{X}_T/T, \Delta_T) \cdot \Pi \supset \text{Mov}(\widetilde{X}_T/T).$$

In fact, let $[D] \in \text{Mov}(\widetilde{X}_T/T)$. Then there exist a $\bar{g} \in \text{PsAut}(X_{\bar{K}}/\bar{K}, \Delta_{\bar{K}})$ and a $[\bar{B}] \in \Pi_{\bar{K}}$ such that $\bar{g} \cdot [\bar{B}] = [D_{\bar{K}}]$. By the construction of \widetilde{X}_T and Proposition 6.4 (1), there exist a $g \in \text{PsAut}(\widetilde{X}_T/T, \Delta_{\bar{K}})$ and a $\Theta \in \Pi$ such that $[\Theta_{\bar{K}}] = [\bar{B}]$ and

$$[(g_*\Theta)_{\bar{K}}] = g_{\bar{K}} \cdot [\Theta_{\bar{K}}] = \bar{g} \cdot [\bar{B}] = [D_{\bar{K}}].$$

By (6.2.3), $g \cdot [\Theta] = [g_*\Theta] = [D] \in \text{Mov}(\widetilde{X}_T/T)$.

Therefore, Conjecture 1.2 (1) is satisfied. As $\overline{\text{Mov}}(X_{\bar{K}}/\bar{K})$ is non-degenerate, $\overline{\text{Mov}}(X_T/T)$ is non-degenerated by (6.2.3). Hence, (1) follows from Theorem 1.3 (2). \square

It is desirable to deduce the cone conjecture of the Calabi-Yau fiber space $(X, \Delta) \rightarrow S$ from $(X_T, \Delta_T) \rightarrow T$, where $T \rightarrow S$ is a dominant étale Galois morphism. This seems to be a difficult problem. The main obstacle is to descent elements from $\text{PsAut}(X_T/T, \Delta_T)$ and $\text{Aut}(X_T/T, \Delta_T)$ to $\text{PsAut}(X/S, \Delta)$ and $\text{Aut}(X/S, \Delta)$. We propose the following question.

Question 6.7. *Let $f : X \rightarrow S$ be a terminal Calabi-Yau fiber space. Let $T \rightarrow S$ be a dominant étale Galois morphism. Possibly shrinking T , there is a natural group homomorphism*

$$\mathrm{PsAut}(X/S) \hookrightarrow \mathrm{PsAut}(X_T/T).$$

Let Γ_S and Γ_T be the images of $\mathrm{PsAut}(X/S)$ and $\mathrm{PsAut}(X_T/T)$ under the group homomorphism $\mathrm{PsAut}(\widetilde{X}_T/T) \rightarrow \mathrm{GL}(N^1(\widetilde{X}_T/T)_{\mathbb{R}})$. Is Γ_S a finite index subgroup of Γ_T ?

A positive answer of Question 6.7 would give that the weak cone conjecture for $\overline{\mathrm{Mov}}^e(X_{\bar{K}}/\bar{K})$ implies that for $\overline{\mathrm{Mov}}^e(X/S)$.

REFERENCES

- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010.
- [Bir11] Caucher Birkar. On existence of log minimal models II. *J. Reine Angew. Math.*, 658:99–113, 2011.
- [Bir12] Caucher Birkar. Existence of log canonical flips and a special LMMP. *Publ. Math. Inst. Hautes Études Sci.*, 115:325–368, 2012.
- [BLvL20] Martin Bright, Adam Logan, and Ronald van Luijk. Finiteness results for K3 surfaces over arbitrary fields. *Eur. J. Math.*, 6(2):336–366, 2020.
- [BM14] Arend Bayer and Emanuele Macrì. MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations. *Invent. math.*, 198:505–590, 2014.
- [FHS21] Stefano Filipazzi, Christopher Hacon, and Roberto Svaldi. Boundedness of elliptic Calabi-Yau threefolds. *arXiv:2112.01352*, 2021.
- [Han87] Masaki Hanamura. On the birational automorphism groups of algebraic varieties. *Compos. Math.*, 63:123–142, 1987.
- [Har77] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer-Verlag, 1977.
- [HX13] Christopher Hacon and Chenyang Xu. Existence of log canonical closures. *Invent. Math.*, 192(1):161–195, 2013.
- [HX16] Christopher Hacon and Chenyang Xu. On finiteness of b-representations and semi-log canonical abundance. In *Minimal models and extremal rays (Kyoto, 2011)*, pages 361–377. Mathematical Society of Japan, 2016.
- [Kaw97] Yujiro Kawamata. On the cone of divisors of Calabi-Yau fiber spaces. *Internat. J. Math.*, 8:665–687, 1997.
- [Kle05] Steven Kleiman. The Picard scheme. In *Fundamental algebraic geometry*, volume 123 of *Math. Surveys Monogr.*, pages 235–321. Amer. Math. Soc., Providence, RI, 2005.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Loo14] Eduard Looijenga. Discrete automorphism groups of convex cones of finite type. *Compos. Math.*, 150(11):1939–1962, 2014.

- [LOP18] Vladimir Lazić, Keiji Oguiso, and Thomas Peternell. The Morrison-Kawamata cone conjecture and abundance on Ricci flat manifolds. In *Uniformization, Riemann-Hilbert correspondence, Calabi-Yau manifolds & Picard-Fuchs equations*, volume 42 of *Adv. Lect. Math. (ALM)*, pages 157–185. Int. Press, Somerville, MA, 2018.
- [LP13] Vladimir Lazić and Thomas Peternell. On the cone conjecture for Calabi-Yau manifolds with Picard number two. *Math. Res. Lett.*, 20(6):1103–1113, 2013.
- [Mar11] Eyal Markman. A survey of Torelli and monodromy results for holomorphic-symplectic varieties. In *Complex and differential geometry*, volume 8 of *Springer Proc. Math.*, pages 257–322. Springer, Heidelberg, 2011.
- [Mor93] David Morrison. Compactifications of moduli spaces inspired by mirror symmetry. Number 218, pages 243–271. 1993. Journées de Géométrie Algébrique d’Orsay (Orsay, 1992).
- [Mor96] David Morrison. Beyond the Kähler cone. In *Proceedings of the Hirzebruch 65 Conference on Algebraic Geometry (Ramat Gan, 1993)*, volume 9 of *Israel Math. Conf. Proc.*, pages 361–376. Bar-Ilan Univ., Ramat Gan, 1996.
- [Mor15] Dave Witte Morris. *Introduction to arithmetic groups*. Deductive Press, 2015.
- [Ogu14] Keiji Oguiso. Automorphism groups of Calabi-Yau manifolds of Picard number 2. *J. Algebraic Geom.*, 23(4):775–795, 2014.
- [Poo17] Bjorn Poonen. *Rational points on varieties*, volume 188. American Mathematical Society, 2017.
- [PS12] Artie Prendergast-Smith. The cone conjecture for abelian varieties. *J. Math. Sci. Univ. Tokyo*, 19(2):243–261, 2012.
- [SC11] Vyacheslav Shokurov and Sung Rak Choi. Geography of log models: theory and applications. *Centr. Eur. J. Math.*, 9(3):489–534, 2011.
- [Sho96] Vyacheslav Shokurov. 3-fold log models. *J. Math. Sci.*, 81(3):2667–2699, 1996.
- [Sta22] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2022.
- [Tak21] Teppi Takamatsu. On the finiteness of twists of irreducible symplectic varieties. *arXiv:2106.11651v2*, 2021.
- [Tot09] Burt Totaro. The cone conjecture for Calabi-Yau pairs in dimension 2. *Duke Math. J.*, 154:241–263, 2009.

(Zhan Li) DEPARTMENT OF MATHEMATICS, SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY, 1088 XUEYUAN RD, SHENZHEN 518055, CHINA

Email address: lizhan@sustech.edu.cn

(Hang Zhao) SCHOOL OF MATHEMATICS AND STATISTICS, YUNNAN UNIVERSITY, KUNMING 650091, CHINA

Email address: zhaoh3@sustech.edu.cn