

OPTIMAL ANALYSIS OF FINITE ELEMENT METHODS FOR THE STOCHASTIC STOKES EQUATIONS

BUYANG LI, SHU MA, AND WEIWEI SUN

ABSTRACT. Numerical analysis for the stochastic Stokes/Navier-Stokes equations is still challenging even though it has been well done for the corresponding deterministic equations. In particular, the existing error estimates of finite element methods for the stochastic equations all suffer from the order reduction with respect to the spatial discretizations. The best convergence result obtained for these fully discrete schemes is only half-order in time and first-order in space, which is not optimal in space in the traditional sense. The purpose of this article is to establish the strong convergence of $O(\tau^{1/2} + h^2)$ and $O(\tau^{1/2} + h)$ in the L^2 norm for the inf-sup stable velocity-pressure finite element approximations, where τ and h denote the temporal stepsize and spatial mesh size, respectively. The error estimates are of optimal order for the spatial discretization considered in this article (with MINI element), and consistent with the numerical experiments. The analysis is based on the fully discrete Stokes semigroup technique and the corresponding new estimates.

1. Introduction

We consider the time-dependent stochastic Stokes equations in a smooth domain $D \subset \mathbb{R}^d$, $d \in \{2, 3\}$, under the stress boundary condition, i.e.,

$$\begin{cases} du = [\nabla \cdot \mathbb{T}(u, p) + f] dt + B(u) dW(t) & \text{in } D \times (0, T], \\ \nabla \cdot u = 0 & \text{in } D \times (0, T], \\ \mathbb{T}(u, p)\mathbf{n} = 0 & \text{on } \partial D \times (0, T], \\ u = u^0 & \text{at } D \times \{0\}, \end{cases} \quad (1.1)$$

where u and p denote the velocity and the pressure of the fluid, respectively, f is a given source field and \mathbf{n} denotes the outward unit normal vector on the boundary ∂D . Moreover, the stress tensor $\mathbb{T}(u, p)$ is defined by

$$\mathbb{T}(u, p) = 2\mathbb{D}(u) - p\mathbb{I} \quad \text{and} \quad \mathbb{D}(u) = \frac{1}{2}(\nabla u + (\nabla u)^T), \quad (1.2)$$

where \mathbb{I} denotes the identity tensor. The stochastic noise is determined by an $L^2(D)^d$ -valued Q -Wiener process $\{W(t); t \geq 0\}$ on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ with respect to the normal filtration $\{\mathcal{F}_t\}_{t \geq 0}$, and a linear operator $B(u) : L^2(D)^d \rightarrow L^2(D)^d$ which depends on the solution nonlinearly.

The numerical approximation of the deterministic Stokes and Navier–Stokes (NS) equations has been in large part well understood nowadays; see [19, 20, 25, 27, 29–31]. For the stochastic Stokes and NS equations driven by a multiplicative noise, several space-time discretizations have been studied.

- For sinusoidal noises, i.e., $B(u)$ maps $L^2(D)^d$ into its divergence-free subspace, Carelli & Prohl [10] studied implicit and semi-implicit time discretizations with a divergence-free finite element approximation in two dimensions under the periodic boundary condition, and proved the strong convergence of velocity approximations with an error bound of $O(\tau^{\frac{1}{2}} + h)$.
- For sinusoidal noises, under the Dirichlet boundary condition, Carelli, Hausenblas & Prohl [9] proposed a Chorin-type splitting method, with certain finite element approximations in space, and proved an error bound of $O(\tau^{\frac{1}{2}} + h + h^2\tau^{-\frac{1}{2}})$ for the velocity.

Key words and phrases. stochastic Stokes equation, multiplicative noise, Wiener process, semi-implicit Euler scheme, mixed FEM, analytic semigroup, error estimate.

- For general non-sinusoidal noises, under the Dirichlet boundary condition, Brzeźniak, Carelli & Prohl [7] proposed two time-stepping schemes based on the mixed FEMs for the stochastic NS equations and proved the convergence of velocity approximations based on compactness arguments.
- For general non-sinusoidal noises, under the periodic boundary condition, the half-order convergence in probability and logarithmic-order strong convergence in L^2 of a time-splitting method were proved by Bessaih, Brzeźniak & Millet [3] and Bessaih & Millet [4], respectively.
- For general non-sinusoidal noises and the Dirichlet boundary condition, Breit & Prohl [6] proved strong convergence of velocity approximations with an error bound of $O(\tau^{\frac{1}{2}} + h)$ in the energy norm for a fully discrete semi-implicit FEM for the 2D stochastic NS equations.
- The convergence of pressure approximations was studied only recently. In particular, for general non-sinusoidal noises and the periodic boundary condition, Feng & Qiu [14] established strong convergence of order $O(\tau^{\frac{1}{2}} + h\tau^{-\frac{1}{2}})$ for both velocity and pressure approximations (the latter is measured in a time-averaged norm) for a fully discrete semi-implicit mixed FEM. The convergence orders for velocity and pressure approximations were improved to $O(\tau^{\frac{1}{2}} + h^2\tau^{-\frac{1}{2}})$ and $O(\tau^{\frac{1}{2}} + h)$, respectively; see [13, 15].

The numerical analysis in the above-mentioned articles is based on different conditions on the noise, which can be viewed as the following Lipschitz continuity and growth conditions in the case that B is an \mathcal{L}_2^0 -valued function on \mathbb{R} :

$$\|B(v) - B(w)\|_{\mathcal{L}_2^0} \leq C\|v - w\|_{L^2} \quad \text{and} \quad \|B(v)\|_{\mathcal{L}_2^0} \leq C(1 + \|v\|_{L^2}) \quad \forall v, w \in L^2(D)^d, \quad (1.3)$$

where \mathcal{L}_2^0 is the space of Hilbert–Schmidt operators from $Q^{1/2}(L^2(D)^d)$ to $L^2(D)^d$. The condition above requires the series expansion of the noise decays faster than general trace-class noises. In order to incorporate trace-class noises into the numerical analysis, (1.3) may be replaced by the following weaker noise condition:

$$\|B(v) - B(w)\|_{\mathcal{L}_2^0} \leq C\|v - w\|_{H^r} \quad \text{and} \quad \|B(v)\|_{\mathcal{L}_2^0} \leq C(1 + \|v\|_{H^r}) \quad \text{with} \quad r > \frac{d}{2}. \quad (1.4)$$

The error analysis under this weaker condition is more challenging. To the best of our knowledge, the error analysis of fully discrete semi-implicit FEMs based on condition (1.4) was proved only recently by Carelli & Prohl [5] under the periodic boundary condition, where the strong convergence of velocity approximations was proved with an error bound of $O(\tau^{\frac{1}{2}} + h)$. Their proof is based on a stochastic pressure decomposition technique.

In the presence of stochastic noises, the half-order temporal convergence shown in the analyses mentioned above is consistent with the numerical experiments, while the first-order spatial convergence is not optimal and inconsistent with the numerical experiments, mainly due to the limitation of the traditional energy approach. On the other hand, the numerical analysis of the stochastic heat equation has been studied extensively [8, 28, 34, 35] and the second-order convergence in space has been proved. For the stochastic Stokes equations, the existing approach is applicable only to a simple case with a sinusoidal noise and a pointwise divergence-free finite element space, for which the error analysis actually reduces to the analysis for an abstract parabolic equation. The second-order convergence for the stochastic Stokes equations under the common setting with a non-sinusoidal noise and frequently used inf-sup stable mixed FEMs has not been proved. In addition, the convergence order of pressure approximations under the weaker noise condition in (1.4) also has not been investigated yet.

In this article, we establish optimal convergence of fully discrete mixed methods with standard inf-sup stable finite element pairs for the stochastic Stokes equations driven by a non-sinusoidal multiplicative noise under the weaker condition in (1.4) and the regularity of the mild solution (see Proposition 3.1). In particular, the strong convergence in L^2 of order $O(\tau^{\frac{1}{2}} + h^2)$ and $O(\tau^{\frac{1}{2}} + h)$ is proved for the velocity and pressure approximations, respectively, where τ and h denote the temporal stepsize and spatial mesh size, respectively. The error estimates are of optimal order for MINI element in the traditional sense and consistent

with the numerical experiments. The analysis presented in this article is based on the fully discrete Stokes semigroup technique and the corresponding new estimates (see Lemma 4.1 and Remark 4.1), and the H^1 -stability of the orthogonal projection onto the discrete divergence-free finite element subspace (see Section 4.1). The stress boundary condition considered in this article is physically natural.

The rest of this article is organized as follows. In Section 2, we collect the assumptions and describe a fully discrete method with a standard inf-sup stable FE pair for the stochastic Stokes and then, present our main theorem. In Section 3, we present the abstract formulation of the stochastic Stokes equations under the stress boundary condition and define the mild solution based on the abstract formulation. In Section 4, we present some technical estimates for the discrete semigroup associated to the Stokes operator. The results are used in the error analysis of the fully discrete FEMs for the stochastic problem in Section 5. In Section 6, we present numerical experiments to support our theoretical analysis by illustrating the convergence orders of the velocity and pressure approximations.

2. Main results

In this section, we present the notations and assumptions to be used in this article, as well as the numerical scheme for the stochastic Stokes equations to be analyzed. Then we present the main theoretical result on the convergence of the numerical approximations.

2.1. Basic notations

Let D be a bounded smooth domain in \mathbb{R}^d , with $d \in \{2, 3\}$, and let $H^s(D)$, $s \geq 0$, be the conventional Sobolev space of functions defined on D , with $L^2(D) = H^0(D)$. The dual space of $H^s(D)$ is denoted by $\dot{H}^{-s}(D)$.

Let $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ be a filtered probability space with probability measure \mathbb{P} , σ -algebra \mathcal{F} and continuous filtration $\{\mathcal{F}_t\}_{t \geq 0}$. The expectation of a random variable v defined on $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ is denoted by $\mathbb{E}v$.

Let $W(t)$ be a standard $L^2(D)^d$ -valued Q -Wiener process on $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$, with expression

$$W(t) = \sum_{\ell} \sqrt{\mu_{\ell}} \phi_{\ell} W_{\ell}(t) \quad \forall t \in [0, T], \quad (2.1)$$

where $\{W_{\ell}(t)\}_{j \geq 1}$ is a family of independent real-valued Wiener processes and the trace operator $Q : L^2(D)^d \rightarrow L^2(D)^d$ is bounded, self-adjoint, positive semi-definite, with eigenvalues μ_{ℓ} and eigenfunctions ϕ_{ℓ} , $\ell = 1, 2, \dots$.

Let \mathcal{L}_2 and \mathcal{L}_2^0 be the spaces of Hilbert-Schmidt operators from $L^2(D)^d$ to $L^2(D)^d$ and from $Q^{1/2}(L^2(D)^d)$ to $L^2(D)^d$, respectively, satisfying

$$\|\Phi\|_{\mathcal{L}_2^0} := \left(\sum_{\ell} \mu_{\ell} \|\Phi \phi_{\ell}\|_{L^2}^2 \right)^{\frac{1}{2}} = \|\Phi Q^{\frac{1}{2}}\|_{\mathcal{L}_2}.$$

For a progressively measurable process $\Phi : [0, T] \rightarrow \mathcal{L}_2(Q^{1/2}(H), H)$ with $\int_0^T \|\Phi(s)\|_{\mathcal{L}_2^0}^2 ds < \infty$ \mathbb{P} -a.s., the stochastic integral $\int_0^t \Phi(s) dW(s)$ can be defined and Itô's isometry holds, i.e.,

$$\mathbb{E} \left\| \int_0^t \Phi(s) dW(s) \right\|_{L^2}^2 = \mathbb{E} \int_0^t \|\Phi(s)\|_{\mathcal{L}_2^0}^2 ds. \quad (2.2)$$

For the simplicity of notation, we denote by $x \lesssim y$ or $y \gtrsim x$ the statement “ $x \leq Cy$ for some positive constant C (which is independent of the stepsize τ and the mesh size h in the numerical approximation) ”.

2.2. Assumptions on the noise and nonlinearity

For the existence and uniqueness of mild solutions of problem (3.12), as well as the numerical approximation to the mild solutions, we work with the following assumptions on the noise and nonlinearity.

Assumption 2.1. (The stochastic noise) We assume that the Q -Wiener process $W(t)$ has the following property:

$$\|(-\Delta)^{\frac{1}{2}}\|_{\mathcal{L}^2} \lesssim 1, \quad (2.3)$$

where $\Delta : H_N^2(D) \rightarrow L^2(D)$ denotes the Neumann Laplacian operator with domain

$$H_N^2(D) = \{v \in H^2(D) : \partial_n v = 0 \text{ on } \partial D\},$$

and $(-\Delta)^{\frac{1}{2}} : H^1(D) \rightarrow L^2(D)$ denotes the fractional power of $-\Delta$.

Remark 2.1. In the case that Q and $-\Delta$ having the same eigenfunctions, condition (2.3) is equivalent to

$$\sum_{\ell} \mu_{\ell} \lambda_{\ell} \lesssim 1,$$

where λ_{ℓ} are the eigenvalues of $-\Delta$.

Assumption 2.2. (The nonlinearity and source term) We assume that $B(v) : L^2(D)^d \rightarrow L^2(D)^d$ is a bounded linear operator for any $v \in L^2(D)^d$, with the following properties: There exists $\beta \in (\frac{d}{2}, 2)$ such that

$$\|B(v)\phi_{\ell}\|_{H^1} \lesssim \|v\|_{H^{\beta}} \|\phi_{\ell}\|_{H^1} \quad \forall v \in H^{\beta}(D)^d, \quad (2.4)$$

$$\|(B(v_1) - B(v_2))\phi_{\ell}\|_{\dot{H}^{-\frac{1}{2}}} \lesssim \|v_1 - v_2\|_{L^2} \|\phi_{\ell}\|_{H^1} \quad \forall v_1, v_2 \in L^2(D)^d, \quad (2.5)$$

$$\|(B(v_1) - B(v_2))\phi_{\ell}\|_{H^1} \lesssim \|v_1 - v_2\|_{H^{\beta}} \|\phi_{\ell}\|_{H^1} \quad \forall v_1, v_2 \in H^{\beta}(D)^d. \quad (2.6)$$

Moreover, we assume that the function $f : [0, T] \times L^2(D)^d \rightarrow L^2(D)^d$ satisfies

$$\|f(t)\|_{L^2} \lesssim 1 \quad \forall 0 \leq t \leq T, \quad (2.7)$$

$$\|f(t_1) - f(t_2)\|_{L^2} \lesssim (t_1 - t_2)^{\frac{1}{2}} \quad \forall 0 \leq t_1 \leq t_2 \leq T. \quad (2.8)$$

Remark 2.2. In the case that $(B(v)\phi_{\ell})(x) = b_{\ell}(v(x))\phi_{\ell}(x)$ for some functions $b_{\ell} : \mathbb{R} \rightarrow \mathbb{R}$, $\ell = 1, 2, \dots$, conditions (2.4)–(2.6) are satisfied if the functions b_{ℓ} are uniformly Lipschitz continuous with respect to ℓ , i.e.,

$$\begin{aligned} |b_{\ell}(\sigma)| &\lesssim 1 + |\sigma| \quad \forall \sigma \in \mathbb{R}, \\ |b_{\ell}(\sigma_1) - b_{\ell}(\sigma_2)| &\lesssim |\sigma_1 - \sigma_2| \quad \forall \sigma_1, \sigma_2 \in \mathbb{R}. \end{aligned}$$

Assumption 2.3. (Initial value) The initial value $u_0 : \Omega \rightarrow L^2(D)^d$ is an $\mathcal{F}_0/\mathcal{B}(L^2(D)^d)$ -measurable function with $u_0 \in L^2(\Omega, \mathcal{D}(A))$.

2.3. The numerical method and its convergence

Let $V_h \times Q_h \subset H^1(D)^d \times L^2(D)$ be a pair of finite element spaces subject to a quasi-uniform triangulation of D with mesh size $h > 0$, with the following properties (see [17, Chapter II]):

- (1) There exists a projection operator $\Pi_h : H^1(D)^d \rightarrow V_h$, called the Fortin projection, satisfying

$$(\nabla \cdot (v - \Pi_h v), q_h) = 0 \quad \forall v \in H^1(D)^d \text{ and } q_h \in Q_h, \quad (2.9)$$

$$\|v - \Pi_h v\|_{H^m} \lesssim h^{1-m} \|v\|_{H^1} \quad \forall v \in H^1(D)^d \text{ and } m = 0, 1, \quad (2.10)$$

$$\|\Pi_h v\|_{H^1} \lesssim \|v\|_{H^1} \quad \forall v \in H^1(D)^d, \quad (2.11)$$

- (2) The following approximation properties hold:

$$\inf_{q_h \in Q_h} \|q - q_h\|_{L^2} \lesssim h^m \|q\|_{H^m} \quad \forall q \in H^m(D) \text{ and } m = 1, 2, \quad (2.12)$$

Theorem 2.4. *Let Assumptions 2.1–2.3 be fulfilled and assume that the finite element space $V_h \times Q_h$ has Properties (1)–(4). Then the numerical solution (u_h^n, p_h^n) , $n = 1, \dots, N$, determined by (2.19) has the following error bounds:*

$$\max_{1 \leq n \leq N} \left(\mathbb{E} \|u(t_n) - u_h^n\|_{L^2}^2 \right)^{\frac{1}{2}} \lesssim \tau^{\frac{1}{2}} + h^2, \quad (2.24)$$

$$\max_{1 \leq n \leq N} \left(\mathbb{E} \left\| \int_0^{t_m} p(s) \, ds - \tau \sum_{n=1}^m p_h^n \right\|_{L^2}^2 \right)^{\frac{1}{2}} \lesssim \tau^{\frac{1}{2}} + h. \quad (2.25)$$

Remark 2.3. The half-order convergence in time is optimal for the proposed semi-implicit Euler scheme. If we consider a prototypical mini element for spatial discretization, then the error estimate in (2.24) is of optimal order.

The proof of Theorem 2.4 is presented in the next three sections based on the techniques of continuous and discrete analytic semigroups.

3. The abstract formulation under the stress boundary condition

In this section, we present the abstract formulation and functional setting of the stochastic Stokes equations under the stress boundary condition, and define the mild solution of the stochastic Stokes equations to be approximated by the numerical solutions.

The natural function spaces associated to incompressible flow is the divergence-free subspaces of $L^2(D)^d$ and $H^1(D)^d$, defined by

$$X = \{v \in L^2(D)^d : \nabla \cdot v = 0\} \quad \text{and} \quad V = X \cap H^1(D)^d. \quad (3.1)$$

It is known that the following orthogonal decomposition holds:

$$L^2(D)^d = X \oplus U \quad \text{with} \quad U = \{\nabla q : q \in H_0^1(D)\}.$$

In particular, any $v \in L^2(D)^d$ can be decomposed as $v = P_X v + \nabla \eta$, where

$$P_X v = v - \nabla \eta$$

denotes the L^2 -orthogonal projection from $L^2(D)^d$ onto X , with η being the solution of the equation

$$\begin{cases} \Delta \eta = \nabla \cdot v & \text{in } D, \\ \eta = 0 & \text{on } \partial D. \end{cases} \quad (3.2)$$

Since the solution η of the Poisson equation satisfies

$$\|\eta\|_{H^{s+1}} \leq C \|v\|_{H^s} \quad \text{for } s \in [0, 2],$$

it follows that the L^2 projection $P_X v = v - \nabla \eta$ has the following properties:

$$\|P_X v\|_{H^s} \lesssim \|v\|_{H^s} \quad \text{for } s \in [0, 2]. \quad (3.3)$$

Since the L^2 projection operator $P_X : L^2(D)^d \rightarrow X$ is self-adjoint, it follows that (by a duality argument)

$$\|P_X v\|_{\dot{H}^{-s}} \lesssim \|v\|_{\dot{H}^{-s}} \quad \text{for } s \in [0, 2]. \quad (3.4)$$

Let $E(t) : X \rightarrow X$ be the semigroup of bounded linear operators defined by $E(t)v^0 := v(t)$ as the solution of the linear Stokes equations

$$\begin{cases} \frac{\partial v}{\partial t} - \nabla \cdot \mathbb{T}(v, q) = 0 & \text{in } D \times \mathbb{R}_+ \\ \nabla \cdot v = 0 & \text{in } D \times \mathbb{R}_+ \\ T(v, q)n = 0 & \text{on } \partial D \times \mathbb{R}_+ \\ v(0) = v^0 & \text{in } D. \end{cases}$$

Let $-A$ be the generator of the semigroup $E(t)$ with domain

$$D(A) = \left\{ v^0 \in X : \lim_{t \rightarrow 0} \frac{E(t)v^0 - v^0}{t} \text{ exists in } X \right\},$$

which is a dense subspace of X . Then

$$\begin{aligned} v^0 \in D(A) &\Leftrightarrow \partial_t v \in C([0, T]; X) \\ &\Leftrightarrow \nabla \cdot \mathbb{T}(v, q) \in C([0, T]; X) \text{ and } \nabla \cdot v = 0 \\ &\Leftrightarrow -\Delta v + \nabla q \in C([0, T]; X) \text{ and } \nabla \cdot v = 0 \\ &\quad 2\mathbb{D}(v)\mathbf{n} - q\mathbf{n} = 0 \text{ on } \partial D \\ &\quad q \text{ is a harmonic function with boundary condition } q = 2\mathbb{D}(v)\mathbf{n} \cdot \mathbf{n} \end{aligned} \quad (3.5)$$

$$\Leftrightarrow v \in C([0, T]; H^2(D)^d), \quad \nabla \cdot v = 0 \text{ and } \mathbb{D}(v)\mathbf{n} \times \mathbf{n} = 0 \text{ on } \partial D, \quad (3.6)$$

where the last equivalence relation follows from the H^2 regularity of the stationary Stokes equations. Since $v^0 \in D(A) \Leftrightarrow Av \in C([0, T]; X)$, it follows from (3.6) that

$$D(A) = \dot{H}^2(D)^d := \{v \in H^2(D)^d : \nabla \cdot v = 0 \text{ and } \mathbb{D}(v)\mathbf{n} \times \mathbf{n} = 0 \text{ on } \partial D\}.$$

Moreover, from (3.5) we see that the operator $A : D(A) \rightarrow X$ can be written as

$$Av = -\nabla \cdot \mathbb{T}(v, q_v) = -\Delta v + \nabla q_v, \quad (3.7)$$

where q_v is determined by v through the following equation:

$$\begin{cases} \Delta q_v = 0 & \text{in } D, \\ q_v = 2\mathbb{D}(v)\mathbf{n} \cdot \mathbf{n} & \text{on } \partial D. \end{cases} \quad (3.8)$$

For $v \in D(A)$, testing (3.7) with w and using integration by parts, we obtain by utilizing the boundary condition $q_v = 2\mathbb{D}(v)\mathbf{n} \cdot \mathbf{n}$ and $\mathbb{D}(v)\mathbf{n} \times \mathbf{n} = 0$ (which imply $2\mathbb{D}(v)\mathbf{n} - q_v\mathbf{n} = 0$ on ∂D)

$$(Av, w) = (2\mathbb{D}(v), \mathbb{D}(w)) \quad \forall w \in V. \quad (3.9)$$

Therefore, the Stokes operator $A : D(A) \rightarrow X$ has an extension $A : V \rightarrow V'$ defined by (3.9).

The boundary condition $\mathbb{T}(u, p)\mathbf{n} = 0$ in (1.1) implies that $\mathbb{D}(u)\mathbf{n} \times \mathbf{n} = 0$ and $p = 2\mathbb{D}(u)\mathbf{n} \cdot \mathbf{n}$, and therefore

$$-P_X \mathbb{T}(u, p) = -\Delta u + P_X \nabla p = -\Delta u + \nabla(p - \eta),$$

where η is the solution of

$$\begin{cases} \Delta \eta = \Delta p & \text{in } D \\ \eta = 0 & \text{on } \partial D. \end{cases} \quad (3.10)$$

This implies that

$$-P_X \mathbb{T}(u, p) = -\Delta u + \nabla q_u = Au, \quad (3.11)$$

where $q_u = p - \eta$ is the harmonic function satisfying $q_u = 2\mathbb{D}(u)\mathbf{n} \cdot \mathbf{n}$ on ∂D .

As a result, applying P_X to (1.1) yields the following abstract formulation of the stochastic Stokes problem in (1.1):

$$\begin{cases} du = [-Au + P_X f] dt + P_X [B(u) dW] & \text{in } D \times (0, T], \\ u = u^0 & \text{at } D \times \{0\}. \end{cases} \quad (3.12)$$

A predictable process $u \in C([0, T]; L^2(\Omega; X))$ is called a mild solution of problem (3.12) if

$$u(t) = E(t)u^0 + \int_0^t E(t-s)P_X f(s)ds + \int_0^t E(t-s)P_X B(u(s))dW(s) \quad \mathbb{P}\text{-a.s.} \quad (3.13)$$

The proof of the following proposition is presented in Appendix A.

Proposition 3.1. (Well-posedness and regularity) *Under Assumptions 2.1–2.3, problem (3.12) has a unique mild solution in the sense of (3.13). Moreover, the mild solution satisfies the following regularity estimates:*

$$\sup_{t \in [0, T]} \mathbb{E} \|u(t)\|_{H^2}^2 \lesssim (1 + \mathbb{E} \|u^0\|_{H^2}^2), \quad (3.14)$$

$$\mathbb{E}\|u(t) - u(s)\|_{H^1}^2 \lesssim (1 + \mathbb{E}\|u^0\|_{H^2}^2)(t - s) \quad \forall 0 \leq s \leq t \leq T \quad (3.15)$$

$$\mathbb{E}\|u(t) - u(s)\|_{H^\beta}^2 \lesssim (1 + \mathbb{E}\|u^0\|_{H^2}^2)(t - s)^{2-\beta} \quad \forall 0 \leq s \leq t \leq T, \quad \forall \beta \in (\frac{d}{2}, 2). \quad (3.16)$$

The error estimates for the numerical approximations will be proved based on the regularity results in Proposition 3.1.

Since the mild solution has the regularity $u \in C([0, T]; L^2(\Omega, H^1(D)^d))$, the inf-sup condition [16, Theorem 4.1] implies that there exists $\int_0^t p(s) ds \in C([0, T]; L^2(\Omega, L^2(D)))$ satisfying following relation:

$$\begin{aligned} \left(\int_0^t p(s) ds, \nabla \cdot v \right) &= (u(t) - u^0, v) + 2 \int_0^t (\mathbb{D}(u(s)), \mathbb{D}(v)) ds \\ &\quad - \int_0^t (f(s), v) ds - \left(\int_0^t B(u(s)) dW(s), v \right) \quad \forall v \in H^1(D)^d. \end{aligned} \quad (3.17)$$

4. Estimates for the discrete semigroups

In this section, we establish some technical estimates of the discrete analytic semigroup associated to the Stokes operator. The results will be used to prove the optimal-order convergence of the numerical solution in Section 5.

The main results of this section are the following three types of error estimates about approximating the semigroup $E(t)P_X$ by the discrete semigroup $\bar{E}_{h,\tau}^n P_{X_h}$, $n = 1, \dots, N$, defined in (2.23). These results are the key to the error analysis for the stochastic Stokes problem.

Lemma 4.1. *For any $v \in H^1(D)^d$, there holds*

$$\sum_{j=1}^n \int_{t_{j-1}}^{t_j} \| [E(s)P_X - \bar{E}_{h,\tau}^j P_{X_h}] v \|_{L^2}^2 ds \leq C(\tau + h^4) \|v\|_{H^1}^2. \quad (4.1)$$

In addition, for all $v \in L^2(D)^d$, there holds

$$\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E(s)P_X - \bar{E}_{h,\tau}^j P_{X_h}] v ds \right\|_{L^2}^2 \leq C(\tau + h^4) \|v\|_{L^2}^2, \quad (4.2)$$

and

$$\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \nabla [E(s)P_X - \bar{E}_{h,\tau}^j P_{X_h}] v ds \right\|_{L^2}^2 \leq C(\tau + h^2) \|v\|_{L^2}^2. \quad (4.3)$$

Remark 4.1. Note that the error estimates of $E(t)P_X - \bar{E}_{h,\tau}^n P_{X_h}$ in [33] holds for an abstract parabolic equation, including the Stokes equations, but requires the condition $v \in X$. Since we only require $B(u) : L^2(D)^d \rightarrow L^2(D)^d$ rather than $B(u) : L^2(D)^d \rightarrow X$, we cannot apply the results in Lemma 4.1 to the stochastic Stokes equations under the condition $v \in X$. By dropping this condition in Lemma 4.1, we manage to prove second-order convergence in space for the numerical solution of the stochastic Stokes equations.

In order to prove Lemma 4.1, we need to extend the H^1 -stability of the L^2 projection P_{X_h} from $v \in X \cap H^1(D)^d$ to $v \in H^1(D)^d$. This is obtained by characterizing the orthogonal complement of X_h in V_h , as discussed in subsection 4.1. The proof of Lemma 4.1 is presented at the end of this section after introducing the orthogonal decomposition and some properties of the discrete semigroup.

4.1. Orthogonal complement of X_h in V_h

Let X_h^\perp be the orthogonal complement of X_h in V_h , i.e., $V_h = X_h \oplus X_h^\perp$, namely, any $v_h \in V_h$ has an orthogonal decomposition

$$v_h = w_h + z_h \quad \text{with} \quad w_h \in X_h \quad \text{and} \quad z_h \in X_h^\perp \quad \text{satisfying} \quad (w_h, z_h) = 0. \quad (4.4)$$

This decomposition is stable in the H^1 norm, as shown in the following lemma.

Lemma 4.2. *The orthogonal decomposition in (4.4) is stable in the H^1 norm, i.e.,*

$$\|w_h\|_{H^1} + \|z_h\|_{H^1} \lesssim \|v_h\|_{H^1} \quad \forall v_h \in V_h. \quad (4.5)$$

Proof. Let $P_{Q_h} : L^2(D) \rightarrow Q_h$ be the L^2 -orthogonal projection onto Q_h . For any given $v_h \in V_h$, the inf-sup condition (2.18) implies that there exists a unique solution $(w_h, q_h) \in V_h \times Q_h$ of the following equations:

$$\begin{aligned} (w_h, a_h) - (q_h, \nabla \cdot a_h) &= (v_h, a_h) & \forall a_h \in V_h, \\ (\nabla \cdot w_h, \eta_h) &= 0 & \forall \eta_h \in Q_h. \end{aligned}$$

Then $w_h \in X_h$ and

$$(w_h, a_h) = (v_h, a_h) \quad \forall a_h \in X_h.$$

Let $z_h = v_h - w_h$, then w_h and z_h are the functions in the orthogonal decomposition (4.4). Before estimating (w_h, q_h) directly, we first introduce $(w, q) \in H^1(D)^d \times H^1(D)$ to be the solution of the continuous problem

$$\begin{aligned} (w, a) - (q, \nabla \cdot a) &= (v_h, a) & \forall a \in H^1(D)^d, \\ (\nabla \cdot w, \eta) &= 0 & \forall \eta \in L^2(D). \end{aligned}$$

Via integration by parts in the equation of w , one can obtain that $w = v_h - \nabla q$, with $q \in H^1(D)$ being the weak solution of

$$\begin{cases} \Delta q = \nabla \cdot v_h & \text{in } D, \\ q = 0 & \text{on } \partial D. \end{cases}$$

The equation above has the standard H^2 elliptic regularity, i.e.,

$$\|q\|_{H^2} \lesssim \|\nabla \cdot v_h\|_{L^2} \lesssim \|v_h\|_{H^1}, \quad (4.6)$$

which implies that

$$\|w\|_{H^1} = \|v_h - \nabla q\|_{H^1} \lesssim \|v_h\|_{H^1} \quad (4.7)$$

Denote $\theta_h = w_h - \Pi_h w$ and $\phi_h = q_h - P_{Q_h} q$, and consider the difference between the equations of w_h and w , i.e.,

$$\begin{aligned} (\theta_h, a_h) - (\phi_h, \nabla \cdot a_h) &= (w - \Pi_h w, a_h) + (P_{Q_h} q - q, \nabla \cdot a_h) & \forall a_h \in V_h, \\ (\nabla \cdot \theta_h, \eta_h) &= 0 & \forall \eta_h \in Q_h. \end{aligned}$$

Substituting $a_h = \theta_h$ and $\eta_h = \phi_h$ into the equations above and using the inverse inequality (2.17) we obtain

$$\begin{aligned} \|\theta_h\|_{L^2}^2 &= (w - \Pi_h w, \theta_h) + (P_{Q_h} q - q, \nabla \cdot \theta_h) \\ &\leq \|w - \Pi_h w\|_{L^2} \|\theta_h\|_{L^2} + \|P_{Q_h} q - q\|_{L^2} \|\nabla \cdot \theta_h\|_{L^2} \\ &\lesssim h \|w\|_{H^1} \|\theta_h\|_{L^2} + h^2 \|q\|_{H^2} \|\theta_h\|_{H^1} \\ &\lesssim h (\|w\|_{H^1} + \|q\|_{H^2}) \|\theta_h\|_{L^2}, \end{aligned}$$

which together with (4.6) and (4.7) implies

$$\|\theta_h\|_{H^1} \lesssim h^{-1} \|\theta_h\|_{L^2} \lesssim \|w\|_{H^1} + \|q\|_{H^2} \leq \|v_h\|_{H^1}.$$

Therefore, the H^1 -stability (2.11) of the Fortin projection operator gives

$$\|w_h\|_{H^1} = \|\theta_h + \Pi_h w\|_{H^1} \lesssim \|v_h\|_{H^1},$$

which yields

$$\|z_h\|_{H^1} = \|v_h - w_h\|_{H^1} \lesssim \|v_h\|_{H^1}.$$

This proves the desired H^1 -stability result in (4.5). \square

Remark 4.2. The H^1 stability in Lemma 4.2 implies the following properties:

$$\|P_{X_h} v\|_{H^1} \leq C \|v\|_{H^1} \quad \text{for } v \in H^1(D)^d, \quad (4.8)$$

$$\|v - P_{X_h} v\|_{H^1} \lesssim \inf_{v_h \in X_h} \|v - v_h\|_{H^1} \leq Ch \|v\|_{H^2} \quad \text{for } v \in \mathcal{D}(A), \quad (4.9)$$

$$\|v - P_{X_h} v\|_{L^2} \lesssim \inf_{v_h \in X_h} \|v - v_h\|_{L^2} \leq Ch^2 \|v\|_{H^2} \quad \text{for } v \in \mathcal{D}(A). \quad (4.10)$$

For any $v \in \mathcal{D}(A)$, we denote by q_v the solution of (3.8), and denote by $R_{X_h} : V \rightarrow X_h$ the Stokes–Ritz projection defined by

$$(v - R_{X_h} v, w_h) + (A_h(v - R_{X_h} v), w_h) - (q_v, \nabla \cdot w_h) = 0 \quad \forall w_h \in X_h, \quad (4.11)$$

which is equivalent to finding $(R_{X_h} v, q_{v,h}) \in V_h \times Q_h$ satisfying

$$\begin{aligned} (v - R_{X_h} v, w_h) + 2(\mathbb{D}(v - R_{X_h} v), \mathbb{D}(w_h)) - (q_v - q_{v,h}, \nabla \cdot w_h) &= 0 \quad \forall w_h \in V_h, \\ (\nabla \cdot (v - R_{X_h} v), q_h) &= 0 \quad \forall q_h \in Q_h. \end{aligned} \quad (4.12)$$

The Stokes–Ritz projection has the following error bound (cf. [12, Lemma 2.44 and Lemma 2.45] and [12, Proposition 4.18]):

$$\|v - R_{X_h} v\|_{L^2} + h\|v - R_{X_h} v\|_{H^1} \lesssim h^2 (\|v\|_{H^2} + \|q_v\|_{H^1}) \leq Ch^2 \|v\|_{H^2} \quad \forall v \in \mathcal{D}(A). \quad (4.13)$$

4.2. Fractional powers of $I + A$

Since the Stokes operator A defined in (3.7) is self-adjoint and positive semi-definite, it follows that the operator $z + A$ is invertible on X for $z \in \mathbb{C} \setminus (-\infty, 0]$, and $-A$ generates a bounded analytic semigroup $E(t) = e^{-tA}$ on X ; see [1, Example 3.7.5]. The fractional powers of the positive definite operator $I + A$ (with compact inverse) can be defined by means of the spectral decomposition (see [23, Appendix B.2]), and the following norm equivalence holds.

Lemma 4.3. *The following equivalence relations hold for $s \in [0, 2]$:*

$$\|v\|_{H^s} \lesssim \|(I + A)^{\frac{s}{2}} v\|_{L^2} \lesssim \|v\|_{H^s} \quad \forall v \in \mathcal{D}(A^{\frac{s}{2}}), \quad (4.14)$$

$$\|v\|_{\dot{H}^{-s}} \lesssim \|(I + A)^{-\frac{s}{2}} v\|_{L^2} \lesssim \|v\|_{\dot{H}^{-s}} \quad \forall v \in \mathcal{D}(A^{\frac{s}{2}})', \quad (4.15)$$

where $\mathcal{D}(A^{\frac{s}{2}}) = (X, \mathcal{D}(A))_{[s/2]}$ for $s \in [0, 2]$.

Remark 4.3. $(X, \mathcal{D}(A))_{[\theta]}$, with $\theta \in [0, 1]$, denotes the complex interpolation spaces between X and $\mathcal{D}(A)$. Since $\mathcal{D}(A^{\frac{1}{2}}) = X \cap H^1(D)^d$, via complex interpolation it can be shown that $\mathcal{D}(A^{\frac{s}{2}}) = X \cap H^s(D)^d$ for $s \in [0, 1]$.

The proof of Lemma 4.3 is presented in Appendix B.

The analyticity of the semigroup $E(t) = e^{-tA}$ implies the following results.

Lemma 4.4. *For any $v \in X$, the following results hold for $t \in (0, T]$:*

$$\|(I + A)^\gamma E(t)v\|_{L^2} \lesssim t^{-\gamma} \|v\|_{L^2} \quad \forall \gamma \geq 0, \quad (4.16)$$

$$\|(I + A)^{-\mu} (I - E(t))v\|_{L^2} \lesssim t^\mu \|v\|_{L^2} \quad \forall \mu \in [0, 1]. \quad (4.17)$$

The following results hold for $0 \leq t_1 \leq t_2 \leq T$:

$$\int_{t_1}^{t_2} \|(I + A)^{\frac{\rho}{2}} E(t_2 - s)v\|_{L^2}^2 ds \lesssim (t_2 - t_1)^{1-\rho} \|v\|_{L^2}^2 \quad \forall \rho \in [0, 1], \quad (4.18)$$

$$\left\| (I + A)^\rho \int_{t_1}^{t_2} E(t_2 - s)v ds \right\|_{L^2} \lesssim (t_2 - t_1)^{1-\rho} \|v\|_{L^2} \quad \forall \rho \in [0, 1]. \quad (4.19)$$

The proof of Lemma 4.4 is also presented in Appendix B.

4.3. The discrete semigroup from spatial discretization

Let $0 = \lambda_{h,1} \leq \lambda_{h,2} \leq \dots \leq \lambda_{h,M_h}$ be the eigenvalues of the discrete Stokes operator $A_h : X_h \rightarrow X_h$ with corresponding orthonormal eigenvectors $\{\varphi_{h,j}\}_{j=1}^{M_h} \subset X_h$ and $\dim(X_h) = M_h$. The operator $-A_h$ generates a bounded analytic semigroup on X_h defined by

$$E_h(t) = e^{-tA_h}, \quad (4.20)$$

which can be expressed as $E_h(t)v_h = \sum_{j=1}^{M_h} e^{-t\lambda_{h,j}}(v_h, \varphi_{h,j})\varphi_{h,j}$ for all $v_h \in X_h$. Hence, the following two properties can be derived based on the proof of [33, Lemma 3.9] and [24, Lemma 3.2 (iii)], respectively:

$$\|(I + A_h)^{\frac{\gamma}{2}} E_h(t)v_h\|_{L^2} \lesssim t^{-\frac{\gamma}{2}} \|v_h\|_{L^2} \quad \text{for } v_h \in X_h, \gamma \in [0, 1], \quad (4.21)$$

$$\int_0^t \|(I + A_h)^{\frac{1}{2}} E_h(s)v_h\|_{L^2}^2 ds \lesssim \|v_h\|_{L^2}^2, \quad \text{for } v_h \in X_h. \quad (4.22)$$

Let

$$\Phi_h(t) := E(t)P_X - E_h(t)P_{X_h} \quad \text{for } t \in [0, T], \quad (4.23)$$

which is the error in approximating the continuous semigroup. The main results of this subsection are the estimates of $\Phi_h(t)$ presented in the following lemma.

Lemma 4.5. *For $v \in H^1(D)^d$, the following estimates hold:*

$$\|\Phi_h(t)v\|_{L^2} \leq Ct^{-\frac{1}{2}}h^2\|v\|_{H^1}, \quad (4.24)$$

and

$$\left(\int_0^t \|\Phi_h(s)v\|_{L^2}^2 ds \right)^{\frac{1}{2}} \leq Ch^2\|v\|_{H^1}, \quad (4.25)$$

For $v \in L^2(D)^d$, the following estimates hold:

$$\|\Phi_h(t)v\|_{L^2} \leq Ct^{-1}h^2\|v\|_{L^2}, \quad (4.26)$$

and

$$\left\| \int_0^t \Phi_h(s)v ds \right\|_{L^2} + h \left\| \int_0^t \nabla \Phi_h(s)v ds \right\|_{L^2} \leq Ch^2\|v\|_{L^2}. \quad (4.27)$$

If $v \in \mathcal{D}(A^{\frac{1+\rho}{2}})$ then

$$\|\Phi_h(t)v\|_{L^2} + h\|\nabla \Phi_h(t)v\|_{L^2} \leq Ct^{-\frac{1-\rho}{2}}h^2\|v\|_{H^{1+\rho}} \quad \text{for } \rho \in [0, 1]. \quad (4.28)$$

Remark 4.4. Lemma 4.5 is proved later based on an orthogonal decomposition of $v = P_X v + \nabla q$ for $v \in L^2(D)^d$, and it does not require $\nabla \cdot v = 0$ in the cases $v \in L^2(D)^d$ and $v \in H^1(D)^d$. This is important for the error analysis of the stochastic Stokes equations.

The proof of Lemma 4.5 is based on error estimates for the semi-discrete FEM for the deterministic linear Stokes problem

$$\begin{cases} \partial_t u - \nabla \cdot \mathbb{T}(u, p) = 0 & \text{in } D \times [0, T], \\ \nabla \cdot u = 0 & \text{in } D \times [0, T], \\ \mathbb{T}(u, p)\mathbf{n} = 0 & \text{on } \partial D \times [0, T], \\ u(0) = u^0 & \text{in } D. \end{cases} \quad (4.29)$$

Applying P_X to the first equation of (4.29) yields the following abstract formulation:

$$\partial_t u + Au = 0 \quad \text{with} \quad u(0) = u^0 \in X, \quad (4.30)$$

of which the solution can be represented $u(t) = E(t)P_X u^0$ in terms of the semigroup generated by the Stokes operator. The FEM for (4.29) reads: for given $u_h(0) = P_{X_h} u^0$, find $(u_h(t), p_h(t)) \in V_h \times Q_h$ such that

$$\begin{cases} (\partial_t u_h(t), v_h) + 2(\mathbb{D}(u_h(t)), \mathbb{D}(v_h)) - (p_h(t), \nabla \cdot v_h) = 0 & \forall v_h \in V_h, \\ (\nabla \cdot u_h(t), q_h) = 0 & \forall q_h \in Q_h. \end{cases} \quad (4.31)$$

This can be written into the following abstract form by choosing $v_h \in X_h$:

$$\frac{d}{dt} u_h(t) + A_h u_h(t) = 0 \quad \text{with} \quad u_h(0) = P_{X_h} u^0, \quad (4.32)$$

of which the solution can be represented as $u_h(t) = E_h(t)P_{X_h} u^0$.

We shall estimate the error between the solutions of (4.29) and (4.31) by estimating the difference between the continuous and discrete resolvent operators, i.e., $(z + A)^{-1}$ and $(z + A_h)^{-1}$. This is presented in the following lemma. Its proof is presented in Appendix B.

Lemma 4.6. *For any $f \in L^2(D)^d$, let w and w_h be the solution and finite element solution of the Stokes equations*

$$\begin{cases} zw - \nabla \cdot \mathbb{T}(w, p) = f & \text{in } D, \\ \nabla \cdot w = 0 & \text{in } D, \\ \mathbb{T}(w, p)n = 0 & \text{on } \partial D, \end{cases} \quad (4.33)$$

and

$$\begin{cases} (zw_h, v_h) + 2(\mathbb{D}(w_h), \mathbb{D}(v_h)) - (p_h, \nabla \cdot v_h) = (f, v_h) & \forall v_h \in V_h, \\ (\nabla \cdot w_h, q_h) = 0 & \forall q_h \in Q_h, \end{cases} \quad (4.34)$$

where $z \in \Sigma_\phi := \{1 + z' \in \mathbb{C} : |\arg(z')| < \phi\}$ for some $\phi \in (0, \pi)$. Then the following results hold:

$$\|w - w_h\|_{L^2} + h\|\nabla(w - w_h)\|_{L^2} \leq Ch^2(\|w\|_{H^2} + \|p\|_{H^1}), \quad (4.35)$$

$$\|w\|_{H^2} + \|p\|_{H^1} \leq C\|f\|_{L^2}, \quad (4.36)$$

where the constant C is independent of $z \in \Sigma_\phi$ (but may depend on ϕ).

Remark 4.5. The first equation in (4.33) can be written as

$$\nabla p = f - (z - \Delta)w.$$

By applying P_X to the first equation in (4.33) and using relation $-P_X \mathbb{T}(w, p) = -\Delta w + \nabla q_w$, as shown in (3.11), we obtain

$$\nabla q_w = P_X f - (z - \Delta)w.$$

Combining the two equations above yields

$$\nabla p = f - P_X f + \nabla q_w.$$

In the case $f \in X$, we obtain $p = q_w$ and therefore (4.35) reduces to

$$\|w - w_h\|_{L^2} + h\|\nabla w - \nabla w_h\|_{L^2} \leq Ch^2\|w\|_{H^2}, \quad (4.37)$$

where we have used the inequality $\|q_w\|_{H^1} \lesssim \|w\|_{H^2}$.

Before proving Lemma 4.5 for the general case $v \in H^1(D)^d$, we consider the simpler case $g \in X \cap H^1(D)^d$ in the following lemma.

Lemma 4.7. *It holds that*

$$\int_0^t \|\Phi_h(s)g\|_{L^2}^2 ds \leq Ch^4\|g\|_{H^1}^2 \quad \forall g \in X \cap H^1(D)^d. \quad (4.38)$$

Proof. Let (u, p) and (u_h, p_h) be the solution of (4.29) and (4.31) with $u^0 = g \in X \cap H^1(D)^d$, respectively. Then the definition of $\Phi_h(s) = E(s)P_X - E_h(s)P_{X_h}$ in (4.23) implies that

$$\Phi_h(s)g = u(s) - u_h(s). \quad (4.39)$$

Subtracting (4.31) from (4.29) yields

$$\begin{cases} (\partial_t(u - u_h), v_h) + 2(\mathbb{D}(u - u_h), \mathbb{D}(v_h)) - (p - p_h, \nabla \cdot v_h) = 0 & \forall v_h \in V_h, \\ (\nabla \cdot (u - u_h), \eta_h) = 0 & \forall \eta_h \in Q_h, \end{cases}$$

and therefore, choosing $v_h \in X_h$ and using the Ritz projection R_{X_h} defined in (4.11), we have

$$\begin{aligned} & (\partial_t(u - u_h), v_h) + ((I + A_h)(R_{X_h}u - u_h), v_h) \\ &= (u - u_h, v_h) + (q - P_{Q_h}q, \nabla \cdot v_h) - (q_u - P_{Q_u}q_u, \nabla \cdot v_h). \end{aligned} \quad (4.40)$$

By denoting $\tilde{e}_h = P_{X_h}u - u_h \in X_h$ and choosing $v_h = (I + A_h)^{-1}\tilde{e}_h \in X_h$ in (4.40), we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|(I + A_h)^{-\frac{1}{2}}\tilde{e}_h\|_{L^2}^2 + \|\tilde{e}_h\|_{L^2}^2 \\ &= \|(I + A_h)^{-\frac{1}{2}}\tilde{e}_h\|_{L^2}^2 + (P_{X_h}u - R_{X_h}u, \tilde{e}_h) + R_q(\tilde{e}_h) + R_{q_u}(\tilde{e}_h) \end{aligned} \quad (4.41)$$

$$\begin{aligned}
&\leq \|(I + A_h)^{-1}\tilde{e}_h\|_{L^2}^2 + C\|R_{X_h}u - P_{X_h}u\|_{L^2}^2 + \frac{1}{4}\|\tilde{e}_h\|_{L^2}^2 + |R_q(\tilde{e}_h)| + |R_{q_u}(\tilde{e}_h)| \\
&\leq \|(I + A_h)^{-1}\tilde{e}_h\|_{L^2}^2 + Ch^4\|u\|_{H^2}^2 + \frac{1}{4}\|\tilde{e}_h\|_{L^2}^2 + |R_q(\tilde{e}_h)| + |R_{q_u}(\tilde{e}_h)|,
\end{aligned}$$

where

$$R_q(\tilde{e}_h) = (q - P_{Q_h}q, \nabla \cdot [(I + A_h)^{-1}\tilde{e}_h]) \quad \text{and} \quad R_{q_u}(\tilde{e}_h) = -(q_u - P_{Q_h}q_u, \nabla \cdot [(I + A_h)^{-1}\tilde{e}_h]).$$

The remainders $R_q(\tilde{e}_h)$ and $R_{q_u}(\tilde{e}_h)$ can be estimated by considering $w = (I + A)^{-1}P_X\tilde{e}_h$ and $w_h = (I + A_h)^{-1}P_{X_h}\tilde{e}_h$, which are the solutions of (4.33) and (4.34), respectively, with $z = 1$ and $f = \tilde{e}_h$. Then applying Lemma 4.6 yields

$$\|[(I + A)^{-1}P_X - (I + A_h)^{-1}P_{X_h}]\tilde{e}_h\|_{H^1} \lesssim h\|\tilde{e}_h\|_{L^2},$$

which implies that

$$\begin{aligned}
|R_q(\tilde{e}_h)| &= |(q - P_{Q_h}q, \nabla \cdot [(I + A)^{-1}P_X\tilde{e}_h])| \\
&\quad + |(q - P_{Q_h}q, \nabla \cdot [(I + A)^{-1}P_X - (I + A_h)^{-1}P_{X_h}]\tilde{e}_h)| \\
&\lesssim \|q - P_{Q_h}q\|_{\dot{H}^{-1}}\|\nabla \cdot [(I + A)^{-1}P_X\tilde{e}_h]\|_{H^1} \\
&\quad + \|q - P_{Q_h}q\|_{L^2}\|[(I + A)^{-1}P_X - (I + A_h)^{-1}P_{X_h}]\tilde{e}_h\|_{H^1} \\
&\lesssim h^2\|q\|_{H^1}\|(I + A)^{-1}P_X\tilde{e}_h\|_{H^2} + h\|q\|_{H^1}h\|\tilde{e}_h\|_{L^2} \\
&\lesssim h^2\|q\|_{H^1}\|\tilde{e}_h\|_{L^2} \\
&\leq Ch^4\|q\|_{H^1}^2 + \frac{1}{8}\|\tilde{e}_h\|_{L^2}^2,
\end{aligned} \tag{4.42}$$

where we have used the inequality $\|q - P_{Q_h}q\|_{\dot{H}^{-1}} \lesssim h^2\|q\|_{H^1}$; see Property (2) of Section 2.3. Similarly, since $\|q_u\|_{H^1}^2 \lesssim \|u\|_{H^2}^2$, it follows that

$$|R_{q_u}(\tilde{e}_h)| \leq Ch^4\|q_u\|_{H^1}^2 + \frac{1}{8}\|\tilde{e}_h\|_{L^2}^2 \leq Ch^4\|u\|_{H^2}^2 + \frac{1}{8}\|\tilde{e}_h\|_{L^2}^2. \tag{4.43}$$

Substituting (4.42)–(4.43) into (4.41), and using Gronwall's inequality with $\tilde{e}_h(0) = 0$, we obtain

$$\begin{aligned}
\|(I + A_h)^{-\frac{1}{2}}\tilde{e}_h(t)\|_{L^2}^2 + \int_0^t \|\tilde{e}_h(s)\|_{L^2}^2 ds &\lesssim h^4 \int_0^t (\|u\|_{H^2}^2 + \|q\|_{H^1}^2) ds \lesssim h^4\|u^0\|_{H^1}^2 \\
&= h^4\|g\|_{H^1}^2,
\end{aligned} \tag{4.44}$$

where the last inequality is the basic energy inequality for the Stokes equations, which can be obtained by testing (4.30) with Au , respectively. Since $u(s) - u_h(s) = u(s) - P_{X_h}u(s) + \tilde{e}_h$, by using the inequality above and the triangle inequality we obtain

$$\int_0^t \|u(s) - u_h(s)\|_{L^2}^2 ds \lesssim h^4\|g\|_{H^1}^2.$$

This proves the result of Lemma 4.7 in view of (4.39). \square

Now we prove Lemma 4.5 by utilizing the result of Lemma 4.7.

Proof of Lemma 4.5. For $v \in L^2(D)^d$ we denote

$$w = (z + A)^{-1}P_Xv \quad \text{and} \quad w_h = (z + A_h)^{-1}P_{X_h}v,$$

which are the solutions of (4.33) and (4.34) with $f = v$.

Since $-A$ generates a bounded analytic semigroup on X , there exists an angle $\phi \in (\frac{\pi}{2}, \pi)$ such that the operator $(z + A)^{-1}$ is analytic with respect to z in the sector $\Sigma_\phi = \{z \in \mathbb{C} \setminus \{0\} : |\arg(z)| < \phi\}$. Moreover, the semigroup $E(t) = e^{-tA}$ can be expressed in terms of the resolvent operator $(z + A)^{-1}$ through a contour integral on the complex plane, i.e.,

$$E(t)P_Xv = \frac{1}{2\pi i} \int_\Gamma (z + A)^{-1}P_Xve^{zt} dz, \quad \text{with } \Gamma = \{1 + z : |\arg(z)| = \phi\} \subset \Sigma_\phi. \tag{4.45}$$

Similarly,

$$E_h(t)P_{X_h}v = \frac{1}{2\pi i} \int_{\Gamma} (z + A_h)^{-1} P_{X_h} v e^{zt} dz. \quad (4.46)$$

Therefore,

$$\begin{aligned} \|\Phi_h(t)v\|_{L^2} &= \|E(t)P_X v - E_h(t)P_{X_h}v\|_{L^2} \\ &= \left\| \frac{1}{2\pi i} \int_{\Gamma} [(z + A)^{-1} P_X v - (z + A_h)^{-1} P_{X_h} v] e^{zt} dz \right\|_{L^2} \\ &\lesssim \int_{\Gamma} \|w - w_h\|_{L^2} e^{\operatorname{Re}(z)t} |dz| \\ &\lesssim \int_{\Gamma} h^2 \|v\|_{L^2} e^{\operatorname{Re}(z)t} |dz| \quad (\text{This follows from Lemma 4.6 with } f = v) \\ &\lesssim t^{-1} h^2 \|v\|_{L^2} \quad \text{for } t \in (0, T]. \end{aligned} \quad (4.47)$$

This proves (4.26).

If $v \in \mathcal{D}(A^{\frac{1+\rho}{2}})$ then we use inequality (4.37), which implies that

$$\begin{aligned} \|w - w_h\|_{L^2} + h\|\nabla(w - w_h)\|_{L^2} &\lesssim h^2 \|w\|_{H^2} \\ &= h^2 \|(z + A)^{-1} v\|_{H^2} \\ &\lesssim h^2 \|(I + A)(z + A)^{-1} v\|_{L^2} \\ &= h^2 \|(I + A)^{\frac{1-\rho}{2}} (z + A)^{-1} (I + A)^{\frac{1+\rho}{2}} v\|_{L^2} \\ &\lesssim h^2 |z|^{-\frac{1+\rho}{2}} \|(I + A)^{\frac{1+\rho}{2}} v\|_{L^2} \quad \forall \rho \in [0, 1], \end{aligned} \quad (4.48)$$

where we have used the interpolation inequality to get

$$\|(I + A)^{\theta} (z + A)^{-1} v\|_{L^2} \leq C_{\theta} |z|^{-(1-\theta)} \|v\|_{L^2} \quad \forall \theta \in [0, 1], v \in X. \quad (4.49)$$

Substituting (4.48) into (4.47) and using (4.14) gives

$$\begin{aligned} \|\Phi_h(t)v\|_{L^2} + h\|\nabla\Phi_h(t)v\|_{L^2} &\lesssim h^2 \|(I + A)^{\frac{1+\rho}{2}} v\|_{L^2} \int_{\Gamma} |z|^{-\frac{1+\rho}{2}} e^{\operatorname{Re}(z)t} |dz|. \\ &\lesssim t^{-\frac{1-\rho}{2}} h^2 \|v\|_{H^{1+\rho}} \quad \forall \rho \in [0, 1], v \in \mathcal{D}(A^{\frac{1+\rho}{2}}). \end{aligned} \quad (4.50)$$

This proves (4.28).

In order to prove (4.24)–(4.25), we consider the orthogonal decomposition $v = P_X v + \nabla q$ for a function $v \in H^1(D)^d$, where $q \in H^1(D)$ is the weak solution of

$$\begin{cases} \Delta q = \nabla \cdot v & \text{in } D \\ q = 0 & \text{on } \partial D, \end{cases}$$

which has the classical $H_0^1(D) \cap H^2(D)$ elliptic regularity, i.e.,

$$\|q\|_{H^2} \lesssim \|\nabla \cdot v\|_{L^2} \lesssim \|v\|_{H^1}. \quad (4.51)$$

Since $\Phi_h(t)\nabla q = E(t)P_X \nabla q - E_h(t)P_{X_h} \nabla q = -E_h(t)P_{X_h} \nabla q \in X_h$, by using the self-adjointness of $E_h(t)$ we have

$$\begin{aligned} (\Phi_h(t)\nabla q, a_h) &= -(P_{X_h} \nabla q, E_h(t)a_h) \\ &= (q, \nabla \cdot [E_h(t)a_h]) \quad (\text{since } E_h(t)a_h \in X_h \text{ and } q = 0 \text{ on } \partial D) \\ &= (q - q_h, \nabla \cdot [E_h(t)a_h]) \quad (\text{for any } q_h \in H_0^1(D) \cap Q_h) \\ &= -(E_h(t)P_{X_h} \nabla(q - q_h), a_h) \\ &= -((I + A_h)^{\frac{1}{2}} E_h(t)(I + A_h)^{-\frac{1}{2}} P_{X_h} \nabla(q - q_h), a_h), \end{aligned}$$

which implies that

$$\Phi_h(t)\nabla q = -(I + A_h)^{\frac{1}{2}} E_h(t)(I + A_h)^{-\frac{1}{2}} P_{X_h} \nabla(q - q_h) \quad \forall q_h \in Q_h \cap H_0^1(D). \quad (4.52)$$

On the one hand, (4.52) can be combined with (4.21) to yield

$$\begin{aligned} \|\Phi_h(t)\nabla q\|_{L^2} &\lesssim \inf_{q_h \in Q_h \cap H_0^1(D)} t^{-\frac{1}{2}} \|(I + A_h)^{-\frac{1}{2}} P_{X_h} \nabla(q - q_h)\|_{L^2} \lesssim \inf_{q_h \in Q_h \cap H_0^1(D)} t^{-\frac{1}{2}} \|q - q_h\|_{L^2} \\ &\lesssim t^{-\frac{1}{2}} h^2 \|q\|_{H^2}, \end{aligned}$$

where (2.13) is used. On the other hand, inequality (4.52) can be combined with (4.22) to yield

$$\begin{aligned} \left(\int_0^T \|\Phi_h(t)\nabla q\|_{L^2}^2 dt \right)^{\frac{1}{2}} &\lesssim \inf_{q_h \in Q_h \cap H_0^1(D)} \|(I + A_h)^{-\frac{1}{2}} P_{X_h} \nabla(q - q_h)\|_{L^2} \lesssim \inf_{q_h \in Q_h \cap H_0^1(D)} \|q - q_h\|_{L^2} \\ &\lesssim h^2 \|q\|_{H^2}. \end{aligned}$$

Substituting (4.51) into the two inequalities above, we obtain

$$\|\Phi_h(t)\nabla q\|_{L^2} \lesssim t^{-\frac{1}{2}} h^2 \|v\|_{H^1} \quad \text{and} \quad \left(\int_0^t \|\Phi_h(s)\nabla q\|_{L^2}^2 ds \right)^{\frac{1}{2}} \lesssim h^2 \|v\|_{H^1}. \quad (4.53)$$

By using inequality (4.50) with $\rho = 0$ (with v replaced by $P_X v$ therein) and Lemma 4.7 with $g = P_X v$, we have

$$\|\Phi_h(t)P_X v\|_{L^2} \lesssim t^{-\frac{1}{2}} h^2 \|v\|_{H^1} \quad \text{and} \quad \left(\int_0^t \|\Phi_h(s)P_X v\|_{L^2}^2 ds \right)^{\frac{1}{2}} \lesssim h^2 \|v\|_{H^1}. \quad (4.54)$$

Then, combining (4.53) and (4.54) in the decomposition $v = P_X v + \nabla q$, we obtain

$$\|\Phi_h(t)v\|_{L^2} \lesssim t^{-\frac{1}{2}} h^2 \|v\|_{H^1} \quad \text{and} \quad \left(\int_0^t \|\Phi_h(s)v\|_{L^2}^2 ds \right)^{\frac{1}{2}} \lesssim h^2 \|v\|_{H^1}. \quad (4.55)$$

This proves (4.24) and (4.25).

It remains to prove (4.27). To this end, we use the expressions in (4.45) and (4.46), which imply that

$$\begin{aligned} \int_0^t \Phi_h(s)v ds &= \int_0^t \frac{1}{2\pi i} \int_{\Gamma} [(z + A)^{-1} P_X v - (z + A_h)^{-1} P_{X_h} v] e^{zs} dz ds \\ &= \frac{1}{2\pi i} \int_{\Gamma} z^{-1} [(z + A)^{-1} P_X v - (z + A_h)^{-1} P_{X_h} v] e^{zs} dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} z^{-1} [w - w_h] e^{zt} dz. \end{aligned}$$

By applying Lemma 4.6 with $f = v$, we have

$$\begin{aligned} &\left\| \int_0^t \Phi_h(s)v ds \right\|_{L^2} + h \left\| \nabla \int_0^t \Phi_h(s)v ds \right\|_{L^2} \\ &\lesssim \int_{\Gamma} |z|^{-1} (\|w - w_h\|_{L^2} + h \|\nabla w - \nabla w_h\|_{L^2}) e^{\operatorname{Re}(z)t} |dz| \\ &\lesssim h^2 \|v\|_{L^2} \int_{\Gamma} |z|^{-1} e^{\operatorname{Re}(z)t} |dz| \lesssim h^2 \|v\|_{L^2}. \end{aligned} \quad (4.56)$$

This proves (4.27). \square

4.4. The discrete semigroup in the full discretization

Let $\lambda_j^* \geq 0$ and ϕ_j^* , $j = 1, 2, \dots$, be the eigenvalues and eigenfunctions of the operator $A : D(A) \rightarrow X$. Similarly, let $\lambda_{h,j}^* \geq 0$ and $\phi_{h,j}^*$, $j = 1, \dots, M_h$, be the eigenvalues and eigenfunctions of the operator $A_h : X_h \rightarrow X_h$.

We denote by $R(\tau A_h) : X_h \rightarrow X_h$ the linear operator defined by

$$R(\tau A_h)v_h = \sum_{j=1}^{M_h} R(\tau \lambda_{h,j}^*)(v_h, \phi_{h,j}^*) \phi_{h,j}^* \quad \text{with} \quad R(z) = \frac{1}{1+z} \quad \text{for} \quad z \in \mathbb{R}, \quad z \neq -1. \quad (4.57)$$

The discrete semigroup $\bar{E}_{h,\tau}$ defined in (2.23) can be written as

$$\bar{E}_{h,\tau}v_h = R(\tau A_h)v_h \quad \text{for } v_h \in X_h.$$

As a temporal version of (4.21) and (4.22), the following estimates hold for any $v_h \in X_h$,

$$\|(I + A_h)^{\frac{\gamma}{2}} \bar{E}_{h,\tau}^n v_h\|_{L^2} \lesssim t_n^{-\frac{\gamma}{2}} \|v_h\|_{L^2} \quad \text{for } 1 \leq n \leq N \text{ and } \gamma \in [0, 1], \quad (4.58)$$

$$\tau \sum_{j=1}^n \|(I + A_h)^{\frac{1}{2}} \bar{E}_{h,\tau}^j v_h\|_{L^2}^2 \lesssim C \|v_h\|_{L^2}^2 \quad \text{for } 1 \leq n \leq N. \quad (4.59)$$

Remark 4.6. Inequality (4.58) is equivalent to

$$|(1 + \lambda_{h,j}^*)^{\frac{\gamma}{2}} R(\tau \lambda_{h,j}^*)^n| \lesssim t_n^{-\frac{\gamma}{2}} \quad \text{for } \lambda_{h,j}^* \geq 0 \text{ and } t_n = n\tau \in [0, T]. \quad (4.60)$$

The proof of this inequality can be found in [33, Lemma 7.3]). Since the function $w_h^j = \bar{E}_{h,\tau}^j v_h$, $n = 1, 2, \dots$, are solutions of the equation

$$\frac{w_h^j - w_h^{j-1}}{\tau} + A_h w_h^j = 0.$$

Testing the equation with w_h^j and summing up the results for $j = 1, \dots, n$, yield the basic energy inequality

$$\max_{1 \leq j \leq n} \frac{1}{2} \|w_h^j\|_{L^2}^2 + \tau \sum_{j=1}^n \|A_h^{\frac{1}{2}} w_h^j\|_{L^2}^2 \leq \frac{1}{2} \|w_h^0\|_{L^2}^2 = \frac{1}{2} \|v_h\|_{L^2}^2,$$

which implies (4.59). \square

The error of $\bar{E}_{h,\tau}^n P_{X_h}$ in approximating $E_h(t_n)P_{X_h}$ is discussed in the following lemma.

Lemma 4.8. *Let $\bar{\Phi}_{h,\tau}^n v := E_h(t_n)P_{X_h}v - \bar{E}_{h,\tau}^n P_{X_h}v$ for $v \in L^2(D)^d$. Then the following estimates hold:*

$$\|\bar{\Phi}_{h,\tau}^n v\|_{L^2} \lesssim \tau^{\frac{1}{2}} \|v\|_{H^1} \quad \forall v \in H^1(D)^d, \quad (4.61)$$

$$\|\bar{\Phi}_{h,\tau}^n v\|_{L^2} \lesssim t_n^{-\frac{1}{2}} \tau^{\frac{1}{2}} \|v\|_{L^2} \quad \forall v \in L^2(D)^d, \quad (4.62)$$

$$\left(\tau \sum_{j=1}^n \|\bar{\Phi}_{h,\tau}^j v\|_{L^2}^2 \right)^{\frac{1}{2}} + \left\| \tau \sum_{j=1}^n \nabla \bar{\Phi}_{h,\tau}^j v \right\|_{L^2} \lesssim \tau^{\frac{1}{2}} \|v\|_{L^2} \quad \forall v \in L^2(D)^d. \quad (4.63)$$

Proof. It is known that the function $F_n(z) = e^{-nz} - R(z)^n$ has the following upper bound:

$$|F_n(\tau \lambda_{h,j})| \lesssim (\tau \lambda_{h,j}^*)^{\frac{1}{2}} \quad \text{for the eigenvalues } \lambda_{h,j} \geq 0. \quad (4.64)$$

A proof of this result can be found in [33, Theorem 7.1] (with $q = \frac{1}{2}$ therein). Inequality immediately implies the following error estimate:

$$\|(E_h(t_n) - \bar{E}_{h,\tau}^n)v_h\|_{L^2} = \|F_n(\tau A_h)v_h\|_{L^2} \lesssim \tau^{\frac{1}{2}} \|v_h\|_{H^1} \quad \forall v_h \in X_h. \quad (4.65)$$

In view of the H^1 -stability of P_{X_h} in (4.8), we have

$$\|(E_h(t_n) - \bar{E}_{h,\tau}^n)P_{X_h}v\|_{L^2} \lesssim \tau^{\frac{1}{2}} \|P_{X_h}v\|_{H^1} \lesssim \tau^{\frac{1}{2}} \|v\|_{H^1} \quad \forall v \in H^1(D)^d. \quad (4.66)$$

This proves (4.61).

Similarly, the following inequality can be shown (see [33, inequality (7.22)] with $q = \frac{1}{2}$)

$$|F_n(\tau \lambda_{h,j})| \lesssim \tau^{\frac{1}{2}} t_n^{-\frac{1}{2}} \quad \text{for the eigenvalues } \lambda_{h,j} \geq 0,$$

which immediately implies that

$$\|(E_h(t_n) - \bar{E}_{h,\tau}^n)P_{X_h}v\|_{L^2} = \|F_n(\tau A_h)P_{X_h}v\|_{L^2} \lesssim \tau^{\frac{1}{2}} t_n^{-\frac{1}{2}} \|P_{X_h}v\|_{L^2} \quad \forall v \in L^2(D)^d. \quad (4.67)$$

This proves (4.62).

The first and second terms in (4.63) were estimated in [22, inequality (4.21) with $\rho = 0$] and [22, inequality (4.19) with $\rho = 1$, replacing $A_h^{-\frac{1}{2}} P_h x$ by $P_h x$ therein], respectively. \square

Proof of Lemma 4.1. By using the expressions

$$\Phi_h(t) := E(t)P_X - E_h(t)P_{X_h} \quad \text{and} \quad \bar{\Phi}_{h,\tau}^n v := E_h(t_n)P_{X_h}v - \bar{E}_{h,\tau}^n P_{X_h}v,$$

and the triangle inequality, we have

$$\begin{aligned} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(s)P_X - \bar{E}_{h,\tau}^j P_{X_h}]v\|_{L^2}^2 ds &\lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E_h(s) - E_h(t_j)]P_{X_h}v\|_{L^2}^2 ds \\ &+ \int_0^{t_n} \|\Phi_h(s)v\|_{L^2}^2 ds + \tau \sum_{j=1}^n \|\bar{\Phi}_{h,\tau}^j v\|_{L^2}^2. \end{aligned} \quad (4.68)$$

The first term on the right-hand side (4.68) can be estimated as follows:

$$\begin{aligned} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E_h(s) - E_h(t_j)]P_{X_h}v\|_{L^2}^2 ds &= \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|(1 - e^{-(t_j-s)A_h})e^{-sA_h}P_{X_h}v\|_{L^2}^2 ds \\ &\lesssim \tau \int_0^{t_n} \|A_h^{\frac{1}{2}}e^{-sA_h}P_{X_h}v\|_{L^2}^2 ds \\ &\lesssim \tau \|v\|_{L^2}^2 \quad \forall v \in L^2(D)^d. \end{aligned} \quad (4.69)$$

The second and third terms on the right-hand side of (4.68) have already been estimated in (4.25) and (4.63), respectively, which imply (4.1).

Similarly, (4.2) can be proved by using (4.27), (4.63) and (4.69).

Similarly, we have

$$\begin{aligned} \left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \nabla[E(s)P_X - \bar{E}_{h,\tau}^j P_{X_h}]v ds \right\|_{L^2}^2 &\lesssim \left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \nabla[E_h(s) - E_h(t_j)]P_{X_h}v ds \right\|_{L^2}^2 \\ &+ \left\| \int_0^{t_n} \nabla\Phi_h(s)v ds \right\|_{L^2}^2 + \left\| \tau \sum_{j=1}^n \nabla\bar{\Phi}_{h,\tau}^j v \right\|_{L^2}^2 \\ &\lesssim (\tau + h^2) \|v\|_{L^2}^2. \end{aligned} \quad (4.70)$$

The first term on the right-hand side of (4.70) was estimated in [22, p. 236, with $\rho = 1$ and $P_h x$ replaced by $A_h^{\frac{1}{2}}P_h x$], i.e.,

$$\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E_h(s) - E_h(t_j)](I + A_h)^{\frac{1}{2}}P_{X_h}v ds \right\|_{L^2}^2 \lesssim \tau \|P_{X_h}v\|_{L^2}^2 \lesssim \tau \|v\|_{L^2}^2.$$

The second and third terms on the right-hand side of (4.70) have been estimated in (4.27) and (4.63), respectively. This proves (4.3). \square

5. Error analysis for the stochastic problem

In this section, we present error estimates for the fully discrete method (2.19) for the stochastic Stokes equations. Some estimates of the noise term in the Hilbert–Schmidt norm are presented in Subsection 5.1, and the error estimates are presented in Section 5.2.

5.1. Estimates in the Hilbert–Schmidt norm

Lemma 5.1. *Under Assumptions 2.1–2.3, the linear operator $B(v) : L^2(D)^d \rightarrow L^2(D)^d$ satisfies*

$$\|E(t)P_X[B(u) - B(v)]\|_{\mathcal{L}^0}^2 \lesssim t^{-\frac{1}{2}} \mathbb{E} \|u - v\|_{L^2}^2 \quad \forall u, v \in L^2(D)^d, \quad (5.1)$$

and

$$\|B(u)\|_{\mathcal{L}^0}^2 \lesssim 1 + \|u\|_{H^\beta}^2 \quad \forall u \in H^\beta(D)^d, \quad \beta \in \left(\frac{3}{2}, 2\right), \quad (5.2)$$

$$\|(I + A)^{\frac{1}{2}}P_X B(u)\|_{\mathcal{L}_2^0}^2 \lesssim 1 + \|u\|_{H^\beta}^2 \quad \forall u \in H^\beta(D)^d, \quad \beta \in (\frac{3}{2}, 2). \quad (5.3)$$

Proof. By using (4.15) with $s = \frac{1}{2}$, (2.4), (4.16) with $\gamma = \frac{1}{4}$, one can show that

$$\begin{aligned} & \|E(t)P_X(B(u) - B(v))\|_{\mathcal{L}_2^0}^2 \\ &= \sum_{\ell} \mu_{\ell} \|E(t)P_X[(B(u) - B(v))\phi_{\ell}]\|_{L^2}^2 \\ &\lesssim \sum_{\ell} \mu_{\ell} \|(I + A)^{\frac{1}{4}}E(t)(I + A)^{-\frac{1}{4}}P_X[(B(u) - B(v))\phi_{\ell}]\|_{L^2}^2 \\ &\lesssim t^{-\frac{1}{2}} \sum_{\ell} \mu_{\ell} \|(B(u) - B(v))\phi_{\ell}\|_{\dot{H}^{-\frac{1}{2}}}^2 \quad (\text{Here (4.16) with } \gamma = \frac{1}{4} \text{ is used}) \\ &\lesssim t^{-\frac{1}{2}} \sum_{\ell} \mu_{\ell} \|B(u) - B(v)\|_{L^2}^2 \|\phi_{\ell}\|_{H^1}^2 \quad (\text{In addition, (3.4) and (4.15) are used}) \\ &\lesssim t^{-\frac{1}{2}} \|u - v\|_{L^2}^2 (1 + \|(-\Delta)^{\frac{1}{2}}\|_{\mathcal{L}_2^0}^2), \quad (\text{Assumption 2.2 is used}) \end{aligned} \quad (5.4)$$

and

$$\begin{aligned} \|B(u)\|_{\mathcal{L}_2^0}^2 &= \sum_{\ell} \mu_{\ell} \|B(u)\phi_{\ell}\|_{L^2}^2 \leq \sum_{\ell} \mu_{\ell} \|B(u)\phi_{\ell}\|_{H^1}^2 \\ &\lesssim \sum_{\ell} \mu_{\ell} (1 + \|u\|_{H^\beta}^2) \|\phi_{\ell}\|_{H^1}^2 \quad (\text{Assumption 2.2 is used}) \\ &\lesssim (1 + \|u\|_{H^\beta}^2) (1 + \|(-\Delta)^{\frac{1}{2}}\|_{\mathcal{L}_2^0}^2). \end{aligned} \quad (5.5)$$

The last two inequalities and Assumption 2.1 imply (5.1)–(5.2). Similarly,

$$\begin{aligned} \|(I + A)^{\frac{1}{2}}P_X B(u)\|_{\mathcal{L}_2^0}^2 &= \sum_{\ell} \mu_{\ell} \|(I + A)^{\frac{1}{2}}P_X B(u)\phi_{\ell}\|_{L^2}^2 \lesssim \sum_{\ell} \mu_{\ell} \|B(u)\phi_{\ell}\|_{H^1}^2 \\ &\lesssim (1 + \|u\|_{H^\beta}^2) (1 + \|(-\Delta)^{\frac{1}{2}}\|_{\mathcal{L}_2^0}^2). \end{aligned} \quad (5.6)$$

This proves (5.3). \square

Remark 5.1. As a result of the estimates in Lemma 5.1, the regularity results in Theorem 3.1 imply that

$$\sup_{t \in [0, T]} \mathbb{E} \|B(u(t))\|_{\mathcal{L}_2^0}^2 + \sup_{t \in [0, T]} \mathbb{E} \|(I + A)^{\frac{1}{2}}P_X B(u(t))\|_{\mathcal{L}_2^0}^2 \lesssim 1. \quad (5.7)$$

The following stability estimates for the numerical solution can be proved by using Lemma 5.1 and will be used in the error analysis. The proof is omitted here, and the details can be found in Appendix C.

Lemma 5.2. *Under Assumptions 2.1–2.3, the numerical solution determined by the fully discrete method (2.22) satisfies the following energy inequality:*

$$\max_{1 \leq n \leq N} \mathbb{E} \|u_h^n\|_{H^{\frac{1}{2}}}^2 + \sum_{n=1}^N \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2 + \tau \sum_{n=1}^N \mathbb{E} \|u_h^n\|_{H^1}^2 \lesssim 1. \quad (5.8)$$

5.2. Error estimates for the velocity

By iterating (2.22) with respect to n , the full discrete method can be rewritten as

$$u_h^n = \bar{E}_{h,\tau}^n P_{X_h} u_0 + \tau \sum_{i=0}^{n-1} \bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1}) + \sum_{i=0}^{n-1} \bar{E}_{h,\tau}^{n-i} P_{X_h} [B(u_h^i) \Delta W_{i+1}]. \quad (5.9)$$

Then, after subtracting (C.1) from (3.13), we obtain the following error equation:

$$u(t_n) - u_h^n = (E(t_n)P_X - \bar{E}_{h,\tau}^n P_{X_h}) u_0$$

$$\begin{aligned}
& + \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} [E(t_n - s)P_X f(s) - \bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1})] ds \\
& + \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} [E(t_n - s)P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i)] dW(s) \\
& =: T_1 + T_2 + T_3,
\end{aligned} \tag{5.10}$$

which implies that

$$\mathbb{E}\|u(t_n) - u_h^n\|_{L^2}^2 \lesssim \sum_{j=1}^3 \mathbb{E}\|T_j\|_{L^2}^2. \tag{5.11}$$

The three terms are estimated below separately below.

Since $T_1 = \Phi_h(t_n)u^0 + \bar{\Phi}_{h,\tau}^n u^0$, by applying (4.28) and (4.61) with $\rho = 1$ and $v = u_0 \in \mathcal{D}(A)$, we obtain

$$\mathbb{E}\|T_1\|_{L^2}^2 \lesssim \mathbb{E}\|\Phi_h(t_n)u^0\|_{L^2}^2 + \mathbb{E}\|\bar{\Phi}_{h,\tau}^n u^0\|_{L^2}^2 \lesssim (\tau + h^4)\mathbb{E}\|u^0\|_{H^2}^2. \tag{5.12}$$

By using the triangle inequality, T_2 can be further decomposed into the following two parts:

$$\begin{aligned}
\mathbb{E}\|T_2\|_{L^2}^2 & \lesssim \mathbb{E}\left\| \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} E(t_n - s)P_X [f(s) - f(t_{i+1})] ds \right\|_{L^2}^2 \\
& + \mathbb{E}\left\| \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} [E(t_n - s)P_X - \bar{E}_{h,\tau}^{n-i} P_{X_h}] f(t_{i+1}) ds \right\|_{L^2}^2 \\
& =: T_{21} + T_{22}.
\end{aligned} \tag{5.13}$$

By using the Hölder continuity of $f(t)$ in (2.8), we have

$$T_{21} \lesssim \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \mathbb{E}\|f(s) - f(t_{i+1})\|_{L^2}^2 ds \lesssim \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} (t_{i+1} - s) ds \lesssim \tau. \tag{5.14}$$

Through a change of variables $\sigma = t_n - s$ and $j = n - i$, we obtain by using the triangle inequality

$$\begin{aligned}
T_{22} & = \mathbb{E}\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E(\sigma)P_X - \bar{E}_{h,\tau}^j P_{X_h}] f(t_{n-j+1}) d\sigma \right\|_{L^2}^2 \\
& \lesssim \mathbb{E}\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E(\sigma) - E(t_j)] P_X [f(t_{n-j+1}) - f(t_n)] d\sigma \right\|_{L^2}^2 \\
& + \mathbb{E}\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E(t_j)P_X - \bar{E}_{h,\tau}^j P_{X_h}] [f(t_{n-j+1}) - f(t_n)] d\sigma \right\|_{L^2}^2 \\
& + \mathbb{E}\left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} [E(\sigma)P_X - \bar{E}_{h,\tau}^j P_{X_h}] f(t_n) d\sigma \right\|_{L^2}^2 \\
& =: T_{22}^a + T_{22}^b + T_{22}^c.
\end{aligned} \tag{5.15}$$

The term T_{22}^a can be estimated by using (4.16) with $\gamma = \frac{1}{2}$, (4.17) with $\mu = \frac{1}{2}$, and the Hölder continuity of f in (2.8), i.e.,

$$\begin{aligned}
T_{22}^a & \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|(I + A)^{\frac{1}{2}} E(\sigma) (I + A)^{-\frac{1}{2}} [I - E(t_j - \sigma)] P_X [f(t_{n-j+1}) - f(t_n)]\|_{L^2}^2 d\sigma \\
& \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-1} (t_j - \sigma) \|f(t_{n-j+1}) - f(t_n)\|_{L^2}^2 d\sigma
\end{aligned} \tag{5.16}$$

$$\lesssim \tau \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-1} (t_n - t_{n-j+1}) d\sigma \lesssim \tau \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-1} d\sigma \lesssim \tau.$$

The term T_{22}^b can be estimated by using (4.26), (4.62) and the Assumption 2.8 shows

$$\begin{aligned} T_{22}^b &\lesssim \mathbb{E} \left(\sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[\Phi_h(t_j) + \bar{\Phi}_{h,\tau}^j][f(t_{n-j+1}) - f(t_n)]\|_{L^2} d\sigma \right)^2 \\ &\lesssim \mathbb{E} \left(\sum_{j=1}^n \int_{t_{j-1}}^{t_j} (t_j^{-1} h^2 + t_j^{-\frac{1}{2}} \tau^{\frac{1}{2}}) \|f(t_{n-j+1}) - f(t_n)\|_{L^2} d\sigma \right)^2 \\ &\lesssim (\tau + h^4) \left(\sum_{j=1}^n \int_{t_{j-1}}^{t_j} t_j^{-1} (t_n - t_{n-j+1})^{\frac{1}{2}} d\sigma \right)^2 \\ &\lesssim \tau + h^4. \end{aligned} \quad (5.17)$$

The term T_{22}^c can be estimated by applying (4.2) directly, i.e.,

$$T_{22}^c \lesssim (\tau + h^4) \|f(t_n)\|_{L^2}^2 \lesssim \tau + h^4. \quad (5.18)$$

Substituting (5.14)–(5.18) into (5.13) yields

$$\mathbb{E} \|T_2\|_{L^2}^2 \lesssim \tau + h^4. \quad (5.19)$$

It remains to estimate the term T_3 in (5.10). By using (2.2) and a change of variables $\sigma = t_n - s$ and $j = n - i$, we can decompose T_3 into three parts as follows:

$$\begin{aligned} \mathbb{E} \|T_3\|_{L^2}^2 &= \mathbb{E} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \|E(t_n - s) P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i)\|_{\mathcal{L}_2^0}^2 ds \\ &= \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(\sigma) P_X B(u(t_n - \sigma)) - \bar{E}_{h,\tau}^j P_{X_h} B(u_h^{n-j})]\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\lesssim \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|E(\sigma) P_X [B(u(t_n - \sigma)) - B(u(t_{n-j}))]\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\quad + \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(\sigma) P_X - \bar{E}_{h,\tau}^j P_{X_h}] B(u(t_{n-j}))\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\quad + \tau \mathbb{E} \sum_{j=1}^n \|\bar{E}_{h,\tau}^j P_{X_h} [B(u(t_{n-j})) - B(u_h^{n-j})]\|_{\mathcal{L}_2^0}^2 \\ &=: T_{31} + T_{32} + T_{33}. \end{aligned} \quad (5.20)$$

The three terms T_{31} , T_{32} and T_{33} are estimated separately below.

The term T_{31} can be estimated by using (5.1) and Hölder continuity (3.15), which imply that

$$\begin{aligned} T_{31} &\leq \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-\frac{1}{2}} \mathbb{E} \|u(t_n - \sigma) - u(t_{n-j})\|_{L^2}^2 d\sigma \\ &\leq C(1 + \mathbb{E} \|u^0\|_{H^2}^2) \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-\frac{1}{2}} (t_j - \sigma) d\sigma \\ &\leq C\tau(1 + \mathbb{E} \|u^0\|_{H^2}^2). \end{aligned} \quad (5.21)$$

The term T_{32} can be estimated by decomposing it into three parts, i.e.,

$$T_{32} \lesssim \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(\sigma) - E(t_j)] P_X [B(u(t_{n-j})) - B(u(t_{n-1}))]\|_{\mathcal{L}_2^0}^2 d\sigma \quad (5.22)$$

$$\begin{aligned}
& + \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(t_j)P_X - \bar{E}_{h,\tau}^j P_{X_h}][B(u(t_{n-j})) - B(u(t_{n-1}))]\|_{\mathcal{L}_2^0}^2 d\sigma \\
& + \mathbb{E} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|[E(\sigma)P_X - \bar{E}_{h,\tau}^j P_{X_h}]B(u(t_{n-1}))\|_{\mathcal{L}_2^0}^2 d\sigma \\
& =: T_{32}^a + T_{32}^b + T_{32}^c.
\end{aligned}$$

The term T_{32}^a can be estimated by using (4.16) with $\gamma = \frac{1}{2}$, (4.17) with $\mu = \frac{1}{2}$, (5.2) and Hölder continuity (3.15), i.e.,

$$\begin{aligned}
T_{32}^a & \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \mathbb{E} \|E(\sigma)[I - E(t_j - \sigma)]P_X[B(u(t_{n-j})) - B(u(t_{n-1}))]\|_{\mathcal{L}_2^0}^2 d\sigma \\
& \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \mathbb{E} \sum_{\ell} \mu_{\ell} \|(I + A)^{\frac{1}{4}} E(\sigma)[I - E(t_j - \sigma)](I + A)^{-\frac{1}{4}} P_X[B(u(t_{n-j})) - B(u(t_{n-1}))]\phi_{\ell}\|_{L^2}^2 d\sigma \\
& \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-\frac{5}{4}}(t_j - \sigma) \mathbb{E} \sum_{\ell} \mu_{\ell} \|B(u(t_{n-j})) - B(u(t_{n-1}))\phi_{\ell}\|_{\dot{H}^{-\frac{1}{2}}}^2 d\sigma \\
& \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-\frac{5}{4}}(t_j - \sigma) \mathbb{E} \|u(t_{n-j}) - u(t_{n-1})\|_{L^2}^2 d\sigma \quad (\text{Assumption 2.2 is used}) \\
& \lesssim \tau(1 + \mathbb{E}\|u^0\|_{H^2}^2) \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \sigma^{-\frac{5}{4}}(t_{n-1} - t_{n-j}) d\sigma \\
& \lesssim \tau(1 + \mathbb{E}\|u^0\|_{H^2}^2). \tag{5.23}
\end{aligned}$$

The term T_{32}^b can be estimated by using the expression $E(t_j)P_X - \bar{E}_{h,\tau}^j P_{X_h} = \Phi_h(t_j) + \bar{\Phi}_{h,\tau}^j$, i.e.,

$$\begin{aligned}
T_{32}^b & = \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \mathbb{E} \|[\Phi_h(t_j) + \bar{\Phi}_{h,\tau}^j][B(u(t_{n-j})) - B(u(t_{n-1}))]\|_{\mathcal{L}_2^0}^2 d\sigma \tag{5.24} \\
& \lesssim \sum_{j=1}^n \int_{t_{j-1}}^{t_j} t_j^{-1} h^4 \mathbb{E} \|(I + A)^{\frac{1}{2}} P_X[B(u(t_{n-j})) - B(u(t_{n-1}))]\|_{\mathcal{L}_2^0}^2 d\sigma \quad (\text{here (4.24) is used}) \\
& \quad + \sum_{j=1}^n \int_{t_{j-1}}^{t_j} t_j^{-1} \tau \mathbb{E} \|B(u(t_{n-j})) - B(u(t_{n-1}))\|_{\mathcal{L}_2^0}^2 d\sigma \quad (\text{here (4.62) is used}) \\
& \lesssim (\tau + h^4) \sum_{j=1}^n \int_{t_{j-1}}^{t_j} t_j^{-1} \mathbb{E} \|u(t_{n-j}) - u(t_{n-1})\|_{H^{\beta}}^2 d\sigma \quad (\text{Assumption 2.2 is used}) \\
& \lesssim (\tau + h^4) (1 + \mathbb{E}\|u^0\|_{H^2}^2) \sum_{j=1}^n \int_{t_{j-1}}^{t_j} t_j^{-1} (t_{n-1} - t_{n-j})^{2-\beta} d\sigma \quad (\text{here (3.16) is used}) \\
& \lesssim (\tau + h^4) (1 + \mathbb{E}\|u^0\|_{H^2}^2).
\end{aligned}$$

The T_{32}^c term can be estimated by directly applying (4.1) and (5.7), we have

$$T_{32}^c \lesssim (\tau + h^4) \sup_{t \in [0, T]} \mathbb{E} \|(I + A)^{\frac{1}{2}} P_X B(u(t))\|_{\mathcal{L}_2^0}^2 \lesssim \tau + h^4. \tag{5.25}$$

The term T_{33} in (5.20) can be estimated by using (4.58) with $\gamma = \frac{1}{2}$, i.e.,

$$\mathbb{E} \|T_{33}\|_{L^2}^2 \lesssim \tau \sum_{j=1}^n \mathbb{E} \|(I + A_h)^{\frac{1}{4}} \bar{E}_{h,\tau}^j (I + A_h)^{-\frac{1}{4}} P_{X_h} [B(u(t_{n-j})) - B(u_h^{n-j})]\|_{\mathcal{L}_2^0}^2$$

$$\begin{aligned}
&\lesssim \tau \sum_{j=1}^n t_j^{-\frac{1}{2}} \mathbb{E} \sum_{\ell} \mu_{\ell} \|P_{X_h}[(B(u(t_{n-j})) - B(u_h^{n-j}))\phi_{\ell}]\|_{\dot{H}^{-\frac{1}{2}}}^2 \\
&\lesssim \tau \sum_{j=1}^n t_j^{-\frac{1}{2}} \mathbb{E} \|u(t_{n-j}) - u_h^{n-j}\|_{L^2}^2 \|(-\Delta)^{\frac{1}{2}}\|_{\mathcal{L}^0}^2 \\
&\lesssim \tau \sum_{i=0}^{n-1} (t_n - t_i)^{-\frac{1}{2}} \mathbb{E} \|u(t_i) - u_h^i\|_{L^2}^2.
\end{aligned} \tag{5.26}$$

In view of (5.20)–(5.26), we have

$$\mathbb{E} \|T_3\|_{L^2}^2 \lesssim \tau + h^4 + \tau \sum_{i=0}^{n-1} (t_n - t_i)^{-\frac{1}{2}} \mathbb{E} \|u(t_i) - u_h^i\|_{L^2}^2. \tag{5.27}$$

Substituting (5.12), (5.19) and (5.27) into (5.11) yields

$$\mathbb{E} \|u(t_n) - u_h^n\|_{L^2}^2 \lesssim \tau + h^4 + \tau \sum_{i=0}^{n-1} (t_n - t_i)^{-\frac{1}{2}} \mathbb{E} \|u(t_i) - u_h^i\|_{L^2}^2. \tag{5.28}$$

By using the discrete Gronwall inequality, we obtain the desired error estimate in (2.24) for the velocity.

5.3. Error estimates for the pressure.

From the numerical scheme in (2.19) we can derive that

$$\begin{aligned}
\tau \sum_{n=1}^m (p_h^n, \nabla \cdot v_h) &= (u_h^m, v_h) - (u_h^0, v_h) + 2\tau \sum_{n=1}^m (\mathbb{D}(u_h^n), \mathbb{D}(v_h)) \\
&\quad - \tau \sum_{n=1}^m (f(t_n), v_h) - \tau \sum_{n=1}^m (B(u_h^{n-1}) \Delta W_n, v_h) \quad v_h \in V_h.
\end{aligned} \tag{5.29}$$

Subtracting (5.29) from (3.17) yields

$$\begin{aligned}
\left(\int_0^{t_m} p(t) dt - \tau \sum_{n=1}^m p_h^n, \nabla \cdot v_h \right) &= ([u(t_m) - u_h^m] - [u^0 - u_h^0], v_h) \\
&\quad + 2 \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}(u(t) - u_h^n) dt, \mathbb{D}(v_h) \right) \\
&\quad - \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} [f(t) - f(t_n)] dt, v_h \right) \\
&\quad - \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} [B(u(t)) - B(u_h^{n-1})] dW(t), v_h \right) \\
&=: J_1(v_h) + J_2(v_h) + J_3(v_h) + J_4(v_h) \quad \forall v_h \in V_h,
\end{aligned}$$

where

$$\begin{aligned}
|J_1(v_h)| &\lesssim (\|u(t_m) - u_h^m\|_{L^2} + \|u^0 - P_{X_h} u^0\|_{L^2}) \|v_h\|_{L^2} =: J_1^* \|v_h\|_{L^2}, \\
|J_2(v_h)| &\leq \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}(u(t) - u_h^n) dt \right\|_{L^2} \|\nabla v_h\|_{L^2} =: J_2^* \|\nabla v_h\|_{L^2} \\
|J_3(v_h)| &\lesssim \int_0^{t_m} \|f(t) - f(t_n)\|_{L^2} dt \|v_h\|_{L^2} =: J_3^* \|v_h\|_{L^2} \\
|J_4(v_h)| &\lesssim \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} P_{X_h} [B(u(t)) - B(u_h^{n-1})] dW(t) \right\|_{\dot{H}^{-1}} \|v_h\|_{H^1} =: J_4^* \|v_h\|_{H^1},
\end{aligned}$$

which imply that (by using the inf-sup condition (2.18) of the finite element space)

$$\mathbb{E} \left\| \int_0^{t_m} p(t) dt - \tau \sum_{n=1}^m p_h^n \right\|_{L^2}^2 = \mathbb{E}|J_1^*|^2 + \mathbb{E}|J_2^*|^2 + \mathbb{E}|J_3^*|^2 + \mathbb{E}|J_4^*|^2. \quad (5.30)$$

By using (2.24) and (4.10), we have

$$\mathbb{E}|J_1^*|^2 \lesssim (\tau + h^4). \quad (5.31)$$

and

$$\begin{aligned} J_2^* &\leq \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}(u(t) - u(t_n)) dt \right\|_{L^2} + \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}(u(t_n) - u_h^n) dt \right\|_{L^2} \\ &\leq \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}(u(t) - u(t_n)) dt \right\|_{L^2} \quad (\text{using (3.13) with } t = t_n \text{ and (C.1)}) \\ &\quad + \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{D}([E(t_n)P_X - \bar{E}_{h,\tau}^n P_{X_h}]u^0) dt \right\|_{L^2} \\ &\quad + \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \mathbb{D}(E(t_n - s)P_X f(s) - \bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1})) ds dt \right\|_{L^2} \\ &\quad + \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \mathbb{D} \left(\int_{t_i}^{t_{i+1}} E(t_n - s)P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i) dW(s) \right) dt \right\|_{L^2} \\ &=: J_{21}^* + J_{22}^* + J_{23}^* + J_{24}^*, \end{aligned} \quad (5.32)$$

where

$$\begin{aligned} \mathbb{E}|J_{21}^*|^2 &\lesssim \int_0^{t_n} \mathbb{E} \|u(t) - u(t_n)\|_{H^1}^2 dt \\ &\lesssim \sum_{n=1}^m \int_{t_{n-1}}^{t_n} (1 + \mathbb{E} \|u^0\|_{H^2}^2) (t - t_n) dt \quad (\text{here (3.15) is used}) \\ &\lesssim \tau, \\ \mathbb{E}|J_{22}^*|^2 &\lesssim \mathbb{E} \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \nabla [E(t)P_X - \bar{E}_{h,\tau}^n P_{X_h}] u^0 dt \right\|_{L^2}^2 + \mathbb{E} \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \nabla [E(t_n) - E(t)] P_X u^0 dt \right\|_{L^2}^2 \\ &\lesssim \tau + h^2 \quad (\text{here (4.3) is used}) \\ &\quad + \mathbb{E} \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \|(I + A)^{-\frac{1}{2}} [E(t_n - t) - I] (I + A)^{\frac{1}{4}} E(t) (I + A)^{\frac{3}{4}} u^0\|_{L^2}^2 dt \\ &\lesssim \tau + h^2 + \mathbb{E} \int_0^{t_m} \tau \|(I + A)^{\frac{1}{4}} E(t) (I + A)^{\frac{3}{4}} u^0\|_{L^2}^2 dt \quad (\text{here (4.17) is used}) \\ &\lesssim \tau + h^2 + \mathbb{E} \int_0^{t_m} \tau t^{-\frac{1}{2}} \|(I + A)^{\frac{3}{4}} u^0\|_{L^2}^2 dt \\ &\lesssim \tau + h^2, \\ \mathbb{E}|J_{23}^*|^2 &\lesssim \mathbb{E} \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \nabla E(t_n - s) P_X [f(s) - f(t_{i+1})] ds dt \right\|_{L^2}^2 \\ &\quad + \mathbb{E} \left\| \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \nabla [E(t_n - s) P_X - \bar{E}_{h,\tau}^{n-i} P_{X_h}] f(t_{i+1}) ds dt \right\|_{L^2}^2 \\ &=: \mathbb{E}|J_{231}^*|^2 + \mathbb{E}|J_{232}^*|^2. \end{aligned} \quad (5.33)$$

The two terms $\mathbb{E}|J_{231}^*|^2$ and $\mathbb{E}|J_{232}^*|^2$ can be estimated as follows:

$$\begin{aligned}
\mathbb{E}|J_{231}^*|^2 &\lesssim \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \|(I+A)^{\frac{1}{2}} E(t_n-s) P_X [f(s) - f(t_{i+1})]\|_{L^2} ds dt \right)^2 \quad (5.34) \\
&\lesssim \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} (t_n-s)^{-\frac{1}{2}} \mathbb{E} \|f(t_{i+1}) - f(s)\|_{L^2} ds dt \right)^2 \\
&\lesssim \left(\int_0^{t_m} \int_0^{t_n} (t_n-s)^{-\frac{1}{2}} \tau^{\frac{1}{2}} ds dt \right)^2 \\
&\lesssim \tau
\end{aligned}$$

and, by using a change of variables $\sigma = t_n - s$ and $j = n - i$,

$$\begin{aligned}
\mathbb{E}|J_{232}^*|^2 &= \mathbb{E} \left\| \tau \sum_{i=0}^{m-1} \sum_{n=i+1}^m \int_{t_i}^{t_{i+1}} \nabla [E(t_n-s) P_X - \bar{E}_{h,\tau}^{n-i} P_{X_h}] f(t_{i+1}) ds \right\|_{L^2}^2 \quad (5.35) \\
&= \mathbb{E} \left\| \tau \sum_{i=0}^{m-1} \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla [E(\sigma) P_X - \bar{E}_{h,\tau}^j P_{X_h}] f(t_{i+1}) d\sigma \right\|_{L^2}^2 \\
&\lesssim \tau \sum_{i=0}^{m-1} \mathbb{E} \left\| \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla [E(\sigma) P_X - \bar{E}_{h,\tau}^j P_{X_h}] f(t_{i+1}) d\sigma \right\|_{L^2}^2 \\
&\lesssim \tau \sum_{i=0}^{m-1} (\tau + h^2) \mathbb{E} \|f(t_{i+1})\|_{L^2}^2 \quad (\text{here (4.3) is used}) \\
&\lesssim \tau + h^2.
\end{aligned}$$

It remains to estimate the term J_{24}^* in (5.32). Since the integrand in the expression of J_{24}^* is independent of t for $t \in (t_{n-1}, t_n]$, it follows that the integration with respect to t is actually a summation times τ . By interchanging the order of integration in the expression of J_{24}^* and using Itô's isometry in (2.2), we obtain

$$\begin{aligned}
\mathbb{E}|J_{24}^*|^2 &= \mathbb{E} \left\| \mathbb{D} \left(\sum_{n=1}^m \int_{t_{n-1}}^{t_n} \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} E(t_n-s) P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i) dW(s) dt \right) \right\|_{L^2}^2 \\
&= \mathbb{E} \left\| \mathbb{D} \left(\sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \sum_{n=i+1}^m \int_{t_{n-1}}^{t_n} [E(t_n-s) P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i)] dt dW(s) \right) \right\|_{L^2}^2 \\
&\lesssim \mathbb{E} \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left\| \sum_{n=i+1}^m \int_{t_{n-1}}^{t_n} \nabla [E(t_n-s) P_X B(u(s)) - \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i)] dt \right\|_{\mathcal{L}_2^0}^2 ds.
\end{aligned}$$

By using a change of variables $\sigma = t - t_i$ and $j = n - i$, we split $\mathbb{E}|J_{24}^*|^2$ into three parts as follows, by using the triangle inequality,

$$\begin{aligned}
\mathbb{E}|J_{24}^*|^2 &= \mathbb{E} \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left\| \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla [E(t_{i+j}-s) P_X B(u(s)) - \bar{E}_{h,\tau}^j P_{X_h} B(u_h^i)] d\sigma \right\|_{\mathcal{L}_2^0}^2 ds \\
&\lesssim \mathbb{E} \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left\| \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla [E(t_{i+j}-s) - E(\sigma)] P_X B(u(s)) d\sigma \right\|_{\mathcal{L}_2^0}^2 ds \\
&\quad + \mathbb{E} \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left\| \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla E(\sigma) P_X [B(u(s)) - B(u_h^i)] d\sigma \right\|_{\mathcal{L}_2^0}^2 ds \\
&\quad + \tau \mathbb{E} \sum_{i=0}^{m-1} \left\| \sum_{j=1}^{m-i} \int_{t_{j-1}}^{t_j} \nabla [E(\sigma) P_X - \bar{E}_{h,\tau}^j P_{X_h}] B(u_h^i) d\sigma \right\|_{\mathcal{L}_2^0}^2
\end{aligned}$$

$$=: \mathbb{E}|J_{241}^*| + \mathbb{E}|J_{242}^*| + \mathbb{E}|J_{243}^*|.$$

In order to estimate the term $\mathbb{E}|J_{241}^*|$, we first note that for $\xi_j, \sigma \in (t_{j-1}, t_j)$ we have $|\xi_j - \sigma| \leq \tau$ and therefore, by using the triangle inequality,

$$\begin{aligned} & \sum_{j=1}^m \int_{t_{j-1}}^{t_j} \|[E(\xi_j) - E(\sigma)]\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma \\ &= \int_0^{2\tau} \|[E(\xi_j) - E(\sigma)]\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma + \sum_{j=3}^m \int_{t_{j-1}}^{t_j} \|[E(\xi_j) - E(\sigma - \tau)] - [E(\sigma) - E(\sigma - \tau)]\| \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\lesssim \tau \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 + \sum_{j=3}^m \int_{t_{j-1}}^{t_j} \|(I+A)^{-1}[E(\xi_j + \tau - \sigma) - I](I+A)E(\sigma - \tau)\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\quad + \sum_{j=3}^m \int_{t_{j-1}}^{t_j} \|(I+A)^{-1}[E(\tau) - I](I+A)E(\sigma - \tau)\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma \\ &\lesssim \tau \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 + \tau^2 \int_{2\tau}^{t_m} \|(I+A)E(\sigma - \tau)\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 d\sigma \quad (\text{here (4.17) with } \mu = 1 \text{ is used}) \\ &\lesssim \tau \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 + \tau^2 \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2 \int_{2\tau}^{t_m} (\sigma - \tau)^{-2} d\sigma \quad (\text{here (4.16) with } \gamma = 1 \text{ is used}) \\ &\lesssim \tau \|\widetilde{v}(s)\|_{\mathcal{L}_2^0}^2. \end{aligned}$$

Then, the term $\mathbb{E}|J_{241}^*|$ can be estimated by applying the above estimate with $\xi = t_{i+j} - s$ and $\widetilde{v}(s) = (I+A)^{\frac{1}{2}}P_X B(u(s))$, i.e.,

$$\mathbb{E}|J_{241}^*| \lesssim \tau \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \mathbb{E}\|(I+A)^{\frac{1}{2}}P_X B(u(s))\|_{\mathcal{L}_2^0}^2 ds \lesssim \tau. \quad (\text{here (5.7) is used})$$

The term $\mathbb{E}|J_{242}^*|$ can be estimated by using a variable transform $t_{m-i} - \sigma = t$, i.e.,

$$\begin{aligned} \mathbb{E}|J_{242}^*| &\lesssim \mathbb{E} \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left\| (I+A)^{\frac{3}{4}} \int_0^{t_{m-i}} E(t_{m-i} - t)(I+A)^{-\frac{1}{4}} P_X [B(u(s)) - B(u_h^i)] dt \right\|_{\mathcal{L}_2^0}^2 ds \\ &\lesssim \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \left(\int_0^{t_{m-i}} (t_{m-i} - t)^{-\frac{3}{4}} dt \right)^2 \mathbb{E}\|(I+A)^{-\frac{1}{4}} P_X [B(u(s)) - B(u_h^i)]\|_{\mathcal{L}_2^0}^2 ds \\ &\hspace{15em} (\text{here (4.19) with } \rho = \frac{3}{4} \text{ is used}) \\ &\lesssim \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \mathbb{E}\|(I+A)^{-\frac{1}{4}} P_X [B(u(s)) - B(u_h^i)]\|_{\mathcal{L}_2^0}^2 ds \\ &\lesssim \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \mathbb{E} \sum_{\ell} \mu_{\ell} \| [B(u(s)) - B(u_h^i)] \phi_{\ell} \|_{\dot{H}^{-\frac{1}{2}}}^2 ds \quad (\text{here (3.4) with } s = \frac{1}{2} \text{ is used}) \\ &\lesssim \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} \mathbb{E}\|u(s) - u_h^i\|_{L^2}^2 ds \quad (\text{here (2.3) and (2.5) are used}) \\ &\lesssim \sum_{i=0}^{m-1} \int_{t_i}^{t_{i+1}} (\mathbb{E}\|u(s) - u(t_i)\|_{L^2}^2 + \mathbb{E}\|u(t_i) - u_h^i\|_{L^2}^2) ds \\ &\lesssim \tau + \max_{1 \leq n \leq N} \mathbb{E}\|u(t_n) - u_h^n\|_{L^2}^2 \quad (\text{here (3.15) is used}) \\ &\lesssim \tau. \quad (\text{here (2.24) is used, which is already proved}) \end{aligned} \tag{5.36}$$

The term $\mathbb{E}|J_{243}^*|$ can be estimated by directly using (4.3), we obtain

$$\begin{aligned} \mathbb{E}|J_{243}^*| &\lesssim \tau \sum_{i=0}^{m-1} (\tau + h^2) \mathbb{E} \|B(u_h^i)\|_{\mathcal{L}_2^0}^2 \\ &\lesssim (\tau + h^2) \sum_{i=0}^{m-1} \tau (1 + \mathbb{E} \|u_h^i\|_{H^{\frac{1}{2}}}^2) \quad (\text{here (5.5) is used}) \\ &\lesssim \tau + h^2. \quad (\text{here (5.8) is used}) \end{aligned} \quad (5.37)$$

The estimates for J_{21}^* , J_{22}^* , J_{23}^* , J_{24}^* above imply that

$$\mathbb{E}|J_2^*|^2 \lesssim \tau + h^2. \quad (5.38)$$

The term J_3 can be estimated directly by using the Hölder continuity of f in time, i.e.,

$$\mathbb{E}|J_3^*|^2 \lesssim \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{E} \|f(t) - f(t_n)\|_{L^2}^2 dt \lesssim \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \tau dt \lesssim \tau. \quad (5.39)$$

The term J_4 can be estimated directly by using Assumptions 2.1 and 2.2, i.e.,

$$\begin{aligned} \mathbb{E}|J_4^*|^2 &\lesssim \mathbb{E} \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \|(I + A_h)^{-\frac{1}{2}} P_{X_h} [B(u(t)) - B(u_h^{n-1})]\|_{\mathcal{L}_2^0}^2 dt \\ &\lesssim \sum_{n=1}^m \int_{t_{n-1}}^{t_n} \mathbb{E} \|u(t) - u_h^{n-1}\|_{L^2}^2 \quad (\text{here (2.3) and (2.5) are used}) \\ &\lesssim \sum_{n=1}^m \int_{t_{n-1}}^{t_n} (\mathbb{E} \|u(t) - u(t_{n-1})\|_{L^2}^2 + \mathbb{E} \|u(t_{n-1}) - u_h^{n-1}\|_{L^2}^2) \\ &\lesssim \tau \quad (\text{here (3.15) and (2.24) are used}). \end{aligned} \quad (5.40)$$

By substituting (5.31) and (5.38)–(5.40) into (5.30), we obtain

$$\mathbb{E} \left\| \int_0^{t_m} p(t) dt - \tau \sum_{n=1}^m p_h^n \right\|_{L^2}^2 \lesssim \tau + h^2.$$

This proves desired error estimate in (2.25) for the pressure. \square

6. Numerical experiments

In this section, we present numerical tests to support the theoretical analysis in Theorems 2.4 by illustrating the convergence of the fully discrete finite element solutions. For a stable discretization in space, we use the prototypical MINI element; cf. [2, 18] for details. All the computations are performed using the software package NGSolve (<https://ngsolve.org/>).

We solve the stochastic Stokes equations in the two-dimensional square $[0, 1] \times [0, 1]$ under the stress boundary condition by the proposed scheme (2.19), up to time $T = 2$, with $u_0 = (0, 0)$ and $f = (1, 1)^\top$. The noise term $B(u)dW$ is determined by

$$B(u) = \begin{pmatrix} \sqrt{u_1^2 + 1} & 0 \\ \sqrt{u_2^2 + 1} & 0 \end{pmatrix}, \quad (6.1)$$

$$W(t, x_1, x_2) = \sum_{\ell_1=1}^{\infty} \sum_{\ell_2=1}^{\infty} \sqrt{\mu_{\ell_1 \ell_2}} \begin{pmatrix} \phi_{\ell_1 \ell_2}(x_1, x_2) \\ 0 \end{pmatrix} w_{\ell_1 \ell_2}(t) \quad \forall t \in [0, T], \quad (6.2)$$

where $w_{\ell_1 \ell_2}(t)$, with $\ell_1, \ell_2 = 1, 2, \dots$, are independent \mathbb{R} -valued Wiener processes, $\mu_{\ell_1 \ell_2} = (\ell_1^2 + \ell_2^2)^{-(2+\varepsilon)}$ with $\varepsilon = 0.1$, and

$$\phi_{\ell_1 \ell_2} = 2 \sin(\ell_1 \pi x_1) \sin(\ell_2 \pi x_2), \quad \ell_1, \ell_2 = 1, 2, \dots,$$

form an orthonormal basis of $L^2(D)$. In particular, the noise term $B(u)dW$ is non-solenoidal and the Q -Wiener process $W(t)$ satisfies Assumption 2.1.

Since the exact solution of the problem is unknown, we compute the errors of the numerical solutions approximately by

$$\text{Error in time} = \left(\mathbb{E} \|v_{h,\tau}^N - v_{h,\tau/2}^N\|_{L^2}^2 \right)^{\frac{1}{2}} \quad \text{for } v = u \text{ and } p,$$

$$\text{Error in space} = \left(\mathbb{E} \|v_{h,\tau}^N - v_{h/2,\tau}^N\|_{L^2}^2 \right)^{\frac{1}{2}} \quad \text{for } v = u \text{ and } p,$$

where $v_{h,\tau}^N$ denotes the one-path simulation at time $t_N = T$ computed by using stepsize τ and mesh size h . The series in (6.1) is truncated to a finite sum of L^d terms (with $L = h^{-1}$), i.e., its temporal increment is computed by

$$W^L(t_{n+1}, x_1, x_2) - W^L(t_n, x_1, x_2) = \tau \sum_{\ell_1=1}^L \sum_{\ell_2=1}^L \sqrt{\mu_{\ell_1 \ell_2}} \begin{pmatrix} \phi_{\ell_1 \ell_2}(x_1, x_2) \\ 0 \end{pmatrix} \xi_{\ell_1 \ell_2}^n,$$

where $\xi_{\ell_1 \ell_2}^n \sim N(0, 1)$. The expectations of the errors are approximated by computing the averages over M samples, with $M = 1024, 2048, 4096$, respectively.

The errors of the numerical solutions are presented in Figures 1 and 2, where we can see that the proposed method has half-order convergence in time for both velocity and pressure approximations, second-order convergence in space for the velocity approximation, and first-order convergence in space for the pressure approximation. Therefore, the convergence orders observed in the numerical experiments are consistent with the theoretical results proved in Theorem 2.4.




FIGURE 1. Time errors of the numerical solutions at $T = 2$ with $h = 2^{-6}$.




FIGURE 2. Spatial errors of the numerical solutions at $T = 2$ with $\tau = 2^{-8}$.

7. Conclusion

We have proved higher-order strong convergence of fully discrete mixed FEMs for the stochastic Stokes equations under the stress boundary condition driven by a stochastic noise satisfying condition (1.4). The error estimates of $O(\tau^{\frac{1}{2}} + h^2)$ and $O(\tau^{\frac{1}{2}} + h)$ are proved for the velocity and pressure approximations, respectively, for a semi-implicit mixed finite element method. The analysis is based on new estimates of the semi-discrete and fully discrete semi-groups associated to the Stokes operator, and the H^1 -stability of the orthogonal projection onto the discrete divergence-free finite element subspace, as shown in Section 4. The improved convergence orders are consistent with the numerical experiments.

For the simplicity of illustration, we have focused on the stochastic Stokes equations in this article. However, the methodology introduced in this article may also be extended to the stochastic NS equations to obtain higher-order convergence in space.

Appendix A: Well-posedness and regularity of the mild solution

In this Appendix we prove Proposition 3.1, including existence and uniqueness of a mild solution to the stochastic Stokes equations Assumptions 2.1–2.3, and the regularity of the mild solution.

Existence and uniqueness: Let $Y = C([0, T]; L^2(\Omega; H)) \cap L^2(\Omega, \{\mathcal{F}_t\}_{t \geq 0}; L^2(0, T; H))$, i.e., the predictable subspace of $C([0, T]; L^2(\Omega; H))$. For any $v \in Y$ we denote by Mv the function defined by

$$Mv(t) = E(t)u^0 + \int_0^t E(t-s)P_X f(s)ds + \int_0^t E(t-s)P_X B(v(s))dW(s) \quad \text{for } t \in [0, T], \quad (\text{A.1})$$

Under Assumptions (2.1)–(2.3), the following results hold, which are simple modifications of (5.1):

$$\|E(t)P_X B(v)\|_{\mathcal{L}_0^2}^2 \lesssim t^{-\frac{1}{2}}(1 + \mathbb{E}\|v\|_{L^2}^2) \quad t > 0, \quad (\text{A.2})$$

$$\|E(t)P_X [B(u) - B(v)]\|_{\mathcal{L}_0^2}^2 \lesssim t^{-\frac{1}{2}} \mathbb{E}\|u - v\|_{L^2}^2 \quad t > 0. \quad (\text{A.3})$$

Then Mv is predictable and

$$\begin{aligned} \mathbb{E}\|Mv(t)\|_{L^2}^2 &\lesssim \mathbb{E}\|u^0\|_{L^2}^2 + \mathbb{E} \int_0^t \|f(s)\|_{L^2}^2 ds + \mathbb{E} \int_0^t \|E(t-s)P_X B(v(s))\|_{\mathcal{L}_0^2}^2 ds \\ &\lesssim (1 + \mathbb{E}\|u^0\|_{L^2}^2) + \int_0^t (t-s)^{-\frac{1}{2}} \mathbb{E}\|v(s)\|_{L^2}^2 ds \\ &\lesssim 1 + \|v\|_Y, \end{aligned}$$

which implies that $Mv \in L^\infty(0, T; L^2(\Omega; L^2(D)^d))$. By considering $\mathbb{E}\|Mv(t_2) - Mv(t_1)\|_{L^2}^2$ one can also prove the continuity of Mv in time. As a result, $Mv \in C([0, T]; L^2(\Omega; L^2(D)^d))$ and therefore the map $M : Y \rightarrow Y$ is well defined.

Clearly, a mild solution of (3.12) is equivalent to a fixed point of M .

To prove the existence and uniqueness of a fixed point of M , we define Y_λ to be Y equipped with an equivalent norm

$$\|w\|_{Y_\lambda} := \max_{t \in [0, T]} e^{-\lambda t} (\mathbb{E}\|w\|_{L^2}^2)^{\frac{1}{2}} \quad \text{for } w \in Y_\lambda. \quad (\text{A.4})$$

If $v_1, v_2 \in Y$ then

$$Mv_1(t) - Mv_2(t) = \int_0^t E(t-s)P_X [B(v_1(s)) - B(v_2(s))]dW(s) \quad \text{for } t \in [0, T].$$

By applying (A.3) we obtain

$$\begin{aligned}
e^{-2\lambda t} \mathbb{E} \|Mv_1(t) - Mv_2(t)\|_{L^2}^2 &\lesssim \int_0^t e^{-2\lambda t} (t-s)^{-\frac{1}{2}} \mathbb{E} \|v_1(s) - v_2(s)\|_{L^2}^2 ds \\
&= \int_0^t e^{-2\lambda(t-s)} (t-s)^{-\frac{1}{2}} e^{-2\lambda s} \mathbb{E} \|v_1(s) - v_2(s)\|_{L^2}^2 ds \\
&\lesssim \int_0^t e^{-2\lambda\sigma} \sigma^{-\frac{1}{2}} d\sigma \|v_1 - v_2\|_{Y_\lambda}, \\
&\lesssim \lambda^{-\frac{1}{2}} \|v_1 - v_2\|_{Y_\lambda},
\end{aligned}$$

which implies that

$$\|Mv_1 - Mv_2\|_{Y_\lambda} \lesssim \lambda^{-\frac{1}{4}} \|v_1 - v_2\|_{Y_\lambda}. \quad (\text{A.5})$$

As a result, by choosing a sufficiently large λ , the map $M : Y_\lambda \rightarrow Y_\lambda$ is a contraction. By the Banach fixed point theorem, M has a unique fixed point in Y_λ (which consists of the same elements as Y). This proves the existence of a unique mild solution satisfying (3.13).

Stability: We first consider the case H^β with parameter $\beta \in (\frac{3}{2}, 2)$ appeared in (5.3). Itô's isometry (2.2) together with (4.14) and (4.16) yields

$$\begin{aligned}
\mathbb{E} \|u(t)\|_{H^\beta}^2 &\lesssim \mathbb{E} \|E(t)(I+A)^{\frac{\beta}{2}} u^0\|_{L^2}^2 + \mathbb{E} \left(\int_0^t \|(I+A)^{\frac{\beta}{2}} E(t-s) P_X f(s)\|_{L^2} ds \right)^2 \\
&\quad + \mathbb{E} \left\| \int_0^t (I+A)^{\frac{\beta}{2}} E(t-s) P_X B(u(s)) dW(s) \right\|_{L^2}^2 \\
&\lesssim \mathbb{E} \|u^0\|_{H^\beta}^2 + \mathbb{E} \left(\int_0^t (t-s)^{-\frac{\beta}{2}} \|f(s)\|_{L^2} ds \right)^2 \\
&\quad + \mathbb{E} \int_0^t \|(I+A)^{\frac{\beta-1}{2}} E(t-s) (I+A)^{\frac{1}{2}} P_X B(u(s))\|_{\mathcal{L}_2^0}^2 ds \\
&\lesssim (1 + \mathbb{E} \|u^0\|_{H^\beta}^2) + \int_0^t (t-s)^{-(\beta-1)} \mathbb{E} \|(I+A)^{\frac{1}{2}} P_X B(u(s))\|_{\mathcal{L}_2^0}^2 ds \\
&\lesssim (1 + \mathbb{E} \|u^0\|_{H^\beta}^2) + \int_0^t (t-s)^{-(\beta-1)} \mathbb{E} \|u(s)\|_{H^\beta}^2 ds \quad (\text{here (5.3) is used}).
\end{aligned}$$

By using the Gronwall's inequality, we arrive at

$$\sup_{t \in [0, T]} \mathbb{E} \|u(t)\|_{H^\beta}^2 \lesssim 1 + \mathbb{E} \|u^0\|_{H^\beta}^2 \quad \text{for all } \frac{3}{2} \leq \beta < 2. \quad (\text{A.6})$$

For the regularity in time, it follows from (5.2), (5.3) and (A.6) that

$$\sup_{t \in [0, T]} \mathbb{E} \|B(u(t))\|_{\mathcal{L}_2^0}^2 + \sup_{t \in [0, T]} \mathbb{E} \|(I+A)^{\frac{1}{2}} P_X B(u(t))\|_{\mathcal{L}_2^0}^2 \lesssim 1 + \mathbb{E} \|u^0\|_{H^2}^2,$$

which together with (4.16)–(4.17) shows

$$\begin{aligned}
\mathbb{E} \|u(t) - u(s)\|_{H^1}^2 &\lesssim \mathbb{E} \|(I+A)^{-\frac{1}{2}} (E(t-s) - I) (I+A) u^0\|_{L^2}^2 \\
&\quad + \mathbb{E} \left(\int_s^t \|(I+A)^{\frac{1}{2}} E(t-\sigma) P_X f(\sigma)\|_{L^2} d\sigma \right)^2 \\
&\quad + \mathbb{E} \int_s^t \|(I+A)^{\frac{1}{2}} E(t-\sigma) P_X B(u(\sigma))\|_{\mathcal{L}_2^0}^2 d\sigma \\
&\lesssim (t-s) (1 + \mathbb{E} \|u^0\|_{H^2}^2) + \int_s^t \mathbb{E} \|(I+A)^{\frac{1}{2}} P_X B(u(\sigma))\|_{\mathcal{L}_2^0}^2 d\sigma \\
&\lesssim (1 + \mathbb{E} \|u^0\|_{H^2}^2) (t-s).
\end{aligned} \quad (\text{A.7})$$

and

$$\mathbb{E} \|u(t) - u(s)\|_{H^\beta}^2 \lesssim \mathbb{E} \|(I+A)^{-\frac{2-\beta}{2}} (E(t-s) - I) (I+A) u^0\|_{L^2}^2 \quad (\text{A.8})$$

$$\begin{aligned}
& + \mathbb{E} \left(\int_s^t \|(I+A)^{\frac{\beta}{2}} E(t-\sigma) P_X f(\sigma)\|_{L^2} d\sigma \right)^2 \\
& + \mathbb{E} \int_s^t \|(I+A)^{\frac{\beta-1}{2}} E(t-\sigma) (I+A)^{\frac{1}{2}} P_X B(u(\sigma))\|_{\mathcal{L}_2^0}^2 d\sigma \\
& \lesssim (t-s)^{2-\beta} (1 + \mathbb{E}\|u^0\|_{H^2}^2) + \int_s^t (t-\sigma)^{1-\beta} \mathbb{E}\|(I+A)^{\frac{1}{2}} P_X B(u(\sigma))\|_{\mathcal{L}_2^0}^2 d\sigma \\
& \lesssim (1 + \mathbb{E}\|u^0\|_{H^2}^2) (t-s)^{2-\beta} \quad \text{for all } \frac{3}{2} < \beta < 2.
\end{aligned}$$

This proves (3.15) and (3.16).

With the help of the above estimates, the H^2 stability follows from

$$\begin{aligned}
\mathbb{E}\|u(t)\|_{H^2}^2 & \lesssim \mathbb{E}\|u^0\|_{H^2}^2 + \mathbb{E} \left\| \int_0^t (I+A)E(t-s) P_X f(s) ds \right\|_{L^2}^2 \\
& \quad + \mathbb{E} \left\| \int_0^t (I+A)^{\frac{1}{2}} E(t-s) (I+A)^{\frac{1}{2}} P_X B(u(s)) dW(s) \right\|_{L^2}^2 \\
& =: \mathbb{E}\|u^0\|_{H^2}^2 + \mathbb{E}|S_1|^2 + \mathbb{E}|S_2|^2,
\end{aligned} \tag{A.9}$$

where we use the triangle inequality to get

$$\begin{aligned}
\mathbb{E}|S_1|^2 & \lesssim \mathbb{E} \left(\int_0^t \|(I+A)E(t-s) P_X [f(s) - f(t)]\|_{L^2} ds \right)^2 + \mathbb{E} \left\| (I+A) \int_0^t E(t-s) P_X f(t) ds \right\|_{L^2}^2 \\
& \lesssim \mathbb{E} \left(\int_0^t (t-s)^{-1} \|f(s) - f(t)\|_{L^2} ds \right)^2 + \mathbb{E}\|f(t)\|_{L^2}^2 \quad (\text{here (4.16) and (4.19) are used}) \\
& \lesssim 1, \quad (\text{here (2.7)–(2.8) are used})
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}|S_2|^2 & \lesssim \mathbb{E} \int_0^t \|(I+A)^{\frac{1}{2}} E(t-s) (I+A)^{\frac{1}{2}} P_X [B(u(s)) - B(u(t))]\|_{\mathcal{L}_2^0}^2 ds \\
& \quad + \mathbb{E} \int_0^t \|(I+A)^{\frac{1}{2}} E(t-s) (I+A)^{\frac{1}{2}} P_X B(u(t))\|_{\mathcal{L}_2^0}^2 ds \\
& \lesssim \int_0^t (t-s)^{-1} \mathbb{E}\|(I+A)^{\frac{1}{2}} P_X [B(u(t)) - B(u(s))]\|_{\mathcal{L}_2^0}^2 ds \quad (\text{here (4.16) is used}) \\
& \quad + \mathbb{E}\|(I+A)^{\frac{1}{2}} P_X B(u(t))\|_{\mathcal{L}_2^0}^2 \quad (\text{here (4.18) with } \rho = 1 \text{ is used}) \\
& \lesssim \int_0^t (t-s)^{-1} \mathbb{E}\|u(t) - u(s)\|_{H^\beta}^2 ds + 1 + \mathbb{E}\|u^0\|_{H^2}^2 \quad (\text{here (5.6) is used}) \\
& \lesssim 1 + \mathbb{E}\|u^0\|_{H^2}^2 + \int_0^t (t-s)^{1-\beta} ds \quad (\text{here (3.16) for } \beta \in (\frac{3}{2}, 2) \text{ is used}) \\
& \lesssim 1 + \mathbb{E}\|u^0\|_{H^2}^2.
\end{aligned}$$

This proves (3.14). The proof of Proposition 3.1 is completed. \square

Appendix B: Proof of several technical lemmas

Proof of Lemma 4.3. By Korn's inequality [21, Theorem 2.4], the following equivalence relation holds:

$$\|v\|_{L^2}^2 + \|\mathbb{D}(v)\|_{L^2}^2 \sim \|v\|_{H^1}^2 \quad \forall v \in H^1(D)^d \cap X. \tag{B.1}$$

This implies the coercitivity of the operator $I + A$ and the existence of an inverse operator $(I + A)^{-1} : X \rightarrow X \cap H^1(D)^d$ (see the Lax–Milgram Lemma in [11, Theorem 6.2-1]). Since $X \cap H^1(D)^d$ is compactly embedded into X , it follows that $(I + A)^{-1}$ is compact. For such a

symmetric positive operator $I + A$ with compact inverse, its fractional powers can be defined by means of the spectral decomposition [23, Appendix B.2].

The H^2 elliptic regularity of the Stokes equations implies that $(I + A)^{-1} : X \rightarrow X \cap H^2(D)^d$, which implies $\|u\|_{H^2} \lesssim \|(I + A)u\|_{L^2}$ for $u \in D(A)$. Since $\|(I + A)u\|_{L^2} \lesssim \|u\|_{H^2}$ for $u \in D(A)$, it follows that the statement in (4.14) holds for $s = 2$. The intermediate case for $s \in (0, 2)$ follows by the real interpolation between the two endpoint cases $s = 0$ and $s = 2$.

(B.1) implies $\|(I + A)^{\frac{1}{2}}v\|_{L^2} \sim \|v\|_{H^1}$ for $v \in D(A)$, i.e., the two norms $\|\cdot\|_{D(A^{\frac{1}{2}})}$ and $\|\cdot\|_{H^1}$ are equivalent on $D(A)$. Since $D(A)$ is dense in $H^1(D)^d \cap X$, it follows that

$$D(A^{\frac{1}{2}}) = \text{closure of } D(A) \text{ under the norm } \|\cdot\|_{D(A^{\frac{1}{2}})} \sim \|\cdot\|_{H^1} = H^1(D)^d \cap X.$$

Therefore, for $s \in (0, 1)$ we have

$$\begin{aligned} D(A^{\frac{s}{2}}) &= \text{closure of } D(A^{\frac{1}{2}}) \text{ under the norm } \|\cdot\|_{D(A^{\frac{s}{2}})} \\ &= \text{closure of } H^1(D)^d \cap X \text{ under the norm } \|\cdot\|_{H^s} = H^s(D)^d \cap X. \end{aligned}$$

The statement in (4.15) follows from the duality argument. \square

Proof of Lemma 4.4. Let $\tilde{E}(t) := e^{-t(I+A)}$ be the semigroup generated by $I + A$, which has a compact inverse operator. Then [23, Lemma 2.5] is applicable and gives the estimates in (4.16)–(4.19) with $E(t)$ replaced by $\tilde{E}(t)$ therein. Since $E(t) = e^t \tilde{E}(t)$ and $t \in (0, T]$, it follows that (4.16)–(4.19) also hold for $E(t)$. \square

Proof of Lemma 4.6. Subtracting (4.34) from (4.33) yields

$$\begin{cases} (z(w - w_h), v_h) + 2(\mathbb{D}(w - w_h), \mathbb{D}(v_h)) - (p - p_h, \nabla \cdot v_h) = 0 & \forall v_h \in V_h, \\ (\nabla \cdot (w - w_h), q_h) = 0 & \forall q_h \in Q_h. \end{cases} \quad (\text{B.2})$$

Let $e_h = P_{X_h} w - w_h \in X_h$. Choosing $(v_h, q_h) = (e_h, P_{Q_h} p - p_h) \in X_h \times Q_h$ in (B.2), where P_{Q_h} denotes the L^2 -orthogonal projection onto Q_h , we have

$$z\|e_h\|_{L^2}^2 + 2\|\mathbb{D}(e_h)\|_{L^2}^2 = -2(\mathbb{D}(w - P_{X_h} w), \mathbb{D}(e_h)) + (p - P_{Q_h} p, \nabla \cdot e_h). \quad (\text{B.3})$$

By considering the imaginary and the real parts of the above equality, separately, we obtain

$$\begin{cases} |\text{Im}(z)|\|e_h\|_{L^2}^2 \lesssim \varepsilon^{-1}\|w - P_{X_h} w\|_{H^1}^2 + \varepsilon\|\mathbb{D}(e_h)\|_{L^2}^2 + \|p - P_{Q_h} p\|_{L^2}, \\ |\text{Re}(z)|\|e_h\|_{L^2}^2 + 2\|\mathbb{D}(e_h)\|_{L^2}^2 \lesssim \varepsilon^{-1}\|w - P_{X_h} w\|_{H^1}^2 + \varepsilon\|\mathbb{D}(e_h)\|_{L^2}^2 + \|p - P_{Q_h} p\|_{L^2}, \end{cases}$$

where $\varepsilon > 0$ can be arbitrarily small. For $z \in \Sigma_\phi := \{1 + z' \in \mathbb{C} : |\arg(z')| < \phi\}$, with some $\phi \in (0, \pi)$, we have $|z| \gtrsim 1$ and $|\text{Re}(z)| \lesssim |\text{Im}(z)|$. By using (2.10), (2.14), (4.9) and Korn's inequality $\|u\|_{H^1}^2 \lesssim \|u\|_{L^2}^2 + \|\mathbb{D}(u)\|_{L^2}^2$, choosing a sufficiently small ε in the above estimates, we can derive the following result:

$$\begin{aligned} |z|^{\frac{1}{2}}\|P_{X_h} w - w_h\|_{L^2} + \|\nabla(P_{X_h} w - w_h)\|_{L^2} &\lesssim \|w - P_{X_h} w\|_{H^1} + \|p - P_{Q_h} p\|_{L^2} \\ &\lesssim h(\|w\|_{H^2} + \|p\|_{H^1}), \end{aligned} \quad (\text{B.4})$$

which, together with triangle inequality and (4.9), implies that

$$\|\nabla(w - w_h)\|_{L^2} \lesssim h(\|w\|_{H^2} + \|p\|_{H^1}). \quad (\text{B.5})$$

In order to obtain an estimate for $\|w - w_h\|_{L^2}$, we consider the following duality argument. Let $\phi \in \mathcal{D}(A)$ and $\phi_h \in X_h$ be the solution of the linear Stokes equations

$$\begin{cases} z\phi - \nabla \cdot \mathbb{T}(\phi, \eta) = w - w_h & \text{in } D, \\ \nabla \cdot \phi = 0 & \text{in } D, \\ \mathbb{T}(\phi, \eta)\mathbf{n} = 0 & \text{on } \partial D, \end{cases} \quad (\text{B.6})$$

and its finite element weak formulation

$$\begin{cases} (z\phi_h, v_h) + 2(\mathbb{D}(\phi_h), \mathbb{D}(v_h)) - (\eta_h, \nabla \cdot v_h) = (w - w_h, v_h) & \forall v_h \in V_h, \\ (\nabla \cdot \phi_h, q_h) = 0 & \forall q_h \in Q_h. \end{cases} \quad (\text{B.7})$$

Testing (B.6) by $\phi \in X$ and considering the imaginary and real parts of the results, we can obtain the following result similarly as (B.4):

$$|z|\|\phi\|_{L^2}^2 + 2\|\nabla\phi\|_{L^2}^2 \leq C|z|^{-1}\|w - w_h\|_{L^2}^2 + \frac{1}{2}|z|\|\phi\|_{L^2}^2,$$

which implies

$$|z|\|\phi\|_{L^2} \lesssim \|w - w_h\|_{L^2}.$$

Through the above estimate and the H^2 estimate of the linear Stokes equations (B.6) (cf. [26, Corollary 2.1]), we obtain

$$\|\phi\|_{H^2} + \|\eta\|_{H^1} \lesssim \|w - w_h\|_{L^2} + |z|\|\phi\|_{L^2} \lesssim \|w - w_h\|_{L^2}. \quad (\text{B.8})$$

Similar to (B.4)–(B.5), we have

$$|z|^{\frac{1}{2}}\|P_{X_h}\phi - \phi_h\|_{L^2} + \|\nabla(\phi - \phi_h)\|_{L^2} \lesssim h(\|\phi\|_{H^2} + \|\eta\|_{H^1}) \lesssim h\|w - w_h\|_{L^2}, \quad (\text{B.9})$$

where the last inequality follows from (B.8).

Since $(\nabla \cdot v_h, P_{Q_h}\eta) = 0$ for $v_h \in X_h$, it follows that

$$-(\eta, \nabla \cdot v_h) = -(\eta - P_{Q_h}\eta, \nabla \cdot v_h) \quad \forall v_h \in X_h.$$

Since $\phi_h \in X_h$, we can derive the following equation from (B.2):

$$(z\phi_h, w - w_h) + 2(\mathbb{D}(\phi_h), \mathbb{D}(w - w_h)) - (\nabla \cdot \phi_h, p - P_{Q_h}p) = 0.$$

Then, testing the first relation of (B.6) by $w - w_h$ and using the two equations above, by utilizing (2.14) and (4.9)–(4.10) we obtain

$$\begin{aligned} \|w - w_h\|_{L^2}^2 &= z(\phi - \phi_h, w - w_h) + 2(\mathbb{D}(\phi - \phi_h), \mathbb{D}(w - w_h)) \\ &\quad - (\nabla \cdot (\phi - \phi_h), p - P_{Q_h}p) - (\eta - P_{Q_h}\eta, \nabla \cdot (w - w_h)) \\ &\lesssim |z|^{\frac{1}{2}}\|\phi - \phi_h\|_{L^2}|z|^{\frac{1}{2}}\|w - w_h\|_{L^2} + \|\nabla(\phi - \phi_h)\|_{L^2}\|\nabla(w - w_h)\|_{L^2} \\ &\quad + h^2\|\phi\|_{H^2}\|p\|_{H^1} + h\|\eta\|_{H^1}\|\nabla(w - w_h)\|_{L^2} \\ &\lesssim (h^2z + h|z|^{\frac{1}{2}})\|w - w_h\|_{L^2}^2 + h\|w - w_h\|_{L^2}\|\nabla(w - w_h)\|_{L^2} \\ &\quad + h^2(\|w\|_{H^2} + \|p\|_{H^1})\|w - w_h\|_{L^2} \quad (\text{here (B.8) and (B.9) are used}) \\ &\lesssim (h^2 + h^3|z|^{\frac{1}{2}})\|w - w_h\|_{L^2}(\|w\|_{H^2} + \|p\|_{H^1}). \quad (\text{here (B.4) are used}) \end{aligned}$$

When $|z| \leq h^{-2}$, the inequality above reduces to

$$\|w - w_h\|_{L^2} \lesssim h^2(\|w\|_{H^2} + \|p\|_{H^1}). \quad (\text{B.10})$$

When $|z| \geq h^{-2}$, (B.4) immediately implies (B.10).

Combining the estimates (B.4) and (B.10), we obtain (4.35). The proof of (4.36) is similar as that for (B.8). \square

Appendix C: Proof of Lemma 5.2

Proof. By iterating with respect to the time levels, the full discrete method in (2.22) can be rewritten as

$$u_h^n = \bar{E}_{h,\tau}^n P_{X_h} u_0 + \tau \sum_{i=0}^{n-1} \bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1}) + \sum_{i=0}^{n-1} \bar{E}_{h,\tau}^{n-i} P_{X_h} [B(u_h^i) \Delta W_{i+1}]. \quad (\text{C.1})$$

The stability of $\bar{E}_{h,\tau}^n$, $1 \leq n \leq N$ together with (2.2) and (4.58) yields

$$\begin{aligned} \mathbb{E}\|u_h^n\|_{H^{\frac{1}{2}}}^2 &\lesssim \mathbb{E}\|\bar{E}_{h,\tau}^n (I + A_h)^{\frac{1}{4}} P_{X_h} u^0\|_{L^2}^2 \\ &\quad + \mathbb{E}\left(\tau \sum_{i=0}^{n-1} \|(I + A_h)^{\frac{1}{4}} \bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1})\|_{L^2}\right)^2 \end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} \left\| \sum_{i=0}^{n-1} (I + A_h)^{\frac{1}{4}} \bar{E}_{h,\tau}^{n-i} P_{X_h} [B(u_h^i) \Delta W_{i+1}] \right\|_{L^2}^2 \\
& \lesssim \mathbb{E} \|u^0\|_{H^{\frac{1}{2}}}^2 + \mathbb{E} \left(\tau \sum_{i=0}^{n-1} t_n^{-\frac{1}{4}} \|\bar{E}_{h,\tau}^{n-i} P_{X_h} f(t_{i+1})\|_{L^2} \right)^2 \\
& \quad + \tau \sum_{i=0}^{n-1} \mathbb{E} \|(I + A_h)^{\frac{1}{4}} \bar{E}_{h,\tau}^{n-i} P_{X_h} B(u_h^i)\|_{\mathcal{L}_2^0}^2 \\
& \lesssim (1 + \mathbb{E} \|u^0\|_{H^{\frac{1}{2}}}^2) + \tau \sum_{i=0}^{n-1} t_{n-i}^{-\frac{1}{2}} \mathbb{E} \|B(u_h^i)\|_{\mathcal{L}_2^0}^2 \\
& \lesssim (1 + \mathbb{E} \|u^0\|_{H^{\frac{1}{2}}}^2) + \tau \sum_{i=0}^{n-1} (t_n - t_i)^{-\frac{1}{2}} (1 + \mathbb{E} \|u_h^i\|_{H^{\frac{1}{2}}}^2). \quad (\text{here (5.2) is used}) \quad (\text{C.2})
\end{aligned}$$

By using the discrete Gronwall inequality, we obtain

$$\max_{1 \leq n \leq N} \mathbb{E} \|u_h^n\|_{H^{\frac{1}{2}}}^2 \lesssim 1 + \mathbb{E} \|u^0\|_{H^{\frac{1}{2}}}^2. \quad (\text{C.3})$$

This proves the desired estimate for the first term in (5.8).

Setting $v_h = u_h^n \in X_h$ in (2.19) and applying the identity $2(a-b)a = |a|^2 - |b|^2 + |a-b|^2$ for $a, b \in \mathbb{R}$ yield

$$\begin{aligned}
& \frac{1}{2} \|u_h^n\|_{L^2}^2 + \frac{1}{2} \|u_h^n - u_h^{n-1}\|_{L^2}^2 + 2\tau \|\mathbb{D}(u_h^n)\|_{L^2}^2 \\
& \leq \frac{1}{2} \|u_h^{n-1}\|_{L^2}^2 + \frac{1}{2} \tau \|f(t_n)\|_{L^2}^2 + \frac{1}{2} \tau \|u_h^n\|_{L^2}^2 + (B(u_h^{n-1}) \Delta W_n, u_h^n),
\end{aligned} \quad (\text{C.4})$$

where the last term can be estimated by applying the property of a martingale

$$\mathbb{E}(B(u_h^{n-1}) \Delta W_n, u_h^{n-1}) = 0$$

and using (5.2) and (C.3) as follows:

$$\begin{aligned}
\mathbb{E}(B(u_h^{n-1}) \Delta W_n, u_h^n) & = \mathbb{E}(B(u_h^{n-1}) \Delta W_n, u_h^n - u_h^{n-1}) \\
& \leq C \mathbb{E} \|B(u_h^{n-1}) \Delta W_n\|_{L^2}^2 + \frac{1}{4} \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2 \\
& \leq C \tau \mathbb{E} \|B(u_h^{n-1})\|_{\mathcal{L}_2^0}^2 + \frac{1}{4} \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2 \\
& \leq C \tau (1 + \mathbb{E} \|u_h^{n-1}\|_{H^{\frac{1}{2}}}^2) + \frac{1}{4} \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2 \\
& \leq C \tau (1 + \mathbb{E} \|u_h^0\|_{H^1}^2) + \frac{1}{4} \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2.
\end{aligned} \quad (\text{C.5})$$

By using Korn's inequality (cf. [21, Theorem 2.4]) $\|u\|_{L^2}^2 + \|\mathbb{D}(u)\|_{L^2}^2 \geq \theta(\|u\|_{L^2}^2 + \|\nabla u\|_{L^2}^2)$, where $\theta > 0$ is some constant, we obtain

$$\mathbb{E} \|u_h^N\|_{L^2}^2 + \sum_{n=1}^N \mathbb{E} \|u_h^n - u_h^{n-1}\|_{L^2}^2 + \tau \sum_{n=1}^N \mathbb{E} \|\nabla u_h^n\|_{L^2}^2 \lesssim 1 + \mathbb{E} \|u_h^0\|_{H^1}^2 + \tau \sum_{n=1}^N \mathbb{E} \|u_h^n\|_{L^2}^2. \quad (\text{C.6})$$

Then applying the discrete Gronwall inequality yields the desired estimate for the last two terms in (5.8). This proves Lemma 5.2. \square

References

- [1] W. Arendt, C. J. Batty, M. Hieber, and F. Neubrander. *Vector-valued Laplace transforms and Cauchy problems*. Birkhäuser, second edition, 2011.
- [2] D. N. Arnold, F. Brezzi, and M. Fortin. A stable finite element for the Stokes equations. *Calcolo*, 21(4):337–344, 1984.

- [3] H. Bessaih, Z. Brzeźniak, and A. Millet. Splitting up method for the 2D stochastic Navier–Stokes equations. *Stoch. Partial Differ. Equ.-Anal. Comput.*, 2(4):433–470, 2014.
- [4] H. Bessaih and A. Millet. Strong L^2 convergence of time numerical schemes for the stochastic two-dimensional Navier–Stokes equations. *IMA J. Numer. Anal.*, 39(4):2135–2167, 2019.
- [5] D. Breit and A. Dodgson. Convergence rates for the numerical approximation of the 2D stochastic Navier–Stokes equations. *Numer. Math.*, 147(3):553–578, 2021.
- [6] D. Breit and A. Prohl. Error analysis for 2D stochastic Navier–Stokes equations in bounded domains. *arXiv preprint arXiv:2109.06495*, 2021.
- [7] Z. Brzeźniak, E. Carelli, and A. Prohl. Finite-element-based discretizations of the incompressible Navier–Stokes equations with multiplicative random forcing. *IMA J. Numer. Anal.*, 33(3):771–824, 2013.
- [8] Y. Cao, J. Hong, and Z. Liu. Approximating Stochastic evolution equations with additive white and rough noises. *SIAM J. Numer. Anal.*, 55(4):1958–1981, 2017.
- [9] E. Carelli, E. Hausenblas, and A. Prohl. Time-splitting methods to solve the stochastic incompressible Stokes equation. *SIAM J. Numer. Anal.*, 50(6):2917–2939, 2012.
- [10] E. Carelli and A. Prohl. Rates of convergence for discretizations of the stochastic incompressible Navier–Stokes equations. *SIAM J. Numer. Anal.*, 50(5):2467–2496, 2012.
- [11] P. G. Ciarlet. *Linear and nonlinear functional analysis with applications*, volume 130. Siam, 2013.
- [12] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159. Springer, 2004.
- [13] X. Feng, A. Prohl, and L. Vo. Optimally convergent mixed finite element methods for the stochastic Stokes equations. *IMA J. Numer. Anal.*, 41(3):2280–2310, 2021.
- [14] X. Feng and H. Qiu. Analysis of fully discrete mixed finite element methods for time-dependent stochastic Stokes equations with multiplicative noise. *J. Sci. Comput.*, 88(2):1–25, 2021.
- [15] X. Feng and L. Vo. Analysis of chorin-type projection methods for the stochastic Stokes equations with general multiplicative noise. *Stoch. Partial Differ. Equ.-Anal. Comput.*, pages 1–38, 2022.
- [16] V. Girault and P.-A. Raviart. *Finite element approximation of the Navier-Stokes equations*, volume 749. Springer Berlin, 1979.
- [17] V. Girault and P.-A. Raviart. *Finite element methods for Navier-Stokes equations: theory and algorithms*, volume 5. Springer Science & Business Media, 1986.
- [18] V. Girault and L. R. Scott. A quasi-local interpolation operator preserving the discrete divergence. *Calcolo*, 40(1):1–19, 2003.
- [19] J. G. Heywood and R. Rannacher. Finite element approximation of the nonstationary Navier-Stokes problem IV: Error analysis for second-order time discretization. *SIAM J. Numer. Anal.*, 27:353–384, 1990.
- [20] R. Ingram. A new linearly extrapolated Crank-Nicolson time-stepping scheme for the Navier-Stokes equations. *Math. Comp.*, 82(284):1953–1973, 2013.
- [21] V. John and W. J. Layton. Analysis of numerical errors in large eddy simulation. *SIAM J. Numer. Anal.*, 40(3):995–1020, 2002.
- [22] R. Kruse. Optimal error estimates of Galerkin finite element methods for stochastic partial differential equations with multiplicative noise. *IMA J. Numer. Anal.*, 34(1):217–251, 2014.
- [23] R. Kruse. *Strong and weak approximation of semilinear stochastic evolution equations*. Springer, 2014.
- [24] R. Kruse and S. Larsson. Optimal regularity for semilinear stochastic partial differential equations with multiplicative noise. *Electron. J. Probab.*, 17:1–19, 2012.
- [25] W. Layton, N. Mays, M. Neda, and C. Trenchea. Numerical analysis of modular regularization methods for the BDF2 time discretization of the Navier-Stokes equations. *ESAIM: M2AN*, 48(3):765–793, 2014.
- [26] B. Li. An explicit formula for corner singularity expansion of the solutions to the Stokes equations in a polygon. *International Journal of Numerical Analysis and Modeling*, 17(6):900–928, 2020.
- [27] M. Marion and R. Temam. Navier-Stokes equations: Theory and approximation. *Handbook of numerical analysis*, 6:503–689, 1998.
- [28] J. D. Mukam and A. Tambue. Strong convergence of the linear implicit Euler method for the finite element discretization of semilinear non-autonomous SPDEs driven by multiplicative or additive noise. *Appl. Numer. Math.*, 147:222–253, 2020.
- [29] R. H. Nochetto and J.-H. Pyo. Error estimates for semi-discrete gauge methods for the Navier–Stokes equations. *Math. Comp.*, 74:521–542, 2016.
- [30] J. Shen. On error estimates of projection methods for Navier–Stokes equations: first-order schemes. *SIAM J. Numer. Anal.*, 29(1):57–77, 1992.
- [31] J. Shen. On error estimates of some higher order projection and penalty-projection methods for Navier–Stokes equations. *Numer. Math.*, 62(1):49–73, 1992.
- [32] C. Taylor and P. Hood. A numerical solution of the Navier-Stokes equations using the finite element technique. *Comput. Fluids*, 1(1):73–100, 1973.
- [33] V. Thomée. *Galerkin finite element methods for parabolic problems*, volume 25. Springer Science & Business Media, 2007.
- [34] X. Wang. Strong convergence rates of the linear implicit Euler method for the finite element discretization of SPDEs with additive noise. *IMA. J. Numer. Anal.*, 37:965–984, 2017.
- [35] Y. Yan. Galerkin finite element methods for stochastic parabolic partial differential equations. *SIAM J. Numer. Anal.*, 43:1363–1384, 2005.

BUYANG LI AND SHU MA: DEPARTMENT OF APPLIED MATHEMATICS, THE HONG KONG POLYTECHNIC UNIVERSITY, HONG KONG. *E-mail address:* `buyang.li@polyu.edu.hk` AND `maisie.ma@connect.polyu.hk`

WEIWEI SUN: DIVISION OF SCIENCE AND TECHNOLOGY, BNU-HKBU UNITED INTERNATIONAL COLLEGE; ADVANCED INSTITUTE OF NATURAL SCIENCE, BEIJING NORMAL UNIVERSITY AT ZHUHAI; ZHUHAI, P.R. CHINA. *E-mail address:* `maweiw@uic.edu.cn`