

ROBUST UTILITY MAXIMIZATION WITH NONLINEAR CONTINUOUS SEMIMARTINGALES

DAVID CRIENS AND LARS NIEMANN

ABSTRACT. In this paper we study a robust utility maximization problem in continuous time under model uncertainty. The model uncertainty is governed by a continuous semimartingale with uncertain local characteristics. Here, the differential characteristics are prescribed by a set-valued function that depends on time and path. We show that the robust utility maximization problem is in duality with a conjugate problem, and study the existence of optimal portfolios for logarithmic, exponential and power utilities.

1. INTRODUCTION

1.1. **The purpose of the current article.** An important problem for a portfolio manager is to maximize the expected utility of his terminal wealth. For complete markets, this problem can be solved by the martingale method developed in [6, 7, 18, 35]. The case of incomplete markets is considerably more difficult. By now, the classical approach to compute the maximized utility (called *value function*) is to use a *duality argument* which was formalized in an abstract manner in the seminal paper [21], see also [15, 19] for other pioneering work. The key idea is to pass to a *dual optimization problem*, which is typically of reduced complexity, and to recover, via a bidual relation, the value function of the original problem as the conjugate of the value function corresponding to the dual problem.

In this paper we are interested in a robust framework, where, instead of a single financial model, a whole family of models is taken into consideration. Financially speaking, we think of a portfolio manager who is uncertain about the real-world measure, but who thinks that it belongs to a certain set of probabilities.

Recently, an abstract duality theory for possibly nondominated robust frameworks was developed in [3], see also [11] for another approach to a robust duality theory under drift and volatility uncertainty for bounded utilities. Compared to the classical case treated in [21], the theory from [3] relies on a measure-independent dual pairing which requires a suitable topological structure. A natural choice for an underlying space is the Wiener space of continuous functions, which can be seen as a *canonical framework*.

In their fundamental work [21], the authors use their abstract theory to establish duality theorems for general semimartingale market models. When it comes to robust duality theorems in nondominated settings, it seems that only Lévy type settings with deterministic uncertainty sets have been studied in detail, see [3, 11]. The purpose of this paper is to reduce the gap between the robust and non-robust case in terms of a robust duality theory for nondominated canonical continuous semimartingale markets with path dependent uncertainty sets. In the following we explain our setting in more detail.

Date: June 29, 2022.

2020 Mathematics Subject Classification. 60G65, 91B16, 93E20.

Key words and phrases. robust utility maximization; robust market price of risk; duality theory; nonlinear continuous semimartingales; semimartingale characteristics; Knightian uncertainty.

DC acknowledges financial support from the DFG project SCHM 2160/15-1 and LN acknowledges financial support from the DFG project SCHM 2160/13-1.

1.2. The setting. Consider the robust utility maximization problem given by

$$(1.1) \quad u(x) := \sup_{g \in \mathcal{C}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)],$$

where $U: (0, \infty) \rightarrow \mathbb{R}$ is a utility function, \mathcal{P} is a set of (possibly nondominated) probability measures on the Wiener space $\Omega := C([0, T]; \mathbb{R}^d)$, with finite time horizon $T > 0$, and $\mathcal{C}(x) := x\mathcal{C}$ is the set of claims that can be \mathcal{P} -quasi surely superreplicated with initial capital $x > 0$, i.e.,

$$\mathcal{C} := \{g: \Omega \rightarrow [0, \infty]: g \text{ is } \mathcal{F}_T^* \text{-measurable and } \exists H \in \mathcal{H}^{\mathcal{P}} \text{ with } 1 + (H \cdot X)_T \geq g \text{ } \mathcal{P}\text{-q.s.}\}.$$

We focus on logarithmic, exponential and power utility, i.e., U is assumed to be one of the following

$$U(x) = \log(x), \quad U(x) = -e^{-\lambda x} \text{ for } \lambda > 0, \quad U(x) = \frac{x^p}{p} \text{ for } p \in (-\infty, 0) \cap (0, 1).$$

The model uncertainty in this framework is introduced through the set \mathcal{P} , which consists of semimartingale laws on the Wiener space. We parameterize \mathcal{P} via a compact parameter space F and drift and volatility coefficients $b: F \times [0, T] \times \Omega \rightarrow \mathbb{R}^d$ and $a: F \times [0, T] \times \Omega \rightarrow \mathbb{S}_+^d$ such that

$$\mathcal{P} = \{P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}: P \circ X_0^{-1} = \delta_{x_0}, (\lambda \otimes P)\text{-a.e. } (dB^P/d\lambda, dC^P/d\lambda) \in \Theta\},$$

where $\mathfrak{P}_{\text{sem}}^{\text{ac}}$ denotes the set of semimartingale laws with absolutely continuous characteristics, X is the coordinate process, $x_0 \in \mathbb{R}^d$ is the initial value, (B^P, C^P) are the P -characteristics of X , and

$$\Theta(t, \omega) := \{(b(f, t, \omega), a(f, t, \omega)): f \in F\}, \quad (t, \omega) \in [0, T] \times \Omega.$$

In words, the set \mathcal{P} of possible real-world measures consists of all continuous semimartingale models whose coefficients take uncertain values in the fully path-dependent set Θ .

Throughout the paper, we assume that b and a are continuous and of linear growth, and that Θ is convex-valued. Further, we will introduce a robust market price of risk, which seems to be a novel object. In case we deal with unbounded utility functions, we additionally assume either a certain uniform boundedness condition or that the volatility coefficient a is uniformly bounded and elliptic. In general, however, we do not impose any ellipticity assumption and thence allow the portfolio manager to take incomplete markets into consideration. It seems to us that this feature is new for robust semimartingale frameworks.

1.3. Comments on the literature. There is already a vast literature on the robust utility maximization problem (1.1) with respect to nondominated probability measures. Nonlinear Lévy frameworks with constant Θ were for instance considered in [3, 11, 24, 28] and nonlinear time-inhomogeneous Lévy settings with time-dependent Θ were studied in [22, 23, 34]. Further, [4] investigated a robust Merton problem with uncertain volatility of Lévy type and volatility state dependent drift. Compared to these papers, we allow for uncertain drift and volatility with fully path dependent coefficients. In particular, our work includes diffusion models with uncertain parameters, such as real-valued nonlinear affine models as studied in [13], as well as stochastic volatility models with uncertain volatility processes.

In contrast, the literature on the robust dual problem in continuous time is less extensive. Indeed, we are only aware of the papers [3, 11], where the problem is investigated from a theoretical perspective, and the only concrete examples we know of are the continuous Lévy frameworks with uniformly bounded and elliptic coefficients as discussed in these papers.

1.4. Main contributions. As in the classical case with one real-world measure ([21]), we are interested in the following question of quantitative nature.

(A) How can we calculate the value function u as given in (1.1)?

Further, we are interested in a qualitative question.

(B) Does an optimal portfolio exist for the problem (1.1)?

As in the non-robust case, we answer Question (A) via a duality argument. Denote the set of equivalent separating measures by

$$\mathcal{D} := \{Q \in \mathfrak{P}_e(\mathcal{P}) : E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C}\}, \quad \mathcal{D}(y) := y\mathcal{D}, \quad y > 0,$$

where $\mathfrak{P}_e(\mathcal{P}) := \{Q \in \mathfrak{P}(\Omega) : \exists P \in \mathcal{P} \text{ with } Q \sim P\}$. The robust dual problem is given by

$$(1.2) \quad v(y) := \inf_{Q \in \mathcal{D}(y)} \sup_{P \in \mathcal{P}} E^P \left[V \left(\frac{dQ}{dP} \right) \right],$$

where V denotes the conjugate of the utility function U . We show that the functions u and v are conjugates, i.e.,

$$u(x) = \inf_{y \geq 0} [v(y) + xy], \quad x > 0, \quad v(y) = \sup_{x \geq 0} [u(x) - xy], \quad y > 0,$$

which constitutes a satisfactory answer to Question (A).

To provide an answer to Question (B), we prove the existence of an optimal portfolio to a relaxed version of the optimization problem (1.1), which accounts for the obstacle that in the nondominated case one cannot rely on classical tools like Komlós' lemma for the approximation scheme of an optimal portfolio. For utility functions which are bounded from below, we also provide conditions for the existence of an optimal portfolio for the original problem (1.1).

Let us comment on our mathematical proofs. In the main body of this paper, we prove that the sets \mathcal{P} and \mathcal{D} are convex and compact, and that the sets \mathcal{C} and \mathcal{D} are in duality, i.e.,

$$(1.3) \quad \{Q \in \mathfrak{P}_e(\mathcal{P}) : E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C} \cap C_b(\Omega; \mathbb{R})\} = \mathcal{D},$$

and

$$(1.4) \quad \{g \in C_b^+(\Omega; \mathbb{R}) : E^Q[g] \leq 1 \text{ for all } Q \in \mathcal{D}\} = \mathcal{C} \cap C_b(\Omega; \mathbb{R}).$$

Once these properties are established we can follow [3] to establish our main results.

For the first duality (1.3), we show that \mathcal{D} coincides with the robust analogue of the set of equivalent local martingale measures, i.e.,

$$\mathcal{M} := \{Q \in \mathfrak{P}_{\text{sem}}^{\text{ac}} : Q \circ X_0^{-1} = \delta_{x_0}, (\lambda \otimes Q)\text{-a.e. } (dB^Q/d\lambda, dC^Q/d\lambda) \in \tilde{\Theta}\},$$

where

$$\tilde{\Theta}(t, \omega) := \{0\}^d \times \{a(f, t, \omega) : f \in F\} \subset \mathbb{R}^d \times \mathbb{S}_+^d, \quad (t, \omega) \in [0, T] \times \Omega.$$

The equality $\mathcal{M} = \mathcal{D}$ relies on a characterization of local martingale measures on the Wiener space, and it resembles the fact that for continuous paths, the set of separating measures coincides with the set of local martingale measures. Regarding the second duality (1.4), the equality $\mathcal{M} = \mathcal{D}$ further allows us to use the robust superhedging duality

$$\sup_{Q \in \mathcal{M}} E^Q[f] = \min \{x \in \mathbb{R} : \exists H \in \mathcal{H} \text{ with } x + (H \cdot S)_T \geq f \text{ } Q\text{-a.s. for all } Q \in \mathcal{M}\},$$

from [31], to show that there is a superhedging strategy for every function in the polar of \mathcal{D} .

Evidently, by virtue of (1.2), to deal with unbounded utility functions, we require some integrability of the Radon–Nikodym derivative dQ/dP . Under suitable ellipticity and linear growth conditions, we establish finite polynomial moments for certain stochastic exponentials. We think that this result is of independent interest.

1.5. Structure of the article. In Section 2.1 we lay out our setting, in Section 2.2 we introduce nonlinear continuous semimartingales, in Section 2.3 we provide the superhedging duality for nonlinear continuous semimartingales and in Section 2.4 we discuss the duality relation between \mathcal{C} and \mathcal{D} . In Section 2.5 we study the robust utility maximization problem, and its dual problem. The proofs for our main results are given in the remaining sections. More precisely, the superhedging duality is proved in Section 3.1. In Section 3.2 we show compactness and convexity of \mathcal{P} and \mathcal{M} ,

while the duality between \mathcal{C} and \mathcal{D} is proved in Section 3.3. The duality relation for the robust utility maximization and the finite polynomial moments of certain stochastic exponentials are proved in Section 3.4.

2. MAIN RESULT

2.1. The Setting. Fix a dimension $d \in \mathbb{N}$ and a finite time horizon $T > 0$. We define Ω to be the space of continuous functions $[0, T] \rightarrow \mathbb{R}^d$ endowed with the uniform topology. The canonical process on Ω is denoted by X , i.e., $X_t(\omega) = \omega(t)$ for $\omega \in \Omega$ and $t \in [0, T]$. It is well-known that $\mathcal{F} := \mathcal{B}(\Omega) = \sigma(X_t, t \in [0, T])$. We define $\mathbf{F} := (\mathcal{F}_t)_{t \in [0, T]}$ as the canonical filtration generated by X , i.e., $\mathcal{F}_t := \sigma(X_s, s \in [0, t])$ for $t \in [0, T]$. Notice that we do not make the filtration \mathbf{F} right-continuous. The set of probability measures on (Ω, \mathcal{F}) is denoted by $\mathfrak{P}(\Omega)$ and endowed with the usual topology of convergence in distribution. Moreover, we denote by $\mathcal{F}^* := \bigcap_{P \in \mathfrak{P}(\Omega)} \mathcal{F}^P$ the universal σ -field. Further, we denote the space of symmetric, positive semidefinite real-valued $d \times d$ matrices by \mathbb{S}_+^d , and by $\mathbb{S}_{++}^d \subset \mathbb{S}_+^d$ the set of all positive definite matrices in \mathbb{S}_+^d .

Let F be a metrizable space and let $b: F \times \llbracket 0, T \rrbracket \rightarrow \mathbb{R}^d$ and $a: F \times \llbracket 0, T \rrbracket \rightarrow \mathbb{S}_+^d$ be two Borel functions such that $(t, \omega) \mapsto b(f, t, \omega)$ and $(t, \omega) \mapsto a(f, t, \omega)$ are predictable for all $f \in F$.

Standing Assumption 2.1.

- (i) F is compact.
- (ii) b and a are continuous.
- (iii) The set $\{(b(f, t, \omega), a(f, t, \omega)): f \in F\} \subset \mathbb{R}^d \times \mathbb{S}_+^d$ is convex for every $(t, \omega) \in \llbracket 0, T \rrbracket$.
- (iv) There exists a constant $C > 0$ such that

$$\|b(f, t, \omega)\|^2 + \|a(f, t, \omega)\| \leq C \left(1 + \sup_{s \in [0, t]} \|\omega(s)\|^2\right)$$

for all $(f, t, \omega) \in F \times \llbracket 0, T \rrbracket$.

We define the correspondences, i.e., the set-valued maps, $\Theta, \tilde{\Theta}: \llbracket 0, T \rrbracket \rightarrow \mathbb{R}^d \times \mathbb{S}_+^d$ by

$$\begin{aligned} \Theta(t, \omega) &:= \{(b(f, t, \omega), a(f, t, \omega)): f \in F\} \subset \mathbb{R}^d \times \mathbb{S}_+^d, \\ \tilde{\Theta}(t, \omega) &:= \{0\}^d \times \{a(f, t, \omega): f \in F\} \subset \mathbb{R}^d \times \mathbb{S}_+^d. \end{aligned}$$

Remark 2.2. Thanks to [9, Lemma 2.11], both correspondences Θ and $\tilde{\Theta}$ have a measurable graph. Notice also that part (iii) of Standing Assumption 2.1 states that Θ is convex-valued.

We denote the set of laws of continuous semimartingales by $\mathfrak{P}_{\text{sem}} \subset \mathfrak{P}(\Omega)$. For $P \in \mathfrak{P}_{\text{sem}}$, we denote the semimartingale characteristics of the coordinate process X by (B^P, C^P) , and we set

$$\mathfrak{P}_{\text{sem}}^{\text{ac}} := \{P \in \mathfrak{P}_{\text{sem}}: P\text{-a.s. } (B^P, C^P) \ll \lambda\},$$

where λ denotes the Lebesgue measure. For $x \in \mathbb{R}^d$, we set

$$\mathcal{R}(x) := \{P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}: P \circ X_0^{-1} = \delta_x, (\lambda \otimes P)\text{-a.e. } (dB^P/d\lambda, dC^P/d\lambda) \in \Theta\}.$$

We further define, for fixed $x_0 \in \mathbb{R}^d$,

$$\begin{aligned} \mathcal{P} &:= \{P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}: P \circ X_0^{-1} = \delta_{x_0}, (\lambda \otimes P)\text{-a.e. } (dB^P/d\lambda, dC^P/d\lambda) \in \Theta\} = \mathcal{R}(x_0), \\ \mathcal{M} &:= \{Q \in \mathfrak{P}_{\text{sem}}^{\text{ac}}: Q \circ X_0^{-1} = \delta_{x_0}, (\lambda \otimes Q)\text{-a.e. } (dB^Q/d\lambda, dC^Q/d\lambda) \in \tilde{\Theta}\}. \end{aligned}$$

Let $\mathcal{Q} \subset \mathfrak{P}(\Omega)$ be a set of probability measures. Recall that a \mathcal{Q} -polar set is a set that is \mathcal{Q} -null under every $Q \in \mathcal{Q}$, and that a property holds \mathcal{Q} -quasi surely, if it holds outside a \mathcal{Q} -polar set. For two sets of probability measures $\mathcal{Q}, \mathcal{R} \subset \mathfrak{P}(\Omega)$, we write $\mathcal{Q} \ll \mathcal{R}$ if every \mathcal{R} -polar set is \mathcal{Q} -polar. If $\mathcal{Q} \ll \mathcal{R}$ and $\mathcal{R} \ll \mathcal{Q}$, we denote this by $\mathcal{Q} \sim \mathcal{R}$.

Remark 2.3. In terms of Standing Assumption 2.9 below, we provide a condition for $\mathcal{P} \sim \mathcal{M}$, see Lemma 3.4 below. Additionally, recall that a continuous semimartingale is a local martingale if and only if its first characteristic vanishes. Hence, in case Standing Assumption 2.9 is in force, \mathcal{M} consists of local martingale measures, and it can be seen as the robust analogue of the set of equivalent local martingale measures. In fact, thanks to the linear growth condition (iv) from Standing Assumption 2.1, the set \mathcal{M} then even consists of (true) martingale measures.

2.2. Example: nonlinear diffusions. Suppose that $b(f, t, \omega) = b'(f, \omega(t))$ and $a(f, t, \omega) = a'(f, \omega(t))$ for all $(f, t, \omega) \in F \times \llbracket 0, T \rrbracket$ and Borel functions $b': F \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $a': F \times \mathbb{R}^d \rightarrow \mathbb{S}^d$. This setting corresponds to a *nonlinear diffusion framework*. In particular, in this case the correspondences Θ and $\tilde{\Theta}$ have *Markovian structure*, i.e., for every $(t, \omega) \in \llbracket 0, T \rrbracket$, the sets $\Theta(t, \omega), \tilde{\Theta}(t, \omega)$ only depend on $\omega(t)$. Here, (ii) – (iv) in Standing Assumption 2.1 are implied by the following conditions:

(ii)' b' and a' are continuous.

(iii)' The set $\{(b'(f, x), a'(f, x)): f \in F\} \subset \mathbb{R}^d \times \mathbb{S}_+^d$ is convex for every $x \in \mathbb{R}^d$.

(iv)' There exists a constant $C > 0$ such that

$$\|b'(f, x)\|^2 + \|a'(f, x)\| \leq C(1 + \|x\|^2)$$

for all $(f, x) \in F \times \mathbb{R}^d$.

In the following, we assume these conditions (and the compactness of F).

For each $x \in \mathbb{R}^d$, we define the sublinear operator \mathcal{E}^x on the convex cone of upper semianalytic functions $\psi: \Omega \rightarrow \mathbb{R}$ by

$$\mathcal{E}^x(\psi) := \sup_{P \in \mathcal{R}(x)} E^P[\psi].$$

For every $x \in \mathbb{R}^d$, we have by construction that $\mathcal{E}^x(\psi(X_0)) = \psi(x)$ for every upper semianalytic function $\psi: \mathbb{R}^d \rightarrow \mathbb{R}$.

Denote, for $t \in [0, T]$, the shift operator $\theta_t: \Omega \rightarrow \Omega$ by $\theta_t(\omega) := \omega((\cdot + t) \wedge T)$ for all $\omega \in \Omega$. As in [10, Proposition 2.8], we obtain the following result.

Proposition 2.4. *For every upper semianalytic function $\psi: \Omega \rightarrow \mathbb{R}$, the equality*

$$\mathcal{E}^x(\psi \circ \theta_t) = \mathcal{E}^x(\mathcal{E}^{X_t}(\psi))$$

holds for every $x \in \mathbb{R}^d$, and every $t \in [0, T]$.

Proposition 2.4 confirms the intuition that the coordinate process is a nonlinear Markov process under the family $\{\mathcal{E}^x: x \in \mathbb{R}\}$, as it implies the equality

$$\mathcal{E}^x(\psi(X_{s+t})) = \mathcal{E}^x(\mathcal{E}^{X_t}(\psi(X_s)))$$

for every upper semianalytic function $\psi: \mathbb{R}^d \rightarrow \mathbb{R}$, $s, t \in [0, T]$ with $s + t \leq T$, and $x \in \mathbb{R}^d$.

Summing up, this setting provides a robust counterpart to classical continuous Markovian financial frameworks. In particular, it allows to combine different Markovian models such as, for instance, Cox–Ingersoll–Ross and Vasiček models. As explained in [13], the latter example is interesting to account for the possibility that in certain scenarios interest rates can take negative values as after the financial crises 2007 – 2008.

2.3. Superhedging duality. In what follows, we define the filtration $\mathbf{G} = (\mathcal{G}_t)_{t \in [0, T]}$ via

$$(2.1) \quad \mathcal{G}_t := \mathcal{F}_t^* \vee \mathcal{N}^{\mathcal{M}}, \quad t \in [0, T],$$

where \mathcal{F}_t^* is the universal completion of \mathcal{F}_t , and $\mathcal{N}^{\mathcal{M}}$ is the collection of sets which are (\mathcal{F}_T, Q) -null for all $Q \in \mathcal{M}$. The set \mathcal{H} consists of all \mathbf{G} -predictable processes $H \in L(X, Q)$ ¹ such that the stochastic integral process $H \cdot X$ is a \mathbf{G} - Q -supermartingale for all $Q \in \mathcal{M}$.²

¹see, for instance, [17, Section III.6.c] for more details

²In the paper [31] the strategies from \mathcal{H} are called admissible. Below, using the classical terminology, we reserve the name admissible for strategies whose wealth process is bounded from below.

Theorem 2.5. *Let $f: \Omega \rightarrow \mathbb{R}$ be an upper semianalytic function such that*

$$\sup_{Q \in \mathcal{M}} E^Q[|f|] < \infty.$$

Then, the superhedging duality holds, i.e.,

$$\sup_{Q \in \mathcal{M}} E^Q[f] = \min \{x \in \mathbb{R}: \exists H \in \mathcal{H} \text{ with } x + (H \cdot S)_T \geq f \text{ } Q\text{-a.s. for all } Q \in \mathcal{M}\}.$$

With Theorem 2.5 we provide a version of [31, Theorem 3.2] which is tailored to our nonlinear semimartingale framework. In particular, with Standing Assumption 2.1 we give explicit parameterized conditions for the superhedging duality. Let us highlight that we do *not* require any ellipticity assumption. Consequently, similar to the abstract result in [31], Theorem 2.5 covers incomplete market situations.

The next corollary will turn out to be useful in Section 3. It shows that for payoffs bounded from below, the optimal superhedging strategy provided by Theorem 2.5 is *admissible* in a robust sense.

Corollary 2.6. *Let $f: \Omega \rightarrow \mathbb{R}_+$ be an upper semianalytic function such that*

$$\sup_{Q \in \mathcal{M}} E^Q[f] < \infty.$$

Then, there exist $H \in \mathcal{H}$ and $C > 0$ with

$$H \cdot X \geq -C \quad Q\text{-a.s. for all } Q \in \mathcal{M},$$

such that

$$\sup_{Q \in \mathcal{M}} E^Q[f] + (H \cdot X)_T \geq f, \quad Q\text{-a.s. for all } Q \in \mathcal{M}.$$

Proof. Set $\pi := \sup_{Q \in \mathcal{M}} E^Q[f] \in \mathbb{R}_+$. The superhedging duality given in Theorem 2.5 grants the existence of a \mathbf{G} -predictable process H such that $H \cdot X$ is a Q -supermartingale for every $Q \in \mathcal{M}$, and

$$\pi + (H \cdot X)_T \geq f \quad Q\text{-a.s. for all } Q \in \mathcal{M}.$$

In particular, as f is non-negative, this implies

$$(H \cdot X)_t \geq E^Q[(H \cdot X)_T | \mathcal{G}_t] \geq -\pi, \quad \text{for all } t \in [0, T], \quad Q\text{-a.s. for all } Q \in \mathcal{M}.$$

This completes the proof. □

2.4. Separating duality for nonlinear continuous semimartingales. Following [3], we define

$$\begin{aligned} \mathcal{C} &:= \{g: \Omega \rightarrow [0, \infty]: g \text{ is } \mathcal{F}_T^*\text{-measurable and } \exists H \in \mathcal{H}^{\mathcal{P}} \text{ with } 1 + (H \cdot X)_T \geq g \text{ } \mathcal{P}\text{-q.s.}\}, \\ \mathfrak{P}_e(\mathcal{P}) &:= \{Q \in \mathfrak{P}(\Omega): \exists P \in \mathcal{P} \text{ with } Q \sim P\}, \\ \mathcal{D} &:= \{Q \in \mathfrak{P}_e(\mathcal{P}): E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C}\}. \end{aligned}$$

Here, the set $\mathcal{H}^{\mathcal{P}}$ of admissible integrands is slightly smaller than the set \mathcal{H} introduced in Section 3.1. To wit, for any collection $\mathcal{R} \subset \mathfrak{P}_{\text{sem}}$, we first define the filtration $\mathbf{G}^{\mathcal{R}} = (\mathcal{G}_t^{\mathcal{R}})_{t \in [0, T]}$ via

$$(2.2) \quad \mathcal{G}_t^{\mathcal{R}} := \mathcal{F}_t^* \vee \mathcal{N}^{\mathcal{R}}, \quad t \in [0, T],$$

where \mathcal{F}_t^* is the universal completion of \mathcal{F}_t , and $\mathcal{N}^{\mathcal{R}}$ is the collection of sets which are (\mathcal{F}_T, P) -null for all $P \in \mathcal{R}$. Then, let $\mathcal{H}^{\mathcal{R}}$ be the set of all $\mathbf{G}^{\mathcal{R}}$ -predictable processes H with $H \in L(X, P)$ for all $P \in \mathcal{R}$ and such that $H \cdot X \geq -C$ P -a.s. for all $P \in \mathcal{R}$ for some $C > 0$, where the constant C may depend on both, H and P .

Remark 2.7. Note that $\mathbf{G}^{\mathcal{M}}$ as defined in (2.2) coincides with the filtration \mathbf{G} introduced in (2.1). Provided Standing Assumption 2.9 below is in force, Lemma 3.4 below shows that $\mathcal{M} \sim \mathcal{P}$. In particular, in this case we have the identities

$$(2.3) \quad \mathbf{G}^{\mathcal{M}} = \mathbf{G}^{\mathcal{P}} = \mathbf{G}.$$

After having established Lemma 3.4, we will use (2.3) throughout Section 3 to ease the notation.

The following theorem was established in [10, Propositions 3.9, 5.7] for one-dimensional nonlinear diffusions, but the argument transfers to our multidimensional path dependent framework. Its compactness part extends [16, Theorem 4.41] and [25, Theorem 2.5] beyond the case where b and a are of Markovian structure, uniformly bounded and globally Lipschitz continuous.

Theorem 2.8. *The sets \mathcal{P} and \mathcal{M} are both convex and compact.*

To get the separating dualities, we need to establish a relation between the sets \mathcal{P} and \mathcal{M} in the sense that we need to change between measures from \mathcal{P} and \mathcal{M} . In the spirit of the linear case, it turns out that this is possible if a certain type of market price of risk exists.

For the remainder of Section 2 we impose the following:

Standing Assumption 2.9 (Existence of Robust Market Price of Risk). There exists a Borel function $\theta: F \times \llbracket 0, T \rrbracket \rightarrow \mathbb{R}^d$ such that $b = a\theta$. Moreover, for every $N > 0$, there exists a constant $C = C_N > 0$ such that

$$\sup \{ \langle (\theta, a\theta)(f, t, \omega) \rangle : f \in F, t \leq T_N(\omega) \} \leq C$$

for all $\omega \in \Omega$, where $T_N(\omega) := \inf(t \in [0, T] : \|\omega(t)\| \geq N) \wedge T$.

It seems to us that the structure of Standing Assumption 2.9 is new in the literature on robust dualities. The results from [3, 11] require a uniform ellipticity assumption. A weak form of such a condition is explained in the following remark.

Remark 2.10. In case $a(F \times \llbracket 0, T \rrbracket) \subset \mathbb{S}_{++}^d$, we can decompose $b = a\theta$ with $\theta = a^{-1}b$.

We are in the position to state the main result of this section.

Theorem 2.11. *The equality $\mathcal{M} = \mathcal{D}$ holds and the sets \mathcal{C} and \mathcal{D} are in duality, i.e.,*

$$(2.4) \quad \mathcal{D} = \{ Q \in \mathfrak{P}_e(\mathcal{P}) : E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C} \cap C_b(\Omega; \mathbb{R}) \},$$

and

$$(2.5) \quad \mathcal{C} \cap C_b(\Omega; \mathbb{R}) = \{ g \in C_b^+(\Omega; \mathbb{R}) : E^Q[g] \leq 1 \text{ for all } Q \in \mathcal{D} \}.$$

Theorem 2.11 can be seen as a generalization of [3, Propositions 5.7, 5.9] from a Lévy setting with constant $\Theta(t, \omega) \equiv \Theta \subset \mathbb{R}^d \times \mathbb{S}_{++}^d$ to a general nonlinear semimartingale framework with path dependent uncertainty set $(t, \omega) \mapsto \Theta(t, \omega)$. Further, it seems to be the first concrete result without an ellipticity assumption, i.e., which also covers incomplete market scenarios, see [3, Remark 3.2]. In particular, the above theorem can be applied for the nonlinear diffusion framework from Section 2.2, which includes, for instance, the affine framework from [13].

2.5. Duality theory for robust utility maximization. Let $U: (0, \infty) \rightarrow [-\infty, \infty)$ be a utility function, i.e., a concave and non-decreasing function. We define $U(0) := \lim_{x \searrow 0} U(x)$, and consider the conjugate function

$$V(y) := \begin{cases} \sup_{x \geq 0} [U(x) - xy], & y > 0, \\ \lim_{y \searrow 0} V(y), & y = 0, \\ \infty, & y < 0. \end{cases}$$

Remark 2.12. A *medial limit* is a positive linear functional $\lim \text{med}: \ell^\infty \rightarrow \mathbb{R}$ satisfying

$$\liminf_{n \rightarrow \infty} x_n \leq \lim \text{med}_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n$$

for every bounded sequence $(x_n)_{n \in \mathbb{N}} \in \ell^\infty$, and for every bounded sequence $(X_n)_{n \in \mathbb{N}}$ of universally measurable functions $X_n: \Omega \rightarrow \mathbb{R}$, the medial limit $\lim \text{med}_{n \rightarrow \infty} X_n$ is again universally measurable with

$$E^P \left[\lim \text{med}_{n \rightarrow \infty} X_n \right] = \lim \text{med}_{n \rightarrow \infty} E^P[X_n],$$

for every $P \in \mathfrak{P}(\Omega)$. Note that a medial limit extends naturally to sequences with values in $[-\infty, \infty]$.

In the realm of robust finance, medial limits have been a suitable substitute for classical arguments such as Komlós' lemma. We refer to [2, 3, 30, 32] for recent applications of medial limits. Let us also mention that the existence of medial limits is guaranteed when working under ZFC together with Martin's axiom, see [27, 29].

Condition 2.13. *Medial limits exist.*

Provided Condition 2.13 is in force, we set

$$\bar{\mathcal{C}} := \left\{ g: \Omega \rightarrow [0, \infty] : g = \lim \text{med}_{n \rightarrow \infty} g_n \text{ with } (g_n)_{n \in \mathbb{N}} \subset \mathcal{C} \right\},$$

and, for $x, y > 0$, we define

$$\mathcal{C}(x) := x\mathcal{C}, \quad \bar{\mathcal{C}}(x) := x\bar{\mathcal{C}}, \quad \mathcal{D}(y) := y\mathcal{D},$$

and

$$u(x) := \sup_{g \in \mathcal{C}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)], \quad \bar{u}(x) := \sup_{g \in \bar{\mathcal{C}}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)],$$

and

$$v(y) := \inf_{Q \in \mathcal{D}(y)} \sup_{P \in \mathcal{P}} E^P \left[V \left(\frac{dQ}{dP} \right) \right],$$

with the convention $\frac{dQ}{dP} := -\infty$ in case Q is not absolutely continuous with respect to P .

Recall that Standing Assumption 2.9 is in force and that θ denotes the robust market price of risk. The following two conditions are used when we consider unbounded utility functions. We only require that *one* of them holds.

Condition 2.14. *The function $\langle \theta, a\theta \rangle$ is uniformly bounded.*

Condition 2.15 (Uniform Ellipticity and Boundedness of Volatility). *There exists a constant $K > 0$ such that*

$$\frac{1}{K} \leq \|a(f, t, \omega) - I_d\| \leq K$$

for all $(f, t, \omega) \in F \times [0, T]$, where I_d denotes the identity matrix in \mathbb{S}_+^d .

Remark 2.16. From a technical point of view, we need one of the Conditions 2.14 and 2.15 to prove that certain stochastic exponentials have polynomial moments, which are needed to treat unbounded utility functions. We emphasize that the scope of the assumptions is different. On one hand, Condition 2.14 is a boundedness condition but it covers incomplete market models. On the other hand, Condition 2.15 allows the drift coefficient b to be unbounded but it enforces a robust version of market completeness.

Both conditions are satisfied for the Lévy framework studied in [3, Section 3]. Beyond the Lévy case, Condition 2.15 holds e.g. for real-valued nonlinear affine processes as studied in [13]. In general, ellipticity assumptions (not necessarily uniform) are quite standard in the literature on linear multidimensional diffusions (see, e.g., [40]).

We are in the position to state the main results of this paper. The first of the following two theorems deals with utility functions which are bounded from below and the second deals with utility functions which are unbounded from below.

Theorem 2.17. *Let U be either a power utility $U(x) = x^p/p$ with exponent $p \in (0, 1)$, or an exponential utility $U(x) = -e^{-\lambda x}$ with parameter $\lambda > 0$. In case U is a power utility assume also that either Condition 2.14 or Condition 2.15 holds. Then,*

- (i) u is nondecreasing, concave and real-valued on $(0, \infty)$,
- (ii) v is nonincreasing, convex, and proper,
- (iii) the functions u and v are conjugates, i.e.,

$$u(x) = \inf_{y \geq 0} [v(y) + xy], \quad x > 0, \quad v(y) = \sup_{x \geq 0} [u(x) - xy], \quad y > 0,$$

- (iv) for every $x > 0$ we have

$$u(x) = \sup_{g \in \mathcal{C}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)] = \sup_{g \in \mathcal{C}(x) \cap \mathcal{C}_b} \inf_{P \in \mathcal{P}} E^P[U(g)].$$

If additionally Condition 2.13 holds, then

- (v) for every $x > 0$ we have $u(x) = \bar{u}(x)$,
- (vi) for every $x > 0$ there exists $g^* \in \bar{\mathcal{C}}(x)$ such that

$$\bar{u}(x) = \sup_{g \in \bar{\mathcal{C}}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)] = \inf_{P \in \mathcal{P}} E^P[U(g^*)].$$

Theorem 2.18. *Suppose that Condition 2.13 holds. Let U be either a power utility $U(x) = x^p/p$ with exponent $p \in (-\infty, 0)$, or the log utility $U(x) = \log(x)$. In case U is the log utility assume also that either Condition 2.14 or Condition 2.15 holds. Then,*

- (i) \bar{u} is nondecreasing, concave and real-valued on $(0, \infty)$,
- (ii) v is nonincreasing, convex, and real-valued on $(0, \infty)$,
- (iii) the functions \bar{u} and v are conjugates, i.e.,

$$\bar{u}(x) = \inf_{y \geq 0} [v(y) + xy], \quad x > 0, \quad v(y) = \sup_{x \geq 0} [\bar{u}(x) - xy], \quad y > 0,$$

- (iv) for every $x > 0$ there exists $g^* \in \bar{\mathcal{C}}(x)$ with

$$\bar{u}(x) = \sup_{g \in \bar{\mathcal{C}}(x)} \inf_{P \in \mathcal{P}} E^P[U(g)] = \inf_{P \in \mathcal{P}} E^P[U(g^*)].$$

The above Theorems 2.17 and 2.18 extend [3, Theorems 3.4, 3.5] from a Lévy setting to a general nonlinear semimartingale framework. In particular, as explained in Section 2.2, our setup includes nonlinear diffusion settings, which already proved to be interesting for financial applications, see, e.g., [13] for an affine setting.

Remark 2.19. It is interesting to consider the case with drift *uncertainty* and volatility *certainly*, which is closely related to the treatment of robust utility maximization with a dominated set of probability measures as studied in [37, 38, 39].

To get a rather concrete situation, suppose that there exists a function $a': [0, T] \times \mathbb{R}^d \rightarrow \mathbb{S}_+^d$ such that $a(f, t, \omega) = a'(t, \omega(t))$ for all $(f, t, \omega) \in F \times \llbracket 0, T \rrbracket$, and a' is continuous, bounded and uniformly elliptic. Then, it is well-known ([40, Theorem 7.2.1]) that $\mathcal{M} = \{Q\}$, where Q is the unique law of a solution process to the SDE

$$dY_t = \sqrt{a'}(t, Y_t) dW_t, \quad Y_0 = x_0,$$

where W is a d -dimensional Brownian motion. In this case, since $\mathcal{M} = \mathcal{D}$ by Theorem 2.11, the value function v of the robust dual problem is given by

$$v(y) = \sup_{P \in \mathcal{P}} E^P \left[V \left(y \frac{dQ}{dP} \right) \right], \quad y > 0.$$

Notice that the structure of this representation is similar to those for the value function of the dual problem in the classical complete case (see [21, Theorem 2.0]) with the difference that v is the supremum over (equivalent) real-world measures in our robust framework. Further, part (iii) of Theorem 2.17 shows that

$$(2.6) \quad u(x) = \inf_{y \geq 0} \left[\sup_{P \in \mathcal{P}} E^P \left[V \left(y \frac{dQ}{dP} \right) \right] + xy \right], \quad x > 0,$$

which is again a similar representation as given by [21, Theorem 2.0] for the classical complete case.

The structure of (2.6) has similarities to the classical incomplete case (see part (iv) from [21, Theorem 2.2]) with the important difference that v is the supremum over real-world measures in our robust framework, while the value function is an infimum over risk-neutral measures in the classical incomplete case.

In the above discussion we allow for a state dependent volatility coefficient. As mentioned on p. 756 in [38], it is not clear whether a so-called least favorable measure exists in this setup. The existence of such is crucial to apply the main theorems from [38].

3. PROOFS

In this section we present the proofs of our main results. The structure is chronological with the appearance of the results in the previous sections.

3.1. Superhedging duality: proof of Theorem 2.5. We start with some auxiliary preparations and then finalize the proof. For any probability measure $P \in \mathfrak{P}(\Omega)$ and any $[0, T]$ -valued stopping time $\tau: \Omega \rightarrow [0, T]$, there exists a version of the regular conditional P -probability given \mathcal{F}_τ , denoted by $P(\cdot | \mathcal{F}_\tau)$. Notice that $\mathcal{F}_\tau = \sigma(X_{t \wedge \tau}, t \in \mathbb{Q}_+)$ is countably generated ([40, Lemma 1.3.3]). Hence, there exists a P -null set $N \in \mathcal{F}_\tau$ such that $P(A | \mathcal{F}_\tau)(\omega) = \mathbb{1}_A(\omega)$ for all $A \in \mathcal{F}_\tau$ and $\omega \notin N$. In particular, we may and will choose a version $P(\cdot | \mathcal{F}_\tau)$ such that

$$P(\{\omega' \in \Omega: \omega = \omega' \text{ on } [0, \tau(\omega)]\} | \mathcal{F}_\tau)(\omega) = 1 \text{ for all } \omega \in \Omega.$$

Hence, for every $[0, T]$ -valued stopping time τ and every $\omega \in \Omega$ the shifted measure

$$(3.1) \quad P^{\tau, \omega} := P(\omega \tilde{\otimes}_{\tau(\omega)} \cdot | \mathcal{F}_\tau)(\omega)$$

defines a probability measure on (Ω, \mathcal{F}) . Here, we define for $\omega, \omega' \in \Omega$ and $t \in [0, T]$ the concatenation

$$\omega \tilde{\otimes}_t \omega' := \omega \mathbb{1}_{[0, t)} + (\omega(t) + \omega'(\cdot - t) - \omega'(0)) \mathbb{1}_{[t, T]}.$$

For $(t, \omega) \in \llbracket 0, T \rrbracket$, we define

$$\mathcal{M}(t, \omega) := \left\{ Q \in \mathfrak{P}_{\text{sem}}^{\text{ac}} : Q \circ X_0^{-1} = \delta_{x_0}, (\lambda \otimes Q)\text{-a.e. } (dB^Q/d\lambda, dC^Q/d\lambda) \in \tilde{\Theta}(\cdot + t, \omega \tilde{\otimes}_t X) \right\}.$$

Note that $\mathcal{M}(0, \omega) = \mathcal{M}$ for every $\omega \in \Omega$.

The next two results can be found in [16, Proposition 4.31], or [9, Section 3] in combination with [9, Lemma 3.5].

Lemma 3.1. *The set $\{(t, \omega, Q) \in \llbracket 0, T \rrbracket \times \mathfrak{P}(\Omega) : Q \in \mathcal{M}(t, \omega)\}$ is Borel.*

Lemma 3.2. *Let $(t, \omega) \in \llbracket 0, T \rrbracket$, let τ be a $[0, T]$ -valued stopping time with $\tau \geq t$, and let $Q \in \mathcal{M}(t, \omega)$. Set $\rho := \tau(\omega \tilde{\otimes}_t X) - t$. Then,*

- (i) *P -a.s. $P^{\rho, X} \in \mathcal{M}(\tau(\omega \tilde{\otimes}_t X), \omega \tilde{\otimes}_t X)$, where $P^{\rho, X}$ is defined in (3.1),*
- (ii) *for every \mathcal{F}_ρ -measurable kernel $\kappa: \Omega \rightarrow \mathfrak{P}(\Omega)$ with P -a.s.*

$$\kappa \in \mathcal{M}(\tau(\omega \tilde{\otimes}_t X), \omega \tilde{\otimes}_t X),$$

the measure \bar{P} defined by

$$\bar{P}(A) := \iint \mathbb{1}_A(\omega' \tilde{\otimes}_{\rho(\omega')} \omega'') \kappa(\omega'; d\omega'') P(d\omega')$$

is an element of $\mathcal{M}(t, \omega)$.

We are in the position to give a proof for Theorem 2.5.

Proof of Theorem 2.5. Thanks to Lemmata 3.1 and 3.2, the family $\{\mathcal{M}(t, \omega) : (t, \omega) \in \llbracket 0, T \rrbracket\}$ satisfies (in particular) the so-called Condition (A) in [31]. Additionally, \mathcal{M} is a nonempty, saturated (in the sense of [31]) set of local martingale measures for X , see Remark 2.3. In order to finish the proof, we need a slight extension of [31, Theorem 3.2] to the space of continuous paths starting in x_0 . This requires a minor extension of [33, Theorem 2.3], again to the space of continuous paths starting in x_0 . This can be found, for example, in [16, Theorem 4.29]. Having this result at hand, one can follow the arguments as in the proof of [31, Theorem 3.2]. \square

3.2. \mathcal{P} and \mathcal{M} are convex and compact: proof of Theorem 2.8. It suffices to prove the claims for \mathcal{P} as \mathcal{M} is the special case with $b \equiv 0$. The convexity follows from [17, Lemma III.3.38, Theorem III.3.40], see the proof of [10, Lemma 5.8] for more details. Next, we prove compactness. First, we show that \mathcal{P} is closed, which follows along the lines of the proof of [10, Proposition 3.8], and then we explain that \mathcal{P} is also relatively compact.

Let $(P^n)_{n \in \mathbb{N}} \subset \mathcal{P}$ be such that $P^n \rightarrow P$ weakly. We have to show that $P \in \mathcal{P}$, i.e., we have to prove that $P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}$ with differential characteristics in Θ . For each $n \in \mathbb{N}$, denote the P^n -characteristics of X by (B^n, C^n) .

Step 1. We start by showing that $\{P^n \circ (X, B^n, C^n)^{-1} : n \in \mathbb{N}\}$ is tight. Since $P^n \rightarrow P$, it suffices to prove tightness of $\{P^n \circ (B^n, C^n)^{-1} : n \in \mathbb{N}\}$. We use Aldous' tightness criterion ([17, Theorem VI.4.5]), i.e., we show the following two conditions:

(a) for every $\varepsilon > 0$, there exists a $K \in \mathbb{R}_+$ such that

$$\sup_{n \in \mathbb{N}} P^n \left(\sup_{s \in [0, T]} \|B_s^n\| + \sup_{s \in [0, T]} \|C_s^n\| \geq K \right) \leq \varepsilon;$$

(b) for every $\varepsilon > 0$,

$$\lim_{\theta \searrow 0} \limsup_{n \rightarrow \infty} \sup \{P^n(\|B_L^n - B_S^n\| + \|C_L^n - C_S^n\| \geq \varepsilon)\} = 0,$$

where the sup is taken over all stopping times $S, L \leq T$ such that $S \leq L \leq S + \theta$.

By the linear growth assumptions on b and a , a standard Gronwall argument (see, e.g., [20, Problem 5.3.15]) shows that

$$\sup_{P \in \mathcal{P}} E^P \left[\sup_{s \in [0, T]} \|X_s\|^2 \right] < \infty.$$

Thus, we get

$$(3.2) \quad \sup_{n \in \mathbb{N}} E^{P^n} \left[\sup_{s \in [0, T]} \|X_s\|^2 \right] \leq \sup_{P \in \mathcal{P}} E^P \left[\sup_{s \in [0, T]} \|X_s\|^2 \right] < \infty.$$

Using the linear growth assumption once again, we obtain that P^n -a.s.

$$\sup_{s \in [0, T]} \|B_s^n\| + \sup_{s \in [0, T]} \|C_s^n\| \leq C \left(1 + \sup_{s \in [0, T]} \|X_s\|^2 \right),$$

where the constant $C > 0$ is independent of n . By virtue of (3.2), this bound yields (a). For (b), take two stopping times $S, L \leq T$ such that $S \leq L \leq S + \theta$ for some $\theta > 0$. Then, using again the linear growth assumptions, we get P^n -a.s.

$$\|B_L^n - B_S^n\| + \|C_L^n - C_S^n\| \leq C(L - S) \left(1 + \sup_{s \in [0, T]} \|X_s\|^2 \right) \leq C\theta \left(1 + \sup_{s \in [0, T]} \|X_s\|^2 \right),$$

which yields (b) by virtue of (3.2). We conclude that $\{P^n \circ (X, B^n, C^n)^{-1} : n \in \mathbb{N}\}$ is tight. Up to passing to a subsequence, from now on we assume that $P^n \circ (X, B^n, C^n)^{-1} \rightarrow P \circ (X, B, C)^{-1}$ weakly.

Step 2. Next, we show that B and C are P -a.s. locally absolutely continuous. For $M > 0$ and $\omega = (\omega^{(1)}, \omega^{(2)}) \in \Omega \times \Omega$, define

$$\tau_M(\omega) := \inf(t \in [0, T] : \|\omega(t)\| \geq M) \wedge T,$$

and

$$\zeta_M(\omega) := \sup \left\{ \frac{\|\omega^{(2)}(t) - \omega^{(2)}(s)\|}{t-s} : 0 \leq s < t \leq \tau_M(\omega) \right\}.$$

Thanks to [10, Lemma 3.6], there exists a dense set $D \subset [0, T]$ such that for every $M \in D$ the map ζ_M is $P \circ (X, B)^{-1}$ -a.s. lower semicontinuous. By the linear growth conditions and the definition of τ_M , for every $M \in D$ there exists a constant $C = C(M) > 0$ such that $P^n(\zeta_M(X, B^n) \leq C) = 1$ for all $n \in \mathbb{N}$. As ζ_M is $P \circ (X, B)^{-1}$ -a.s. lower semicontinuous, [36, Example 17, p. 73] yields that

$$0 = \liminf_{n \rightarrow \infty} P^n(\zeta_M(X, B^n) > C) \geq P(\zeta_M(X, B) > C).$$

Further, since D is dense in $[0, T]$, this yields that P -a.s. B is locally Lipschitz continuous, i.e., in particular locally absolutely continuous. Similarly, we get that C is P -a.s. locally Lipschitz and hence, locally absolutely continuous. Finally, thanks to [17, Theorem IX.2.4], under P the coordinate process X is a semimartingale with characteristics (B, C) . We conclude that $P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}$.

Step 3. In this last step, we prove that $(dB/d\lambda, dC/d\lambda) \in \Theta$. By [10, Lemma 3.2], the correspondence $(t, \omega) \mapsto \Theta(t, \omega)$ is continuous, as F is compact and b and a are continuous. Hence, [10, Lemma 3.4] implies that

$$(3.3) \quad \bigcap_{m \in \mathbb{N}} \overline{\text{conv}} \Theta([t, t + 1/m], \omega) \subset \Theta(t, \omega)$$

for all $(t, \omega) \in \llbracket 0, T \rrbracket$. Here, $\overline{\text{conv}}$ denotes the closure of the convex hull. By virtue of [12, Corollary 8, p. 48], P^n -a.s. for λ -a.a. $t \in [0, T - 1/m]$, we have

$$(3.4) \quad \begin{aligned} m(B_{t+1/m}^n - B_t^n, C_{t+1/m}^n - C_t^n) &\in \overline{\text{conv}}(dB^n/d\lambda, dC^n/d\lambda)([t, t + 1/m]) \\ &\subset \overline{\text{conv}} \Theta([t, t + 1/m], X). \end{aligned}$$

Thanks to Skorokhod's coupling theorem, with little abuse of notation, there exist random variables

$$(X^0, B^0, C^0), (X^1, B^1, C^1), (X^2, B^2, C^2), \dots$$

defined on some probability space (Σ, \mathcal{G}, Q) such that (X^0, B^0, C^0) has distribution $P \circ (X, B, C)^{-1}$, (X^n, B^n, C^n) has distribution $P^n \circ (X, B^n, C^n)^{-1}$ and Q -a.s. $(X^n, B^n, C^n) \rightarrow (X^0, B^0, C^0)$ in the local uniform topology. For every $n \in \mathbb{N}$, we have Q -a.s.

$$(dB^n/d\lambda, dC^n/d\lambda) = \lim_{m \rightarrow \infty} m(B_{\cdot+1/m}^n - B_{\cdot}^n, C_{\cdot+1/m}^n - C_{\cdot}^n).$$

Now, with (3.3) and (3.4), we get that Q -a.s. for λ -a.a. $t < T$

$$(dB^0/d\lambda, dC^0/d\lambda)(t) \in \bigcap_{m \in \mathbb{N}} \overline{\text{conv}} \Theta([t, t + 1/m], X^0) \subset \Theta(t, X^0).$$

This means that $P \in \mathcal{P}$ and therefore \mathcal{P} is closed.

To finish the proof, it remains to show that \mathcal{P} is relatively compact. Thanks to Prohorov's theorem, it suffices to prove tightness, which follows from an application of Aldous' tightness criterion as in Step 1 above. We omit the details. The proof is complete. \square

3.3. Separating duality for nonlinear semimartingales: proof of Theorem 2.11. Recall that Standing Assumption 2.9 is in force. The following lemma, which is an important ingredient of the proof of Theorem 2.11, can be seen as an extension of Beneš' condition ([20, Corollary 3.5.16]). To prove the lemma we use a local change of measure in combination with a Gronwall type argument (see, e.g., [5, 8] for related strategies).

Lemma 3.3. *Take $P \in \mathcal{P}$ and denote the differential characteristics of X under P by (b^P, a^P) . There exists a predictable function $f' : \llbracket 0, T \rrbracket \rightarrow F$ such that that $(\lambda \otimes P)$ -a.e.*

$$(b^P, a^P) = (a(f')\theta(f'), a(f')),$$

where θ is as in Standing Assumption 2.9. Moreover, for $c^P := \theta(f')$, the stochastic exponential

$$Z^P := \exp \left(- \int_0^\cdot \langle c_s^P, dX_s^c \rangle - \frac{1}{2} \int_0^\cdot \langle c_s^P, a_s^P c_s^P \rangle ds \right)$$

is well-defined and a P -martingale. Here,

$$X^c := X - X_0 - \int_0^\cdot b_s^P ds$$

is the continuous local P -martingale part of X .

Proof. Let \mathcal{P} be the predictable σ -field on $\llbracket 0, T \rrbracket$. Thanks to [9, Lemma 2.12], the graph $\text{gr } \Theta$ is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{B}(\mathbb{S}_+^d)$ -measurable. Thus,

$$\begin{aligned} G &:= \{(t, \omega) \in \llbracket 0, T \rrbracket : (b_t^P(\omega), a_t^P(\omega)) \notin \Theta(t, \omega)\} \\ &= \{(t, \omega) \in \llbracket 0, T \rrbracket : (t, \omega, b_t^P(\omega), a_t^P(\omega)) \notin \text{gr } \Theta\} \in \mathcal{P}. \end{aligned}$$

We define

$$\pi(t, \omega) := \begin{cases} (b(f_0, t, \omega), a(f_0, t, \omega)), & \text{if } (t, \omega) \in G, \\ (b_t^P(\omega), a_t^P(\omega)), & \text{if } (t, \omega) \notin G, \end{cases}$$

where $f_0 \in F$ is arbitrary but fixed. Thanks to the measurable implicit function theorem [1, Theorem 18.17], as (b, a) is a Carathéodory function on $F \times \llbracket 0, T \rrbracket$ in the sense that it is continuous in the F and \mathcal{P} -measurable in the $\llbracket 0, T \rrbracket$ variable, the correspondence $\gamma: \llbracket 0, T \rrbracket \rightarrow F$ defined by

$$\gamma(t, \omega) := \{f \in F : (b(f, t, \omega), a(f, t, \omega)) = \pi(t, \omega)\}$$

is \mathcal{P} -measurable and it admits a measurable selector, i.e., there exists a \mathcal{P} -measurable function $f': \llbracket 0, T \rrbracket \rightarrow F$ such that $\pi(t, \omega) = (b(f'(t, \omega), t, \omega), a(f'(t, \omega), t, \omega))$ for all $(t, \omega) \in \llbracket 0, T \rrbracket$. Since $P \in \mathcal{P}$, we have $(\lambda \otimes P)$ -a.e. $\pi = (b^P, a^P)$, and further $b = a\theta$ by Standing Assumption 2.9. Putting these pieces together, we conclude that f' has the properties from in the statement of the lemma.

Let us now prove the martingale property of Z^P . For every $n \in \mathbb{N}$, define the stopping time

$$T_n := \inf\{t \in [0, T] : \|X_t\| \geq n\}.$$

Thanks to Standing Assumption 2.9, up to an evanescence set, the process $\langle c^P, a^P c^P \rangle$ is bounded on $\llbracket 0, T \wedge T_n \rrbracket$. Consequently, the stopped process $Z_{\cdot \wedge T_n}^P$ is a P -martingale and the global process Z^P is a well-defined, non-negative local P -martingale. Recall that latter implies that Z^P is a P -supermartingale. Thus, Z^P is a P -martingale if and only if $E^P[Z_T^P] = 1$. In the following we verify this condition. We define the probability measure Q_n via the Radon–Nikodym density $dQ_n/dP = Z_{T \wedge T_n}^P$. Clearly, since P -a.s. $Z_{T \wedge T_n}^P > 0$, we have $P \sim Q_n$ and Girsanov's theorem ([17, Theorem III.3.24]) yields that X is a Q_n -semimartingale with absolutely continuous characteristics. More precisely, the differential characteristics (b^{Q_n}, a^{Q_n}) of the coordinate process X under Q_n are given by

$$b^{Q_n} = b^P - a^P c^P \mathbf{1}_{[0, T_n]} = b^P \mathbf{1}_{]T_n, T]}, \quad a^{Q_n} = a^P.$$

Using the Burkholder–Davis–Gundy inequality, the equivalence $P \sim Q_n$ and the linear growth assumption on a , for all $t \in [0, T]$, we get

$$\begin{aligned} E^{Q_n} \left[\sup_{s \in [0, t \wedge T_n]} \|X_s\|^2 \right] &\leq C \left(1 + E^{Q_n} \left[\int_0^{t \wedge T_n} \text{tr} [a_s^P] ds \right] \right) \\ &\leq C \left(1 + E^{Q_n} \left[\int_0^{t \wedge T_n} \left(1 + \sup_{r \in [0, s]} \|X_r\|^2 \right) ds \right] \right) \\ &\leq C + C \int_0^t E^{Q_n} \left[\sup_{r \in [0, s \wedge T_n]} \|X_r\|^2 \right] ds, \end{aligned}$$

where C is a constant independent of n . Now, Gronwall's lemma yields that

$$E^{Q_n} \left[\sup_{s \in [0, t \wedge T_n]} \|X_s\|^2 \right] \leq C e^{CT}, \quad t \in [0, T].$$

Thus, using the monotone convergence theorem for the first equality, we obtain

$$E^P[Z_T^P] = \lim_{n \rightarrow \infty} E^P[Z_t^P \mathbf{1}_{\{T_n > T\}}] = \lim_{n \rightarrow \infty} Q_n(T_n > T).$$

Further, we get that

$$Q_n(T_n \leq T) = Q_n\left(\sup_{s \in [0, T \wedge T_n]} \|X_s\| \geq n\right) \leq \frac{C e^{CT}}{n^2}.$$

As the final term converges to zero as $n \rightarrow \infty$, we conclude that $E^P[Z_T^P] = 1$, which completes the proof. \square

Adapting the proof of Lemma 3.3, it is straightforward to establish a version of Lemma 3.3 for a starting measure $Q \in \mathcal{M}$. We omit a precise statement.

Lemma 3.4. *For every $P \in \mathcal{P}$, there exists $Q \in \mathcal{M}$ with $P \sim Q$. Conversely, for every $Q \in \mathcal{M}$, there exists $P \in \mathcal{P}$ with $Q \sim P$. In particular, $\mathcal{P} \sim \mathcal{M}$.*

Proof. Take $P \in \mathcal{P}$, and denote by (b^P, a^P) the differential characteristics of X under P . Let Z^P be the P -martingale from Lemma 3.3 and define a probability measure $Q \sim P$ via the Radon–Nikodym derivative $dQ/dP = Z_T^P$. Then, Girsanov’s theorem ([17, Theorem III.3.24]) shows that $Q \in \mathcal{M}$.

Conversely, let $Q \in \mathcal{M}$, and denote by a^Q the second differential characteristic of X under Q . As in the first part of the proof of Lemma 3.3, the measurable implicit function theorem [1, Theorem 18.17] implies the existence of a predictable map $f': \llbracket 0, T \rrbracket \rightarrow F$ such that $a_t^Q(\omega) = a(f'(t, \omega), t, \omega)$ for $(\lambda \otimes Q)$ -a.a. $(t, \omega) \in \llbracket 0, T \rrbracket$. Let θ be as in Standing Assumption 2.9 and define a predictable process via $c_t^Q(\omega) := \theta(f'(t, \omega), t, \omega)$ for all $(t, \omega) \in \llbracket 0, T \rrbracket$. We can argue in the same manner as in the second part of the proof of Lemma 3.3 and obtain that the process Z^Q defined by

$$Z^Q := \exp\left(\int_0^\cdot \langle c_s^Q, dX_s \rangle - \frac{1}{2} \int_0^\cdot \langle c_s^Q, a_s^Q c_s^Q \rangle ds\right),$$

is a Q -martingale. Hence, we can define a measure $P \sim Q$ via the Radon–Nikodym derivative $dP/dQ = Z_T^Q$. By Girsanov’s theorem, $P \in \mathfrak{P}_{\text{sem}}^{\text{ac}}$ and the differential characteristics of X under P are given by $(a^Q c^Q, a^Q)$. Since

$$\begin{aligned} (a_t^Q(\omega) c_t^Q(\omega), a_t^Q(\omega)) &= (a(f'(t, \omega), t, \omega) \theta(f'(t, \omega), t, \omega), a(f'(t, \omega), t, \omega)) \\ &= (b(f'(t, \omega), t, \omega), a(f'(t, \omega), t, \omega)) \in \Theta(t, \omega) \end{aligned}$$

for $(\lambda \otimes Q)$ -a.a. $(t, \omega) \in \llbracket 0, T \rrbracket$, and $P \sim Q$, we conclude that $P \in \mathcal{P}$. \square

Recall from Remark 2.7, that Lemma 3.4 entails the equality

$$\mathbf{G}^{\mathcal{M}} = \mathbf{G}^{\mathcal{P}} = \mathbf{G}.$$

We will use this throughout the remainder of this section without further mentioning.

Lemma 3.5. *Let H be a \mathbf{G} -predictable process. Then, $H \in L(X, P)$ for every $P \in \mathcal{P}$ if and only if $H \in L(X, Q)$ for every $Q \in \mathcal{Q}$. In particular, $\mathcal{H}^{\mathcal{P}} = \mathcal{H}^{\mathcal{M}}$.*

Proof. It suffices to establish the first part of the statement, since Lemma 3.4 then provides the equality $\mathcal{H}^{\mathcal{P}} = \mathcal{H}^{\mathcal{M}}$. In this regard, suppose $H \in L(X, P)$ for every $P \in \mathcal{P}$, and let $Q \in \mathcal{M}$. By Lemma 3.4, there exists a measure $P \in \mathcal{P}$ with $P \sim Q$. Hence, [26, Lemma V.2] implies that $H \in L(X, Q)$. The reverse implication follows by symmetry. \square

Proposition 3.6 (First Duality).

$$\mathcal{M} = \mathcal{D} = \{Q \in \mathfrak{P}_e(\mathcal{P}) : E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C} \cap C_b\}.$$

Proof. Replacing [3, Lemma 5.2] by Lemma 3.4, one can proceed similarly as in the proof of [3, Proposition 5.7]. For completeness, let us outline the argument. First, we set

$$(\mathcal{C} \cap C_b)^\circ := \{Q \in \mathfrak{P}_e(\mathcal{P}) : E^Q[g] \leq 1 \text{ for all } g \in \mathcal{C} \cap C_b\},$$

and $\mathcal{D} \subset (\mathcal{C} \cap C_b)^\circ$ by definition. To see $\mathcal{M} \subset \mathcal{D}$, let $g \in \mathcal{C}$ and $Q \in \mathcal{M}$. Then, by Lemma 3.5, there exists $H \in \mathcal{H}^{\mathcal{M}}$ such that

$$g \leq 1 + (H \cdot X)_T \quad Q\text{-a.s.}$$

As $H \cdot X$ is a Q -supermartingale, this implies $E^Q[g] \leq 1$. It remains to show $(\mathcal{C} \cap C_b)^\circ \subset \mathcal{M}$. To this end, we define

$$\Gamma := \{g \in C_b(\Omega) : \exists H \in \mathcal{H}^P \text{ such that } g \leq (H \cdot X)_T\}.$$

Let $Q \in (\mathcal{C} \cap C_b)^\circ$. This implies $E^Q[g] \leq 0$ for every $g \in \Gamma$. Hence, by virtue of [3, Lemma 5.6], Q is a local martingale measure. As $Q \in \mathfrak{P}_e(\mathcal{P})$, there exists $P \in \mathcal{P}$ with $Q \sim P$. Girsanov's theorem ([17, Theorem III.3.24]) together with Remark 2.3 then yield that $Q \in \mathcal{M}$. \square

Proposition 3.7 (Second Duality).

$$\mathcal{C} \cap C_b(\Omega; \mathbb{R}) = \{g \in C_b^+(\Omega; \mathbb{R}) : E^Q[g] \leq 1 \text{ for all } Q \in \mathcal{D}\}.$$

Proof. As, by definition, $\mathcal{C} \cap C_b(\Omega; \mathbb{R}) \subset \{g \in C_b^+(\Omega; \mathbb{R}) : E^Q[g] \leq 1 \text{ for all } Q \in \mathcal{D}\}$, it suffices to prove the converse inclusion $\mathcal{C} \cap C_b(\Omega; \mathbb{R}) \supset \{g \in C_b^+(\Omega; \mathbb{R}) : E^Q[g] \leq 1 \text{ for all } Q \in \mathcal{D}\}$. Let $g \in C_b^+(\Omega; \mathbb{R})$ be such that $E^Q[g] \leq 1$ for all $Q \in \mathcal{D}$. As g is bounded and continuous, the superhedging duality given in Corollary 2.6, together with the equality $\mathcal{M} = \mathcal{D}$ (Proposition 3.6), grants the existence of an $\mathbf{G}^{\mathcal{M}}$ -predictable process H such that $H \in \mathcal{H}^{\mathcal{M}}$, and

$$1 + (H \cdot X)_T \geq g \quad Q\text{-a.s. for all } Q \in \mathcal{M}.$$

As $H \in \mathcal{H}^P$ by Lemma 3.5, this shows $g \in \mathcal{C} \cap C_b$. The proof is complete. \square

Now, Theorem 2.11 follows directly from the previous two propositions.

Proof of Theorem 2.11. The duality (2.4) follows from Proposition 3.6, and the duality (2.5) follows from Proposition 3.7. \square

3.4. Duality theory for robust utility maximization: proofs of Theorems 2.17 and 2.18.

Recall that Standing Assumption 2.9 is in force. The idea of proof is to apply the abstract duality results given by [3, Theorems 2.10 and 2.16]. This requires some care to account for the lack of boundedness from above of the power utility $U(x) = \frac{x^p}{p}$, $p \in (0, 1)$, and the log utility $U(x) = \log(x)$.

3.4.1. *Some preparations.* The following lemma is a generalization of [14, Corollary 2] in the sense that instead of Brownian motion we consider a continuous local martingale with uniformly elliptic volatility. The novelty in our proof is the application of time change and comparison arguments to deduce certain moment bounds for the driving local martingale from those of Brownian motion.

Proposition 3.8. *Suppose that Condition 2.15 holds. Let $Q \in \mathcal{M}$ and denote the differential characteristics of X under P by $(b^Q = 0, a^Q)$. Furthermore, take a predictable process c^Q of linear growth, i.e., there exists a constant $C > 0$ such that*

$$\|c_t^Q(\omega)\| \leq C \left(1 + \sup_{s \in [0, t]} \|\omega(s)\|\right),$$

for $(\lambda \otimes Q)$ -a.a. $(t, \omega) \in \llbracket 0, T \rrbracket$. Define a continuous local P -martingale by

$$Z^Q := \exp \left(\int_0^\cdot \langle c_s^Q, dX_s \rangle - \frac{1}{2} \int_0^\cdot \langle c_s^Q, a_s^Q c_s^Q \rangle ds \right).$$

For every $p \geq 1$ we have

$$E^Q[(Z_T^Q)^p] < \infty.$$

Proof. Throughout the proof, fix $p \geq 1$. By virtue of [14, Corollary 1], it suffices to prove that there exists a partition $0 = t_0 < t_1 < \dots < t_m = T$ of the interval $[0, T]$ such that

$$E^Q \left[\exp \left(C_p \int_{t_{n-1}}^{t_n} \langle c_s^Q, a_s^Q c_s^Q \rangle ds \right) \right] < \infty, \quad n = 1, 2, \dots, m.$$

Fix $n \in \{1, \dots, m\}$. By the linear growth assumption on c^Q and the a.e. boundedness assumption on a^Q (which stems from the definition of \mathcal{M} and Condition 2.15), we have

$$E^Q \left[\exp \left(\mathbf{C}_p \int_{t_{n-1}}^{t_n} \langle c_s^Q, a_s^Q c_s^Q \rangle ds \right) \right] \leq CE^Q \left[\exp \left(\mathbf{C}_p (t_n - t_{n-1}) \sup_{s \in [0, T]} \|X_s\|^2 \right) \right],$$

where $\mathbf{C}_p > 0$ depends on $T > 0$ and the power p . Itô's formula shows that

$$d\|X_t\|^2 = 2\langle X_t, dX_t \rangle + \text{tr} [a_s^Q] ds.$$

Thanks to Condition 2.15, there exists a constant $\mathbf{K} > 0$ such that

$$(3.5) \quad \frac{\|\xi\|^2}{\mathbf{K}} \leq \langle \xi, a(f, t, \omega)\xi \rangle \leq \mathbf{K}\|\xi\|^2$$

for all $(\xi, f, t, \omega) \in \mathbb{R}^d \times F \times [0, T]$. Next, define

$$L := \int_0^\cdot \frac{\langle X_s, a_s^Q X_s \rangle ds}{\mathbf{K}\|X_s\|^2}.$$

Notice that the second inequality in (3.5) implies that $L_T \leq T$. Set $S_t := \inf(s \in [0, T] : L_s \geq t)$ for $t \in [0, T]$. It can be shown as in the proof of [10, Lemma 6.1] that, under Q up to time $L_T \leq T$, we have

$$d\|X_{S_t}\|^2 = 2\sqrt{\mathbf{K}}\|X_{S_t}\|dW_t + \frac{\mathbf{K}\|X_{S_t}\|^2 \text{tr} [a_{S_t}^Q]}{\langle X_{S_t}, a_{S_t}^Q X_{S_t} \rangle} dt,$$

where W is a one-dimensional Brownian motion (possibly defined on an extended probability space which we ignore in our notation for simplicity). Let Y be a $d\mathbf{K}^{5/2}$ -dimensional square Bessel process with dynamics

$$dY_t = 2\sqrt{Y_t}dW_t + d\mathbf{K}^{5/2} dt, \quad Y_0 = \|x_0\|^2/\sqrt{\mathbf{K}}.$$

It is classical that this SDE satisfies strong existence and pathwise uniqueness (see, e.g., [20, Chapter 5]). By virtue of (3.5), we get that Q -a.s. for λ -a.a. $t \in [0, L_T]$

$$\frac{\mathbf{K}\|X_{S_t}\|^2 \text{tr} [a_{S_t}^Q]}{\langle X_{S_t}, a_{S_t}^Q X_{S_t} \rangle} \leq \frac{\mathbf{K}\|X_{S_t}\|^2 \text{tr} [a_{S_t}^Q]}{\frac{1}{\mathbf{K}}\|X_{S_t}\|^2} \leq d\mathbf{K}^3,$$

and it follows as in the proof of [8, Lemma 5.6] that Q -a.s. $\|X_{S_t}\|^2/\sqrt{\mathbf{K}} \leq Y_t$ for all $t \in [0, L_T]$. Recall the well-known fact that $Y =_{\text{law}} \|B\|^2$, where B is a $d\mathbf{K}^{5/2}$ -dimensional standard Brownian motion. Thus, provided $t_{n+1} - t_n$ is taken small enough, the process

$$V := \exp \left(\frac{\sqrt{\mathbf{K}}\mathbf{C}_p(t_n - t_{n-1})Y}{2} \right)$$

is square Q -integrable and consequently, a Q -submartingale ([20, Proposition 1.3.6]). Finally, using that $L_T \leq T$ and Doob's inequality, we get

$$\begin{aligned} E^Q \left[\exp \left(\mathbf{C}_p(t_n - t_{n-1}) \sup_{s \in [0, T]} \|X_s\|^2 \right) \right] &= E^Q \left[\exp \left(\mathbf{C}_p(t_n - t_{n-1}) \sup_{s \in [0, T]} \|X_{S_{L_s}}\|^2 \right) \right] \\ &\leq E^Q \left[\sup_{s \in [0, L_T]} V_s^2 \right] \leq E^Q \left[\sup_{s \in [0, T]} V_s^2 \right] \leq 4E^Q [V_T^2] < \infty. \end{aligned}$$

This completes the proof. \square

Lemma 3.9. *Suppose that either Condition 2.14 or Condition 2.15 holds. Then, for every $P \in \mathcal{P}$ there exists a probability measure $Q \in \mathcal{M}$ such that*

$$E^Q \left[\left(\frac{dP}{dQ} \right)^p \right] < \infty, \quad \forall p > 0.$$

Proof. Take $P \in \mathcal{P}$. Similar to the proof of Lemma 3.4, we can define a particular measure $Q \in \mathcal{M}$ via the Radon–Nikodym density

$$\frac{dQ}{dP} = Z_T^P := \exp \left(- \int_0^T \langle c_s^P, dX_s^c \rangle - \frac{1}{2} \int_0^T \langle c_s^P, a_s^P c_s^P \rangle ds \right),$$

where $X^c = X - X_0 - \int_0^\cdot b_s^P ds$ is the continuous local P -martingale part of X and c^P is as in Lemma 3.3. A simple computation yields that

$$\frac{dP}{dQ} = \left(\frac{dQ}{dP} \right)^{-1} = \exp \left(\int_0^T \langle c_s^P, dX_s \rangle - \frac{1}{2} \int_0^T \langle c_s^P, a_s^P c_s^P \rangle ds \right).$$

In case Condition 2.14 holds, $\int_0^T \langle c_s^P, a_s^P c_s^P \rangle ds$ is Q -a.s. bounded and the claim follows from [14, Theorem 1]. Suppose now that Condition 2.15 holds. Notice that c^P is $(\lambda \otimes Q)$ -a.e. of linear growth by the definition of \mathcal{P} , Standing Assumption 2.1 and Condition 2.15. Therefore, Proposition 3.8 yields the claim. \square

Lemma 3.10. *Suppose that either Condition 2.14 or Condition 2.15 holds. Let $x > 0$ and $(g_n)_{n \in \mathbb{N}} \subset \mathcal{C}(x)$. Then, for every $P \in \mathcal{P}$ and every $\varepsilon \in (0, 1)$, we have*

$$\sup_{n \in \mathbb{N}} E^P [(g_n)^\varepsilon] < \infty.$$

Proof. The lemma follows similar to [3, Lemma 5.11]. Fix $\varepsilon \in (0, 1)$, $P \in \mathcal{P}$ and let $(g_n)_{n \in \mathbb{N}} \subset \mathcal{C}(x)$. Set $p := \frac{1}{1-\varepsilon}$. By virtue of Lemma 3.9, there exists a probability measure $Q \in \mathcal{M}$ with

$$E^Q \left[\left(\frac{dP}{dQ} \right)^p \right] < \infty.$$

Hence, Hölder's inequality together with Proposition 3.6 implies

$$\sup_{n \in \mathbb{N}} E^P [(g_n)^\varepsilon] \leq E^Q \left[\left(\frac{dP}{dQ} \right)^p \right]^{1-\varepsilon} E^Q [g_n]^\varepsilon \leq E^Q \left[\left(\frac{dP}{dQ} \right)^p \right]^{1-\varepsilon} x^\varepsilon < \infty,$$

which gives the claim. \square

The following estimate can be extracted from the proof of [3, Theorem 2.10].

Lemma 3.11. *Let $P \in \mathcal{P}$, and let $Q \in \mathcal{D}$ be such that $Q \ll P$. Then, for every $x, y > 0$, we have*

$$(3.6) \quad u(x) \leq \bar{u}(x) \leq v(y) + xy \leq E^P \left[\max \left\{ V_1 \left(y \frac{dQ}{dP} \right), 0 \right\} \right].$$

Finally, we present two lemmata which deal with the power and the log utility separately.

Lemma 3.12. *Suppose that either Condition 2.14 or Condition 2.15 holds. Let U be a power utility function $U(x) = \frac{x^p}{p}$ with exponent $p \in (0, 1)$. Then,*

- (i) *there exists an $x > 0$ such that $\bar{u}(x) < \infty$,*
- (ii) *for every $x > 0$ and $(g_n)_{n \in \mathbb{N}} \subset \mathcal{C}(x)$, the sequence of random variables*

$$\max \left\{ U \left(g_n + \frac{1}{n} \right), 0 \right\}, \quad n \in \mathbb{N},$$

is uniformly integrable for every $P \in \mathcal{P}$.

Proof. We start with (i). Let $P \in \mathcal{P}$. By virtue of Lemma 3.11, it suffices to construct $Q \in \mathcal{D}$ such that $Q \ll P$ and

$$E^P \left[\max \left\{ V_1 \left(y \frac{dQ}{dP} \right), 0 \right\} \right] < \infty, \quad V_1(y) = \sup_{x \geq 0} \left[\frac{(x+1)^p}{p} - xy \right].$$

This follows as in [3, Lemma 5.16], when replacing [3, Lemma 5.10] by Lemma 3.9. Regarding (ii), this can be shown as [3, Lemma 5.13] by using Lemma 3.10 instead of [3, Lemma 5.11]. \square

Lemma 3.13. *Suppose that either Condition 2.14 or Condition 2.15 holds. Let U be the log utility $U(x) = \log(x)$. Then,*

- (i) *there exists an $x > 0$ such that $\bar{u}(x) < \infty$,*

(ii) for every $x > 0$ and $(g_n)_{n \in \mathbb{N}} \subset \mathcal{C}(x)$, the sequence of random variables

$$\max \left\{ U\left(g_n + \frac{1}{n}\right), 0 \right\}, \quad n \in \mathbb{N},$$

is uniformly integrable for every $P \in \mathcal{P}$,

(iii) for each $y > 0$, and each $P \in \mathcal{P}$, there exists $Q \in \mathcal{D}$ with $Q \ll P$ such that

$$E^P \left[\max \left\{ V_1 \left(y \frac{dQ}{dP} \right), 0 \right\} \right] < \infty,$$

where

$$V_1(y) := \sup_{x \geq 0} [\log(x+1) - xy].$$

Proof. By virtue of Lemma 3.11, (iii) implies (i). To see (iii), one argues as in [3, Lemma 5.15], replacing [3, Lemma 5.10] by Lemma 3.9. Regarding (ii), this can be shown as [3, Lemma 5.12], using Lemma 3.10 instead of [3, Lemma 5.11]. \square

3.4.2. *Duality for utilities bounded from below: proof of Theorem 2.17.* Recall that in this section U is either a power utility $U(x) = \frac{x^p}{p}$, $p \in (0, 1)$, or an exponential utility $U(x) = -e^{-\lambda x}$, $\lambda > 0$. Note that both utilities are bounded from below, and the power utility is unbounded from above. Hence, by virtue of [3, Theorem 2.10], the claim follows directly from Theorems 2.8 and 2.11 in case of an exponential utility. To handle the power utility, we additionally apply Lemma 3.12. \square

3.4.3. *Duality for utilities unbounded from below: proof of Theorem 2.18.* Recall that in this section U is either a power utility $U(x) = \frac{x^p}{p}$, $p \in (-\infty, 0)$, or the log utility $U(x) = \log(x)$. Note that both utilities are unbounded from below, i.e., $U(0) = \lim_{x \rightarrow 0} U(x) = -\infty$, and the log utility is unbounded from above. Hence, by virtue of [3, Theorem 2.16], the claim follows directly from Theorems 2.8 and 2.11 in case of a power utility. To handle the log utility, we additionally apply Lemma 3.13. \square

REFERENCES

- [1] C. D. Aliprantis and K. B. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer Berlin Heidelberg, 3rd ed., 2006.
- [2] D. Bartl, P. Cheridito and M. Kupper. Robust expected utility maximization with medial limits. *Journal of Mathematical Analysis and Applications*, 471(3):752–775, 2019.
- [3] D. Bartl, M. Kupper and A. Neufeld. Duality theory for robust utility maximisation. *Finance and Stochastics*, 25(3):469–503, 2021.
- [4] S. Biagini and M. Ç. Pınar. The robust Merton problem of an ambiguity averse investor. *Mathematics and Financial Economics*, 11(1):1–24, 2017.
- [5] P. Cheridito, D. Filipović and M. Yor. Equivalent and absolutely continuous measure changes for jump-diffusion processes. *The Annals of Applied Probability*, 15(3):1713–1732, 2005.
- [6] J. C. Cox and C. F. Huang. Optimal consumption and portfolio policies when asset prices follow a diffusion process. *Journal of Economic Theory*, 49:33–83, 1989.
- [7] J. C. Cox and C. F. Huang. A variational problem arising in financial economics. *Journal of Mathematical Economics*, 20:465–487, 1991.
- [8] D. Criens. No arbitrage in continuous financial markets. *Mathematics and Financial Economics*, 14:461–506, 2020.
- [9] D. Criens and L. Niemann. Nonlinear continuous semimartingales. arXiv:2204.07823v2, 2022.
- [10] D. Criens and L. Niemann. Markov selections and Feller properties of nonlinear diffusions. arXiv:2205.15200v2, 2022.
- [11] L. Denis and M. Kervarec. Optimal investment under model uncertainty in nondominated models. *SIAM Journal on Control and Optimization*, 51(3):1803–1822, 2013.
- [12] J. Diestel and J. J. Uhl, Jr. *Vector Measures*. American Mathematical Society, 1977.

- [13] T. Fadina, A. Neufeld, and T. Schmidt. Affine processes under parameter uncertainty. *Probability, Uncertainty and Quantitative Risk*, 4(5), 2019.
- [14] B. Grigelionis and V. Mackevičius. The finiteness of moments of a stochastic exponential. *Statistics & Probability Letters*, 64:243–248, 2003.
- [15] H. He and N. D. Pearson. Consumption and portfolio policies with incomplete markets and short-sale constraints: the infinite-dimensional case. *Journal of Economic Theory*, 54(2):259–304, 1991.
- [16] J. Hollender. *Lévy-Type Processes under Uncertainty and Related Nonlocal Equations*. PhD thesis, TU Dresden, 2016.
- [17] J. Jacod and A. N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer Berlin Heidelberg, 2nd ed., 2003.
- [18] I. Karatzas, J. P. Lehoczky and S. E. Shreve. Optimal portfolio and consumption decisions for a “small investor” on a finite horizon. *SIAM Journal on Control and Optimization*, 25:1557–1586, 1987.
- [19] I. Karatzas, J. P. Lehoczky, S. E. Shreve and G. L. Xu. Martingale and duality methods for utility maximisation in an incomplete market. *SIAM Journal on Control and Optimization*, 29:702–730, 1991.
- [20] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer New York, 2nd edition, 1991.
- [21] D. Kramkov and W. Schachermayer. The asymptotic elasticity of utility functions and optimal investment in incomplete markets. *The Annals of Applied Probability* 9(3):904–950, 1999.
- [22] Z. Liang and M. Ma. Consumption–investment problem with pathwise ambiguity under logarithmic utility. *Mathematics and Financial Economics*, 13(4):519–541, 2019.
- [23] Z. Liang and M. Ma. Robust consumption-investment problem under CRRA and CARA utilities with time-varying confidence sets. *Mathematical Finance*, 30(3):1035–1072, 2020.
- [24] Q. Lin and F. Riedel. Optimal consumption and portfolio choice with ambiguous interest rates and volatility. *Economic Theory*, 71(3):1189–1202, 2021.
- [25] C. Liu and A. Neufeld. Compactness criterion for semimartingale laws and semimartingale optimal transport. *Transactions of the American Mathematical Society*, 372(1):187–231, 2019.
- [26] J. Mémin. Espaces de semi martingales et changement de probabilité. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 52(1):9–39, 1980.
- [27] P. A. Meyer. Limites médiales, d’après Mokobodzki. In *Séminaire de Probabilités VII*, pages 198–204, Springer Berlin Heidelberg, 1973.
- [28] A. Neufeld and M. Nutz. Robust utility maximization with Lévy processes. *Mathematical Finance*, 28(1):82–105, 2018.
- [29] D. Normann. Martin’s axiom and medial functions. *Mathematica Scandinavica*, 38(1):167–176, 1976.
- [30] M. Nutz. Pathwise construction of stochastic integrals. *Electronic Communications in Probability*, 17(24):1–7, 2012.
- [31] M. Nutz. Robust superhedging with jumps and diffusion. *Stochastic Processes and their Applications*, 125(12):4543–4555, 2015.
- [32] M. Nutz. Superreplication under model uncertainty in discrete time. *Finance and Stochastics*, 18(4):791–803, 2014.
- [33] M. Nutz and R. van Handel. Constructing sublinear expectations on path space. *Stochastic Processes and their Applications*, 123(8):3100–3121, 2013.
- [34] K. Park and H. Y. Wong. Robust Consumption-Investment With Return Ambiguity: A Dual Approach With Volatility Ambiguity. forthcoming in *SIAM Journal on Financial Mathematics*, 2022.
- [35] S. R. Pliska. A stochastic calculus model of continuous trading: optimal portfolio. *Mathematics of Operations Research*, 11:371–382, 1986.

- [36] D. Pollard. *Convergence of Stochastic Processes*. Springer New York, 1984.
- [37] M.-C.-Quenez Optimal portfolio in a multiple-priors model. In *Seminar on Stochastic Analysis, Random Fields and Applications IV*, vol. 58 of Progress in Probability, pages 291–321, Birkhäuser, Basel, 2004.
- [38] A. Schied. Optimal investment for robust utility functionals in complete market models. *Mathematics of Operations Research*, 30(3):750–764, 2005.
- [39] A. Schied and C.-T. Wu. Duality theory for optimal investments under model uncertainty. *Stochastics & Decisions*, 23:199-217, 2005.
- [40] D. W. Stroock and S. R. S. Varadhan. *Multidimensional Diffusion Processes*. Springer Berlin Heidelberg, reprint of 1997 ed., 2006.

ALBERT-LUDWIGS UNIVERSITY OF FREIBURG, ERNST-ZERMELO-STR. 1, 79104 FREIBURG, GERMANY

Email address: david.criens@stochastik.uni-freiburg.de

Email address: lars.niemann@stochastik.uni-freiburg.de