

# Exact sum rules for spectral zeta functions of homogeneous 1D quantum oscillators, revisited

**André Voros**

Université Paris–Saclay, CNRS, CEA, Institut de Physique Théorique  
91191 Gif-sur-Yvette, France  
E-mail: [andre.voros@ipht.fr](mailto:andre.voros@ipht.fr)

*dedicated to Professor Michael V. BERRY for his 80-th birthday*

July 12, 2022

## Abstract

We survey sum rules for spectral zeta functions of homogeneous 1D Schrödinger operators, that mainly result from the exact WKB method.

It is a great honour and pleasure to write this note for Professor Michael Berry, to celebrate his major contributions to asymptotic analysis and wave mechanics, particularly quantum and semiclassical theory, throughout his career. I have been very close to the semiclassical world since the age of 10 (Fig. 1), and thus I have greatly benefited from his generous and inspiring guidance, warm encouragements and friendship in our numerous interactions. It all started with his major survey of semiclassical theory with K. Mount, [7] just when I began my PhD. Another highlight was when he offered me (with N. Balazs and M. Tabor) to coauthor an article on quantum maps [6] which means very much to me; and he inspired my activity throughout on quantum chaology and exponential asymptotics (with more details below). Indirectly, the group around him at Bristol and the whole British school of asymptotics have also proved very stimulating, and my gratitude goes to them as well.



Figure 1: My walk to and from high school took me every day - unknowingly at the time - alongside the house where L. Brillouin was born, in the city of Sèvres, France.

We present an overview of a specific early result in exact Wentzel–Kramers–Brillouin (WKB) analysis: *exact sum rules for the spectral zeta functions* of homogeneous 1D Schrödinger operators,  $\hat{H}_N = -d^2/dq^2 + |q|^N$  on  $\mathbb{R}$  for  $N = 1, 2, \dots$  (Spectral zeta functions have been studied by Prof. Berry as well, in actually harder 2D cases. [5]) Our sum rules generalize explicit formulae obeyed by the Riemann zeta function  $\zeta$ ,

$$\zeta(2m) = \frac{(2\pi)^{2m} |B_{2m}|}{2(2m)!}, \quad m = 1, 2, \dots \quad (B_{2m} : \text{Bernoulli numbers}). \quad (1)$$

The Riemann  $\zeta$  function is indeed (up to an elementary factor) the spectral zeta function of one operator  $\hat{H}_N$ : the harmonic oscillator ( $N = 2$ ), whose spectrum is at the same time exactly solved by the basic semiclassical WKB method. This is no coincidence: the extension of (1) to other homogeneous cases is a by-product of exact WKB handling generalized to higher-degree potentials. [19][22] We have much detailed the case of the quartic potential  $q^4$  before [20, § 10]; here we sketch a more global picture disclosing some new exact identities: (30)–(35), among which (30) and (34) have fully closed forms quite similar to (1). We will skip all but minimal background and details, pointing to references instead.

## 1 The potentials $|q|^N$ and exact WKB analysis

### 1.1 Spectral functions, and notations

The (parity-symmetric) Schrödinger operator  $\hat{H}_N$  is known to have a purely discrete positive energy spectrum  $\{E_k\}_{k=0,1,\dots}$  ( $N$ -dependence is implied), with  $E_k \uparrow +\infty$ , and  $E_k \propto k^{2N/(N+2)}$  asymptotically, for  $k \rightarrow +\infty$ . It then admits full, resp. parity-twisted, *spectral generalized-zeta functions* [22, § 1]

$$Z_N(s, \lambda) \stackrel{\text{def}}{=} \sum_k (E_k + \lambda)^{-s}, \quad \text{resp. } Z_N^P(s, \lambda) \stackrel{\text{def}}{=} \sum_k (-1)^k (E_k + \lambda)^{-s} \quad (2)$$

(the sums converge for  $\text{Re } s > \frac{N+2}{2N}$ ); and definite-parity variants as well,

$$Z_N^\pm(s, \lambda) \stackrel{\text{def}}{=} \sum_{\substack{k \\ \text{even} \\ \text{odd}}} (E_k + \lambda)^{-s} = \frac{1}{2}[Z_N \pm Z_N^P](s, \lambda) \quad (3)$$

(+/- amounts to imposing a Neumann/Dirichlet condition at the center  $q = 0$ ).

We write  $Z_N^\#$  for any of  $(Z_N, Z_N^P, Z_N^+, Z_N^-)$  (and likewise for all spectral functions). Either pair  $(Z_N, Z_N^P)$  or  $(Z_N^+, Z_N^-)$  suffices to generate the quartet, but formulae may favour any of the four types, so we keep them all.

The functions  $Z_N^\#(s, \lambda)$  are *meromorphic in the whole  $s$ -plane and regular at  $s = 0$* . For us they mainly serve to generate two special (sets of) spectral functions in one variable,

$$D_N^\#(\lambda) \stackrel{\text{def}}{=} \exp -Z_N^{\#'}(s, \lambda)|_{s=0} \quad (' = \partial_s) \quad (\text{spectral determinants}), \quad (4)$$

$$Z_N^\#(s) \stackrel{\text{def}}{=} Z_N^\#(s, \lambda = 0) \quad (\text{spectral zeta functions}). \quad (5)$$

The latter zeta functions go back to [12, § 7] in general, and to [14] for our  $|q|^N$  case. They enter these fundamental expansion formulae for the determinants:

$$\log D_N^\#(\lambda) = -Z_N^\#(0) - \sum_{n=1}^{\infty} \frac{Z_N^\#(n)}{n} (-\lambda)^n. \quad (6)$$

The radius of convergence here is the lowest relevant eigenvalue, always  $> 0$ . And the values  $Z_N(1)$ ,  $Z_N^\pm(1)$  for  $N = 1, 2$  (for which the defining series diverge) are meant as regularized values or finite parts.

Being homogeneous,  $\hat{H}_N$  is invariant modulo dilations under complex rotations  $q \mapsto e^{i\varphi_N} q$  with  $\varphi_N \stackrel{\text{def}}{=} 4\pi/(N+2)$ ; the order  $L_N$  of that symmetry (the lowest integer  $L > 0$  making  $L\varphi_N \equiv 0 \pmod{2\pi}$ ) is

$$L_N = \frac{1}{2}N + 1 \quad \text{for } N \text{ even,} \quad N + 2 \quad \text{for } N \text{ odd.} \quad (7)$$

The degree  $N$  will most often occur through the combination

$$\nu_N \stackrel{\text{def}}{=} 1/(N+2) = \varphi_N/(4\pi) \quad (\text{to be denoted } \nu \text{ for maximum brevity}). \quad (8)$$

## 1.2 Exact WKB approaches

Exact WKB analysis uses the idea that for 1D potentials analytic in the complex  $q$ -plane, the ordinary complex-WKB equations for the Schrödinger solution can be reinterpreted to describe its analytical continuation in  $q$ , an operation that gives *fully exact* results. The topic much grew around 1980, but quite a few earlier works as quoted in [23, § 1.2] were crucial to it; we owe Prof. Berry our awareness of an essential building block, Dingle's treatment of asymptotics. [10] It is much beyond the scope of this note (limited to a special early application) to go into the overall subject - which now has several variants, and we know of no truly global reference either.

Our own use of an exact WKB scheme was in part triggered by the Berry–Mount survey quoted above, as it very clearly exposed the complex WKB method and highlighted its limitations: thus, for the simplest (i.e., parabolic) quantum 1D barrier problem, even that most efficient semiclassical method left the *phase* of the transmission coefficient *undetermined*. [7, § 3.3] We were at the time aiming that complex WKB method at a quartic oscillator, [4] but clearly the above issue was standing in our way and had to be settled first. The upshot was that using *complex*  $\hbar$ , and borrowing from Balian–Bloch, [3] Sato–Kawai–Kashiwara, [15] and Dingle, [10] the complex WKB method does describe the full transmission coefficient (or Jost function) *exactly* for a quadratic barrier [20, § 7] - an approach which then at once extended to a quartic potential and beyond.

In the end, the primary concrete output of an exact WKB analysis of the operator  $\hat{H}_N$  is an exact bilinear functional equation in the style of Sibuya [16, chap. 5] constraining the spectral determinants  $D_N^\pm$ : [19][21][22]

$$\begin{aligned} e^{i\nu\pi} D_N^+(\lambda) D_N^-(e^{4i\nu\pi} \lambda) - e^{-i\nu\pi} D_N^+(e^{4i\nu\pi} \lambda) D_N^-(\lambda) &= 2i \quad (N \neq 2) \quad (9) \\ &= 2i e^{-i\pi\lambda/4} \quad (N = 2). \quad (10) \end{aligned}$$

Then, by enforcing the compatibility of (9) with the expansions (6), concrete polynomial identities for the *exact* values  $Z_N^{\#'}(0), Z_N^{\#}(n)$  ( $n = 1, 2, \dots$ ) will arise.

Such results do not fall into the same class as earlier (also exact)  $Z$ -value formulae but at the *negative* integers  $(-n)$ ,  $n = 0, 1, \dots$ : *trace identities*. [14] The latter arise through expanding the partition function  $\sum_k e^{-tE_k}$  for  $t \rightarrow 0^+$ , an operation wholly belonging to conventional (not exact) asymptotic analysis.  $Z$ -values at *positive* integers have an unrelated, more transcendental, nature. (For the Riemann  $\zeta$  function,  $\zeta(-n)$  and  $\zeta(1+n)$  are connected by the Riemann Functional Equation, but *this* is an arithmetic feature special to this case.)

## 2 Special known cases

### 2.1 The harmonic oscillator ( $N = 2$ )

$\hat{H}_2$  has the explicit spectrum  $\{E_k = 2k + 1\}_{k=0,1,\dots}$  giving the spectral determinants

$$D_2(\lambda) = 2^{-\lambda/2} \sqrt{2\pi} / \Gamma(\frac{1+\lambda}{2}), \quad D_2^P(\lambda) = 2 \Gamma(\frac{3+\lambda}{4}) / \Gamma(\frac{1+\lambda}{4}), \quad (11)$$

and spectral zeta functions which are the Dirichlet lambda and beta functions respectively: [1, eqs. (23.2.20–21)]

$$Z_2(s) = (1 - 2^{-s})\zeta(s), \quad Z_2^P(s) = \beta(s). \quad (12)$$

The functional equation (10) obeyed here by the determinants amounts to the *reflection formula for the Gamma function* and can be split as [23, § 4.1]

$$D_2(\lambda)D_2(-\lambda) = 2 \cos \frac{1}{2}\pi\lambda, \quad D_2^P(\lambda)/D_2^P(-\lambda) = \cot \frac{1}{4}\pi(1 - \lambda). \quad (13)$$

In order to make use of (6), we take the logarithms and expand around  $\lambda = 0$ , then identify the resulting sides order by order.

With  $D_2$  we thus get

$$2 \left[ -Z_2'(0) - \sum_{m=1}^{\infty} \frac{Z_2(2m)}{2m} \lambda^{2m} \right] \equiv \log 2 - \sum_{m=1}^{\infty} \frac{|G_{2m}|}{4m(2m)!} (\pi\lambda)^{2m}, \quad (14)$$

with the right-hand side using the suitable primitive of a classic Taylor formula,  $\tan \frac{1}{2}\pi\lambda = \sum_{m=1}^{\infty} \frac{|G_{2m}|}{(2m)!} (\pi\lambda)^{2m-1}$  ( $G_{2m}$ : Genocchi numbers). [13]

With  $D_2^P$  we likewise get

$$2 \sum_{m=0}^{\infty} \frac{Z_2^P(2m+1)}{2m+1} \lambda^{2m+1} \equiv \sum_{m=0}^{\infty} \frac{|E_{2m}|}{(2m+1)!} (\frac{1}{2}\pi\lambda)^{2m+1} : \quad (15)$$

the right-hand side now came by integration from  $\sec \frac{1}{2}\pi\lambda = \sum_{m=0}^{\infty} \frac{|E_{2m}|}{(2m)!} (\frac{1}{2}\pi\lambda)^{2m}$  ( $E_{2m}$ : Euler numbers). Overall, we have thus reached:  $Z_2'(0) = -\frac{1}{2} \log 2$ , and

[1, chap. 23]

$$Z_2(2m) = \frac{\pi^{2m}}{4(2m)!} |G_{2m}| \quad (m > 0), \quad Z_2^P(2m+1) = \frac{(\frac{1}{2}\pi)^{2m+1}}{2(2m)!} |E_{2m}|. \quad (16)$$

The identities:  $G_{2m} \equiv 2(1 - 2^{2m})B_{2m}$ , [13] and (12) for  $Z_2$ , together make the formula (16) for  $Z_2(2m)$  equivalent to (1) for  $\zeta(2m)$  with Bernoulli numbers. But (16) with Genocchi numbers is more *symmetric*: the  $G_{2m}$  are *integers*  $(-1, 1, -3, 17, \dots)$ , just like the  $E_{2m}$ . Moreover, we deliberately drew (16) (or (1)) straight from the Gamma-function reflection formula: not only this approach is most elementary, but it specifically extends to anharmonic cases.

## 2.2 The case $N = 1$

The spectral determinants for the potential  $|q|$  use the Airy function, [22, § 3]

$$D_1^+(\lambda) = -2\sqrt{\pi}\text{Ai}'(\lambda), \quad D_1^-(\lambda) = 2\sqrt{\pi}\text{Ai}(\lambda) \quad (17)$$

(just for this §:  $' \stackrel{\text{def}}{=} d/d\lambda$ ). A special property here, not seen for spectral determinants  $D_N^\#$  with  $N > 1$ , is that  $\text{Ai}(\lambda)$  obeys a *linear differential equation* (Airy's equation), making all its Taylor coefficients at 0 explicit:  $\text{Ai}^{(n)}(0) = 3^{(n-2)/3}\pi^{-1} \sin \frac{2}{3}(n+1)\pi \Gamma(\frac{n+1}{3})$ . [1, chap. 10.4] Then,  $\log D_1^\pm$  can in turn be expanded to any order in closed form, giving

$$Z_1^{+'}(0) = \log \frac{3^{1/3}\Gamma(1/3)}{2\sqrt{\pi}}, \quad Z_1^{-'}(0) = \log \frac{3^{2/3}\Gamma(2/3)}{2\sqrt{\pi}} \quad (\text{regularized sums}), \quad (18)$$

$$Z_1^+(1) = 0, \quad Z_1^-(1) = -\rho \quad ( \quad " \quad " \quad ) \quad (19)$$

$$Z_1^+(2) = 1/\rho, \quad Z_1^-(2) = \rho^2, \quad (20)$$

$$Z_1^+(3) = 1, \quad Z_1^-(3) = \frac{1}{2} - \rho^3, \quad (21)$$

$\vdots$

$\vdots$

$$\text{with } \rho \stackrel{\text{def}}{=} D_1^P(0) = -\text{Ai}'(0)/\text{Ai}(0) = 3^{5/6}(2\pi)^{-1}\Gamma(2/3)^2 (\approx 0.729011133). \quad (22)$$

Here the main exact-WKB functional equation (9) reduces to a classic Wronskian relation [1, eq. (10.4.12)].

The spectral zeta function  $Z_1^-$  (aka Airy zeta function) is covered in earlier literature [9, § 4], but not its partner  $Z_1^+$  until [22], to our knowledge. In particular we find no previous mention of the exceptional rational value  $Z_1^+(3) = 1$  (the inverse cubes of the unsigned zeros of  $\text{Ai}'$  sum up to unity!).

## 3 The general 1D homogeneous ( $N$ integer) case

### 3.1 General results

For general potentials  $|q|^N$ , the extension of the Gamma-function reflection formula from the case  $N = 2$  is thus the functional relation (9)–(10). If we now

expand it in powers  $\lambda^n$  using (6) then identify sides order by order, we attain a countable sequence of *exact identities* or *sum rules* (where  $' = d/ds$  again):

$$(n = 0 :) \quad Z'_N(0) = \log \sin \nu\pi \quad (23)$$

$$- \cot \nu\pi \sin 2\nu\pi Z_N^P(1) + \cos 2\nu\pi Z_N(1) = 0 \quad [\text{indeterminacy for } N = 2] \quad (24)$$

$$- \cot \nu\pi \sin 4\nu\pi Z_N^P(2) + \cos 4\nu\pi Z_N(2) = -4 \cos^2 \nu\pi Z_N^P(1)^2 \quad (25)$$

$$- \cot \nu\pi \sin 6\nu\pi Z_N^P(3) + \cos 6\nu\pi Z_N(3) = 4 \cos^2 \nu\pi [2 \cos^2 \nu\pi Z_N^P(1)^3 \\ - 3 \cos 2\nu\pi Z_N^P(1) Z_N^P(2)] \quad (26)$$

$\vdots$

$$- \cot \nu\pi \sin 2n\nu\pi Z_N^P(n) + \cos 2n\nu\pi Z_N(n) = \mathcal{P}_{N,n}\{Z_N^P(m)\}_{1 \leq m < n} \quad (27)$$

with  $\mathcal{P}_{N,n}$  being homogeneous polynomials of degree  $n$  if  $Z_N^P(m)$  are assigned degree  $m$ . (A path linking (9) to (23)–(27) is best detailed in [21, § IV.2–3].)

For  $N = 2$ , (27) expresses  $Z_2(n)$  for  $n > 1$  even and  $Z_2^P(n)$  for  $n$  odd, like (16). So, (27) does generalize the harmonic-case result (16) - but with two shortcomings:

a) the left-hand side of (27) is not a basic zeta-value  $Z_N^\#$  (i.e.,  $Z_N$  or  $Z_N^P$  or  $Z_N^\pm$ ) in general, but rather one weird linear combination thereof (in which the weights recur with  $n$ -period  $L_N$ );

b) the right-hand side in (27) is not explicit in general, because hardly any values  $\{Z_N^P(m)\}_{m < n}$  are determined by the above recursion in  $n$ : only one linear combination gains an expression at each step, when two would be needed.

To mitigate a), the left-hand side of (27) is *sometimes* one of the basic  $Z_N^\#$ :

- $Z_N$  if  $n = \ell L_N$  for  $\ell$  integer;
- $Z_N^P$  if  $N \equiv 2 \pmod{4}$  and  $n = \frac{1}{2} \ell L_N$  for  $\ell$  odd;
- $Z_N^\pm$  if  $N$  is odd and  $n = (\ell + \frac{1}{2}) L_N \mp \frac{1}{2}$  for  $\ell$  integer.

To mitigate b), *all values for  $n = 0$  and 1* are actually known *in closed form*, by earlier extraneous (i.e., not exact-WKB) arguments: [18]

$$n = 0: \quad (Z_N^P)'(0) = \log[\nu^{N\nu} \Gamma(\nu)/\Gamma(1-\nu)], \quad Z'_N(0) = \log \sin \nu\pi \quad (28)$$

(curiously, the exact-WKB method recovers  $Z'_N(0)$  as (23), but not at all  $Z_N^{P'}(0)$ );

$$n = 1: \quad Z_N^P(1) = \frac{\sin \nu\pi}{2\sqrt{\pi}} (2\nu)^{2N\nu} \frac{\Gamma(\nu)\Gamma(2\nu)\Gamma(3\nu)}{\Gamma(2\nu + \frac{1}{2})}, \quad Z_N(1) = \frac{\tan 2\nu\pi}{\tan \nu\pi} Z_N^P(1) \quad (29)$$

(by Weber–Schafheitlin integral formulae; the exact-WKB method still misses  $Z_N^P$  at  $n = 1$ , but it does catch the *ratio*  $Z_N(1)/Z_N^P(1)$  this time, through (24)).

### 3.2 Examples

For convenience, Tables 1–2 list the closed-form expressions (28)–(29) and the exact identities (23)–(26) for some values of  $N$  (splitting even vs odd  $N$  because the parameter  $L_N$  given by (7) will recur later).

We then present a synoptic view in Tables 3–4.

$N :$	2	4	6
$Z_N^{\text{P}'(0)}$	$\log \frac{\Gamma(1/4)^2}{2^{3/2}\pi}$	$\log \frac{3^{1/3}\Gamma(1/3)^4}{2^{7/3}\pi^2}$	$\log \frac{2^{-9/4}\Gamma(1/8)}{\Gamma(7/8)}$
$Z_N^{\text{P}(1)}$	$\pi/4$	$\frac{\Gamma(1/3)^5}{2^{11/3}3^{1/3}\pi^2}$	$\frac{\pi\Gamma(5/4)}{2^{7/4}\Gamma(7/8)^2}$
$Z_N'(0)$	$-\frac{1}{2}\log 2$	$-\log 2$	$\frac{1}{2}\log[(2-\sqrt{2})/4]$
$Z_N(1)$	$\infty$	$3Z_4^{\text{P}(1)}$	$(1+\sqrt{2})Z_6^{\text{P}(1)}$
$n = 2$	$Z_2(2) = 2Z_2^{\text{P}(1)^2}$	$3Z_4^{\text{P}(2)} + Z_4(2) = 6Z_4^{\text{P}(1)^2}$	$Z_6^{\text{P}(2)} = \sqrt{2}Z_6^{\text{P}(1)^2}$
$n = 3$	$Z_2^{\text{P}(3)} = 2Z_2^{\text{P}(1)^3}$	$Z_4(3) = \frac{9}{2}[-Z_4^{\text{P}(1)^3} + Z_4^{\text{P}(1)}Z_4^{\text{P}(2)}] = \frac{1}{6}Z_4(1)^3 - \frac{1}{2}Z_4(1)Z_4(2)$	$(\sqrt{2}+1)Z_6^{\text{P}(3)} + Z_6(3) = -(3\sqrt{2}+4)Z_6^{\text{P}(1)^3} + 3(2+\sqrt{2})Z_6^{\text{P}(1)}Z_6^{\text{P}(2)}$

Table 1: For  $N$  even, top 4 rows ( $n = 0, 1$ ): closed-form values from (28)–(29); bottom 4 rows ( $n = 0$  to 3): exact-WKB sum rules from (23)–(26); the two sets share the middle 2 rows.

$N :$	1	3
$Z_N^{\text{P}'(0)}$	$-\log \rho$	$\log \frac{5^{-3/5}\Gamma(1/5)}{\Gamma(4/5)}$
$Z_N^{\text{P}(1)}$	$\rho$	$(\frac{2}{5})^{1/5} \frac{\sqrt{\pi}\Gamma(6/5)}{\phi\Gamma(9/10)}$
$Z_N'(0)$	$\log[\sqrt{3}/2]$	$\frac{1}{2}\log[(5-\sqrt{5})/8]$
$Z_N(1)$	$-Z_1^{\text{P}(1)}$	$(2+\sqrt{5})Z_3^{\text{P}(1)}$
$n = 2$	$Z_1^-(2) = Z_1^{\text{P}(1)^2}$	$Z_3^+(2) = \phi Z_3^{\text{P}(1)^2}$
$n = 3$	$Z_1(3) = \frac{1}{2}Z_1^{\text{P}(1)^3} + \frac{3}{2}Z_1^{\text{P}(1)}Z_1^{\text{P}(2)} = \frac{5}{2}Z_1(1)^3 - \frac{3}{2}Z_1(1)Z_1(2)$	$Z_3^-(3) = -(\phi + \frac{1}{2})Z_3^{\text{P}(1)^3} + \frac{3}{2}Z_3^{\text{P}(1)}Z_3^{\text{P}(2)}$

Table 2: As Table 1 but for  $N$  odd. In the  $N = 1$  column:  $\rho$  as in (22); in the  $N = 3$  column:  $\phi = \frac{1}{2}(\sqrt{5} + 1)$ , the golden ratio.

Both Tables 3–4 highlight as *unshaded* the cells with values admitting *fully explicit* closed forms. That includes: the leftmost column in each Table, from (16) and (18)–(21) respectively (here, exact-WKB sum rules alone determine only half the data for  $N = 2$ , and  $Z_1'(0)$  for  $N = 1$ ); then the top 2 rows ( $n = 0, 1$ ) from (28)–(29), determining complete data ( $Z_N$  and  $Z_N^{\text{P}}$ ). Further closed forms fill the third rows ( $n = 2$ ): these are now specifically *exact-WKB results* from the sum rule (25) helped by (29), but that only covers one specific linear combination of  $Z_N(2)$  and  $Z_N^{\text{P}}(2)$  at each  $N$ . ( $Z_N(n)$ ,  $Z_N^{\text{P}}(n)$  values taken separately are known only in terms of generalized hypergeometric  ${}_4F_3$  infinite series for  $n = 2$ , and of  $n$ -uple definite integrals for general  $n$ : [20, App. C] those we do not call fully closed forms.)

Two, and only two, exceptional basic spectral-zeta values  $Z_N^{\#}(n)$  then lie

$n \setminus N$	2	4	6	8	10	...
0	$Z'_2, Z_2^{\# \prime}$	$Z'_4, Z_4^{\# \prime}$	$Z'_6, Z_6^{\# \prime}$	$Z'_8, Z_8^{\# \prime}$	$Z'_{10}, Z_{10}^{\# \prime}$	...
1	$Z_2^P$	←	$*, Z_N^{\#}$	→		...
2	$Z_2$	*	$Z_6^P$	*	*	*
3	$Z_2^P$	$Z_4$	*	*	$Z_{10}^P$	*
4	$Z_2$	*	$Z_6$	*	*	...
5	$Z_2^P$	*	*	$Z_8$	*	
6	$Z_2$	$Z_4$	$Z_6^P$	*	$Z_{10}$	
⋮	⋮	⋮	⋮	⋮	⋮	⋮

Table 3: In black: the entries in the exact-WKB spectral sum rules (23)–(27) for  $N$  even; \* stands for left-hand sides in generic form. (Save for the derivative signs  $'$  in the top row, each  $N$ -column has a pattern of  $n$ -period  $L_N = \frac{1}{2}N + 1$  as given by (7)). In grey: additional exact data in closed form, from (28)–(29). In shaded cells: data *not* ending up in fully closed form.

$n \setminus N$	1	3	5	7	9	...
0	$Z'_1, Z_1^{\# \prime}$	$Z'_3, Z_3^{\# \prime}$	$Z'_5, Z_5^{\# \prime}$	$Z'_7, Z_7^{\# \prime}$	$Z'_9, Z_9^{\# \prime}$	...
1	$Z_1^+, Z_1^{\#}$	←	$*, Z_N^{\#}$	→		...
2	$Z_1^-, Z_1^{\#}$	$Z_3^+$	*	*	*	*
3	$Z_1, Z_1^{\#}$	$Z_3^-$	$Z_5^+$	*	*	*
4	$Z_1^+, Z_1^{\#}$	*	$Z_5^-$	$Z_7^+$	*	*
5	$Z_1^-, Z_1^{\#}$	$Z_3$	*	$Z_7^-$	$Z_9^+$	*
6	$Z_1, Z_1^{\#}$	*	*	*	$Z_9^-$	⋮
7	$Z_1^+, Z_1^{\#}$	$Z_3^+$	$Z_5$	*	*	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮

Table 4: As with Table 1, but for  $N$  odd hence now  $L_N = N + 2$  by (7), and adding exact data from (18)–(21) (also in grey).



among those new ( $N > 2$  and  $n = 2$ ) closed forms from (25), thus creating the closest anharmonic analogs of (16):

- with  $N$  even (Table 3), our results for  $N = 4$  are not new [19]; for the *sextic* oscillator on the other hand, using  $Z_6^P(1)$  from Table 1:

$$Z_6^P(2) = \sqrt{2} Z_6^P(1)^2 = \frac{1}{8} [\pi \Gamma(\frac{5}{4})]^2 / \Gamma(\frac{7}{8})^4 (\approx 0.71895230), \quad (30)$$

which allows (26) to add one more closed-form result at  $n = 3$ ,

$$\begin{aligned} (1 + \sqrt{2}) Z_6^P(3) + Z_6(3) &= (3\sqrt{2} + 2) Z_6^P(1)^3 \\ &= (3 + \sqrt{2}) 2^{-19/4} [\pi \Gamma(\frac{5}{4})]^3 / \Gamma(\frac{7}{8})^6 (\approx 2.26279887). \end{aligned} \quad (31)$$

The next identity, for  $n = 4$ , is also new and can be written as

$$Z_6(4) = \frac{1}{3}(248 - 175\sqrt{2}) Z_6(1)^4 - \frac{4}{3}(2 - \sqrt{2}) Z_6(1) Z_6(3) \quad (32)$$

(but no longer in fully closed form, because we do not know  $Z_6(3)$  analytically). For completeness, we also specify the last marked ( $N = 6$ )-cell in Table 3:

$$\begin{aligned} Z_6^P(6) &= -\frac{1}{30}(210 + 151\sqrt{2}) Z_6^P(1)^6 + \frac{1}{2}(34 + 23\sqrt{2}) Z_6^P(1)^4 Z_6^P(2) \\ &\quad - \frac{1}{2}(18 + 15\sqrt{2}) Z_6^P(1)^2 Z_6^P(2)^2 + \frac{1}{2}(2 + \sqrt{2}) Z_6^P(2)^3 \\ &\quad - \frac{2}{3}(6 + 5\sqrt{2}) Z_6^P(1)^3 Z_6^P(3) + 2(2 + \sqrt{2}) Z_6^P(1) Z_6^P(2) Z_6^P(3) \\ &\quad - \frac{1}{3}\sqrt{2} Z_6^P(3)^2 + \frac{6}{5}\sqrt{2} Z_6^P(1) Z_6^P(5). \end{aligned} \quad (33)$$

Numerical confirmations can be obtained from [19, Table] (or [20, § 10 and Table 5], but together with the *errata* on the quoted Web page).

- with  $N$  odd (Table 4), for the *cubic* oscillator, [16, chap. 5][22, § 3] using  $Z_3^P(1)$  from Table 2 and the golden ratio  $\phi = \frac{1}{2}(1 + \sqrt{5})$ :

$$Z_3^+(2) = \phi Z_3^P(1)^2 = (\frac{2}{5})^{2/5} \phi^{-1} \pi [\Gamma(\frac{6}{5}) / \Gamma(\frac{9}{10})]^2 (\approx 0.99352218); \quad (34)$$

unlike (31), we see no closed-form result for  $n = 3$  as we lack a value for  $Z_3^P(2)$ , which our sum rules leave undetermined. The next identity for a basic zeta-value (Table 4), also new, can be written as

$$\begin{aligned} Z_3(5) &= \frac{1}{48}(369163 - 165095\sqrt{5}) Z_3(1)^5 + \frac{5}{24}(2503 - 1119\sqrt{5}) Z_3(1)^3 Z_3(2) \\ &\quad + \frac{5}{16}(23 - 11\sqrt{5}) Z_3(1) Z_3(2)^2 + \frac{5}{6}(-31 + 14\sqrt{5}) Z_3(1)^2 Z_3(3) \\ &\quad - \frac{5}{6} Z_3(2) Z_3(3) + \frac{5}{8}(-7 + 3\sqrt{5}) Z_3(1) Z_3(4) \quad (\approx 0.894912). \end{aligned} \quad (35)$$

Note: this and (32) exemplify the general fact that for all  $n$  multiple of  $L_N$ , the  $Z_N(n)$  also express in terms of  $Z_N$ -values alone; that is because the full determinant  $D_N(\lambda)$  inherits from (9) an *autonomous* functional equation with a symmetry of order  $L_N$ . [22] (For  $N = 1$  and 4, both having  $L_N = 3$ ,  $Z_1(3)$  is thus recalled in Table 2,  $Z_4(3)$  in Table 1, and  $Z_4(6)$  in [19, eq. (6)].)

All those examples very concretely testify to the capacity of the WKB framework to yield *analytically exact results* for anharmonic oscillators.

### 3.3 Concluding remarks

As further results, spectral sum rules have been found for more general potentials [17][9], 2D billiards [11][5] and compact hyperbolic surfaces (Selberg-zeta zeros), [24, App. B] plus analogous sum rules for the Riemann zeros as well. [24, Tables]

Now, the early exact-WKB results we have reviewed don't close the story; fortunately so, because they appear heavily underdetermined (one functional equation (9) for two unknown functions  $D_N^\pm$ , one sum rule (27) for two independent  $Z$ -values at each  $n$ ). But actually, obvious analyticity constraints on  $D_N^\pm$  suffice to obtain (from (9) alone) separate *fixed-point equations* for  $D_N^+$  and  $D_N^-$ , [21][22] proven to be *contracting* in this homogeneous-potential case: [2] thereby, both  $D_N^\pm$  end up *fully determined*, which means exact and effective quantization of the spectrum. However, that extra information beyond (9) seems of a more global nature, and not reducible to a new set of finite spectral sum rules completing (27) (except for  $N = 1$ , as shown by (18)–(21)).

Finally, exact WKB treatments also work to solve general (inhomogeneous polynomial) potentials (as reviewed in [23], and more lately [25]).

**Acknowledgment:** we used Mathematica [26] to attain (32), (33), (35).

## References

- [1] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions*, Dover, New York (1965).
- [2] A. Avila, *Convergence of an exact quantization scheme*, Commun. Math. Phys. **249** (2004) 305–318.
- [3] R. Balian and C. Bloch, *Solutions of the Schrödinger equation in terms of classical paths*, Ann. Phys. (NY) **85** (1974) 514–545.
- [4] R. Balian, G. Parisi, A. Voros, *Quartic oscillator*, in: S. Albeverio et al. (eds.), *Feynman Path Integrals* [(Proceedings, Marseille 1978), Lecture Notes in Physics **106**, Springer, Berlin (1979) 337–360 (errata: [18, footnote p. 209] or [8, p. 203–204])].
- [5] M.V. Berry, *Spectral zeta functions for Aharonov–Bohm quantum billiards*, J. Phys. **A19** (1986) 2281–2296.
- [6] M.V. Berry, N.L. Balazs, M. Tabor and A. Voros, *Quantum maps*, Ann. Phys. (NY) **122** (1979) 26–63.
- [7] M.V. Berry and K.E. Mount, *Semiclassical approximations in wave mechanics*, Rep. Prog. Phys. **35** (1972) 315–397.
- [8] D. Chudnovsky and G. Chudnovsky (eds.), *The Riemann Problem, Complete Integrability and Arithmetic Applications* (Proceedings, IHÉS and Columbia University, 1979–1980), Lecture Notes in Mathematics **925**, Springer, Berlin (1982).

- [9] R. E. Crandall, *On the Quantum Zeta Function*, J. Phys. **A29** (1996) 6795–6816.
- [10] R.B. Dingle, *Asymptotic Expansions: their Derivation and Interpretation*, Academic Press, London (1973).
- [11] C. Itzykson, P. Moussa and J.-M. Luck, *Sum rules for quantum billiards*, J. Phys. **A19** (1986) L111-L115.
- [12] S. Minakshisundaram, Å. Pleijel, *Some properties of the eigenfunctions of the Laplace-operator on Riemannian manifolds*, Can. J. Math. **1** (1949) 242–256.
- [13] The OEIS Foundation Inc. (2022), Entry A001469 in The On-Line Encyclopedia of Integer Sequences, <http://oeis.org/A001469> .
- [14] G. Parisi, *Trace identities for the Schrödinger operator and the WKB method*, Preprint LPTENS 78/9 (École Normale Supérieure, Paris, March 1978), in [8, p.178–183].
- [15] M. Sato, T. Kawai and M. Kashiwara, *Microfunctions and pseudo-differential equations*, in: H. Komatsu (ed.), *Hyperfunctions and pseudo-differential equations* (Proceedings, Katata 1971), Lecture Notes in Mathematics **287**, Springer, Berlin (1973) 265–529.
- [16] Y. Sibuya, *Global theory of a second order linear ordinary differential operator with a polynomial coefficient*, North-Holland, Amsterdam (1975).
- [17] F. Steiner, *Magic sum rules for confinement potentials*, Phys. Lett. **B159** (1985) 397–402.
- [18] A. Voros, *The zeta function of the quartic oscillator*, Nucl. Phys. **B165** (1980) 209–236 (errata : [www.ipht.fr/Docspht/articles/t79/046/public/erratum.pdf](http://www.ipht.fr/Docspht/articles/t79/046/public/erratum.pdf)), augmented version: *Zeta functions of the quartic (and homogeneous anharmonic) oscillators*, in [8] p.184–208.
- [19] A. Voros, *Correspondance semi-classique et résultats exacts : cas des spectres d'opérateurs de Schrödinger homogènes*, C.R. Acad. Sci., Paris, Sér. I **293** (1981) 709–712 (<https://gallica.bnf.fr/ark:/12148/bpt6k64461526/f351.image>), English translation: *Semiclassical correspondence and exact results: the case of the spectra of homogeneous Schrödinger operators*, J. Physique Lettres **43** (1982) L-1–L-4 (erratum: ibid. p.159).
- [20] A. Voros, *The return of the quartic oscillator. The complex WKB method*, Ann. Inst. H. Poincaré **A39** (1983) 211–338 ([eudml.org/doc/76217](http://eudml.org/doc/76217) with errata in § “Notes”).

- [21] A. Voros, *Exact anharmonic quantization condition (in one dimension)*, in: *Quasiclassical methods* (Proceedings, IMA, Minneapolis 1995), eds. J. Rauch and B. Simon, IMA series vol. **95**, Springer-Verlag, New York (1997) 189–224.
- [22] A. Voros, *Airy function (exact WKB results for potentials of odd degree)*, J. Phys. **A32** (1999) 1301–1311 [Errata: in eqs. (18),  $D^\pm(e^{-i\varphi} \lambda)$  should read  $D^\pm(-e^{-i\varphi} \lambda)$  (twice); and underneath,  $[0, e^{-i\varphi} \infty)$  should read  $[0, -e^{-i\varphi} \infty)$ ];
- [23] A. Voros, *The general 1D Schrödinger equation as an exactly solvable problem*, in: Y. Takei (ed.), *Recent trends in exponential asymptotics* (Proceedings, Kyoto 2004), RIMS Kôkyûroku **1424** (2005) 214–231.
- [24] A. Voros, *Zeta functions over zeros of zeta functions*, Lecture Notes of the Unione Matematica Italiana **8**, Springer-Verlag (2010), and refs. therein.
- [25] A. Voros, *Zeta-regularization for exact-WKB resolution of a general 1D Schrödinger equation*, J. Phys. **A45** (2012) 374007.
- [26] S. Wolfram, *Mathematica*, 3rd ed., Wolfram Media/Cambridge University Press, New York (1996).