

Rabi coupled fermions in the BCS-BEC crossover

L. Dell'Anna^{1,2}, F. De Bettin¹, and L. Salasnich^{1,2,3}

¹*Dipartimento di Fisica e Astronomia "Galileo Galilei" and QTech Center,
Università di Padova, Via Marzolo 8, 35131 Padova, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Padova, Via Marzolo 8, 35131 Padova, Italy*

³*Istituto Nazionale di Ottica del Consiglio Nazionale delle Ricerche,
via Nello Carrara 2, 50019 Sesto Fiorentino, Italy*

(Dated: July 1, 2022)

We investigate the three-dimensional BCS-BEC crossover in the presence of a Rabi coupling which strongly affects several properties of the system, such as the chemical potential, the pairing gap and the superfluid density. We determine the critical interaction strength, below which the system is normal also at zero temperature. Finally, we calculate the effect of the Rabi coupling on the critical temperature of the superfluid-to-normal phase transition by using different theoretical schemes.

PACS numbers: 67.85.Lm, 74.20.Fg, 47.37.+q

I. INTRODUCTION

An extremely important achievement in this field of ultracold atoms has been the realization of the crossover from the Bardeen-Cooper-Schrieffer (BCS) superfluid phase of loosely bound pairs of fermions to the Bose-Einstein condensate (BEC) of tightly bound composite bosons [1]. One of the most intriguing types of interactions is the so-called spin-orbit coupling (SOC), that is the coupling between the spin of a particle and its momentum. Although SOC is ubiquitous in Nature, it was not possible to control it in any area of physics until it was demonstrated in a breakthrough experiment [2] that the spin of an atom could be coupled to its center-of-mass motion by dressing two atomic spin states with a pair of laser beams. This technique has been then adopted in other experimental investigations of bosonic [3] and fermionic [4] atomic gases with artificial SOC, and also Rabi coupling. Triggered by the pioneering remarkable experiments of atoms with SOC, in the last few years a large number of theoretical papers have analyzed, within a mean field approach, the effect of SOC of Rashba [5] and Dresselhaus [6] type (often with the inclusion of a Rabi term) in Bose-Einstein condensates [7–12] and in the BCS-BEC crossover of superfluid fermions [13–29].

The aim of this paper is the study of an ultracold gas of Rabi coupled fermionic atoms interacting via a two body contact potential, but without the SOC term. The itinerant ferromagnetism of repulsive fermions with Rabi coupling was studied both in two [30] and three [31] spatial dimensions. Here, instead, we want to investigate the interplay of Rabi coupling and attractive interaction for fermions in the three-dimensional BCS-BEC crossover. In Section II we introduce Rabi coupling in the model of attractive fermions. In Section III we investigate the problem at the mean field level, also considering an improved determination [32] of the critical temperature of the superfluid-to-normal phase transition. In Section IV we consider beyond-mean-field corrections and use them to calculate the critical temperature with the inclusion of Gaussian fluctuations [28, 33].

II. THE MODEL

Our ultracold Fermi gas model is enriched with the addition of Rabi coupling, which enables the spin of the particles involved to flip. The Euclidean action, omitting the explicit dependence of the fermionic fields $\psi_\sigma(\mathbf{r}, \tau)$ ($\sigma = \uparrow, \downarrow$) on space \mathbf{r} and imaginary time τ , then, reads

$$S[\bar{\psi}, \psi] = \int_0^\beta d\tau \int_V d^3\mathbf{r} \left[\sum_\sigma \bar{\psi}_\sigma \left(\partial_\tau - \frac{\nabla^2}{2m} - \mu \right) \psi_\sigma - g \bar{\psi}_\uparrow \bar{\psi}_\downarrow \psi_\downarrow \psi_\uparrow + \omega_R (\bar{\psi}_\uparrow \psi_\downarrow + \bar{\psi}_\downarrow \psi_\uparrow) \right], \quad (1)$$

where $\beta = 1/(k_B T)$ with T the temperature and k_B the Boltzmann constant, μ is the chemical potential and V is the volume. Notice that in this paper we set $\hbar = 1$. Within the path integral formalism, the partition function of the system is given by

$$Z = \int D[\bar{\psi}, \psi] e^{-S[\bar{\psi}, \psi]}, \quad (2)$$

and from the partition function Z all the thermodynamical quantities can be derived.

We perform an Hubbard-Stratonovich transformation, so that the model can be rewritten in terms of a new action depending also on a new spinless complex field $\Delta(\mathbf{r}, \tau)$:

$$S[\bar{\Delta}, \Delta, \bar{\psi}, \psi] = \int_0^\beta d\tau \int_V d^3\mathbf{r} \left[\frac{|\Delta(\mathbf{r}, \tau)|^2}{g} - \frac{1}{2} \bar{\Psi}(\mathbf{r}, \tau) G^{-1} \Psi(\mathbf{r}, \tau) \right] + \beta \sum_{\mathbf{p}} \xi_{\mathbf{p}}, \quad (3)$$

where $\xi_{\mathbf{p}} = \mathbf{p}^2/(2m) - \mu$ is the shifted single-particle energy and the Nambu spinors take the form

$$\Psi(\mathbf{r}, \tau) = \begin{pmatrix} \psi_\uparrow(\mathbf{r}, \tau) \\ \psi_\downarrow(\mathbf{r}, \tau) \\ \bar{\psi}_\downarrow(\mathbf{r}, \tau) \\ \bar{\psi}_\uparrow(\mathbf{r}, \tau) \end{pmatrix} \quad (4)$$

while the inverse fermionic propagator is

$$G^{-1} = \begin{pmatrix} -\partial_\tau + \frac{\nabla^2}{2m} + \mu & \Delta(\mathbf{r}, \tau) & -\omega_R & 0 \\ \bar{\Delta}(\mathbf{r}, \tau) & -\partial_\tau - \frac{\nabla^2}{2m} - \mu & 0 & \omega_R \\ -\omega_R & 0 & -\partial_\tau + \frac{\nabla^2}{2m} + \mu & -\Delta(\mathbf{r}, \tau) \\ 0 & \omega_R & -\bar{\Delta}(\mathbf{r}, \tau) & -\partial_\tau - \frac{\nabla^2}{2m} - \mu \end{pmatrix}. \quad (5)$$

The action is now Gaussian in the fermionic degrees of freedom, therefore, we can integrate over them obtaining an effective theory for the complex field Δ , whose action reads

$$S_{\text{eff}}[\bar{\Delta}, \Delta] = \int_0^\beta d\tau \int_V d^3\mathbf{r} \frac{|\Delta(\mathbf{r}, \tau)|^2}{g} - \frac{1}{2} \text{Tr} \ln(G^{-1}) + \beta \sum_{\mathbf{p}} \xi_{\mathbf{p}}. \quad (6)$$

It is also convenient and useful for what follows to write G^{-1} in Fourier space, after denoting the four momenta by capital letters, like $P = (i\nu_n, \mathbf{p})$, introducing the fermionic Matsubara frequencies $\nu_n = \frac{(2n+1)\pi}{\beta}$, with $n \in \mathbb{Z}$. From Eq. (5) we find that the Fourier components of the inverse fermionic propagator G_{KP}^{-1} reads, therefore,

$$\begin{pmatrix} (i\nu_n - \xi_{\mathbf{p}})\delta_{K,P} & \Delta_{K+P} & -\omega_R \delta_{K,-P} & 0 \\ \bar{\Delta}_{K+P} & (i\nu_n + \xi_{\mathbf{p}})\delta_{K,P} & 0 & \omega_R \delta_{K,-P} \\ -\omega_R \delta_{(K,-P)} & 0 & (i\nu_n - \xi_{\mathbf{p}})\delta_{K,P} & -\Delta_{K+P} \\ 0 & \omega_R \delta_{K,-P} & -\bar{\Delta}_{K+P} & (i\nu_n + \xi_{\mathbf{p}})\delta_{K,P} \end{pmatrix}. \quad (7)$$

III. MEAN FIELD APPROACH

We can now performe, from Eq. (6), the saddle point approximation, choosing $\Delta = \Delta_0$, namely homogeneous in space and time and, without lack of generality, fixing it real. The mean field action, then, reads

$$S_{\text{mf}} = \beta V \frac{\Delta_0^2}{g} - \frac{1}{2} \text{Tr} \ln(G^{-1}) + \beta \sum_{\mathbf{p}} \xi_{\mathbf{p}}, \quad (8)$$

where G_0^{-1} is equal to G^{-1} where we take $\Delta = \Delta_0$. It is convenient working in Matsubara representation so that

$$\det(G_0^{-1}) = (\nu_n^2 + (\omega_{\mathbf{p}}^-)^2) (\nu_n^2 + (\omega_{\mathbf{p}}^+)^2), \quad (9)$$

with

$$\omega_{\mathbf{p}}^+ = \sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2} + \omega_R, \quad (10)$$

$$\omega_{\mathbf{p}}^- = \sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2} - \omega_R. \quad (11)$$

These two energies correspond to the poles of the fermionic propagator after a Wick rotation, meaning that they are the single particle excitation energies of the theory, and they differ from the case without Rabi coupling only by a constant shift ω_R . The presence of Rabi coupling splits the excitation energies into two different energy levels separated by a shift $2\omega_R$. It is immediately

clear that $\omega_{\mathbf{p}}^-$ may take negative values, which is somewhat unexpected. This may happen for $\Delta_0 < \omega_R$, a regime which is unphysical, as we will see, unless $\Delta_0 = 0$.

The mean-field grand potential is given by

$$\Omega_{\text{mf}} = k_B T S_{\text{mf}}. \quad (12)$$

Explicitly we have

$$\Omega_{\text{mf}} = V \frac{\Delta_0^2}{g} - \frac{1}{2} \sum_{\nu_n, \mathbf{p}} \left[\ln(\nu_n^2 + (\omega_{\mathbf{p}}^+)^2) + \ln(\nu_n^2 + (\omega_{\mathbf{p}}^-)^2) \right], \quad (13)$$

where we used $\text{Tr} \ln(G_0^{-1}) = \ln \det(G_0^{-1})$ and Eq. (9). After summing over the Matsubara frequencies, we get

$$\frac{\Omega_{\text{mf}}}{V} = \frac{\Delta_0^2}{g} - \frac{k_B T}{2V} \sum_{\mathbf{p}} \left\{ \ln \left[2(1 + \cosh(\beta\omega_{\mathbf{p}}^+)) \right] + \ln \left[2(1 + \cosh(\beta\omega_{\mathbf{p}}^-)) \right] \right\}. \quad (14)$$

A. Gap and number equations

Minimizing the mean-field grand potential Ω_{mf} with respect to Δ_0 we obtain the so-called gap equation

$$\frac{1}{g} = \frac{1}{4V} \sum_{\mathbf{p}} \left[\frac{\tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^+)}{\sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2}} + \frac{\tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^-)}{\sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2}} \right]. \quad (15)$$

This equation is divergent in the ultraviolet and requires a regularization of the interaction strength g , namely

$$\frac{1}{g} = -\frac{m}{4\pi a_F} + \frac{1}{V} \sum_{\mathbf{p}} \frac{m}{\mathbf{p}^2}, \quad (16)$$

with a_F being the physical s-wave scattering length.

The total number density n is instead obtained by

$$n = -\frac{1}{V} \frac{\partial \Omega_{\text{mf}}}{\partial \mu}, \quad (17)$$

which is the so-called number equation. The sum over Matsubara frequencies has the same form as the one of the gap equation. After some manipulations, the renormalized gap equation and the number equation read

$$-\frac{mV}{4\pi a_F} = \sum_{\mathbf{p}} \left[\frac{\tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^+) + \tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^-)}{4\sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2}} - \frac{m}{\mathbf{p}^2} \right] \quad (18)$$

$$nV = \sum_{\mathbf{p}} \left[1 - \frac{\xi_{\mathbf{p}}}{2} \frac{\tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^+) + \tanh(\frac{\beta}{2}\omega_{\mathbf{p}}^-)}{\sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2}} \right]. \quad (19)$$

The difference with respect to the case without Rabi coupling is a shift of $\pm\omega_R$ in the arguments of the hyperbolic tangents, which makes the derivation of analytic results more demanding.




FIG. 1: Gap and chemical potential obtained solving Eqs. (18) and (19) in the zero-temperature limit. Upper panel: adimensional energy gap Δ_0/ω_F vs inverse adimensional scattering length $y = 1/(k_F a_F)$. Lower panel: adimensional energy gap Δ_0/ω_F vs inverse adimensional scattering length $1/(k_F a_F)$. Three values of the adimensional Rabi frequency: $\omega_R/\omega_F = 0$ (dashed curve); $\omega_R/\omega_F = 0.5$ (solid curve); $\omega_R/\omega_F = 1$ (dotted curve). Here $k_F = (3\pi^2 n)^{1/3}$ is the Fermi wavenumber and $\omega_F = k_F^2/(2m)$ is the Fermi frequency.

B. Zero temperature

At zero temperature Eqs. (18) and (19) simplify because the hyperbolic tangent goes to one for $T \rightarrow 0^+$. However, careful attention should be paid studying the sign of $\omega_{\mathbf{p}}^-$, which affects the form of the equations. In fact, if $\omega_{\mathbf{p}}^- > 0$ for any value of the momentum \mathbf{p} , the number and gap equations will take the same form as the ones with no Rabi interaction, while if for some values of \mathbf{p} the energy $\omega_{\mathbf{p}}^- < 0$, the equations will take a different form, as we will show below. The main result of this zero-temperature analysis is the following: i) if $\Delta_0 < \omega_R$ the Rabi frequency ω_R change the momenta domain of integration in the equations in such a way that the gap equation has no finite solutions; ii) if $\Delta_0 > \omega_R$, instead, the Rabi frequency ω_R does not affect the gap and the number equations.

In Fig. 1 we report the plots of the energy gap Δ_0 (upper panel) and chemical potential μ (lower panel) as functions of the s-wave scattering length a_F . There re-




FIG. 2: Critical strength y_c vs adimensional Rabi frequency ω_R/ω_F at zero temperature. For $y \leq y_c$ the fermionic system is normal, i.e. the energy gap $\Delta_0 = 0$. Here $y = 1/(k_F a_F)$, where a_F is the s-wave scattering length and $k_F = (3\pi^2 n)^{1/3}$ is the Fermi wavenumber, with $\omega_F = k_F^2/(2m)$ the Fermi frequency.

sults are then analogous to the ones in the case without Rabi coupling for $\Delta_0 > \omega_R$, but exhibit a different behaviour below such a threshold. In particular, both these quantities, the energy gap Δ_0 and the chemical potential μ , for a small range of $1/(k_F a_F)$, have two branches. However, the stable branch (associated to the minima of the grand potential Ω) is the upper branch shown in the figures, superimposed to the curves of $\omega_R = 0$. The unstable is not reported.

At zero temperature ($T = 0$) the main effect of the Rabi coupling ω_R is, therefore, to make the system normal, i.e. with $\Delta_0 = 0$, for $1/(k_F a_F) \leq y_c$, where $y_c = -\infty$ for $\omega_R = 0$. The critical strength y_c grows by increasing ω_R . Specifically, given $\Delta_0(y)$ with $1/(k_F a_F)$ for $\omega_R = 0$, y_c is obtained from the condition $\Delta_0(y_c) = \omega_R$. This means that inverting the plot of $\Delta_0(y)$ (obtained for $\omega_R = 0$) one gets immediately y_c vs ω_R , as shown, for the sake of completeness, in Fig. 2.

C. Critical temperature

We now investigate the behaviour of the system at the critical temperature T_c , at which the energy gap $\Delta_0(T_c) = 0$. Let us define, for simplicity, the following adimensional quantities: $\tilde{\mu} = \mu/\omega_F$, $\tilde{T}_c = k_B T_c/\omega_F$, $\tilde{\omega}_R = \omega_R/\omega_F$ and $y = 1/(k_F a_F)$. In this case the gap and the number equations can be written as follows

$$y = \frac{2\tilde{T}_c^2}{\pi} J_3(\tilde{\mu}, \tilde{T}_c, \tilde{\omega}_R), \quad (20)$$

$$\tilde{T}_c = \left[\frac{4}{J_4(\tilde{\mu}, \tilde{T}_c, \tilde{\omega}_R)} \right]^{2/3}, \quad (21)$$




FIG. 3: Adimensional critical temperature $k_B T_c / \omega_F$ vs inverse adimensional scattering length $y = 1/(k_F a_F)$, with a_F the s-wave scattering length, $k_F = (3\pi^2 n)^{1/3}$ the Fermi wavenumber and $\omega_F = k_F^2 / (2m)$ the Fermi frequency. Thin curves are the mean-field ones, obtained solving Eqs. (20) and (21), while thick curves are obtained from Eqs. (24) and (25). Three values of the adimensional Rabi frequency: $\omega_R / \omega_F = 0$ (dashed curve); $\omega_R / \omega_F = 0.3$ (solid curve); $\omega_R / \omega_F = 0.5$ (dot-dashed curve).

where

$$J_3 = \int_0^{+\infty} dx x^2 \left[\frac{\tanh\left(\frac{1}{2}\left(x^2 - \frac{\tilde{\mu}}{T_c} + \frac{\tilde{\omega}_R}{T_c}\right)\right)}{2\left(x^2 - \frac{\tilde{\mu}}{T_c}\right)} + \frac{\tanh\left(\frac{1}{2}\left(x^2 - \frac{\tilde{\mu}}{T_c} - \frac{\tilde{\omega}_R}{T_c}\right)\right)}{2\left(x^2 - \frac{\tilde{\mu}}{T_c}\right)} - \frac{1}{x^2} \right] \quad (22)$$

and

$$J_4 = \int_0^{+\infty} dx x^4 \left[\frac{1}{\cosh^2\left[\frac{1}{2}\left(x^2 - \frac{\tilde{\mu}}{T_c} + \frac{\tilde{\omega}_R}{T_c}\right)\right]} + \frac{1}{\cosh^2\left[\frac{1}{2}\left(x^2 - \frac{\tilde{\mu}}{T_c} - \frac{\tilde{\omega}_R}{T_c}\right)\right]} \right]. \quad (23)$$

Solving the coupled Eqs. (20) and (21) we obtain the critical temperature T_c as a function of the inverse scattering length $1/a_F$ for different values of the Rabi coupling ω_R . The results are shown as thin curves in Fig. 3. As expected, the Rabi coupling inhibits the formation of Cooper pairs: the stronger the Rabi coupling, the higher is the threshold of the scattering rate above which superfluidity can occur at the mean field level.

IV. BEYOND MEAN FIELD

We are now presenting a couple of techniques which allow us to go beyond the mean field analysis. The first approach is a method based on the determination of the superfluid density, while the second one is based on the inclusion of the Gaussian fluctuations. In the latter case we will show some explicit results in the so-called bosonic approximation.

A. By superfluid density

An improved determination of the critical temperature T_c can be obtained with the method proposed by Babaev and Kleinert [32], which is the three-dimensional analog of the Nelson-Kosterlitz criterion. In particular,

$$k_B T_c = \alpha \frac{n_s(T_c)}{4m} \left(\frac{2}{n}\right)^{1/3}, \quad (24)$$

where $n_s(T)$ is the mean-field superfluid density and n is the total fermionic number density. Actually $J = n_s / (4m)$ is the stiffness in an effective XY model, $H = \frac{J}{2} \int d\mathbf{r} \nabla \theta(\mathbf{r})$, where θ is the local phase of the pairing Δ .

The constant α is fixed to the value $\alpha = 2\pi / \zeta(3/2)^{2/3}$ such that T_c turns out to be the exact value $k_B T_c = \pi / m(n / (2\zeta(3/2)))^{2/3}$ for $n/2$ non-interacting bosons with mass $2m$ in the deep BEC regime. The superfluid density $n_s(T)$ can be calculated, following the Landau's approach [34], getting

$$n_s(T) = n + \frac{1}{6} \int \frac{d^3 p}{(2\pi)^3} \frac{p^2}{m} \left[\frac{df_B(\omega_{\mathbf{p}}^+)}{d\omega_{\mathbf{p}}^+} + \frac{df_B(\omega_{\mathbf{p}}^-)}{d\omega_{\mathbf{p}}^-} \right] \quad (25)$$

where

$$f_B(\omega) = \frac{1}{e^{\beta\omega} - 1} \quad (26)$$

is the Bose-Einstein distribution and n is the total number density. The thick curves of Fig. 3 are obtained using Eq. (24) with Eq. (25), where Δ_0 and μ are numerically determined from Eqs. (18) and (19).

As done previously, it is convenient to introduce adimensional quantities, $\tilde{\mu} = \mu / \omega_F$, $\tilde{T}_c = k_B T_c / \omega_F$, $\tilde{\omega}_R = \omega_R / \omega_F$ and $y = 1 / (k_F a_F)$ together with $\tilde{\Delta}_0 = \Delta_0 / \omega_F$, where we recall that $\omega_F = k_F^2 / (2m) = (3\pi^2 n)^{2/3} / (2m)$. We can, therefore, rewrite Eq. (24) as

$$\tilde{T}_c = \frac{2}{(6\sqrt{\pi} \zeta(3/2))^{2/3}} \frac{n_s(\tilde{T}_c)}{n} \quad (27)$$

which, in the deep BEC where all the fermions contribute to the superfluid density, gives $\tilde{T}_c \approx 0.218$, and where

$$\frac{n_s(\tilde{T}_c)}{n} = 1 - \frac{1}{2\tilde{T}_c} \int_0^{+\infty} dx x^4 \left[\frac{\sum_{s=\pm 1} \frac{e^{\frac{1}{\tilde{T}_c} \left(\sqrt{(x^2 - \tilde{\mu})^2 + \tilde{\Delta}_0^2} + s \tilde{\omega}_R \right)}}{\left(e^{\frac{1}{\tilde{T}_c} \left(\sqrt{(x^2 - \tilde{\mu})^2 + \tilde{\Delta}_0^2} + s \tilde{\omega}_R \right)} - 1 \right)^2} \right].$$

B. Gaussian fluctuations

We now introduce Gaussian fluctuations in the partition function of the system adopting the Nozières-Schmitt-Rink approach [33]. The aim is to derive a more

precise form for the number equation in order to understand the role of quantum fluctuations on the relation between the chemical potential μ and the density n of fermions.

In order to introduce fluctuations we separate the field $\Delta(\mathbf{r}, \tau)$ in its homogeneous part Δ_0 , obtained minimizing the grand potential at the mean field level, and its small fluctuations $\eta(\mathbf{r}, \tau)$ around the saddle point solution

$$\Delta(\mathbf{r}, \tau) = \Delta_0 + \eta(\mathbf{r}, \tau). \quad (28)$$

After some calculations, at the leading order in η , the effective action Eq. (6) at the Gaussian level reads

$$S_{\text{eff}} \simeq S_{\text{mf}} + \frac{1}{2} \sum_Q (\bar{\eta}_Q \ \eta_{-Q}) M_Q \begin{pmatrix} \eta_Q \\ \bar{\eta}_{-Q} \end{pmatrix} \quad (29)$$

with

$$M_Q = \frac{1}{g} \mathbb{I} + \chi_Q, \quad (30)$$

where χ_Q is the contribution coming from the expansion of $\text{Tr} \ln(G^{-1})$ and \mathbb{I} denotes the 2×2 identity matrix in Nambu space. The components of χ_Q , shown in Appendix, depend on the the quadrivector $Q = (iv_n, \mathbf{q})$ where v_n are bosonic Matsubara frequencies, $v_n = \frac{2\pi n}{\beta}$, with $n \in \mathbb{Z}$.

Our objective is to find an expression for the grand canonical potential from which we can recover a treatable expression for the contribution of the Gaussian fluctuations to the number equation. The effective theory we obtained is Gaussian, meaning that it can be integrated explicitly, getting the following partition function

$$Z = e^{-S_{\text{mf}}} \prod_Q [\det(M_Q)]^{-1}. \quad (31)$$

The grand potential, then, reads

$$\Omega = \Omega_{\text{mf}} + \Omega_{\text{g}}, \quad (32)$$

where Ω_{mf} is given by Eq. (12) and

$$\Omega_{\text{g}} = k_B T \sum_Q \ln[\det(M_Q)]. \quad (33)$$

To compute the sum over Matsubara frequencies in (33) one may analytically continue the argument of the sum by setting $iv_n \rightarrow \omega$ and transforming the sum into an integral. In this way we eventually find

$$\Omega_{\text{g}} = -\frac{1}{\pi} \sum_{\mathbf{q}} \int_{-\infty}^{+\infty} d\omega f_B(\omega) \tilde{\delta}(\omega, \mathbf{q}) \quad (34)$$

with $f_B(\omega)$ the Bose-Einstein distribution and

$$\tilde{\delta}(\omega, \mathbf{q}) = \arctan \left[\frac{\text{Im}[\det[M_{\omega, \mathbf{q}}]]}{\text{Re}[\det[M_{\omega, \mathbf{q}}]]} \right] \quad (35)$$




FIG. 4: Adimensional critical temperature $k_B T_c / \omega_F$ vs inverse adimensional scattering length $y = 1/(k_F a_F)$ by including Gaussian fluctuations within the bosonic approximation, i.e. solving Eqs. (18) and (38). Also here a_F is the s-wave scattering length, $k_F = (3\pi^2 n)^{1/3}$ is the Fermi wavenumber and $\omega_F = k_F^2 / (2m)$ is the Fermi frequency. Four values of the adimensional Rabi frequency: $\omega_R / \omega_F = 0$ (dashed curve); $\omega_R / \omega_F = 0.3$ (solid curve); $\omega_R / \omega_F = 0.5$ (dot-dashed curve); $\omega_R / \omega_F = 1$ (dot-dot-dashed curve).

the phase of the complex matrix element $M_{\omega, \mathbf{q}}$ derived from (30). Thus, the Gaussian correction to the number density reads

$$n_{\text{g}} = -\frac{1}{V} \frac{\partial \Omega_{\text{g}}}{\partial \mu} = \frac{1}{\pi V} \sum_{\mathbf{q}} \int_{-\infty}^{+\infty} d\omega f_B(\omega) \frac{\partial \tilde{\delta}(\omega, \mathbf{q})}{\partial \mu}. \quad (36)$$

In the strong coupling limit $a_F \rightarrow 0^+$ the system becomes a gas of free bosonic dimers, made of two fermions with opposite spin and binding energy $E_B = -1/(ma_F^2)$. These bosons have mass $m_B = 2$ and chemical potential $\mu_B = 2\mu - E_B$. Indeed, as shown in Ref. [28], in this regime $\mu_B \rightarrow 0$ and

$$\frac{\partial \tilde{\delta}(\omega, \mathbf{q})}{\partial \mu} \rightarrow \pi \delta\left(\omega - \frac{q^2}{2m_B} + \mu_B\right) \quad (37)$$

where $\delta(x)$ is the Dirac delta function. It follows immediately that, in the strong coupling approximation, also called bosonic approximation [28], the number equation with Gaussian fluctuations becomes

$$nV = \sum_{\mathbf{p}} \left[1 - \frac{\xi_{\mathbf{p}}}{2} \frac{\tanh\left(\frac{\beta}{2}\omega_{\mathbf{p}}^+\right) + \tanh\left(\frac{\beta}{2}\omega_{\mathbf{p}}^-\right)}{\sqrt{\xi_{\mathbf{p}}^2 + \Delta_0^2}} \right] + \sum_{\mathbf{q}} \frac{1}{e^{\beta\left(\frac{q^2}{4m} - \mu_B\right)} - 1}. \quad (38)$$

In Fig. 4 we report the critical temperature T_c as a function of the inverse scattering length $1/a_F$ obtained by solving Eqs. (18) and (38), setting $\mu_B = 0$ and $\Delta_0 = 0$.

V. CONCLUSIONS

The BCS-BEC crossover has been studied in the presence of Rabi coupling by using the finite-temperature path integral formalism. The behavior of many physical quantities has been studied along the whole crossover, including the mean-field chemical potential and energy gap at zero temperature, and the critical temperature at and beyond mean-field level. We have found that only in the deep BEC regime the physical properties of the system are not affected by the Rabi coupling. In general, also at zero temperature it exists a critical interaction strength, below which the system is normal. We have determined this critical strength as a function of the Rabi coupling. In the last part of the paper we have calculated the critical temperature of the superfluid-to-normal phase transition for different values of the Rabi coupling. The treatment beyond mean-field level has been carried out following two different procedures: the Babaev-Kleinert [32] and the Nozieres-Schmitt-Rink [33] approaches: the first being more heuristic and the second being more rigorous but computationally demanding. Indeed, the Nozieres-Schmitt-Rink scheme has been used but adopting the bosonic pair approximation [28], which makes the scheme more feasible numerically.

Acknowledgements. LS acknowledges the BIRD project "Ultracold atoms in curved geometries" of the University of Padova for partial support.

Appendix

Let us write the fermionic propagator, including quantum fluctuations, as follows

$$G^{-1} = G_0^{-1} + \hat{\eta} = G_0^{-1}(1 + G_0\hat{\eta}) \quad (39)$$

where we introduce the matrix

$$\hat{\eta}_Q = \begin{pmatrix} 0 & \eta_Q & 0 & 0 \\ \bar{\eta}_Q & 0 & 0 & 0 \\ 0 & 0 & 0 & -\eta_Q \\ 0 & 0 & -\bar{\eta}_Q & 0 \end{pmatrix}. \quad (40)$$

so that, in the action, the $\text{Tr} \ln(G^{-1})$ can be written in the following way

$$\frac{1}{2} \text{Tr} \ln(G^{-1}) = \frac{1}{2} \text{Tr} \ln(G_0^{-1}) + \frac{1}{2} \text{Tr} \ln(1 + G_0\hat{\eta}). \quad (41)$$

We can now expand the last term, getting the following leading term

$$\frac{1}{2} \text{Tr} (G_0\hat{\eta}G_0\hat{\eta}) = \frac{k_B T}{V} \sum_{Q,P} \text{tr}(G_{0P}\hat{\eta}_Q G_{0P+Q}\hat{\eta}_{-Q}) \quad (42)$$

which can be recasted, performing the matrix products, as follows

$$\frac{1}{2} \text{Tr} (G_0\hat{\eta}G_0\hat{\eta}) = \frac{1}{2} \sum_Q (\bar{\eta}_Q \ \eta_{-Q}) \chi_Q \begin{pmatrix} \eta_Q \\ \bar{\eta}_{-Q} \end{pmatrix} \quad (43)$$

where χ_Q is a 2×2 matrix, introduced in the main text in Eq. (30), which, after performing the inverse of G_0^{-1} and making the matrix products, can be written explicitly. This matrix is composed by the following diagonal terms

$$\begin{aligned} (\chi_Q)_{11} = (\chi_{-Q})_{22} &= \frac{k_B T}{V} \sum_{\mathbf{p}, \nu_n} \frac{[(i\nu_n - \xi_{\mathbf{p}})((i\nu_n)^2 - \xi_{\mathbf{p}}^2 - \Delta_0^2) - \omega_R^2(i\nu_n + \xi_{\mathbf{p}})]}{(\nu_n^2 + (\omega_{\mathbf{p}}^+)^2)(\nu_n^2 + (\omega_{\mathbf{p}}^-)^2)} \\ &\times \frac{[(i\nu_m + i\nu_n + \xi_{\mathbf{q}+\mathbf{p}})((i\nu_m + i\nu_n)^2 - \xi_{\mathbf{q}+\mathbf{p}}^2 - \Delta_0^2) - \omega_R^2(i\nu_m + i\nu_n - \xi_{\mathbf{q}+\mathbf{p}})]}{((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^+)^2)((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^-)^2)} \\ &+ \omega_R^2 \frac{k_B T}{V} \sum_{\mathbf{p}, \nu_n} \frac{[\Delta_0^2 - \omega_R^2 + (i\nu_n - \xi_{\mathbf{p}})^2]}{(\nu_n^2 + (\omega_{\mathbf{p}}^+)^2)(\nu_n^2 + (\omega_{\mathbf{p}}^-)^2)} \frac{[\Delta_0^2 - \omega_R^2 + (i\nu_m + i\nu_n + \xi_{\mathbf{p}+\mathbf{q}})^2]}{((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^+)^2)((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^-)^2)} \end{aligned} \quad (44)$$

and off-diagonal terms

$$\begin{aligned} (\chi_Q)_{12} = (\chi_Q)_{21} &= \Delta_0^2 \frac{k_B T}{V} \sum_{\mathbf{p}, \nu_n} \frac{[(\nu_n^2 + \xi_{\mathbf{p}}^2) - \omega_R^2 + \Delta_0^2]}{(\nu_n^2 + (\omega_{\mathbf{p}}^+)^2)(\nu_n^2 + (\omega_{\mathbf{p}}^-)^2)} \frac{[((\nu_m + \Omega_{n_F})^2 + \xi_{\mathbf{p}+\mathbf{q}}^2) - \omega_R^2 + \Delta_0^2]}{((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^+)^2)((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^-)^2)} \\ &- 4\omega_R^2 \Delta_0^2 \frac{k_B T}{V} \sum_{\mathbf{p}, \nu_n} \frac{\nu_n}{(\nu_n^2 + (\omega_{\mathbf{p}}^+)^2)(\nu_n^2 + (\omega_{\mathbf{p}}^-)^2)} \frac{(\nu_m + \nu_n)}{((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^+)^2)((\nu_m + \nu_n)^2 + (\omega_{\mathbf{q}+\mathbf{p}}^-)^2)}. \end{aligned} \quad (45)$$

-
- [1] W. Zwerger, *The BCS-BEC Crossover and the Unitary Fermi Gas* (Springer, 2012).
- [2] Y. J. Lin, K. Jimenez-Garcia, and I. B. Spielman, *Nature* **471**, 83 (2011).
- [3] J.-Y. Zhang *et al.*, *Phys. Rev. Lett.* **109**, 115301 (2012).
- [4] P. Wang *et al.*, *Phys. Rev. Lett.* **109**, 095301 (2012).
- [5] Yu.A. Bychkov, and E.I. Rashba, *J. Phys. C* **17**, 6039 (1984).
- [6] G. Dresselhaus, *Phys. Rev.* **100**, 580 (1955).
- [7] Y. Li, L.P. Pitaevskii, and S. Stringari, *Phys. Rev. Lett.* **108**, 225301 (2012).
- [8] G.I. Martone, Yun Li, L.P. Pitaevskii, and S. Stringari, *Phys. Rev. A* **86**, 063621 (2012).
- [9] M. Merkl, A. Jacob, F. E. Zimmer, P. Ohberg, and L. Santos, *Phys. Rev. Lett.* **104**, 073603 (2010).
- [10] V. Achilleos *et al.*, *Phys. Rev. Lett.* **110**, 264101 (2013).
- [11] S. Sinha, R. Nath, and L. Santos, *Phys. Rev. Lett.* **107**, 270401 (2011).
- [12] Y. Deng, J. Cheng, H. Jing, C. P. Sun, and S. Yi, *Phys. Rev. Lett.* **108**, 125301 (2012).
- [13] J. P. Vyasankere and V. B. Shenoy, *Phys. Rev. B* **83**, 094515 (2011).
- [14] J. P. Vyasankere, S. Zhang, and V. B. Shenoy, *Phys. Rev. B* **84**, 014512 (2011).
- [15] M. Gong, S. Tewari, and C. Zhang, *Phys. Rev. Lett.* **107**, 195303 (2011).
- [16] H. Hu, L. Jiang, X.-J. Liu, and H. Pu, *Phys. Rev. Lett.* **107**, 195304 (2011).
- [17] Z.-Q. Yu and H. Zhai, *Phys. Rev. Lett.* **107**, 195305 (2011).
- [18] M. Iskin and A. L. Subasi, *Phys. Rev. Lett.* **107**, 050402 (2011).
- [19] M. Iskin and A. L. Subasi, *Phys. Rev. A* **84**, 043621 (2011).
- [20] L. Jiang, X.-J. Liu, H. Hu, and H. Pu, *Phys. Rev. A* **84**, 063618 (2011).
- [21] Li Han and C.A.R. Sa de Melo, *Phys. Rev. A* **85**, 011606(R) (2012).
- [22] K. Zhou and Z. Zhang, *Phys. Rev. Lett.* **108**, 025301 (2012).
- [23] M. Iskin, *Phys. Rev. A* **85**, 013622 (2012).
- [24] L. He and X.-G. Huang, *Phys. Rev. Lett.* **108**, 145302 (2012).
- [25] X.J. Liu, M.F. Borunda, X. Liu, and J. Sinova, *Phys. Rev. Lett.* **102**, 046402 (2009).
- [26] P. Wang *et al.*, *Phys. Rev. Lett.* **109**, 095301 (2012).
- [27] L. Dell'Anna, G. Mazzaella, and L. Salasnich, *Phys. Rev. A* **84**, 033633 (2011); *Phys. Rev. A* **86**, 053632 (2012).
- [28] L. Dell'Anna and S. Grava, *Condens. Matter* **6**, 16 (2021).
- [29] P.D. Powell, G. Baym, and C. A. R. Sa de Melo, *Phys. Rev. A* **105**, 063304 (2022).
- [30] L. Salasnich and V. Penna, *New J. Phys.* **19**, 043018 (2017).
- [31] V. Penna and L. Salasnich, *J. Phys. B: At. Mol. Opt. Phys.* **52**, 035301 (2019).
- [32] E. Babaev and H. Kleinert, *Phys. Rev. B* **59**, 12083 (1999).
- [33] P. Nozieres and S. Schmitt-Rink, *J. Low Temp. Phys.* **59**, 195 (1985).
- [34] L.D. Landau, E.M. Lifshitz, *Statistical Physics - Part 2: Theory of the Condensed State*, *Course of Theoretical Physics*, vol. **9** (Pergamon Press, 1980).