

# Approximation Algorithms for Continuous Clustering and Facility Location Problems

Deeparnab Chakrabarty  
deeparnab@dartmouth.edu  
Department of Computer Science  
Dartmouth College, NH, USA

Maryam Negahbani  
maryam.negahbani.gr@dartmouth.edu  
Department of Computer Science  
Dartmouth College, NH, USA

Ankita Sarkar  
ankita.sarkar.gr@dartmouth.edu  
Department of Computer Science  
Dartmouth College, NH, USA

## Abstract

In this paper, we consider center-based clustering problems where  $C$ , the set of points to be clustered, lies in a metric space  $(X, d)$ , and the set  $X$  of candidate centers is potentially infinite-sized. We call such problems *continuous* clustering problems to differentiate them from the *discrete* clustering problems where the set of candidate centers is explicitly given. It is known that for many objectives, when one restricts the set of centers to  $C$  itself and applies an  $\alpha_{\text{dis}}$ -approximation algorithm for the discrete version, one obtains a  $\beta \cdot \alpha_{\text{dis}}$ -approximation algorithm for the continuous version via the triangle inequality property of the distance function. Here  $\beta$  depends on the objective, and for many objectives such as  $k$ -median,  $\beta = 2$ , while for some others such as  $k$ -means,  $\beta = 4$ . The motivating question in this paper is *whether this gap of factor  $\beta$  between continuous and discrete problems is inherent, or can one design better algorithms for continuous clustering than simply reducing to the discrete case as mentioned above?* In a recent SODA 2021 paper, Cohen-Addad, Karthik, and Lee prove a factor-2 and a factor-4 hardness, respectively, for the continuous versions of the  $k$ -median and  $k$ -means problems, even when the number of cluster centers is a constant. The discrete problem for a constant number of centers is easily solvable exactly using enumeration, and therefore, in certain regimes, the “ $\beta$ -factor loss” seems unavoidable.

In this paper, we describe a technique based on the *round-or-cut* framework to approach continuous clustering problems. We show that, for the continuous versions of some clustering problems, we can design approximation algorithms attaining a better factor than the  $\beta$ -factor blow-up mentioned above. In particular, we do so for: the uncapacitated facility location problem with uniform facility opening costs ( $\lambda$ -UFL); the  $k$ -means problem; the individually fair  $k$ -median problem; and the  $k$ -center with outliers problem. Notably, for  $\lambda$ -UFL, where  $\beta = 2$  and the discrete version is NP-hard to approximate within a factor of 1.27, we describe a 2.32-approximation for the continuous version, and indeed  $2.32 < 2 \times 1.27$ . Also, for  $k$ -means, where  $\beta = 4$  and the best known approximation factor for the discrete version is 9, we obtain a 32-approximation for the continuous version, which is better than  $4 \times 9 = 36$ .

The main challenge one faces is that most algorithms for the discrete clustering problems, including the state of the art solutions, depend on Linear Program (LP) relaxations that become infinite-sized in the continuous version. To overcome this, we design new linear program relaxations for the continuous clustering problems which, although having exponentially many constraints, are amenable to the round-or-cut framework.

**Related version.** *ESA 2022 version:* <https://drops.dagstuhl.de/opus/volltexte/2022/16971> [15]

**Funding.** The authors were supported by NSF award #2041920.

## 1 Introduction

Clustering is a ubiquitous problem arising in various areas ranging from data analysis to operations research. One popular class of clustering problems are the so-called *center-based clustering* problems where the quality of the clustering is determined by a function of the distances of every point in  $C$  to the “centers” of the clusters they reside in. Two extensively studied measures are the sum of these distances, with the resulting problem called the  $k$ -median problem, and the sum of squares of these distances, with the resulting problem called the  $k$ -means problem.

In most settings, these center-based clustering problems are NP-hard and one considers approximation algorithms for the same. Traditionally, however, approximation algorithms for these problems have been studied in *finite/discrete* metric spaces and, in fact, usually under the constraint that the set of centers, aka facilities, can be selected only from a prescribed subset  $F \subseteq X$  of the metric space. Indeed, this model makes perfect sense when considering applications in operations research where the possible depot-locations may

be constrained. These discrete problems have been extensively studied [37, 50, 17, 33, 18, 16, 39, 38, 5, 41, 44, 45, 9] over the last three decades. For instance, for the  $k$ -median problem, the best known approximation algorithm is a 2.675-approximation [10], while the best known hardness is  $(1 + 2/e) \approx 1.74$  [38]. For the  $k$ -means problem, the best known approximation algorithm is a  $(9 + \varepsilon)$ -approximation algorithm [1, 34, 41], while the best hardness is  $1 + 8/e \approx 3.94$  [38].

Restricting to a finite metric space, however, makes the problem easier, and indeed many of the above algorithms in the papers mentioned above would be infeasible to implement if  $X$  were extremely large – for instance, if  $X$  were  $\mathbb{R}^m$  for some large dimension  $m$ , and the distance function were the  $\ell_p$ -metric, for some  $p$ . On the other hand, it is reasonably easy to show using triangle inequality that if one considers opening centers from  $C$  itself, and thus reduces the problem to its discrete version, then one incurs a hit of a factor  $\beta$  in the approximation factor, where  $\beta$  is a constant depending on the objective function. In particular, if we look at the sum-of-distances objectives such as in  $k$ -median, then  $\beta = 2$ , while if one looks at the sum-of-squared-distances such as in  $k$ -means, then  $\beta = 4$ . Therefore, one immediately gets a 5.35-approximation for the *continuous  $k$ -median* problem and a  $36 + \varepsilon$ -approximation for the *continuous  $k$ -means* problem. The question we investigate is

*Is this factor  $\beta$  hit necessary between the continuous and discrete versions of center based clustering problems, or can one design better approximation algorithms for the continuous case?*

It is crucial to note that when considering designing algorithms, we do not wish to make *any* assumptions on the underlying metric space  $(X, d)$ . For instance, we do not wish to assume  $X = \mathbb{R}^m$  for some  $m$ . This is important, for we really want to compare ourselves with the  $\beta$  which is obtained using only the triangle-inequality and symmetry property of  $d$ . On the other hand, to exhibit that a certain algorithm does not work, any candidate metric space suffices.

Recently, in a thought-provoking paper [22], Cohen-Addad, Karthik, and Lee show that, unless  $P = NP$ , the  $k$ -median and  $k$ -means problem defined on  $(\mathbb{R}^m, \ell_\infty)$  cannot have an approximation ratio better than 2 and 4, respectively, *even when  $k$  is a constant!* Since the discrete problems have trivial exact algorithms via enumeration when  $k$  is a constant, this seems to indicate that in certain cases the above factor  $\beta$  hit is unavoidable. Is it possible that the *inapproximability* of the continuous problem is indeed  $\beta$  times the inapproximability of the discrete version?

## 1.1 Our Results

Our main contribution is a direct approach towards the continuous versions of clustering problems. We apply this to the following clustering problems where we obtain a factor better than  $\beta \cdot \alpha_{\text{dis}}$ , where  $\alpha_{\text{dis}}$  is the best known factor for the discrete version of the problem.

- In the continuous  $\lambda$ -UFL problem, a “soft” version of the continuous  $k$ -median problem, one is allowed to pick any number of centers but has to pay a parameter  $\lambda$  for each picked center. The objective is to minimize the sum of distances of points in  $C$  to picked centers plus the cost for opening these centers. Again, note that the centers can be opened anywhere in  $X$ . For the discrete version, where the only possible center locations are in  $C$ , there is a 1.488-approximation due to Li [44], and a hardness of approximation within a factor of 1.278 is known due to Guha and Khuller [33]. We describe a 2.32-approximation algorithm. Note that  $2.32 < 2 \cdot 1.278$  and thus, for this problem, the inapproximability is *not*  $\beta$  times that of the discrete case. We also show how the reduction of [22] carries over, to prove a hardness of 2 for this problem.
- In the continuous  $k$ -means problem, we wish to minimize the sum of squares of distances of clients to the closest open center. Recall that for this problem we have  $\beta = 4$ , and thus one gets a 36-factor algorithm for the continuous  $k$ -means using the best known 9-factor [1, 34, 41] algorithm for the discrete problem. We describe an improved 32-approximation for the continuous  $k$ -means problem.
- For the continuous  $k$ -median problem, our techniques fall short of improving  $2 \times$  the best known approximation factor for the discrete  $k$ -median problem. On the other hand, we obtain better algorithms for the *individually fair* or *priority* version of the continuous  $k$ -median problem. In this problem, every point  $v \in C$  has a specified radius  $r(v)$  and desires a center opened within this distance. The

objective is the same as the  $k$ -median problem: minimize the sum of all the distances. This problem arises as a possible model [40, 14, 46, 53, 48, 7] in the study of fair clustering solutions, since the usual  $k$ -median algorithms may place certain clients inordinately far away. At a technical level, this problem is a meld of the  $k$ -median and the  $k$ -center problems; the latter is NP-hard, which forces one to look at *bicriterion* approximations. An  $(\alpha, \gamma)$ -approximation would return a solution within  $\alpha$  times the optimum but may connect  $v$  to a point as far as  $\gamma r(v)$  away. Again, any  $(\alpha, \gamma)$ -approximation for the *discrete* version where  $X = C$  would imply a  $(2\alpha, 2\gamma)$ -approximation for the continuous version.

The best discrete approximation is (8, 3) due to Vakilian and Yalçiner [53] which would imply an (16, 3)-approximation for the continuous version. We describe an (8, 8)-approximation for the continuous version of the problem.

- In the  $k$ -center with outliers (kCwO) problem, we are given a parameter  $m \leq |C|$ , and we need to serve only  $m$  of the clients. The objective is the maximum distance of a served client to its center. The  $k$ -center objective is one of the objectives for which most existing discrete algorithms can compare themselves directly with the continuous optimum. The 3-approximation algorithm in [18] for the kCwO problem is one such example. However, the best known algorithm for kCwO for the discrete case (when  $X = C$ ) is a 2-approximation by Chakrabarty, Goyal, and Krishnaswamy [12] which proceeds via LP rounding, and does not give a 2-approximation for continuous kCwO. This was explicitly noted in a work by Ding, Yu, and Wang (“... *unclear of the resulting approximation ratio for the problem in Euclidean space.*”) [27], that describes a 2-approximation for kCwO in Euclidean space, however, violating the number of clients served. We give a proper 2-approximation for the continuous kCwO problem (with no assumptions on the metric space) with no violations.

## 1.2 Our Technical Insight

Most state of the art approximation algorithms for center-based clustering problems are based on LP relaxations where one typically has variables  $y_i$  for *every* potential location of a center. When the set  $X$  is large, this approach becomes infeasible. Our main technical insight, underlying all our results, is to use a different style of linear program with polynomially many variables but exponentially many constraints. We then use the *round-or-cut* framework to obtain our approximation factor. More precisely, given a *potential* solution to our program, we either “round” it to get a desired solution within the asserted approximation factor, or we find a *separating hyperplane* proving that this potential solution is infeasible. Once this hyperplane is fed to the ellipsoid algorithm [32], the latter generates another potential solution, and the process continues. Due to the ellipsoid method’s guarantees, we obtain our approximation factor in polynomial time.

For every client  $v \in C$ , our LP relaxation has variables of the form  $y(v, r)$ , indicating whether there is some point  $x \in X$  in an  $r$ -radius around  $v$  which is “open” as a center. Throughout the paper we use  $r$  as a quantity varying “continuously”, but it can easily be discretized, with a loss of at most  $\frac{1}{\text{poly}(n)}$ , to arise from a set of size  $\leq \text{poly}(n)$ . Thus there are only polynomially many such variables. We add the natural “monotonicity” constraints:  $y(v, r) \leq y(v, s)$  whenever  $r \leq s$ . Interestingly, for one of the applications, we also need the monotonicity constraints for non-concentric balls: if  $B(v, r) \subseteq B(u, s)$ , then we need  $y(v, r) \leq y(u, s)$ .

We have a variable  $C_v$  indicating the cost the client  $v$  pays towards the optimal solution. Next, we connect the  $C_v$ ’s and the  $y(v, r)$ ’s in the following ways (when  $\beta = 1$ , and something similar when  $\beta = 2$ ). One connection states that for any  $r$ ,  $C_v \geq r \cdot (1 - y(v, r))$  and we add these to our LP. For the last two applications listed above, this suffices. However, one can also state the stronger condition of  $C_v \geq \int_0^\infty (1 - y(v, r)) dr$ . Indeed, the weaker constraint is the “Markov-style inequality” version of the stronger constraint.

Our second set of constraints restrict the  $y(v, r)$ ’s to be “not too large”. For instance, for the fair  $k$ -median or kCwO problems where we are only allowed  $k$  points from  $X$ , we assert that for *any* set  $\mathcal{B}$  of disjoint balls  $B(v, r)$ , we must have the sum of the respective  $y(v, r)$ ’s to be at most  $k$ . This set of constraints is exponentially many, and this is the set of constraints that need the round-or-cut machinery. For the  $\lambda$ -UFL problem, we have that the sum of the  $y(v, r)$ ’s scaled by  $\lambda$  plus the sum of the  $C_v$ ’s should be at most  $\text{opt}_g$ , which is a running guess of  $\text{opt}$ .

Once we set up the framework above, then we can *port many existing rounding* algorithms for the discrete clustering problems without much hassle. In particular, this is true for rounding algorithms which use the

$C_v$ 's as the core driving force. For the continuous  $\lambda$ -UFL problem, we port the rounding algorithm from the paper [50] by Shmoys, Tardos, and Aardal. For the continuous  $k$ -means problem, we port the rounding algorithm from the paper [17] by Charikar, Guha, Shmoys, and Tardos. For the continuous fair  $k$ -median problem, we port the rounding algorithm from the paper [14] by Chakrabarty and Negahbani, which itself builds on the algorithm present in the paper [2] by Alamdari and Shmoys. For the continuous kCwO problem, we port the rounding algorithm present in the paper [12] by Chakrabarty, Goyal, and Krishnaswamy.

Our results fall short for the continuous  $k$ -median problem (without fairness), where we can port the rounding algorithm from the paper [17] and get a 6.67-approximation. This, however, does not improve upon the 5.35-factor mentioned earlier.

### 1.3 Other Related Works and Discussion

The continuous  $k$ -means and median problems have been investigated quite a bit in the specific setting when  $X = \mathbb{R}^m$  and when  $d(u, v)$  is the  $\ell_2$  distance. The paper [47] by Matoušek describes an  $(1 + \varepsilon)$ -approximation (PTAS) that runs in time  $O(n \log^k n \cdot \varepsilon^{-O(k^2 m)})$ . This led to a flurry of results [35, 26, 43, 19, 30] on obtaining PTASes with better dependencies on  $k$  and  $m$  via the applications of coresets. There is a huge and growing literature on coresets, and we refer the interested reader to the paper [25] by Cohen-Addad, Saulpic, and Schwiegelshohn, and the references within, for more information. Another approach to the continuous  $k$ -means problem has been local search. The paper [41] which describes a  $9 + \varepsilon$ -approximation was first stated for the geometric setting, however it also went via the discretization due to Matoušek [47] and suffered a running time of exponential dependency on the dimension. More recent papers [31, 23] described local-search based PTASes for metrics with doubling dimension  $D$ , with running time exponentially depending on  $D$ . These doubling metrics generalize  $(\mathbb{R}^m, \ell_2)$ -metrics. However, none of the above ideas seem to suggest better constant factor approximations for the continuous  $k$ -median/means problem in the general case, and indeed even when  $X = \mathbb{R}^m$  but  $m$  is part of the input.

The  $k$ -means problem in the metric space  $(\mathbb{R}^m, \ell_2)$ , where  $m$  and  $k$  are not constants, has been studied extensively [52, 6, 20, 24, 1, 21], and is called the Euclidean  $k$ -means problem. The discrete version of this problem was proved APX-hard in 2000 [52], but the APX-hardness of the continuous version was proved much later, in 2015 [6]. More recently, the hardness results for both versions have been improved: the discrete Euclidean  $k$ -means problem is hard to approximate to factor 1.17, while the continuous problem is hard to approximate to factor 1.07 [20]. Moreover, under assumption of a complexity theoretic hypothesis called the Johnson coverage hypothesis, these numbers have been improved to 1.73 and 1.36, respectively [24]. On the algorithmic side, the discrete Euclidean  $k$ -means problem admits a better approximation ratio than the general case: a 6.36 approximation was described in 2017 [1], which was very recently improved to 5.912 [21].

We believe that our paper takes the first stab at getting approximation ratios better than  $\beta \times$  the best discrete factor for the *continuous* clustering problems. Round-or-cut is a versatile framework for approximation algorithm design with many recent applications [49, 11, 3, 13, 4], and the results in our paper is yet another application of this paradigm. However, many questions remain. We believe that the most interesting question to tackle is the continuous  $k$ -median problem. The best known discrete  $k$ -median algorithms are, in fact, combinatorial in nature, and are obtained via applying the primal-dual/dual-fitting based methods [39, 38, 45, 10] on the discrete LP. However, their application still needs an explicit description of the facility set, and it is interesting to see if they can be *directly* ported to the continuous setting.

All the algorithms in our paper, actually *still* open centers from  $C$ . Even then, we are able to do better than simply reducing to the discrete case, because we do not commit to the  $\beta$  loss upfront, and instead round from a fractional solution that can open centers anywhere in  $X$ . This raises an interesting question for the  $k$ -median problem (or any other center based clustering problem): consider the potentially infinite-sized LP which has variables  $y_i$  for all  $i \in X$ , but restrict to the optimal solution which only is allowed to open centers from  $C$ . How big is this “integrality gap”? It is not too hard to show that for the  $k$ -median problem this is between 2 and 4. The upper bound gives hope we can get a true 4-approximation for the continuous  $k$ -median problem, but it seems one would need new ideas to obtain such a result.

**Organization of this Paper** In the main body, we focus on the continuous  $\lambda$ -UFL and the continuous fair  $k$ -median results, since we believe that they showcase the technical ideas in this paper. Proofs of certain

statements have been deferred to the appendix. The description of the results on continuous  $k$ -means and continuous  $k$ -center with outliers can be found in Appendix B and Appendix C, respectively.

## 2 Preliminaries

Given a metric space  $(X, d)$  on points  $X$  with pairwise distances  $d$ , we use the notation  $d(v, S) = \min_{i \in S} d(v, i)$  for  $v \in X$  and  $S \subseteq X$  to denote  $v$ 's distance to the set  $S$ .

**Definition 1** (Continuous  $k$ -median (Cont- $k$ -Med)). *The input is a metric space  $(X, d)$ , clients  $C \subseteq X, |C| = n \in \mathbb{N}$ , and  $k \in \mathbb{N}$ . The goal is to find  $S \subseteq X, |S| = k$  minimizing  $\text{cost}(S) := \sum_{v \in C} d(v, S)$ .*

**Definition 2** (Continuous Fair  $k$ -median (ContFair- $k$ -Med)). *Given the Cont- $k$ -Med input, plus fairness radii  $r : C \rightarrow [0, \infty)$ , the goal is to find  $S \subseteq X, |S| = k$  such that  $\forall v \in C, d(v, S) \leq r(v)$ , minimizing  $\text{cost}(S)$ .*

In the Uncapacitated Facility Location (UFL) problem, the restriction of opening only  $k$  facilities is replaced by having a cost associated with opening each facility. When these costs are equal to the same value  $\lambda$  for all facilities, the problem is called  $\lambda$ -UFL.

**Definition 3** (Continuous  $\lambda$ -UFL (Cont- $\lambda$ -UFL)). *Given a metric space  $(X, d)$ , clients  $C \subseteq X, |C| = n \in \mathbb{N}$ , and  $\lambda \geq 0$ , find  $S \subseteq X$  that minimizes the sum of “connection cost”  $\text{cost}_C(S) := \sum_{v \in C} d(v, S)$  and “facility opening cost”  $\text{cost}_F(S) := \lambda|S|$ .*

Let  $\Delta = \max_{u, v \in C} d(u, v)$  denote the diameter of a metric  $(X, d)$ . For  $x \in X, 0 \leq r \leq \Delta$ , the ball of radius  $r$  around  $x$  is  $B(x, r) := \{x' \in X \mid d(x', x) \leq r\}$ . Throughout the paper, we use balls of the form  $B(v, r)$  where  $v$  is a client and  $r \in \mathbb{R}$ . To circumvent the potentially infinite number of radii, the radii can be discretized into  $I_\varepsilon = \{\varepsilon, 2\varepsilon, \dots, \lceil \Delta/\varepsilon \rceil \varepsilon\}$  for a small constant  $\varepsilon = O(1/n^2)$ . Thereupon, we can appeal to the following lemma to bound the size of  $I_\varepsilon$  by  $O(n^5)$ .

**Lemma 4** (Rewording of Lemma 4.1, [1]). *Losing a factor of  $(1 + \frac{100}{n^2})$ , we can assume that for any  $u, v \in C, 1 \leq d(u, v) \leq n^3$ .*

For simplicity of exposition, we present our techniques using radii in  $\mathbb{R}$ , and observe that discretizing to  $I_\varepsilon$  incurs an additive loss of at most  $O(n\varepsilon) = O(1/n)$  in our guarantees. We also note that  $\log \text{opt} \leq \log(n\Delta) = O(\log n)$  by the above, which enables us to efficiently binary-search over our guesses  $\text{opt}_g$ .

## 3 Continuous $\lambda$ -UFL

We start this section with our 2.32-approximation for Cont- $\lambda$ -UFL (Theorem 5). For this, we introduce a new linear programming formulation, and adapt the rounding algorithm of Shmoys-Tardos-Aardal to the new program. The resulting procedure exhibits our main ideas, and serves as a warm-up for the remaining sections. Also, in Section 3.2, we prove that it is NP-hard to approximate Cont- $\lambda$ -UFL within a factor of  $2 - o(1)$ , using ideas due to Cohen-Addad, Karthik, and Lee [22]. This shows that the continuous version cannot be approximated as well as the discrete version, which has a best-known approximation factor of 1.463 [44].

### 3.1 Approximation algorithm

This subsection is dedicated to proving the following theorem:

**Theorem 5.** *There is a polynomial time algorithm that, for an instance of Cont- $\lambda$ -UFL with optimum  $\text{opt}$ , yields a solution with cost at most  $(\frac{2}{1-\varepsilon} + \varepsilon)\text{opt} < 2.32 \cdot \text{opt}$ . Here  $\varepsilon = O(\frac{1}{n^2})$ .*

We design the following linear program for Cont- $\lambda$ -UFL. We use variables  $C_v$  for the connection cost of each client  $v$ , and  $y(v, r)$  for the number of facilities opened within each ball of the form  $B(v, r)$ . We also

use a guess of the optimum  $\text{opt}_g$ , which we will soon discuss how to obtain. Throughout, we use  $y(B)$  as shorthand for  $y(v, r)$  where  $B = B(v, r)$ .

$$\lambda \sum_{B \in \mathcal{B}} y(B) + \sum_{v \in C} C_v \leq \text{opt}_g \quad \forall \mathcal{B} \subseteq \{B(v, r)\}_{\substack{v \in C \\ r \in \mathbb{R}}} \text{ pairwise disjoint} \quad (\text{UFL})$$

$$\int_0^\infty (1 - y(v, r)) dr \leq C_v \quad \forall v \in C, r \in \mathbb{R} \quad (\text{UFL-1})$$

$$\begin{aligned} y(v, r) &\leq y(v, r') & \forall v \in C, r, r' \in \mathbb{R} \text{ s.t. } r \leq r' & (\text{UFL-2}) \\ y(v, r) &\geq 0, C_v \geq 0 & \forall v \in C, r \in \mathbb{R} \end{aligned}$$

Observe that, given a solution  $S \subseteq X$  of cost at most  $\text{opt}_g$ , we can obtain a feasible solution of UFL as follows. For client  $v \in C$ , we set  $C_v = d(v, S)$ . For  $v \in C, r \in \mathbb{R}$ , we set  $y(v, r) = 0$  for  $r < d(v, S)$  and  $y(v, r) = 1$  for  $r \geq d(v, S)$ .

Our approach is to round a solution  $(C, y)$  of UFL. Observe that there are polynomially many constraints of the form (UFL-1) and (UFL-2); hence, we can efficiently obtain a solution  $(C, y)$  that satisfies them. So for the remainder of this section, we assume that those constraints are satisfied. On the other hand, there are infinitely many constraints of type (UFL). This is why we employ a round-or-cut framework via the ellipsoid algorithm [32]. We begin with an arbitrary  $\text{opt}_g$ , and when ellipsoid asks us if a proposed solution  $(C, y)$  is feasible, we run the following algorithm.

The algorithm inputs  $\alpha < 1$ , and defines  $r_\alpha(v)$  as the minimum radius at which client  $v \in C$  has at least  $\alpha$  mass of open facilities around it. First, all clients are deemed *uncovered* ( $U = C$ ). Iteratively, the algorithm picks the  $j$ , i.e. the uncovered client, with the smallest  $r_\alpha(j)$ .  $j$  is put into the set  $\text{Reps}_{(C, y), \alpha}$ . Any client  $v$  within distance  $r_\alpha(j) + r_\alpha(v)$  of  $j$  is considered a *child* of  $j$  and is now *covered*. When all clients are covered, i.e.  $U = \emptyset$ , the algorithm outputs  $\text{Reps}_{(C, y), \alpha}$ .

---

**Algorithm 1** Filtering for Cont- $\lambda$ -UFL

---

**Input:** A proposed solution  $(\{C_v\}_{v \in C}, \{y(v, r)\}_{v \in C, r \in \mathbb{R}})$  for UFL, parameter  $\alpha \in (e^{-2}, 1)$

- 1:  $r_\alpha(v) \leftarrow \min\{r \in \mathbb{R} \mid y(v, r) \geq \alpha\}$  for all  $v \in C$
- 2:  $\text{Reps}_{(C, y), \alpha} \leftarrow \emptyset$  ▷ “representative” clients
- 3:  $U \leftarrow C$  ▷ “uncovered” clients
- 4: **while**  $U \neq \emptyset$  **do**
- 5:     Pick  $j \in U$  with minimum  $r_\alpha(j)$
- 6:      $\text{child}(j) \leftarrow \{v \in U \mid d(v, j) \leq r_\alpha(v) + r_\alpha(j)\}$
- 7:      $U \leftarrow U \setminus \text{child}(j)$
- 8:      $\text{Reps}_{(C, y), \alpha} \leftarrow \text{Reps}_{(C, y), \alpha} \cup j$
- 9: **end while**

**Output:**  $\text{Reps}_{(C, y), \alpha}$

---

Notice that, by construction, the collection of balls  $\{B(j, r_\alpha(j))\}_{j \in \text{Reps}_{(C, y), \alpha}}$  is pairwise disjoint. Hence, the following constraint, which we call  $\text{Sep}_{(C, y), \alpha}$ , is of the form (UFL):

$$\lambda \sum_{j \in \text{Reps}_{(C, y), \alpha}} y(j, r_\alpha(j)) + \sum_{v \in C} C_v \leq \text{opt}_g \quad (\text{Sep}_{(C, y), \alpha})$$

We will show that

**Lemma 6.** *If  $(C, y)$  satisfies (UFL-1), (UFL-2), and  $\text{Sep}_{(C, y), \alpha}$ , then there exists a suitable  $\alpha \in (e^{-2}, 1)$  for which the output of Algorithm 1 has cost at most  $\frac{2}{1-e^{-2}} \text{opt}_g$ .*

Thus, if we find that the desired approximation ratio is not attained, then it must be that  $\text{Sep}_{(C, y), \alpha}$  was not satisfied, and we can pass it to ellipsoid as a separating hyperplane. If ellipsoid finds that the feasible region of our linear program is empty, then we increase  $\text{opt}_g$  and try again. Otherwise, we obtain a solution  $\text{Reps}_{(C, y), \alpha}$  that attains the desired guarantees.

We now analyze Algorithm 1 to prove Lemma 6.

*Proof of Lemma 6.* For this proof, we will fix  $(C, y)$ , and refer to  $\text{Reps}_{(C,y),\alpha}$  as  $\text{Reps}_\alpha$ .

To prove a suitable  $\alpha$  exists, assume  $\alpha$  is picked uniformly at random from  $(\beta, 1)$  for some  $0 < \beta < 1$ ; we will see later that  $\beta = e^{-2}$  is optimal. Take  $\text{Reps}_\alpha$ , the output of Algorithm 1 on  $(C, y)$ . By definition of  $r_\alpha$ ,  $\sum_{j \in \text{Reps}_\alpha} y(j, r_\alpha(j)) \geq \alpha |\text{Reps}_\alpha|$ . Thus  $\text{cost}_F(\text{Reps}_\alpha) = \lambda |\text{Reps}_\alpha| \leq \frac{1}{\alpha} \cdot \lambda \sum_{j \in \text{Reps}_\alpha} y(j, r_\alpha(j))$ , which implies

$$\mathbf{Exp}[\text{cost}_F(\text{Reps}_\alpha)] \leq \frac{\ln(1/\beta)}{(1-\beta)} \lambda \sum_{j \in \text{Reps}_\alpha} y(j, r_\alpha(j)). \quad (1)$$

To bound the expected connection cost, take  $v \in C$  and observe that, since all the clients are ultimately covered in Algorithm 1, there has to exist  $j \in \text{Reps}_\alpha$  for which  $v \in \text{child}(j)$ . By construction of  $\text{child}$ ,  $d(v, j) \leq r_\alpha(v) + r_\alpha(j)$ , which is at most  $2r_\alpha(v)$  by our choice of  $j$  in Line 5. Thus, for any client  $v$ , we get  $d(v, \text{Reps}_\alpha) \leq d(v, j) \leq 2r_\alpha(v)$ . So we are left to bound  $\mathbf{Exp}[r_\alpha(v)]$  for an arbitrary client  $v \in C$ .

We have that  $\mathbf{Exp}[r_\alpha(v)] = \frac{1}{1-\beta} \int_\beta^1 r_\alpha(v) d\alpha \leq \frac{1}{1-\beta} \int_0^1 r_\alpha(v) d\alpha$ . We notice that at  $\alpha = y(v, r)$ ,  $r_\alpha(v) = r$ . Also,  $r_0(v) = 0$ . So given (UFL-2) for all balls  $B(v, r)$  with  $r \in \mathbb{R}$ , we can apply a change of variable to the integral to get  $\int_0^1 r_\alpha(v) d\alpha = \int_{r_0(v)}^{r_1(v)} (y(v, r_1(v)) - y(v, r)) dr \leq \int_0^\infty (1 - y(v, r)) dr \leq C_v$ , where the last inequality is by (UFL-1). Thus we have  $\mathbf{Exp}[r_\alpha(v)] \leq \frac{C_v}{1-\beta}$ . Summing  $d(v, \text{Reps}_\alpha)$  over all  $v \in C$  we have

$$\mathbf{Exp}[\text{cost}_C(\text{Reps}_\alpha)] = \sum_{v \in C} \mathbf{Exp}[d(v, \text{Reps}_\alpha)] \leq 2 \sum_{v \in C} \mathbf{Exp}[r_\alpha(v)] \leq \frac{2}{1-\beta} \sum_{v \in C} C_v. \quad (2)$$

To balance  $\text{cost}_F$  from (1) and  $\text{cost}_C$  from (2), we set  $\beta = e^{-2}$ . The expected  $\text{Cont-}\lambda\text{-UFL}$  cost of  $\text{Reps}_\alpha$  is, using  $\text{Sep}_{(C,y),\alpha}$ ,

$$\mathbf{Exp}[\text{cost}_F(\text{Reps}_\alpha) + \text{cost}_C(\text{Reps}_\alpha)] \leq \frac{2}{1-e^{-2}} \left( \lambda \sum_{j \in \text{Reps}_\alpha} y(j, r_\alpha(j)) + \sum_{v \in C} C_v \right) \leq \frac{2 \cdot \text{opt}_g}{1-e^{-2}}.$$

Since the bound holds in expectation over a random  $\alpha$ , there must exist an  $\alpha \in (\beta, 1)$  that satisfies it deterministically.  $\square$

To obtain a suitable  $\alpha$ , we can adapt the derandomization procedure from the discrete version [50]. The procedure relies on having polynomially many interesting radii; for this, we recall that while we have used  $r \in \mathbb{R}$  for simplicity, our radii are actually  $r \in I_\varepsilon$ ,  $|I_\varepsilon| = O(n^5)$ .

### 3.2 Hardness of approximation

Our hardness result for this problem is as follows:

**Theorem 7.** *Given an instance of  $\text{Cont-}\lambda\text{-UFL}$  and  $\varepsilon > 0$ , it is NP-hard to distinguish between the following:*

- *There exists  $S \subseteq X$  such that  $\text{cost}_F(S) + \text{cost}_C(S) \leq (1 + 6\varepsilon)n$*
- *For any  $S \subseteq X$ ,  $\text{cost}_F(S) + \text{cost}_C(S) \geq (2 - \varepsilon)n$*

Thus we exhibit hardness of approximation up to a factor of  $\frac{2-\varepsilon}{1+6\varepsilon}$ , which tends to 2 as  $\varepsilon \rightarrow 0$ . Our reduction closely follows the hardness proof for  $\text{Cont-}k\text{-Med}$  [22]. We relegate the details to Appendix D.

## 4 Continuous Fair $k$ -Median

The main result of this section is the following theorem.

**Theorem 8.** *There exists a polynomial time algorithm for  $\text{ContFair-}k\text{-Med}$  that, for an instance with optimum cost  $\text{opt}$ , yields a solution with cost at most  $8\text{opt} + \varepsilon$ , in which, each client  $v \in C$  is provided an open facility within distance  $8r(v) + \varepsilon$  of itself. Here  $\varepsilon = O\left(\frac{1}{n^2}\right)$ .*

We create a round-or-cut framework, via the ellipsoid algorithm [32], that adapts the Chakrabarty-Negahbani algorithm [14] to the continuous setting. For this, we will modify the UFL linear program to suit ContFair- $k$ -Med. As before,  $\text{opt}_g$  is a guessed optimum,  $C_v$  is the cost share of a client  $v \in C$ , and  $y(v, r)$  represents the number of facilities opened in  $B(v, r)$ . There are two key modifications. First, we expand the monotonicity constraints of the form (UFL-2) to include *non-concentric* balls, which are crucial for adapting the fairness guarantee of Chakrabarty and Negahbani [14]. Second, we enforce the fairness constraints by requiring  $y(v, r(v)) \geq 1$  for each client  $v \in C$ .

$$\sum_{v \in C} C_v \leq \text{opt}_g \quad (\text{LP})$$

$$\sum_{B \in \mathcal{B}} y(B) \leq k \quad \forall \mathcal{B} \subseteq \{B(v, r)\}_{v \in C, r \in \mathbb{R}} \text{ pairwise disjoint} \quad (\text{LP-1})$$

$$\int_0^\infty (1 - y(v, r)) dr \leq C_v \quad \forall v \in C \quad (\text{LP-2})$$

$$y(u, r) \leq y(v, r') \quad \forall u, v \in C, r, r' \in \mathbb{R}, B(u, r) \subseteq B(v, r') \quad (\text{LP-3})$$

$$y(v, r(v)) \geq 1 \quad \forall v \in C \quad (\text{LP-4})$$

$$y(v, r) \geq 0, C_v \geq 0 \quad \forall v \in C, r \in \mathbb{R}$$

We will frequently use the following property of LP. See Appendix A.2 for the proof.

**Lemma 9.** *Consider a solution  $(C, y)$  of LP. If for a client  $v$ ,  $(C, y)$  satisfies all constraints of the form (LP-2) and (LP-3) involving  $v$ , then for any  $r_0 \in \mathbb{R}$ ,  $C_v \geq r_0 (1 - y(v, r_0))$ .*

As before, we will only worry about the constraints that are exponentially many. These are (LP-1). For this, we use ellipsoid [32]. Given a proposed solution  $(C, y)$  of LP, we construct  $\text{Reps}_{(C, y)} \subseteq C$ , as follows.

We first perform a filtering step. For each  $v \in C$ , we define  $R(v) := \min r(v), 2C_v$ . In the beginning, all clients are “uncovered” (i.e.  $U = C$ ). In each iteration, let  $j \in U$  be the uncovered client with the minimum  $R(j)$ ; and add  $j$  to our set of “representatives”  $\text{Reps}_{(C, y)}$ . Any  $v \in U$  within distance  $2R(v)$  of  $j$  (including  $j$  itself) will be added to the set  $\text{child}(j)$ , and will be removed from  $U$ . After all clients are covered, i.e.  $U = \emptyset$ , the algorithm outputs  $\text{Reps}_{(C, y)}$ . For a formal description of this algorithm, see Appendix A.1.

For a  $j \in \text{Reps}_{(C, y)}$ , let  $s(j)$  be the closest client to  $j$  in  $\text{Reps}_{(C, y)} \setminus \{j\}$ . Let  $a(j) := \frac{d(j, s(j))}{2}$ . So the collection of balls  $\{B(j, a(j))\}_{j \in \text{Reps}_{(C, y)}}$  is pairwise disjoint, and the following constraint, which we call  $\text{Sep}_{(C, y)}$ , is of the form (LP-1).

$$\sum_{j \in \text{Reps}_{(C, y)}} y(j, a(j)) \leq k \quad (\text{Sep}_{(C, y)})$$

We have that

**Lemma 10.** *If  $(C, y)$  satisfies (LP-2)-(LP-4) and  $\text{Sep}_{(C, y)}$ , then*

1.  $\forall j \in \text{Reps}_{(C, y)}, y(j, a(j)) \geq \frac{1}{2}$

2.  $|\text{Reps}_{(C, y)}| \leq 2k$

*Proof.* Fix  $j \in \text{Reps}_{(C, y)}$ . By Line 6 of Algorithm 3,

$$a(j) = \frac{d(j, s(j))}{2} \geq R(j) \quad (3)$$

So if  $R(j) = r(j)$ , then by (LP-3) and (LP-4),  $y(j, a(j)) \geq y(j, r(j)) \geq 1$ . Else  $R(j) = 2C_j$ . By Lemma 9,  $C_j \geq a(j) (1 - y(j, a(j)))$ .

If  $C_j = 0$ , then this implies  $y(j, a(j)) \geq 1$ . Otherwise, substituting  $a(j)$  by  $R(j)$  from (3) and setting  $R(j) = 2C_j$  gives  $C_j \geq 2C_j (1 - y(j, a(j)))$ , i.e.  $y(j, a(j)) \geq \frac{1}{2}$ . Now, by  $\text{Sep}_{(C, y)}$ , we have  $k \geq \sum_{j \in \text{Reps}_{(C, y)}} y(j, a(j)) \geq \frac{1}{2} \cdot |\text{Reps}_{(C, y)}|$ .  $\square$



So if we find that  $|\text{Reps}_{(C,y)}| > 2k$ , then  $\text{Sep}_{(C,y)}$  must be violated, and we can pass it to ellipsoid as a separating hyperplane. Hence in polynomial time, we either find that our feasible region is empty, or we get  $(C, y)$  and  $\text{Reps}_{(C,y)}$  such that  $(C, y)$  satisfies (LP), (LP-2)-(LP-4), and  $\text{Sep}_{(C,y)}$ . In the first case, we increase  $\text{opt}_g$  and try again. In the latter case, we round  $(C, y)$  further to attain our desired approximation ratios, via a rounding algorithm that we will now describe. This algorithm focuses on  $\text{Reps}_{(C,y)}$  and ignores other clients, as justified by the following lemma.

**Lemma 11.**  $S \subseteq X$  be a solution to *ContFair- $k$ -Med*. Consider a proposed solution  $(C, y)$  of LP that satisfies (LP). Then  $\sum_{v \in C} d(v, S) \leq \sum_{j \in \text{Reps}_{(C,y)}} |\text{child}(j)| d(j, S) + 4\text{opt}_g$ .

The proof closely follows from a standard technique for the discrete version [17, 14]. We provide the proof in Appendix A.3.

Our algorithm will also ignore facilities outside  $\text{Reps}_{(C,y)}$ , so our solution will be a subset of  $\text{Reps}_{(C,y)}$ . For the remainder of this section, we fix  $(C, y)$ , and refer to  $\text{Reps}_{(C,y)}$  as  $\text{Reps}$ . We write the following polynomial-sized linear program, DLP, where  $\text{Reps}$  are the only clients and the only facilities. The objective function of DLP is a lower bound on  $\sum_{j \in \text{Reps}} |\text{child}(j)| d(j, S)$ , so hereafter we compare our output with DLP. We do not include fairness constraints in this program, and we will see later that it is not necessary to do so.

In DLP, the variables  $z_i$  for each  $i \in \text{Reps}$  denote whether  $i$  is open as a facility. The variables  $x_{ij}$  for  $i, j \in \text{Reps}$  denote whether the client  $j$  uses the facility  $i$ .

$$\text{minimize } \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in \text{Reps}} x_{ij} d(j, i) \quad (\text{DLP})$$

$$\sum_{i \in \text{Reps}} z_i \leq k \quad (\text{DLP-1})$$

$$\sum_{i \in \text{Reps}} x_{ij} = 1 \quad \forall j \in \text{Reps} \quad (\text{DLP-2})$$

$$x_{ij} \leq y_i \quad \forall i, j \in \text{Reps} \quad (\text{DLP-3})$$

$$x_{ij} \geq 0, z_i \geq 0 \quad \forall i, j \in \text{Reps}$$

We will now round  $(C, y)$  to an integral solution of DLP. Our first step is to convert  $(C, y)$  to a fractional solution  $(\bar{x}, \bar{z})$  of DLP. To do this, for each  $j \in \text{Reps}$ , we consolidate the  $y$ -mass in  $B(j, a(j))$  onto  $j$ , i.e. we set  $\bar{z}_j = y(j, a(j))$ . By Lemma 10.1, each  $\bar{z}_j$  is then at least  $\frac{1}{2}$ . This allows  $j$  to use only itself and  $s(j)$  as its fractional facilities.

---

**Algorithm 2** Consolidation for *ContFair- $k$ -Med*

---

**Input:** A proposed solution  $(\{C_v\}_{v \in C}, \{y(v, r)\}_{v \in C, r \in I_\varepsilon})$  for LP, and  $\text{Reps}$  from Algorithm 3

- 1: **for**  $j \in \text{Reps}$  **do**
- 2:    $s(j) \leftarrow \arg \min_{v \in \text{Reps} \setminus j} d(j, v)$
- 3:    $a(j) \leftarrow d(j, s(j))/2$
- 4:    $\bar{z}_j \leftarrow \min \{y(j, a(j)), 1\}$
- 5:    $\bar{x}_{jj} \leftarrow \bar{z}_j$
- 6:    $\bar{x}_{js(j)} \leftarrow 1 - \bar{z}_j$
- 7: **end for**

**Output:**  $(\bar{x}, \bar{z})$

---

**Lemma 12.**  $\forall j \in \text{Reps}, \bar{z}_j \geq \frac{1}{2}$ , and  $(\bar{x}, \bar{z})$  is a feasible solution of DLP with cost at most  $2\text{opt}_g$ .

*Proof.* For a  $j \in \text{Reps}$ , if  $y(j, a(j)) = 1$  then  $\bar{z}_j = 1$ . Otherwise, by Lemma 10.1,  $\bar{z}_j = y(j, a(j)) \geq \frac{1}{2}$ .

Hence  $1 - \bar{z}_j \leq \frac{1}{2} \leq \bar{z}_{s(j)}$ , which implies feasibility by construction and  $\text{Sep}_{(C,y)}$ . It also implies that  $\sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i) = (1 - \bar{z}_j) d(j, s(j))$ . If  $y(j, a(j)) = 1$ , then the RHS above is  $0 \leq 2C_j$ . Otherwise

$\sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i) = (1 - y(j, a(j))) d(j, s(j)) = 2(1 - y(j, a(j))) a(j) \leq 2C_j$  where the last inequality follows from Lemma 9. Multiplying by  $|\text{child}(j)|$  and summing over all  $j \in \text{Reps}$ , we have by Line 5 in Algorithm 3,

$$\sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i) \leq 2 \sum_{j \in \text{Reps}} |\text{child}(j)| C_j \leq 2 \sum_{j \in \text{Reps}} \sum_{v \in \text{child}(j)} C_v = 2 \sum_{v \in C} C_v$$

which is at most  $2\text{opt}_g$  by (LP).  $\square$

Now, to round  $(\bar{x}, \bar{z})$  to an integral solution, we appeal to an existing technique [17, 14]. We state the relevant result here, and provide the proof in Appendix A.4.

**Lemma 13** ([17, 14]). *Let  $(\bar{x}, \bar{z})$  be a feasible solution of DLP with cost at most  $2\text{opt}_g$ , such that  $\forall j \in \text{Reps}$ ,  $\bar{z}_j \geq \frac{1}{2}$ . Then there exists a polynomial time algorithm that produces  $S \subseteq \text{Reps}$  such that (1)  $|S| = k$ ; (2) If  $\bar{z}_j = 1$ , then  $j \in S$ ; (3)  $\forall j \in \text{Reps}$ , at least one of  $j, s(j)$  is in  $S$ ; and (4)  $\sum_{j \in \text{Reps}} |\text{child}(j)| d(j, S) \leq 4\text{opt}_g$ .*

Thus, by Lemma 11, we have shown that  $\sum_{v \in C} d(v, S) \leq \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in S} d(j, S) + 4\text{opt}_g \leq 8\text{opt}_g$ . Now we show the fairness ratio, adapting a related result [14, Lemma 3] from the discrete version. This is where we crucially require the monotonicity constraints (LP-3) for *non-concentric* balls.

**Lemma 14.**  $\forall v \in C, d(v, S) \leq 8r(v)$ .

*Proof.* Fix  $v \in C$ , and let  $j \in \text{Reps}$  such that  $v \in \text{child}(j)$ . By construction of  $\text{child}(j)$  in Algorithm 3,

$$d(v, j) \leq 2R(v) \leq 2r(v) \tag{4}$$

So if  $j \in S$ , then we are done. Otherwise, by Lemma 13.2,  $\bar{z}_j < 1$ , i.e. by Algorithm 2,  $y(j, a(j)) < 1$ . But by the fairness constraints (LP-4),  $y(v, r(v)) \geq 1$ . So by the monotonicity constraints (LP-3),  $B(v, r(v)) \not\subseteq B(j, a(j))$ , as otherwise we would have  $1 \leq y(v, r(v)) \leq y(j, a(j)) < 1$ , a contradiction.

So fix  $w \in B(v, r(v)) \setminus B(j, a(j))$ . We have

$$a(j) < d(j, w), \quad d(v, w) \leq r(v) \tag{5}$$

By Lemma 13.3, either  $j \in S$  or  $s(j) \in S$ , so

$$\begin{aligned} d(j, S) &\leq d(j, s(j)) = 2a(j) < 2d(j, w) && \dots \text{ by (5)} \\ &\leq 2(d(v, j) + d(v, w)) \leq 2(2r(v) + r(v)) && \dots \text{ by (4) and (5)} \\ &= 6r(v) \end{aligned}$$

So, by (4),  $d(v, S) \leq d(v, j) + d(j, S) \leq 2r(v) + 6r(v) = 8r(v)$ .  $\square$

Thus we have proved Theorem 8.

We observe here that, by the simple reduction of setting all  $r(v)$ 's to  $\infty$ , Theorem 8 implies a solution of cost  $8\text{opt} + \varepsilon$  for **Cont- $k$ -Med**. We improve this ratio via an improved rounding procedure by Charikar, Guha, Shmoys, and Tardos [17], which rounds  $(\bar{x}, \bar{z})$  such that  $\sum_{j \in \text{Reps}} |\text{child}(j)| d(j, S) \leq \frac{8}{3}\text{opt}_g$ , instead of the  $4\text{opt}_g$  that we obtain above. This yields:

**Corollary 15.** *There exists a polynomial time algorithm for Cont- $k$ -Med that, on an instance with optimum cost  $\text{opt}$ , yields a solution of cost at most  $6\frac{2}{3}\text{opt} + \varepsilon$ .*

This improved rounding, however, no longer guarantees to open either  $j$  or  $s(j)$  for each  $j \in \text{Reps}$ . Such a guarantee (Lemma 13.3) is crucial to our fairness bound in Lemma 14. So the improvement is not naively adaptable to **ContFair- $k$ -Med**.

## Acknowledgements

We thank the reviewers for their comments, which helped us improve our exposition and exhibit a stronger connection to existing work.

## References

- [1] Sara Ahmadian, Ashkan Norouzi-Fard, Ola Svensson, and Justin Ward. Better guarantees for k-means and euclidean k-median by primal-dual algorithms. *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 61–72, 2017.
- [2] Soroush Alamdari and David B. Shmoys. A bicriteria approximation algorithm for the  $k$ -center and  $k$ -median problems. In *Proc., Workshop on Approximation and Online Algorithms (WAOA)*, pages 66–75. Springer, 2017.
- [3] Hyung-Chan An, Mohit Singh, and Ola Svensson. LP-based algorithms for capacitated facility location. *SIAM Journal on Computing (SICOMP)*, 46(1):272–306, 2017. Preliminary version in FOCS 2014.
- [4] Georg Anegg, Haris Angelidakis, Adam Kurpisz, and Rico Zenklusen. A technique for obtaining true approximations for k-center with covering constraints. *Math. Program.*, 192(1):3–27, 2022.
- [5] V. Arya, N. Garg, R. Khandekar, A. Meyerson, K. Munagala, and V. Pandit. Local Search Heuristics for the  $k$ -median and Facility Location Problems. *Proc., ACM Symposium on Theory of Computing (STOC)*, 2001.
- [6] Pranjali Awasthi, Moses Charikar, Ravishankar Krishnaswamy, and Ali Kemal Sinop. The hardness of approximation of euclidean k-means. *CoRR*, abs/1502.03316, 2015.
- [7] Tanvi Bajpai, Deeparnab Chakrabarty, Chandra Chekuri, and Maryam Negahbani. Revisiting Priority k-Center: Fairness and Outliers. In *Proc., International Colloquium on Automata, Languages and Programming (ICALP)*, pages 21:1–21:20, 2021.
- [8] Boaz Barak, Pravesh K. Kothari, and David Steurer. Small-Set Expansion in Shortcode Graph and the 2-to-2 Conjecture. In *Proc., Innovations in Theoretical Computer Science (ITCS)*, 2018.
- [9] Jaroslaw Byrka, Fabrizio Grandoni, Thomas Rothvoss, and Laura Sanità. Steiner tree approximation via iterative randomized rounding. *Journal of the ACM*, 60(1):1–33, 2013.
- [10] Jaroslaw Byrka, Thomas Pensyl, Bartosz Rybicki, Aravind Srinivasan, and Khoa Trinh. An improved approximation for  $k$ -median and positive correlation in budgeted optimization. *ACM Trans. Algorithms*, 2017.
- [11] Deeparnab Chakrabarty, Chandra Chekuri, Sanjeev Khanna, and Nitish Korula. Approximability of capacitated network design. *Algorithmica*, 72(2):493–514, 2015.
- [12] Deeparnab Chakrabarty, Prachi Goyal, and Ravishankar Krishnaswamy. The non-uniform  $k$ -center problem. *ACM Trans. Algorithms*, 2020. Preliminary version in ICALP 2016.
- [13] Deeparnab Chakrabarty and Maryam Negahbani. Generalized center problems with outliers. *ACM Trans. Algorithms*, 2019.
- [14] Deeparnab Chakrabarty and Maryam Negahbani. Better algorithms for individually fair k-clustering. *Adv. in Neu. Inf. Proc. Sys. (NeurIPS)*, 2021.
- [15] Deeparnab Chakrabarty, Maryam Negahbani, and Ankita Sarkar. Approximation Algorithms for Continuous Clustering and Facility Location Problems. In Shiri Chechik, Gonzalo Navarro, Eva Rotenberg, and Grzegorz Herman, editors, *30th Annual European Symposium on Algorithms (ESA 2022)*, volume 244 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 33:1–33:15, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [16] Moses Charikar and Sudipto Guha. Improved Combinatorial Algorithms for the Facility Location and k-Median Problems. In *Proc., IEEE Symposium on Foundations of Computer Science (FOCS)*, 1999.
- [17] Moses Charikar, Sudipto Guha, Éva Tardos, and David B. Shmoys. A constant-factor approximation algorithm for the k-median problem. *Journal of Computer and System Sciences*, 65(1):129–149, 2002.

- [18] Moses Charikar, Samir Khuller, David M. Mount, and Giri Narasimhan. Algorithms for facility location problems with outliers. In *Proc., ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 642–651, 2001.
- [19] Ke Chen. On k-median clustering in high dimensions. In *Proc., ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1177–1185, 2006.
- [20] Vincent Cohen-Addad and Karthik C.S. Inapproximability of clustering in lp metrics. In *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 519–539, 2019.
- [21] Vincent Cohen-Addad, Hossein Esfandiari, Vahab Mirrokni, and Shyam Narayanan. Improved approximations for Euclidean k-means and k-median, via nested quasi-independent sets. In *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2022*, page 1621–1628, New York, NY, USA, 2022. Association for Computing Machinery.
- [22] Vincent Cohen-Addad, C. S. Karthik, and Euiwoong Lee. On approximability of clustering problems without candidate centers. In *Proc., ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2635–2648, 2021.
- [23] Vincent Cohen-Addad, Philip N Klein, and Claire Mathieu. Local search yields approximation schemes for k-means and k-median in euclidean and minor-free metrics. *SIAM Journal on Computing (SICOMP)*, 48(2):644–667, 2019.
- [24] Vincent Cohen-Addad, Karthik C. S., and Euiwoong Lee. Johnson coverage hypothesis: Inapproximability of k-means and k-median in  $\ell_p$ -metrics. In *Proceedings of the 2022 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1493–1530, 2022.
- [25] Vincent Cohen-Addad, David Saulpic, and Chris Schwiegelshohn. A new coresets framework for clustering. In *Proc., ACM Symposium on Theory of Computing (STOC)*, pages 169–182, 2021.
- [26] W Fernandez De La Vega, Marek Karpinski, Claire Kenyon, and Yuval Rabani. Approximation schemes for clustering problems. In *Proc., ACM Symposium on Theory of Computing (STOC)*, pages 50–58, 2003.
- [27] Hu Ding, Haikuo Yu, and Zixiu Wang. Greedy strategy works for k-center clustering with outliers and coresets construction. In *Proc., European Symposium on Algorithms*, pages 40:1–40:16, 2019.
- [28] Irit Dinur, Subhash Khot, Guy Kindler, Dor Minzer, and Muli Safra. On non-optimally expanding sets in grassmann graphs. In *Proc., ACM Symposium on Theory of Computing (STOC)*, 2018.
- [29] Irit Dinur, Subhash Khot, Guy Kindler, Dor Minzer, and Muli Safra. Towards a proof of the 2-to-1 games conjecture? In *Proc., ACM Symposium on Theory of Computing (STOC)*, 2018.
- [30] Dan Feldman, Morteza Monemizadeh, and Christian Sohler. A PTAS for k-means clustering based on weak coresets. In *Proceedings of the 5th Annual Symposium on Computational Geometry (SoCG)*, pages 11–18, 2007.
- [31] Zachary Friggstad, Mohsen Rezapour, and Mohammad R Salavatipour. Local search yields a PTAS for k-means in doubling metrics. *SIAM Journal on Computing (SICOMP)*, 48(2):452–480, 2019.
- [32] Martin Grötschel, László Lovász, and Alexander Schriver. *Geometric Algorithms and Combinatorial Optimization*. Springer, Berlin, Heidelberg, 1993.
- [33] Sudipto Guha and Samir Khuller. Greedy strikes back: Improved facility location algorithms. *Journal of Algorithms*, 31(1):228–248, 1999.
- [34] Anupam Gupta and Kanat Tangwongsan. Simpler analyses of local search algorithms for facility location. *arXiv preprint arXiv:0809.2554*, 2008.

- [35] S. Har-Peled and S. Mazumdar. Coresets for  $k$ -means and  $k$ -median clustering and their applications. In *Proc., ACM Symposium on Theory of Computing (STOC)*, pages 291–300, 2004.
- [36] D. Hochbaum and D. Shmoys. A best possible heuristic for the  $k$ -center problem. *Math. Oper. Res.*, 10:180–184, 1985.
- [37] Dorit S. Hochbaum and David B. Shmoys. A best possible heuristic for the  $k$ -center problem. *Math. Oper. Res.*, 10(2):180–184, 1985.
- [38] Kamal Jain, Mohammad Mahdian, Evangelos Markakis, Amin Saberi, and Vijay V Vazirani. Greedy facility location algorithms analyzed using dual fitting with factor-revealing LP. *Journal of the ACM*, 50(6):795–824, 2003.
- [39] Kamal Jain and Vijay V. Vazirani. Approximation algorithms for metric facility location and  $k$ -median problems using the primal-dual schema and Lagrangean relaxation. *Journal of the ACM*, 48(2):274–296, 2001.
- [40] Christopher Jung, Sampath Kannan, and Neil Lutz. Service in your neighborhood: Fairness in center location. In *Proceedings, Foundations of Responsible Computing, FORC 2020*, volume 156, pages 5:1–5:15, 2020.
- [41] Tapas Kanungo, David M Mount, Nathan S Netanyahu, Christine D Piatko, Ruth Silverman, and Angela Y Wu. A local search approximation algorithm for  $k$ -means clustering. *Computational Geometry*, 28(2-3):89–112, 2004.
- [42] Subhash Khot, Dor Minzer, and Muli Safra. On independent sets, 2-to-2 games, and grassmann graphs. *Proc., ACM Symposium on Theory of Computing (STOC)*, 2017.
- [43] A. Kumar, Y. Sabharwal, and S. Sen. A simple linear time  $(1 + \epsilon)$ -approximation algorithm for  $k$ -means clustering in any dimensions. In *Proc., IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 454–462, 2004.
- [44] Shi Li. A 1.488 approximation algorithm for the uncapacitated facility location problem. *Information and Computation*, 222:45–58, 2013.
- [45] Shi Li and Ola Svensson. Approximating  $k$ -median via pseudo-approximation. *SIAM Journal on Computing (SICOMP)*, 45(2):530–547, 2016.
- [46] Sepideh Mahabadi and Ali Vakilian. (individual) fairness for  $k$ -clustering. In *Proc., International Conference on Machine Learning (ICML)*, pages 7925–7935, 2020.
- [47] Jiri Matoušek. On approximate geometric  $k$ -clustering. *Discrete & Computational Geometry*, 24(1):61–84, 2000.
- [48] Ján Plesník. A heuristic for the  $p$ -center problems in graphs. *Discrete Applied Mathematics*, 17(3):263 – 268, 1987.
- [49] Robert D. Carr, Toshihiro Fujito, Goran Konjevod, and Ojas Parekh. A  $2\frac{1}{10}$ -approximation Algorithm for a Generalization of the Weighted Edge-Dominating Set Problem. *J. Comb. Optim.*, 2001. Preliminary version appeared in *Proc. 8th ESA*, pages 132–142, 2000.
- [50] David B. Shmoys, Éva Tardos, and Karen Aardal. Approximation algorithms for facility location problems (extended abstract). In *Proc., ACM Symposium on Theory of Computing (STOC)*, 1997.
- [51] Khot Subhash, Dor Minzer, and Muli Safra. Pseudorandom sets in Grassmann graph have near-perfect expansion. In *Proc., IEEE Symposium on Foundations of Computer Science (FOCS)*, 2018.
- [52] Luca Trevisan. When hamming meets euclid: The approximability of geometric tsp and steiner tree. *SIAM Journal on Computing*, 30(2):475–485, 2000.

- [53] Ali Vakilian and Mustafa Yalciner. Improved approximation algorithms for individually fair clustering. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera, editors, *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pages 8758–8779. PMLR, 28–30 Mar 2022.

## A Details of the ContFair- $k$ -Med results

### A.1 Filtering algorithm

---

**Algorithm 3** Filtering for ContFair- $k$ -Med

---

**Input:** A proposed solution  $(\{C_v\}_{v \in C}, \{y(v, r)\}_{v \in C, r \in \mathbb{R}})$  for LP

- 1:  $R(v) \leftarrow \min\{r(v), 2C_v\}$  for all  $v \in C$
- 2:  $\text{Reps}_{(C, y)} \leftarrow \emptyset$  ▷ “representative” clients
- 3:  $U \leftarrow C$  ▷ “uncovered” clients
- 4: **while**  $U \neq \emptyset$  **do**
- 5:     Pick  $j \in U$  with minimum  $R(j)$
- 6:      $\text{child}(j) \leftarrow \{v \in U \mid d(v, j) \leq 2R(v)\}$
- 7:      $U \leftarrow U \setminus \text{child}(j)$
- 8:      $\text{Reps}_{(C, y)} \leftarrow \text{Reps}_{(C, y)} \cup \{j\}$
- 9: **end while**

**Output:**  $\text{Reps}_{(C, y)}$

---

### A.2 Proof of Lemma 9

**Lemma 9.** Consider a solution  $(C, y)$  of LP. If for a client  $v$ ,  $(C, y)$  satisfies all constraints of the form (LP-2) and (LP-3) involving  $v$ , then for any  $r_0 \in \mathbb{R}$ ,  $C_v \geq r_0(1 - y(v, r_0))$ .

*Proof.*

$$\begin{aligned}
C_v &\geq \int_0^\infty (1 - y(v, r)) \, dr && \dots \text{(LP-2)} \\
&\geq \int_0^{r_0} (1 - y(v, r)) \, dr \\
&\geq (1 - y(v, r_0)) \int_0^{r_0} \, dr && \dots \text{by (LP-3), } y(v, r) \leq y(v, r_0) \\
&= r_0(1 - y(v, r_0))
\end{aligned}$$

□

### A.3 Proof of Lemma 11

**Lemma 11.**  $S \subseteq X$  be a solution to ContFair- $k$ -Med. Consider a proposed solution  $(C, y)$  of LP that satisfies (LP). Then  $\sum_{v \in C} d(v, S) \leq \sum_{j \in \text{Reps}_{(C, y)}} |\text{child}(j)| d(j, S) + 4\text{opt}_g$ .

*Proof.* Consider a client  $v \in C$ . In Algorithm 3, all clients are eventually covered, so there must exist  $j \in \text{Reps}_{(C, y)}$  such that  $v \in \text{child}(j)$ . Thus

$$d(v, S) \leq d(j, S) + d(v, j) \leq d(j, S) + 2R(v)$$

where the last inequality follows by Line 6. Summing this over all  $v \in C$ ,

$$\begin{aligned}
\sum_{v \in C} d(v, S) &\leq \sum_{j \in \text{Reps}_{(C, y)}} \sum_{v \in \text{child}(j)} (d(j, S) + 2R(v)) \\
&\leq \sum_{j \in \text{Reps}_{(C, y)}} |\text{child}(j)| d(j, S) + 4 \sum_{v \in C} C_v
\end{aligned}$$

where the last inequality follows because  $R(v) \leq 2C_v$ . The lemma follows by (LP).  $\square$

#### A.4 Proof of Lemma 13

**Lemma 13** ([17, 14]). *Let  $(\bar{x}, \bar{z})$  be a feasible solution of DLP with cost at most  $2\text{opt}_g$ , such that  $\forall j \in \text{Reps}$ ,  $\bar{z}_j \geq \frac{1}{2}$ . Then there exists a polynomial time algorithm that produces  $S \subseteq \text{Reps}$  such that (1)  $|S| = k$ ; (2) If  $\bar{z}_j = 1$ , then  $j \in S$ ; (3)  $\forall j \in \text{Reps}$ , at least one of  $j, s(j)$  is in  $S$ ; and (4)  $\sum_{j \in \text{Reps}} |\text{child}(j)| d(j, S) \leq 4\text{opt}_g$ .*

We first round  $(\bar{x}, \bar{z})$  to a  $\{\frac{1}{2}, 1\}$ -integral solution  $(\tilde{x}, \tilde{z})$  of the same cost. Then, we round  $(\tilde{x}, \tilde{z})$  to an integral solution  $(x, z)$  with twice the cost.

##### Rounding to a $\{\frac{1}{2}, 1\}$ -integral solution

We first exhibit this procedure for Lemma 13, and then modify it slightly for Lemma 24. We will assume that  $\sum_{j \in \text{Reps}} \bar{z}_j$  is an integer because, if it is not, then it is strictly less than  $k$ , and we can distribute the value  $k - \sum_{j \in \text{Reps}} \bar{z}_j$  among the  $\bar{z}$ -values of some arbitrary representatives, without increasing the quantity  $\sum_{i \in X} \bar{x}_{ij} d(j, i)$  for any  $j \in \text{Reps}$ . Hence

**Assumption 1.**  $\sum_{j \in \text{Reps}} \bar{z}_j \in \mathbb{Z}$ .

Given the assumption, we round to a  $\{\frac{1}{2}, 1\}$ -solution as follows: we start with  $\tilde{z}$  being  $\bar{z}$ , and then we find two representatives  $j_1, j_2$  that violate  $\{\frac{1}{2}, 1\}$ -integrality. We will pick the one with a lower value of  $|\text{child}(j)| d(j, s(j))$ ; say this is  $j_1$ . Then we will decrease  $\tilde{z}_{j_1}$  and increase  $\tilde{z}_{j_2}$  until one of them hits  $\frac{1}{2}$  or 1. We will repeat this until  $\tilde{z}$  becomes  $\{\frac{1}{2}, 1\}$ -integral. Alongside,  $\tilde{x}$  values are adjusted accordingly.

---

##### Algorithm 4 Rounding to a $\{\frac{1}{2}, 1\}$ -integral solution

---

**Input:** A solution  $(\{\bar{x}_{ij}\}_{i,j \in \text{Reps}}, \{\bar{z}_j\}_{j \in \text{Reps}})$  of DLP; other  $\tilde{x}, \tilde{z}$  values implicitly zero

```

1:  $(\tilde{x}, \tilde{z}) \leftarrow (\bar{x}, \bar{z})$ 
2: while  $\exists j_1, j_2 \in \text{Reps}, j_1 \neq j_2, \frac{1}{2} < \tilde{z}_{j_1}, \tilde{z}_{j_2} < 1$  do
3:                                      $\triangleright$  wlog assume  $|\text{child}(j_1)| d(j_1, s(j_1)) \leq |\text{child}(j_2)| d(j_2, s(j_2))$ 
4:      $\delta(j_1, j_2) \leftarrow \min \{ \tilde{z}_{j_2} - \frac{1}{2}, 1 - \tilde{z}_{j_1} \}$ 
5:      $\tilde{z}_{j_1} \leftarrow \tilde{z}_{j_1} - \delta(j_1, j_2)$ 
6:      $\tilde{z}_{j_2} \leftarrow \tilde{z}_{j_2} + \delta(j_1, j_2)$ 
7:     for  $j \in \{j_1, j_2\}$  do
8:          $\tilde{x}_{jj} \leftarrow \tilde{z}_j$ 
9:          $\tilde{x}_{s(j)j} \leftarrow \tilde{1} - \tilde{z}_j$ 
10:    end for
11: end while
Output:  $(\tilde{x}, \tilde{z})$ 

```

---

**Lemma 16** ([14, 17]). *Let  $(\bar{x}, \bar{z})$  be a feasible solution of DLP with cost at most  $2\text{opt}_g$ , such that  $\forall j \in \text{Reps}, \bar{z}_j \geq \frac{1}{2}$ . Then  $(\tilde{x}, \tilde{z})$  is a feasible  $\{\frac{1}{2}, 1\}$ -integral solution of DLP of cost at most  $2\text{opt}_g$ , such that  $\forall j \in \text{Reps}, \bar{z}_j = 1 \implies \tilde{z}_j = 1$ .*

*Proof.* Observe that Algorithm 4 never alters  $\tilde{z}$ -values that are 0 or 1 in Line 1. Moreover, it never lowers a  $\tilde{z}$ -value that is at least  $\frac{1}{2}$  to a value strictly below  $\frac{1}{2}$ . So the property

$$\forall j \in \text{Reps}, \tilde{z}_j \geq \frac{1}{2}$$

as initially ensured, is preserved throughout. Thus, our construction never violates (DLP-3); and, by Assumption 1, we can find  $j_1, j_2$  as in Line 2 iff  $\tilde{z}$  is not  $\{\frac{1}{2}, 1\}$ -integral.

Now consider the quantity

$$\psi(\tilde{x}, \tilde{z}) := \sum_{j \in \text{Reps}} \max \left\{ \tilde{z}_j - \frac{1}{2}, 1 - \tilde{z}_j \right\}$$

Observe that the while-loop in Algorithm 4 strictly decreases  $\psi(\tilde{x}, \tilde{z})$  in every iteration, but never makes it negative. Also,  $\psi(\tilde{x}, \tilde{z}) = 0$  iff  $(\tilde{x}, \tilde{z})$  is  $\{\frac{1}{2}, 1\}$ -integral. So the while-loop terminates and yields a  $\{\frac{1}{2}, 1\}$ -integral solution. Note that every iteration of the while-loop preserves  $\sum_{j \in \text{Reps}} \tilde{z}_j$ . So we preserve (DLP-1). We satisfy (DLP-2) by construction.

Finally, for the cost bound:

$$\sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in X} \tilde{x}_{ij} d(j, i) = \sum_{j \in \text{Reps}} \tilde{x}_{s(j)j} d(j, s(j))$$

In one iteration of the while-loop,  $\tilde{x}_{s(j_1)j_1}$  decreases by  $\delta(j_1, j_2)$ , while  $\tilde{x}_{s(j_2)j_2}$  increases by the same. So the RHS above, in one iteration, increases by

$$\delta(j_1, j_2) (|\text{child}(j_1)| d(j_1, s(j_1)) - |\text{child}(j_2)| d(j_2, s(j_2)))$$

which is designed to be non-positive in Line 2. Thus by Line 1 and Lemma 12 we have

$$\sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in X} \tilde{x}_{ij} d(j, i) \leq \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in X} \bar{x}_{ij} d(j, i) \leq 2 \text{opt}_g$$

□

### Rounding to an integral solution

Given  $(\tilde{x}, \tilde{z})$  as constructed above, we first open all  $j \in \text{Reps}$  that have  $\tilde{z}_j = 1$ . We discard these representatives, and also any representatives that now have  $\tilde{z}_{s(j)} = 1$ . On the remaining  $\text{Reps}$ , we construct a forest, as follows: we draw directed edges of the form  $(j, s(j))$ , and arbitrarily discard any antiparallel edges that occur when  $s(s(j)) = j$ . Then we open alternate levels of every tree in this forest, choosing either odd or even levels to get the lower cost.



---

**Algorithm 5** Rounding to an integral solution
 

---

**Input:** A  $\{\frac{1}{2}, 1\}$ -integral solution  $(\{\tilde{x}_{ij}\}_{i,j \in \text{Reps}}, \{\tilde{z}_j\}_{j \in \text{Reps}})$  of DLP, other  $\tilde{x}, \tilde{z}$  implicitly zero

```

1:  $(x, z) \leftarrow$  all zeros
2:  $V \leftarrow \text{Reps}$  ▷ vertex set
3: for  $j \in \text{Reps}$  such that  $\tilde{z}_j = 1$  do
4:    $z_j \leftarrow 1, x_{jj} \leftarrow 1$ 
5:    $V \leftarrow V \setminus j$ 
6: end for
7: for  $j \in \text{Reps}$  such that  $z_j = 1$  do
8:    $x_{s(j)j} \leftarrow 1$ 
9:    $V \leftarrow V \setminus j$ 
10: end for
11:  $E \leftarrow \{(j, s(j)) \mid j \in V\}$  ▷ edge set, has antiparallel edges
12: for Each connected component  $T$  in  $(V, E)$  do
13:   Pick  $j \in T$  such that  $s(s(j)) = j$  ▷ unique, corresponds to closest pair in  $T$ 
14:    $E \leftarrow E \setminus (s(j), j)$  ▷ removing antiparallel edges
15:    $\text{root}(T) \leftarrow s(j)$  ▷ now all edges point upwards in the tree, and levels well-defined
16:    $T_{\text{odd}} \leftarrow$  odd level vertices in  $T$ 
17:    $T_{\text{even}} \leftarrow$  even level vertices in  $T$ 
18:   if  $\sum_{j \in T} d(j, T_{\text{odd}}) \leq \sum_{j \in T} d(j, T_{\text{even}})$  then
19:      $\forall j \in T_{\text{odd}}, z_j \leftarrow 1, x_{jj} \leftarrow 1$ 
20:      $\forall j \in T_{\text{even}}, x_{s(j)j} \leftarrow 1$ 
21:   else
22:      $\forall j \in T_{\text{even}}, z_j \leftarrow 1, x_{jj} \leftarrow 1$ 
23:      $\forall j \in T_{\text{odd}}, x_{s(j)j} \leftarrow 1$ 
24:   end if
25: end for
Output:  $(x, z)$ 

```

---

**Lemma 17** ([17]). *Let  $(\tilde{x}, \tilde{z})$  be a feasible  $\{\frac{1}{2}, 1\}$ -integral solution of DLP with cost at most  $2\text{opt}_g$ . Then  $(x, z)$  is a feasible integral solution of DLP with cost at most  $4\text{opt}_g$ , such that  $\forall j \in \text{Reps}, \tilde{z}_j = 1 \implies z_j = 1$ ; and either  $z_j = 1$  or  $z_{s(j)} = 1$ .*

*Proof.* By construction,  $(x, z)$  is integral, and satisfies (DLP-3) and (DLP-2). The construction also ensures that for each  $j \in \text{Reps}$ , at least one of  $z_j, z_{s(j)}$  is set to 1.

Now we show that  $(x, z)$  satisfies (DLP-1). Let  $\text{Reps}_{\frac{1}{2}} = \{j \in \text{Reps} \mid \tilde{z}_j = \frac{1}{2}\}$ . Algorithm 5 sets all  $z$  values in  $\text{Reps} \setminus \text{Reps}_{\frac{1}{2}}$  to 1, and sets at most  $\frac{\lceil \text{Reps}_{\frac{1}{2}} \rceil}{2}$  of the  $z$  values in  $\text{Reps}_{\frac{1}{2}}$  to 1. By Assumption 1, and since Algorithm 4 maintains  $\sum_{i \in X} \tilde{z}_i = \sum_{i \in X} \bar{z}_i$ , we have that  $|\text{Reps}_{\frac{1}{2}}|$  is even, so  $\left\lceil \frac{|\text{Reps}_{\frac{1}{2}}|}{2} \right\rceil = \frac{|\text{Reps}_{\frac{1}{2}}|}{2}$ . Thus

$$\sum_{i \in X} z_i = |\text{Reps} \setminus \text{Reps}_{\frac{1}{2}}| + \frac{|\text{Reps}_{\frac{1}{2}}|}{2} = \sum_{j \in \text{Reps}} \tilde{z}_j$$

which is at most  $k$  since  $(\tilde{x}, \tilde{z})$  is feasible.

Finally, for the cost bound, it suffices to show the following: for each  $j \in \text{Reps}$ ,

$$\sum_{i \in X} x_{ij} d(j, i) \leq 2 \sum_{i \in X} \tilde{x}_{ij} d(j, i).$$

If  $z_j = \tilde{z}_j = 1$ , then LHS = RHS = 0 above. Otherwise  $\tilde{z}_j = \frac{1}{2} = \tilde{x}_{s(j)j}$ . So the RHS is exactly  $2 \cdot \frac{1}{2} \cdot d(j, s(j))$ , and since either  $z_j = 1$  or  $z_{s(j)} = 1$ , the LHS is at most  $d(j, s(j))$ .  $\square$

Setting  $S := \{j \in \text{Reps} \mid z_j = 1\}$  yields Lemma 13.

## B 32-approximation for Cont- $k$ -Means

**Definition 18** (Continuous  $k$ -means (Cont- $k$ -Means)). *The input is a metric space  $(X, d)$ , clients  $C \subseteq X, |C| = n \in \mathbb{N}$ , and  $k \in \mathbb{N}$ . The goal is to find  $S \subseteq X, |S| = k$  minimizing  $\text{cost}(S) := \sum_{v \in C} d(v, S)^2$ .*

**Theorem 19.** *There exists a polynomial time algorithm that, for an instance of Cont- $k$ -Means with optimum  $\text{opt}$ , yields a solution of cost at most  $32\text{opt} + \varepsilon$ . Here  $\varepsilon = O\left(\frac{1}{n^2}\right)$ .*

To prove this, we proceed with a round-or-cut framework similar to the one in Section 4 using a linear program similar to LP (except the fairness constraints). Here the variables are  $\{C_v^2\}_{v \in C}$  for the cost shares and  $\{y(v, r)\}_{v \in C, r \in \mathbb{R}}$  for the facility masses. The constraints involving only the  $y(v, r)$ 's remain the same as in LP, while those that involve  $C_v$ 's are now adapted to the squared costs.

$$\sum_{v \in C} C_v^2 \leq \text{opt}_g \quad (\text{LP}^2)$$

$$\sum_{B \in \mathcal{B}} y(B) \leq k \quad \forall \mathcal{B} \subseteq \{B(v, r)\}_{v \in C, r \in \mathbb{R}} \text{ pairwise disjoint} \quad (\text{LP}^2-1)$$

$$2 \int_0^\infty (1 - y(v, r)) r dr \leq C_v^2 \quad \forall v \in C \quad (\text{LP}^2-2)$$

$$\begin{aligned} y(u, r) &\leq y(v, r') & \forall u, v \in C, r, r' \in \mathbb{R}, B(u, r) \subseteq B(v, r') & (\text{LP}^2-3) \\ y(v, r) &\geq 0, C_v^2 \geq 0 & \forall v \in C, r \in \mathbb{R} \end{aligned}$$

The rounding algorithm and its analysis are analogous to our results in Section 4. We first prove the following analogue of Lemma 9 for  $\text{LP}^2$ .

**Lemma 20.** *Consider a solution  $(C^2, y)$  of  $\text{LP}^2$ . If for a client  $v$ ,  $(C^2, y)$  satisfies all constraints of the form (LP<sup>2</sup>-2) and (LP<sup>2</sup>-3) involving  $v$ , then for any  $r_0 \in \mathbb{R}$ ,  $C_v^2 \geq r_0^2 (1 - y(v, r_0))$ .*

*Proof.*

$$\begin{aligned} C_v^2 &\geq 2 \int_0^\infty (1 - y(v, r)) r dr && \dots (\text{LP}^2-2) \\ &\geq 2 \int_0^{r_0} (1 - y(v, r)) r dr \\ &\geq (1 - y(v, r_0)) \cdot 2 \int_0^{r_0} r dr && \dots \text{by (LP}^2\text{-3), } y(v, r) \leq y(v, r_0) \\ &= r_0^2 (1 - y(v, r_0)) \end{aligned}$$

□

The filtering procedure is the same as Algorithm 3, except  $R(v)$  now equals  $\sqrt{2}C_v$ . Also, for clarity, we call the output  $\text{Reps}_{(C^2, y)}$ . For a  $j \in \text{Reps}_{(C^2, y)}$ ,  $s(j)$  and  $a(j)$  are defined as before, and we use the same separating constraint, which we now call  $\text{Sep}_{(C^2, y)}$ .

$$\sum_{j \in \text{Reps}_{(C^2, y)}} y(j, a(j)) \leq k \quad (\text{Sep}_{(C^2, y)})$$

As before, we will focus on satisfying  $\text{Sep}_{(C^2, y)}$  and argue that this is sufficient for our rounding algorithm. We prove an analogue of Lemma 10.

**Lemma 21.** *If  $(C^2, y)$  satisfies (LP<sup>2</sup>-2)-(LP<sup>2</sup>-3) and  $\text{Sep}_{(C^2, y)}$ , then*

1.  $\forall j \in \text{Reps}_{(C^2, y)}, y(j, a(j)) \geq \frac{1}{2}$
2.  $|\text{Reps}_{(C^2, y)}| \leq 2k$

*Proof.* Fix  $j \in \text{Reps}_{(C^2, y)}$ . By construction of child sets, we have

$$a(j) = \frac{d(j, s(j))}{2} \geq R(v) = \sqrt{2}C_j \quad (6)$$

By Lemma 20,

$$C_j^2 \geq a(j)^2 (1 - y(j, a(j)))$$

If  $C_j = 0$ , since  $a(j) > 0$  by definition, this implies  $y(j, a(j)) \geq 1$ . Otherwise, substituting  $a(j)$  by  $R(v) = \sqrt{2}C_j$  from (6) gives

$$C_j^2 \geq 2C_j^2 (1 - y(j, a(j))) \implies y(j, a(j)) \geq \frac{1}{2}.$$

Now, by  $\text{Sep}_{(C^2, y)}$ ,

$$k \geq \sum_{j \in \text{Reps}_{(C^2, y)}} y(j, a(j)) \geq \frac{1}{2} \cdot |\text{Reps}_{(C^2, y)}|$$

□

So if we find that  $|\text{Reps}_{(C^2, y)}| > 2k$ , we can pass  $\text{Sep}_{(C^2, y)}$  to ellipsoid as a separating hyperplane. Hence for our rounding algorithm, we can now use a  $(C^2, y)$  that satisfies  $\text{Sep}_{(C^2, y)}$ . We fix such a  $(C^2, y)$ , and hereafter refer and  $\text{Reps}_{(C^2, y)}$  as simply  $\text{Reps}$ . We now prove an analog of Lemma 11, which allows us to ignore clients outside of  $\text{Reps}$  hereafter.

**Lemma 22.**  *$S \subseteq X$  be a solution to Cont- $k$ -Means. Consider a proposed solution  $(C^2, y)$  of  $\text{LP}^2$  that satisfies  $(\text{LP}^2)$ . Then*

$$\sum_{v \in C} d(v, S)^2 \leq 2 \sum_{j \in \text{Reps}_{(C, y)}} |\text{child}(j)| d(j, S)^2 + 16\text{opt}_g$$

*Proof.* Consider a client  $v \in C$ . In the filtering algorithm, all clients are eventually covered, so there must exist  $j \in \text{Reps}_{(C^2, y)}$  such that  $v \in \text{child}(j)$ . Thus

$$d(v, S) \leq d(j, S) + d(v, j) \leq d(j, S) + 2R(v) = d(j, S) + 2\sqrt{2}C_v$$

where the last inequality follows by Line 6. Squaring, we get

$$d(v, S)^2 \leq (d(j, S) + 2\sqrt{2}C_v)^2 \leq 2(d(j, S)^2 + (2\sqrt{2}C_v)^2) = 2(d(j, S)^2 + 8C_v^2)$$

Summing this over all  $v \in C$ ,

$$\begin{aligned} \sum_{v \in C} d(v, S)^2 &\leq 2 \sum_{j \in \text{Reps}_{(C, y)}} \sum_{v \in \text{child}(j)} (d(j, S)^2 + 8C_v^2) \\ &\leq 2 \sum_{j \in \text{Reps}_{(C, y)}} |\text{child}(j)| d(j, S)^2 + 16 \sum_{v \in C} C_v^2 \end{aligned}$$

The lemma follows by  $(\text{LP}^2)$ . □

Next, following our algorithm for ContFair- $k$ -Med closely, we will round  $(C^2, y)$  to an integral solution of  $(\text{DLP}^2)$ , which has the same constraints as DLP, but the objective function changes to suit the  $k$ -means problem:

$$\text{minimize } \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in \text{Reps}} x_{ij} d(j, i)^2 \quad (\text{DLP}^2)$$

This is achieved via the same consolidation procedure as in Algorithm 2. We prove an analogue of Lemma 12.

**Lemma 23.**  $(\bar{x}, \bar{z})$  is a feasible solution of  $\text{DLP}^2$  with cost at most  $4\text{opt}_g$  and,  $\forall j \in \text{Reps}$ ,  $\bar{z}_j \geq \frac{1}{2}$ .

*Proof.* For a  $j \in \text{Reps}$ , if  $y(j, a(j)) = 1$  then  $\bar{z}_j = 1$ . Otherwise, by Lemma 21.1,  $\bar{z}_j = y(j, a(j)) \geq \frac{1}{2}$ . Hence  $1 - \bar{z}_j \leq \frac{1}{2} \leq \bar{z}_{s(j)}$ , which implies feasibility by construction and Sep. We also have

$$\sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i)^2 = (1 - \bar{z}_j) d(j, s(j))^2$$

If  $y(j, a(j)) = 1$ , then the RHS above is  $0 \leq 4C_j^2$ . Otherwise

$$\sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i)^2 = (1 - y(j, a(j))) d(j, s(j))^2 = 4(1 - y(j, a(j))) a(j)^2 \leq 4C_j^2$$

where the last inequality follows from Lemma 20. Multiplying by  $|\text{child}(j)|$  and summing over all  $j \in \text{Reps}$ , we have by construction of  $\text{Reps}$ ,

$$\begin{aligned} \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in \text{Reps}} \bar{x}_{ij} d(j, i)^2 &\leq 4 \sum_{j \in \text{Reps}} |\text{child}(j)| C_j^2 \leq 4 \sum_{j \in \text{Reps}} \sum_{v \in \text{child}(j)} C_v^2 \\ &= 4 \sum_{v \in C} C_v^2 \leq 4\text{opt}_g \end{aligned}$$

□

Finally, we prove an analog of Lemma 13.

**Lemma 24** (Modified from [17, 14]). *Let  $(\bar{x}, \bar{z})$  be a feasible solution of  $\text{DLP}^2$  with cost at most  $4\text{opt}_g$ , such that  $\forall j \in \text{Reps}$ ,  $\bar{z}_j \geq \frac{1}{2}$ . Then there exists a polynomial time algorithm that produces  $S \subseteq \text{Reps}$  such that*

1.  $|S| = k$
2. If  $\bar{z}_j = 1$ , then  $j \in S$ .
3.  $\forall j \in \text{Reps}$ , at least one of  $j, s(j)$  is in  $S$
4.  $\sum_{j \in \text{Reps}} |\text{child}(j)| d(j, S)^2 \leq 8\text{opt}_g$

*Proof.* The proof is similar to that of Lemma 13. We first apply Algorithm 4, except we now take  $j_1, j_2$  such that  $|\text{child}(j_1)| d(j_1, s(j_1))^2 \leq |\text{child}(j_2)| d(j_2, s(j_2))^2$ . This, via similar reasoning as the proof of Lemma 16, gives us a  $\{\frac{1}{2}, 1\}$ -integral solution  $(\tilde{x}, \tilde{z})$  of  $\text{DLP}^2$  such that  $\forall j \in \text{Reps}$ ,  $\tilde{z}_j = 1 \implies \bar{z}_j = 1$ , and

$$\sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in X} \tilde{x}_{ij} d(j, i)^2 \leq \sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in X} \bar{x}_{ij} d(j, i)^2 \leq 4\text{opt}_g$$

We then apply Algorithm 4. Similarly to the proof of Lemma 17, we get an integral solution  $(x, z)$  satisfying the necessary  $\text{DLP}^2$  constraints. We also have that  $\forall j \in \text{Reps}$ ,  $\tilde{z}_j = 1 \implies z_j = 1$  and either  $z_j = 1$  or  $z_{s(j)} = 1$ . So we need to show that

$$\sum_{i \in X} x_{ij} d(j, i)^2 \leq 2 \sum_{i \in X} d(j, i)^2$$

If  $z_j = \tilde{z}_j = 1$ , then both the LHS and the RHS are 0. Otherwise  $\tilde{z}_j = \frac{1}{2} = \tilde{x}_{s(j)j}$ . So the RHS is exactly  $2 \cdot \frac{1}{2} \cdot d(j, s(j))^2$ , and the LHS is at most  $d(j, s(j))^2$ . □

So by Lemma 22, we have, for the  $S$  constructed above,

$$\sum_{v \in C} d(v, S)^2 \leq 2 \cdot 8\text{opt}_g + 16\text{opt}_g = 32\text{opt}_g$$

## Generalizing further to Continuous $(k, p)$ -clustering

Our approach extends to the generalization where the objective function is the sum of some  $p^{\text{th}}$  power of the distances.

**Definition 25** (Continuous  $(k, p)$ -clustering (Cont- $(k, p)$ -Clustering)). *The input is a metric space  $(X, d)$ , clients  $C \subseteq X, |C| = n \in \mathbb{N}$ , and  $k, p \in \mathbb{N}$ . The goal is to find  $S \subseteq X, |S| = k$  minimizing  $\text{cost}(S) := \sum_{v \in C} d(v, S)^p$ .*

**Theorem 26.** *There exists a polynomial time algorithm that, for an instance of Cont- $(k, p)$ -Clustering with optimum  $\text{opt}$ , yields a solution of cost at most  $2^{2p+1}\text{opt} + \varepsilon$ . Here  $\varepsilon = O\left(\frac{1}{n^2}\right)$ .*

This follows by the same framework as for Cont- $k$ -Med and Cont- $k$ -Means, with the following alterations:

**LP formulation.** The cost-share variables are  $\{C_v^p\}_{v \in C}$ . (LP<sup>2</sup>) is replaced by  $\sum_{v \in C} C_v^p \leq \text{opt}_g$ , and (LP<sup>2</sup>-2) is replaced by  $p \int_0^1 (1 - y(v, r)) r^{p-1} dr \leq C_v^p$ .

**Filtering.**  $R(v) := 2^{1/p} C_v$ .

**Linear program on Reprs.** The objective function in (DLP<sup>2</sup>) is replaced by

$$\sum_{j \in \text{Reps}} |\text{child}(j)| \sum_{i \in \text{Reps}} x_{ij} d(j, i)^p.$$

**Rounding to  $\{\frac{1}{2}, 1\}$ -integral solution.** In Algorithm 4, we use  $j_1, j_2$  such that

$$|\text{child}(j_1)| d(j_1, s(j_1))^p \leq |\text{child}(j_2)| d(j_2, s(j_2))^p.$$

The guarantees are obtained using the relaxed triangle inequality  $(a + b)^p \leq 2^{p-1}(a^p + b^p)$ . Via the filtering step, we get

$$\sum_{v \in C} d(v, S)^2 \leq 2^{p-1} \sum_{j \in S} |\text{child}(j)| d(j, S) + 2^{2p} \text{opt}_g$$

and the consolidation step incurs a factor of  $2^p$ . The remaining algorithm incurs a factor of 2 irrespective of the value of  $p$ . So we get an approximation factor of  $2^{p-1} \cdot 2^p \cdot 2 + 2^{2p} = 2^{2p+1}$ .

## C Continuous $k$ -center with Outliers

In the  $k$ -center problem, the input is the same as that for  $k$ -median, but the objective is now the *maximum* distance that a client has to the open facilities, rather than the sum. Furthermore, we consider a version here that allows *outliers*, i.e. only some  $m$  clients need to be considered when computing the cost.

**Definition 27** (Continuous  $k$ -center with Outliers (Cont- $k$ CwO)). *The input is a metric space  $(X, d)$ , clients  $C \subseteq X, |C| = n \in \mathbb{N}$ , and  $k, m \in \mathbb{N}$ . The goal is to find  $S \subseteq X, |S| = k$  and  $D \subseteq C, |D| \geq m$  minimizing  $\max_{v \in D} d(v, S)$ .*

In this section, we provide an algorithm for Cont- $k$ CwO that matches the state of the art for discrete  $k$ -CwO up to a small additive factor. The best approximation algorithm for the discrete version is a 2-approximation primal-rounding approach [12]. Later, Chakrabarty and Negahbani [13] extended this result to allow for generalized constraints on facilities, using a round-or-cut framework. Both of those results are tight due to a hardness of approximation by Hochbaum and Shmoys [36]. Via our idea of creating  $y$ -variables for balls around clients, we adapt the discrete results [14, 12] to the Cont- $k$ CwO problem.

**Theorem 28.** *There exists a polynomial time algorithm that, for an instance of Cont- $k$ CwO with optimum  $\text{opt}$ , yields a solution of cost at most  $2\text{opt} + \varepsilon$ . Here  $\varepsilon = O\left(\frac{1}{n^2}\right)$ .*

As before, we proceed with a round-or-cut framework with a guessed optimum  $\text{opt}_g$ , and variables  $y(v, r) = y(B)$  representing the number of open facilities in each ball  $B = B(v, r)$ . We will also have variables  $\text{cov}_v$  for each  $v \in C$  representing whether or not  $v$  has a facility open within distance  $\text{opt}_g$  of it. Our linear program  $k\text{-CwO}$  is as follows:

$$\begin{aligned} \sum_{B \in \mathcal{B}} y(B) &\leq k & \forall \mathcal{B} = \{B(v, r)\}_{\substack{v \in C \\ r \in \mathbb{R}}} \text{ pairwise disjoint} & (k\text{-CwO-1}) \\ y(v, \text{opt}_g) &\geq \text{cov}_v & \forall v \in C & (k\text{-CwO-2}) \\ y(v, r) &\leq y(v, r') & \forall v \in C, r, r' \in \mathbb{R} & (k\text{CwO-3}) \\ \sum_{v \in C} \text{cov}_v &\geq m & & (k\text{CwO-4}) \end{aligned}$$

All constraints except for  $(k\text{-CwO-1})$  are polynomially many, and we use ellipsoid for  $(k\text{-CwO-1})$ . Given a proposed solution  $(\text{cov}, y)$  of  $k\text{-CwO}$ , we construct  $\text{Reps}_{\text{cov}} \subseteq C$  and  $S_{\text{cov}} \subseteq \text{Reps}_{\text{cov}}$  as follows. We first produce  $\text{Reps}_{\text{cov}}$  such that all clients are within  $2\text{opt}_g$  of it, but it is potentially of size larger than  $k$ . Then, we pick  $S_{\text{cov}} \subseteq \text{Reps}_{\text{cov}}, |S_{\text{cov}}| = k$  by choosing the  $k$  largest child sets.

Constructing  $\text{Reps}_{\text{cov}}$  is simple: initially all clients are uncovered ( $U = C$ ), and we pick the uncovered client  $i \in U$  with maximum  $\text{cov}_i$ . We put  $i$  in  $\text{Reps}_{\text{cov}}$ , and consider all uncovered clients within distance  $2\text{opt}_g$  of  $i$  to be in  $\text{child}(i)$ , which are then covered. We repeat this until all clients are covered.

---

**Algorithm 6** Rounding for Cont- $k\text{CwO}$

---

**Input:**  $\{\text{cov}_v\}_{v \in C}, \text{opt}_g$

- 1:  $\text{Reps}_{\text{cov}} \leftarrow \emptyset$
- 2:  $U \leftarrow C$
- 3: **while**  $U \neq \emptyset$  **do**
- 4: Pick  $i \in U$  with maximum  $\text{cov}_i$
- 5:  $\text{Reps}_{\text{cov}} \leftarrow \text{Reps}_{\text{cov}} + i$
- 6:  $\text{child}(i) \leftarrow U \cap B(i, 2\text{opt}_g)$
- 7:  $U \leftarrow U \setminus B(i, 2\text{opt}_g)$
- 8: **end while**
- 9:  $S_{\text{cov}} \leftarrow \emptyset$
- 10: **while**  $|S_{\text{cov}}| < k$  **do**  $\triangleright$  construct  $S_{\text{cov}}$  to be the  $k$  points that cover the most
- 11: Pick  $i \in \text{Reps}_{\text{cov}} \setminus S_{\text{cov}}$  with maximum  $|\text{child}(i)|$
- 12:  $S_{\text{cov}} \leftarrow S_{\text{cov}} + i$
- 13: **end while**

**Output:**  $S_{\text{cov}}$

---

Observe that, by construction of  $\text{Reps}_{\text{cov}}$  in Algorithm 6, the collection of balls  $\mathcal{B}_{\text{cov}} := \{B(i, \text{opt}_g)\}_{i \in \text{Reps}_{\text{cov}}}$  is pairwise disjoint. So the following constraint, called  $\text{Sep}_{\text{cov}}$  is of the form  $(k\text{-CwO-1})$ :

$$\sum_{i \in \text{Reps}_{\text{cov}}} y(i, \text{opt}_g) \leq k \quad (\text{Sep}_{\text{cov}})$$

We will show that

**Lemma 29.** *If  $(\text{cov}, y)$  satisfies  $(k\text{-CwO-2})$ - $(k\text{CwO-4})$  and  $\text{Sep}_{\text{cov}}$ , then*

$$\sum_{i \in S_{\text{cov}}} |\text{child}(i)| \geq m$$

So if we get  $\sum_{i \in S_{\text{cov}}} |\text{child}(i)| < m$ , we know that  $\text{Sep}_{\text{cov}}$  must be violated, so we return it to ellipsoid as a separating hyperplane. Hence, in polynomial time, we can either conclude that our guess of  $\text{opt}_g$  should be larger, or obtain  $S_{\text{cov}}$  such that  $\sum_{i \in S_{\text{cov}}} |\text{child}(i)| \geq m$ . Since  $\text{child}(i)$  sets are pairwise disjoint by construction, putting  $D = \sqcup_{i \in S_{\text{cov}}} \text{child}(i)$  yields the 2-approximation.

*Proof of Lemma 29.* For a fixed  $\text{opt}_g$  and  $(\text{cov}, y)$ , let  $\text{Reps} = \text{Reps}_{\text{cov}}, S = S_{\text{cov}}$ . By  $(k\text{-CwO-2})$  and  $\text{Sep}_{\text{cov}}$ ,

$$\sum_{i \in \text{Reps}} \text{cov}_i \leq \sum_{i \in \text{Reps}} y(i, \text{opt}_g) \leq k \quad (7)$$

Also, by Algorithm 6 and  $(k\text{CwO-4})$ ,

$$\sum_{i \in \text{Reps}} |\text{child}(i)| \text{cov}_i \geq \sum_{i \in \text{Reps}} \sum_{v \in \text{child}(i)} \text{cov}_v = \sum_{v \in C} \text{cov}_v \geq m \quad \dots \text{ by } (k\text{CwO-4}) \quad (8)$$

By (7) and (8), we can view  $\sum_{i \in \text{Reps}} |\text{child}(i)| \cdot \frac{\text{cov}_i}{k}$  as an average of quantities  $|\text{child}(i)|$  with weights  $\frac{\text{cov}_i}{k}$ . Since this average is at least  $m/k$ , the  $k$  largest  $|\text{child}(i)|$ 's must sum up to at least  $m$ , as desired.  $\square$

## D Hardness of approximation for Cont- $\lambda$ -UFL

**Theorem 7.** *Given an instance of Cont- $\lambda$ -UFL and  $\varepsilon > 0$ , it is NP-hard to distinguish between the following:*

- *There exists  $S \subseteq X$  such that  $\text{cost}_F(S) + \text{cost}_C(S) \leq (1 + 6\varepsilon)n$*
- *For any  $S \subseteq X$ ,  $\text{cost}_F(S) + \text{cost}_C(S) \geq (2 - \varepsilon)n$*

We will reduce from the following result, that was used [22] to show a hardness of  $(2 - \varepsilon)$  for Cont- $k$ -Med.

**Theorem 30** ([42, 28, 29, 8, 51]). *Given a graph  $G = (V, E)$  and an  $\varepsilon > 0$ , it is NP-hard to distinguish between the following:*

- *There exists pairwise disjoint independent sets  $V_1, V_2, V_3, V_4 \subseteq V$  such that  $\forall i \in [4], |V_i| \geq \frac{1-\varepsilon}{4} |V|$ .*
- *There is no independent set in  $G$  of size  $\varepsilon |V|$ .*

*Proof of Theorem 7.* Given a graph  $G = (V, E)$  and  $\varepsilon > 0$ , we construct the following instance of the Cont- $\lambda$ -UFL problem.

For  $m := |E|$ , let  $X = \mathbb{R}^m$ , and let  $d$  be the distance induced in  $\mathbb{R}^m$  by the  $\ell_\infty$ -norm. Fix an arbitrary orientation on  $E$ , and let the clients be  $C := \{A(v) \mid v \in V\}$  where, for a  $v \in V$  and  $e = (u', v') \in E$ ,

$$A(v)(e) := \begin{cases} 2 & \text{if } v = u' \\ -2 & \text{if } v = v' \\ 0 & \text{otherwise} \end{cases}$$

So we can hereafter say  $n = |V| = |C|$ . Now let  $\lambda = \varepsilon n$ . Also, fix an arbitrary optimal solution of this Cont- $\lambda$ -UFL instance that opens  $r$  facilities for some  $r \geq 1$ , and set  $\varepsilon' = \frac{\varepsilon}{r}$ .

The proof follows by Lemma 31, which analyzes the two directions of this reduction.  $\square$

**Lemma 31.** *Consider a Cont- $\lambda$ -UFL instance constructed from the graph  $G = (V, E)$ , as per the construction described earlier. Suppose the optimum in this instance opens  $r$  facilities, and let  $\varepsilon' = \varepsilon/r$ .*

*If there are pairwise disjoint independent sets  $V_1, V_2, V_3, V_4 \subseteq V$ , each of size at least  $\frac{1-\varepsilon'}{4}n$ , then there is a solution in the Cont- $\lambda$ -UFL instance with cost at most  $(1 + 6\varepsilon)n$ . Conversely, if there is no independent set of size  $\varepsilon'n$  in  $G$ , then the optimum in the Cont- $\lambda$ -UFL instance is at least  $(2 - \varepsilon)n$ .*

*Proof. Large independent sets imply low-cost UFL:* Given pairwise disjoint independent sets  $V_1, V_2, V_3, V_4 \subseteq V$ , each of size  $\frac{1-\varepsilon'}{4}n \geq \frac{1-\varepsilon}{4}n$  in  $G$ , we can produce a solution in the Cont- $\lambda$ -UFL instance that has cost at most  $(1 + 6\varepsilon)n$ . To construct this solution, open the facilities  $s_1, s_2, s_3, s_4$ , defined as follows. For  $i \in [4]$  and each coordinate  $e = (u', v') \in E$ ,

$$s_i(e) := \begin{cases} 1 & \text{if } u' \in V_i \\ -1 & \text{if } v' \in V_i \\ 0 & \text{otherwise} \end{cases}$$

Since coordinates in any  $A(v)$  are in  $\{\pm 2, 0\}$  and coordinates in each  $s_i$  are in  $\{\pm 1, 0\}$ , we get the simple upper bound of  $d(A(v), s_i) \leq 3$  for any  $i \in [4], v \in V$ .

For some  $i \in [4], v \in V_i$  and  $e = (u', v') \in E$ , we study the quantity  $|A(v)(e) - s_i(e)|$ . If  $u' = v$ , then it is  $|2 - 1| = 1$ . If  $v' = v$ , then it is  $|-2 - (-1)| = 1$ . If  $u' \in V_i - v$ , then it is  $|0 - 1| = 1$ . If  $v' \in V_i - v$ , then it is  $|0 - (-1)| = 1$ . When  $e$  lies outside  $V_i$ , then it is zero. So when  $v \in V_i$ , we get a tighter bound of  $d(A(v), s_i) \leq 1$ . Thus the cost of connecting clients to facilities is at most

$$\sum_{i=1}^4 |V_i| + 3 \left| \left( V \setminus \bigcup_{i=1}^4 V_i \right) \right| \leq 4 \cdot \frac{1-\varepsilon}{4} n + 3\varepsilon n \leq (1 + 2\varepsilon)n$$

We open 4 facilities, so the facility opening cost is  $4\lambda = 4\varepsilon n$ . Thus the total cost of this solution is  $(1 + 6\varepsilon)n$ , as promised.

**Small maximum independent set implies high-cost UFL:** Suppose there is no independent set in  $G$  of size  $\varepsilon'n$ . In the optimum solution of **Cont- $\lambda$ -UFL** that we had fixed, let the facilities opened be  $S = \{s_1, \dots, s_r\}$  and, for  $i \in [r]$ , let the set  $C_i$  be the clients assigned to  $s_i$  in this solution in other words,  $d(v, S) = d(v, s_i)$  for all  $i \in [r], v \in C_i$ . Also let  $V_i := \{v \in V \mid A(v) \in C_i\}$ . We have the following claim (Claim 1, [22]):

**Claim 32.** *If  $G$  has no independent set of size  $\varepsilon'n$ , then the induced subgraph of  $G$  on  $V_i$  contains a matching  $M_i$  of size at least  $\frac{|V_i| - \varepsilon'n}{2}$ .*

We prove the claim at the end of this section. Given the claim, we have that

$$\begin{aligned} \sum_{v \in V_i} d(A(v), s_i) &\geq \sum_{(u,v) \in M_i} (d(A(v), s_i) + d(A(u), s_i)) \\ &\geq \sum_{(u,v) \in M_i} d(A(u), A(v)) \\ &= \sum_{(u,v) \in M_i} |A(u)((u,v)) - A(v)((u,v))| \\ &= 4|M_i| \\ &\geq 2(|V_i| - \varepsilon'n) \end{aligned}$$

and thus  $\text{cost}_C(S)$  is

$$\sum_{j \in C} d(j, S) = \sum_{i=1}^r \sum_{v \in V_i} d(A(v), s_i) \geq 2n - 2\varepsilon'rn = 2n - 2\varepsilon n$$

Since  $r$  facilities are open,  $\text{cost}_F(S)$  is  $\lambda r = r\varepsilon n \geq \varepsilon n$ . Thus  $\text{cost}_F(S) + \text{cost}_C(S)$  is at least  $n(2 - 2\varepsilon + \varepsilon) \geq n(2 - \varepsilon)$ .  $\square$

*Proof of Claim 32 [22].* If  $|V_i| \geq \varepsilon'n$ , then it cannot be independent. So pick an edge in the subgraph induced by  $V_i$ , include it in the matching  $M_i$ , and discard its endpoints from  $V_i$ . Repeat this process until  $|V_i| < \varepsilon'n$  to obtain the desired  $M_i$ .  $\square$