

DIAMETER ESTIMATION OF (m, ρ) -QUASI EINSTEIN MANIFOLDS

ABSOS ALI SHAIKH^{1*}, PROSENJIT MANDAL², CHANDAN KUMAR MONDAL³

ABSTRACT. This paper aims to study the (m, ρ) -quasi Einstein manifold. This article shows that a complete and connected Riemannian manifold under certain conditions becomes compact. Also, we have determined an upper bound of the diameter for such a manifold. It is also exhibited that the potential function acquiesces to the Hodge-de Rham potential up to a real constant in an (m, ρ) -quasi Einstein manifold. Later, some triviality and integral conditions are established for a non-compact complete (m, ρ) -quasi Einstein manifold having finite volume. Finally, it is proved that with some certain constraints, a complete Riemannian manifold admits finite fundamental group. Furthermore, some conditions for compactness criteria have also been deduced.

1. INTRODUCTION AND PRELIMINARIES

A non-gradient generalized m -quasi-Einstein manifold ([3]) is an $n(> 2)$ -dimensional Riemannian manifold (N, g) such that it preserves a smooth function $\alpha \in C^\infty(N)$, with

$$(1) \quad \alpha g + \frac{1}{m} W^b \otimes W^b = \frac{1}{2} \mathcal{L}_W g + Ric,$$

where $W \in \chi(N)$, it's dual 1-form W^b , $\mathcal{L}_W g$ indicates the Lie derivative of g along W and m is a scalar with $0 < m \leq \infty$. In (1), if we replace α by $\lambda + \rho R$, for some real constants λ, ρ and the scalar curvature R of N , then (N, g) is called a non-gradient (m, ρ) -quasi Einstein manifold ([16]), and in this case equation (1) reduces to

$$(2) \quad \lambda g + \rho R g + \frac{1}{m} W^b \otimes W^b = \frac{1}{2} \mathcal{L}_W g + Ric.$$

If W is the gradient of a real valued smooth function f on N , then the (m, ρ) -quasi Einstein manifold is said to be a (gradient) (m, ρ) -quasi Einstein manifold (for details see [7, 10]), with

* Corresponding author.

2020 Mathematics Subject Classification: 53C20; 53C21; 53C25; 53E20.

Key words and phrases: Riemannian manifold; (m, ρ) -Quasi Einstein Manifold; scalar curvature; index form; diameter estimation.

potential function f . In this case, (2) reduces to

$$(3) \quad \lambda g + \rho Rg + \frac{1}{m} df \otimes df = \nabla^2 f + Ric.$$

If $m = \infty$ and $\rho = 0$ (resp., $m = \infty$), then (3) compresses to the Ricci soliton equation (see e.g. [6, 8, 9, 14, 18, 20]) (resp., ρ -Einstein soliton equation (see e.g. [17, 19])). In (3), if m and λ is taken from $\chi(N)$, then (N, g) is known as the generalized m -quasi Einstein manifold [5]. Furthermore, tracing (3), we obtain

$$(4) \quad \lambda n + n\rho R + \frac{1}{m} |\nabla f|^2 = \Delta f + R.$$

For a fixed $\phi \in C^\infty(N)$, the weighted Laplacian is defined by (see, [12])

$$\Delta_\phi f = \Delta f - \langle \nabla \phi, \nabla f \rangle.$$

Catino [5] introduced the notion of a generalized quasi-Einstein manifold and characterized such type of manifolds with harmonic Weyl tensor. For a compact m -quasi Einstein metric having constant scalar curvature, a triviality result is proved by Case et al. [4]. They also determined that all compact 2-dimensional m -quasi Einstein manifolds are trivial. For a compact gradient ρ -Einstein soliton with some conditions, Shaikh et al. [19] have provided a lower bound of the diameter. Huang and Wei [10] have deduced some rigidity results on a compact (m, ρ) -quasi Einstein manifolds, in particular, by using some conditions on scalar curvature or on the constant ρ , they have showed that such a manifold is trivial. Demirbağ and Güler [7] have characterized an (m, ρ) -quasi Einstein manifold admitting closed conformal or parallel vector field. Also, they have established some rigidity results giving some new examples of the (m, ρ) -quasi Einstein manifold.

López and Rio [13] have proved a compactness theorem for a complete Riemannian manifold satisfying $\mathcal{L}_W g + Ric \geq \lambda g$, with potential vector field W having bounded norm. Wylie [22] has proved that the fundamental group of a complete Riemannian manifold satisfying $\mathcal{L}_W g + Ric \geq \lambda g$, for $\lambda > 0$, is finite.

Hence inspiring by the above studies and the study of [2], in this article, we have showed the following:

In the first theorem we have generalized the work of Limoncu [11], and obtained an upper bound of the diameter.

Theorem 1.1. *Let (N, g) be a complete and connected Riemannian manifold satisfying $Ric + \nabla^2 f - \frac{1}{m}df \otimes df \geq \lambda g + \rho Rg$. If $|f| \leq K$ and $\rho R \geq K_1$, for some real constants K, K_1 choosing in such a way that $(\lambda + K_1) > 0$, then N is compact and the diameter satisfies*

$$(5) \quad diam(N) \leq \pi\sqrt{2}\sqrt{\frac{(n-1) + K\sqrt{2}}{\lambda + K_1}}.$$

Next, we have generalized the work of Li and Wenyi [12], and obtained the following:

Theorem 1.2. *Assume that $(N, g, e^{-f}dvol)$ is a smooth metric measure space satisfying $Ric + \nabla^2 f - \frac{1}{m}df \otimes df \geq \lambda g + \rho Rg$, and u is a positive smooth f -harmonic function on N , then*

$$(6) \quad \frac{1}{2}\Delta F \geq (\lambda + \rho R)F + \frac{1}{n}F^2 - \frac{2}{n}|\nabla f|F^{\frac{3}{2}} - |\nabla F|F^{\frac{1}{2}} + \frac{1}{m}\{df(\nabla\omega)\}^2 + \frac{1}{2}\langle\nabla f, \nabla F\rangle,$$

where we denote $F = |\nabla \log u|^2$.

Theorem 1.3. *If (N, g) is a compact and oriented (m, ρ) -quasi Einstein manifold, then its potential function is given by $f = \sigma + C_1$, where C_1 is a constant and σ is the Hodge-de Rham potential .*

Let (N, g) be an oriented Riemannian manifold and $A^k(N)$ be the collection of all k -th differential forms in N . Now for given integer $k \geq 0$, the global inner product in $A^k(N)$ is defined by (see, [15])

$$\langle \zeta, \omega \rangle = \int_N \zeta \wedge * \omega,$$

for $\zeta, \omega \in A^k(N)$. Here “ $*$ ” denotes the Hodge star operator. The global norm of $\zeta \in A^k(N)$ is given by $\|\zeta\|^2 = \langle \zeta, \zeta \rangle$ with $\|\zeta\|^2 \leq \infty$.

Theorem 1.4. *Let (N, g) be a complete (m, ρ) -quasi Einstein manifold which is non-compact and of finite volume. If $\rho > \frac{1}{n}$ and also W admits finite global norm, then the following holds:*

- (i) *If $R \geq \frac{\lambda n}{1-\rho n}$, then N is trivial,*
- (ii) *If $(\lambda + \rho)R \geq 0$, then $\frac{1}{m} \int_N |W|^2 dV \leq \int_N R dV$, and*
- (iii) *$\int_N R dV \leq \frac{\lambda n}{1-\rho n} Vol(N)$, where $Vol(N)$ represents the volume of N .*

By virtue of Theorem 1.4, we deduce the following corollary:

Corollary 1.4.1. *Let (N, g) be a complete (m, ρ) -quasi Einstein manifold which is non-compact and of finite volume. If $\rho < \frac{1}{n}$ and W is of finite global norm, then the following holds:*

- (i) *If $R \leq \frac{\lambda n}{1-\rho n}$, then N is trivial,*
- (ii) *If $(\lambda + \rho)R \leq 0$, then $\int_N R dV \leq \frac{1}{m} \int_N |X|^2 dV$, and*
- (iii) *$\int_N R dV \geq \frac{\lambda n}{1-\rho n} \text{Vol}(N)$.*

Next, we are interested in studying a complete Riemannian manifold (N, g) with a vector field W such that,

$$(7) \quad \lambda g + \rho Rg + \frac{1}{m} W^b \otimes W^b \leq \frac{1}{2} \mathcal{L}_W g + Ric,$$

In (7) the equality gives the fundamental equation of an (m, ρ) -quasi Einstein manifold. Here the main aim is to prove the following results:

Theorem 1.5. *If (N, g) is a complete Riemannian manifold satisfying (7) along with $\int_0^r \rho R = K_3$, for some constant K_3 , then the fundamental group of N is finite.*

Theorem 1.6. *Let (N, g) be a complete Riemannian manifold satisfying (7) with $\|W\|$ bounded and $\int_0^r \rho R = K_3$, for some constant K_3 with $(\lambda + K_3) > 0$. Then N becomes compact.*

2. PROOF OF THE RESULTS

Proof of Theorem 1.1. Let $p, q \in N$ and let Γ be a minimizing unit speed geodesic segment from p to q of length d . Considering a parallel orthonormal frame $\{\xi_1 = \Gamma', \xi_2, \dots, \xi_n\}$ along Γ and a smooth function $h \in C^\infty([0, d])$ such that $h(0) = h(d) = 0$, we get (see, [11]),

$$(8) \quad \sum_{i=2}^n \eta(h\xi_i, h\xi_i) = \int_0^d \{(n-1)h'^2 - h^2 Ric(\Gamma', \Gamma')\} dt,$$

where η denotes the index form of Γ .

With our assumption the relation (8), yields

$$(9) \quad \begin{aligned} \sum_{i=2}^n \eta(h\xi_i, h\xi_i) &\leq \int_0^d \left[(n-1)h'^2 + h^2 \left\{ \nabla^2 f - \frac{1}{m} df \otimes df - \lambda g - \rho Rg \right\}(\Gamma', \Gamma') \right] dt \\ &= \int_0^d \left[(n-1)h'^2 + h^2 \nabla^2 f(\Gamma', \Gamma') - \frac{h^2}{m} (df(\Gamma'))^2 - h^2(\lambda + \rho R) \right] dt. \end{aligned}$$

Again, since $|f| \leq K$, we obtain (see, [11]),

$$\begin{aligned}
\int_0^d h^2 \nabla^2 f(\Gamma', \Gamma') dt &= \int_0^d h^2 g(\nabla_{\Gamma'} \nabla f, \Gamma') dt \\
&= \int_0^d h^2 \Gamma'(g(\nabla f, \Gamma')) dt \\
(10) \qquad \qquad \qquad &\leq 2K\sqrt{d} \left(\int_0^d \left(\frac{d}{dt}(hh') \right)^2 dt \right)^{\frac{1}{2}}.
\end{aligned}$$

The equations (9) and (10) together implies

$$\begin{aligned}
\sum_{i=2}^n \eta(h\xi_i, h\xi_i) &\leq \int_0^d (n-1)h'^2 dt + 2K\sqrt{d} \left(\int_0^d \left(\frac{d}{dt}(hh') \right)^2 dt \right)^{\frac{1}{2}} \\
(11) \qquad \qquad \qquad &- \int_0^d \frac{h^2}{m} (df(\Gamma'))^2 dt - \int_0^d h^2(\lambda + \rho R) dt.
\end{aligned}$$

If we take $h(t) = \sin(\frac{\pi}{d}t)$, then (11), yields

$$\begin{aligned}
\sum_{i=2}^n \eta(h\xi_i, h\xi_i) &\leq (n-1) \int_0^d \frac{\pi^2}{d^2} \cos^2 \left(\frac{\pi}{d}t \right) dt + 2K\sqrt{d} \left(\int_0^d \left(\frac{\pi^2}{d^2} \cos \left(\frac{2\pi}{d}t \right) \right)^2 dt \right)^{\frac{1}{2}} \\
&\quad - \lambda \int_0^d \sin^2 \left(\frac{\pi}{d}t \right) dt - \int_0^d \sin^2 \left(\frac{\pi}{d}t \right) \rho R dt \\
&\leq (n-1) \frac{\pi^2}{2d} + \frac{K\pi^2\sqrt{2}}{d} - \frac{\lambda}{2}d - \frac{K_1}{2}d.
\end{aligned}$$

Since Γ is a minimizing geodesic, hence $\eta(h\xi_i, h\xi_i) \geq 0$. Consequently,

$$0 \leq (n-1) \frac{\pi^2}{2d} + \frac{K\pi^2\sqrt{2}}{d} - \frac{\lambda}{2}d - \frac{K_1}{2}d.$$

The above relation entails

$$d \leq \pi\sqrt{2} \sqrt{\frac{(n-1) + K\sqrt{2}}{\lambda + K_1}}.$$

This completes the proof. □

Proof of Theorem 1.2. Set $\omega = \log u$, then $F = |\nabla\omega|^2$ and ω satisfies the equation:

$$\Delta_f \omega + |\nabla\omega|^2 = 0.$$

Thus,

$$\Delta_f \omega = -F.$$

Therefore,

$$(12) \quad |\nabla^2 \omega|^2 \geq \frac{1}{n} |\Delta \omega|^2 = \frac{1}{n} |F - \langle \nabla f, \nabla \omega \rangle|^2 \geq \frac{1}{n} F^2 - \frac{2}{n} \langle \nabla f, \nabla \omega \rangle F,$$

and

$$(13) \quad \langle \nabla \omega, \nabla \Delta_f \omega \rangle = -\langle \nabla \omega, \nabla F \rangle.$$

Now, the weighted Bochner formula:

$$\frac{1}{2} \Delta_f |\nabla \omega|^2 = |\nabla^2 \omega|^2 + \langle \nabla \omega, \nabla \Delta_f \omega \rangle + Ric_f(\nabla \omega, \nabla \omega),$$

together with (12), (13) and our assumption, yields

$$\frac{1}{2} \Delta_f F \geq \frac{1}{n} F^2 - \frac{2}{n} \langle \nabla f, \nabla \omega \rangle F - \langle \nabla \omega, \nabla F \rangle + (\lambda + \rho R) |\nabla \omega|^2 + \frac{1}{m} df \otimes df(\nabla \omega, \nabla \omega).$$

This implies

$$(14) \quad \frac{1}{2} \Delta F - \langle \nabla f, \nabla F \rangle \geq \frac{1}{n} F^2 - \frac{2}{n} \langle \nabla f, \nabla \omega \rangle F - \langle \nabla \omega, \nabla F \rangle + (\lambda + \rho R) F + \frac{1}{m} \{df(\nabla \omega)\}^2.$$

Applying the Cauchy-Schwarz inequality, we obtain

$$\langle \nabla \omega, \nabla f \rangle \leq |\nabla \omega| |\nabla f| = |\nabla f| F^{\frac{1}{2}}.$$

Using the last inequality in (14), we get the estimation. \square

Proof of Theorem 1.3. If $W \in \chi(N)$, then as a consequence of Hodge-de Rham decomposition theorem, (see e.g. [21]), W can be written as

$$(15) \quad W - \nabla \sigma = Y,$$

where $\sigma \in \chi(N)$ is the Hodge-de Rham potential and $div Y = 0$. By considering an (m, ρ) -quasi Einstein manifold (N, g) so that the equation (2) holds, then we obtain

$$(16) \quad R + div W = \lambda n + \frac{1}{m} |W|^2 + \rho R n.$$

Therefore (15) concludes that $div W = \Delta \sigma$ and hence (16) entails

$$(17) \quad \lambda n + \frac{1}{m} |W|^2 + \rho R n = R + \Delta \sigma.$$

Again (3) indicates that

$$(18) \quad \lambda n + \frac{1}{m} |\nabla f|^2 + \rho R n = R + \Delta f.$$

From (17) and (18), we have

$$\Delta(f - \sigma) = 0,$$

which follows the result. \square

Proof of Theorem 1.4. We have, for $l > 0$

$$\begin{aligned} \frac{1}{l} \int_{\mathcal{B}(p, 2l)} |W| dV &\leq \left(\int_{\mathcal{B}(p, 2l)} \left(\frac{1}{l}\right)^2 dV \right)^{1/2} \left(\int_{\mathcal{B}(p, 2l)} \langle W, W \rangle dV \right)^{1/2} \\ &\leq \frac{1}{l} \left(Vol(N) \right)^{1/2} \|W\|_{\mathcal{B}(p, 2l)}, \end{aligned}$$

where $\mathcal{B}(p, l)$ is the open ball of radius l and center at p . Thus

$$\liminf_{l \rightarrow \infty} \frac{1}{l} \int_{\mathcal{B}(p, 2l)} |W| dV = 0.$$

Again, a Lipschitz continuous function ω_l exists, (see [23]) such that for some real constant $K_2 > 0$,

$$0 \leq \omega_l(x) \leq 1 \quad \forall x \in N,$$

$$\text{supp } \omega_l \subset \mathcal{B}(p, 2l),$$

$$|d\omega_l| \leq \frac{K_2}{l} \quad \text{almost everywhere on } N \text{ and}$$

$$\omega_l(x) = 1 \quad \forall x \in \mathcal{B}(p, l).$$

As $\lim_{l \rightarrow \infty} \omega_l = 1$, using the function ω_l , we get

$$\frac{C}{l} \int_{\mathcal{B}(p, 2l)} |W| dV \geq \left| \int_{\mathcal{B}(p, 2l)} \omega_l \text{div} W dV \right|.$$

From the defining relation of (m, ρ) -quasi Einstein manifold, we obtain

$$(19) \quad \int_N \left\{ \lambda n + \frac{1}{m} |W|^2 + (\rho n - 1) R \right\} dV = 0.$$

By virtue of equation (19) and our assumption, we get the desired results. \square

To prove Theorem 1.5, we need the following results:

Lemma 2.1 ([22]). *Let (N, g) be a complete Riemannian manifold and $p, q \in N$ with $r = d(p, q) > 1$. If Γ is a minimal geodesic joining from p to q and parametrized by the arc length s , then*

$$(20) \quad 2(n-1) + \mathcal{E}_p + \mathcal{E}_q \geq \int_0^r Ric(\Gamma'(s), \Gamma'(s)) ds,$$

where $\mathcal{E}_q = \max\{0, \sup\{Ric_y(v, v) : y \in B(q, 1), \|v\| = 1\}\}$, for $q \in N$.

Lemma 2.2. *If (N, g) is a complete Riemannian manifold satisfying (7), then for any $p, q \in N$,*

$$(21) \quad \max\left\{1, \frac{1}{K_3 + \lambda} (2(n-1) + \mathcal{E}_p + \mathcal{E}_q + \|W_p\| + \|W_q\|)\right\} \geq d(p, q).$$

Proof. Suppose that $d(p, q) > 1$ and Γ is a minimal geodesic joining from p to q and parametrized by arc length s . Then along Γ , we can easily get

$$\mathcal{L}_W g(\Gamma', \Gamma') = 2 \frac{d}{ds} [g(W, \Gamma')].$$

Applying Lemma 2.1, we obtain

$$(22) \quad 2(n-1) + \mathcal{E}_p + \mathcal{E}_q \geq \int_0^r Ric(\Gamma'(s), \Gamma'(s)) ds.$$

Using the Cauchy-Schwarz inequality and (7), we have

$$\begin{aligned} \int_0^r Ric(\Gamma'(s), \Gamma'(s)) ds &\geq g_p(W, \Gamma'(0)) - g_q(W, \Gamma'(r)) + \int_0^r (W^\flat(\Gamma'(s)))^2 ds + \lambda d(p, q) + \int_0^r \rho R ds \\ &\geq \lambda d(p, q) - \|W_p\| - \|W_q\| + K_3 d(p, q). \end{aligned}$$

In view of (22) and solving for $d(p, q)$, the last inequality entails (21). \square

Proof of Theorem 1.5. We omit the proof of Theorem 1.5, as it is straightforward from the Lemma 2.2 and following the technique of [22, Theorem 1.1]. \square

Proof of Theorem 1.6. Let $p \in N$ and $\Gamma : [0, \infty) \rightarrow N$ be any geodesic starting from p and parametrized by the arc length s . Hence (7) and the Cauchy-Schwarz inequality together imply that

$$\begin{aligned} \int_0^r Ric(\Gamma'(s), \Gamma'(s)) ds &\geq \int_0^r (W^\flat(\Gamma'(s)))^2 + \lambda r + \int_0^r \rho R g + 2g(W_p, \Gamma'(0)) - 2g(W_{\Gamma(r)}, \Gamma'(r)) ds \\ &\geq 2g(W_p, \Gamma'(0)) - 2\|W_{\Gamma(r)}\| + \lambda r + K_3 r. \end{aligned}$$

The boundedness of $\|W\|$ provides

$$\int_0^{+\infty} Ric(\Gamma'(s), \Gamma'(s)) = +\infty,$$

and hence Ambrose's compactness criteria [1] concludes that N is compact. \square

ACKNOWLEDGMENT

The second author gratefully acknowledges to the CSIR(File No.:09/025(0282)/2019-EMR-I), Govt. of India for the award of JRF. Also the third author conveys sincere thanks to the Netaji Subhas Open University for partial financial assistance (Project No.: AC/140/2021-22).

REFERENCES

- [1] Ambrose, W., *A theorem of Myers*, Duke Math. J., **24** (1957), 345–348.
- [2] Aquino, C., Barros, A. and Ribeiro, E. Jr., *Some applications of the Hodge-de Rham decomposition to Ricci solitons*, Results. Math., **60** (2011), 245–254.
- [3] Barros, A. A. and Gomes, J. N., *Triviality of compact m -quasi-Einstein manifolds*, Results Math., **71(1-2)** (2016), 241–250.
- [4] Case, J., Shu, Y. J. and Wei, G., *Rigidity of quasi-Einstein metrics*, Diff. Geom. Appl., **29(1)** (2011), 93–100.
- [5] Catino, G., *Generalized quasi Einstein manifolds with harmonic Weyl tensor*, Math. Z., **271** (2012), 751–756.
- [6] Chow, B. and Knopf, D., *The Ricci flow: an introduction*, Amer. Math. Soc., 2004.
- [7] Demirbağ, S. A. and Güler, S., *Rigidity of (m, ρ) -quasi Einstein manifolds*, Math. Nach., **290** (2017), 2100–2110.
- [8] Deshmukh, S. and Al-Sodais, H., *A note on almost Ricci solitons*, Anal. Math. Phys., **10(4)** (2020), 75–86.
- [9] Hamilton, R. S., *Three-manifolds with positive Ricci curvature*, J. Diff. Geom., **17** (1982), 255–306.
- [10] Huang, G. and Wei, Y., *The classification of (m, ρ) -quasi-Einstein manifolds*, Ann. Glob. Anal. Geom., **44** (2013), 269–282.
- [11] Limoncu, M., *The Bakry-Émery Ricci tensor and its applications to some compactness theorems*, Math. Z., **271** (2012), 715–722.
- [12] Li, C. and Wenyi, C., *Gradient estimates for positive smooth f -harmonic functions*, Acta Math. Scientia, **30(5)** (2010), 1614–1628.
- [13] López, M. F. and Rio, E. G., *A remarks on compact Ricci solitons*, Math. Ann., **340** (2008), 893–896.

- [14] Mondal, C. K. and Shaikh, A. A., *On Ricci solitons whose potential is convex*, Proc. Indian Acad. Sci. (Math. Sci.), **130(55)** (2020), 1–7.
- [15] Morita, S., *Geometry of differential forms*, Amer. Math. Soc., **2001**.
- [16] Rovenski, V. and Patra, D. S., *On non-gradient (m, ρ) -quasi-Einstein contact metric manifolds*, J. Geom., **112(14)** (2021), 1–17.
- [17] Shaikh, A. A., Cunha, A. W. and Mandal, P., *Some characterizations of ρ -Einstein solitons*, J. Geom. Phys., **166** (2021), 104270.
- [18] Shaikh, A. A., Mondal, C. K. and Mandal, P., *Compact gradient ρ -Einstein soliton is isometric to the Euclidean sphere*, Indian J. Pure Appl. Math., **52** (2021), 335–339.
- [19] Shaikh, A. A., Mandal, P. and Mondal, C. K., *Diameter estimation of gradient ρ -Einstein soliton*, J. Geom. Phys., **177** (2022), 104518.
- [20] Sharma, R. and Deshmukh, S., *Ricci almost solitons with associated projective vector field*, Adv. Geom., **22(1)** (2022), 1–8.
- [21] Warner, F., *Foundations of differentiable manifolds and Lie groups*, Springer-Verlag, New York, 1983.
- [22] Wylie, W., *Complete shrinking Ricci solitons have finite fundamental group*, Proc. Amer. Math. Soc., **136** (2008), 1803–1806.
- [23] Yau, S. T., *Some function-theoretic properties of complete Riemannian manifold and their applications to geometry*, Indiana Univ. Math. J., **25(7)** (1976), 659–670.

^{1,2}DEPARTMENT OF MATHEMATICS,
THE UNIVERSITY OF BURDWAN, GOLAPBAG,
PURBA BARDHAMAN-713101,
WEST BENGAL, INDIA

³SCHOOL OF SCIENCES,
NETAJI SUBHAS OPEN UNIVERSITY,
DURGAPUR REGIONAL CENTER, DURGAPUR-713214
PASCHIM BARDHAMAN,
WEST BENGAL, INDIA

Email address: ¹aask2003@yahoo.co.in, aashaikh@math.buruniv.ac.in

Email address: ²prosenjitmandal235@gmail.com

Email address: ³chan.alge@gmail.com, chandanmondal@wbnsou.ac.in