

ON REGULARITY AND PROJECTIVE DIMENSION UP TO SYMMETRY

DINH VAN LE AND HOP D. NGUYEN

ABSTRACT. Ideals in infinite dimensional polynomial rings that are invariant under the action of a certain monoid have been extensively studied recently. Of particular interest is the asymptotic behavior of truncations of such an ideal in finite dimensional polynomial subrings. It has been conjectured that the Castelnuovo–Mumford regularity and projective dimension are eventual linear functions along such truncations. In the present paper we provide evidence for these conjectures. We show that for monomial ideals the projective dimension is eventually linear, while the regularity is asymptotically linear.

1. INTRODUCTION

Ideals in infinite dimensional polynomial rings that are invariant under the action of the infinite symmetric group Sym or the monoid of increasing functions Inc have recently been intensively studied, with motivations from group theory [4], algebraic statistics [10], and representation theory [3]; see [6] for a nice survey. A fascinating research direction in this theory is to explore whether existing results related to ideals in Noetherian polynomial rings can be extended to the infinite dimensional case. Examples of successful extensions include equivariant Hilbert’s basis theorem [1, 4, 5, 10, 22], equivariant Hilbert-Serre theorem [12, 20, 21], equivariant Buchberger algorithm [5, 9], and equivariant Hochster’s formula [19]. See, e.g., also [7, 23, 26, 27] for related results. It should be noted that these results have inspired further work in combinatorics [15] and polyhedral geometry [11, 16].

One challenging open problem in the mentioned research direction is to extend the classical Hilbert’s syzygy theorem to the equivariant setting. More specifically, let $R = K[x_i \mid i \in \mathbb{N}]$ be a polynomial ring in infinitely many variables over a field K and let $I \subseteq R$ be a Π -invariant ideal (with $\Pi = \text{Sym}$ or $\Pi = \text{Inc}$) that is equivariantly finitely generated, i.e., I is generated by finitely many Π -orbits. It is of great interest to understand the minimal free resolution of I over R , and in particular, to understand how the finiteness property of I is reflected in that resolution.

An approach to this problem is as follows. One considers a Π -invariant chain $\mathcal{I} = (I_n)_{n \geq 1}$ with I_n an ideal in the Noetherian polynomial ring $R_n = K[x_1, \dots, x_n]$ such that $I = \bigcup_{n \geq 1} I_n$ (e.g., one can take the *saturated* chain $\overline{\mathcal{I}} = (\overline{I}_n)_{n \geq 1}$ with $\overline{I}_n = I \cap R_n$ for $n \geq 1$). The ideal I may be regarded as a *global* object and the chain \mathcal{I} a *local* one. By studying free resolutions of the local ideals I_n one has an opportunity to understand a free resolution of the global ideal I . This approach is undertaken in [22], where the chain \mathcal{I} is viewed as an FI- or OI-module and a projective resolution of this module is obtained with interesting finiteness properties (see [22, Theorems 7.1 and 7.7]).

2020 *Mathematics Subject Classification.* 13A50, 13C15, 13D02, 13F20, 16P70, 16W22.

Key words and phrases. Invariant ideal, monoid, polynomial ring, symmetric group.

Numerical aspects of free resolutions of the local ideals I_n , namely their Castelnuovo–Mumford regularity and projective dimension, are further studied in [13, 14], where the following expectations are proposed:

Conjecture 1.1. *Let $\Pi = \text{Sym}$ or $\Pi = \text{Inc}$ and let $\mathcal{I} = (I_n)_{n \geq 1}$ be a Π -invariant chain of proper homogeneous ideals. Then the following hold:*

- (i) $\text{pd}I_n$ is eventually a linear function in n .
- (ii) $\text{reg}I_n$ is eventually a linear function in n .

In fact, this conjecture is stated in [13, 14] for more general polynomial rings. See also [20] for a further generalization of the conjecture to the context of FI- and OI-modules.

Conjecture 1.1 is verified for Sym-invariant chains of monomial ideals by Murai [18]. Raicu [25] independently also proves Conjecture 1.1(ii) for such chains. Recently, Murai’s results have been refined in his joint work with Raicu on equivariant Hochster’s formula [19].

In the present paper we study Conjecture 1.1 in the case that \mathcal{I} is an Inc-invariant chain of monomial ideals. Our main result reads as follows:

Theorem 1.2. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals. Then the following hold:*

- (i) $\text{pd}I_n$ is eventually a linear function in n , provided that \mathcal{I} is a saturated chain.
- (ii) $\lim_{n \rightarrow \infty} \frac{\text{reg}I_n}{n}$ exists.

The first part of this theorem generalizes Murai’s result on Sym-invariant chains [18, Corollary 3.7], and our proof improves a key step in his argument (see Lemma 3.2).

For the second part of Theorem 1.2 we actually show that $\lim_{n \rightarrow \infty} (\text{reg}I_n/n)$ has the expected value that is predicted by [13, Conjecture 4.12] (see Theorem 4.2). This is done by using an induction on the so-called q -invariant introduced in [21]. There are two new ingredients in our proof. First, we introduce a certain chain of ideals to reduce the q -invariant (see Definition 4.5). This chain is somewhat similar to the one employed in [13, 14, 21], but has an additional property that it is Inc-invariant and thus suitable for our inductive argument. Second, we define a function λ that records the minimal last exponent of a generator of a monomial ideal (see Section 2.3). This function can be extended to Inc-invariant chains and will be exploited to give a lower bound on regularity along those chains. Such a bound is essential for a base case in the inductive argument (see Lemma 4.12).

Most positive results so far on Conjecture 1.1 deal with saturated chains. It is therefore worth mentioning that the constructions employed in the proof of Theorem 1.2(ii) are useful for studying non-saturated chains. Note that for those chains of ideals, the initial behavior of the functions $\text{reg}I_n$ and $\text{pd}I_n$ can be rather wild, even after the point where the chains stabilize (see Example 3.4).

Let us now describe the structure of the paper. In the next section we set up notation and review relevant facts on invariant chains and monomial ideals. The proof of Theorem 1.2(i) is given in Section 3. In Section 4 we prove that $\lim_{n \rightarrow \infty} (\text{reg}I_n/n)$ exists and has the expected integral value. In addition, we show that Conjecture 1.1(ii) is true for any Inc-invariant chains of monomial ideals up to an appropriate saturation (see Corollary 4.18). Some open problems are discussed in Section 5. Finally, the Appendix provides the proofs of two technical results in Section 4.

2. PRELIMINARIES

In this section we provide necessary notions and basic facts concerning invariant chains and monomial ideals.

2.1. Invariant chains of ideals. Let \mathbb{N} denote the set of positive integers. For $n \in \mathbb{N}$ let $\text{Sym}(n)$ be the symmetric group on $[n] = \{1, \dots, n\}$. It is convenient to identify each element of $\text{Sym}(n)$ with a permutation of \mathbb{N} that fixes every $k \geq n+1$. In this manner, $\text{Sym}(n)$ is exactly the stabilizer subgroup of $n+1$ in $\text{Sym}(n+1)$. Set

$$\text{Sym} = \bigcup_{n \geq 1} \text{Sym}(n).$$

Then Sym is the group of *finite permutations* of \mathbb{N} , i.e. permutations that fix all but finitely many elements of \mathbb{N} . In the literature Sym is sometimes also denoted by $\text{Sym}(\infty)$.

The *monoid of strictly increasing maps on \mathbb{N}* is defined as

$$\text{Inc} = \{\pi: \mathbb{N} \rightarrow \mathbb{N} \mid \pi(n) < \pi(n+1) \text{ for all } n \geq 1\}.$$

For $m \leq n$ set

$$\text{Inc}_{m,n} = \{\pi \in \text{Inc} \mid \pi(m) \leq n\}.$$

Throughout let $R = K[x_i \mid i \in \mathbb{N}]$ be an infinite dimensional polynomial ring over a field K . For $n \in \mathbb{N}$ the Noetherian polynomial ring $R_n = K[x_1, \dots, x_n]$ is regarded as a subring of R . Thus, one may view $R = \bigcup_{n \geq 1} R_n$ as the limit of the chain of increasing subrings $R_1 \subset R_2 \subset \dots$.

Let $\Pi = \text{Sym}$ or $\Pi = \text{Inc}$. Consider the action of Π induced by

$$\pi \cdot x_i = x_{\pi(i)} \quad \text{for every } \pi \in \Pi \text{ and } i \geq 1.$$

In what follows we often write $\pi(f)$ instead of $\pi \cdot f$ whenever $\pi \in \Pi$ and $f \in R$. An ideal $I \subseteq R$ is Π -invariant if

$$\Pi(f) = \{\pi(f) \mid \pi \in \Pi\} \subseteq I \quad \text{for every } f \in I.$$

A fundamental result of Cohen [4, 5] (see also [1, 10]) says that the ring R is Π -equivariant Noetherian, in the sense that every Π -invariant $I \subseteq R$ is generated by finitely many orbits, i.e. there exist $f_1, \dots, f_m \in R$ such that $I = \langle \Pi(f_1), \dots, \Pi(f_m) \rangle$. In this form, Cohen's result is also known as an equivariant Hilbert's basis theorem. And like the classical result of Hilbert, it also has an equivalent version in terms of chains of ideals, stating that every Π -invariant chain of ideals stabilizes (see [10] and also [11]). Here, a chain of ideals $\mathcal{I} = (I_n)_{n \geq 1}$ with $I_n \subseteq R_n$ for $n \geq 1$ is called Π -invariant if

$$(2.1) \quad \langle \Pi_{m,n}(I_m) \rangle_{R_n} \subseteq I_n \quad \text{for all } n \geq m \geq 1,$$

where

$$\Pi_{m,n} = \begin{cases} \text{Sym}(n) & \text{if } \Pi = \text{Sym}, \\ \text{Inc}_{m,n} & \text{if } \Pi = \text{Inc}, \end{cases}$$

and this chain *stabilizes* if there exists $r \in \mathbb{N}$ such that the inclusion in (2.1) becomes an equality for all $n \geq m \geq r$. The smallest number r with this property is called the *stability index* of \mathcal{I} and denoted by $\text{ind}(\mathcal{I})$.

The two versions of Cohen's result indicate that each Π -invariant ideal $I \subseteq R$ (viewed as a global object) is intimately related to any Π -invariant chain of ideals $\mathcal{I} = (I_n)_{n \geq 1}$ (viewed as a local object) with limit I , i.e. $I = \bigcup_{n \geq 1} I_n$. This kind of local-global relations has been further explored in the context of polyhedral geometry in [11, 16].

Among all Π -invariant chains with the same limit $I \subseteq R$, the *saturated* chain $\overline{\mathcal{I}} = (\bar{I}_n)_{n \geq 1}$ defined by $\bar{I}_n = I \cap R_n$ for $n \geq 1$ is evidently the largest one. It should be noted that any Sym-invariant chain eventually coincides with a saturated one (see [11, Corollary 6.1]). However, this is not true for Inc-invariant chains, and in fact, there are subtle differences between a saturated chain and non-saturated ones (see, e.g. Example 3.4). Note also that Sym-invariant chains are always Inc-invariant because $\text{Inc}_{m,n}(f) \subseteq \text{Sym}(n)(f)$ for all $n \geq m$ and $f \in R_m$.

2.2. Betti numbers of monomial ideals. Multigraded Betti numbers of monomial ideals can be computed via simplicial homology. We recall here this formula and a few related facts. Let J be a monomial ideal in $R_n = K[x_1, \dots, x_n]$ with minimal set of monomial generators $G(J)$. For $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$ let $x^{\mathbf{a}} = \prod_{i=1}^n x_i^{a_i}$. The *support* of $x^{\mathbf{a}}$ is denoted by

$$\text{supp}(x^{\mathbf{a}}) = \{i \in [n] \mid a_i > 0\}.$$

Let $\text{Msupp}(x^{\mathbf{a}}) = \max\{i \mid i \in \text{supp}(x^{\mathbf{a}})\}$ be the maximal index in $\text{supp}(x^{\mathbf{a}})$. We define

$$\text{Msupp}(J) = \max\{\text{Msupp}(u) \mid u \in G(J)\}.$$

Endow R_n with the standard \mathbb{Z}^n -grading. Then J admits a minimal \mathbb{Z}^n -graded free resolution over R_n . For $i \geq 0$ and $\mathbf{a} \in \mathbb{Z}_{\geq 0}^n$ let $\beta_{i,\mathbf{a}}(J) = \dim_K \text{Tor}_i(J, K)_{\mathbf{a}}$ denote the i th *Betti number* of J in degree \mathbf{a} . Moreover, let $x^F = \prod_{j \in F} x_j$ for any $F \subseteq [n]$. The *upper Koszul simplicial complex* of J in degree \mathbf{a} is defined as

$$\Delta_{\mathbf{a}}^J = \left\{ F \subseteq [n] \mid \frac{x^{\mathbf{a}}}{x^F} \in J \right\}.$$

Let $\text{Lcm}(J)$ be the *lcm lattice* of J , i.e. the lattice of all least common multiple of monomials in $G(J)$. The following result is well-known (see, e.g. [17, Theorem 1.34] and [24, Theorem 58.8]).

Lemma 2.1. *Let $J \subseteq R_n$ be a monomial ideal. Then for all $i \geq 0$ and $\mathbf{a} \in \mathbb{Z}_{\geq 0}^n$ it holds that*

$$\beta_{i,\mathbf{a}}(J) = \dim_K \tilde{H}_{i-1}(\Delta_{\mathbf{a}}^J),$$

where $\tilde{H}_{i-1}(\Delta_{\mathbf{a}}^J)$ denotes the $(i-1)$ th reduced homology group of $\Delta_{\mathbf{a}}^J$ with coefficients in K . Moreover, $\beta_{i,\mathbf{a}}(J) = 0$ if $x^{\mathbf{a}} \notin \text{Lcm}(J)$.

The functorial property of reduced homology implies the following (see [18, Lemma 2.5]).

Lemma 2.2. *Let $\Delta \subseteq \Sigma \subseteq \Gamma$ be simplicial complexes. If $\tilde{H}_i(\Sigma) = 0$ for some integer i , then the natural map $\tilde{H}_i(\Delta) \rightarrow \tilde{H}_i(\Gamma)$ induced by the inclusion $\Delta \subseteq \Gamma$ is zero.*

Recall that the *projective dimension* $\text{pd}J$ and the (*Castelnuovo-Mumford*) *regularity* $\text{reg}J$ of J measure the width and the height of the Betti table of J , respectively. More precisely,

$$\begin{aligned} \text{pd}J &= \max\{i \mid \beta_{i,\mathbf{a}}(J) \neq 0 \text{ for some } \mathbf{a} \in \mathbb{Z}_{\geq 0}^n\}, \\ \text{reg}J &= \max\{|\mathbf{a}| - i \mid \beta_{i,\mathbf{a}}(J) \neq 0\}, \end{aligned}$$

where $|\mathbf{a}| = \sum_{j=1}^n a_j$ if $\mathbf{a} = (a_1, \dots, a_n)$.

We will need the following formula for regularity. It is a consequence of [2, Theorem 4.7]; see [14, Corollary 5.2] for a proof.

Lemma 2.3. *Let $J \subseteq R_n$ be a monomial ideal. Consider a variable x_k of R_n . Let $d \geq 0$ be an integer such that $J : x_k^d = J : x_k^{d+1}$. Then*

$$\max\{\text{reg}\langle J : x_k^e, x_k \rangle \mid 0 \leq e \leq d\} \leq \text{reg} J \in \{\text{reg}\langle J : x_k^e, x_k \rangle + e \mid 0 \leq e \leq d\}.$$

2.3. Weight functions. We introduce two invariants of monomial ideals that play an important role in this paper. Let $J \subsetneq R_n$ be a nonzero monomial ideal with minimal set of monomial generators $G(J)$. For $\mathbf{a} \in \mathbb{Z}_{\geq 0}^n$ set $\lambda(x^{\mathbf{a}}) = a_k$ with $k = \text{Msupp}(x^{\mathbf{a}})$ and $w(x^{\mathbf{a}}) = \max\{a_i \mid i \in \text{supp}(x^{\mathbf{a}})\}$. Define

$$\begin{aligned} \lambda(J) &= \min\{\lambda(u) \mid u \in G(J)\}, \\ w(J) &= \min\{w(u) \mid u \in G(J)\}. \end{aligned}$$

Evidently, $\lambda(u) \leq w(u)$ for all $u \in G(J)$, hence $\lambda(J) \leq w(J)$.

Example 2.4. If $J = \langle x_1^4 x_2, x_1^3 x_3^2 \rangle \subseteq R_3$, then

$$\lambda(J) = \lambda(x_1^4 x_2) = 1, \quad w(J) = w(x_1^3 x_3^2) = 3.$$

The function w is introduced in [14] (in a more general context) as a measure of the growth of regularity (see Conjecture 4.1). The function λ , on the other hand, will be used in the next section to estimate the non-vanishing degrees of Betti numbers, resulting in lower bounds for regularity and projective dimension. For our purposes, it is convenient to extend these functions to invariant chains of monomial ideals. To this end, we need the following.

Lemma 2.5. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals with $r = \text{ind}(\mathcal{I})$. Then*

$$\lambda(I_n) \leq \lambda(I_{n+1}) \leq w(I_{n+1}) = w(I_r) \quad \text{for all } n \geq r.$$

Proof. Assume $n \geq r$ and let $u \in G(I_{n+1})$ with $\lambda(u) = \lambda(I_{n+1})$. Then there exist $v \in G(I_n)$ and $\pi \in \text{Inc}_{n, n+1}$ such that $u = \pi(v)$. Since $\lambda(u) = \lambda(\pi(v)) = \lambda(v)$, one gets

$$\lambda(I_n) \leq \lambda(v) = \lambda(u) = \lambda(I_{n+1}).$$

As noted above, the inequality $\lambda(I_{n+1}) \leq w(I_{n+1})$ follows at once from definition. The last equality is essentially shown in [14, Lemma 3.5], but we include its proof here for convenience of the reader. Arguing similarly as for the function λ one obtains $w(I_r) \leq w(I_{n+1})$. Now take $u' \in G(I_r)$ with $w(u') = w(I_r)$. Since $u' \in I_r \subseteq I_{n+1}$, there exists $v' \in G(I_{n+1})$ that divides u' . It follows that

$$w(I_{n+1}) \leq w(v') \leq w(u') = w(I_r). \quad \square$$

The preceding lemma implies that the sequence $(w(I_n))_{n \geq r}$ is constant, while the sequence $(\lambda(I_n))_{n \geq r}$ is eventually constant. This permits the following definition.

Definition 2.6. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals with $r = \text{ind}(\mathcal{I})$. Define

$$\begin{aligned} \lambda(\mathcal{I}) &= \max\{\lambda(I_n) \mid n \geq r\}, \\ w(\mathcal{I}) &= w(I_r). \end{aligned}$$

By Lemma 2.5 it is clear that $\lambda(\mathcal{I}) \leq w(\mathcal{I})$. The next result simplifies the computation of $\lambda(\mathcal{I})$ in the case of saturated chains.

Lemma 2.7. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals with $r = \text{ind}(\mathcal{I})$. If \mathcal{I} is saturated, then*

$$\lambda(\mathcal{I}) = \lambda(I_n) \quad \text{for all } n \geq r.$$

Proof. By Lemma 2.5, it suffices to show that $G(I_r) \subseteq G(I_n)$ for all $n \geq r$. Let $u \in G(I_r)$. If $u \notin G(I_n)$, there would exist $v \in I_r$ and $\pi \in \text{Inc}_{r,n}$ such that $\pi(v)$ is a proper divisor of u . Since $u \in R_r$, this gives $\pi(v) \in R_r$. Hence, $\pi(v) \in I_n \cap R_r = I_r$ because \mathcal{I} is saturated. But this contradicts the fact that $u \in G(I_r)$. \square

Note that this lemma is not true without the assumption that \mathcal{I} is saturated.

Example 2.8. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain with $I_3 = \langle x_1^2, x_2^2 x_3, x_3^2 \rangle$ and $\text{ind}(\mathcal{I}) = 3$. Then $I_n = \langle x_i^2 \mid 1 \leq i \leq n \rangle$ for $n \geq 4$. Thus, $\lambda(\mathcal{I}) = 2 > 1 = \lambda(I_3)$.

We conclude this section with a simple criterion for eventual linearity of numerical functions.

Lemma 2.9. *Let $f : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$ be a function and let $s \in \mathbb{Z}_{\geq 0}$. Then there exists $t \in \mathbb{Z}$ such that $f(n) = sn + t$ for $n \gg 0$ if one of the following conditions is satisfied:*

- (i) *For $n \gg 0$ it holds that $f(n+1) \leq f(n) + s$ and $f(n) \geq sn + r$ for some $r \in \mathbb{Z}$.*
- (ii) *For $n \gg 0$ it holds that $f(n+1) \geq f(n) + s$ and $f(n) \leq sn + r$ for some $r \in \mathbb{Z}$.*

Proof. We only prove (i), since the proof of (ii) is similar. Setting $t_n = f(n) - sn$ one deduces from (i) that $r \leq t_{n+1} \leq t_n$ for $n \gg 0$. So the sequence $(t_n)_{n \geq 1}$ must be eventually constant. \square

3. PROJECTIVE DIMENSION

The main goal of this section is to verify Conjecture 1.1(i) for saturated Inc-invariant chains of monomial ideals.

Theorem 3.1. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero saturated Inc-invariant chain of proper monomial ideals. Then there exists a constant $d \geq 1$ such that $\text{pd} I_n = n - d$ for all $n \gg 0$.*

Recall that any Sym-invariant chain of ideals is also Inc-invariant and eventually saturated (see [11, Corollary 6.1]). Thus, Theorem 3.1 generalizes Murai's result on Sym-invariant chains [18, Corollary 3.7]. To prove this theorem we follow his strategy, relating the non-vanishing of certain Betti numbers of I_n and I_{n+1} when $n \gg 0$. More specifically, the main ingredient in the proof is the following generalization and slight improvement of [18, Proposition 3.6].

Lemma 3.2. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero saturated Inc-invariant chain of proper monomial ideals with $r = \text{ind}(\mathcal{I})$. Let $\mathbf{a} = (a_1, \dots, a_t, 0, \dots, 0) \in \mathbb{Z}_{\geq 0}^n$ with $t \geq 1$, $a_t \geq 1$, and $n \geq r$. For each $p \in \mathbb{Z}_{\geq 0}$ denote $(\mathbf{a}, p) = (a_1, \dots, a_t, p, 0, \dots, 0) \in \mathbb{Z}_{\geq 0}^{n+1}$. Assume that $\beta_{i,\mathbf{a}}(I_n) \neq 0$ for some $i \geq 0$. Then the following hold:*

- (i) $a_t \geq \lambda(\mathcal{I})$;
- (ii) *There exists some $\lambda(\mathcal{I}) \leq p \leq a_t$ such that $\beta_{i+1,(\mathbf{a},p)}(I_{n+1}) \neq 0$.*

Proof. (i) Since $\beta_{i,\mathbf{a}}(I_n) \neq 0$, Lemma 2.1 implies $x^{\mathbf{a}} \in \text{Lcm}(I_n)$, i.e. $x^{\mathbf{a}} = \text{lcm}(u_1, \dots, u_k)$ for some $u_1, \dots, u_k \in G(J)$. It follows that $a_t = \lambda(x^{\mathbf{a}}) = \lambda(u_i)$ for some $i \in [k]$. Hence,

$$a_t \geq \lambda(I_n) = \lambda(\mathcal{J}),$$

where the last equality is due to Lemma 2.7.

(ii) Here the proof of [18, Proposition 3.6] carries over with some adjustments. We include the details for the sake of completeness. For $p \geq 0$ set

$$\Delta^{(p)} = \left\{ F \subseteq [t] : x_{t+1}^p \frac{x^{\mathbf{a}}}{x^F} \in I_{n+1} \right\}$$

and denote $\Delta = \Delta^{(0)}$. Also, let $\Sigma = \Delta^{[t]} := \Delta \cup \{F \cup \{t\} : F \in \Delta\}$ be the cone over Δ with apex t . We first prove the following:

- (a) $\Delta^{(0)} \subseteq \Delta^{(1)} \subseteq \dots \subseteq \Delta^{(a_t)}$;
- (b) $\Delta_{\mathbf{a}}^{I_n} = \Delta = \Delta^{(\lambda-1)}$, where $\lambda = \lambda(\mathcal{J})$;
- (c) $\Delta_{(\mathbf{a},p)}^{I_{n+1}} = \Delta^{(p)} \cup \{\{t+1\} \cup F : F \in \Delta^{(p-1)}\}$;
- (d) $\Delta^{(a_t)} \supseteq \Sigma$.

For (a): This is immediate.

For (b): We first show that $\Delta_{\mathbf{a}}^{I_n} = \Delta$. If $F \in \Delta_{\mathbf{a}}^{I_n}$, then $F \subseteq \text{supp } x^{\mathbf{a}} \subseteq [t]$. This implies the second equality in the chain

$$\Delta_{\mathbf{a}}^{I_n} = \left\{ F \subseteq [n] : \frac{x^{\mathbf{a}}}{x^F} \in I_n \right\} = \left\{ F \subseteq [t] : \frac{x^{\mathbf{a}}}{x^F} \in I_n \right\} = \left\{ F \subseteq [t] : \frac{x^{\mathbf{a}}}{x^F} \in I_{n+1} \right\} = \Delta.$$

The third equality holds since $I_n = I_{n+1} \cap R_n$, which is because \mathcal{J} is saturated. To prove $\Delta = \Delta^{(\lambda-1)}$ let $F \in \Delta^{(\lambda-1)}$. Then

$$x_{t+1}^{\lambda-1} \frac{x^{\mathbf{a}}}{x^F} \in I_{n+1}.$$

This means that $x_{t+1}^{\lambda-1} \frac{x^{\mathbf{a}}}{x^F}$ is divisible by some $u \in G(I_{n+1})$. Since $\lambda(u) \geq \lambda(I_{n+1}) = \lambda$ by Lemma 2.7, it follows that u divides $\frac{x^{\mathbf{a}}}{x^F}$. Hence, $\frac{x^{\mathbf{a}}}{x^F} \in I_{n+1}$, and therefore $F \in \Delta^{(0)} = \Delta$.

For (c): If $F \in \Delta_{(\mathbf{a},p)}^{I_{n+1}}$, then $F \subseteq \text{supp } x^{(\mathbf{a},p)} \subseteq [t+1]$. Assume that $F \subseteq [t+1]$. If $t+1 \notin F$, then

$$F \in \Delta_{(\mathbf{a},p)}^{I_{n+1}} \Leftrightarrow F \subseteq [t] \text{ and } x_{t+1}^p \frac{x^{\mathbf{a}}}{x^F} \in I_{n+1} \Leftrightarrow F \in \Delta^{(p)}.$$

On the other hand, if $t+1 \in F$, then $F = \{t+1\} \cup G$ with $G \subseteq [t]$. Hence,

$$\begin{aligned} F \in \Delta_{(\mathbf{a},p)}^{I_{n+1}} &\Leftrightarrow G \subseteq [t] \text{ and } x_{t+1}^p \frac{x^{\mathbf{a}}}{x^{G \cup \{t+1\}}} \in I_{n+1} \\ &\Leftrightarrow G \subseteq [t] \text{ and } x_{t+1}^{p-1} \frac{x^{\mathbf{a}}}{x^G} \in I_{n+1} \\ &\Leftrightarrow G \in \Delta^{(p-1)}. \end{aligned}$$

Thus we obtain (c).

For (d): It suffices to show that if $F \in \Delta$ and $t \notin F$, then $F \cup \{t\} \in \Delta^{(a_t)}$. Since $F \in \Delta$, it follows from (a) that $F \subseteq [t-1]$ and

$$(3.1) \quad \frac{x^{\mathbf{a}}}{x^F} = \frac{x_1^{a_1} \cdots x_{t-1}^{a_{t-1}}}{x^F} x_t^{a_t} \in I_n.$$

Let $\sigma_{t-1} : \mathbb{N} \rightarrow \mathbb{N}$ be the map defined by

$$\sigma_{t-1}(i) = \begin{cases} i & \text{if } 1 \leq i \leq t-1, \\ i+1 & \text{if } i \geq t. \end{cases}$$

Evidently, $\sigma_{t-1} \in \text{Inc}_{n,n+1}$. Thus $\sigma_{t-1}(I_n) \subseteq I_{n+1}$, and so (3.1) implies that

$$I_{n+1} \ni x_t^{a_t-1} \sigma_{t-1} \left(\frac{x^{\mathbf{a}}}{x^F} \right) = x_t^{a_t-1} \frac{x_1^{a_1} \cdots x_{t-1}^{a_{t-1}}}{x^F} x_{t+1}^{a_t} = x_{t+1}^{a_t} \frac{x^{\mathbf{a}}}{x^{F \cup \{t\}}}.$$

Therefore $F \cup \{t\} \in \Delta^{(a_t)}$, as wanted.

Next we prove the desired assertion on Betti numbers. By Lemma 2.1 and Claim (b), one gets

$$0 < \beta_{i,\mathbf{a}}(I_n) = \dim_K \tilde{H}_{i-1}(\Delta_{\mathbf{a}}^{I_n}) = \dim_K \tilde{H}_{i-1}(\Delta^{(\lambda-1)}).$$

Take a non-zero element $\alpha \in \tilde{H}_{i-1}(\Delta^{(\lambda-1)})$. Consider the maps

$$\tilde{H}_{i-1}(\Delta^{(\lambda-1)}) \xrightarrow{\iota_\lambda} \tilde{H}_{i-1}(\Delta^{(\lambda)}) \xrightarrow{\iota_{\lambda+1}} \cdots \xrightarrow{\iota_{a_t}} \tilde{H}_{i-1}(\Delta^{(a_t)})$$

induced by the chain in Claim (a). Denote $\alpha_{\lambda-1} = \alpha$ and $\alpha_k = (\iota_k \circ \cdots \circ \iota_\lambda)(\alpha)$ for $\lambda \leq k \leq a_t$. Note that $\iota_{a_t} \circ \cdots \circ \iota_\lambda$ is the natural map induced by the inclusion $\Delta^{(\lambda-1)} \subseteq \Delta^{(a_t)}$. Note also that $\tilde{H}_{i-1}(\Sigma) = 0$ since Σ is a cone. So Claim (d) and Lemma 2.2 yield

$$\iota_{a_t} \circ \cdots \circ \iota_\lambda = 0.$$

In particular, $\alpha_{a_t} = 0$. Thus there exists an integer $\lambda \leq p \leq a_t$ such that $\alpha_{p-1} \neq 0$ but $\alpha_p = 0$.

Set $\Gamma = \Delta^{(p-1)}$ and let $\Gamma^{[t+1]}$ be the cone over Γ with apex $t+1$. By Claim (c),

$$\Delta_{(\mathbf{a},p)}^{I_{n+1}} = \Delta^{(p)} \cup \{\{t+1\} \cup F : F \in \Delta^{(p-1)}\} = \Delta^{(p)} \cup \Gamma^{[t+1]}.$$

One has $\Delta^{(p)} \cap \Gamma^{[t+1]} = \Gamma$ since $t+1 \notin \Delta^{(p)}$. Consider the Mayer-Vietoris exact sequence

$$\cdots \rightarrow \tilde{H}_i(\Delta_{(\mathbf{a},p)}^{I_{n+1}}) \xrightarrow{\delta} \tilde{H}_{i-1}(\Gamma) \xrightarrow{\eta} \tilde{H}_{i-1}(\Delta^{(p)}) \oplus \tilde{H}_{i-1}(\Gamma^{[t+1]}) = \tilde{H}_{i-1}(\Delta^{(p)}) \rightarrow \cdots,$$

where $\tilde{H}_{i-1}(\Gamma^{[t+1]}) = 0$ because $\Gamma^{[t+1]}$ is a cone. One can check that $\eta(\alpha_{p-1}) = \pm \alpha_p = 0$. Hence, $0 \neq \alpha_{p-1} \in \text{Ker } \eta = \text{Im } \delta$. It follows that $\tilde{H}_i(\Delta_{(\mathbf{a},p)}^{I_{n+1}}) \neq 0$, and therefore

$$\beta_{i+1,(\mathbf{a},p)}(I_{n+1}) = \dim_K \tilde{H}_i(\Delta_{(\mathbf{a},p)}^{I_{n+1}}) \neq 0. \quad \square$$

Now we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. Note that $\text{pd } I_n \leq n-1$ for all $n \geq 1$ by Hilbert's syzygy theorem. On the other hand, it follows from Lemma 3.2(ii) that

$$\text{pd } I_{n+1} \geq \text{pd } I_n + 1 \quad \text{for all } n \geq r.$$

Now the proof is complete by using Lemma 2.9. □

Remark 3.3. Keep the assumption of Theorem 3.1.

- (i) It is natural to ask when $\text{pd}I_n$ becomes a linear function, or slightly less specifically, can one find a bound s such that

$$\text{pd}I_{n+1} = \text{pd}I_n + 1 \quad \text{for all } n \geq s?$$

If \mathcal{I} is a Sym-invariant chain of monomial ideals, then s can be simply chosen to be the stability index $\text{ind}(\mathcal{I})$, as shown in [18, Corollary 3.7]. However, this bound is not true for Inc-invariant chains; see Example 3.4.

- (ii) By the Auslander–Buchsbaum formula and Theorem 3.1, we see that

$$d = 1 + \lim_{n \rightarrow \infty} \text{depth}(R_n/I_n).$$

It would be interesting to have a combinatorial interpretation of d in terms of the chain \mathcal{I} .

To close this section we note that the proofs of Theorem 3.1 and Lemma 3.2 cannot be directly extended to arbitrary non-saturated chains. This can be seen by the following example, where the projective dimension and regularity may decrease after the stability index.

Example 3.4. Consider the Inc-invariant chain $\mathcal{I} = (I_n)_{n \geq 1}$ with $\text{ind}(\mathcal{I}) = r \geq 6$ and

$$I_r = \langle x_1, x_3, x_4, \dots, x_{r-1} \rangle_{x_1 x_3 x_4 \cdots x_{r-1}} + \langle x_2 x_3, x_{r-1} x_r \rangle.$$

Then one can check that

$$\text{pd}I_r = r - 1, \quad \text{pd}I_{r+1} = 3, \quad \text{pd}I_{r+2} = 5.$$

In fact, it is clear that $x_1 x_3 x_4 \cdots x_{r-1}$ is an element in the socle of R_r/I_r . Hence, $\text{depth}(R_r/I_r) = 0$. So by the Auslander–Buchsbaum formula, $\text{pd}I_r = \text{pd}(R_r/I_r) - 1 = r - 1$. Since

$$I_{r+1} = \langle x_2 x_3, x_2 x_4, x_3 x_4 \rangle + \langle x_{r-1} x_r, x_{r-1} x_{r+1}, x_r x_{r+1} \rangle,$$

one gets

$$\text{pd}I_{r+1} = \text{pd}\langle x_2 x_3, x_2 x_4, x_3 x_4 \rangle + \text{pd}\langle x_{r-1} x_r, x_{r-1} x_{r+1}, x_r x_{r+1} \rangle + 1 = 3.$$

Similarly, $\text{pd}I_{r+2} = 5$. Thus, although \mathcal{I} has stability index r , the sequence $(\text{pd}I_n)_n$ may fluctuate for a while from r before the stability pattern sets in.

It is also not hard to show that $\text{reg}I_r = r - 1 > \text{reg}I_{r+1} = \text{reg}I_{r+2} = 3$. Note that for saturated chains, the sequence $(\text{reg}I_n)_n$ is always non-decreasing after the stability index (see [14, Proof of Proposition 4.14]).

4. REGULARITY

For chains of monomial ideals Conjecture 1.1(ii) has the following more specific form (see [14, Conjecture 4.12]):

Conjecture 4.1. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals. Then there exists a constant D such that*

$$\text{reg}I_n = (w(\mathcal{I}) - 1)n + D \quad \text{for all } n \gg 0.$$

The main goal of this section is to prove the following evidence for the previous conjecture. We also verify this conjecture for a special class of chains of monomial ideals (see Lemma 4.12) and show that it is true up to a certain saturation (see Corollary 4.18).

Theorem 4.2. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals. Then

$$\lim_{n \rightarrow \infty} \frac{\text{reg } I_n}{n} = w(\mathcal{I}) - 1.$$

We prove this result using induction on the q -invariant of \mathcal{I} , defined as follows.

Definition 4.3. Let $J \subseteq R_n$ be a nonzero monomial ideal with $\text{Msupp}(J) = p$. Let $\delta(J)$ denote the maximal degree of a minimal monomial generator of J . Then the number

$$q(J) = \sum_{j=0}^{\delta(J)} \dim_K \left(\frac{R_p}{J \cap R_p} \right)_j.$$

is called the q -invariant of J .

For an Inc-invariant chain of nonzero monomial ideals $\mathcal{I} = (I_n)_{n \geq 1}$ with $r = \text{ind}(\mathcal{I})$, we set $q(\mathcal{I}) = q(I_r)$ and call it the q -invariant of \mathcal{I} .

The q -invariant is introduced in [21] (in a slightly different form). It is very useful for inductive arguments on Inc-invariant chains of monomial ideals, as illustrated in [13, 14, 21].

Example 4.4.

- (i) Evidently, $q(J) = 0$ if and only if $J = R_n$ for some $n \geq 1$.
- (ii) Let $J = \langle x_1^2 \rangle \subseteq R_2$. Then $\text{Msupp}(J) = 1$, $\delta(J) = 2$ and

$$q(J) = \sum_{j=0}^2 \dim_K \left(\frac{R_1}{J \cap R_1} \right)_j = \sum_{j=0}^2 \dim_K \left(\frac{K[x_1]}{\langle x_1^2 \rangle} \right)_j = 2.$$

The following construction, inspired by an analogous construction in [21, Lemma 6.10], will be used to decrease the q -invariant.

Definition 4.5. Let $e \geq 0$ be an integer and $\mathcal{I} = (I_n)_{n \geq 1}$ a nonzero Inc-invariant chain of monomial ideals with $\text{ind}(\mathcal{I}) = r$. Denote $p = \text{Msupp}(I_r)$. We define a chain $\mathcal{I}_e = (I_{e,n})_{n \geq 1}$ as follows:

$$I_{e,n} = \begin{cases} 0 & \text{if } n \leq r, \\ \langle (I_n : x_{n-r+p}^e) \cap R_{n-r+p-1} \rangle_{R_n} & \text{if } n \geq r+1. \end{cases}$$

Remark 4.6. Keep the assumption of Definition 4.5.

- (i) It is evident that $\text{Msupp}(I_n) \leq n - r + p$ for all $n \geq r$ since $I_n = \langle \text{Inc}_{r,n}(I_r) \rangle_{R_n}$, and one can show that equality occurs if the chain \mathcal{I} is saturated.
- (ii) The chain \mathcal{I}_e in Definition 4.5 is different from the one considered in [21, Lemma 6.10]. The latter is defined by taking colon ideals by fixed variables, while in Definition 4.5 we essentially take colon ideals by last variables in the supports. In this paper it is more convenient to work with the chain in Definition 4.5 because it is also Inc-invariant (see Lemma 4.7 below). The chain in [21], on the other hand, is only invariant with respect to a certain submonoid of Inc.

Basic properties of the chain \mathcal{I}_e are collected in the next result. We postpone its technical proof to the Appendix. See [14, Lemmas 5.3 and 5.10] for analogous properties of the chain considered in [21, Lemma 6.10].

Lemma 4.7. Let $e \geq 0$ be an integer and $\mathcal{I} = (I_n)_{n \geq 1}$ a nonzero Inc-invariant chain of monomial ideals with $\text{ind}(\mathcal{I}) = r$. Employ the notation of Definition 4.5. Then the following hold:

- (i) \mathcal{I}_e is a nonzero Inc-invariant chain of monomial ideals with $\text{ind}(\mathcal{I}_e) = r + 1$.
- (ii) One has $q(\mathcal{I}_e) \leq q(\mathcal{I})$, and equality occurs if and only if $I_{e,r+1} = \langle I_r \rangle_{R_{r+1}}$.
- (iii) One has $w(\mathcal{I}_e) \leq w(\mathcal{I})$, with equality if $0 \leq e \leq w(\mathcal{I}) - 1$.
- (iv) It holds that $\text{reg } I_n \geq \text{reg } I_{e,n}$ for all $n \geq r + 1$.

To prove Theorem 4.2 by induction we replace the chain \mathcal{I} with \mathcal{I}_e , employing the inequality $q(\mathcal{I}_e) \leq q(\mathcal{I})$ in Lemma 4.7(ii). For the induction step it is thus necessary to distinguish the case $q(\mathcal{I}) = q(\mathcal{I}_e)$. In fact, one only needs to consider this case when $e = w(\mathcal{I}) - 1$. This leads to the following notions. Recall that $\lambda(\mathcal{I}) \leq w(\mathcal{I})$ by Lemma 2.5.

Definition 4.8. A nonzero Inc-invariant chain $\mathcal{I} = (I_n)_{n \geq 1}$ of proper monomial ideals is called λ -maximal if $\lambda(\mathcal{I}) = w(\mathcal{I})$.

Definition 4.9. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain of ideals with $r = \text{ind}(\mathcal{I})$ and let $p = \text{Msupp}(I_r)$. We say that \mathcal{I} is *quasi-saturated* if $I_{n+1} \cap R_{n-r+p} = I_n \cap R_{n-r+p}$ for all $n \geq r$.

Remark 4.10. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a quasi-saturated chain with $r = \text{ind}(\mathcal{I})$ and $p = \text{Msupp}(I_r)$. Denote by $I = \bigcup_{n \geq 1} I_n$ the limit ideal of the chain. Let $\bar{\mathcal{I}} = (\bar{I}_n)_{n \geq 1}$ be the saturated chain of I , i.e. $\bar{I}_n = I \cap R_n$ for $n \geq 1$. Then $\bar{I}_{n-r+p} = I \cap R_{n-r+p} = \bigcup_{m \geq n} (I_m \cap R_{n-r+p}) = I_n \cap R_{n-r+p}$ for all $n \geq r$. This yields $I_n = \langle \bar{I}_{n-r+p} \rangle_{R_n}$ for all $n \geq r$. Thus the chain \mathcal{I} is a shift of the saturated chain $\bar{\mathcal{I}}$ (up to finitely many ideals in the beginning of the chain), which justifies the name ‘‘quasi-saturated’’. For this reason, Lemma 3.2 also holds for any quasi-saturated chain of monomial ideals.

We need the following characterization of the equality $q(\mathcal{I}) = q(\mathcal{I}_e)$.

Lemma 4.11. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals with $r = \text{ind}(\mathcal{I})$. For $e \geq 0$ consider the chain \mathcal{I}_e as in Definition 4.5. The following are equivalent:

- (i) $q(\mathcal{I}) = q(\mathcal{I}_e)$;
- (ii) The chain \mathcal{I} is quasi-saturated and $e \leq \lambda(\mathcal{I}) - 1$.

In particular, if $q(\mathcal{I}) = q(\mathcal{I}_e)$ with $e = w(\mathcal{I}) - 1$, then \mathcal{I} is quasi-saturated and λ -maximal.

Proof. (i) \Rightarrow (ii): Assume that $q(\mathcal{I}) = q(\mathcal{I}_e)$. Then $I_{e,r+1} = \langle I_r \rangle_{R_{r+1}}$ by Lemma 4.7(ii). This yields $I_{e,n+1} = \langle I_n \rangle_{R_{n+1}}$ for all $n \geq r$ since $\text{ind}(\mathcal{I}_e) = r + 1$. Note that $\text{Msupp}(I_n) \leq n - r + p$ for all $n \geq r$, with $p = \text{Msupp}(I_r)$. Hence, $\langle I_n \rangle_{R_{n+1}} \subseteq \langle I_{n+1} \cap R_{n-r+p} \rangle_{R_{n+1}} \subseteq \langle (I_{n+1} : x_{n-r+p+1}^e) \cap R_{n-r+p} \rangle_{R_{n+1}} = I_{e,n+1}$ for all $n \geq r$, and equalities must hold throughout. It follows that

$$I_n \cap R_{n-r+p} = \langle I_n \rangle_{R_{n+1}} \cap R_{n-r+p} = \langle I_{n+1} \cap R_{n-r+p} \rangle_{R_{n+1}} \cap R_{n-r+p} = I_{n+1} \cap R_{n-r+p}$$

for all $n \geq r$, i.e. \mathcal{I} is quasi-saturated.

Next we show that $e \leq \lambda(\mathcal{I}) - 1$. If this were not the case, then by Lemma 2.7 there would exist a minimal generator $u = x_1^{a_1} \cdots x_k^{a_k}$ of I_r with $1 \leq a_k \leq e$. Note that $k \leq p$. Since \mathcal{I} is quasi-saturated, one has $v = x_1^{a_1} \cdots x_{k-1}^{a_{k-1}} x_p^{a_k} \in I_r$, and hence $w = x_1^{a_1} \cdots x_{k-1}^{a_{k-1}} x_{p+1}^{a_k} \in I_{r+1}$. It follows that

$$u/x_k^{a_k} = w/x_{p+1}^{a_k} \in \langle (I_{r+1} : x_{p+1}^e) \cap R_p \rangle_{R_{r+1}} = I_{e,r+1}.$$

On the other hand, $u/x_k^{ak} \notin \langle I_r \rangle_{R_{r+1}}$ since u is a minimal generator of I_r . This, however, contradicts the fact that $I_{e,r+1} = \langle I_r \rangle_{R_{r+1}}$.

(ii) \Rightarrow (i): By Lemma 4.7(ii), it suffices to show that $I_{e,r+1} = \langle I_r \rangle_{R_{r+1}}$. Since the inclusion “ \supseteq ” is trivial, we only need to verify the reverse one. Let $u \in G(I_{e,r+1})$. Then $u \in R_p$ and $ux_{p+1}^e \in I_{r+1}$ with $p = \text{Msupp}(I_r)$. The latter implies $u \in I_{r+1}$ since $e \leq \lambda(\mathcal{I}) - 1 = \lambda(I_{r+1}) - 1$, where the last equality is due to Lemma 2.7. Hence, $u \in I_{r+1} \cap R_p \subseteq I_r$ because \mathcal{I} is quasi-saturated.

Finally, the equivalence of (i) and (ii) together with Lemma 2.5 yields the last statement. \square

The next result verifies Conjecture 4.1 for quasi-saturated chains that are λ -maximal.

Lemma 4.12. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be a nonzero Inc-invariant chain of proper monomial ideals. If \mathcal{I} is quasi-saturated and λ -maximal, then there exists $D \in \mathbb{Z}$ such that*

$$\text{reg } I_n = (w(\mathcal{I}) - 1)n + D \quad \text{for all } n \gg 0.$$

Proof. We can apply Lemma 3.2 because the chain \mathcal{I} is quasi-saturated (see Remark 4.10). Since $w(\mathcal{I}) = \lambda(\mathcal{I})$, Lemma 3.2(ii) yields

$$\text{reg } I_{n+1} \geq \text{reg } I_n + w(\mathcal{I}) - 1 \quad \text{for all } n \gg 0.$$

On the other hand, [14, Corollary 4.8] says that there exists $D' \in \mathbb{Z}$ such that

$$\text{reg } I_n \leq (w(\mathcal{I}) - 1)n + D' \quad \text{for all } n \gg 0.$$

The desired conclusion now follows from Lemma 2.9. \square

We are now in position to prove Theorem 4.2.

Proof of Theorem 4.2. Let $w = w(\mathcal{I})$. By [14, Corollary 4.8], it suffices to show that there exists a constant c such that

$$\text{reg } I_n \geq (w - 1)n + c \quad \text{for all } n \gg 0.$$

We proceed by induction on $q(\mathcal{I})$. If $q(\mathcal{I}) = 0$, then $I_n = R_n$ for all $n \geq \text{ind}(\mathcal{I})$, and there is nothing to prove. So assume that $q(\mathcal{I}) > 0$. Set $e_0 = w - 1$. Recall that $q(\mathcal{I}_{e_0}) \leq q(\mathcal{I})$ by Lemma 4.7(ii). We distinguish two cases:

Case 1: $q(\mathcal{I}_{e_0}) < q(\mathcal{I})$. As $e_0 < w$, one has $w(\mathcal{I}_{e_0}) = w$ by Lemma 4.7(iii). So by the induction hypothesis there exists a constant c such that

$$\text{reg } I_{e_0,n} \geq (w - 1)n + c \quad \text{for all } n \gg 0.$$

Since $\text{reg } I_n \geq \text{reg } I_{e_0,n}$ for all $n \geq \text{ind}(\mathcal{I}) + 1$ by Lemma 4.7(iv), the desired conclusion follows.

Case 2: $q(\mathcal{I}) = q(\mathcal{I}_{e_0})$. In this case, \mathcal{I} is quasi-saturated and λ -maximal by Lemma 4.11. Therefore, Lemma 4.12 allows us to conclude the proof. \square

Using a previous result in [14], an immediate consequence of Theorem 4.2 is that Conjecture 4.1 is true for Inc-invariant chains of Artinian monomial ideals.

Corollary 4.13. *If $\mathcal{I} = (I_n)_{n \geq 1}$ is an Inc-invariant chain of monomial ideals with $\dim(R_n/I_n) = 0$ for all $n \gg 0$, then there exists an integer D such that $\text{reg } I_n = (w(\mathcal{I}) - 1)n + D$ for all $n \gg 0$.*

Proof. Since $\text{reg } I_n$ is eventually a linear function by [14, Corollary 6.5], the conclusion follows easily from Theorem 4.2. \square

Remark 4.14. When $\text{reg } I_n$ is a linear function (as in Lemma 4.12 and Corollary 4.13), it would be interesting to understand the constant D . In case \mathcal{I} is a Sym-invariant chain of monomial ideals with $r = \text{ind}(\mathcal{I})$ and $w = w(\mathcal{I})$, Raicu [25, Theorem 6.1] (see also [18, Remark 3.11]) shows that $D = \text{reg}(I_r : (x_1 \cdots x_r)^{w-1})$. However, this is not true for Inc-invariant chains, as illustrated by the following example. Consider the Inc-invariant chain $\mathcal{I} = (I_n)_{n \geq 1}$ with $I_r = \langle x_1 x_r^w, x_r^s \rangle$ and $\text{ind}(\mathcal{I}) = r$, where $r \geq 2$ and $s \geq w + 1$. Then $w(\mathcal{I}) = \lambda(\mathcal{I}) = w$, and $I_r : (x_1 \cdots x_r)^{w-1} = \langle x_r \rangle$ has regularity 1. On the other hand, one can show that

$$\text{reg } I_n = (w-1)n + 1 + (r-1)(s-2w+1) \quad \text{for all } n \geq 2r-2.$$

Thus, depending on w and s , the constant $D = 1 + (r-1)(s-2w+1)$ can be either smaller or bigger than $\text{reg}(I_r : (x_1 \cdots x_r)^{w-1})$.

In the remainder of this section we show that Conjecture 4.1 is true for any chain of monomial ideals up to an appropriate saturation. To this end, let us first introduce the following construction.

Definition 4.15. Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain of ideals and let m be a positive integer. The m -saturation of \mathcal{I} is the chain $\mathcal{I}^{[m]} = (J_n)_{n \geq 1}$ determined by $J_1 = \langle 0 \rangle$ and

$$J_n = \left\langle \sum_{k=2}^n x_k^m I_{k-1} \right\rangle_{R_n} \quad \text{for } n \geq 2.$$

As shown below, $\mathcal{I}^{[m]}$ is a saturated chain, justifying the name. Nevertheless, one should not confuse $\mathcal{I}^{[m]}$ with the saturated chain $\overline{\mathcal{I}}$ considered so far.

Example 4.16. Consider again the chain \mathcal{I} in Example 2.8. Let $\overline{\mathcal{I}} = (\bar{I}_n)_{n \geq 1}$ and $\mathcal{I}^{[m]} = (J_n)_{n \geq 1}$ be the saturated chain and m -saturation of \mathcal{I} , respectively. Recall that $I_n = \langle x_i^2 \mid 1 \leq i \leq n \rangle$ for $n \geq 4$, hence $I = \bigcup_{n \geq 1} I_n = \langle x_i^2 \mid i \geq 1 \rangle \subset R$. This gives $\bar{I}_n = I \cap R_n = \langle x_i^2 \mid 1 \leq i \leq n \rangle$ for all $n \geq 1$. On the other hand,

$$\begin{aligned} J_4 &= \langle x_4^m I_3 \rangle_{R_4} = \langle x_1^2 x_4^m, x_2^2 x_3 x_4^m, x_3^2 x_4^m \rangle_{R_4}, \\ J_5 &= \langle x_4^m I_3 + x_5^m I_4 \rangle_{R_5} = \langle x_1^2 x_4^m, x_2^2 x_3 x_4^m, x_3^2 x_4^m \rangle_{R_5} + \langle x_i^2 x_5^m \mid 1 \leq i \leq 4 \rangle_{R_5}, \text{ etc.} \end{aligned}$$

The next result gathers some properties of m -saturation. Its proof will be given in the Appendix.

Lemma 4.17. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain of ideals. For $m \geq 1$ let $\mathcal{I}^{[m]} = (J_n)_{n \geq 1}$ be the m -saturation of \mathcal{I} . Then $\mathcal{I}^{[m]}$ is a saturated Inc-invariant chain. If, moreover, \mathcal{I} is a chain of monomial ideals, then the following hold:*

- (i) $\lambda(\mathcal{I}^{[m]}) = m$ and $w(\mathcal{I}^{[m]}) = \max\{w(\mathcal{I}), m\}$.
- (ii) For $n \geq 2$, $i \geq 0$ and $\mathbf{a} = (a_1, \dots, a_{n-1}) \in \mathbb{Z}_{\geq 0}^{n-1}$ one has

$$\beta_{i,(\mathbf{a},m)}(J_n) = \beta_{i,\mathbf{a}}(I_{n-1}) + \beta_{i-1,\mathbf{a}}(J_{n-1}), \quad \text{where } (\mathbf{a}, m) = (a_1, \dots, a_{n-1}, m).$$

- (iii) $\text{reg } J_n = \max\{\text{reg } I_{n-1} + m, \text{reg } J_{n-1} + m - 1\}$ for all $n \geq 2$.

Let us now show that Conjecture 4.1 is true for the m -saturation of an arbitrary Inc-invariant chain of monomial ideals, given that m is large enough.

Corollary 4.18. *Let \mathcal{J} be a nonzero Inc-invariant chain of proper monomial ideals. For $m \geq 1$ let $\mathcal{J}^{[m]} = (J_n)_{n \geq 1}$ be the m -saturation of \mathcal{J} . Then for $m \geq w(\mathcal{J})$ there exists an integer D such that*

$$\operatorname{reg} J_n = (m-1)n + D \quad \text{for all } n \gg 0.$$

Proof. If $m \geq w(\mathcal{J})$, then the chain $\mathcal{J}^{[m]}$ is saturated and λ -maximal by Lemma 4.17. So the result follows from Lemma 4.12. \square

We conclude this section with the following remark, showing in particular that the verification of Conjecture 4.1 can be reduced to the case of saturated chains if $w(\mathcal{J}) \geq 2$.

Remark 4.19. Let \mathcal{J} be a nonzero Inc-invariant chain of proper monomial ideals with $w(\mathcal{J}) \geq 2$. In order to verify Conjecture 4.1 for \mathcal{J} , it is enough to show that this conjecture is true for an m -saturation $\mathcal{J}^{[m]}$ of \mathcal{J} for some $1 \leq m \leq w(\mathcal{J}) - 1$. Indeed, suppose that Conjecture 4.1 holds for such an m -saturation. Then from Lemma 4.17(i) we get $w(\mathcal{J}^{[m]}) = w(\mathcal{J})$, hence

$$\operatorname{reg} J_n = \operatorname{reg} J_{n-1} + w(\mathcal{J}) - 1 > \operatorname{reg} J_{n-1} + m - 1 \quad \text{for all } n \gg 0.$$

So Lemma 4.17(iii) yields $\operatorname{reg} J_n = \operatorname{reg} I_{n-1} + m$ for all $n \gg 0$, and therefore $\operatorname{reg} I_n = \operatorname{reg} J_n - m$ must be eventually a linear function.

5. OPEN PROBLEMS

In this section we briefly discuss two open problems that are related to Conjecture 1.1.

As mentioned in the introduction, Conjecture 1.1 is verified for Sym-invariant chains of monomial ideals by Murai [18] and Raicu [25]. This result has been covered in their joint work on equivariant Hochster's formula [19], where they give a combinatorial description of the $\operatorname{Sym}(n)$ -module structure of $\operatorname{Tor}_i(I_n, K)$ when $I_n \subseteq R_n$ is a $\operatorname{Sym}(n)$ -invariant monomial ideal. It would be interesting to extend such an equivariant Hochster's formula to Inc-invariant chains.

Problem 5.1. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain of monomial ideals. Give an equivariant description of $\operatorname{Tor}_i(I_n, K)$ for $n \gg 0$.*

A possible approach to Conjecture 1.1 is to study primary decompositions of the ideals I_n . Experiments with Macaulay2 [8] lead to the following.

Conjecture 5.2. *Let $\mathcal{I} = (I_n)_{n \geq 1}$ be an Inc-invariant chain of monomial ideals. Then the number of associated primes of I_n is eventually a linear function.*

This conjecture is closely related to a recent work of Draisma, Eggermont, and Farooq [7]. There it is shown that if $\mathcal{I} = (I_n)_{n \geq 1}$ is a Sym-invariant chain of ideals (in more general polynomial rings), then the number of $\operatorname{Sym}(n)$ -orbits of minimal primes of I_n is eventually a quasipolynomial.

6. APPENDIX

We provide here the proofs of Lemmas 4.7 and 4.17. For $j \geq 0$ let $\sigma_j : \mathbb{N} \rightarrow \mathbb{N}$ be the increasing map defined by

$$\sigma_j(i) = \begin{cases} i & \text{if } 1 \leq i \leq j, \\ i+1 & \text{if } i \geq j+1. \end{cases}$$

Obviously, $\sigma_j \in \text{Inc}_{n,n+1}$ for all $n \geq 1$. Moreover, for any $f \in R_n$ it holds that

$$(6.1) \quad \text{Inc}_{n,n+1}(f) = \bigcup_{j \geq 0} \sigma_j(f) = \bigcup_{j=0}^n \sigma_j(f).$$

Proof of Lemma 4.7. Denote $p = \text{Msupp}(I_r)$.

(i) It is easily seen that the chain \mathcal{I}_e is Inc-invariant. Let us show that $\text{ind}(\mathcal{I}_e) = r + 1$. Since $I_{e,n} = 0$ for $n \leq r$ by definition, it suffices to check that $I_{e,n+1} = \langle \text{Inc}_{n,n+1}(I_{e,n}) \rangle_{R_{n+1}}$ for $n \geq r + 1$. Take $u \in G(I_{e,n+1})$ with $n \geq r + 1$. Then $ux_{n-r+p+1}^e \in I_{n+1}$, and so $ux_{n-r+p+1}^e = \sigma_j(v)z$ for some $j \geq 0$, $v \in G(I_n)$, and $z \in R_{n+1}$. Note that $\text{Msupp}(v) \leq \text{Msupp}(I_n) \leq n - r + p$. By definition of σ_j , it is clear that $\text{Msupp}(v) \leq \text{Msupp}(\sigma_j(v)) \leq \text{Msupp}(v) + 1$. Consider the following cases:

Case 1: $\text{Msupp}(v) \leq n - r + p - 1$. Then $\text{Msupp}(\sigma_j(v)) \leq n - r + p$. From $ux_{n-r+p+1}^e = \sigma_j(v)z$ we deduce that $\sigma_j(v)$ divides u . Since $v \in I_n \cap R_{n-r+p-1} \subseteq I_{e,n}$, we get $u \in \text{Inc}_{n,n+1}(I_{e,n})$.

Case 2: $\text{Msupp}(v) = \text{Msupp}(\sigma_j(v)) = n - r + p$. Then $\sigma_j(v) = v$, and v divides u . Note that $I_n = \langle \text{Inc}_{n-1,n}(I_{n-1}) \rangle_{R_n}$ since $n \geq r + 1$. Hence $v = \sigma_k(v')$ for some $k \geq 0$ and $v' \in I_{n-1}$. Evidently, $I_{n-1} \subseteq I_{e,n}$. Thus $u \in \langle \text{Inc}_{n,n+1}(I_{e,n}) \rangle_{R_{n+1}}$, because u is divisible by $\sigma_k(v') \in \text{Inc}_{n,n+1}(I_{e,n})$.

Case 3: $\text{Msupp}(v) = n - r + p$ and $\text{Msupp}(\sigma_j(v)) = n - r + p + 1$. This means that $j < n - r + p$. We can write $v = v'x_{n-r+p}^d$ with $d \geq 1$ and $\text{Msupp}(v') \leq n - r + p - 1$. Then from

$$ux_{n-r+p+1}^e = \sigma_j(v)z = \sigma_j(v')x_{n-r+p+1}^d z$$

we deduce that $d \leq e$ and $\sigma_j(v')$ divides u . Hence, $v' \in \langle (I_n : x_{n-r+p}^e) \cap R_{n-r+p-1} \rangle_{R_n} = I_{e,n}$ and again $u \in \langle \text{Inc}_{n,n+1}(I_{e,n}) \rangle_{R_{n+1}}$, as desired.

(ii) We first observe the following isomorphism of graded rings

$$\frac{R_p}{I_r \cap R_p} \cong \frac{R_{r+1}}{\langle I_r, x_{p+1}, \dots, x_{r+1} \rangle}.$$

Denote $p' = \text{Msupp}(I_{e,r+1})$. Since $I_{e,r+1} = \langle (I_{r+1} : x_{p+1}^e) \cap R_p \rangle_{R_{r+1}}$, it is obvious that $p' \leq p$. Hence

$$\langle I_r, x_{p+1}, \dots, x_{r+1} \rangle \subseteq \langle I_{e,r+1}, x_{p'+1}, \dots, x_{r+1} \rangle.$$

Combining this with the fact that $\delta(I_r) \geq \delta(I_{r+1}) \geq \delta(I_{e,r+1})$ we get

$$q(\mathcal{I}_e) = \sum_{j=0}^{\delta(I_{e,r+1})} \dim_K \left(\frac{R_{r+1}}{\langle I_{e,r+1}, x_{p'+1}, \dots, x_{r+1} \rangle} \right)_j \leq \sum_{j=0}^{\delta(I_r)} \dim_K \left(\frac{R_{r+1}}{\langle I_r, x_{p+1}, \dots, x_{r+1} \rangle} \right)_j = q(\mathcal{I}).$$

One checks easily that the equality occurs if and only if $I_{e,r+1} = \langle I_r \rangle_{R_{r+1}}$.

(iii) Choose $u \in G(I_r)$ with $w(u) = w(I_r)$. Since $I_r \subseteq I_{e,r+1}$, there exists $v \in G(I_{e,r+1})$ that divides u . It follows that

$$w(\mathcal{I}) = w(I_r) = w(u) \geq w(v) \geq w(I_{e,r+1}) = w(\mathcal{I}_e).$$

Let us show that the equality holds if $0 \leq e \leq w(\mathcal{I}) - 1$. Indeed, take any monomial $\tilde{v} \in G(I_{e,r+1})$. It suffices to prove that $w(\tilde{v}) \geq w(\mathcal{I})$. Since $\tilde{v}x_{p+1}^e \in I_{r+1}$, it is divisible by $\sigma_j(\tilde{u})$ for some $j \geq 0$ and $\tilde{u} \in G(I_r)$. This implies

$$\max\{w(\tilde{v}), e\} = w(\tilde{v}x_{p+1}^e) \geq w(\sigma_j(\tilde{u})) = w(\tilde{u}) \geq w(I_r) = w(\mathcal{I}).$$

Hence, $w(\tilde{v}) \geq w(\mathcal{I})$ because $e < w(\mathcal{I})$.

(iv) Denote $L = \langle I_n : x_{n-r+p}^e, x_{n-r+p} \rangle_{R_n}$. Then $\text{reg } L \leq \text{reg } I_n$ by Lemma 2.3. On the other hand, $L = I_{e,n} + \langle x_{n-r+p} \rangle_{R_n}$ and x_{n-r+p} is a non-zero-divisor on $R_n/I_{e,n}$. Hence, $\text{reg } I_{e,n} = \text{reg } L \leq \text{reg } I_n$. \square

Next we prove Lemma 4.17.

Proof of Lemma 4.17. We first show that $\mathcal{S}^{[m]}$ is an Inc-invariant chain. Since \mathcal{S} is Inc-invariant, it suffices to verify the following equalities for all $n \geq 2$:

$$\begin{aligned} J_n &= x_n^m \langle I_{n-1} \rangle_{R_n} + \langle J_{n-1} \rangle_{R_n}, \\ \langle \text{Inc}_{n-1,n}(J_{n-1}) \rangle_{R_n} &= x_n^m \langle \text{Inc}_{n-2,n-1}(I_{n-2}) \rangle_{R_n} + \langle J_{n-1} \rangle_{R_n}. \end{aligned}$$

The former is immediate from definition. To prove the latter we compute the orbit $\text{Inc}_{n-1,n}(x_k^m f)$ for $f \in R_{k-1}$ with $k \leq n-1$. Using (6.1) one gets

$$\text{Inc}_{n-1,n}(x_k^m f) = \bigcup_{j=0}^{k-1} \sigma_j(x_k^m f) \cup \bigcup_{j=k}^{n-1} \sigma_j(x_k^m f) = x_{k+1}^m \text{Inc}_{k-1,k}(f) \cup \{x_k^m f\}.$$

It follows that

$$\begin{aligned} \langle \text{Inc}_{n-1,n}(J_{n-1}) \rangle_{R_n} &= \sum_{k=2}^{n-1} \langle \text{Inc}_{n-1,n}(x_k^m I_{k-1}) \rangle_{R_n} = \sum_{k=2}^{n-1} \langle x_{k+1}^m \text{Inc}_{k-1,k}(I_{k-1}) + x_k^m I_{k-1} \rangle_{R_n} \\ &= x_n^m \langle \text{Inc}_{n-2,n-1}(I_{n-2}) \rangle_{R_n} + \sum_{k=2}^{n-1} x_k^m \langle \text{Inc}_{k-2,k-1}(I_{k-2}) + I_{k-1} \rangle_{R_n} \\ &= x_n^m \langle \text{Inc}_{n-2,n-1}(I_{n-2}) \rangle_{R_n} + \sum_{k=2}^{n-1} x_k^m \langle I_{k-1} \rangle_{R_n} \\ &= x_n^m \langle \text{Inc}_{n-2,n-1}(I_{n-2}) \rangle_{R_n} + \langle J_{n-1} \rangle_{R_n}, \end{aligned}$$

where the second-to-last equality is due to the fact that the chain \mathcal{S} is Inc-invariant.

To see that the chain $\mathcal{S}^{[m]}$ is saturated, one only needs to note that

$$J_{n+1} \cap R_n = (x_{n+1}^m \langle I_n \rangle_{R_{n+1}} + \langle J_n \rangle_{R_{n+1}}) \cap R_n = J_n \quad \text{for all } n \geq 1.$$

From now on, assume that \mathcal{S} is a chain of monomial ideals.

(i) Let $n \geq 2$. It is clear from the definition that $u \in G(J_n)$ precisely when $u = x_k^m v$ for some $v \in G(I_{k-1})$ and $2 \leq k \leq n$. This yields the claim since $\lambda(x_k^m v) = m$ and $w(x_k^m v) = \max\{m, w(v)\}$.

(ii) We exclude the trivial case $\mathbf{a} = \mathbf{0}$ and suppose that $\mathbf{a} = (a_1, \dots, a_t, 0, \dots, 0)$ with $a_t \neq 0$ for some $t \in [n-1]$. Denote $\Delta = \Delta_{\mathbf{a}}^{I_{n-1}}$, $\Gamma = \Delta_{\mathbf{a}}^{J_{n-1}}$, and $\Gamma^{[k]}$ the cone over Γ with apex k . We claim that

- (a) $\Gamma^{[t]} \subseteq \Delta$,
- (b) $\Delta_{(\mathbf{a}, m)}^{J_n} = \Delta \cup \Gamma^{[n]}$.

Evidently, $J_{n-1} \subseteq I_{n-1}$, hence $\Gamma \subseteq \Delta$. So to prove (a) it suffices to show that $F \cup \{t\} \in \Delta$ for any $F \in \Gamma$ with $t \notin F$. Set $\mathbf{a}' = (a_1, \dots, a_{t-1}, 0, \dots, 0)$. As the chain $\mathcal{S}^{[m]}$ is saturated, one has

$$\frac{x^{\mathbf{a}'}}{x^F} x_t^{a_t} = \frac{x^{\mathbf{a}}}{x^F} \in J_{n-1} \cap R_t = J_t.$$

Since $J_t : x_t^{a_t} = (x_t^m \langle I_{t-1} \rangle_{R_t} + \langle J_{t-1} \rangle_{R_t}) : x_t^{a_t} \subseteq \langle I_{t-1} \rangle_{R_t}$, this yields

$$\frac{x^{a'}}{x^F} \in J_t : x_t^{a_t} \subseteq \langle I_{t-1} \rangle_{R_t} \subseteq I_{n-1}.$$

It follows that

$$\frac{x^a}{x^{F \cup \{t\}}} = \frac{x^{a'}}{x^F} x_t^{a_t-1} \in I_{n-1},$$

and hence $F \cup \{t\} \in \Delta$.

Next we prove (b). Let $p \geq 0$ and consider, as in the proof of Lemma 3.2, the simplicial complex

$$\Gamma^{(p)} = \left\{ F \subseteq [n-1] \mid \frac{x^a}{x^F} x_n^p \in J_n \right\} = \left\{ F \subseteq [n-1] \mid \frac{x^a}{x^F} \in (J_n : x_n^p) \cap R_{n-1} \right\}.$$

One has $(J_n : x_n^m) \cap R_{n-1} = I_{n-1}$, hence $\Gamma^{(m)} = \Delta$. On the other hand, $\Gamma^{(m-1)} = \Gamma$ since

$$(J_n : x_n^{m-1}) \cap R_{n-1} = (x_n \langle I_{n-1} \rangle_{R_n} + \langle J_{n-1} \rangle_{R_n}) \cap R_{n-1} = J_{n-1}.$$

Thus Claim (c) in the proof of Lemma 3.2 gives

$$\Delta_{(\mathbf{a}, m)}^{J_n} = \Gamma^{(m)} \cup \{\{n\} \cup F \mid F \in \Gamma^{(m-1)}\} = \Delta \cup \Gamma^{[n]}.$$

Let us now verify the stated formula for Betti numbers. One has $\Delta \cap \Gamma^{[n]} = \Gamma$ since $n \notin \Delta$. So using (b) we obtain the Mayer-Vietoris exact sequence

$$\cdots \rightarrow \tilde{H}_{i-1}(\Gamma) \xrightarrow{\delta_{i-1}=(\psi_{i-1}, \varphi_{i-1})} \tilde{H}_{i-1}(\Delta) \oplus \tilde{H}_{i-1}(\Gamma^{[n]}) \rightarrow \tilde{H}_{i-1}(\Delta_{(\mathbf{a}, m)}^{J_n}) \rightarrow \tilde{H}_{i-2}(\Gamma) \xrightarrow{\delta_{i-2}} \cdots.$$

Since $\Gamma^{[n]}$ is a cone, $\varphi_k : \tilde{H}_k(\Gamma) \rightarrow \tilde{H}_k(\Gamma^{[n]}) = 0$ is a zero map for all k . On the other hand, it follows from (a) and Lemma 2.2 that $\psi_k : \tilde{H}_k(\Gamma) \rightarrow \tilde{H}_k(\Delta)$ is also a zero map for all k . Therefore, δ_k is a zero map for all k . The above exact sequence thus induces the following short exact sequence

$$0 \rightarrow \tilde{H}_{i-1}(\Delta) \rightarrow \tilde{H}_{i-1}(\Delta_{(\mathbf{a}, m)}^{J_n}) \rightarrow \tilde{H}_{i-2}(\Gamma) \rightarrow 0,$$

which provides

$$\beta_{i, (\mathbf{a}, m)}(J_n) = \dim \tilde{H}_{i-1}(\Delta_{(\mathbf{a}, m)}^{J_n}) = \dim \tilde{H}_{i-1}(\Delta) + \dim \tilde{H}_{i-2}(\Gamma) = \beta_{i, \mathbf{a}}(I_{n-1}) + \beta_{i-1, \mathbf{a}}(J_{n-1}).$$

(iii) From (ii) it follows that $\text{reg } J_n \geq \max\{\text{reg } I_{n-1} + m, \text{reg } J_{n-1} + m - 1\}$. On the other hand, using Lemma 2.3 one obtains

$$\text{reg } J_n \in \{\text{reg } \langle J_n : x_n^e, x_n \rangle + e \mid 0 \leq e \leq m\} = \{\text{reg } I_{n-1} + m, \text{reg } J_{n-1} + e \mid 0 \leq e \leq m - 1\}.$$

Therefore, it must hold that $\text{reg } J_n = \max\{\text{reg } I_{n-1} + m, \text{reg } J_{n-1} + m - 1\}$. \square

ACKNOWLEDGMENTS

This work is partially supported by the Simons Foundation Targeted Grant for the Institute of Mathematics - VAST (Award number: 558672), and by the Vietnam Academy of Science and Technology (grants CSCL01.01/22-23 and NCXS02.01/22-23). Parts of this work were carried out during a stay of the second author at the Vietnam Institute for Advanced Study in Mathematics (VIASM). He would like to thank VIASM for its hospitality and generous support. He also thanks the participants of the MFO-RIMS Tandem Workshop on ‘‘Symmetries on Polynomial Ideals and Varieties’’ for useful discussions related to this work.

REFERENCES

- [1] M. Aschenbrenner and C.J. Hillar, *Finite generation of symmetric ideals*. Trans. Amer. Math. Soc. **359** (2007), no. 11, 5171–5192.
- [2] G. Caviglia, H.T. Hà, J. Herzog, M. Kummini, N. Terai and N.V. Trung, *Depth and regularity modulo a principal ideal*. J. Algebr. Comb. **49** (2019), 1–20.
- [3] T. Church, J.S. Ellenberg, and B. Farb, *FI-modules and stability for representations of symmetric groups*. Duke Math. J. **164** (2015), no. 9, 1833–1910.
- [4] D.E. Cohen, *On the laws of a metabelian variety*. J. Algebra **5** (1967), 267–273.
- [5] D.E. Cohen, *Closure relations, Buchberger’s algorithm, and polynomials in infinitely many variables*. In: *Computation theory and logic*, 78–87, Lecture Notes in Comput. Sci., **270**, Springer, Berlin, 1987.
- [6] J. Draisma, *Noetherianity up to symmetry*. In: *Combinatorial algebraic geometry*, Lecture Notes in Mathematics **2108**, pp. 33–61, Springer, 2014.
- [7] J. Draisma, R.H. Eggermont, and Azhar Farooq, *Components of symmetric wide-matrix varieties*. Preprint, 2021, available at [arXiv:2103.05415](https://arxiv.org/abs/2103.05415).
- [8] D. Grayson and M. Stillman, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [9] C.J. Hillar, R. Krone, and A. Leykin, *Equivariant Gröbner bases*. In: *The 50th anniversary of Gröbner bases*, 129–154, Adv. Stud. Pure Math. **77**, Math. Soc. Japan, Tokyo, 2018.
- [10] C.J. Hillar and S. Sullivant, *Finite Gröbner bases in infinite dimensional polynomial rings and applications*. Adv. Math. **229** (2012), no. 1, 1–25.
- [11] T. Kahle, D.V. Le, and T. Römer, *Invariant chains in algebra and discrete geometry*. SIAM J. Discrete Math. **36** (2022), no. 2, 975–999.
- [12] R. Krone, A. Leykin, and A. Snowden, *Hilbert series of symmetric ideals in infinite polynomial rings via formal languages*. J. Algebra **485** (2017), 353–362.
- [13] D.V. Le, U. Nagel, H.D. Nguyen, and T. Römer, *Codimension and projective dimension up to symmetry*. Math. Nachr. **293** (2020), 346–362.
- [14] D.V. Le, U. Nagel, H.D. Nguyen, and T. Römer, *Castelnuovo-Mumford regularity up to symmetry*. Int. Math. Res. Not. Vol. 2021, No. **14**, 11010–11049.
- [15] D.V. Le and T. Römer, *A Kruskal-Katona type theorem and applications*. Discrete Math. **343** (5) (2020).
- [16] D.V. Le and T. Römer, *Theorems of Carathéodory, Minkowski-Weyl, and Gordan up to symmetry*. Preprint, 2021, available at [arXiv:2110.10657](https://arxiv.org/abs/2110.10657).
- [17] E. Miller and B. Sturmfels, *Combinatorial Commutative Algebra*. Graduate Texts in Mathematics, vol. **227**, Springer-Verlag, New York (2005).
- [18] S. Murai, *Betti tables of monomial ideals fixed by permutations of the variables*. Trans. Amer. Math. Soc. **373** (2020), 7087–7107.
- [19] S. Murai and C. Raicu, *An equivariant Hochster’s formula for \mathfrak{S}_n -invariant monomial ideals*. J. London Math. Soc. **105** (2022), 1974–2010.
- [20] U. Nagel, *Rationality of equivariant Hilbert series and asymptotic properties*. Trans. Amer. Math. Soc. **374** (2021), 7313–7357.
- [21] U. Nagel and T. Römer, *Equivariant Hilbert series in non-noetherian polynomial rings*. J. Algebra **486** (2017), 204–245.
- [22] U. Nagel and T. Römer, *FI- and OI-modules with varying coefficients*. J. Algebra **535** (2019), 286–322.
- [23] R. Nagpal and A. Snowden, *Symmetric subvarieties of infinite affine space*. Preprint, 2020, available at [arXiv:2011.09009](https://arxiv.org/abs/2011.09009).
- [24] I. Peeva, *Graded Syzygies*. Algebra and Applications. Volume **14**, Springer, London (2011).
- [25] C. Raicu, *Regularity of \mathfrak{S}_n -invariant monomial ideals*. J. Combin. Theory Ser. A **177**, 2021.
- [26] S. Sam and A. Snowden, *GL-equivariant modules over polynomial rings in infinitely many variables*. Trans. Amer. Math. Soc. **368** (2016), no. 2, 1097–1158.
- [27] S. Sam and A. Snowden, *Gröbner methods for representations of combinatorial categories*. J. Amer. Math. Soc. **30** (2017), no. 1, 159–203.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT OSNABRÜCK, 49069 OSNABRÜCK, GERMANY

Email address: dlevan@uos.de

INSTITUTE OF MATHEMATICS, VIETNAM ACADEMY OF SCIENCE AND TECHNOLOGY, 18 HOANG QUOC
VIET, 10307 HANOI, VIETNAM

Email address: ngdhop@gmail.com