

TORSION GROUPS OF ELLIPTIC CURVES OVER $\mathbb{Q}(\mu_{p^\infty})$

TOMISLAV GUŽVIĆ AND BORNA VUKOREPA

ABSTRACT. Let E/\mathbb{Q} be an elliptic curve and $p \in \{5, 7, 11\}$ be a prime. We determine the possibilities for $E(\mathbb{Q}(\zeta_p))_{tors}$. Additionally, we determine all the possibilities for $E(\mathbb{Q}(\zeta_{16}))_{tors}$ and $E(\mathbb{Q}(\zeta_{27}))_{tors}$. Using these results we are able to determine the possibilities for $E(\mathbb{Q}(\mu_{p^\infty}))_{tors}$.

1. INTRODUCTION

Let K be a number field such that $[K : \mathbb{Q}] = d$ and let E/K be an elliptic curve. A celebrated theorem of Mordell and Weil shows that $E(K)$ is a finitely generated abelian group. Therefore this group can be decomposed as $E(K) = E(K)_{tors} \oplus \mathbb{Z}^r$, $r \geq 0$. It is known that $E(K)_{tors}$ is of the form $C_m \oplus C_n$ for two positive integers m, n such that m divides n , where C_m and C_n denote cyclic groups of order m and n , respectively.

One of the goals in the theory of elliptic curves is the classification of torsion groups of elliptic curves defined over various fields.

Let d be a positive integer. Define $\Phi(d)$ to be the set of possible isomorphism classes of groups $E(K)_{tors}$, where K runs through all number fields K of degree d and E runs through all elliptic curves over K . In [28], Merel proved that $\Phi(d)$ is finite for all positive integers d . The set $\Phi(1)$ can be seen in Theorem 1.1 and was determined by Mazur [26].

Theorem 1.1 (Mazur, [26]). *Let E/\mathbb{Q} be an elliptic curve. Then*

$$E(\mathbb{Q})_{tors} \cong \begin{cases} C_m, & m = 1, \dots, 10, 12, \\ C_2 \oplus C_{2m}, & m = 1, \dots, 4. \end{cases}$$

The set $\Phi(2)$ has been determined by Kenku, Momose and Kamienny [23], [17]. Derickx, Etropolski, Hoeij, Morrow and Zureick-Brown have determined $\Phi(3)$ in [8].

Define $\Phi^{\text{CM}}(d)$ to be the set of possible isomorphism classes of groups $E(K)_{tors}$, where K runs through all number fields K of degree d and E runs through all elliptic curves with complex multiplication (CM). The set $\Phi^{\text{CM}}(1)$ has been determined by Olson in [33] and $\Phi^{\text{CM}}(d)$ for $d = 2, 3$ by Zimmer and his collaborators in [9], [29] and [34]. The sets $\Phi^{\text{CM}}(d)$, for $4 \leq d \leq 13$ have been

2010 *Mathematics Subject Classification.* 11G05.

Key words and phrases. Elliptic curves.

determined by Clark, Corn, Rice and Stankewicz in [7]. Bourdon, Pollack and Stankewicz have determined torsion groups of CM elliptic curves over odd degree number fields in [3].

Define $\Phi_{\mathbb{Q}}(d) \subseteq \Phi(d)$ to be the set of possible isomorphism classes of groups $E(K)_{tors}$, where K runs through all number fields K of degree d and E runs through all elliptic curves defined over \mathbb{Q} . For $d = 2, 3$, the sets $\Phi_{\mathbb{Q}}(d)$ have been determined by Najman [32].

Theorem 1.2. *Let E/\mathbb{Q} be an elliptic curve and K/\mathbb{Q} a quadratic extension. Then $E(K)_{tors}$ is isomorphic to the one of the following groups:*

$$C_m, \quad m = 1, 2, \dots, 9, 10, 12, 15, 16$$

$$C_2 \oplus C_{2m}, \quad m = 1, 2, 3, 4, 5, 6$$

$$C_3 \oplus C_{3m}, \quad m = 1, 2$$

$$C_4 \oplus C_4.$$

C_{15} is the only group which appears in only finitely many cases, and only over the extensions $\mathbb{Q}(\sqrt{5})$ and $\mathbb{Q}(\sqrt{-15})$.

Theorem 1.3. *Let E/\mathbb{Q} be an elliptic curve and K/\mathbb{Q} a cubic extension. Then $E(K)_{tors}$ is isomorphic to the one of the following groups:*

$$C_m, \quad m = 1, 2, \dots, 10, 12, 13, 14, 18, 21$$

$$C_2 \oplus C_{2m}, \quad m = 1, 2, 3, 4, 7.$$

C_{21} is the only group which appears in only finitely many cases, and only over the extension $\mathbb{Q}(\zeta_9)^+$.

The set $\Phi_{\mathbb{Q}}(4)$ has been determined by Chou [5] and González-Jiménez and Najman [12]. The set $\Phi_{\mathbb{Q}}(5)$ has been determined by González-Jiménez in [10]. González-Jiménez and Najman have also proved that $\Phi_{\mathbb{Q}}(p) = \Phi(1)$ for primes $p \geq 7$ in [12]. For an odd prime ℓ and a positive integer d , Propp [35] has determined when there exists a degree d number field K and an elliptic curve E/K with $j(E) \in \mathbb{Q} \setminus \{0, 1728\}$ such that $E(K)_{tors}$ contains a point of order ℓ .

Let μ_n , for positive integer n , be the set of all complex numbers ω such that $\omega^n = 1$. Note that for a prime number p we have that $\mathbb{Q}(\mu_p) = \mathbb{Q}(\zeta_p)$, where ζ_p is, as usual, p^{th} primitive root of unity.

For a prime number p , we define a set μ_{p^∞} as the set of all complex numbers ω for which there exists non-negative integer k such that $\omega^{p^k} = 1$. Note that $\mathbb{Q}(\mu_{p^\infty})$ is the set \mathbb{Q} extended with all $p^{n^{\text{th}}}$ primitive roots of unity.

In [13], the authors considered the following problem: assume that E/\mathbb{Q} is an elliptic curve, p a prime number and $K = \mathbb{Q}(\mu_p^\infty)$. They show that the torsion subgroup of E grows only over small subfields of K . More precisely, they showed the following:

Theorem 1.4. *Let E/\mathbb{Q} be an elliptic curve, then for a prime number $p \geq 5$ it holds that*

$$E(\mathbb{Q}(\mu_{p^\infty}))_{\text{tors}} = E(\mathbb{Q}(\mu_p))_{\text{tors}}.$$

Furthermore,

$$E(\mathbb{Q}(\mu_{3^\infty}))_{\text{tors}} = E(\mathbb{Q}(\mu_{3^3}))_{\text{tors}} \quad \text{and} \quad E(\mathbb{Q}(\mu_{2^\infty}))_{\text{tors}} = E(\mathbb{Q}(\mu_{2^4}))_{\text{tors}}.$$

Remark. This result is “the best possible”. For $E = 27a4$ we have that

$$E(\mathbb{Q}(\mu_{3^2}))_{\text{tors}} = C_9 \subsetneq C_{27} = E(\mathbb{Q}(\mu_{3^3}))_{\text{tors}}$$

and for $E = 32a4$ it holds that

$$E(\mathbb{Q}(\mu_{2^3}))_{\text{tors}} = C_2 \oplus C_4 \subsetneq C_2 \oplus C_8 = E(\mathbb{Q}(\mu_{2^4}))_{\text{tors}}.$$

It becomes natural to ask how can the torsion group of E/\mathbb{Q} grow when we consider the base change $E/\mathbb{Q}(\zeta_p)$. This becomes much harder than it seems as p grows because our methods sometimes rely on pure computation.

Our results are the following theorems.

Theorem 1.5. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{16}))_{\text{tors}}$ is one of the following groups (apart from those in Mazur’s theorem):*

$$C_4 \oplus C_4 \quad (15a1), \quad C_2 \oplus C_{10} \quad (2112bd2).$$

Theorem 1.6. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{27}))_{\text{tors}}$ can be only one of the next two groups (apart from those in Mazur’s theorem):*

$$C_3 \oplus C_3, \quad C_3 \oplus C_6, \quad C_3 \oplus C_9, \quad C_{21}, \quad C_{27}.$$

Additionally, in Lemma 5.1 we give a description of the set of possible group structures $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ can be isomorphic to for $p = 5, 7$ and 11 , where E/\mathbb{Q} is an elliptic curve.

We discuss further attempts to classify $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$, for arbitrary prime number p .

Magma [2] code used in this paper can be found [here](#).

2. NOTATION AND AUXILIARY RESULTS

Let E/F be an elliptic curve defined over a number field F . There exists an F -rational cyclic isogeny $\phi: E \rightarrow E'$ of degree n if and only if $\langle P \rangle$, where $P \in E(\overline{F})$ is a point of order n , is a $\text{Gal}(\overline{F}/F)$ -invariant group; in this case we say that E has an F -rational n -isogeny. When $F = \mathbb{Q}$, the possible degrees of n -isogenies of elliptic curves over \mathbb{Q} are known by the following theorem.

Theorem 2.1 (Mazur [27], Kenku [19], [21], [20], [22]). *Let E/\mathbb{Q} be an elliptic curve with a rational n -isogeny. Then*

$$n \in \{1, \dots, 19, 21, 25, 27, 37, 43, 67, 163\}.$$

There are infinitely many elliptic curves (up to $\overline{\mathbb{Q}}$ -isomorphism) with a rational n -isogeny over \mathbb{Q} for

$$n \in \{1, \dots, 10, 12, 13, 16, 18, 25\}$$

and only finitely many for all the other n . If E does not have complex multiplication, then $n \leq 18$ with $n \neq 14$ or $n \in \{21, 25, 37\}$.

Galois representations. Let E/\mathbb{Q} be an elliptic curve and let n a positive integer. The field $\mathbb{Q}(E[n])$ is the number field obtained by adjoining to \mathbb{Q} all the x and y -coordinates of the points of $E[n]$. The absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ acts on $E[n]$ by its action on the coordinates of the points, inducing a mod n Galois representation attached to E :

$$\rho_{E,n} : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(E[n]).$$

After we fix a basis for the n -torsion, we can identify $\text{Aut}(E[n])$ with $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$. This means that we can consider $\rho_{E,n}(\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}))$ as a subgroup of $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$, uniquely determined up to conjugacy. We shall denote $\rho_{E,n}(\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}))$ by $G_E(n)$. Moreover, since $\mathbb{Q}(E[n])$ is a Galois extension of \mathbb{Q} and $\ker \rho_{E,n} = \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(E[n]))$, by the first isomorphism theorem we have $G_E(n) \cong \text{Gal}(\mathbb{Q}(E[n])/\mathbb{Q})$.

We would like to know what are the possibilities for $G_E(n)$ as a subgroup of $\text{GL}(C_n)$. For some values of n , this can be seen in Tables 1 and 2. For most values of n we do not have a list of possibilities of $G_E(n)$, but we have a result that helps us see if for a given matrix subgroup M of $\text{GL}(C_n)$ there exists an elliptic curve E/\mathbb{Q} such that $\rho_{E,n}(\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})) = M$ (up to conjugation). The following lemma is well-known and it will be useful in cases where we analyze torsion group of elliptic curve E/\mathbb{Q} over a maximal real subfield of $\mathbb{Q}(\zeta_p)$.

Lemma 2.2. *Let E/\mathbb{Q} be an elliptic curve and L/K a quadratic extension of number fields with $L = K(\sqrt{d})$. Let $E(K)_{(2')}$ be the group of K -rational points of E of odd order. Then we have:*

$$E(K(\sqrt{d}))_{(2')} \cong E(K)_{(2')} \oplus E^d(K)_{(2')}.$$

Since all cyclotomic extensions are Galois over \mathbb{Q} , the following result imposes restrictions on the possibilities for torsion subgroup of E/\mathbb{Q} over a cyclotomic fields.

Lemma 2.3. *Let E/\mathbb{Q} be an elliptic curve, $m, n \in \mathbb{N}$ and K a finite Galois extension of \mathbb{Q} . Let $E(K)[mn] \cong C_m \oplus C_{mn}$ and $P \in E(K)$ point of order mn . Then we have:*

$$[\mathbb{Q}(mP) : \mathbb{Q}] \mid M(\phi(n), [K : \mathbb{Q}]),$$

where $M(\cdot, \cdot)$ is the greatest common divisor and ϕ is the Euler function.

Proof. Let P be a point of order mn with coordinates in K . Then we can take $Q \in E[mn]$ such that $\{P, Q\}$ is a basis for $E[mn]$. Consider the Galois representation modulo n with respect to E :

$$\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Z}/n\mathbb{Z}).$$

Let $\sigma \in \text{Gal}(K/\mathbb{Q})$. Then we have $P^\sigma = \alpha P + \beta Q$ for some $\alpha, \beta \in C_{mn}$ because the action of σ on P preserves the order of a point.

Now we have $P^\sigma - \alpha P = \beta Q$, so $\beta Q \in E(K)$. From that follows $m\beta \equiv 0 \pmod{mn}$. Multiplying by m gives us $(mP)^\sigma = \alpha(mP)$ so $(mP)^\sigma \in \langle mP \rangle$ for all $\sigma \in \text{Gal}(K/\mathbb{Q})$. Because of preserving the order, α has to be in $(\mathbb{Z}/n\mathbb{Z})^\times$.

Since by considering the restriction map we get $\text{Gal}(K/\mathbb{Q}) \cong \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})/\text{Gal}(\overline{\mathbb{Q}}/K)$, we have $(mP)^\sigma \in \langle mP \rangle$ for all $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Therefore, for all $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$:

$$\rho(\sigma) = \begin{pmatrix} \varphi(\sigma) & \tau(\sigma) \\ 0 & \psi(\sigma) \end{pmatrix},$$

where $\varphi, \psi, \tau : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow C_n$, and φ, ψ are homomorphisms with image in $(C_n)^\times$. We know that $(mP)^\sigma = g(mP) \Leftrightarrow \varphi(\sigma) = g$, for all $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Therefore, we have:

$$|Im(\varphi)| = |\{(mP)^\sigma : \sigma \in \text{Gal}(K/\mathbb{Q})\}| = |Orb(mP)|.$$

It is clear that $\text{Stab}(mP) = \text{Gal}(K/\mathbb{Q}(mP))$, so by orbit and stabilizer theorem we have:

$$|Im(\varphi)| = \frac{|\text{Gal}(K/\mathbb{Q})|}{|\text{Gal}(K/\mathbb{Q}(mP))|} = [\mathbb{Q}(mP) : \mathbb{Q}].$$

On the other hand, we have $Im(\varphi) \leq (\mathbb{Z}/n\mathbb{Z})^\times$, so we have:

$$[\mathbb{Q}(mP) : \mathbb{Q}] \mid \phi(n).$$

$[\mathbb{Q}(mP) : \mathbb{Q}] \mid [K : \mathbb{Q}]$ is obvious and the proof is complete. □

One of the crucial results that we will need is the main result from [6]:

Theorem 2.4. *Let E/\mathbb{Q} be a rational elliptic curve. Then $E(\mathbb{Q}^{ab})_{tors}$ is isomorphic to the one of the following groups:*

$$C_m, \quad m = 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 25, 27, 37, 43, 67, 163$$

$$C_2 \oplus C_{2m}, \quad m = 1, 2, \dots, 8, 9$$

$$C_3 \oplus C_{3m}, \quad m = 1, 3$$

$$C_4 \oplus C_{4m}, \quad m = 1, 2, 3, 4$$

$$C_5 \oplus C_5,$$

$$C_6 \oplus C_6,$$

$$C_8 \oplus C_8.$$

This means that all of our candidate torsion subgroups are the subgroups of the groups in the above list. Our approach will mainly consist of eliminating a certain set of possibilities from the list above in order to classify torsion groups of elliptic curves over a specific cyclotomic field.

3. TORSION GROWTH OVER $\mathbb{Q}(\zeta_{16})$

Assume that E/\mathbb{Q} is an elliptic curve and that $C_m \oplus C_{mn} \subseteq E(\mathbb{Q}(\zeta_{16}))_{tors}$. By the properties of the Weil pairing, we have $\mathbb{Q}(\zeta_m) \subseteq \mathbb{Q}(\zeta_{16})$. It follows that $m \in \{1, 2, 4, 8\}$. We first eliminate a certain amount of cyclic groups listed in Theorem 2.4.

Lemma 3.1. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{16}))_{tors}$ is not isomorphic to C_n if*

$$n \in \{11, 14, 17, 18, 19, 21, 25, 27, 37, 43, 67, 163\}.$$

Proof. Lemma 2.3 gives us that if P_n is a point of order $n \notin \{17, 21, 25, 37\}$, we have $[\mathbb{Q}(P_n) : \mathbb{Q}] \mid 2$, which is impossible by Theorem 1.2. By the same lemma we get that if P_n is a point of order $n \in \{21, 25, 37\}$, then we have $[\mathbb{Q}(P_n) : \mathbb{Q}] \mid 4$, which is impossible by [5, Theorem 1.4].

It remains to consider the case $n = 17$. By [12, Theorem 5.8] we conclude that the point P_{17} of order 17 cannot be defined over some strictly smaller subfield of $\mathbb{Q}(\zeta_{16})$. That means that all $\sigma \in \text{Gal}(\mathbb{Q}(\zeta_{16})/\mathbb{Q})$ act differently on P_{17} . Since $\text{Gal}(\mathbb{Q}(\zeta_{16}))$ has four elements σ such that $\sigma^2 = id$, we have that $P_{17}^{\sigma^2} = k^2 P_{17} = P_{17}$ for four different σ . That means that we have $k^2 \equiv 1 \pmod{17}$ for four different k , a contradiction. \square

After eliminating plenty of cyclic groups, we discuss the cases when E obtains full 2-torsion over $\mathbb{Q}(\zeta_{16})$. This is done by the following lemmas:

Lemma 3.2. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{16}))_{tors}$ is not isomorphic to $C_2 \oplus C_{14}$ or $C_2 \oplus C_{18}$.*

Proof. We will prove the result for $C_2 \oplus C_{14}$ and the proof for the case $C_2 \oplus C_{18}$ is identical.

Let $P_{14} \in E(\mathbb{Q}(\zeta_{16}))$ be the point of order 14. From Lemma 2.3 we get that $[\mathbb{Q}(2P_{14}) : \mathbb{Q}] \mid 2$. It is also well-known that $[\mathbb{Q}(E[2]) : \mathbb{Q}] \in \{1, 2, 3, 6\}$. Since $E[2]$ is defined over $\mathbb{Q}(\zeta_{16})$, we have $[\mathbb{Q}(E[2]) : \mathbb{Q}] \in \{1, 2\}$.

Let Q_2 be a point of order 2 different from $7P_{14}$. We now have $[\mathbb{Q}(2P_{14}, 7P_{14}, Q_2) : \mathbb{Q}] \mid 4$. Since $2P_{14}, 7P_{14}$ and Q_2 generate our torsion subgroup $C_2 \oplus C_{14}$, we now know that this torsion subgroup appears over some strictly smaller subfield of $\mathbb{Q}(\zeta_{16})$.

Now we get a contradiction by using Theorem 1.2 and [5, Theorem 1.4]. \square

Lemma 3.3. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{16}))_{tors}$ is not isomorphic to C_{15} , $C_2 \oplus C_{12}$, $C_4 \oplus C_{12}$, $C_4 \oplus C_8$, $C_4 \oplus C_{16}$ or $C_8 \oplus C_8$.*

Proof. Both $X_1(15)$ and $X_1(2, 12)$ are elliptic curves. A computation in Magma shows that $X_1(15)(\mathbb{Q}(\zeta_{16}))$ and $X_1(15)(\mathbb{Q})$ have the same Mordell-Weil group structure. Therefore, since $C_{15} \notin \Phi(1)$, it also cannot appear over $\mathbb{Q}(\zeta_{16})$.

The curves $X_1(2, 12)(\mathbb{Q}(\zeta_{16}))$ and $X_1(2, 12)(\mathbb{Q}(i))$ have the same Mordell-Weil group structure. It was proven in [31, Lemma 7] that $C_2 \oplus C_{12}$ does not appear as a torsion subgroup over $\mathbb{Q}(i)$. Therefore, it also cannot appear over $\mathbb{Q}(\zeta_{16})$. This also covers the case $C_4 \oplus C_{12}$.

We consider the modular curves $X_1(4, 8)(\mathbb{Q}(\zeta_{16}))$ and $X_1(4, 8)(\mathbb{Q}(\zeta_8))$ which are actually elliptic curves. A computation in Magma shows that $X_1(4, 8)(\mathbb{Q}(\zeta_{16}))$ has rank 0 and the same torsion as $X_1(4, 8)(\mathbb{Q}(\zeta_8))$, which contains only cusps, see [4, Case 6.11]. This also covers the cases $C_8 \oplus C_8$ and $C_4 \oplus C_{16}$. \square

The following lemma is a bit more complicated than the previous ones. The idea is to consider the corresponding modular curve $X_1(16)$ and its Jacobian $J_1(16)$ over some cyclotomic fields in order to determine that the Jacobian has rank 0. After that, we determine torsion of $J_1(\mathbb{Q}(\zeta_{16}))$ and consequently the number of points on $X_1(16)(\mathbb{Q}(\zeta_{16}))$, all of which turn out to be cusps.

Lemma 3.4. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{16}))_{tors}$ is not isomorphic to C_{16} , $C_2 \oplus C_{16}$ or C_{13} .*

Proof. We consider the modular curve $X_1(16)(\mathbb{Q}(\zeta_{16}))$ and its Jacobian $J_1(16)(\mathbb{Q}(\zeta_{16}))$. We will demonstrate the use of standard methods for determining points on $X_1(16)(\mathbb{Q}(\zeta_{16}))$.

A computation in Magma shows that $r(J_1(16)(\mathbb{Q}(\zeta_{16}))) = 0$. Since

$$r(J_1(16)(\mathbb{Q}(\zeta_8)(\sqrt{\zeta_8}))) = r(J_1(16)(\mathbb{Q}(\zeta_8))) + r(J_1^{58}(16)(\mathbb{Q}(\zeta_8))),$$

the computation of the rank becomes shorter and we obtain that the rank of our Jacobian is 0.

Now we determine $J_1(16)(\mathbb{Q}(\zeta_{16}))_{tors}$. Rational prime $p = 17$ splits completely in $\mathbb{Q}(\zeta_{16})$ so by

reducing modulo some \mathfrak{p} that lies above p we get an injection

$$\text{red}_{\mathfrak{p}} : J_1(16)(\mathbb{Q}(\zeta_{16}))_{\text{tors}} \rightarrow J_1(16)(\mathbb{F}_{17}).$$

This map is injective due to the result of Katz [18].

A computation in Magma shows that $|J_1(16)(\mathbb{F}_{17})| = 400$. It follows that $|J_1(16)(\mathbb{Q}(\zeta_{16}))| \leq 400$. By using the generators of the 2-torsion subgroup of $J_1(16)(\mathbb{Q}(\zeta_{16}))$ and some elements of $J_1(16)(\mathbb{Q}(\zeta_{16}))$ that we get from some known points on $X_1(16)(\mathbb{Q}(\zeta_{16}))$, we are able to generate a group with 400 elements. Therefore, we know exactly how $J_1(16)(\mathbb{Q}(\zeta_{16}))$ looks like.

Now we are able to determine all points on $X_1(16)(\mathbb{Q}(\zeta_{16}))$ by considering the Mumford representations of the elements of $J_1(16)(\mathbb{Q}(\zeta_{16}))$. We easily get that $|X_1(16)(\mathbb{Q}(\zeta_{16}))| = 14$ with all points being cusps. Therefore, we can conclude that there are no elliptic curves $E/\mathbb{Q}(\zeta_{16})$ (and consequently E/\mathbb{Q}) with a point of order 16 over $\mathbb{Q}(\zeta_{16})$.

It remains to show that $E(\mathbb{Q}(\zeta_{16}))_{\text{tors}}$ can't be C_{13} . We consider the modular curve $X_1(13)(\mathbb{Q}(\zeta_{16}))$ and its Jacobian $J_1(13)(\mathbb{Q}(\zeta_{16}))$.

A computation in Magma shows that $r(J_1(13)(\mathbb{Q}(\zeta_{16}))) = 0$. As in the previous lemma, we obtain:

$$r(J_1(13)(\mathbb{Q}(\zeta_{16}))) = r(J_1(16)(\mathbb{Q}(\zeta_8))) + r(J_1^{58}(16)(\mathbb{Q}(\zeta_8))) = 0.$$

The next step is to determine $J_1(13)(\mathbb{Q}(\zeta_{16}))_{\text{tors}}$. We determine the two-torsion subgroup, which turns out to be trivial. Using the result of Katz [18], we get an injection:

$$\text{red}_{\mathfrak{p}} : J_1(13)(\mathbb{Q}(\zeta_{16}))_{\text{tors}} \rightarrow J_1(13)(\mathbb{F}_{17}).$$

We also get that rational prime $q = 41$ has inertia degree 2 in $\mathbb{Q}(\zeta_{16})$ so we have another injection:

$$\text{red}_q : J_1(13)(\mathbb{Q}(\zeta_{16}))_{\text{tors}} \rightarrow J_1(13)(\mathbb{F}_{41^2}).$$

We notice that $\gcd(\#J_1(13)(\mathbb{F}_{17}), \#J_1(13)(\mathbb{F}_{41^2})) = 76$, so $\#J_1(13)(\mathbb{Q}(\zeta_{16})) \mid 76$.

Since the two torsion subgroup is trivial, we get that $\#J_1(13)(\mathbb{Q}(\zeta_{16})) \mid 19$. We can find a point of order 19 on our Jacobian. By checking the Mumford representations of those divisors, we find that all of the points on the Jacobian come from cusps on $X_1(13)(\mathbb{Q}(\zeta_{16}))$ (and actually $X_1(13)(\mathbb{Q})$).

Therefore, we can conclude that there are no elliptic curves $E/\mathbb{Q}(\zeta_{16})$ (and consequently E/\mathbb{Q}) such that $E(\mathbb{Q}(\zeta_{16}))_{\text{tors}} \cong C_{13}$. \square

4. TORSION GROWTH OVER $\mathbb{Q}(\zeta_{27})$

In this section we prove Theorem 1.6 using a series of lemmas. First we eliminate some possibilities for a cyclic group to appear as the subgroup of $E(\mathbb{Q}(\zeta_{27}))$. Assume that E/\mathbb{Q} is an elliptic curve and that $C_m \oplus C_{mn} \subseteq E(\mathbb{Q}(\zeta_{27}))_{\text{tors}}$. By the properties of the Weil pairing and taking the Theorem 2.4 into account, we have $\mathbb{Q}(\zeta_m) \subseteq \mathbb{Q}(\zeta_{27})$. It follows that $m \in \{1, 2, 3, 6\}$.

Lemma 4.1. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{27}))_{tors}$ is not isomorphic to C_n if $n \in \{11, 13, 14, 15, 16, 17, 19, 25, 37, 43, 67, 163\}$.*

Proof. $\boxed{C_{11}, C_{25}}$: If $n \in \{11, 25\}$ is a point of order n , then then by Lemma 2.3 we have $[\mathbb{Q}(P_n) : \mathbb{Q}] \mid 2$, which is impossible by Theorem 1.2.

$\boxed{C_{13}}$: Let $P_{13} \in E(\mathbb{Q}(\zeta_{27}))$ be the point of order 13. By Lemma 2.3 we have $[\mathbb{Q}(P_{13}) : \mathbb{Q}] \mid 6$. Therefore, this torsion subgroup is defined over $\mathbb{Q}(\zeta_9)$. Theorem 1.2 tells us that this torsion subgroup cannot be defined over quadratic field. Therefore, it is defined over sextic or cubic field. Assume it is defined over sextic field (entire $\mathbb{Q}(\zeta_9)$). Then we can use Lemma 2.2 to get:

$$C_{13} \cong E(\mathbb{Q}(\zeta_9))_{(2')} \cong E(\mathbb{Q}(\zeta_9^+))_{(2')} \oplus E^{-3}(\mathbb{Q}(\zeta_9^+))_{(2')}.$$

This means that either E or E^{-3} has torsion subgroup $\mathbb{Z}/13\mathbb{Z}$ defined over $\mathbb{Q}(\zeta_9)^+$.

Now we will be finished if we prove that torsion subgroup $\mathbb{Z}/13\mathbb{Z}$ cannot appear over $\mathbb{Q}(\zeta_9)^+$. To do this, we consider $X_1(13)(\mathbb{Q}(\zeta_9)^+)$.

As before, we use Magma to determine that $J_1(13)(\mathbb{Q}(\zeta_9)^+) \cong J_1(13)(\mathbb{Q})$ and that all points on the Jacobian come from cusps, which completes the proof.

$\boxed{C_{14}}$: Let $P_{14} \in E(\mathbb{Q}(\zeta_{27}))$ be a point of order 14. By Lemma 2.3 it follows that $[\mathbb{Q}(2P_{14}) : \mathbb{Q}]$ divides 6, so $\mathbb{Q}(2P_{14})$ is contained in $\mathbb{Q}(\zeta_9)$. The point $7P_{14}$ of order 2 satisfies $[\mathbb{Q}(7P_{14}) : \mathbb{Q}] \in \{1, 2, 3\}$, which means that it is also contained in $\mathbb{Q}(\zeta_9)$. It follows that $P_{14} \in E(\mathbb{Q}(\zeta_9))$. Consider the modular curve $X_1(14)$. It is an elliptic curve with LMFDB label 14.a5. On the LMFDB page of the mentioned curve we can see that its torsion subgroup does not grow in any number field contained in $\mathbb{Q}(\zeta_9)$. It remains to show that $r(E(\mathbb{Q})) = r(E(\mathbb{Q}(\zeta_9))) = 0$, which turns out to be true by a computation in Magma [2]. Therefore $X_1(14)(\mathbb{Q}) = X_1(14)(\mathbb{Q}(\zeta_9))$ and there does not exist an elliptic curve over \mathbb{Q} with a point of order 14 over $\mathbb{Q}(\zeta_9)$ and consequently over $\mathbb{Q}(\zeta_{27})$.

$\boxed{C_{15}}$: Let $P_{15} \in E(\mathbb{Q}(\zeta_{27}))$ be a point of order 15. Then $3P_{15}$ is a point of order 5 and $[\mathbb{Q}(3P_{15}) : \mathbb{Q}]$ is a divisor of $[\mathbb{Q}(\zeta_{27}) : \mathbb{Q}] = 18$. By Table 1 we see that $[\mathbb{Q}(3P_{15}) : \mathbb{Q}] \in \{1, 2\}$. The same way as in the Lemma 4.3, $C_2 \oplus C_{12}$ case, we see that the point $5P_{15}$ of order 3 is also defined over at most quadratic extension contained in $\mathbb{Q}(\zeta_{27})$. Since there is only one quadratic extension contained in $\mathbb{Q}(\zeta_{27})$, namely $\mathbb{Q}(\zeta_3)$, we have $P_{15} \in E(\mathbb{Q}(\zeta_3))$, which contradicts the Theorem 1.2.

$\boxed{C_{16}}$: Let $P_{16} \in E(\mathbb{Q}(\zeta_{27}))$ be a point of order 16. Then the point $8P_{16}$ has order 2 and we have $[\mathbb{Q}(8P_{16}) : \mathbb{Q}] \in \{1, 2, 3\}$. By [12, Proposition 4.8.] we have $[\mathbb{Q}(P_{16}) : \mathbb{Q}] = 2^a 3^b$, where $a \geq 0$ is an integer and $b \in \{0, 1\}$. Since the field $\mathbb{Q}(P_{16})$ is contained in $\mathbb{Q}(\zeta_{27})$, it follows that $2^a 3^b$ divides $[\mathbb{Q}(\zeta_{27}) : \mathbb{Q}] = 18$. We conclude that $[\mathbb{Q}(P_{16}) : \mathbb{Q}] \in \{1, 2, 3, 6\}$. Assume that $[\mathbb{Q}(8P_{16}) : \mathbb{Q}] = 3$. This means that $\mathbb{Q}(8P_{16})$ is cyclic, so the entire 2-torsion is contained in this field and P_{16} is defined over a number field of degree 3 or 6, but such a field is contained in $\mathbb{Q}(\zeta_9)$. Therefore we have $C_2 \oplus C_{16} \subseteq E(\mathbb{Q}(\zeta_9))$, but this is impossible by [14, Theorem 1.1]. It remains to consider the case when $[\mathbb{Q}(8P_{16}) : \mathbb{Q}] \in \{1, 2\}$. It follows that $[\mathbb{Q}(P_{16}) : \mathbb{Q}] \in \{1, 2\}$ again by [12, Proposition 4.8.].

Since $\mathbb{Q}(P_{16})$ is at most quadratic extension of \mathbb{Q} contained in $\mathbb{Q}(\zeta_{27})$, it follows that $\mathbb{Q}(P_{16}) \subseteq \mathbb{Q}(\zeta_3)$. By [30, Theorem 1] this turns out to be impossible.

$\boxed{C_{17}, C_{37}}$: Assume that $n \in \{17, 37\}$ and that $P_n \in E(\mathbb{Q}(\zeta_{27}))$ is a point of order n . By [12, Theorem 5.8] it follows that $[\mathbb{Q}(P_n) : \mathbb{Q}]$ is divisible by 4, but since $\mathbb{Q}(P_n) \subseteq \mathbb{Q}(\zeta_{27})$, this is impossible.

$\boxed{C_{19}}$: Let us consider the case when $n = 19$. If $P_{19} \in E(\mathbb{Q}(\zeta_{27}))$ is a point of order 19, then E has a rational 19-isogeny. By [25], we have $j(E) = -2^{15} \cdot 3^3$. The 19th division polynomial $f_{E,19}$ must have a root over $\mathbb{Q}(\zeta_{27})$. Using Magma, we check that this is not the case and therefore we arrive at the contradiction.

$\boxed{C_{43}, C_{67}, C_{163}}$: Assume that $n \in \{43, 67, 163\}$ and $P_n \in E(\mathbb{Q}(\zeta_{27}))[n]$. By [25, Theorem 2.1] it follows that $[\mathbb{Q}(P_n) : \mathbb{Q}] \geq \frac{n-1}{2} > [\mathbb{Q}(\zeta_{27}) : \mathbb{Q}] = 18$, a contradiction. \square

Lemma 4.2. *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}(\zeta_{27}))_{tors}$ is not isomorphic to C_{18} or $C_2 \oplus C_{18}$.*

Proof. If $E(\mathbb{Q}(\zeta_{27}))_{tors} \cong C_{18}$, then Lemma 2.3 directly gives us that this torsion subgroup is defined over a number field of degree 6 which can only be $\mathbb{Q}(\zeta_9)$.

If $E(\mathbb{Q}(\zeta_{27}))_{tors} \cong C_2 \oplus C_{18}$, then Lemma 2.3 gives us that if $P_{18} \in E(\mathbb{Q}(\zeta_{27}))_{tors}$ is a point of order 18, then $2P_{18}$ is defined over $\mathbb{Q}(\zeta_9)$. We know that $[\mathbb{Q}(E[2]) : \mathbb{Q}] \in \{1, 2, 3, 6\}$, but all unique subextensions of $\mathbb{Q}(\zeta_{27})$ of those degrees are contained in $\mathbb{Q}(\zeta_9)$, so again our torsion subgroup is defined over $\mathbb{Q}(\zeta_9)$. Now from Lemma 2.2 we get that

$$C_9 \cong E(\mathbb{Q}(\zeta_9))_{(2')} \cong E(\mathbb{Q}(\zeta_9^+))_{(2')} \oplus E^{-3}(\mathbb{Q}(\zeta_9^+))_{(2')}.$$

Therefore, one of $E(\mathbb{Q}(\zeta_9^+))$ and $E^{-3}(\mathbb{Q}(\zeta_9^+))$ has a point of order 9. Let $P_2 \in E(\mathbb{Q}(\zeta_9))_{tors}$ be a point of order 2. If $[\mathbb{Q}(P_2) : \mathbb{Q}] \in \{1, 3\}$, then P_2 is on both $E(\mathbb{Q}(\zeta_9^+))$ and $E^{-3}(\mathbb{Q}(\zeta_9^+))$. If $[\mathbb{Q}(P_2) : \mathbb{Q}] = 2$, then there is another point Q_2 of order 2 on E defined over \mathbb{Q} . In any case, both $E(\mathbb{Q}(\zeta_9^+))$ and $E^{-3}(\mathbb{Q}(\zeta_9^+))$ have a point of order 2. Finally, one of them has a point of order 18. However, it was proved in [24, Lemma 3.4.7] that all the points on $X_1(18)(\mathbb{Q}(\zeta_9)^+)$ are cusps, which completes the proof. \square

Lemma 4.3. *Let E/\mathbb{Q} be an elliptic curve. Assume that $C_2 \oplus C_{2n} \cong E(\mathbb{Q}(\zeta_{27}))_{tors}$. Then $n \in \{1, 2, 3, 4\}$. Additionally, $C_6 \oplus C_6 \not\subseteq E(\mathbb{Q}(\zeta_{27}))$.*

Proof. From Theorem 2.4 it follows that $n \leq 9$. We have shown that $E(\mathbb{Q}(\zeta_{27}))$ cannot contain a point of order 18 in Lemma 4.2.

$\boxed{C_2 \oplus C_{10}}$: Let $P_5 \in E(\mathbb{Q}(\zeta_{27}))$ be the point of order 5. It follows that E has a rational 5-isogeny and $[\mathbb{Q}(P_5) : \mathbb{Q}] \in \{1, 2\}$. If $G_E(2) \subseteq 2B$, then by Table 1 we see that $[\mathbb{Q}(E[2]) : \mathbb{Q}] \in \{1, 2\}$. Thus we have found two at most quadratic fields contained in $\mathbb{Q}(\zeta_{27})$. Since $\mathbb{Q}(\zeta_{27})$ has an unique quadratic subextension $F = \mathbb{Q}(\sqrt{-3})$, it follows that $F = \mathbb{Q}(E[2]) = \mathbb{Q}(P_5)$. We conclude that $C_2 \oplus C_{10} \subseteq E(F)_{tors}$, which is impossible by [30, Theorem 1].

Assume that $G_E(2) = 2Cn$. By Theorem [38, Theorem 1.1] it follows that $j(E) = t^2 + 1728$, for some $t \in \mathbb{Q}$. Since E has a rational 5-isogeny, by [38, Theorem 1.3] we have $j(E) = \frac{25(s^2+10s+5)^3}{s^5}$, for some $s \in \mathbb{Q} \setminus \{0\}$. It remains to find rational points on the induced curve. By [16, Page 61] we see that such rational points do not exist.

$C_2 \oplus C_{12}$: Let $P_3 \in E(\mathbb{Q}(\zeta_{27}))$ be the point of order 3. The extension $\mathbb{Q}(P_3)$ is cyclic over \mathbb{Q} since it is a subfield of $\mathbb{Q}(\zeta_{27})$. By Table 1 we see that $G_E(3)$ must be contained in the Borel subgroup of $\text{GL}_2(\mathbb{Z}/3\mathbb{Z})$. Applying Theorem [25, Theorem 9.3] it follows that $[\mathbb{Q}(P_3) : \mathbb{Q}] \in \{1, 2\}$, so we have $\mathbb{Q}(P_3) \subseteq \mathbb{Q}(\zeta_3)$. Assume that $G_E(2) \subseteq 2B$. By [12, Proposition 4.6] it follows that $C_2 \oplus C_4 \subseteq E(\mathbb{Q}(\zeta_3))$. We conclude that $C_2 \oplus C_{12} \subseteq E(\mathbb{Q}(\zeta_3))$, which is impossible by [30, Theorem 1].

Consider the case when $G_E(2) = 2Cn$. By [12, Proposition 4.8] it follows that the point P_4 of order 4 is defined over cubic or sextic subfield contained in $\mathbb{Q}(\zeta_{27})$. A computation in Magma [2] shows that if a point of order 4 is defined over cubic or sextic number field, then $G_E(2) = \text{GL}_2(\mathbb{Z}/2\mathbb{Z})$, a contradiction.

$C_2 \oplus C_{16}$: By [11, Corollary 3.5] it follows that this is impossible.,

$C_6 \oplus C_6$: Since $\mathbb{Q}(E[6])$ is contained in $\mathbb{Q}(\zeta_{27})$, we have that $|G_E(6)|$ divides $[\mathbb{Q}(\zeta_{27}) : \mathbb{Q}] = 18$. Additionally, the group $G_E(6)$ is cyclic. If $|G_E(6)| < 6$, then it follows that E obtains a full 6-torsion over a number field of degree 1, 2 or 3, but this is impossible by Theorem 1.2 and Theorem 1.3. Assume that $|G_E(6)| \in \{6, 9\}$. A search in Magma [2] shows that there exists only one cyclic group G with such property, namely:

$$G := \left\langle \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} \right\rangle.$$

Reducing G modulo 2 and modulo 3 we see that $G_E(2) = 2Cn$ and $G_E(3) = 3Cs.1.1$. By [38, Theorem 1.1] we have that $j(E) = t^2 + 1728$, for some $t \in \mathbb{Q}$. Similar from Theorem [38, Theorem 1.2] we have that $j(E) = f(s)^3$, for some rational function $f(s)$ and $s \in \mathbb{Q}$. A computation in Magma [2] shows that the only affine point on the elliptic curve

$$t^2 + 1728 = x^3$$

is $(t, x) = (0, 12)$. A direct computation shows that the equation $12 = f(s)$ does not have a rational solution in \mathbb{Q} . Therefore E cannot have $C_6 \oplus C_6$ torsion over $\mathbb{Q}(\zeta_{27})$. □

5. TORSION GROWTH OVER $\mathbb{Q}(\zeta_5), \mathbb{Q}(\zeta_7)$ AND $\mathbb{Q}(\zeta_{11})$

We note that if p is a prime number and E/\mathbb{Q} an elliptic curve such that $E(\mathbb{Q}(\zeta_p))_{tors}$ contains a subgroup isomorphic to $C_n \oplus C_{mn}$, then by the properties of Weil pairing we have $\mathbb{Q}(\zeta_n) \subseteq \mathbb{Q}(\zeta_p)$ which forces $n \leq 2$.

Lemma 5.1. *Let E/\mathbb{Q} be an elliptic curve and let $p \in \{5, 7, 11\}$ be a prime number. Apart from the groups in Mazur's theorem, the group $E(\mathbb{Q}(\zeta_p))_{tors}$ can only be isomorphic to one of the following groups:*

- If $p = 5$,

$$C_5 \oplus C_5 \text{ (550k2)}, \quad C_{15} \text{ (50a2)} \quad \text{and} \quad C_{16} \text{ (15a7)}.$$

- If $p = 7$,

$$C_{13} \text{ (147c2)}, \quad C_{14} \text{ (49a1)}, \\ C_{18} \text{ (14a4)}, \quad C_2 \oplus C_{14} \text{ (49a4)}, \quad C_2 \oplus C_{18} \text{ (14a5)}.$$

- If $p = 11$,

$$C_{11} \text{ (121b2)}, \quad C_{25} \text{ (11a3)}, \quad C_2 \oplus C_{10} \text{ (10230bg2)}.$$

Proof. Assume that $p = 5$. By [4, Theorem 5] we conclude that the only possibilities are the ones listed in this Lemma and C_{17} . It is easy to rule out C_{17} by using [12, Theorem 5.8].

Consider the case when $p = 11$. If E/\mathbb{Q} is an elliptic curve and if $C_n \oplus C_{mn} \in E(\mathbb{Q}(\zeta_{11}))$, then by the properties of the Weil pairing we have $\mathbb{Q}(\zeta_n) \subseteq \mathbb{Q}(\zeta_{11})$, which forces $n \in \{1, 2, 11\}$. Applying the Theorem 2.4 we eliminate the possibility $n = 11$. By [16, Lemma 6.0.8], it remains to show that the groups C_{15}, C_{16} and $C_2 \oplus C_{12}$ do not occur. We note that the set $\Phi_{\mathbb{Q}}(2)$ is described by Theorem 1.2 and the description of the set $\Phi_{\mathbb{Q}}(5)$ can be found in [10, Theorem 1].

- C_{15} : Assume that $P_{15} \in E(\mathbb{Q}(\zeta_{11}))$ is a point of order 15. Obviously we have

$$E(\mathbb{Q}(\zeta_{11}))[15] \cong C_{15}.$$

By Lemma 2.3 it follows that $[\mathbb{Q}(P_{15}) : \mathbb{Q}] \in \{1, 2\}$, but this contradicts the Theorem 1.2.

- C_{16} : If $P_{16} \in E(\mathbb{Q}(\zeta_{11}))$ is a point of order 16, then $8P_{16}$ has order 2 and is defined over at most quadratic extension contained in $\mathbb{Q}(\zeta_{11})$, which is $\mathbb{Q}(\sqrt{-11})$. By [12, Proposition 4.8] it follows that $[\mathbb{Q}(P_{16}) : \mathbb{Q}] \in \{1, 2\}$. A computation in Magma [2] shows that $X_1(16)(\mathbb{Q}(\sqrt{-11}))$ contains only cusps.
- $C_2 \oplus C_{12}$: As in the previous case, we show that $C_2 \oplus C_{12} \subseteq E(\mathbb{Q}(\sqrt{-11}))$. The modular curve $X_1(2, 12)(\mathbb{Q}(\sqrt{-11}))$ has rank 0 and same torsion as over \mathbb{Q} , which means that there does not exist an elliptic curve with $C_2 \oplus C_{12}$ torsion over $\mathbb{Q}(\sqrt{-11})$.

It remains to consider $p = 7$. Assume that $n \in \{11, 15, 16, 17, 21, 25, 19, 37, 43, 67, 163, 27\}$ and that $E(\mathbb{Q}(\zeta_7)) \cong C_n$.

- $n \in \{11, 15, 17, 25\}$: Lemma 2.3 gives us that if P_n is a point of order n , we have $[\mathbb{Q}(P_n) : \mathbb{Q}] \mid 2$. Now Theorem 1.2 gives us the contradiction.
- $n \in \{19, 37, 43, 67, 163\}$: From [12, Theorem 5.8], we get that the point of order n cannot be defined over the field $\mathbb{Q}(\zeta_7)$ (a degree 6 extension).
- $n = 27$: This follows from [14, Theorem 1.1].

- $n = 16$: Lemma 2.3 gives us that if $P_{16} \in E(\mathbb{Q}(\zeta_7))$ is a point of order 16, we have $[\mathbb{Q}(P_{16}) : \mathbb{Q}] \mid 2$. That means that $P_{16} \in E(\mathbb{Q}(\sqrt{-7}))$. We can use the similar methods as before in Magma to consider $X_1(16)(\mathbb{Q}(\sqrt{-7}))$ and prove that E cannot have a point of order 16 defined over $\mathbb{Q}(\sqrt{-7})$.
- $n = 21$: For C_{21} , we conclude from [5, Lemma 2.7] that E has a \mathbb{Q} -rational 21-isogeny. There are 4 elliptic curves (up to $\overline{\mathbb{Q}}$ -isomorphism) with a rational 21-isogeny (see [1, p.78-80]). Therefore, we can use the division polynomial method since the elliptic curves with the same j -invariant have identical division polynomials, up to scalar. We will consider the seventh division polynomials. We can use Magma [2] to factor those polynomials in the field $\mathbb{Q}(\zeta_7)$ and see that they have no zeroes there. Hence, this case is impossible.

It remains to eliminate only three non-cyclic groups. For $C_2 \oplus C_{16}$, we can use Lemma 2.3 to show that if $P_{16} \in E(\mathbb{Q}(\zeta_7))$ is of order 16, then $[\mathbb{Q}(2P_{16}) : \mathbb{Q}] \mid 2$. By [12, Proposition 4.6], we can conclude that $[\mathbb{Q}(P_{16}) : \mathbb{Q}] \in \{1, 2\}$. Hence, we have a point of order 16 defined over $\mathbb{Q}(\sqrt{-7})$. However, we already proved that this can't happen when we considered C_{16} . For $C_2 \oplus C_{10}$ and $C_2 \oplus C_{12}$, we consider modular curves $X_1(2, 10)(\mathbb{Q}(\zeta_7))$ and $X_1(2, 12)(\mathbb{Q}(\zeta_7))$ and use Magma to show that they don't have non-cuspidal points, which completes the proof. \square

Remark. Ideally, one would like to give a useful description of possible isomorphism classes of $E/\mathbb{Q}(\zeta_p)$, where E/\mathbb{Q} is an elliptic curve and p is a prime number. One can start with the following question that seems to be out of reach for the authors at the time of writing this paper.

Let $n \in \{13, 16, 18, 25\}$. Under what conditions on the prime number p does there exist an elliptic curve E/\mathbb{Q} with a point $P_n \in E(\mathbb{Q}(\zeta_p))$ of order n ?

5.1. Appendix: Images of Mod p Galois representations associated to elliptic curves over \mathbb{Q} .

For each possible known subgroup $G_E(p) \subsetneq \mathrm{GL}_2(\mathbb{F}_p)$ where E/\mathbb{Q} is a non-CM elliptic curve and p is a prime, Tables 1 and 2 list in the first and second column the corresponding labels in Sutherland and Zywina notations, and the following data:

- $d_v = [G_E(p) : G_E(p)_v] = |G_E(p) \cdot v|$ for $v \in \mathbb{F}_p^2$, $v \neq (0,0)$; equivalently, the degrees of the extensions $\mathbb{Q}(P)$ over \mathbb{Q} for points $P \in E(\overline{\mathbb{Q}})$ of order p .
- $d = |G_E(p)|$; equivalently, the degree $\mathbb{Q}(E[p])$ over \mathbb{Q} .

Note that Tables 1 and 2 are partially extracted from Table 3 of [36]. The difference is that [36, Table 3] only lists the minimum of d_v , which is denoted by d_1 therein.

Sutherland	Zywina	d_v	d	Sutherland	Zywina	d_v	d
2Cs	G_1	1	1	5Cs.4.1	G_1	2, 4, 8	8
2B	G_2	1, 2	2	5Ns.2.1	G_3	8, 16	16
2Cn	G_3	3	3	5Cs	G_2	4, 4	16
3Cs.1.1	$H_{1,1}$	1, 2	2	5B.1.1	$H_{6,1}$	1, 20	20
3Cs	G_1	2, 4	4	5B.1.2	$H_{5,1}$	4, 5	20
3B.1.1	$H_{3,1}$	1, 6	6	5B.1.4	$H_{6,2}$	2, 20	20
3B.1.2	$H_{3,2}$	2, 3	6	5B.1.3	$H_{5,2}$	4, 10	20
3Ns	G_2	4	8	5Ns	G_4	8, 16	32
3B	G_3	2, 6	12	5B.4.1	G_6	2, 20	40
3Nn	G_4	8	16	5B.4.2	G_5	4, 10	40
5Cs.1.1	$H_{1,1}$	1, 4	4	5Nn	G_7	24	48
5Cs.1.3	$H_{1,2}$	2, 4	4	5B	G_8	4, 20	80
				5S4	G_9	24	96

Acknowledgments. *The authors gratefully acknowledge support from the QuantiXLie Center of Excellence, a project co-financed by the Croatian Government and European Union through the European Regional Development Fund - the Competitiveness and Cohesion Operational Programme (Grant KK.01.1.1.01.0004) and by the Croatian Science Foundation under the project no. IP-2018-01-1313.*

REFERENCES

- [1] B. J. Birch and W. Kuyk (editors): *Modular functions of one variable IV*, volume 476. Springer Berlin, Heidelberg, 1975. ↑ 13.
- [2] W. Bosma, J. Cannon, and C. Playoust: *The Magma Algebra System I: The User Language*. *J. Symbolic Comput.*, 24(3-4):235–265, 1997, ISSN 0747-7171. <http://dx.doi.org/10.1006/jsco.1996.0125>, Computational algebra and number theory (London, 1993). ↑ 3, 9, 11, 12, 13.

Sutherland	Zywina	d_v	d	Sutherland	Zywina	d_v	d
7Ns.2.1	$H_{1,1}$	6, 9, 18	18	7Nn	G_6	48	96
7Ns.3.1	G_1	12, 18	36	7B.2.1	$H_{7,2}$	3, 42	126
7B.1.1	$H_{3,1}$	1, 42	42	7B.2.3	$H_{7,1}$	6, 21	126
7B.1.3	$H_{4,1}$	6, 7	42	7B	G_7	6, 42	252
7B.1.2	$H_{5,2}$	3, 42	42	11B.1.4	$H_{1,1}$	5, 110	110
7B.1.5	$H_{5,1}$	6, 21	42	11B.1.5	$H_{2,1}$	5, 110	110
7B.1.6	$H_{3,2}$	2, 21	42	11B.1.6	$H_{2,2}$	10, 55	110
7B.1.4	$H_{4,2}$	3, 14	42	11B.1.7	$H_{1,2}$	10, 55	110
7Ns	G_2	12, 36	72	11B.10.4	G_1	10, 110	220
7B.6.1	G_3	2, 42	84	11B.10.5	G_2	10, 110	220
7B.6.3	G_4	6, 14	84	11Nn	G_3	120	240
7B.6.2	G_5	6, 42	84				

TABLE 1. Possible images $G_E(p) \neq \text{GL}_2(\mathbb{F}_p)$, for $p \leq 11$, for non-CM elliptic curves E/\mathbb{Q} .

Sutherland	Zywina	d_v	d	Sutherland	Zywina	d_v	d
13S4	G_7	72, 96	288	13B.4.1	G_5	6, 156	936
13B.3.1	$H_{5,1}$	3, 156	468	13B.4.2	G_4	12, 78	936
13B.3.2	$H_{4,1}$	12, 39	468	13B	G_6	12, 156	1872
13B.3.4	$H_{5,2}$	6, 156	468	17B.4.2	G_1	8, 272	1088
13B.3.7	$H_{4,2}$	12, 78	468	17B.4.6	G_2	16, 136	1088
13B.5.1	G_2	4, 156	624	37B.8.1	G_1	12, 1332	15984
13B.5.2	G_1	12, 52	624	37B.8.2	G_2	36, 444	15984
13B.5.4	G_3	12, 156	624				

TABLE 2. Known images $G_E(p) \neq \text{GL}_2(\mathbb{F}_p)$, for $p = 13, 17$ or 37 , for non-CM elliptic curves E/\mathbb{Q} .

[3] A. Bourdon and P. Pollack: *Torsion subgroups of CM elliptic curves over odd degree number fields*. Int. Math. Res. Not. IMRN, 16:4923–4961, 2017. <https://doi.org/10.1093/imrn/rnw163>. ↑ 2.

[4] P. Bruin and F. Najman: *A criterion to rule out torsion groups for elliptic curves over number fields*. Res. Number Theory, 2(3), 2016. <https://doi.org/10.1007/s40993-015-0031-5>. ↑ 7, 12.

[5] M. Chou: *Torsion of rational elliptic curves over quartic Galois number fields*. J. Number Theory, 160:603–628, 2016. <https://doi.org/10.1016/j.jnt.2015.09.013>. ↑ 2, 6, 7, 13.

[6] M. Chou: *Torsion of rational elliptic curves over the maximal abelian extension of \mathbb{Q}* . Pacific J. Math., 302(2):481–509, 2019. <https://doi.org/10.2140/pjm.2019.302.481>. ↑ 5.

[7] P. L. Clark, P. Corn, A. Rice, and J. Stankiewicz: *Computation on elliptic curves with complex multiplication*. LMS J. Comput. Math., 17:509–539, 2014. <https://doi.org/10.1112/S1461157014000072>. ↑ 2.

- [8] M. Derickx, A. Etropolski, M. V. Hoeij, J. S. Morrow, and D. Zureick-Brown: *Sporadic cubic torsion*. Algebra Number Theory, 15 (7):1837–1864, 2021. <https://doi.org/10.2140/ant.2021.15.1837>. ↑ 1.
- [9] G. Fung, H. Ströher, H. Williams, and H. Zimmer: *Torsion groups of elliptic curves with integral j -invariant over pure cubic fields*. J. Number Theory, 36:12–45, 1990. [https://doi.org/10.1016/0022-314X\(90\)90003-A](https://doi.org/10.1016/0022-314X(90)90003-A). ↑ 1.
- [10] E. González-Jiménez: *Complete classification of the torsion structures of rational elliptic curves over quintic number fields*. J. Algebra, 478:484–505, 2017. <https://doi.org/10.1016/j.jalgebra.2017.01.012>. ↑ 2, 12.
- [11] E. González-Jiménez and Á. Lozano-Robledo: *On the minimal degree of definition of p -primary torsion subgroups of elliptic curves*. Math. Res. Lett., 24:1067–1096, 2017. <https://dx.doi.org/10.4310/MRL.2017.v24.n4.a7>. ↑ 11.
- [12] E. González-Jiménez and F. Najman: *Growth of torsion groups of elliptic curves upon base change*. Math. Comp., 89:1457–1485, 2020. <https://doi.org/10.1090/mcom/3478>. ↑ 2, 6, 9, 10, 11, 12, 13.
- [13] T. Gužvić and I. Krijan: *Torsion groups of elliptic curves over some infinite abelian extensions of \mathbb{Q}* . Submitted. ↑ 2.
- [14] T. Gužvić: *Torsion growth of rational elliptic curves in sextic number fields*. Journal of Number Theory, 220:330–345, 2021. <https://doi.org/10.1016/j.jnt.2020.09.010>. ↑ 9, 12.
- [15] T. Gužvić: *Torsion of elliptic curves with rational j -invariant defined over number fields of prime degree*. Proc. Amer. Math. Soc., 149:3261–3275, 2021. <https://doi.org/10.1090/proc/15500>.
- [16] T. Gužvić: *Torsion of elliptic curves with rational j -invariant over number fields*. Phd thesis, 2021. ↑ 11, 12.
- [17] S. Kamienny: *Torsion points on elliptic curves and q -coefficients of modular forms*. Invent. Math., 109:221–229, 1992. <https://doi.org/10.1007/BF01232025>. ↑ 1.
- [18] N.M. Katz: *Galois properties of torsion points on abelian varieties*. Invent. Math., 62:481–502, 1981. <https://doi.org/10.1007/BF01394256>. ↑ 8.
- [19] M. A. Kenku: *The modular curve $X_0(39)$ and rational isogeny*. Math. Proc. Cambridge Philos. Soc., 85:21–23, 1979. <https://doi.org/10.1017/S0305004100055444>. ↑ 4.
- [20] M. A. Kenku: *The modular curve $X_0(169)$ and rational isogeny*. J. London Math. Soc., s2-22:239–244, 1980. <https://doi.org/10.1112/jlms/s2-22.2.239>, Corrigendum: <https://doi.org/10.1112/jlms/s2-23.3.428-s>. ↑ 4.
- [21] M. A. Kenku: *The modular curves $X_0(65)$ and $X_0(91)$ and rational isogeny*. Math. Proc. Cambridge Philos. Soc., 87:15–20, 1980. <https://doi.org/10.1017/S0305004100056462>. ↑ 4.
- [22] M. A. Kenku: *The modular curve $X_0(125)$, $X_1(25)$ and $X_1(49)$* . J. London Math. Soc., s2-23:415–427, 1981. <https://doi.org/10.1112/jlms/s2-23.3.415>. ↑ 4.
- [23] M. A. Kenku and F. Momose: *Torsion points on elliptic curves defined over quadratic fields*. Nagoya Math. J., 109:125–149, 1988. <https://doi.org/10.1017/S0027763000002816>. ↑ 1.
- [24] I. Krijan: *Torsion groups of elliptic curves over infinite abelian extensions of \mathbb{Q}* . <https://web.math.pmf.unizg.hr/~ikrijan/pdfs/disertacija.pdf>, Phd thesis, 2020. ↑ 10.
- [25] Á. Lozano-Robledo: *On the field of definition of p -torsion points on elliptic curves over the rationals*. Math. Ann., 357:279–305, 2013. <https://doi.org/10.1007/s00208-013-0906-5>. ↑ 10, 11.
- [26] B. Mazur: *Modular curves and the Eisenstein ideal*. Inst. Hautes Etudes Sci. Publ. Math., 47:33–186, 1978. <https://doi.org/10.1007/BF02684339>. ↑ 1.

- [27] B. Mazur: *Rational isogenies of prime degree*. Invent. Math., 44:129–162, 1978. <https://doi.org/10.1007/BF01390348>. ↑ 4.
- [28] L. Merel: *Bornes pour la torsion des courbes elliptiques sur les corps de nombres*. Invent. Math., 124:437–449, 1996. <https://doi.org/10.1007/s002220050059>. ↑ 1.
- [29] H. Müller, H. Ströher, and H. Zimmer: *Torsion groups of elliptic curves with integral j -invariant over quadratic fields*. J. Reine Angew. Math., 397:100–161, 1989. <https://doi.org/10.1515/crll.1989.397.100>. ↑ 1.
- [30] F. Najman: *Complete classification of torsion of elliptic curves over quadratic cyclotomic fields*. J. Number Theory, 130:1964–1968, 2010. <https://doi.org/10.1016/j.jnt.2009.12.008>. ↑ 10, 11.
- [31] F. Najman: *Torsion of elliptic curves over cyclotomic quadratic fields*. Math. J. Okayama Univ., 53:75–82, 2011. https://web.math.pmf.unizg.hr/~fnajman/tors_kon.pdf. ↑ 7.
- [32] F. Najman: *Torsion of rational elliptic curves over cubic fields and sporadic points on $X_1(n)$* . Math. Res. Lett., 23(1):245–272, 2016. <https://dx.doi.org/10.4310/MRL.2016.v23.n1.a12>. ↑ 2.
- [33] L. D. Olson: *Points of finite order on elliptic curves with complex multiplication*. Manuscripta Math., 14:195–205, 1974. <https://doi.org/10.1007/BF01171442>. ↑ 1.
- [34] A. Pethő, T. Weis, and H. Zimmer: *Torsion groups of elliptic curves with integral j -invariant over general cubic number fields*. Internat. J. Algebra Comput., 7:353–413, 1997. <https://doi.org/10.1142/S0218196797000174>. ↑ 1.
- [35] O. Propp: *Cartan images and ℓ -torsion points of elliptic curves with rational j -invariant*. Res. Number Theory, 4, 2018. <https://doi.org/10.1007/s40993-018-0097-y>. ↑ 2.
- [36] A. V. Sutherland: *Computing images of Galois representations attached to elliptic curves*. Forum Math. Sigma, 4:41–79, 2016. <https://doi.org/10.1017/fms.2015.33>. ↑ 14.
- [37] The LMFDB Collaboration: *The L -functions and modular forms database*, 2019. <http://www.lmfdb.org>, [Online; accessed 30 October 2019].
- [38] D. Zywna: *On the possible images of the mod ℓ representations associated to elliptic curves over \mathbb{Q}* . <https://arxiv.org/pdf/1508.07660.pdf>, Accepted for publication in Proceedings of the AMS. ↑ 11.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, ZAGREB, CROATIA

Email address: tguzvic@math.hr

URL: <https://web.math.pmf.unizg.hr/~tguzvic/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, ZAGREB, CROATIA

Email address: bovukor@math.hr

URL: <https://www.math.pmf.unizg.hr/hr/borna-vukorepa>