

# NOTE ON TAMAGAWA NUMBERS OF TORI ATTACHED TO CM ALGEBRAS

YASUHIRO OKI

ABSTRACT. We prove that any integer power of 2 can be realized as the Tamagawa number of a torus attached to a CM algebra considered by Guo–Sheu–Yu and Liang–Yang–Yu. Such a result is obtained by Liang–Yang–Yu under assuming “a generalized Landau conjecture”. The main contribution of this paper is to remove the above assumption.

## CONTENTS

1.	Introduction	1
2.	Galois extensions with Galois groups the quaternion groups	2
3.	Tori and their Tamagawa numbers	3
4.	Proof of the main theorem	5
	References	8

## 1. INTRODUCTION

Let  $T$  be a torus over a global field  $L_0$ . Then Ono introduced in [Ono63a] the Tamagawa number  $\tau(T)$  of  $T$ , which is important in the arithmetic of tori. He proved that Tamagawa numbers are always positive rational numbers, and constructed some examples of tori whose Tamagawa numbers are non-integral. Moreover, he conjectured in [Ono63b] that any positive rational number should be realized as the Tamagawa number of a torus. This conjecture was proved by Katayama ([Kat85]) in the case that  $L_0$  is a number field. See also [LYY21, Appendix A].

Here we consider the Tamagawa numbers of tori over  $\mathbb{Q}$  attached to CM algebras, which are investigated by Guo–Sheu–Yu ([GSY22]) and Liang–Yang–Yu ([LYY21])<sup>1</sup>. Let  $L$  is a CM algebra, that is,

$$L := L_1 \times \cdots \times L_n$$

where  $L_1, \dots, L_n$  is a CM field. Write  $L_\nu^+$  for the maximum totally real subfield of  $L_\nu$  for each  $\nu \in \{1, \dots, n\}$ , and put

$$L^+ := L_1^+ \times \cdots \times L_n^+.$$

Denote by  $N_{L/L^+}$  the norm map of  $L/L^+$ . Then we define a  $\mathbb{Q}$ -torus  $T_{L/L^+}$  as

$$T_{L/L^+} := \left\{ t \in \prod_{\nu=1}^n \text{Res}_{L_\nu/\mathbb{Q}} \mathbb{G}_{m, L_\nu} \mid N_{L/L^+}(t) \in \mathbb{G}_{m, \mathbb{Q}} \right\}.$$

In [LYY21], the authors considered a variant of the result of [Kat85] for  $T_{L/L^+}$ , and obtained the following:

---

<sup>1</sup>Such tori are called “CM tori” in [LYY21]. However, this terminology could be confused with the Serre group, which is a pro-torus defined in [DMOS82]. This is pointed out by Tetsushi Ito and Teruhisa Koshikawa.

**Theorem 1.1.** (i) ([GSY22, Theorem 1.1]; see also Corollary 3.2) *For any CM algebra  $L$ , the Tamagawa number  $\tau(T_{L/L^+})$  is an integer power of 2.*

(ii) ([LYY21, Corollary 7.6]) *For any  $n \in \mathbb{Z}_{\geq 0}$ , there is a CM algebra  $L$  such that*

$$\tau(T_{L/L^+}) = 2^n.$$

(iii) ([LYY21, Theorem 7.12]) *Assume the following.*

(L) (“a generalized Landau conjecture”) *For any  $a \in \mathbb{Z}_{\geq 0}$ , the set*

$$S(a) := \{ax^2 + 1 \in \mathbb{Z}_{>0} \mid x \in \mathbb{Z}_{>0}\}$$

*contains infinitely many prime numbers.*

*Then, for any  $n \in \mathbb{Z}_{<0}$ , there is a CM algebra  $L$  such that*

$$\tau(T_{L/L^+}) = 2^n.$$

Note that the conjecture (L) for  $a = 1$  is called the *Landau conjecture*, which is still open. Furthermore, by considering the set  $S(a)$  in the case that  $a$  is a certain prime number, it was also derived that (L) is not needed for  $n \geq -709,617$ .

In this short note, we remove the assumption in Theorem 1.1 (iii).

**Theorem 1.2** (Theorem 4.1). *For any  $n \in \mathbb{Z}$ , there is a CM algebra  $L$  such that*

$$\tau(T_{L/L^+}) = 2^n.$$

We sketch the proof of Theorem 1.2. We may assume  $n < 0$  by Theorem 1.1 (ii). Put  $n' := -n$ . The most important point is finding  $n'$  distinct CM fields  $L_1, \dots, L_{n'}$  satisfying the following:

- for any  $\nu \in \{1, \dots, n'\}$ ,  $L_\nu/\mathbb{Q}$  is Galois with Galois group the quaternion group  $Q_8$ ,
- the decomposition groups of  $L_\nu/\mathbb{Q}$  at all finite places are cyclic for each  $\nu \in \{1, \dots, n'\}$ ,
- there is an isomorphism  $\text{Gal}(\tilde{L}/\mathbb{Q}) \cong \prod_{\nu=1}^{n'} \text{Gal}(L_\nu/\mathbb{Q})$ , where  $\tilde{L}$  is the composite field of  $L_1, \dots, L_{n'}$ .

Note that [LYY21] uses the conjecture (L) to accomplish it. The main innovation of this paper is to provide another solution to the above problem, which is based on the Dirichlet prime number theorem and a variant of the inverse Galois problem. More precisely, we use the result of Kiming ([Kim90]), which gives a criterion for existence of a quadratic extension of a (real) biquadratic field which is Galois over  $\mathbb{Q}$  with Galois group  $Q_8$ .

**Organization of this paper.** In Section 2, we state the result on Galois extensions of Galois groups  $Q_8$  obtained by Kiming. Section 3 recalls the results given by [GSY22] and [LYY21] on the Tamagawa numbers of tori attached to CM algebras. Finally, we give a proof of Theorem 1.2 in Section 4.

**Acknowledgment.** The author would like to thank Tetsushi Ito, Teruhisa Koshikawa and Seidai Yasuda for their valuable comments.

This work was carried out with the support from the JSPS Research Fellowship for Young Scientists and KAKENHI Grant Number 22J00570.

## 2. GALOIS EXTENSIONS WITH GALOIS GROUPS THE QUATERNION GROUPS

Let  $k_0$  be a field of characteristic different from 2, and denote by  $\text{Br}(k_0)$  the Brauer group of  $k_0$ . Then we define a pairing

$$(\cdot, \cdot)_{k_0} : k_0^\times \times k_0^\times \rightarrow \text{Br}(k_0)$$

by sending  $(a, b) \in k_0^\times \times k_0^\times$  to the Brauer class of the quaternion algebra

$$\left(\frac{a, b}{k_0}\right) := k_0 \oplus k_0\alpha \oplus k_0\beta \oplus k_0\alpha\beta,$$

where  $\alpha^2 = a$ ,  $\beta^2 = b$  and  $\beta\alpha = -\alpha\beta$ . By definition, the pairing  $(, )_{k_0}$  is symmetric, and the image of  $(, )_{k_0}$  is contained in the 2-torsion group of  $\text{Br}(k_0)$ .

For  $a \in k_0^\times$ , put

$$k_{0,a} := k[T]/(T^2 - a).$$

We denote by  $N_{k_{0,a}/k_0} : k_{0,a}^\times \rightarrow k_0^\times$  the norm map of  $k_{0,a}/k_0$ .

**Proposition 2.1** ([GS06, Proposition 1.1.7]). *Let  $a, b \in k_0^\times$ . Then the following are equivalent:*

- (i)  $(a, b)_{k_0} = 1$ ,
- (ii)  $a \in N_{k_{0,b}/k_0}(k_{0,b}^\times)$ ,
- (iii)  $b \in N_{k_{0,a}/k_0}(k_{0,a}^\times)$ .

Let  $Q_8$  be the quaternion group of order 8, that is,

$$Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$$

satisfying  $i^2 = j^2 = k^2 = -1$ ,  $ji = -ij$ ,  $kj = -jk$  and  $ik = -ki$ . The following can be proved by direct computation.

**Lemma 2.2.** *A subgroup of  $Q_8$  is either one of the following:*

$$\{1\}, \quad \langle -1 \rangle, \quad \langle i \rangle, \quad \langle j \rangle, \quad \langle k \rangle, \quad Q_8.$$

*In particular, a subgroup  $H$  of  $Q_8$  is cyclic if and only if so is its image in  $Q_8/\langle -1 \rangle$ .*

We give a criterion for existence of certain  $Q_8$ -extension, which is one of the most important keys to prove Theorem 1.2.

**Theorem 2.3** ([Kim90, Theorem 4]). *Let  $k^+ = k_0(\sqrt{a}, \sqrt{b})$  be a biquadratic extension of  $k_0$ . Then the following are equivalent.*

- (i) *There is a quadratic extension of  $k/k^+$  such that  $k/k_0$  is Galois with Galois group  $Q_8$ ,*
- (ii)  $(a, a)_{k_0}(b, b)_{k_0}(a, b)_{k_0} = 1$ .

### 3. TORI AND THEIR TAMAGAWA NUMBERS

Let  $L_0$  be a number field, and fix an algebraic closure  $\bar{L}_0$  of  $L_0$ . We write  $\Gamma_{L_0}$  for the Galois group  $\text{Gal}(\bar{L}_0/L_0)$ .

For a place  $v$  of  $L_0$ , let  $L_{0,v}$  be the completion of  $L_0$  at  $v$ . Moreover, if  $v$  is finite, denote by  $\text{val}_v$  the discrete valuation on  $L_{0,v}$  satisfying  $\text{val}_v(\varpi) = 1$  for any uniformizer  $\varpi$  of  $L_{0,v}$ . In addition, we write  $q_v$  for the order of the residue field of  $L_{0,v}$ . On the other hand,  $|\cdot|_{\mathbb{R}}$  and  $|\cdot|_{\mathbb{C}}$  denote the usual absolute values on  $\mathbb{R}$  and  $\mathbb{C}$  respectively. Then put

$$|\cdot|_v : L_{0,v}^\times \rightarrow \mathbb{R}_{>0}; x \mapsto \begin{cases} q_v^{-\text{val}_v(x)} & \text{if } v \text{ is finite,} \\ |x|_{\mathbb{R}} & \text{if } L_{0,v} \cong \mathbb{R}, \\ |x|_{\mathbb{C}}^2 & \text{if } L_{0,v} \cong \mathbb{C}. \end{cases}$$

On the other hand, let  $\mathbb{A}_{L_0}$  and  $\mathbb{A}_{L_0}^\times$  be the adèle ring and the idèle group of  $L_0$  respectively. Put

$$|\cdot|_{\mathbb{A}_{L_0}^\times} : \mathbb{A}_{L_0}^\times \rightarrow \mathbb{R}_{>0}; (a_v)_v \mapsto \prod_v |a_v|_v,$$

where  $v$  runs through all places of  $L_0$ .

Let  $\mathbb{G}_m$  be the multiplicative group over  $L_0$ , that is,  $\mathbb{G}_m := \text{Spec } L_0[X, X^{-1}]$ . Recall that a *torus* over  $L_0$  is an algebraic group  $T$  over  $L_0$  satisfying  $T \otimes_{L_0} \overline{L_0} \cong \mathbb{G}_{m, \overline{L_0}}^d$  for some  $d \in \mathbb{Z}_{\geq 0}$ .

Let  $T$  be a torus over  $L_0$ . In the following, we recall the definition of the Tamagawa number of  $T$ . See [Ono63a, 3.2] for details. Let

$$X^*(T) := \text{Hom}_{\overline{L_0}}(T \otimes_{L_0} \overline{L_0}, \mathbb{G}_{m, \overline{L_0}}),$$

which is a finite free  $\mathbb{Z}$ -module equipped with continuous left action of  $\Gamma_{L_0}$  on  $X^*(T)$ . Define a subgroup  $T(\mathbb{A}_{L_0})^1$  of  $T(\mathbb{A}_{L_0})$  as follows:

$$T(\mathbb{A}_{L_0})^1 := \bigcap_{\chi \in X^*(T)^{\Gamma_{L_0}}} \text{Ker}(|\cdot|_{\mathbb{A}_{L_0}^\times} \circ \chi).$$

Then there is a unique Haar measure on it which is characterized by special values of the local Artin  $L$ -factors of the complex representation  $X^*(T) \otimes_{\mathbb{Z}} \mathbb{C}$  of  $\Gamma_{L_0}$  at all finite places of  $L_0$ . See [Ono63a, (3.2.1)]. We denote by  $\omega_{T, \mathbb{A}_{L_0}}^1$  the Haar measure on the quotient  $T(\mathbb{A}_{L_0})^1/T(L_0)$  induced by the above measure. Then the total volume

$$\tau(T) := \int_{T(\mathbb{A}_{L_0})^1/T(L_0)} \omega_{T, \mathbb{A}_{L_0}}^1$$

of  $T(\mathbb{A}_{L_0})^1/T(L_0)$  with respect to  $\omega_{T, \mathbb{A}_{L_0}}^1$  is known to be finite. We call  $\tau(T)$  the *Tamagawa number* of  $T$ .

In the sequel, we consider the Tamagawa number of the following torus. Let  $L$  be a CM algebra, that is,

$$L := L_1 \times \cdots \times L_n,$$

where  $L_1, \dots, L_n$  is a CM field. For  $\nu \in \{1, \dots, n\}$ , denote by  $L_\nu^+$  the maximum totally real subfield of  $L_\nu$ , and put

$$L^+ := L_1^+ \times \cdots \times L_n^+.$$

Then the norm map of  $L/L^+$  induces a homomorphism

$$N_{L/L^+} : \prod_{\nu=1}^n \text{Res}_{L_\nu/\mathbb{Q}} \mathbb{G}_{m, L_\nu} \rightarrow \prod_{\nu=1}^n \text{Res}_{L_\nu^+/\mathbb{Q}} \mathbb{G}_{m, L_\nu^+}.$$

We regard  $\mathbb{G}_{m, \mathbb{Q}}$  as a subgroup of  $\prod_{\nu=1}^n \text{Res}_{L_\nu^+/\mathbb{Q}} \mathbb{G}_m$  by the diagonal embedding. Then we set

$$T_{L/L^+} := \left\{ t \in \prod_{\nu=1}^n \text{Res}_{L_\nu/\mathbb{Q}} \mathbb{G}_{m, L_\nu} \mid N_{L/L^+}(t) \in \mathbb{G}_{m, \mathbb{Q}} \right\},$$

which is a torus over  $\mathbb{Q}$ .

**Proposition 3.1** ([GSY22, Theorem 1.1]). *Let  $L$  and  $L^+$  be as above.*

- (i) *We have  $\tau(T_{L/L^+}) = 2^n / (\mathbb{A}_{\mathbb{Q}} : \mathbb{Q}^\times \cdot N_{L/L^+}(T_{L/L^+}(\mathbb{A}_{\mathbb{Q}})))$ .*
- (ii) *Let  $S(L/L^+)$  be the set of prime numbers  $\ell$  for which  $N_{L/L^+}(T_{L/L^+}(\mathbb{Q}_\ell))$  does not contain  $\mathbb{Z}_\ell^\times$ . Then we have*

$$(\mathbb{A}_{\mathbb{Q}} : \mathbb{Q}^\times \cdot N_{L/L^+}(T_{L/L^+}(\mathbb{A}_{\mathbb{Q}}))) \mid \prod_{\ell \in S(L/L^+)} (\mathbb{Z}_\ell^\times : \mathbb{Z}_\ell^\times \cap N_{L/L^+}(T_{L/L^+}(\mathbb{Q}_\ell))).$$

**Corollary 3.2.** *For any CM algebra  $L$ , the Tamagawa number  $\tau(T_{L/L^+})$  is an integer power of 2.*

*Proof.* By Proposition 3.1, it suffices to prove that  $(\mathbb{Z}_\ell^\times : \mathbb{Z}_\ell^\times \cap N_{L/L^+}(T_{L/L^+}(\mathbb{Q}_\ell)))$  is a power of 2 for any prime number  $\ell$ . However, this follows from the inclusion  $(\mathbb{Z}_\ell^\times)^2 \subset N_{L/L^+}(T_{L/L^+}(\mathbb{Q}_\ell))$ . ■

**Proposition 3.3** ([LYY21, Corollary 7.6]). *For  $n \in \mathbb{Z}_{\geq 0}$ , there is a CM algebra  $L$  such that*

$$\tau(T_{L/L^+}) = 2^n.$$

**Proposition 3.4.** *Let  $L$  be a CM field which is Galois over  $\mathbb{Q}$  with Galois group  $Q_8$ .*

(i) *Assume that the decomposition group at  $w$  is cyclic for any finite place  $w$  of  $L$ . Then we have*

$$\tau(T_{L/L^+}) = 1/2.$$

(ii) *If  $L$  does not satisfy the assumption in (i), then we have*

$$\tau(T_{L/L^+}) = 2.$$

*Proof.* This is essentially the same as [LYY21, Proposition 6.7]. Note that the authors assume that  $L$  is of particular form. However, they only use the assumption on the decomposition groups of  $L/\mathbb{Q}$ . ■

**Proposition 3.5** ([LYY21, Proposition 1.2]; see also [LYY21, Proposition 7.7], [Rüd20, Proposition 7.21]). *Let  $n' \in \mathbb{Z}_{>0}$ . For  $\nu \in \{1, \dots, n'\}$ , let  $L_\nu$  be a CM field which is Galois over  $\mathbb{Q}$  with Galois group  $Q_8$ . We further assume the following:*

(i) *if we write  $\tilde{L}$  for the composite field of  $L_\nu$  for  $m \in \{1, \dots, n'\}$ , then there is an isomorphism*

$$\text{Gal}(\tilde{L}/\mathbb{Q}) \cong \prod_{\nu=1}^{n'} \text{Gal}(L_\nu/\mathbb{Q}),$$

(ii)  $\tau(T_{L_\nu/L_\nu^+}) = 1/2$  for all  $\nu \in \{1, \dots, n'\}$ .

Put  $L := L_1 \times \dots \times L_{n'}$  and  $L^+ := L_1^+ \times \dots \times L_{n'}^+$ . Then we have

$$\tau(T_{L/L^+}) = 1/2^{n'}.$$

#### 4. PROOF OF THE MAIN THEOREM

The goal of this section is to give a proof of the following:

**Theorem 4.1.** *For any  $n \in \mathbb{Z}$ , there is a CM algebra  $L$  such that  $\tau(T_{L/L^+}) = 2^n$ .*

In the following, we give a proof of Theorem 4.1. First, we recall the properties on the pairings  $(, )_0 := (, )_{\mathbb{Q}}$  and  $(, )_v := (, )_{\mathbb{Q}_v}$  for (possibly infinite) places  $v$  of  $\mathbb{Q}$ .

**Proposition 4.2.** *Let  $v$  be a place of  $\mathbb{Q}$ , and  $a, b \in \mathbb{Q}_v^\times$ .*

(i) *If  $v$  is infinite, then we have  $(a, b)_v = 1$  if and only if either  $a$  or  $b$  is positive.*

(ii) *Assume that  $v = \ell$  is a non-dyadic finite place, and  $a, b \in \mathbb{Z}_\ell^\times$ . Then we have  $(a, b)_v = 1$ .*

(iii) *Under the assumption on  $v$  in (ii), let  $a = \ell$  and  $b \in \mathbb{Z}_\ell^\times$ . Then we have  $(a, b)_\ell = (\frac{b}{\ell})$ , where  $(\frac{*}{\ell})$  is the Legendre symbol.*

(iv) *If  $v = 2$  and  $a, b \in 1 + 4\mathbb{Z}_2$ , then we have  $(a, b)_2 = 1$ .*

*Proof.* (i): This follows from Proposition 2.1 and the equality  $N_{\mathbb{C}/\mathbb{R}}(\mathbb{C}^\times) = \mathbb{R}_{>0}$ .

(ii): Since  $\ell \neq 2$  and  $a \in \mathbb{Z}_\ell^\times$ , we have

$$\mathbb{Q}_{\ell, a} \cong \begin{cases} \mathbb{Q}_\ell \times \mathbb{Q}_\ell & \text{if } a \in (\mathbb{Z}_\ell^\times)^2, \\ \mathbb{Q}_{\ell^2} & \text{if } a \notin (\mathbb{Z}_\ell^\times)^2. \end{cases}$$

Hence  $N_{\mathbb{Q}_{\ell,a}/\mathbb{Q}_{\ell}}(\mathbb{Q}_{\ell,a}^{\times})$  contains  $\mathbb{Z}_{\ell}^{\times}$ . Hence the assertion follows from Proposition 2.1 and the assumption  $b \in \mathbb{Z}_{\ell}^{\times}$ .

(iii): Since  $\ell$  is not equal to 2, we have

$$N_{\mathbb{Q}_{\ell}(\sqrt{\ell})/\mathbb{Q}}(\mathbb{Q}_{\ell}(\sqrt{\ell})^{\times}) = \langle -\ell \rangle \times (\mathbb{Z}_{\ell}^{\times})^2.$$

Therefore the assertion follows from Proposition 2.1.

(iv): Since  $1 + 8\mathbb{Z}_2 = (\mathbb{Z}_2^{\times})^2$ , there is an isomorphism

$$\mathbb{Q}_{2,a} \cong \begin{cases} \mathbb{Q}_2 \times \mathbb{Q}_2 & \text{if } a \in (\mathbb{Z}_2^{\times})^2, \\ \mathbb{Q}_4 & \text{if } a \notin (\mathbb{Z}_2^{\times})^2. \end{cases}$$

In particular,  $N_{\mathbb{Q}_{2,a}/\mathbb{Q}_{\ell}}(\mathbb{Q}_{2,a}^{\times})$  contains  $\mathbb{Z}_2^{\times}$ . Hence the assertion follows from  $b \in 1 + 4\mathbb{Z}_2$  and Proposition 2.1.  $\blacksquare$

**Proposition 4.3.** *Let  $a, b \in \mathbb{Q}^{\times}$ . Then we have  $(a, b)_0 = 1$  if and only if  $(a, b)_v = 1$  for all places  $v$  of  $\mathbb{Q}$ .*

*Proof.* This is contained in the Albert–Brauer–Hasse–Noether theorem.  $\blacksquare$

**Lemma 4.4.** *Let  $L^+$  be a totally real field which is Galois over  $\mathbb{Q}$ .*

- (i) *Let  $L/L^+$  be a quadratic extension such that  $L/\mathbb{Q}$  is Galois. Then  $L$  is either totally real or CM.*
- (ii) *If there is an  $L$  satisfying (i), then there is a totally imaginary quadratic extension  $L'/L^+$  such that  $L'/\mathbb{Q}$  is Galois and  $\text{Gal}(L'/\mathbb{Q}) \cong \text{Gal}(L/\mathbb{Q})$ .*

*Proof.* (i): If  $L$  is not totally real, then  $L$  is totally complex since  $L/\mathbb{Q}$  is Galois. Fix an embedding  $\iota: L \hookrightarrow \mathbb{C}$ , and let  $c$  be the element of  $\text{Gal}(L/\mathbb{Q})$  induced by the composite of the complex conjugation and  $\iota$ . Then  $c$  is the unique non-trivial element of  $\text{Gal}(L/L^+) \subset \text{Gal}(L/\mathbb{Q})$  since  $L^+$  is totally real. On the other hand, the assumption that  $L^+/\mathbb{Q}$  is totally real implies that  $\text{Gal}(L/L^+)$  is central in  $\text{Gal}(L/\mathbb{Q})$ . Hence  $c$  is contained in the center of  $\text{Gal}(L/\mathbb{Q})$ , which implies that  $L$  is a CM field.

(ii): By (i), we may assume that  $L$  is totally real. Write  $L = L^+(\sqrt{\alpha})$ , where  $\alpha \in (L^+)^{\times}$ , and put  $L' := L^+(\sqrt{-\alpha})$ . Note that  $L'$  corresponds to  $\langle (c_0, 1) \rangle$  under the isomorphism

$$\text{Gal}(L(\sqrt{-1})/\mathbb{Q}) \cong \text{Gal}(L/\mathbb{Q}) \times \text{Gal}(\mathbb{Q}(\sqrt{-1})/\mathbb{Q}) \cong \text{Gal}(L/\mathbb{Q}) \times \mathbb{Z}/2.$$

Here  $c_0$  is the unique non-trivial element of  $\text{Gal}(L/L^+)$ , which is central in  $\text{Gal}(L/\mathbb{Q})$ . Then  $L'$  is not totally real  $L'$  is Galois over  $\mathbb{Q}$ , and hence it is CM by (i). Moreover, the composite

$$\text{Gal}(L/\mathbb{Q}) \xrightarrow{g \mapsto (g, 0)} \text{Gal}(L/\mathbb{Q}) \times \mathbb{Z}/2 \cong \text{Gal}(L(\sqrt{-1})/\mathbb{Q}) \rightarrow \text{Gal}(L'/\mathbb{Q})$$

is an isomorphism by the definition of  $L'$ .  $\blacksquare$

Let  $\Lambda$  be the set of unordered pairs of prime numbers  $\{\ell, \ell'\}$  satisfying

$$\ell \equiv \ell' \equiv 1 \pmod{4}, \quad \left(\frac{\ell'}{\ell}\right) = 1.$$

**Lemma 4.5.** *For any  $n' \in \mathbb{Z}_{>0}$ , there are  $n'$ -pairs  $\{\ell_1, \ell'_1\}, \dots, \{\ell_{n'}, \ell'_{n'}\}$  in  $\Lambda$  such that*

$$\#\{\ell_1, \dots, \ell_{n'}, \ell'_1, \dots, \ell'_{n'}\} = 2n'.$$

*Proof.* By the Dirichlet prime number theorem, there are  $n'$  distinct prime numbers  $\ell_1, \dots, \ell_{n'}$  which are congruent to 1 modulo 4. Moreover, for  $\nu \in \{1, \dots, n'\}$ , the Dirichlet prime number theorem implies the existence of prime numbers  $\ell'_1, \dots, \ell'_{n'}$  satisfying

$$\begin{cases} \ell'_\nu \equiv 1 \pmod{4}, \left(\frac{\ell'_\nu}{\ell'_\nu}\right) = 1 & \text{if } \nu = 1, \\ \ell'_\nu \equiv 1 \pmod{4}, \left(\frac{\ell'_\nu}{\ell'_\nu}\right) = 1, \ell'_\nu \notin \{\ell'_1, \dots, \ell'_{\nu-1}\} & \text{if } \nu \geq 2. \end{cases}$$

Therefore the assertion holds.  $\blacksquare$

For  $\lambda := \{\ell, \ell'\} \in \Lambda$ , put  $L_\lambda^+ := \mathbb{Q}(\sqrt{\ell}, \sqrt{\ell'})$ .

**Lemma 4.6.** *For any  $\lambda \in \Lambda$ , the decomposition groups of  $L_\lambda^+/\mathbb{Q}$  at all finite places are cyclic.*

*Proof.* Write  $\Lambda = \{\ell, \ell'\}$ . Let  $v$  be a finite place of  $L_\lambda^+$  which lies above a prime number  $\ell_0$ . If  $\ell_0 \notin \{\ell, \ell'\}$ , then  $L_\lambda^+/\mathbb{Q}$  is unramified at  $v$ , and hence the assertion holds for  $v$ . The assertion for  $\ell_0 = \ell$  follows from the assumption  $\left(\frac{\ell'}{\ell}\right) = 1$ . Finally, in the case  $\ell_0 = \ell'$ , the statement is a consequence of the equality  $\left(\frac{\ell}{\ell'}\right) = \left(\frac{\ell'}{\ell}\right) = 1$ .  $\blacksquare$

**Proposition 4.7.** *For any  $\lambda \in \Lambda$ , there is a CM field  $L$  containing  $L_\lambda^+$  such that  $L/\mathbb{Q}$  is Galois with Galois group  $Q_8$ .*

*Proof.* By Lemma 4.4, it suffices to prove the existence of  $Q_8$ -extension  $L$  of  $\mathbb{Q}$  containing  $L_\lambda^+$ . Write  $\Lambda = \{\ell, \ell'\}$ . It is equivalent to the equality

$$(\ell, \ell)_0(\ell', \ell')_0(\ell, \ell')_0 = 1,$$

which is a consequence of Theorem 2.3. Moreover, Proposition 4.3 implies that it is sufficient to prove the following.

**Claim.** We have  $(\ell, \ell)_v(\ell', \ell')_v(\ell, \ell')_v = 1$  for all place  $v$  of  $\mathbb{Q}$ .

**Case 1.  $v$  is infinite.** In this case, the assertion follows from Proposition 4.2 since  $\ell, \ell' > 0$ .

**Case 2.  $v$  is a finite place distinct from 2,  $\ell$  and  $\ell'$ .** Since  $v \notin \{\ell, \ell'\}$ ,  $\ell$  and  $\ell'$  are units in  $\mathbb{Q}_v$ . Hence we have

$$(\ell, \ell)_v = (\ell', \ell')_v = (\ell, \ell')_v = 1,$$

which implies the assertion.

**Case 3.  $v = 2$ .** Since  $\ell \equiv \ell' \equiv 1 \pmod{4}$ , Proposition 4.2 (iv) implies

$$(\ell, \ell)_2 = (\ell', \ell')_2 = (\ell, \ell')_2 = 1.$$

Hence the assertion holds.

**Case 4.  $v = \ell$ .** The assumption  $\ell \equiv 1 \pmod{4}$  is equivalent to the equality  $\left(\frac{-1}{\ell}\right) = -1$ . Hence we have

$$(\ell, \ell)_\ell = (\ell, -1)_\ell = 1.$$

On the other hand, the assumption  $\left(\frac{\ell'}{\ell}\right) = 1$  implies

$$(\ell, \ell')_\ell = (\ell', \ell')_\ell = 1.$$

Therefore we obtain the desired equality.

**Case 5.  $v = \ell'$ .** Since  $\ell' \equiv 1 \pmod{4}$  and

$$\left(\frac{\ell}{\ell'}\right) = \left(\frac{\ell'}{\ell}\right) = 1,$$

the assertion follows from the same argument as Case 4.  $\blacksquare$

*Proof of Theorem 4.1.* By Proposition 3.3, we may assume  $n < 0$ . Put  $n' := -n \in \mathbb{Z}_{>0}$ . Take  $\lambda_1 = \{\ell_1, \ell'_1\}, \dots, \lambda_{n'} = \{\ell_{n'}, \ell'_{n'}\} \in \Lambda$  satisfying

$$\#\{\ell_1, \dots, \ell_{n'}, \ell'_1, \dots, \ell'_{n'}\} = 2n'.$$

Note that it is possible by Lemma 4.5. Then, Proposition 4.7 implies that there is a CM field  $L_{\lambda_\nu}$  containing  $L_{\lambda_\nu}^+$  satisfying  $\text{Gal}(L_{\lambda_\nu}/\mathbb{Q}) \cong Q_8$  for any  $\nu \in \{1, \dots, n'\}$ .

**Claim.** The  $n'$ -tuple of CM fields  $L_{\lambda_1}, \dots, L_{\lambda_{n'}}$  satisfies (i)–(iii) in Proposition 3.5.

It is already mentioned that  $L_{\lambda_\nu}$  satisfies (i) for any  $\nu$ . Moreover, since the composite of  $L_{\lambda_1}^+, \dots, L_{\lambda_{n'}}^+$  has degree  $4^{n'}$  over  $\mathbb{Q}$ , the CM fields  $L_{\lambda_1}, \dots, L_{\lambda_{n'}}$  satisfies (ii) by the same argument as the proof of [LYY21, Theorem 7.12]. On the other hand, for (iii), take a finite places  $w$  of  $L_{\lambda_\nu}$  and  $v$  of  $L_{\lambda_\nu}^+$  satisfying  $w \mid v$ . Then Lemma 4.6 implies the cyclicity of the decomposition group of  $L_{\lambda_\nu}^+/\mathbb{Q}$  at  $v$ . Since  $\text{Gal}(L_{\lambda_\nu}/\mathbb{Q}) \cong Q_8$ , (iii) follows from Lemma 2.2.

Since Claim is proved, if we set

$$L := L_{\lambda_1} \times \cdots \times L_{\lambda_{n'}},$$

then we obtain the desired equality  $\tau(T_{L/L^+}) = 2^{-n'} = 2^n$ . ■

#### REFERENCES

- [DMOS82] P. Deligne, J. S. Milne, A. Ogus, K. Shih, *Hodge cycles, Motives and Shimura varieties*, Lect. Notes in Math. 900, Springer, 1982.
- [GS06] P. Pille, T. Szamuely, *Central Simple Algebras and Galois Cohomology*, Cambridge Studies in Advanced Mathematics, Vol. 101, 2006.
- [GSY22] J.-W. Guo, N.-H. Sheu, C.-F. Yu, *Class numbers of CM algebraic tori, CM abelian varieties and components of unitary Shimura varieties*, Nagoya Math. J. **245** (2022), 74–99.
- [Kat85] S. Katayama, *On the Tamagawa number of algebraic tori*, Sūgaku, **37** no. 1 (1985), 81–83.
- [Kim90] I. Kiming, *Explicit classifications of some 2-extensions of a field of characteristic different from 2*, Can. J. Math. **42**, no. 5 (1990), 825–855.
- [LYY21] P.-X. Liang, H.-Y. Yang, C.-F. Yu, *On Tamagawa numbers of CM tori*, preprint, arXiv:2109.04121, 2021, with an appendix by J. Li and C.-F. Yu.
- [Ono63a] T. Ono, *On the Tamagawa number of algebraic tori*, Ann. of Math. (2) **78** (1963), 47–73.
- [Ono63b] T. Ono, *On Tamagawa numbers*, Sūgaku, **15** (1963), 72–81.
- [Rüd20] T. Rüd, *Explicit Tamagawa numbers for certain algebraic tori over number fields*, preprint, arXiv:2009.04431, 2020.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCENCE, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-KU, SAPPORO, HOKKAIDO, 060-0810, JAPAN.

*Email address:* oki@math.sci.hokudai.ac.jp