

UNIFORM IN TIME CONVERGENCE TO BOSE-EINSTEIN CONDENSATION FOR A WEAKLY INTERACTING BOSE GAS WITH EXTERNAL POTENTIALS

CHARLOTTE DIETZE AND JINYEOP LEE

ABSTRACT. We consider a gas of weakly interacting bosons in three dimensions subject to an external potential in the mean field regime. Assuming that the initial state of our system is a product state, we show that in the trace topology of one-body density matrices, the dynamics of the system can be described by the solution to the corresponding Hartree type equation. Using a dispersive estimate for the Hartree type equation, we obtain an error term that is uniform in time. Moreover, the dependence of the error term on the particle number is optimal. We also consider a class of intermediate regimes between the mean field regime and the Gross–Pitaevskii regime, where the error term is uniform in time but not optimal in the number of particles.

1. INTRODUCTION

Bose-Einstein condensation was proposed by Bose and Einstein [5, 25] in 1924. Since then, the topic has gained great interest in both physics and mathematics [4, 43, 56, 59, 76, 77], in particular after the first experimental observation by Wieman, Cornell and Ketterle [1, 18] in 1995.

We consider the dynamics of Bose gas of N particles in three dimensions interacting through a symmetric two-body potential in the presence of an external potential. We assume our initial state $\psi_{N,0} \in L^2(\mathbb{R}^{3N})$ to be fully factorised, i.e.,

$$(1.1) \quad \psi_{N,0}(x_1, \dots, x_N) = \prod_{j=1}^N \varphi_0(x_j)$$

for some $\varphi_0 \in L^2(\mathbb{R}^3)$ with $\|\varphi_0\|_2 = 1$. The time evolution of our state is governed by the Hamiltonian

$$(1.2) \quad H_N = \sum_{j=1}^N (-\Delta_{x_j} + V(x_j)) + \frac{\lambda}{N} \sum_{i < j}^N w_N(x_i - x_j).$$

The time-evolution $\psi_{N,t}$ is the solution to

$$(1.3) \quad \begin{cases} i\partial_t \psi_{N,t} &= H_N \psi_{N,t} \\ \psi_{N,t}|_{t=0} &= \psi_{N,0}. \end{cases}$$

Alternatively, one can write,

$$\psi_{N,t} = e^{-iH_N t} \psi_{N,0}$$

for every $t \in \mathbb{R}$. Here, $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ denotes the external potential and

$$w_N(x) := N^{3\beta} w(N^\beta x) \quad \text{for } 0 \leq \beta < \frac{1}{3}$$

is the interaction potential, where $w \in L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^2)$ is independent of N .¹ We consider weakly interacting bosons, so the coupling constant $\lambda \in \mathbb{R}$ is small in the sense that $|\lambda| \leq \lambda_0$ for some λ_0 depending on φ_0 , $\|w\|_1$ and V . We will give the precise assumptions on the external potential V , on φ_0 , and on λ_0 later.

We prepare a product state that is close to the ‘ground state of an interacting Bose gas in a trap’, which is called the Bose-Einstein condensation state. Then we want to investigate the time evolution of the state after removing the trap and replacing it by a small external potential V .

Since the initial state $\psi_{N,0}$ is a product state, see (1.1), we expect that, in an appropriate sense, the time-evolved state $\psi_{N,t}$ is also approximately given by a product state $\prod_{j=1}^N \varphi_t(x_j)$ for some $\varphi_t \in L^2(\mathbb{R}^3)$, where φ_t is the solution to the corresponding Hartree-type equation (Hartree)

$$(Hartree) \quad \begin{cases} i\partial_t \varphi_t &= (-\Delta + V)\varphi_t + \lambda(w * |\varphi_t|^2)\varphi_t \\ \varphi_t|_{t=0} &= \varphi_0, \end{cases}$$

for $\beta = 0$, and to the nonlinear Schrödinger equation (NLS)

$$(NLS) \quad \begin{cases} i\partial_t \varphi_t &= (-\Delta + V)\varphi_t + \lambda a |\varphi_t|^2 \varphi_t \\ \varphi_t|_{t=0} &= \varphi_0, \end{cases}$$

for $\beta \in (0, 1/3)$, where $a = \int_{\mathbb{R}^3} dx w(x)$.

We cannot expect $\prod_{j=1}^N \varphi_t(x_j)$ and $\psi_{N,t}$ to be close in the L^2 -norm sense, see [58]. However, we can prove that they are close in the trace topology of one-body density matrices as $N \rightarrow \infty$. More precisely, we define the marginal one-particle density $\gamma_{N,t}^{(1)}$ by its operator kernel

$$(1.4) \quad \gamma_{N,t}^{(1)}(x; y) = \int_{\mathbb{R}^{3(N-1)}} dZ \overline{\psi_{N,t}(y, Z)} \psi_{N,t}(x, Z).$$

Our main result is the following.

Theorem 1.1. *Let $0 \leq \beta < 1/3$ and let $N \in \mathbb{N}$. Suppose that $w \in L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ and*

$$w_N(x) := N^{3\beta} w(Nx).$$

Let $V \in W^{2,\infty}(\mathbb{R}^3)$ be real-valued and such that there exists a constant $C^V > 0$ such that

$$\|e^{it(-\Delta+V)} f\|_\infty \leq C^V \|f\|_1$$

for all $f \in L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$. Let $\varphi_0 \in H^2(\mathbb{R}^3)$ with $\|\varphi_0\|_2 = 1$. We let our system have fully factorized initial data, i.e.,

$$\psi_{N,0}(\mathbf{x}) = \prod_{j=1}^N \varphi_0(x_j)$$

for $\mathbf{x} = (x_1, x_2, \dots, x_N) \in \mathbb{R}^{3N}$. Let

$$\psi_{N,t} = e^{-iH_N t} \psi_{N,0}$$

for $t > 0$ where H_N is defined in (1.2). Here, we assume that $|\lambda| \leq \lambda_0$ with $0 < \lambda_0 \leq 1$ depending only on $\|V\|_{W^{2,\infty}}$, C^V , $\|w\|_1$, $\|\varphi_0\|_{H^2}$, and $\|e^{i(-\Delta+V)} \varphi_0\|_1$. Then there exists a constant $C > 0$ depending only on β , $\|V\|_{W^{2,\infty}}$, C^V , $\|w\|_1$, and $\|w\|_2$, such that

$$(1.5) \quad \text{Tr} \left| \gamma_{N,t}^{(1)} - |\varphi_t\rangle\langle\varphi_t| \right| \leq \begin{cases} CN^{-1} & \text{if } \beta = 0, \\ CN^{(3\beta-1)/2} & \text{if } 0 < \beta < 1/3 \end{cases}$$

¹The choice $\beta = 0$ is called mean-field regime, and $\beta = 1$ is called Gross-Pitaevskii regime. Moreover, $\beta \in (0, 1)$ is called intermediate regime between the mean-field regime and the Gross-Pitaevskii regime.

for all $t > 0$. In particular, the constant C does not depend on t or N .

Remarks 1.1.

- (1) A special case of Theorem 1.1, namely in the mean field regime and without external potential, i.e. $\beta = 0$ and $V = 0$, was proved in [51, Theorem 1.3]. Theorem 1.1 is new because we can also treat external potentials.
- (2) In many occasions, relevant physical quantities can be obtained from the one-particle density matrix, so in these occasions we can obtain a good approximation by replacing the one particle density $\gamma_{N,t}^{(1)}$ by $|\varphi_t\rangle\langle\varphi_t|$.
- (3) A similar theorem can be obtained for general λ (without smallness assumption) with time dependent constant C .
- (4) The paper covers the mean field regime ($\beta = 0$) and a part of the intermediate regime, namely $0 < \beta < 1/3$. The next goal would be to consider the remaining part of the intermediate regime ($1/3 \leq \beta < 1$) and the final goal would be the Gross-Pitaevskii (GP) regime ($\beta = 1$).
- (5) Note that the scaling in N conserves the L^1 -norm of w , i.e.

$$\|w\|_{L^1(\mathbb{R}^3)} = \|w_N\|_{L^1(\mathbb{R}^3)}$$

but not the L^2 -norm

$$\|w_N\|_{L^2(\mathbb{R}^3)} = N^{3\beta/2} \|w\|_{L^2(\mathbb{R}^3)}.$$

Since we use some estimates in terms of $\|w_N\|_{L^2(\mathbb{R}^3)}$ in our proof, the rate of convergence in Theorem 1.1 for $\beta > 0$ is drastically different than for $\beta = 0$.

Our proof is based on [12, 13, 80] we derive the mean-field limit with interaction potential $w \in L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ under the existence of external potential V . We will explain the details later in Section 1.2.

We track dependence of $\|\varphi_t\|_\infty$, which helps us get a better estimate after using the Grönwall inequality. Applying the time decay estimate obtained in [21], we obtain a time independent rate of convergence.

1.1. History.

Many-Body Convergence. Now we are going to review the results on the derivation of the effective one-body Schrödinger equation from the many-body Schrödinger equation. To be more precise, a typical result in this direction is that, if one has an initial many-body wave function $\psi_{N,0}$ with

$$(1.6) \quad \lim_{N \rightarrow \infty} \text{Tr} \left| \frac{1}{N} \gamma_{N,0} - |\varphi_0\rangle\langle\varphi_0| \right| = 0$$

for some given $\varphi_0 \in L^2(\mathbb{R}^3)$, then at later times $t > 0$,

$$(1.7) \quad \lim_{N \rightarrow \infty} \text{Tr} \left| \frac{1}{N} \gamma_{N,t} - |\varphi_t\rangle\langle\varphi_t| \right| = 0,$$

where φ_t is a solution of an effective partial differential equation.

For the mean-field regime, i.e., $\beta = 0$, with differentiable interaction potential w , Hepp [46] showed (1.7). Moreover, Spohn [81] obtained (1.7) for a bounded interaction potential w . Ginibre and Velo provided a series of works in this direction [33, 34, 38].

Later, for singular potentials, the derivation of (1.7) was obtained by Bardos, Golse, and Mauer [2] for $L^2(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$ potentials in one dimension, and by Erdős and Yau [30] for the Coulomb potential in three dimensions. These works are based on the BBGKY² hierarchy method. Therefore, the rate of convergence

²Bogoliubov–Born–Green–Kirkwood–Yvon

in these works was not given explicitly. Moreover, the convergence is only established for any time of order 1.

For factorized initial states, Rodnianski and Schlein developed a scheme to obtain the rate of convergence in [80] and they showed the rate

$$(1.8) \quad \text{Tr} \left| \frac{1}{N} \gamma_{N,0} - |\varphi\rangle\langle\varphi| \right| \leq \frac{Ce^{Ct}}{\sqrt{N}}$$

for interaction potentials w including the Coulomb interaction potential. Knowles and Pickl [49] provided a scheme to cover more singular potentials than Coulomb with the same rate based on the techniques in [74].

For bounded and integrable interaction potentials, the optimal rate in N was provided in [27]. This work was extended in [11]. In [13], Chen, Lee, and Schlein provided the rate Ce^{Ct}/N , which is optimal in N , for the Coulomb interaction potential. This work was generalized in [12] to more singular interaction potentials using Strichartz estimates for the Hartree equation. The time dependence was investigated in [51].

Elgart, Schlein [26] and Michelangeli, Schlein [62] proved convergence for the mean-field Boson star equation, i.e.,

$$H_N = \sum_{j=1}^N (1 - \Delta_{x_j})^{1/2} + \frac{\lambda}{N} \sum_{i<j}^N \frac{1}{x_i - x_j}.$$

Later Lee provided the rate of convergence [55].

For the intermediate regime and GP regime $0 < \beta \leq 1$, Erdős, Yau, and Schlein proved (1.7) in [28, 29]. Moreover, for the GP regime $\beta = 1$, Benedikter, de Oliveira and Schlein [3], Pickl [75], and Brennecke and Schlein [6] provided the convergence rate.

It is important to have a time independent bound to show the stability of Bose-Einstein condensation. There have been a few works in the direction of improving time dependence. In [42], the authors provided a norm approximation with a bound of the form $(1+t)^{1/2}/N$ for a cut-off Coulomb interaction potential. In [50], for $\beta = 0$ with a cut-off Coulomb potential, the trace norm was bounded by $(1+t)/N$. For $\beta \in (0, 1/3)$ with an integrable interaction function w , the trace norm was bounded by $(1+t)^2 \log^{16}(1+t)/N^\epsilon$ for some small parameter ϵ depending on β . In [51], for interaction potential in the form of $\lambda e^{-\mu|x|}|x|^{-\gamma}$ without cut-off, trace norm bounds have been proved. Here, in this paper, we provide time independent bound of the trace norm by using the time decay estimate for one-body nonlinear Schrödinger equations.

The second order correction was obtained by a norm approximation by a series of works by Grillakis, Machedon and Margetis [39–42]. Pickl [73] also considered this regime. Recently, Napiórkowski provided a nice review paper about this topic [69]. There are many directions to investigate further. For example, there are at least two directions. Regarding mixture condensation, one can find results in [19, 22, 53, 60, 61]. Results on three-body interactions can be found in [14, 15, 52, 67] and in the references therein. In [13], they mentioned in a remark that their approach could cover external potentials. This was tracked in [54].

There are also several results [39–42, 58, 64, 65] on a norm approximation, which is much more precise than the approximation of the one particle marginal density in trace norm. As we have already mentioned earlier, it is *not* possible to show that $\psi_{N,t}$ is close to a product state in the norm sense, and instead, one has to take excitations into account. This can be done by using Bogoliubov theory, which requires a more detailed analysis of the Hamiltonian in Fock space than the method, which is used in this paper. Other results including external potentials are [7, 8, 23, 57, 66, 68].

Since there is a correlation among particles, it might not be natural to expect a time independent bound for the norm approximation. Hence, in this paper, we provide the convergence in terms of a trace norm approximation of 1-particle density matrices instead of a norm approximation of wave functions.

Time decay Estimate for one-body nonlinear Schrödinger equations. Let us first review several results on one-body nonlinear Schrödinger equations without external potentials. Ginibre and Velo [35, Theorem 6.1(1)] considered Hartree type nonlinearities with repulsive interaction potentials w and they proved the decay estimate

$$(1.9) \quad \|\varphi_t\|_q \leq C(1 + |t|)^{-\frac{d}{2}\left(\frac{1}{q'} - \frac{1}{q}\right)} \text{ for all } t \neq 0$$

for all q with $\left[\frac{1}{2} - \frac{1}{d}\right]_+ \leq \frac{1}{q} \leq \frac{1}{2}$, where $[\cdot]_+$ denotes the positive part. Note that $\left(\frac{1}{2} - \frac{1}{d}\right)^{-1} = \left(\frac{d-2}{2d}\right)^{-1} = \frac{2d}{d-2} = 2^*$, so their result corresponds to the energy critical and energy subcritical case. In particular, the case $q = \infty$ is not covered in dimension $d = 3$. Hayashi and Naumkin [44] looked at critical nonlinearities, namely the local nonlinearity

$$(1.10) \quad \lambda|\varphi_t|^{\frac{2}{d}}\varphi_t \text{ for } \lambda \in \mathbb{R} \text{ and } d \in \{1, 2, 3\}$$

and the nonlocal nonlinearity

$$(1.11) \quad \lambda(|\cdot|^{-1} * |\varphi_t|^2)\varphi_t \text{ for } \lambda \in \mathbb{R} \text{ and } d \geq 2$$

and they can also add non-critical nonlinearities. They showed a dispersive estimate of the form

$$(1.12) \quad \|\varphi_t\|_\infty \leq C(1 + |t|)^{-\frac{d}{2}} \text{ for all } t \neq 0.$$

Later Kato and Pusateri [48] gave an alternative proof of the result in [44], which was based on a careful analysis of the equation in Fourier space.

Grillakis and Machedon [39, Corollary 3.4] considered Hartree type nonlinearities with nonnegative, radial, decreasing interaction potential $w \in L^1(\mathbb{R}^d) \cap C_0^1(\mathbb{R}^d)$ for sufficiently regular but possibly large initial data. They showed a decay estimate of the form (1.12) using a Grönwall type argument after proving suitable a-priority estimates. Their result was applied in [65], where they showed a norm approximation for the dynamics of many-body quantum systems in the context of Bose-Einstein condensation.

Other results without external potentials, we would like to mention are [10, 16, 20, 24, 32, 36, 37, 45, 72].

Next, let us turn to results on nonlinear Schrödinger equations with external potentials. In dimension $d = 1$, Cuccagna, Georgiev and Visciglia [17] proved a decay estimate of the form (1.12) for subcritical nonlinearities $\pm|\varphi_t|^{p-1}\varphi_t$ with $3 < p < 5$ and small initial data. The corresponding result for the critical nonlinearities $\pm|\varphi_t|^{p-1}\varphi_t$ with $p = 3$ was proved by Germain, Pusateri and Rousset [31]. Their proof relies on the use of the distorted Fourier transform and the analysis of an oscillatory integral. A similar result was proved by Naumkin [70, 71]. In dimension $d = 3$, Pusateri and Soffer [78] considered the nonlinearity $-u^2$ and small initial data. They showed that

$$(1.13) \quad \|\varphi_t\|_\infty \leq C(1 + |t|)^{-(1+\alpha)} \text{ for all } t \neq 0$$

for some $\alpha > 0$. Two other results in dimension $d = 3$ with external potentials, which we would like to mention, are scattering in H^1 for an external potential V with small negative part by Hong [47], and a classification of the dynamics of solutions to the cubic nonlinear Schrödinger equation with small initial data and a radial external potential V , which is such that the operator $-\Delta + V$ has exactly one negative eigenvalue, by Nakanishi [63].

1.2. Strategy of the Proof. In order to prove the main result, we will need good control on the time evolution. To this end, we will define a truncated dynamics in Section 4, for which we have good bounds. In Section 5, we then compare the dynamics with its truncated version, for which we deduce us good bounds on the dynamics. Moreover, we divide the dynamics into parity-conserving and non-parity-conserving parts. In order to obtain these bounds, we estimate the time-derivative and then use the Grönwall lemma. Since the

estimate for the time derivative contains a factor of $\|\varphi_t\|_\infty$ and by Proposition 2.1, we know from (2.4) that $\|\varphi_t\|_\infty \in L^1(\mathbb{R})$. Thus the bound we get after applying the Grönwall lemma will be the time independent.

1.3. Structure of the Paper. In Section 2, we review the necessary preliminaries. In particular, we review the result on time decay estimate with external potential, see 2.1, and the Fock Space formalism for the many-body problem. In Section 3, we prove Theorem 1.1 assuming Propositions 3.1 and 3.2. Section 5 is devoted to proving Propositions 3.1 and 3.2. To this end, we show several results on comparison dynamics. In Section 4, we introduce the truncated dynamics and prove a bound for it. Section 5 is devoted to proving Propositions 3.1 and 3.2. To this end, we show several results on comparison dynamics.

1.4. Notation.

- (1) The L^p -norm in \mathbb{R}^3 can be denoted by $\|\cdot\|_{L^p(\mathbb{R}^3)}$ or $\|\cdot\|_p$.
- (2) Sometimes, when there can be no confusion, we omit the subscript for norms, for example,

$$\|\cdot\|_{L^2(\mathbb{R}^3)}, \|\cdot\|_{\mathcal{F}}, \text{ or } \langle \cdot, \cdot \rangle_{\mathcal{F}}$$

are just written as

$$\|\cdot\|, \|\cdot\|, \text{ or } \langle \cdot, \cdot \rangle,$$

respectively.

- (3) Our constant C may change from line to line.
- (4) The indicator function is denoted by χ , for example

$$\chi(N \leq M) = \begin{cases} 1 & \text{if } N \leq M, \\ 0 & \text{if } N > M. \end{cases}$$

Acknowledgements. The authors would like to express their deepest gratitude to Phan Thành Nam for his continued support and very helpful discussions. The authors also thank Arnaud Triay and Alessandro Michelangeli for helpful discussions. Partial support from the Deutsche Forschungsgemeinschaft (DFG project Nr. 426365943) and Istituto Nazionale di Alta Matematica “F. Severi,” through the Intensive Period “INdAM Quantum Meetings (IQM22)” is acknowledged.

2. PRELIMINARIES

2.1. Time Decay Estimate under the Existence of External Potential. This subsection is devoted to reviewing the time decay estimate under the existence of external potential.

Proposition 2.1 (Theorem 1.1. of [21], Dispersive estimates for the Hartree type equation in $d \geq 3$ for small initial data). *Let $d \geq 3$ and let $k \in \mathbb{N}$ be the smallest even number with $k > \frac{d}{2}$. Let $V \in W^{k,\infty}(\mathbb{R}^d)$ be a real-valued function and satisfy*

$$(2.1) \quad \|e^{-it(-\Delta+V)} f\|_\infty \leq C^V |t|^{-\frac{d}{2}} \|f\|_1$$

for every $f \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ and some constant $C^V \geq 1$. Let the interaction potential $w \in L^1(\mathbb{R}^d) \cap L^{\frac{d}{2}}(\mathbb{R}^d)$ be an even, real-valued function. Let $u_0 \in H^k(\mathbb{R}^d)$ and let $u \in C(\mathbb{R}, H^2(\mathbb{R}^d)) \cap C^1(\mathbb{R}, H^{-1}(\mathbb{R}^d))$ be the unique global strong solution to the Hartree type equation

$$(2.2) \quad \begin{cases} i\partial_t u &= (-\Delta + V)u + (w * |u|^2)u \\ u(0) &= u_0. \end{cases}$$

Assume that the initial data is sufficiently small, that is,

$$(2.3) \quad \|e^{i(-\Delta+V)}u_0\|_1, \|u_0\|_{H^k} \leq \epsilon_0$$

for some $\epsilon_0 = \epsilon_0(d, \|V\|_{W^{k,\infty}}, C^V, \|w\|_1) > 0$. Then there exists a constant $C_0 = C_0(d, \|V\|_{W^{k,\infty}}, C^V, \|w\|_1) \geq 1$ such that

$$(2.4) \quad \|u(t)\|_\infty \leq \frac{C_0}{(1+|t|)^{\frac{d}{2}}}$$

for all $t \geq 0$. Furthermore, if we assume that

$$(2.5) \quad \|e^{i(-\Delta+V)}(\partial_t u)(0)\|_1, \|(\partial_t u)(0)\|_{H^k} \leq \tilde{\epsilon}_0$$

for some $\tilde{\epsilon}_0 = \tilde{\epsilon}_0(d, \|V\|_{W^{k,\infty}}, C^V, \|w\|_1) > 0$, then

$$(2.6) \quad \|\partial_t u(t)\|_\infty \leq \frac{\tilde{C}_0}{(1+|t|)^{\frac{d}{2}}}$$

for all $t \geq 0$, where $\tilde{C}_0 = \tilde{C}_0(d, \|V\|_{W^{k,\infty}}, C^V, \|w\|_1) > 0$.

Remarks 2.1.

(1) Let $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ and assume that

$$(2.7) \quad \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy \frac{|V(x)||V(y)|}{|x-y|^2} < (4\pi)^2$$

and

$$(2.8) \quad \sup_{x \in \mathbb{R}^3} \int_{\mathbb{R}^3} dy \frac{|V(y)|}{|x-y|} < 4\pi.$$

Then [79, Theorem 1.1] there exists a constant $C^V > 0$ depending only on V such that

$$(2.9) \quad \|e^{-itH}f\|_\infty \leq C^V |t|^{-\frac{3}{2}} \|f\|_1 \text{ for all } f \in L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3), t \in \mathbb{R} \setminus \{0\}.$$

(2) Proposition 2.1 remains true for any global solution to the cubic non-linear Schrödinger equation:

We may replace the interaction potential by $w = \pm \delta_0$, that is, the nonlinearity is given by $\pm |u|^2 u$.

Idea of the proof. The proof uses ideas from Grillakis and Machedon [39, Corollary 3.4] and Kato and Pusateri [48]. We only describe the proof strategy for

$$\|u(t)\|_\infty \leq \frac{C_0}{(1+|t|)^{\frac{d}{2}}} \text{ for all } t \geq 0$$

as the proof of the corresponding estimate for $\|\partial_t u(t)\|_\infty$ is similar. Define

$$M(T) := \sup_{0 \leq t \leq T} (1+|t|)^{\frac{d}{2}} \|u(t)\|_\infty + \sup_{0 \leq t \leq T} \|D^k u(t)\|_2 + \|u_0\|_2,$$

and note that it suffices to show

$$M(T) \leq C_0 \text{ for every } T \geq 0$$

for some $C_0 > 0$ to be determined. Using Duhamel's formula and the dispersive estimate

$$\|e^{-it(-\Delta+V)}f\|_\infty \leq C^V |t|^{-\frac{d}{2}} \|f\|_1 \text{ for all } f \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$$

for the operator $-\Delta + V$, which was part of our assumptions on the external potential V , we obtain after a computation

$$M(T) \leq \epsilon + CM(T)^3,$$

where $\epsilon, C > 0$ do not depend on T and ϵ is small if the initial data is small. If $\epsilon > 0$ is small enough, then the graph of the function

$$f : [0, \infty) \rightarrow \mathbb{R}, \quad f(x) := \epsilon + Cx^3 - x$$

has two distinct non-negative zeros, see Figure 2.1




FIGURE 2.1. Graph of f

Denote by C_0 the first zero of f . By a blow-up criterion for H^k -solutions similar to [9, Theorem 4.10.1], there exists $T_{\max} \in (0, \infty]$ such that $M \in C([0, T_{\max}))$. Furthermore, if $T_{\max} < \infty$, then $\lim_{T \uparrow T_{\max}} M(T) = \infty$. If $M(0) \leq C_0$, then by the smallness of the initial data, we deduce that $T_{\max} = \infty$ and

$$M(T) \leq C_0 \text{ for every } T \geq 0.$$

□

Remark 2.2. By writing $u = \alpha v$ for some $\alpha > 0$ to be chosen, (2.2) becomes

$$(2.10) \quad \begin{cases} i\partial_t v &= (-\Delta + V)v + \alpha^2(w * |v|^2)v \\ v(0) &= \frac{1}{\alpha}u_0 =: v_0. \end{cases}$$

If α is small, this is an equation with weak coupling for large initial data. Hence, the results in Proposition 2.1 can be converted to the corresponding results for weakly interacting Hartree/NLS equations with large initial data.

2.2. Fock Space. To investigate the dynamics of Bose gas with N particles, we embed our problem into a bigger space, so-called the Fock space. We briefly review the well-known Fock space formalism. The bosonic Fock space is a Hilbert space given by

$$\mathcal{F} = \bigoplus_{n \geq 0} L^2(\mathbb{R}^3)^{\otimes_s n},$$

where $L^2(\mathbb{R}^3)^{\otimes_s n}$ is the symmetric subspace of $L^2(\mathbb{R}^{3n})$, that is, such that, for

$$g(x_1, \dots, x_n) = g(x_{\sigma(1)}, \dots, x_{\sigma(n)})$$

for all $g(x_1, \dots, x_n) \in L^2(\mathbb{R}^3)^{\otimes_s n}$ and all $\sigma \in S_n$, where S_n denotes the set of all permutations of $\{1, \dots, n\}$. We denote an element (or state) $\psi \in \mathcal{F}$ by

$$\psi = \psi^{(0)} \oplus \psi^{(1)} \oplus \psi^{(2)} \oplus \dots = \left(\psi^{(n)} \right)_{n \geq 0}$$

where $L^2(\mathbb{R}^3)^{\otimes_s n}$ for all $n \geq 0$. The inner product on \mathcal{F} is defined by

$$\begin{aligned} \langle \psi_1, \psi_2 \rangle &:= \sum_{n \geq 0} \langle \psi_1^{(n)}, \psi_2^{(n)} \rangle_{L^2(\mathbb{R}^{3n})} \\ (2.11) \quad &= \overline{\psi_1^{(0)}} \psi_2^{(0)} + \sum_{n \geq 1} \int_{\mathbb{R}^{3n}} dx_1 \dots dx_n \overline{\psi_1^{(n)}(x_1, \dots, x_n)} \psi_2^{(n)}(x_1, \dots, x_n). \end{aligned}$$

For $f \in L^2(\mathbb{R}^3)$, the creation operator $a^*(f)$ and the annihilation operator $a(f)$ on \mathcal{F} are defined by

$$(2.12) \quad (a^*(f)\psi)^{(n)}(x_1, \dots, x_n) = \frac{1}{\sqrt{n}} \sum_{j=1}^n f(x_j) \psi^{(n-1)}(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n)$$

and

$$(2.13) \quad (a(f)\psi)^{(n)}(x_1, \dots, x_n) = \sqrt{n+1} \int_{\mathbb{R}^3} dx \overline{f(x)} \psi^{(n+1)}(x, x_1, \dots, x_n).$$

Define

$$(2.14) \quad \phi(f) := a^*(f) + a(f).$$

Note that $\phi(f)$ is self-adjoint, while the creation operator $a^*(f)$ and the annihilation operator $a(f)$ are not self-adjoint. We also use operator-valued distributions a_x^* and a_x satisfying

$$(2.15) \quad a^*(f) = \int_{\mathbb{R}^3} dx f(x) a_x^*, \quad a(f) = \int_{\mathbb{R}^3} dx \overline{f(x)} a_x$$

for any $f \in L^2(\mathbb{R}^3)$. The annihilation and creation operator satisfy the canonical commutation relation (CCR), which are given by

$$[a(f), a^*(g)] = \langle f, g \rangle_{L^2(\mathbb{R}^3)}, \quad [a(f), a(g)] = [a^*(f), a^*(g)] = 0,$$

and

$$[a_x, a_y^*] = \delta(x - y), \quad [a_x, a_y] = [a_x^*, a_y^*] = 0.$$

The number operator \mathcal{N} is defined by

$$(2.16) \quad \mathcal{N} := \int_{\mathbb{R}^3} dx a_x^* a_x$$

and it satisfies that $(\mathcal{N}\psi)^{(n)} = n\psi^{(n)}$. The domain of the number operator \mathcal{N} is given by

$$(2.17) \quad D(\mathcal{N}) = \left\{ \psi \in \mathcal{F} : \sum_{n \geq 1} n^2 \|\psi^{(n)}\|^2 < \infty \right\}$$

For any bounded operator $J : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$, its second quantization $d\Gamma(J) : D(\mathcal{N}) \rightarrow \mathcal{F}$ is given by

$$(d\Gamma(J)\psi)^{(n)} = \sum_{j=1}^n J_j \psi^{(n)}$$

where $J_j = 1 \otimes \dots \otimes J \otimes \dots \otimes 1$ is the operator J acting only on the j -th variable. For any compact operator $J : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$ with kernel $J(x; y)$, we can write its second quantization $d\Gamma(J)$ in terms of operator-valued distributions as

$$(2.18) \quad d\Gamma(J) = \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy J(x; y) a_x^* a_y.$$

Intuitively speaking, both creation and annihilation operators behave like $\mathcal{N}^{1/2}$ as the following lemma shows:

Lemma 2.1 ([80, Lemma 2.1]). *For $\alpha > 0$, let $D(\mathcal{N}^\alpha) = \{\psi \in \mathcal{F} : \sum_{n \geq 1} n^{2\alpha} \|\psi^{(n)}\|^2 < \infty\}$ denote the domain of the operator \mathcal{N}^α . For any $f \in L^2(\mathbb{R}^3)$ and any $\psi \in D(\mathcal{N}^{1/2})$, we have*

$$(2.19) \quad \begin{aligned} \|a(f)\psi\| &\leq \|f\| \|\mathcal{N}^{1/2}\psi\|, \\ \|a^*(f)\psi\| &\leq \|f\| \|(\mathcal{N} + 1)^{1/2}\psi\|, \\ \|\phi(f)\psi\| &\leq 2\|f\| \|(\mathcal{N} + 1)^{1/2}\psi\|. \end{aligned}$$

Moreover, for any bounded one-particle operator J on $L^2(\mathbb{R}^3)$ and for every $\psi \in D(\mathcal{N})$, we find

$$(2.20) \quad \|d\Gamma(J)\psi\| \leq \|J\| \|\mathcal{N}\psi\|.$$

For $f \in L^2(\mathbb{R}^3)$, the Weyl operator $W(f)$ is defined by

$$W(f) := \exp(a^*(f) - a(f)),$$

and it satisfies

$$W(f) = e^{-\|f\|^2/2} \exp(a^*(f)) \exp(-a(f)).$$

Coherent states $\psi(f)$ can be defined in terms of the Weyl operator as

$$(2.21) \quad \psi(f) := W(f)\Omega = e^{-\|f\|^2/2} \exp(a^*(f))\Omega = e^{-\|f\|^2/2} \sum_{n \geq 0} \frac{1}{\sqrt{n!}} f^{\otimes n}.$$

The following lemmata indicate useful properties of the Weyl operator and coherent states:

Lemma 2.2 ([80, Lemma 2.2]). *Let $f, g \in L^2(\mathbb{R}^3)$.*

(1) *The commutation relation between the Weyl operators is given by*

$$W(f)W(g) = W(g)W(f)e^{-2i \cdot \text{Im}\langle f, g \rangle} = W(f + g)e^{-i \cdot \text{Im}\langle f, g \rangle}.$$

(2) *The Weyl operator is unitary and satisfies*

$$W(f)^* = W(f)^{-1} = W(-f).$$

(3) *The coherent states are eigenvectors of annihilation operators, i.e.,*

$$a_x \psi(f) = f(x) \psi(f) \quad \rightarrow \quad a(g) \psi(f) = \langle g, f \rangle_{L^2} \psi(f).$$

(4) *The commutation relation between the Weyl operator and the annihilation operator (or the creation operator) is*

$$W^*(f) a_x W(f) = a_x + f(x) \quad \text{and} \quad W^*(f) a_x^* W(f) = a_x^* + \overline{f(x)}.$$

(5) *The distribution of \mathcal{N} with respect to the coherent state $\psi(f)$ is Poisson. In particular,*

$$\langle \psi(f), \mathcal{N} \psi(f) \rangle = \|f\|^2, \quad \langle \psi(f), \mathcal{N}^2 \psi(f) \rangle - \langle \psi(f), \mathcal{N} \psi(f) \rangle^2 = \|f\|^2.$$

Let

$$(2.22) \quad d_N := \frac{\sqrt{N!}}{N^{N/2}e^{-N/2}}.$$

Note that $C^{-1}N^{1/4} \leq d_N \leq CN^{1/4}$ for some constant $C > 0$ independent of N , which can be easily checked by using Stirling's formula.

Lemma 2.3 ([11, Lemma 6.3]). *There exists a constant $C > 0$ independent of N such that, for any $\varphi_0 \in L^2(\mathbb{R}^3)$ with $\|\varphi_0\| = 1$, we have*

$$\left\| (\mathcal{N} + 1)^{-1/2} W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \leq \frac{C}{d_N}.$$

Lemma 2.4 ([55, Lemma 7.2]). *Let P_m be the projection onto the m -particle sector of the Fock space \mathcal{F} for a non-negative integer m . Then, for any non-negative integer $k \leq (1/2)N^{1/3}$ and any $\varphi_0 \in L^2(\mathbb{R}^3)$ with $\|\varphi_0\| = 1$,*

$$\left\| P_{2k} W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \leq \frac{2}{d_N}$$

and

$$\left\| P_{2k+1} W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \leq \frac{2(k+1)^{3/2}}{d_N \sqrt{N}}.$$

3. PROOF OF THE MAIN RESULT

To consider the problem in the Fock space formalism, we extend the Hamiltonian H_N in (1.2) to the Fock space by

$$(3.1) \quad \mathcal{H}_N := \int_{\mathbb{R}^3} dx a_x^* (-\Delta_x + V(x)) a_x + \frac{\lambda}{2N} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) a_x^* a_y^* a_y a_x.$$

With this definition, we have $(\mathcal{H}_N \psi)^{(N)} = H_N \psi^{(N)}$ for any $\psi \in \mathcal{F}$. The kernel of the one-particle marginal density $\gamma_\psi^{(1)}$ associated with ψ is

$$(3.2) \quad \gamma_\psi^{(1)}(x; y) = \frac{1}{\langle \psi, \mathcal{N} \psi \rangle} \langle \psi, a_y^* a_x \psi \rangle.$$

Note that $\gamma_\psi^{(1)}$ is a trace class operator on $L^2(\mathbb{R}^3)$ and $\text{Tr} \gamma_\psi^{(1)} = 1$. (3.2) is equivalent to (1.4) in the sense that $\gamma_\psi^{(1)} = \gamma_{N,t}^{(1)}$ for every $\psi = (1_{n=N} \psi_{N,t})_{n \geq 0} \in \mathcal{F}$ and $\psi_{N,t} \in L^2(\mathbb{R}^3)^{\otimes_s N}$.

Let $\gamma_{N,t}^{(1)}(x; y)$ for $x, y \in \mathbb{R}^3$ be the kernel of the one-particle marginal density associated with the time evolution of the factorized state $\varphi_0^{\otimes N}$ with respect to the Hamiltonian \mathcal{H}_N . By the equivalence of (3.2) and (1.4),

$$(3.3) \quad \begin{aligned} \gamma_{N,t}^{(1)} &= \frac{\langle e^{-i\mathcal{H}_N t} \varphi_0^{\otimes N}, a_y^* a_x e^{-i\mathcal{H}_N t} \varphi_0^{\otimes N} \rangle}{\langle e^{-i\mathcal{H}_N t} \varphi_0^{\otimes N}, \mathcal{N} e^{-i\mathcal{H}_N t} \varphi_0^{\otimes N} \rangle} = \frac{1}{N} \langle \varphi_0^{\otimes N}, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} \varphi_0^{\otimes N} \rangle \\ &= \frac{1}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\rangle. \end{aligned}$$

If we use the coherent states $W^*(\sqrt{N}\varphi_s)\Omega$ instead of the factorized state $\varphi_0^{\otimes N}$ in (3.3) and expand $a_y^*a_x$ around $N\overline{\varphi_t(y)}\varphi_t(x)$, then we are lead to consider the operator

$$(3.4) \quad \begin{aligned} & W^*(\sqrt{N}\varphi_s)e^{i\mathcal{H}_N(t-s)}(a_x - \sqrt{N}\varphi_t(x))e^{-i\mathcal{H}_N(t-s)}W(\sqrt{N}\varphi_s) \\ & = W^*(\sqrt{N}\varphi_s)e^{i\mathcal{H}_N(t-s)}W(\sqrt{N}\varphi_t)a_xW^*(\sqrt{N}\varphi_t)e^{-i\mathcal{H}_N(t-s)}W(\sqrt{N}\varphi_s). \end{aligned}$$

After a computation, we obtain

$$(3.5) \quad i\partial_t W^*(\sqrt{N}\varphi_t)e^{-i\mathcal{H}_N(t-s)}W(\sqrt{N}\varphi_s) =: \left(\sum_{k=0}^4 \mathcal{L}_k(t; s) \right) W^*(\sqrt{N}\varphi_t)e^{-i\mathcal{H}_N(t-s)}W(\sqrt{N}\varphi_s),$$

where \mathcal{L}_k is an operator with k creation and/or annihilation operators. The exact formulas for \mathcal{L}_k are as follows:

$$(3.6) \quad \mathcal{L}_0(t; s) := \frac{N\lambda}{2} \int_s^t d\tau \int_{\mathbb{R}^3} dx (w_N * |\varphi_\tau|^2)(x) |\varphi_\tau(x)|^2,$$

$$(3.7) \quad \mathcal{L}_1(t; s) := \mathcal{L}_1 = 0,$$

$$(3.8) \quad \begin{aligned} \mathcal{L}_2(t; s) := \mathcal{L}_2(t) &= \int_{\mathbb{R}^3} dx a_x^* (-\Delta_x + V(x))a_x + \lambda \int_{\mathbb{R}^3} dx (w_N * |\varphi_t|^2)(x) a_x^* a_x \\ &+ \lambda \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \overline{\varphi_t(x)} \varphi_t(y) a_y^* a_x \\ &+ \frac{\lambda}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) (\varphi_t(x) \varphi_t(y) a_x^* a_y^* + \overline{\varphi_t(x)} \overline{\varphi_t(y)} a_x a_y), \end{aligned}$$

$$(3.9) \quad \mathcal{L}_3(t; s) := \mathcal{L}_3(t) = \frac{\lambda}{\sqrt{N}} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) (\varphi_t(y) a_x^* a_y^* + \overline{\varphi_t(y)} a_x^* a_y) a_x,$$

$$(3.10) \quad \mathcal{L}_4(t; s) := \mathcal{L}_4 = \frac{\lambda}{2N} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) a_x^* a_y^* a_x a_y.$$

Defining the unitary operator $\mathcal{U}(t; s)$ by

$$(3.11) \quad \mathcal{U}(t; s) := e^{i\omega(t; s)} W^*(\sqrt{N}\varphi_t) e^{-i\mathcal{H}_N(t-s)} W(\sqrt{N}\varphi_s),$$

with the phase factor

$$\omega(t; s) = \int_s^t d\tau \mathcal{L}_0(\tau; s),$$

we get

$$(3.12) \quad i\partial_t \mathcal{U}(t; s) = (\mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4) \mathcal{U}(t; s) =: \mathcal{L}_N(t) \mathcal{U}(t; s) \quad \text{and} \quad \mathcal{U}(s; s) = I$$

and

$$(3.13) \quad W^*(\sqrt{N}\varphi_s) e^{i\mathcal{H}_N(t-s)} (a_x - \sqrt{N}\varphi_t(x)) e^{-i\mathcal{H}_N(t-s)} W(\sqrt{N}\varphi_s) = \mathcal{U}^*(t; s) a_x \mathcal{U}(t; s),$$

where we used (3.4) and the fact that the operator $e^{i\omega(t; s)}$ is just a multiplication by a complex number.

Let

$$\begin{aligned}
(3.14) \quad \widehat{\mathcal{L}}_2(t) &:= \int_{\mathbb{R}^3} dx a_x^* (-\Delta_x + V(x)) a_x + \lambda \int_{\mathbb{R}^3} dx (w_N * |\varphi_t|^2)(x) a_x^* a_x \\
&+ \lambda \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \overline{\varphi_t(x)} \varphi_t(y) a_y^* a_x \\
&+ \frac{\lambda}{2N^{3\beta/2}} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) (\varphi_t(x) \varphi_t(y) a_x^* a_y^* + \overline{\varphi_t(x)} \overline{\varphi_t(y)} a_x a_y).
\end{aligned}$$

Also we let $\widehat{\mathcal{L}} := \widehat{\mathcal{L}}_2 + \mathcal{L}_4$ and define the unitary operator $\widehat{\mathcal{U}}(t; s)$ by

$$(3.15) \quad i\partial_t \widehat{\mathcal{U}}(t; s) = \widehat{\mathcal{L}}(t) \widehat{\mathcal{U}}(t; s) \quad \text{and} \quad \widehat{\mathcal{U}}(s; s) = 1.$$

Since $\widehat{\mathcal{L}}$ does not change the parity of the number of particles,

$$(3.16) \quad \langle \Omega, \widehat{\mathcal{U}}^*(t; 0) a_y \widehat{\mathcal{U}}(t; 0) \Omega \rangle = \langle \Omega, \widehat{\mathcal{U}}^*(t; 0) a_x^* \widehat{\mathcal{U}}(t; 0) \Omega \rangle = 0.$$

Now, we have the following bounds for $E_t^{(1)}(J)$ and $E_t^{(2)}(J)$, which will be defined and Proposition 3.1 and Proposition 3.2:

Proposition 3.1. *Suppose that the assumptions in Theorem 1.1 hold. For any compact Hermitian operator J on $L^2(\mathbb{R}^3)$, let*

$$E_t^{(1)}(J) := \frac{d_N}{N} \left\langle W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega \right\rangle.$$

Then, there exist constants C depending only on $\|w\|_{L^1(\mathbb{R}^3)}$, $\|w\|_{L^2(\mathbb{R}^3)}$, $\|V\|_{W^{2,\infty}}$ and C^V such that

$$|E_t^{(1)}(J)| \leq \frac{C\|J\|}{N}.$$

Proposition 3.2. *Suppose that the assumptions in Theorem 1.1 hold. For any compact Hermitian operator J on $L^2(\mathbb{R}^3)$, let*

$$E_t^{(2)}(J) := \frac{d_N}{\sqrt{N}} \left\langle W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \mathcal{U}^*(t; 0) \phi(J\varphi_t) \mathcal{U}(t; 0) \Omega \right\rangle.$$

Then, there exist constants C depending only on $\|w\|_{L^1(\mathbb{R}^3)}$, $\|w\|_{L^2(\mathbb{R}^3)}$, $\|V\|_{W^{2,\infty}}$ and C^V such that

$$|E_t^{(2)}(J)| \leq \begin{cases} C\|J\|N^{-1} & \text{if } \beta = 0 \\ C\|J\|N^{-\frac{1+3\beta}{2}} & \text{if } 0 < \beta < 1/3 \end{cases}$$

The proofs of these propositions will be given in Section 5.

Proof of Theorem 1.1. Recall that

$$(3.17) \quad \gamma_{N,t}^{(1)}(x; y) = \frac{1}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\rangle.$$

From the definition of the creation operator in (2.12) and the definition of d_N in (2.22), we can easily find that

$$(3.18) \quad \{0, 0, \dots, 0, \varphi_0^{\otimes N}, 0, \dots\} = \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega,$$

where the $\varphi_0^{\otimes N}$ on the left-hand side is in the N -th sector of the Fock space. Recall that P_N is the projection onto the N -particle sector of the Fock space. From (2.21), we find that

$$\frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega = \frac{\sqrt{N!}}{N^{N/2}e^{-N/2}}P_N W(\sqrt{N}\varphi_0)\Omega = d_N P_N W(\sqrt{N}\varphi_0)\Omega.$$

Since \mathcal{H}_N does not change the number of particles, we also have that

$$\begin{aligned} \gamma_{N,t}^{(1)}(x; y) &= \frac{1}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega \right\rangle \\ &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} P_N W(\sqrt{N}\varphi_0)\Omega \right\rangle \\ &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, P_N e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} W(\sqrt{N}\varphi_0)\Omega \right\rangle \\ &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} W(\sqrt{N}\varphi_0)\Omega \right\rangle, \end{aligned}$$

where we used that $P_N \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega = \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega$ in the last step. To simplify it further, we use the relation

$$(3.19) \quad e^{i\mathcal{H}_N t} a_x e^{-i\mathcal{H}_N t} = W(\sqrt{N}\varphi_0)\mathcal{U}^*(t; 0)(a_x + \sqrt{N}\varphi_t(x))\mathcal{U}(t; 0)W^*(\sqrt{N}\varphi_0),$$

which follows from the first equality in Lemma 2.2(4), the definition of \mathcal{U} in (3.11) and the unitarity of the Weyl operator, see Lemma 2.2(2). By (3.19) and an analogous result for the creation operator, we obtain that

$$\begin{aligned} \gamma_{N,t}^{(1)}(x; y) &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, e^{i\mathcal{H}_N t} a_y^* a_x e^{-i\mathcal{H}_N t} W(\sqrt{N}\varphi_0)\Omega \right\rangle \\ &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, W(\sqrt{N}\varphi_0)\mathcal{U}^*(t; 0) \left(a_y^* + \sqrt{N}\overline{\varphi_t(y)} \right) \left(a_x + \sqrt{N}\varphi_t(x) \right) \mathcal{U}(t; 0)\Omega \right\rangle. \end{aligned}$$

Thus,

$$(3.20) \quad \begin{aligned} \gamma_{N,t}^{(1)}(x; y) - \overline{\varphi_t(y)}\varphi_t(x) &= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, W(\sqrt{N}\varphi_0)\mathcal{U}^*(t; 0)a_y^* a_x \mathcal{U}(t; 0)\Omega \right\rangle \\ &\quad + \overline{\varphi_t(y)} \frac{d_N}{\sqrt{N}} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, W(\sqrt{N}\varphi_0)\mathcal{U}^*(t; 0)a_x \mathcal{U}(t; 0)\Omega \right\rangle \\ &\quad + \varphi_t(x) \frac{d_N}{\sqrt{N}} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}}\Omega, W(\sqrt{N}\varphi_0)\mathcal{U}^*(t; 0)a_y^* \mathcal{U}(t; 0)\Omega \right\rangle. \end{aligned}$$

Recall the definition of $E_t^{(1)}(J)$ and $E_t^{(2)}(J)$ in Propositions 3.1 and 3.2. For any compact one-particle Hermitian operator J on $L^2(\mathbb{R}^3)$, we have

$$\text{Tr} \left(J \left(\gamma_{N,t}^{(1)} - |\varphi_0\rangle\langle\varphi_0| \right) \right) = \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy J(x; y) \left(\gamma_{N,t}^{(1)}(y; x) - \varphi_t(y)\overline{\varphi_t(x)} \right)$$

$$\begin{aligned}
&= \frac{d_N}{N} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, W(\sqrt{N}\varphi_0) \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega \right\rangle \\
&\quad + \frac{d_N}{\sqrt{N}} \left\langle \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, W(\sqrt{N}\varphi_0) \mathcal{U}^*(t; 0) \phi(J\varphi_t) \mathcal{U}(t; 0) \Omega \right\rangle \\
&= E_t^{(1)}(J) + E_t^{(2)}(J).
\end{aligned}$$

The second step can be seen by using the expression for $d\Gamma(J)$ in terms of operator-valued distributions in (2.18), (3.20), the definition of annihilation and creation operators in terms of operator-valued distributions in (2.15) and the definition of ϕ in (2.14). Thus, from Propositions 3.1 and 3.2, we find that

$$(3.21) \quad \left| \text{Tr} \left(J(\gamma_{N,t}^{(1)} - |\varphi_t\rangle \langle \varphi_t|) \right) \right| \leq \begin{cases} C \|J\| N^{-1} & \text{if } \beta = 0 \\ C \|J\| N^{-\frac{1+3\beta}{2}} & \text{if } 0 < \beta < 1/3. \end{cases}$$

Since the space of compact operators is the dual to the space of the trace class operators, and since $\gamma_{N,t}^{(1)}$ and $|\varphi_t\rangle \langle \varphi_t|$ are Hermitian and trace class, there exists an orthonormal basis $(f_k)_{k \in \mathbb{N}} \subset L^2(\mathbb{R}^3)$ and a sequence of real numbers $(s_k)_{k \in \mathbb{N}} \subset \mathbb{R}$ with $\sum_{k \in \mathbb{N}} s_k < \infty$ such that

$$(3.22) \quad \gamma_{N,t}^{(1)} - |\varphi_t\rangle \langle \varphi_t| = \sum_{k \in \mathbb{N}} s_k |f_k\rangle \langle f_k|.$$

Now for any $K \in \mathbb{N}$ choose the Hermitian compact operator

$$(3.23) \quad J_K := \sum_{k=1}^K \text{sign}(s_k) |f_k\rangle \langle f_k|,$$

and note that $\|J_K\| \leq 1$. We have

$$(3.24) \quad \text{Tr} \left| \gamma_{N,t}^{(1)} - |\varphi_t\rangle \langle \varphi_t| \right| = \sum_{k \in \mathbb{N}} s_k = \lim_{K \rightarrow \infty} \sum_{k=1}^K s_k = \lim_{K \rightarrow \infty} \left| \text{Tr} \left(J_K (\gamma_{N,t}^{(1)} - |\varphi_t\rangle \langle \varphi_t|) \right) \right|.$$

Combining this with (3.21), we obtain

$$\text{Tr} \left| \gamma_{N,t}^{(1)} - |\varphi_t\rangle \langle \varphi_t| \right| \leq \begin{cases} CN^{-1} & \text{if } \beta = 0, \\ CN^{(3\beta-1)/2} & \text{if } 0 < \beta < 1/3, \end{cases}$$

which concludes the proof of Theorem 1.1. \square

4. TRUNCATION DYNAMICS

First, we introduce a truncated time-dependent generator with fixed $M > 0$ as follows:

$$\begin{aligned}
\mathcal{L}_N^{(M)}(t) &= \int_{\mathbb{R}^3} dx a_x^* (-\Delta_x + V(x)) a_x \\
&\quad + \lambda \int_{\mathbb{R}^3} dx (w_N * |\varphi_t|^2)(x) a_x^* a_x + \lambda \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \overline{\varphi_t(x)} \varphi_t(y) a_y^* a_x \\
&\quad + \frac{\lambda}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \left(\varphi_t(x) \varphi_t(y) a_x^* a_y^* + \overline{\varphi_t(x)} \overline{\varphi_t(y)} a_x a_y \right)
\end{aligned}$$

$$\begin{aligned}
& + \lambda \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) a_x^* \left(\overline{\varphi_t(y)} a_y \chi(\mathcal{N} \leq M) + \varphi_t(y) \chi(\mathcal{N} \leq M) a_y^* \right) a_x \\
& + \frac{\lambda}{2N} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) a_x^* a_y^* a_y a_x.
\end{aligned}$$

We remark that M will be chosen to be $M = N^{1-3\beta}$ later in the proof of Lemma 5.1. Define a unitary operator $\mathcal{U}_N^{(M)}$ by

$$(4.1) \quad i\partial_t \mathcal{U}_N^{(M)}(t; s) = \mathcal{L}_N^{(M)}(t) \mathcal{U}_N^{(M)}(t; s) \quad \text{and} \quad \mathcal{U}_N^{(M)}(s; s) = 1.$$

Lemma 4.1. *Suppose that the assumptions in Theorem 1.1 hold and let $\mathcal{U}_N^{(M)}$ be the unitary operator defined in (4.1). Then, for any $j \in \mathbb{N}$ there exists a constant $K = K(\|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2, j) > 0$ such that for all $N \in \mathbb{N}$, $M > 0$, $\psi \in \mathcal{F}$, and $t, s \in \mathbb{R}$,*

$$\left\langle \mathcal{U}_N^{(M)}(t; s) \psi, (\mathcal{N} + 1)^j \mathcal{U}_N^{(M)}(t; s) \psi \right\rangle \leq C \langle \psi, (\mathcal{N} + 1)^j \psi \rangle \exp \left(K (1 + \sqrt{M/N^{1-3\beta}}) \right).$$

Proof. Following the proof of [80, Lemma 3.5], see [80, (3.15)], we have

$$(4.2) \quad \frac{d}{dt} \langle \mathcal{U}_N^{(M)}(t; 0) \psi, (\mathcal{N} + 1)^j \mathcal{U}_N^{(M)}(t; 0) \psi \rangle = \lambda(A + B),$$

where

$$\begin{aligned}
A & := 2 \sum_{k=0}^{j-1} \binom{j}{k} (-1)^k \operatorname{Im} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \varphi_t(x) \varphi_t(y) \\
& \quad \times \langle \mathcal{U}_N^{(M)}(t; 0) \psi, (\mathcal{N}^{k/2} a_x^* a_y^* (\mathcal{N} + 2)^{k/2} + (\mathcal{N} + 1)^{k/2} a_x^* a_y^* (\mathcal{N} + 3)^{k/2}) \mathcal{U}_N^{(M)}(t; 0) \psi \rangle \\
B & := \frac{2}{\sqrt{N}} \sum_{k=0}^{j-1} \binom{j}{k} \operatorname{Im} \int_{\mathbb{R}^3} dx \langle \mathcal{U}_N^{(M)}(t; 0) \psi, a_x^* a(w_N(x-\cdot) \varphi_t) \chi(\mathcal{N} \leq M) (\mathcal{N} + 1)^{k/2} a_x \mathcal{N}^{k/2} \mathcal{U}_N^{(M)}(t; 0) \psi \rangle.
\end{aligned}$$

To control the contribution from the first term on the right-hand side of (4.2), we use bounds of the form

$$\begin{aligned}
& \left| \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \varphi_t(x) \varphi_t(y) \langle \mathcal{U}_N^{(M)}(t; 0) \psi, (\mathcal{N} + 1)^{\frac{k}{2}} a_x^* a_y^* (\mathcal{N} + 3)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi \rangle \right| \\
& \leq \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy |w_N(x-y) \varphi_t(x) \varphi_t(y)| \|a_x (\mathcal{N} + 1)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\| \|a_y^* (\mathcal{N} + 3)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\| \\
& \leq \|w_N\|_1 \|\varphi_t\|_\infty^2 \left(\int_{\mathbb{R}^3} dx \|a_x (\mathcal{N} + 1)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\|^2 \right)^{1/2} \left(\int_{\mathbb{R}^3} dy \|a_y (\mathcal{N} + 3)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\|^2 \right)^{1/2} \\
& \leq \|w\|_1 \|\varphi_t\|_\infty^2 \|(\mathcal{N} + 3)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\|^2.
\end{aligned}$$

Here we used Young's inequality in the second step and the definition of the number operator in (2.16) and $\|w_N\|_1 = \|w\|_1$ in the last step.

On the other hand, to control the second integral on the right-hand side of (4.2), we use that

$$\begin{aligned}
& \left| \int_{\mathbb{R}^3} dx \langle \mathcal{U}_N^{(M)}(t; 0) \psi, a_x^* a(w_N(x-\cdot) \varphi_t) \chi(\mathcal{N} \leq M) (\mathcal{N} + 1)^{k/2} a_x \mathcal{N}^{k/2} \mathcal{U}_N^{(M)}(t; 0) \psi \rangle \right| \\
& \leq \int_{\mathbb{R}^3} dx \|a_x (\mathcal{N} + 1)^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\| \|a(w_N(x-\cdot) \varphi_t) \chi(\mathcal{N} \leq M)\| \|a_x \mathcal{N}^{\frac{k}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\| \\
& \leq M^{1/2} \sup_x \|w_N(x-\cdot) \varphi_t\| \|(\mathcal{N} + 1)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0) \psi\|^2
\end{aligned}$$

$$\begin{aligned} &\leq M^{1/2} \|w_N\|_2 \|\varphi_t\|_\infty \|(\mathcal{N} + 1)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi\|^2 \\ &\leq M^{1/2} N^{3\beta/2} \|w\|_2 \|\varphi_t\|_\infty \|(\mathcal{N} + 1)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi\|^2. \end{aligned}$$

Using $|\lambda| \leq 1$, this gives us that

$$\begin{aligned} &\left| \frac{d}{dt} \langle \mathcal{U}_N^{(M)}(t; 0)\psi, (\mathcal{N} + 1)^j \mathcal{U}_N^{(M)}(t; 0)\psi \rangle \right| \\ &\leq 2 \sum_{k=0}^{j-1} \binom{j}{k} 2 \|w\|_1 \|\varphi_t\|_\infty^2 \|(\mathcal{N} + 3)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi\|^2 \\ &\quad + \frac{2}{\sqrt{N}} \sum_{k=0}^{j-1} \binom{j}{k} M^{1/2} N^{3\beta/2} \|w\|_2 \|\varphi_t\|_\infty \|(\mathcal{N} + 1)^{\frac{k+1}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi\|^2 \\ &\leq 2 \sum_{k=0}^{j-1} \binom{j}{k} \left(2 \|w\|_1 \|\varphi_t\|_\infty^2 + \frac{1}{\sqrt{N}} M^{1/2} N^{3\beta/2} \|w\|_2 \|\varphi_t\|_\infty \right) \|(\mathcal{N} + 3)^{\frac{j}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi\|^2 \\ &\leq 2 \cdot 2^j \left(2 \|w\|_1 \|\varphi_t\|_\infty^2 + \left(\frac{M}{N^{1-3\beta}} \right)^{1/2} \|w\|_2 \|\varphi_t\|_\infty \right) \left\| (3(\mathcal{N} + 1))^{\frac{j}{2}} \mathcal{U}_N^{(M)}(t; 0)\psi \right\|^2 \\ &\leq 2 \cdot 6^j \left(2 \|w\|_1 \|\varphi_t\|_\infty^2 + \left(\frac{M}{N^{1-3\beta}} \right)^{1/2} \|w\|_2 \|\varphi_t\|_\infty \right) \langle \mathcal{U}_N^{(M)}(t; 0)\psi, (\mathcal{N} + 1)^j \mathcal{U}_N^{(M)}(t; 0)\psi \rangle. \end{aligned}$$

Applying the Grönwall lemma together with Proposition 2.1, we get

$$\begin{aligned} &\langle \mathcal{U}_N^{(M)}(t; 0)\psi, (\mathcal{N} + 1)^j \mathcal{U}_N^{(M)}(t; 0)\psi \rangle \\ &\leq \langle \psi, (\mathcal{N} + 1)^j \psi \rangle \exp \left(\int_0^t ds 2 \cdot 6^j \left(2 \|w\|_1 \|\varphi_s\|_\infty^2 + \left(\frac{M}{N^{1-3\beta}} \right)^{1/2} \|w\|_2 \|\varphi_s\|_\infty \right) \right) \\ &\leq \langle \psi, (\mathcal{N} + 1)^j \psi \rangle \exp \left(\int_0^t ds 2 \cdot 6^j \left(2 \|w\|_1 C_0^2 (1 + |s|)^{-3} + \left(\frac{M}{N^{1-3\beta}} \right)^{1/2} \|w\|_2 C_0 (1 + |s|)^{-\frac{3}{2}} \right) \right) \\ &\leq \langle \psi, (\mathcal{N} + 1)^j \psi \rangle \exp \left(K \left(1 + \left(\frac{M}{N^{1-3\beta}} \right)^{1/2} \right) \right). \end{aligned}$$

□

5. COMPARISON OF DYNAMICS

The main goal of this section is to provide important lemmata to prove Proposition 3.1 and 3.2.

Lemma 5.1. *Suppose that the assumptions in Theorem 1.1 hold. Let $\mathcal{U}(t; s)$ be the unitary evolution defined in (3.12). Then for any $\psi \in \mathcal{F}$ and $j \in \mathbb{N}$, there exists a constant $C \equiv C(j, \|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2) > 0$ such that*

$$\langle \mathcal{U}(t; s)\psi, \mathcal{N}^j \mathcal{U}(t; s)\psi \rangle \leq C \langle \psi, (\mathcal{N} + 1)^{2L_{j,\beta} + 2} \psi \rangle,$$

where

$$L_{j,\beta} := \left\lceil \frac{j + (3\beta/2)}{1 - 3\beta} \right\rceil.$$

In this section we modify the proof given in previous articles, for example, [80] or [12] and the references therein.

We now begin the proof of Lemma 5.1. To prove the lemma, we compare the dynamics of \mathcal{U} and $\mathcal{U}^{(M)}$ in Lemma 5.3. To do so, we recall weak bounds on the \mathcal{U} dynamics.

Lemma 5.2 ([80, Lemma 3.6]). *For arbitrary $t, s \in \mathbb{R}$ and $\psi \in \mathcal{F}$, we have*

$$\langle \psi, \mathcal{U}(t; s) \mathcal{N} \mathcal{U}^*(t; s) \psi \rangle \leq 6 \langle \psi, (\mathcal{N} + N + 1) \psi \rangle.$$

Moreover, for every $\ell \in \mathbb{N}$, there exists a constant $C(\ell)$ such that

$$\begin{aligned} \langle \psi, \mathcal{U}(t; s) \mathcal{N}^{2\ell} \mathcal{U}^*(t; s) \psi \rangle &\leq C(\ell) \langle \psi, (\mathcal{N} + N)^{2\ell} \psi \rangle, \\ \langle \psi, \mathcal{U}(t; s) \mathcal{N}^{2\ell+1} \mathcal{U}^*(t; s) \psi \rangle &\leq C(\ell) \langle \psi, (\mathcal{N} + N)^{2\ell+1} (\mathcal{N} + 1) \psi \rangle \end{aligned}$$

for all $t, s \in \mathbb{R}$ and $\psi \in \mathcal{F}$.

Proof. The proof can be found in [80]. □

Now we are ready to compare the dynamics of \mathcal{U} and $\mathcal{U}_N^{(M)}$.

Lemma 5.3. *Suppose that the assumptions in Theorem 1.1 hold. Then, for every $j \in \mathbb{N}$ and $\psi \in \mathcal{F}$, there exists a constant $C \equiv C(j, \|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2) > 0$ such that for all $t, s \in \mathbb{R}$*

$$(5.1) \quad \begin{aligned} &\left| \langle \mathcal{U}(t; s) \psi, \mathcal{N}^j (\mathcal{U}(t; s) - \mathcal{U}_N^{(M)}(t; s)) \psi \rangle \right| \\ &\leq C(N^{1-3\beta}/M)^{L_{j,\beta}} \|(\mathcal{N} + 1)^{L_{j,\beta}+1} \psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right) \end{aligned}$$

and

$$(5.2) \quad \begin{aligned} &\left| \langle \mathcal{U}_N^{(M)}(t; 0) \psi, \mathcal{N}^j (\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle \right| \\ &\leq C(N^{1-3\beta}/M)^{L_{j,\beta}} \|(\mathcal{N} + 1)^{L_{j,\beta}+1} \psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right). \end{aligned}$$

Proof. To simplify the notation, we consider the case $s = 0$ and $t > 0$ only; other cases can be treated in a similar manner. To prove the first inequality of the lemma, we expand the difference of the two evolutions as follows:

$$\begin{aligned} &\langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j (\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle = \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j \mathcal{U}(t; 0) (1 - \mathcal{U}^*(t; 0) \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle \\ &= -i \int_0^t ds \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j \mathcal{U}(t; 0) (\mathcal{L}_N(s) - \mathcal{L}_N^{(M)}(s)) \mathcal{U}_N^{(M)}(s; 0) \psi \rangle \\ &= -\frac{i\lambda}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x - y) \\ &\quad \times \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j \mathcal{U}(t; s) a_x^* (\overline{\varphi_s(y)}) a_y \chi(\mathcal{N} > M) + \varphi_s(y) \chi(\mathcal{N} > M) a_y^* a_x \mathcal{U}_N^{(M)}(s; 0) \psi \rangle \\ &= -\frac{i\lambda}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \langle a_x \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi, a(w_N(x - \cdot) \varphi_s) \chi(\mathcal{N} > M) a_x \mathcal{U}_N^{(M)}(s; 0) \psi \rangle \\ &\quad - \frac{i\lambda}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \langle a_x \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi, \chi(\mathcal{N} > M) a^*(w_N(x - \cdot) \varphi_s) a_x \mathcal{U}_N^{(M)}(s; 0) \psi \rangle. \end{aligned}$$

Hence, using $|\lambda| \leq 1$,

$$\left| \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j (\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle \right|$$

$$\begin{aligned}
&\leq \frac{1}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \|a_x \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi\| \\
&\quad \times \|a(w_N(x - \cdot) \varphi_s) a_x \chi(\mathcal{N} > M + 1) \mathcal{U}_N^{(M)}(s; 0) \psi\| \\
&+ \frac{1}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \|a_x \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi\| \\
&\quad \times \|a^*(w_N(x - \cdot) \varphi_s) a_x \chi(\mathcal{N} > M) \mathcal{U}_N^{(M)}(s; 0) \psi\| \\
&\leq \frac{2}{\sqrt{N}} \int_0^t ds \sup_x \|w_N(x - \cdot) \varphi_s\| \\
&\quad \times \int_{\mathbb{R}^3} dx \|a_x \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi\| \|a_x (\mathcal{N} + 1)^{1/2} \chi(\mathcal{N} > M) \mathcal{U}_N^{(M)}(s; 0) \psi\| \\
&\leq \frac{2}{\sqrt{N}} \int_0^t ds \sup_x \|w_N(x - \cdot) \varphi_s\| \\
(5.3) \quad &\quad \times \|\mathcal{N}^{1/2} \mathcal{U}^*(t; s) \mathcal{N}^j \mathcal{U}(t; 0) \psi\| \|(\mathcal{N} + 1) \chi(\mathcal{N} > M) \mathcal{U}_N^{(M)}(s; 0) \psi\|.
\end{aligned}$$

Since $\chi(\mathcal{N} > M) \leq (\mathcal{N}/M)^L$ for any $L > 1$, we find that, from [80, (3.30)],

$$\begin{aligned}
&\left| \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j (\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle \right| \\
&\leq CN^j \|(\mathcal{N} + 1)^{j+1} \psi\| \\
&\quad \times \int_0^t ds \sup_x \|w_N(x - \cdot) \varphi_s\|_2 \langle \mathcal{U}_N^{(M)}(s; 0) \psi, (\mathcal{N} + 1)^2 \chi(\mathcal{N} > M) \mathcal{U}_N^{(M)}(s; 0) \psi \rangle^{1/2} \\
&\leq CN^j \|(\mathcal{N} + 1)^{j+1} \psi\| \\
&\quad \times \int_0^t ds N^{3\beta/2} \|w\|_2 \|\varphi_s\|_\infty \langle \mathcal{U}_N^{(M)}(s; 0) \psi, (\mathcal{N} + 1)^2 \chi(\mathcal{N} > M) \mathcal{U}_N^{(M)}(s; 0) \psi \rangle^{1/2} \\
&\leq CN^{j+(3\beta/2)} \|(\mathcal{N} + 1)^{j+1} \psi\| \langle \mathcal{U}_N^{(M)}(s; 0) \psi, \frac{(\mathcal{N} + 1)^{2+2L_{j,\beta}}}{M^{2L_{j,\beta}}} \mathcal{U}_N^{(M)}(s; 0) \psi \rangle^{1/2}
\end{aligned}$$

where

$$C = C(j, \|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2)$$

and

$$L_{j,\beta} := \left\lceil \frac{j + (3\beta/2)}{1 - 3\beta} \right\rceil.$$

Thus, we conclude that

$$\begin{aligned}
&\left| \langle \mathcal{U}(t; 0) \psi, \mathcal{N}^j (\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0)) \psi \rangle \right| \\
&\leq C(N/M)^{j+(3\beta/2)} \|(\mathcal{N} + 1)^{L_{j,\beta}+1} \psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right) \\
&\leq C(N^{1-3\beta}/M)^{L_{j,\beta}} \|(\mathcal{N} + 1)^{L_{j,\beta}+1} \psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right).
\end{aligned}$$

To prove (5.2), we proceed similarly; analogously to (5.3) we find

$$\begin{aligned} & \langle \mathcal{U}_N^{(M)}(t; 0)\psi, \mathcal{N}^j \left(\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0) \right) \psi \rangle \\ &= -\frac{i\lambda}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \langle a_x \mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi, a(w_N(x - \cdot)\varphi_t) \chi(\mathcal{N} > M) a_x \mathcal{U}_N^{(M)}(s; 0)\psi \rangle \\ & \quad - \frac{i\lambda}{\sqrt{N}} \int_0^t ds \int_{\mathbb{R}^3} dx \langle a_x \mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi, \chi(\mathcal{N} > M) a^*(w_N(x - \cdot)\varphi_t) a_x \mathcal{U}_N^{(M)}(s; 0)\psi \rangle \end{aligned}$$

and thus, by $|\lambda| \leq 1$, $\chi(\mathcal{N} > M) \leq \left(\frac{\mathcal{N}}{M}\right)^L$ for any $L > 1$, $\|w_N\|_2 = N^{3\beta/2}\|w\|_2$ and Proposition 2.1, (5.4)

$$\begin{aligned} & \left| \langle \mathcal{U}_N^{(M)}(t; 0)\psi, \mathcal{N}^j \left(\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0) \right) \psi \rangle \right| \\ & \leq \frac{C}{\sqrt{N}} \int_0^t ds \sup_x \|w_N(x - \cdot)\varphi_s\|_2 \|\mathcal{N}^{1/2}\mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi\| \|\mathcal{N}\chi(\mathcal{N} > M)\mathcal{U}_N^{(M)}(s; 0)\psi\| \\ & \leq \frac{C}{\sqrt{N}} \int_0^t ds \|w_N\|_2 \|\varphi_s\|_\infty \|\mathcal{N}^{1/2}\mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi\| \|\mathcal{N}\left(\frac{\mathcal{N}}{M}\right)^{L_{j,\beta}} \mathcal{U}_N^{(M)}(s; 0)\psi\| \\ & \leq \frac{C}{M^{L_{j,\beta}}\sqrt{N^{1-3\beta}}} \|w\|_2 \int_0^t ds C_0(1+|s|)^{-3/2} \|\mathcal{N}^{1/2}\mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi\| \|\mathcal{N}^{L_{j,\beta}+1}\mathcal{U}_N^{(M)}(s; 0)\psi\|. \end{aligned}$$

By Lemma 5.2, Lemma 4.1 and $j + 1/2 \leq L_{j,\beta} + 1$, we have

$$\begin{aligned} & \|\mathcal{N}^{1/2}\mathcal{U}(t; s)^* \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi\| \leq 6\|(\mathcal{N} + N + 1)^{1/2} \mathcal{N}^j \mathcal{U}_N^{(M)}(t; 0)\psi\| \\ & \leq 12N^{j+1/2}\|(\mathcal{N} + 1)^{j+1/2} \mathcal{U}_N^{(M)}(t; 0)\psi\| \\ (5.5) \quad & \leq CN^{j+1/2}\|(\mathcal{N} + 1)^{j+1/2}\psi\| \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right) \\ & \leq CN^{j+1/2}\|(\mathcal{N} + 1)^{L_{j,\beta}+1}\psi\| \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right). \end{aligned}$$

By Lemma 4.1, we have

$$(5.6) \quad \|\mathcal{N}^{L_{j,\beta}+1}\mathcal{U}_N^{(M)}(s; 0)\psi\| \leq C\|(\mathcal{N} + 1)^{L_{j,\beta}+1}\psi\| \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right).$$

Combining (5.4), (5.5) and (5.6), we obtain

$$\begin{aligned} & \left| \langle \mathcal{U}_N^{(M)}(t; 0)\psi, \mathcal{N}^j \left(\mathcal{U}(t; 0) - \mathcal{U}_N^{(M)}(t; 0) \right) \psi \rangle \right| \\ (5.7) \quad & \leq \frac{C}{M^{L_{j,\beta}}\sqrt{N^{1-3\beta}}} N^{j+1/2}\|(\mathcal{N} + 1)^{L_{j,\beta}+1}\psi\|^2 \exp\left(2K(1 + \sqrt{M/N^{1-3\beta}})\right) \\ & \leq \frac{C}{M^{L_{j,\beta}}} N^{j+3\beta/2}\|(\mathcal{N} + 1)^{L_{j,\beta}+1}\psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right) \\ & \leq C(N^{1-3\beta}/M)^{L_{j,\beta}}\|(\mathcal{N} + 1)^{L_{j,\beta}+1}\psi\|^2 \exp\left(K(1 + \sqrt{M/N^{1-3\beta}})\right), \end{aligned}$$

where we used $j + 3\beta/2 \leq (1 - 3\beta)L_{j,\beta}$ in the last step. Again, we have

$$C = C(j, \|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2).$$

□

Let us now prove Lemma 5.1.

Proof of Lemma 5.1. Let $M = N^{1-3\beta}$. Then by Lemmata 4.1 and 5.3, we get

$$\begin{aligned} & \langle \mathcal{U}(t; s)\psi, \mathcal{N}^j \mathcal{U}(t; s)\psi \rangle \\ &= \langle \mathcal{U}(t; s)\psi, \mathcal{N}^j (\mathcal{U} - \mathcal{U}_N^{(M)})(t; s)\psi \rangle + \langle (\mathcal{U} - \mathcal{U}_N^{(M)})(t; s)\psi, \mathcal{N}^j \mathcal{U}_N^{(M)}(t; s)\psi \rangle \\ & \quad + \langle \mathcal{U}_N^{(M)}(t; s)\psi, \mathcal{N}^j \mathcal{U}_N^{(M)}(t; s)\psi \rangle \\ & \leq C \langle \psi, (\mathcal{N} + 1)^{2L_{j, \beta} + 2} \psi \rangle. \end{aligned}$$

□

Recall the definition of $\widehat{\mathcal{U}}(t; s)$ in (3.15). In the next lemma, we prove an estimate for the evolution with respect to $\widehat{\mathcal{U}}$.

Lemma 5.4. *Suppose that the assumptions in Theorem 1.1 hold. Then, for any $\psi \in \mathcal{F}$ and $j \in \mathbb{N}$, there exists a constant $C = C(\|V\|_{W^{2, \infty}}, C^V, \|w\|_1) > 0$ such that*

$$\langle \widehat{\mathcal{U}}(t; s)\psi, \mathcal{N}^j \widehat{\mathcal{U}}(t; s)\psi \rangle \leq C \langle \psi, (\mathcal{N} + 1)^j \psi \rangle.$$

Proof. Let $\widehat{\psi} = \widehat{\mathcal{U}}(t; s)\psi$ and assume without loss of generality that $t \geq s$. We have

$$\begin{aligned} \frac{d}{dt} \langle \widehat{\psi}, (\mathcal{N} + 1)^j \widehat{\psi} \rangle &= \langle \widehat{\psi}, [i(\widehat{\mathcal{L}}_2 + \mathcal{L}_4), (\mathcal{N} + 1)^j] \widehat{\psi} \rangle \\ &= -\frac{1}{N^{3\beta/2}} \operatorname{Im} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \varphi_t(x) \varphi_t(y) \langle \widehat{\psi}, [a_x^* a_y^*, (\mathcal{N} + 1)^j] \widehat{\psi} \rangle \\ &= \frac{1}{N^{3\beta/2}} \operatorname{Im} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \varphi_t(x) \varphi_t(y) \langle \widehat{\psi}, a_x^* a_y^* ((\mathcal{N} + 3)^j - (\mathcal{N} + 1)^j) \widehat{\psi} \rangle \\ &= \frac{1}{N^{3\beta/2}} \operatorname{Im} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x-y) \varphi_t(x) \varphi_t(y) \\ & \quad \times \langle (\mathcal{N} + 3)^{(j+1)/2} a_x \widehat{\psi}, a_y (\mathcal{N} + 3)^{(1-j)/2} ((\mathcal{N} + 3)^j - (\mathcal{N} + 1)^j) \widehat{\psi} \rangle. \end{aligned}$$

Then, using Young's inequality, one gets

$$\begin{aligned} \frac{d}{dt} \langle \widehat{\psi}, (\mathcal{N} + 1)^j \widehat{\psi} \rangle &\leq \frac{1}{N^{3\beta/2}} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy |w_N(x-y)| |\varphi_t(x)| |\varphi_t(y)| \|(\mathcal{N} + 3)^{(j-1)/2} a_x \widehat{\psi}\| \\ & \quad \times \|a_y (\mathcal{N} + 3)^{(1-j)/2} ((\mathcal{N} + 3)^j - (\mathcal{N} + 1)^j) \widehat{\psi}\| \\ &\leq \frac{1}{N^{3\beta/2}} \|\varphi_t\|_\infty^2 \left(\int_{\mathbb{R}^3} dx \|(\mathcal{N} + 3)^{(j-1)/2} a_x \widehat{\psi}\|^2 \right)^{1/2} \\ & \quad \times \left(\int_{\mathbb{R}^3} dy \|a_y (\mathcal{N} + 3)^{(1-j)/2} ((\mathcal{N} + 3)^j - (\mathcal{N} + 1)^j) \widehat{\psi}\|^2 \right)^{1/2} \end{aligned}$$

Since, for all $j \in \mathbb{N}$, $|(\mathcal{N} + 3)^j - (\mathcal{N} + 1)^j| \leq C_j (\mathcal{N} + 1)^{j-1}$ for some $C_j > 0$, we have that

$$\begin{aligned} \frac{d}{dt} \langle \widehat{\mathcal{U}}(t; s)\psi, (\mathcal{N} + 1)^j \widehat{\mathcal{U}}(t; s)\psi \rangle &\leq \frac{C_j}{N^{3\beta/2}} \|\varphi_t\|_\infty^2 \|(\mathcal{N} + 1)^{j/2} \widehat{\mathcal{U}}(t; s)\psi\|^2 \\ &= \frac{C_j}{N^{3\beta/2}} \|\varphi_t\|_\infty^2 \langle \widehat{\mathcal{U}}(t; s)\psi, (\mathcal{N} + 1)^j \widehat{\mathcal{U}}(t; s)\psi \rangle. \end{aligned}$$

Here, note that C_j can change from line to line. Applying Grönwall's lemma with Proposition 2.1, we conclude that

$$\begin{aligned} \langle \widehat{\mathcal{U}}(t; s)\psi, (\mathcal{N} + 1)^j \widehat{\mathcal{U}}(t; s)\psi \rangle &\leq \langle \psi, (\mathcal{N} + 1)^j \psi \rangle \exp\left(\int_s^t dr \frac{C_j}{N^{3\beta/2}} C_0^2 \frac{1}{(1 + |r|)^3}\right) \\ &\leq C \langle \psi, (\mathcal{N} + 1)^j \psi \rangle. \end{aligned}$$

□

Lemma 5.5. *For all $\psi \in \mathcal{F}$ and $f \in L^2(\mathbb{R}^3)$, we have the following inequalities with a constant $C = C(\|V\|_{W^{2,\infty}}, C^V, \|w\|_1, \|w\|_2) > 0$.*

If $\beta = 0$,

$$(5.8) \quad \|(\mathcal{N} + 1)^{j/2} \mathcal{L}_3(t)\psi\| \leq \frac{C}{N} \|\varphi_t\|_{L^\infty} \|(\mathcal{N} + 1)^{(j+3)/2} \psi\|,$$

$$(5.9) \quad \|(\mathcal{N} + 1)^{j/2} (\mathcal{L}_2(t) - \widehat{\mathcal{L}}_2(t))\psi\| = 0, \text{ and}$$

$$(5.10) \quad \|(\mathcal{N} + 1)^{j/2} (\mathcal{U}^*(t; 0)\phi(f)\mathcal{U}(t; 0) - \widetilde{\mathcal{U}}^*(t; 0)\phi(f)\widetilde{\mathcal{U}}(t; 0))\Omega\| \leq \frac{C\|f\|}{N}.$$

If $0 < \beta < 1/3$,

$$(5.11) \quad \|(\mathcal{N} + 1)^{j/2} \mathcal{L}_3(t)\psi\| \leq \frac{C}{\sqrt{N^{1-3\beta}}} \|\varphi_t\|_{L^\infty} \|(\mathcal{N} + 1)^{(j+3)/2} \psi\|,$$

$$(5.12) \quad \|(\mathcal{N} + 1)^{j/2} (\mathcal{L}_2(t) - \widehat{\mathcal{L}}_2(t))\psi\| \leq C \|\varphi_t\|_{L^\infty} \|(\mathcal{N} + 1)^{(j+3)/2} \psi\|, \text{ and}$$

$$(5.13) \quad \|(\mathcal{N} + 1)^{j/2} (\mathcal{U}^*(t; 0)\phi(f)\mathcal{U}(t; 0) - \widetilde{\mathcal{U}}^*(t; 0)\phi(f)\widetilde{\mathcal{U}}(t; 0))\Omega\| \leq \frac{C\|f\|}{\sqrt{N^{1-3\beta}}}.$$

Proof. For (5.8) and (5.11), we copy the proof from [55, Lemma 5.3] with $\|w_N\|_2 = N^{3\beta/2}\|w\|_2$.

For (5.9) and (5.12), we follow the proof from [55, Lemma 5.3] with $\|w_N\|_1 = \|w\|_1$, and we replace \mathcal{L}_3 by

$$\mathcal{L}_2 - \widehat{\mathcal{L}}_2 = \lambda \frac{1 - N^{-3\beta/2}}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} dx dy w_N(x - y) (\varphi_t(x)\varphi_t(y)a_x^*a_y^* + \overline{\varphi_t(x)}\overline{\varphi_t(y)}a_x a_y).$$

For (5.10) and (5.13), we follow the proof from [55, Lemma 5.4] with Lemma 5.4, (5.8), (5.11), (5.9), and (5.12). □

Proof of Proposition 3.1. Recall that

$$E_t^{(1)}(J) = \frac{d_N}{N} \left\langle W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega \right\rangle.$$

We begin by noting that

$$\begin{aligned} |E_t^{(1)}(J)| &= \left| \frac{d_N}{N} \left\langle W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega \right\rangle \right| \\ &\leq \frac{d_N}{N} \left\| (\mathcal{N} + 1)^{-\frac{1}{2}} W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \left\| (\mathcal{N} + 1)^{\frac{1}{2}} \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega \right\|. \end{aligned}$$

Recall that, by Lemma 2.3,

$$\|(\mathcal{N} + 1)^{-\frac{1}{2}} W^*(\sqrt{N}\varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega\| \leq \frac{C}{d_N}$$

and, by applying Lemma 5.1 twice,

$$\begin{aligned} \|(\mathcal{N} + 1)^{\frac{1}{2}} \mathcal{U}^*(t; 0) d\Gamma(J) \mathcal{U}(t; 0) \Omega\| &\leq C \|(\mathcal{N} + 1)^{L_{j, \beta} + 1} d\Gamma(J) \mathcal{U}(t; 0) \Omega\| \\ &\leq C \|J\| \|(\mathcal{N} + 1)^{L_{j, \beta} + 2} \mathcal{U}(t; 0) \Omega\| \leq C \|J\| \|(\mathcal{N} + 1)^{L_{(L_{j, \beta} + 2), \beta} + 1} \Omega\| = C \|J\|, \end{aligned}$$

we obtain

$$|E_t^{(1)}(J)| \leq \frac{C \|J\|}{N}.$$

□

Proof of Proposition 3.2. Let

$$\mathcal{R}(f) = \mathcal{U}^*(t; 0) \phi(f) \mathcal{U}(t; 0) - \widehat{\mathcal{U}}^*(t; 0) \phi(f) \widehat{\mathcal{U}}(t; 0).$$

Then

$$\begin{aligned} |E_t^{(2)}(J)| &= \frac{d_N}{\sqrt{N}} \left\langle W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \widehat{\mathcal{U}}^*(t) \phi(J \varphi_t) \widehat{\mathcal{U}}(t; 0) \Omega \right\rangle \\ &\quad + \frac{d_N}{\sqrt{N}} \left\langle W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega, \mathcal{R}(J \varphi_t) \Omega \right\rangle \\ &\leq \frac{d_N}{\sqrt{N}} \left\| \sum_{k=0}^{\infty} (\mathcal{N} + 1)^{-\frac{5}{2}} P_{2k+1} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \left\| (\mathcal{N} + 1)^{\frac{5}{2}} \widehat{\mathcal{U}}^*(t) \phi(J \varphi_t) \widehat{\mathcal{U}}(t; 0) \Omega \right\| \\ (5.14) \quad &\quad + \frac{d_N}{\sqrt{N}} \left\| (\mathcal{N} + 1)^{-\frac{1}{2}} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\| \left\| (\mathcal{N} + 1)^{\frac{1}{2}} \mathcal{R}(J \varphi_t) \Omega \right\|. \end{aligned}$$

Let $K = \frac{1}{2} N^{1/3}$. By Lemmata 2.3, we have 2.4 show that

$$\begin{aligned} &\left\| \sum_{k=0}^{\infty} (\mathcal{N} + 1)^{-\frac{5}{2}} P_{2k+1} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\|^2 \\ &\leq \sum_{k=0}^K \left\| (\mathcal{N} + 1)^{-\frac{5}{2}} P_{2k+1} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\|^2 \\ &\quad + \frac{1}{K^4} \sum_{k=K}^{\infty} \left\| (\mathcal{N} + 1)^{-1/2} P_{2k+1} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\|^2 \\ &\leq \left(\sum_{k=0}^K \frac{C}{(k+1)^2 d_N^2 N} \right) + \frac{C}{N^{4/3}} \left\| (\mathcal{N} + 1)^{-1/2} W^*(\sqrt{N} \varphi_0) \frac{(a^*(\varphi_0))^N}{\sqrt{N!}} \Omega \right\|^2 \leq \frac{C}{d_N^2 N^{4/3}}. \end{aligned}$$

Using Lemma 5.4,

$$\begin{aligned} \|(\mathcal{N} + 1)^{\frac{5}{2}} \widehat{\mathcal{U}}^*(t) \phi(J \varphi_t) \widehat{\mathcal{U}}(t) \Omega\| &\leq C \|(\mathcal{N} + 1)^{\frac{5}{2}} \phi(J \varphi_t) \widehat{\mathcal{U}}(t) \Omega\| \\ &\leq C \|(\mathcal{N} + 1)^3 \phi(J \varphi_t) \widehat{\mathcal{U}}(t) \Omega\| = C \|\phi(J \varphi_t) (\mathcal{N} + 2)^3 \widehat{\mathcal{U}}(t) \Omega\| \\ &\leq C \|J \varphi_t\| \|(\mathcal{N} + 2)^{7/2} \widehat{\mathcal{U}}(t; 0) \Omega\| \leq C \|J\| \|(\mathcal{N} + 2)^{7/2} \Omega\| = C \|J\|. \end{aligned}$$

For the second term on the right-hand side of (5.14), we use Lemma 2.3, (5.10), and (5.13), for $f = J \varphi_t$. Altogether, we have

$$\|(\mathcal{N} + 1)^{j/2} \mathcal{R}(f) \Omega\| \leq \begin{cases} C \|J\| N^{-1} & \text{if } \beta = 0 \\ C \|J\| N^{-\frac{1+3\beta}{2}} & \text{if } 0 < \beta < 1/3 \end{cases}$$

which is the desired conclusion. □

REFERENCES

- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell. Observation of Bose-Einstein condensation in a dilute atomic vapor. *Science*, 269(5221):198–201, 1995.
- [2] C. Bardos, F. Golse, and N. J. Mauser. Weak coupling limit of the n -particle Schrödinger equation. *Methods and Applications of Analysis*, 7(2):275–294, 2000.
- [3] N. Benedikter, G. de Oliveira, and B. Schlein. Quantitative derivation of the Gross-Pitaevskii equation. *Comm. Pure Appl. Math.*, 68(8):1399–1482, 2015.
- [4] N. Benedikter, M. Porta, and B. Schlein. *Effective evolution equations from quantum dynamics*. Springer, 2016.
- [5] Bose. Plancks gesetz und lichtquantenhypothese. *ZeitSchrift für Physik*, 26(1):178–181, 1924.
- [6] C. Brennecke and B. Schlein. Gross-pitaevskii dynamics for Bose-Einstein condensates. *Analysis & PDE*, 12(6):1513–1596, 2019.
- [7] C. Brennecke, B. Schlein, and S. Schraven. Bogoliubov theory for trapped bosons in the Gross-Pitaevskii regime. *Ann. Henri Poincaré*, 23(5):1583–1658, 2022.
- [8] C. Brennecke, B. Schlein, and S. Schraven. Bose-Einstein condensation with optimal rate for trapped bosons in the Gross-Pitaevskii regime. *Math. Phys. Anal. Geom.*, 25(2):71, 2022. Id/No 12.
- [9] T. Cazenave. *Semilinear Schrödinger Equations*. American Mathematical Society, 2003.
- [10] T. Cazenave and F. Weissler. Rapidly decaying solutions of the nonlinear Schrödinger equation. *Communications in Mathematical Physics*, 147:75–100, 1992.
- [11] L. Chen and J. O. Lee. Rate of convergence in nonlinear Hartree dynamics with factorized initial data. *J. Math. Phys.*, 52(5):052108, 25, 2011.
- [12] L. Chen, J. O. Lee, and J. Lee. Rate of convergence toward Hartree dynamics with singular interaction potential. *J. Math. Phys.*, 59(3):031902, 2018.
- [13] L. Chen, J. O. Lee, and B. Schlein. Rate of convergence towards Hartree dynamics. *J. Stat. Phys.*, 144(4):872–903, 2011.
- [14] T. Chen and N. Pavlović. The quintic NLS as the mean field limit of a boson gas with three-body interactions. *Journal of Functional Analysis*, 260(4):959–997, 2011.
- [15] X. Chen. Second order corrections to mean field evolution for weakly interacting bosons in the case of three-body interactions. *Archive for Rational Mechanics and Analysis*, 203(2):455–497, 2012.
- [16] Y. Cho and T. Ozawa. Small data scattering of Hartree type fractional Schrödinger equations in dimension 2 and 3. *Journal of the Korean Mathematical Society*, 55(2), 2018.
- [17] S. Cuccagna, V. Georgiev, and N. Visciglia. Decay and scattering of small solutions of pure power NLS in \mathbb{R} with $p > 3$ and with a potential. <https://arxiv.org/abs/1209.5863>, 2012.
- [18] K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle. Bose-Einstein condensation in a gas of sodium atoms. *Physical Review Letters*, 75(22):3969–3973, 1995.
- [19] G. de Oliveira and A. Michelangeli. Mean-field dynamics for mixture condensates via fock space methods. *Reviews in Mathematical Physics*, 31(08):1950027, 2019.
- [20] P. Deift and X. Zhou. Long-time asymptotics for solutions of the NLS equation with initial data in a weighted sobolev space. *Communications on Pure and Applied Mathematics*, 56(8):1029–1077, 2003.
- [21] C. Dietze. Dispersive estimates for nonlinear Schrödinger equations with external potentials. *J. Math. Phys.*, 62(11):111502, 2021.
- [22] D. Dimonte, M. Falconi, and A. Olgiatei. On some rigorous aspects of fragmented condensation. *Nonlinearity*, 34(1):1, 2020.
- [23] D. Dimonte and E. L. Giacomelli. On Bose-Einstein condensates in the Thomas-Fermi regime. <https://arxiv.org/abs/2112.02343>, 2021.
- [24] T. Duyckaerts, J. Holmer, and S. Roudenko. Scattering for the non-radial 3d cubic nonlinear Schrödinger equation. <https://arxiv.org/abs/0710.3630>, 2007.
- [25] Einstein. Quantentheorie des einatomigen idealen gases. *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 1(3), 1925.
- [26] A. Elgart and B. Schlein. Mean field dynamics of boson stars. *Comm. Pure Appl. Math.*, 60(4):500–545, 2007.
- [27] L. Erdős and B. Schlein. Quantum dynamics with mean field interactions: a new approach. *J. Stat. Phys.*, 134(5-6):859–870, 2009.
- [28] L. Erdős, B. Schlein, and H.-T. Yau. Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems. *Inventiones mathematicae*, 167(3):515–614, 2007.
- [29] L. Erdős, B. Schlein, and H.-T. Yau. Rigorous derivation of the Gross-Pitaevskii equation with a large interaction potential. *Journal of the American Mathematical Society*, 22(4):1099–1156, 2009.

- [30] L. Erdős and H.-T. Yau. Derivation of the nonlinear Schrödinger equation from a many body Coulomb system. *Adv. Theor. Math. Phys.*, 5(6):1169–1205, 2001.
- [31] P. Germain, F. Pusateri, and F. Rousset. The nonlinear Schrödinger equation with a potential. *Annales de l'Institut Henri Poincaré. Analyse non linéaire*, 35(6):1477–1530, 2018.
- [32] J. Ginibre and T. Ozawa. Long range scattering for nonlinear Schrödinger and Hartree equations in space dimension $n \geq 2$. *Communications in Mathematical Physics*, 151(3), 1993.
- [33] J. Ginibre and G. Velo. The classical field limit of scattering theory for nonrelativistic many-boson systems. I. *Comm. Math. Phys.*, 66(1):37–76, 1979.
- [34] J. Ginibre and G. Velo. The classical field limit of scattering theory for nonrelativistic many-boson systems. II. *Comm. Math. Phys.*, 68(1):45–68, 1979.
- [35] J. Ginibre and G. Velo. On a class of non linear Schrödinger equations with non local interaction. *Mathematische Zeitschrift*, 170(2):109–136, 1980.
- [36] J. Ginibre and G. Velo. Time decay of finite energy solutions of the non linear Klein-Gordon and Schrödinger equations. *Annales de l'Institut Henri Poincaré. Physique théorique*, 43(4):399–442, 1985.
- [37] J. Ginibre and G. Velo. Scattering theory in the energy space for a class of Hartree equations. <https://arxiv.org/abs/math/9809183>, 1998.
- [38] J. Ginibre and G. Velo. *Scattering theory in the energy space for a class of Hartree equations*, volume 263 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 2000.
- [39] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, I. *Comm. Math. Phys.*, 324(2):601–636, 2013.
- [40] M. Grillakis and M. Machedon. Pair excitations and the mean field approximation of interacting bosons, II. *Commun. Partial Differ. Equ.*, 42(1):24–67, 2017.
- [41] M. G. Grillakis, M. Machedon, and D. Margetis. Second-order corrections to mean field evolution of weakly interacting bosons. I. *Comm. Math. Phys.*, 294(1):273–301, 2010.
- [42] M. G. Grillakis, M. Machedon, and D. Margetis. Second-order corrections to mean field evolution of weakly interacting bosons. II. *Adv. Math.*, 228(3):1788–1815, 2011.
- [43] E. P. Gross. Structure of a quantized vortex in boson systems. *Nuovo Cimento*, 20:454–477, 1961.
- [44] N. Hayashi and P. I. Naumkin. Asymptotics for large time of solutions to the nonlinear Schrödinger and Hartree equations. *Amer. J. Math.*, 120(2):369–389, 1998.
- [45] N. Hayashi and Y. Tsutsumi. Scattering theory for Hartree type equations. *Annales de l'Institut Henri Poincaré. Physique théorique*, 1987.
- [46] K. Hepp. The classical limit for quantum mechanical correlation functions. *Comm. Math. Phys.*, 35:265–277, 1974.
- [47] Y. Hong. Scattering for a nonlinear Schrödinger equation with a potential. <https://arxiv.org/abs/1403.3944>, 2014.
- [48] J. Kato and F. Pusateri. A new proof of long range scattering for critical nonlinear Schrödinger equations. <https://arxiv.org/abs/1004.0721>, 2010.
- [49] A. Knowles and P. Pickl. Mean-field dynamics: singular potentials and rate of convergence. *Comm. Math. Phys.*, 298(1):101–138, 2010.
- [50] E. Kuz. Rate of convergence to mean field for interacting bosons. *Communications in Partial Differential Equations*, 40(10):1831–1854, 2015.
- [51] J. Lee. On the time dependence of the rate of convergence towards Hartree dynamics for interacting bosons. *J. Stat. Phys.*, 176(2):358–381, 2019.
- [52] J. Lee. Rate of convergence towards mean-field evolution for weakly interacting bosons with singular three-body interactions. <https://arxiv.org/abs/2006.13040>, 2020.
- [53] J. Lee. Rate of convergence toward Hartree type equations for mixture condensates with factorized initial data. *J. Math. Phys.*, 62(9):091901, 2021.
- [54] J. Lee and A. Michelangeli. *in preparation*.
- [55] J. O. Lee. Rate of convergence towards semi-relativistic Hartree dynamics. *Ann. Henri Poincaré*, 14(2):313–346, 2013.
- [56] A. J. Leggett. Bose-Einstein condensation in the alkali gases: Some fundamental concepts. *Reviews of Modern Physics*, 73:307–356, 2001.
- [57] M. Lewin, P. T. Nam, and N. Rougerie. Derivation of 's theory for generic mean-field Bose systems. *Adv. Math.*, 254:570–621, 2014.
- [58] M. Lewin, P. T. Nam, and B. Schlein. Fluctuations around Hartree states in the mean-field regime. *American Journal of Mathematics*, 137(6):1613–1650, 2015.
- [59] E. H. Lieb, R. Seiringer, J. P. Solovej, and J. Yngvason. *The mathematics of the Bose gas and its condensation*, volume 34. Springer Science & Business Media, 2005.

- [60] A. Michelangeli, P. T. Nam, and A. Olgiati. Ground state energy of mixture of Bose gases. *Reviews in Mathematical Physics*, 31(02):1950005, 2019.
- [61] A. Michelangeli and A. Olgiati. Mean-field quantum dynamics for a mixture of Bose–Einstein condensates. *Analysis and Mathematical Physics*, 7(4):377–416, 2017.
- [62] A. Michelangeli and B. Schlein. Dynamical collapse of boson stars. *Comm. Math. Phys.*, 311(3):645–687, 2012.
- [63] K. Nakanishi. Global dynamics below excited solitons for the nonlinear Schrödinger equation with a potential. *Journal of the Mathematical Society of Japan*, 69(4):1353–1401, 2017.
- [64] P. T. Nam and M. Napiórkowski. Bogoliubov correction to the mean-field dynamics of interacting bosons. <https://arxiv.org/abs/1509.04631>, 2015.
- [65] P. T. Nam and M. Napiórkowski. A note on the validity of bogoliubov correction to mean-field dynamics. *Journal de mathématiques pures et appliquées*, 108(5):662–688, 2017.
- [66] P. T. Nam, M. Napiórkowski, J. Ricaud, and A. Triay. Optimal rate of condensation for trapped bosons in the Gross-Pitaevskii regime. <https://arxiv.org/abs/2001.04364>.
- [67] P. T. Nam, J. Ricaud, and A. Triay. The condensation of a trapped dilute Bose gas with three-body interactions. <https://arxiv.org/abs/2110.08195>, 2021.
- [68] P. T. Nam and A. Triay. Bogoliubov excitation spectrum of trapped Bose gases in the Gross-Pitaevskii regime. <https://arxiv.org/abs/2106.11949>, 2021.
- [69] M. Napiórkowski. Dynamics of interacting bosons: a compact review. <https://arxiv.org/abs/2101.04594>, 2021.
- [70] I. P. Naumkin. Sharp asymptotic behavior of solutions for cubic nonlinear Schrödinger equations with a potential. *Journal of Mathematical Physics*, 57(5), 2016.
- [71] I. P. Naumkin. Nonlinear Schrödinger equations with exceptional potentials. *Journal of Differential Equations*, 265(9):4575–4631, 2018.
- [72] T. Ozawa. Long range scattering for nonlinear Schrödinger equations in one space dimension. *Communications in Mathematical Physics*, 139(3):479–493, 1991.
- [73] P. Pickl. On the time dependent gross pitaevskii- and Hartree equation. <https://arxiv.org/abs/0808.1178>, 2008.
- [74] P. Pickl. A simple derivation of mean field limits for quantum systems. *Lett. Math. Phys.*, 97(2):151–164, 2011.
- [75] P. Pickl. Derivation of the time dependent Gross-Pitaevskii equation with external fields. *Rev. Math. Phys.*, 27(1):1550003, 45, 2015.
- [76] L. Pitaevskii. Vortex lines in an imperfect Bose gas. *Journal of Experimental and Theoretical Physics*, 13(2):451–454, 1961.
- [77] Y. Pomeau and S. Rica. Model of superflow with rotons. *Physical Review Letters*, 71(2):247–250, 1993.
- [78] F. Pusateri and A. Soffer. Bilinear estimates in the presence of a large potential and a critical NLS in 3d. <https://arxiv.org/abs/2003.00312>, 2020.
- [79] I. Rodnianski and W. Schlag. Time decay for solutions of Schrödinger equations with rough and time-dependent potentials. *Interventiones Mathematicae*, 155(3):451–513, 2004.
- [80] I. Rodnianski and B. Schlein. Quantum fluctuations and rate of convergence towards mean field dynamics. *Comm. Math. Phys.*, 291(1):31–61, 2009.
- [81] H. Spohn. Kinetic equations from Hamiltonian dynamics: Markovian limits. *Rev. Modern Phys.*, 52(3):569–615, 1980.

(C. Dietze) DEPARTMENT OF MATHEMATICS, LMU MUNICH, THERESIENSTRASSE 39, 80333 MUNICH, GERMANY
 Email address: dietze@math.lmu.de

(J. Lee) DEPARTMENT OF MATHEMATICS, LMU MUNICH, THERESIENSTRASSE 39, 80333 MUNICH, GERMANY
 Email address: lee@math.lmu.de