

On the blow-up analysis at collapsing poles for solutions of singular Liouville type equations.

Gabriella Tarantello

July 1, 2022

Abstract

We analyse a blow-up sequence of solutions for Liouville type equations involving Dirac measures with "collapsing" poles. We consider the case where blow-up occurs exactly at a point where the poles coalesce.

After proving that a "quantization" property still holds for the "blow-up mass", we obtain precise pointwise estimates when blow-up occurs with the least blow-up mass.

Interestingly, such estimates express the exact analogue of those obtained in [17] for solutions of "regular" Liouville equations, where the "collapsing" Dirac measures are neglected. Such information will be used in [25] to describe the asymptotic behaviour of minimizers of the Donaldson functional introduced in [10], yielding to mean curvature 1-immersions of surfaces into hyperbolic 3-manifolds.

1 Introduction

In this note we analyze the behaviour of a sequence of solutions for Liouville type equations involving Dirac measures whose poles coalesce at a point where the solutions blow-up. Namely, we have blow-up at a point of "collapsing" singularities. This situation naturally arises in the study of Toda systems of Liouville type in the analysis of the so called "shadow" system discussed in [15]. It has motivated the work in [19],[14], where (after [24]) the phenomenon of "blow-up without concentration" was recorded and illustrated by various examples.

More conveniently, for the following discussion, we notice that the role of the poles may be replaced by the zeroes of the weight functions appearing in the equations governing the "regular" part of the given solution sequence. In this formulation, the issue is to understand the blow-up behavior of a (regular) sequence at a point of "collapsing" zeroes.

MSC: 35J75 35J61. Keywords: Blow-up analysis, Singular Liouville equations, Blow-up at collapsing poles, Mass quantization and pointwise estimate

In fact, this is exactly what may occur when studying the asymptotic behavior of constant mean curvature (CMC) c -immersions of a closed oriented surface S of genus $g \geq 2$ into hyperbolic 3-manifolds. Following [10], it was recently established in [13] that for $|c| < 1$, the moduli space of all such (CMC) c -immersions can be parametrized by elements of the tangent bundle $T(\mathcal{T}_g(S))$. Recall that $\mathcal{T}_g(S)$ is the space of all conformal structures on S identified up to biholomorphisms in the same homotopic class of the identity. Actually, for $|c| < 1$, the authors in [13] could label any such (CMC) c -immersion with the minimizer (and unique critical point) of the Donaldson functional associated to a given element of $T(\mathcal{T}_g(S))$, as introduced in [10].

In [25], we initiated our investigation about the existence of analogous (CMC) 1-immersions. As established in [25], those immersions can be detected only as "limits", as $c \rightarrow 1^-$, of the (CMC) c -immersions obtained in [13]. In view of the work of Bryant [6] about (CMC) 1-immersions into the hyperbolic space \mathbb{H}^3 , we expect that those (CMC) c -immersions develop at the limit: $c \rightarrow 1^-$, finitely many "punctures", corresponding to possible ramification points. We hope to capture such behaviour by means of our blow-up analysis, with the "punctures" being realized by the blow-up points. But we face a delicate situation, exactly when, at the limit, the pull-back metrics of the (CMC) c -immersions blow-up at points of "collapsing" zeroes of the holomorphic quadratic differentials identified by the second fundamental form of the immersion. Indeed, recall that any holomorphic quadratic differential admits exactly $4(g-1)$ zeroes in X , counted with multiplicity (see [11]).

The pointwise estimates we establish here (see Theorem 5) have helped us to obtain in [25] the first existence result about (regular) (CMC) 1-immersions of surfaces of genus 2 into 3-manifolds of sectional curvature -1, see [25] for details.

More precisely, we start by showing that even in the "collapsing" case, the blow-up mass (see (1.12) below) is "quantized" and must take values in $8\pi\mathbb{N}$. This is a somewhat expected property, as it has been worked out already in the context of systems in [15],[16] and in [14] only for two collapsing zeroes. Nevertheless, for the sake of completeness, we have chosen to provide in Section 2 a detailed proof (see Theorem 1), together with other useful extensions of known facts.

More importantly, we obtain sharp pointwise estimates when blow-up occur with the least blow-up mass 8π . Interestingly, such estimates are exactly the analogue of the estimates obtained by Li in [17] for solution-sequences of Liouville equations with non-vanishing weight functions, where none of the "collapsing" issues discussed here arise.

In order to state our result, (after a translation) we localize our analysis around the origin. Therefore we consider a solution sequence u_k satisfying:

$$\begin{cases} -\Delta u_k = h_k e^{u_k} - 4\pi \sum_{j=1}^s \alpha_j \delta_{p_{j,k}} & (1.1) \\ \int B_r h_k e^{u_k} \leq C & (1.2) \end{cases}$$

with

$$s \geq 2 \text{ and } \underline{\text{distinct}} \text{ points } p_{j,k} \rightarrow 0, \text{ as } k \rightarrow +\infty, j = 1, \dots, s \quad (1.3)$$

$$0 < a \leq h_k \leq b, |\nabla h_k| \leq A \text{ in } B_r. \quad (1.4)$$

In addition, we require the following bounded oscillation property:

$$\max_{\partial B_r} u_k - \min_{\partial B_r} u_k \leq C \quad (1.5)$$

which is always verified when u_k is actually a "localization" of a globally defined function, for example over a Riemann surface (see e.g. Theorem 4).

We define the "regular" part ξ_k of u_k as given by

$$u_k(x) = \xi_k(x) + \sum_{j=1}^n 2\alpha_j \ln |x - p_{j,k}| \quad (1.6)$$

which defines a smooth function in B_r and satisfies

$$\begin{cases} -\Delta \xi_k = (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k e^{\xi_k} & \text{in } B_r \end{cases} \quad (1.7)$$

$$\begin{cases} \int_{B_r} W_k e^{\xi_k} \leq C \text{ with } W_k(x) = (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k(x) \end{cases} \quad (1.8)$$

$$\begin{cases} \max_{\partial B_r} \xi_k - \min_{\partial B_r} \xi_k \leq C \end{cases} \quad (1.9)$$

We assume that ξ_k admits a blow-up point at the origin, that is

$$\xi_k(0) = \max_{B_r} \xi_k \rightarrow +\infty, \text{ as } k \rightarrow +\infty. \quad (1.10)$$

Since by our assumptions we know that ξ_k can admit at most finitely many blow-up points (see Remark 2.1), by taking $r > 0$ smaller if necessary, we can further assume that the origin is the only blow-up point for ξ_k in B_r , namely:

$$\forall \varepsilon \in (0, r) \exists C_\varepsilon > 0 : \max_{\overline{B_r} \setminus B_\varepsilon} \xi_k \leq C_\varepsilon. \quad (1.11)$$

Finally, we define the "blow-up mass" of ξ_k at the origin as follows

$$\sigma = \lim_{\delta \rightarrow 0^+} \lim_{k \rightarrow +\infty} \int_{B_\delta} W_k e^{\xi_k}. \quad (1.12)$$

The main results contained in this note can be summarized as follows:

Theorem. *Assume that ξ_k satisfies (1.7)-(1.11). If $\alpha_j \in \mathbb{N}$ and (1.3),(1.4) hold then, for σ in (1.12) we have:*

$$\sigma \in 8\pi\mathbb{N}.$$

Furthermore, if $\sigma = 8\pi$ then:

$$p_{j,k} \neq 0 \quad \forall j = 1, \dots, s, \quad W_k(0) = \prod_{j=1}^s |p_{j,k}|^{2\alpha_j} h_k(0) > 0 \quad \text{and} \quad W_k(0) \xrightarrow{k \rightarrow \infty} 0^+$$

$$\xi_k(x) = \ln \frac{e^{\xi_k(0)}}{(1 + W_k(0)e^{\xi_k(0)}|x|^2)^2} + O(1) \quad \text{in } B_r; \quad (1.13)$$

$$\int_{B_r} |\nabla \xi_k|^2 = 16\pi(\xi_k(0) + \ln(W_k(0))) + O(1);$$

$$\xi_k(0) + \min_{\partial B_r} \xi_k + 2 \ln W_k(0) = O(1)$$

$$u_k(0) + \min_{\partial B_r} u_k \longrightarrow +\infty, \quad \text{as } k \longrightarrow +\infty, \quad u_k \text{ in (1.6)}. \quad (1.14)$$

By (1.14), we see that our original sequence u_k satisfying (1.1),(1.2) and (1.5) blows-up at the origin if and only if its regular part ξ_k (see (1.6)) blows-up there as well. Furthermore, as already observed, it is remarkable that the pointwise estimate (1.13) is the exact analogue of the one established in [17], when $W_k(0)$ is bounded uniformly away from zero, and none of the issues (discussed here) about "collapsing of zeros" arise.

Our results are just a first contribution towards the understanding of the blow-up phenomena at "collapsing" zeroes (or poles). We hope they open the way to the description of "multiple" blow-up profiles, already present in the "non-collapsing" case, according to the analysis in [4],[27] and [28].

2 Blow-up at collapsing zeroes: local analysis.

Let $\Omega \subset \mathbb{R}^2$ be an open, bounded and regular set. We consider the sequence: $\eta_k \in C^2(\Omega) \cap C^0(\overline{\Omega})$, satisfying the following Liouville type problem:

$$\begin{cases} -\Delta \eta_k = W_k e^{\eta_k} \text{ in } \Omega & (2.1) \end{cases}$$

$$\begin{cases} \max_{\partial \Omega} \eta_k - \min_{\partial \Omega} \eta_k \leq C & (2.2) \end{cases}$$

$$\begin{cases} \int_{\Omega} W_k e^{\eta_k} \leq C & (2.3) \end{cases}$$

with a weight function $W_k \in L^\infty(\Omega)$.

After the work of Brezis-Merle [5], a vast literature is now available, concerning the asymptotic behavior of η_k (possibly along a subsequence), as $k \longrightarrow +\infty$, according to various assumptions on W_k and its vanishing behavior, see for example [3],[8],[9],[18],[24],[26].

Motivated by our applications in [25], where we describe the asymptotic behaviour of minimizers of the Donaldson functional introduced in [10] (see [13]), here we shall take W_k to satisfy:

$$W_k \geq 0 \quad \text{and} \quad \|W_k\|_{L^\infty(\Omega)} + \int_{\Omega} \frac{1}{(W_k)^{\varepsilon_0}} \leq C, \quad \text{for some } \varepsilon_0 > 0. \quad (2.4)$$

A first important information about the sequence η_k is the following well known result, stemming from [5]:

Proposition A. *Assume (2.1)-(2.3) and (2.4). If $\overline{\lim}_{k \rightarrow +\infty} \int_{\Omega} W_k e^{\eta_k} < 4\pi$, then η_k^+ is uniformly bounded in $C_{loc}^0(\Omega)$.*

Proof. See Proposition 5.3.13 in [26]. □

Consequently, as in [5], it is natural to define $z_0 \in \Omega$ a blow-up point for the sequence η_k , if the following holds

$$\exists z_k \rightarrow z_0 \text{ with } \eta_k(z_k) \rightarrow +\infty \text{ as } k \rightarrow \infty.$$

Moreover, we call the blow-up mass of η_k at z_0 the value:

$$\sigma(z_0) = \lim_{r \rightarrow 0} \underline{\lim}_{k \rightarrow +\infty} \int_{B_r(z_0)} W_k e^{\eta_k}. \quad (2.5)$$

Remark 2.1. *By virtue of Proposition A we know that $\sigma(z_0) \geq 4\pi$, and so, by the assumption (2.3), we know that η_k can admit at most a finite number of blow-up points.*

From now on we shall denote by \mathcal{S} the finite set (possibly empty) of all blow-up points of η_k in Ω , and refer to \mathcal{S} as the blow-up set.

The following holds.

Proposition B. *Under the assumptions of Proposition A, η_k^+ is uniformly bounded in $C_{loc}^0(\Omega \setminus \mathcal{S})$, along a subsequence.*

Proof. See Proposition 5.3.17 in [26]. □

The information above allows us to provide the following "rough" description about the asymptotic behaviour of (a subsequence of) η_k . It is a general version of analogous statements available in literature (under stronger assumptions on the vanishing properties of W_k) starting with [5],[18],[3] and then [24],[19],[14]. See also [15],[16],[20],[21],[22],[23] for analogous results in the context of Liouville type systems.

Proposition 2.1. *Let η_k satisfy (2.1)-(2.3) with $W_k \rightarrow W$ uniformly in $C_{loc}^0(\Omega)$, and assume that (2.4) holds. Then (along a subsequence) η_k satisfies one of the following alternatives, as $k \rightarrow +\infty$:*

(i) $\eta_k \rightarrow -\infty$ uniformly on compact sets of Ω ,

(ii) $\eta_k \rightarrow \eta_0$ in $C_{loc}^2(\Omega)$, with η_0 satisfying:
$$\begin{cases} -\Delta \eta_0 = W e^{\eta_0} & \text{in } \Omega; \\ \int_{\Omega} W e^{\eta_0} \leq C, \end{cases}$$

(iii) (blow-up): There exists a finite set $\mathcal{S} \neq \emptyset$ of blow-up points of η_k in Ω such that, as $k \rightarrow +\infty$:

a) either (blow-up with "concentration")

$$\begin{aligned} \eta_k &\longrightarrow -\infty \text{ uniformly on compact sets of } \Omega \setminus \mathcal{S} \\ W_k e^{\eta_k} &\rightharpoonup \sum_{q \in \mathcal{S}} \sigma(q) \delta_q, \text{ weakly in the sense of measures,} \end{aligned} \quad (2.6)$$

b) or (blow-up without concentration)

$$\begin{aligned} \eta_k &\longrightarrow \eta_0 \text{ in } C_{loc}^2(\Omega \setminus \mathcal{S}); \\ W_k e^{\eta_k} &\rightharpoonup \sum_{q \in \mathcal{S}} \sigma(q) \delta_q + W e^{\eta_0}, \text{ weakly in the sense of measures,} \end{aligned}$$

and

$$\begin{cases} -\Delta \eta_0 = W e^{\eta_0} + \sum_{q \in \mathcal{S}} \sigma(q) \delta_q & \text{in } \Omega; \\ \int_{\Omega} W e^{\eta_0} \leq C. \end{cases}$$

Moreover, the blow-up mass $\sigma(q) \geq 4\pi$, $\forall q \in \mathcal{S}$.

Proof. As already mentioned, the claimed results are available in literature under various assumptions on W_k , and for completeness we highlight here the main arguments involved. Firstly we observe that the sequence:

$$\phi_k = \eta_k - \min_{\partial\Omega} \eta_k$$

is uniformly bounded in $\partial\Omega$ and by the Green representation formula we have:

$$\phi_k(x) = \frac{1}{2\pi} \int_{\Omega} \ln\left(\frac{1}{|x-y|}\right) W_k e^{\eta_k} dy + \Psi_k \quad (2.7)$$

with suitable $\Psi_k \rightarrow \Psi$ uniformly in $C^2(\Omega)$.

Recalling that \mathcal{S} is the (possibly empty) blow-up set of η_k in Ω , by Proposition B we know that η_k^+ is uniformly bounded on compact sets of $\Omega \setminus \mathcal{S}$. Therefore, we can use well known potential estimates to conclude that (along a subsequence):

$$\phi_k \longrightarrow \phi_0 \text{ uniformly in } C_{loc}^{1,\alpha}(\Omega \setminus \mathcal{S}). \quad (2.8)$$

Next, letting $s_0 = \frac{\varepsilon_0}{1+\varepsilon_0} \in (0, 1)$ with $\varepsilon_0 > 0$ in (2.4), we check:

$$\begin{aligned} \min_{\partial\Omega} \eta_k &= \min_{\Omega} \eta_k \leq \int_{\Omega} \eta_k^+ = \frac{1}{s_0} \int_{\Omega} s_0 \eta_k^+ \leq C \int_{\Omega} e^{s_0 \eta_k} \\ &\leq C \left(\int_{\Omega} W_k e^{\eta_k} \right)^{s_0} \left(\int_{\Omega} \frac{1}{W_k^{\varepsilon_0}} \right)^{1-s_0} \leq C. \end{aligned}$$

So (along a subsequence)

$$\text{either } \min_{\partial\Omega} \eta_k \longrightarrow -\infty \text{ on compact sets of } \Omega \setminus \mathcal{S} \text{ or } \left| \min_{\partial\Omega} \eta_k \right| \leq C. \quad (2.9)$$

If the first alternative holds in (2.9), then (along a subsequence) we deduce alternative (i) in case $\mathcal{S} = \emptyset$, or in case $\mathcal{S} \neq \emptyset$ then *blow-up with concentration* in alternative (iii).

On the contrary in case the second alternative holds in (2.9), then by virtue of (2.8), we find (along a subsequence) that

$$\eta_k \longrightarrow \eta_0 \text{ uniformly in } C_{loc}^0(\Omega \setminus \mathcal{S}).$$

Hence we conclude that either (ii) or alternative b) of (iii) hold according to whether $\mathcal{S} = \emptyset$ or $\mathcal{S} \neq \emptyset$, and in view of Remark 2.1 the proof is completed. \square

As discussed in [19] and [14], all the alternatives of Proposition 2.1 can actually occur. When alternative (iii) holds, then to better understand the behavior of η_k around a blow-up point $q \in \mathcal{S}$, it is crucial to identify the specific value of the blow-up mass $\sigma(q)$ in (2.5). To this purpose, for the weight function W_k , we shall work under the following assumption,

$$|\nabla W_k| \leq A \text{ and } W_k \longrightarrow W \text{ in } C_{loc}^0(\Omega). \quad (2.10)$$

According to the results in [18] and [3], the value of $\sigma(q)$ depends on whether the limiting function W in (2.10) vanishes or not at $q \in \mathcal{S}$. If locally around $q \in \mathcal{S}$ there holds:

$$W_k(x) = |x - p_k|^{2\alpha} h_k(x) \text{ in } B_r(q), \alpha \geq 0, p_k \longrightarrow q, \quad (2.11)$$

$$0 < a \leq h_k \leq b \text{ and } |\nabla h_k| \leq A \text{ in } B_r \quad (2.12)$$

then we know the following:

Theorem A. ([3],[18]) *If η_k in Proposition 2.1 satisfies alternative (iii) and for $q \in \mathcal{S}$ the weight function W_k satisfies (2.11) and (2.12) in $B_r(q)$, then only alternative a) occurs, namely we have blow-up with "concentration" as described in (2.6) and*

(i) *if $\alpha = 0$ in (2.11) then $\sigma(q) = 8\pi$,*

(ii) *if $\alpha > 0$ in (2.11) then $\sigma(q) = 8\pi(1 + \alpha)$.*

In this note, we shall focus to the case where, for $q \in \mathcal{S}$, we have $W(q) = 0$ and q is the accumulation point of different zeroes of W_k (collapsing of zeroes).

In view of the applications we have in mind, we consider the case where the zeroes of W_k have integral multiplicity, and more precisely for $q \in \mathcal{S}$ and $r > 0$ sufficiently small, we assume:

$$W_k(x) = \left(\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j} \right) h_k(x), \quad x \in B_r(q), \quad \alpha_j \in \mathbb{N}, \quad s \geq 2 \quad (2.13)$$

$$p_{j,k} \neq p_{l,k} \text{ for } j \neq l \text{ and } p_{j,k} \longrightarrow q, \text{ as } k \longrightarrow +\infty, \quad \forall j = 1, \dots, s. \quad (2.14)$$

In this case we start by proving a *quantization* property for the blow-up mass in (2.5), which completes the result in [14] (where only two zeroes of W_k coalesce at q) and follows as in [15] and [16], where analogous information were deduced in the context of systems.

Theorem 1. *Suppose that η_k in Proposition 2.1 satisfy alternative (iii) and for $q \in \mathcal{S}$ assume (2.12)-(2.14). Then $\sigma(q) \in 8\pi\mathbb{N}$.*

In order to prove Theorem 1 we need some preliminaries. First of all, we can "localize" our analysis around the blow-up point q . Indeed by means of (2.7), we have

$$\eta_k(x) = \min_{\partial\Omega} \eta_k + \frac{1}{2\pi} \int_{\Omega} \ln\left(\frac{1}{|x-y|}\right) W_k e^{\eta_k} dy + \Psi_k$$

with Ψ_k uniformly bounded in $C^2(\Omega)$. Hence we easily check (as in [2]) that η_k satisfies the bounded oscillation property around q .

Thus, after a translation, we can take $q = 0$, and for $r > 0$ sufficiently small, we need to analyze a sequence $\xi_k \in C^2(B_r) \cap C^0(\overline{B_r})$ satisfying:

$$\begin{cases} -\Delta \xi_k = (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k(x) e^{\xi_k} & \text{in } B_r \end{cases} \quad (2.15)$$

$$\begin{cases} \max_{\partial B_r} \xi_k - \min_{\partial B_r} \xi_k \leq C \end{cases} \quad (2.16)$$

$$\begin{cases} \max_{\overline{B_r}} \xi_k = \xi_k(x_k) \rightarrow +\infty, \text{ and } x_k \rightarrow 0, \text{ as } k \rightarrow +\infty, \end{cases} \quad (2.17)$$

where $s \geq 2$, $\alpha_j \in \mathbb{N}$, h_k satisfies (2.12) in B_r ,

$$p_{j,k} \neq p_{l,k} \text{ for } j \neq l, p_{j,k} \rightarrow 0 \text{ as } k \rightarrow +\infty, \forall j = 1, \dots, s, \quad (2.18)$$

and

$$\int_{B_r} W_k e^{\xi_k} \leq C, \text{ where } W_k(x) = (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k(x). \quad (2.19)$$

Finally, without loss of generality, in view of (2.12), we can assume also that,

$$h_k \rightarrow h, \text{ in } C_{loc}^0(B_r) \text{ with } h(0) = 1 \quad (2.20)$$

and zero is the only blow-up point of ξ_k in B_r , that is:

$$\forall 0 < \delta < r \exists C_\delta > 0 : \max_{\overline{B_r} \setminus B_\delta} \xi_k \leq C_\delta. \quad (2.21)$$

Clearly, under the above assumptions, we have that (2.10) holds with

$$W(x) = |x|^{2\alpha} h(x), h(0) = 1 \text{ and } \alpha = \sum_{j=1}^s \alpha_j \in \mathbb{N}. \quad (2.22)$$

Furthermore, Proposition 2.1 applies to ξ_k in B_r , and by (2.17) and (2.21) we know that (along a subsequence) ξ_k satisfies alternative (iii) with $\mathcal{S} = \{0\}$, and we have

$$m := \lim_{\delta \searrow 0} \liminf_{k \rightarrow +\infty} \frac{1}{2\pi} \int_{B_\delta(0)} W_k e^{\xi_k} \geq 2. \quad (2.23)$$

Since, as above, by Green representation formula, we can write:

$$\xi_k(x) = \min_{\overline{B_r}} \xi_k + \frac{1}{2\pi} \int_{B_r} \ln\left(\frac{1}{|x-y|}\right) W_k e^{\xi_k} dy + \psi_k, x \in B_r, \quad (2.24)$$

with ψ_k uniformly bounded in $C^2(B_r)$, we can use the information in part (iii) of Proposition 2.1, to deduce that (along a subsequence) the following holds:

$$\nabla \xi_k \longrightarrow -m \frac{x}{|x|^2} + \nabla \phi, \quad \text{uniformly in } C_{loc}^1(B_r \setminus \{0\}), \quad \text{as } k \longrightarrow +\infty, \quad (2.25)$$

with a suitable ϕ smooth in B_r .

At this point, to establish Theorem 1, it suffices to prove the following:

Theorem 2. *Under the above assumptions, in (2.23) we have that $m \in 4\mathbb{N}$.*

Proof. We set $\alpha = \sum_{j=1}^s \alpha_j \in \mathbb{N}$ and note that $\alpha \geq 2$. Letting

$$\tau_k = \max_{j=1, \dots, s} |p_{j,k}| \longrightarrow 0, \quad \text{as } k \longrightarrow +\infty, \quad (2.26)$$

we define

$$q_{j,k} = \frac{p_{j,k}}{\tau_k}, \quad j = 1, \dots, s,$$

and since $|q_{j,k}| \leq 1$, along a subsequence, we can assume:

$$q_{j,k} \longrightarrow q_j \quad \text{as } k \longrightarrow +\infty, \quad j = 1, \dots, s.$$

Therefore, (recalling the normalization $h(0) = 1$)

$$0 \leq W_{1,k}(x) := (\prod_{j=1}^s |x - q_{j,k}|^{2\alpha_j}) h_k(\tau_k x) \longrightarrow \prod_{j=1}^s |x - q_j|^{2\alpha_j} = W_1(x), \quad (2.27)$$

as $k \longrightarrow +\infty$, uniformly on compact sets of \mathbb{R}^2 . We should keep in mind that $W_1(x)$ vanishes exactly at the set

$$Z_1 := \{q_1, \dots, q_s\}, \quad (2.28)$$

and the q_j 's may not be distinct. We define

$$\varphi_k(x) = \xi_k(\tau_k x) + 2(\alpha + 1) \ln \tau_k \quad \text{in } D_k = \{|x| \leq \frac{r}{\tau_k}\} \quad (2.29)$$

satisfying:

$$-\Delta \varphi_k = W_{1,k}(x) e^{\varphi_k} \quad \text{in } D_k$$

$$\int_{D_k} W_{1,k} e^{\varphi_k} \leq C.$$

By scaling (2.24), for any given $x_1, x_2 \in \mathbb{R}^2$ and k large, we have:

$$\varphi_k(x_1) - \varphi_k(x_2) = \frac{1}{2\pi} \int_{D_k} \ln \left(\frac{|x_2 - y|}{|x_1 - y|} \right) W_{1,k} e^{\varphi_k} + T_k(x_1, x_2) \quad (2.30)$$

with T_k uniformly bounded in C_{loc}^2 . Therefore (as in Lemma 2.2 of [2]) we find $R_0 > 1$ sufficiently large, such that $\forall R \geq R_0$ there holds,

$$\max_{\partial B_R} \varphi_k - \min_{\partial B_R} \varphi_k \leq C_R,$$

with a suitable constant C_R , depending on R only.

Since $W_{1,k}$ in (2.27) satisfies (2.4) and (2.10) in any open and bounded set of \mathbb{R}^2 (for large k), we can therefore apply Proposition 2.1 to φ_k in any ball B_R with $R \geq R_0$.

Furthermore, by a diagonalization process, along a subsequence, we can set

$$\mu = \lim_{R \rightarrow +\infty} \liminf_{k \rightarrow +\infty} \frac{1}{2\pi} \int_{B_R} W_{1,k} e^{\varphi_k} \leq m, \quad (2.31)$$

and we can describe the asymptotic behavior of φ_k , as $k \rightarrow +\infty$, by one of the following alternatives:

$$(i) \quad \varphi_k \rightarrow -\infty \text{ uniformly on compact sets of } \mathbb{R}^2 \text{ and } \mu = 0, \quad (2.32)$$

$$(ii) \quad \varphi_k \rightarrow \varphi_0 \text{ in } C_{loc}^2(\mathbb{R}^2), \quad -\Delta\varphi_0 = W_1(x)e^{\varphi_0} \text{ in } \mathbb{R}^2, \quad (2.33)$$

$$\mu = \frac{1}{2\pi} \int_{\mathbb{R}^2} W_1(x)e^{\varphi_0} \quad (W_1 \text{ is given in (2.27)}) \quad (2.34)$$

(iii) There exists a finite blow-up set

$$S_\varphi := \{q : \exists z_k \rightarrow q \text{ and } \varphi_k(z_k) \rightarrow +\infty\}$$

such that

a) either $\varphi_k \rightarrow -\infty$ uniformly on compact sets of $\mathbb{R}^2 \setminus S_\varphi$,

$$W_{1,k} e^{\varphi_k} \rightarrow \sum_{q \in S_\varphi} \sigma(q) \delta_q \text{ weakly in the sense of measures,} \quad (2.35)$$

$$\mu = \frac{1}{2\pi} \sum_{q \in S} \sigma(q) \text{ with } \sigma(q) \geq 4\pi. \quad (2.36)$$

In addition we know that,

$$\begin{aligned} \sigma(q) &= 8\pi \text{ if } q \neq q_j, \forall j = 1, \dots, s, \\ \sigma(q) &= 8\pi(1 + \alpha_j) \text{ if } q = q_j \text{ and } q \neq q_k \text{ for } k \neq j; \end{aligned} \quad (2.37)$$

b) or $\varphi_k \rightarrow \varphi_0$ in $C_{loc}^2(\mathbb{R}^2 \setminus S_\varphi)$,

$$W_{1,k} e^{\varphi_k} \rightarrow \sum_{q \in S} \sigma(q) \delta_q + W_1 e^{\varphi_0} \text{ weakly in the sense of measures,} \quad (2.38)$$

$$-\Delta\varphi_0 = W_1(x)e^{\varphi_0} + \sum_{q \in S_\varphi} \sigma(q) \delta_q \text{ in } \mathbb{R}^2, \quad (2.39)$$

$$\mu = \frac{1}{2\pi} \left(\int_{\mathbb{R}^2} W_1(x)e^{\varphi_0} + \sum_{q \in S_\varphi} \sigma(q) \right). \quad (2.40)$$

Furthermore, when alternative b) holds in (ii) then necessarily: $S_\varphi \subset Z_1$ (see (2.28)), and blow-up takes place at points in Z_1 where different zero points of $W_{1,k}$ collapse together. Namely, for $q \in S_\varphi$ there holds:

$$\begin{aligned} q &= q_{j_1} = q_{j_2} = \dots = q_{j_m} \in Z_1 \\ \text{with } m &\geq 2, \text{ and } 1 \leq j_1 < \dots < j_m \leq s. \end{aligned} \quad (2.41)$$

Moreover, in this case $\sigma(q)$ must satisfy:

$$4\pi \leq \sigma(q) < 4\pi(\alpha(q) + 1) \text{ for } \alpha(q) := \sum_{k=1}^m \alpha(q_{j_k}). \quad (2.42)$$

Indeed, by (2.39), we have that φ_0 admits a logarithmic singularity at q of order $\frac{\sigma(q)}{2\pi}$. At the same time we must ensure the integrability of $|x - q|^{2\alpha(q)} e^{\varphi_0}$ around q (recall (2.27) and see (2.40)), and this is possible only if the inequality given in (2.42) holds.

When either alternative (ii) or b) in (iii) hold, then we can argue similarly around infinity. Therefore, by (2.33) and (2.34), or by (2.39) and (2.40) respectively, we find that,

$$\varphi_0(x) = -\mu \ln(|x|) + O(1), \text{ for } |x| \geq 1; \quad (2.43)$$

(see e.g. [7]), and again, by the given integrability condition, we derive that

$$\mu = 2(\alpha + 1 + \sigma) \text{ for some } \sigma > 0. \quad (2.44)$$

With the help of the information above we can establish the following:

Claim 1: If φ_k satisfies alternative (iii), then $\sigma(q) \in 8\pi\mathbb{N}$, $\forall q \in S_\varphi$ (2.45)

Since $\alpha_j \in \mathbb{N}$, $\forall j = 1, \dots, s$, by virtue of (2.37) to establish the claim we only have to consider the case where $q \in S_\varphi$ satisfies (2.41).

To this purpose, we go back to the original sequence ξ_k . Actually we replace $\xi_k(x)$ with $\xi_k(x + \frac{p_{1,k} + p_{2,k}}{2})$ and $\frac{p_{1,k} + p_{2,k}}{2} \rightarrow 0$, so we can assume, without loss of generality, that,

$$p_{1,k} = -p_{2,k}, \forall k \in \mathbb{N}. \quad (2.46)$$

We proceed by induction on s , the number of different zeroes of W_k collapsing at the origin, the only blow-up point of ξ_k (see (2.15),(2.17),(2.18),(2.21)).

In case $s = 2$, then as shown in [14], we easily reach the desired conclusion. Indeed, by (2.26) and (2.46) we have: $\tau_k = |p_{1,k}| = |p_{2,k}|$, so by letting: $q_k = \frac{p_{1,k}}{\tau_k}$ we have $|q_k| = 1$ and $\frac{p_{2,k}}{\tau_k} = -q_k$. Thus, (along a subsequence) we have:

$$\begin{aligned} -\Delta\varphi_k &= |x - q_k|^{2\alpha_1} |x + q_k|^{2\alpha_2} h_k(\tau_k x) e^{\varphi_k} \text{ in } D_k \\ q_k &\rightarrow q_0 \neq 0, \text{ with } q_1 = q_0 \text{ and } q_2 = -q_0. \end{aligned}$$

So, for the scaled sequence φ_k we do not have to face the possibility of a further "collapsing" of zeroes, and the desired conclusion follows simply by (2.37).

Next, let $s \geq 3$, and assume that $q \in S_\varphi$ satisfies

$$q = q_{j_1} = q_{j_2} = \dots = q_{j_m}, \quad j_i \in \{1, \dots, s\} \quad \text{and} \quad m \geq 2.$$

It suffices to show that $m \leq s - 1$, since then by the induction assumption we could conclude that $\sigma(q) \in 8\pi\mathbb{N}$, as desired. As above, for suitable $q_0 \in \mathbb{R}^2$, we have: $q_1 = q_0$ and $q_2 = -q_0$. In case $q_0 \neq 0$, we see that, if $q \neq \pm q_0$, then $q \in \{q_3, \dots, q_s\}$ and necessarily $m \leq s - 2$. While if $q = q_0$ or $q = -q_0$, then $q \neq q_2$ or $q \neq q_1$ respectively, and again $m \leq s - 1$. Next we suppose that $q_0 = 0$. In case $q \neq 0$, then we conclude as above that $m \leq s - 2$. Hence suppose that $q = q_0 = 0$, and let $j_0 \in \{1, \dots, s\}$ so that (along a subsequence): $\tau_k = |p_{j_0, k}|$. Consequently $|q_{j_0}| = 1$ and so $q \neq q_{j_0}$ and we can still conclude that $m \leq s - 1$. So, if q coincides with m -collapsing zeros in Z_1 , then we have shown that, $2 \leq m \leq s - 1$, and as already mentioned, the desired conclusion follows by the induction hypothesis.

To proceed further, we recall the following:

Lemma A (Lemma 2.1 of [20]). *Let u satisfy*

$$\begin{cases} \Delta u + e^u = 4\pi \sum_{j=1}^N \beta_j \delta_{q_j} & \text{in } \mathbb{R}^2 \\ \int_{\mathbb{R}^2} e^u < +\infty \end{cases}$$

with $\beta_j \in \mathbb{N} \cup \{0\}$ and $q_j \in \mathbb{R}^2$, $j = 1, \dots, N$. Then

$$\int_{\mathbb{R}^2} e^u = 4\pi \left(\sum_{j=1}^N \beta_j + 1 + \sigma \right) \in 8\pi\mathbb{N} \quad \text{with } \sigma > 1.$$

By combining Claim 1 (see (2.45)) and Lemma A we obtain:

$$\text{Claim 2: } \mu \in 4\mathbb{N} \cup \{0\}. \quad (2.47)$$

To establish (2.47), notice that in case φ_k satisfies (i) then $\mu = 0$. While if φ_k satisfies (ii), then we can apply Lemma A to the function,

$$u := \varphi_0 + \sum_{j=1}^s 2\alpha_j \ln |x - q_j|, \quad \alpha_j \in \mathbb{N} \quad (2.48)$$

to conclude, that

$$\mu = \frac{1}{2\pi} \int_{\mathbb{R}^2} \prod_{j=1}^s |x - q_j|^{2\alpha_j} e^{\varphi_0} = \frac{1}{2\pi} \int_{\mathbb{R}^2} e^u \in 4\mathbb{N}.$$

On the other hand, if φ_k satisfies alternative a) of (iii), then by (2.36) and (2.45), still we find: $\mu \in 4\mathbb{N}$.

Finally, if alternative b) in (iii) holds, then with the notation in (2.41),(2.42) we see that u in (2.48) satisfies:

$$\Delta u + e^u = 4\pi \sum_{q_j \in Z_1 \setminus S_\varphi} \alpha_j \delta_{q_j} + 4\pi \sum_{q \in S_\varphi} \left(\alpha(q) - \frac{\sigma(q)}{4\pi} \right) \delta_q$$

and in view of (2.42), we can check that Lemma A applies to u . As a consequence

$$\int_{\mathbb{R}^2} W_1(x) e^{\varphi_0} = \int_{\mathbb{R}^2} \prod_{j=1}^s |x - q_j|^{2\alpha_j} e^{\varphi_0} = \int_{\mathbb{R}^2} e^u \in 8\pi\mathbb{N}.$$

Therefore, by (2.40) and (2.45), we derive that $\mu \in 4\mathbb{N}$ as desired.

Notice in particular that, when φ_k satisfies either (ii) or alternative b) of (iii), then μ must satisfy (2.44) with $\sigma \in \mathbb{N}$.

Finally to conclude the proof of Theorem 2, we recall the following relation between m and μ established e.g. in [15]:

$$m^2 - \mu^2 = 4(1 + \alpha)(m - \mu). \quad (2.49)$$

For completeness we have included a proof of (2.49) in the Appendix.

At this point we see that, if φ_k satisfies (i) then $\mu = 0$ and from (2.49) we find $m = 4(\alpha + 1) \in 4\mathbb{N}$, as $\alpha = \sum_{j=1}^s \alpha_j \in \mathbb{N}$. In case φ_k satisfies (ii) then by (2.44) and (2.47), $\mu \in 4\mathbb{N}$ and $\mu > 2(\alpha + 1)$. Therefore, from (2.49) we must have that necessarily $m = \mu \in 4\mathbb{N}$. Finally, if φ_k satisfies alternative (iii), then again from (2.49) we have that, either $m = \mu \in 4\mathbb{N}$ or $m = 4(\alpha + 1) - \mu$, with $\alpha \in \mathbb{N}$ and $\mu \in 4\mathbb{N}$. Thus, in any case, $m \in 4\mathbb{N}$, and the proof is completed. \square

The "local" results above can be used to describe the asymptotic behaviour of solutions for Liouville-type equations on a compact Riemann surface (X, g) . Denote by $d_g(\cdot, \cdot)$ the distance in (X, g) . We consider the sequence $v_k \in C^{2,\alpha}(X)$ to satisfy:

$$-\Delta v_k = R_k e^{v_k} - f_k \text{ in } X, \quad (2.50)$$

$$R_k(z) = \left(\prod_{j=1}^N (d_g(z, z_{j,k}))^{2\alpha_j} \right) g_k(z), \quad z \in X; \quad (2.51)$$

with

$$g_k \in C^1(X) : a \leq g_k \leq b, \quad |\nabla g_k| \leq A \text{ and } g_k \rightarrow g_0 \text{ in } C^0(X); \quad (2.52)$$

$$z_{j,k} \in X : z_{j,k} \neq z_{l,k}, \quad j \neq l \text{ and } z_{j,k} \rightarrow z_j, \quad j = 1, \dots, N; \quad (2.53)$$

$$f_k \in C^{0,\alpha}(X), \quad f_k \rightarrow f_0 \text{ in } L^p(X) \text{ for some } p > 1, \quad \int_X f_0 dA \neq 0; \quad (2.54)$$

and so,

$$R_k(z) \rightarrow R_0(z) := \left(\prod_{j=1}^N (d_g(z, z_j))^{2\alpha_j} \right) g_0(z) \text{ in } C^0(X).$$

As before we assume that,

$$\alpha_j \in \mathbb{N}, \quad j = 1, \dots, N. \quad (2.55)$$

We denote by $Z = \{z \in X : R_0(z) = 0\} = \{z_1, \dots, z_N\}$ the zero set of R_0 , formed by the points z_j given in (2.53), for $j = 1, \dots, N$. Again we must keep in mind that such points may not be distinct, since, at the limit, different zeroes of R_k could coalesce at the same zero of R_0 . Therefore, we let $Z_0 \subset Z$ be the subset (possibly empty) of Z , given by such "collapsing" zeroes, namely:

$$Z_0 = \{z \in Z : \exists s \geq 2, 1 \leq j_1 < \dots < j_s \leq N \text{ such that} \\ z = z_{j_1} = \dots = z_{j_s} \text{ and } z \notin Z \setminus \{z_{j_1}, \dots, z_{j_s}\}\}.$$

By combining the "local" results obtained above, we can establish the following:

Theorem 3. *Let v_k satisfy (2.50) and assume (2.51)-(2.55). Then, along a subsequence, one of the following alternatives holds:*

(i) (compactness) : $v_k \rightarrow v_0$ in $C^2(X)$ with

$$-\Delta v_0 = R_0 e^{v_0} - f_0, \quad \text{in } X \quad (2.56)$$

(ii) (blow-up) : There exists a finite blow-up set

$$\mathcal{S} = \{q \in X : \exists q_k \rightarrow q \text{ and } v_k(q_k) \rightarrow +\infty, \text{ as } k \rightarrow +\infty\}$$

such that, v_k^+ is uniformly bounded in $C_{loc}^0(X \setminus \mathcal{S})$ and, as $k \rightarrow +\infty$,

(a) either (blow-up with concentration) :

$$v_k \rightarrow -\infty \text{ uniformly on compact sets of } X \setminus \mathcal{S}, \\ R_k e^{v_k} \rightharpoonup \sum_{q \in \mathcal{S}} \sigma(q) \delta_q \text{ weakly in the sense of measures, } \sigma(q) \in 8\pi\mathbb{N}.$$

In particular, $\int_X f_0 dA \in 8\pi\mathbb{N}$ in this case.

(b) or (blow-up without concentration) :

$$v_k \rightarrow v_0 \text{ in } C_{loc}^2(X \setminus \mathcal{S}), \\ R_k e^{v_k} \rightharpoonup R_0 e^{v_0} + \sum_{q \in \mathcal{S}} \sigma(q) \delta_q \text{ weakly in the sense of measures,} \\ -\Delta v_0 = R_0 e^{v_0} + \sum_{q \in \mathcal{S}} \sigma(q) \delta_q - f_0 \text{ in } X, \sigma(q) \in 8\pi\mathbb{N}.$$

Furthermore, if alternative b) of (ii) holds then $\mathcal{S} \subset Z_0$ and so, in this case, blow-up occurs only at points where different zeroes of R_k coalesce at the limit.

Proof. If $\max_X v_k \leq C$, then the right hand side of (2.50) is uniformly bounded in $L^p(X)$, $p > 1$. Hence, by setting $v_k = w_k + d_k$ with $d_k = \int_X v_k$, then by elliptic estimates (see [1]), we see that w_k is uniformly bounded in $C^{1,\alpha}(X)$, and

so along a subsequence, we obtain that, $w_k \rightarrow w_0$ in $C^1(X)$, as $k \rightarrow +\infty$. After integration of (2.50), we have:

$$\int_X e^{v_k} R_k = \int_X f_k \rightarrow \int_X f_0, \text{ as } k \rightarrow +\infty. \quad (2.57)$$

Since by assumption: $\int_X f_0 \neq 0$, then from (2.57) we see that necessarily, $\int_X f_0 > 0$, and so $d_k \rightarrow \log(\frac{\int_X f_0}{\int_X R_0 e^{w_0}}) = d_0$. Consequently, $v_k \rightarrow v_0 = w_0 + d_0$ in $C^1(X)$ and v_0 satisfies (2.56). Thus alternative (i) holds in this case.

Next assume that, (along a subsequence)

$$\max_X v_k \rightarrow +\infty, \text{ as } k \rightarrow +\infty.$$

This implies that the blow-up set of (a subsequence of) v_k is not empty, that is:

$$\mathcal{S} = \{q \in X \mid \exists q_k \in X : q_k \rightarrow q \text{ and } v_k(q_k) \rightarrow +\infty\} \neq \emptyset.$$

At this point, around any $q \in X$, we introduce local conformal coordinates centered at the origin, and (with abuse of notation) we shall denote in the same way the local expressions of v_k and f_k in such coordinates, as defined in a small ball B_r . In this way, in B_r we may consider the sequence θ_k satisfying:

$$-\Delta\theta_k = f_k \text{ in } B_r \text{ and } \theta_k = 0 \text{ in } \partial B_r.$$

Hence, θ_k is uniformly bounded in $C^{1,\alpha}(X)$ and (along a subsequence) we may assume that, $\theta_k \rightarrow \varphi_0$ in $C^1(X)$, as $k \rightarrow +\infty$, with $\Delta\theta_0 = f_0$ in B_r and $\theta_0 = 0$ in ∂B_r . Therefore, the new sequence $\xi_k = v_k - \theta_k$ satisfies:

$$-\Delta\xi_k = W_k e^{\xi_k} \text{ in } B_r$$

with suitable $W_k \geq 0$ such that, in B_r the following holds:

$$|\nabla W_k| \leq A \text{ and } W_k \rightarrow W_0 \text{ uniformly.}$$

From (2.57) also we have:

$$\int_{B_r} W_k e^{\xi_k} \leq C,$$

with suitable constant $C > 0$. Furthermore,

$$\begin{aligned} &\text{if } W_0(0) = 0 \text{ then } W_k(z) = \prod_{j=1}^s |z - p_{j,k}|^{2\alpha_j} h_k(z) \text{ for } s \geq 2, \\ &\text{and } h_k \text{ satisfies (2.12) in } B_r, \alpha_j \in \mathbb{N} \text{ and } p_{j,k} \rightarrow 0, \text{ as } k \rightarrow \infty. \end{aligned}$$

In any case, W_k satisfies (2.4) in B_r (with suitable $\varepsilon_0 > 0$) and therefore, by Remark 2.1 and (2.57) we conclude that the blow-up set \mathcal{S} is finite. Also, by using the Green representation formula for v_k in X then, in the usual way (see e.g. [2] or [26]), for every compact set $K \subset X \setminus \mathcal{S}$ we can check that,

$$\max_K v_k - \min_K v_k \leq C, \quad (2.58)$$

with suitable $C > 0$ depending only on K .

Thus, by (2.58), we can apply alternative (iii) of Proposition 2.1 and Theorem 1 to obtain (in conformal coordinates) the local blow-up description of v_k around any blow-up point $q \in \mathcal{S}$. At the same time the property (2.58) also allows us to patch together such local information and arrive at the desired statement in (ii). □

Furthermore, as a direct consequence of Theorem 3, we can recover a compactness result, well known in the non-collapsing case:

Corollary 2.1. *Under the assumptions of Theorem 3, if*

$$\limsup_{k \rightarrow +\infty} \int_X R_k e^{\xi_k} < 8\pi$$

then, along a subsequence, $\xi_k \rightarrow \xi_0$ in $C^{1,\alpha}(X)$ with ξ_0 satisfying (2.56).

3 Local estimates in case of least blow-up mass.

This section is devoted to provide a more detailed description about a sequence ξ_k which satisfies the local problem (2.15) and admits a blow-up point at the origin and (2.18) holds. Namely, ξ_k blows-up at a point where different zeroes of the weight function W_k "collapse" together.

We focus to the case of least blow-up mass 8π , namely when (2.23) holds as follows:

$$m = \lim_{\delta \searrow 0} \liminf_{k \rightarrow +\infty} \frac{1}{2\pi} \int_{B_\delta} W_k e^{\xi_k} = 4. \quad (3.1)$$

This is a first important step towards the more involved description of "multiple" blow-up profiles, and we refer to [14] for some progress in this direction, $s = 2$ in (2.18).

With the notation of the previous section, a first consequence of (3.1) is the following:

Proposition 3.1. *Let ξ_k satisfy (2.15)-(2.21) and suppose that (3.1) holds. Then $m = \mu = 4$ and the sequence φ_k in (2.29) must satisfy alternative a) of (iii) in Proposition 2.1, with a **unique** blow-up point q_0 .*

Furthermore if $W_1(q_0) = 0$ (i.e. $q_0 \in Z_1 = \{q_1, \dots, q_s\}$) then q_0 must satisfy (2.41), namely different zeroes of $W_{1,k}$ collapse at q_0 .

Proof. First of all we can exclude that φ_k satisfies alternative (i). Indeed, in this case $\mu = 0$, and so by (2.49) we would have $m = 4(1 + \alpha) > 4$, in contradiction with (3.1). Therefore, $0 < \mu \leq m$, and since by (2.47) we have $\mu \in 4\mathbb{N}$, we see that necessarily $m = \mu = 4$. This fact allows us to conclude also that φ_k cannot satisfy alternative (ii) or alternative b) of (iii). In fact, in this situation, by (2.44), we would have: $\mu = 2(\alpha + 1 + \sigma)$, with α and σ positive integers, a

contradiction to (3.1). So φ_k can only satisfy alternative a) of (iii) with a single blow-up point q_0 . Furthermore, if $q_0 \in Z_1 = \{q_1, \dots, q_s\}$, then (2.37) together with (3.1) allow us to conclude that, $s \geq 2$ and different zeroes of $W_{1,k}$ must converge to q_0 . \square

To proceed further, we use a different more convenient normalization for ξ_k from the previous section. In fact, we observe that, the translated function: $\xi_k(x + x_k)$ (with the point x_k defined in (2.17)) satisfies an analogous problem, possibly in a smaller ball around the origin, where properties (2.15)-(2.21) continue to hold, simply with the point $p_{j,k}$ replaced by $(p_{j,k} - x_k) \rightarrow 0$, as $k \rightarrow +\infty$. More importantly, since $x_k \rightarrow 0$ for $k \rightarrow +\infty$, then both sequences $\xi_k(x)$ and its translated $\xi_k(x + x_k)$ admit the same value of m in (2.23). Thus, without loss of generality, we can assume that,

$$\xi_k(0) = \max_{B_r} \xi_k \rightarrow +\infty, \text{ as } k \rightarrow +\infty. \quad (3.2)$$

Also, after relabeling the indices, (along a subsequence) we may take:

$$0 \leq |p_{1,k}| \leq |p_{2,k}| \leq \dots \leq |p_{s,k}|, \text{ with } s \geq 2; \quad (3.3)$$

so that,

$$\tau_k = |p_{s,k}| > 0, \quad (3.4)$$

$$q_{j,k} = \frac{p_{j,k}}{\tau_k} \rightarrow q_j, \text{ as } k \rightarrow +\infty, \forall j = 1, \dots, s, \quad (3.5)$$

$$0 \leq |q_1| \leq |q_2| \leq \dots \leq |q_s| = 1. \quad (3.6)$$

It is important to notice that, with this new normalization, we no longer expect (2.46) to hold. We have:

Theorem 4. *Let ξ_k satisfy (2.15),(2.16),(2.18)-(2.21) and assume (3.1)-(3.6). Then the points $p_{j,k} \neq 0$, $1 \leq j \leq s$ and there exists $s_1 \in \{2, \dots, s\}$ such that (along a subsequence) the following holds, as $k \rightarrow +\infty$:*

$$z_{j,k} := \frac{p_{j,k}}{|p_{s_1,k}|} \rightarrow z_j \neq 0, \forall j = 1, \dots, s_1, \quad (3.7)$$

and if $s_1 < s$ then

$$\frac{p_{j,k}}{|p_{s,k}|} \rightarrow q_j \neq 0 \text{ and } \frac{|p_{j,k}|}{|p_{s_1,k}|} \rightarrow +\infty, \forall j = s_1 + 1, \dots, s. \quad (3.8)$$

Moreover

$$\xi_k(0) + 2 \ln |p_{s_1,k}| + 2 \sum_{j=1}^s \alpha_j \ln |p_{j,k}| \rightarrow +\infty. \quad (3.9)$$

Proof. If

$$q_{j,k} = \frac{p_{j,k}}{|p_{s,k}|} \rightarrow q_j \neq 0, \forall j = 1, \dots, s \quad (3.10)$$

then it suffices to choose $s = s_1$. Indeed, we only need to check (3.9). To this purpose, by the normalization (3.2), we see that $q_0 = 0$ in Proposition 3.1. So for the sequence φ_k in (2.29) we have

$$\varphi_k(0) = \sup_{D_k} \varphi_k \longrightarrow +\infty. \quad (3.11)$$

On the other hand, by (2.29), (3.4) and (3.10), we find:

$$\begin{aligned} \varphi_k(0) &= \xi_k(0) + 2 \ln \tau_k + 2\alpha \ln |p_{s,k}| \\ &= \xi_k(0) + 2 \ln |p_{s,k}| + 2 \sum_{j=1}^s \alpha_j \ln |p_{j,k}| + O(1), \end{aligned}$$

and so, (3.9) follows with $s_1 = s$ in this case. Since $|q_s| = 1$, let us now assume that there exists $\bar{s} \in \{1, \dots, s-1\}$ such that,

$$q_1 = q_2 = \dots = q_{\bar{s}} = 0, \quad \text{and} \quad q_j \neq 0 \quad \text{for} \quad j = \bar{s} + 1, \dots, s. \quad (3.12)$$

Since the origin is a blow-up point for φ_k , by Proposition 3.1 we know that (3.12) must hold with $\bar{s} \geq 2$. Clearly, Theorem 1 applies to φ_k around the origin, and it implies that,

$$m_\varphi = \lim_{\delta \searrow 0} \lim_{k \rightarrow +\infty} \frac{1}{2\pi} \int_{B_\delta} W_{1,k} e^{\varphi_k} \in 4\mathbb{N}.$$

Since $m_\varphi \leq m = 4$, we see that necessarily $m_\varphi = 4$. This information allows us to apply Proposition 3.1 to φ_k (around the origin), with $0 \leq |q_{1,k}| \leq |q_{2,k}| \leq \dots \leq |q_{\bar{s},k}| \rightarrow 0$, as $k \rightarrow +\infty$. Therefore, by setting:

$$\tau_{1,k} = |q_{\bar{s},k}| \quad \text{and} \quad \bar{\alpha} = \sum_{j=1}^{\bar{s}} \alpha_j \in \mathbb{N}, \quad \bar{\alpha} \geq 2, \quad (3.13)$$

for the new "scaled" function:

$$\varphi_{1,k}(x) := \varphi_k(\tau_{1,k}x) + 2(\bar{\alpha} + 1) \ln \tau_{1,k}$$

we know that (along a subsequence), $\varphi_{1,k}(0) \rightarrow \infty$, as $k \rightarrow +\infty$, and actually the origin is the only blow-up point of $\varphi_{1,k}$, where "concentration" occurs, as described by alternative a) of (iii) in Proposition 2.1.

Furthermore (along a subsequence), as $k \rightarrow +\infty$, we have:

$$\frac{q_{j,k}}{\tau_{1,k}} \longrightarrow q_j^{(1)}, \quad \forall j = 1, \dots, \bar{s};$$

with $0 \leq |q_1^{(1)}| \leq |q_2^{(1)}| \leq \dots \leq |q_{\bar{s}}^{(1)}| = 1$. If

$$q_j^{(1)} \neq 0, \quad \forall j = 1, \dots, \bar{s}; \quad (3.14)$$

then from (3.12) we deduce that, as $k \rightarrow +\infty$,

$$\begin{aligned} \frac{|p_{j,k}|}{|p_{\bar{s},k}|} &= \frac{|q_{j,k}|}{\tau_{1,k}} \rightarrow |q_j^{(1)}| \neq 0, \quad \forall j = 1, \dots, \bar{s}, \\ \frac{|p_{j,k}|}{|p_{s,k}|} &\rightarrow |q_j| \neq 0, \quad \frac{|p_{j,k}|}{|p_{\bar{s},k}|} = \frac{|q_{j,k}|}{\tau_{1,k}} \rightarrow +\infty, \quad \forall j = \bar{s} + 1, \dots, s. \end{aligned} \quad (3.15)$$

So, if (3.14) holds then (3.7) and (3.8) are verified with $s_1 = \bar{s}$.

Moreover, to show that also (3.9) holds with $s_1 = \bar{s}$, simply note that $\varphi_{1,k}(0) \rightarrow +\infty$, as $k \rightarrow +\infty$, and in view of (2.29),(3.13),(3.15) we have

$$\begin{aligned} \varphi_{1,k}(0) &= \varphi_k(0) + 2(\bar{\alpha} + 1) \ln \tau_{1,k} \\ &= \xi_k(0) + 2(\alpha + 1) \ln |p_{k,s}| + 2(\bar{\alpha} + 1) \ln \frac{|p_{k,\bar{s}}|}{|p_{k,s}|} \\ &= \xi_k(0) + 2(\bar{\alpha} + 1) \ln |p_{k,\bar{s}}| + 2(\alpha - \bar{\alpha}) \ln |p_{k,s}| \\ &= \xi_k(0) + 2 \ln |p_{k,\bar{s}}| + \sum_{j=1}^s 2\alpha_j \ln |p_{k,j}| + O(1). \end{aligned}$$

Hence the desired conclusion follows in this case.

On the contrary, if $q_j^{(1)}$ vanishes for some $j \in \{1, \dots, \bar{s}\}$, then we can repeat the argument above for the sequence $\varphi_{1,k}$, and eventually continue in this way, by taking further scalings. However, at each step the number of "collapsing" zeroes decreases, and moreover Proposition 3.1 applies to each of the new scaled sequences. Therefore, this procedure must stop after finitely many steps.

In case we assume "by contradiction" that $p_{1,k} = 0$ identically, then we would end up with a sequence blowing-up at zero and satisfying a Liouville type problem with a weight function vanishing at the origin with the same order of $|x|^{2\alpha_1}$. Thus, to such a sequence, we can apply (2.42) (see [3]) and obtain a blow-up mass $m = \frac{\sigma(0)}{2\pi} = 4(1 + \alpha_1) > 4$, a contradiction. Hence $p_{1,k} \neq 0$, and so by (2.18) and (3.3) we conclude that $p_{j,k} \neq 0$ for all $1 \leq j \leq s$, as claimed. Consequently, after finitely many steps, we must arrive at sequence which blows-up at zero and satisfying a Liouville-type problem with a weight function never vanishing around the origin. In other words we have found $s_1 \in \{2, \dots, s\}$ for which (3.15) holds with \bar{s} replaced by s_1 . And as before, we check that (3.9) is also satisfied. \square

Theorem 4 identifies the appropriate scale to use in order to gain good control on ξ_k in a tiny neighborhood around the origin. Indeed, according to Theorem 4, we define:

$$\varepsilon_k = |p_{s_1,k}| \rightarrow 0, \quad \text{as } k \rightarrow +\infty \quad (3.16)$$

and let

$$z_{j,k} = \frac{p_{j,k}}{\varepsilon_k} \rightarrow z_j \neq 0 \quad \text{for } j = 1, \dots, s_1.$$

If $s_1 < s$, we set:

$$\varepsilon_{j,k} := \frac{\varepsilon_k}{|p_{j,k}|} \rightarrow 0, \quad \hat{p}_{j,k} := \frac{p_{j,k}}{|p_{j,k}|} \rightarrow \hat{p}_j, \quad |\hat{p}_j| = 1 \quad \forall j = s_1 + 1, \dots, s.$$

We define

$$v_k(x) = \xi_k(\varepsilon_k x) + 2 \ln \varepsilon_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}|,$$

satisfying:

$$-\Delta v_k(x) = \prod_{j=1}^{s_1} |x - z_{j,k}|^{2\alpha_j} V_k(x) e^{v_k(x)} \quad \text{in } D_k = B_{\frac{r}{\varepsilon_k}}$$

with

$$V_k(x) = \prod_{j=s_1+1}^s |\varepsilon_{j,k} x - \hat{p}_{j,k}|^{2\alpha_j} h_k(\varepsilon_k x) A_k \quad (3.17)$$

(in case $s_1 = s$ the product term in (3.17) is omitted), and

$$A_k = \prod_{j=1}^{s_1} \left(\frac{\varepsilon_k}{|p_{j,k}|} \right)^{2\alpha_j} \longrightarrow \prod_{j=1}^{s_1} \left(\frac{1}{|z_j|} \right)^{2\alpha_j} = A_0 > 0, \quad \text{as } k \longrightarrow +\infty.$$

Recall that, h_k satisfies (2.12) and (2.20) in B_r , and therefore:

$$R_k(x) := \prod_{j=1}^{s_1} |x - z_{j,k}|^{2\alpha_j} V_k(x) \longrightarrow R_0(x) := A_0 \prod_{j=1}^{s_1} |x - z_j|^{2\alpha_j}. \quad (3.18)$$

Clearly, the zero set of R_0 in (3.18) is given by:

$$Z = \{z_1, \dots, z_{s_1}\},$$

where the z_j 's may not be necessarily distinct, but they satisfy:

$$0 < \delta_0 < |z_1| \leq |z_2| \leq \dots \leq |z_{s_1}| < \frac{L_0}{4} \quad (3.19)$$

with suitable $\delta_0 > 0$ and $L_0 \geq 1$. Also we have:

$$\int_{D_k} R_k(x) e^{v_k} \leq C.$$

More importantly, in view of (3.7),(3.8),(3.9), we can check that,

$$v_k(0) = \max_{D_k} v_k \longrightarrow +\infty, \quad \text{as } k \longrightarrow +\infty.$$

By (3.1), we see that the origin is the only blow-up point of v_k , where the well known blow-up analysis of [5],[18],[3] applies. As a consequence we find:

$$R_k e^{v_k} \rightharpoonup 8\pi\delta_0 \quad \text{weakly in the sense of measures, as } k \longrightarrow +\infty, \quad (3.20)$$

locally on compact sets. Furthermore (see Corollary 5.6.57 of [26]), for every $R > 0$ the following well known estimate holds:

$$\left| v_k(x) - \ln \frac{e^{v_k(0)}}{\left(1 + \frac{e^{v_k(0)}}{8} R_k(0) |x|^2\right)^2} \right| \leq C_R \quad \text{in } \overline{B}_R, \quad (3.21)$$

with a suitable constant $C_R > 0$ depending on R only. In particular, from (3.21) we derive:

$$|v_k(x) + v_k(0)| \leq C_R, \quad \forall x \in \partial B_R. \quad (3.22)$$

$$\max_{\partial B_R} v_k - \min_{\partial B_R} v_k \leq C_R. \quad (3.23)$$

Lemma 3.1. *Let $L_0 > 1$ fixed to satisfy (3.19). Then for every $R > L_0$ there exists C_R such that,*

$$\max_{\frac{1}{2}L_0 \leq |y| \leq R} \{v_k(y) + 2(\bar{\alpha} + 1) \ln |y|\} \leq C_R \quad (3.24)$$

with $\bar{\alpha} = \sum_{j=1}^{s_1} \alpha_j \in \mathbb{N}$.

Proof. We start by observing the following:

Claim: For all $\varepsilon > 0$ there exist $k_\varepsilon \in \mathbb{N}$ and $r_\varepsilon > 0$ such that, for $\delta \in (0, r_\varepsilon)$ and $R \geq \frac{L_0}{4}$ we have:

$$\int_{B_\delta \setminus B_{R\varepsilon_k}} W_k e^{\xi_k} = \int_{B_{\frac{\delta}{\varepsilon_k}} \setminus B_R} R_k e^{v_k} < \varepsilon, \quad \forall k \geq k_\varepsilon; \quad (3.25)$$

with ε_k given in (3.16).

To establish (3.25), recall that by (3.1) we find $k_\varepsilon \in \mathbb{N}$ and $r_\varepsilon > 0$ such that

$$\int_{B_\delta} W_k e^{\xi_k} \leq 8\pi + \frac{\varepsilon}{2}, \quad \forall k \geq k_\varepsilon \quad \text{and} \quad \delta \in (0, r_\varepsilon).$$

On the other hand, by taking k_ε larger if necessary, from (3.20), also we have that

$$\int_{B_{\frac{L_0\varepsilon_k}{4}}} W_k e^{\xi_k} = \int_{B_{\frac{L_0}{4}}} R_k e^{v_k} \geq 8\pi - \frac{\varepsilon}{2}, \quad \forall k \geq k_\varepsilon$$

and we immediately derive (3.25).

To establish (3.24), we argue by contradiction and assume there exists $R_1 > L_0$, such that

$$\exists y_k \in B_{R_1} \setminus B_{\frac{L_0}{2}} : v_k(y_k) + 2(\bar{\alpha} + 1) \ln |y_k| \longrightarrow \infty, \quad \text{as } k \longrightarrow +\infty.$$

Define:

$$\psi_k(x) = v_k(|y_k|x) + 2(\bar{\alpha} + 1) \ln |y_k|, \quad \text{for } x \in \Omega := \left\{ \frac{1}{2} < |x| < 2 \right\}$$

satisfying:

$$-\Delta \psi_k(x) = \left(\prod_{j=1}^{s_1} \left| x - \frac{z_{j,k}}{|y_k|} \right|^{2\alpha_j} \right) V_k(|y_k|x) e^{\psi_k} \quad \text{in } \Omega,$$

$$\psi_k\left(\frac{y_k}{|y_k|}\right) \longrightarrow +\infty, \quad \text{as } k \longrightarrow +\infty.$$

Setting:

$$R_{1,k}(x) = \prod_{j=1}^{s_1} \left| x - \frac{z_{j,k}}{|y_k|} \right|^{2\alpha_j} V_k(|y_k|x),$$

in view of (3.19), we check easily that there exist $0 < a_1 \leq b_1$ and $A > 0$ such that

$$0 < a_1 \leq R_{1,k}(x) \leq b_1 \quad \text{and} \quad |\nabla R_{1,k}|(x) \leq A \quad x \in \Omega, \quad \int_{\Omega} R_{1,k} e^{\psi_k} \leq C.$$

Therefore, if along a subsequence, we assume that,

$$\frac{y_k}{|y_k|} \longrightarrow y_0, \quad \text{as } k \longrightarrow \infty,$$

then $|y_0| = 1$, and so y_0 is a blow-up point of ψ_k in Ω . As above, from [5],[18],[3] we have (along a subsequence)

$$R_{1,k} e^{\psi_k} \rightharpoonup 8\pi \delta_{y_0}, \quad \text{weakly in the sense of measures in } \Omega.$$

and therefore, for $\delta > 0$ sufficiently small, there holds (along a subsequence):

$$\int_{\{|z-|y_k|y_0| < \delta|y_k|\}} R_k(z) e^{v_k(z)} dz = \int_{B_\delta(y_0)} R_{1,k} e^{\psi_k} \geq 4\pi, \quad \text{as } k \longrightarrow +\infty.$$

Consequently,

$$\int_{\{(1-\delta)\frac{L_0}{2} \leq |z| \leq (1+\delta)R_1\}} R_k(z) e^{v_k(z)} dz \geq 4\pi, \quad \text{as } k \longrightarrow +\infty,$$

a contradiction to (3.25). \square

We can reformulate (3.24) in terms of ξ_k as follows:

$$\xi_k(x) + 2(\bar{\alpha} + 1) \ln |x| + \sum_{j=s_1+1}^s 2\alpha_j \ln |p_{j,k}| \leq C_k, \quad \frac{L_0}{2} \varepsilon_k \leq |x| \leq R\varepsilon_k, \quad (3.26)$$

where the summation term in (3.26) should be dropped in case $s_1 = s$. But since for large k we have:

$$0 \leq W_k(x) \leq C_R |x|^{2\bar{\alpha}} \prod_{j=s_1+1}^s |p_{j,k}|^{2\alpha_j} \quad \text{for } \frac{L_0}{2} \varepsilon_k \leq |x| \leq R\varepsilon_k,$$

then from (3.26) we find:

$$0 \leq W_k(x) e^{\xi_k} \leq \frac{C_R}{|x|^2} \quad \text{for } \frac{L_0}{2} \varepsilon_k \leq |x| \leq R\varepsilon_k \quad (3.27)$$

with suitable $C_R > 0$ and k large.

Lemma 3.2. *For $\varepsilon > 0$ sufficiently small there exist $k_\varepsilon \in \mathbb{N}$ and $C_\varepsilon > 0$ such that,*

$$\xi_k(x) \leq \min_{\partial B_r} \xi_k + (4 + \varepsilon) \ln \frac{1}{|x|} + C_\varepsilon, \quad \text{for } x \in B_r \setminus B_{L_0 \varepsilon_k} \quad \text{and } k \geq k_\varepsilon. \quad (3.28)$$

Proof. Firstly, let us fix $\delta \in (0, r)$ sufficiently small, so that for large k there holds:

$$M_k(\delta) := \frac{1}{2\pi} \int_{B_\delta(0)} W_k e^{\xi_k} < 4 + \frac{\varepsilon}{2} \quad \text{and} \quad \int_{B_\delta \setminus B_{L_0 \varepsilon_k}} W_k e^{\xi_k} < 4\pi\varepsilon. \quad (3.29)$$

Since (3.28) clearly holds in $B_r \setminus B_\delta$, we are left to establish it in the set

$$\Omega_{k,\delta} = \{x : R_0 \varepsilon_k \leq |x| \leq \delta\}.$$

In view of (3.29) we can establish the following:

$$\textbf{Claim} : \quad \text{The inequality (3.28) holds for } x : |x| = L_0 \varepsilon_k. \quad (3.30)$$

To obtain (3.30) we use (2.24) to write:

$$\begin{aligned} \xi_k(x) &= \min_{\partial B_r} \xi_k + M_k(\delta) \ln \frac{1}{|x|} + \frac{1}{2\pi} \int_{\{|x| \leq \delta\}} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} \\ &\quad + \int_{\{\delta \leq |y| \leq r\}} \ln\left(\frac{1}{|x-y|}\right) W_k(y) e^{\xi_k(y)} + O(1). \end{aligned}$$

But, for $\delta \leq |y| \leq r$ and $|x| = L_0 \varepsilon_k$, we see that,

$$|\ln|x-y|| \leq |\ln|y|| + C, \quad \text{and} \quad W_k(y) e^{\xi_k(y)} \leq C \text{ (by (3.27))},$$

and we deduce:

$$\xi_k(x) \leq \min_{\partial B_r} \xi_k + M_k(\delta) \ln \frac{1}{|x|} + \frac{1}{2\pi} \int_{\{|y| \leq \delta\}} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} + C. \quad (3.31)$$

In order to estimate the integral term in (3.31), we let

$$D(x) = \{|y| \leq \frac{|x|}{2}\} \cup \{|x-y| \geq \frac{|x|}{2} \text{ and } |y| \leq 2|x|\}$$

and observe that, if $y \in D(x)$, then $|\ln \frac{|x|}{|x-y|}| \leq 4$ and consequently

$$\left| \int_{D(x)} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} \right| \leq C.$$

Moreover, if $2|x| \leq |y| \leq \delta$, then $\frac{|x|}{|x-y|} \leq \frac{|x|}{|y|} \frac{1}{|\frac{y}{|y|} - \frac{x}{|x|}|} \leq 1$ and therefore

$$\int_{\{2|x| \leq |y| \leq \delta\}} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} \leq 0.$$

So we are left to estimate from above the given integral term on $B_{\frac{|x|}{2}}(x)$.

Actually, we can easily check (as above) that, for any $\sigma \in (0, \frac{1}{2})$, we can find a suitable constant $C_\sigma > 0$ such that

$$\left| \int_{\{\sigma|x| \leq |y-x| \leq \frac{|x|}{2}\}} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} dy \right| \leq C_\sigma.$$

Next notice that, if $y \in B_{\sigma|x|}(x)$ then $(1-\sigma)L_0\varepsilon_k \leq |y| \leq (1+\sigma)L_0\varepsilon_k$, and so we can use (3.27) to estimate

$$\begin{aligned} \int_{B_{\sigma|x|}(x)} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y) e^{\xi_k(y)} dy &\leq C \int_{\{|\frac{x}{|x|} - \frac{y}{|y|}| \leq \sigma\}} \ln\left(\frac{1}{|\frac{x}{|x|} - \frac{y}{|y|}|\right) \frac{1}{|y|^2} dy \\ &= C \int_{\{|\frac{x}{|x|} - z| \leq \sigma\}} \ln\left(\frac{1}{|\frac{x}{|x|} - z|\right) \frac{1}{|z|^2} dz \leq C. \end{aligned}$$

This information together with (3.29), implies (3.30).

To proceed further, we define:

$$\phi_k(x) = \xi_k(x) - \min_{\partial B_r} \xi_k - (4 + \varepsilon) \ln \frac{1}{|x|}, \quad x \in \Omega_{k,\delta},$$

and in view of (3.30) we know that, ϕ_k is uniformly bounded from above on $\partial\Omega_{k,\delta}$, and consequently it satisfies:

$$\begin{cases} -\Delta\phi_k = W_k e^{\xi_k} & \text{in } \Omega_{k,\delta} \\ \phi_k \leq C & \text{in } \partial\Omega_{k,\delta}. \end{cases}$$

We are going to apply a well known lemma from [5] (see e.g. Lemma 5.2.1 of [26]) to the function ϕ_k satisfying:

$$\begin{cases} -\Delta\tilde{\phi}_k = \tilde{f}_k & \text{in } B_\delta \\ \tilde{\phi}_k = C & \text{in } \partial B_\delta \end{cases} \quad \text{with } \tilde{f}_k = \begin{cases} W_k e^{\xi_k} & \text{in } \Omega_{k,\delta} \\ 0 & \text{otherwise} \end{cases} \quad (3.32)$$

Thus, as a consequence of the second inequality in (3.29) and Lemma 5.2.1 of [26], for any $1 \leq q < \frac{1}{\varepsilon}$ we find a constant $c_\varepsilon = c_\varepsilon(q) > 0$ such that, $\|e^{\tilde{\phi}_k}\|_{L^q(B_\delta)} \leq c_\varepsilon$. Moreover, by the maximum principle, we know that

$$\phi_k \leq \tilde{\phi}_k \quad \text{in } \Omega_{k,\delta} \quad \text{and so } \|e^{\phi_k}\|_{L^q(\Omega_{k,\delta})} \leq c_\varepsilon \quad (3.33)$$

for $1 \leq q \leq \frac{1}{\varepsilon}$. Since $0 \leq W_k(x) \leq C|x|^{2\bar{\alpha}}$ in $\Omega_{k,\delta}$ with $\bar{\alpha} \geq 2$, we find that, $0 \leq W_k e^{\xi_k} \leq C|x|^{2(\bar{\alpha}-2)-\varepsilon} e^{\phi_k}$ and, by virtue of (3.33), we conclude that,

$$\|\tilde{f}_k\|_{L^p(B_\delta)} = \|W_k e^{\xi_k}\|_{L^p(\Omega_{k,\delta})} \leq C_\varepsilon = C_\varepsilon(p), \quad \text{for } 1 \leq p < \frac{2}{3\varepsilon}.$$

Therefore, by (3.32), we can use elliptic estimates to conclude that $\tilde{\phi}_k$ is uniformly bounded in B_δ . In turn, from (3.33), we deduce:

$$\phi_k \leq C \quad \text{in } \Omega_{k,\delta}$$

and (3.28) is established. \square

The estimate (3.28) implies in particular that, for any $\varepsilon > 0$ sufficiently small, we have:

$$0 \leq W_k(x)e^{\xi_k(x)} \leq C|x|^{2(\bar{\alpha}-2)-\varepsilon}, \text{ for } L_0\varepsilon_k \leq |x| \leq r \text{ and } k \geq k_\varepsilon$$

with $\bar{\alpha} \geq 2$. As a consequence, we have:

$$\int_{L_0\varepsilon_k \leq |x| \leq r} |\ln|x||W_k(x)e^{\xi_k(x)}dx \leq C, \quad (3.34)$$

and we shall take advantage of (3.34) to refine the estimate (3.28) as follows.

Proposition 3.2. *We have:*

$$\xi_k(x) = \min_{\partial B_r} \xi_k + 4 \ln \frac{1}{|x|} + O(1), \text{ for } L_0\varepsilon_k \leq |x| \leq r. \quad (3.35)$$

Proof. As before, we first establish (3.35) for $|x| = L_0\varepsilon_k$. To this purpose set,

$$\mu_k = \frac{1}{2\pi} \int_{|y| \leq 2L_0\varepsilon_k} W_k(y)e^{\xi_k(y)}dy = \frac{1}{2\pi} \int_{|z| \leq 2L_0} R_k(z)e^{v_k(z)}dz \longrightarrow 4, \quad (3.36)$$

as $k \longrightarrow +\infty$. Well known estimates established in [7] (see also [26]) allow us to conclude that,

$$|\mu_k - 4|v_k(0) = O(1). \quad (3.37)$$

Let $x = \varepsilon_k x'$ with $|x'| = L_0$ and write

$$\begin{aligned} \xi_k(x) &= \min_{\partial B_r} \xi_k + \mu_k \ln \frac{1}{|x|} + \frac{1}{2\pi} \int_{\{|y| \leq 2L_0\varepsilon_k\}} \ln\left(\frac{|x|}{|x-y|}\right) W_k(y)e^{\xi_k(y)} dy \\ &\quad + \frac{1}{2\pi} \int_{\{2L_0\varepsilon_k \leq |y| < r\}} \ln\left(\frac{1}{|x-y|}\right) W_k(y)e^{\xi_k(y)} dy + O(1) \\ &= \min_{\partial B_r} \xi_k + \mu_k \ln \frac{1}{\varepsilon_k} + \frac{1}{2\pi} \int_{\{|y| \leq 2L_0\}} \ln\left(\frac{|x'|}{|x'-y|}\right) R_k(y)e^{v_k(y)} dy \\ &\quad + \frac{1}{2\pi} \int_{\{2L_0\varepsilon_k \leq |y| < r\}} \ln\left(\frac{1}{|y|}\right) W_k(y)e^{\xi_k(y)} dy + O(1). \end{aligned} \quad (3.38)$$

By virtue of (3.20) and (3.21) we find: $\int_{\{|y| \leq L_0\}} \ln\left(\frac{|x'|}{|x'-y|}\right) R_k(y)e^{v_k(y)} dy \longrightarrow 0$, as $k \longrightarrow +\infty$, while (3.34) implies that the last integral in (3.38) is uniformly bounded. In conclusion, we have obtained:

$$\xi_k(x) = \min_{\partial B_r} \xi_k + \mu_k \ln \frac{1}{\varepsilon_k} + O(1), \text{ for } |x| = \varepsilon_k L_0. \quad (3.39)$$

As a consequence, for $|y| = L_0$, we have:

$$\begin{aligned} v_k(y) &= \xi_k(\varepsilon_k y) + 2 \ln \varepsilon_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| \\ &= \min_{\partial B_r} \xi_k + (2\bar{\alpha} - \mu_k) \ln \varepsilon_k + \sum_{j=s_1+1}^s 2\alpha_j \ln |p_{j,k}| + O(1) \end{aligned}$$

At this point, we can use (3.22) to deduce the following crucial information:

$$v_k(0) = -\min_{\partial B_r} \xi_k + (2(\bar{\alpha} + 1) - \mu_k) \ln \frac{1}{\varepsilon_k} + \sum_{j=s_1+1}^s 2\alpha_j \ln \frac{1}{|p_{j,k}|} + O(1). \quad (3.40)$$

Since $2(\bar{\alpha} + 1) - \mu_k \rightarrow 2(\bar{\alpha} - 1) \geq 2$, as $k \rightarrow +\infty$, from (3.37) and (3.40) we obtain that,

$$0 < \ln \frac{1}{\varepsilon_k} \leq C v_k(0) \quad \text{and} \quad |\mu_k - 4| \ln \frac{1}{\varepsilon_k} \leq C. \quad (3.41)$$

for suitable $C > 0$. Consequently, by (3.39),

$$|\xi_k(x) - 4 \ln \frac{1}{\varepsilon_k} - \min_{\partial B_r} \xi_k| \leq C, \quad \text{for } |x| = L_0 \varepsilon_k$$

and (3.35) is established for $|x| = L_0 \varepsilon_k$.

Clearly, by (2.16), (3.35) also holds for $|x| = r$, and we can argue exactly as above for the function $\phi_k = \xi_k(x) - \min_{\partial B_r} \xi_k + 4 \ln |x|$, in order to show that $\|\phi_k\|_{L^\infty\{R_0 \varepsilon_k \leq |x| < r\}} \leq C$, and conclude that (3.35) holds. \square

The estimate (3.35) allows us to show the following additional estimates.

Proposition 3.3. *Under the above assumptions there holds*

$$v_k(0) = -\min_{\partial B_r} \xi_k + 2 \ln \varepsilon_k - \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| + O(1) \quad (3.42)$$

$$|v_k(y) + v_k(0) + 4 \ln |y|| \leq C \quad \text{for } L_0 \leq |y| \leq \frac{r}{\varepsilon_k} \quad (3.43)$$

$$\xi_k(0) = -\min_{\partial B_r} \xi_k - 2 \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| + O(1) \quad (3.44)$$

$$\begin{aligned} \int_{B_r} |\nabla \xi_k|^2 &= 16\pi(v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) + O(1) \\ &= -16\pi(\min_{\partial B_r} \xi_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}|) + O(1) \\ &= 16\pi(\xi_k(0) + 2 \sum_{j=1}^s \alpha_j \ln |p_{j,k}|) + O(1) \end{aligned} \quad (3.45)$$

Proof. From (3.35) we derive that,

$$\begin{aligned} v_k(y) &= \min_{\partial B_r} \xi_k - 2 \ln \varepsilon_k \\ &\quad + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| + 4 \ln \frac{1}{|y|} + O(1), \quad \text{for } L_0 \leq |y| \leq \frac{r}{\varepsilon_k}. \end{aligned} \quad (3.46)$$

Hence, by using (3.46) with $|y| = L_0\varepsilon_k$ together with (3.22), we find:

$$-v_k(0) = \min_{\partial B_r} \xi_k - 2 \ln \varepsilon_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| + O(1) \quad (3.47)$$

and (3.42) is established. At this point, by inserting (3.47) into (3.46), we readily obtain (3.43). Also we obtain (3.44) from (3.42), once we recall that, $v_k(0) = \xi_k(0) + 2 \ln \varepsilon_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}|$. Finally, to obtain (3.45), we multiply both sides of the equation (2.15) by: $\xi_k - \min_{\partial B_r} \xi_k \geq 0$, and then integrate over B_r . We find

$$\begin{aligned} \int_{B_r} |\nabla \xi_k|^2 - \int_{\partial B_r} \frac{\partial \xi_k}{\partial \nu} (\xi_k - \min_{\partial B_r} \xi_k) &= \int_{B_r} W_k e^{\xi_k} (\xi_k - \min_{\partial B_r} \xi_k) \\ &= \int_{\{|x| \leq L_0\varepsilon_k\}} W_k e^{\xi_k} (\xi_k - \min_{\partial B_r} \xi_k) + \int_{\{L_0\varepsilon_k \leq |x| \leq r\}} W_k e^{\xi_k} (\xi_k - \min_{\partial B_r} \xi_k). \end{aligned}$$

Therefore, by means of (3.35), we obtain:

$$\int_{\{L_0\varepsilon_k \leq |x| \leq r\}} W_k e^{\xi_k} (\xi_k - \min_{\partial B_r} \xi_k) \leq C \int_{B_r} |x|^{2(\bar{\alpha}-2)} \ln \frac{1}{|x|} dx \leq C$$

(recall that $\bar{\alpha} \geq 2$). Since ξ_k admits the origin as its only blow-up point on B_r , then it is uniformly bounded in C^1 -norm on ∂B_r , and in view (2.16) we deduce: $|\int_{\partial B_r} \frac{\partial \xi_k}{\partial \nu} (\xi_k - \min_{\partial B_r} \xi_k)| \leq C$. Finally, we compute:

$$\begin{aligned} &\int_{\{|x| \leq L_0\varepsilon_k\}} W_k(x) e^{\xi_k(x)} (\xi_k(x) - \min_{\partial B_r} \xi_k) dx \\ &= \int_{\{|x| \leq L_0\varepsilon_k\}} W_k(x) (\xi_k(x) - \xi_k(0)) dx \\ &\quad + (\xi_k(0) - \min_{\partial B_r} \xi_k) \int_{\{|x| \leq L_0\varepsilon_k\}} W_k(x) e^{\xi_k(x)} dx \\ &= \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} (v_k(y) - v_k(0)) dy \\ &\quad + (\xi_k(0) - \min_{\partial B_r} \xi_k) \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy \\ &= \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} (v_k(y) - v_k(0)) dy + \\ &\quad - 2(\min_{\partial B_r} \xi_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}|) \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy + O(1). \end{aligned} \quad (3.48)$$

Since by (3.36) and (3.41) we have:

$$(v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy - 8\pi \leq C,$$

then, by (3.42), we can estimate the last term in (3.48) as follows:

$$\begin{aligned}
& -2(\min_{\partial B_r} \xi_k + \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}|) \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy \\
& = 2(v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) \int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy \\
& = (v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) 16\pi \\
& \quad + 2(v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) (\int_{\{|y| \leq L_0\}} R_k(y) e^{v_k(y)} dy - 8\pi) \\
& = (v_k(0) + 2 \ln \frac{1}{\varepsilon_k}) 16\pi + O(1).
\end{aligned}$$

Finally, we use (3.21) to estimate

$$\begin{aligned}
& \left| \int_{\{|x| \leq L_0\}} R_k(x) e^{v_k(x)} (v_k(x) - v_k(0)) dx \right| \\
& \leq C \int_{\{|x| \leq L_0\}} \frac{e^{v_k(0)}}{(1 + \frac{e^{v_k(0)}}{8} R_k(0) |x|^2)^2} \ln(1 + \frac{e^{v_k(0)}}{8} R_k(0) |x|^2) dx \\
& \leq C \int_{\mathbb{R}^2} \frac{dy}{(1 + |y|^2)^2} \ln(1 + |y|^2) \leq C
\end{aligned}$$

and so (3.45) follows. \square

Since $R_k(0) = h_k(0)$, we can combine (3.21) and (3.45) to find

$$\left| v_k(y) - \ln \frac{e^{v_k(0)}}{(1 + \frac{e^{v_k(0)}}{8} h_k(0) |y|^2)^2} \right| \leq C, \quad \text{for } |y| \leq \frac{r}{\varepsilon_k}. \quad (3.49)$$

and obtain in particular that: $\int_{\{|y| \leq \frac{r}{\varepsilon_k}\}} e^{v_k(y)} dy \leq C$.

In view of Theorem 4, we know that $p_{j,k} \neq 0 \forall k \in \mathbb{N}$, and so $W_k(0) = \prod_{j=1}^s |p_{j,k}|^{2\alpha_j} h_k(0) > 0$. Hence we can formulate (3.49) in terms of ξ_k as follows:

Corollary 3.1. *Under the above assumptions we have, $0 < W_k(0) \rightarrow 0$, as $k \rightarrow +\infty$, and*

$$\xi_k(x) = \ln \left(\frac{e^{\xi_k(0)}}{(1 + \frac{e^{\xi_k(0)}}{8} W_k(0) |x|^2)^2} \right) + O(1) \quad \text{in } B_r. \quad (3.50)$$

It is interesting to compare (3.50) with the analogous one "bubble" estimate established in [7] and [17] (see e.g. Theorem 0.3 in [17]) concerning the profile

of blow-up solutions of (2.15)-(2.17), when $W_k \xrightarrow{k \rightarrow +\infty} W$ uniformly in B_r and $W(0) > 0$, in which case (3.1) is automatically satisfied (see (i) of Theorem A) and no collapsing issues arise. Indeed, our pointwise estimate in (3.50) is the striking exact analogue of the one provided in Theorem 0.3 of [17], carried over to the case where W satisfies (2.22) and in particular $W(0) = 0$.

Furthermore, for the sequence:

$$u_k(x) = \xi_k(x) + \sum_{j=1}^s 2\alpha_j \log |x - p_{j,k}|,$$

satisfying:

$$-\Delta u_k = h_k e^{u_k} - 4\pi \sum_{j=1}^s \alpha_j \delta_{p_{j,k}} \quad \text{in } B_r, \quad (3.51)$$

we realize that, the estimate (3.44) stated for ξ_k in Proposition 3.3 reduces, in terms of u_k , to the following:

$$u_k(0) + \min_{\partial B_r} u_k = -\ln W_k(0) \longrightarrow +\infty, \quad \text{as } k \longrightarrow +\infty,$$

as stated in (1.14), and implying (by (1.11)) that blow-up for solutions of the singular problem (3.51) is equivalent to blow-up for its "regular" part.

We shall use the estimates established here to describe the asymptotic behaviour of minimizers of the Donaldson functional considered in [10],[13], and to obtain in particular that for Riemann surfaces of genus 2, it is always bounded from below, although not coercive. Furthermore, we shall provide rather precise information on when the infimum is attained. In this way we obtain the first existence result about (CMC) 1-immersions of a closed orientable surface of genus 2 into hyperbolic 3-manifolds.

For this purpose, we conveniently summarize the results established above for a sequence ξ_k satisfying:

$$\left\{ \begin{array}{l} -\Delta \xi_k(x) = (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k(x) + g_k(x) \quad \text{in } B_r \\ s \geq 2 \quad \text{and } \alpha_j \in \mathbb{N} \quad \text{for } j = 1, \dots, s \\ \xi_k(0) = \max_{B_r} \xi_k \longrightarrow +\infty, \quad \text{as } k \longrightarrow +\infty, \\ \forall 0 < \delta < r \quad \exists C_\delta > 0 : \max_{\overline{B_r} \setminus B_\delta} \xi_k \leq C_\delta \\ \max_{\partial B_r} \xi_k - \min_{\partial B_r} \xi_k \leq C \\ \int_{B_r} W_k e^{\xi_k} \leq C, \quad W_k(x) := (\prod_{j=1}^s |x - p_{j,k}|^{2\alpha_j}) h_k(x) \end{array} \right. \quad (3.52)$$

Theorem 5. *Suppose ξ_k satisfies (3.52) with the points $p_{j,k}$ satisfying (2.18) and (3.3), h_k satisfying (2.12) and (2.20), and g_k a convergent sequence in $L^p(B_r)$, $p > 1$.*

If (3.1) holds, then the points $p_{j,k} \neq 0$, $j = 1, \dots, s$; and there exists $s_1 \in \{2, \dots, s\}$ such that (along a subsequence)

$$z_{j,k} := \frac{p_{j,k}}{|p_{s_1,k}|} \longrightarrow z_j \neq 0, \quad \forall j = 1, \dots, s_1,$$

if $s_1 < s$ then $\frac{p_{j,k}}{|p_{s,k}|} \rightarrow q_j \neq 0$ and $\frac{|p_{j,k}|}{|p_{s_1,k}|} \rightarrow +\infty, \forall j = s_1 + 1, \dots, s.$

Moreover,

$$\begin{aligned} \xi_k(0) + 2 \ln |p_{s_1,k}| + \ln(W_k(0)) &\rightarrow +\infty, \text{ as } k \rightarrow +\infty, \\ \xi_k(0) + \left(\min_{\partial B_r} \xi_k + 2 \ln(W_k(0)) \right) &= O(1) \\ \xi_k(x) = \ln \left(\frac{e^{\xi_k(0)}}{\left(1 + \frac{W_k(0)}{8} e^{\xi_k(0)} |x|^2\right)^2} \right) &+ O(1), \text{ } x \in B_r \\ \int_{B_r} |\nabla \xi_k|^2 dx &= 16\pi (\xi_k(0) + \ln(W_k(0))) + O(1). \end{aligned}$$

Proof. It suffices to observe that the results established above apply to the sequence: $\hat{\xi}_k := \xi_k + \chi_k$, where χ_k is the unique solution of the Dirichlet problem:

$$\begin{cases} \Delta \chi_k = g_k & \text{in } B_r \\ \chi_k = 0 & \text{in } \partial B_r. \end{cases}$$

Indeed, by elliptic estimates, χ_k converges strongly in $C^{1,\alpha}(B_r)$ and moreover, $\ln W_k(0) = \sum_{j=1}^s 2\alpha_j \ln |p_{j,k}| + O(1)$. Then it is easy to check that in terms of the original sequence ξ_k , we get exactly the claimed estimates. \square

4 Appendix: The Proof of (2.49).

By virtue of the properties pointed out in Section 2 for the sequence φ_k defined in (2.29), to establish (2.49) we can follow word by word the arguments used in [16] in order to show the same identity (i.e.(4.9) in [16]), see also [14]. To this purpose we start with the following:

Lemma 4.1. *Let μ be defined in (2.31). Then there exists $R_0 > 0$ sufficiently large, such that, as $k \rightarrow +\infty$,*

$$\nabla \varphi_k(x) \rightarrow -\mu \frac{x}{|x|^2} + O\left(\frac{1}{|x|^2}\right) \quad (4.1)$$

uniformly on compact sets of $\mathbb{R}^2 \setminus B_{R_0}$.

Proof. From (2.30) we have

$$\begin{aligned} \nabla \varphi_k(x) &= -\frac{\tau_k}{2\pi} \int_{B_r} \left(\frac{\tau_k x - y}{|\tau_k x - y|^2} \right) W_k(y) e^{\xi_k(y)} dy \\ &= -\frac{\tau_k}{2\pi} \int_{\{|y| \leq \tau_k |x|^2\}} \left(\frac{\tau_k x - y}{|\tau_k x - y|^2} \right) W_k(y) e^{\xi_k(y)} dy \\ &\quad - \frac{\tau_k}{2\pi} \int_{\{\tau_k |x|^2 \leq |y| \leq r\}} \left(\frac{\tau_k x - y}{|\tau_k x - y|^2} \right) W_k(y) e^{\xi_k(y)} dy \\ &=: I_{1,k}(x) + I_{2,k}(x). \end{aligned}$$

Now, for $R_0 > 1$ (to be fixed later) let $|x| \geq R_0$ and $\tau_k|x|^2 \leq |y| \leq r$, then we have:

$$(1 - \frac{1}{R_0})|y| \leq |\tau_k x - y| \leq (1 + \frac{1}{R_0})|y|$$

and therefore

$$|I_{2,k}| \leq C \int_{\{\tau_k|x|^2 \leq |y| \leq r\}} \frac{\tau_k}{|y|} e^{\xi_k(y)} W_k(y) dy \leq \frac{C}{|x|^2}, \quad |x| \geq R_0 > 1.$$

By recalling (2.27) and (2.29), we find:

$$I_{1,k}(x) = -\frac{1}{2\pi} \int_{\{|z| \leq |x|^2\}} \frac{(x-z)}{|x-z|^2} W_{1,k}(z) e^{\varphi_k(z)} dz.$$

Therefore, in case φ_k satisfies alternative (2.32), then (4.1) readily follows. Indeed, $I_{1,k} \rightarrow 0$, as $k \rightarrow +\infty$, uniformly on compact sets of $\mathbb{R}^2 \setminus B_{R_0}$ and $\mu = 0$ in this case. While, in case φ_k satisfies alternative (2.35), then for $R_0 > 1$ sufficiently large so that, $S_\varphi \subset\subset B_{R_0}$, we have, as $k \rightarrow +\infty$:

$$I_{1,k}(x) \rightarrow -\frac{1}{2\pi} \sum_{q \in S_\varphi} \sigma(q) \frac{(x-q)}{|x-q|^2} \quad \text{uniformly on compact sets of } \mathbb{R}^2 \setminus B_{R_0},$$

and, in this case (see (2.36)), we have :

$$\begin{aligned} -\frac{1}{2\pi} \sum_{q \in S_\varphi} \sigma(q) \frac{(x-q)}{|x-q|^2} &= -\mu \frac{x}{|x|^2} + \frac{1}{2\pi} \sum_{q \in S_\varphi} \left(\frac{x}{|x|^2} - \frac{(x-q)}{|x-q|^2} \right) \\ &= -\mu \frac{x}{|x|^2} + O\left(\frac{1}{|x|^2}\right), \quad \text{for } |x| \geq R_0. \end{aligned}$$

So also in this case (4.1) is established. Finally, in case φ_k satisfies alternative (2.33) or (2.38), then by virtue of (2.43) and (2.44) we find that,

$$W_1(x) e^{\varphi_0} \leq \frac{C}{|x|^{2(\rho+1)}} \quad \text{for } |x| \geq R_0 \quad (4.2)$$

with $R_0 \gg 1$ sufficiently large and suitable constants $C > 0$ and $\rho \geq 1$. In this case, as $k \rightarrow +\infty$, we find:

$$I_{1,k}(x) \rightarrow -\mu \frac{x}{|x|^2} + R_1(x), \quad \text{uniformly on compact sets of } \mathbb{R}^2 \setminus B_{R_0}$$

with

$$\begin{aligned} R_1(x) &= \frac{1}{2\pi} \int_{\{|y| \geq |x|^2\}} W_1(x) e^{\varphi_0} \\ &\quad + \frac{1}{2\pi} \int_{\{|y| < |x|^2\}} \left(\frac{x}{|x|^2} - \frac{x-y}{|x-y|^2} \right) W_1(y) e^{\varphi_0(y)} dy + R_2(x), \end{aligned}$$

where $R_2(x) = 0$ in case alternative (2.33) holds (i.e. $S_\varphi = \emptyset$) or

$$R_2(x) = \frac{1}{2\pi} \sum_{q \in S_\varphi} \sigma(q) \left(\frac{x}{|x|^2} - \frac{x-q}{|x-q|^2} \right) = O\left(\frac{1}{|x|^2}\right)$$

in case alternative (2.38) holds (i.e. $S_\varphi \neq \emptyset$). Clearly, in view of (4.2), we easily check that,

$$\int_{\{|y| \geq |x|^2\}} W_1(y) e^{\varphi_0(y)} \leq \frac{C}{|x|^2}, \quad \forall |x| \geq R_0.$$

By recalling the following easy facts:

$$\begin{aligned} \text{if } |x-y| \geq \frac{|x|}{2} \text{ then } \left| \frac{x}{|x|^2} - \frac{x-y}{|x-y|^2} \right| &\leq 4 \frac{|y|}{|x|^2} \\ \text{if } |x-y| \leq \frac{|x|}{2} \text{ then } \left| \frac{x}{|x|^2} - \frac{x-y}{|x-y|^2} \right| &\leq \frac{4}{|x-y|}, \end{aligned}$$

and by using (4.2), for $|x| \geq R_0$ we derive:

$$\begin{aligned} & \left| \int_{\{|y| \leq |x|^2\}} \left(\frac{x}{|x|^2} - \frac{x-y}{|x-y|^2} \right) W_{1,k}(y) e^{\varphi_0} dy \right| \\ & \leq \frac{C}{|x|^2} \int_{\mathbb{R}^2} |y| W_1(y) e^{\varphi_0} dy + C \int_{\{|x-y| \leq \frac{|x|}{2}\}} \frac{1}{|x-y|} W_1(y) e^{\varphi_0} dy \\ & \leq \frac{C}{|x|^2} + C \int_{\{|x-y| \leq \frac{|x|}{2}\}} \frac{1}{|x-y|} \frac{1}{|y|^{2(\rho+1)}} dy \\ & = \frac{C}{|x|^2} + \frac{C}{|x|^{2\rho+1}} \int_{\{|\frac{x}{|x|} - z| \leq \frac{1}{2}\}} \frac{1}{|\frac{x}{|x|} - z|} \frac{dz}{|z|^{2(\rho+1)}} \leq \frac{C}{|x|^2} \quad \text{for } |x| \geq R_0 \geq 1. \end{aligned}$$

By combining the estimates above, we obtain (4.1) also when φ_k satisfies either alternative (2.33) or alternative (2.38), and the proof is completed. \square

By using (2.25) and (4.1), we can easily show that, as $k \rightarrow +\infty$:

$$r \int_{\partial B_r} (|\partial_\nu \xi_k| - \frac{1}{2} |\nabla \xi_k|^2) d\sigma \rightarrow 2\pi \frac{m^2}{2} + o_r(1)$$

with m in (2.23) and $o_r(1) \rightarrow 0$, as $r \rightarrow 0$, and

$$R \int_{\partial B_R} (|\partial_\nu \xi_k| - \frac{1}{2} |\nabla \xi_k|^2) d\sigma \rightarrow 2\pi \frac{\mu^2}{2} + o_R(1)$$

with $o_R(1) \rightarrow 0$, as $R \rightarrow +\infty$.

Lemma 4.2. *There holds:*

$$m^2 - \mu^2 = 4(\alpha + 1)(m - \mu).$$

Proof. As in [16],[14], we use the Pohozaev identity (see e.g. (5.2.14) in [26]) for ξ_k (satisfying (2.15)) in $B_r \setminus B_{R\tau_k}$ and obtain:

$$\begin{aligned}
& r \int_{\partial B_r} (|\partial_\nu \xi_k|^2 - \frac{1}{2} |\nabla \xi_k|^2 + W_k e^{\xi_k}) d\sigma \\
& \quad - R \int_{\partial B_R} (|\partial_\nu \varphi_k|^2 - \frac{1}{2} |\nabla \varphi_k|^2 + W_{1,k} e^{\varphi_k}) d\sigma \\
& = \int_{B_{\frac{r}{\tau_k}} \setminus B_R} 2(1 + \sum_{j=1}^s \alpha_j \frac{y(y - q_{j,k})}{|y - q_{j,k}|^2}) W_{1,k}(y) e^{\varphi_k} dy \\
& \quad + \int_{B_{\frac{r}{\tau_k}} \setminus B_R} (\tau_k y \cdot \frac{\nabla h_k(\tau_k y)}{h_k(\tau_k y)}) W_{1,k}(y) e^{\varphi_k(y)} dy.
\end{aligned} \tag{4.3}$$

By using (2.21) and (2.25), and the convergence (4.1) for φ_k , we easily check that, for the left hand side $(LHS)_k$ of (4.3), we find

$$(LHS)_k \longrightarrow 2\pi(\frac{m^2}{2} - \frac{\mu^2}{2}) + o_r(1) + o_R(1), \text{ as } k \longrightarrow +\infty.$$

While for the right hand side $(RHS)_k$ of (4.3), we have:

$$\begin{aligned}
(RHS)_k & = 2(1 + \alpha) \int_{B_{\frac{r}{\tau_k}} \setminus B_R} W_{1,k}(y) e^{\varphi_k} \\
& \quad + \sum_{j=1}^s 2\alpha_j \int_{B_{\frac{r}{\tau_k}} \setminus B_R} q_{k,j} \cdot \frac{y - q_{k,j}}{|y - q_{k,j}|^2} W_{1,k} e^{\varphi_k} + o_r(1)
\end{aligned}$$

and we conclude that, as $k \longrightarrow +\infty$,

$$(RHS)_k \longrightarrow 4\pi(1 + \alpha)(m - \mu) + o_r(1) + o_R(1).$$

Then we obtain the desired conclusion, by letting $r \longrightarrow 0^+$ and $R \longrightarrow +\infty$. \square

Acknowledgments

We thank Dr. Martin Mayer for his precious assistance and useful comments during the preparation of the manuscript.

This work was partially supported by: MIUR excellence project: Department of Mathematics, University of Rome "Tor Vergata" CUP E83C18000100006, by "Beyond Borders" research project: "Variational approaches to PDE" and Fondi di Ricerca Scientifica d'Ateneo, 2021: Research Project "Geometric Analysis in Nonlinear PDEs", CUP E83C22001810005.

References

- [1] Aubin, T., *Some Nonlinear Problems in Riemannian Geometry*. Springer-Verlag Berlin Heidelberg, 1998

- [2] Bartolucci D., Chen C.C., Lin C.S., Tarantello G., Profile of blow-up solutions to mean field equations with singular data. *Comm. Partial Differential Equations* **29** (2004), no. 7-8, 1241–1265
- [3] Bartolucci D., Tarantello G., Liouville type equations with singular data and their applications to periodic multivortices for the electroweak theory. *Comm. Math. Phys.* **229** (2002), no. 1, 3–47
- [4] Bartolucci D., Tarantello G., Asymptotic blow-up analysis for singular Liouville type equations with applications. *J. Diff. Eq.* **262** (2017), no. 7, 3887–3931
- [5] Brezis H., Merle F., Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions. *Comm. Partial Differential Equations* **16** (1991), no. 8-9, 1223–1253.
- [6] Bryant R., Surfaces of mean curvature one in hyperbolic space, Théorie des variétés minimales et applications *Astérisque* No. 154-155 (1987), 12, 321–347, 353 (1988)
- [7] Chen C.C., Lin C.S., Sharp estimates for solutions of multi-bubbles in compact Riemann surfaces. *Comm. Pure Appl. Math.* **55** (2002), no. 6, 728–771.
- [8] Chen C.C., Lin C.S., Topological degree for a mean field equation on Riemann surfaces. *Comm. Pure Appl. Math.* **56** (2003), no. 12, 1667–1727
- [9] Chen C.C., Lin C.S., Mean field equation of Liouville type with singular data: topological degree. *Comm. Pure Appl. Math.* **68**, (2015), no. 6, 887–947.
- [10] Goncalves K., Uhlenbeck K., Moduli space theory for constant mean curvature surfaces immersed in space-forms. *Comm. Anal. Geom.* **15** (2007), 299–305.
- [11] Griffiths P., Harris J., *Principles of Algebraic Geometry*, Wiley Classics Library (2014), John Wiley & Sons
- [12] Huang Z., Lucia M., Tarantello G., Bifurcation for minimal surface equation in hyperbolic 3-manifolds. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **38** (2021), no. 2, 243–279
- [13] Huang Z., Lucia M., Tarantello G., Donaldson Functional in Teichmüller Theory, *Int. Math. Res. Notes*, to appear.
- [14] Lee Y., Lin C.S., Tarantello G., Yang W., Sharp estimates for solutions of mean field equations with collapsing singularity. *Comm. Partial Differential Equations* **42** (2017), no. 10, 1549–1597

- [15] Lee Y., Lin C.S., Wei J., Yang W., Degree counting and shadow system for Toda system of rank two: one bubbling. *J. Diff. Eq.* **264** (2018), no. 7, 4343-4401.
- [16] Lee Y., Lin C.S., Yang W., Zhang L., Degree counting for Toda system with simple singularity: one point blow up. *J. Diff. Eq.* **268** (2020), no. 5, 2163-2209
- [17] Li Y.Y., Harnack type inequality: the method of moving planes. *Comm. Math. Phys.* **200** (1999), no. 2, 421-444
- [18] Li Y.Y., Shafrir I., Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two. *Indiana Univ. Math. J.* **43** (1994), no. 4, 1255-1270
- [19] Lin C.S, Tarantello G., When "blow-up" does not imply "concentration": a detour from Brezis-Merle's result. *C.R. Math. Acad. Sci. Paris* **354** (2016), no. 5, 493-498
- [20] Lin C.S., Wei J.C., Yang W., Zhang L., On rank-2 Toda systems with arbitrary singularities: local mass and new estimates. *Anal. PDE* **11** (2018), no. 4, 873-898
- [21] Lin C.S., Wei J.C., Zhang L., Classification of blowup limits for SU(3) singular Toda systems. *Anal. PDE* **8** (2015), no. 4, 807-837.
- [22] Lin C.S., Wei J., Zhao C., Asymptotic behavior of SU(3) Toda system in a bounded domain. *Manuscripta Math.* **137** (2012), no. 1-2, 1-18.
- [23] Lin C.S., Wei J., Zhao C., Sharp estimates for fully bubbling solutions of a SU(3) Toda system. *Geom. Funct. Anal.* **22** (2012), no. 6, 1591-1635.
- [24] Ohtsuka H., Suzuki T., Blow-up analysis for Liouville type equation in self-dual gauge field theories. *Commun. Contemp. Math.* **7** (2005), no. 2, 177-205.
- [25] Tarantello G., Asymptotics for minimizers of a Donaldson functional and mean curvature 1-immersions of surfaces into hyperbolic 3-manifolds. *Preprint 2022*
- [26] Tarantello G., *Selfdual Gauge Field Vortices An Analytical Approach*. Progress in Nonlinear Diff. Eq. and Their Applications **72**, Birkhäuser, Basel, 2008.
- [27] Wei J., Zhang L., Estimates for Liouville equation with quantized singularities. *Advances in Mathematics* **380** (2021), 1-45.
- [28] Wei J., Zhang L., Laplacian vanishing theorem for quantized singular Liouville equation, preprint 2022.