

DEFINABILITY AND DECIDABILITY FOR RINGS OF INTEGERS IN TOTALLY IMAGINARY FIELDS

CALEB SPRINGER

ABSTRACT. We show that the ring of integers of \mathbb{Q}^{tr} is existentially definable in the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$, where \mathbb{Q}^{tr} denotes the field of all totally real numbers. This implies that the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$ is undecidable and first-order non-definable in $\mathbb{Q}^{\text{tr}}(i)$. More generally, when L is a totally imaginary quadratic extension of a totally real field K , we use the unit groups R^\times of orders $R \subseteq \mathcal{O}_L$ to produce existentially definable totally real subsets $X \subseteq \mathcal{O}_L$. Under certain conditions on K , including the so-called JR-number of \mathcal{O}_K being the minimal value $\text{JR}(\mathcal{O}_K) = 4$, we deduce the undecidability of \mathcal{O}_L . This extends previous work which proved an analogous result in the opposite case $\text{JR}(\mathcal{O}_K) = \infty$. In particular, unlike prior work, we do not require that L contains only finitely many roots of unity.

1. INTRODUCTION

1.1. A motivating example. This paper is motivated by the desire to prove the following theorem concerning definability and decidability. For background and an overview of decidability and definability for infinite algebraic extensions of \mathbb{Q} , we refer readers to the introduction of [Shl18]; see also the prequel to our work here [Spr20]. Recall that $\alpha \in \overline{\mathbb{Q}}$ is *totally real* if the roots of its minimal polynomial are all real numbers. We write \mathbb{Q}^{tr} for the field of all totally real algebraic numbers and \mathbb{Z}^{tr} for its ring of integers, and we set $i = \sqrt{-1}$.

Theorem 1.1. *The ring of integers \mathbb{Z}^{tr} of \mathbb{Q}^{tr} is existentially definable in the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$. In particular, the first-order theory of the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$ is undecidable.*

Remark 1.2. The paragraph below Corollary 3.19 in [Koe14] mistakenly declared the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$ to be decidable.¹ The reference given for this claim is [Dar00, Theorem 10.7], which actually proves the decidability of the overring $\mathcal{O}_{\mathbb{Q}^{\text{tr}}(i)}[\frac{1}{p}]$ where p is any prime number. As seen above, the ring of integers $\mathcal{O}_{\mathbb{Q}^{\text{tr}}(i)}$ itself is undecidable.

Theorem 1.1 is proven by extending and generalizing the methods which were first used to prove the first-order undecidability of rings of integers in fields such as $\mathbb{Q}^{(2)}$ [MRUV20] and $\mathbb{Q}_{\text{ab}}^{(d)}$ for $d \geq 2$ [Spr20]. Here, $F^{(d)}$ denotes the compositum of all extensions of degree at most d of a field F , and $F_{\text{ab}}^{(d)} \subseteq F^{(d)}$ is the maximal abelian subfield. However, notice that the papers [MRUV20, Spr20] additionally proved that the fields $\mathbb{Q}_{\text{ab}}^{(d)}$ themselves are undecidable by appealing to results of Videla [Vid00a] and Shlapentokh [Shl18] which imply that each $\mathbb{Q}_{\text{ab}}^{(d)}$ has a first-order definable ring of integers.

The situation is entirely the opposite in the setting of Theorem 1.1. Indeed, Fried, Haran, and Völklein proved that \mathbb{Q}^{tr} is decidable [FHV94], which implies that $\mathbb{Q}^{\text{tr}}(i)$ is also decidable. It has been previously observed that, because J. Robinson proved that \mathbb{Z}^{tr} is undecidable

Date: September 30, 2022.

¹We thank Aharon Razon for pointing this out after reading an earlier draft of this paper.

[Rob62], the field \mathbb{Q}^{tr} is thus an example of a subfield of $\overline{\mathbb{Q}}$ whose ring of integers \mathbb{Z}^{tr} does not admit any first-order definition in \mathbb{Q}^{tr} . In the framework of Shlapentokh [Shl18, §2.1], this is intuitively understood as a result of the fact that the field \mathbb{Q}^{tr} is too “close” to $\overline{\mathbb{Q}}$, so it does not have enough “expressive power” to have a definable ring of integers. By Theorem 1.1, we immediately see that $\mathbb{Q}^{\text{tr}}(i)$ also has a first-order non-definable ring of integers, as expected.

We pause to note that $\mathbb{Q}^{\text{tr}}(i)$ is known to be an ω -free PAC field; see [Jar11, Example 5.10.7]. Therefore, the non-definability of the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$ can also be deduced from the following result of Dittman and Fehm: If R is a first-order definable subring of an ω -free PAC field L , then R is a field [DF21, Proposition 3]. Their method, which builds upon work of Chatzidakis [Cha19], is entirely different from the techniques that we use below.

1.2. Main results. In general, this paper considers totally imaginary quadratic extensions of totally real fields. We begin with a definability result. Say that a totally real field $K \subseteq \overline{\mathbb{Q}}$ is *closed under square roots* if $\alpha \in K$ whenever α is a totally real number such that $\alpha^2 \in K$.

Theorem 1.3 (Theorem 3.4). *If K is a totally real field which is closed under square roots and L is any quadratic imaginary extension of K , then \mathcal{O}_K is existentially definable in \mathcal{O}_L .*

This theorem implies Theorem 1.1 immediately, given J. Robinson’s proof that \mathbb{Z}^{tr} is undecidable [Rob62]. However, we can actually prove first-order undecidability for rings of integers \mathcal{O}_L in a more general context where \mathcal{O}_K is not known to be existentially definable in \mathcal{O}_L . For any $n \geq 1$, let ζ_n denote a primitive n -th root of unity.

Theorem 1.4 (Theorem 4.5). *Let $S \subseteq \mathbb{N}$ be infinite. Define $K_0 = \mathbb{Q}(\{\zeta_n + \bar{\zeta}_n : n \in S\})$ and let K_1 be the maximal totally real subfield of $K_0^{(2)}$. If $K \supseteq K_1$ is a totally real field and L is any totally imaginary quadratic extension of K , then \mathcal{O}_L is undecidable.*

In this theorem, the so-called JR-number is $\text{JR}(\mathcal{O}_K) = 4$, i.e., the minimal possible value, while the totally real fields K considered in [MRUV20, Spr20] have the maximal possible value $\text{JR}(\mathcal{O}_K) = \infty$; see Section 4.1 for definitions and notation. We also provide a version of our undecidability result, namely Theorem 4.4, which does not place a restriction on the value of $\text{JR}(\mathcal{O}_K)$.

1.3. Proof method: Leveraging unit groups. Within the body of literature concerning undecidability results for rings of integers of algebraic extensions of \mathbb{Q} , the totally real subfields of $\overline{\mathbb{Q}}$ and their totally imaginary quadratic extensions have received special attention. There are multiple methods which can be used in this context, but we focus on the work stemming from J. Robinson for now, and delay the discussion of elliptic curves until the following section.

J. Robinson [Rob62] provided a general sufficient condition for proving the undecidability of a ring of algebraic integers \mathcal{O} , and we follow Videla [Vid00b] by using an improved version credited to Henson [VDD88]: If there is a parametrized family \mathcal{F} of definable subsets of \mathcal{O} which contains finite sets of arbitrarily large cardinality, then the first-order theory of \mathcal{O} is undecidable. This sufficient condition, along with a theorem of Siegel [Sie21], leads to a strategy that can be applied to \mathcal{O}_K when K is a totally real field. Indeed, we use the parametrized family of subsets $\{X_t\}_{t \in \mathbb{Q}}$, where X_t contains the elements $\alpha \in \mathcal{O}_K$ whose conjugates all lie in the real interval $(0, t)$. Determining whether $\{X_t\}_{t \in \mathbb{Q}}$ contains finite sets of arbitrarily large size is related to the JR-number of \mathcal{O}_K ; see Section 4.1 for more details.

This strategy enabled J. Robinson to prove the undecidability of the rings of integers of both \mathbb{Q}^{tr} and $\mathbb{Q}_{\text{tr}}^{(2)} = \mathbb{Q}(\{\sqrt{n} : n \in \mathbb{N}\})$.

After using this method to prove \mathcal{O}_K is undecidable for a totally real field K , we wish to do the same for the rings of integers \mathcal{O}_L for all totally imaginary quadratic extensions L of K . To extend the JR-number argument to such a field, a key ingredient is the fact that $[\mathcal{O}_L^\times : \mu(L)\mathcal{O}_K^\times] \leq 2$ where $\mu(L)$ is the set of roots of unity in L . Therefore, if $N = \#\mu(L) < \infty$ is finite, then $(\mathcal{O}_L^\times)^{2N} \subseteq K$ is an existentially definable totally real subset of \mathcal{O}_L . By using sums and difference of powers of units, we can produce a useful parametrized family of subsets of \mathcal{O}_L analogous to the sets X_t defined above. Undecidability is thereby proved for \mathcal{O}_L in many cases, including $L = \mathbb{Q}^{(2)}$ or more generally $L = \mathbb{Q}_{\text{ab}}^{(d)}$ for $d \geq 2$; see [MRUV20, Spr20].

However, the assumption $\#\mu(L) < \infty$ above implies that this method cannot yet apply when $L = \mathbb{Q}^{\text{tr}}(i)$. To remedy this, we choose to instead work with the unit group R^\times of an existentially definable non-maximal order $R \subseteq \mathcal{O}_L$ with $\mu(R) = \{\pm 1\}$ trivial. In this case, we can show that $(R^\times)^2$ is totally real set. By choosing a suitable ring $R \subseteq \mathcal{O}_L$, we obtain the foundation for developing a unit group-based argument similar to the ones appearing in [MRUV20, Spr20] which allows L to contain infinitely many roots of unity.

1.4. A comparison with abelian varieties. We conclude this section by comparing our main theorems to similar results which leverage elliptic curves instead of unit groups. There are many papers which use of elliptic curves, or abelian varieties in general, to prove various undecidability results, including [Den80, Vid00b, Poo02, MRS22]. We refer to [Shl09] for additional background. As an example which is relevant to the consideration of totally imaginary extensions of totally real fields, consider the following theorem of Shlapentokh.

Theorem 1.5 (Main Theorem B, [Shl09]). *Let K be a totally real algebraic extension of \mathbb{Q} which has a totally real extension of degree 2, and let K' be a finite extension of K such that there exists an elliptic curve E defined over K' with $E(K')$ finitely generated and of positive rank. If L is a quadratic totally imaginary extension of K , then \mathbb{Z} is existentially definable in the ring of integers \mathcal{O}_K and Hilbert's Tenth Problem is unsolvable over \mathcal{O}_K .*

Although Theorems 1.4 and 1.5 are similar insofar as they both apply in the context of totally imaginary quadratic extensions of totally real subfields of $\overline{\mathbb{Q}}$, it is instructive to also note the differences and complementary strengths. Heuristically, it is easiest to apply the elliptic curve-based methods to relatively “small” algebraic extensions of \mathbb{Q} over which it is easy to find elliptic curves with a finitely generated group of rational points. In contrast, the unit group-based methods work best for “bigger” fields in which there is an abundance of units available for manipulation. We can make this more precise with a couple of examples.

The prototypical example of an infinite algebraic extension $K \supseteq \mathbb{Q}$ which satisfies the hypotheses of Theorem 1.5 is a \mathbb{Z}_p -extension of \mathbb{Q} ; see [Shl09, §10]. These \mathbb{Z}_p -extensions are difficult to handle with the unit group-based methods because there is at most one subextension of K of any given degree, hence a paucity of units. We also note that Theorem 1.5 proves that Hilbert's Tenth Problem is unsolvable, rather than only showing that the first-order theory is undecidable, and it also applies when $[K : \mathbb{Q}] < \infty$.

On the other hand, when deploying elliptic curves, we emphasize that it is necessary for the group of rational points to be a finitely generated group. This is a core requirement of the proof method, and the restriction would remain for any straightforward variant or

generalization which uses abelian varieties instead of elliptic curves, such as [MRS22, Theorem 1.1]. Therefore, the following theorem of Fehm and Petersen shows that this general method is not useable when the field K is *large*, in the sense of Pop [Pop96].

Theorem 1.6 (Theorem 1.2, [FP10]). *If $L \subseteq \overline{\mathbb{Q}}$ is a large field and A/L is an abelian variety, then $A(L)$ has infinite rank.*

The first proof of this theorem in the case of elliptic curves is credited to Tamagawa; see Kobayashi [Kob06, Proposition 1]. We refer to [FJ74, LR08, MR18, Pet06] for some additional results on finitely and non-finitely generated groups of rational points.

Because any algebraic extension of a large field is itself large, Theorem 1.6 shows that if $K \subseteq \overline{\mathbb{Q}}$ is a large totally real field, then abelian variety-based methods such as Theorem 1.5 cannot handle K or its quadratic totally imaginary extensions. However, it is clear that the unit group-based methods presented in this paper can work for large fields because the field \mathbb{Q}^{tr} is large. It is also conjectured that \mathbb{Q}^{ab} is a large field (see [BSF13, §3] for background and an overview), and it is easy to construct extensions $L \supseteq \mathbb{Q}^{\text{ab}}$ which are covered by Theorem 1.4. It would be interesting to determine whether or not the methods of this paper can be refined to prove the undecidability of the ring of integers of \mathbb{Q}^{ab} itself.

Acknowledgments. The author thanks Arno Fehm for assistance with references, and Aharon Razon for sharing Remark 1.2. Additional thanks to Kirsten Eisenträger, Jochen Koenigsmann, Alexandra Shlapentokh, and Carlos Videla for their helpful comments. This work was supported by the Additional Funding Programme for Mathematical Sciences, delivered by EPSRC (EP/V521917/1) and the Heilbronn Institute for Mathematical Research.

2. NON-MAXIMAL ORDERS AND UNITS

As indicated in Section 1.3, given a totally real field K and a totally imaginary quadratic extension L , we want a subring $R \subseteq \mathcal{O}_L$ which does not contain any nontrivial roots of unity. We start by defining the desired ring, then proceed to analyze its group of units.

2.1. A useful non-maximal subring.

Definition 2.1. *Given an integer $m \geq 1$ and a field $L \subseteq \overline{\mathbb{Q}}$, let $R_{m,L}$ be the subset of \mathcal{O}_L defined by the positive existential formula $\varphi_m(x)$, given as follows:*

$$\exists a \quad (x - ma)(x - 1 - ma) \dots (x - (m - 1) - ma) = 0$$

The following properties of $R_{m,L}$ are immediate from the definition.

Lemma 2.2. *Let $L \subseteq \overline{\mathbb{Q}}$ be a field and let $m \geq 1$ be an integer.*

- (a) *If $K \subseteq L$ is a subfield, then $R_{m,L} \cap K = R_{m,K}$.*
- (b) *$R_{m,L}$ is an existentially definable subring of \mathcal{O}_L which contains $m\mathcal{O}_L$.*
- (c) *$R_{m,L} \subseteq \mathcal{O}_L$ is non-maximal if and only if $m \geq 2$ and $L \neq \mathbb{Q}$.*

Proof. By the definition of $R_{m,L}$ and the formula $\varphi_m(x)$ above, the first claim is obvious. Moreover, $R_{m,L}$ is the preimage of $\mathbb{Z}/m\mathbb{Z}$ under the natural surjective map $\mathcal{O}_L \rightarrow \mathcal{O}_L/m\mathcal{O}_L$. Therefore, $R_{m,L}$ is a subring of \mathcal{O}_L and non-maximal precisely when $\mathbb{Z}/m\mathbb{Z} \neq \mathcal{O}_L/m\mathcal{O}_L$, i.e., when $m \geq 2$ and $L \neq \mathbb{Q}$. \square

Before we analyze the unit group $R_{m,L}^\times$, we recall some general elementary facts. In essence, this theorem clarifies that any ring containing an algebraic unit u also contains u^{-1} , and computes the rank of the unit groups of any ring of algebraic integers. Given a set $S \subseteq \overline{\mathbb{Q}}$, let $\mu(S)$ denote the set of roots of unity contained in S .

Theorem 2.3. *Let $L \subseteq \overline{\mathbb{Q}}$ be a number field with r real and $2s$ imaginary embeddings.*

- (a) *If $u \in \mathcal{O}_L^\times$, then $u^{-1} \in \mathbb{Z}[u]$.*
- (b) *If $R \subseteq \mathcal{O}_L$ is any subring, then $R^\times = \mathcal{O}_L^\times \cap R$.*
- (c) *If $\mathcal{O} \subseteq \mathcal{O}_L$ is any suborder, then $\mathcal{O}^\times \cong \mu(\mathcal{O}) \times \mathbb{Z}^{r+s-1}$.*

Proof. An algebraic unit u has minimal polynomial $m(x) = x^n + c_{n-1}x^{n-1} \cdots + c_1x \pm 1$ with integer coefficients. Evaluating this polynomial at u , along with rearranging terms, shows that $u(u^{n-1} + c_{n-1}u^{n-2} + \cdots + c_1) = \pm 1$, which proves the first claim. The second claim follows immediately from the first.

By Dirichlet's unit theorem, $\mathcal{O}_L^\times \cong \mu(L) \times \mathbb{Z}^{r+s-1}$. Thus, we only need to check the rank of \mathcal{O}^\times . Because $\mathcal{O} \subseteq \mathcal{O}_L$ is a suborder, the index $N = [\mathcal{O}_L : \mathcal{O}]$ is finite and $N\mathcal{O}_L \subseteq \mathcal{O}$. If $u \in \mathcal{O}_L^\times$ then u reduces modulo N to an element of $(\mathcal{O}_L/N\mathcal{O}_L)^\times$, and $u^t \equiv 1 \pmod{N\mathcal{O}_L}$ for some $1 \leq t \leq \#(\mathcal{O}_L/N\mathcal{O}_L)^\times < N^{[L:\mathbb{Q}]}$. We conclude $u^t \in \mathcal{O}^\times$. Therefore, \mathcal{O}^\times has the same rank as \mathcal{O}_L^\times and we are done. \square

As an application, this shows the structure of the unit group of the ring $R_{m,L}$.

Proposition 2.4. *Let $m \geq 2$ be an integer. If L is a number field with r real embeddings and $2s$ imaginary embeddings, then*

$$R_{m,L}^\times \cong \{\pm 1\} \times \mathbb{Z}^{r+s-1}.$$

In particular, the only roots of unity contained in $R_{m,L}$ are trivial.

Proof. By Theorem 2.3, we only need to show that $R_{m,L}$ has no roots of unity other than ± 1 . If $\zeta \in R_{m,L}^\times$ is a nontrivial root of unity, then L contains the nontrivial cyclotomic subfield $L_0 = \mathbb{Q}(\zeta) \supsetneq \mathbb{Q}$, and thus $R_{m,L_0} = R_{m,L} \cap L_0 \supseteq \mathbb{Z}[\zeta]$ by Lemma 2.2. But $\mathbb{Z}[\zeta]$ is the maximal order of L_0 , while R_{m,L_0} is a non-maximal order, which is a contradiction. \square

2.2. Totally imaginary extensions of totally real fields. We now restrict our attention to the main focus of this paper: totally imaginary quadratic extensions of totally real fields. The following is a generalization of [Was97, Theorem 4.12] to the case of non-maximal orders. Given a field L and a subring $\mathcal{O} \subseteq L$, we write $\mu(\mathcal{O})$ for the set of roots of unity in \mathcal{O} .

Theorem 2.5. *Let K be a totally real field, let L be a totally imaginary quadratic extension, and let $\mathcal{O} \subseteq L$ be an order which is stable under complex conjugation. If $u \in \mathcal{O}^\times$, then its complex conjugate is $\bar{u} = \zeta u$ where $\zeta \in \mu(\mathcal{O})$. In particular, writing $\mathcal{O}_{\text{tr}} = \mathcal{O} \cap K$,*

$$[\mathcal{O}^\times : \mu(\mathcal{O}) \cdot \mathcal{O}_{\text{tr}}^\times] \leq 2.$$

Proof. Notice that every conjugate of u/\bar{u} has absolute value 1 because $\sigma(\bar{u}) = \overline{\sigma(u)}$ for every embedding $\sigma : L \hookrightarrow \mathbb{C}$; see [Was97, p.39]. Therefore, $\frac{u}{\bar{u}}$ is a root of unity [Was97, Lemma 1.6]. In other words, $\frac{u}{\bar{u}} \in \mu(L) \cap \mathcal{O} = \mu(\mathcal{O})$.

Thus, we define a group homomorphism $\varphi : \mathcal{O}^\times \rightarrow \mu(\mathcal{O})/\mu(\mathcal{O})^2$ induced by $u \mapsto u/\bar{u}$ and compute its kernel. First, we check that if $u = \zeta u_1$ for $\zeta \in \mu(\mathcal{O})$ and $u_1 \in \mathcal{O}^\times \cap K$, then $\varphi(u) = u/\bar{u} = \zeta u_1/\overline{\zeta u_1} = \zeta^2$. Conversely, if $u \in \mathcal{O}^\times$ and $\varphi(u) = u/\bar{u} = \zeta^2$, then we rearrange

to see $\bar{\zeta}u = \zeta\bar{u} \in \mathcal{O}_{\text{tr}}$ is totally real. Thus, $\ker \varphi = \mu(\mathcal{O}) \cdot \mathcal{O}_{\text{tr}}^\times$. Since we know the index $[\mu(\mathcal{O}) : \mu(\mathcal{O})^2] \leq 2$, the proof is done. \square

We now wish to apply this theorem to the non-maximal subrings defined in the previous section. In our context, this theorem is helpful because we define subrings with no nontrivial roots of unity. Before we move on, we note that the application of Theorem 2.5 in the case when $\mathcal{O} = \mathcal{O}_L$ is a central ingredient in [MRUV20, Spr20]. The following corollary was also inspired by an analogous fact for maximal orders [Was97, Proof of Proposition 1.5].

Remark 2.6. *It is easy to show, e.g., using Magma [BCP97], that it is possible to have $R_{2,L}^\times \not\subseteq K$ in Corollary 2.7. Indeed, this occurs for the cyclotomic field $L = \mathbb{Q}(\zeta_{15})$. Thus, squaring the group of units is necessary in the statement of the corollary when $m = 2$.*

Corollary 2.7. *Let K be a totally real field with totally imaginary quadratic extension L .*

- (a) $(R_{2,L}^\times)^2 \subseteq R_{2,K}^\times$
- (b) *If $m \geq 3$, then $R_{m,L}^\times = R_{m,K}^\times$.*

Proof. Let $u \in R_{m,L}^\times$. Clearly $R_{m,L}$ is stable under complex conjugation by definition. Combining Proposition 2.4 and Theorem 2.5, either $\bar{u} = u$ or $\bar{u} = -u$. To finish the proof, assume $m \geq 3$ and suppose that $\bar{u} = -u$. Writing $u = mb + j$ for $b \in \mathcal{O}_L$ and $j \in \mathbb{Z}$ we have $\bar{u} = m\bar{b} + j$. Therefore,

$$2u = u - \bar{u} = mb - m\bar{b} \in m\mathcal{O}_L.$$

However, this implies that m divides 2 because u is a unit, and this is impossible because $m \geq 3$. We conclude that $\bar{u} = u$ is totally real. \square

3. DEFINING THE TOTALLY REAL SUBRING.

We now show how to deploy the unit groups described in the previous section. To do this, we explicitly write certain totally real algebraic integers as the sum of units.

Lemma 3.1. *Let $K_0 \subseteq K_1$ be totally real fields. If $a = 2(2d + 1)$ for some $d \in \mathcal{O}_{K_0}$ such that $\sqrt{d^2 + d} \in K_1$, then there is a unit $u \in R_{2,K_1}^\times$ such that $a = u + 1/u$. In particular,*

$$a^2 = u^2 + 1/u^2 + 2.$$

Proof. Define $u_j = 2b_j + 1 \in R_{2,K_1}$ where $b_1 = d + \sqrt{d^2 + d}$ and $b_2 = d - \sqrt{d^2 + d}$. It is easy to compute directly that u_1 and u_2 are inverses of each other, and hence units from R_{2,K_1} :

$$\begin{aligned} u_1 u_2 &= (2b_1 + 1)(2b_2 + 1) = (2d + 1 + 2\sqrt{d^2 + d})(2d + 1 - 2\sqrt{d^2 + d}) \\ &= (2d + 1)^2 - 4(d^2 + d) = 1. \end{aligned}$$

Moreover, $u_1 + u_2 = 2(b_1 + b_2 + 1) = 2(2d + 1) = a$. We take $u = u_1$, in which case $1/u = u_2$. It follows immediately that $a^2 = (u + 1/u)^2 = u^2 + 1/u^2 + 2$. \square

Lemma 3.2. *Let K_0 be a totally real field. If $d \in \mathcal{O}_{K_0}$ and all conjugates of d are outside the open interval $(-2, 0) \subseteq \mathbb{R}$, then*

$$32d = u^2 + 1/u^2 - v^2 - 1/v^2$$

for units $u, v \in R_{2,K_1}^\times$ where $K_1 = K_0(\sqrt{d^2 + d}, \sqrt{(d-1)^2 + (d-1)})$ is totally real.

Proof. It is easy to see that the values of the function $f(x) = x^2 + x$ are negative precisely when $x \in (-1, 0)$. Therefore, because all conjugates of d are outside $(-2, 0)$, the elements $d^2 + d$ and $(d-1)^2 + (d-1)$ are both totally nonnegative, hence their square roots are totally real. Thus, applying Lemma 3.1 twice,

$$32d = 4(2d+1)^2 - 4(2(d-1)+1)^2 = u^2 + 1/u^2 - v^2 - 1/v^2.$$

for some $u, v \in R_{2, K_1}^\times$. □

Now we are ready to define the desired totally real subsets X within the totally imaginary fields. This can be seen as a generalization of [MRUV20, Lemma 7] and [Spr20, Lemma 2.7].

Theorem 3.3. *Let K_0 be a totally real field and let K_1 be the maximal totally real subfield of $K_0^{(2)}$. If $K \supseteq K_1$ is a totally real field, then for every totally imaginary quadratic extension L of K , there is an existentially definable subset $X \subseteq \mathcal{O}_L$ satisfying*

$$\mathcal{O}_{K_0} \subseteq X \subseteq \mathcal{O}_K.$$

Proof. We write

$$X_0 = \{u_1^2 + u_2^2 - u_3^2 - u_4^2 : u_1, \dots, u_4 \in R_{2, L}^\times\}.$$

We have $X_0 \subseteq \mathcal{O}_K$ by Corollary 2.7, and the set

$$X_1 = \{d \in \mathcal{O}_L : 32d \in X_0\}.$$

contains all elements $d \in \mathcal{O}_{K_0}$ whose conjugates are all outside $(-2, 0)$ by Lemma 3.2. We repeat the same trick from before to finish the proof: If $d \in \mathcal{O}_{K_0}$, then $(d-1)^2$ and $(d+1)^2$ are totally nonnegative. Thus $(d-1)^2, (d+1)^2 \in X_1$, and $4d = (d+1)^2 - (d-1)^2$.

In summary, $\mathcal{O}_{K_0} \subseteq X \subseteq \mathcal{O}_K$, where

$$X = \{\alpha \in \mathcal{O}_L : \exists x_1, x_2 \in X_1, 4\alpha = x_1 - x_2\}.$$

Because the ring $R_{2, L}$ is existentially definable in \mathcal{O}_L , this completes the proof. □

By applying the previous theorem in the case where $K_0 = K$, we find that the ring of integers \mathcal{O}_K itself is existentially definable in \mathcal{O}_L , as written in the following theorem. Recall that K is *closed under square roots* if $\alpha \in K$ whenever $\alpha \in \overline{\mathbb{Q}}$ is totally real and $\alpha^2 \in K$.

Theorem 3.4. *If K is a totally real field which is closed under square roots and L is any quadratic imaginary extension of K , then \mathcal{O}_K is existentially definable in \mathcal{O}_L .*

Since \mathbb{Q}^{tr} is closed under square roots, we obtain the definability portion of Theorem 1.1.

Corollary 3.5. *\mathbb{Z}^{tr} is existentially definable in the ring of integers of $\mathbb{Q}^{\text{tr}}(i)$.*

Throughout this section, we have exploited the fact that, within the fields K that we consider, many algebraic integers can be written as the sum of a bounded number of units. For contrast, note that Frey and Jarden showed the following when F is a number field: There is no integer $n \geq 1$ such that every element of \mathcal{O}_F is the sum of at most n units in \mathcal{O}_F^\times [JN07, Theorem 1]. In the same paper, they construct many infinite algebraic extensions of \mathbb{Q} in which every element of the ring of integers is the sum of at most 2 units; see [JN07, Theorem 8]. Although the fields they consider are not used in this paper, their complementary results illustrate how we are exploiting a phenomenon that exists only for (some) infinite algebraic extensions of \mathbb{Q} , and not for number fields.

4. UNDECIDABILITY

The definability results in the preceding section have immediate consequences for decidability. Namely, in the context of Theorem 3.4, if the existential or first-order theory of \mathcal{O}_K is undecidable then the same is true for \mathcal{O}_L . However, this is not the end of the story because the implications for decidability go beyond the hypotheses of the theorem. Indeed, we can deduce a much more general undecidability result with a further development of the ideas used in previous unit group-based methods [MRUV20, Spr20]. Specifically, we only need to define “enough” of the maximal totally real subring to realize the JR-number.

To start, we recall the following lemma, which is C.W. Henson’s extension [VDD88, §3.3] of a result of J. Robinson [Rob62, Theorem 2]; see also the presentation in [MRUV20, Lemma 2.2]. This will be the basis for all of our undecidability results.

Lemma 4.1. *Let \mathcal{O} be a ring of algebraic integers. If there is a family \mathcal{F} of subsets of \mathcal{O} , parametrized by an \mathcal{L}_{ring} -formula, which contains finite sets of arbitrarily large cardinality, then \mathcal{O} has undecidable first-order theory.*

4.1. Undecidability for totally real \mathcal{O}_K . Before considering totally imaginary fields, we briefly sketch the JR-number method for proving the undecidability of the ring of integers of a totally real field; see [Rob62, VV15a, VV15b] for more details.

Definition 4.2. *Given a totally real number $\alpha \in \overline{\mathbb{Q}}$ and $t \in \mathbb{R}$, we write $0 \ll \alpha \ll t$ if every conjugate of α lies in the interval $(0, t)$. For a totally real subset $X \subseteq \overline{\mathbb{Q}}$, write $X_t = \{\alpha \in X : 0 \ll \alpha \ll t\}$. The JR-number of X is*

$$\text{JR}(X) = \inf\{t \in \mathbb{R} : \#X_t = \infty\}.$$

In particular, when K is totally real, J. Robinson [Rob62] observed that if $\text{JR}(\mathcal{O}_K)$ is either ∞ or a minimum, then Lemma 4.1 implies that \mathcal{O}_K is undecidable. Indeed, $\{X_t\}_{t \in \mathbb{Q}}$ can be realized as a parametrized family of existentially definable sets when $X = \mathcal{O}_K$ and $t \in \mathbb{Q}$, as written below. This follows from Siegel’s proof that the totally nonnegative elements of K are precisely the elements which are sums of four squares [Sie21]. Consequently, J. Robinson deduced that the ring of totally real algebraic integers \mathbb{Z}^{tr} is undecidable.

Theorem 4.3. *Let K be a totally real field. If $X = \mathcal{O}_K$ and $t = \frac{a}{b} \in \mathbb{Q}$, then X_t is defined in \mathcal{O}_K by the formula:*

$$\exists y_0, \dots, y_8 \in \mathcal{O}_K [bxy_0^2 \neq 0 \wedge bxy_0^2 \neq a \wedge xy_0^2 = y_1^2 + \dots + y_4^2 \wedge (a - bxy_0^2) = y_5^2 + \dots + y_8^2]$$

Note that, in general, algebraic integers might be written as the sum of squares of algebraic non-integers, so y_0 plays the role of a common denominator.

4.2. Undecidability for totally imaginary \mathcal{O}_L . We are now ready to deploy a modified version of the JR-method for totally imaginary fields; compare with [MRUV20, Spr20].

Theorem 4.4. *Let K_0 be a totally real field for which $\text{JR}(\mathcal{O}_{K_0})$ is either an attained minimum or ∞ , and let K_1 be the maximal totally real subfield of $K_0^{(2)}$. If $K \supseteq K_1$ is any totally real field with $\text{JR}(\mathcal{O}_K) = \text{JR}(\mathcal{O}_{K_0})$, and L is any totally imaginary quadratic extension of K , then the first order theory of \mathcal{O}_L is undecidable.*

Proof. By Theorem 3.3, there is an existentially definable subset $X \subseteq \mathcal{O}_L$ satisfying

$$\mathcal{O}_{K_0} \subseteq X \subseteq \mathcal{O}_K.$$

Therefore, the parametrized formula $\phi_X(x; a, b)$ defined by

$$\exists y_0, \dots, y_8 \in X [bxy_0^2 \neq 0 \wedge bxy_0^2 \neq a \wedge xy_0^2 = y_1^2 + \dots + y_4^2 \wedge (a - bxy_0^2) = y_5^2 + \dots + y_8^2]$$

defines sets which satisfy the containments

$$\left\{ x \in \mathcal{O}_{K_0} : 0 \ll x \ll \frac{a}{b} \right\} \subseteq \{x \in \mathcal{O}_L : \phi_X(x; a, b)\} \subseteq \left\{ x \in \mathcal{O}_K : 0 \ll x \ll \frac{a}{b} \right\}.$$

The containments follow from Theorem 4.3. Because $\text{JR}(\mathcal{O}_K) = \text{JR}(\mathcal{O}_{K_0})$ is either infinite or an attained minimum, the sets on the lefthand and righthand sides above are finite sets of arbitrarily large size as $\frac{a}{b}$ ranges over positive rational numbers approaching $\text{JR}(\mathcal{O}_K)$ from the left. Therefore, by Lemma 4.1, we have proved first-order undecidability. \square

There are two extremes for the JR-number of a ring of totally real algebraic integers, namely 4 and ∞ . For example, the fields $K_0 = \mathbb{Q}$ and $K_1 = \mathbb{Q}(\{\sqrt{n} : n \geq 1\})$ have $\text{JR}(\mathcal{O}_{K_0}) = \text{JR}(\mathcal{O}_{K_1}) = \infty$. Applying Theorem 4.4 in this case leads to a recovery of some of the results that appeared in [MRUV20, Spr20]. At the other extreme, we write the following.

Theorem 4.5. *Let S be any infinite set of positive integers, let $K_0 = \mathbb{Q}(\{\zeta_n + \bar{\zeta}_n : n \in S\})$, and let K_1 be the maximal totally real subfield of $K_0^{(2)}$. If K is any totally real extension of K_1 and L is any totally imaginary quadratic extension of K , then the first order theory of \mathcal{O}_L is undecidable.*

Proof. Under the given hypotheses, $\text{JR}(\mathcal{O}_{K_0}) = 4$ is the smallest possible JR-number and is realized as a minimum. Indeed, we have $0 \ll \zeta_n + \bar{\zeta}_n + 2 \ll 4$ for all $n \in S$, and these are the only totally real elements of $\overline{\mathbb{Q}}$ with this property; see Kronecker [Kro57] and also [Rob64]. Therefore, $\text{JR}(\mathcal{O}_{K_0}) = \text{JR}(\mathcal{O}_K)$ is clear and Theorem 4.4 applies. \square

In recent years, examples of totally real fields K have been discovered for which $\text{JR}(\mathcal{O}_K)$ is neither 4 nor ∞ , i.e., $\text{JR}(\mathcal{O}_K)$ is non-extremal; see [Cas18, CVV20, GR17, VV15a]. A better understanding of the behavior of JR-numbers under field extension would be required before applying Theorem 4.4 to fields K_0 for which $\text{JR}(\mathcal{O}_{K_0}) \in (4, \infty)$ is non-extremal.

REFERENCES

- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [BSF13] Lior Bary-Soroker and Arno Fehm. Open problems in the theory of ample fields. In *Geometric and differential Galois theories*, volume 27 of *Sémin. Congr.*, pages 1–11. Soc. Math. France, Paris, 2013.
- [Cas18] Marianela Castillo Fernández. *On the Julia Robinson number of rings of totally real algebraic integers in some towers of Nested Square Roots*. PhD thesis, Universidad de Concepción, 2018. URL: <http://repositorio.udec.cl/handle/11594/3003>.
- [Cha19] Zoé Chatzidakis. Amalgamation of types in pseudo-algebraically closed fields and applications. *J. Math. Log.*, 19(2):1950006, 28, 2019.
- [CVV20] Marianela Castillo Fernández, Xavier Vidaux, and Carlos R. Videla. Julia Robinson numbers and arithmetical dynamic of quadratic polynomials. *Indiana Univ. Math. J.*, 69(3):873–885, 2020.
- [Dar00] L. Darnière. Decidability and local-global principles. In *Hilbert’s tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999)*, volume 270 of *Contemp. Math.*, pages 145–167. Amer. Math. Soc., Providence, RI, 2000.
- [Den80] J. Denef. Diophantine sets over algebraic integer rings. II. *Trans. Amer. Math. Soc.*, 257(1):227–236, 1980.

- [DF21] Philip Dittmann and Arno Fehm. Nondefinability of rings of integers in most algebraic fields. *Notre Dame J. Form. Log.*, 62(3):589–592, 2021.
- [FHV94] Michael D. Fried, Dan Haran, and Helmut Völklein. Real Hilbertianity and the field of totally real numbers. In *Arithmetic geometry (Tempe, AZ, 1993)*, volume 174 of *Contemp. Math.*, pages 1–34. Amer. Math. Soc., Providence, RI, 1994.
- [FJ74] Gerhard Frey and Moshe Jarden. Approximation theory and the rank of abelian varieties over large algebraic fields. *Proc. London Math. Soc. (3)*, 28:112–128, 1974.
- [FP10] Arno Fehm and Sebastian Petersen. On the rank of abelian varieties over ample fields. *Int. J. Number Theory*, 6(3):579–586, 2010.
- [GR17] Pierre Gillibert and Gabriele Ranieri. Julia robinson’s numbers. *arXiv e-prints*, Oct 2017.
- [Jar11] Moshe Jarden. *Algebraic patching*. Springer Monographs in Mathematics. Springer, Heidelberg, 2011.
- [JN07] Moshe Jarden and Władysław Narkiewicz. On sums of units. *Monatsh. Math.*, 150(4):327–332, 2007.
- [Kob06] Emi Kobayashi. A remark on the Mordell-Weil rank of elliptic curves over the maximal abelian extension of the rational number field. *Tokyo J. Math.*, 29(2):295–300, 2006.
- [Koe14] Jochen Koenigsmann. Undecidability in number theory. In *Model theory in algebra, analysis and arithmetic*, volume 2111 of *Lecture Notes in Math.*, pages 159–195. Springer, Heidelberg, 2014.
- [Kro57] L. Kronecker. Zwei Sätze über Gleichungen mit ganzzahligen Coefficienten. *J. Reine Angew. Math.*, 53:173–175, 1857.
- [LR08] Álvaro Lozano-Robledo. Ranks of abelian varieties over infinite extensions of the rationals. *Manuscripta Math.*, 126(3):393–407, 2008.
- [MR18] Barry Mazur and Karl Rubin. Diophantine stability. *Amer. J. Math.*, 140(3):571–616, 2018. With an appendix by Michael Larsen.
- [MRS22] Barry Mazur, Karl Rubin, and Alexandra Shlapentokh. Existential definability and diophantine stability. *arXiv e-prints*, page arXiv:2208.09963, August 2022.
- [MRUV20] Carlos Martínez-Ranero, Javier Utreras, and Carlos R. Videla. Undecidability of $\mathbb{Q}^{(2)}$. *Proc. Amer. Math. Soc.*, 148(3):961–964, 2020.
- [Pet06] Sebastian Petersen. On a question of Frey and Jarden about the rank of abelian varieties. *J. Number Theory*, 120(2):287–302, 2006.
- [Poo02] Bjorn Poonen. Using elliptic curves of rank one towards the undecidability of Hilbert’s tenth problem over rings of algebraic integers. In *Algorithmic number theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 33–42. Springer, Berlin, 2002.
- [Pop96] Florian Pop. Embedding problems over large fields. *Ann. of Math. (2)*, 144(1):1–34, 1996.
- [Rob62] Julia Robinson. On the decision problem for algebraic rings. In *Studies in mathematical analysis and related topics*, pages 297–304. Stanford Univ. Press, Stanford, Calif, 1962.
- [Rob64] Raphael M. Robinson. Intervals containing infinitely many sets of conjugate algebraic units. *Ann. of Math. (2)*, 80:411–428, 1964.
- [Shl09] Alexandra Shlapentokh. Rings of algebraic numbers in infinite extensions of \mathbb{Q} and elliptic curves retaining their rank. *Arch. Math. Logic*, 48(1):77–114, 2009.
- [Shl18] Alexandra Shlapentokh. First-order decidability and definability of integers in infinite algebraic extensions of the rational numbers. *Israel Journal of Mathematics*, 226(2):579–633, 2018.
- [Sie21] Carl Siegel. Darstellung total positiver Zahlen durch Quadrate. *Math. Z.*, 11(3-4):246–275, 1921.
- [Spr20] Caleb Springer. Undecidability, unit groups, and some totally imaginary infinite extensions of \mathbb{Q} . *Proc. Amer. Math. Soc.*, 148(11):4705–4715, 2020.
- [VDD88] Lou Van Den Dries. Elimination theory for the ring of algebraic integers. *Journal für die Reine und Angewandte Mathematik*, 1988(388):189–205, 1988.
- [Vid00a] Carlos R. Videla. Definability of the ring of integers in pro-p galois extensions of number fields. *Israel Journal of Mathematics*, 118(1):1–14, Dec 2000.
- [Vid00b] Carlos R. Videla. The undecidability of cyclotomic towers. *Proc. Amer. Math. Soc.*, 128(12):3671–3674, 2000.

- [VV15a] Xavier Vidaux and Carlos R. Videla. Definability of the natural numbers in totally real towers of nested square roots. *Proceedings of the American Mathematical Society*, 143(10):4463–4477, 2015.
- [VV15b] Xavier Vidaux and Carlos R. Videla. A note on the northcott property and undecidability. *Bulletin of the London Mathematical Society*, 48(1):58–62, 2015.
- [Was97] Lawrence C. Washington. *Introduction to cyclotomic fields*, volume 83. Springer, New York, 2nd edition, 1997.

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, UK,
AND THE HEILBRONN INSTITUTE FOR MATHEMATICAL RESEARCH, BRISTOL, UK
Email address: `c.springer@ucl.ac.uk`