

HYPERBOLIC FIBERED SLICE KNOTS WITH RIGHT-VEERING MONODROMY

DONGTAI HE

ABSTRACT. We construct a hyperbolic fibered slice knot with right-veering monodromy, which disproves a conjecture posed in [HKK⁺21].

1. INTRODUCTION

We give a negative answer to Question 8.2 posed in [HKK⁺21].

Question 1.1. If K is a hyperbolic fibered slice knot, does the fractional Dehn twist coefficient (FDTC) of the monodromy vanish?

We construct a hyperbolic slice fibered knot K' with positive FDTC.

1.1. Motivation. The authors [HKK⁺21] observe that many low-crossing slice fibered knots have zero FDTC. Any $(p, 1)$ -cable of slice fibered knot is still slice fibered, whereas FDTC of $(p, 1)$ -cable equals to $1/p$. The authors therefore ask the above question about hyperbolic fibered slice knots.

Baldwin, Ni and Sivek [BNS22, Corollary 1.7] prove the following related proposition in terms of the τ -invariant in Heegaard Floer homology:

Proposition 1.2. If $K \subset S^3$ is a fibered knot with thin knot Floer homology such that $\tau(K) < g(K)$, then FDTC vanishes.

The τ -invariant vanishes for slice knots. Proposition 1.2 explains the case for low-crossing fibered slice knots because many of those are thin.

We have an immediate corollary:

Corollary 1.3. The knot Floer homology of K' is not thin.

1.2. Organization. We follow the recipe of Kazez and Roberts [KR13] to construct hyperbolic fibered knots with positive FDTC. The search for ribbon knot is inspired by the work of Hitt and Silver [HS91]. In section 2 we review Nielson-Thurston classification of surface automorphism and examples from Kazez and Roberts. We construct our example K' in section 3.

2. MONODROMIES OF FIBERED KNOTS IN S^3

2.1. Surface automorphism. We first recall Nielson-Thurston classification of surface automorphisms:

Theorem 2.1. [CCB88, Thu88] Let S be an oriented hyperbolic surface with geodesic boundary, and let $h \in \text{Aut}(S, \partial S)$. Then h is freely isotopic to either

- (1) a pseudo-Anosov homeomorphism ϕ that preserves a pair of geodesic laminations λ^s and λ^u ,
- (2) a periodic homeomorphism ϕ such that $\phi^n = id$ for some n ,
- (3) a reducible homeomorphism h' that preserves a maximal collection of simple closed geodesic curves in S . To avoid overlap, we consider h reducible only when it is not periodic.

In particular, we only regard h as reducible only if it is not periodic to avoid overlap. Let $\Phi : S \times [0, 1] \rightarrow S$ be an isotopy from h to its Thurston representative ϕ . Considering the restriction of Φ to the boundary ∂S , we have a homeomorphism:

$$\Phi_{\partial} : \partial S \times [0, 1] \rightarrow \partial S \times [0, 1]$$

defined by $\Phi_{\partial}(x, t) = (\Phi_t(x), t)$. The fractional Dehn twist coefficient $c(h)$ can be defined as the winding number of the arc $\Phi_{\partial}(\theta \times [0, 1])$. Nielson-Thurston classification guarantees that $c(h) \in \mathbb{Q}$.

Thurston proved that a fibered knot is hyperbolic if and only if its monodromy is pseudo-Anosov. Fractional Dehn twist coefficient is closely related to the following notion of right-veeringness.

Definition 2.2. [HKM07] A homeomorphism $h \in \text{Aut}(S, \partial S)$ is called right-veering if for every based point $x \in \partial S$ and every properly embedded arc α starting at x , $h(\alpha)$ is to the right of α , after isotoping $h(\alpha)$ so that it intersects α minimally. Similarly, h is called left-veering if $h(\alpha)$ is to the left of α .

Proposition 2.3. [HKM07] h is right-veering if and only if $c(h) > 0$ for every component of ∂S , and h is left-veering if and only if $c(h) < 0$ for every component of ∂S .

If $c(h) = 0$, one can find two arcs such that one is moved by h to the right and the other to the left. The significance of right-veeringness is highlighted by the following theorem of Honda, Kazez and Matić:

Theorem 2.4. [HKM09] Every open book that is compatible with a tight contact structure is right-veering.

A large source of examples of reducible right-veering homeomorphism comes from the class of fibered cable knots. Indeed, if h is the monodromy of a fibered (p, q) -cable knot $K_{p,q}$ with Seifert surface S , then $c(h) = 1/pq$ and h is reducible. Let $\{C_i\}$ be the collection of curves preserved by h' . $\{C_i\}$ partitions S into subsurfaces $\{S_j\}$ permuted by h' . Let S_0 be the subsurface containing $\partial S = K_{p,q}$, then $h'|_{S_0}$ is periodic. Kazez and Roberts characterize the monodromy h of a fibered knot K in S^3 in the following theorem:

Theorem 2.5. [KR13]

- (1) If h is periodic, then K is the unknot or a (p, q) -torus knot.
- (2) If h has a reducible Thurston representative h' with periodic $h'|_{S_0}$, then K is a (p, q) -cable knot, and $c(h) = 1/pq$.
- (3) [Gab97] If h is either pseudo-Anosov or reducible with $h'|_{S_0}$ pseudo-Anosov. Then either $c(h) = 0$ or $c(h) = 1/r$, where $2 \leq |r| \leq 4g(K) - 2$.

Corollary 2.6. $c(h) = 0$ or $1/r$ for some integer r , $|r| \geq 2$. In particular, $|c(h)| \leq 1/2$.

In particular, the $(2, 1)$ -cable of a fibered knot in S^3 has its monodromy attaining maximum FDTC. We review hyperbolic case in the next section.

2.2. **Stallings’ twist and (2,1)-cable.** Let U be an unknot properly embedded in a surface F . We say U is untwisted relative to F if U bounds a disk transverse to F along U . A Stallings’ twist [Sta78] is a surgery along such an untwisted U . Kazez and Roberts apply Stallings’ twist on $(2, 1)$ –cables to produce hyperbolic fibered knots with maximum FDTC = $1/2$.

Let (S, h) be an open book decomposition of S^3 with connected binding K , where h is pseudo-Anosov and $c(h) = 0$. Let $K_{2,1}$ be the $(2, 1)$ –cable of K . The fibered surface Σ of $K_{2,1}$ can be viewed as the union of two copies S_0, S_1 of S connected by a 1-handle. Let H be the monodromy of this new open book.

We choose a simple closed curve C in Σ such that $C_0 = C \cap S_0$ and $C_1 = C \cap S_1$ are two essential arcs. Moreover, we require C_i to be nonseparating in S_i . Let T_C be the right-handed Dehn twist along C and $H' = T_C \circ H$.

Theorem 2.7. [KR13] H' is pseudo-Anosov and $c(H') = 1/2$.

3. RIBBON FIBERED KNOT

We are ready to construct a hyperbolic ribbon fibered knot with positive FDTC. Let K be the knot 10_{153} from Rolfsen’s knot table. K is a hyperbolic ribbon fibered knot with 3-genus 3. Figure 1 is a ribbon diagram for 10_{153} .




FIGURE 1. A ribbon diagram for the knot $K = 10_{153}$

Let h denote the monodromy. According to [CL], h can be presented as described in Figure 2. One can see that h is neither right-veering nor left-veering by choosing different endpoints of γ . Therefore, $c(h) = 0$.




FIGURE 2. Monodromy of the fibered knot 10_{153} . h can be presented as the word **abcBEGhcd**, where x denotes a right-handed Dehn twist about x and X denotes a left-handed Dehn twist about x . A word is read from right to left so that aB means perform a left-handed Dehn twist about b then perform a right-handed Dehn twist about a .

A Seifert surface S of K can be obtained by Seifert's algorithm as explained in Figure 3. The genus of F is 3 so that F is the fibered surface.




FIGURE 3. The surface obtained by Seifert's algorithm has genus 3. c is a non-separating properly embedded arc on the surface.

Let $K_{2,1}$ be the $(2, 1)$ -cable of K (Figure 4). The twisted band connecting the two copies of K is added at p . c is a nonseparating properly embedded arc on the fibered surface of K . $K_{2,1}$ is also fibered whose fibered surface Σ can be obtained by connecting two copies of S

with the same twisted band at p . Then define a simple closed curve C to be a band sum of the two copies of c along an arc running across the twisted band.




FIGURE 4. $(2, 1)$ -cable of K .

Let T_C denote the right-handed Dehn twist along C , and denote the resulting fibered knot K' . By [KR13, Corollary 4.6], the monodromy $T_C \circ H$ is pseudo-Anosov and right-veering with $c(T_C \circ H) = \frac{1}{2}$.

Recall that $K = 10_{153}$ is a ribbon knot, so is $K_{2,1}$. C is an unknotted untwisted curve. Performing a right-handed Dehn twist along C has the same effect on $(S^3, K_{2,1})$ applying a (-1) -surgery along C . The resulting manifold is still S^3 and we have a new knot K' . C winds around two copies of a ribbon band (Figure 5).




FIGURE 5. The curve C winds around two copies of a ribbon band. This figure shows one of the ribbon. The other ribbon is on the other copy from the $(2, 1)$ -cable

Figure 6 illustrates the effect of (-1) -surgery along C to the ribbon bands. The resulting knot K' is still a ribbon knot.

Theorem 3.1. K' is a hyperbolic ribbon fibered knot with $\text{FDTC} = 1/2$. Hence, the monodromy is right-veering.




FIGURE 6. The effect of (-1) - surgery along C after isotopy.

REFERENCES

- [BNS22] John A Baldwin, Yi Ni, and Steven Sivek. Floer homology and right-veering monodromy, 2022. preprint, arXiv:2204.04093.
- [CCB88] Andrew J Casson, Andrew J Casson, and Steven A Bleiler. *Automorphisms of surfaces after Nielsen and Thurston*. Number 9. Cambridge University Press, 1988.
- [CL] Jae Choon Cha and Charles Livingston. Knotinfo: Table of knot invariants. <http://www.indiana.edu/~knotinfo>.
- [Gab97] David Gabai. Problems in foliations and laminations. *Stud. in Adv. Math. AMS/IP*, 2:1–34, 1997.
- [HKK⁺21] Diana Hubbard, Keiko Kawamuro, Feride Ceren Kose, Gage Martin, Olga Plamenevskaya, Katherine Raoux, Linh Truong, and Hannah Turner. Braids, fibered knots, and concordance questions. In *Research Directions in Symplectic and Contact Geometry and Topology*, pages 293–324. Springer, 2021.
- [HKM07] Ko Honda, William H. Kazez, and Gordana Matić. Right-veering diffeomorphisms of compact surfaces with boundary. *Invent. Math.*, 169(2):427–449, 2007.
- [HKM09] Ko Honda, William H. Kazez, and Gordana Matić. On the contact class in Heegaard Floer homology. *J. Differential Geom.*, 83(2):289–311, 2009.
- [HS91] L Richard Hitt and Daniel S Silver. Ribbon knot families via Stallings’ twists. *Journal of the Australian Mathematical Society*, 50(3):356–372, 1991.
- [KR13] William H. Kazez and Rachel Roberts. Fractional Dehn twists in knot theory and contact topology. *Algebr. Geom. Topol.*, 13(6):3603–3637, 2013.
- [Sta78] John R. Stallings. Constructions of fibered knots and links. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2*, Proc. Sympos. Pure Math., XXXII, pages 55–60. Amer. Math. Soc., Providence, R.I., 1978.
- [Thu88] William P Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bulletin of the American mathematical society*, 19(2):417–431, 1988.